

FKN, first proof, rewritten

Ehud Friedgut and Gil Kalai and Assaf Naor

Abstract

About twenty years ago we wrote a paper, "Boolean Functions whose Fourier Transform is Concentrated on the First Two Levels", [1]. In it we offered several proofs of the statement that Boolean functions $f(x_1, x_2, \dots, x_n)$, whose Fourier coefficients are concentrated on the lowest two levels are close to a constant function or to a function of the form $f = x_k$ or $f = 1 - x_k$.

Returning to the paper lately, we noticed that the presentation of the first proof is rather cumbersome, and includes several typos. In this note we rewrite that proof, as a service to the public.

Here is the main theorem of FKN [1].

Theorem 0.1. *Let $f : \{0, 1\}^n \rightarrow \{0, 1\}$, $\|f\|_2^2 = p$. If $\sum_{|S|>1} \widehat{f}^2(S) = \delta$ then either $p < K'\delta$ or $p > 1 - K'\delta$ or $\|f(x_1, x_2, \dots, x_n) - x_i\|_2^2 \leq K\delta$ for some i or $\|f(x_1, x_2, \dots, x_n) - (1 - x_i)\|_2^2 \leq K\delta$ for some i . Here, K' and K are absolute constants.*

Proof: First observe that we may assume that $p = 1/2$: replace f by a function $g : \{0, 1\}^{n+1} \rightarrow \{0, 1\}$ defined by

$$g(x_1, \dots, x_n, x_{n+1}) = f(x_1, \dots, x_n) \cdot \frac{1 + \chi_{n+1}}{2} + (1 - f(1 - x_1, \dots, 1 - x_n)) \cdot \frac{1 - \chi_{n+1}}{2}$$

and note that $|g|_2^2 = 1/2$, and $\sum_{|S|>1} \widehat{f}^2(S) = \sum_{|S|>1} \widehat{g}^2(S)$. Then prove that g is close to a dictator (or anti-dictator). If the influential coordinate for g is $n + 1$ then f is necessarily close to a constant, and if the influential coordinate for g is some other i , then f too is necessarily close to a dictator or anti-dictator of the i th variable.

Let

$$f^{=1} := \sum \widehat{f}(i) \chi_i.$$

Note that $f^{=1}$ is an odd function, i.e. $f^{=1}(S) = -f^{=1}(S^c)$, and that

$$|f^{=1}|_2^2 = \sum \widehat{f}(i)^2 = 1/4 - \delta.$$

Note also that

$$\begin{aligned}
(1/4 - \delta) &= \langle f^{\neq 1}, f^{\neq 1} \rangle = \sum \widehat{f}(i)^2 = \langle f, f^{\neq 1} \rangle \\
&= \sum_{S \subset [n]} 2^{-n} f(S) f^{\neq 1}(S) = \sum_{S \subset [n]} 2^{-n} f(S) \frac{f^{\neq 1}(S) - f^{\neq 1}(S^c)}{2} = \\
&\sum_{S \subset [n]} 2^{-n} f^{\neq 1}(S) \frac{f(S) - f(S^c)}{2} \leq \sum_{S \subset [n]} 2^{-n} \frac{1}{2} |f^{\neq 1}(S)| = \frac{1}{2} |f^{\neq 1}|_1.
\end{aligned}$$

So

$$\frac{1}{2} - 2\delta \leq |f^{\neq 1}|_1, |f^{\neq 1}|_2 = \sqrt{1/4 - \delta} \leq \frac{1}{2} - \delta.$$

Normalizing, let $h = \frac{f^{\neq 1}}{|f^{\neq 1}|_2}$, then

$$|h|_2 = 1, |h|_1 \geq \frac{1/2 - 2\delta}{1/2 - \delta} \geq 1 - 4\delta$$

So $h = \sum a_i \chi_i$, with $\sum a_i^2 = 1$, and we wish to prove that one of the a_i 's is close to 1 or -1. This follows immediately from the following result of König, Schütt and Tomczak-Jaegermann, [2].

Theorem 0.2. *Let a_1, a_2, \dots, a_n be real numbers such that $\sum a_i^2 = 1$. Let E be the expected value of $|\sum a_i \chi_i|$. Then*

$$\left| E - \sqrt{\frac{2}{\pi}} \right| \leq \left(1 - \sqrt{\frac{2}{\pi}} \right) \max_{1 \leq i \leq n} |a_i|. \quad (0.1)$$

In our case this yields

$$\max\{|a_i|\} \geq \frac{1 - \sqrt{\frac{2}{\pi}} - 4\delta}{1 - \sqrt{\frac{2}{\pi}}} > 1 - 20\delta.$$

References

- [1] E. Friedgut, G. Kalai, A. Naor, Boolean functions whose Fourier transform is concentrated on the first two levels, *Advances in Applied Mathematics* 29 (2002), pp. 427–437.
- [2] König, H., C. Schütt, N. Tomczak-Jaegermann, Projection constants of symmetric spaces and variants of Khintchine's inequality. *J. Reine Angew. Math.* 511 (1999), 1–42.