

EXISTENCE OF PHYSICAL MEASURES IN SOME EXCITATION-INHIBITION NETWORKS

MATTEO TANZI AND LAI-SANG YOUNG

ABSTRACT. In this paper we present a rigorous analysis of a class of coupled dynamical systems in which two distinct types of components, one excitatory and the other inhibitory, interact with one another. These network models are finite in size but can be arbitrarily large. They are inspired by real biological networks, and possess features that are idealizations of those in biological systems. Individual components of the network are represented by simple, much studied dynamical systems. Complex dynamical patterns on the network level emerge as a result of the coupling among its constituent subsystems. Appealing to existing techniques in (nonuniform) hyperbolic theory, we study their Lyapunov exponents and entropy, and prove that large time network dynamics are governed by physical measures with the SRB property.

1. INTRODUCTION

As dynamical systems come in too many flavors to be classified or even systematically described, when studying the subject one usually learns from paradigms. In the mathematical theory of chaotic systems, a great deal of intuition has been derived from classical examples such as expanding circle maps [KS69], geodesic flows on manifolds of negative curvature [Hop39, Ano67], hyperbolic billiards [Sin70, Bun74] [BSC90, CM06], the Lorenz attractor [Lor62, GW79], logistic maps [Jak81, CE80, BC85], Hénon attractors and generalizations [BC91, YB93, WY08], and so on. Our notion of what a chaotic dynamical system looks like has been intimately tied to these examples. Textbook pictures of chaotic behavior are those with identifiable expanding and contracting directions in their phase spaces, and where most pairs of nearby orbits can be seen to diverge quickly, at exponential rates.

Chaotic behavior in the sense of hyperbolicity in high dimensional systems can have a different appearance, however. Independent of dimension, the positivity of Lyapunov exponents must, by definition, entail the same local geometry of invariant subspaces and invariant manifolds, but visualizing this picture for a system with high dimensional phase space can be challenging. For high dimensional systems, it may be profitable to turn to other more salient characteristics. For example, in a network in which a large number of simpler constituent dynamical systems are coupled together, dynamics of the constituent subsystems may be more readily observable. Properties such as degree of synchronization, the coherence or incoherence of dynamical behavior among constituent subsystems, the number of distinguishable states and the predictability of dynamical patterns — all are known to be of consequence in real-world examples, see e.g. [BB66, TW82, Kur03, BTLT06]. But how are these behaviors related to hyperbolicity? What are the signatures of chaotic

MT acknowledges funding from the H2020 Marie Skłodowska-Curie Actions, project: “Ergodic Theory of Complex Systems” (project no. 843880).

LSY was supported in part by NSF Grant DMS 1901009.

behaviors in networks? How is the number of positive Lyapunov exponents reflected? And do concepts like SRB measures and physical measures continue to make sense? These and other questions are part of a broad swath of open territory in the theory of high dimensional dynamical systems that remains to be explored, and are most likely going to be (at least partially) answered through the introduction of new paradigms for high-dimensional systems.

In the belief that examples will contribute to our understanding of hyperbolicity in high dimensional systems, we present in this paper a new class of models — new in the sense that their ergodic theory has not been studied — and demonstrate how they can be analyzed using existing tools. These models are examples of Excitation-Inhibition networks. They consist of arbitrary numbers of interacting components some of which excite and others suppress. The interaction is bi-directional and the model is more general than a skew-product: an excitatory subsystem excites inhibitory components which when activated send feedback inhibition to the excitatory subsystem. These examples are inspired by biology, but we do not pretend that they are depictions of any specific biological system. In order to make connections with hyperbolic theory, it was necessary to make the models analyzable, and to do that, we have had to introduce a fair amount of idealization.

Though far from the first such examples to be studied, we mention two ways in which our models differ from many previous works. The first is that they include interactions of components that are not identical. While homogeneous materials in physics (e.g. [SL77]) have inspired the study of systems in which identical maps are coupled (for a random sample see [BS88, Kan93, BK96, KY07, KL09, SB16] and the articles in [CF05]), examples from biological networks typically involve different substances or agents with distinct characteristics and functions (see [BMBY15, FT14, MS08, WM15, YFB⁺12, LKYJW20]; see also [PvST20]). The network models studied here are an addition to the relatively small collection of examples of the latter kind. A second characteristic of our example networks is that their hyperbolicity is not apparent; that is, expanding and contracting directions are not obvious to the eye, unlike, e.g., weakly coupled lattices of expanding circle maps, as in e.g. [BS88, KL06] or even more strongly coupled networks, as in e.g. [KY10]. It is our hope that examples in which hyperbolic properties are emergent may offer greater insight into how chaotic (or nonchaotic) behaviors would manifest in these large dynamical systems.

Finally, though the mathematical models studied here are not intended to be models of realistic physical or biological systems, they are inspired by real biological phenomena. We have incorporated into our models features such as the competition and balancing of excitatory and inhibitory forces, the activation of certain processes upon threshold crossing, and relaxation to equilibria in the absence of inputs. This paper is a small step in the direction of promoting biology-inspired models, which are, in our view, underrepresented among the current pool of examples in dynamical systems. More generally, we wish to demonstrate that dynamical systems theory has the language and the tools to shed light on natural phenomena, to offer insight on a conceptual level even when analysis of the system exactly as defined is out of reach.




FIGURE 1. Network architecture: An Excitatory (E) environment interacting with an arbitrary number of Inhibitory (I) components. When activated, the I-components slow down the dynamics of the E-environment.

2. MODEL DESCRIPTION AND MAIN RESULTS

Sect 2.1 contains an informal overview of the model; details are given in Sect 2.2. Our main results are stated in Sect 3 below.

2.1. Overview. The network studied in this paper is made up of arbitrary numbers of *excitatory* and *inhibitory* components. Simplifying, we group the excitatory components together to form a single “excitatory environment” (E-environment). Inhibitory components (I-components) are modeled individually.

When decoupled from the rest of the network, the E-environment is modeled as an Anosov flow that we take as exemplification of chaotic behavior in continuous time. A global cross-section σ_0 to this flow is fixed. Excitatory signals are sent to I-components each time the trajectory returns to σ_0 .

When decoupled from the E-environment, I-components are modeled as North-South flows on the unit circle, i.e., there are two fixed points, to be thought of as N and S-poles; all nonstationary trajectories go from N-pole to S-pole. Each excitatory signal produces a fairly abrupt rotation of the circle. For each I-component, the amount rotated depends on the location of σ_0 visited by the Anosov trajectory; consecutive visits produce what resembles a sequence of random rotations. Different I-components are affected differently at each return to σ_0 .

Finally, there is feedback from the I-components to the E-environment. For each I-component, there is a designated region of its phase space with the property that while there, the component has a suppressive effect on the E-environment, modeled as a reduction in speed for the Anosov flow. The I-components do not interact with one another directly, but only through the E-environment: collectively they determine when the next excitatory signal will be.

This completes our model description in words. A precise description is given below. As we will show, our model is a piecewise smooth flow on a $(3 + N)$ -dimensional phase space where $N \in \mathbb{Z}^+$ is the number of I-components. Our main result is that the dynamics of networks of this type can be described by natural invariant measures that are physical measures with the SRB property; see [ER85]. Numerical illustrations are presented in Sect. 3.2.

2.2. Precise model description. For $k \in \mathbb{Z}^+$, let $\mathbb{T}^k := \mathbb{R}^k / \mathbb{Z}^k$ denote the k -dimensional torus, which we also identify with $\mathbb{T} \times \cdots \times \mathbb{T}$, the k -fold product of $\mathbb{T} = \mathbb{R}/\mathbb{Z}$. Euclidean norms on \mathbb{T}^k, \mathbb{R} and products thereof are denoted by $|\cdot|$. Below

we present the full model in four steps (A-D), describing first the dynamics of the E-environment and I-components in isolation, i.e. when each is decoupled from the rest of the system, before describing how they interact.

A. *Excitatory environment in isolation.* Consider a flow f^t that is a time-1-suspension of a linear hyperbolic toral automorphism $A : \mathbb{T}^2 \curvearrowright$.

More precisely, let $A : \mathbb{T}^2 \curvearrowright$ be an Anosov diffeomorphism, which for simplicity we assume to be linear, i.e. there is a constant splitting $T\mathbb{T}^2 = E_A^u \oplus E_A^s$ such that for all $p \in \mathbb{T}^2$,

$$DA_p E_A^u = E_A^u, \quad DA_p E_A^s = E_A^s;$$

and there exists $\lambda > 0$ such that

$$|DA_p v| = e^\lambda |v| \quad \forall v \in E_A^u, \quad |DA_p v| = e^{-\lambda} |v| \quad \forall v \in E_A^s.$$

Letting (x, y, w) denote the coordinates in $\mathbb{T}^2 \times [0, 1]$, the flow f^t is defined on the set $M_f := (\mathbb{T}^2 \times [0, 1]) / \sim$ where $(x, y, 1)$ is identified with $(A(x, y), 0)$. To introduce a Riemannian metric on M_f , we assign to the tangent space at $(x, y, w) \in \mathbb{T}^2 \times [0, 1]$ an inner product with respect to which E_A^u, E_A^s and the w -axis are orthogonal and the resulting norm $\|\cdot\|$ has $\|\cdot\| = |\cdot|$ in the w -direction and

$$\|v\| = e^{\lambda w} |v| \quad \text{for } v \in E_A^u, \quad \|v\| = e^{-\lambda w} |v| \quad \text{for } v \in E_A^s.$$

It is easy to see that

- (i) M_f equipped with $\|\cdot\|$ is a C^∞ Riemannian manifold;
- (ii) if f^t is the flow generated by the vector field ∂_w , the positive unit vector in the w -direction, then f^t is a C^∞ Anosov flow for which $\|Df^t(v)\| = e^{\lambda t} \|v\|$ for $v \in E_A^u$ and $\|Df^t(v)\| = e^{-\lambda t} \|v\|$ for $v \in E_A^s$ everywhere on M_f .
- (iii) Finally, $\sigma_0 := \mathbb{T}^2 \times \{0\}$ is a cross-section to the flow with first return map $= A$ and return time $\equiv 1$.

The E-environment of our model when operating in isolation is given by the flow f^t on $(M_f, \|\cdot\|)$.

B. *Inhibitory components when decoupled from the E-environment.* Fix arbitrary $N \in \mathbb{Z}^+$. For $i = 1, 2, \dots, N$, we let g_i^t be the flow on \mathbb{T} generated by a C^2 vector field v_{g_i} with the following properties:

- (i) $v_{g_i}(0) = v_{g_i}(\frac{1}{2}) = 0$, and $v_{g_i}(z) \neq 0$ for $z \neq 0, \frac{1}{2}$;
- (ii) $v'_{g_i}(0) = \lambda_i^-$ and $v'_{g_i}(\frac{1}{2}) = \lambda_i^+$ for some $\lambda_i^- < 0 < \lambda_i^+$.

It follows from (i) and (ii) above that g_i^t is a North-South (N-S) flow, with S-pole at $z_i = 0$ and N-pole at $z_i = \frac{1}{2}$. All other trajectories go from N-pole to S-pole.

The **phase space** of the full model is $M := M_f \times \mathbb{T}^N$.

C. *Action of E-environment on I-components without feedback.* For each $i \in \{1, \dots, N\}$, we let $r_i : \mathbb{T} \rightarrow \mathbb{T}$ be a C^2 order-preserving local diffeomorphism of degree ≥ 1 and, identifying \mathbb{T} with $[0, 1)$, we let $\hat{r}_i : [0, 1) \rightarrow \mathbb{R}$ be the lift of r_i with $\hat{r}_i(0) \in [0, 1)$, i.e., \hat{r}_i is the continuous function with the property that for all $x \in [0, 1)$, $\pi \hat{r}_i(x) = r_i(x)$ where $\pi : \mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z}$ is the usual projection. Fix $b \in (0, 1)$. Consider the vector field v_F on $M_f \times \mathbb{T}^N$ given by

$$v_F(x, y, w; z_1, \dots, z_N) = \begin{cases} (\partial_w; b^{-1} \hat{r}_1(x), \dots, b^{-1} \hat{r}_N(x)) & \text{for } w \in [0, b) \\ (\partial_w; v_{g_1}, \dots, v_{g_N}) & \text{for } w \in [b, 1) \end{cases}$$

Let F^t be the flow defined by v_F . Then F^t is a skew-product with the Anosov flow f^t in the base driving the dynamics on N circle fibers with coordinates z_1, \dots, z_N . For $w \in [0, b)$, the action on each z_i -fiber is a rotation, the total amount rotated at the end of b units of time being $\hat{r}_i(x)$, while on the rest of M_f , fiber dynamics are N-S flows independent of the base.

Observe that v_F is discontinuous at the two cross-sections

$$\Sigma_0 := \{w = 0\} \quad \text{and} \quad \Sigma_b := \{w = b\}$$

to the flow F^t . These discontinuities pose no issues in the definition of the flow: a flowline starting from Σ_0 simply follows one vector field until it reaches Σ_b , where it switches to the other vector field until it returns to Σ_0 . On $\{w \in (0, b)\}$, v_F is also discontinuous at $\{x = 0\}$, though starting from Σ_0 , the time- b -map F^b is well defined and is a C^2 diffeomorphism from Σ_0 to Σ_b because r_i is a local diffeomorphism.

D. *The full model: E-I interaction with feedback.* We have described above an E-to-I action that takes place on $\{w \in [0, b)\}$. For simplicity we will assume that the feedback from I to E takes place only on $\{w \in (b, 1]\}$. To define the latter, we modify the vector field ∂_w in a z_i -dependent way: Fix $\Phi : \{0, 1, \dots, N\} \rightarrow \mathbb{R}$ an increasing function such that $\Phi(0) = 0$ and $\Phi < 1$. Calling $\chi_{(1/2, 1)}$ the characteristic function of the arc $(1/2, 1) \subset \mathbb{T} = \mathbb{R}/\mathbb{Z}$, we define

$$(1) \quad \tilde{v}_f = \tilde{v}_{f; z_1, \dots, z_N} = \left[1 - \Phi \left(\sum_{i=1}^N \chi_{(1/2, 1)}(z_i) \right) \right] \partial_w := c_\Phi(z_1, \dots, z_N) \partial_w .$$

That is to say, the presence of each z_i in the region $(1/2, 1)$ of \mathbb{T} causes the Anosov flow to reduce its speed by some amount determined by the function Φ . For example one could take $\Phi(n) = \eta n$ with $\eta \in [0, 1/N)$, where the inhibitory effect of multiple z_i being in $(1/2, 1)$ is additive.

The vector field for the full model is then given by

$$v_{\mathbf{F}} = \begin{cases} v_F & \text{for } w \in [0, b) \\ (\tilde{v}_f; v_{g_1}, \dots, v_{g_N}) & \text{for } w \in [b, 1] . \end{cases}$$

The flow associated with the vector field $v_{\mathbf{F}}$ is denoted \mathbf{F}^t .

Observe that on $\{w \in [b, 1]\}$, the vector field $v_{\mathbf{F}}$ is discontinuous at $\cup_{i=1, \dots, N} \{z_i = 0, \frac{1}{2}\}$, the discontinuity set of the function c_Φ in (1). This set is the union of a finite number of codimension one surfaces. Because every g_i^t leaves the interval $(\frac{1}{2}, 1)$ invariant, flowlines do not cross from one side of these surfaces to the other.

While F^t , the flow without feedback, is a skew product with the Anosov flow f^t in the base and z_i -dynamics in the fibers, this skew-product structure is not preserved with the introduction of feedback. However, because the I-units affect the E-component through time changes only, the first return map $H : \Sigma_0 \rightarrow \Sigma_0$ of \mathbf{F}^t continues to be a skew-product. It can be written as $H = H_2 \circ H_1$ where $H_1 : \Sigma_0 \rightarrow \Sigma_b$ is given by

$$H_1(x, y, 0; z_1, \dots, z_N) = (x, y, b; z_1 + \hat{r}_1(x), \dots, z_N + \hat{r}_N(x))$$

and $H_2 : \Sigma_b \rightarrow \Sigma_0$ is defined as follows:

$$(2) \quad H_2(x, y, b; z_1, \dots, z_N) = (A(x, y), 0, G(z_1, \dots, z_N))$$

where

$$G(z_1, \dots, z_N) = (g_1^{\tau(z_1, \dots, z_N)}(z_1), \dots, g_N^{\tau(z_1, \dots, z_N)}(z_N))$$

and $\tau(z_1, \dots, z_N) = (1 - b)c_\Phi(z_1, \dots, z_N)^{-1}$.

The formula for τ , the time to go from Σ_b to Σ_0 , can be understood as follows: Without feedback, $\tau \equiv 1 - b$. With feedback, it is lengthened by the factor indicated, so that $1 - b \leq \tau \leq (1 - b)(1 - \Phi(N))^{-1} := \tau_{\max}$. The skew-product structure of H will play a crucial role in our arguments.

To summarize, then, the E-environment acts on the I-units with possibly large rotations every time the Anosov flow returns to the cross-section Σ_0 . After these rotations, the I-units collectively determine the subsequent speed of the Anosov flow depending on the positions to which they are rotated. The I-units do not interact with one another directly but do so by influencing the time to the next activation, that is, the time for the Anosov flow to reach Σ_0 .

2.3. Technical assumptions. The following conditions are assumed throughout:

(a) *The fiber flows g_i^t .* Recall that for all i , g_i^t has an attractive fixed point at $z = 0$ and a repelling fixed point at $z = \frac{1}{2}$. We now assume additionally that for each i , there are numbers $\delta_i^+, \delta_i^- > 0$ and $c \in (0, 1)$ such that if

$$I_i^+ = \left[\frac{1}{2} - \delta_i^+, \frac{1}{2} + \delta_i^+ \right] \quad \text{and} \quad I_i^- = [0 - \delta_i^-, 0 + \delta_i^-] ,$$

then for all $(1 - b) \leq t \leq \tau_{\max}$:

(A1) $(g_i^t)'|_{I_i^+} > e^\lambda$, $(g_i^t)'|_{I_i^-} \leq c$, and

(A2) $g_i^t(I_i^+) \supset \mathbb{T} \setminus I_i^-$.

(b) *The functions $r_i(x)$.* Recall that the total amount z_i is rotated on $\{w \in [0, b]\}$ depends on the x -coordinate of the Anosov trajectory as explained in Paragraph C. We now impose some additional conditions on the rotation function $r_i(x)$. Define

$$d_i := \text{dist}(\partial(I_i^-), \partial g_i^{(1-b)}(I_i^+)).$$

Then there exist $\varepsilon > 0$ small enough and $c' > 0$ such that

(A3) $r_i' > \varepsilon^{-1}$ except where $r_i \in (d_i, 1 - d_i)$,

(A4) $r_i' > c'$ everywhere on \mathbb{T} .

The existence of c' satisfying (A4) follows from the fact that r_i is a local diffeomorphism. In Figure 2 we illustrate the technical assumptions (A1)-(A4) and give some examples of r_i .

Example of N-S flow and function r_i satisfying assumptions (A1)-(A4).

A 2×2 invertible matrix M defines an action $\hat{g} : \mathbb{R} \times \mathbb{R}^2 \setminus \{0\} \rightarrow \mathbb{R}^2 \setminus \{0\}$ given by $g(t, v) = e^{tM}v$. Let $\mathbb{P}^1(\mathbb{R}) \cong \mathbb{T}$ be the 1-D real projective space. It is easy to check that \hat{g} projects to a flow g^t on \mathbb{T} , and that picking M diagonal with entries $\pm\alpha$ gives rise to a North-South flow on \mathbb{T} where $(g^t)' = e^{\pm\alpha t}$ at the two poles. Given λ, b , and τ_{\max} , (A1)-(A2) are satisfied by choosing α sufficiently large.

Now fix $\kappa =$ the degree of r_i , the N-S flow (so that d_i is fixed), and $\varepsilon \ll \kappa^{-1}$. Pick κ disjoint closed arcs on \mathbb{T} , $\{[w_{1,\ell}, w_{2,\ell}]\}_{\ell=0}^{\kappa-1}$, of length $\varepsilon(1 - 2d_i)$ and assume they are ordered. Define $\tilde{r}_i : \mathbb{T} \rightarrow \mathbb{R}$ to be a C^2 increasing function with the following




FIGURE 2. The left panel above shows an example of the time- $(1-b)$ -map, $g^{(1-b)}$, for a North-South flow. The diagram shows how $g^{1-b}(I^+)$ should overlap with I^- for (A2) to be satisfied, where I^- and I^+ satisfy (A1). The other two panels show examples of functions r_i . These functions have the characteristic of being very steep for those x such that $d \leq r_i(x) \leq 1-d$, and, although possibly very small, its slope is never equal to zero. Here r_1 has degree 1 and r_2 has degree two.

properties: (i) it is affine on $[w_{1,\ell}, w_{2,\ell}]$ with constant slope ε^{-1} and $\tilde{r}_i([w_{1,\ell}, w_{2,\ell}]) = [d_i + \ell, 1 - d_i + \ell]$; (ii) on $(w_{2,\ell}, w_{1,\ell+1})$, it increases monotonically from $1 - d_i + \ell$ to $d_i + \ell + 1$ with $\tilde{r}'_i > 2d_i$ (independent of $\varepsilon > 0$). Finally let r_i be the projection of \tilde{r}_i to \mathbb{T} .

3. STATEMENT OF THE MAIN RESULTS AND AN ILLUSTRATIVE EXAMPLE

3.1. Main results. The model described in Sect. 2.2 is a piecewise smooth flow on a $(3+N)$ -dimensional manifold M where $N \in \mathbb{Z}^+$ is arbitrary. In finite dimensional systems, observable events are often equated with positive Lebesgue measure sets, and *physical measures* are natural invariant measures. We begin by recalling the definition of a physical measure. Below, m denotes the Riemannian measure on M .

Definition 3.1. Let φ^t be a smooth or piecewise smooth flow on M . A φ^t -invariant Borel probability measure μ is called a **physical measure** if there is a Borel set $U \subset M$ with $m(U) > 0$ such that for every continuous function $h : M \rightarrow \mathbb{R}$,

$$\frac{1}{T} \int_0^T h(\varphi^t(x)) dt \rightarrow \int h d\mu \quad \text{as } T \rightarrow \infty$$

for m -a.e. $x \in U$.

If φ^t possesses an ergodic probability measure μ absolutely continuous with respect to Lebesgue, then μ is a physical measure by the Birkhoff Ergodic Theorem. But the notion of physical measure is meaningful even when φ^t is dissipative, i.e., when all invariant probability measures are singular with respect to m .

We will assert below that under suitable conditions, the flow \mathbf{F}^t defined in Sect. 2.2 possesses a physical measure. To state these conditions, we first clarify the relations among the system's constants:

- (i) chosen freely are $N = \#$ I-components and $\kappa =$ maximum degree of the r_i ;
- (ii) λ , the expanding coefficient of the Anosov flow, depends on N (see below);

- (iii) the g_i^t are then chosen to satisfy (A1) and (A2) (note (A1) depends on λ);
- (iv) c' is chosen depending on κ and g_i ;
- (v) ε depends on everything in (i)-(iii) (see below); and
- (vi) the r_i are then chosen to satisfy (A3)-(A4) (which depends on ε).

Theorem 3.2. *Let \mathbf{F}^t have the form in Sect. 2.2. There are two functions $\lambda(N)$ and $\varepsilon(N, \lambda, g_i^t, \kappa)$ such that if in addition to satisfying (A1)-(A4), λ and ε are chosen so that*

$$\lambda \geq \lambda(N) \quad \text{and} \quad \varepsilon \leq \varepsilon(N, \lambda, g_i^t, \kappa),$$

then \mathbf{F}^t admits a physical measure μ .

For systems with hyperbolic properties, a standard way to produce physical measures is to construct SRB measures.

Definition 3.3. Let φ^t be a smooth or piecewise smooth flow on M . A φ^t -invariant Borel probability measure μ is called an **SRB measure** if

- (i) φ^t has a positive Lyapunov exponent μ -a.e.;
- (ii) unstable manifolds are defined μ -a.e.
- (iii) conditional probabilities of μ on unstable manifolds have densities.

For definiteness, let us use the term “unstable manifolds” to refer to weak unstable manifolds, which includes flowlines, to be distinguished from strong unstable manifolds, which are one dimension lower. For flows without singularities, (ii) follows automatically from (i). In the presence of discontinuities or singularities, additional conditions are needed to ensure that unstable manifolds are defined; that is why we have included that as part of the definition.

Theorem 3.4. *Under the assumptions of Theorem 3.2, the flow \mathbf{F}^t has an ergodic SRB measure μ with exactly one positive Lyapunov exponent and no zero Lyapunov exponent aside from the one in the flow direction.*

Theorem 3.2 then follows from Theorem 3.4 and the absolute continuity of the strong stable foliation.

The proofs of Theorems 3.2 and 3.4 are contained in Sects. 4-6. For clarity of exposition, *the proof we present is written for $N = 2$* . The case $N = 1$ is simpler, and for $N \geq 3$, the proof is conceptually identical to that for $N = 2$. However, some choices of constants depend on N , the main reason being that the singularity set grows in complexity as N increases. We will identify the dependence on N as we go along.

3.2. Modeling and simulations. As remarked in the Introduction, the models considered are not intended to be realistic models of specific physical or biological systems, but they exhibit a few characteristics typical of such systems: Excitatory-inhibitory relations among constituent components, the activation of certain processes upon crossing of thresholds, and the relaxation to equilibrium in the absence of excitatory input — these properties appear often in biological systems.

Under (A1)–(A4), the system is hyperbolic (though not uniformly so). In many dynamical systems that have come to symbolize chaotic behavior, such as Anosov diffeomorphisms, billiards or standard maps, the landscape is dominated by clearly




FIGURE 3. Results for numerical simulations of a population of $N = 20$ I-units with $v_g(z) = -7 \sin(2\pi z)$, $A = (10, 3; 3, 1)$, $\Phi(n) = n^{-3/10}$, and where the functions r_i are all different having degrees between 2 and 3. The x -axis is time in all the panels. The first panel shows the Raster plot of the “activation” events. The y -axis is the I-unit number. A marker is added at (t, i) when I-unit number i has coordinate crossing the value 0.5 at time t . The second panel shows the w coordinate of the E-environment. The last three panels shows the orbits of I-units number 10 and 11 and 20. Because their r_i are different, distinct units exhibit different activation patterns.

expanding and contracting directions – defined everywhere or at least on large portions of the phase space. This is not the case in the network models described in Section 2. In these models, individual I-component are described by simple N-S flows with large perturbations mediated by other network components occurring at seemingly random times. These events inject unpredictability into the system, leading to very rich and varied patterns of collective dynamical behaviors. This is quite typical of high dimensional hyperbolic systems with few unstable directions. Such systems are not uncommon in real-world settings, and they may be relatively amenable to analysis.

In Figure 3 we present an illustrative example of a network model of the type studied in this paper, to give a sense of its time evolution. Shown are snapshots of the E-environment and three of the I-components on a randomly chosen time interval. These snapshots were “typical” to the degree that they were not chosen with specific characteristics in mind; in large dynamical systems, the patterns that can arise are infinitely rich and it is not clear what constitutes “typical” behavior. The character of the dynamics depends on parameters, including the choices of Φ , v_g , A , and especially $\{r_i\}$, the collection of functions describing the amount the I-component is rotated.

Qualitative properties of some of these dependences can be analyzed, e.g. since z_i is expected to be near 0 a good fraction of the time (this is what we will prove), rotating by $> \frac{1}{2}$ is more likely to cause z_i to cross 0.5. In this way, one can identify which I-component is likely to activate more often. Moreover components with

similar r_i tend to activate simultaneously. Another observation is that if Φ is linear, then a constraint on the impact of each I-component is that it should scale like $\sim \frac{1}{N}$ as system size N tends to infinity, to ensure that $\Phi(\sum_{i=1}^N \chi_{(\frac{1}{2}, 1)}(z_i)) < 1$; see Equation (1) in Sect. 2.2, Paragraph D. An interesting question is if, in the limit $N \rightarrow \infty$, the system can be described by a self-consistent operator (as defined e.g. in [ST21] for systems in discrete time) and what are the properties of this operator.

While a complete characterization of the very rich dynamical behaviors of this model is likely out of reach, many aspects are amenable to analysis. We will not delve into such an investigation here, however, as that would take us too far from our goal of proving the results stated in Section 3.

4. DISTRIBUTION OF PUSHED-FORWARD MASS: A PRELIMINARY ESTIMATE

In Sections 4–6, unless declared otherwise, N is assumed to be 2.

We will, for the most part, be working with the first return map $H : \Sigma_0 \circlearrowleft$ of the flow \mathbf{F}^t defined in Sect. 2.2. Here, $\Sigma_0 \cong \mathbb{T}^2 \times \mathbb{T}^2$ (with Euclidean norm); we use $(x, y; z_1, z_2)$ to denote its coordinates. Guessing that the bulk of pushed-forward mass is likely to accumulate near $\{z_1 = z_2 = 0\}$, the S-poles of the fiber flows (g_i^t, g_2^t) in (z_1, z_2) -space, so that the Lyapunov exponents of H in (z_1, z_2) -space are likely both negative, we seek to construct an SRB measure with 1D unstable manifolds. Our guess is informed by assumptions (A1)–(A4), which were designed to ensure that mass could not collect around the N-poles. Guided by this intuition, we push forward Lebesgue measure on 1D curves roughly aligned with E_A^u , hoping to obtain an SRB measure in the limit.

The aim of this section is to confirm the hypothesis on mass concentration.

4.1. Setup and main proposition. Let $W_A^u \subset \mathbb{T}^2$ be a piece of unstable manifold for $A : \mathbb{T}^2 \circlearrowleft$, and let $\gamma_0 : [0, a] \rightarrow \Sigma_0$ be a parametrization of $W_A^u \times \{0\}$ by arc length. For $n \geq 1$, let $\gamma_n : [0, e^{\lambda n} a] \rightarrow \Sigma_0$ be defined by

$$\gamma_n(u) = H^n(\gamma_0(e^{-\lambda n} u)) ,$$

so that if π_{xy} denotes the projection of $\Sigma_0 \cong \mathbb{T}^2 \times \mathbb{T}^2$ to its first \mathbb{T}^2 -component, then $\pi_{xy} \circ \gamma_n$ is a parametrization of $A^n(W_A^u)$ by arc length.

Action of H in (u, z_1, z_2) -space. In the analysis to follow, it will be convenient to view the action of H in the 3D space $\mathbb{R} \times \mathbb{T}^2$ with coordinates (u, z_1, z_2) , where u is arclength parametrization of $A^n(W_A^u)$. Abusing notation slightly (to avoid excessive numbers of symbols), we view γ_n as a curve in this space parametrized by

$$\gamma_n = (u, \pi_{z_1}(\gamma_n(u)), \pi_{z_2}(\gamma_n(u))), \quad u \in [0, e^{\lambda n} a],$$

where π_{z_i} has the obvious meaning, and study the evolution of γ_n in this setting under the iteration of H .

To define the action of H , we need to express the amount of rotation as functions of u . We introduce a γ_0 -dependent family of functions

$$\iota_n : [0, e^{\lambda n} a] \rightarrow \mathbb{T} , \quad \iota_n(u) = \pi_x(\gamma_n(u)) .$$

Then $\iota_n(u)$ has the form $\iota_n(u) = \beta u + \pi_x(\gamma_n(0))$ where β is the angle between the unstable direction E_A^u and the x direction and $\pi_x(\gamma_n(0))$ is a constant depending on γ_n .

The action of H on (u, z_1, z_2) -space can now be written as $\Phi := \Phi_3 \circ \Phi_2 \circ \Phi_1$ where

$$\begin{aligned}\Phi_1 &: (u; z_1, z_2) \mapsto (u; z_1 + r_1(\iota_n(u)), z_2 + r_2(\iota_n(u))) \\ \Phi_2 &: (u; z_1, z_2) \mapsto (u; g_1^{\tau_1(z_1, z_2)}(z_1), g_2^{\tau_2(z_1, z_2)}(z_2)) \\ \Phi_3 &: (u; z_1, z_2) \mapsto (e^\lambda u; z_1, z_2)\end{aligned}$$

This is the setup we will be working with.

Observe that in this setup, the curves of interest are all graphs of functions in the variable u : The curve γ_0 is the graph of the function $\theta_0(u) \equiv (0, 0)$. Suppose we show inductively that

$$\gamma_n = \text{graph}(\theta_n) \quad \text{for some} \quad \theta_n = (\theta_n^1, \theta_n^2) : [0, e^{\lambda n} a] \rightarrow \mathbb{T}^2.$$

Then Φ_1 transforms the graph of θ_n to the graph of another function $\psi_n = (\psi_n^1, \psi_n^2)$, i.e. $\Phi_1(\text{graph}(\theta_n)) = \text{graph}(\psi_n)$, where

$$\psi_n(u) = \theta_n(u) + (r_1(\iota_n(u)), r_2(\iota_n(u))) .$$

Likewise, Φ_2 transforms the graph of ψ_n to the graph of another function $\zeta_n = (\zeta_n^1, \zeta_n^2)$ with

$$\zeta_n(u) = (g_1^{\tau(\psi_n(u))}(\psi_n^1(u)), g_2^{\tau(\psi_n(u))}(\psi_n^2(u))),$$

and Φ_3 transforms the graph of ζ_n to the graph of θ_{n+1} with

$$\theta_{n+1}(u) = \zeta_n(e^{-\lambda} u) .$$

As discussed at the beginning of this section, we are interested in where mass accumulates in (z_1, z_2) -space starting from uniform measure on γ_0 and transporting mass forward by H . Below we translate this statement into one about ψ_n . The symbols I_i^- and ε refer to those in Sect. 2.3.

Proposition 4.1. *Assume $e^\lambda > 3$ and $\varepsilon > 0$ is sufficiently small. Then there exists $C_\# > 0$ (dependent on κ, c', a, β , and the N - S flows g_i) such that for all $n \in \mathbb{N}$ and $i = 1, 2$,*

$$\frac{1}{e^{\lambda n} a} \cdot m\{u \in [0, e^{\lambda n} a] : \psi_n^i(u) \in (I_i^-)^c\} < C_\# \varepsilon$$

where m denotes Lebesgue measure on \mathbb{R} .

This is the main result of this section. Had ψ_n^i been smooth and monotonically increasing, a natural way to estimate $m\{\psi_n^i(u) \in (I_i^-)^c\}$ for each i would be to examine the derivative of ψ_n^i and to count how many times its graph wraps around \mathbb{T} as u varies over the interval $[0, e^{\lambda n} a]$. But as we will show, the functions ψ_n^i are neither continuous nor monotonic due to discontinuities in our model. In Sect. 4.2, we will investigate the extent to which they fail to be continuous and monotonic. The proof of Proposition 4.1 is given in Sect. 4.3.

4.2. Geometry of iterated curves. Starting from $\theta_0^1 \equiv 0$, we see from the definition of $\Phi := \Phi_3 \circ \Phi_2 \circ \Phi_1$ that Φ_1 and Φ_3 will take a smooth piece of graph that is monotonically increasing to another such piece of graph, and new singularities can only be created as we pass from ψ_n to ζ_n , so we focus below on the action of Φ_2 .

The following language will be used to facilitate the discussion: We say a function $h : \mathbb{R} \rightarrow \mathbb{R}$ is “monotonic at x ” if there is a neighborhood of x on which h is monotonic, C^1 at x if there is a neighborhood on which h is C^1 , and “ C^{1*} at x ” if there is a neighborhood on which h is C^1 except possibly at x . We call x a “singularity” of h if it is not C^1 at x , and use $S(h)$ to denote the set of singularities.

Lemma 4.2. *The following hold for all $n \in \mathbb{N}$ and $i = 1, 2$:*

(i) *The set $S(\psi_n^i)$ is finite, and $\exists c_1 > 0$ depending only on c' and the N - S flow g_i (independent of n) such that $(\psi_n^i)'(u) \geq c_1$ for all $u \notin S(\psi_n^i)$.*

Suppose ψ_n^i , $i = 1, 2$, is C^1 (resp. C^{1}) and monotonic at \bar{u} .*

(ii) *If $\psi_n^i(\bar{u}) \neq 0, \frac{1}{2}$ for $i = 1, 2$, then ζ_n^i is C^1 (resp. C^{1*}) and monotonic at \bar{u} .*

(iii) *If $\psi_n^1(\bar{u}) = 0$ or $\frac{1}{2}$, then ζ_n^1 is C^{1*} and monotonic at \bar{u} .*

(iv) *If $\psi_n^1(\bar{u}) \neq 0, \frac{1}{2}$ and $\psi_n^2(\bar{u}) = 0$ or $\frac{1}{2}$, then ζ_n^1 is discontinuous and not necessarily monotonic at \bar{u} .*

Statements (iii) and (iv) hold with $i = 1, 2$ interchanged.

Once Item (i) is proved, it will follow that for $\bar{u} \notin S(\psi_n^i)$, if ζ_n^i is continuous at \bar{u} , then it is automatically monotonic; if ζ_n^i is discontinuous at \bar{u} , then it either jumps up or down, and a violation of monotonicity is created if and only if ζ_n^i jumps down.

Proof. The dynamics on $\mathbb{T}^2 = (z_1, z_2)$ -space are generated by a single C^2 vector field (v_{g_1}, v_{g_2}) , but \mathbb{T}^2 is divided into 4 quadrants by the lines $z_i = 0, \frac{1}{2}$, $i = 1, 2$, and singularities of Φ_2 arise from the fact that the flow times τ in g_i^r are different on these 4 quadrants.

The assertion in (ii) follows immediately. Item (i) is proved inductively, using (ii), (A4), which guarantees a lower bound on r'_i , and the fact that Φ_2 and Φ_3 can only decrease the derivative of ψ_n^i by a finite amount.

For (iii) and (iv), it is useful to picture a small piece of curve $u \mapsto \psi_n^i(u)$ in \mathbb{T}^2 defined for u in a neighborhood of \bar{u} , and see how it is transformed to $u \mapsto \zeta_n^i(u)$ by applying the appropriate flow-maps. In the scenario of (iii), different flow-times are applied to $u < \bar{u}$ and $u > \bar{u}$, but no discontinuity is created because 0 and $\frac{1}{2}$ are fixed points of the flow g_t^i . A jump discontinuity in ζ_n^1 is created in the scenario in (iv). This jump can be up or down depending on the location of $\psi_n(\bar{u})$: For example, if $\psi_n^2(\bar{u}) = \frac{1}{2}$ and $\psi_n^1(\bar{u}) \in (0, \frac{1}{2})$, then $\tau(\psi_n(u))$ for $u < \bar{u}$ is smaller than $\tau(\psi_n(u))$ for $u > \bar{u}$, and since g_t^1 is decreasing at \bar{u} , the jump is downwards, i.e., ζ_n^1 loses monotonicity at \bar{u} . Similar reasoning will show that the jump is upwards (hence ζ_n^1 retains its monotonicity) if we take $\psi_n^2(\bar{u}) = 0$ and $\psi_n^1(\bar{u})$ as before. \square

As it turns out, isolated points of nondifferentiability are of no concern to us for as long as the function is continuous at those points, i.e., the singularities created in scenario (iii) are harmless. Our concern is (iv). Note also that discontinuities of ζ_n^1 are determined not by ψ_n^1 but by ψ_n^2 .

As downward jumps are unavoidable, we try to control the distances jumped. Let $h : [0, b] \rightarrow \mathbb{T}$ be a piecewise continuous, monotonically increasing function, i.e., there exist $0 = d_0 < d_1 < \dots < d_k = b$ such that h is continuous and monotonically increasing on each interval (d_{j-1}, d_j) , $j = 1, \dots, k$, and it has jump discontinuities at d_1, \dots, d_{k-1} . Notice that no conditions are imposed on the relation between $h^-(d_j)$ and $h^+(d_j)$ where $h^\pm(d_j)$ denote the left and right limits of h at d_j . Given such an h , we let $\Gamma h : [0, b] \rightarrow \mathbb{R}$ be the function with the following properties:

- (i) $(\Gamma h)(u) = h(u) \bmod 1$ for all $u \in [0, b]$;
- (ii) Γh is monotonically increasing on $[0, b]$, i.e. $u_1 \leq u_2 \implies (\Gamma h)(u_1) \leq (\Gamma h)(u_2)$ for all $u_1, u_2 \in [0, b]$;
- (iii) $(\Gamma h)^-(d_j) < (\Gamma h)^+(d_j) < (\Gamma h)^-(d_j) + 1$;
- (iv) $(\Gamma h)(0) \in [0, 1)$.

That is to say, Γh changes h into a monotonically increasing function with jump sizes < 1 while preserving the values of $h \bmod 1$. It is obvious that Γh is uniquely defined and has jump discontinuities at the same locations as h .

For $i = 1, 2$ and all n , let

$$\hat{\theta}_n^i = \Gamma(\theta_n^i), \quad \hat{\psi}_n^i = \Gamma(\psi_n^i), \quad \hat{\zeta}_n^i = \Gamma(\zeta_n^i).$$

We let $\mathcal{R}(\hat{\theta}_n^i)$ denote the length of the shortest interval containing the image of $\hat{\theta}_n^i([0, e^{\lambda n} a])$; $\mathcal{R}(\hat{\psi}_n^i)$ and $\mathcal{R}(\hat{\zeta}_n^i)$ are defined analogously. We are interested especially in $\mathcal{R}(\hat{\psi}_n^i)$, which as we will see is an overcount for the number of times ψ_n^i wraps around \mathbb{T} (see the last paragraph of Sect. 4.1).

Proposition 4.3. *Assume that $e^\lambda > 3$. Then there is $c_2 > 0$ (depending on κ, β, a , and λ) such that for every $n \in \mathbb{N}$ and $i \in \{1, 2\}$,*

$$(3) \quad \mathcal{R}(\hat{\psi}_n^i) \leq c_2 e^{\lambda n}.$$

Proof. Let κ be such that the degree of r_i is $\leq \kappa$ for every i .

We claim that for $i = 1, 2$ and for $n \geq 1$,

$$\begin{aligned} \mathcal{R}(\hat{\theta}_n^i) &= \mathcal{R}(\hat{\zeta}_{n-1}^i), & \mathcal{R}(\hat{\psi}_n^i) &\leq \mathcal{R}(\hat{\theta}_n^i) + \kappa e^{\lambda n} a \beta + \kappa, \\ \mathcal{R}(\hat{\zeta}_n^i) &\leq \mathcal{R}(\hat{\psi}_n^i) + 1 + X_n^i \end{aligned}$$

where X_n^i is the number of new discontinuities created as we go from ψ_n^i to ζ_n^i . The first equality is clear, as θ_n^i is simply a rescaling of ζ_n^i in u . The quantity $\kappa e^{\lambda n} a \beta$ comes from the fact that the rotation functions r_i have degree bounded by κ , and β is the modulus of the cosine of the angle between W_A^u and the x -axis. The “ $+\kappa$ ” in $\mathcal{R}(\hat{\psi}_n^i)$ and “ $+1$ ” in $\mathcal{R}(\hat{\zeta}_n^i)$ are end point corrections: the amount rotated can be arbitrarily close to $\kappa \lceil e^{\lambda n} a \beta \rceil$, i.e. the smallest integer $\geq e^{\lambda n} a \beta$ times κ , while g_i^t can increase the length of an arc by $\frac{1}{2}$ at either end.

We claim that $X_n^1 \leq 2(\mathcal{R}(\hat{\psi}_n^2) + 1)$. This is because by Lemma 4.2 (iv), discontinuities are created only when ψ_n^2 crosses $z_2 = 0, \frac{1}{2}$, and the number of times that can happen is $\leq 2(\mathcal{R}(\hat{\psi}_n^2) + 1)$. Similarly, $X_n^2 \leq 2(\mathcal{R}(\hat{\psi}_n^1) + 1)$.

Letting $\mathcal{R}(\hat{\psi}_n) = \max_{i=1,2} \mathcal{R}(\hat{\psi}_n^i)$ and similarly for $\mathcal{R}(\hat{\zeta}_n)$, we deduce from the relations above that

$$\mathcal{R}(\hat{\zeta}_n) \leq 3\mathcal{R}(\hat{\psi}_n) + 3 \leq 3(\mathcal{R}(\hat{\zeta}_{n-1}) + \kappa e^{\lambda n} a \beta + \kappa) + 3.$$

Applying this relation recursively, we obtain

$$\begin{aligned} \mathcal{R}(\hat{\zeta}_n) &\leq 3\mathcal{R}(\hat{\zeta}_{n-1}) + 3\kappa e^{\lambda n} a \beta + 3 + 3\kappa \\ &\leq 3^2 \mathcal{R}(\hat{\zeta}_{n-2}) + \kappa a \beta (3e^{\lambda n} + 3^2 e^{\lambda(n-1)}) + (3+1)(3+3\kappa) \\ &\leq \dots \\ &\leq 3^n \mathcal{R}(\hat{\zeta}_0) + 3\kappa a \beta \sum_{i=0}^{n-1} 3^i e^{\lambda(n-i)} + (3+3\kappa) \sum_{i=0}^{n-1} 3^i. \end{aligned}$$

The first and third terms are $\mathcal{O}(3^n)$ as $n \rightarrow \infty$. Since $e^\lambda > 3$ is assumed, the middle term is $\mathcal{O}(e^{\lambda n})$. It follows that $\mathcal{R}(\hat{\zeta}_n) = \mathcal{O}(e^{\lambda n})$, from which the asserted statement follows. \square

Remark 4.4. Notice that while the condition $e^\lambda > 3$ in Proposition 4.3 is sufficient for $N = 2$, more stringent lower bounds for λ are needed as N , the number of I-components, increases. This is because going from ψ_n^1 to ζ_n^1 , jump discontinuities

are created when ψ_n^i crosses $0, \frac{1}{2}$ for all $i \neq 1$. Thus more jump discontinuities are created at each step for larger N , and a larger lower bound for λ is required to beat the growth in number of singularities.

4.3. Proof of mass concentration. In addition to the results from Sect. 4.2, the proof below will rely heavily on Assumptions (A1)-(A4) in Sect. 2.3, and the notation will be as in that section.

Proof of Proposition 4.1. The proof consists of two steps.

To ensure that mass is concentrated near S-pole, our first step is to show that for $i = 1, 2$ and all $n \in \mathbb{N}$,

$$(4) \quad \text{if } u \notin S(\psi_n^i) \text{ is such that } \psi_n^i(u) \in (I_i^-)^c, \text{ then } (\psi_n^i)'(u) \geq \beta\varepsilon^{-1} .$$

We fix i , and prove the claim by induction on n .

The idea is to look at u such that $\psi_n^i(u)$ is near N-pole and compare it with $\psi_{n-1}^i(e^{-\lambda}u)$. If $\psi_{n-1}^i(e^{-\lambda}u)$ was near N-pole, then $(\psi_{n-1}^i)'(e^{-\lambda}u)$ was large by induction, and there are no mechanisms that can decrease this derivative by too much going to $\psi_n^i(u)$; if it was far from N-pole, then it must have experienced a large rotation to get there at step n , and (A3) ensures that r_i creates a very steep slope for such rotations.

More precisely, assume that (4) is true for $n-1$, and let u be as in (4). Then

$$\psi_n^i(u) = g_i^{\tau(\psi_{n-1}(e^{-\lambda}u))}(\psi_{n-1}^i(e^{-\lambda}u)) + \hat{r}_i(\iota_n(u)) ,$$

and since $u \mapsto \tau(\psi_{n-1}(e^{-\lambda}u))$ is locally constant, we have

$$(\psi_n^i)'(u) = (g_i^{\tau(\psi_{n-1}(e^{-\lambda}u))})'(\psi_{n-1}^i(e^{-\lambda}u)) \cdot (\psi_{n-1}^i)'(e^{-\lambda}u) \cdot e^{-\lambda} + (\hat{r}_i \circ \iota_n)'(u) .$$

We consider separately the following two cases.

Case (a): $\psi_{n-1}^i(e^{-\lambda}u) \in I_i^+$. Here we claim to have

$$(\psi_n^i)'(u) \geq e^\lambda \cdot \beta\varepsilon^{-1} \cdot e^{-\lambda} = \beta\varepsilon^{-1} .$$

That $(g_i^{\tau(\dots)})'(\psi_{n-1}^i(e^{-\lambda}u)) > e^\lambda$ follows from (A1) together with $\psi_{n-1}^i(e^{-\lambda}u) \in I_i^+$. That $(\psi_{n-1}^i)'(e^{-\lambda}u) \geq \beta\varepsilon^{-1}$ follows from the induction hypothesis, and the \hat{r}_i -term can be dropped because it is > 0 .

Case (b): $\psi_{n-1}^i(e^{-\lambda}u) \in (I_i^+)^c$. It follows from (A2) that $g_i^{\tau(\psi_{n-1}(e^{-\lambda}u))}(\psi_{n-1}^i(e^{-\lambda}u))$ is at least distance d_i away from ∂I_i^- . In order for $\psi_n^i(u)$ to belong to $(I_i^-)^c$, $\hat{r}_i(\iota_n(u))$ must be $> d_i$, so $r_i'(\iota_n(u)) > \varepsilon^{-1}$ by (A3). Since $(\psi_{n-1}^i)' > 0$, we have

$$(\psi_n^i)'(u) \geq (\hat{r}_i \circ \iota_n)'(u) = \hat{r}_i'(\iota_n(u))\iota_n'(u) > \beta\varepsilon^{-1} .$$

Our second step is to use the derivative information above to bound the number of times the graph of ψ_n^i meets the interval $(I_i^-)^c$. Here we have taken care of the fact that ψ_n^i is not monotonic by replacing it with $\hat{\psi}_n^i$ (see Sect. 4.2). Define

$$G_n^i = \{u \in [0, e^{\lambda n}] : \hat{\psi}_n^i(u) \in (I_i^-)^c + \mathbb{Z}\} .$$

Thinking of the gaps at jumps as pieces of graph with infinitely large slopes, we see that the number of components of $(I_i^-)^c + \mathbb{Z}$ reached by the range of $\hat{\psi}_n^i$ is $\leq \mathcal{R}(\hat{\psi}_n^i) + 2$, and we have shown that $\mathcal{R}(\hat{\psi}_n^i) < c_2 e^{\lambda n}$ (Proposition 4.3). If J is a

component of $(I_i^-)^c + \mathbb{Z}$, then $|(\hat{\psi}_n^i)^{-1}(J)| \leq |(I^-)^c| \beta^{-1} \varepsilon$ from Step 1. Altogether, we have

$$\frac{1}{e^{\lambda n a}} m(G_n^i) < \frac{1}{e^{\lambda n a}} \cdot c_2 e^{\lambda n} \cdot |(I_i^-)^c| \beta^{-1} \varepsilon < C_{\#} \varepsilon$$

for some $C_{\#}$, completing the proof. \square

5. A CANDIDATE SRB MEASURE FOR THE RETURN MAP

Assumptions (A1)–(A4) are in effect throughout. We continue to develop the ideas outlined at the beginning of Sect. 4, namely to construct a candidate SRB measure for the first return map $H : \Sigma_0 \circlearrowleft$ of the flow \mathbf{F}^t by pushing forward Lebesgue measure on curves with a component in the unstable direction. Let γ_0 be as defined in Sect. 4.1, and let m_{γ_0} be Lebesgue measure on γ_0 . Assuming $a = 1$ so m_{γ_0} is a probability measure, we let

$$\mu_n := \frac{1}{n} \sum_{i=0}^{n-1} H_*^i(m_{\gamma_0}) .$$

5.1. Limit points of μ_n and relation to singularities. The main obstacle to concluding any limit point of μ_n is an H -invariant probability measure with desirable properties is the presence of singularities, so that is what we will focus on.

Recall that $H = H_2 \circ H_1$ (Sect. 2.2), and that $H_1 : \Sigma_0 \rightarrow \Sigma_b$ is a diffeomorphism whereas $H_2 : \Sigma_b \rightarrow \Sigma_0$ is piecewise smooth with discontinuities. Let $\mathcal{S}(\cdot)$ denote the singularity set of a map. Then

$$\mathcal{S}(H_2) = \sigma_b \times S$$

where $S := S_1 \cup S_2$, $S_i := \{z_i = 0, \frac{1}{2}\} \subset \mathbb{T}^2$, and $\sigma_b := \mathbb{T}^2 \times \{b\} \subset M_f$. Because $\mathcal{S}(H_2)$ has a simpler geometry than $\mathcal{S}(H) = H_1^{-1} \mathcal{S}(H_2)$, and $\mathcal{S}(\bar{H}) = \mathcal{S}(H_2)$ for $\bar{H} = H_1 \circ H_2$, it is simpler to work with \bar{H} . In the rest of Sect. 5.1, we will consider

$$(5) \quad \bar{\mu}_n := \frac{1}{n} \sum_{i=0}^{n-1} \bar{H}_*^i(H_{1*} m_{\gamma_0}) , \quad n = 1, 2, \dots .$$

The $(H_1^{-1})_*$ -image of any weak*-accumulation point of $\{\bar{\mu}_n\}$ is clearly an accumulation point of $\{\mu_n\}$.

We show below that weak*-accumulation points of $\{\bar{\mu}_n\}$, which exist in the space of all Borel probability measures by compactness, are invariant measures of \bar{H} with controlled properties near its singularity set. For $X \subset \mathbb{T}^m$ and $\varepsilon > 0$, we denote by X_ε the ε -neighborhood of X .

Lemma 5.1. *There exists $c_3 > 0$ (with the same dependencies of c_1 and c_2) such that if $\{\bar{\mu}_n\}_{n \in \mathbb{N}}$ is as defined in (5), then for all $\xi > 0$,*

$$(6) \quad \bar{\mu}_n(\sigma_b \times S_\xi) \leq c_3 \xi .$$

Proof. We estimate $\bar{\mu}_n(\sigma_b \times (S_i)_\xi)$ for $i = 1$. The case $i = 2$ is analogous.

Recalling the definition of ψ_n , one can see that

$$\bar{H}_*^i(H_{1*} m_{\gamma_0})(\sigma_b \times (S_1)_\xi) = \frac{1}{e^{\lambda i}} m \left((\hat{\psi}_i^1)^{-1}(S_1)_\xi \right) .$$

By point i) of Lemma 4.2 $(\psi_i^1)' \geq c_1 > 0$ where ψ_i^1 is differentiable, therefore the preimage of each copy of $(S_1)_\xi$ in one fundamental domain on the lift has measure at

most $2\xi c_1^{-1} e^{-\lambda i}$. By Proposition 4.3, $\hat{\psi}_i^1$ spans at most $c_2 e^{\lambda i}$ fundamental domains. The result is proved with $c_3 = 4c_2 c_1^{-1}$. \square

Proposition 5.2. *Under Assumptions (A1)–(A4), any accumulation point $\bar{\mu}$ of the sequence in (5) is an invariant probability measure for \bar{H} , and it satisfies*

$$(7) \quad \bar{\mu}(\sigma_b \times S_\xi) \leq c_3 \xi$$

for every $\xi > 0$, where c_3 is as in Lemma 5.1.

Proof. Let \mathcal{M} be the set of all Borel probability measures on Σ_b , and let

$$\mathcal{M}_S := \{\nu \in \mathcal{M} : \nu(\sigma_b \times S_\xi) \leq c_3 \xi \ \forall \xi > 0\}.$$

Then by Lemma 5.1, $\bar{\mu}_n \in \mathcal{M}_S$. To use the standard Krylov-Bogolyubov argument to prove that any accumulation point of $\{\bar{\mu}_n\}$ is fixed by \bar{H}_* , it suffices to show that \bar{H}_* acts continuously on \mathcal{M}_S . Let $\nu_n \in \mathcal{M}_S$ be such that ν_n converges to ν in the weak* topology, and fix a continuous function $\varphi : \Sigma_b \rightarrow \mathbb{R}$. We will show $\int \varphi d(\bar{H}_* \nu_n) \rightarrow \int \varphi d(\bar{H}_* \nu)$.

Given an arbitrarily small $\delta > 0$, let $\xi > 0$ be small enough so that

$$(8) \quad c_3 \xi \sup_{p \in \Sigma_b} |\varphi(p)| \leq \frac{\delta}{2}.$$

Let $U_1 = (\sigma_b \times S_{\xi/2})^c$, $U_2 := (\sigma_b \times S_\xi)^c$, $U'_1 := \bar{H}U_1$ and $U'_2 := \bar{H}U_2$. Then U'_1 and U'_2 are closed sets with $U'_2 \subset \text{int}(U'_1)$. Let $\rho : \Sigma_b \rightarrow [0, 1]$ be a continuous function with $\rho|_{U'_2} = 1$ and $\rho|_{U'_1{}^c} = 0$. Since the support of ρ is contained in U'_1 and that of $(1 - \rho)$ is contained in $(U'_2)^c$, we have, by the injectivity of \bar{H} ,

$$(9) \quad \text{supp}(\rho \circ \bar{H}) \subset U_1, \quad \text{supp}((1 - \rho) \circ \bar{H}) \subset U_2^c = \sigma_b \times S_\xi.$$

Writing

$$\int_{\Sigma_b} \varphi d\bar{H}_* \nu_n = \int_{\Sigma_b} (((1 - \rho)\varphi) \circ \bar{H}) d\nu_n + \int_{\Sigma_b} ((\rho\varphi) \circ \bar{H}) d\nu_n,$$

we have that the first integral on the right is $< \frac{1}{2}\delta$ by (8) and (9). Since the support of $\rho \circ \bar{H}$ is bounded away from $\sigma_b \times S$, the second integral converges to $\int (\rho\varphi) \circ \bar{H} d\nu$ as $n \rightarrow \infty$. It follows that

$$\left| \int \varphi d(\bar{H}_* \nu_n) - \int \varphi d(\bar{H}_* \nu) \right| \leq \delta$$

for all large n , proving the convergence claimed.

To complete the proof, observe that \mathcal{M}_S is closed and therefore every accumulation point belongs to \mathcal{M}_S . \square

The results above imply the following: Any limit point μ of μ_n where μ_n is as defined at the beginning of Sect. 5 is an invariant probability measure of H . From here on let $\mathcal{S} = \mathcal{S}(H)$ denote the singularity set of H . Then it follows from Proposition 5.2 that $\mu(\mathcal{S}_{\varepsilon_1}) < \text{const } \varepsilon_1$ for every $\varepsilon_1 > 0$.

5.2. Lyapunov exponents of (H, μ) . In this section we study the Lyapunov exponents (LE) of (H, μ) where $H : \Sigma_0 \circlearrowleft$ is the first return map and μ is a limit point of the sequence of measures μ_n . Recall that H has a skew-product structure: the base is $A : \mathbb{T}^2 \circlearrowleft$, and fiber variables are $(z_1, z_2) \in \mathbb{T}^2$. Let $E_{z_i} = \langle \partial_{z_i} \rangle$ denote the 1-dimensional space spanned by ∂_{z_i} , and let $E_{z_1 z_2} = \langle \partial_{z_1}, \partial_{z_2} \rangle$ be the space spanned by ∂_{z_1} and ∂_{z_2} . We claim that for $p \in \Sigma_0 \setminus \mathcal{S}$, $DH_p(E_{z_i}) = E_{z_i}$ for $i = 1, 2$: Writing

$$DH_p|_{E_{z_1 z_2}} = D(H_2)_{H_1(p)}|_{E_{z_1 z_2}} \circ D(H_1)_p|_{E_{z_1 z_2}},$$

we have $D(H_1)_p|_{E_{z_1 z_2}} = \text{Id}$ because on $\{\pi_{xy}(p)\} \times \mathbb{T}^2$, H_1 is a rigid translation, and writing $q = H_1(p)$,

$$D(H_2)_q|_{E_{z_1 z_2}} \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} = \begin{pmatrix} D(g_1^\tau)_{\pi_{z_1} q} w_1 \\ D(g_2^\tau)_{\pi_{z_2} q} w_2 \end{pmatrix},$$

where $\tau = \tau(\pi_{z_1} q, \pi_{z_2} q)$ is a locally constant function. This proves the claim.

Recalling that μ gives zero measure to the singularity set, so LEs are defined μ -a.e., we let $\lambda_{z_i}(p)$ denote the LE at p in the E_{z_i} -direction. Below $\varepsilon > 0$ is the number in Sect 2.2, Technical assumptions (b).

Lemma 5.3. *Assuming that $\varepsilon > 0$ is small enough, there is an H -invariant measure $\hat{\mu} = \mu|_B$, the restriction of μ to a Borel subset B , such that for $\hat{\mu}$ -a.e. p , $\lambda_{z_i}(p) < 0$ for $i = 1, 2$.*

Proof. It follows from the assumptions in Sect 2.3 that there exist $C > 1$ and $c < 1$ independent of ε such that for $i = 1, 2$,

- (i) $\|DH_p|_{E_{z_i}}\| \leq C$ and
- (ii) $\|DH_p|_{E_{z_i}}\| \leq c$ whenever $z_i \in I_i^-$

for all $p \in \Sigma_0 \setminus \mathcal{S}$. Let i be fixed. Assume ε is small enough that if μ_e is an ergodic component of μ with $\mu_e\{z_i \in I_i^-\} > 1 - \sqrt{\varepsilon}$, then by the Ergodic Theorem,

$$\lambda_{z_i} \leq \sqrt{\varepsilon} \log C + (1 - \sqrt{\varepsilon}) \log c < \frac{1}{2} \log c < 0 \quad \mu_e\text{-a.e.}$$

By Proposition 4.1 and the way we constructed μ , it follows that

$$(10) \quad \mu(\{z_i \in (I_i^-)^c\}) < C_{\#} \varepsilon.$$

Let $\{\mu_e\}_{e \in \mathcal{E}}$ be an ergodic decomposition of the invariant measure μ , and let $\mathcal{E}' \subset \mathcal{E}$ consist of those μ_e satisfying $\mu_e\{z_i \in I_i^-\} > 1 - \sqrt{\varepsilon}$ that by equation (10) is nonempty. Then we may take $\hat{\mu}$ to be $\int_{\mathcal{E}'} \mu_e$. \square

Let us assume from here on that $\hat{\mu}$ has been normalized so that $\hat{\mu}(\Sigma_0) = 1$. Recall that λ is the positive LE of the Anosov map A .

Corollary 5.4. *There exists $\lambda_{\min} < 0$ such that at $\hat{\mu}$ -a.e. p , λ is a LE and the other three LE are $\leq \lambda_{\min}$.*

Proof. Let $v^u = (\partial_A^u; 0, 0)$ be a tangent vector at p where $\partial_A^u \in E_A^u$ is vector along the unstable direction of A . Notice that $\forall n \in \mathbb{N}$, $DH_p^n v^u \in E_A^u \oplus E_{z_1 z_2}$ and $\pi_{xy} DH_p^n v^u = A^n \partial_A^u$, so $\|DH_p^n v^u\| \geq \|A^n \partial_A^u\| \geq e^{\lambda n} \|v^u\|$. This means that there is a DH -invariant 1D subspace $E^u \subset E_A^u \oplus E_{z_1 z_2}$ in which the LE is $\geq \lambda$. This LE is in fact $= \lambda$, since by the Multiplicative Ergodic Theorem, $\lim_{n \rightarrow \infty} \frac{1}{n} \log \sin \angle(E^u(H^n(p)), E_{z_1 z_2}) = 0$ $\hat{\mu}$ -almost everywhere.

A similar argument shows that $v^s := (\partial_A^s; 0, 0)$ grows exponentially under DH^{-n} with $\|DH_p^{-n}v^s\| \geq e^{\lambda n}\|v^s\|$. This implies that every vector $v \in E_A^s \oplus E_{z_1z_2}$ with a component in E_A^s grows exponentially under DH^{-n} . This together with Lemma 5.3 proves that all three LE on $E_A^s \oplus E_{z_1z_2}$ are $\leq \lambda_{\min} := \min\{-\lambda, \frac{1}{2}\log c\} < 0$. \square

5.3. $(H, \hat{\mu})$ as a nonuniformly hyperbolic system. The map $H : \Sigma_0 \circlearrowright$ with the invariant measure $\hat{\mu}$ is, *a priori*, mildly nonuniformly hyperbolic: At $\hat{\mu}$ -a.e. p , there is a splitting of its tangent space $T_p\Sigma_0 = E^u(p) \oplus E^s(p)$ into a 1D unstable subspace $E^u(p)$ (which varies with p) and a 3D stable subspace $E^s(p) \equiv E_A^s \oplus E_{z_1, z_2}$. Restricted to E_{z_i} , DH_p is sometimes expanding and sometimes contracting, depending on whether the z_i -coordinate of $H_1(p)$ is closer to $\frac{1}{2}$ or to 0. When the z_i -coordinate of $H_1(p)$ is closer to $\frac{1}{2}$, the expansion is stronger than e^λ by (A1). A stronger expansion in E_{z_i} than along ∂_A^u decreases the angle between E^u and E_{z_1, z_2} , and the repeated occurrence of such a scenario can potentially cause $\angle(E^u, E_{z_1, z_2})$ to come arbitrarily close to 0.

We do not know that $(H, \hat{\mu})$ is genuinely nonuniformly hyperbolic, but have to treat it as such unless proven otherwise. A standard technique for dealing with nonuniformly hyperbolic systems is through the use of certain point-dependent coordinate changes called Lyapunov charts (see e.g. [Pes76, Kat80, LY85, You95, KS06, BY17]). We review briefly below, in nontechnical terms, what these charts can do for us; details are provided in Appendix A.

For a piecewise smooth diffeomorphism equipped with an invariant probability measure that is not too concentrated near the singularity set, such as $H : \Sigma_0 \circlearrowright$, singularity set \mathcal{S} , and an invariant measure $\hat{\mu}$ with the property in Proposition 5.2, the following are known to hold:

- (1) Let $B(r) \subset \mathbb{R}^4$ be the ball of radius r centered at 0. Then at $\hat{\mu}$ -a.e. $p \in \Sigma_0$, there is a diffeomorphism

$$\Phi_p : B(r(p)) \rightarrow \Sigma_0 \quad \Phi_p(0) = p ,$$

with the property that the maps

$$\mathcal{H}_p = (\Phi_{H(p)})^{-1} \circ H \circ \Phi_p$$

that go from one chart to the next are *uniformly* hyperbolic with controlled second derivatives. In fact, \mathcal{H}_p is C^1 -near a linear map with diagonal entries equal to the exponentials of the Lyapunov exponents at p .

- (2) In exchange for uniform hyperbolicity, we have given up on
 - (i) uniform chart sizes: $r(\cdot)$ is measurable and can be arbitrarily near zero;
 - (ii) uniform regularity for the chart maps Φ_p .
- (3) Chart sizes can be chosen to vary slowly along orbits, with $r(Hp)/r(p) \sim 1$. This ensures the overflowing property that is crucial for establishing the existence of local stable and unstable manifolds. Distortion estimates along unstable manifolds are easily deduced from Lyapunov charts.
- (4) For $\hat{\mu}$ satisfying the condition in Proposition 5.2, it can be arranged that $\Phi_p(B(r(p))) \cap \mathcal{S} = \emptyset$ for a.e. p , so that for as long as one works within the domains of charts, one does not “see” the singularities.

6. PROOF OF SRB PROPERTY

Let ν be an ergodic component of the measure $\hat{\mu}$ defined in the last section. We assume in particular that ν possesses all the properties of $\hat{\mu}$ found in Sects. 5.2 and 5.3. In this section, we will (i) show that ν is an SRB measure for the first return map $H : \Sigma_0 \circlearrowleft$ of the flow \mathbf{F}^t ; this is carried out in Sects. 6.1 and 6.2, and (ii) build an invariant measure for \mathbf{F}^t out of ν ; this is carried out in Sect. 6.3.

6.1. Entropy of (H, ν) . Recall that λ is the positive Lyapunov exponent of (H, ν) . Our next result concerns $h_\nu(H)$, the metric entropy of H with respect to ν .

Proposition 6.1.

$$(11) \quad h_\nu(H) = \lambda.$$

To prove this proposition, recall that $H : \sigma_0 \times \mathbb{T}^2 \rightarrow \sigma_0 \times \mathbb{T}^2$ is a skew product, which we may write as

$$H(x, y; z_1, z_2) = (A(x, y); T_{(x,y)}(z_1, z_2))$$

where $\mathbb{T}_{(x,y)}^2$ is the vertical fiber over (x, y) and $T_{(x,y)} : \mathbb{T}_{(x,y)}^2 \rightarrow \mathbb{T}_{A(x,y)}^2$ is the fiber map. Let $m_{\mathbb{T}^2}$ denote Lebesgue measure on the base. Since $(\pi_{(x,y)})_* \hat{\mu} = m_{\mathbb{T}^2}$ and $(A, m_{\mathbb{T}^2})$ is ergodic, it follows that $(\pi_{(x,y)})_* \nu = m_{\mathbb{T}^2}$. Let $\{\nu_{(x,y)}\}_{(x,y) \in \mathbb{T}^2}$ be a disintegration of ν on vertical fibers, i.e.

$$d\nu(x, y; z_1, z_2) = d\nu_{(x,y)}(z_1, z_2) dm_{\mathbb{T}^2}(x, y).$$

Lemma 6.2. $\nu_{(x,y)}$ is atomic for $m_{\mathbb{T}^2}$ -a.e. (x, y) .

The idea of the proof goes back to [Kat80], who proved that if all the Lyapunov exponents of a diffeomorphism with respect to an ergodic measure are strictly negative, then the measure is supported on a periodic orbit. A fiber version of Katok's result, meaning the corresponding result for skew products when all the Lyapunov exponents of the fiber maps are strictly negative, is proved in [RW01]. Singularities aside, our setup fits this setting, as both of the exponents of $\{T_{(x,y)}\}$ are strictly negative. Our proof follows that in [RW01] nearly verbatim. The presence of singularities is immaterial because the proof uses Lyapunov charts, and for as long as one works within Lyapunov charts, the singularities of H are not “visible” by Property (ii) in Sect. 5.3.

Proof of Proposition 6.1. The assertion follows from Lemma 6.2 and a general result (see [BC92] Corollary 2 or [Kif12]) which asserts that the entropy of a skew-product map is equal to the sum of the entropy of the base and fiber entropy. (For the definition of fiber entropy, see [BC92] or [Kif12].) In our setting, denoting the fiber entropy by $h_\nu(\{T_{(x,y)}\})$, we have

$$h_\nu(H) = h_{m_{\mathbb{T}^2}}(A) + h_\nu(\{T_{(x,y)}\}).$$

Because the conditional measures on fibers are purely atomic, $h_\nu(\{T_{(x,y)}\}) = 0$. \square

6.2. Proof of SRB property for (H, ν) . The definition of SRB measure requires the almost-everywhere existence of unstable manifolds, a fact guaranteed by Proposition 6.6 for (H, μ) .

One way to build SRB measures is to push forward Lebesgue measure on a curve or disk having the dimension of E^u and roughly aligned with E^u (e.g. a piece of local unstable manifold), and to show that for large n , a positive fraction of the

pushed-forward measure accumulates on a stack of unstable manifolds of uniform length, with uniformly bounded conditional densities on unstable leaves. This is an option, but one that would have to control the lengths of the connected components of γ_n (for which techniques are well developed in the billiards literature, see [CM06]) and distortion along these curves (see Sect. 4). Another possibility, which we have chosen to adopt, is to appeal to a known result, namely converse to the entropy formula.

We recall this result, first proved for diffeomorphisms of compact manifolds (without singularities). Notations in the statement of Theorems 6.3 and 6.4 and their sketches of proofs are independent of those in the rest of this paper.

Theorem 6.3. *[[Led84],[LY85]] Let $f : M \circlearrowleft$ be a C^2 diffeomorphism of a compact Riemannian manifold M , and let θ be an f -invariant Borel probability measure. Let λ_i be the distinct Lyapunov exponents of (f, θ) and let m_i be the multiplicity of λ_i . In the setting above, if*

$$(12) \quad h_\theta(f) = \int \sum_{\lambda_i > 0} \lambda_i m_i \, d\theta ,$$

then θ is an SRB measure.

This result was first proved in [Led84] assuming $\lambda_i \neq 0$ for all i ; it was extended in [LY85] to allow zero Lyapunov exponents.

Theorem 6.4. *The setting of Theorem 6.3 can be extended to the following. Assume there is*

- (i) a set $\mathcal{S} \subset M$ that is the finite union of codimension one submanifolds,
- (ii) a C^2 -bounded map $f|_{M \setminus \mathcal{S}} : M \setminus \mathcal{S} \rightarrow M$ that is a diffeomorphism between $M \setminus \mathcal{S}$ and its image;
- (iii) an f -invariant Borel probability measure θ on $M \setminus \mathcal{S}$ with the property that for some $C > 0$, $\theta(\mathcal{S}_\varepsilon) < C\varepsilon$ for all small $\varepsilon > 0$.

Then the assertion in Theorem 6.3 continues to be valid, that is, Eq (12) implies θ is an SRB measure.

Requirement (iii) for $H : \Sigma_0 \circlearrowleft$ is provided by Proposition 5.2. The statement above is sufficient for our purposes, though the boundedness of second derivatives can be relaxed as long as f is controlled in a neighborhood of \mathcal{S} , and the measure can be more concentrated near \mathcal{S} than in Condition (iii) (see e.g. the conditions treated in [KS06]).

The proof of Theorem 6.4 is nearly identical to that of Theorem 6.3. We include in Appendix B a very brief outline to show how similar the two results are and how the presence of the singularity set is dealt with.

Corollary 6.5. *(H, ν) is an SRB measure.*

Ingredients of the proof include (i) the entropy formula (12), proved in Corollary 5.4 and Proposition 6.1; (ii) Lyapunov charts with the properties in Appendix A; and (iii) Theorem 6.4, a direct application of which gives the desired result.

6.3. SRB and physical measures for the flow. Passing of results of this type from cross-section map to flow is standard, but we include it for completeness. To distinguish between objects associated with the flow \mathbf{F}^t and those associated with the return map H , we will write $W_{\mathbf{F};\text{loc}}^u$, resp. $W_{H;\text{loc}}^u$, and so on.

Proof of Theorem 3.4. Let ν be as above, and let $T_0 : \Sigma_0 \rightarrow (0, \infty)$ be the return time from the cross-section Σ_0 to itself under the flow \mathbf{F}^t . We let $\nu_{\mathbf{F}}$ be the normalization of the pushforward of ν up to return time, i.e.,

$$\nu_{\mathbf{F}} := \frac{1}{\int T_0 d\nu} \int_0^\infty \mathbf{F}_*^t(\nu|_{\{t < T_0\}}) dt$$

where $\nu|_{\{t < T_0\}}(X) = \nu(X \cap \{t < T_0\})$ for any measurable set $X \subset \Sigma_0$. Then $\nu_{\mathbf{F}}$ is clearly an \mathbf{F}^t -invariant Borel probability measure on M .

To prove the SRB property of $\nu_{\mathbf{F}}$, we need to show that its disintegration on local weak unstable manifolds $W_{\mathbf{F};\text{loc}}^u$ of \mathbf{F}^t have conditional densities. For $q \in M$ let $p \in \Sigma_0$ be such that $q = \mathbf{F}^{t_q}(p)$ for $0 < t_q < T_0(p)$. Then, for $\nu_{\mathbf{F}}$ -a.e. $q \in M$ there exists r such that

$$W_{\mathbf{F};\text{loc}}^u(q) = \bigcup_{\{t: |t-t_q| < r\}} \mathbf{F}^t(W_{H;\text{loc}}^u(p))$$

is a piece of local weak unstable manifold of the flow. From the definition of $\nu_{\mathbf{F}}$, conditional probabilities on these objects are clearly equivalent to $m_{W_{\mathbf{F};\text{loc}}^u}$, the Lebesgue measure on $W_{\mathbf{F};\text{loc}}^u$. \square

We say a point $q \in M$ is *future-generic* with respect to $\nu_{\mathbf{F}}$ if for all continuous observables $h : M \rightarrow \mathbb{R}$,

$$\frac{1}{T} \int_0^T h(\mathbf{F}^t(q)) dt \rightarrow \int h d\nu_{\mathbf{F}} \quad \text{as } T \rightarrow \infty .$$

The invariant measure $\nu_{\mathbf{F}}$ is a physical measure if the set of points future-generic with respect to $\nu_{\mathbf{F}}$ has positive Lebesgue measure (m) on M .

Proof of Theorem 3.2. Let $W = W_{\mathbf{F};\text{loc}}^u(q)$, and assume m_W -a.e. $q' \in W$ is future-generic wrt $\nu_{\mathbf{F}}$. Defining $W_{\mathbf{F};\text{loc}}^{ss}(q')$ to be the strong stable manifold at q' and letting

$$(13) \quad \Lambda(q) := \bigcup_{q' \in W} W_{\mathbf{F};\text{loc}}^{ss}(q') ,$$

we have that $m(\Lambda(q)) > 0$ by the absolute continuity of the $W_{\mathbf{F};\text{loc}}^{ss}$ -foliation. Since $d(\mathbf{F}^t(q'), \mathbf{F}^t(q'')) \rightarrow 0$ exponentially fast for $q'' \in W_{\mathbf{F};\text{loc}}^{ss}(q')$, q'' is future-generic wrt $\nu_{\mathbf{F}}$ when q' is. This proves that m -a.e. $q'' \in \Lambda(q)$ is future-generic wrt $\nu_{\mathbf{F}}$. \square

The structures here are in fact so simple one does not need to invoke the absolute continuity of $W_{\mathbf{F};\text{loc}}^{ss}$. We claim that for $q' \in W$ in (13), $W_{\mathbf{F};\text{loc}}^{ss}(q') \subset \{w = \text{const}\}$. To see this, let $p \in \Sigma_0$, and consider $W_{H;\text{loc}}^s(p)$, the 3D local stable manifold for H at p . Then $W_{H;\text{loc}}^s(p) \subset \Sigma_0$, and because the return time T_0 is locally constant,

$$\mathbf{F}^{T_0(p)}(W_{H;\text{loc}}^s(p)) \subset W_{H;\text{loc}}^s(Hp) .$$

Letting $T_n(p) = T_0(p) + T_0(Hp) + \dots + \dots + T_0(H^{n-1}p)$, the same argument gives

$$\mathbf{F}^{T_n(p)}(W_{H;\text{loc}}^s(p)) \subset W_{H;\text{loc}}^s(H^n p)$$

for all $n \in \mathbb{Z}^+$. This implies that $W_{\mathbf{F};\text{loc}}^{ss}(p) \subset \{w = 0\}$, and the claim follows.

APPENDIX

A. Lyapunov charts and related results

First some notation: We fix a number $\lambda_0 < \min\{\lambda, -\lambda_{\min}\}$, and small numbers $0 < \delta_1, \delta_2 \ll \lambda_0$. The domains of Lyapunov charts are subsets of $\mathbb{R} \times \mathbb{R}^3$, with norm $|(v, w)|' := \max\{|v|, |w|\}$ where $|\cdot|$ denotes Euclidean norm on \mathbb{R} or \mathbb{R}^3 , and $B(r) := \{q \in \mathbb{R} \times \mathbb{R}^3 : |q|' \leq r\}$. Norms on Σ_0 are denoted by $\|\cdot\|$ as before. We first state – without proof – their properties, postponing explanation for some aspects to the end.

- On a set $\Gamma \subset \Sigma_0$ of full $\hat{\mu}$ -measure are defined

(a) a measurable family of linear maps $L : \Gamma \rightarrow \mathcal{L}(\mathbb{R} \times \mathbb{R}^3, T_p\Sigma_0)$ with

$$L(p)(\mathbb{R} \times \{0\}) = E^u(p), \quad L(p)(\{0\} \times \mathbb{R}^3) = E^s(p),$$

(b) a measurable function $\ell : \Gamma \rightarrow [0, \infty)$ with

$$(14) \quad e^{-\delta_1} \ell(p) \leq \ell(H(p)) \leq e^{\delta_1} \ell(p) \quad \text{for a.e. } p.$$

- The Lyapunov chart at p is given by

$$\Phi_p : B(\delta_2 \ell(p)^{-1}) \rightarrow \Sigma_0, \quad \Phi_p = L(p)|_{B(\delta_2 \ell(p)^{-1})}.$$

Here we have identified neighborhoods of 0 in $T_p\Sigma_0$ with neighborhoods of p in Σ_0 via the exponential map. Connecting maps between charts

$$\mathcal{H}_p : B(\delta_2 \ell(p)^{-1}) \rightarrow \mathbb{R} \times \mathbb{R}^3 \quad \text{are} \quad \mathcal{H}_p = (\Phi_{H(p)})^{-1} \circ H \circ \Phi_p.$$

- The linear maps $L(p)$, hence $(D\Phi_p)_0$ and $(D\mathcal{H}_p)_0$, are designed to produce the following one-step hyperbolicity:

$$(i) \quad \begin{aligned} &\text{for } v \in \mathbb{R} \times \{0\}, |D(\mathcal{H}_p)_0(v)| \geq e^{\lambda_0} |v|, \\ &\text{for } v \in \{0\} \times \mathbb{R}^3, |D(\mathcal{H}_p)_0(v)| \leq e^{-\lambda_0} |v|. \end{aligned}$$

- The restriction of Φ_p to $B(\delta_2 \ell(p)^{-1})$ ensures the following:

- (ii) $\Phi_p(B(\delta_2 \ell(p)^{-1})) \cap \mathcal{S} = \emptyset$;
- (iii) for all $q, q' \in B(\delta_2 \ell(p)^{-1})$,

$$\|\Phi_p(q) - \Phi_p(q')\| \leq |q - q'|' \leq \ell(p) \|\Phi_p(q) - \Phi_p(q')\|;$$

- (iv) $\text{Lip}(\mathcal{H}_p - (D\mathcal{H}_p)_0) \leq \delta_2$, and
- (v) $\text{Lip}(D\mathcal{H}_p) \leq \ell(p)$.

Property (ii) and the slowly varying property of ℓ

Property (ii) and property (14) of ℓ are used to ensure the overflowing condition needed in the proof of local unstable manifolds for the connecting maps \mathcal{H}_p . Charts for maps with singularities were treated in [KS06], but since the setting of [KS06] is more complicated than the one here and this part of the theory is less standard we review the main ideas on how to arrange for (14) and property (ii).

Given a measurable function $\ell_1 : \Gamma \rightarrow [1, \infty)$, we construct ℓ with the property in (14) by letting

$$(15) \quad \ell(p) = \sup_{n \in \mathbb{Z}} e^{-\delta_1 |n|} \ell_1(H^n p)$$

provided the right side is finite for a.e. p . Let us assume, without proof, that a function ℓ_0 satisfying

$$e^{-\delta_1} \ell_0(p) \leq \ell_0(H(p)) \leq e^{\delta_1} \ell_0(p)$$

and all the properties above except for Item (ii) have been constructed. Let

$$\ell_1(p) = \max\{\ell_0(p), d(p, \mathcal{S})^{-1}\}$$

where $d(p, \mathcal{S})$ is distance to the singularity set, and define $\ell(p)$ as in (15). To check the finiteness of the right side, we observe that

$$\ell_1(H^n p) \leq \max\{\ell_0(H^n(p)), e^{-\delta_1|n|} d(H^n p, \mathcal{S})^{-1}\},$$

and the quantities $e^{-\delta_1|n|} d(H^n p, \mathcal{S})^{-1}$, $n \in \mathbb{Z}$, are uniformly bounded because

$$\sum_{n=0}^{\infty} \hat{\mu}(\mathcal{S}_{e^{-n\delta_1/2}}) < \infty$$

by Proposition 5.2, so by the Borel-Cantelli Lemma, for $\hat{\mu}$ -a.e. p , there is $n(p)$ such that $H^{|n|}(p) \notin \mathcal{S}_{e^{-n\delta_1/2}}$ for all $n \geq n(p)$.

It remains to check that ℓ so defined satisfies Item (ii): Since $\ell(p) \geq d(p, \mathcal{S})^{-1}$, it follows that for all $p' \in \Phi_p(B(\delta_2 \ell(p)^{-1}))$, $\|p - p'\| \leq \delta_2 \ell(p)^{-1} \leq \delta_2 d(p, \mathcal{S})$.

Local unstable manifolds and distortion

The following results gleaned from Lyapunov charts are central to the definition of SRB measures. We state them without proof as they are standard; see the references above. Below we write the domain of charts as $B(r) = B^u(r) \times B^s(r)$ where $B^u(r) = \{v \in \mathbb{R} : |v| < r\}$ and $B^s(r) = \{v \in \mathbb{R}^3 : |v| < r\}$.

Proposition 6.6. *For $\delta_2 > 0$ small enough, there are constants D_ℓ (depending on ℓ) with respect to which the following holds at $\hat{\mu}$ -a.e. p :*

(a) (Existence of local unstable manifolds) *There is a C^2 function*

$$\beta_p^u : B^u(\delta_2 \ell(p)^{-1}) \rightarrow B^s(\delta_2 \ell(p)^{-1})$$

with the property that

- $\beta_p^u(0) = 0$, $D\beta_p^u(0) = 0$ and $\|D\beta_p^u\| < \frac{1}{10}$;
- $(\mathcal{H}_{H^{-1}p})^{-1} \text{graph}(\beta_p^u) \subset \text{graph}(\beta_{H^{-1}p}^u)$
- for all $q_1, q_2 \in \text{graph}(\beta_p^u)$, $|(\mathcal{H}_{H^{-1}p})^{-1}q_1 - (\mathcal{H}_{H^{-1}p})^{-1}q_2|' < e^{-\lambda_0}|q_1 - q_2|'$.

An analogous statement holds for local stable manifolds.

(b) (Distortion estimate) *For $q \in \text{graph}(\beta_p^u)$, let $E^u(q)$ denote the tangent space of $\text{graph}(\beta_p^u)$ at q . Then for all $q_1, q_2 \in \text{graph}(\beta_p^u)$ and all $n \geq 1$,*

$$\left| \log \frac{D\mathcal{H}_{H^{-n}p}^n|_{E^u(q_1)}}{D\mathcal{H}_{H^{-n}p}^n|_{E^u(q_2)}} \right| \leq D_{\ell(p)} |q_1 - q_2|.$$

Moreover, the function

$$\Delta_p(q) := \lim_{n \rightarrow \infty} \frac{D\mathcal{H}_{H^{-n}p}^n|_{E^u(p)}}{D\mathcal{H}_{H^{-n}p}^n|_{E^u(q)}}$$

is Lipschitz-continuous and bounded from above and below.

We define the Φ_p -image of $\text{graph}(\beta_p^u)$, denoted $W_{\text{loc}}^u(p)$, as the *local unstable manifold* at p . Local stable and unstable manifolds vary in size and can be arbitrarily small in diameter, but for p in *uniformity sets*, i.e., sets of the form $\Gamma_{\bar{\ell}} := \{\ell \leq \bar{\ell}\}$ where $\bar{\ell}$ is a fixed number, they contain disks of fixed radii depending on $\bar{\ell}$.

B. Outline of proof of Theorem 6.4 following [Led84]

Let (f, θ) be as in the statement of the theorem. We construct Lyapunov charts with the properties in Appendix A. Note in particular Item (ii), which ensures that the images of charts do not meet \mathcal{S} . This, we claim, is all that is needed to ensure that the argument in [Led84] will go through. We summarize very briefly this argument to give some idea of what it entails:

Step 1. One constructs a measurable partition η with the properties that

- (i) it is subordinate to unstable manifolds, i.e., for ν -a.e. p , $\eta(p) \subset W^u(p)$ and contains a neighborhood of p in $W^u(p)$, and
 - (ii) η is a Markov partition, i.e., $\eta \leq f^{-1}\eta$,
- and shows that $h_\theta(f) = h_\theta(f, \eta)$ via an auxiliary finite-entropy partition (see also [LS82]).

The partition η is constructed by taking a stack of local unstable disks $\{D_\alpha^u\}$ through points in a uniformity set Γ_{ℓ_0} (as defined at the end of Sect. 5.3), iterating forward and taking intersections.

Step 2. Consider the quotient M/η . Let θ_T be the quotient measure on this space and $\{\theta_\eta\}$ a family of conditional probabilities on elements of η . We introduce a new measure Θ so that $\Theta_T = \theta_T$ and for θ -a.e. p , Θ_η on $\eta(p)$ has a density equal to $\Delta_p(\cdot)$ normalized where $\Delta_p(\cdot)$ is as defined in Proposition 6.6(b). One then proves, using the equality in (12) and an argument relying on the convexity of $-\log$, that $\Theta = \theta$, so θ is an SRB measure.

As can be seen from the outline above, all the structures involved in the proof originate from within Lyapunov charts: The D_α^u are local unstable manifolds obtained from charts. The putative conditional densities $\Delta_p(\cdot)$ are defined on elements of η (which can be much larger than charts), but for θ -a.e. p , $f^{-n}(\eta(p)) \subset W_{\text{loc}}^u(q)$ for some $q \in \Gamma_{\ell_0}$, so again it suffices to have distortion estimates for local manifolds within charts as in Proposition 6.6(b). In particular, once property (ii) in Sect. 5.3 is ensured, the singularity set does not appear in these constructions, except to render the elements of η more cut up, but that is immaterial.

REFERENCES

- [Ano67] Dmitry Victorovich Anosov, *Geodesic flows on closed Riemannian manifolds of negative curvature*, Trudy Matematicheskogo Instituta Imeni VA Steklova **90** (1967), 3–210.
- [BB66] John Buck and Elisabeth Buck, *Biology of synchronous flashing of fireflies*, Nature **211** (1966), no. 5049, 562–564.
- [BC85] Michael Benedicks and Lennart Carleson, *On iterations of $1 - ax^2$ on $(-1, 1)$* , Annals of Mathematics **122** (1985), 1–25.
- [BC91] ———, *The dynamics of the Hénon map*, Annals of Mathematics **133** (1991), no. 1, 73–169.
- [BC92] Thomas Bogenschütz and Hans Crauel, *The Abramov-Rokhlin formula*, Ergodic Theory and Related Topics III, Springer, 1992, pp. 32–35.

- [BK96] Jean Bricmont and Antti Kupiainen, *High temperature expansions and dynamical systems*, Communications in Mathematical Physics **178** (1996), no. 3, 703–732.
- [BMBY15] Nathan Breitsch, Gregory Moses, Erik Boczko, and Todd Young, *Cell cycle dynamics: clustering is universal in negative feedback systems*, Journal of mathematical biology **70** (2015), no. 5, 1151–1175.
- [BS88] Leonid A. Bunimovich and Yakov G. Sinai, *Spacetime chaos in coupled map lattices*, Nonlinearity **1** (1988), no. 4, 491.
- [BSC90] Leonid A. Bunimovich, Yakov G. Sinai, and Nikolai Ivanovich Chernov, *Markov partitions for two-dimensional hyperbolic billiards*, Russian Mathematical Surveys **45** (1990), no. 3, 105.
- [BTLT06] Frede Blaabjerg, Remus Teodorescu, Marco Liserre, and Adrian V Timbus, *Overview of control and grid synchronization for distributed power generation systems*, IEEE Transactions on industrial electronics **53** (2006), no. 5, 1398–1409.
- [Bun74] Leonid A. Bunimovich, *On ergodic properties of certain billiards*, Funktsional. Anal. i Prilozhen **8** (1974), no. 3, 73–74.
- [BY17] Alex Blumenthal and Lai-Sang Young, *Entropy, volume growth and srb measures for banach space mappings*, Inventiones mathematicae **207** (2017), no. 2, 833–893.
- [CE80] Pierre Collet and J-P Eckmann, *On the abundance of aperiodic behaviour for maps on the interval*, Communications in Mathematical Physics **73** (1980), no. 2, 115–160.
- [CF05] Jean-René Chazottes and Bastien Fernandez, *Dynamics of coupled map lattices and of related spatially extended systems*, vol. 671, Springer Science & Business Media, 2005.
- [CM06] Nikolai Chernov and Roberto Markarian, *Chaotic billiards*, no. 127, American Mathematical Soc., 2006.
- [ER85] J-P Eckmann and David Ruelle, *Ergodic theory of chaos and strange attractors*, The theory of chaotic attractors, Springer, 1985, pp. 273–312.
- [FT14] Bastien Fernandez and Lev S Tsimring, *Typical trajectories of coupled degrade-and-fire oscillators: from dispersed populations to massive clustering*, Journal of mathematical biology **68** (2014), no. 7, 1627–1652.
- [GW79] John Guckenheimer and Robert F Williams, *Structural stability of lorenz attractors*, Publications Mathématiques de l’Institut des Hautes Études Scientifiques **50** (1979), no. 1, 59–72.
- [Hop39] E Hopf, *Statistik der geoddtischen Linien in Mannigfaltigkeiten negativer Krümmung*, Ber Verh. Sachs. Akad. Wiss. Leipzig. Math.-Nat. Kl. **51** (1939), 261–304.
- [Jak81] Michael V Jakobson, *Absolutely continuous invariant measures for one-parameter families of one-dimensional maps*, Communications in Mathematical Physics **81** (1981), no. 1, 39–88.
- [Kan93] Kunihiko Kaneko, *Theory and applications of coupled map lattices*, John Wiley & Sons, 1993.
- [Kat80] Anatole Katok, *Lyapunov exponents, entropy and periodic orbits for diffeomorphisms*, Publications Mathématiques de l’Institut des Hautes Études Scientifiques **51** (1980), no. 1, 137–173.

- [Kif12] Yuri Kifer, *Ergodic theory of random transformations*, vol. 10, Springer Science & Business Media, 2012.
- [KL06] Gerhard Keller and Carlangelo Liverani, *Uniqueness of the SRB measure for piecewise expanding weakly coupled map lattices in any dimension*, *Communications in Mathematical Physics* **262** (2006), no. 1, 33–50.
- [KL09] ———, *Map lattices coupled by collisions*, *Communications in Mathematical Physics* **291** (2009), no. 2, 591–597.
- [KS69] K. Krzyżewski and W. Szlenk, *On invariant measures for expanding differentiable mappings*, *Studia Mathematica* **33** (1969), no. 1, 83–92.
- [KS06] Anatole Katok and Jean-Marie Strelcyn, *Invariant manifolds, entropy and billiards. smooth maps with singularities*, vol. 1222, Springer, 2006.
- [Kur03] Yoshiki Kuramoto, *Chemical oscillations, waves, and turbulence*, Courier Corporation, 2003.
- [KY07] Elisha Kobre and Lai-Sang Young, *Extended systems with deterministic local dynamics and random jumps*, *Communications in Mathematical Physics* **275** (2007), no. 3, 709–720.
- [KY10] José Koiller and Lai-Sang Young, *Coupled map networks*, *Nonlinearity* **23** (2010), no. 5, 1121.
- [Led84] François Ledrappier, *Propriétés ergodiques des mesures de Sinai*, *Publications Mathématiques de l’IHÉS* **59** (1984), 163–188.
- [LKYJW20] Wei-Hsiang Lin, Edo Kussell, Lai-Sang Young, and Christine Jacobs-Wagner, *Origin of exponential growth in nonlinear reaction networks*, *Proceedings of the National Academy of Sciences* **117** (2020), no. 45, 27795–27804.
- [Lor62] Edward N Lorenz, *The statistical prediction of solutions of dynamical equations*, *Proceedings of the International Symposium on Numerical Weather Prediction, 1962*, Meteor. Soc. Japan, 1962.
- [LS82] François Ledrappier and Jean-Marie Strelcyn, *A proof of the estimation from below in Pesin’s entropy formula*, *Ergodic Theory and Dynamical Systems* **2** (1982), no. 2, 203–219.
- [LY85] François Ledrappier and L-S Young, *The metric entropy of diffeomorphisms: Part I: Characterization of measures satisfying Pesin’s entropy formula*, *Annals of Mathematics* (1985), 509–539.
- [MS08] Alexandre Mauroy and Rodolphe Sepulchre, *Clustering behaviors in networks of integrate-and-fire oscillators*, *Chaos: An interdisciplinary journal of nonlinear science* **18** (2008), no. 3, 037122.
- [Pes76] Ja B Pesin, *Families of invariant manifolds corresponding to nonzero characteristic exponents*, *Mathematics of the USSR-Izvestiya* **10** (1976), no. 6, 1261.
- [PvST20] Tiago Pereira, Sebastian van Strien, and Matteo Tanzi, *Heterogeneously coupled maps: hub dynamics and emergence across connectivity layers*, *Journal of the European Mathematical Society* **22** (2020), 2183–2252.
- [RW01] David Ruelle and Amie Wilkinson, *Absolutely singular dynamical foliations*, *Communications in Mathematical Physics* **219** (2001), no. 3, 481–487.

- [SB16] Fanni Sélley and Péter Bálint, *Mean-field coupling of identical expanding circle maps*, Journal of Statistical Physics **164** (2016), no. 4, 858–889.
- [Sin70] Yakov G Sinai, *Dynamical systems with elastic reflections*, Russian Mathematical Surveys **25** (1970), 137–91.
- [SL77] Herbert Spohn and Joel L Lebowitz, *Stationary non-equilibrium states of infinite harmonic systems*, Communications in Mathematical Physics **54** (1977), no. 2, 97–120.
- [ST21] Fanni M Sélley and Matteo Tanzi, *Linear response for a family of self-consistent transfer operators*, Communications in Mathematical Physics **382** (2021), no. 3, 1601–1624.
- [TW82] Roger D Traub and RK Wong, *Cellular mechanism of neuronal synchronization in epilepsy*, Science **216** (1982), no. 4547, 745–747.
- [WM15] Dan Wilson and Jeff Moehlis, *Clustered desynchronization from high-frequency deep brain stimulation*, PLoS computational biology **11** (2015), no. 12, e1004673.
- [WY08] Qiudong Wang and Lai-Sang Young, *Toward a theory of rank one attractors*, Annals of Mathematics (2008), 349–480.
- [YB93] Lai-Sang Young and Michael Benedicks, *Sinai-Bowen-Ruelle measures for certain Hénon maps*, Inventiones mathematicae **112** (1993), no. 3, 541–576.
- [YFB⁺12] Todd R Young, Bastien Fernandez, Richard Buckalew, Gregory Moses, and Erik M Boczko, *Clustering in cell cycle dynamics with general response/signaling feedback*, Journal of theoretical biology **292** (2012), 103–115.
- [You95] Lai-Sang Young, *Ergodic theory of differentiable dynamical systems*, ed. branner and hjorth ed., NATO ASI series, vol. Real and Complex Dynamics, pp. 293–336, Kluwer Academic Publishers, 1995.

MATTEO TANZI: COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, NEW YORK, NY 10012, USA

Email address: `matteo.tanzi@nyu.edu`

LAI-SANG YOUNG: COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, NEW YORK, NY 10012, USA, AND INSTITUTE FOR ADVANCED STUDY, PRINCETON, NEW JERSEY 08540, USA

Email address: `lsy@cims.nyu.edu`