

ON THE COEFFICIENTS IN AN ASYMPTOTIC EXPANSION OF $(1 + 1/x)^x$

T. M. DUNSTER* AND JESSICA M. PEREZ†

Abstract. The function $g(x) = (1+1/x)^x$ has the well-known limit e as $x \rightarrow \infty$. The coefficients c_j in an asymptotic expansion for $g(x)$ are considered. A simple recursion formula is derived, and then using Cauchy's integral formula the coefficients are approximated for large j . From this it is shown that $|c_j| \rightarrow 1$ as $j \rightarrow \infty$.

Key words. Series expansions, Asymptotic representations in the complex plane, Integrals of Cauchy type

AMS subject classifications. 41A58, 30E15, 30E20

1. Introduction and main results. Chen and Choi [3] presented a method to estimate Euler's constant e , accurate to as many decimal places as desired. Their starting point was the well known limit $\lim_{x \rightarrow \infty} g(x) = e$, where

$$(1.1) \quad g(x) = \left(1 + \frac{1}{x}\right)^x.$$

Their method was based on an asymptotic expansion of $g(x)$ for large values of x , namely

$$(1.2) \quad g(x) \sim \sum_{j=0}^{\infty} \frac{c_j}{x^j}.$$

We shall show that this asymptotic expansion converges for $x > 1$.

In [3] they proved that $c_j = ea_j$, where a_j are rational numbers which alternate in sign, and are given explicitly by

$$(1.3) \quad a_j = (-1)^j \sum \frac{1}{k_1! k_2! \cdots k_j!} \left(\frac{1}{2}\right)^{k_1} \left(\frac{1}{3}\right)^{k_2} \cdots \left(\frac{1}{j+1}\right)^{k_j}.$$

Here for each j the sum is taken over all possible combinations of nonnegative integers $k_1, k_2, k_3, \dots, k_j$ that satisfy the relation $\sum_{l=1}^j l k_l = j$. We remark that Ponomorenko [7] gave a much simpler proof of (1.3) using Faà di Bruno's formula generalizing the chain rule to higher derivatives.

The number of terms in the sum (1.3) is the partition function $P(j)$ [4, Sect. 27.14(i)]. Hardy and Ramanujan [5] gave the asymptotic formula

$$(1.4) \quad P(j) \sim \frac{\exp(\pi\sqrt{2j/3})}{4j\sqrt{3}}$$

as $j \rightarrow \infty$. It is therefore evident that the number of terms in the formula (1.3) grows exponentially in \sqrt{j} , and so is only practicable for small or moderate values of j .

*Department of Mathematics and Statistics, San Diego State University, 5500 Campanile Drive, San Diego, CA 92182, USA. (mdunster@sdsu.edu, <https://tmdunster.sdsu.edu>).

†Department of Mathematics and Statistics, San Diego State University, 5500 Campanile Drive, San Diego, CA 92182, USA. (jessica.perez.aof@gmail.com).

We derive a new way of computing the coefficients c_j , by a simple recursion formula. We also provide a simple asymptotic approximation for the coefficients as $j \rightarrow \infty$, and this shows that in absolute value they approach the value 1.

Our main results read as follows. In [section 2](#) we prove [\(1.5\)](#), and in [section 3](#) we prove [\(1.6\)](#).

THEOREM 1.1. *For $x > 1$ the expansion [\(1.2\)](#) converges, where the coefficients c_j are given recursively by $c_0 = e$ and*

$$(1.5) \quad c_{j+1} = \frac{1}{j+1} \sum_{l=0}^j (-1)^{j-l+1} \binom{j-l+1}{j-l+2} c_l \quad (j = 0, 1, 2, \dots).$$

Moreover as $j \rightarrow \infty$

$$(1.6) \quad c_j = (-1)^j \left(1 + \frac{1}{j} \right) + \mathcal{O} \left(\frac{\ln(j)}{j^2} \right).$$

REMARK 1. *Since $\lim_{x \rightarrow \infty} g(x) = e$ it is clear from [\(1.2\)](#) that $c_0 = e$.*

2. Proof of the recursion formula [\(1.5\)](#). Define $z = 1/x$ and $f(z) = g(1/z)$ so that [\(1.2\)](#) is written in the new form

$$(2.1) \quad f(z) = (1+z)^{1/z} = \sum_{j=0}^{\infty} c_j z^j.$$

and in this we consider z complex. Moreover, on writing it as

$$(2.2) \quad f(z) = \exp \{ z^{-1} \ln(1+z) \},$$

we note that it has a removable singularity at $z = 0$, and therefore can be considered analytic at $z = 0$ by assuming $f(0) = \lim_{z \rightarrow 0} f(z) = c_0 = e$. We also see it has one finite singularity (a logarithmic branch point) at $z = -1$. Therefore the radius of convergence of the series [\(2.1\)](#) is 1, i.e. it converges for $|z| < 1$. Thus [\(1.2\)](#) converges for $x > 1$, as asserted. By taking the principal branch of the logarithm in [\(2.2\)](#) we have that $f(z)$ is analytic on the cut plane $\mathbb{C} \setminus (-\infty, -1]$.

Next, on taking the natural logarithm of both sides of [\(2.1\)](#), we get

$$(2.3) \quad \frac{1}{z} \ln(1+z) = \ln \left\{ \sum_{j=0}^{\infty} c_j z^j \right\}.$$

Then expand the $\ln(1+z)$ term by its Maclaurin expansion valid for $|z| < 1$ and we arrive at

$$(2.4) \quad \sum_{j=1}^{\infty} (-1)^j \frac{z^j}{j+1} = \ln \left\{ \sum_{j=0}^{\infty} c_j z^j \right\}.$$

Next differentiate both sides with respect to z to yield

$$(2.5) \quad \sum_{j=1}^{\infty} (-1)^j \frac{j z^{j-1}}{j+1} = \sum_{j=1}^{\infty} j c_j z^{j-1} \left[\sum_{j=0}^{\infty} c_j z^{j-1} \right]^{-1}.$$

By shifting indices of the series starting at $j = 1$ to start at $j = 0$, and taking the series $\sum_{j=0}^{\infty} c_j z^j$ to the left-hand side, we see this is equivalent to

$$(2.6) \quad \sum_{j=0}^{\infty} c_j z^j \sum_{j=0}^{\infty} d_j z^j = \sum_{j=0}^{\infty} (j+1) c_{j+1} z^j,$$

where d_j is given by

$$(2.7) \quad d_j = (-1)^{j+1} \frac{j+1}{j+2}.$$

We now use the Cauchy product, which is the discrete convolution of two infinite series. It is given by the formula [2, Sect. 73]

$$(2.8) \quad \sum_{j=0}^{\infty} C_j \sum_{j=0}^{\infty} D_j = \sum_{j=0}^{\infty} \left[\sum_{l=0}^j C_l D_{j-l} \right].$$

Applying this to the left-hand side of (2.6) we combine both power series to the following single power series

$$(2.9) \quad \sum_{j=0}^{\infty} c_j z^j \sum_{j=0}^{\infty} d_j z^j = \sum_{j=0}^{\infty} \left[\sum_{l=0}^j c_l d_{j-l} \right] z^j.$$

Finally substitute this into the left-hand side of (2.6), and equate coefficients of z^j , to obtain

$$(2.10) \quad \sum_{l=0}^j c_l d_{j-l} = (j+1) c_{j+1},$$

and then using (2.7) this leads to (1.5).

3. Proof of the asymptotic approximation (1.6). We will use the famous Cauchy integral formula [4, Eq. 1.9.31] to obtain an integral representation for the coefficients c_j . If C_r is the positively orientated circle $\{z : |z| = r\}$ for arbitrary $r \in (0, 1)$ then from (2.1) and (2.2)

$$(3.1) \quad c_j = \frac{1}{2\pi i} \oint_{C_r} \frac{(1+z)^{1/z}}{z^{j+1}} dz = \frac{1}{2\pi i} \oint_{C_r} \frac{\exp\{z^{-1} \ln(1+z)\}}{z^{j+1}} dz.$$

In the second integral of (3.1) we rewrite the exponential term using the geometric series

$$(3.2) \quad \frac{1}{z} = -\frac{1}{1-(1+z)} = -(1 + \delta + \delta^2 + \delta^3 + \dots),$$

where $\delta = 1+z$, assuming $0 < |\delta| < 1$. So from (2.2)

$$(3.3) \quad f(z) = \exp \left\{ -\ln(\delta) \left(1 + \sum_{j=1}^{\infty} \delta^j \right) \right\} = \frac{1}{\delta} \exp \left\{ -\ln(\delta) \sum_{j=1}^{\infty} \delta^j \right\}.$$

Note that $\delta^j \ln(\delta) \rightarrow 0$ as $\delta \rightarrow 0$ for $j = 1, 2, 3, \dots$ by L'Hopital's rule. So using the Maclaurin expansion of the exponential function along with (3.3) this function has the expansion

$$(3.4) \quad f(z) = \frac{1}{\delta} \left(1 - v + \frac{v^2}{2!} - \frac{v^3}{3!} + \dots \right),$$

for $0 < |\delta| < 1$, where $v = \ln(\delta) \sum_{j=1}^{\infty} \delta^j$. From this one deduces for small δ that

$$(3.5) \quad v = -\delta \ln(\delta) - \delta^2 \ln(\delta) + \mathcal{O}\{\delta^3 \ln(\delta)\}, \quad v^2 = \delta^2 \ln^2(\delta) + \mathcal{O}\{\delta^3 \ln^2(\delta)\},$$

and

$$(3.6) \quad v^j = \mathcal{O}\{\delta^3 \ln^3(\delta)\} \quad (j = 3, 4, 5, \dots).$$

Recalling $\delta = 1 + z$ we consequently have from (3.4) - (3.6)

$$(3.7) \quad f(z) = (1+z)^{-1} - \ln(1+z) + R(z),$$

where

$$(3.8) \quad R(z) = (1+z)^{1/z} - (1+z)^{-1} + \ln(1+z) \\ = \left[\frac{1}{2} \ln(1+z)^2 - \ln(1+z) \right] (1+z) + \mathcal{O}\left\{ \ln(1+z)^3 (1+z)^2 \right\},$$

as $z \rightarrow -1$.

Now substitute (3.7) into (3.1) to get

$$(3.9) \quad c_j = I_{1,j} + I_{2,j} + \eta_j,$$

where

$$(3.10) \quad I_{1,j} = \frac{1}{2\pi i} \oint_{C_r} \frac{1}{z^{j+1}(1+z)} dz, \quad I_{2,j} = -\frac{1}{2\pi i} \oint_{C_r} \frac{\ln(1+z)}{z^{j+1}} dz,$$

and

$$(3.11) \quad \eta_j = \frac{1}{2\pi i} \oint_{C_r} \frac{R(z)}{z^{j+1}} dz.$$

The integrals in (3.10) can readily be evaluated by residue theory. For the first we have by the geometric series expansion


$$(3.12) \quad I_{1,j} = \operatorname{Res}_{z=0} \left\{ \frac{1}{z^{j+1}(z+1)} \right\} = \operatorname{Res}_{z=0} \left\{ \sum_{s=0}^{\infty} (-1)^s z^{s-j-1} \right\} = (-1)^j.$$

Likewise for $I_{2,j}$ one finds that

$$(3.13) \quad I_{2,j} = -\operatorname{Res}_{z=0} \left\{ \frac{\ln(1+z)}{z^{j+1}} \right\} = \frac{(-1)^j}{j}.$$

For the integral (3.11) we make a change of variable $w = -(z+1)$ to obtain the following

$$(3.14) \quad \eta_j = \frac{(-1)^{j+1}}{2\pi i} \oint_{C_r} \frac{R(-1-w)}{(1+w)^{j+1}} dw.$$

FIG. 1. Contour $\Gamma_{\epsilon,\rho}$

The contour C'_r in the w plane is now the circle $\{w : |w + 1| = r\}$ for $0 < r < 1$, and is positively orientated. This lies in the left half plane and encircles $w = -1$. The integrand of (3.14) has a branch point at $w = 0$ and a pole at $w = -1$, and is analytic elsewhere in the w plane having a cut along the non-negative real axis. So we can deform the contour to a new one, called $\Gamma_{\epsilon,\rho}$, as seen in Figure 1.

This contour consists of circles γ_ϵ and γ_ρ centered at $w = 0$, of radius ϵ and ρ (respectively), where $0 < \epsilon < 1 < \rho$, and horizontal line segments l_1 and l_2 with end points $w = \epsilon \pm i0$ and $w = \rho \pm i0$ above and below the cut.

Now from (3.8) we see that the integrand of (3.14) is $\mathcal{O}\{w \ln(w)^2\}$ as $w \rightarrow 0$, and $\mathcal{O}\{w^{-j-1} \ln(w)\}$ as $w \rightarrow \infty$. Hence the contributions of γ_ϵ and γ_ρ vanish as $\epsilon \rightarrow 0$ and $\rho \rightarrow \infty$. Then the only contribution will be along $l_1 \cup l_2$, where now these lines extend from 0 to ∞ . We are therefore left with

$$(3.15) \quad \eta_j = \frac{(-1)^{j+1}}{2\pi i} \int_{l_1 \cup l_2} \frac{R(-1-w)}{(1+w)^{j+1}} dw.$$

Next, the contributions of real terms in the integrand of (3.15) cancel. Hence, using $\Im\{\ln(z)^2\} = 2 \arg(z) \ln(|z|)$ ($z \in \mathbb{C} \setminus \{0\}$), we have from (3.8) $\Im\{R(-1-w)\} =$

$\mathcal{O}\{w \ln(w)\}$ uniformly for $w \in l_1 \cup l_2$, and so from (3.15) for unbounded j

$$(3.16) \quad \eta_j = \mathcal{O} \left\{ \int_0^\infty \frac{w \ln(w)}{(1+w)^{j+1}} dw \right\}.$$

We then let $1+w = e^t$, and consequently from (3.16) obtain

$$(3.17) \quad \eta_j = \mathcal{O} \left\{ \int_0^\infty e^{-jt}(e^t - 1) \ln(e^t - 1) dt \right\}.$$

Now split the integral in (3.17) into two integrals, one from $t = 0$ to $t = 1$ and the other from $t = 1$ to $t = \infty$. For the first use $e^t - 1 = \mathcal{O}(t)$ for $0 \leq t \leq 1$, and for the second use $\ln(e^t - 1) = \mathcal{O}(t)$ for $1 \leq t < \infty$. Thus as $j \rightarrow \infty$ we deduce that

$$(3.18) \quad \eta_j = \mathcal{O} \left\{ \int_0^1 e^{-jt} t \ln(t) dt \right\} + \mathcal{O} \left\{ \int_1^\infty e^{-(j-1)t} t dt \right\} \\ = \mathcal{O} \left(\frac{\ln(j)}{j^2} \right) + \mathcal{O} \left(\frac{1}{je^j} \right) = \mathcal{O} \left(\frac{\ln(j)}{j^2} \right),$$

where the third \mathcal{O} term comes from [6, Chap. 9, Thm. 1.1], and the fourth \mathcal{O} term came from integration by parts. Finally, from (3.9), (3.12), (3.13) and (3.18) we arrive at (1.6).

Addendum. Christian Berg kindly made us aware of his recent joint paper [1]. In this they study a more general function $h_\alpha(z) = (1 + 1/z)^{\alpha z}$ where $\alpha > 0$. They obtain a number of results for the coefficients in a Maclaurin series in z for $e^{-\alpha} h_\alpha(-1/z)$, which are polynomials in α . In particular they use properties of the exponential Bell partition polynomials to obtain a recursion relation, which for $\alpha = 1$ reduces to our formula (1.5). In comparison we only require elementary techniques to obtain (1.5).

Acknowledgments. TMD acknowledges financial support from Ministerio de Ciencia e Innovación, Spain, projects MTM2015-67142-P (MINECO/FEDER, UE) and PGC2018-098279-B-I00 (MCIU/AEI/FEDER, UE).

REFERENCES

- [1] C. BERG, E. MASSA, AND A. P. PERON, *A family of entire functions connecting the Bessel function J_1 and the Lambert W function*, *Constr. Approx.*, 53 (2021), pp. 121–154.
- [2] J. W. BROWN AND R. V. CHURCHILL, *Complex variables and applications*, Boston: McGraw-Hill Higher Education, 9 ed., 2014.
- [3] C.-P. CHEN AND J. CHOI, *An asymptotic formula for $(1 + 1/x)^x$ based on the partition function*, *Amer. Math. Monthly*, 44 (2014), pp. 338–343.
- [4] *NIST Digital Library of Mathematical Functions*. <http://dlmf.nist.gov/>, Release 1.1.1 of 2021-03-15, <http://dlmf.nist.gov/>. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [5] G. H. HARDY AND S. RAMANUJAN, *Asymptotic formulae in combinatory analysis*, *Proc. London Math. Soc.*, 17 (1918), pp. 75–115.
- [6] F. W. J. OLVER, *Asymptotics and special functions*, AKP Classics, A K Peters Ltd., Wellesley, MA, 1997. Reprint of the 1974 original [Academic Press, New York].
- [7] V. PONOMARENKO, *Asymptotic formula for $(1 + 1/x)^x$, revisited*, *Amer. Math. Monthly*, 122 (2015), p. 587.