

ON THE BETTI NUMBERS OF THE TANGENT CONES FOR GORENSTEIN MONOMIAL CURVES

PINAR METE

ABSTRACT. The aim of the article is to study the Betti numbers of the tangent cone of Gorenstein monomial curves in affine 4-space. If C_S is a non-complete intersection Gorenstein monomial curve whose tangent cone is Cohen-Macaulay, we show that the possible Betti sequences are $(1,5,5,1)$, $(1,5,6,2)$ and $(1,6,8,3)$.

1. INTRODUCTION

Let S denote the numerical semigroup generated by the positive integers $n_1 < n_2 < \dots < n_d$ with $\gcd(n_1, \dots, n_d) = 1$. Consider the polynomial rings $R = k[x_1, \dots, x_d]$ and $k[t]$ over the field k . The semigroup ring $k[S] = k[t^{n_1}, \dots, t^{n_d}]$ is the k -subalgebra of $k[t]$. $\varphi : R \rightarrow k[S] \subset k[t]$ with $\varphi(x_i) = t^{n_i}$ is the k -algebra homomorphism for $i = 1, \dots, d$ and its kernel I_S is called the toric ideal of S .

Let $m = (t^{n_1}, \dots, t^{n_d})$ be the maximal ideal of the one-dimensional local ring $k[[t^{n_1}, \dots, t^{n_d}]]$. When k is algebraically closed field, the semigroup ring $k[S] = k[t^{n_1}, \dots, t^{n_d}]$ is isomorphic to the coordinate ring R/I_S of C_S and the coordinate ring $gr_m(k[[t^{n_1}, \dots, t^{n_d}]])$ of the tangent cone of C_S at the origin is isomorphic to the ring R/I_{S_*} . Here, I_{S_*} is generated by the polynomials f_* which is the homogeneous summand of $f \in I_S$ and is called the defining ideal of the tangent cone of C_S . A monomial curve C_S is Gorenstein if the associated local ring $k[[t^{n_1}, \dots, t^{n_d}]]$ is Gorenstein. $k[[t^{n_1}, \dots, t^{n_d}]]$ is Gorenstein if and only if the semigroup S is symmetric [10]. We recall that the numerical semigroup S is symmetric if and only if for all $x \in \mathbb{Z}$ either $x \in S$ or $F(S) - x \in S$, where $F(S)$ denote the Frobenius number of S .

Finding an explicit minimal free resolution of a standard k -algebra is one of the core areas in commutative algebra. Since it is very difficult to obtain a description of the differential in the resolution, we can get some information about the numerical invariants of the resolution such as Betti numbers. The i -th Betti number of an R -module M , $\beta_i(M)$, is the rank of the free modules appearing in the minimal free resolution of M where

$$0 \rightarrow R^{\beta_{k-1}} \rightarrow \dots \rightarrow R^{\beta_1} \rightarrow R^{\beta_0}$$

and the Betti sequence of M , $\beta(M)$, is $(\beta_0(M), \beta_1(M), \dots, \beta_{k-1}(M))$. Stamate [12] gave a broad survey on the Betti numbers of the numerical semigroup rings and stated the problem of describing the Betti numbers and the minimal free resolution for the tangent cone when S is 4-generated semigroup which is symmetric, or

Date: May 11, 2021.

2020 Mathematics Subject Classification. primary 13H10; secondary 13P10, 14H20.

Key words and phrases. Gorenstein monomial curves, tangent cones, Betti numbers.

equivalently, C_S is a Gorenstein monomial curve in affine 4-space [see [12], Problem 9.9.]. In this paper, we solve this problem for Cohen-Macaulay tangent cone of a monomial curve in $\mathbb{A}^4(k)$ corresponding to a symmetric numerical semigroup. All computations have been done using SINGULAR [6],[7].

2. THE NON-COMPLETE INTERSECTION GORENSTEIN MONOMIAL CURVES

For the rest of the paper, we assume that C_S is a Gorenstein non-complete intersection monomial curve in \mathbb{A}^4 . Now, we recall Bresinsky's theorem, which gives the explicit description of the defining ideal of C_S .

Theorem 2.1. [4] *Let C_S be a monomial curve having the parametrization*

$$x_1 = t^{n_1}, x_2 = t^{n_2}, x_3 = t^{n_3}, x_4 = t^{n_4}$$

where $S = \langle n_1, n_2, n_3, n_4 \rangle$ is a numerical semigroup minimally generated by n_1, n_2, n_3, n_4 . The semigroup $\langle n_1, n_2, n_3, n_4 \rangle$ is symmetric and C_S is a non-complete intersection curve if and only if I_S is generated by the set

$$\{f_1 = x_1^{\alpha_1} - x_3^{\alpha_{13}} x_4^{\alpha_{14}}, f_2 = x_2^{\alpha_2} - x_1^{\alpha_{21}} x_4^{\alpha_{24}}, f_3 = x_3^{\alpha_3} - x_1^{\alpha_{31}} x_2^{\alpha_{32}}, \\ f_4 = x_4^{\alpha_4} - x_2^{\alpha_{42}} x_3^{\alpha_{43}}, f_5 = x_3^{\alpha_{43}} x_1^{\alpha_{21}} - x_2^{\alpha_{32}} x_4^{\alpha_{14}}\}$$

where the polynomials f_i 's are unique up to isomorphism with $0 < \alpha_{ij} < \alpha_j$ with $\alpha_i n_i \in \langle n_1, \dots, \hat{n}_i, \dots, n_4 \rangle$ such that α_i 's are minimal for $1 \leq i \leq 4$, where \hat{n}_i denotes that $n_i \notin \langle n_1, \dots, \hat{n}_i, \dots, n_4 \rangle$.

Theorem 2.1 implies that for any non-complete intersection Gorenstein monomial curve C_S , the variables can be renamed to obtain generators exactly of the given form, and this means that there are six isomorphic possible permutations which can be considered within three cases:

- (1) $f_1 = (1, (3, 4))$
 - (a) $f_2 = (2, (1, 4)), f_3 = (3, (1, 2)), f_4 = (4, (2, 3)), f_5 = ((1, 3), (2, 4))$
 - (b) $f_2 = (2, (1, 3)), f_3 = (3, (2, 4)), f_4 = (4, (1, 2)), f_5 = ((1, 4), (2, 3))$
- (2) $f_1 = (1, (2, 3))$
 - (a) $f_2 = (2, (3, 4)), f_3 = (3, (1, 4)), f_4 = (4, (1, 2)), f_5 = ((2, 4), (1, 3))$
 - (b) $f_2 = (2, (1, 4)), f_3 = (3, (2, 4)), f_4 = (4, (1, 3)), f_5 = ((1, 3), (4, 2))$
- (3) $f_1 = (1, (2, 4))$
 - (a) $f_2 = (2, (1, 3)), f_3 = (3, (1, 4)), f_4 = (4, (2, 3)), f_5 = ((1, 2), (3, 4))$
 - (b) $f_2 = (2, (3, 4)), f_3 = (3, (1, 2)), f_4 = (4, (1, 3)), f_5 = ((2, 3), (1, 4))$

Here, the notations $f_i = (i, (j, k))$ and $f_5 = ((i, j), (k, l))$ denote the generators $f_i = x_i^{\alpha_i} - x_j^{\alpha_{ij}} x_k^{\alpha_{ik}}$ and $f_5 = x_i^{\alpha_{ki}} x_j^{\alpha_{lj}} - x_k^{\alpha_{jk}} x_l^{\alpha_{il}}$. Thus, given a Gorenstein monomial curve C_S , if we have the extra condition $n_1 < n_2 < n_3 < n_4$, then the generator set of its defining ideal I_S is exactly given by one of these six permutations.

The study of the Cohen-Macaulayness of tangent cones of monomial curves constitutes an important problem [1],[2]. In [3], Arslan and Mete determined the common arithmetic conditions satisfied by the generators of the defining ideals of C_S and under these conditions they found the generators of the tangent cone of C_S . In [2], they provided necessary and sufficient conditions for the Cohen-Macaulayness of the tangent cone of C_S in all six permutations and gave the following theorem:

Theorem 2.2. [2] (1) *Suppose that I_S is given as in the Case 1(a). Then R/I_S is Cohen-Macaulay if and only if $\alpha_2 \leq \alpha_{21} + \alpha_{24}$.*

(2) Suppose that I_S is given as in Case 1(b). (i) Assume that $\alpha_{32} < \alpha_{42}$ and $\alpha_{14} \leq \alpha_{34}$. Then R/I_{S_*} is Cohen-Macaulay if and only if

1. $\alpha_2 \leq \alpha_{21} + \alpha_{23}$
2. $\alpha_{42} + \alpha_{13} \leq \alpha_{21} + \alpha_{34}$ and
3. $\alpha_3 + \alpha_{13} \leq \alpha_1 + \alpha_{32} + \alpha_{34} - \alpha_{14}$.

(ii) Assume that $\alpha_{42} \leq \alpha_{32}$. Then R/I_{S_*} is Cohen-Macaulay if and only if

1. $\alpha_2 \leq \alpha_{21} + \alpha_{23}$
2. $\alpha_{42} + \alpha_{13} \leq \alpha_{21} + \alpha_{34}$ and
3. either $\alpha_{34} < \alpha_{14}$ and $\alpha_3 + \alpha_{13} \leq \alpha_{21} + \alpha_{32} - \alpha_{42} + 2\alpha_{34}$
or $\alpha_{14} \leq \alpha_{34}$ and $\alpha_3 + \alpha_{13} \leq \alpha_1 + \alpha_{32} + \alpha_{34} - \alpha_{14}$.

(3) Suppose that I_S is given as in Case 2(a). (i) Assume that $\alpha_{24} < \alpha_{34}$ and $\alpha_{13} \leq \alpha_{23}$. Then R/I_{S_*} is Cohen-Macaulay if and only if

1. $\alpha_3 \leq \alpha_{31} + \alpha_{34}$
2. $\alpha_{12} + \alpha_{34} \leq \alpha_{41} + \alpha_{23}$ and
3. $\alpha_2 + \alpha_{12} \leq \alpha_1 + \alpha_{23} - \alpha_{13} + \alpha_{24}$.

(ii) Assume that $\alpha_{34} \leq \alpha_{24}$. Then R/I_{S_*} is Cohen-Macaulay if and only if

1. $\alpha_3 \leq \alpha_{31} + \alpha_{34}$
2. $\alpha_{12} + \alpha_{34} \leq \alpha_{41} + \alpha_{23}$ and
3. either $\alpha_{23} < \alpha_{13}$ and $\alpha_2 + \alpha_{12} \leq \alpha_{41} + 2\alpha_{23} + \alpha_{24} - \alpha_{34}$
or $\alpha_{13} \leq \alpha_{23}$ and $\alpha_2 + \alpha_{12} \leq \alpha_1 + \alpha_{23} - \alpha_{13} + \alpha_{24}$.

(4) Suppose that I_S is given as in Case 2(b). (i) Assume that $\alpha_{34} < \alpha_{24}$ and $\alpha_{12} \leq \alpha_{32}$. Then R/I_{S_*} is Cohen-Macaulay if and only if

1. $\alpha_2 \leq \alpha_{21} + \alpha_{24}$ and
2. $\alpha_3 + \alpha_{13} \leq \alpha_1 + \alpha_{32} - \alpha_{12} + \alpha_{34}$.

(ii) Assume that $\alpha_{24} \leq \alpha_{34}$. Then R/I_{S_*} is Cohen-Macaulay if and only if

1. $\alpha_2 \leq \alpha_{21} + \alpha_{24}$ and
2. either $\alpha_{32} < \alpha_{12}$ and $\alpha_3 + \alpha_{13} \leq \alpha_{41} + 2\alpha_{32} + \alpha_{34} - \alpha_{24}$
or $\alpha_{12} \leq \alpha_{32}$ and $\alpha_3 + \alpha_{13} \leq \alpha_1 + \alpha_{32} - \alpha_{12} + \alpha_{34}$.

(5) Suppose that I_S is given as in the Case 3(a). Then R/I_{S_*} is Cohen-Macaulay tangent cone if and only if $\alpha_2 \leq \alpha_{21} + \alpha_{23}$ and $\alpha_3 \leq \alpha_{31} + \alpha_{34}$.

(6) Suppose that I_S is given as in Case 3(b). (i) Assume that $\alpha_{23} < \alpha_{43}$ and $\alpha_{14} \leq \alpha_{24}$. Then R/I_{S_*} is Cohen-Macaulay if and only if

1. $\alpha_{12} + \alpha_{43} \leq \alpha_{31} + \alpha_{24}$ and
2. $\alpha_2 + \alpha_{12} \leq \alpha_1 + \alpha_{23} + \alpha_{24} - \alpha_{14}$.

(ii) Assume that $\alpha_{43} \leq \alpha_{23}$. Then R/I_{S_*} is Cohen-Macaulay if and only if

1. $\alpha_{12} + \alpha_{43} \leq \alpha_{31} + \alpha_{24}$ and
2. either $\alpha_{24} < \alpha_{14}$ and $\alpha_2 + \alpha_{12} \leq \alpha_{31} + 2\alpha_{24} + \alpha_{23} - \alpha_{43}$
or $\alpha_{14} \leq \alpha_{24}$ and $\alpha_2 + \alpha_{12} \leq \alpha_1 + \alpha_{23} + \alpha_{24} - \alpha_{14}$.

Recently, Katsabekis gave the remaining standard bases for I_S in [9] using above conditions for the Cohen-Macaulayness of the tangent cone of C_S .

3. BETTI SEQUENCES OF COHEN-MACAULAY TANGENT CONES

In this section, we determine the Betti sequences of the tangent cone of C_S whose tangent cone is Cohen-Macaulay. Buchsbaum-Eisenbud criterion [5] and the following Lemma (see also [8]) will be used in the all proofs.

Lemma 3.1. [13] *Assume that the multiplicity of the monomial curve is n_i . Suppose that the k -algebra homomorphism $\pi_i : R \rightarrow \bar{R} = k[x_1, \dots, \bar{x}_i, \dots, x_d]$ is defined by $\pi_i(x_i) = \bar{x}_i = 0$ and $\pi_j(x_j) = x_j$ for $i \neq j$, and set $\bar{I} = \pi_i(I_{S_*})$. If the tangent cone $gr_m(k[[t^{n_1}, \dots, t^{n_d}]])$ is Cohen-Macaulay, then the Betti sequences of $gr_m(k[[t^{n_1}, \dots, t^{n_d}]])$ and of \bar{R}/\bar{I} are the same.*

This is a very effective result to reduce the number of cases for finding the Betti numbers of the tangent cones.

Case 1(a) : Suppose that I_S is minimally generated by the polynomials

$$\begin{aligned} f_1 &= x_1^{\alpha_1} - x_3^{\alpha_{13}} x_4^{\alpha_{14}}, & f_2 &= x_2^{\alpha_2} - x_1^{\alpha_{21}} x_4^{\alpha_{24}}, & f_3 &= x_3^{\alpha_3} - x_1^{\alpha_{31}} x_2^{\alpha_{32}}, \\ f_4 &= x_4^{\alpha_4} - x_2^{\alpha_{42}} x_3^{\alpha_{43}}, & f_5 &= x_1^{\alpha_{51}} x_3^{\alpha_{53}} - x_2^{\alpha_{52}} x_4^{\alpha_{54}} \end{aligned}$$

where $\alpha_1 = \alpha_{21} + \alpha_{31}$, $\alpha_2 = \alpha_{32} + \alpha_{42}$, $\alpha_3 = \alpha_{13} + \alpha_{43}$, $\alpha_4 = \alpha_{14} + \alpha_{24}$. The condition $n_1 < n_2 < n_3 < n_4$ implies $\alpha_1 > \alpha_{13} + \alpha_{14}$, $\alpha_4 < \alpha_{42} + \alpha_{43}$ and $\alpha_3 < \alpha_{31} + \alpha_{32}$.

C_S has Cohen-Macaulay tangent cone only under the condition $\alpha_2 \leq \alpha_{21} + \alpha_{24}$ by [2]. Mete and Zengin [11] gave the explicit minimal free resolution for the tangent cone of C_S and showed that the Betti sequence of its tangent cone is $(1, 5, 6, 2)$.

Case 1(b) : Suppose that I_S is minimally generated by the polynomials

$$\begin{aligned} f_1 &= x_1^{\alpha_1} - x_3^{\alpha_{13}} x_4^{\alpha_{14}}, & f_2 &= x_2^{\alpha_2} - x_1^{\alpha_{21}} x_3^{\alpha_{23}}, & f_3 &= x_3^{\alpha_3} - x_2^{\alpha_{32}} x_4^{\alpha_{34}}, \\ f_4 &= x_4^{\alpha_4} - x_1^{\alpha_{41}} x_2^{\alpha_{42}}, & f_5 &= x_1^{\alpha_{51}} x_4^{\alpha_{54}} - x_2^{\alpha_{52}} x_3^{\alpha_{53}} \end{aligned}$$

where $\alpha_1 = \alpha_{21} + \alpha_{41}$, $\alpha_2 = \alpha_{32} + \alpha_{42}$, $\alpha_3 = \alpha_{13} + \alpha_{23}$, $\alpha_4 = \alpha_{14} + \alpha_{34}$. The condition $n_1 < n_2 < n_3 < n_4$ implies $\alpha_1 > \alpha_{13} + \alpha_{14}$ and $\alpha_4 < \alpha_{41} + \alpha_{42}$. Under the restriction $\alpha_2 \leq \alpha_{21} + \alpha_{23}$ by Theorem 2.2 and one possible condition $\alpha_3 \leq \alpha_{32} + \alpha_{34}$, the Betti sequences for the Cohen-Macaulay tangent cone of C_S are $(1, 5, 5, 1)$ and $(1, 5, 6, 2)$, as given in [11]. Here, the other possible condition $\alpha_3 > \alpha_{32} + \alpha_{34}$ will be considered.

Theorem 3.2. *The Betti sequence of the tangent cone R/I_{S_*} of C_S is $(1, 6, 8, 3)$, if I_S is given as in Case 1(b) when $\alpha_3 > \alpha_{32} + \alpha_{34}$.*

Proof. Suppose that I_S is given as in the Case 1(b) when $\alpha_3 > \alpha_{32} + \alpha_{34}$.

(i) By Proposition 2.7 in [9], if $\alpha_{32} < \alpha_{42}$ and $\alpha_{14} \leq \alpha_{34}$, then,

$$G = \{f_1, f_2, f_3, f_4, f_5, f_6 = x_3^{\alpha_3 + \alpha_{13}} - x_1^{\alpha_1} x_2^{\alpha_{32}} x_4^{\alpha_{34} - \alpha_{14}}\},$$

(ii) By Proposition 2.9 in [9],

(1) if $\alpha_{42} \leq \alpha_{32}$ and $\alpha_{34} < \alpha_{14}$, then

$$G = \{f_1, f_2, f_3, f_4, f_5, f_6 = x_3^{\alpha_3 + \alpha_{13}} - x_1^{\alpha_{21}} x_2^{\alpha_{32} - \alpha_{42}} x_4^{2\alpha_{34}}\},$$

(2) if $\alpha_{42} \leq \alpha_{32}$ and $\alpha_{14} \leq \alpha_{34}$, then

$$G = \{f_1, f_2, f_3, f_4, f_5, f_6 = x_3^{\alpha_3 + \alpha_{13}} - x_1^{\alpha_1} x_2^{\alpha_{32}} x_4^{\alpha_{34} - \alpha_{14}}\}$$

are standard bases for I_S . Since $\bar{I} = \pi_i(I_{S_*})$ which sends x_1 to 0, the generators of the defining ideal of \bar{I} is generated by

$$G_* = (x_3^{\alpha_{13}} x_4^{\alpha_{14}}, x_2^{\alpha_2}, x_2^{\alpha_{32}} x_4^{\alpha_{34}}, x_4^{\alpha_4}, x_2^{\alpha_{42}} x_3^{\alpha_{43}}, x_3^{\alpha_3 + \alpha_{13}}).$$

Now, consider the case (i). Since the Betti sequences of R/I_{S_*} and \bar{R}/\bar{I} are the same which follows from Lemma 3.1, we will show that the sequence,

$$0 \rightarrow R^3 \xrightarrow{\phi_3} R^8 \xrightarrow{\phi_2} R^6 \xrightarrow{\phi_1} R^1 \rightarrow 0$$

is a minimal free resolution of $\overline{R}/\overline{I}$, where

$$\phi_1 = \begin{pmatrix} x_3^{\alpha_{13}} x_4^{\alpha_{14}} & x_2^{\alpha_2} & x_2^{\alpha_{32}} x_4^{\alpha_{34}} & x_4^{\alpha_4} & x_2^{\alpha_{42}} x_3^{\alpha_{13}} & x_3^{\alpha_3 + \alpha_{13}} \end{pmatrix},$$

$$\phi_2 = \begin{pmatrix} 0 & -x_2^{\alpha_{32}} x_4^{\alpha_{34} - \alpha_{14}} & 0 & -x_4^{\alpha_{34}} & x_2^{\alpha_{42}} & x_3^{\alpha_3} & 0 & 0 \\ 0 & 0 & -x_4^{\alpha_{34}} & 0 & 0 & 0 & x_3^{\alpha_{13}} & 0 \\ x_4^{\alpha_{14}} & x_3^{\alpha_{13}} & x_2^{\alpha_{42}} & 0 & 0 & 0 & 0 & 0 \\ -x_2^{\alpha_{32}} & 0 & 0 & x_3^{\alpha_{13}} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -x_4^{\alpha_{14}} & 0 & -x_2^{\alpha_{32}} & x_3^{\alpha_3} \\ 0 & 0 & 0 & 0 & 0 & -x_4^{\alpha_{14}} & 0 & -x_2^{\alpha_{42}} \end{pmatrix},$$

$$\phi_3 = \begin{pmatrix} x_3^{\alpha_{13}} & 0 & 0 \\ -x_4^{\alpha_{14}} & x_2^{\alpha_{42}} & 0 \\ 0 & -x_3^{\alpha_{13}} & 0 \\ x_2^{\alpha_{32}} & 0 & 0 \\ 0 & x_2^{\alpha_{32}} x_4^{\alpha_{34} - \alpha_{14}} & x_3^{\alpha_3} \\ 0 & 0 & -x_2^{\alpha_{42}} \\ 0 & x_4^{\alpha_{34}} & 0 \\ 0 & 0 & x_4^{\alpha_{14}} \end{pmatrix}.$$

Since $\phi_1 \phi_2 = \phi_2 \phi_3 = 0$, the sequence above is a complex. One can easily check that $\text{rank}(\phi_1) = 1$, $\text{rank}(\phi_2) = 5$ and $\text{rank}(\phi_3) = 3$. As the 5-minors of ϕ_2 , we get $x_3^{2\alpha_3 + 3\alpha_{13}}$ by deleting the row 6, and the columns 1, 3, 5, and $-x_4^{2\alpha_4 + \alpha_{14}}$ by deleting the row 4 and the columns 2, 7, 8. These two determinants are relatively prime, so $I(\phi_2)$ contains a regular sequence of length 2. As the 3-minors of ϕ_3 , we have $x_2^{\alpha_2 + \alpha_{42}}$ by deleting the rows 1, 3, 5, 7, 8, and $-x_3^{\alpha_3 + 2\alpha_{13}}$ by deleting the rows 2, 4, 6, 7, 8, and finally, $-x_4^{\alpha_4 + \alpha_{14}}$ by deleting the rows 1, 3, 4, 5, 6. Since these are relatively prime, $I(\phi_3)$ contains a regular sequence of length 3.

The other cases (ii)-(1) and (2) can be done similarly and are, hence, omitted. \blacksquare

Case 2(a) : Suppose that I_S is minimally generated by the polynomials

$$f_1 = x_1^{\alpha_1} - x_2^{\alpha_{12}} x_3^{\alpha_{13}}, \quad f_2 = x_2^{\alpha_2} - x_3^{\alpha_{23}} x_4^{\alpha_{24}}, \quad f_3 = x_3^{\alpha_3} - x_1^{\alpha_{31}} x_4^{\alpha_{34}},$$

$$f_4 = x_4^{\alpha_4} - x_1^{\alpha_{41}} x_2^{\alpha_{42}}, \quad f_5 = x_1^{\alpha_{41}} x_3^{\alpha_{23}} - x_2^{\alpha_{12}} x_4^{\alpha_{34}}$$

where $\alpha_1 = \alpha_{31} + \alpha_{41}$, $\alpha_2 = \alpha_{12} + \alpha_{42}$, $\alpha_3 = \alpha_{13} + \alpha_{23}$, $\alpha_4 = \alpha_{24} + \alpha_{34}$. The condition $n_1 < n_2 < n_3 < n_4$ implies $\alpha_1 > \alpha_{12} + \alpha_{13}$, $\alpha_2 > \alpha_{23} + \alpha_{24}$ and $\alpha_4 < \alpha_{41} + \alpha_{42}$. Note that the assumption $\alpha_3 \leq \alpha_{31} + \alpha_{34}$ is one of the necessary and sufficient conditions for the Cohen-Macaulayness of the tangent cone by Theorem 2.2.

Theorem 3.3. *The Betti sequence of the tangent cone R/I_{S^*} of C_S is $(1, 6, 8, 3)$, if I_S is given as in Case 2(a).*

Proof. Suppose that I_S is given as in the Case 2(a).

(i) By Proposition 2.11 in [9], if $\alpha_{24} < \alpha_{34}$ and $\alpha_{13} \leq \alpha_{23}$, then

$$G = \{f_1, f_2, f_3, f_4, f_5, f_6 = x_2^{\alpha_2 + \alpha_{12}} - x_1^{\alpha_1} x_3^{\alpha_{23} - \alpha_{13}} x_4^{\alpha_{24}}\}$$

(ii) By Proposition 2.13 in [9],

(1) if $\alpha_{34} \leq \alpha_{24}$ and $\alpha_{23} < \alpha_{13}$, then

$$G = \{f_1, f_2, f_3, f_4, f_5, f_6 = x_2^{\alpha_2 + \alpha_{12}} - x_1^{\alpha_{41}} x_3^{2\alpha_{23}} x_4^{\alpha_{24} - \alpha_{34}}\}$$

(2) if $\alpha_{34} \leq \alpha_{24}$ and $\alpha_{13} \leq \alpha_{23}$, then

$$G = \{f_1, f_2, f_3, f_4, f_5, f_6 = x_2^{\alpha_2 + \alpha_{12}} - x_1^{\alpha_1} x_3^{\alpha_{23} - \alpha_{13}} x_4^{\alpha_{24}}\}$$

are standard bases for I_S . Since $\bar{I} = \pi_i(I_{S_*})$ which sends x_1 to 0, the generators of the defining ideal of \bar{I} is generated by

$$G_* = (x_2^{\alpha_{12}} x_3^{\alpha_{13}}, x_3^{\alpha_{23}} x_4^{\alpha_{24}}, x_3^{\alpha_3}, x_4^{\alpha_4}, x_2^{\alpha_{12}} x_4^{\alpha_{34}}, x_2^{\alpha_2 + \alpha_{12}}).$$

Now, consider the case (i). Since the Betti sequences of R/I_{S_*} and \bar{R}/\bar{I} are the same which follows from Lemma 3.1, we will show that the sequence,

$$0 \rightarrow R^3 \xrightarrow{\phi_3} R^8 \xrightarrow{\phi_2} R^6 \xrightarrow{\phi_1} R^1 \rightarrow 0$$

is a minimal free resolution of \bar{R}/\bar{I} , where

$$\phi_1 = \begin{pmatrix} x_2^{\alpha_{12}} x_3^{\alpha_{13}} & x_3^{\alpha_{23}} x_4^{\alpha_{24}} & x_3^{\alpha_3} & x_4^{\alpha_4} & x_2^{\alpha_{12}} x_4^{\alpha_{34}} & x_2^{\alpha_2 + \alpha_{12}} \end{pmatrix},$$

$$\phi_2 = \begin{pmatrix} x_3^{\alpha_{23}} & x_2^{\alpha_{23} - \alpha_{13}} x_4^{\alpha_{24}} & x_4^{\alpha_{34}} & x_2^{\alpha_2} & 0 & 0 & 0 & 0 \\ 0 & -x_2^{\alpha_{12}} & 0 & 0 & -x_3^{\alpha_{13}} & x_4^{\alpha_{34}} & 0 & 0 \\ -x_2^{\alpha_{12}} & 0 & 0 & 0 & x_4^{\alpha_{24}} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -x_3^{\alpha_{23}} & -x_2^{\alpha_{12}} & 0 \\ 0 & 0 & -x_3^{\alpha_{13}} & 0 & 0 & 0 & x_4^{\alpha_{24}} & x_2^{\alpha_2} \\ 0 & 0 & 0 & -x_3^{\alpha_{13}} & 0 & 0 & 0 & -x_4^{\alpha_{34}} \end{pmatrix},$$

$$\phi_3 = \begin{pmatrix} x_4^{\alpha_{24}} & 0 & 0 \\ -x_3^{\alpha_{13}} & x_4^{\alpha_{34}} & 0 \\ 0 & -x_3^{\alpha_{23} - \alpha_{13}} x_4^{\alpha_{24}} & x_2^{\alpha_2} \\ 0 & 0 & -x_4^{\alpha_{34}} \\ x_2^{\alpha_{12}} & 0 & 0 \\ 0 & x_2^{\alpha_{12}} & 0 \\ 0 & -x_3^{\alpha_{23}} & 0 \\ 0 & 0 & x_3^{\alpha_{13}} \end{pmatrix}.$$

Since $\phi_1 \phi_2 = \phi_2 \phi_3 = 0$, the sequence above is a complex. One can easily check that $rank(\phi_1) = 1$, $rank(\phi_2) = 1$ and $rank(\phi_3) = 3$.

As the 5-minors of ϕ_2 , we have $-x_3^{2\alpha_3 + \alpha_{13}}$ by removing the columns 2, 7, 8 and the row 3, and $-x_4^{2\alpha_4 + \alpha_{34}}$ by removing the columns 1, 2, 4 and the row 4. These two determinants are relatively prime, so $I(\phi_2)$ contains a regular sequence of length 2. As the 3-minors of ϕ_3 , we get $x_2^{2\alpha_2 + 2\alpha_{12}}$ by removing the rows 1, 2, 4, 7, 8, and $x_3^{\alpha_3 + \alpha_{13}}$ by removing the rows 1, 3, 4, 5, 6, and finally $-x_4^{\alpha_4 + \alpha_{34}}$ by removing the rows 3, 5, 6, 7, 8. These three determinants constitute a regular sequence. Thus, $I(\phi_3)$ contains a regular sequence of length 3.

The other cases (ii)-(1) and (2) can be done similarly and are, hence, omitted. \blacksquare

Case 2(b) : Suppose that I_S is minimally generated by the polynomials

$$f_1 = x_1^{\alpha_1} - x_2^{\alpha_{12}} x_3^{\alpha_{13}}, \quad f_2 = x_2^{\alpha_2} - x_1^{\alpha_{21}} x_4^{\alpha_{24}}, \quad f_3 = x_3^{\alpha_3} - x_2^{\alpha_{32}} x_4^{\alpha_{34}},$$

$$f_4 = x_4^{\alpha_4} - x_1^{\alpha_{41}} x_3^{\alpha_{43}}, \quad f_5 = x_1^{\alpha_{41}} x_2^{\alpha_{32}} - x_3^{\alpha_{13}} x_4^{\alpha_{24}}.$$

where $\alpha_1 = \alpha_{21} + \alpha_{41}$, $\alpha_2 = \alpha_{12} + \alpha_{32}$, $\alpha_3 = \alpha_{13} + \alpha_{43}$, $\alpha_4 = \alpha_{24} + \alpha_{34}$. The condition $n_1 < n_2 < n_3 < n_4$ implies $\alpha_1 > \alpha_{12} + \alpha_{13}$ and $\alpha_4 < \alpha_{41} + \alpha_{43}$.

Under the restriction $\alpha_2 \leq \alpha_{21} + \alpha_{24}$ by Theorem 2.2 and one possible condition $\alpha_3 \leq \alpha_{32} + \alpha_{34}$, the Betti sequences for the Cohen-Macaulay tangent cone of C_S

are $(1, 5, 5, 1)$ and $(1, 5, 6, 2)$, as given in [11]. Here, the other possible condition $\alpha_3 > \alpha_{32} + \alpha_{34}$ will be considered.

Theorem 3.4. *The Betti sequence of the tangent cone R/I_{S^*} of C_S is $(1, 6, 8, 3)$, if I_S is given as in Case 2(b) when $\alpha_3 > \alpha_{32} + \alpha_{34}$.*

Proof. Suppose that I_S is given as in the Case 2(b).

(i) By Proposition 2.16 in [9], if $\alpha_{34} < \alpha_{24}$ and $\alpha_{12} \leq \alpha_{32}$, then

$$G = \{f_1, f_2, f_3, f_4, f_5, f_6 = x_3^{\alpha_3 + \alpha_{13}} - x_1^{\alpha_1} x_2^{\alpha_{32} - \alpha_{12}} x_4^{\alpha_{34}}\}$$

(ii) By Proposition 2.18 in [9],

(1) if $\alpha_{24} \leq \alpha_{34}$ and $\alpha_{32} < \alpha_{12}$, then

$$G = \{f_1, f_2, f_3, f_4, f_5, f_6 = x_3^{\alpha_3 + \alpha_{13}} - x_1^{\alpha_{41}} x_2^{2\alpha_{32}} x_4^{\alpha_{34} - \alpha_{24}}\},$$

(2) if $\alpha_{24} \leq \alpha_{34}$ and $\alpha_{12} \leq \alpha_{32}$, then

$$G = \{f_1, f_2, f_3, f_4, f_5, f_6 = x_3^{\alpha_3 + \alpha_{13}} - x_1^{\alpha_1} x_2^{\alpha_{32} - \alpha_{12}} x_4^{\alpha_{34}}\}$$

are standard bases for I_S . Since $\bar{I} = \pi_i(I_{S^*})$ which sends x_1 to 0, then the generators of the defining ideal of \bar{I} is generated by

$$G_* = (x_2^{\alpha_{12}} x_3^{\alpha_{13}}, x_2^{\alpha_2}, x_2^{\alpha_{32}} x_4^{\alpha_{34}}, x_4^{\alpha_4}, x_3^{\alpha_{13}} x_4^{\alpha_{24}}, x_3^{\alpha_3 + \alpha_{13}}).$$

Now, consider the case (i). Since the Betti sequences of R/I_{S^*} and \bar{R}/\bar{I} are the same which follows from Lemma 3.1, we will show that the sequence,

$$0 \rightarrow R^3 \xrightarrow{\phi_3} R^8 \xrightarrow{\phi_2} R^6 \xrightarrow{\phi_1} R^1 \rightarrow 0$$

is a minimal free resolution of \bar{R}/\bar{I} , where

$$\phi_1 = \begin{pmatrix} x_2^{\alpha_{12}} x_3^{\alpha_{13}} & x_2^{\alpha_2} & x_2^{\alpha_{32}} x_4^{\alpha_{34}} & x_4^{\alpha_4} & x_3^{\alpha_{13}} x_4^{\alpha_{24}} & x_3^{\alpha_3 + \alpha_{13}} \end{pmatrix},$$

$$\phi_2 = \begin{pmatrix} x_4^{\alpha_{24}} & x_2^{\alpha_{32}} & x_3^{\alpha_3} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -x_3^{\alpha_{13}} & 0 & 0 & 0 & 0 & x_4^{\alpha_{34}} & 0 & 0 \\ 0 & 0 & 0 & 0 & -x_3^{\alpha_{13}} & 0 & -x_2^{\alpha_{12}} & -x_4^{\alpha_{24}} & 0 \\ 0 & 0 & 0 & -x_3^{\alpha_{13}} & 0 & 0 & 0 & 0 & x_2^{\alpha_{32}} \\ -x_2^{\alpha_{12}} & 0 & 0 & x_4^{\alpha_{34}} & x_2^{\alpha_{32}} x_4^{\alpha_{34} - \alpha_{24}} & x_3^{\alpha_3} & 0 & 0 & 0 \\ 0 & 0 & -x_2^{\alpha_{12}} & 0 & 0 & -x_4^{\alpha_{24}} & 0 & 0 & 0 \end{pmatrix},$$

$$\phi_3 = \begin{pmatrix} x_3^{\alpha_3} & -x_2^{\alpha_{32}} x_4^{\alpha_{34} - \alpha_{24}} & 0 \\ 0 & x_4^{\alpha_{34}} & 0 \\ -x_4^{\alpha_{24}} & 0 & 0 \\ 0 & 0 & x_2^{\alpha_{32}} \\ 0 & -x_2^{\alpha_{12}} & -x_4^{\alpha_{24}} \\ x_2^{\alpha_{12}} & 0 & 0 \\ 0 & x_3^{\alpha_{13}} & 0 \\ 0 & 0 & x_3^{\alpha_{13}} \end{pmatrix}.$$

Since $\phi_1 \phi_2 = \phi_2 \phi_3 = 0$, the sequence above is a complex. One can easily check that $\text{rank}(\phi_1) = 1$, $\text{rank}(\phi_2) = 5$ and $\text{rank}(\phi_3) = 3$. As the 5-minors of ϕ_2 , we get $-x_3^{2\alpha_3 + 3\alpha_{13}}$ by removing the row 6 and the columns 1, 7, 8, and $x_4^{2\alpha_4 + \alpha_{24}}$ by removing the row 4 and the columns 2, 3, 5. These two determinants are relatively prime, so $I(\phi_2)$ contains a regular sequence of length 2. As the 3-minors of ϕ_3 , we have $-x_2^{\alpha_2 + \alpha_{12}}$ by removing the rows 1, 2, 3, 7, 8, and $x_3^{\alpha_3 + 2\alpha_{13}}$ by removing the rows 2, 3, 4, 5, 6, and finally, $-x_4^{\alpha_4 + \alpha_{24}}$ by removing the rows 1, 4, 6, 7, 8. These

three determinants constitute a regular sequence. Thus, $I(\phi_3)$ contains a regular sequence of length 3.

The other cases (ii)-(1) and (2) can be done similarly and are, hence, omitted. \blacksquare

Case 3(a) : Suppose that I_S is minimally generated by the polynomials

$$\begin{aligned} f_1 &= x_1^{\alpha_1} - x_2^{\alpha_{12}} x_4^{\alpha_{14}}, & f_2 &= x_2^{\alpha_2} - x_1^{\alpha_{21}} x_3^{\alpha_{23}}, & f_3 &= x_3^{\alpha_3} - x_1^{\alpha_{31}} x_4^{\alpha_{34}}, \\ f_4 &= x_4^{\alpha_4} - x_2^{\alpha_{42}} x_3^{\alpha_{43}}, & f_5 &= x_1^{\alpha_{31}} x_2^{\alpha_{42}} - x_3^{\alpha_{23}} x_4^{\alpha_{14}} \end{aligned}$$

where $\alpha_1 = \alpha_{21} + \alpha_{31}$, $\alpha_2 = \alpha_{12} + \alpha_{42}$, $\alpha_3 = \alpha_{23} + \alpha_{43}$, $\alpha_4 = \alpha_{14} + \alpha_{34}$. The condition $n_1 < n_2 < n_3 < n_4$ implies $\alpha_1 > \alpha_{12} + \alpha_{14}$ and $\alpha_4 < \alpha_{42} + \alpha_{43}$.

C_S has Cohen-Macaulay tangent cone only under the conditions $\alpha_2 \leq \alpha_{21} + \alpha_{23}$ and $\alpha_3 \leq \alpha_{31} + \alpha_{34}$ by [2]. Mete and Zengin [11] gave the explicit minimal free resolution for the tangent cone of C_S and showed that the Betti sequence of its tangent cone is $(1, 5, 6, 2)$.

Case 3(b) : Suppose that I_S is minimally generated by the polynomials

$$\begin{aligned} f_1 &= x_1^{\alpha_1} - x_2^{\alpha_{12}} x_4^{\alpha_{14}}, & f_2 &= x_2^{\alpha_2} - x_3^{\alpha_{23}} x_4^{\alpha_{24}}, & f_3 &= x_3^{\alpha_3} - x_1^{\alpha_{31}} x_2^{\alpha_{32}}, \\ f_4 &= x_4^{\alpha_4} - x_1^{\alpha_{41}} x_3^{\alpha_{43}}, & f_5 &= x_2^{\alpha_{12}} x_3^{\alpha_{43}} - x_1^{\alpha_{31}} x_4^{\alpha_{24}}. \end{aligned}$$

Here, $\alpha_1 = \alpha_{31} + \alpha_{41}$, $\alpha_2 = \alpha_{12} + \alpha_{32}$, $\alpha_3 = \alpha_{23} + \alpha_{43}$, $\alpha_4 = \alpha_{14} + \alpha_{24}$. The condition $n_1 < n_2 < n_3 < n_4$ implies $\alpha_1 > \alpha_{12} + \alpha_{14}$, $\alpha_2 > \alpha_{23} + \alpha_{24}$, $\alpha_3 < \alpha_{31} + \alpha_{32}$ and $\alpha_4 < \alpha_{41} + \alpha_{43}$.

Theorem 3.5. *The Betti sequence of the tangent cone R/I_{S^*} of C_S is $(1, 6, 8, 3)$, if I_S is given as in the Case 3(b).*

Proof. Suppose that I_S is given as in the Case 3(b).

(i) By Proposition 2.21 in [9], if $\alpha_{23} < \alpha_{43}$ and $\alpha_{14} \leq \alpha_{24}$, then

$$\{f_1, f_2, f_3, f_4, f_5, f_6 = x_2^{\alpha_2 + \alpha_{12}} - x_1^{\alpha_1} x_3^{\alpha_{23}} x_4^{\alpha_{24} - \alpha_{14}}\},$$

(ii) By Proposition 2.23 in [9],

(1) if $\alpha_{43} \leq \alpha_{23}$ and $\alpha_{24} < \alpha_{14}$ then

$$\{f_1, f_2, f_3, f_4, f_5, f_6 = x_2^{\alpha_2 + \alpha_{12}} - x_1^{\alpha_{31}} x_3^{\alpha_{23} - \alpha_{43}} x_4^{2\alpha_{24}}\},$$

(2) if $\alpha_{43} \leq \alpha_{23}$ and $\alpha_{14} \leq \alpha_{24}$ then

$$\{f_1, f_2, f_3, f_4, f_5, f_6 = x_2^{\alpha_2 + \alpha_{12}} - x_1^{\alpha_1} x_3^{\alpha_{23}} x_4^{\alpha_{24} - \alpha_{14}}\}$$

are standard bases for I_S . Since $\bar{I} = \pi_i(I_{S^*})$ which sends x_1 to 0, then the generators of the defining ideal of \bar{I} is generated by

$$G_* = (x_2^{\alpha_{12}} x_4^{\alpha_{14}}, x_3^{\alpha_{23}} x_4^{\alpha_{24}}, x_3^{\alpha_3}, x_4^{\alpha_4}, x_2^{\alpha_{12}} x_3^{\alpha_{43}}, x_2^{\alpha_2 + \alpha_{12}}).$$

Now, consider the case (i). Since the Betti sequences of R/I_{S^*} and \bar{R}/\bar{I} are the same which follows from Lemma 3.1, we will show that the sequence,

$$0 \rightarrow R^3 \xrightarrow{\phi_3} R^8 \xrightarrow{\phi_2} R^6 \xrightarrow{\phi_1} R^1 \rightarrow 0$$

is a minimal free resolution of \bar{R}/\bar{I} , where

$$\phi_1 = \begin{pmatrix} x_2^{\alpha_{12}} x_4^{\alpha_{14}} & x_3^{\alpha_{23}} x_4^{\alpha_{24}} & x_3^{\alpha_3} & x_4^{\alpha_4} & x_2^{\alpha_{12}} x_3^{\alpha_{43}} & x_2^{\alpha_2 + \alpha_{12}} \end{pmatrix},$$

$$\phi_2 = \begin{pmatrix} x_3^{\alpha_{43}} & x_3^{\alpha_{23}} x_4^{\alpha_{24}-\alpha_{14}} & x_4^{\alpha_{24}} & x_2^{\alpha_2} & 0 & 0 & 0 & 0 \\ 0 & -x_2^{\alpha_{12}} & 0 & 0 & 0 & 0 & -x_3^{\alpha_{43}} & x_4^{\alpha_{14}} \\ 0 & 0 & 0 & 0 & -x_2^{\alpha_{12}} & 0 & x_4^{\alpha_{24}} & 0 \\ 0 & 0 & -x_2^{\alpha_{12}} & 0 & 0 & 0 & 0 & -x_3^{\alpha_{23}} \\ -x_4^{\alpha_{14}} & 0 & 0 & 0 & x_3^{\alpha_{23}} & x_2^{\alpha_2} & 0 & 0 \\ 0 & 0 & 0 & -x_4^{\alpha_{14}} & 0 & -x_3^{\alpha_{43}} & 0 & 0 \end{pmatrix},$$

$$\phi_3 = \begin{pmatrix} x_2^{\alpha_2} & 0 & x_3^{\alpha_{23}} x_4^{\alpha_{24}-\alpha_{14}} \\ 0 & x_4^{\alpha_{14}} & -x_3^{\alpha_{43}} \\ 0 & -x_3^{\alpha_{23}} & 0 \\ -x_3^{\alpha_{43}} & 0 & 0 \\ 0 & 0 & x_4^{\alpha_{24}} \\ x_4^{\alpha_{14}} & 0 & 0 \\ 0 & 0 & x_2^{\alpha_{12}} \\ 0 & x_2^{\alpha_{12}} & 0 \end{pmatrix}.$$

Since $\phi_1\phi_2 = \phi_2\phi_3 = 0$, the sequence above is a complex. It is easy to show that $\text{rank}(\phi_1) = 1$, $\text{rank}(\phi_2) = 5$ and $\text{rank}(\phi_3) = 3$. As the 5-minors of ϕ_2 , we have $-x_3^{2\alpha_3+\alpha_{43}}$ by removing the row 3 and the columns 2, 3, 4, and $x_4^{2\alpha_4+\alpha_{14}}$ by removing the row 4 and the columns 2, 5, 6. These two determinants are relatively prime, so $I(\phi_2)$ contains a regular sequence of length 2. As the 3-minors of ϕ_3 , we get $-x_2^{\alpha_2+2\alpha_{12}}$ by removing the rows 2, 3, 4, 5, 6, and $x_3^{\alpha_3+\alpha_{43}}$ by removing the rows 1, 5, 6, 7, 8, and finally, $x_4^{\alpha_4+\alpha_{14}}$ by removing the rows 1, 3, 4, 7, 8. These three determinants constitute a regular sequence. Thus, $I(\phi_3)$ contains a regular sequence of length 3.

The other cases (ii)-(1) and (2) can be done similarly and are, hence, omitted. ■

REFERENCES

1. F. Arslan, Cohen-Macaulayness of tangent cones, Proc. Amer. Math. Soc. 128 (2000) 2243-2251.
2. F. Arslan, A. Katsabekis, M. Nalbandiyan, On the Cohen-Macaulayness of tangent cones of monomial curves in $\mathbb{A}^4(K)$, Turkish Journal of Math. 43 (2019) 1425-1446.
3. F. Arslan and P. Mete, Hilbert functions of Gorenstein monomial curves, Proc. Amer. Math. Soc. 135 (2007) 1993-2002.
4. H. Bresinsky, Symmetric semigroups of integers generated by four elements, Manuscripta Math. 17 (1975) 205-219.
5. D. Buchsbaum and D. Eisenbud, What makes a complex exact?, Journal of Algebra 323 (1973) 259-268.
6. G.-M. Greuel and G. Pfister, A Singular Introduction to Commutative Algebra (Springer-Verlag, 2002).
7. G.-M. Greuel, G. Pfister, H. Schönemann, (2001). SINGULAR 2.0. A Computer Algebra System for Polynomial Computations. Centre for Computer Algebra, University of Kaiserslautern. Available at <http://www.singular.uni-kl.de>.
8. J. Herzog and D. I. Stamate, On the defining equations of the tangent cone of a numerical semigroup ring, Journal of Algebra 418 (2014) 8-28.
9. A. Katsabekis, Hilberts series of tangent cones for Gorenstein monomial curves in $\mathbb{A}^4(K)$, Turkish Journal of Math. 45 (2021) 597-616.
10. E. Kunz, The value-semigroup of a one-dimensional Gorenstein ring. Proc. Amer. Math. Soc. 25 (1970) 748-751.
11. P. Mete, E. E. Zengin, Minimal free resolutions of the tangent cones for Gorenstein monomial curves, Turkish Journal of Math. 43 (2019) 2782-2793.

12. D. Stamate, Betti numbers for numerical semigroup rings, *Multigraded Algebra and Applications*, 133-157, Springer Proceedings in Mathematics and Statistics, 238 (Springer, Cham, 2018).
13. M. Şahin and N. Şahin, Betti numbers for certain Cohen-Macaulay tangent cones, *Bulletin of the Australian Mathematical Soc.* 99 (2019) 68-77.

DEPARTMENT OF MATHEMATICS, BALIKESIR UNIVERSITY, BALIKESIR, 10145 TURKEY
Email address: `pinarm@balikesir.edu.tr`