Lyapunov–Krasovskii functionals for some classes of nonlinear time delay systems

Gerson Portilla¹, Irina V. Alexandrova² and Sabine Mondié¹

Abstract— In this contribution, we study an homogeneous class of nonlinear time delay systems with time-varying perturbations. Using the Lyapunov-Krasovskii approach, we introduce a functional that leads to perturbation conditions matching those obtained previously in the Razumikhin framework. The functionals are applied to the estimation of the domain of attraction and of the system solutions. An illustrative example is given.

I. INTRODUCTION

The homogeneous approximation is the top choice for the stability analysis for nonlinear systems when the first approximation is zero. This strategy along with the Lyapunov framework have produced several results on stability [1], [2], [3], robustness [4] and controller design [5]. The Lyapunov-Razumikhin approach has had a key role in the development of the main results on homogeneous approximations of timedelay systems: delay-independent stability [1], [2], [3] and applications such as estimates of the solutions [1], [6] and of the region of attraction [2]. The Lyapunov-Razumikhin framework allowed establishing an outstanding result: if the delay-free system is asymptotically stable, then the trivial solution of the homogeneous delay system is asymptotically stable for all delays when the homogeneity degree is strictly greater than one [1].

A common practice is to use the Lyapunov function of the delay-free system to determine the stability of the system in presence of delays, as proposed in the Lyapunov-Razumikhin approach. The asymptotic stability of delay free systems with time-varying perturbations has been addressed in [7] in the framework of the Lyapunov direct method. There, a method for constructing a Lyapunov functions was given, and conditions insuring the system subject to such time varying perturbations remains asymptotically stable was presented. In [1], the above Lyapunov function for a delay free system was used in the Lyapunov-Razumikhin framework to determine conditions for the asymptotic stability of time-delay systems with perturbations.

In recent years, the study, in the Lyapunov-Krasovskii framework, of homogeneous systems with delays has received some attention thanks to the Lyapunov-Krasovskii functional

*The work of the first and third authors was supported by project CONACYT A1-S-24796, Mexico. The work of the second author was supported by the Russian Science Foundation, Project 19-71-00061.

¹Gerson Portilla and Sabine Mondié are with Department of Automatic Control, CINVESTAV-IPN, 07360 Mexico D.F., Mexico gportilla, smondie@ctrl.cinvestav.mx

²Irina V. Alexandrova is with Department of Applied Mathematics and Control Processes, St.Petersburg State University, 199034 St.Petersburg, Russia i.v.aleksandrova@spbu.ru for systems of homogeneity degree strictly greater than one introduced in [8], [9]. Its construction, whose starting point is the homogeneous delay free system Lyapunov function, is inspired by the so-called complete type functionals approach [10].

In [1], [6], the following class of nonlinear systems with time-varying perturbations

$$\dot{x}(t) = f(x(t), x(t-h)) + B(t)Q(x(t), x(t-h))$$
(1)

is studied. Here, $x(t) \in \mathbb{R}^n$, h > 0 is a constant delay, $f(\mathbf{x}_1, \mathbf{x}_2)$ is a homogeneous vector function of degree $\mu > 1$, that is

$$f(c\mathbf{x}_1, c\mathbf{x}_2) = c^{\mu} f(\mathbf{x}_1, \mathbf{x}_2) \quad \forall c > 0, \ \mathbf{x}_1, \mathbf{x}_2 \in \mathbb{R}^n,$$

satisfies the Lipschitz condition and is continuously differentiable with respect to \mathbf{x}_2 , B(t) is a matrix with continuous and bounded elements, $||B(t)|| \leq \hat{b}$, and the vector function $Q(\mathbf{x}_1, \mathbf{x}_2)$ is continuously differentiable and satisfies

$$\|Q(\mathbf{x}_{1},\mathbf{x}_{2})\| \leq p_{1} \|\mathbf{x}_{1}\|^{\sigma} + p_{2} \|\mathbf{x}_{2}\|^{\sigma}, \quad p_{1}, p_{2} \geq 0, \quad (2)$$
$$\left\|\frac{\partial Q(\mathbf{x}_{1},\mathbf{x}_{2})}{\partial \mathbf{x}_{j}}\right\| \leq q_{j1} \|\mathbf{x}_{1}\|^{\sigma-1} + q_{j2} \|\mathbf{x}_{2}\|^{\sigma-1},$$

 $q_{jk} \ge 0, \ j, k = 1, 2$. The following result is known.

Theorem 1. [1], [6] If the delay free system

$$\dot{x} = f(x(t), x(t)) \tag{3}$$

is asymptotically stable, then the asymptotic stability is preserved for the trivial solution of system (1) *with perturbations of the following classes:* (a) *the integral*

$$I(t) = \int_0^t B(s)ds \tag{4}$$

is bounded and $\sigma > \frac{\mu+1}{2}$; (b) integral (4) is unbounded but

$$\frac{1}{T} \int_{t}^{t+T} B(s) ds \xrightarrow[T \to +\infty]{} 0 \tag{5}$$

uniformly with respect to $t \ge 0$, and $\sigma \ge \mu$.

In case (a), the elements of B(t) may describe periodic oscillations with zero mean values, without any restriction on their amplitudes, whereas assumption (b) holds, in particular, when the elements of B(t) are almost periodic functions with zero mean values. It is also known [2] that if system (3) is asymptotically stable, then the trivial solution of system (1) with $B(t) \equiv \mathbf{0}$ is asymptotically stable for all values of the delay. The aim of this paper is to construct the Lyapunov-Krasovskii functionals for system (1), which on the one hand, allow to verify the results of [1], [6], and on the other hand, are useful in practical applications such as the construction of estimates for the solutions, etc. The functionals are based on the Lyapunov functions for delay free systems constructed by the method introduced in [7]. It is worth mentioning that they are not directly derived from an application of the construction [9] with the Lyapunov function [7].

The contribution is organised as follows. In Section II, the main properties of homogeneous systems are introduced. The construction of the functional is presented in Section III. In Section IV, estimates of the attraction region and of the solutions achieved by using this functional are presented. Section V is devoted to an illustrative example. The paper ends with conclusions.

II. PRELIMINARIES

In this paper, we assume that the initial functions belong to the space of \mathbb{R}^n -valued continuous functions on [-h, 0], which is denoted by $C_{[-h,0]}$. This space is endowed with the norm $\|\varphi\|_h = \max_{\theta \in [-h,0]} \|\varphi(\theta)\|$, where $\|\cdot\|$ stands for the Euclidean norm. Function $x(t, \varphi)$ denotes the solution of system (1) with an initial function φ , whereas x_t is the state of system (1):

$$x_t: \quad \theta \to x(t+\theta), \quad \theta \in [-h, 0].$$

Assume that the vector function $f(\mathbf{x}_1, \mathbf{x}_2)$ is continuously differentiable with respect to \mathbf{x}_1 . Due to the homogeneity, there exist $m_1, m_2, \eta_{11}, \eta_{12} \ge 0$ such that

$$\|f(\mathbf{x}_{1}, \mathbf{x}_{2})\| \leq m_{1} \|\mathbf{x}_{1}\|^{\mu} + m_{2} \|\mathbf{x}_{2}\|^{\mu},$$

$$\left\|\frac{\partial f(\mathbf{x}_{1}, \mathbf{x}_{2})}{\partial \mathbf{x}_{1}}\right\| \leq \eta_{11} \|\mathbf{x}_{1}\|^{\mu-1} + \eta_{12} \|\mathbf{x}_{2}\|^{\mu-1}.$$
(6)

We emphasize that, for the development of the Lyapunov– Krasovskii approach, differentiability of f with respect to x_1 , instead of that with respect to x_2 in the Lyapunov– Razumikhin approach [1], [6], is required.

Throughout the paper, we assume that the delay free system (3) is asymptotically stable. It is known [11], [4] that there exists a positive definite and twice continuously differentiable Lyapunov function $V(\mathbf{x})$, which is homogeneous of degree $\gamma \geq 2$ and satisfies together with its derivatives the following inequalities

$$\left(\frac{\partial V(\mathbf{x})}{\partial \mathbf{x}}\right)^T f(\mathbf{x}, \mathbf{x}) \le -\mathbf{w} \|\mathbf{x}\|^{\gamma+\mu-1}, \quad \mathbf{w} > 0,$$

$$\alpha_0 \|\mathbf{x}\|^{\gamma} \le V(\mathbf{x}) \le \alpha_1 \|\mathbf{x}\|^{\gamma}, \qquad (7)$$

$$\left\|\frac{\partial V(\mathbf{x})}{\partial \mathbf{x}}\right\| \le \beta \|\mathbf{x}\|^{\gamma-1}, \quad \left\|\frac{\partial^2 V(\mathbf{x})}{\partial \mathbf{x}^2}\right\| \le \psi \|\mathbf{x}\|^{\gamma-2},$$

where α_0 , α_1 , β , $\psi > 0$. Given $\alpha > 1$, introduce the set

$$S_{\alpha} = \Big\{ \varphi \in C_{[-h,0]} \, \Big| \, \|\varphi(\theta)\| \le \alpha \|\varphi(0)\|, \ \theta \in [-h,0] \Big\}.$$

It was shown in [12] that to analyse the stability it is enough to compute a lower bound for the Lyapunov–Krasovskii functional on the set S_{α} , and to ensure the negativity of the time derivative along the solutions of system (1). The set S_{α} also plays an important role in the construction of the estimates for solutions in Section IV.

The following constants will be used in the sequel:

$$m = m_1 + m_2, \quad \eta = \eta_{11} + \eta_{12}, \quad p = p_1 + p_2,$$

$$q = q_{11} + q_{12}, \quad \kappa_1 = \psi m_2 + \beta \eta_{12}, \quad \kappa_2 = \psi p_2 + \beta q_{12},$$

$$L_1 = \psi m + \beta \eta, \quad L_2 = \psi p + \beta q,$$

$$L_3 = L_2 + \beta (q_{21} + q_{22}).$$

III. CONSTRUCTION OF THE FUNCTIONAL

In this section, we construct a Lyapunov–Krasovskii functional for system (1) with perturbations either of class (a) or (b). Following [1], we introduce the integral

$$L(t,\varepsilon) = \int_0^{t+h} e^{-\varepsilon(t+h-s)} B(s) ds, \qquad (8)$$

where $\varepsilon = 0$ in case (a), and $\varepsilon > 0$ in case (b). It is known [13] that in case (b) integral (8) satisfies the following property: there exists a function $\omega(\varepsilon)$ such that $\omega(\varepsilon) \to 0$ as $\varepsilon \to 0$, and

$$\varepsilon \| L(t,\varepsilon) \| \le \omega(\varepsilon)$$

for all $t \ge 0$. In case (a), there is $l_0 > 0$ such that $||L(t,0)|| \le l_0$, thus we set $\omega(\varepsilon) = l_0 \varepsilon$.

Inspired by the Lyapunov function for system (1) presented in [1] and using the construction of Lyapunov-Krasovskii functionals for homogeneous systems introduced in [8], [9], we propose the following functional for system (1):

$$v(t,\varphi) = V(\varphi(0)) + \left(\frac{\partial V(\mathbf{x})}{\partial \mathbf{x}}\right)^T \Big|_{\mathbf{x}=\varphi(0)}$$
(9)

$$\times \left(\int_{-h}^0 \left(f(\varphi(0),\varphi(\theta)) + B(t+\theta+h)Q(\varphi(0),\varphi(\theta))\right) d\theta - L(t,\varepsilon)Q(\varphi(0),\varphi(0))\right)$$

$$+ \int_{-h}^0 (\mathbf{w}_1 + (h+\theta)\mathbf{w}_2) \|\varphi(\theta)\|^{\gamma+\mu-1} d\theta.$$

Here, $w_1, w_2 > 0$ are such that $w_0 = w - w_1 - hw_2 > 0$. Notice that functional (9) does not coincide with the construction from [9] applied with the Lyapunov function of [1]. We prove that the Lyapunov–Krasovskii functional (9) is suitable for the stability analysis of system (1) below.

Lemma 1. There exist $\delta > 0$ and $c_0, c_1, c_2 > 0$ such that the time derivative of functional (9) along the solutions of system (1), admits a bound of the form

$$\frac{dv(t,x_t)}{dt} \le -c_0 \|x(t)\|^{\gamma+\mu-1} - c_1 \|x(t-h)\|^{\gamma+\mu-1} \quad (10)$$
$$-c_2 \int_{-h}^0 \|x(t+\theta)\|^{\gamma+\mu-1} d\theta,$$

in the neighbourhood $||x_t||_h \leq \delta$.

Proof. Differentiating each of the three summands of (9) along the solutions of (1), we obtain

$$\frac{dv(t,x_t)}{dt} = -\mathbf{w}_0 \|x(t)\|^{\gamma+\mu-1} - \mathbf{w}_1 \|x(t-h)\|^{\gamma+\mu-1} - \mathbf{w}_2 \int_{-h}^0 \|x(t+\theta)\|^{\gamma+\mu-1} d\theta + \sum_{j=1}^5 \Lambda_j,$$

where

$$\begin{split} \Lambda_{1} &= \left(\frac{\partial V(\mathbf{x})}{\partial \mathbf{x}}\right)^{T} \varepsilon L(t,\varepsilon)Q(\mathbf{x},\mathbf{x})\Big|_{\mathbf{x}=x(t)}, \\ \Lambda_{2} &= \left(\frac{\partial V(\mathbf{x})}{\partial \mathbf{x}}\right)^{T} \\ &\times \int_{t-h}^{t} \left[\frac{\partial f(\mathbf{x},x(s))}{\partial \mathbf{x}} + B(s+h)\frac{\partial Q(\mathbf{x},x(s))}{\partial \mathbf{x}}\right] ds\Big|_{\mathbf{x}=x(t)} \\ &\times \left(f(x(t),x(t-h)) + B(t)Q(x(t),x(t-h))\right), \\ \Lambda_{3} &= -\left(\frac{\partial V(\mathbf{x}_{1})}{\partial \mathbf{x}_{1}}\right)^{T} L(t,\varepsilon) \sum_{j=1}^{2} \frac{\partial Q(\mathbf{x}_{1},\mathbf{x}_{2})}{\partial \mathbf{x}_{j}}\Big|_{\mathbf{x}_{1}=\mathbf{x}_{2}=x(t)} \\ &\times \left(f(x(t),x(t-h)) + B(t)Q(x(t),x(t-h))\right), \\ \Lambda_{4} &= \left(f(x(t),x(t-h)) + B(t)Q(x(t),x(t-h))\right)^{T} \\ &\times \left(\frac{\partial^{2}V(\mathbf{x})}{\partial \mathbf{x}^{2}}\right)\Big|_{\mathbf{x}=x(t)} \int_{-h}^{0} \left(f(x(t),x(t+\theta)) \\ &+ B(t+\theta+h)Q(x(t),x(t+\theta))\right) d\theta, \\ \Lambda_{5} &= -\left(f(x(t),x(t-h)) + B(t)Q(x(t),x(t-h))\right)^{T} \\ &\times \left(\frac{\partial^{2}V(\mathbf{x})}{\partial \mathbf{x}^{2}}\right)\Big|_{\mathbf{x}=x(t)} L(t,\varepsilon)Q(x(t),x(t)). \end{split}$$

Next, in concordance with (2), (6), (7) and using the inequality

$$\begin{aligned} \gamma_1^{l_1} \gamma_2^{l_2} \gamma_3^{l_3} &\leq \gamma_1^{l_1+l_2+l_3} + \gamma_2^{l_1+l_2+l_3} + \gamma_3^{l_1+l_2+l_3}, \\ \gamma_1, \gamma_2, \gamma_3 &\geq 0, \quad l_1, l_2, l_3 > 0, \end{aligned}$$

we estimate the terms Λ_j separately:

$$\begin{split} |\Lambda_{1}| &\leq \beta p \omega(\varepsilon) \|x(t)\|^{\gamma+\sigma-1}, \\ |\Lambda_{2}| &\leq \beta m \eta h \|x(t)\|^{\gamma+2\mu-2} + \beta m_{2} \eta h \|x(t-h)\|^{\gamma+2\mu-2} \\ &+ \beta m \eta_{12} \int_{-h}^{0} \|x(t+\theta)\|^{\gamma+2\mu-2} d\theta \\ &+ \beta \hat{b}h(qm+p\eta) \|x(t)\|^{\gamma+\mu+\sigma-2} \\ &+ \beta \hat{b}h(qm_{2}+p_{2}\eta) \|x(t-h)\|^{\gamma+\mu+\sigma-2} \\ &+ \beta \hat{b}(q_{12}m+p\eta_{12}) \int_{-h}^{0} \|x(t+\theta)\|^{\gamma+\mu+\sigma-2} d\theta \\ &+ \beta \hat{b}^{2} p q h \|x(t)\|^{\gamma+2\sigma-2} + \beta \hat{b}^{2} p_{2} q h \|x(t-h)\|^{\gamma+2\sigma-2} \\ &+ \beta \hat{b}^{2} p q_{12} \int_{-h}^{0} \|x(t+\theta)\|^{\gamma+2\sigma-2} d\theta, \end{split}$$

$$\begin{split} |\Lambda_4| &\leq \psi m^2 h \|x(t)\|^{\gamma+2\mu-2} + \psi m m_2 h \|x(t-h)\|^{\gamma+2\mu-2} \\ &+ \psi m m_2 \int_{-h}^0 \|x(t+\theta)\|^{\gamma+2\mu-2} d\theta \\ &+ 2 \hat{b} \psi m p h \|x(t)\|^{\gamma+\mu+\sigma-2} \\ &+ \hat{b} \psi (m p_2 + m_2 p) h \|x(t-h)\|^{\gamma+\mu+\sigma-2} d\theta \\ &+ \psi \hat{b}^2 p^2 h \|x(t)\|^{\gamma+2\sigma-2} + \psi \hat{b}^2 p p_2 h \|x(t-h)\|^{\gamma+2\sigma-2} \\ &+ \psi \hat{b}^2 p p_2 \int_{-h}^0 \|x(t+\theta)\|^{\gamma+2\sigma-2} d\theta, \\ |\Lambda_3 + \Lambda_5| &\leq L_3 \frac{\omega(\varepsilon)}{\varepsilon} \Big(m \|x(t)\|^{\gamma+\mu+\sigma-2} \\ &+ \hat{b} p \|x(t)\|^{\gamma+2\sigma-2} + m_2 \|x(t-h)\|^{\gamma+\mu+\sigma-2} \\ &+ \hat{b} p_2 \|x(t-h)\|^{\gamma+2\sigma-2} \Big). \end{split}$$

Notice that $\gamma+2\mu-2>\gamma+\mu-1$ due to $\mu>1.$ Moreover, $\sigma>1$ leads to

$$\begin{aligned} \gamma+\mu+\sigma-2 > \gamma+\mu-1, \\ \gamma+2\sigma-2 > \gamma+\mu-1 \end{aligned}$$

both in cases (a) and (b). The term Λ_1 is absent in case (a) when $\varepsilon = 0$. In case (b), the degree $\gamma + \sigma - 1 \ge \gamma + \mu - 1$ whilst the coefficient $\omega(\varepsilon)$ can be done arbitrarily small by the choice of ε . Hence, bound (10) holds in a neighbourhood $||x_t||_h \le \delta$ with

$$\begin{aligned} c_0 &= \mathbf{w}_0 - \beta p \omega(\varepsilon) \delta^{\sigma-\mu} - mhL_1 \delta^{\mu-1} \\ &- \left(\hat{b}h(pL_1 + mL_2) + mL_3 \frac{\omega(\varepsilon)}{\varepsilon} \right) \delta^{\sigma-1} \\ &- p \hat{b} \left(\hat{b}hL_2 + L_3 \frac{\omega(\varepsilon)}{\varepsilon} \right) \delta^{2\sigma-\mu-1}, \\ c_1 &= \mathbf{w}_1 - m_2 hL_1 \delta^{\mu-1} \\ &- \left(\hat{b}h(p_2L_1 + m_2L_2) + m_2L_3 \frac{\omega(\varepsilon)}{\varepsilon} \right) \delta^{\sigma-1} \\ &- p_2 \hat{b} \left(\hat{b}hL_2 + L_3 \frac{\omega(\varepsilon)}{\varepsilon} \right) \delta^{2\sigma-\mu-1}, \\ c_2 &= \mathbf{w}_2 - m\kappa_1 \delta^{\mu-1} - \hat{b}(p\kappa_1 + m\kappa_2) \delta^{\sigma-1} \\ &- p \hat{b}^2 \kappa_2 \delta^{2\sigma-\mu-1}. \end{aligned}$$

Here, $\delta > 0$ is chosen in such a way that the constants c_0, c_1, c_2 are positive. Notice that if we replace $\omega(\varepsilon)$ with zero and $\omega(\varepsilon)/\varepsilon$ with l_0 in the above expressions, we recover case (a).

Lemma 2. There exist $\delta > 0$ and $a_1(\alpha) > 0$ such that functional (9) admits on the set S_{α} a lower bound of the form

$$v(t,\varphi) \ge a_1(\alpha) \|\varphi(0)\|^{\gamma} + \mathbf{w}_1 \int_{-h}^0 \|\varphi(\theta)\|^{\gamma+\mu-1} d\theta, \quad (11)$$

where $\|\varphi\|_h \leq \delta$.

Proof. Using bounds (2), (6) and (7), we estimate the second summand of functional (9). First,

$$\left| \left(\frac{\partial V(\mathbf{x})}{\partial \mathbf{x}} \right)^T \right|_{\mathbf{x} = \varphi(0)} L(t, \varepsilon) Q(\varphi(0), \varphi(0)) \right|$$

$$\leq \beta p \frac{\omega(\varepsilon)}{\varepsilon} \|\varphi(0)\|^{\gamma + \sigma - 1}.$$

Second, the remaining term is estimated on the set S_{α} :

$$\left| \left(\frac{\partial V(\mathbf{x})}{\partial \mathbf{x}} \right)^T \right|_{\mathbf{x}=\varphi(0)} \int_{-h}^0 \left(f(\varphi(0),\varphi(\theta)) + B(t+\theta+h) \right) \\ \times Q(\varphi(0),\varphi(\theta)) d\theta \leq \beta h(m_1+m_2\alpha^\mu) \|\varphi(0)\|^{\gamma+\mu-1} \\ + \beta \hat{b}h(p_1+p_2\alpha^\sigma) \|\varphi(0)\|^{\gamma+\sigma-1}, \quad \varphi \in S_\alpha.$$

Hence, the required bound (11) holds with

$$a_1(\alpha) = \alpha_0 - \beta h(m_1 + m_2 \alpha^{\mu}) \delta^{\mu - 1} - \left(\beta \hat{b} h(p_1 + p_2 \alpha^{\sigma}) + \beta p \frac{\omega(\varepsilon)}{\varepsilon}\right) \delta^{\sigma - 1},$$

where $\delta > 0$ is chosen in such a way that $a_1(\alpha) > 0$. \Box

Lemma 3. Functional (9) admits the upper bound of the form

$$v(t,\varphi) \le b_0 \|\varphi(0)\|^{\gamma} + b_1 \int_{-h}^0 \|\varphi(\theta)\|^{\gamma} d\theta, \quad b_0, b_1 > 0,$$
(12)

in the neighbourhood $\|\varphi\|_h \leq \delta$. Moreover, there exist $b_2, b_3 > 0$ such that

$$v(t,\varphi) \le \alpha_1 \|\varphi(0)\|^{\gamma} + b_2 \|\varphi\|_h^{\gamma+\mu-1} + b_3 \|\varphi\|_h^{\gamma+\sigma-1}.$$
 (13)

Proof. Making the estimations in the same way as in Lemma 2, we arrive at the required bounds with

$$b_{0} = \alpha_{1} + \beta m h \delta^{\mu - 1} + \beta p \left(\hat{b}h + \frac{\omega(\varepsilon)}{\varepsilon} \right) \delta^{\sigma - 1},$$

$$b_{1} = (\beta m_{2} + w_{1} + h w_{2}) \delta^{\mu - 1} + \beta p_{2} \hat{b} \delta^{\sigma - 1},$$

$$b_{2} = (\beta m + w_{1} + h w_{2})h, \quad b_{3} = \beta p \left(\hat{b}h + \frac{\omega(\varepsilon)}{\varepsilon} \right).$$

Lemmas 1–3 imply that functional (9) allows verifying Theorem 1 [12], with slightly different assumptions on the right-hand sides.

IV. ESTIMATES

A. Estimates for the attraction region

Based on the classical ideas [14], [15] and the bounds presented in Lemmas 1–3, we obtain the following estimate for the attraction region of the trivial solution of system (1), see [9] for the use of the set S_{α} in this context.

Theorem 2. Let system (3) be asymptotically stable and Δ be a positive root of equation

$$\alpha_1 \Delta^{\gamma} + b_2 \Delta^{\gamma+\mu-1} + b_3 \Delta^{\gamma+\sigma-1} = a_1(\alpha) \delta^{\gamma}, \qquad (14)$$

where δ is defined in Lemmas 1–2. Then, the set of initial functions $\|\varphi\|_h < \Delta$ is the estimate of the attraction region of the trivial solution of (1).

Remark 1. It follows from the proof of Theorem 2 that $\|\varphi\|_h < \Delta$ implies $\|x_t\|_h \leq \delta$ for all $t \geq 0$.

B. Estimates for the solutions

In [16], a novel approach for the calculation of the estimates of the solutions is presented, which combines the use of Lyapunov-Krasovskii functionals with ideas of the Razumikhin framework. In this section, we apply this approach to system (1) using functional (9). First, as in [16], [17], we connect functional $v(t, \varphi)$ with its time derivative based on Lemmas 1 and 3:

$$\frac{dv(t,x_t)}{dt} \le -\rho v(t,x_t)^{\frac{\gamma+\mu-1}{\gamma}}, \quad t \ge 0,$$
(15)

where

$$\rho = \frac{c}{b^{\frac{\gamma+\mu-1}{\gamma}} \left(2 \max\{1,h\}\right)^{\frac{\mu-1}{\gamma}}},$$

 $b = \max\{b_0, b_1\}$ and $c = \min\{c_0, c_2\}$. Second, following the idea of [16], we consider a comparison equation of the form

$$\frac{du(t)}{dt} = -\tilde{\rho}u^{\frac{\gamma+\mu-1}{\gamma}}(t),$$
$$u(0) = u_0 = (\alpha_1 + b_2\Delta^{\mu-1} + b_3\Delta^{\sigma-1}) \|\varphi\|_h^{\gamma},$$

where $\tilde{\rho} < \rho$, and

$$1 + \tilde{\rho}h\left(\frac{\mu - 1}{\gamma}\right) \times (\alpha_1 + b_2\Delta^{\mu - 1} + b_3\Delta^{\sigma - 1})^{\frac{\mu - 1}{\gamma}}\Delta^{\mu - 1} \le \alpha^{\mu - 1}$$

The latter condition is necessary, since the lower bound in Lemma 2 holds on the set of functions S_{α} only. The solution for the comparison equation is

$$u(t) = u_0 \left[1 + \tilde{\rho} \left(\frac{\mu - 1}{\gamma} \right) u_0^{\frac{\mu - 1}{\gamma}} t \right]^{-\frac{\gamma}{\mu - 1}}.$$

Finally, we arrive at the following result.

Theorem 3. Let the trivial solution of system (1) be homogeneous and asymptotically stable. The solutions of system (1) with initial functions satisfying $\|\varphi\|_h < \Delta$, where Δ is defined in Theorem 2, admit an estimate of the form

$$\|x(t,\varphi)\| \le \hat{c}_1 \|\varphi\|_h \left[1 + \hat{c}_2 \|\varphi\|_h^{\mu-1} t\right]^{-\frac{1}{\mu-1}}$$

where

$$\hat{c}_1 = \left(\frac{\alpha_1 + b_2 \Delta^{\mu-1} + b_3 \Delta^{\sigma-1}}{a_1(\alpha)}\right)^{\frac{1}{\gamma}} = \frac{\delta}{\Delta},$$
$$\hat{c}_2 = \tilde{\rho} \left(\frac{\mu - 1}{\gamma}\right) \left(\alpha_1 + b_2 \Delta^{\mu-1} + b_3 \Delta^{\sigma-1}\right)^{\frac{\mu-1}{\gamma}}$$

Consider the system

$$\begin{pmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{pmatrix} = \begin{pmatrix} x_2^{\mu}(t) \\ -x_1^{\mu}(t) - x_2^{\mu}(t-h) \end{pmatrix}$$
(16)
+
$$\begin{pmatrix} \cos t + \sin(\sqrt{2}t) & 0 \\ 0 & \cos t + \cos(\sqrt{2}t) \end{pmatrix} \begin{pmatrix} x_1^{\sigma}(t-h) \\ x_2^{\sigma}(t) \end{pmatrix},$$

where $x_1(t)$, $x_2(t) \in \mathbb{R}$, $h \in \mathbb{R}^+$, and $\mu > 1$ is a number with odd numerator and denominator. In [18], the following Lyapunov function for the unperturbed and delay free part of system (16) was introduced:

$$V(x) = \frac{1}{\mu+1} \left(x_1^{\mu+1} + x_2^{\mu+1} \right) + \zeta x_1^{\mu} x_2, \quad \zeta > 0.$$
 (17)

It was shown that if

$$\zeta < \min\left\{\frac{1}{\mu+1}, \frac{4}{(\mu+1)^2}\right\},\,$$

then the Lyapunov function (17) allows to prove the asymptotic stability of the unperturbed system (16) with h = 0. Furthermore, the time-derivative of function (17) along the trajectories of system (16), when h = 0, admits an upper bound of the form

$$\begin{aligned} \frac{dV(x(t))}{dt} &\leq -\mathbf{w} \|x(t)\|^{2\mu}, \\ \text{where } \mathbf{w} &= \frac{\kappa}{2^{\mu-1}}, \text{ and} \\ \kappa &= \min\left\{1 - \zeta(\mu+1), \zeta, \frac{\zeta}{1+\zeta}\left(1 - \frac{\zeta(1+\mu)^2}{4}\right)\right\}. \end{aligned}$$

We take the system parameters $\sigma = \mu = 5$, h = 10, and set $\zeta = 0.0001$, $\alpha = 1.1$. Compute w = $6.2 \cdot 10^{-6}$. The constants characterising the Lyapunov function and its derivatives are the following:

$$\begin{split} &\alpha_0 = (1/2)^{\frac{\mu-1}{2}} \left(\frac{1}{\mu+1} - \zeta \right), \\ &\alpha_1 = \frac{1}{\mu+1} + \zeta, \\ &\beta = \sqrt{(1+\zeta\mu)^2 + (1+\zeta)^2}, \quad \psi = 2(\mu+\mu^2\zeta). \end{split}$$

The constants from bounds (2) and (6) are

$$m_1 = m_2 = \sqrt{2}, \quad \eta_{11} = \mu, \quad \eta_{12} = 0, \quad p_1 = p_2 = 1,$$

 $q_{11} = q_{22} = \sigma, \quad q_{12} = q_{21} = 0.$

Now, we calculate integral (8) and compute the function

$$\omega(\varepsilon) = \varepsilon \left(\frac{2\varepsilon + 1}{\varepsilon^2 + 1} + \frac{\max\{\varepsilon + 2\sqrt{2}, 2\varepsilon + \sqrt{2}\}}{\varepsilon^2 + 2} \right) \xrightarrow[\varepsilon \to 0]{} 0.$$

Choosing the initial function $\varphi(\theta) = [4.9 \cdot 10^{-4}, 4.9 \cdot 10^{-4}]^T$, $\theta \in [-10, 0]$, we compute the estimate for the solutions of (16) from Theorem 3. The constants characterising the estimate are shown in Table I. The system response and the computed bound are depicted in Fig. 1 as continuous and dashed lines, respectively.

 TABLE I

 CONSTANTS FOR THE ESTIMATES OF SOLUTIONS



Fig. 1. Estimation of the solution of system (16) in log scale

VI. CONCLUDING REMARKS

A Lyapunov-Krasovskii functional allowing the analysis of the solutions of homogeneous time-delay systems subject to time varying perturbations allows presenting estimates of the system solutions and of the domain of attraction. The classes of periodic and almost periodic perturbations with zero mean values are studied. An illustrative example validates the results.

REFERENCES

- A. Y. Aleksandrov and A. P. Zhabko, "On the asymptotic stability of solutions of nonlinear systems with delay," *Siberian Mathematical Journal*, vol. 53, no. 3, pp. 393–403, 2012.
- [2] A. Y. Aleksandrov, G.-D. Hu, and A. P. Zhabko, "Delay-independent stability conditions for some classes of nonlinear systems," *IEEE Transactions on Automatic Control*, vol. 59, no. 8, pp. 2209–2214, 2014.
- [3] D. Efimov, A. Polyakov, W. Perruquetti, and J.-P. Richard, "Weighted homogeneity for time-delay systems: finite-time and independent of delay stability," *IEEE Transactions on Automatic Control*, vol. 61, no. 1, pp. 210–215, 2016.
- [4] L. Rosier, "Homogeneous Lyapunov function for homogeneous continuous vector field," *Systems and Control Letters*, vol. 19, no. 6, pp. 467–473, 1992.
- [5] H. Hermes, "Homogeneous coordinates and continuous asymptotically stabilizing feedback controls," *Differential Equations, Stability and Control*, vol. 109, no. 1, pp. 249–260, 1991.
- [6] A. Y. Aleksandrov, E. Aleksandrova, and A. P. Zhabko, "Asymptotic stability conditions and estimates of solutions for nonlinear multiconnected time-delay systems," *Circuits, Systems, and Signal Processing*, vol. 35, no. 10, pp. 3531–3554, 2016.
- [7] A. Y. Aleksandrov, "The stability of equilibrium of non-stationary systems," *Journal of Applied Mathematics and Mechanics*, vol. 60, no. 2, pp. 199–203, 1996.
- [8] A. Y. Aleksandrov, A. P. Zhabko, and V. Pecherskiy, "Complete type functionals for some classes of homogeneous differential-difference systems," *Proc. 8th International Conference "Modern methods of applied mathematics, control theory and computer technology*", pp. 5–8 (in Russian), 2015.

- [9] A. P. Zhabko and I. V. Alexandrova, "Complete type functionals for homogeneous time delay systems," *Automatica*, vol. 125, p. 109456, 2021.
- [10] V. Kharitonov, *Time-delay systems: Lyapunov functionals and matrices*. Basel: Birkhäuser, 2013.
- [11] V. I. Zubov, *Methods of A.M. Lyapunov and their application*. P. Noordhoff, 1964.
- [12] I. V. Alexandrova and A. P. Zhabko, "At the junction of Lyapunov-Krasovskii and Razumikhin approaches," *IFAC-PapersOnLine*, vol. 51, no. 14, pp. 147–152, 2018.
- [13] A. M. Fink, Almost periodic differential equations. Springer, 1974.
- [14] N. N. Krasovskii, Certain Problems of Stability Theory of Motion. Moscow: Fizmatgiz, 1959. In Russian.
- [15] D. Melchor-Aguilar and S.-I. Niculescu, "Estimates of the attraction region for a class of nonlinear time-delay systems," *IMA J. Math. Control Information*, vol. 24, no. 4, pp. 523–550, 2007.
- [16] G. Portilla, I. V. Alexandrova, and S. Mondié, "Estimates for weighted homogeneous delay systems: a Lyapunov-Krasovskii-Razumikhin approach," *American Control Conference, Accepted*, 2021.
- [17] G. Portilla, I. V. Alexandrova, S. Mondié, and A. P. Zhabko, "Estimates for solutions of homogeneous time-delay systems: comparison of Lyapunov–Krasovskii and Lyapunov–Razumikhin techniques," *International Journal of Control, Submitted*, 2021.
- [18] A. Y. Aleksandrov, E. B. Aleksandrova, A. V. Ekimov, and N. V. Smirnov, A collection of tasks and exercises on the theory of stability: a handbook. Publishing House "Lan", 2016. In Russian.