

# An uniform version of Dvir and Moran's theorem

Gábor Hegedüs

Óbuda University

Bécsi út 96/B, Budapest, Hungary, H-1032

hegedus.gabor@uni-obuda.hu

May 11, 2021

## Abstract

Dvir and Moran proved the following upper bound for the size of a family  $\mathcal{F}$  of subsets of  $[n]$  with  $\text{Vdim}(\mathcal{F}\Delta\mathcal{F}) \leq d$ .

Let  $d \leq n$  be integers. Let  $\mathcal{F}$  be a family of subsets of  $[n]$  with  $\text{Vdim}(\mathcal{F}\Delta\mathcal{F}) \leq d$ . Then

$$|\mathcal{F}| \leq 2 \sum_{k=0}^{\lfloor d/2 \rfloor} \binom{n}{k}.$$

Our main result is the following uniform version of Dvir and Moran's result.

Let  $d \leq n$  be integers. Let  $\mathcal{F}$  be an uniform family of subsets of  $[n]$  with  $\text{Vdim}(\mathcal{F}\Delta\mathcal{F}) \leq d$ . Then

$$|\mathcal{F}| \leq 2 \binom{n}{\lfloor d/2 \rfloor}.$$

Denote by  $\mathbf{v}_F \in \{0, 1\}^n$  the characteristic vector of a set  $F \subseteq [n]$ .

Our proof is based on the following uniform version of Croot-Lev-Pach Lemma:

Let  $0 \leq d \leq n$  be integers. Let  $\mathcal{H}$  be a  $k$ -uniform family of subsets of  $[n]$ . Let  $\mathbb{F}$  be a field. Suppose that there exists a polynomial  $P(x_1, \dots, x_n, y_1, \dots, y_n) \in \mathbb{F}[x_1, \dots, x_n, y_1, \dots, y_n]$  with  $\deg(P) \leq d$

such that  $P(\mathbf{v}_F, \mathbf{v}_F) \neq 0$  for each  $F \in \mathcal{H}$  and  $P(\mathbf{v}_F, \mathbf{v}_G) = 0$  for each  $F \neq G \in \mathcal{H}$ . Then

$$|\mathcal{H}| \leq 2^{\binom{n}{\lfloor d/2 \rfloor}}.$$

## 1 Introduction

Throughout this paper  $n$  denotes a positive integer, and  $[n]$  stands for the set  $\{1, 2, \dots, n\}$ . We denote by  $2^{[n]}$  the set of all subsets of  $[n]$ . Subsets of  $2^{[n]}$  are called *set families*. Let  $\binom{[n]}{m}$  denote the family of all subsets of  $[n]$  which have cardinality  $m$ , and  $\binom{[n]}{\leq m}$  of all subsets that have size at most  $m$ .

A family  $\mathcal{F}$  of subsets of  $[n]$  is *k-uniform*, if  $|F| = k$  for each  $F \in \mathcal{F}$ .

Let  $\mathbb{F}$  be a field.  $\mathbb{F}[x_1, \dots, x_n] = \mathbb{F}[\mathbf{x}]$  denotes the ring of polynomials in the variables  $x_1, \dots, x_n$  over  $\mathbb{F}$ . For a subset  $F \subseteq [n]$  we write  $\mathbf{x}_F = \prod_{j \in F} x_j$ . In particular,  $\mathbf{x}_\emptyset = 1$ .

We denote by  $\mathbf{v}_F \in \{0, 1\}^n$  the characteristic vector of a set  $F \subseteq [n]$ .

It is a challenging old problem to find strong upper bounds for the size of progression-free subsets in finite Abelian groups. Croot, Lev and Pach achieved recently a breakthrough in this research area and proved a new exponential upper bound for the size of three-term progression-free subsets in the groups  $(\mathbb{Z}_4)^n$  (see [4]), where  $n \geq 1$  is an arbitrary integer. They based their proof on the following simple statement (see [4] Lemma 1).

**Proposition 1.1** *Suppose that  $n \geq 1$  and  $d \geq 0$  are integers,  $P$  is a multilinear polynomial in  $n$  variables of total degree at most  $d$  over a field  $\mathbb{F}$ , and  $A \subseteq \mathbb{F}^n$  is a subset with*

$$|A| > 2 \sum_{i=0}^{d/2} \binom{n}{i}.$$

*If  $P(\mathbf{a} - \mathbf{b}) = 0$  for all  $\mathbf{a}, \mathbf{b} \in A$ ,  $\mathbf{a} \neq \mathbf{b}$ , then  $P(\mathbf{0}) = 0$ .*

Consider a family  $\mathcal{F}$  of subsets of  $[n]$ . We say that  $\mathcal{F}$  *shatters*  $M \subseteq [n]$  if

$$\{F \cap M : F \in \mathcal{F}\} = 2^M.$$

Define

$$\text{sh}(\mathcal{F}) = \{M \subseteq [n] : \mathcal{F} \text{ shatters } M\}.$$

We say that a family  $\mathcal{F}$  has *VC-dimension*  $m$ , if  $m$  is the maximum of the size of sets shattered by  $\mathcal{F}$ . We denote by  $\text{Vdim}(\mathcal{F})$  the *VC-dimension* of a family  $\mathcal{F}$ .

The following result is fundamental in the theory of shattering.

**Theorem (Sauer[10], Perles, Shelah[11], Vapnik, Chervonenkis[12])**  
*Let  $\mathcal{F}$  be a family of subsets of  $[n]$  with  $\text{Vdim}(\mathcal{F}) \leq d$ . Then*

$$|\mathcal{F}| \leq \sum_{k=0}^d \binom{n}{k}$$

*and the upper bound is sharp.*

Let  $\mathcal{F}$  and  $\mathcal{G}$  be families of subsets of  $[n]$ . We denote by  $\mathcal{F}\Delta\mathcal{G}$  the symmetric difference of these families:

$$\mathcal{F}\Delta\mathcal{G} := \{A\Delta B : A \in \mathcal{F}, B \in \mathcal{G}\}.$$

Dvir and Moran proved an upper bound for the size of a family  $\mathcal{F}$  of subsets of  $[n]$  with  $\text{Vdim}(\mathcal{F}\Delta\mathcal{F}) \leq d$ . Their proof based on Proposition 1.1.

**Theorem 1.2** *Let  $0 \leq d \leq n$  be integers. Let  $\mathcal{F}$  be a family of subsets of  $[n]$  with  $\text{Vdim}(\mathcal{F}\Delta\mathcal{F}) \leq d$ . Then*

$$|\mathcal{F}| \leq 2 \sum_{k=0}^{\lfloor d/2 \rfloor} \binom{n}{k}.$$

Cambie, Girão and Kang proved the following improved version of Theorem 1.2 in [2].

**Theorem 1.3** *Let  $d < n$  be positive integers with  $d \equiv r \pmod{2}$  for some  $r \in \{0, 1\}$ . Let  $\mathcal{F}$  be a family of subsets of  $[n]$  with  $\text{Vdim}(\mathcal{F}\Delta\mathcal{F}) \leq d$ . Then*

$$|\mathcal{F}| \leq 2^r \sum_{k=0}^{\lfloor d/2 \rfloor} \binom{n-r}{k}.$$

Kleitman's theorem (see [7]) is an immediate consequence of Theorem 1.3.

**Corollary 1.4** *Let  $d < n$  be positive integers with  $d \equiv r \pmod{2}$  for some  $r \in \{0, 1\}$ . Let  $\mathcal{F}$  be a family of subsets of  $[n]$  with  $\mathcal{F}\Delta\mathcal{F} \subseteq \binom{[n]}{\leq d}$ . Then*

$$|\mathcal{F}| \leq 2^r \sum_{k=0}^{\lfloor d/2 \rfloor} \binom{n-r}{k}.$$

Our main result is the following uniform version of Theorem 1.2.

**Theorem 1.5** *Let  $d \leq n$  be integers. Let  $\mathcal{F}$  be an uniform family of subsets of  $[n]$  with  $\text{Vdim}(\mathcal{F}\Delta\mathcal{F}) \leq d$ . Then*

$$|\mathcal{F}| \leq 2 \binom{n}{\lfloor d/2 \rfloor}.$$

We prove here a new uniform version of Proposition 1.1. The proof of Theorem 1.5 is based completely on this result.

**Theorem 1.6** *Let  $0 \leq d \leq n$  be integers. Let  $\mathcal{H}$  be a  $k$ -uniform family of subsets of  $[n]$ . Let  $\mathbb{F}$  be a field. Suppose that there exists a polynomial  $P(\mathbf{x}, \mathbf{y}) \in \mathbb{F}[\mathbf{x}, \mathbf{y}]$  with  $\deg(P) \leq d$  (here  $\mathbf{x} = (x_1, \dots, x_n)$ ,  $\mathbf{y} = (y_1, \dots, y_n)$ ) such that  $P(\mathbf{v}_F, \mathbf{v}_F) \neq 0$  for each  $F \in \mathcal{H}$  and  $P(\mathbf{v}_F, \mathbf{v}_G) = 0$  for each  $F \neq G \in \mathcal{H}$ . Then*

$$|\mathcal{H}| \leq 2 \binom{n}{\lfloor d/2 \rfloor}.$$

In Section 2 we collected the preliminaries about Gröbner basis theory and standard monomials. In Section 3 we present our proofs. In Section 4 we give an interesting conjecture which strengthens our main result.

## 2 Preliminaries

Define  $V(\mathcal{F})$  as the subset  $\{\mathbf{v}_F : F \in \mathcal{F}\} \subseteq \{0, 1\}^n \subseteq \mathbb{F}^n$  for any family of subsets  $\mathcal{F} \subseteq 2^{[n]}$ .

It is natural to consider the ideal  $I(V(\mathcal{F}))$ :

$$I(V(\mathcal{F})) := \{f \in \mathbb{F}[\mathbf{x}] : f(\mathbf{v}) = 0 \text{ whenever } \mathbf{v} \in V(\mathcal{F})\}.$$

It is easy to verify that we can identify the algebra  $\mathbb{F}[\mathbf{x}]/I(V(\mathcal{F}))$  and the algebra of  $\mathbb{F}$  valued functions on  $V(\mathcal{F})$ . Consequently

$$\dim_{\mathbb{F}} \mathbb{F}[\mathbf{x}]/I(V(\mathcal{F})) = |\mathcal{F}|.$$

We recall some basic facts about Gröbner basis theory and standard monomials. We refer to [1], [3] for details.

We say that a linear order  $\prec$  on the monomials is a *term order*, if 1 is the minimal element of  $\prec$ , and  $\mathbf{u}\mathbf{w} \prec \mathbf{v}\mathbf{w}$  holds for any monomials  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  with  $\mathbf{u} \prec \mathbf{v}$ . The two most important term orders are the lexicographic order  $\prec_l$  and the deglex order  $\prec_d$ . Recall the definition of the deglex order: we have  $\mathbf{u} \prec_d \mathbf{v}$  iff either  $\deg \mathbf{u} < \deg \mathbf{v}$ , or  $\deg \mathbf{u} = \deg \mathbf{v}$ , and  $\mathbf{u} \prec_l \mathbf{v}$ .

The *leading monomial*  $\text{lm}(f)$  of a nonzero polynomial  $f \in \mathbb{F}[\mathbf{x}]$  is the  $\prec$ -largest monomial which appears with nonzero coefficient in  $f$ .

Let  $I$  be an ideal of  $\mathbb{F}[\mathbf{x}]$ . A finite subset  $\mathcal{G} \subseteq I$  is a *Gröbner basis* of  $I$  if for every  $f \in I$  there exists a  $g \in \mathcal{G}$  such that  $\text{lm}(g)$  divides  $\text{lm}(f)$ . It can be shown that  $\mathcal{G}$  is actually a basis of  $I$ , i.e.  $\mathcal{G}$  generates  $I$  as an ideal of  $\mathbb{F}[\mathbf{x}]$  (cf. [3] Corollary 2.5.6). A well-known fact is (cf. [1, Corollary 1.6.5, Theorem 1.9.1]) that every nonzero ideal  $I$  of  $\mathbb{F}[\mathbf{x}]$  has a Gröbner basis.

A monomial  $\mathbf{z} \in \mathbb{F}[\mathbf{x}]$  is a *standard monomial for  $I$*  if it is not a leading monomial for any  $f \in I$ . We denote by  $\text{sm}(I)$  the set of standard monomials of  $I$ .

Let  $\mathcal{F} \subseteq 2^{[n]}$  be a set family. It is easy to check that the standard monomials of the ideal  $I(\mathcal{F}) := I(V(\mathcal{F}))$  are square-free monomials.

It is a fundamental fact that  $\text{sm}(I)$  gives a basis of the  $\mathbb{F}$ -vector-space  $\mathbb{F}[\mathbf{x}]/I$ . This means that every polynomial  $g \in \mathbb{F}[\mathbf{x}]$  can be uniquely written in the form  $h + f$  where  $f \in I$  and  $h$  is a unique  $\mathbb{F}$ -linear combination of monomials from  $\text{sm}(I)$ . Consequently if  $g \in \mathbb{F}[\mathbf{x}]$  is an arbitrary polynomial and  $\mathcal{G}$  is a Gröbner basis of  $I$ , then we can reduce  $g$  with  $\mathcal{G}$  into a linear combination of standard monomials for  $I$ .

### 3 Proofs

Let  $0 \leq k \leq n/2$ , where  $k$  and  $n$  are integers. Let  $\mathcal{M}_{k,n}$  stand for the set of all monomials  $\mathbf{x}_G$  such that  $G = \{s_1 < s_2 < \dots < s_j\} \subset [n]$  for which  $j \leq k$  and  $s_i \geq 2i$  holds for every  $i$ ,  $1 \leq i \leq j$ . We write  $\mathcal{M}_k$  instead of the more

precise  $\mathcal{M}_{k,n}$ , if  $n$  is clear from the context. It is easy to check that

$$|\mathcal{M}_k| = \binom{n}{k}.$$

Let  $\mathcal{D}_{k,n}$  denote the set of all sets  $H = \{s_1 < s_2 < \dots < s_j\} \subset [n]$  for which  $j \leq k$  and  $s_i \geq 2i$  holds for every  $i$ ,  $1 \leq i \leq j$ .

We described completely the standard monomials of the complete uniform families of all  $k$  element subsets of  $[n]$  in [6].

**Theorem 3.1** *Let  $\prec$  an arbitrary term order such that  $x_n \prec \dots \prec x_1$ . Let  $0 \leq k \leq n$  be integers and define  $j := \min(k, n - k)$ . Then*

$$\text{sm}(V\binom{[n]}{k}) = \mathcal{M}_{j,n}.$$

**Corollary 3.2** *Let  $0 \leq k \leq n$  be integers and define  $j := \min(k, n - k)$ . Suppose that  $d \leq j$ . Then*

$$\mathcal{D}_{k,n} \cap \binom{[n]}{\leq d} = \mathcal{D}_{d,n}.$$

Let  $0 \leq k \leq n$  be arbitrary integers. Define the vector system

$$\mathcal{F}(n, k, 2) := V\binom{[n]}{k} \times V\binom{[n]}{k} \subseteq \{0, 1\}^{2n}.$$

It is easy to verify the following Corollary from Theorem 3.1.

**Corollary 3.3** *Let  $\prec$  an arbitrary term order such that  $x_n \prec \dots \prec x_1$ . Let  $0 \leq k \leq n$  be integers and define  $j := \min(k, n - k)$ . Then*

$$\text{sm}(\mathcal{F}(n, k, 2)) = \{x_{M_1} \cdot y_{M_2} : M_1, M_2 \in \mathcal{D}_{j,n}\} \subseteq \mathbb{F}[\mathbf{x}, \mathbf{y}].$$

Mészáros and Rónyai proved the following result in [8] Lemma 1 (see also [9] Theorem 7).

**Theorem 3.4** *Let  $\prec$  an arbitrary term order such that  $x_n \prec \dots \prec x_1$ . Let  $\mathcal{F}$  be a family of subsets of  $[n]$ . Then  $\text{sm}(V(\mathcal{F})) \subseteq \{x_U : U \in \text{sh}(\mathcal{F})\}$ .*

**Proof of Theorem 1.6:**

Consider the matrix  $M \in \mathbb{F}^{\mathcal{H} \times \mathcal{H}}$ , where  $M_{(F,G)} := P(\mathbf{v}_F, \mathbf{v}_G)$  for each  $F, G \in \mathcal{H}$ .

It follows from the assumptions that  $M$  is a diagonal matrix, where nonzero elements stand in the diagonal, hence

$$\text{rank}(M) = |\mathcal{H}|.$$

Let  $Q$  denote the reduction of  $P$  via the deglex Gröbner basis of  $I(\mathcal{F}(n, k, 2))$ . Then  $\deg(Q) \leq \deg(P) \leq d$  and  $M_{(F,G)} = Q(\mathbf{v}_F, \mathbf{v}_G)$  for each  $F, G \in \mathcal{H}$ .

Let  $j := \min(k, n - k)$ . It follows from Corollary 3.3 that we can write the polynomial  $Q$  into the form

$$Q(\mathbf{x}, \mathbf{y}) = \sum_{M_1, M_2 \in \mathcal{D}_{j,n}} c_{M_1, M_2} x_{M_1} \cdot y_{M_2} \in \mathbb{F}_2[\mathbf{x}, \mathbf{y}],$$

where  $c_{M_1, M_2} \in \mathbb{F}_2$  for each  $M_1, M_2 \in \mathcal{D}_{j,n}$ . After grouping the terms of the polynomial  $Q(\mathbf{x}, \mathbf{y})$  we get that

$$Q(\mathbf{x}, \mathbf{y}) = \sum_{M \in \mathcal{D}_{j,n} \cap \{U: |U| \leq \lfloor d/2 \rfloor\}} c_{M\mathbf{x}} g_M(\mathbf{y}) + \sum_{J \in \mathcal{D}_{j,n} \cap \{U: |U| \leq \lfloor d/2 \rfloor\}} d_J \mathbf{y}_J h_J(\mathbf{x}),$$

where  $c_M, d_J \in \mathbb{F}_2$ ,  $h_J(\mathbf{x}) \in \mathbb{F}_2[\mathbf{x}]$ ,  $g_M(\mathbf{y}) \in \mathbb{F}_2[\mathbf{y}]$  for each  $J, M \in \mathcal{D}_{j,n}$ .

Then it follows from Corollary 3.2 that

$$Q(\mathbf{x}, \mathbf{y}) = \sum_{M \in \mathcal{D}_{\lfloor d/2 \rfloor, n}} c_{M\mathbf{x}} g_M(\mathbf{y}) + \sum_{J \in \mathcal{D}_{\lfloor d/2 \rfloor, n}} d_J \mathbf{y}_J h_J(\mathbf{x}),$$

Since

$$|\mathcal{D}_{\lfloor d/2 \rfloor, n}| = \binom{n}{\lfloor d/2 \rfloor},$$

hence we get that

$$\text{rank}(M) \leq 2 \binom{n}{\lfloor d/2 \rfloor}.$$

It follows from the equality  $\text{rank}(M) = |\mathcal{H}|$  that

$$|\mathcal{H}| \leq 2 \binom{n}{\lfloor d/2 \rfloor}.$$

□

**Proof of Theorem 1.5:**

Let  $\mathbb{F} := GF(2)$ . It follows from Theorem 3.4 that

$$\text{sm}(V(\mathcal{F}\Delta\mathcal{F}), \prec_d) \subseteq \{x_U : U \in \text{sh}(\mathcal{F}\Delta\mathcal{F})\}.$$

Since  $\text{Vdim}(\mathcal{F}\Delta\mathcal{F}) \leq d$ , hence

$$\text{sm}(V(\mathcal{F}\Delta\mathcal{F}), \prec_d) \subseteq \{x_U : |U| \leq d\}.$$

Let  $\mathcal{G}$  denote a fixed deglex Gröbner basis of  $I(V(\mathcal{F}\Delta\mathcal{F}))$ . Denote by  $g : V(\mathcal{F}\Delta\mathcal{F}) \rightarrow \mathbb{F}$  the function where  $g(\mathbf{0}) = 1$  and  $g(\mathbf{v}_T) = 0$  for each  $T \in \mathcal{F}\Delta\mathcal{F} \setminus \{\emptyset\}$ .

If we reduce  $g$  with the Gröbner basis  $\mathcal{G}$ , we get the polynomial  $g' \in \mathbb{F}[\mathbf{x}]$ . Clearly  $\deg(g') \leq d$ , because  $g'$  is a linear combination of deglex standard monomials of  $I(V(\mathcal{F}\Delta\mathcal{F}))$  and  $\text{sm}(V(\mathcal{F}\Delta\mathcal{F}), \prec_d) \subseteq \{x_U : |U| \leq d\}$ . Since  $\mathcal{G}$  is a Gröbner basis of  $I(V(\mathcal{F}\Delta\mathcal{F}))$ , hence

$$g(\mathbf{v}_G) = g'(\mathbf{v}_G)$$

for each  $G \in \mathcal{F}\Delta\mathcal{F}$ .

Define the polynomial function  $f : V(\mathcal{F}) \times V(\mathcal{F}) \rightarrow \mathbb{F}$  by

$$f(\mathbf{x}, \mathbf{y}) := g'(\mathbf{x} + \mathbf{y}).$$

Then

$$f(\mathbf{v}_F, \mathbf{v}_F) = g'(\mathbf{0}) = g(\mathbf{0}) = 1$$

for each  $F \in \mathcal{F}$  and

$$f(\mathbf{v}_F, \mathbf{v}_G) = g'(\mathbf{v}_F + \mathbf{v}_G) = g'(\mathbf{v}_{F\Delta G}) = g(\mathbf{v}_{F\Delta G}) = 0$$

for each  $F, G \in \mathcal{F}$ , where  $F \neq G$ .

We can apply Theorem 1.6 with the choices  $\mathcal{H} := \mathcal{F}$  and  $P(\mathbf{x}, \mathbf{y}) := f(\mathbf{x}, \mathbf{y})$ . □

## 4 Concluding remarks

We think that the next conjecture is the best form of Theorem 1.5.

**Conjecture 1** *Let  $d < n$  be positive integers with  $d \equiv r \pmod{2}$  for some  $r \in \{0, 1\}$ . Let  $\mathcal{F}$  be an uniform family of subsets of  $[n]$  with  $\text{Vdim}(\mathcal{F}\Delta\mathcal{F}) \leq d$ . Then*

$$|\mathcal{F}| \leq 2^r \binom{n-r}{\lfloor d/2 \rfloor}.$$

## References

- [1] W. W. Adams and P. Lounstaunau, *An Introduction to Gröbner bases*. AMS (1994).
- [2] S. Cambie, A. Girão and R. J. Kang, VC dimension and a union theorem for set systems. *Elect. Journal of Comb.*, **26**, 1-8 (2019).
- [3] D. Cox, J. Little and D. O’Shea, *Ideals, Varieties, and Algorithms*. (Springer-Verlag, Berlin, Heidelberg, 1992)
- [4] E. Croot, V. Lev and P. Pach, Progression-free sets in  $\mathbb{Z}_4^n$ . *Annals of Math.*, **185**, 331-337 (2017)
- [5] Z. Dvir and S. Moran, A Sauer-Shelah-Perles Lemma for Sumsets. *Elect. Journal of Comb.*, **25(4)**, 4-8 (2018).
- [6] G. Hegedűs and L. Rónyai, Gröbner bases for complete uniform families, *Journal of Alg. Comb.* **17** (2003) 171–180 .
- [7] D. J. Kleitman, On a combinatorial conjecture of Erdős. *Journal of Comb. Theory*, **1(2)**, 209-214 (1966).
- [8] T. Mészáros and L. Rónyai, Some combinatorial applications of Gröbner bases. In: Algebraic Informatics (Winkler, F. ed.), 4th International Conference, CAI 2011, Linz, Proceedings. 65–83, Springer-Verlag, Heidelberg (2011)
- [9] S. Moran and C. Rashtchian, Shattered Sets and the Hilbert Function. In 41st International Symposium on Mathematical Foundations of Computer Science (MFCS 2016) (Vol. 58, p. 70). Schloss Dagstuhl–Leibniz-Zentrum für Informatik.
- [10] N. Sauer, On the density of families of sets, *J. Combin. Theory A* **13**, 145–147 (1972).
- [11] S. Shelah, A combinatorial problem; stability and order for models and theories in infinitary languages, *Pacific Journal of Math.* **41** 247–261 (1972)

- [12] V. N. Vapnik, A. Ya. Chervonenkis, On the uniform convergence of relative frequencies of events to their probabilities, *Theory of Probability and Appl.* **XVI** , 264–280.(1971)