REPRESENTATIVES FOR UNIPOTENT CLASSES AND NILPOTENT ORBITS

MIKKO KORHONEN^{*†}, DAVID I. STEWART[†], AND ADAM R. THOMAS[†]

ABSTRACT. Let G be a simple algebraic group over an algebraically closed field k of characteristic p. The classification of the conjugacy classes of unipotent elements of G(k) and nilpotent orbits of G on Lie(G) is well-established. One knows there are representatives of every unipotent class as a product of root group elements and every nilpotent orbit as a sum of root elements. We give explicit representatives in terms of a Chevalley basis for the *eminent* classes. A unipotent (resp. nilpotent) element is said to be eminent if it is not contained in any subsystem subgroup (resp. subalgebra), or a natural generalisation if G is of type D_n . From these representatives, it is straightforward to generate representatives for any given class. Along the way we also prove recognition theorems for identifying both the unipotent classes and nilpotent orbits of exceptional algebraic groups.

1. INTRODUCTION

Let G be a connected reductive algebraic group over an algebraically closed field k of characteristic p. Fix a maximal torus T of G, and positive system of roots; then it is well-known that every unipotent class has a representative which is a product of root group elements and that every nilpotent orbit has a representative which is a sum of root elements. However, it is not easy to extract such representatives from the existing literature and there are a few errors (some perpetrated by the second author). This note gives a definitive list when G is simple. For compactness, we provide the representatives of certain distinguished classes we call *eminent* classes. An eminent unipotent (resp. nilpotent) element is one which is not contained in any proper subsystem subgroup (resp. subalgebra) of G, or a natural generalisation if G has a factor of type D_n —the precise definition is given in the next section. Given representatives of the eminent classes in the simple case, it is then a routine task to find representatives of all classes for G. (See Section 2.4.)

Theorem 1.1. Let G be a simple k-group over a field $k = \overline{k}$ of characteristic p with Lie algebra Lie(G).

- (i) A nilpotent element $e \in \text{Lie}(G)$ is eminent if and only if it is G-conjugate to one of the representatives in Table 1 (G of classical type) or Table 2 (G of exceptional type). Moreover, no two representatives in Tables 1 and 2 are G-conjugate.
- (ii) A unipotent element u ∈ G is eminent if and only if it is G-conjugate to one of the representatives in Table 3 (G of classical type) or Table 4 (G of exceptional type). Moreover, no two representatives in Tables 3 and 4 are G-conjugate.

We note that the only time a regular element is not eminent is when G is of type D_n . Indeed, such elements are contained in generalised subsystem subgroups or subalgebras of type B_{n-1} .

We make a few remarks on the tables. An explanation of the notation used is given in Section 2.1; we observe Bourbaki doctrine when exhibiting representatives. When G = SL(V), SO(V) or Sp(V) is a classical group, we also supply the action of the elements on the natural *G*-module *V*, which characterises them up to conjugacy in *G*. Lastly, since we use the same representatives as [LS12, Tables 13.3, 14.1], one can invoke [LS12, Thms 19.1,19.2] to generate representatives for the classes in Table 4 from those in Table 2 as follows: a representative of the unipotent class with label *X* is $\prod_{j \in J} x_{\alpha_j}(1)$, where $\sum_{j \in J} e_{\alpha_j}$ is the representative of the nilpotent element with label *X*.

²⁰¹⁰ Mathematics Subject Classification. 17B45, 20G99.

^{*} Partially supported by SNSF grant 200021_146223 and NSFC grants 11771200 and 11931005.

[†] We thank the Isaac Newton Institute for Mathematical Sciences for support and hospitality during the programme Groups, Representations and Applications: New perspectives, when some of the work on this paper was undertaken. (EPSRC grant number EP/R014604/1.)

Type	Char.	Representation	Representative
A_n	any	V(n+1)	$e_{\alpha_1} + \dots + e_{\alpha_n}$
$B_n; n \ge 2$	$p \neq 2$	V(2n+1)	$e_{\alpha_1} + \dots + e_{\alpha_n}$
	p=2	D(n+1)	$e_{\alpha_1} + \dots + e_{\alpha_n}$
$C_n; n \ge 2$	any	V(2n)	$e_{\alpha_1} + \dots + e_{\alpha_n}$
	p=2	$W_l(n); 0 < l < \frac{n}{2}$	$\left(\sum_{i=1}^{n-1} e_{\alpha_i}\right) + e_{2\alpha_l + \dots + 2\alpha_{n-1} + \alpha_n}$
$D_n; n \ge 4$	p = 2	$W_l(n); \ \frac{n+1}{2} < l < n$	$\sum_{i=1}^{n-1} e_{\alpha_i} + e_{\beta_l}, \text{ where } \beta_l = \alpha_n + \sum_{i=2l-n-1}^{n-2} \alpha_i$

 Table 1. Eminent nilpotent representatives in classical types

Type	Char.	Label	Representative
G_2	any	G_2	$e_{10} + e_{01}$
	p = 3	$(\tilde{A}_1)_3$	$e_{21} + e_{32}$
F_4	any	F_4	$e_{1000} + e_{0100} + e_{0010} + e_{0001}$
	p=2	$F_4(a_2)$	$e_{0001} + e_{0011} + e_{1100} + e_{0120}$
	p=2	$(C_3)_2$	$e_{0001} + e_{1110} + e_{0120} + e_{1222}$
	p = 2	$(A_2A_1)_2$	$e_{0111} + e_{1121} + e_{0122} + e_{1220}$
E_6	any	E_6	$e_{10000} + e_{00000} + e_{01000} + e_{00100} + e_{00010} + e_{00001} \\ 0 $
	any	$E_6(a_1)$	$e_{10000} + e_{01000} + e_{00010} + e_{00001} + e_{00100} + e_{01100}$
	p=2	$E_{6}(a_{3})$	$e_{10000} + e_{00100} + e_{01100} + e_{00011} + e_{00110} + e_{01110}$
E_7	any	E_7	$e_{100000} + e_{000000} + e_{010000} + e_{001000} + e_{000100} + e_{000010} + e_{000010} + e_{000001}$
	any	$E_7(a_1)$	$e_{100000} + e_{010000} + e_{000100} + e_{000010} + e_{000001} + e_{001000} + e_{011000}$
	any	$E_7(a_2)$	$e_{100000} + e_{00000} + e_{01000} + e_{00100} + e_{00110} + e_{00011} + e_{00001}$
	p=2	$E_7(a_3)$	$e_{100000} + e_{000001} + e_{001000} + e_{011000} + e_{000110} + e_{001100} + e_{011100}$
	p=2	$(A_6)_2$	$e_{000110} + e_{000011} + e_{111000} + e_{011000} + e_{001100} + e_{011100} + e_{122100}$
E_8	any	E_8	$e_{1000000} + e_{0000000} + e_{0100000} + e_{0010000} + e_{0001000} + e_{0000100} + e_{0000010} + e_{0000001} + e_{0000001} + e_{0000001} + e_{0000001} + e_{0000001} + e_{0000001} + e_{00000001} + e_{00000001} + e_{000000001} + e_{000000001} + e_{000000000000000000000000000000000000$
	any	$E_8(a_1)$	$e_{1000000} + e_{0000000} + e_{0001000} + e_{0000100} + e_{0000010} + e_{0000001} + e_{0010000} + e_{0110000}$
	any	$E_8(a_2)$	$e_{1000000} + e_{000000} + e_{010000} + e_{000001} + e_{0011000} + e_{001100} + e_{0001100} + e_{0001100}$
	p=2	$E_{8}(a_{3})$	$e_{0000010} + e_{0000001} + e_{1100000} + e_{0110000} + e_{0110000} + e_{0011000} + e_{0001100} + e_{0111000}$
	p = 2	$E_8(a_4)$	$e_{1100000} + e_{0010000} + e_{0110000} + e_{0011000} + e_{0001100} + e_{0000110} + e_{0000011} + e_{0111000}$
	p = 2	$E_8(a_5)$	$e_{1100000} + e_{0000110} + e_{0000011} + e_{0110000} + e_{0011000} + e_{0111000} + e_{0011100} + e_{0011100}$
	p=2	$(D_7(a_1))_2$	$e_{0001000} + e_{0000001} + e_{1100000} + e_{0011000} + e_{0000011} + e_{011100} + e_{0111100} + e_{0121110}$
	p=2	$(D_7)_2$	$e_{1000000} + e_{0110000} + e_{0011000} + e_{0111000} + e_{0011100} + e_{0001110} + e_{0000111} + e_{1111111}$
	p=2	$(D_5A_2)_2$	$e_{0000011} + e_{0000111} + e_{1111000} + e_{111100} + e_{0121000} + e_{0111100} + e_{0011110} + e_{0111100} + e_{011100} + e_{011100} + e_{011100} + e_{011100} + e_{0111100} + e_{011100} + e_{011100} + e_{011100} + e_{01100} + e_{0100} + e_{000} + e_{000} + e_{000} + e_{000} + e$
	p = 3	$(A_7)_3$	$e_{0001110} + e_{0000111} + e_{1110000} + e_{1111000} + e_{0011100} + e_{0111100} + e_{0121000} + e_{0011111} \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0$

 Table 2. Eminent nilpotent representatives in exceptional types

Type	Char.	Representation	Representative
A_n	any	V(n+1)	$x_{\alpha_1}(1)\cdots x_{\alpha_n}(1)$
$B_n; n \ge 2$	$p \neq 2$	V(2n+1)	$x_{\alpha_1}(1)\cdots x_{\alpha_n}(1)$
	p=2	V(2n) + R	$x_{\alpha_1}(1)\cdots x_{\alpha_n}(1)$
$C_n; n \ge 2$	any	V(2n)	$x_{\alpha_1}(1)\cdots x_{\alpha_n}(1)$
$D_n; n \ge 4$	p = 2	V(2n-2l+2) + V(2l-2)	$\left(\prod_{i=1}^{n-1} x_{\alpha_i}(1)\right) x_{\beta_l}(1), \text{ where } \beta_l = \alpha_n + \sum_{i=2l-n-1}^{n-2} \alpha_i$
		$\frac{n+1}{2} < l < n$	

Table 3. Eminent unipotent representatives in classical types

Type	Char.	Label	Type	Char.	Label
G_2	any	G_2	E_8	any	E_8
	p = 3	$(\tilde{A}_1)_3$		any	$E_8(a_1)$
F_4	any	F_4		any	$E_8(a_2)$
	p = 2	$F_4(a_2)$		p=2	$E_8(a_3)$
	p = 2	$(\tilde{A}_2 A_1)_2$		p=2	$E_8(a_4)$
E_6	any	E_6		p=2	$E_8(a_5)$
	any	$E_{6}(a_{1})$		p=2	$(D_5A_2)_2$
	p = 2	$E_{6}(a_{3})$		p=3	$(A_7)_3$
E_7	any	E_7			
	any	$E_7(a_1)$			
	any	$E_7(a_2)$			
	p = 2	$E_7(a_3)$			

Table 4. Eminent unipotent representatives in exceptional types

It is often useful to have a means of recognising unipotent and nilpotent elements in exceptional types from easily computable invariants. For unipotent classes, this was accomplished by Lawther [Law09, Section 3]. Let $C = C_{\text{Lie}(G)}(x)$ for x a unipotent or nilpotent element. We use the following abbreviations.

JBS := List of Jordan block sizes

DS := Dimensions of the terms in the derived series of C

LS := Dimensions of the terms in the lower central series of C

ALG := Dimension of the Lie algebra of derivations of C

ALG' := Dimension of the derived subalgebra of the Lie algebra of derivations of C

- NIL := Dimension of the Nilradical of C
- NDS := Dimensions of the Lie normaliser of each term in the derived series of C

Type	Char.	Data for nilpotent orbits	Type	Char.	Data for unipotent classes
G_2	$p \neq 3$	JBS on V_{\min}	G_2	$p \neq 3$	JBS on V_{\min}
	p = 3	JBS on $\operatorname{Lie}(G)$		p = 3	JBS on $\operatorname{Lie}(G)$
F_4	p = 2	JBS on V_{\min} and $\operatorname{Lie}(G)$; DS	F_4	$p \leq 3$	JBS on V_{\min} and $\operatorname{Lie}(G)$
	p = 3	JBS on V_{\min} and $\operatorname{Lie}(G)$		$p \ge 5$	JBS on V_{\min}
	$p \ge 5$	JBS on V_{\min}	E_6	$p \neq 3$	JBS on V_{\min}
E_6	p = 2	JBS on $\text{Lie}(G)$; DS		p = 3	JBS on $\operatorname{Lie}(G)$
	p = 3	JBS on $\operatorname{Lie}(G)$; ALG	E_7	p = 2	JBS on $\text{Lie}(G)$; LS
	$p \ge 5$	JBS on V_{\min}		p = 3	JBS on V_{\min} and $\operatorname{Lie}(G)$
E_7	p = 2	JBS on $\text{Lie}(G)$; DS; ALG		$p \ge 5$	JBS on V_{\min}
	p = 3	JBS on V_{\min} and $\operatorname{Lie}(G)$; DS	E_8	p = 2	JBS on $\text{Lie}(G)$; LS
	$p \ge 5$	JBS on V_{\min}		$p \geq 3$	JBS on $\operatorname{Lie}(G)$
E_8	p=2	JBS on $\text{Lie}(G)$; DS; ALG; ALG'			
	p=3	JBS on $\text{Lie}(G)$; NDS; NIL			
	p = 5	JBS on $\text{Lie}(G)$; DS			
	$p \ge 5$	JBS on $\operatorname{Lie}(G)$			

Table 5. Recognition data for nilpotent orbits and unipotent classes.

Theorem 1.2. Let G be an exceptional k-group. Then Table 5 gives a list of data which is sufficient to identify the class of any nilpotent or unipotent element of G.

Remarks 1.3. (i) One important application of Theorem 1.2 is its use in the proof of Theorem 1.1(i) for G of exceptional type. To find the eminent nilpotent elements in Lie(G) the strategy is to list all maximal subsystem subalgebras up to conjugacy; for each \mathfrak{m} in that list, construct representatives of all distinguished orbits; and then use Theorem 1.2 to identify their class in G.

(ii) We supply auxiliary tables in Section 5; whenever at least two nilpotent orbits have the same Jordan block structure on V_{\min} (when it exists) and Lie(G) we list them in a table along with the data required to distinguish them, using the same abbreviations given before Proposition 1.2.

2. Preliminaries

2.1. Notation. Throughout G is a reductive k-group, and most often G is simple. Fix a Borel subgroup $B \leq G$ containing a maximal torus T. This defines a root system Φ with base $\Delta = \{\alpha_1, \ldots, \alpha_l\}$ of simple roots generating the positive roots Φ^+ as positive integral sums. We use the Bourbaki ordering for a Dynkin diagram a corresponding roots and follow [Bou82] in denoting roots by a string of numbers in the form of a Dynkin diagram corresponding to coefficients of a root expressed in terms of the simple roots. For example, the highest short root of an F_4 system is denoted 1232.

The adjoint action of G on Lie(G) defines a Cartan decomposition $\text{Lie}(G) \cong \text{Lie}(T) \oplus \bigoplus_{\alpha \in \Phi} ke_{\alpha}$. Corresponding to the root spaces ke_{α} one has the one-parameter subgroups $U_{\alpha} < G$, where each $U_{\alpha} \cong \mathbb{G}_a$ is a root groups. In case G is simple and simply connected, we may insist that the e_{α} form a Chevalley basis; usually, this is only recognised as canonical up to a choice of the sign of a coefficient λ for certain triples of roots (α, β, γ) such that $[e_{\alpha}, e_{\beta}] = \lambda e_{\gamma}$. These choices determine the structure constants for Lie(G) [Car89, Proposition 4.2.2]. Through exponentiation one can fix an isomorphism $ke_{\alpha} \to U_{\alpha}$ in which $ce_{\alpha} \mapsto x_{\alpha}(c)$ for all $c \in k$.

The computations done in Section 3 were performed with GAP4 [gro14] in the simply connected case, using the representatives for the nilpotent orbits given in [Ste16] (see the latest arXiv version for a correction in type F_4). In the adjoint case we used MAGMA [BCP97]. Although MAGMA uses a different choice of signs in the structure constants compared to GAP4, the same representatives can still be used and this is justified as follows.

All representatives listed in [Ste16] are of the form $e = \sum_{i=1}^{t} e_{\beta_i}$, where $\beta_1, \ldots, \beta_t \in \Phi^+$ are \mathbb{Z} -independent roots. Thus for any choice of signs $c_1, \ldots, c_t \in \{1, -1\}$ there exists a semisimple element in G that conjugates e to the nilpotent element $\sum_{i=1}^{t} c_i e_{\beta_i}$; see for example [Hum75, Lemma 16.2C]. In particular, the representatives in [Ste16] do not depend on the signs of structure constants in the Chevalley basis of Lie(G).

2.2. Eminent elements and generalised subsystem subgroups and subalgebras. Recall that a subgroup M of G is a *subsystem subgroup* if it is semisimple (the definition of which we assume to include connected) and normalised by a maximal torus.¹ Note that the derived subgroup of any Levi subgroup is a subsystem subgroup. Define also a *subsystem subalgebra* to be a subalgebra $\mathfrak{m} \leq \text{Lie}(G)$ with $\mathfrak{m} = \text{Lie}(M)$ for some subsystem subgroup M.

Define a generalised subsystem subgroup as follows. If M is a subsystem subgroup of G containing some simple factors M_1, \ldots, M_t of type D_{r_1}, \ldots, D_{r_t} where $t \ge 0$, then there is a semisimple subgroup $M_{i,l}$ of M_i of type $B_l B_{r_i-l-1}$ with $0 \le l \le r_i - 1$ (see the proof of Lemma 2.2). Then the generalised subsystem subgroups are obtained by replacing some subset of the M_i with the subgroups $M_{i,l}$. Accordingly, we define a generalised subsystem subalgebra \mathfrak{m} to be Lie(M) for a generalised subsystem subgroup M.

Recall that a unipotent or nilpotent element x is *distinguished* if it is contained in no proper Levi of G. The following definition is more restrictive.

Definition 2.1. A unipotent element $u \in G$ is called *eminent* if it is not contained in any proper generalised subsystem subgroup of G. A nilpotent element $e \in \text{Lie}(G)$ is called *eminent* if it is not contained in any proper generalised subsystem subalgebra of Lie(G).

For use in the proof of Theorem 1.1 we record the conjugacy classes of maximal generalised subsystem subgroups which are not Levi subgroups. The maximal generalised subsystem subalgebras follow immediately.

Lemma 2.2. Suppose that G is a classical group. Let M be a generalised subsystem subgroup which is maximal amongst proper ones, such that M is not a Levi subgroup. Then M is conjugate to precisely one subgroup with root system as given in the following table. Moreover, every subgroup in the table is a maximal generalised subsystem subgroup.

¹Unless G has a factor with root system Φ such that the pair (Φ, p) is $(B_n, 2)$, $(C_n, 2)$, $(F_4, 2)$ or $(G_2, 3)$, this implies it has a symmetric root system which is closed under sums. In any case, the relevant root systems can be obtained from the Dynkin diagram using a version of the Borel-de Siebenthal algorithm; see [MT11, Prop. 13.15].

G	Char.	М
A_n		
$B_n; n \ge 2$	$p \neq 2$	$B_m D_{n-m}; 1 \le m \le n$
	p=2	$B_m B_{n-m}; 1 \le m \le \frac{n}{2}$
	any	D_n
$C_n; n \ge 2$	any	$C_m C_{n-m}; 1 \le m \le \frac{n}{2}$
	p=2	D_n
$D_n; n \ge 4$	any	$D_m D_{n-m}; 2 \le m \le \frac{n}{2}$
	any	B_{n-1}
	$p \neq 2$	$B_m B_{n-m-1}; 1 \le m \le \frac{n}{2}$

 Table 6. Maximal generalised subsystem subgroups

Proof. For the subsystem subgroups this is a routine use of the Borel–de Siebenthal algorithm for closed subsystems [MT11, Prop. 13.12] and [MT11, Prop. 13.15(i),(ii)] for the 2-closed non-closed subsystems. When G has type D_n further consideration is required. The definition of generalised subsystem subgroups immediately implies that they will not be maximal unless we replace D_n itself with $B_m B_{n-m-1}$ with $0 \le m \le n-1$. When p = 2, the subgroups of type $B_m B_{n-m-1}$ with $1 \le m \le n-1$ are contained in B_{n-1} (indeed they are subsystem subgroups of B_{n-1} since p = 2) and so they are not maximal. We need to check that these subgroups are unique up to conjugacy and maximal amongst proper generalised subsystem subgroups. When $p \ne 2$, the subgroups of type $B_m B_{n-m-1}$ are the connected stabiliser of a decomposition of V into non-degenerate subspaces of dimensions 2m + 1 and 2n - 2m - 1 and thus are unique up to conjugacy since G acts transitively on non-degenerate subspaces of dimension k. Moreover, this implies they are maximal amongst all connected proper subgroups of G and thus amongst all proper generalised subsystem subgroups. When p = 2, a subgroup of type B_{n-1} is the stabiliser of a non-singular vector and is again unique up to conjugacy and maximal amongst all connected proper subgroups of G. □

We observe that eminence is invariant under isogenies.

Lemma 2.3. Let $\phi : G \to H$ be an isogeny of reductive k-groups.

- (i) The map ϕ induces a bijection between the unipotent varieties of G and H, and the set of generalised subsystem subgroups of G and H.
- (ii) If ϕ is a central isogeny then the map $d\phi$ induces a bijection between the nilpotent cones of G and H, and the set of generalised subsystem subalgebras of G and H.

Proof. The bijections of unipotent varieties and nilpotent cones follow from [Car93, Proposition 5.1.1] and [Jan04, 2.7]. Since the kernel of ϕ is central, it is contained in any maximal torus; as ϕ is surjective with finite kernel it follows that ϕ is a bijection on maximal tori. Thus M is a subsystem subgroup if and only if $\phi(M)$ is. If M is a generalised subsystem subgroup, the result follows from the definition.

2.3. Unipotent classes and nilpotent orbits in classical groups. Let G be simple of classical type. In light of Lemma 2.3, for the purposes of this paper it does no harm for us to assume that G = SL(V), Sp(V), or SO(V).

We will choose forms preserved by G on V consistent with [Jan73]. Let V have k-basis $\{v_1, v_2, \ldots, v_m\}$. For $G = \operatorname{Sp}(V)$ and m = 2n, we choose a non-degenerate alternating bilinear form on V defined by $(v_i, v_{n+i}) = 1 = -(v_{n+i}, v_i)$ and $(v_i, v_j) = 0$ for all other i, j. For $G = \operatorname{SO}(V)$ with V of odd dimension m = 2n+1, choose a non-degenerate symmetric bilinear form on V defined by $(v_i, v_{n+i}) = 1 = (v_{n+i}, v_i)$, $(v_{2n+1}, v_{2n+1}) = 2$ and $(v_i, v_j) = 0$ for all other i, j. When p = 2, we also define a quadratic form on V by $Q(v_i) = 0$ for $i \leq 2n$, $Q(v_{2n+1}) = 1$, such that Q has polarization (-, -) on V. Finally, for $\operatorname{SO}(V)$ with V of even dimension m = 2n choose a non-degenerate symmetric bilinear form on V defined by $(v_i, v_{n+i}) = 1 = (v_{n+i}, v_i)$ and $(v_i, v_j) = 0$ for all other i, j. And in this case for p = 2, define a quadratic form on V by $Q(v_i) = 0$ for all i, with polarization (-, -) on V.

If G = SL(V), the class of a unipotent or nilpotent element $x \in G$ is determined by its Jordan block sizes on V. In what follows, we will describe the unipotent classes and nilpotent orbits for G = Sp(V) and G = SO(V). In this case for x a unipotent element or a nilpotent element, we have a decomposition $V \downarrow k[x] = V_1 \perp \cdots \perp V_t$, where the V_i are orthogonally indecomposable k[x]-modules. Here orthogonally indecomposable means that if $V_i = W \perp W'$ as k[x]-modules, then W = 0 or W' = 0. Hesselink [Hes79, 3.5] determined the orthogonally indecomposable k[x]-modules that can occur, and thus classified the unipotent classes and nilpotent orbits for Sp(V) and O(V). The possibilities for orthogonally indecomposable k[x]-modules are as follows:

- The k[x]-module V(m) is non-degenerate of dimension m and x acts on V(m) with a single Jordan block of size m.
- The k[x]-module W(m) has dimension 2m, with $W(m) = W_1 \oplus W_2$ where W_i are totally singular of dimension m and x acts on each W_i with a single Jordan block of size m.
- The k[x]-module $W_l(m)$ only occurs when p = 2 and x is a nilpotent element. It is nondegenerate of dimension 2m and x acts with two Jordan blocks of size m. If $G = \operatorname{Sp}(V)$ then 0 < l < m/2 and $l = \min\{n : (x^{n+1}(v), x^n(v)) = 0 \text{ for all } v \in W_l(m)\}$. If $G = \operatorname{SO}(V)$ then $(m+1)/2 < l \le m$ and $l = \min\{n : Q(x^n(v)) = 0 \text{ for all } v \in W_l(m)\}$.
- The k[x]-module D(m) only occurs when G = SO(V) with $\dim(V)$ odd, p = 2 and x is a nilpotent element. It is degenerate of dimension 2m 1 and x acts with two Jordan blocks of sizes m and m 1.
- The k[x]-module R only occurs when G = SO(V) with $\dim(V)$ odd, p = 2 and x is a unipotent element. It is the 1-dimensional radical of V and so x acts trivially on it.

It is clear that the decomposition of $V \downarrow k[x]$ into orthogonally indecomposable summands determines the class of x. There are various ways of writing down the decomposition $V \downarrow k[x]$, we will use the *distinguished normal form* introduced in [LS12]. Since we are interested in the simple group SO(V), we state when a class in O(V) meets the subgroup SO(V) and if so, whether it splits into two SO(V)classes; see [LS12, Lemma 3.11, Propositions 5.25, 6.22]. The following lemma is a combination of results from [LS12, Chs. 5.2, 5.3, 5.6, 6.2, 6.8].

Lemma 2.4. Let G = Sp(V) or O(V) and let x be unipotent or nilpotent element of G or Lie(G), respectively. The following list states which orthogonal decompositions $V \downarrow k[x]$ can occur. Conversely, the multiset of orthogonal factors determines x uniquely up to conjugacy.

Furthermore, x is distinguished if and only if r = 0.

(i) If $p \neq 2$, then

$$V \downarrow k[x] = \sum_{i=1}^{r} W(m_i)^{a_i} \oplus \sum_{j=1}^{s} V(n_j),$$

where the n_j are distinct, and all are even (resp. odd) when G = Sp(V) (resp. O(V)). For G = O(V) and x unipotent, the class meets SO(V). For G = O(V), a class splits into two SO(V)-classes if and only if s = 0 and $a_i = 0$ for all odd m_i .

(ii) If x is unipotent, p = 2 and V is of even dimension, then

$$V \downarrow k[x] = \sum_{i=1}^{r} W(m_i)^{a_i} \oplus \sum_{j=1}^{t} V(2n_j)^{b_j}$$

where the n_j are distinct and $b_j \leq 2$ for all j. Such a class meets SO(V) if $\sum_j b_j$ is even and it splits if and only if t = 0.

(iii) If x is unipotent, p = 2 and V is of odd dimension, then

$$V \downarrow k[x] = \sum_{i=1}^{r} W(m_i)^{a_i} \oplus \sum_{j=1}^{t} V(2n_j)^{b_j} \oplus R$$

where the n_j are distinct and $b_j \leq 2$.

(iv) If x is nilpotent, p = 2 and G = Sp(V), then

$$V \downarrow k[x] = \sum_{i=1}^{r} W(m_i)^{a_i} \oplus \sum_{j=1}^{s} W_{l_j}(n_j) \oplus \sum_{k=1}^{t} V(2q_k)^{b_k}$$

where the sequences (n_j) , (l_j) and $(n_j - l_j)$ are strictly decreasing, $b_k \leq 2$ and for all j, k either $q_k > n_j - l_j$ or $q_k < l_j$.

(v) If x is nilpotent, p = 2 and G = O(V) with V of even dimension, then

$$V \downarrow k[x] = \sum_{i=1}^{r} W(m_i)^{a_i} \oplus \sum_{j=1}^{s} W_{l_j}(n_j)$$

where the sequences (n_j) , (l_j) and $(n_j - l_j)$ are strictly decreasing. Each class splits into two SO(V)-classes if and only if s = 0.

(vi) If x is nilpotent, p = 2 and V is of odd dimension, then

$$V \downarrow k[x] = \sum_{i=1}^{r} W(m_i)^{a_i} \oplus \sum_{j=1}^{s} W_{l_j}(n_j) \oplus D(m)$$

where the sequences (n_j) , (l_j) and $(n_j - l_j)$ are strictly decreasing and $m < l_s$.

Remark 2.5. Each simple k-group of classical type has a unique class of nilpotent and unipotent elements of highest dimension called *regular*. A representative of the regular class is well-known to be $\prod_i x_{\alpha_i}(1)$ for unipotent elements [Ste65, 3.2] and $\sum_i e_{\alpha_i}$ for nilpotent elements by [Spr66, 5.9] and [Ken87], where the product and sum are supported on all simple roots. From [LS12, 3.3.6 and pp. 60–61] we see that the regular elements are usually the only ones which have a Jordan block of largest possible size. In type A, this means that x acts with a single Jordan block. For the other classical cases from the lemma, with $m = \dim(V)$:

- (i) $V \downarrow k[x] = V(m-1) + V(1)$ if G = SO(V) and m is even, otherwise $V \downarrow k[x] = V(m)$;
- (ii) $V \downarrow k[x] = V(m)$ if G = Sp(V), and $V \downarrow k[x] = V(m-2) + V(2)$ if G = SO(V);
- (iii) $V \downarrow k[x] = V(m-1) + R;$
- (iv) $V \downarrow k[x] = V(m);$
- (v) $V \downarrow k[x] = W_{m/2}(m/2);$
- (vi) $V \downarrow k[x] = D((m+1)/2).$

2.4. Recovering all classes from the eminent ones. We present a brief discussion of how one can find a full set of representatives of the unipotent classes and nilpotent orbits iteratively from a set of representatives of the eminent classes and orbits. (If G is of exceptional type, one can simply look at the full list in [Ste16].) Since an element x is eminent in some generalised subsystem it suffices to give a set of Chevalley generators for maximal generalised subsystem subgroups. The Borel-de Siebenthal algorithm solves the problem in the case of subsystems. Hence the following lemma completes the picture.

Lemma 2.6. Let G be a simple k-group of type D_n with $p \neq 2$. Let $\beta = \alpha_m + \alpha_{m+1} + \ldots + \alpha_{n-2}$ and define:

$$H_{1} = \langle U_{\pm\alpha_{1}}, \dots, U_{\pm\alpha_{m-1}}, x_{\beta+\alpha_{n-1}}(t) x_{\beta+\alpha_{n}}(-t), x_{-\beta-\alpha_{n-1}}(t) x_{-\beta-\alpha_{n}}(-t) \mid t \in k \rangle,$$
$$H_{2} = \langle U_{\pm\alpha_{m+1}}, \dots, U_{\pm\alpha_{n-2}}, x_{\alpha_{n-1}}(t) x_{\alpha_{n}}(t), x_{-\alpha_{n-1}}(t) x_{-\alpha_{n}}(t) \mid t \in k \rangle.$$

Then the subgroups H_1 and H_2 commute, H_1 has type B_m and H_2 has type B_{n-m-1} .

Proof. This is proved in [Tes95, pp.67–68], with the generators amended to reflect our choice of stucture constants. \Box

If $p \neq 2$, then Theorem 1.1 implies that the unique eminent class of x in type B_m is regular. Then in view of the lemma, an eminent unipotent element in the subsystem subgroup $H := H_1H_2$ can be represented by $u = u_1u_2$, where

$$u_1 = x_{\alpha_1}(1) \cdots x_{\alpha_{m-1}}(1) x_{\beta+\alpha_{n-1}}(1) x_{\beta+\alpha_n}(-1) \quad \text{and} \quad u_2 = x_{\alpha_{m+1}}(1) \cdots x_{\alpha_{n-2}}(1) x_{\alpha_{n-1}}(1) x_{\alpha_n}(1).$$

Similarly, if e is a regular nilpotent element of Lie(H), then e is conjugate to $e_{\beta+\alpha_{n-1}} - e_{\beta+\alpha_n} + \sum_{i=1,i\neq m}^{n} (e_{\alpha_i})$. For even more representatives that work regardless of the choice of signs of the structure constants, see Remark 4.6.

2.5. Representatives for classical groups in small dimensions. To illustrate our main theorem, Tables 7–13 provide representatives for the unipotent classes and nilpotent orbits in classical algebraic groups of small dimension, as considered in [LS12, Section 8].

Type A_n is easily described. The class of x corresponds to a partition $d_1 + \cdots + d_t = n+1$. Viewed as an element of SL_{n+1} and put in Jordan normal form, the elements of the superdiagonal that x is supported on correspond to the simple roots that it is supported on. More specifically, a unipotent representative is $\prod_{k=1}^n x_{\alpha_k}(\varepsilon_k)$, and a nilpotent representative is $\sum_{k=1}^n \varepsilon_k e_{\alpha_k}$, where $\varepsilon_k = 1$ if $\sum_{j=1}^{i-1} d_j < k < \sum_{j=1}^i d_j$ for some $1 \leq i \leq t$, and $\varepsilon_k = 0$ otherwise.

In the remaining explicit examples we give, there is an injective map from the set of unipotent classes to the set of nilpotent orbits and we have chosen representatives in such a way that a representative for a unipotent class is given by $\prod_{i=1}^{t} x_{\beta_i}(1)$, where $\sum_{i=1}^{t} e_{\beta_i}$ is the corresponding representative for the corresponding nilpotent orbit. Therefore we only list representatives for nilpotent orbits, apart from SO₇(k) where the injection is not as natural. When $p \neq 2$, the characteristic is good for G, and so there is a Springer map, hence a fortiori a bijection from the set of unipotent classes to the set of nilpotent orbits. When p = 2 and $G = \text{Sp}_{2n}$, a unipotent class is mapped to a nilpotent orbit such that the decompositions in Lemma 2.4 (ii) and (iv) coincide. For a unipotent element of $\text{SO}_{2n}(k)$, the decomposition of the corresponding nilpotent element is described in [LS12, 6.3]. In the tables for the orthogonal groups, some representatives have two rows of decompositions. The first row corresponds to the case where $p \neq 2$, and the second row corresponds to the case where p = 2.

Representative x	Char.	$V \downarrow k[x]$
0	any	$W(1)^{2}$
e_{10}	any	W(2)
e_{01}	any	W(1) + V(2)
$e_{01} + e_{21}$	p = 2	$V(2)^{2}$
$e_{10} + e_{01}$	any	V(4)

Table 7. Representatives for nilpotent orbits and unipotent classes of $\text{Sp}_4(k)$.

Representative x	Char.	$V \downarrow k[x]$
0	any	$W(1)^{3}$
$e_{100} + e_{010}$	any	W(3)
e_{100}	any	W(2) + W(1)
$e_{100} + e_{001}$	any	W(2) + V(2)
$e_{010} + e_{001}$	any	W(1) + V(4)
$e_{001} + e_{221}$	p=2	$W(1) + V(2)^2$
e ₀₀₁	any	$W(1)^2 + V(2)$
$e_{010} + e_{001} + e_{221}$	any	V(4) + V(2)
$e_{100} + e_{010} + e_{001}$	any	V(6)
$e_{100} + e_{010} + e_{221}$ (nilp. only)	p = 2	$W_1(3)$

Table 8. Representatives for nilpotent orbits and unipotent classes of $\mathrm{Sp}_6(k).$

Representative x	Char.	$V \downarrow k[x]$
0	any	$W(1)^4$
$e_{1000} + e_{0100} + e_{0010}$	any	W(4)
$e_{1000} + e_{0100}$	any	W(3) + W(1)
$e_{1000} + e_{0100} + e_{0001}$	any	W(3) + V(2)
$e_{1000} + e_{0010}$	any	$W(2)^{2}$
e_{1000}	any	$W(2) + W(1)^2$
$e_{1000} + e_{0001}$	any	W(2) + W(1) + V(2)
$e_{0100} + e_{0001} + e_{2221}$	p=2	$W(2) + V(2)^2$
$e_{1000} + e_{0010} + e_{0001}$	any	W(2) + V(4)
e_{0001}	any	$W(1)^3 + V(2)$
$e_{0001} + e_{2221}$	p = 2	$W(1)^2 + V(2)^2$
$e_{0010} + e_{0001}$	any	$W(1)^2 + V(4)$
$e_{0010} + e_{0001} + e_{2221}$	any	W(1) + V(4) + V(2)
$e_{0100} + e_{0010} + e_{0001}$	any	W(1) + V(6)
$e_{0010} + e_{0001} + e_{0221} + e_{2221}$	p = 2	$V(4) + V(2)^2$
$e_{1000} + e_{0010} + e_{0001} + e_{0221}$	p=2	$V(4)^2$

$e_{0100} + e_{0010} + e_{0001} + e_{2221}$	any	V(6) + V(2)
$e_{1000} + e_{0100} + e_{0010} + e_{0001}$	any	V(8)
$e_{1000} + e_{0100} + e_{0010} + e_{2221}$ (nilp. only)	p = 2	$W_1(4)$
$e_{0100} + e_{0010} + e_{0221}$ (nilp. only)	p = 2	$W_1(3) + W(1)$

Table 9. Representatives for nilpotent orbits and unipotent classes of $\mathrm{Sp}_8(k).$

Representative x	Char.	$V \downarrow k[u]$
1	any	$W(1)^3 + V(1)$
		$W(1)^{3} + R$
$x_{100}(1)x_{010}(1)$	any	W(3) + V(1)
		W(3) + R
$x_{100}(1)$	any	W(2) + W(1) + V(1)
		W(2) + W(1) + R
$x_{100}(1)x_{001}(1)$	any	W(2) + V(3)
		W(2) + V(2) + R
$x_{001}(1)$	any	$W(1)^2 + V(3)$
		$W(1)^2 + V(2) + R$
$x_{010}(1)x_{001}(1)$	any	W(1) + V(5)
		W(1) + V(4) + R
$x_{100}(1)x_{122}(1)$	p = 2	$W(1) + V(2)^2 + R$
$x_{100}(1)x_{010}(1)x_{012}(1)$	p = 2	V(4) + V(2) + R
$x_{100}(1)x_{010}(1)x_{001}(1)$	any	V(7)
		V(6) + R

Table 10. Representatives for unipotent classes of $SO_7(k)$.

Representative x	Char.	$V \downarrow k[e]$
0	any	$W(1)^3 + V(1)$
		$W(1)^3 + D(1)$
$e_{100} + e_{010}$	any	W(3) + V(1)
		W(3) + D(1)
e_{100}	any	W(2) + W(1) + V(1)
		W(2) + W(1) + D(1)
$e_{100} + e_{001}$	any	W(2) + V(3)
		W(2) + D(2)
e_{001}	any	$W(1)^2 + V(3)$
		$W(1)^2 + D(2)$
$e_{010} + e_{001}$	any	W(1) + V(5)
		W(1) + D(3)
$e_{100} + e_{122}$	p = 2	$W_2(2) + W(1) + D(1)$
$e_{100} + e_{010} + e_{012}$	p = 2	$W_3(3) + D(1)$
$e_{100} + e_{010} + e_{001}$	any	V(7)
		D(4)

Table 11. Representatives for nilpotent orbits of $SO_7(k)$.

Representative x	$V \downarrow k[e]$	$V \downarrow k[u]$
0	$W(1)^{4}$	$W(1)^4$
$e_{10_0^0} + e_{01_0^0} + e_{00_0^1}$	W(4)	W(4)
$e_{10_0^0} + e_{01_0^0} + e_{00_1^0}$	W(4)	W(4)
$e_{100}^{}_0 + e_{010}^{}_0$	W(3) + W(1)	W(3) + W(1)
$e_{100}^{}_0 + e_{001}^{}_0$	$W(2)^{2}$	$W(2)^{2}$
$e_{10_0^0} + e_{00_1^0}$	$W(2)^{2}$	$W(2)^{2}$
e_{100}^{0}	$W(2) + W(1)^2$	$W(2) + W(1)^2$
$e_{10_0^0} + e_{00_0^1} + e_{00_1^0}$	W(2) + V(3) + V(1)	W(2) + V(3) + V(1)
	$W(2) + W_2(2)$	$W(2) + V(2)^2$
$e_{00_0^1} + e_{00_1^0}$	$W(1)^2 + V(3) + V(1)$	$W(1)^2 + V(3) + V(1)$
	$W(1)^2 + W_2(2)$	$W(1)^2 + V(2)^2$
$e_{01_0^0} + e_{00_0^1} + e_{00_1^0}$	W(1) + V(5) + V(1)	W(1) + V(5) + V(1)
	$W(1) + W_3(3)$	W(1) + V(4) + V(2)
$e_{10_0^0} + e_{01_0^0} + e_{00_0^1} + e_{11_1^0}$	V(5) + V(3)	V(5) + V(3)
	$W_{3}(4)$	$V(4)^{2}$
$e_{100}^{0} + e_{010}^{0} + e_{001}^{0} + e_{001}^{0}$	V(7) + V(1)	V(7) + V(1)
	$W_4(4)$	V(6) + V(2)

Table 12. Representatives for nilpotent orbits and unipotent classes of $SO_8(k)$.

Representative x	$V \downarrow k[e]$	$V \downarrow k[u]$
0	$W(1)^{5}$	$W(1)^{5}$
$e_{1000} + e_{0100} + e_{0010} + e_{0001} + e_{0001}$	W(5)	W(5)
$e_{100_0^0} + e_{010_0^0} + e_{001_0^0}$	W(4) + W(1)	W(4) + W(1)
$e_{1000}^{0} + e_{0100}^{0} + e_{0001}^{0}$	W(3) + W(2)	W(3) + W(2)
$e_{1000} + e_{0100}$	$W(3) + W(1)^2$	$W(3) + W(1)^2$
$e_{1000} + e_{0010}$	$W(2)^2 + W(1)$	$W(2)^2 + W(1)$
e_{1000}	$W(2) + W(1)^3$	$W(2) + W(1)^3$
$e_{1000} + e_{0100} + e_{0001} + e_{0001}$	W(3) + V(3) + V(1)	W(3) + V(3) + V(1)
	$W(3) + W_2(2)$	$W(3) + V(2)^2$
$e_{100_0^0} + e_{000_0^1} + e_{000_1^0}$	W(2) + W(1) + V(3) + V(1)	W(2) + W(1) + V(3) + V(1)
	$W(2) + W(1) + W_2(2)$	$W(2) + W(1) + V(2)^2$
$e_{000_0^1} + e_{000_1^0}$	$W(1)^3 + V(3) + V(1)$	$W(1)^3 + V(3) + V(1)$
	$W(1)^3 + W_2(2)$	$W(1)^3 + V(2)^2$
$e_{1000}^{0} + e_{0010}^{0} + e_{0001}^{0} + e_{0001}^{0} + e_{0001}^{0}$	W(2) + V(5) + V(1)	W(2) + V(5) + V(1)
	$W(2) + W_3(3)$	W(2) + V(4) + V(2)
$e_{001_0^0} + e_{000_0^1} + e_{000_1^0}$	$W(1)^2 + V(5) + V(1)$	$W(1)^2 + V(5) + V(1)$
	$W(1)^2 + W_3(3)$	$W(1)^2 + V(4) + V(2)$
$e_{010_0^0} + e_{001_0^0} + e_{000_0^1} + e_{011_1^0}$	W(1) + V(5) + V(3)	W(1) + V(5) + V(3)
	$W(1) + W_3(4)$	$W(1) + V(4)^2$
$e_{010_0^0} + e_{001_0^0} + e_{000_0^1} + e_{000_1^0}$	W(1) + V(7) + V(1)	W(1) + V(7) + V(1)
	$W(1) + W_4(4)$	W(1) + V(6) + V(2)
$e_{1000} + e_{0100} + e_{0010} + e_{0001} + e_{0001} + e_{0111}$	V(7) + V(3)	V(7) + V(3)
	$W_4(5)$	W(1) + V(6) + V(4)
$e_{1000} + e_{0100} + e_{0010} + e_{0001} + e_{0001} + e_{0001}$	W(1) + V(9) + V(1)	W(1) + V(9) + V(1)
	$W_{5}(5)$	V(8) + V(2)

Table 13. Representatives for nilpotent orbits and unipotent classes of $SO_{10}(k)$.

3. Proof of Theorem 1.2

For unipotent elements, the required result is given in [Law09, Section 3]. We describe the calculations in the nilpotent case. First we need the following lemma, which shows that for the action of nilpotent elements on Lie(G), the coincidences of Jordan block sizes for nilpotent elements on different orbits are the same in the adjoint case and the simply connected case.

Lemma 3.1. Let G_{sc} be a simply connected simple algebraic group over k of exceptional type, and let G_{ad} be the corresponding simple algebraic group of adjoint type. Furthermore, let x be either a unipotent

element of G_{sc} or a nilpotent element of $\text{Lie}(G_{sc})$. Then the Jordan block sizes of x on $\text{Lie}(G_{sc})$ and $\text{Lie}(G_{ad})$ are the same.

Proof. For types G_2 , F_4 and E_8 we have $G_{sc} = G_{ad}$, so it suffices to consider the case where G_{sc} is of type E_6 or E_7 . In this case, we show that $\text{Lie}(G_{ad}) \cong \text{Lie}(G_{sc})^*$ as G_{sc} -modules, from which the lemma follows. If $\text{Lie}(G_{sc})$ is simple, then $\text{Lie}(G_{sc})$ is an irreducible Weyl module and thus $\text{Lie}(G_{sc})^* \cong \text{Lie}(G_{sc})$ and $\text{Lie}(G_{ad}) \cong \text{Lie}(G_{sc})$ by [Jan03, Lemma II.2.13 b)].

If $\text{Lie}(G_{sc})$ is not simple $(p = 3 \text{ for } E_6 \text{ and } p = 2 \text{ for } E_7)$, then by [Hog82] there is a nonsplit short exact sequence

$$0 \to k \to \text{Lie}(G_{sc}) \to W \to 0$$

of G_{sc} -modules, where W is irreducible. Since W is self-dual, it follows that we have a nonsplit short exact sequence

$$0 \to W \to \text{Lie}(G_{sc})^* \to k \to 0.$$

Because $\text{Lie}(G_{sc})$ is a Weyl module with highest weight the highest root, by [Jan03, II.2.12 (4), Proposition II.2.14] we have $\text{Ext}_{G_{sc}}^1(k, W) \cong \text{Ext}_{G_{sc}}^1(W, k) \cong k$. Furthermore, the group $\text{Ext}_{G_{sc}}^1(k, W)$ is a k-vector space, and equivalence classes of extensions which are scalar multiples of each other correspond to isomorphic G_{sc} -modules. It follows then from $\text{Ext}_{G_{sc}}^1(k, W) \cong k$ that a nonsplit extension of k by W is unique up to isomorphism of G_{sc} -modules, and so must be isomorphic to $\text{Lie}(G_{sc})^*$.

By [Hog82] there is a nonsplit short exact sequence

$$0 \to W \to \text{Lie}(G_{ad}) \to k \to 0$$

of G_{sc} -modules, so we conclude that $\operatorname{Lie}(G_{ad}) \cong \operatorname{Lie}(G_{sc})^*$ as G_{sc} -modules.

The representatives and tables of Jordan blocks on the minimal and adjoint modules is given in [Ste16]. The remaining data is supplied via in-built functions in GAP4 [gro14] and MAGMA such as LieDerivedSeries. It is then straightforward to confirm there are no coincidences.

4. Proof of Theorem 1.1

We first treat the case where G is classical, making heavy use of Lemma 2.4. Throughout the section x and x' are unipotent elements of G or nilpotent elements of Lie(G). In light of Lemma 2.3, we take G to be SL(V), Sp(V), or SO(V) accordingly with natural module V. The proof is completed in the following two lemmas.

Lemma 4.1. Suppose that x is not conjugate to an element in Table 1 or Table 3. Then x is not eminent.

Proof. If x is not distinguished, then x is contained in a Levi subgroup or a Levi subalgebra and thus is not eminent. Therefore in what follows, we will assume that x is distinguished. This already completes the proof for G = SL(V), since the only distinguished elements are the regular ones. Suppose then that G = Sp(V) or G = SO(V). Comparing Tables 1 and 3 with Lemma 2.4, we can assume that $V \downarrow k[x]$ has at least two summands, and r = 0 in the notation of Lemma 2.4.

If $p \neq 2$, this means that $V \downarrow k[x] = \sum_{j=1}^{s} V(n_j)$ with $s \geq 2$. It follows that x is contained in the stabiliser of a non-trivial decomposition of V into non-degenerate subspaces, and thus x is contained in a proper generalised subsystem subgroup or subalgebra.

For p = 2, we must consider the unipotent and nilpotent cases separately. Suppose first that x is unipotent. If G is of type C_n , then the argument from the previous paragraph applies.

If G has type D_n , then by Lemma 2.4(ii) we have $V \downarrow k[x] = \sum_{j=1}^s V(2n_j)$, where s is even. When

there are at least four summands, it follows that x is contained in a maximal subsystem subgroup of type $D_m D_{n-m}$ (one may choose $m = n_1 + n_2$). For s = 2 there is one case remaining which is not in Table 3, namely $V \downarrow k[x] = V(2n-2) + V(2)$, corresponding to the regular unipotent class. It follows from the description of unipotent classes of B_{n-1} in [LS12, Section 6.8] that x is contained in the generalised subsystem subgroup of type B_{n-1} (conjugate to the element x' in SO(W) with $W \downarrow k[x'] = V(2n-2) \oplus R$).

If G has type B_n , then in the notation of Lemma 2.4(iii) we have $V \downarrow k[x] = V \downarrow k[x] = \sum_{j=1}^{s} V(2n_j) + R$, where s > 1. Similarly to the previous cases, it follows that x is contained in a subsystem subgroup of

where s > 1. Similarly to the previous cases, it follows that x is contained in a subsystem subgroup of type $D_m B_{n-m-1}$ (again one may choose $m = n_1 + n_2$).

It remains to consider the case where x is nilpotent. If G has type C_n , then $V \downarrow k[x]$ has at least two summands, each of them isomorphic to $V(2q_k)$ or $W_{l_j}(n_j)$. Therefore x is contained in the stabiliser of the non-trivial decomposition of V into two non-degenerate subspaces, which is a subsystem subalgebra of type $C_m C_{n-m}$. If G has type D_n , then $V \downarrow k[x]$ either has at least two summands isomorphic to $W_{l_j}(n_j)$, or is isomorphic to $W_n(n)$. In the first case, x is contained in a subsystem subalgebra of type $D_{n_j}D_{n-n_j}$. In the second case, it follows from [LS12, Section 5.6] that x is contained in the generalised subsystem subalgebra of type B_{n-1} (conjugate to the element x' in $\mathfrak{so}(W)$ with $W \downarrow k[x'] = D(n)$). Finally if G has type B_n , then $V \downarrow k[x]$ has at least one summand isomorphic to $W_{l_j}(n_j)$, and so x is contained in a subsystem subalgebra of type $D_{n_j}B_{n-n_j}$.

Lemma 4.2. The elements in Tables 1 and 3 are eminent.

Proof. Let x be such an element. It follows from Lemma 2.4 that x is distinguished. Therefore if x is not eminent, then it must be contained one of the maximal generalised subsystem subgroups or subalgebras described in Lemma 2.2.

Suppose first that x acts on V with a single Jordan block. Elements in the subgroups and subalgebras of Lemma 2.2 act on V with at least two Jordan blocks, so it follows that x is eminent. The remaining cases occur when p = 2 and in all of them x acts with two Jordan blocks on V. We treat them in turn.

Consider first the case where G = SO(V) is of type B_n , and let R be the 1-dimensional radical of V. We will show that elements from the subgroups and subalgebras of Lemma 2.2 act on V with at least three Jordan blocks, which proves that x must be eminent. For M < G of type D_n , we have $V \downarrow M = W \oplus R$, where W is the natural module for M = SO(W). Unipotent elements of M and nilpotent elements of Lie(M) act on W with an even number of Jordan blocks by Lemma 2.4 (ii) and (v), so in particular they have at least three Jordan blocks on V. The other possibility in Lemma 2.2 is M < G of type $B_m B_{n-m}$ for $1 \le m \le n/2$. In this case $(V/R) \downarrow M = V_1 \oplus V_2$, where V_1 is 2m-dimensional and V_2 is 2(n-m)-dimensional. Thus unipotent elements of M and nilpotent elements of Lie(M) act on V/R with at least three Jordan blocks by Lemma 2.4 (ii) and (vi), and therefore they also act on V with at least three Jordan blocks by Lemma 2.4 (iii) and (vi), and therefore they also act on V with at least three Jordan blocks by Lemma 2.4 (iii) and (vi), and therefore they also act on V with at least three Jordan blocks.

If $G = \operatorname{Sp}(V)$ of type C_n , then x is nilpotent and $V \downarrow k[x] = W_l(n)$ for some $0 < l < \frac{n}{2}$. Such an element x acts with two Jordan blocks of size n on V. By considering each of the maximal subsystem subalgebras, we see that only those of type $C_{\frac{n}{2}}C_{\frac{n}{2}}$ (if n is even) and D_n contain elements that act with two Jordan blocks of size n. In type $C_{\frac{n}{2}}C_{\frac{n}{2}}$ the only element which does that acts as V(n) + V(n) on V,

which is already in canonical form and thus not conjugate to x. In the second case, the elements x' which act with two Jordan blocks in SO(V') have $V' \downarrow k[x'] = W_l(n)$ for some $\frac{n+1}{2} < l \leq n$. The definition of $W_l(n)$ for orthogonal groups in even dimension, given in [LS12, p. 66], shows that all of these elements x' are conjugate to a nilpotent element acting via W(n) in Lie(G), and thus not conjugate to x.

If G = SO(V) of type D_n , then there are both unipotent classes and nilpotent orbits to consider. In all of the cases the elements x being considered act with two Jordan blocks on V. Using Lemma 2.4 we see that every element contained in subsystem subgroups and subalgebras of type $D_m D_{n-m}$ $(1 \le m \le \frac{m}{2})$ will act with at least four Jordan blocks on V. Thus they are not conjugate to x. By Lemma 2.2, what remains is to show that x is not conjugate to an element of the maximal generalised subsystem subgroup M (or subalgebra \mathfrak{m}) of type B_{n-1} . When x is unipotent, this follows from [Kor18, Lemma 3.8], which shows that the only unipotent elements $x' \in M$ that act with two Jordan blocks on V are those with $V \downarrow k[x'] = V(2) + V(2n-2)$. Such elements x' are regular and not conjugate to x. Now let x be nilpotent. We consider the distinguished nilpotent elements x' in $\mathfrak{m} = \mathfrak{so}(W)$. By Lemma 2.4 (vi), we have $W \downarrow k[x'] = \sum_{j=1}^{s} W_{l_j}(n_j) \oplus D(m)$ for some integers l_j, n_j and m. If s = 0, then $W \downarrow k[x'] = D(n)$ and x' is a regular nilpotent element of Lie(G), and therefore is not conjugate to x. When $s \ge 1$, the elements x' are contained in generalised subsystem subalgebras of type $D_{n-m}B_{m-1}$, which are contained in maximal subsystem subalgebras of type $D_{n-m}D_m$. We have already seen that any nilpotent element contained in such a subalgebra will act with at least four Jordan blocks on V, and thus cannot be conjugate to x. This completes the proof of the lemma. \square

The remaining task for G of classical type is to prove that the representatives of the eminent elements do indeed act with the claimed decompositions on V. When x is regular, a representative is provided by Remark 2.5. The non-regular eminent classes occur in characteristic p = 2 for types C_n and D_n , the representatives for these classes are constructed in the following sections.

4.1. Nilpotent representatives in type C_n . Let G = Sp(V) with V a k-vector space of dimension 2n. We need to prove that the nilpotent elements

$$e_l = \left(\sum_{i=1}^{n-1} e_{\alpha_i}\right) + e_{2\alpha_l + \dots + 2\alpha_{n-1} + \alpha_n}$$

in Table 1 act as $W_l(n)$ on the natural module V. We do this in the next lemma, but first present our choice of Chevalley basis for $\mathfrak{sp}(V)$.

Recall the form described at the beginning of Section 2.3. For this choice of form, one checks that there is a Cartan subalgebra \mathfrak{h} of diagonal matrices in $\mathfrak{sp}(V) \subset \mathfrak{gl}(V)$ of the form $\operatorname{diag}(h_1, \ldots, h_n, -h_n, \ldots, -h_1)$. For $1 \leq i \leq n$, define maps $\varepsilon_i : \mathfrak{h} \to k$ by $\varepsilon_i(h) = h_i$.

For all i, j let $E_{i,j}$ be the linear endomorphism on V such that $E_{i,j}(e_j) = e_i$ and $E_{i,j}(e_k) = 0$ for $k \neq j$. Then one checks that the endomorphisms e_{λ} in Table 14 are elements of $\mathfrak{sp}(V)$, and are simultaneous eigenvectors for ad \mathfrak{h} .

From a dimension count, we deduce that concatenating the elements in Table 14 together with a basis of \mathfrak{h} gives a basis of \mathfrak{sp}_{2n} . Moreover, $\Phi = \{\pm(\varepsilon_i \pm \varepsilon_j) : 1 \leq i < j \leq n\} \cup \{\pm 2\varepsilon_i : 1 \leq i \leq n\}$, and $\Phi^+ = \{\varepsilon_i \pm \varepsilon_j : 1 \leq i < j \leq n\} \cup \{2\varepsilon_i : 1 \leq i \leq n\}$ is a system of positive roots. We let $\Delta = \{\alpha_1, \ldots, \alpha_n\}$ be our base of Φ corresponding to Φ^+ , where $\alpha_i = \varepsilon_i - \varepsilon_{i+1}$ for $1 \leq i < n$ and $\alpha_n = 2\varepsilon_n$. We give the expressions of roots $\alpha \in \Phi^+$ in terms of Δ in Table 15. Lastly, checking commutator relations reveals the basis in Table 14 is in fact a Chevalley basis, with positive structure constants for extraspecial pairs.

In particular, $e_{\alpha_i} = E_{i,i+1} - E_{n+i+1,n+i}$ for $1 \le i \le n-1$, $e_{\alpha_n} = E_{n,2n}$ and $e_{2\alpha_l + \dots + 2\alpha_{n-1} + \alpha_n} = E_{l,n+l}$ for l < n.

$e_{\varepsilon_i - \varepsilon_j} = E_{i,j} - E_{n+j,n+i}$	for all $i \neq j$,
$e_{\varepsilon_i + \varepsilon_j} = E_{j,n+i} + E_{i,n+j}$	for all $i \neq j$,
$e_{-(\varepsilon_i + \varepsilon_j)} = E_{n+j,i} + E_{n+i,j}$	for all $i \neq j$,
$e_{2\varepsilon_i} = E_{i,n+i}$	for all i ,
$e_{-2\varepsilon_i} = E_{n+i,i}$	for all i .

Table 14. Chevalley basis for \mathfrak{sp}_{2n}

Root		$\alpha = \sum_{k=1}^{n} c_k \alpha_i$
$\varepsilon_i - \varepsilon_j,$	$1 \leq i < j \leq n$	$\sum_{i \le k \le j-1} \alpha_k$
$\varepsilon_i + \varepsilon_j,$	$1 \leq i < j \leq n$	$\sum_{i \le k \le j-1} \alpha_k + \sum_{j \le k \le n-1} 2\alpha_k + \alpha_n$
$2\varepsilon_i,$	$1 \leq i \leq n$	$\sum_{i \le k \le n-1} 2\alpha_k + \alpha_n$

Table 15.	Type C_n ,	expressions	for	positive	roots	in	terms	of	base	Δ .
-----------	--------------	-------------	-----	----------	------------------------	----	------------------------	----	------	------------

Lemma 4.3. For 0 < l < n/2, the element $e_l = \sum_{i=1}^{n-1} e_{\alpha_i} + e_{2\alpha_l + \dots + 2\alpha_{n-1} + \alpha_n} \in \mathfrak{sp}_{2n}(k)$ acts on V as $V \downarrow k[e_l] = W_l(n)$.

Proof. For ease of notation we set $e = e_l$. From the description of the k[e]-module $W_l(n)$ in Section 2.3, it suffices to show that e acts on the natural 2*n*-dimensional module V with two Jordan blocks of size n and that $l = \min\{m : (e^{m+1}(v), e^m(v)) = 0 \text{ for all } v \in V\}$.

Using Tables 14 and 15, we write

$$e = E_{l,n+l} + \sum_{i=1}^{n-1} (E_{i,i+1} + E_{n+i+1,n+i}),$$

and therefore the action of e on V is described as follows. We have

$$e(v_{1}) = 0$$

$$e(v_{i}) = v_{i-1} \qquad (2 \le i \le n)$$

$$e(v_{n+l}) = v_{l} + v_{n+l+1}$$

$$e(v_{j}) = v_{j+1} \qquad (n+1 \le j < 2n, j \ne n+l)$$

$$e(v_{2n}) = 0.$$

We immediately see that $e^n(v_i) = 0$ for all $1 \le i \le 2n$. Furthermore, the relations above readily imply that the kernel of e is the 2-dimensional subspace of V generated by v_1 and v_{2n} . Thus the Jordan normal form of e has two Jordan blocks of size n.

To show that $l = \min\{m : (e^{m+1}(v), e^m(v)) = 0 \text{ for all } v \in V\}$ we start by observing

$$(e^{j+1}(v_{n+l-j}), e^j(v_{n+l-j})) = (v_l + v_{n+l+1}, v_{n+l}) = 1$$
 for all $j < l$.

It remains to show that $(e^{l+1}(v), e^{l}(v)) = 0$ for all $v \in V$. Since $(e^{l+1}(v_i), e^{l}(v_j)) + (e^{l+1}(v_j), e^{l}(v_i)) = 0$ for all i, j, by definition of $\mathfrak{sp}(V)$ preserving the alternating form, it suffices to show that $(e^{l+1}(v_i), e^{l}(v_i)) = 0$ for all basis vectors v_i .

If $1 \leq i \leq n$ or $n+l < i \leq 2n$ we immediately have that $(e^{l+1}(v_i), e^l(v_i)) = 0$ since *e* stabilises the totally isotropic subspaces $\langle v_1, \ldots, v_n \rangle$ and $\langle v_{n+l+1}, \ldots, v_{2n} \rangle$. Now suppose that $1 \leq j \leq l$. Then $e^{l}(v_{n+j}) = v_{l-j+1} + v_{n+l+j}$ and so $e^{l+1}(v_{n+j}) = v_{l-j} + v_{n+l+j+1}$ and so $(e^{l+1}(v_{n+j}), e^{l}(v_{n+j})) = 0$, as required.

4.2. Nilpotent and unipotent representatives in type D_n . Let G = SO(V) with V a k-vector space of dimension 2n. We need to establish the correctness of the representatives in Tables 3 and 5. Recall we define the root $\beta_l = \alpha_n + \sum_{i=2l-n-1}^{n-2} \alpha_i$ for $(n+1)/2 < l \leq n$.

We mimic the process from the last section. Using the form from the beginning of Section 2.3 we check we may choose a Cartan subalgebra \mathfrak{h} of the form $\operatorname{diag}(h_1, \ldots, h_n, -h_n, \ldots, -h_1)$ with $h_i \in k$. For $1 \leq i \leq n$, define maps $\varepsilon_i : \mathfrak{h} \to k$ by $\varepsilon_i(h) = h_i$. A Chevalley basis for $\mathfrak{so}(V)$ with positive structure constants for extraspecial pairs is given in Table 16.

$$e_{\varepsilon_i - \varepsilon_j} = E_{i,j} - E_{n+j,n+i} \qquad \text{for all } i \neq j,$$

$$e_{\varepsilon_i + \varepsilon_j} = E_{j,n+i} - E_{i,n+j} \qquad \text{for all } i < j,$$

$$e_{-(\varepsilon_i + \varepsilon_j)} = E_{n+i,j} - E_{n+j,i} \qquad \text{for all } i < j.$$

Table 16. Chevalley basis of $\mathfrak{so}(2n)$.

Now $\Phi = \{\pm(\varepsilon_i \pm \varepsilon_j) : 1 \le i < j \le n\}$ is the root system of $\mathfrak{so}(V)$, and $\Phi^+ = \{\varepsilon_i \pm \varepsilon_j : 1 \le i < j \le n\}$ is a system of positive roots. Here the base Δ of Φ corresponding to Φ^+ is $\Delta = \{\alpha_1, \ldots, \alpha_n\}$, where $\alpha_i = \varepsilon_i - \varepsilon_{i+1}$ for $1 \le i < n$ and $\alpha_n = \varepsilon_{n-1} + \varepsilon_n$. We give the expressions of roots $\alpha \in \Phi^+$ in terms of Δ in Table 17.

Root	$\alpha = \sum_{k=1}^{n} c_k \alpha_i$
$\varepsilon_i - \varepsilon_j, 1 \le i < j \le n$	$\sum_{i \le k \le j-1} \alpha_k$
$\varepsilon_i + \varepsilon_j, 1 \le i < j \le n-1$	$\sum_{i \le k \le j-1} \alpha_k + \sum_{j \le k \le n-2} 2\alpha_k + \alpha_{n-1} + \alpha_n$
$\varepsilon_i + \varepsilon_n, \ 1 \le i \le n-1$	$\sum_{i \le k \le n-2} \alpha_k + \alpha_n$

Table 17. Type D_n , expressions for positive roots in terms of base Δ .

Lemma 4.4. The element $u_l = \prod_{i=1}^{n-1} x_{\alpha_i}(1) \cdot x_{\beta_l}(1)$ acts on V as $V \downarrow k[u_l] = V(2l-2) \oplus V(2n-2l+2)$.

Proof. From Tables 16 and 17, we have $x_{\alpha_i}(1) = I + E_{i,i+1} + E_{n+i+1,n+i}$ for all $1 \le i \le n-1$, where I is the identity and $E_{i,j}$ is defined as before. Furthermore, we see that $x_{\beta_l}(1) = I + E_{2l-n-1,2n} + E_{n,2l-1}$. Now $u := u_l$ acts on the basis elements of V as follows:

$$u(v_i) = v_i + v_{i-1} + \dots + v_1 \quad (1 \le i \le n)$$

$$u(v_j) = v_j + v_{j+1} \quad (n+1 \le j < 2n, \ j \ne 2l-1)$$

$$u(v_{2l-1}) = v_{2l-1} + v_{2l} + v_n + v_{n-1} + \dots + v_1$$

$$u(v_{2n}) = v_{2n} + v_{2l-n-1} + v_{2l-n-2} + \dots + v_1.$$

A calculation shows that the fixed point space of u has dimension 2, and that it is spanned by v_1 and $v_{2l-n} + v_{2n}$. Therefore the Jordan normal form of u has two Jordan blocks. To see that the Jordan block sizes are 2l-2 and 2n-2l+2, it suffices to show that $(u-1)^{2l-2} = 0$ and $(u-1)^{2l-3} \neq 0$, as then the largest Jordan block size in u is 2l-2. To this end, a calculation shows that $(u-1)^{2l-2}(v_i) = 0$ for all i and

$$(u-1)^{2l-3}(v_{n+1}) = \begin{cases} v_1, & \text{if } 2l-2 > n, \\ v_{2n}+v_2+v_1, & \text{if } 2l-2 = n. \end{cases}$$

It is now clear that $((u-1)^{2l-3}(v_{n+1}), v_{n+1}) = 1$ and $(u-1)^{2l-2}(v_{n+1}) = 0$. Therefore V(2l-2) occurs as summand of $V \downarrow k[u]$ by [Kor20, Lemma 6.9], and we must have $V \downarrow k[u] = V(2l-2) \oplus V(2n-2l+2)$, as required.

Lemma 4.5. The element $e_l = e_{\beta_l} + \sum_{i=1}^{n-1} e_{\alpha_i}$ acts on V via $V \downarrow k[e_l] = W_l(n)$.

Proof. The tables above yield $e_{\alpha_i} = E_{i,i+1} + E_{n+i+1,n+i}$ for i < n and $e_{\beta_l} = E_{2l-n-1,n} + E_{n,2l-1}$, so that

$$e := e_l = E_{2l-n-1,2n} + E_{n,2l-1} + \sum_{i=1}^{n-1} (E_{i,i+1} + E_{n+i+1,n+i}).$$

We need to show that e has two Jordan blocks of size n and that $l = \min\{m \mid Q(e^m(v)) = 0 \forall v \in V\}$. We start by calculating the action of e on the basis vectors of V:

$$e(v_1) = 0$$

$$e(v_i) = v_{i-1} \ (2 \le i \le n)$$

$$e(v_j) = v_{j+1} \ (n+1 \le j < 2n, j \ne 2l-1)$$

$$e(v_{2l-1}) = v_{2l} + v_n$$

$$e(v_{2n}) = v_{2l-n-1}.$$

One sees that the kernel of e has dimension 2, and it is spanned by v_1 and $v_{2l-n} + v_{2n}$. Therefore the Jordan normal form of e has two Jordan blocks and a routine calculation with the basis $\{v_i\}$ shows that $e^n(v) = 0$ for all $v \in V$.

Finally, we must show that $l = \min\{m \mid Q(e^m(v)) = 0 \forall v \in V\}$. Since $e^{l-1}(v_{n+1}) = v_l + v_{n+l}$, it follows that $Q(e^{l-1}(v_{m+1})) = 1$ and so $Q(e^j(e^{l-1-j}(v_{m+1}))) = 1$ for all $j \leq l-1$. It remains to prove that $Q(e^l(v)) = 0$ for all $v \in V$. Since $2l \geq n$, it follows that $(e^l(v), e^l(w)) = (v, e^{2l}(w)) = 0$. Therefore, it suffices to show that $Q(e^l(v_i)) = 0$ for all basis vectors v_i . If $1 \leq i \leq n$ or $2l \leq i \leq 2n$ then $e^l(v_i) = v_j$ for some j and so $Q(e^l(v_i)) = 0$. When $n+1 \leq i \leq 2l-1$, we have $e^l(v_i) = v_{l+i} + v_{n-i+l}$ and so again, $Q(e^l(v_i)) = 0$.

Remark 4.6. Interpreting the coefficients of the unipotent element u_l of Lemma 4.4 as integers, reduction mod p for p > 2 gives an element acting on the natural module with Jordan block sizes (2l-1, 2n-2l+1). Indeed, a calculation shows that the fixed point space is spanned by v_1 and $v_{2l-n} + v_{2n}$, and furthermore

$$(u-1)^{2l-2}(v_j) = \begin{cases} (-1)^n 2v_1, & \text{if } j = n+1\\ 0, & \text{if } j \neq n+1 \end{cases}$$

so the largest Jordan block size of u_l is 2l - 1.

Therefore u_l is regular in the subsystem $B_m B_{n-m+1}$ of D_n and is therefore another representative alongside that provided in Lemma 2.6. Similarly one can calculate that the nilpotent element e_l of Lemma 4.5 has Jordan block sizes (2l - 1, 2n - 2l + 1) in characteristic $p \neq 2$. Since the roots involved in u_l and e_l are Z-independent, by [Hum75, Lemma 16.2C] these representatives have the advantage that they work regardless of the choice of signs of the structure constants.

4.3. Exceptional types. In this section we prove Theorem 1.1 for G of exceptional type. If x is unipotent, this follows from work of Lawther in [Law09]. It is routine to refine his lists to remove those distinguished unipotent classes of G which are not eminent. For each non-eminent unipotent class we list its maximal subsystem overgroups in Table 18, for completeness. We note that a representative of a unipotent class with label X is simply $\prod_{j \in J} x_{\alpha_j}(1)$, where $\sum_{j \in J} e_{\alpha_j}$ is the representative of the nilpotent element with label X given in [Ste16]. This follows from the proofs in [LS12] (see the introduction of Sections 17 and 18 in ibid.) since the representatives in [Ste16] are deduced from [LS12].

In good characteristic, [LT11, p.24] provides enough information to find the non-eminent distinguished nilpotent orbits. As might be expected, these classes are in bijection with the non-eminent distinguished unipotent classes (sending a unipotent class to the nilpotent orbit of the same label). It remains to consider the nilpotent orbits of Lie(G) in bad characteristic. We follow Lawther's approach, which entails constructing representatives of each distinguished nilpotent orbit of Lie(M) for M a maximal subsystem subgroup of G and determining its G-class. To do this we make heavy use of use Theorem 1.2.

Carrying out this calculation provides a proof of Theorem 1.1. Lastly, we mention Table 19. It provides a list of distinguished, non-eminent nilpotent orbits, giving their maximal subsystem overalgebras, for all characteristics. Specialising to good characteristic, we recover the analogous result in [Law09].

Type	Char.	Label	Maximal subsystem overgroups
G_2	any	$G_2(a_1)$	$A_2, A_1 \tilde{A}_1 \ (p \neq 2), \ \tilde{A}_2 \ (p = 3)$
F_4	any	$F_4(a_1)$	$B_4, C_4 \ (p=2)$
	$p \neq 2$	$F_4(a_2)$	A_1C_3
	any	$F_4(a_3)$	$B_4, A_2 \tilde{A}_2 \ (p \neq 3), A_1 C_3 \ (p \neq 2), C_4 \ (p = 2)$
	p = 2	$(C_3(a_1))_2$	B_4,C_4
E_6	$p \neq 2$	$E_{6}(a_{3})$	A_1A_5
E_7	$p \neq 2$	$E_7(a_3)$	A_1D_6
	any	$E_7(a_4)$	$A_1D_6, A_7 \ (p=2)$
	any	$E_7(a_5)$	$A_1D_6 \ (p \neq 2), \ A_2A_5 \ (p \neq 3)$
E_8	$p \neq 2$	$E_8(a_3)$	A_1E_7
	$p \neq 2$	$E_8(a_4)$	D_8
	$p \neq 2$	$E_{8}(a_{5})$	D_8
	any	$E_8(a_6)$	$A_8 \ (p \neq 3), \ D_8 \ (p \neq 2)$
	any	$E_{8}(a_{7})$	$A_4A_4 \ (p \neq 5), \ A_2E_6 \ (p \neq 3), \ A_1E_7 \ (p \neq 2), \ D_8 \ (p \neq 2)$
	any	$E_{8}(b_{4})$	$A_1 E_7, D_8 \ (p=2)$
	any	$E_8(b_5)$	$A_2E_6 \ (p \neq 3), \ A_1E_7 \ (p \neq 2)$
	any	$E_8(b_6)$	$A_2E_6, D_8 \ (p \neq 2), A_8 \ (p = 3)$
	p = 2	$(D_7(a_1))_2$	$A_1 E_7, D_8$

 Table 18. Maximal subsystem overgroups of non-eminent distinguished unipotent elements in exceptional types

Type	Char.	Label	Maximal subsystem overalgebras
G_2	any	$G_2(a_1)$	$A_2, A_1\tilde{A}_1 \ (p \neq 2)$
F_4	any	$F_4(a_1)$	B_4
	$p \neq 2$	$F_4(a_2)$	A_1C_3
	any	$F_4(a_3)$	$B_4, A_2 \tilde{A}_2 \ (p \neq 3), A_1 C_3 \ (p \neq 2)$
	p = 2	$(C_3(a_1))_2$	B_4
E_6	$p \neq 2$	$E_{6}(a_{3})$	A_1A_5
E_7	$p \neq 2$	$E_7(a_3)$	A_1D_6
	any	$E_{7}(a_{4})$	A_1D_6
	any	$E_7(a_5)$	$A_1D_6 \ (p \neq 2), \ A_2A_5 \ (p \neq 3)$
E_8	$p \neq 2$	$E_8(a_3)$	A_1E_7
	$p \neq 2$	$E_8(a_4)$	D_8
	$p \neq 2$	$E_{8}(a_{5})$	D_8
	any	$E_8(a_6)$	$A_8 \ (p \neq 3), \ D_8 \ (p \neq 2)$
	any	$E_8(a_7)$	$A_4A_4 \ (p \neq 5), \ A_2E_6 \ (p \neq 3), \ A_1E_7 \ (p \neq 2), \ D_8 \ (p \neq 2)$
	any	$E_8(b_4)$	A_1E_7
	any	$E_{8}(b_{5})$	$A_2E_6 \ (p \neq 3), \ A_1E_7 \ (p \neq 2)$
	any	$E_8(b_6)$	$A_2E_6, D_8 \ (p \neq 2)$

 Table 19. Maximal subsystem overalgebras of non-eminent distinguished nilpotent elements in exceptional types

5. Appendix: Auxiliary tables for Theorem 1.2

Below follows some auxiliary data for Theorem 1.2. Let G be expectional. In each table we use a horizontal line to separate the sets of nilpotent orbits (represented by their labels) for which the Jordan block structure on V_{\min} (when it exists) and Lie(G) coincide. For each class we then provide the additional data required to distinguish them. See the introduction for the terminology (DS, ALG, etc.).

Label	DS data
$F_4(a_1)$	10, 5, 2, 0
$F_4(a_2)$	10, 3, 0
B_3	14, 9, 6
$F_4(a_3)$	14, 7, 3, 0
$(C_3(a_1))_2$	16, 8, 4, 0
$(\tilde{A}_2 A_1)_2$	16, 8, 3, 0
$C_3(a_1)$	20, 16, 15, 7, 4, 0
$\tilde{A}_2 A_1$	20, 16, 15, 7, 1, 0

Table 20. $G = F_4, p = 2$

Label	DS data	
D_5	12, 8, 6, 2, 0	
$E_{6}(a_{3})$	12, 6, 0	
D_4	20, 17, 16	
$D_4(a_1)$	20, 15, 8, 0	
A_3A_1	24, 21, 20, 14, 12, 4, 0	
$A_2^2 A_1$	24, 21, 20, 12, 1, 0	

Table 21. $G = E_6, p = 2$

Label	ALG data
E_6	31 (sc) / 24 (ad)
$E_6(a_1)$	60 (sc) / 29 (ad)

Table 22. $G = E_6, p = 3$

Label	ALG data
E_7	62
$E_{7}(a_{1})$	67
D_6	58
$E_{7}(a_{4})$	63
$D_6(a_1)$	70
$(A_6)_2$	75
D_5A_1	73
$E_7(a_5)$	91
D_5	70
$E_{6}(a_{3})$	84
D_4A_1	63
$A_3A_2A_1$	69
$(A_3A_2)_2$	74
$D_4(a_1)A_1$	83
D_4	66
$D_4(a_1)$	82

Label	DS data
E_6	15, 12, 9
$E_6(a_1)$	15, 9, 4, 0
$D_6(a_1)$	19, 15
A_6	19, 17

Table 23. $G = E_7, p = 3$

Label	DS data
$D_6(a_2)$	26, 21, 20, 12, 2, 0
A_5A_1	26, 21, 20, 12, 1, 0
$(A_3A_1)^{(1)}$	43, 39, 37, 29, 21, 20, 4, 0
$A_{2}^{2}A_{1}$	43, 39, 37, 27, 8, 0

Table 24. $G = E_7$ (simply connected), p = 2

Label	ALG data
E_7	56
$E_{7}(a_{1})$	58
D_6	52
$E_{7}(a_{4})$	54
$D_6(a_1)$	56
$(A_6)_2$	57
D_5A_1	60
$E_{7}(a_{5})$	71
D_5	42
$E_{6}(a_{3})$	50
A_4A_2	54
$D_5(a_1)$	39
D_4A_1	57
$A_3A_2A_1$	58
$(A_3A_2)_2$	60
$D_4(a_1)A_1$	65
A_{2}^{2}	56
A_3	55

Label	DS data
$D_6(a_2)$	26, 21, 20, 12, 2, 0
A_5A_1	26, 21, 20, 12, 1, 0
A_3A_2	39, 35, 26, 23, 7, 0
D_4	39, 35, 28
$D_4(a_1)$	39, 35, 28, 26, 14, 0
$(A_3A_1)^{(1)}$	43, 40, 37, 29, 21, 20, 4, 0
$A_2^2 A_1$	43, 40, 37, 27, 8, 0

Table 25. $G = E_7$ (adjoint), p = 2

Label	NDS data
E_8	15, 18, 248
$E_8(a_1)$	16, 18, 248
E_6A_1	32, 33, 39
$E_8(b_6)$	33, 34, 39, 60, 165, 248
$E_6(a_1)A_1$	34, 40, 53, 132, 248
$(A_7)_3$	33, 35, 47, 105, 248
A_7	36, 36, 36
E_6	36, 37, 59
$E_{6}(a_{1})$	39, 56, 60
$E_7(a_4)$	40, 47, 63
A_6A_1	38, 38, 38
$D_6(a_1)$	43,49
A_6	41,41

Label	NIL data
$E_6(a_3)A_1$	36
$D_5(a_1)A_2$	43

Label	DS data
E_8	10, 7, 0
$E_8(a_1)$	10, 1, 0

Table 26. $G = E_8, p = 5$

Table 27.
$$G = E_8, p = 3$$

Label	ALG, ALG'	Label	ALG, ALG'	
	data		data	
E_8	79,74	D_5A_2	76, 67	
$E_8(a_1)$	79,73	$E_{8}(a_{7})$	102,99	
$E_8(a_2)$	89,85	$D_6(a_2)$	73,71	
$E_8(a_3)$	108, 103	A_5A_1	73, 58	
$E_8(a_4)$	139, 136	D_5	64,60	
E_7	68, 64	$E_{6}(a_{3})$	70, 64	
$E_{8}(b_{4})$	70, 63	$(D_4A_2)_2$	64,58	
$E_7(a_1)$	72, 69	$A_4 A_2 A_1$	64, 57	
$E_8(a_5)$	79,72	$D_5(a_1)A_1$	71, 64	
$E_{8}(b_{5})$	91,86	D_4A_2	83,73	
$(D_7)_2$	102,98	A_{3}^{2}	88,76	
$E_8(a_6)$	107, 101	$D_4(a_1)A_2$	99,72	
D_7	69, 65	D_4A_1	91,88	
$D_7(a_1)$	77,70	$A_3A_2A_1$	92,86	
A_7	85,72	$(A_3A_2)_2$	101,95	
$D_7(a_2)$	101, 93	$D_4(a_1)A_1$	109, 104	
$E_7(a_2)$	61, 57	D_4	88,85	
E_6A_1	61, 53	$D_4(a_1)$	92,90	
$(D_7(a_1))_2$	69, 63	A_3A_1	94,92	
$E_7(a_3)$	82,78	$2A_2A_1$	94,87	
D_6	71,68			
$(D_5A_2)_2$	72,65			
$E_7(a_4)$	74,67			
A_6A_1	77,66			
$D_6(a_1)$	82,78			
$(A_6)_2$	85,76			

Label	DS data
D_5A_1	44, 39, 37, 29, 19, 18, 2, 0
$E_{7}(a_{5})$	44, 39, 37, 29, 17, 1, 0
$E_6(a_3)A_1$	44, 39, 37, 27, 8, 0
$D_5(a_1)A_2$	48, 43, 37, 21, 1, 0
A_4A_3	48, 43, 37, 19, 1, 0
$A_{3}A_{1}^{2}$	80, 76, 75, 71, 70, 54, 39, 38, 6, 0
$A_{2}^{2}A_{1}^{2}$	80, 76, 75, 71, 70, 50, 14, 0

Table 28. $G = E_8, p = 2$

References

- [BCP97] W. Bosma, J. Cannon, and C. Playoust, The Magma algebra system. I. The user language, vol. 24, 1997, Computational algebra and number theory (London, 1993), pp. 235–265. MR 1484478
- [Bou82] N. Bourbaki, Éléments de mathématique: groupes et algèbres de Lie, Masson, Paris, 1982, Chapitre 9. Groupes de Lie réels compacts. [Chapter 9. Compact real Lie groups]. MR 682756 (84i:22001)
- [Car89] R. W. Carter, Simple groups of Lie type, Wiley Classics Library, John Wiley & Sons Inc., New York, 1989, Reprint of the 1972 original, A Wiley-Interscience Publication. MR MR1013112 (90g:20001)
- [Car93] _____, Finite groups of Lie type, Wiley Classics Library, John Wiley & Sons Ltd., Chichester, 1993, Conjugacy classes and complex characters, Reprint of the 1985 original, A Wiley-Interscience Publication. MR MR1266626 (94k:20020)
- [gro14] The GAP group, Gap groups, algorithms, and programming, Version 4.7.5 (2014), http://www.gap-system.org.
- [Hes79] W. H. Hesselink, Nilpotency in classical groups over a field of characteristic 2, Math. Z. 166 (1979), no. 2, 165–181. MR 525621
- [Hog82] G. M. D. Hogeweij, Almost-classical Lie algebras. I, II, Nederl. Akad. Wetensch. Indag. Math. 44 (1982), no. 4, 441–452, 453–460. MR 683531 (84f:17007)
- [Hum75] J. E. Humphreys, *Linear algebraic groups*, Springer-Verlag, New York-Heidelberg, 1975, Graduate Texts in Mathematics, No. 21. MR 0396773
- [Jan73] J. C. Jantzen, Darstellungen halbeinfacher algebraischer Gruppen und zugeordnete kontravariante Formen, Bonn. Math. Schr. (1973), no. 67, v+124. MR 0401935
- [Jan03] _____, *Representations of algebraic groups*, second ed., Mathematical Surveys and Monographs, vol. 107, American Mathematical Society, Providence, RI, 2003. MR MR2015057 (2004h:20061)
- [Jan04] _____, Nilpotent orbits in representation theory, Lie theory, Progr. Math., vol. 228, Birkhäuser Boston, Boston, MA, 2004, pp. 1–211. MR 2042689 (2005c:14055)
- [Ken87] S. V. Keny, Existence of regular nilpotent elements in the Lie algebra of a simple algebraic group in bad characteristics, J. Algebra 108 (1987), no. 1, 195–201. MR 887203
- [Kor18] M. Korhonen, Unipotent elements forcing irreducibility in linear algebraic groups, J. Group Theory 21 (2018), no. 3, 365–396. MR 3794921
- [Kor20] _____, Hesselink normal forms of unipotent elements in some representations of classical groups in characteristic two, J. Algebra **559** (2020), 268–319. MR 4097906
- [Law09] R. Lawther, Unipotent classes in maximal subgroups of exceptional algebraic groups, J. Algebra 322 (2009), no. 1, 270–293. MR 2526390
- [LT11] R. Lawther and D. M. Testerman, Centres of centralizers of unipotent elements in simple algebraic groups, Mem. Amer. Math. Soc. 210 (2011), no. 988, vi+188. MR 2780340 (2012c:20127)
- [LS12] M. W. Liebeck and G. M. Seitz, Unipotent and nilpotent classes in simple algebraic groups and Lie algebras, Mathematical Surveys and Monographs, vol. 180, American Mathematical Society, Providence, RI, 2012. MR 2883501
- [MT11] G. Malle and D. M. Testerman, Linear algebraic groups and finite groups of Lie type, Cambridge Studies in Advanced Mathematics, vol. 133, Cambridge University Press, Cambridge, 2011. MR 2850737 (2012i:20058)
- [Spr66] T. A. Springer, Some arithmetical results on semi-simple Lie algebras, Inst. Hautes Études Sci. Publ. Math. (1966), no. 30, 115–141. MR 206171
- [Ste65] R. Steinberg, Regular elements of semisimple algebraic groups, Inst. Hautes Études Sci. Publ. Math. (1965), no. 25, 49–80. MR 180554
- [Ste16] D. I. Stewart, On the minimal modules for exceptional Lie algebras: Jordan blocks and stabilizers, LMS J. Comput. Math. 19 (2016), no. 1, 235–258. MR 3530500
- [Tes95] D. M. Testerman, A_1 -type overgroups of elements of order p in semisimple algebraic groups and the associated finite groups, J. Algebra **177** (1995), no. 1, 34–76. MR 1356359 (96j:20067)

Department of Mathematics, Southern University of Science and Technology, Shenzhen 518055, Guangdong, P. R. China

Email address: korhonen_mikko@hotmail.com (Korhonen)

School of Mathematics and Statistics, Herschel Building, Newcastle, NE1 7RU, UK

Email address: david.stewart@ncl.ac.uk (Stewart)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WARWICK, COVENTRY, CV4 7AL, UK

Email address: adam.r.thomas@warwick.ac.uk (Thomas)