

The coarse Novikov conjecture for extensions of coarsely embeddable groups *

Qin Wang and Yazhou Zhang

Abstract

Let $(1 \rightarrow N_n \rightarrow G_n \rightarrow Q_n \rightarrow 1)_{n \in \mathbb{N}}$ be a sequence of extensions of countable discrete groups. Endow $(G_n)_{n \in \mathbb{N}}$ with metrics associated to proper length functions on $(G_n)_{n \in \mathbb{N}}$ respectively such that the sequence of metric spaces $(G_n)_{n \in \mathbb{N}}$ have uniform bounded geometry. We show that if $(N_n)_{n \in \mathbb{N}}$ and $(Q_n)_{n \in \mathbb{N}}$ are coarsely embeddable into Hilbert space, then the coarse Novikov conjecture holds for the sequence $(G_n)_{n \in \mathbb{N}}$, which may not admit a coarse embedding into Hilbert space.

1 Introduction

M. Gromov introduced the notion of coarse embeddings of metric spaces into Hilbert space, and suggested that finitely generated discrete groups that are coarsely embeddable into Hilbert space, when viewed as metric spaces, might satisfy the Novikov conjecture [13]. G. Yu proved that this is indeed the case [33]. Recall that a map $f : X \rightarrow H$ from a metric space X to a Hilbert space H is a *coarse embedding* if there exist two non-decreasing functions $\rho_1, \rho_2 : [0, \infty) \rightarrow [0, \infty)$ such that $\lim_{t \rightarrow \infty} \rho_i(t) = +\infty$ for $i = 1, 2$ and such that $\rho_1(d(x, y)) \leq \|f(x) - f(y)\| \leq \rho_2(d(x, y))$ for all $x, y \in X$.

Let $(1 \rightarrow N_n \rightarrow G_n \rightarrow Q_n \rightarrow 1)_{n \in \mathbb{N}}$ be a sequence of extensions of countable discrete groups. For each $n \in \mathbb{N}$, let l_{G_n} be a proper length function on G_n . We can define a family of left invariant metrics d_{G_n} on G_n by $d_{G_n}(g, h) = l_{G_n}(g^{-1}h)$. Let $\pi : G_n \rightarrow Q_n$ be the quotient map. Define length functions on N_n and Q_n according to $l_{N_n}(g) = l_{G_n}(g)$ for all $g \in N_n$, and $l_{Q_n}(x) = \min\{l_{G_n}(g) : g \in G_n \text{ and } \pi(g) = x\}$ for all $x \in Q_n$. Then these length functions induce left invariant metrics d_{N_n} and d_{Q_n} on N_n and Q_n , respectively. It follows that we have three sequences discrete metric spaces $(N_n, d_{N_n})_{n \in \mathbb{N}}$, $(G_n, d_{G_n})_{n \in \mathbb{N}}$ and $(Q_n, d_{Q_n})_{n \in \mathbb{N}}$. The sequence of metric spaces $(G_n, d_{G_n})_{n \in \mathbb{N}}$ is said to have uniform bounded geometry if for every $r > 0$, the number of the elements in $B_{G_n}(x, r) = \{y \in G_n : d_{G_n}(x, y) \leq r\}$ is at most N_r for some $N_r > 0$ independent of n . The sequence of metric spaces $(N_n, d_{N_n})_{n \in \mathbb{N}}$ is said to be coarsely embeddable into Hilbert space if there exists a family of maps $f_n : N_n \rightarrow H_n$, and two non-decreasing functions $\rho_1, \rho_2 : [0, \infty) \rightarrow [0, \infty)$ such that $\lim_{t \rightarrow \infty} \rho_i(t) = +\infty$ for $i = 1, 2$ and $\rho_1(d_{N_n}(x, y)) \leq \|f_n(x) - f_n(y)\| \leq \rho_2(d_{N_n}(x, y))$ for all $x, y \in N_n$.

*Supported in part by NSFC (No. 11831006, 11771143).

The main purpose of this paper is to prove the following result:

Theorem 1.1. *Let $(1 \rightarrow N_n \rightarrow G_n \rightarrow Q_n \rightarrow 1)_{n \in \mathbb{N}}$ be a sequence of extensions of countable discrete groups. Endow $(G_n)_{n \in \mathbb{N}}$ with metrics associated to proper length functions on $(G_n)_{n \in \mathbb{N}}$ respectively such that the sequence of metric spaces $(G_n)_{n \in \mathbb{N}}$ have uniform bounded geometry. If $(N_n)_{n \in \mathbb{N}}$ and $(Q_n)_{n \in \mathbb{N}}$ are coarsely embeddable into Hilbert space, then the coarse Novikov conjecture holds for the sequence $(G_n)_{n \in \mathbb{N}}$.*

The coarse Novikov conjecture is the injectivity part of the coarse Baum-Connes conjecture [16, 25, 29], and a geometric analogue of the strong Novikov conjecture [4]. In the case of a noncompact complete Riemannian manifold, the coarse Novikov conjecture provides an algorithm to determine when the higher index of an elliptic operator on the noncompact complete Riemannian manifold is nonzero. In particular, it implies that the higher index of the Dirac operator on a uniformly contractible Riemannian manifold is nonzero. By Proposition 4.33 in [25], the coarse Novikov conjecture implies Gromov's conjecture that a uniformly contractible Riemannian manifold with bounded geometry cannot have uniformly positive scalar curvature, and the zero-in-the-spectrum conjecture stating that the Laplace operator acting on the space of all L^2 -forms of a uniformly contractible Riemannian manifold has zero in its spectrum. See [34] for a comprehensive survey on the coarse Novikov conjecture.

A remarkable progress was achieved by G. Yu [33] who proved the coarse Baum-Connes conjecture, and consequently, the coarse Novikov conjecture for metric space with bounded geometry which admits a coarse embedding into Hilbert space. A fundamental idea underlining the approach in [33] is that the index of a Dirac operator is more computable if the Dirac operator is twisted by a family of "almost flat Bott bundles". This approach inspires several later progresses on the coarse Novikov conjecture for coarse embeddings into certain Banach spaces [19, 6] or non-positively curved manifolds [12, 27]. See also [9, 11, 20, 24, 32] for closely related developments.

In a recent paper, G. Arzhantseva and R. Tessera [3] answer in the negative a long standing open question raised in [7, 15]: Does coarse embeddability into Hilbert space is preserved under group extensions of finitely generated groups? Their counterexamples are certain restricted permutational wreath products $\mathbb{Z}_2 \wr_G H$ and $\mathbb{Z}_2 \wr_G (H \times \mathbb{F}_n)$, where G is a *Gromov monster group* [14, 1, 22], i.e., a finitely generated group which contains in its Cayley graph an isometrically embedded expander [14], and H is a *Haagerup monster group* [1, 22], i.e., a finitely generated group with the Haagerup property but without Yu's property A [33, 21]. The reason why the groups $\mathbb{Z}_2 \wr_G H$ and $\mathbb{Z}_2 \wr_G (H \times \mathbb{F}_n)$ do not coarsely embed into Hilbert space is that both groups contain isometrically in their Cayley graphs a *relative expander*, an innovative notion introduced by G. Arzhantseva and R. Tessera in [2]. Consequently, the group $\mathbb{Z}_2 \wr_G H$ or $\mathbb{Z}_2 \wr_G (H \times \mathbb{F}_n)$ also provide the first example of an extension of countable discrete groups or an extension of finitely generated groups which does not coarsely embed into any L^p -space for any $1 \leq p < \infty$, nor into any uniformly curved Banach space, and yet does not contain any weakly embedded expander [2].

Very recently, J. Deng, Q. Wang and G. Yu [10] show that if a sequence of group extensions $(1 \rightarrow N_n \rightarrow G_n \rightarrow Q_n \rightarrow 1)_{n \in \mathbb{N}}$ has "*A-by-CE*" structure, namely, the coarse disjoint union [21] of the sequence $(N_n)_{n \in \mathbb{N}}$ with the induced metric from the word metrics of $(G_n)_{n \in \mathbb{N}}$ has Yu's property A, and the coarse disjoint union of the sequence $(Q_n)_{n \in \mathbb{N}}$ with the quotient metrics coarsely embeds into Hilbert space

(denoted briefly, CE), then the coarse Baum-Connes conjecture holds for the coarse disjoint union of the sequence $(G_n)_{n \in \mathbb{N}}$. It follows that the coarse Baum-Connes conjecture, and hence the coarse Novikov conjecture, holds for the relative expanders in [2], the group extensions in [3] mentioned above (see [5, 23] for an alternative proof for these group extensions), and certain box spaces of free groups in [8], which do not coarsely embed into Hilbert space and yet do not coarsely contain any weakly embedded expander. In [10], J. Deng, Q. Wang and G. Yu raise an open problem: Does "CE-by-CE" imply the coarse Baum-Connes conjecture?

Our main result Theorem 1.1 provides an answer to the injectivity part of this problem. Namely, "CE-by-CE" implies the coarse Novikov conjecture. Moreover, we have the following result associated to [10, Theorem 3.13], which is related to relative expanders.

Theorem 1.2. *Let $(1 \rightarrow N_n \rightarrow G_n \rightarrow Q_n \rightarrow 1)_{n \in \mathbb{N}}$ be a sequence of extensions of finite groups with uniformly finite generating subsets. Endow $(N_n)_{n \in \mathbb{N}}$ with the induced metrics from the word metrics of $(G_n)_{n \in \mathbb{N}}$ and $(Q_n)_{n \in \mathbb{N}}$ with the quotient metrics from $(G_n)_{n \in \mathbb{N}}$, respectively. If the coarse disjoint unions $(N_n)_{n \in \mathbb{N}}$ and $(Q_n)_{n \in \mathbb{N}}$ are coarsely embeddable into Hilbert space, then the coarse Novikov conjecture holds for the coarse disjoint union $(G_n)_{n \in \mathbb{N}}$.*

As a special case of Theorem 1.1, we have the following result for a group extension.

Theorem 1.3. *Let $1 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 1$ be a short exact sequence of countable discrete groups. If N and Q are coarsely embeddable into Hilbert space, then the coarse Novikov conjecture holds for G .*

Note that J. Deng [9] has proved that the strong Novikov conjecture with arbitrary coefficient G - C^* -algebras holds for CE-by-CE groups. His result basically covers this case because G. Yu [30] proved that the Baum-Connes conjecture for a discrete group G with coefficients in $\ell^\infty(G, K(H))$ is equivalent to the coarse Baum-Connes conjecture for G as a metric space with a length metric. Here our paper provides a different method for this special case.

The paper is organized as follows. In Section 2, we briefly recall the concept of the Roe algebra, the coarse Novikov conjecture and Yu's localization algebra techniques. In Section 3, we introduce the twisted Roe algebra and the twisted localization algebra for the sequence of extensions of countable discrete groups associated to the coarse embeddings of the quotient groups $(G_m/N_m)_{m \in \mathbb{N}}$ into a Hilbert spaces. In section 4, we show that the evaluation map from the K -theory of the twisted localization algebra to the K -theory of the twisted Roe algebra is an isomorphism. In Section 5, we give the constructions of the Bott maps, and show that the Bott map from the K -theory of the uniform localization algebra to the K -theory of the twisted localization algebra is an isomorphism. Consequently, we prove the main result by recalling the result in the previous section.

We should mention that the surjectivity part of the open question raised in [10] still remains open:

Problem. Does "CE-by-CE" imply the *surjectivity* of the coarse Baum-Connes assembly map for group extensions?

2 The coarse Novikov conjecture for a sequence of metric spaces

In this section, we shall briefly recall the Roe algebra, Yu's localization algebra and the coarse Novikov conjecture.

Let X be a proper metric space. Recall that a metric space is *proper* if every closed bounded subset is compact. An X -*module* is a separable Hilbert space equipped with a faithful and non-degenerate $*$ -representation π of $C_0(X)$, the algebra of all complex-valued continuous functions on X vanishing at infinity. An X -module is said to be *standard* if no nonzero functions in $C_0(X)$ acts as a compact operator. When H_X is an X -module, for each $f \in C_0(X)$ and $h \in H_X$, we simply denote $(\pi(f))h$ by fh .

Definition 2.1. [25, 26] Let X be a proper metric space, and let H_X be an X -module.

- (1) The *support* $\text{Supp}(T)$ of a bounded linear operator $T : H_X \rightarrow H_X$ is the complement in $X \times X$ of the set of points $(x, y) \in X \times X$ for which there exist functions $f \in C_0(X), g \in C_0(X)$ with $f(x) \neq 0, g(y) \neq 0$ such that $gTf = 0$.
- (2) The *propagation* of a bounded linear operator $T : H_X \rightarrow H_X$ is defined to be

$$\sup\{d(x, y) : (x, y) \in \text{Supp}(T)\}.$$

We say that T has *finite propagation* if $\text{propagation}(T) < \infty$.

- (3) A bounded linear operator $T : H_X \rightarrow H_X$ is said to be *locally compact* if the operators fT and Tf are compact for all $f \in C_0(X)$.

Denote by $\mathbb{C}[X, H_X]$, or simply $\mathbb{C}[X]$, the set of all locally compact, finite propagation operators on H_X . It is straightforward to check that $\mathbb{C}[X]$ is a $*$ -algebra which, up to non-canonical isomorphisms, does not depend on the choice of standard non-degenerate X -modules [26, 31].

Definition 2.2. [25, 26] The *Roe algebra* $C^*(X, H_X)$, or simply denoted by $C^*(X)$, is defined to be the operator norm closure of $\mathbb{C}[X, H_X]$ on a standard non-degenerate X -module H_X .

We may use a specific standard non-degenerate X -module. Let Z be a countable dense subset of X , and let H be a fixed separable and infinite dimensional Hilbert space. Let $K := K(H)$ denote the algebra of compact operators on H . The algebra $C_0(X)$ acts on $\ell^2(Z)$ by point-wise multiplications. Thus we can choose H_X to be $\ell^2(Z) \otimes H$, where $C_0(X)$ acts on H_X by $f(\eta \otimes h) = f\eta \otimes h$ for $f \in C_0(X), \eta \in \ell^2(Z), h \in H$.

Definition 2.3. Define $\mathbb{C}_f[X]$ to be the $*$ -algebra of all bounded functions $T : Z \times Z \rightarrow K := K(H)$, viewed as $Z \times Z$ -matrices, such that:

- (1) for any bounded subset $B \subset X$, the set $\{(x, y) \in (B \times B) \cap (Z \times Z) | T(x, y) \neq 0\}$ is finite;
- (2) there exists $L > 0$ such that $\#\{y \in Z | T(x, y) \neq 0\} < L$ and $\#\{y \in Z | T(y, x) \neq 0\} < L$ for all $x \in Z$;
- (3) there exists $R \geq 0$ such that $T(x, y) = 0$ whenever $d(x, y) > R$ for all $x, y \in Z$.

Note that $\mathbb{C}_f[X]$ is $*$ -isomorphic to $\mathbb{C}[X, \ell^2(Z, H)]$ by a non-canonical $*$ -isomorphism [31, 33]. Hence, we can use $\mathbb{C}_f[X]$ to replace $\mathbb{C}[X]$ to define the Roe algebra $C^*(X)$ of X .

Definition 2.4. Let X be a discrete metric space with *bounded geometry*, i.e., for any $R > 0$ there exists $N > 0$ such that the number of elements in every ball of radius R is less than N . For each $d \geq 0$, the *Rips complex* $P_d(X)$ at scale d is defined to be the simplicial polyhedron where the set of vertices is X , and a finite subset $\{x_0, x_1, \dots, x_q\}$ in X spans a simplex if and only if $d(x_i, x_j) \leq d$ for all $0 \leq i, j \leq q$.

Endow $P_d(X)$ with the *spherical metric*. A detailed description for the spherical metric on a Rips complex can be found in [28, Definition 7.2.8]. In particular, for each $d' \geq d \geq 0$, the canonical inclusion $i_{d'd} : P_d(X) \rightarrow P_{d'}(X)$ is a coarse equivalence, and a homeomorphism onto its image [28, Proposition 7.2.11].

The coarse Novikov conjecture. For any discrete metric space X with bounded geometry, the coarse Baum-Connes assembly map

$$\mu : \lim_{d \rightarrow \infty} K_*(P_d(X)) \rightarrow \lim_{d \rightarrow \infty} K_*(C^*(P_d(X))) \cong K_*(C^*(X))$$

is injective, where $K_*(P_d(X)) = KK_*(C_0(P_d(X)), \mathbb{C})$ is the K -homology group [18] of the proper metric space $P_d(X)$ for each $d \geq 0$.

Yu's localization algebra [31] will play an important role in the proof of the main result of this paper.

Definition 2.5. [31, 28] Let X be a proper metric space. Define $\mathbb{C}_L[X]$ to be the $*$ -algebra of all bounded and uniformly continuous functions

$$g : [0, \infty) \rightarrow \mathbb{C}_f[X] \subset C^*(X)$$

such that there exists a bounded function $R : [0, \infty) \rightarrow [0, \infty)$ with $\lim_{t \rightarrow \infty} R(t) = 0$ and such that $(g(t))(x, y) = 0$ whenever $d(x, y) > R(t)$ for all $t \in [0, \infty)$, $x, y \in Z$.

The localization algebra $C_L^*(X)$ is defined to be the norm completion of $\mathbb{C}_L[X]$ with respect to the norm $\|g\|_\infty := \sup_{t \in [0, \infty)} \|g(t)\|$.

The K -theory of the localization algebras is invariant under strong Lipschitz homotopy invariance [31].

Definition 2.6. [31] Let X and Y be two proper metric spaces, and let f and g be two Lipschitz maps from X to Y . A continuous homotopy $F : [0, 1] \times X \rightarrow Y$ between f and g is said to be *strongly Lipschitz* [31] if

- (1) there exists $C > 0$ such that $d(F(t, x), F(t, y)) \leq Cd(x, y)$ for all $x, y \in X$ and $t \in [0, 1]$;
- (2) for any $\varepsilon > 0$, there exists $\delta > 0$ such that $d(F(t_1, x), F(t_2, x)) < \varepsilon$ for all $x \in X$ if $|t_1 - t_2| < \delta$;
- (3) $F(0, x) = f(x)$ and $F(1, x) = g(x)$ for all $x \in X$.

A metric space X is said to be *strongly Lipschitz homotopy equivalent* to Y , if there exist two Lipschitz maps $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that $f \circ g$ and $g \circ f$ are strongly Lipschitz homotopy equivalent to the identity maps id_X and id_Y , respectively.

Proposition 2.7. [31] *If X is strongly Lipschitz homotopy equivalent to Y , then $K_*(C_L^*(X))$ is naturally isomorphic to $K_*(C_L^*(Y))$.*

The following Mayer-Vietoris sequence is proved in [31].

Proposition 2.8. [31] *Let X be a simplicial complex endowed with the spherical metric, and let X_1 and X_2 be its two simplicial subcomplexes endowed with the subspace metric. Then we have the following six term exact sequence*

$$\begin{array}{ccccc} K_0(C_L^*(X_1 \cap X_2)) & \longrightarrow & K_0(C_L^*(X_1)) \oplus K_0(C_L^*(X_2)) & \longrightarrow & K_0(C_L^*(X_1 \cup X_2)) \\ & \uparrow & & & \downarrow \\ K_1(C_L^*(X_1 \cup X_2)) & \longleftarrow & K_1(C_L^*(X_1)) \oplus K_1(C_L^*(X_2)) & \longleftarrow & K_1(C_L^*(X_1 \cap X_2)) \end{array}$$

One can define a local assembly map $\mu_L : K_*(X) \rightarrow K_*(C_L^*(X))$ [31].

Proposition 2.9. [31] *For a finite dimensional simplicial polyhedron endowed with the spherical metric, the local assembly map μ_L is an isomorphism.*

The *evaluation-at-zero map* $e : C_L^*(X) \rightarrow C^*(X)$ is defined by $e(g) = g(0)$. It induces a homomorphism at the K -theory level $e_* : K_*(C_L^*(X)) \rightarrow K_*(C^*(X))$. For a discrete metric space X with bounded geometry, we have the following commutative diagram:

$$\begin{array}{ccc} & \lim_{d \rightarrow \infty} K_*(P_d(X)) & \\ \mu_L \swarrow & & \searrow \mu \\ \lim_{d \rightarrow \infty} K_*(C_L^*(P_d(X))) & \xrightarrow{e_*} & \lim_{d \rightarrow \infty} K_*(C^*(P_d(X))) \end{array}$$

\cong is placed between the two arrows from the top node to the bottom nodes.

Proposition 2.10. *Let X be a discrete metric space with bounded geometry. To prove the coarse Novikov conjecture for X , it suffices to show that e_* is injective.*

In the rest of this section, we introduce the coarse Baum-Connes conjecture for a sequence of metric spaces.

Let $(X_n)_{n \in \mathbb{N}}$ be a sequence of metric spaces with uniform bounded geometry. For each $d > 0, n \in \mathbb{N}$, we choose a countable dense subset $X_d^n \subset P_d(X_n)$ such that $X_d^n \subset X_{d'}^n$ if $d < d'$.

Definition 2.11. For each $d > 0$, the algebraic uniform Roe algebra $\mathbb{C}_u[(P_d(X_n))_{n \in \mathbb{N}}]$ is the collection of tuples $T = (T^{(n)})_{n \in \mathbb{N}}$ where each $T^{(n)} : X_d^n \times X_d^n \rightarrow K$ is a bounded function satisfying the following conditions:

- (1) there exists $M > 0$ such that $\|T^{(n)}(x, y)\| \leq M$ for each $n \in \mathbb{N}$ and $x, y \in Z_d^n$;
- (2) there exists $r > 0$ such that for each $n \in \mathbb{N}$, $T^{(n)}(x, y) = 0$ for all $x, y \in X_d^n$ satisfying $d(x, y) \geq r$;
- (3) there exists $L > 0$ such that for each $n \in \mathbb{N}$ and $x \in X_d^n$,

$$\#\{y \in X_d^n : T^{(n)}(x, y) \neq 0\} \leq L, \text{ and } \#\{y \in X_d^n : T^{(n)}(y, x) \neq 0\} \leq L;$$

(4) for any sequence of uniformly bounded subsets $(B_n)_{n \in \mathbb{N}}$ where $B_n \subset P_d(X_n)$, the set

$$\left\{ (x, y) \in (B_n \times B_n) \cap (X_d^n \times X_d^n) : T^{(n)}(x, y) \neq 0 \right\}$$

is uniformly finite.

The algebraic uniform Roe algebra $\mathbb{C}_u[(P_d(X_n))_{n \in \mathbb{N}}]$ is a $*$ -algebra and admits a faithful representation on $\bigoplus \ell^2(X_d^n) \otimes H$ via multiplication. Denote $E = \bigoplus \ell^2(X_d^n) \otimes H$. The uniform Roe algebra for the sequence $(P_d(X_n))_n$, denoted by $C_u^*((P_d(X_n))_{n \in \mathbb{N}})$, the completion of $\mathbb{C}_u[(P_d(X_n))_{n \in \mathbb{N}}]$ under the operator norm on $\mathcal{B}(E)$.

Definition 2.12. The algebraic uniform localization algebra, $\mathbb{C}_{u,L}[(P_d(X_n))_{n \in \mathbb{N}}]$ is the $*$ -algebra of all bounded and uniformly continuous functions

$$g : [0, \infty) \rightarrow \mathbb{C}_u[(P_d(X_n))_{n \in \mathbb{N}}]$$

such that $g(t)$ is of the form $g(t) = (g^{(n)}(t))_{n \in \mathbb{N}}$, where the tuples $(g^{(n)}(t))_{n \in \mathbb{N}}$ satisfies the condition of Definition 2.11 with uniform constants for all $t \in [0, \infty)$ and there is a bounded function $R : [0, \infty) \rightarrow [0, \infty)$ with $\lim_{t \rightarrow \infty} R(t) = 0$, such that for all $n \in \mathbb{N}, x, y \in X_d^n$ and $t \in [1, \infty)$,

$$(g^{(n)}(t))_{n \in \mathbb{N}}(x, y) = 0 \text{ if } d(x, y) > R(t).$$

The uniform localization algebra $C_{u,L}^*((P_d(X_n))_{n \in \mathbb{N}})$ is defined to be the completion of $\mathbb{C}_{u,L}[(P_d(X_n))_{n \in \mathbb{N}}]$ with respect to the norm $\|g\| = \sup_{t \in [0, \infty)} \|g(t)\|$.

Naturally, we have the evaluation-at-zero map $e : C_{u,L}^*((P_d(X_n))_{n \in \mathbb{N}}) \rightarrow C_u^*((P_d(X_n))_{n \in \mathbb{N}})$ defined by $e(g) = g(0)$. It induces a homomorphism at the K-theory level $e_* : K_*(C_{u,L}^*((P_d(X_n))_{n \in \mathbb{N}})) \rightarrow K_*(C_u^*((P_d(X_n))_{n \in \mathbb{N}}))$.

The coarse Novikov conjecture for a sequence of metric spaces. For any sequence of discrete metric spaces $(X_n, d_n)_{n \in \mathbb{N}}$ with uniform bounded geometry, the map

$$e_* : K_*(C_{u,L}^*((P_d(X_n))_{n \in \mathbb{N}})) \rightarrow K_*(C_u^*((P_d(X_n))_{n \in \mathbb{N}}))$$

is injective.

3 Twisted algebras with coefficients from the sequence of quotient groups $(G_m/N_m)_{m \in \mathbb{N}}$

Let $(1 \rightarrow N_m \rightarrow G_m \rightarrow G_m/N_m \rightarrow 1)_{m \in \mathbb{N}}$ be a sequence of extensions of countable discrete groups such that $(N_m)_{m \in \mathbb{N}}$ and $(G_m/N_m)_{m \in \mathbb{N}}$ are coarsely embeddable into Hilbert spaces. In this section, we will define the twisted Roe algebras and the twisted localization algebras for the sequence of Rips complexes $(P_d(G_m))_{m \in \mathbb{N}}$, in which the coefficients come from the coarse embedding of the sequence of quotient groups $(G_m/N_m)_{m \in \mathbb{N}}$ into Hilbert spaces. The construction of these twisted algebras has its origin in [33].

To get started, we first recall the C^* -algebra associated with an infinite-dimensional Euclidean space introduced by Higson, Kasparov and Trout [17]. Let V be a countably infinite dimensional Euclidean space. Denote by V_a, V_b , and so on, the finite dimensional affine subspaces of V . Denote by V_a^0 the finite dimensional linear subspace of V consisting of differences of elements of V_a . Let $\text{Cliff}(V_a^0)$ be the complexified Clifford algebra on V_a^0 , and let $\mathcal{C}(V_a)$ be the graded C^* -algebra of continuous functions from V_a to $\text{Cliff}(V_a^0)$ vanishing at infinity. Let $\mathcal{S} := C_0(\mathbb{R})$ be the C^* -algebra of all continuous functions on \mathbb{R} vanishing at infinity. Then \mathcal{S} is graded according to odd and even functions. Define the graded tensor product

$$\mathcal{A}(V_a) = \mathcal{S} \hat{\otimes} \mathcal{C}(V_a).$$

If $V_a \subseteq V_b$, we have a decomposition $V_b = V_{ba}^0 + V_a$, where V_{ba}^0 is the orthogonal complement of V_a^0 in V_b^0 . For each $v_b \in V_b$, we have a unique decomposition $v_b = v_{ba} + v_a$, where $v_{ba} \in V_{ba}^0$ and $v_a \in V_a$. Every function h on V_a can be extended to a function \tilde{h} on V_b by the formula: $\tilde{h}(v_{ba} + v_a) = h(v_a)$.

Definition 3.1 ([17]). (1) If $V_a \subseteq V_b$, we define C_{ba} to be the Clifford algebra-valued function $V_b \rightarrow \text{Cliff}(V_b^0)$, $v_b \mapsto v_{ba} \in V_{ba}^0 \subset \text{Cliff}(V_b^0)$. Let X be the function of multiplication by x on \mathbb{R} , considered as a degree one and unbounded multiplier of \mathcal{S} . Define a homomorphism $\beta_{ba} : \mathcal{A}(V_a) \rightarrow \mathcal{A}(V_b)$ by

$$\beta_{ba}(g \hat{\otimes} h) = g(X \hat{\otimes} 1 + 1 \hat{\otimes} C_{ba})(1 \hat{\otimes} \tilde{h})$$

for all $g \in \mathcal{S}$ and $h \in \mathcal{C}(V_a)$, where $g(X \hat{\otimes} 1 + 1 \hat{\otimes} C_{ba})$ is defined by the functional calculus of g on the unbounded, essentially self-adjoint multiplier $X \hat{\otimes} 1 + 1 \hat{\otimes} C_{ba}$.

(2) If $V_a \subset V_b \subset V_c$, then we have $\beta_{cb} \circ \beta_{ba} = \beta_{ca}$. Hence, the above homomorphisms give rise to a directed system $(\mathcal{A}(V_a))$ as V_a ranges over finite dimensional affine subspaces of H . We define a C^* -algebra $\mathcal{A}(V)$ by :

$$\mathcal{A}(V) = \varinjlim \mathcal{A}(V_a),$$

where the direct limit is taken over the directed set of all finite-dimensional affine subspaces $V_a \subset V$, using the homomorphisms β_{ba} in (1).

Now let's return back to the case of interest. Let $(1 \rightarrow N_m \rightarrow G_m \rightarrow G_m/N_m \rightarrow 1)_{m \in \mathbb{N}}$ be a sequence of extensions of countable discrete groups such that $(N_m)_{m \in \mathbb{N}}$ and $(G_m/N_m)_{m \in \mathbb{N}}$ are coarsely embeddable into Hilbert spaces. For each $m \in \mathbb{N}$, let l_{G_m} be a proper length function on G_m , so that a left invariant metric d_{G_m} on G_m is defined by $d_{G_m}(g, h) = l_{G_m}(g^{-1}h)$. Define proper length functions on N_m and G_m/N_m according to $l_{N_m}(h) = l_{G_m}(h)$ for all $h \in N_m$, and $l_{G_m/N_m}(gN_m) = \min\{l_{G_m}(gh) : h \in N_m\}$ for all $gN_m \in G_m/N_m$. Denote the associated left-invariant metrics by d_{N_m} and d_{G_m/N_m} , respectively. Observe that the inclusion $N_m \hookrightarrow G_m$ is an isometry, and the quotient map $G_m \rightarrow G_m/N_m$ is contractive for each $m \in \mathbb{N}$. Assume that the sequence of metric spaces $(G_m)_{m \in \mathbb{N}}$ and $(G_m/N_m)_{m \in \mathbb{N}}$ have uniform bounded geometry.

Let $f_m : G_m/N_m \rightarrow H_m$ be a family of coarse embeddings of $(G_m/N_m)_{m \in \mathbb{N}}$ into a Hilbert spaces $\{H_m\}_{m \in \mathbb{N}}$, respectively. For each $m \in \mathbb{N}$, $n \in \mathbb{N}$, $x \in G_m/N_m$, we define $W_n^m(x)$ to be the finite dimensional Euclidean subspace of H_m spanned by $\{f_m(y) : y \in G_m/N_m, d_{G_m/N_m}(x, y) \leq n^2\}$. Let $W^m(x) = \cup_{n \in \mathbb{N}} W_n^m(x)$. Then we have the following: (1) $W_n^m(x) \subseteq W_{n+1}^m(x)$ for all $n \in \mathbb{N}$ and $x \in$

G_m/N_m ; (2) for each $r > 0$, there exists $N > 0$ independent of m such that $W_n^m(x) \subset W_{n+1}^m(y)$ for all $n \geq N$, $x, y \in G_m/N_m$ with $d_{G_m/N_m}(x, y) \leq r$; (3) by the uniform bounded geometry property of the sequence of metric spaces $(G_n/N_n)_{n \in \mathbb{N}}$, for each $n \in \mathbb{N}$, there exists $d_n > 0$ independent of m such that the dimension of $W_n^m(x)$ is less than d_n for all $x \in G_m/N_m$. It follows that $W^m(x)$ is independent of the choice of $x \in G_m/N_m$. We denote

$$V_m := W^m(x)$$

and, without loss of generality, assume that V_m is dense in H_m . For each $m \in \mathbb{N}$, the set $\mathbb{R}_+ \times H_m$ is equipped with the weakest topology for which the projection to H_m is continuous with respect to the weak topology on H_m , and the function $t^2 + \|h\|^2$ is continuous in $(t, h) \in \mathbb{R}_+ \times H_m$. In this way, the space $\mathbb{R}_+ \times H_m$ is a locally compact Hausdorff space. Note that, for all $v \in H_m$ and $r > 0$, the set $\text{Ball}(v, r) := \{(t, h) \in \mathbb{R}_+ \times H_m \mid t^2 + \|h - v\|^2 < r^2\}$ is an open subset of $\mathbb{R}_+ \times H_m$. For finite dimensional affine subspaces $V_a \subset V_b$ of V_m , the map β_{ba} takes $C_0(\mathbb{R}_+ \times V_a)$ into $C_0(\mathbb{R}_+ \times V_b)$. It follows that the C^* -subalgebra $\varinjlim C_0(\mathbb{R}_+ \times V_a)$ of $C_0(\mathbb{R}_+ \times H_m)$ is in fact $*$ -isomorphic to $C_0(\mathbb{R}_+ \times H_m)$. The support of an element $a \in \mathcal{A}(V_m)$ is defined to be the complement in $\mathbb{R}_+ \times H_m$ of the subset of all (t, h) for which there exists $g \in C_0(\mathbb{R}_+ \times H_m)$ such that $ag = 0$ but $g(t, h) \neq 0$. The algebra $C_0(\mathbb{R}_+ \times H_m)$ acts on $\mathcal{A}(V_m) \hat{\otimes} K$ by the formula: $g(a \hat{\otimes} k) = ga \hat{\otimes} k$. Hence we can define the support of an element in $\mathcal{A}(V_m) \hat{\otimes} K$ in a similar way.

For each $d > 0$, $m \in \mathbb{N}$, the map $f_m : G_m/N_m \rightarrow H_m$ can be extended to the Rips complexes $P_d(G_m/N_m)$ as follows. Let $G_m = \sqcup_{g \in \Lambda} gN_m$ be a coset decomposition of G_m , where $\Lambda \subset G_m$ is a set of representatives of the cosets in G_m/N_m . For any point $x = \sum_{g \in \Lambda} c_g gN_m \in P_d(G_m/N_m)$, where all but finitely many coefficients $c_g \geq 0$ are zero and $\sum_{g \in \Lambda} c_g = 1$, we define

$$f_m(x) = \sum_{g \in \Lambda} c_g f(gN_m).$$

For any point $x = \sum_{g \in \Lambda} c_g gN_m \in P_d(G_m/N_m)$ and $n \in \mathbb{N}$, define $W_n^m(x)$ to be the Euclidean subspace of V_m spanned by $W_n^m(gN_m)$ for all gN_m such that $c_g \neq 0$.

Let $\pi : G_m \rightarrow G_m/N_m$ be the quotient map. It induces a map $\pi : P_d(G_m) \rightarrow P_d(G_m/N_m)$ by

$$\pi \left(\sum_{i=0}^k c_i g_i \right) = \sum_{i=0}^k c_i \pi(g_i),$$

where $c_i \geq 0$ and $\sum_{i=0}^k c_i = 1$. For each $m \in \mathbb{N}$, choose a countable dense subset Z_d^m of $P_d(G_m)$ for each $d \geq 0$ such that $\pi(Z_d^m)$ is dense in $P_d(G_m/N_m)$, and $Z_d^m \subseteq Z_{d'}^m$ if $d \leq d'$.

Definition 3.2. For each $d \geq 0$, the *algebraic uniform twisted Roe algebra* $\mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ is defined to be the set of all tuples $T = (T^{(m)})_{m \in \mathbb{N}}$ where each $T^{(m)} : Z_d^m \times Z_d^m \rightarrow \mathcal{A}(V_m) \hat{\otimes} K$ is a bounded function satisfying the following conditions:

- (1) there exists an integer N such that

$$T^{(m)}(x, y) \in (\beta_N(\pi(x)))(\mathcal{A}(W_N^m(\pi(x)))) \hat{\otimes} K \subseteq \mathcal{A}(V_m) \hat{\otimes} K$$

for each $m \in \mathbb{N}$ and $x, y \in Z_d^m$, where $\beta_N(\pi(x)) : \mathcal{A}(W_N^m(\pi(x))) \rightarrow \mathcal{A}(V_m)$ is the $*$ -homomorphism associated to the inclusion of $W_N^m(\pi(x))$ into V_m , and K is the algebra of compact operators;

(2) there exists $M > 0$ such that $\|T^{(m)}(x, y)\| \leq M$ for each $m \in \mathbb{N}$ and $x, y \in Z_d^m$;

(3) there exists $L > 0$ such that, for each $m \in \mathbb{N}$ and $y \in Z_d^m$,

$$\#\{x : T^{(m)}(x, y) \neq 0\} \leq L, \quad \#\{x : T(y, x) \neq 0\} \leq L;$$

(4) for any sequence of uniformly bounded subsets $(B_m)_{m \in \mathbb{N}}$ where $B_m \subset P_d(G_m)$, the set

$$\{(x, y) \in (B_m \times B_m) \cap (Z_d^m \times Z_d^m) : T^{(m)}(x, y) \neq 0\}$$

is uniformly finite.

(5) there exists $r_1 > 0$ such that for each $m \in \mathbb{N}$, if $d_{P_d(G_m)}(x, y) > r_1$, then $T^{(m)}(x, y) = 0$;

(6) there exists $r_2 > 0$ such that $\text{Supp}(T^{(m)}(x, y)) \subseteq \text{Ball}(f(\pi(x)), r_2)$ for each $m \in \mathbb{N}$ and $x, y \in Z_d^m$ where

$$\text{Ball}(f(\pi(x)), r_2) := \{(s, h) \in \mathbb{R}_+ \times H : s^2 + \|h - f(\pi(x))\|^2 < r_2^2\}.$$

We define a product structure on $\mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ by:

$$(T_1 T_2)^{(m)}(x, y) = \sum_{z \in Z_d^m} T_1^{(m)}(x, z) T_2^{(m)}(z, y).$$

Then $\mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ is made into a $*$ -algebra via matrix multiplications and the $*$ -operation on $\mathcal{A}(V_m) \hat{\otimes} K$. Let

$$E^m = \left\{ \sum_{x \in Z_d^m} a_x[x] : a_x \in \mathcal{A}(V_m) \hat{\otimes} K, \sum_{x \in Z_d^m} a_x^* a_x \text{ converges in norm} \right\}.$$

Note that E^m is a Hilbert C^* -module over $\mathcal{A}(V_m) \hat{\otimes} K$ with

$$\left\langle \sum_{x \in Z_d^m} a_x[x], \sum_{x \in Z_d^m} b_x[x] \right\rangle = \sum_{x \in Z_d^m} a_x^* b_x,$$

$$\left(\sum_{x \in Z_d^m} a_x[x] \right) a = \sum_{x \in Z_d^m} a_x a[x]$$

for all $a \in \mathcal{A}(V_m) \hat{\otimes} K$. We can define a $*$ -representation of the $*$ -algebra $\mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ on E^m as follows:

$$T \left(\sum_{x \in Z_d^m} a_x[x] \right) = \sum_{y \in Z_d^m} \left(\sum_{x \in Z_d^m} T^{(m)}(y, x) a_x[x] \right) [y],$$

where $T = (T^{(m)})_{m \in \mathbb{N}} \in \mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ and $\sum_{x \in Z_d^m} a_x[x] \in E^m$. Then by taking sum of these $*$ -representations, we obtain a faithful $*$ -representation for the $*$ -algebra $\mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ on $\bigoplus_m E^m$.

Definition 3.3. The *uniform twisted Roe algebra* $C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$ is defined to be the operator norm closure of $\mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ in $\mathcal{B}(\bigoplus_m E^m)$, the C^* -algebra of all module homomorphisms from $\bigoplus_m E^m$ to $\bigoplus_m E^m$ for which there is an adjoint module homomorphism.

Definition 3.4. Let $\mathbb{C}_{u,L}[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ be the set of all bounded and uniformly norm-continuous functions

$$g : \mathbb{R}_+ \rightarrow \mathbb{C}_{u,L}[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$$

satisfying the following conditions:

- (1) there exists $N > 0$ such that $(g(t))(x, y) \in (\beta_N(\pi(x)))(\mathcal{A}(W_N^m(\pi(x)))) \hat{\otimes} K \subseteq \mathcal{A}(V_m) \hat{\otimes} K$ for all $m \in \mathbb{N}, t \in \mathbb{R}_+, x, y \in Z_d^m$;
- (2) there exists a bounded function $R : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that $\lim_{t \rightarrow \infty} R(t) = 0$ and for all $m \in \mathbb{N}, x, y \in Z_d^m$, if $d_{P_d(G_m/N_m)}(\pi(x), \pi(y)) > R(t)$, then $(g(t))(x, y) = 0$;
- (3) there exists $R > 0$ such that $\text{Supp}(g(t)(x, y)) \subseteq \text{Ball}(f(\pi(x)), R)$ for all $m \in \mathbb{N}, t \in \mathbb{R}_+, x, y \in Z_d^m$.

The *uniform twisted localization algebra* $C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$ is defined to be the norm completion of $\mathbb{C}_{u,L}[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ with respect to the norm $\|g\| = \sup_{t \in \mathbb{R}_+} \|g(t)\|_{C^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})}$.

Remark 3.5. For each $m \in \mathbb{N}$, we can define algebraic twisted Roe algebra $\mathbb{C}[(P_d(G_m), \mathcal{A}(V_m))]$ consisting of all bounded functions $T^{(m)} : Z_d^m \times Z_d^m \rightarrow \mathcal{A}(V_m) \hat{\otimes} K$ satisfying the conditions of Definition 3.2. Then the $*$ -algebra $\mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ can be regarded as $\prod_{m \in \mathbb{N}}^u \mathbb{C}[P_d(G_m), \mathcal{A}(V_m)]$, where we add u on the direct product symbol to emphasize that the latter $*$ -algebra consists of the family of operators $T^{(m)}$, $m \in \mathbb{N}$, satisfy the conditions of Definition 3.2 with uniform constants. Denote by $\prod_{m \in \mathbb{N}}^u C^*(P_d(G_m), \mathcal{A}(V_m))$, is the operator norm closure of $\prod_{m \in \mathbb{N}}^u \mathbb{C}[P_d(G_m), \mathcal{A}(V_m)]$ in $\mathcal{B}(\oplus_m E^m)$. Naturally, the C^* -algebra $\prod_{m \in \mathbb{N}}^u C^*(P_d(G_m), \mathcal{A}(V_m))$ is equal to $C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$. We can similarly define the C^* -algebra $\prod_{m \in \mathbb{N}}^u C_L^*(P_d(G_m), \mathcal{A}(V_m))$ as another form of $C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$.

There is a natural *evaluation-at-zero* homomorphism

$$e : C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}) \rightarrow C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$$

defined by $e(g) = g(0)$. It induces a homomorphism at the K -theory level:

$$e_* : \lim_{d \rightarrow \infty} K_*(C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})) \rightarrow \lim_{d \rightarrow \infty} K_*(C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})).$$

4 Reduction to the sequence of normal subgroups $(N_m)_{m \in \mathbb{N}}$

In this section, we shall prove that the "*twisted coarse Baum-Connes conjecture for the sequence $(G_m)_{m \in \mathbb{N}}$* " holds. Namely, we have the following result.

Theorem 4.1. *Let $(1 \rightarrow N_m \rightarrow G_m \rightarrow G_m/N_m \rightarrow 1)_{m \in \mathbb{N}}$ be an extension of countable discrete groups. Assume that the sequence of metric spaces $(G_m)_{m \in \mathbb{N}}$ have uniform bounded geometry. If $(N_m)_{m \in \mathbb{N}}$ and*

$(G_m/N_m)_{m \in \mathbb{N}}$ are coarsely embeddable into Hilbert space, then the evaluation-at-zero homomorphism on K -theory

$$e_* : \lim_{d \rightarrow \infty} K_*(C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})) \rightarrow \lim_{d \rightarrow \infty} K_*(C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}))$$

is an isomorphism.

The proof proceeds by decomposing the twisted algebras into various smaller ideals or subalgebras, and applying a Mayer-Vietoris sequence argument and the strong Lipschitz homotopy invariance for these subalgebras in the twisted localization algebras to reduce the twisted coarse Baum-Connes conjecture for the sequence $(G_n)_{n \in \mathbb{N}}$ to the coarse Baum-Connes conjecture for sequence of the normal subgroups $(N_m)_{m \in \mathbb{N}}$ with *non-twisted coefficients*, which holds by appealing to Yu's original work in [33] or the recent work [10] of J. Deng, Q. W and G. Yu for the sequence $(N_m)_{m \in \mathbb{N}}$.

To do so, we first discuss ideals of the uniform twisted algebras associated with any family of open subsets of $\{\mathbb{R}_+ \times H_m\}_{m \in \mathbb{N}}$.

Definition 4.2. (1) The support of an element $T = (T^{(m)})_{m \in \mathbb{N}} \in \mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ is defined to be a sequence of subsets

$$\text{Supp}(T) := (\text{Supp}(T^{(m)}))_{m \in \mathbb{N}}$$

where

$$\text{Supp}(T^{(m)}) := \left\{ (x, y, t, h) \in Z_d^m \times Z_d^m \times \mathbb{R}_+ \times H_m : (t, h) \in \text{Supp}(T^{(m)}(x, y)) \right\}.$$

(2) The support of an element g in $\mathbb{C}_{u,L}[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ is defined to be $\bigcup_{t \in \mathbb{R}_+} \text{Supp}(g(t))$.

Let $O = (O_m)_{m \in \mathbb{N}}$ be a sequence of open subsets, where each O_m is an open subset in $\mathbb{R}_+ \times H_m$. Define $\mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]_O$ to be the subalgebra of $\mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$ consisting of all elements $T = (T^{(m)})_{m \in \mathbb{N}}$ satisfying $\text{Supp}(T^{(m)}) \subset Z_d^m \times Z_d^m \times O_m$ for each $m \in \mathbb{N}$, i.e.,

$$\mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]_O = \left\{ T \in \mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}] : \text{Supp}(T^{(m)}(x, y)) \subseteq O_m, \forall m \in \mathbb{N}, x, y \in Z_d^m \right\}.$$

Define C^* -subalgebra $C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_O$ to be the norm closure of $\mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]_O$ in $C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$. similarly, We can also define C^* -subalgebra $C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_O$. Note that $C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_O$ and $C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_O$ are closed two-sided ideals of $C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$ and $C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$, respectively.

We have the following lemma analogous to [33, Lemma 6.3].

Lemma 4.3. For each $m \in \mathbb{N}$, $v_m \in V_m$, $r > 0$, let $\text{Ball}(v_m, r) = \{(t, h) \in \mathbb{R}_+ \times H_m : t^2 + \|h - v_m\| < r^2\}$, and X_i^m , $1 \leq i \leq i_0$, be mutually disjoint subsets of G_m/N_m . Fix arbitrarily $1 < j \leq i_0$. Let $O_r = (O_{m,r})_{m \in \mathbb{N}}$ and $O'_r = (O'_{m,r})_{m \in \mathbb{N}}$ be two sequences of open subsets, in which for each $m \in \mathbb{N}$,

$$O_{m,r} = \bigcup_{i=1}^{j-1} \left(\bigcup_{x \in X_i^m} \text{Ball}(f(x), r) \right) \quad \text{and} \quad O'_{m,r} = \bigcup_{x \in X_j^m} \text{Ball}(f(x), r),$$

where $f_m : G_m/N_m \rightarrow H_m$ is the coarse embedding. Then for each $r_0 > 0$ we have

$$\begin{aligned}
(1) \quad & \lim_{r < r_0, r \rightarrow r_0} C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r \cup O'_r} \\
& = \lim_{r < r_0, r \rightarrow r_0} C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r} + \lim_{r < r_0, r \rightarrow r_0} C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O'_r}; \\
(2) \quad & \lim_{r < r_0, r \rightarrow r_0} C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r \cap O'_r} \\
& = \lim_{r < r_0, r \rightarrow r_0} (C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r} \cap C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O'_r}); \\
(3) \quad & \lim_{r < r_0, r \rightarrow r_0} C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r \cup O'_r} \\
& = \lim_{r < r_0, r \rightarrow r_0} C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r} + \lim_{r < r_0, r \rightarrow r_0} C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O'_r}; \\
(4) \quad & \lim_{r < r_0, r \rightarrow r_0} C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r \cap O'_r} \\
& = \lim_{r < r_0, r \rightarrow r_0} (C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r} \cap C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O'_r}).
\end{aligned}$$

It would be convenient to introduce the following terminology.

Definition 4.4. Let $r > 0$. For each $m \in \mathbb{N}$, a family of open subsets $\{O_{m,r}(i)\}_{i \in J}$ of $\mathbb{R}_+ \times H_m$ is said to be $(G_m/N_m, r)$ -separate if:

- (1) $O_{m,r}(i) \cap O_{m,r}(j) = \emptyset$ if $i \neq j$;
- (2) for each $i \in J$, there exists $x_i^m \in G_m$ such that $O_{m,r}(i) \subseteq \text{Ball}(f_m(\pi(x_i^m)), r)$, where f is the coarse embedding $f : G_m/N_m \rightarrow H_m$ and

$$\text{Ball}(f_m(\pi(x_i^m)), r) = \{(t, h) \in \mathbb{R}_+ \times H_m : t^2 + \|h - f_m(\pi(x_i^m))\|^2 < r^2\}.$$

The following proposition contains a key step in reducing the twisted coarse Baum-Connes conjecture for the sequence $(G_m)_{m \in \mathbb{N}}$ to the coarse Baum-Connes conjecture for the sequence $(N_m)_{m \in \mathbb{N}}$ with non-twisted coefficients.

Proposition 4.5. Let $r > 0$. If $O_r = (O_{m,r})_{m \in \mathbb{N}}$ be a sequence of open subsets, where each $O_{m,r}$ is a family of $(G_m/N_m, r)$ -separate open subsets of $\mathbb{R}_+ \times H_m$. Then

$$e_* : \lim_{d \rightarrow \infty} K_*(C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r}) \rightarrow \lim_{d \rightarrow \infty} K_*(C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r})$$

is an isomorphism.

Proof. We first discuss the structure of the C^* -algebra $C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r}$.

Let $T = (T^{(m)})_{m \in \mathbb{N}} \in \mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}]$. For each $m \in \mathbb{N}$, since $O_{m,r}$ is the union of a family of disjoint open subsets $\{O_{m,r}(i)\}_{i \in J}$ of $\mathbb{R}_+ \times H_m$, the operator $T^{(m)}$ has a formal decomposition $T^{(m)} = \sum_{i \in J} T_i^{(m)}$, where $T_i^{(m)} := T^{(m)}|_{O_{m,r}(i)}$ is the restriction of $T^{(m)}$ on the component $O_{m,r}(i)$, so that $\text{Supp}(T_i^{(m)}) \subseteq Z_d^m \times Z_d^m \times O_{m,r}(i)$. It follows that the correspondence

$$T = (T^{(m)})_{m \in \mathbb{N}} \mapsto ((T_i^{(m)})_{i \in J})_{m \in \mathbb{N}}$$

gives rise to a $*$ -isomorphism

$$\mathbb{C}_u[(P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}] \cong \prod_{m \in \mathbb{N}} \prod_{i \in J} \mathbb{C}[P_d(G_m), \mathcal{A}(V_m)]_{O_{m,r}(i)} \cong \prod_{\substack{i \in J \\ m \in \mathbb{N}}} \mathbb{C}[P_d(G_m), \mathcal{A}(V_m)]_{O_{m,r}(i)}$$

where we add u on the direct product symbol to emphasize that the family of operators $T_i^{(m)}$, $i \in J, m \in \mathbb{N}$, satisfy the conditions of Definition 3.2 with uniform constants.

Since $\{O_{m,r}(i)\}_{i \in J}$ is a family of $(G_m/N_m, r)$ -separate open subsets of $\mathbb{R}_+ \times H_m$, there exists $x_i^m \in G_m$ such that $O_{m,r}(i) \subseteq \text{Ball}(f_m(\pi(x_i^m)), r)$ for each $i \in J$. Note that the condition (6) in Definition 3.2 and the coarse embedding property of the sequence $(G_m/N_m)_{m \in \mathbb{N}}$ imply that for each $m \in \mathbb{N}$, $d > 0$ and $i \in J$,

$$C^*(P_d(G_m), \mathcal{A}(V_m))_{O_{m,r}(i)} = \lim_{R \rightarrow \infty} C^*\left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m)\right)_{O_{m,r}(i)},$$

where $\mathcal{N}_{G_m}(x_i^m N_m, R)$ denotes the R -neighborhood of the coset $x_i^m N_m$ in G_m . More concretely, for each $i \in J$ and $R > 0$, there exists an integer $N_R > 0$, and $g_1^{(i)}, g_2^{(i)}, \dots, g_{N_R}^{(i)}$ in G_m , such that

$$\mathcal{N}_{G_m}(x_i^m N_m, R) = \bigsqcup_{k=1}^{N_R} g_k^{(i)} N_m,$$

where $g_k^{(i)} N_m \in G_m/N_m$ satisfies $d_{G_m/N_m}(\pi(g_k^{(i)}), \pi(x_i^m)) < R$. In other words, for each k there exists $h_k^{(i)} \in N_m$ such that $d_G(g_k^{(i)}, x_i^m h_k^{(i)}) < R$. Note that for each fixed $R > 0$, the uniform bounded geometry property of the sequence $(G_m/N_m)_{m \in \mathbb{N}}$ implies that the integer N_R is independent of $m \in \mathbb{N}$ and $i \in J$. Since the family of coarse embeddings $f_m : G_m/N_m \rightarrow H_m$ are controlled by two common non-decreasing functions, it follows that

$$C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r} = \lim_{R \rightarrow \infty} \prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^*\left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m)\right)_{O_{m,r}(i)}$$

and

$$C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r} = \lim_{R \rightarrow \infty} \prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^*\left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m)\right)_{O_{m,r}(i)}.$$

Hence, to prove the proposition, we only need to prove that

$$e_* : \lim_{d \rightarrow \infty} \lim_{R \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^*\left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m)\right)_{O_{m,r}(i)} \right) \rightarrow \\ \lim_{d \rightarrow \infty} \lim_{R \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^*\left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m)\right)_{O_{m,r}(i)} \right)$$

is an isomorphism.

Note that for any fixed $d > 0$, if $R < R'$ then $\mathcal{N}_{G_m}(x_i^m N_m, R) \subseteq \mathcal{N}_{G_m}(x_i^m N_m, R')$ for each $m \in \mathbb{N}$ and $i \in J$, so that we have natural inclusions

$$\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^*\left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m)\right)_{O_{m,r}(i)} \subseteq \prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^*\left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R')), \mathcal{A}(V_m)\right)_{O_{m,r}(i)}$$

and

$$\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^*\left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m)\right)_{O_{m,r}(i)} \subseteq \prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^*\left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R')), \mathcal{A}(V_m)\right)_{O_{m,r}(i)}.$$

On the other hand, for any fixed $R > 0$, if $d < d'$, we have natural inclusions

$$\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^* \left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m) \right)_{O_{m,r(i)}} \subseteq \prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^* \left(P_{d'}(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m) \right)_{O_{m,r(i)}}$$

and

$$\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^* \left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m) \right)_{O_{m,r(i)}} \subseteq \prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^* \left(P_{d'}(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m) \right)_{O_{m,r(i)}}.$$

Therefore, we can change the order of inductive limits on K -theory groups:

$$\begin{aligned} \lim_{d \rightarrow \infty} \lim_{R \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^* \left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m) \right)_{O_{m,r(i)}} \right) &\cong \\ \lim_{R \rightarrow \infty} \lim_{d \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^* \left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m) \right)_{O_{m,r(i)}} \right) & \end{aligned}$$

and

$$\begin{aligned} \lim_{d \rightarrow \infty} \lim_{R \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^* \left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m) \right)_{O_{m,r(i)}} \right) &\cong \\ \lim_{R \rightarrow \infty} \lim_{d \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^* \left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m) \right)_{O_{m,r(i)}} \right) & \end{aligned}$$

As a result, to prove the proposition, it suffices to prove that, for each fixed $R > 0$,

$$\begin{aligned} e_* : \lim_{d \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^* \left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m) \right)_{O_{m,r(i)}} \right) &\rightarrow \\ \lim_{d \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^* \left(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R)), \mathcal{A}(V_m) \right)_{O_{m,r(i)}} \right) & \end{aligned}$$

is an isomorphism.

Note that when d is large enough, for each $m \in \mathbb{N}$, $P_d(\mathcal{N}_{G_m}(x_i^m N_m, R))$ as a subcomplex of $P_d(G_m)$ is strongly Lipschitz homotopy equivalent to $P_d(x_i^m N_m)$ by a linear homotopy. More precisely, for each $h \in N_m$, there exists $\tilde{h} \in N_m$ such that

$$d_{P_d(G_m)}(g_k^{(i)} h, x_i \tilde{h} h_k^{(i)}) = d_{P_d(G_m)}(g_k^{(i)}, x_i h_k^{(i)}) < R$$

for each $g_k^{(i)} N \subseteq \mathcal{N}_{G_m}(x_i^m N_m, R)$, so that we can connect $g_k^{(i)} h \in g_k^{(i)} N_m$ with $x_i \tilde{h} h_k^{(i)} \in x_i N_m$ by a line segment for each $h \in N_m$. Since $x_i N_m$ is isometric to N_m , it follows that $P_d(\mathcal{N}_{G_m}(x_i^m N_m, R))$ as a

subcomplex of $P_d(G_m)$ is strongly Lipschitz homotopy equivalent to $P_d(N_m)$ for all $m \in \mathbb{N}$ and $i \in J$. Moreover, the constants controlling the strong Lipschitz homotopy equivalences are independent of $m \in \mathbb{N}$ and $i \in J$, because the metric on the Rips complex $P_d(G_m)$ is symmetric/homogeneous. It follows from the strong Lipschitz homotopy invariance for K -theory of Roe algebras and localization algebras that the inclusion maps $P_d(x_i^m N_m) \rightarrow P_d(\mathcal{N}_{G_m}(x_i^m N_m, R))$ for each $m \in \mathbb{N}$ induce the following commutative diagram

$$\begin{array}{ccc}
\lim_{d \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^*(P_d(N_m)), \mathcal{A}(V_m)_{O_{m,r}(i)} \right) & \xrightarrow{e_*} & \lim_{d \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^*(P_d(N_m)), \mathcal{A}(V_m)_{O_{m,r}(i)} \right) \\
\downarrow \cong & & \downarrow \cong \\
\lim_{d \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^*(P_d(\mathcal{N}_{G_m}(x_i^m N_m))), \mathcal{A}(V_m)_{O_{m,r}(i)} \right) & \xrightarrow{e_*} & \lim_{d \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^*(P_d(\mathcal{N}_{G_m}(x_i^m N_m))), \mathcal{A}(V_m)_{O_{m,r}(i)} \right) \\
\downarrow \cong & & \downarrow \cong \\
\lim_{d \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^*(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R))), \mathcal{A}(V_m)_{O_{m,r}(i)} \right) & \xrightarrow{e_*} & \lim_{d \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^*(P_d(\mathcal{N}_{G_m}(x_i^m N_m, R))), \mathcal{A}(V_m)_{O_{m,r}(i)} \right)
\end{array}$$

where the vertical maps are isomorphisms by strongly Lipschitz homotopy invariance.

Therefore, to prove the proposition, it suffices to prove that

$$e_* : \lim_{d \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^*(P_d(N_m)), \mathcal{A}(V_m)_{O_{m,r}(i)} \right) \rightarrow \lim_{d \rightarrow \infty} K_* \left(\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^*(P_d(N_m)), \mathcal{A}(V_m)_{O_{m,r}(i)} \right)$$

is an isomorphism.

Note that we have now arrived at the Roe algebras of $P_d(N_m)$ with non-twisted coefficients $\prod_{i \in J} \mathcal{A}_{O_{m,r}(i)}$. Namely, we have

$$\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C^*(P_d(N_m)), \mathcal{A}(V_m)_{O_{m,r}(i)} \cong \prod_{m \in \mathbb{N}}^u C^*(P_d(N_m)), \prod_{i \in J} \mathcal{A}(V_m)_{O_{m,r}(i)}$$

and

$$\prod_{\substack{i \in J \\ m \in \mathbb{N}}}^u C_L^*(P_d(N_m)), \mathcal{A}(V_m)_{O_{m,r}(i)} \cong \prod_{m \in \mathbb{N}}^u C_L^*(P_d(N_m)), \prod_{i \in J} \mathcal{A}(V_m)_{O_{m,r}(i)}$$

where $\prod_{i \in J} \mathcal{A}_{O_{m,r}(i)} = \left\{ (a_i)_{i \in J} : a_i \in \mathcal{A}_{O_{m,r}(i)}, \sup_{i \in J} \|a_i\| < +\infty \right\}$ for each $m \in \mathbb{N}$.

Since the sequence of normal subgroups $(N_m)_{m \in \mathbb{N}}$ with the proper metric d_{N_m} induced from the sequence $(G_m)_{m \in \mathbb{N}}$ is a sequence of metric spaces with uniform bounded geometry, and admits a coarse embedding into Hilbert space by assumption, we can appeal to the same proof of the celebrate Theorem 1.1 of G. Yu [33] to conclude that

$$e_* : \lim_{d \rightarrow \infty} K_* \left(\prod_{m \in \mathbb{N}}^u C_L^*(P_d(N_m)), \prod_{i \in J} \mathcal{A}(V_m)_{O_{m,r}(i)} \right) \rightarrow \lim_{d \rightarrow \infty} K_* \left(\prod_{m \in \mathbb{N}}^u C^*(P_d(N_m)), \prod_{i \in J} \mathcal{A}(V_m)_{O_{m,r}(i)} \right)$$

is an isomorphism. Consequently, we have that

$$e_* : \lim_{d \rightarrow \infty} K_*(C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r}) \rightarrow \lim_{d \rightarrow \infty} K_*(C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r})$$

is an isomorphism, as desired. The proof is complete. \square

Now we are ready to prove the main result of this section:

Proof of Theorem 4.1: For any $r > 0$, we define $O_{m,r} \subseteq \mathbb{R}_+ \times H_m$ by

$$O_{m,r} = \bigcup_{x \in G_m/N_m} \text{Ball}(f(x), r),$$

where $f_m : G_m/N_m \rightarrow H_m$ is the coarse embedding, and

$$\text{Ball}(f(x), r) := \{(t, h) \in \mathbb{R}_+ \times H_m : t^2 + \|h - f(x)\|^2 < r\}.$$

Denote $O_r = (O_{m,r})_{m \in \mathbb{N}}$. By definition, we have

$$\begin{aligned} C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r}) &= \lim_{r \rightarrow \infty} C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r}, \\ C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r}) &= \lim_{r \rightarrow \infty} C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r}. \end{aligned}$$

It is straightforward to check that

$$\begin{aligned} \lim_{d \rightarrow \infty} \lim_{r \rightarrow \infty} K_*(C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r}) &= \lim_{r \rightarrow \infty} \lim_{d \rightarrow \infty} K_*(C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r}), \\ \lim_{d \rightarrow \infty} \lim_{r \rightarrow \infty} K_*(C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r}) &= \lim_{r \rightarrow \infty} \lim_{d \rightarrow \infty} K_*(C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r}). \end{aligned}$$

So it suffices to show that, for any $r_0 > 0$,

$$e_* : \lim_{d \rightarrow \infty} K_* \left(\lim_{r < r_0, r \rightarrow r_0} C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r} \right) \rightarrow \lim_{d \rightarrow \infty} K_* \left(\lim_{r < r_0, r \rightarrow r_0} C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})_{O_r} \right)$$

is an isomorphism.

Since the sequence $(G_m/N_m)_{m \in \mathbb{N}}$ has uniform bounded geometry and the family of coarse embeddings $f_m : G_m/N_m \rightarrow H_m$ are controlled by two common non-decreasing functions, for any $r_0 > 0$ and $m \in \mathbb{N}$, there exist finitely many, say k_0 independent of m , mutually disjoint subsets $\{X_{m,k}\}_{k=1}^{k_0}$ of G_m/N_m such that $G_m/N_m = \bigsqcup_{k=1}^{k_0} X_{m,k}$, and for each $1 \leq k \leq k_0$, we have $d(f(x), f(x')) > 3r_0$ for any two distinct elements $x, x' \in X_{m,k}$.

For any $0 < r < r_0$ and each $1 \leq k \leq k_0$, let

$$O_{m,r,k} = \bigcup_{x \in X_{m,k}} \text{Ball}(f(x), r).$$

Then $O_{m,r} = \bigcup_{k=1}^{k_0} O_{m,r,k}$. Moreover, every $O_{m,r,k}$ and

$$O_{m,r,k} \cap \left(\bigcup_{j=1}^{k-1} O_{m,r,j} \right)$$

are $(G_m/N_m, r)$ -separate for any $1 < k \leq k_0$. By Proposition 4.5 and a Mayer-Vietoris sequence argument, we have

$$e_* : \lim_{d \rightarrow \infty} K_*(C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})) \rightarrow \lim_{d \rightarrow \infty} K_*(C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}))$$

is an isomorphism. This completes the proof. \square

5 The Bott maps and proof of the main theorem

In this section, we shall prove the main theorem of this paper. We shall construct an asymptotic morphism β from $\mathcal{S} \hat{\otimes} C_u^*((P_d(G_m))_{m \in \mathbb{N}})$ to $C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$, which plays the role of the Bott map in the geometric analogue, as proved by G. Yu [33], of infinite dimensional Bott periodicity introduced by Higson, Kasparov and Trout in [17]. This geometric analogue of the infinite dimensional Bott periodicity is then used to reduce the coarse Novikov conjecture for the sequence of extensions of coarsely embeddable groups $(N_m)_{m \in \mathbb{N}}$ and $(G_m/N_m)_{m \in \mathbb{N}}$ to the twisted coarse Baum-Connes conjecture for the sequence $(G_m)_{m \in \mathbb{N}}$ with coefficients coming from a family of coarse embeddings of quotient groups $(G_m/N_m)_{m \in \mathbb{N}}$ to Hilbert spaces. We thus complete the proof of Theorem 1.1 by recalling Proposition 4.1 of the previous section.

Let $(1 \rightarrow N_m \rightarrow G_m \rightarrow G_m/N_m \rightarrow 1)_{m \in \mathbb{N}}$ be a sequence of extensions of countable discrete groups such that $(N_m)_{m \in \mathbb{N}}$ and $(G_m/N_m)_{m \in \mathbb{N}}$ are coarsely embeddable into Hilbert space. Choose a family of proper length function $(l_{G_m})_{m \in \mathbb{N}}$ on $(G_m)_{m \in \mathbb{N}}$ to endow three sequences of groups left invariant metrics d_{N_m} , d_{G_m} and d_{G_m/N_m} as before. Let $\pi : G_m \rightarrow G_m/N_m$ be the quotient map, and let $f_m : G_m/N_m \rightarrow H_m$ be a coarse embedding of G_m/N_m into a real Hilbert space H_m for each $m \in \mathbb{N}$. Recall that for each $n \in \mathbb{N}$ and $x \in G_m/N_m$, we define

$$W_n^m(x) := \text{span}\{f(y) : y \in G_m/N_m, d_{G_m/N_m}(x, y) \leq n^2\},$$

$$V_m := W_m(x) := \cup_{n \in \mathbb{N}} W_n^m(x)$$

and, without loss of generality, assume that V_m is dense in H_m . For each $d > 0$, the map $f : G_m/N_m \rightarrow H_m$ is extended to the Rips complex $f : P_d(G_m/N_m) \rightarrow H_m$. Choose a countable dense subset Z_d^m of $P_d(G_m)$ such that $\pi(Z_d^m)$ is dense in $P_d(G_m/N_m)$, and $Z_d^m \subseteq Z_{d'}^m$ if $d \leq d'$. For each $z \in Z_d^m$ with $\pi(z) = \sum_{x \in G_m/N_m} c_x x$, we define $W_N^m(\pi(z))$ to be the Euclidean subspace of V_m spanned by $W_n^m(x)$ for those x such that $c_x \neq 0$.

For each $m \in \mathbb{N}$ and $x \in Z_d^m$, and any finite-dimensional affine subspace W_m of V_m containing $f_m(\pi(x))$, recall that in Definition 3.1 we define a *-homomorphism

$$\beta_{W_m}(x) : \mathcal{S} := C_0(\mathbb{R}) \rightarrow \mathcal{A}(W_m)$$

by the formula

$$\beta_{W_m}(x)(g) = g \left(X \hat{\otimes} 1 + 1 \hat{\otimes} C_{W_m, f_m(\pi(x))} \right)$$

for all $g \in \mathcal{S}$, where X is the degree one and unbounded multiplier of \mathcal{S} defined by $(X(g))(x) = xg(x)$ for each $g \in \mathcal{S}$ with compact support and all $x \in \mathbb{R}$, and $C_{W_m, f_m(\pi(x))}$ is the Clifford algebra-valued function on W_m defined by

$$C_{W_m, f_m(\pi(x))}(v) = v - f(\pi(x)) \in W_m^0 \subset \text{Cliff}(W_m^0)$$

for all $v \in W_m$. Taking inductive limit as in Definition 3.1, these maps $\beta_{W_m}(x)$ for all W_m as above induce a *-homomorphism

$$\beta_m(x) : \mathcal{S} \rightarrow \mathcal{A}(V_m).$$

Definition 5.1. Let $d \geq 0$. For each $t \in [1, \infty)$, we define a map

$$\beta_t : \mathcal{S} \hat{\otimes} \mathbb{C}_u[(P_d(G_m))_{m \in \mathbb{N}}] \rightarrow C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$$

by the formula

$$(\beta_t(g \hat{\otimes} T))^{(m)}(x, y) = \beta_m(x)(g_t) \hat{\otimes} T^{(m)}(x, y)$$

for all $g \in \mathcal{S}, T = (T^{(m)})_{m \in \mathbb{N}} \in \mathbb{C}_u[(P_d(G_m))_{m \in \mathbb{N}}]$ and $x, y \in Z_d^m$, where $g_t(\tau) = g(t^{-1}\tau)$ for all $\tau \in \mathbb{R}$, and $\beta_m(x) : \mathcal{S} \rightarrow \mathcal{A}(V_m)$ is the *-homomorphism associated to the inclusion of the zero-dimensional affine space $f_m(\pi(x))$ into V_m defined above.

Similarly, we have the following definition associated to the uniform localization algebras.

Definition 5.2. Let $d \geq 0$. For each $t \in [1, \infty)$, we define a map

$$(\beta_{u,L})_t : \mathcal{S} \hat{\otimes} \mathbb{C}_{u,L}[(P_d(G_m))_{m \in \mathbb{N}}] \rightarrow C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$$

by the formula

$$\left((\beta_{u,L})_t(g \hat{\otimes} h) \right)(\tau) = \beta_t(g \hat{\otimes} h(\tau))$$

for all $g \in \mathcal{S}, h \in \mathbb{C}_{u,L}[(P_d(G_m))_{m \in \mathbb{N}}]$ and $\tau \in [0, \infty)$, where β_t is defined in Definition 5.1 above.

We have the following result analogous to Lemma 7.6 in [33].

Lemma 5.3. For each $d \geq 0$, the maps $(\beta_t)_{t \geq 1}$ and $((\beta_{u,L})_t)_{t \geq 1}$ extend to asymptotic morphisms

$$\begin{aligned} \beta : \mathcal{S} \hat{\otimes} C_u^*((P_d(G_m))_{m \in \mathbb{N}}) &\rightsquigarrow C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}), \\ \beta_{u,L} : \mathcal{S} \hat{\otimes} C_L^*((P_d(G_m))_{m \in \mathbb{N}}) &\rightsquigarrow C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}). \end{aligned}$$

Proof. To show that the maps $(\beta)_{t \geq 1}$ give rise to an asymptotic morphism, we first claim that

$$\|\beta(x)(g_t) - \beta(y)(g_t)\|$$

tends to 0 uniformly on the band $\{(x, y) \in Z_d^m \times Z_d^m : d_{P_d(G_m)}(x, y) \leq r\}_{m \in \mathbb{N}}$ for each fixed $g \in C_0(\mathbb{R})$ and $r > 0$.

Indeed, for each $m \in \mathbb{N}, r > 0$ and $x, y \in Z_d^m$ with $d_{P_d(G_m)}(x, y) \leq r$, take $n \geq 1$ such that the finite-dimensional Euclidean subspace $W_n^m(\pi(x))$ of V contains $0, f_m(\pi(x))$ and $f_m(\pi(y))$. By Definition 3.1, we have

$$\begin{aligned} \beta_m(x)(g_t) &= \left(\beta_{V_m, W_n^m(\pi(x))} \circ \beta_{W_n^m(\pi(x))}(x) \right)(g_t), \\ \beta_m(y)(g_t) &= \left(\beta_{V_m, W_n^m(\pi(x))} \circ \beta_{W_n^m(\pi(x))}(y) \right)(g_t) \end{aligned}$$

where

$$\beta_{V_m, W_n^m(\pi(x))} : \mathcal{A}(W_n^m(\pi(x))) \rightarrow \mathcal{A}(V_m)$$

is the *-homomorphism associated to the inclusion $W_n^m(\pi(x)) \subset V_m$. Thus it suffices to deal with

$$\left(\beta_{W_n^m(\pi(x))}(x) \right)(g_t) - \left(\beta_{W_n^m(\pi(x))}(y) \right)(g_t).$$

Denote $G_x = X \hat{\otimes} 1 + 1 \hat{\otimes} C_{W_n^m(\pi(x)), \pi(x)}$ and $G_y = X \hat{\otimes} 1 + 1 \hat{\otimes} C_{W_n^m(\pi(x)), \pi(y)}$. Note that

$$C_{W_n^m(\pi(x)), \pi(x)}(v) - C_{W_n^m(\pi(x)), \pi(y)}(v) = (v - f_m(\pi(x))) - (v - f_m(\pi(y))) = f_m(\pi(y)) - f_m(\pi(x))$$

for all $v \in W_n^m(x)$. For the generators $g(s) = (s \pm i)^{-1}$ of $C_0(\mathbb{R})$, and for $t \in [1, \infty)$, we have that

$$\begin{aligned} \left\| \left(\beta_{W_n^m(\pi(x))}(x) \right)(g_t) - \left(\beta_{W_n^m(\pi(x))}(y) \right)(g_t) \right\| &= \left\| \left(t^{-1} G_x \pm i \right)^{-1} - \left(t^{-1} G_y \pm i \right)^{-1} \right\| \\ &\leq t \left\| (G_x \pm ti)^{-1} \right\| \|G_x - G_y\| \left\| (G_y \pm ti)^{-1} \right\| \\ &\leq t^{-1} \|f_m(\pi(y)) - f_m(\pi(x))\|. \end{aligned}$$

Since the family of coarse embeddings $f_m : P_d(G_m/N_m) \rightarrow H_m$ are controlled two common non-decreasing functions and the quotient map $\pi : P_d(G_m) \rightarrow P_d(G_m/N_m)$ is contractive, we know that

$$\sup_{m \in \mathbb{N}} \left\{ \|f_m(\pi(x)) - f_m(\pi(y))\| \mid x, y \in Z_d^m, d_{P_d(G_m)}(x, y) \leq r \right\} < \infty.$$

Hence,

$$\left\| \left(\beta_{W_n^m(\pi(x))}(x) \right)(g_t) - \left(\beta_{W_n^m(\pi(x))}(y) \right)(g_t) \right\|$$

tends to 0 uniformly on $\{(x, y) \in Z_d^m \times Z_d^m : d_{P_d(G_m)}(x, y) \leq r\}_{m \in \mathbb{N}}$ for $g(s) = (s \pm i)^{-1}$ as $t \rightarrow \infty$. However, by the Stone-Weierstrass theorem, the algebra generated by these two functions is dense in $\mathcal{S} := C_0(\mathbb{R})$. So we complete the proof of the claim.

Let E^m be the Hilbert C^* -module over $\mathcal{A}(V_m) \hat{\otimes} K$ as defined in Definition 3.3. For every $g \in \mathcal{S}$ and $T = (T^{(m)})_{m \in \mathbb{N}} \in \mathbb{C}_u[(P_d(G_m))_{m \in \mathbb{N}}]$, and for $t \in [1, \infty)$, note that we have the following formula:

$$\beta_t(g \hat{\otimes} T) = N_{g_t} \cdot (1 \hat{\otimes} T),$$

where $N_{g_t} : \bigoplus_m E^m \rightarrow \bigoplus_m E^m$ is defined to be the direct sum $N_{g_t} = \bigoplus_m N_{g_t}^m$, in which the bounded module homomorphisms $N_{g_t}^m$ are defined by

$$N_{g_t}^m \left(\sum_{x \in Z_d^m} a_x[x] \right) = \sum_{x \in Z_d^m} \left(\beta(x)(g_t) \hat{\otimes} 1 \right) a_x[x],$$

and $1 \hat{\otimes} T : \bigoplus_m E^m \rightarrow \bigoplus_m E^m$ is the bounded module homomorphism defined by

$$(1 \hat{\otimes} T) \bigoplus_m \left(\sum_{x \in Z_d^m} a_x[x] \right) = \bigoplus_m \left(\sum_{y \in Z_d^m} \left(\sum_{x \in Z_d^m} (1 \hat{\otimes} T(y, x)) a_x[x] \right) [y] \right)$$

for all $\sum_{x \in Z_d^m} a_x[x] \in E^m$. Since

$$\|N_{g_t}\| = \sup_{x \in Z_d^m} \left\| \beta(x)(g_t) \right\| \leq \|g\|,$$

we have that

$$\left\| \beta_t(g \hat{\otimes} T) \right\| \leq \|g\| \|T\|.$$

It follows that β_t extends to a bounded linear map from the algebraic tensor product $\mathcal{S} \hat{\otimes} C_u^*((P_d(G))_{m \in \mathbb{N}})$ to $C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$. Consequently, the family of maps $(\beta_t)_{t \geq 1}$ extends to an asymptotic morphism from the maximal tensor product $\mathcal{S} \hat{\otimes}_{\max} C_u^*((P_d(G))_{m \in \mathbb{N}})$ to $C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$. Since \mathcal{S} is nuclear, we conclude that $(\beta_t)_{t \geq 1}$ extends to an asymptotic morphism from $\mathcal{S} \hat{\otimes} C_u^*((P_d(G))_{m \in \mathbb{N}})$ to $C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$. The case for the localization algebras is similar. The proof is complete. \square

Note that the asymptotic morphisms

$$\beta : \mathcal{S} \hat{\otimes} C_u^*((P_d(G))_{m \in \mathbb{N}}) \rightsquigarrow C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}),$$

$$\beta_{u,L} : \mathcal{S} \hat{\otimes} C_{u,L}^*((P_d(G))_{m \in \mathbb{N}}) \rightsquigarrow C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})$$

induce homomorphisms on K -theory:

$$\beta_* : K_*(\mathcal{S} \hat{\otimes} C_u^*((P_d(G))_{m \in \mathbb{N}})) \rightarrow K_*(C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}}))$$

$$(\beta_{u,L})_* : K_*(\mathcal{S} \hat{\otimes} C_{u,L}^*((P_d(G))_{m \in \mathbb{N}})) \rightarrow K_*(C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})).$$

Finally we are ready to complete the proof of the main result of this paper.

Proof of Theorem 1.1: Consider the following commutative diagram:

$$\begin{array}{ccc} \lim_{d \rightarrow \infty} K_*(\mathcal{S} \hat{\otimes} C_{u,L}^*((P_d(G))_{m \in \mathbb{N}})) & \xrightarrow{e_*} & \lim_{d \rightarrow \infty} K_*(\mathcal{S} \hat{\otimes} C_u^*((P_d(G))_{m \in \mathbb{N}})) \\ (\beta_{u,L})_* \downarrow \cong & & \beta_* \downarrow \\ \lim_{d \rightarrow \infty} K_*(C_{u,L}^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})) & \xrightarrow[\cong]{e_*} & \lim_{d \rightarrow \infty} K_*(C_u^*((P_d(G_m), \mathcal{A}(V_m))_{m \in \mathbb{N}})). \end{array}$$

We first show that $(\beta_{u,L})_*$ is an isomorphism for any $d \geq 0$. Note that the sequence of groups $(G_m)_{m \in \mathbb{N}}$ have uniform bounded geometry and the K -theory of all the above localization algebras is invariant under the strong Lipschitz homotopy equivalence. Using Mayer-Vietoris sequence argument for uniform localization algebras and uniform twisted localization algebras and induction on the dimension of skeletons of $P_d(G_m)$ for all m , the general case can be reduced to the 0-dimensional case, namely, if $D_m \subset P_d(G_m)$ for each $m \in \mathbb{N}$ is a δ -separated subset (meaning that $d_{P_d(G_m)}(x, y) > \delta$ if $x \neq y \in D_m$) for some $\delta > 0$, then

$$(\beta_{u,L})_* : K_*(\mathcal{S} \hat{\otimes} C_{u,L}^*((D_m)_{m \in \mathbb{N}})) \rightarrow K_*(C_{u,L}^*((D_m, \mathcal{A}(V_m))_{m \in \mathbb{N}}))$$

is an isomorphism. It is clear that this can be reformulated as

$$(\beta_{u,L})_* : \prod_{\substack{x \in D_m \\ m \in \mathbb{N}}}^u K_*(\mathcal{S} \hat{\otimes} C_{u,L}^*({x})) \rightarrow \prod_{\substack{x \in D_m \\ m \in \mathbb{N}}}^u K_*(C_{u,L}^*({x}, \mathcal{A}(V_m))),$$

which is an isomorphism by the celebrated infinite dimensional Bott periodicity theorem of Higson, Kasparov and Trout [17]. It follows that the Bott map $(\beta_{u,L})_*$ in the commutative diagram above is an isomorphism even before the inductive limit in $d \rightarrow \infty$ is taken. Since both $C_u^*((P_d(G))_{m \in \mathbb{N}})$ and

$C_{u,L}^*((P_d(G))_{m \in \mathbb{N}})$ are trivially graded, by taking inductive limit and recalling Theorem 4.1, we conclude that the evaluation map on K -theory

$$e_* : \lim_{d \rightarrow \infty} K_* \left(C_{u,L}^*((P_d(G))_{m \in \mathbb{N}}) \right) \rightarrow \lim_{d \rightarrow \infty} K_* \left(C_u^*((P_d(G))_{m \in \mathbb{N}}) \right)$$

is injective. Now Theorem 1.1 follows from Proposition 2.10. This completes the whole proof. \square

Acknowledgements

We would like to thank Jintao Deng and Guoliang Yu for valuable suggestions and helpful discussions. We also would like to thank Alexander Engel for helpful comments.

References

- [1] G. Arzhantseva, and D. Osajda. Graphical small cancellation groups with the Haagerup property. (2014) arXiv: 1404.6807.
- [2] G. Arzhantseva, and R. Tessera. Relative expanders. *Geom. Funct. Anal.* 25 no. 2 (2015) 317–341.
- [3] G. Arzhantseva, and R. Tessera. Admitting a coarse embedding is not preserved under group extensions. *Int. Math. Res. Not. IMRN* 2019, no. 20, 6480–6498.
- [4] P. Baum, A. Connes and N. Higson. Classifying space for proper actions and K-theory of group C^* -algebras. *C*-Algebras: 1943–1993*, San Antonio, TX, 1993, in: *Contemp. Math.*, vol. 167, Amer. Math. Soc., Providence, RI, 1994, pp. 240–291.
- [5] B. M. Braga, Y. C. Chung, and K. Li. Coarse Baum-Connes conjecture and rigidity for Roe algebras. *J. Funct. Anal.* 279 no. 9, (2020) 108728.
- [6] X. Chen, Q. Wang and G. Yu. The coarse Novikov conjecture and Banach spaces with Property (H) . *J. Funct. Anal.*, 268 (2015) 2754–2786.
- [7] M. Dadarlat, and E. Guentner. Construncions preserving Hilbert space uniform embeddability of discrete groups. *Trans. Amer. Math. Soc.* 355, no. 8 (2003): 3253–75.
- [8] T. Delabie, and A. Khukhro. Box spaces of the free group that neither contain expanders nor embed into a Hilbert space. *Adv. Math.* 336 (2018) 70–96.
- [9] J. Deng. The Novikov conjecture and extensions of coarsely embeddable groups. *J. Noncommutat. Geom.* (2020), forthcoming; to appear in JNCG.
- [10] J. Deng, Q. Wang and G. Yu. The coarse Baum–Connes conjecture for certain group extensions and relative expanders. arXiv:2102.10617.

- [11] G. Gong, Q. Wang, G. Yu. Geometrization of the strong Novikov conjecture for residually finite groups. *J. Reine Angew. Math.* 621 (2008), 159–189.
- [12] S. Gong, J. Wu, and G. Yu, The Novikov conjecture, the group of volume preserving diffeomorphisms and Hilbert-Hadamard spaces. to appear in *Geom. Funct. Anal.* 2021.
- [13] M. Gromov, *Asymptotic invariants for infinite groups*, London Math. Soc. Lecture Notes Ser., vol. 182, Academic Press, New York, 1993, pp. 1–295.
- [14] M. Gromov. Random walks in random groups. *Geom. Funct. Anal.* 13(1) (2003) 73–146.
- [15] E. Guentner, J. Kaminker. Geometric and analytic properties of groups. In: *Noncommutative Geometry*, edited by S. Doplicher and R. Longo. Lecture Notes Math. 1831 (2004): 253-62.
- [16] N. Higson, J. Roe. On the coarse Baum–Connes conjecture, in: S. Ferry, A. Ranicki, J. Rosenberg (Eds.), *Proc. 1993 Oberwolfach Conference in the Novikov Conjecture*. London Math. Soc. Lecture Notes Ser., vol. 227, Cambridge Univ. Press, Cambridge, 1995, pp. 227–254.
- [17] N. Higson, G. Kasparov and J. Trout. A Bott periodicity theorem for infinite dimensional Euclidean space. *Adv. Math.*, 135 (1998), no. 1, 1–40.
- [18] G. Kasparov. Topological invariants of elliptic operators I: K -homology. *Mathematics of the USSR*, 751–792, 1975.
- [19] G. Kasparov, G. Yu. The coarse geometric Novikov conjecture and uniform convexity. *Adv. Math.* 206 (1) (2006) 1–56.
- [20] G. Kasparov, G. Yu. The Novikov conjecture and geometry of Banach spaces. *Geom. Topol.* 16 (2012) 101–999.
- [21] P. Nowak, and G. Yu. *Large space geometry*. Zürich: European Mathematical Society (EMS), 2012.
- [22] D. Osajda. Small cancellation labellings of some infinite graphs and applications. *Acta Math.* 225 (2020), 159–191.
- [23] H. Oyono-Oyono. Baum–Connes conjecture and extensions. *J. Reine Angew. Math.* 532 (2001) 133–149.
- [24] H. Oyono-Oyono, and G. Yu. K -theory for the maximal Roe algebras of certain expanders. *J. Funct. Anal.* (10) 257 (2009), 3239–3292.
- [25] J. Roe. Coarse cohomology and index theory on complete Riemannian manifolds. *Mem. Amer. Math. Soc.* 104 (497)(1993), x+90 pp.
- [26] J. Roe. *Index Theory, Coarse Geometry, and Topology of Manifolds*. CBMS Conf. Proceedings, vol. 90, Amer. Math. Soc., Providence, RI, 1996.
- [27] L. Shan and Q. Wang. The coarse geometric Novikov conjecture for subspaces of non–positively curved manifolds. *J. Funct. Anal.*, 248 (2007), no. 2, 448–471.

- [28] R. Willett and G. Yu. Higher index theory. Cambridge University Press, 2020.
- [29] G. Yu. Coarse Baum–Connes conjecture. *K-Theory*, 9 (3) (1995) 199–221.
- [30] G. Yu. Baum–Connes Conjecture and Coarse Geometry. *K-Theory*, 9 (1995) 223–231.
- [31] G. Yu. Localization algebras and the coarse Baum–Connes conjecture. *K-Theory*, 11 (1997) 307–318.
- [32] G. Yu, The Novikov conjecture for groups with finite asymptotic dimension. *Ann. of Math. (2)* 147 (1998), no. 2, 325–355.
- [33] G. Yu. The coarse Baum–Connes conjecture for spaces which admit a uniform embedding into Hilbert space. *Invent. Math.*, 139 (2000) 201–240.
- [34] G. Yu. Higher index theory of elliptic operators and geometry of groups. *Proceedings of International Congress of Mathematicians, Madrid, 2006, vol. II*, pp. 1623–1639.

Qin Wang

Research Center for Operator Algebras, and Shanghai Key Laboratory of Pure Mathematics and Mathematical Practice, School of Mathematical Sciences, East China Normal University, Shanghai, 200241, P. R. China. E-mail: qwang@math.ecnu.edu.cn

Yazhou Zhang

Research Center for Operator Algebras, School of Mathematical Sciences, East China Normal University, Shanghai, 200241, P. R. China. E-mail: 52185500010@stu.ecnu.edu.cn