

THE RESOLUTION OF EUCLIDEAN MASSLESS FIELD OPERATORS OF HIGHER SPINS ON \mathbb{R}^6 AND THE L^2 METHOD

QIANQIAN KANG, WEI WANG, AND YUCHEN ZHANG

ABSTRACT. The resolution of 4-dimensional massless field operators of higher spins was constructed by Eastwood-Penrose-Wells by using the twistor method. Recently physicists are interested in 6-dimensional physics including the massless field operators of higher spins on Lorentzian space $\mathbb{R}^{5,1}$. Its Euclidean version \mathcal{D}_0 and their function theory are discussed in [14]. In this paper, we construct an exact sequence of Hilbert spaces as weighted L^2 spaces resolving \mathcal{D}_0 :

$$L^2_\varphi(\mathbb{R}^6, \mathcal{V}_0) \xrightarrow{\mathcal{D}_0} L^2_\varphi(\mathbb{R}^6, \mathcal{V}_1) \xrightarrow{\mathcal{D}_1} L^2_\varphi(\mathbb{R}^6, \mathcal{V}_2) \xrightarrow{\mathcal{D}_2} L^2_\varphi(\mathbb{R}^6, \mathcal{V}_3) \longrightarrow 0,$$

with suitable operators \mathcal{D}_l and vector spaces \mathcal{V}_l . Namely, we can solve $\mathcal{D}_l u = f$ in $L^2_\varphi(\mathbb{R}^6, \mathcal{V}_l)$ when $\mathcal{D}_{l+1} f = 0$ for $f \in L^2_\varphi(\mathbb{R}^6, \mathcal{V}_{l+1})$. This is proved by using the L^2 method in the theory of several complex variables, which is a general framework to solve overdetermined PDEs under the compatibility condition. To apply this method here, it is necessary to consider weighted L^2 spaces, an advantage of which is that any polynomial is L^2_φ integrable. As a corollary, we prove that

$$P(\mathbb{R}^6, \mathcal{V}_0) \xrightarrow{\mathcal{D}_0} P(\mathbb{R}^6, \mathcal{V}_1) \xrightarrow{\mathcal{D}_1} P(\mathbb{R}^6, \mathcal{V}_2) \xrightarrow{\mathcal{D}_2} P(\mathbb{R}^6, \mathcal{V}_3) \longrightarrow 0$$

is a resolution, where $P(\mathbb{R}^6, \mathcal{V}_l)$ is the space of all \mathcal{V}_l -valued polynomials. This provides an analytic way to construct a resolution of a differential operator acting on vector valued polynomials.

1. INTRODUCTION

The resolution of massless field operators of higher spins over the complexified Minkowski space \mathbb{C}^4 was constructed by Eastwood-Penrose-Wells [7] by using twistor method. The Euclidean version of massless field operator of spin $k/2$ is also called k -Cauchy-Fueter operator (cf. [4] [24] and references therein). Recently physicists are interested in 6-dimensional physics including the massless field operators of higher spins on Lorentzian space $\mathbb{R}^{5,1}$ (cf. [16, 19] and references there in). The Euclidean version of these operators are

$$\mathcal{D}_0 : C^\infty(\mathbb{R}^6, \odot^k \mathbb{C}^4) \longrightarrow C^\infty(\mathbb{R}^6, \mathbb{C}^4 \otimes \odot^{k-1} \mathbb{C}^4),$$

$k = 1, 2, \dots$, where $\odot^p \mathbb{C}^4$ is p -th symmetric power of \mathbb{C}^4 . A $\odot^k \mathbb{C}^4$ -valued distribution f is called k -monogenic if it satisfies $\mathcal{D}_0 f = 0$. In [14], we proved various properties for k -monogenic functions, e.g. the existence of infinite number of k -monogenic polynomials. In order to study k -monogenic functions, we need to solve the nonhomogeneous equation $\mathcal{D}_0 u = f$, which is overdetermined for $k > 1$. So we need to find the compatibility condition for solvability, and more generally a resolution of \mathcal{D}_0 . Motivated by 4-dimensional and the quaternionic cases (c.f. [1, 2, 3, 22] and references therein), a natural candidate of the resolution is

$$(1.1) \quad C^\infty(\mathbb{R}^6, V_0) \xrightarrow{\mathcal{D}_0} C^\infty(\mathbb{R}^6, V_1) \xrightarrow{\mathcal{D}_1} C^\infty(\mathbb{R}^6, V_2) \xrightarrow{\mathcal{D}_2} C^\infty(\mathbb{R}^6, V_3) \xrightarrow{\mathcal{D}_3} C^\infty(\mathbb{R}^6, V_4) \longrightarrow 0$$

Key words and phrases. resolution; Euclidean massless field operator of high spins; the L^2 method; overdetermined PDEs; the compatibility condition; differential complexes.

The first author is partially supported by National Nature Science Foundation in China (No. 11801523) and the foundation of Zhejiang International Studies University (No. BD2019B9). The second and third authors are partially supported by National Nature Science Foundation in China (No. 11971425).

with

$$(1.2) \quad V_l := \odot^{k-l}\mathbb{C}^4 \otimes \wedge^l \mathbb{C}^4,$$

when $k \geq 4$. But it is already known [14, Section 1] that the image of \mathcal{D}_0 consists of functions only valued in a subspace of V_1 , the kernel of the *contraction* given by

$$(1.3) \quad \mathcal{C} : \odot^p \mathbb{C}^4 \otimes \wedge^q \mathbb{C}^4 \rightarrow \odot^{p-1} \mathbb{C}^4 \otimes \wedge^{q-1} \mathbb{C}^4$$

with $p = k - 1, q = 1$. Denote

$$(1.4) \quad \mathcal{V}_l := \ker \mathcal{C} |_{V_l}.$$

Then $\mathcal{V}_4 = \{0\}$ automatically. Denote by \mathcal{D}_l the restriction of \mathcal{D}_l in (1.1) to $C^\infty(\mathbb{R}^6, \mathcal{V}_l)$. We construct the following differential complex:

$$(1.5) \quad 0 \longrightarrow C^\infty(\mathbb{R}^6, \mathcal{V}_0) \xrightarrow{\mathcal{D}_0} C^\infty(\mathbb{R}^6, \mathcal{V}_1) \xrightarrow{\mathcal{D}_1} C^\infty(\mathbb{R}^6, \mathcal{V}_2) \xrightarrow{\mathcal{D}_2} C^\infty(\mathbb{R}^6, \mathcal{V}_3) \longrightarrow 0,$$

and call it *k-monogenic complex*, $k = 4, 5, \dots$. Note that \mathbb{C}^4 is the spin representation of $\mathfrak{so}(6, \mathbb{C})$ and \mathcal{V}_l as the contraction of V_l is an irreducible representation of $\mathfrak{so}(6, \mathbb{C})$ (cf. [8]).

The L^2 method is a powerful method to solve $\bar{\partial}$ -equation in the theory of several complex variables (cf. e.g. [5, 10, 11]). In fact, it is a general framework to solve overdetermined PDEs under the compatibility condition, which is also given by a system of PDEs. The main difficulty to use this method is to prove the corresponding L^2 estimate. It was applied to the k -Cauchy-Fueter complex over \mathbb{R}^{4n} in [23] and also the Neumann problem associated to the k -Cauchy-Fueter complex on k -pseudoconvex domains in \mathbb{R}^4 [24]. The latter case is restricted to dimension 4 because only over \mathbb{R}^4 the corresponding L^2 estimates was proved.

In this paper we consider the weighted L^2 estimate of the k -monogenic complex as in [23]. We define an inner product $\langle \cdot, \cdot \rangle$ on V_l and \mathcal{V}_l induced from $\otimes^k \mathbb{C}^4$. Let $L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l)$ be the Hilbert space of \mathcal{V}_l -valued L_φ^2 -integrable functions on \mathbb{R}^6 with weighted inner product

$$(1.6) \quad \langle f, h \rangle_\varphi := \int_{\mathbb{R}^6} \langle f(x), h(x) \rangle e^{-\varphi} dx,$$

with weight $\varphi = |x|^2$. Denote weighted norm $\|f\|_\varphi := \langle f, f \rangle_\varphi^{\frac{1}{2}}$.

$Dom(\mathcal{D}_l)$ consists of $f \in L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l)$ such that $\mathcal{D}_l f = u$ in the weak sense for some $u \in L_\varphi^2(\mathbb{R}^6, \mathcal{V}_{l+1})$, i.e.

$$(1.7) \quad \langle u, g \rangle_\varphi = \langle f, \Theta_l g \rangle_\varphi$$

for any $C_0^\infty(\mathbb{R}^6, \mathcal{V}_{l+1})$, where Θ_l is the formal adjoint of \mathcal{D}_l . The differential operator \mathcal{D}_l defines a linear, closed, densely defined operator from $L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l)$ to $L_\varphi^2(\mathbb{R}^6, \mathcal{V}_{l+1})$, which we also denote by \mathcal{D}_l . Denote by \mathcal{D}_l^* the adjoint operator of \mathcal{D}_l between Hilbert spaces $L_\varphi^2(\mathbb{R}^6, \mathcal{V}_{l+1})$ and $L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l)$. The sequence

$$(1.8) \quad L_\varphi^2(\mathbb{R}^6, \mathcal{V}_0) \xrightarrow{\mathcal{D}_0} L_\varphi^2(\mathbb{R}^6, \mathcal{V}_1) \xrightarrow{\mathcal{D}_1} L_\varphi^2(\mathbb{R}^6, \mathcal{V}_2) \xrightarrow{\mathcal{D}_2} L_\varphi^2(\mathbb{R}^6, \mathcal{V}_3) \longrightarrow 0,$$

is a complex of Hilbert spaces, i.e., for any $u \in Dom(\mathcal{D}_l)$ and $\mathcal{D}_l u \in Dom(\mathcal{D}_{l+1})$, we have $\mathcal{D}_{l+1} \mathcal{D}_l u = 0$. To find solution to the equation

$$(1.9) \quad \mathcal{D}_l u = f,$$

for $f \in L_\varphi^2(\mathbb{R}^6, \mathcal{V}_{l+1})$ satisfying

$$(1.10) \quad \mathcal{D}_{l+1} f = 0,$$

we consider the *associated Hodge Laplacian operator* $\square_l : L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l) \longrightarrow L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l)$ given by

$$(1.11) \quad \square_l := \mathcal{D}_{l-1} \mathcal{D}_{l-1}^* + \mathcal{D}_l^* \mathcal{D}_l, \quad l = 1, 2, \quad \text{and} \quad \square_3 := \mathcal{D}_2 \mathcal{D}_2^*.$$

Theorem 1.1. *Suppose that $\varphi(x) = |x|^2$ and $k = 6, 7, \dots$. There exists a constant $C > 0$ only depending on k such that*

(1) \square_l has a bounded self-adjoint inverse N_l such that

$$(1.12) \quad \|N_l f\|_\varphi \leq C \|f\|_\varphi \quad \text{for any } f \in L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l).$$

(2) $\mathcal{D}_l^* N_{l+1} f$ is the canonical solution to the nonhomogeneous equation (1.9)-(1.10), i.e.

$$\mathcal{D}_l(\mathcal{D}_l^* N_{l+1} f) = f,$$

if $\mathcal{D}_{l+1} f = 0$, and $\mathcal{D}_l^* N_{l+1} f$ orthogonal to $\ker \mathcal{D}_l$. Moreover,

$$(1.13) \quad \|\mathcal{D}_l^* N_{l+1} f\|_\varphi \leq C \|f\|_\varphi, \quad \|\mathcal{D}_{l+1} N_{l+1} f\|_\varphi \leq C \|f\|_\varphi.$$

Corollary 1.1. *When $k = 6, 7, \dots$, the sequence (1.8) is exact.*

The key step to prove Theorem 1.1 is to establish the following weighted L^2 estimate.

Theorem 1.2. *Suppose that $\varphi(x) = |x|^2$ and $k = 6, 7, \dots$. There exists a constant $C > 0$ only depending on k such that*

$$(1.14) \quad \|f\|_\varphi^2 \leq C \left(\|\mathcal{D}_l f\|_\varphi^2 + \|\mathcal{D}_{l-1}^* f\|_\varphi^2 \right),$$

for any $f \in \text{Dom}(\mathcal{D}_l) \cap \text{Dom}(\mathcal{D}_{l-1}^*)$, $l = 1, 2, 3$.

The restriction $k \geq 6$ is a technique requirement since we only prove the estimate (1.14) in this case. The massless field operators of higher spins on any dimensional Euclidean space was introduced by Souček earlier [20, 21]. We only consider 6-dimensional case here because we can use spin indices based on $\mathfrak{so}(6, \mathbb{C}) \cong \mathfrak{sl}(4, \mathbb{C})$, as two-component notation in dimension 4 based on $\mathfrak{so}(4, \mathbb{C}) \cong \mathfrak{sl}(2, \mathbb{C}) \oplus \mathfrak{sl}(2, \mathbb{C})$.

An advantage to consider weighted L^2 space is that any polynomial on \mathbb{R}^6 is L_φ^2 integrable. This allow us to deduce a resolution of the operator \mathcal{D}_0 on $P(\mathbb{R}^6, \mathcal{V}_0)$, the module of \mathcal{V}_0 -valued polynomials over \mathbb{R}^6 .

Theorem 1.3. *When $k = 6, 7, \dots$, the sequence*

$$(1.15) \quad P(\mathbb{R}^6, \mathcal{V}_0) \xrightarrow{\mathcal{D}_0} P(\mathbb{R}^6, \mathcal{V}_1) \xrightarrow{\mathcal{D}_1} P(\mathbb{R}^6, \mathcal{V}_2) \xrightarrow{\mathcal{D}_2} P(\mathbb{R}^6, \mathcal{V}_3) \longrightarrow 0,$$

is exact.

To prove this theorem, we need the following proposition.

Proposition 1.1. *When $k = 4, 5, \dots$, the complex (1.5) is an elliptic complex.*

Compared to the quaternionic case [23], the main difficulty comes from the algebraic complexity when passing from vector space $V_l = \odot^{k-l} \mathbb{C}^4 \otimes \wedge^l \mathbb{C}^4$ to its contraction subspace \mathcal{V}_l , although only linear algebra is used to overcome it. In Section 2, we give some basic propositions on symmetrization and antisymmetrization to handle functions valued in the subspace \mathcal{V}_l . To write down the formal adjoint operator \mathcal{D}_l^* explicitly, we introduce the orthogonal projection \mathcal{P}_l from V_l to \mathcal{V}_l^\perp . In Section 3, we give the expression of operator \mathcal{D}_l and formal adjoint operators \mathcal{D}_l^* . Then we prove the L^2 estimate (1.14). In Section 4, we give the canonical solution to the nonhomogeneous equations (1.9)-(1.10) by the general framework to solve nonhomogeneous overdetermined PDEs. As a corollary, we show the sequence (1.8) is exact. Since the \mathcal{V}_l -valued polynomials over \mathbb{R}^6 are L_φ^2 integrable, we show that (1.15) is also exact. In Section 5, we establish the ellipticity of the differential complex (1.5) by showing the exactness of its symbol sequence, based on which we show \square_l is an elliptic differential operator and then prove Theorem 1.3.

2. LINEAR ALGEBRA FOR SYMMETRIC AND EXTERIOR FORMS

2.1. Symmetrization and antisymmetrization. An element $\xi \in \otimes^t \mathbb{C}^4$ is a tuple $(\xi_{A_1 \dots A_t})$ with $\xi_{A_1 \dots A_t} \in \mathbb{C}$, where $A_1, \dots, A_t = 1, 2, 3, 4$. The *symmetric power* $\odot^k \mathbb{C}^4$ is a subspace of $\otimes^k \mathbb{C}^4$, whose element is a tuple $(\xi_{B_1 \dots B_k})$ such that $\xi_{B_1 \dots B_k}$ is invariant under permutations of subscripts. An element of $\odot^{k-l} \mathbb{C}^4 \otimes \wedge^l \mathbb{C}^4$ is denoted by $(\eta_{B_1 \dots B_{k-l}}^{A_1 \dots A_l})$, where $\eta_{B_1 \dots B_{k-l}}^{A_1 \dots A_l}$ is symmetric in B_1, \dots, B_{k-l} and antisymmetric in A_1, \dots, A_l . The norm $\|\xi\|$ for $\xi \in \odot^{k-l} \mathbb{C}^4 \otimes \wedge^l \mathbb{C}^4$ is the norm of ξ as an element of $\otimes^k \mathbb{C}^4$, i.e., $\|\xi\| = \sum_{A_1, \dots, B_1, \dots} \|\xi_{B_1 \dots B_{k-l}}^{A_1 \dots A_l}\|$.

Symmetrization and antisymmetrization of an element $\xi \in \otimes^t \mathbb{C}^4$ is given by

$$(2.1) \quad \xi_{(A_1 \dots A_t)} := \frac{1}{t!} \sum_{\sigma \in S_t} \xi_{A_{\sigma_1} \dots A_{\sigma_t}}, \quad \xi_{[A_1 \dots A_t]} := \frac{1}{t!} \sum_{\sigma \in S_t} \epsilon_{1 \dots t}^{\sigma_1 \dots \sigma_t} \xi_{A_{\sigma_1} \dots A_{\sigma_t}},$$

respectively, where S_t denotes the permutation group of t elements and $\epsilon_{1 \dots t}^{\sigma_1 \dots \sigma_t}$ is the sign of the permutation from $(1, \dots, t)$ to $(\sigma_1, \dots, \sigma_t)$. The symmetrization or antisymmetrization of $\xi \in \otimes^t \mathbb{C}^4$ is an element of $\odot^t \mathbb{C}^4$ or $\wedge^t \mathbb{C}^4$.

Lemma 2.1 (cf. [23]). (1) For any $\xi \in \odot^t \mathbb{C}^4$ and $\zeta \in \otimes^t \mathbb{C}^4$, we have

$$\sum_{B_1, \dots, B_t} \xi_{B_1 \dots B_t} \overline{\zeta_{B_1 \dots B_t}} = \sum_{B_1, \dots, B_t} \xi_{B_1 \dots B_t} \overline{\zeta_{(B_1 \dots B_t)}}.$$

(2) For any $\xi \in \wedge^t \mathbb{C}^4$ and $\eta \in \otimes^t \mathbb{C}^4$, we have

$$\sum_{B_1, \dots, B_t} \xi_{B_1 \dots B_t} \overline{\eta_{B_1 \dots B_t}} = \sum_{B_1, \dots, B_t} \xi_{B_1 \dots B_t} \overline{\eta_{[B_1 \dots B_t]}}.$$

Proof. By definition (2.1), we have

$$\begin{aligned} \sum_{B_1, \dots, B_t} \xi_{B_1 \dots B_t} \overline{\zeta_{(B_1 \dots B_t)}} &= \frac{1}{t!} \sum_{B_1, \dots, B_t} \sum_{\sigma \in S_t} \xi_{B_1 \dots B_t} \overline{\zeta_{B_{\sigma_1} \dots B_{\sigma_t}}} \\ &= \frac{1}{t!} \sum_{\sigma \in S_t} \sum_{B_1, \dots, B_t} \xi_{B_{\sigma_1} \dots B_{\sigma_t}} \overline{\zeta_{B_{\sigma_1} \dots B_{\sigma_t}}} = \sum_{B_1, \dots, B_t} \xi_{B_1 \dots B_t} \overline{\zeta_{B_1 \dots B_t}}, \end{aligned}$$

by $\xi \in \odot^t \mathbb{C}^4$ and relabeling indices. The proof of (2) is analogous to (1). \square

Lemma 2.2. (1) For $(\xi_{B_1 \dots B_t}) \in \otimes^t \mathbb{C}^4$ symmetric in $B_2 \dots B_t$, we have

$$\xi_{(B_1 \dots B_t)} = \frac{1}{t} \sum_{s=1}^t \xi_{B_s B_1 \dots \widehat{B}_s \dots},$$

where \widehat{B}_s means omitting B_s .

(2) For $(\xi_{B_1 \dots B_t}) \in \otimes^t \mathbb{C}^4$ antisymmetric in $B_2 \dots B_t$, we have

$$\xi_{[B_1 \dots B_t]} = \frac{1}{t} \left(\xi_{B_1 \dots B_t} - \sum_{s=2}^t \xi_{B_s B_2 \dots B_1 \dots} \right),$$

(3) For $\xi \in \otimes^t \mathbb{C}^4$, we have

$$\xi_{[B_1 \dots B_t]} = \xi_{[[B_1 \dots B_{t_1}] B_{t_1+1} \dots B_t]}.$$

Proof. (1) Its proof is similar to (2).

(2) By the definition of antisymmetrization (2.1), we have

$$(2.2) \quad \xi_{[B_1 \dots B_t]} = \frac{1}{t!} \sum_{\sigma \in S_t} \epsilon_{12 \dots t}^{\sigma_1 \dots \sigma_t} \xi_{B_{\sigma_1} \dots B_{\sigma_t}} = \frac{1}{t!} \sum_{s=1}^t \sum_{\sigma \in S_t, \sigma_1=s} \epsilon_{12 \dots t}^{s \sigma_2 \dots \sigma_t} \xi_{B_s B_{\sigma_2} \dots B_{\sigma_t}}.$$

Since f is antisymmetric in the last $t - 1$ indices, we get

$$\begin{aligned}\xi_{[B_1 \dots B_t]} &= \frac{(t-1)!}{t!} \xi_{B_1 \dots B_t} + \frac{1}{t!} \sum_{s=2}^t \sum_{\sigma \in S_t, \sigma_1=s} \epsilon_{12 \dots t}^{s \sigma_2 \dots \sigma_t} \epsilon_{s \sigma_2 \dots \sigma_s \dots \sigma_t}^{s 2 \dots 1 \dots t} \xi_{B_s B_2 \dots B_1 \dots} \\ &= \frac{1}{t} \xi_{B_1 \dots B_t} - \frac{1}{t} \sum_{s=2}^t \xi_{B_s B_2 \dots B_1 \dots}.\end{aligned}$$

This completes the proof of (2).

(3) Denote $\Xi_{B_1 \dots B_t} := \xi_{[B_1 \dots B_{t_1}] B_{t_1+1} \dots B_t}$. By definition of antisymmetrization, we have

$$\begin{aligned}(2.3) \quad \Xi_{[B_1 \dots B_t]} &= \frac{1}{t!} \sum_{\sigma \in S_t} \epsilon_{1 \dots t}^{\sigma_1 \dots \sigma_t} \Xi_{B_{\sigma_1} \dots B_{\sigma_t}} = \frac{1}{t! t_1!} \sum_{\tau \in S_{t_1}} \sum_{\sigma \in S_t} \epsilon_{1 \dots t}^{\tau \sigma_1 \dots \tau \sigma_{t_1} \dots \sigma_t} \xi_{B_{\tau \sigma_1} \dots B_{\tau \sigma_{t_1}} \dots B_{\sigma_t}} \\ &= \frac{1}{t! t_1!} \sum_{\tau \in S_{t_1}} \sum_{\kappa \in S_t} \epsilon_{1 \dots t}^{\kappa_1 \dots \kappa_t} \xi_{B_{\kappa_1} \dots B_{\kappa_t}} = \xi_{[B_1 \dots B_t]},\end{aligned}$$

by relabeling indices and permutations in the third identity. \square

2.2. The orthogonal projection $\mathcal{P} : V_l \rightarrow \mathcal{V}_l^\perp$. The contraction \mathcal{C} (1.3) given by

$$(2.4) \quad \mathcal{C}(f)_{B_1 \dots B_{p-1}}^{A_1 \dots A_{q-1}} = \sum_{C=1}^4 f_{B_1 \dots B_{p-1} C}^{C A_1 \dots A_{q-1}},$$

satisfies

$$(2.5) \quad \mathcal{C} \circ \mathcal{C} f = 0.$$

This is because for any fixed $A_1, \dots, A_{q-2}, B_1, \dots, B_{p-2}$,

$$(\mathcal{C} \circ \mathcal{C} f)_{B_1 \dots B_{p-2}}^{A_1 \dots A_{q-2}} = \sum_A \mathcal{C}(f)_{B_1 \dots B_{p-2} A}^{A A_1 \dots A_{q-2}} = \sum_{A, C} f_{B_1 \dots B_{p-2} A C}^{C A A_1 \dots A_{q-2}} = 0,$$

by f symmetric in subscripts A, C and antisymmetric in superscripts A, C .

Let \mathcal{V}_l^\perp be the orthogonal complement of \mathcal{V}_l in V_l . Now we construct a linear transformation \mathcal{P}_l from V_l to \mathcal{V}_l^\perp ($l = 1, 2$)

$$(2.6) \quad \begin{aligned}\mathcal{P}_1(f)_{B_1 B_2 \dots B_{k-1}}^A &:= \frac{k-1}{k+2} \delta_{(B_1}^A \mathcal{C}(f)_{B_2 \dots B_{k-1})}, & f \in V_1, \\ \mathcal{P}_2(f)_{B_1 \dots B_{k-2}}^{A_1 A_2} &:= \frac{2(k-2)}{k} \delta_{(B_1}^{[A_1} \mathcal{C}(f)_{B_2 \dots B_{k-2})}^{A_2]}, & f \in V_2.\end{aligned}$$

Proposition 2.1. \mathcal{P}_l is an orthogonal projection from V_l to \mathcal{V}_l^\perp , $l = 1, 2$.

Proof. We only prove the case $l = 2$ since it is similar for the case $l = 1$. Note that

$$(2.7) \quad \mathcal{C}(\mathcal{P}_2 f) = \mathcal{C}(f)$$

for any $f \in V_2 = \odot^{k-2} \mathbb{C}^4 \otimes \wedge^2 \mathbb{C}^4$, since $\mathcal{C}(\mathcal{P}_2 f)_{B_1 \dots B_{k-3}}^{A_2} = \sum_{A_1} \mathcal{P}_2(f)_{B_1 \dots B_{k-3} A_1}^{A_1 A_2}$ equals to

$$\begin{aligned}& \frac{1}{k} \sum_{A_1} \left(\delta_{B_1}^{A_1} \mathcal{C}(f)_{B_2 \dots B_{k-3} A_1}^{A_2} + \dots + \delta_{B_{k-3}}^{A_1} \mathcal{C}(f)_{B_1 \dots B_{k-4} A_1}^{A_2} + \delta_{A_1}^{A_1} \mathcal{C}(f)_{B_1 \dots B_{k-3}}^{A_2} \right. \\ & \quad \left. - \delta_{B_1}^{A_2} \mathcal{C}(f)_{B_2 \dots B_{k-3} A_1}^{A_1} - \dots - \delta_{B_{k-3}}^{A_2} \mathcal{C}(f)_{B_1 \dots B_{k-4} A_1}^{A_1} - \delta_{A_1}^{A_2} \mathcal{C}(f)_{B_1 \dots B_{k-3}}^{A_1} \right) \\ &= \mathcal{C}(f)_{B_1 \dots B_{k-3}}^{A_2},\end{aligned}$$

by (2.4), (2.5), definition (2.6) and Lemma 2.2 (1). Then \mathcal{P}_2 is a projection since

$$\mathcal{P}_2(\mathcal{P}_2 f)_{B_1 \dots B_{k-2}}^{A_1 A_2} = \frac{2(k-2)}{k} \delta_{(B_1)_{B_2 \dots B_{k-2}}}^{[A_1] (\mathcal{C} \mathcal{P}_2 f)^{A_2}} = \frac{2(k-2)}{k} \delta_{(B_1)_{B_2 \dots B_{k-2}}}^{[A_1] \mathcal{C}(f)^{A_2}} = \mathcal{P}_2(f)_{B_1 \dots B_{k-2}}^{A_1 A_2},$$

by (2.7). For any $h \in \mathcal{V}_2$, we have

$$\begin{aligned} k \langle \mathcal{P}_2 f, h \rangle &= \sum_{A_1, A_2, B_1, \dots, B_{k-2}} \left\langle \sum_{s=1}^{k-2} \left(\delta_{B_s}^{A_1} \mathcal{C}(f)_{B_1 \dots \widehat{B}_s \dots B_{k-2}}^{A_2} - \delta_{B_s}^{A_2} \mathcal{C}(f)_{B_1 \dots \widehat{B}_s \dots B_{k-2}}^{A_1} \right), h_{B_1 \dots B_{k-2}}^{A_1 A_2} \right\rangle \\ &= \sum_{s=1}^{k-2} \left\{ \sum_{A_2, B_1, \dots, \widehat{B}_s, \dots} \left\langle \mathcal{C}(f)_{B_1 \dots \widehat{B}_s \dots B_{k-2}}^{A_2}, \sum_{B_s} h_{B_1 \dots B_{k-2}}^{B_s A_2} \right\rangle \right. \\ &\quad \left. - \sum_{A_1, B_1, \dots, \widehat{B}_s, \dots} \left\langle \mathcal{C}(f)_{B_1 \dots \widehat{B}_s \dots B_{k-2}}^{A_1}, \sum_{B_s} h_{B_1 \dots B_{k-2}}^{A_1 B_s} \right\rangle \right\} = 0, \end{aligned}$$

by definition (2.6) and $\mathcal{C}h = 0$. Hence, $\mathcal{P}_2 f \in \mathcal{V}_2^\perp$, and so it is a projection from V_2 to \mathcal{V}_2^\perp .

For any $f \in \ker \mathcal{P}_2$, we know $f \in \mathcal{V}_2$ since we have $\mathcal{C}(f) = \mathcal{C}(\mathcal{P}_2 f) = 0$. On the other hand, for any $f \in \mathcal{V}_2$, we know $\mathcal{P}_2 f = 0$ by definition (2.6). Then $f \in \ker \mathcal{P}_2$ if and only if $f \in \mathcal{V}_2$. Hence \mathcal{P}_2 is an orthogonal projection from V_2 to \mathcal{V}_2^\perp . \square

We also need to know the norm of $\mathcal{P}_l(\xi)$ for $\xi \in V_l$ in the proof of the L^2 estimate. We have

Proposition 2.2.

$$(2.8) \quad \begin{aligned} \|\mathcal{P}_1(\xi)\|^2 &= \frac{k-1}{k+2} \|\mathcal{C}(\xi)\|^2, & \text{for } \xi \in V_1, \\ \|\mathcal{P}_2(\xi)\|^2 &= \frac{2(k-2)}{k} \|\mathcal{C}(\xi)\|^2, & \text{for } \xi \in V_2. \end{aligned}$$

Proof. For $\xi \in V_1$, we have

$$(2.9) \quad \begin{aligned} (k+2)^2 \|\mathcal{P}_1(\xi)\|^2 &= (k+2)^2 \sum_{A, B_1, \dots, B_{k-1}} \left\| \mathcal{P}_1(\xi)_{B_1 \dots B_{k-1}}^A \right\|^2 \\ &= \sum_{A, B_1, \dots, B_{k-1}} \sum_{j, l=1}^{k-1} \left\langle \delta_{B_j}^A (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_{k-1}}, \delta_{B_l}^A (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_l \dots B_{k-1}} \right\rangle \\ &= \sum_{A, B_1, \dots, B_{k-1}} \left(\sum_{j=1}^{k-1} \left\langle \delta_{B_j}^A (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_{k-1}}, \delta_{B_j}^A (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_{k-1}} \right\rangle \right. \\ &\quad \left. + \sum_{j \neq l} \left\langle \delta_{B_j}^A (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_{k-1}}, \delta_{B_l}^A (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_l \dots B_{k-1}} \right\rangle \right) := S_1 + S_2, \end{aligned}$$

by (2.6). But we have

$$(2.10) \quad S_1 = 4(k-1) \sum_{B_1, \dots, B_{k-2}} \left\| (\mathcal{C}\xi)_{B_1 \dots B_{k-2}} \right\|^2 = 4(k-1) \|\mathcal{C}(\xi)\|^2,$$

by relabeling indices and

$$\begin{aligned}
(2.11) \quad S_2 &= \sum_{j \neq l} \sum_{\dots, \widehat{B}_j, \dots, \widehat{B}_l, \dots} \sum_A \left\langle (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots A \dots B_{k-1}}, (\mathcal{C}\xi)_{B_1 \dots A \dots \widehat{B}_l \dots B_{k-1}} \right\rangle \\
&= (k-1)(k-2) \sum_{B_1, \dots, B_{k-2}} \left\| (\mathcal{C}\xi)_{B_1 \dots B_{k-2}} \right\|^2 = (k-1)(k-2) \|\mathcal{C}(\xi)\|^2,
\end{aligned}$$

by relabeling indices. Then the sum of (2.10) and (2.11) gives us the first identity of (2.8).

For $\xi \in V_2$, we rewrite $k^2 \|\mathcal{P}_2(\xi)\|^2$ as in (2.9)

$$\begin{aligned}
(2.12) \quad k^2 \|\mathcal{P}_2(\xi)\|^2 &= 4 \sum_{\substack{A_1, A_2, \\ B_1, \dots, B_{k-2}}} \left\langle \sum_j \delta_{B_j}^{[A_1} (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_{k-2}}^{A_2]}, \sum_l \delta_{B_l}^{[A_1} (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_l \dots B_{k-2}}^{A_2]} \right\rangle \\
&= 4 \sum_j \sum_{\substack{A_1, A_2, \\ B_1, \dots, B_{k-2}}} \left\langle \delta_{B_j}^{[A_1} (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_{k-2}}^{A_2]}, \delta_{B_j}^{[A_1} (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_{k-2}}^{A_2]} \right\rangle \\
&\quad + 4 \sum_{j \neq l} \sum_{\substack{A_1, A_2, \\ B_1, \dots, B_{k-2}}} \left\langle \delta_{B_j}^{[A_1} (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_{k-2}}^{A_2]}, \delta_{B_l}^{[A_1} (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_l \dots B_{k-2}}^{A_2]} \right\rangle := S_1 + S_2,
\end{aligned}$$

by (2.6). Since $S_1 = S_2 = 0$ if $A_1 = A_2$, we only need to consider the summation over $A_1 \neq A_2$. We have

$$\begin{aligned}
(2.13) \quad S_1 &= \sum_j \sum_{B_1, \dots, B_{k-2}} \sum_{A_1 \neq A_2} \left| \delta_{B_j}^{A_1} (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_{k-2}}^{A_2} - \delta_{B_j}^{A_2} (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_{k-2}}^{A_1} \right|^2 \\
&= \sum_j \sum_{B_1, \dots, \widehat{B}_j, \dots, B_{k-2}} \sum_{A_1 \neq A_2} \left(\left| (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_{k-2}}^{A_2} \right|^2 + \left| (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_{k-2}}^{A_1} \right|^2 \right) \\
&= 6(k-2) \sum_{A, B_1, \dots, B_{k-3}} \left| (\mathcal{C}\xi)_{B_1 \dots B_{k-3}}^A \right|^2 = 6(k-2) \|\mathcal{C}(\xi)\|^2.
\end{aligned}$$

By taking summation over B_j at first and then B_l , we see that S_2 equals to

$$\begin{aligned}
(2.14) \quad &\sum_{j \neq l} \sum_{\substack{A_1, A_2, \\ B_1, \dots, B_{k-2}}} \left\langle \delta_{B_j}^{A_1} (\mathcal{C}\xi)_{\dots \widehat{B}_j \dots B_l \dots}^{A_2} - \delta_{B_j}^{A_2} (\mathcal{C}\xi)_{\dots \widehat{B}_j \dots B_l \dots}^{A_1}, \delta_{B_l}^{A_1} (\mathcal{C}\xi)_{\dots B_j \dots \widehat{B}_l \dots}^{A_2} - \delta_{B_l}^{A_2} (\mathcal{C}\xi)_{\dots B_j \dots \widehat{B}_l \dots}^{A_1} \right\rangle \\
&= \sum_{j \neq l} \sum_{\substack{A_1, A_2, \\ \dots, \widehat{B}_j, \dots}} \left\{ \left\langle (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_l \dots}^{A_2}, \delta_{B_l}^{A_1} (\mathcal{C}\xi)_{B_1 \dots A_1 \dots \widehat{B}_l \dots}^{A_2} \right\rangle + \left\langle (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_l \dots}^{A_1}, \delta_{B_l}^{A_2} (\mathcal{C}\xi)_{B_1 \dots A_2 \dots \widehat{B}_l \dots}^{A_1} \right\rangle \right. \\
&\quad \left. - \left\langle (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_l \dots}^{A_2}, \delta_{B_l}^{A_2} (\mathcal{C}\xi)_{B_1 \dots A_1 \dots \widehat{B}_l \dots}^{A_1} \right\rangle - \left\langle (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots B_l \dots}^{A_1}, \delta_{B_l}^{A_1} (\mathcal{C}\xi)_{B_1 \dots A_2 \dots \widehat{B}_l \dots}^{A_2} \right\rangle \right\} \\
&= 2 \sum_{j \neq l} \sum_{\substack{A_1, A_2, \\ \dots, \widehat{B}_j, \dots, \widehat{B}_l, \dots}} \left\langle (\mathcal{C}\xi)_{B_1 \dots \widehat{B}_j \dots A_1 \dots}^{A_2}, (\mathcal{C}\xi)_{B_1 \dots A_1 \dots \widehat{B}_l \dots}^{A_2} \right\rangle \\
&= 2 \sum_{j \neq l} \sum_{A_2, B_1, \dots, B_{k-3}} \left\| (\mathcal{C}\xi)_{B_1 \dots B_{k-3}}^{A_2} \right\|^2 = 2(k-3)(k-2) \|\mathcal{C}(\xi)\|^2.
\end{aligned}$$

Here last two terms in the right hand side of the first identity vanish by $\mathcal{C} \circ \mathcal{C} f = 0$ in (2.5). We relabel indices in the forth identity. Apply (2.13)-(2.14) to (2.12) to get the second identity of (2.8). \square

3. THE L^2 ESTIMATE

3.1. The Euclidean massless field operator. Sämann, Wolf [19] and Mason et al. [16] used the embedding $\mathbb{R}^{5,1} \hookrightarrow \mathbb{M} \subseteq \mathbb{C}^{4 \times 4}$:

$$(3.1) \quad (x^0, x^1, \dots, x^5) \mapsto \begin{pmatrix} 0 & x^0 + x^5 & -x^3 - ix^4 & -x^1 + ix^2 \\ -x^0 - x^5 & 0 & -x^1 - ix^2 & x^3 - ix^4 \\ x^3 + ix^4 & x^1 + ix^2 & 0 & -x^0 + x^5 \\ x^1 - ix^2 & -x^3 + ix^4 & x^0 - x^5 & 0 \end{pmatrix},$$

to study massless field equation, where $\mathbb{CM} = \wedge^2 \mathbb{C}^4$ is the space of complex antisymmetric 4×4 matrices of dimension 6. This embedding is the generalization of the embedding of the Minkowski space into 2×2 -Hermitian matrix space: $\mathbb{R}^{3,1} \hookrightarrow \mathbb{C}^{2 \times 2}$,

$$(x^0, x^1, x^2, x^3) \mapsto \begin{pmatrix} x^0 + x^1 & x^2 + ix^3 \\ x^2 - ix^3 & x^0 - x^1 \end{pmatrix}.$$

The advantage of this embedding is that ones can use two-component notation generalizing Penrose's two-spinor notation [17, 18] and apply the twistor method to study these operators. On the other hand, we can embed 4-dimensional Euclidean space, the quaternionic space \mathbb{H} , into a real subspace of \mathbb{C}^4 by $\mathbb{H} \hookrightarrow \mathbb{C}^{2 \times 2}$,

$$x^0 + ix^1 + jx^2 + kx^3 \mapsto \begin{pmatrix} x^0 + ix^1 & -x^2 - ix^3 \\ x^2 - ix^3 & x^0 - ix^1 \end{pmatrix},$$

and obtain the elliptic version of the differential operators corresponding to massless field equations of higher spins on \mathbb{R}^4 , which are called k -Cauchy-Fueter operators in [22]. For the higher-dimensional case, we use the embedding $\mathbb{H}^n \hookrightarrow \mathbb{C}^{2n \times 2}$, and also apply the twistor method to study k -Cauchy-Fueter equations, e.g. to find series expansion of k -regular functions on \mathbb{H}^n by Penrose integral formula (cf. [12][13]). Motivated by the quaternionic case, we introduce the embedding of 6-dimensional Euclidean space into $\mathbb{C}^{4 \times 4}$ in [14] by $\iota : \mathbb{R}^6 \hookrightarrow \wedge^2 \mathbb{C}^4 \subseteq \mathbb{C}^{4 \times 4}$ given by

$$(3.2) \quad \mathbb{R}^6 \ni x = (x^0, x^1, \dots, x^5) \mapsto \iota(x) = \begin{pmatrix} 0 & ix^0 + x^5 & x^3 + ix^4 & x^1 + ix^2 \\ -ix^0 - x^5 & 0 & x^1 - ix^2 & -x^3 + ix^4 \\ -x^3 - ix^4 & -x^1 + ix^2 & 0 & -ix^0 + x^5 \\ -x^1 - ix^2 & x^3 - ix^4 & ix^0 - x^5 & 0 \end{pmatrix}.$$

This is essentially the embedding (3.1) with x^0 replaced by ix^0 , up to conjugate and sign of some terms. The Euclidean version \mathcal{D}_0 of these massless field operators are

$$\mathcal{D}_0 : C^\infty(\mathbb{R}^6, \odot^k \mathbb{C}^4) \longrightarrow C^\infty(\mathbb{R}^6, \mathbb{C}^4 \otimes \odot^{k-1} \mathbb{C}^4)$$

with

$$(3.3) \quad \mathcal{D}_0(f)_{B_2 \dots B_k}^A := \sum_{B_1} \nabla^{B_1 A} f_{B_1 B_2 \dots B_k},$$

where ∇^{AB} are complex vector fields and the matrix (∇^{AB}) is just the embedding matrix (3.2) with the coordinate x^j replaced by ∂_{x^j} , i.e.,

$$(3.4) \quad (\nabla^{AB}) := \begin{pmatrix} 0 & i\partial_{x^0} + \partial_{x^5} & \partial_{x^3} + i\partial_{x^4} & \partial_{x^1} + i\partial_{x^2} \\ -i\partial_{x^0} - \partial_{x^5} & 0 & \partial_{x^1} - i\partial_{x^2} & -\partial_{x^3} + i\partial_{x^4} \\ -\partial_{x^3} - i\partial_{x^4} & -\partial_{x^1} + i\partial_{x^2} & 0 & -i\partial_{x^0} + \partial_{x^5} \\ -\partial_{x^1} - i\partial_{x^2} & \partial_{x^3} - i\partial_{x^4} & i\partial_{x^0} - \partial_{x^5} & 0 \end{pmatrix}.$$

Define the differential operator $\mathcal{D}_l : C^\infty(\mathbb{R}^6, V_l) \longrightarrow C^\infty(\mathbb{R}^6, V_{l+1})$ by

$$(3.5) \quad (\mathcal{D}_l f)_{B_2 \dots B_{k-l}}^{A_1 \dots A_{l+1}} := \sum_{B_1} \nabla^{B_1} [A_1 f_{B_1 B_2 \dots B_{k-l}}^{A_2 \dots A_{l+1}}].$$

Proposition 3.1. *The sequence*

$$(3.6) \quad 0 \longrightarrow C^\infty(\mathbb{R}^6, V_0) \xrightarrow{\mathcal{D}_0} C^\infty(\mathbb{R}^6, V_1) \xrightarrow{\mathcal{D}_1} C^\infty(\mathbb{R}^6, V_2) \xrightarrow{\mathcal{D}_2} C^\infty(\mathbb{R}^6, V_3) \xrightarrow{\mathcal{D}_3} C^\infty(\mathbb{R}^6, V_4) \longrightarrow 0$$

is a differential complex, i.e., $\mathcal{D}_{l+1} \mathcal{D}_l = 0$.

Proof. By definition, we have

$$(3.7) \quad \begin{aligned} (\mathcal{D}_{l+1} \mathcal{D}_l f)_{B_3 \dots B_{k-l}}^{A_1 \dots A_{l+2}} &= \sum_{B_2} \nabla^{B_2} [A_1 (\mathcal{D}_l f)_{B_2 \dots B_{k-l}}^{A_2 \dots A_{l+2}}] = \sum_{B_1, B_2} \nabla^{B_2} [A_1 \nabla^{B_1} [A_2 f_{B_1 \dots B_{k-l}}^{A_3 \dots A_{l+2}}]] \\ &= \sum_{B_1, B_2} \nabla^{B_2} [A_1 \nabla^{B_1} [A_2] f_{B_1 \dots B_{k-l}}^{A_3 \dots A_{l+2}}], \end{aligned}$$

by using Lemma 2.2 (3). Here $[\dots | \dots | \dots]$ means we do not antisymmetrize indices inside $| \dots |$. Note that ∇^{BC} commutes with ∇^{DA} since they are differential operators with constant coefficients. So

$$(3.8) \quad \begin{aligned} 2 \sum_{B, D} \nabla^{B[A} \nabla^{D]C} f_{\dots BD} &= \sum_{B, D} (\nabla^{BA} \nabla^{DC} - \nabla^{BC} \nabla^{DA}) f_{\dots BD} \\ &= \sum_{B, D} \nabla^{BA} \nabla^{DC} f_{\dots BD} - \sum_{B, D} \nabla^{BA} \nabla^{DC} f_{\dots DB} = 0, \end{aligned}$$

by relabeling B and D in the second identity and f symmetric in B and D . (3.7) vanishes by (3.8). \square

Recall that $\mathcal{C}(\mathcal{D}_0 f) = 0$ for any $f \in C^1(U, \mathcal{V}_0)$ (cf. [14, Introduction]). This fact is true in general.

Proposition 3.2. (1) For $f \in C^\infty(\mathbb{R}^6, \mathcal{V}_l)$, we have $\mathcal{D}_l f \in C^\infty(\mathbb{R}^6, \mathcal{V}_{l+1})$, $l = 0, 1, 2$;
(2) $\mathcal{V}_4 = \{0\}$.

Proof. (1) This is because

$$(3.9) \quad \begin{aligned} \mathcal{C}(\mathcal{D}_l f)_{B_1 \dots B_{k-l-2}}^{A_1 \dots A_l} &= \sum_C \mathcal{D}_l(f)_{B_1 \dots B_{k-l-2} C}^{C A_1 \dots A_l} = \sum_{C, B} \nabla^{B[C} f_{B_1 \dots B_{k-l-2} C B}^{A_1 \dots A_l]} \\ &= \frac{1}{l+1} \sum_{C, B} \left(\nabla^{BC} f_{B_1 \dots B_{k-l-2} C B}^{A_1 \dots A_l} - \sum_{s=1}^l \nabla^{B A_s} f_{B_1 \dots B_{k-l-2} C B}^{A_1 \dots C \dots A_l} \right) = 0, \end{aligned}$$

by using (2.4), Lemma 2.2 (2), $\mathcal{C}f = 0$ and ∇^{BC} antisymmetric in B and C while f symmetric in B and C .

(2) For $f = (f_{B_1 \dots B_{k-4}}^{A_1 A_2 A_3 A_4}) \in \mathcal{V}_4$, it is obvious that $f_{B_1 \dots B_{k-4}}^{A_1 A_2 A_3 A_4} \neq 0$ only if $\{A_1, A_2, A_3, A_4\} = \{1, 2, 3, 4\}$ and so B_j must equal to one of A_1, \dots, A_4 . Without loss of generality, we assume $A_1 = B_{k-4}$. It follows from $\mathcal{C}(f) = 0$ that

$$f_{B_1 \dots B_{k-5} A_1}^{A_1 A_2 A_3 A_4} = \sum_C f_{B_1 \dots B_{k-5} C}^{C A_2 A_3 A_4} = 0,$$

for any fixed $A_2, \dots, A_4, B_1, \dots, B_{k-5}$. So $f = 0$. \square

Since \mathcal{D}_l is the restriction of \mathcal{D}_l on $C^\infty(\mathbb{R}^6, \mathcal{V}_l)$, we have the following corollary.

Corollary 3.1. (1.5) is a differential complex.

3.2. The formal adjoint operators. Let $\epsilon_{ABCD} = \epsilon^{ABCD}$ be the sign of the permutation from $(1, 2, 3, 4)$ to (A, B, C, D) . Then ϵ_{ABCD} vanishes if $\{A, B, C, D\} \neq \{1, 2, 3, 4\}$. We use ϵ^{ABCD} and ϵ_{ABCD} to raise and low indices respectively. For example,

$$\nabla_{AB} := \frac{1}{2} \sum_{C,D} \epsilon_{ABCD} \nabla^{CD}.$$

Then we have

$$\nabla^{AB} = \frac{1}{2} \sum_{C,D=1}^4 \epsilon^{ABCD} \nabla_{CD},$$

since $\sum_{C,D} \epsilon_{ABCD} \epsilon^{CDEF} = 2(\delta_A^E \delta_B^F - \delta_A^F \delta_B^E)$ by definition (cf. [16, P.6]). We know that [14, Proposition 2.1] the operators ∇^{AB} and ∇_{AB} defined above satisfy

$$(3.10) \quad \overline{\nabla_{AB}} = \nabla^{AB} \quad \text{and} \quad \sum_A \nabla_{AB_1} \nabla^{AB_2} = \delta_{B_1}^{B_2} \Delta.$$

Let Θ_{AB} be a scalar differential operator defined by

$$(3.11) \quad \Theta_{AB} f := -e^\varphi \nabla_{AB} (e^{-\varphi} f) = -\nabla_{AB} f + (\nabla_{AB} \varphi) f.$$

Lemma 3.1. *The formal adjoint of the scalar differential operator ∇^{AB} is Θ_{AB} .*

Proof. For any $u, v \in C_0^1(\mathbb{R}^6, \mathbb{C})$, we have

$$\begin{aligned} \langle \nabla^{AB} u, v \rangle_\varphi &= \int_{\mathbb{R}^6} (\nabla^{AB} u) \bar{v} e^{-\varphi} dV = \int_{\mathbb{R}^6} \left\{ \nabla^{AB} (u \bar{v} e^{-\varphi}) - u \cdot \overline{\nabla_{AB} (v e^{-\varphi})} \right\} dV \\ &= \int_{\mathbb{R}^6} u \cdot \overline{\Theta_{AB} v} e^{-\varphi} dV = \langle u, \Theta_{AB} v \rangle_\varphi, \end{aligned}$$

by using Stocks' formula. □

Proposition 3.3. *The formal adjoint \mathcal{D}_l^* of \mathcal{D}_l is given by*

$$(3.12) \quad (\mathcal{D}_l^* f)_{B_1 \dots B_{k-l}}^{A_1 \dots A_l} = - \sum_E \Theta_{E(B_1} f_{B_2 \dots B_{k-l})}^{EA_1 \dots A_l},$$

for $f \in C_0^\infty(\mathbb{R}^6, V_{l+1})$, $l = 0, 1, 2$.

Proof. By definition of the formal adjoint operator, for any $h \in C_0^\infty(\mathbb{R}^6, V_{l+1})$, we have

$$\begin{aligned} \langle f, \mathcal{D}_l^* h \rangle_\varphi &= \langle \mathcal{D}_l f, h \rangle_\varphi = \sum_{\substack{A_1, \dots, A_{l+1}, \\ B_2, \dots, B_{k-l}}} \left\langle \sum_{B_1} \nabla_{B_1} [A_1 f_{B_1 \dots B_{k-l}}^{A_2 \dots A_{l+1}}], h_{B_2 \dots B_{k-l}}^{A_1 \dots A_{l+1}} \right\rangle_\varphi \\ &= \sum_{A_1, \dots, B_1, \dots} \left\langle \nabla_{B_1 A_1} f_{B_1 \dots B_{k-l}}^{A_2 \dots A_{l+1}}, h_{B_2 \dots B_{k-l}}^{A_1 \dots A_{l+1}} \right\rangle_\varphi = \sum_{A_1, \dots, B_1, \dots} \left\langle f_{B_1 \dots B_{k-l}}^{A_2 \dots A_{l+1}}, \Theta_{B_1 A_1} h_{B_2 \dots B_{k-l}}^{A_1 \dots A_{l+1}} \right\rangle_\varphi \\ &= \sum_{A_1, \dots, B_1, \dots} \left\langle f_{B_1 \dots B_{k-l}}^{A_2 \dots A_{l+1}}, - \sum_{A_1} \Theta_{A_1(B_1} h_{B_2 \dots B_{k-l})}^{A_1 \dots A_{l+1}} \right\rangle_\varphi \end{aligned}$$

by using Lemma 2.1 twice and Lemma 3.1. The result follows. □

Lemma 3.2. *For $\varphi = |x|^2$, we have*

$$[\nabla^{AB}, \Theta_{CD}] = 8\delta_{[C}^A \delta_{D]}^B.$$

Proof. Since ∇^{AB}, ∇_{CD} are scalar differential operators with constant coefficients, we have

$$(3.13) \quad [\nabla^{AB}, \Theta_{CD}] = [\nabla^{AB}, -\nabla_{CD} + \nabla_{CD}\varphi] = [\nabla^{AB}, \nabla_{CD}\varphi] = \nabla^{AB}\nabla_{CD}\varphi,$$

by (3.11). Denote

$$z^{AB} := \begin{pmatrix} 0 & ix_0 + x_5 & x_3 + ix_4 & x_1 + ix_2 \\ -ix_0 - x_5 & 0 & x_1 - ix_2 & -x_3 + ix_4 \\ -x_3 - ix_4 & -x_1 + ix_2 & 0 & -ix_0 + x_5 \\ -x_1 - ix_2 & x_3 - ix_4 & ix_0 - x_5 & 0 \end{pmatrix}.$$

Note that for any fixed $D \in \{1, 2, 3, 4\}$, we have $\varphi = \sum_E z^{ED} \overline{z^{ED}}$, and

$$(3.14) \quad \nabla_{CD} z^{ED} = 2\delta_C^E, \quad \nabla_{CD} \overline{z^{ED}} = 0, \quad \nabla_{AB} z^{CD} = 2(\delta_A^C \delta_B^D - \delta_A^D \delta_B^C),$$

by [14, Lemma 2.1], from which we see that

$$(3.15) \quad \nabla_{CD}\varphi = \sum_E (\nabla_{CD} z^{ED} \cdot \overline{z^{ED}} + z^{ED} \cdot \nabla_{CD} \overline{z^{ED}}) = \sum_E 2\delta_C^E \cdot \overline{z^{ED}} = 2\overline{z^{CD}}.$$

Apply (3.15) to (3.13) to get

$$[\nabla^{AB}, \Theta_{CD}] = 2\nabla^{AB} \overline{z^{CD}} = 2\overline{\nabla_{AB} z^{CD}} = 4(\delta_A^C \delta_B^D - \delta_A^D \delta_B^C).$$

The lemma is proved. \square

By Proposition 3.3, we know $\Theta_l f$ belongs to $C_0^\infty(\mathbb{R}^6, \mathcal{V}_l)$ for $f \in C_0^\infty(\mathbb{R}^6, \mathcal{V}_{l+1})$, and so $\Theta_l f - \mathcal{P}(\Theta_l f) \in C_0^\infty(\mathbb{R}^6, \mathcal{V}_l)$. Then the formal adjoint $\Theta_l : C_0^\infty(\mathbb{R}^6, \mathcal{V}_{l+1}) \rightarrow C_0^\infty(\mathbb{R}^6, \mathcal{V}_l)$ of \mathcal{D}_l satisfies

$$(3.16) \quad \Theta_l f = \mathcal{D}_l^* f - \mathcal{P}(\mathcal{D}_l^* f).$$

This is because for $f \in C_0^\infty(\mathbb{R}^6, \mathcal{V}_{l+1})$,

$$\langle h, \Theta_l f \rangle_\varphi = \langle \mathcal{D}_l h, f \rangle_\varphi = \langle \mathcal{D}_l h, f \rangle_\varphi = \langle h, \mathcal{D}_l^* f \rangle_\varphi = \langle h, \mathcal{D}_l^* f - \mathcal{P}(\mathcal{D}_l^* f) \rangle_\varphi,$$

for any $h \in C_0^\infty(\mathbb{R}^6, \mathcal{V}_l)$, by using Proposition 3.2 (1) and Proposition 2.1.

3.3. Proof of the L^2 estimate. It is a well known fact that differential operator $\mathcal{D}_l : L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l) \rightarrow L_\varphi^2(\mathbb{R}^6, \mathcal{V}_{l+1})$ defines a linear, closed, densely defined operator.

Lemma 3.3. *Suppose that $\eta_n \in C_0^\infty(\mathbb{R}^6, \mathbb{R})$ with $\eta_n \equiv 1$ on $B(0, n)$, $\text{supp } \eta_n \subset B(0, n+2)$ and $|\text{grad } \eta_n| \leq 1$. For $f \in \text{Dom}(\mathcal{D}_l) \cap \text{Dom}(\mathcal{D}_{l-1}^*)$, we have $\eta_n f \in \text{Dom}(\mathcal{D}_l) \cap \text{Dom}(\mathcal{D}_{l-1}^*)$ and*

$$\|f - \eta_n f\|_\varphi + \|\mathcal{D}_l(f) - \mathcal{D}_l(\eta_n f)\|_\varphi + \|\mathcal{D}_{l-1}^*(f) - \eta_n \mathcal{D}_{l-1}^*(\eta_n f)\|_\varphi \rightarrow 0, \quad \text{as } n \rightarrow +\infty.$$

Proof. Let $u = \mathcal{D}_l f$ in the weak sense. In definition (1.7) the operator Θ_l can be replaced by the formal adjoint operator \mathcal{D}_l^* by (3.16). Then

$$\begin{aligned} \langle \eta_n f, \Theta_l g \rangle_\varphi &= \langle f, \eta_n \mathcal{D}_l^* g \rangle_\varphi = \langle f, \mathcal{D}_l^*(\eta_n g) \rangle_\varphi - \sum \left\langle f_{B_1 \dots B_{k-l}}^{A_1 \dots A_l}, -\sum_E \nabla_{EB_1} \eta_n g_{B_2 \dots B_{k-l}}^{EA_1 \dots A_l} \right\rangle_\varphi \\ &= \langle \eta_n u, g \rangle_\varphi + \sum \left\langle \nabla^{EB_1} \eta_n f_{B_1 \dots B_{k-l}}^{A_1 \dots A_l}, g_{B_2 \dots B_{k-l}}^{EA_1 \dots A_l} \right\rangle_\varphi \end{aligned}$$

for any $g \in C_0^\infty(\mathbb{R}^6, \mathcal{V}_{l+1})$. Consequently, we have

$$\begin{aligned} \|\mathcal{D}_l(\eta_n f) - \eta_n \mathcal{D}_l(f)\|_\varphi &= \sum_{E, A_1, \dots, B_2, \dots} \int_{\mathbb{R}^6} \left| \sum_{B_1} \nabla^{B_1} [E \eta_n f_{B_1 \dots B_{k-l}}^{A_1 \dots A_l}] \right|^2 e^{-\varphi} dV \\ &\leq C \sum_{E, A_1, \dots, B_2, \dots} \int_{B(0, n+2) \setminus B(0, n)} \left| f_{B_1 \dots B_{k-l}}^{A_1 \dots A_l} \right|^2 e^{-\varphi} dV \rightarrow 0, \end{aligned}$$

for some absolute constant $C > 0$. Similarly, we have

$$\left(\mathcal{D}_{l-1}^*(\eta_n f) - \eta_n \mathcal{D}_{l-1}^*(f)\right)_{B_0 \dots B_{k-l}}^{A_2 \dots A_l} = \sum_E \nabla_{E(B_0)} \eta_n f_{B_1 \dots B_{k-l}}^{EA_2 \dots A_l},$$

and so $\|\mathcal{D}_{l-1}^*(\eta_n f) - \eta_n \mathcal{D}_{l-1}^*(f)\|_\varphi \rightarrow 0$. The result follows. \square

Proof of Theorem 1.2. We only need to prove the estimate (1.14) for any $f \in C_0^\infty(\mathbb{R}^6, \mathcal{V}_l)$. This is because we can assume $f \in L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l)$ is compactly supported by Lemma 3.3, and can check by definition that δ regularization $f_\delta = f * \psi_\delta$, for nonnegative $\psi \in C_0^\infty(\mathbb{R}^6, \mathbb{R})$ with $\text{supp } \psi \subset B(0, 1)$ and $\int \psi = 1$, satisfies

$$\|f - f_\delta\|_\varphi + \|\mathcal{D}_l(f) - \mathcal{D}_l(f_\delta)\|_\varphi + \|\mathcal{D}_{l-1}^*(f) - \mathcal{D}_{l-1}^*(f_\delta)\|_\varphi \rightarrow 0, \quad \text{as } \delta \rightarrow 0.$$

(1) For $l = 1$, noting that $\mathcal{D}_0^* = \mathcal{D}_0^*$ for any $f \in C_0^\infty(\mathbb{R}^6, \mathcal{V}_1)$, we have

$$\begin{aligned} k \|\mathcal{D}_0^* f\|_\varphi^2 &= k \sum_{B_1, \dots, B_k} \left\langle \sum_C \Theta_{C(B_1} f_{B_2 \dots B_k}^C, \sum_D \Theta_{D(B_1} f_{B_2 \dots B_k}^D \right\rangle_\varphi \\ &= k \sum_{C, D, B_1, \dots, B_k} \left\langle \Theta_{C(B_1} f_{B_2 \dots B_k}^C, \Theta_{DB_1} f_{B_2 \dots B_k}^D \right\rangle_\varphi \\ (3.17) \quad &= \sum_{C, D, B_1, \dots, B_k} \left\langle \Theta_{CB_1} f_{B_2 \dots B_k}^C, \Theta_{DB_1} f_{B_2 \dots B_k}^D \right\rangle_\varphi \\ &\quad + \sum_{s=2}^k \sum_{C, D, B_1, \dots, B_k} \left\langle \Theta_{CB_s} f_{B_1 \dots \widehat{B}_s \dots B_k}^C, \Theta_{DB_1} f_{B_2 \dots B_k}^D \right\rangle_\varphi := \Sigma_1 + \Sigma_2, \end{aligned}$$

by using (3.12) and Lemma 2.1 twice. We find that

$$(3.18) \quad \Sigma_1 = \sum_{B_1, \dots, B_k} \left\| \sum_C \Theta_{CB_k} f_{B_1 \dots B_{k-1}}^C \right\|_\varphi^2 \geq 0,$$

To handle Σ_2 , take adjoint and use commutators to change the order of operators to get

$$\begin{aligned} \Sigma_2 &= \sum_{s=2}^k \sum_{C, D, B_1, \dots, B_k} \left\langle \nabla^{DB_1} \Theta_{CB_s} f_{B_1 \dots \widehat{B}_s \dots B_k}^C, f_{B_2 \dots B_k}^D \right\rangle_\varphi \\ &= \sum_{s=2}^k \sum_{C, D, B_1, \dots, B_k} \left\langle \Theta_{CB_s} \nabla^{DB_1} f_{B_1 \dots \widehat{B}_s \dots B_k}^C, f_{B_2 \dots B_k}^D \right\rangle_\varphi \\ &\quad + \sum_{s=2}^k \sum_{C, D, B_1, \dots, B_k} \left\langle \left[\nabla^{DB_1}, \Theta_{CB_s} \right] f_{B_1 \dots \widehat{B}_s \dots B_k}^C, f_{B_2 \dots B_k}^D \right\rangle_\varphi := \Sigma_3 + \Sigma_4, \end{aligned}$$

and

$$\begin{aligned} \Sigma_4 &= \sum_{s=2}^k \sum_{C, D, B_1, \dots, B_k} \left\langle 8\delta_{[C}^D \delta_{B_s}^{B_1} f_{B_1 \dots \widehat{B}_s \dots B_k}^C, f_{B_2 \dots B_k}^D \right\rangle_\varphi \\ (3.19) \quad &= 4 \sum_{s=2}^k \sum_{C, B_2, \dots, B_k} \left\langle f_{B_2 \dots B_k}^C, f_{B_2 \dots B_k}^C \right\rangle_\varphi - 4 \sum_{s=2}^k \sum_{B_1, \dots, B_k} \left\langle f_{B_1 \dots \widehat{B}_s \dots B_k}^{B_1}, f_{B_2 \dots B_k}^{B_s} \right\rangle_\varphi = 4(k-1) \|f\|_\varphi^2, \end{aligned}$$

by using Lemma 3.2 and $\mathcal{C}f = 0$. This term is the main term that we need to control. To control Σ_3 , let us isolate the term concerning $\mathcal{D}_1 f$, note that

$$\begin{aligned}\Sigma_3 &= \sum_{s=2}^k \sum_{C,D,B_1,\dots,B_k} \left\langle \nabla^{DB_1} f_{B_1 \dots \widehat{B}_s \dots B_k}^C, \nabla^{CB_s} f_{B_2 \dots B_k}^D \right\rangle_{\varphi} \\ &= \sum_{s=2}^k \sum_{C,D,B_1,\dots,B_k} \left\langle \nabla^{DB_1} f_{B_1 \dots \widehat{B}_s \dots B_k}^C, -2\nabla^{B_s[C} f_{B_2 \dots B_k}^{D]} \right\rangle_{\varphi} \\ &\quad + \sum_{s=2}^k \sum_{C,D,B_1,\dots,B_k} \left\langle \nabla^{DB_1} f_{B_1 \dots \widehat{B}_s \dots B_k}^C, \nabla^{DB_s} f_{B_2 \dots B_k}^C \right\rangle_{\varphi} := \Sigma_{31} + \Sigma_{32},\end{aligned}$$

by using Lemma 3.1 and ∇^{AB} antisymmetric in A, B . We see that

$$\Sigma_{32} = \sum_{s=2}^k \sum_{C,D,D_1,\dots,D_{k-2}} \left\| \sum_E \nabla^{DE} f_{ED_1 \dots D_{k-2}}^C \right\|_{\varphi}^2 \geq 0,$$

by relabeling indices and

$$\begin{aligned}\Sigma_{31} &= -2 \sum_{s=2}^k \sum_{C,D,B_1,\dots,B_k} \left\langle \nabla^{B_1[C} f_{B_1 \dots \widehat{B}_s \dots B_k}^{D]}, \nabla^{B_s[C} f_{B_2 \dots B_k}^{D]} \right\rangle_{\varphi} \\ &= -2 \sum_{s=2}^k \sum_{C,D,B_2,\dots,\widehat{B}_s,\dots,B_k} \left\langle \mathcal{D}_1(f)_{B_2 \dots \widehat{B}_s \dots B_k}^{CD}, \mathcal{D}_1(f)_{B_2 \dots \widehat{B}_s \dots B_k}^{CD} \right\rangle_{\varphi} = -2(k-1) \|\mathcal{D}_1 f\|_{\varphi}^2,\end{aligned}$$

by using Lemma 2.1. So we get

$$(3.20) \quad \Sigma_3 \geq -2(k-1) \|\mathcal{D}_1 f\|_{\varphi}^2.$$

By (3.17)-(3.20), we get the L^2 estimate for $l = 1$: $\|f\|_{\varphi}^2 \leq \frac{k}{4(k-1)} \|\mathcal{D}_0^* f\|_{\varphi}^2 + \frac{1}{2} \|\mathcal{D}_1 f\|_{\varphi}^2$.

(2) For $l = 2$, the proof is similar, but is more complicated, because we have to use projection \mathcal{P}_l . Note that

$$(3.21) \quad \|\mathcal{D}_1^* f\|_{\varphi}^2 = \langle \mathcal{D}_1^* f, \mathcal{D}_1^* f \rangle_{\varphi} = \|\mathcal{D}_1^* f\|_{\varphi}^2 + \|\mathcal{P}_1(\mathcal{D}_1^* f)\|_{\varphi}^2,$$

by (3.16). Apply Proposition 2.8 to $\|\mathcal{P}_1 \mathcal{D}_1^*(f)\|_{\varphi}^2$ to get

(3.22)

$$\begin{aligned}\|\mathcal{P}_1(\mathcal{D}_1^* f)\|_{\varphi}^2 &= \frac{k-1}{k+2} \sum_{B_1,\dots,B_{k-2}} \left\| \sum_E \mathcal{D}_1^*(f)_{B_1 \dots B_{k-2} E}^E \right\|_{\varphi}^2 = \frac{k-1}{k+2} \sum_{B_1,\dots,B_{k-2}} \left\| \sum_{C,E} \Theta_{C(B_1} f_{B_2 \dots B_{k-2} E)}^{CE} \right\|_{\varphi}^2 \\ &= \frac{1}{(k-1)(k+2)} \sum_{B_1,\dots,B_{k-2}} \left\| \sum_{C,E} \Theta_{CE} f_{B_1 \dots B_{k-2}}^{CE} \right\|_{\varphi}^2 \\ &\leq \frac{4}{(k-1)(k+2)} \sum_{B_1,\dots,B_{k-2}} \sum_E \left\| \sum_C \Theta_{CE} f_{B_1 \dots B_{k-2}}^{CE} \right\|_{\varphi}^2 \\ &\leq \frac{4}{(k-1)(k+2)} \sum_{A_1, B_1, \dots, B_{k-1}} \left\| \sum_C \Theta_{CB_1} f_{B_2 \dots B_{k-1}}^{CA_1} \right\|_{\varphi}^2,\end{aligned}$$

by (3.12) and $\mathcal{C}f = 0$. We use the inequality $|\sum_E a_E|^2 \leq 4 \sum_E |a_E|^2$ in the first inequality and add extra nonnegative terms in the second inequality. Now we have

$$(3.23) \quad \begin{aligned} (k-1) \|\mathcal{D}_1^* f\|_\varphi^2 &= (k-1) \|\mathcal{D}_1^* f\|_\varphi^2 - (k-1) \|\mathcal{P}_1 \mathcal{D}_1^*(f)\|_\varphi^2 \\ &= \left\{ \sum_{C,D,A_1,B_1,\dots,B_{k-1}} \left\langle \Theta_{CB_1} f_{B_2\dots B_{k-1}}^{CA_1}, \Theta_{DB_1} f_{B_2\dots B_{k-1}}^{DA_1} \right\rangle_\varphi - (k-1) \|\mathcal{P}_1 \mathcal{D}_1^*(f)\|_\varphi^2 \right\} \\ &\quad + \sum_{s=2}^{k-1} \sum_{C,D,A_1,B_1,\dots,B_{k-1}} \left\langle \Theta_{CB_s} f_{B_1\dots \widehat{B}_s \dots B_{k-1}}^{CA_1}, \Theta_{DB_1} f_{B_2\dots B_{k-1}}^{DA_1} \right\rangle_\varphi := \Sigma_1 + \Sigma_2, \end{aligned}$$

by (3.21) and expanding symmetrization as in (3.17). Apply (3.22) to Σ_1 in (3.23) to get

$$(3.24) \quad \Sigma_1 \geq \frac{k-2}{k+2} \sum_{D,A_1,B_1,\dots,B_{k-1}} \left\| \sum_C \Theta_{CB_1} f_{B_2\dots B_{k-1}}^{CA_1} \right\|_\varphi^2 \geq 0,$$

if $k \geq 2$. To control Σ_2 , we use commutator to change order of differential operator again to get

$$(3.25) \quad \begin{aligned} \Sigma_2 &= \sum_{s=2}^{k-1} \sum_{C,D,A_1,B_1,\dots,B_{k-1}} \left\langle \nabla^{DB_1} \Theta_{CB_s} f_{B_1\dots \widehat{B}_s \dots B_{k-1}}^{CA_1}, f_{B_2\dots B_{k-1}}^{DA_1} \right\rangle_\varphi \\ &= \sum_{s=2}^{k-1} \sum_{C,D,A_1,B_1,\dots,B_{k-1}} \left\langle \Theta_{CB_s} \nabla^{DB_1} f_{B_1\dots \widehat{B}_s \dots B_{k-1}}^{CA_1}, f_{B_2\dots B_{k-1}}^{DA_1} \right\rangle_\varphi \\ &\quad + \sum_{s=2}^{k-1} \sum_{C,D,A_1,B_1,\dots,B_{k-1}} \left\langle \left[\nabla^{DB_1}, \Theta_{CB_s} \right] f_{B_1\dots \widehat{B}_s \dots B_{k-1}}^{CA_1}, f_{B_2\dots B_{k-1}}^{DA_1} \right\rangle_\varphi := \Sigma_3 + \Sigma_4, \end{aligned}$$

by using Lemma 3.1. As in the case $l = 1$, we have

$$(3.26) \quad \Sigma_4 = 4(k-2) \|f\|_\varphi^2.$$

To control Σ_3 , let us isolate the term concerning $\mathcal{D}_2 f$. Rewrite Σ_3 as

$$(3.27) \quad \begin{aligned} \frac{1}{k-2} \Sigma_3 &= \frac{1}{k-2} \sum_{s=2}^{k-1} \sum_{C,D,A_1,B_1,\dots,B_{k-1}} \left\langle \nabla^{DB_1} f_{B_1\dots \widehat{B}_s \dots B_{k-1}}^{CA_1}, \nabla^{CB_s} f_{B_2\dots B_{k-1}}^{DA_1} \right\rangle_\varphi \\ &= \sum_{\substack{C,D,A_1,E_1,E_2, \\ B_3,\dots,B_{k-1}}} \left\langle \nabla^{DE_1} f_{B_3\dots B_{k-1}E_1}^{CA_1}, \nabla^{CE_2} f_{B_3\dots B_{k-1}E_2}^{DA_1} \right\rangle_\varphi \\ &= \sum_{\substack{C,D,A_1,E_1,E_2, \\ B_3,\dots,B_{k-1}}} \left\{ -3 \left\langle \nabla^{DE_1} f_{B_3\dots B_{k-1}E_1}^{CA_1}, \nabla^{E_2} [C f_{B_3\dots B_{k-1}E_2}^{DA_1}] \right\rangle_\varphi \right. \\ &\quad \left. + \left\langle \nabla^{DE_1} f_{B_3\dots B_{k-1}E_1}^{CA_1}, \nabla^{A_1 E_2} f_{B_3\dots B_{k-1}E_2}^{DC} \right\rangle_\varphi \right. \\ &\quad \left. + \left\langle \nabla^{DE_1} f_{B_3\dots B_{k-1}E_1}^{CA_1}, \nabla^{DE_2} f_{B_3\dots B_{k-1}E_2}^{CA_1} \right\rangle_\varphi \right\} := \Sigma_{31} + \Sigma_{32} + \Sigma_{33}, \end{aligned}$$

by using Lemma 3.1, Lemma 2.2 (2) and relabeling indices. It is easy to see Σ_{33} is a squared sum, which is nonnegative and

$$\begin{aligned}\Sigma_{31} &= -3 \sum_{\substack{C,D,A_1,E_1,E_2, \\ B_3,\dots,B_{k-1}}} \left\langle \nabla^{E_1[C} f_{B_3\dots B_{k-1}E_1}^{DA_1]}, \nabla^{E_2[C} f_{B_3\dots B_{k-1}E_2}^{DA_1]} \right\rangle_{\varphi} \\ &= -3 \sum_{\substack{C,D,A_1, \\ B_3,\dots,B_{k-1}}} \left\| \mathcal{D}_2(f)_{B_3\dots B_{k-1}}^{CA_1} \right\|_{\varphi}^2 = -3 \|\mathcal{D}_2 f\|_{\varphi}^2.\end{aligned}$$

It follows from the expression of $\frac{1}{k-2}\Sigma_3$ in the second identity in (3.27) that

$$\Sigma_{32} = - \sum_{\substack{D,F,G,E_1,E_2, \\ B_3,\dots,B_{k-1}}} \left\langle \nabla^{DE_1} f_{B_3\dots B_{k-1}E_1}^{GF}, \nabla^{GE_2} f_{B_3\dots B_{k-1}E_2}^{DF} \right\rangle_{\varphi} = \frac{-1}{k-2}\Sigma_3,$$

by relabeling indices A_1 as F and C as G and using f_{\dots}^{AB} antisymmetric in A, B . Hence, we get

$$(3.28) \quad \Sigma_3 \geq -\frac{3(k-2)}{2} \|\mathcal{D}_2 f\|_{\varphi}^2,$$

when $k > 2$. By (3.23)-(3.28), we get

$$\|f\|_{\varphi}^2 \leq \frac{k-1}{4(k-2)} \|\mathcal{D}_1^* f\|_{\varphi}^2 + \frac{3}{8} \|\mathcal{D}_2 f\|_{\varphi}^2.$$

(3) For $l = 3$, since $\mathcal{D}_3 f = 0$, we need to prove $\|f\|^2 \leq C \|\mathcal{D}_2^* f\|^2$. Similar to (3.21), we have

$$(3.29) \quad \|\mathcal{D}_2^* f\|_{\varphi}^2 = \|\mathcal{D}_2^* f\|_{\varphi}^2 + \|\mathcal{P}_2 \mathcal{D}_2^*(f)\|_{\varphi}^2.$$

Apply Proposition 2.2 to $\|\mathcal{P}_2 \mathcal{D}_2^*(f)\|_{\varphi}^2$ to get

$$\begin{aligned}(3.30) \quad (k-2) \|\mathcal{P}_2 \mathcal{D}_2^* f\|_{\varphi}^2 &= \frac{2(k-2)^2}{k} \sum_{A_1, B_1, \dots, B_{k-3}} \left\| \sum_{C,E} \Theta_{C(E} f_{B_1\dots B_{k-3}}^{CA_1E} \right\|_{\varphi}^2 \\ &= \frac{2}{k} \sum_{A_1, B_1, \dots, B_{k-3}} \left\| \sum_{C,E} \Theta_{CE} f_{B_1\dots B_{k-3}}^{CA_1E} \right\|_{\varphi}^2 \leq \frac{6}{k} \sum_{A_1, \dots, B_{k-3}} \sum_{E \neq A_1} \left\| \sum_C \Theta_{CE} f_{B_1\dots B_{k-3}}^{CA_1E} \right\|_{\varphi}^2 \\ &\leq \frac{6}{k} \sum_{A_1, A_2, \dots, B_{k-2}} \left\| \sum_C \Theta_{CB_1} f_{B_2\dots B_{k-2}}^{CA_1A_2} \right\|_{\varphi}^2,\end{aligned}$$

by using $\mathcal{C}f = 0$ again, where we use the inequality $|\sum_{j=1}^3 a_j|^2 \leq 3 \sum_{j=1}^3 |a_j|^2$ in the first inequality and add some nonnegative terms in the second inequality.

As in the case $l = 2$ in (3.23), we have

$$\begin{aligned}(3.31) \quad (k-2) \|\mathcal{D}_2^* f\|_{\varphi}^2 &= (k-2) \|\mathcal{D}_2^* f\|_{\varphi}^2 - (k-2) \|\mathcal{P}_2 \mathcal{D}_2^*(f)\|_{\varphi}^2 \\ &= \left\{ \sum_{\substack{C,D,A_1,A_2, \\ B_1,\dots,B_{k-2}}} \left\langle \Theta_{CB_1} f_{B_2\dots B_{k-2}}^{CA_1A_2}, \Theta_{DB_1} f_{B_2\dots B_{k-2}}^{DA_1A_2} \right\rangle_{\varphi} - (k-2) \|\mathcal{P}_2 \mathcal{D}_2^*(f)\|_{\varphi}^2 \right\} \\ &\quad + \sum_{s=2}^{k-2} \sum_{\substack{C,D,A_1,A_2, \\ B_1,\dots,B_{k-2}}} \left\langle \Theta_{CB_s} f_{B_1\dots \widehat{B_s}\dots B_{k-2}}^{CA_1A_2}, \Theta_{DB_1} f_{B_2\dots B_{k-2}}^{DA_1A_2} \right\rangle_{\varphi} := \Sigma_1 + \Sigma_2.\end{aligned}$$

Apply (3.30) to Σ_1 in (3.31) to get

$$(3.32) \quad \Sigma_1 \geq \left(1 - \frac{6}{k}\right) \sum_{D, A_1, A_2, B_1, \dots} \left\| \sum_C \Theta_{CB_1} f_{B_2 \dots B_{k-2}}^{CA_1 A_2} \right\|_{\varphi}^2 \geq 0,$$

if $k \geq 6$. For Σ_2 , we can rewrite it as

$$\begin{aligned} \Sigma_2 &= \sum_{s=2}^{k-2} \sum_{C, D, A_1, A_2, B_1, \dots} \left\langle \nabla^{DB_1} \Theta_{CB_s} f_{B_1 \dots \widehat{B_s} \dots B_{k-2}}^{CA_1 A_2}, f_{B_2 \dots B_{k-2}}^{DA_1 A_2} \right\rangle_{\varphi} \\ &= \sum_{s=2}^{k-2} \sum_{C, D, A_1, A_2, B_1, \dots} \left\langle \Theta_{CB_s} \nabla^{DB_1} f_{B_1 \dots \widehat{B_s} \dots B_{k-2}}^{CA_1 A_2}, f_{B_2 \dots B_{k-2}}^{DA_1 A_2} \right\rangle_{\varphi} \\ &\quad + \sum_{s=2}^{k-2} \sum_{C, D, A_1, A_2, B_1, \dots} \left\langle [\nabla^{DB_1}, \Theta_{CB_s}] f_{B_1 \dots \widehat{B_s} \dots B_{k-2}}^{CA_1 A_2}, f_{B_2 \dots B_{k-2}}^{DA_1 A_2} \right\rangle_{\varphi} := \Sigma_3 + \Sigma_4, \end{aligned}$$

by Lemma 3.1. Similarly to the case $l = 1$, we have

$$(3.33) \quad \Sigma_4 = 4(k-3) \|f\|_{\varphi}^2.$$

To control Σ_3 , we write

$$\begin{aligned} \frac{1}{k-3} \Sigma_3 &= \frac{1}{k-3} \sum_{s=2}^{k-2} \sum_{C, D, A_1, A_2, B_1, \dots, B_{k-2}} \left\langle \nabla^{DB_1} f_{B_1 \dots \widehat{B_s} \dots B_{k-2}}^{CA_1 A_2}, \nabla^{CB_s} f_{B_2 \dots B_{k-2}}^{DA_1 A_2} \right\rangle_{\varphi} \\ &= \sum_{\substack{C, D, A_1, A_2, E_1, E_2, \\ B_1, \dots, B_{k-4}}} \left\langle \nabla^{DE_1} f_{B_1 \dots B_{k-4} E_1}^{CA_1 A_2}, \nabla^{CE_2} f_{B_1 \dots B_{k-4} E_2}^{DA_1 A_2} \right\rangle_{\varphi} \\ &= \sum_{\substack{C, D, A_1, A_2, E_1, E_2, \\ B_1, \dots, B_{k-4}}} \left\{ -4 \left\langle \nabla^{DE_1} f_{B_1 \dots B_{k-4} E_1}^{CA_1 A_2}, \nabla^{E_2} [C f_{B_1 \dots B_{k-4} E_2}^{DA_1 A_2}] \right\rangle_{\varphi} \right. \\ &\quad + \left\langle \nabla^{DE_1} f_{B_1 \dots B_{k-4} E_1}^{CA_1 A_2}, \nabla^{A_1 E_2} f_{B_1 \dots B_{k-4} E_2}^{DC A_2} \right\rangle_{\varphi} \\ &\quad + \left\langle \nabla^{DE_1} f_{B_1 \dots B_{k-4} E_1}^{CA_1 A_2}, \nabla^{A_2 E_2} f_{B_1 \dots B_{k-4} E_2}^{DA_1 C} \right\rangle_{\varphi} \\ &\quad \left. + \left\langle \nabla^{DE_1} f_{B_1 \dots B_{k-4} E_1}^{CA_1 A_2}, \nabla^{DE_2} f_{B_1 \dots B_{k-4} E_2}^{CA_1 A_2} \right\rangle_{\varphi} \right\} := \Sigma_{31} + \Sigma_{32} + \Sigma_{33} + \Sigma_{34}, \end{aligned}$$

by Lemma 3.1 and relabeling indices. It is easy to see that Σ_{34} is a nonnegative squared norm, and

$$\Sigma_{31} = -4 \sum_{\substack{C, D, A_1, A_2, E_1, E_2, \\ B_1, \dots, B_{k-4}}} \left\langle \nabla^{E_1} [C f_{B_1 \dots B_{k-4} E_1}^{DA_1 A_2}], \nabla^{E_2} [C f_{B_1 \dots B_{k-4} E_2}^{DA_1 A_2}] \right\rangle_{\varphi} = -4 \|\mathcal{Q}_3 f\|_{\varphi}^2 = 0,$$

by Lemma 2.1, while

$$\Sigma_{32} = \Sigma_{33} = \frac{-1}{k-3} \Sigma_3,$$

by relabeling indices again. Hence,

$$(3.34) \quad \Sigma_3 \geq 0.$$

Apply (3.32)-(3.34) to (3.31) to get

$$\|f\|_\varphi^2 \leq \frac{k-2}{4(k-3)} \|\mathcal{D}_2^* f\|_\varphi^2.$$

The estimate (1.14) is proved. \square

4. PROOF OF MAIN THEOREMS

We use a general machine to deduce the existence of solution from the L^2 -estimate (cf. e.g. [5]).

Proposition 4.1. *The \square_l is a densely defined, closed, self-adjoint and non-negative operator with domain*

$$Dom(\square_l) = \{f \in L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l) \mid f \in Dom(\mathcal{D}_l), f \in Dom(\mathcal{D}_{l-1}^*), \mathcal{D}_{l-1}^* f \in Dom(\mathcal{D}_{l-1}), \mathcal{D}_l f \in Dom(\mathcal{D}_l^*)\}.$$

This general fact from functional analysis essentially dues to Gaffney [9] (See also [5, Proposition 4.2.3] [23, Proposition 3.1]). So we omit its proof here.

Proof of Theorem 1.1. (1) Theorem 1.2 implies that

$$\frac{1}{C} \|h\|_\varphi^2 \leq \|\mathcal{D}_{l-1}^* h\|_\varphi^2 + \|\mathcal{D}_l h\|_\varphi^2 = \langle \square_l h, h \rangle_\varphi \leq \|\square_l h\|_\varphi \|h\|_\varphi,$$

for $h \in Dom(\square_l)$. Thus \square_l is bounded from below and injective. Since \square_l is self-adjoint and closed, $Range \square_l$ is a dense subset of $L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l)$ by Proposition 4.1. For fixed $f \in L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l)$, we define the complex anti-linear functional

$$\lambda_f : \square_l h \longrightarrow \langle f, h \rangle_\varphi,$$

which is well defined on the dense subset $Range \square_l$ of $L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l)$, since

$$|\lambda_f(\square_l h)| = |\langle f, h \rangle_\varphi| \leq \|f\|_\varphi \|h\|_\varphi \leq C \|f\|_\varphi \|\square_l h\|_\varphi,$$

for $h \in Dom \square_l$. We see that λ_f is bounded on a dense subset and can be uniquely extended to the whole space $L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l)$. By the Riesz representation theorem, there exists a unique $F \in L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l)$ such that $\lambda_f(G) = \langle F, G \rangle_\varphi$ for any $G \in L_\varphi^2(\mathbb{R}^6, \mathcal{V}_l)$ and $\|F\|_\varphi = |\lambda_f| \leq C \|f\|_\varphi$. So we have $\langle F, \square_l h \rangle_\varphi = \langle f, h \rangle_\varphi$ for any $h \in Dom(\square_l)$. This implies $F \in Dom(\square_l^*)$ and $\square_l^* F = f$. Since \square_l is self-adjoint, $F \in Dom(\square_l)$ and $\square_l F = f$. We write $F = N_l f$. Then $\|N_l f\|_\varphi \leq C \|f\|_\varphi$.

(2) Since $N_{l+1} f \in Dom(\square_{l+1})$, we have $\mathcal{D}_l^* N_{l+1} f \in Dom(\mathcal{D}_l)$, $\mathcal{D}_{l+1} N_{l+1} f \in Dom(\mathcal{D}_{l+1}^*)$ and

$$(4.1) \quad \mathcal{D}_l \mathcal{D}_l^* N_{l+1} f = f - \mathcal{D}_{l+1}^* \mathcal{D}_{l+1} N_{l+1} f,$$

by $\square_{l+1} N_{l+1} f = f$. Because $\mathcal{D}_{l+1} f = 0$ and $\mathcal{D}_{l+1} \mathcal{D}_l H = 0$ for any $H \in Dom(\mathcal{D}_l)$, the above identity implies $\mathcal{D}_{l+1}^* \mathcal{D}_{l+1} N_{l+1} f \in Dom(\mathcal{D}_{l+1})$ and

$$\mathcal{D}_{l+1} \mathcal{D}_{l+1}^* \mathcal{D}_{l+1} N_{l+1} f = 0,$$

by \mathcal{D}_{l+1} acting on both sides of (4.1). Then

$$0 = \langle \mathcal{D}_{l+1} \mathcal{D}_{l+1}^* \mathcal{D}_{l+1} N_{l+1} f, \mathcal{D}_{l+1} N_{l+1} f \rangle_\varphi = \|\mathcal{D}_{l+1}^* \mathcal{D}_{l+1} N_{l+1} f\|_\varphi^2,$$

i.e., $\mathcal{D}_{l+1}^* \mathcal{D}_{l+1} N_{l+1} f = 0$. Hence, by (4.1), we have

$$\mathcal{D}_l \mathcal{D}_l^* N_{l+1} f = f.$$

Moreover, we have $\mathcal{D}_l^* N_{l+1} f \perp \ker \mathcal{D}_l$ since $\langle H, \mathcal{D}_l^* N_{l+1} f \rangle_\varphi = \langle \mathcal{D}_l H, N_{l+1} f \rangle_\varphi = 0$ for any $H \in \ker \mathcal{D}_l$. The estimate (1.13) follows from

$$\|\mathcal{D}_l^* N_{l+1} f\|_\varphi^2 + \|\mathcal{D}_{l+1} N_{l+1} f\|_\varphi^2 = \langle \square_{l+1} N_{l+1} f, N_{l+1} f \rangle_\varphi \leq C \|f\|_\varphi^2.$$

The theorem is proved. \square

Proof of Theorem 1.3. Note that $\|f\|_\varphi^2 < +\infty$ for $f \in P(\mathbb{R}^6, \mathcal{V}_{l+1})$, where $\varphi = |x|^2$. So there exists $u \in L^2(\mathbb{R}^6, \mathcal{V}_l)$, such that $\mathcal{D}_l u = f$ and $\mathcal{D}_{l-1}^* u = 0$ by Theorem 1.1. Consequently,

$$(4.2) \quad \square_l u = \Theta_{l-1} \mathcal{D}_l u + \mathcal{D}_{l-1} \Theta_{l-1} u = \Theta_{l-1} f, \quad l = 1, 2, 3.$$

in the sense of distributions, where $\mathcal{D}_{l+1} f = 0$ and $\Theta_{l-1} f$ is a polynomial by the expression of \mathcal{D}_l^* in (3.12) and \mathcal{P} in (2.6).

On the other hand, \square_l is an elliptic differential operator of second order. This is because

$$\langle \sigma(\square_l) \xi, \xi \rangle = \langle \sigma_l \xi, \sigma_l \xi \rangle + \langle \sigma_{l-1}^* \xi, \sigma_{l-1}^* \xi \rangle,$$

for $\xi \in \mathcal{V}_l$, where the inner product is the Euclidean inner product of \mathcal{V}_l and $\sigma(\square_l)$ and σ_l are symbols of operators \square_l and \mathcal{D}_l (cf. (5.1)), respectively. We see that

$$\ker \sigma(\square_l) = \ker \sigma_l \cap \ker \sigma_{l-1}^* = \text{Im} \sigma_{l-1} \cap \ker \sigma_{l-1}^* = \{0\},$$

by Proposition 1.1. Thus we know the solution u of (4.2) is real analytic by applying Theorem 6.6.1 in [15] to elliptic differential operator \square_l of second order with real analytic coefficients. We write the Taylor expression of u as $u = \sum_{m=0}^{\infty} u_m$, where u_m is a polynomial of homogeneous degree m . Suppose f is a polynomial of degree L . Since \mathcal{D}_l is a first order differential operator with constant coefficients, then $\mathcal{D}_l u_m$ is a polynomial of degree $m - 1$ or vanishes. Hence, $\mathcal{D}_l u = f$ implies that

$$\mathcal{D}_l \left(\sum_{m=0}^{L+1} u_m \right) = f.$$

So we get a polynomial solution to $\mathcal{D}_l u = f$ if $\mathcal{D}_{l+1} f = 0$. The result follows. \square

5. THE ELLIPTICITY OF k -MONOGENIC-COMPLEX

Recall that the symbol of the matrix differential operator $\mathcal{D} = \sum_{|\alpha| \leq m} A_{\alpha_1 \dots \alpha_N}(x) \partial_{x_1}^{\alpha_1} \dots \partial_{x_N}^{\alpha_N} : C^\infty(\Omega, W) \rightarrow C^\infty(\Omega, W')$ at (x, v) is defined to be

$$(5.1) \quad \sigma(\mathcal{D})_{(x,v)} := \sum_{|\alpha|=m} A_{\alpha_1 \dots \alpha_N}(x) \left(\frac{v_1}{i} \right)^{\alpha_1} \dots \left(\frac{v_N}{i} \right)^{\alpha_N} : W \rightarrow W',$$

where Ω is a domain in \mathbb{R}^N and $A_{\alpha_1 \dots \alpha_N}$ is a linear transformation from vector space W to W' , $v \in \mathbb{R}^N$. A differential complex

$$C^\infty(\Omega, W_0) \xrightarrow{\mathcal{D}_0} \dots \xrightarrow{\mathcal{D}_{n-1}} C^\infty(\Omega, W_n)$$

is called *elliptic* if its symbol sequence

$$W_0 \xrightarrow{\sigma(\mathcal{D}_0)_{(x,v)}} \dots \xrightarrow{\sigma(\mathcal{D}_{n-1})_{(x,v)}} W_n$$

is exact for any $x \in \Omega$, $v \in \mathbb{R}^N \setminus \{0\}$, that is $\ker \sigma(\mathcal{D}_l)_{(x,v)} = \text{Im} \sigma(\mathcal{D}_{l-1})_{(x,v)}$.

Proof of Proposition 1.1. Let us prove the symbol sequence

$$(5.2) \quad 0 \rightarrow \mathcal{V}_0 \xrightarrow{\sigma_0} \dots \xrightarrow{\sigma_2} \mathcal{V}_3 \rightarrow 0,$$

is exact for fixed $x \in \mathbb{R}^6$ and $v \in \mathbb{R}^6 \setminus \{0\}$, where $\sigma_l := \sigma(\mathcal{D}_l)_{(x,v)}$. Note that

$$(5.3) \quad (\sigma_l f)_{B_2 \dots B_{k-l}}^{A_1 \dots A_{l+1}} = \sum_{B_1=1}^4 M^{B_1} [A_1 f_{B_1 \dots B_{k-l}}^{A_2 \dots A_{l+1}}],$$

with

$$M^{AB} := \frac{1}{i} \begin{pmatrix} 0 & iv_0 + v_5 & v_3 + iv_4 & v_1 + iv_2 \\ -iv_0 - v_5 & 0 & v_1 - iv_2 & -v_3 + iv_4 \\ -v_3 - iv_4 & -v_1 + iv_2 & 0 & -iv_0 + v_5 \\ -v_1 - iv_2 & v_3 - iv_4 & iv_0 - v_5 & 0 \end{pmatrix}$$

a antisymmetric matrix. Since $\sigma_{l+1} \circ \sigma_l = 0$ follows from $\mathcal{D}_{l+1} \circ \mathcal{D}_l = 0$, we only need to prove σ_0 is injective, $\ker \sigma_l \subseteq \text{Im} \sigma_{l-1}$, $l = 1, 2$, and σ_2 is surjective.

(1) For any $\xi \in \ker \sigma_0$, we have

$$\sigma_0(\xi)_{B_2 \dots B_k}^{A_1} = \sum_{B_1} M^{B_1 A_1} \xi_{B_1 B_2 \dots B_k} = 0,$$

for any fixed A_1, B_2, \dots, B_k . It is known that the determinant of M is nonvanishing for any $v \neq 0$ since $M\overline{M}^T = |v|^2 I_{4 \times 4}$ which can be deduced from [14, (2.5) and Proposition 2.1]. This essentially comes from the fact that \mathcal{D}_0 is the Dirac operator. So we have $\xi_{B_1 \dots B_k} = 0$ for any B_1, \dots, B_k . Hence, σ_0 is injective.

(2) For any $\xi \in \ker \sigma_1$, let $\Xi \in \mathcal{V}_0$ be given by

$$\Xi_{B_1 \dots B_k} := \sum_E M_{E(B_1}^{-1} \xi_{B_2 \dots B_k)}^E,$$

where M^{-1} is the inverse of M . Then

$$\begin{aligned} \sigma_0(\Xi)_{B_2 \dots B_k}^{A_1} &= \sum_{B_1} M^{B_1 A_1} \Xi_{B_1 B_2 \dots B_k} \\ (5.4) \quad &= \frac{1}{k} \sum_{E, B_1} \left[M_{EB_1}^{-1} M^{B_1 A_1} \xi_{B_2 \dots B_k}^E + \sum_{s=2}^k M_{EB_s}^{-1} M^{B_1 A_1} \xi_{B_1 \dots \widehat{B}_s \dots B_k}^E \right], \end{aligned}$$

by using Lemma 2.2 (1). Since $\xi \in \ker \sigma_1$, we have

$$(5.5) \quad 0 = 2\sigma_1(\xi)_{B_2 \dots \widehat{B}_s \dots B_k}^{A_1 E} = \sum_{B_1} M^{B_1 A_1} \xi_{B_1 \dots \widehat{B}_s \dots B_k}^E - \sum_{B_1} M^{B_1 E} \xi_{B_1 \dots \widehat{B}_s \dots B_k}^{A_1},$$

by Lemma 2.2 (2). Apply (5.5) to (5.4) to get

$$\sigma_0(\Xi)_{B_2 \dots B_k}^{A_1} = \frac{1}{k} \left[\sum_E \delta_E^{A_1} \xi_{B_2 \dots B_k}^E + \sum_{s=2}^k \sum_{B_1} \delta_{B_s}^{B_1} \xi_{B_1 \dots \widehat{B}_s \dots B_k}^{A_1} \right] = \xi_{B_2 \dots B_k}^{A_1},$$

since M^{-1} is the inverse of M . Thus $\sigma_0 \Xi = \xi$ and so $\ker \sigma_1 \subseteq \text{Im} \sigma_0$.

(3) For any $\xi \in \ker \sigma_2$, set

$$\Xi_{B_1 \dots B_{k-1}}^A := \sum_E M_{E(B_1}^{-1} \xi_{B_2 \dots B_{k-1}}^{AE}.$$

We claim $\Xi \in \mathcal{V}_1$. Then

$$\begin{aligned} \sigma_1(\Xi)_{B_2 \dots B_{k-1}}^{A_1 A_2} &= \sum_{B_1} M^{B_1 [A_1} \Xi_{B_1 B_2 \dots B_{k-1}}^{A_2]} \\ (5.6) \quad &= \frac{1}{k-1} \sum_{E, B_1} \left[M_{EB_1}^{-1} M^{B_1 [A_1} \xi_{B_2 \dots B_{k-1}}^{A_2]E} + \sum_{s=2}^{k-1} M_{EB_s}^{-1} M^{B_1 [A_1} \xi_{B_1 \dots \widehat{B}_s \dots B_{k-1}}^{A_2]E} \right], \end{aligned}$$

by Lemma 2.2 (1). Since $\xi \in \ker \sigma_2$, then for fixed $s \in \{2, \dots, k-1\}$, we have

$$(5.7) \quad \begin{aligned} 0 &= 3\sigma_2(\xi)_{B_2 \dots \widehat{B_s} \dots B_{k-1}}^{A_1 A_2 E} = 3 \sum_{B_1} M^{B_1 [A_1 \xi_{B_1 \dots \widehat{B_s} \dots B_{k-1}}^{A_2 E}]} \\ &= 2 \sum_{B_1} M^{B_1 [A_1 \xi_{B_1 \dots \widehat{B_s} \dots B_{k-1}}^{A_2} E]} + \sum_{B_1} M^{B_1 E} \xi_{B_1 \dots \widehat{B_s} \dots B_{k-1}}^{A_1 A_2}, \end{aligned}$$

by (5.3) and Lemma 2.2 (2). Apply (5.7) to (5.6) to get

$$\sigma_1(\Xi)_{B_2 \dots B_{k-1}}^{A_1 A_2} = \frac{1}{k-1} \left(\sum_E \delta_E^{[A_1 \xi_{B_2 \dots B_{k-1}}^{A_2} E]} - \frac{1}{2} \sum_{s=2}^{k-1} \sum_{B_1} \delta_{B_s}^{B_1} \xi_{B_1 \dots \widehat{B_s} \dots B_{k-1}}^{A_1 A_2} \right) = \frac{-k}{2(k-1)} \xi_{B_2 \dots B_{k-1}}^{A_1 A_2},$$

by M^{-1} inverse to M again. Thus $\sigma_1 \left(\frac{2(k-1)}{-k} \Xi \right) = \xi$.

It remains to show the claim $\mathcal{C}(\Xi) = 0$. Note that for any fixed B_1, \dots, B_{k-2} ,

$$(5.8) \quad \begin{aligned} (k-1) \mathcal{C}(\Xi)_{B_1 \dots B_{k-2}} &= (k-1) \sum_{A_1, A_2} M_{A_2(B_1 \xi_{B_2 \dots B_{k-2} A_1}^{A_1 A_2})}^{-1} \\ &= \sum_{A_1, A_2} \left[M_{A_2 A_1}^{-1} \xi_{B_1 \dots B_{k-2}}^{A_1 A_2} + \sum_{s=1}^{k-2} M_{A_2 B_s}^{-1} \xi_{B_1 \dots A_1 \dots}^{A_1 A_2} \right] = \sum_{A_1, A_2} M_{A_2 A_1}^{-1} \xi_{B_1 \dots B_{k-2}}^{A_1 A_2}, \end{aligned}$$

by $\mathcal{C}\xi = 0$. Since $\det M \neq 0$, $C(\Xi) = 0$ follows from

$$\begin{aligned} (k-1) \sum_{B_1} M^{E B_1} \mathcal{C}(\Xi)_{B_1 \dots B_{k-2}} &= \sum_{A_1, A_2, B_1} M^{B_1 E} M_{A_2 A_1}^{-1} \xi_{B_1 \dots B_{k-2}}^{A_1 A_2} \\ &= - \sum_{A_1, A_2} \sum_{B_1} \left(M^{B_1 A_1} \xi_{B_1 \dots B_{k-2}}^{A_2 E} - M^{B_1 A_2} \xi_{B_1 \dots B_{k-2}}^{A_1 E} \right) M_{A_2 A_1}^{-1} \\ &= \sum_{A_2, B_1} \delta_{A_2}^{B_1} \xi_{B_1 \dots B_{k-2}}^{A_2 E} + \sum_{A_1, B_1} \delta_{A_1}^{B_1} \xi_{B_1 \dots B_{k-2}}^{A_1 E} = 2 \sum_{B_1} \xi_{B_1 \dots B_{k-2}}^{B_1 E} = 0, \end{aligned}$$

for all indices E, B_2, \dots, B_{k-2} , by using (5.7), M antisymmetric and $\mathcal{C}\xi = 0$. So $\Xi \in \mathcal{V}_1$. $\ker \sigma_2 \subseteq \text{Im} \sigma_1$ is proved.

(4) For any $\xi \in \ker \sigma_3 = \mathcal{V}_3$, we do not know whether $\sum_E M_{E(B_1 \xi_{B_2 \dots B_{k-2}}^{A_1 A_2 E})}^{-1}$ belongs to \mathcal{V}_2 or not. But note that the diagram

$$\begin{array}{ccccc} \odot^{k-1} \mathbb{C}^4 \otimes \wedge^3 \mathbb{C}^4 & \xrightarrow{4\tilde{\sigma}} & \odot^{k-2} \mathbb{C}^4 \otimes \wedge^4 \mathbb{C}^4 & \longrightarrow & 0 \\ \downarrow \mathcal{C} & & \downarrow \mathcal{C} & & \\ \mathcal{V}_2 & \xrightarrow{-3\sigma_2} & \mathcal{V}_3 & \longrightarrow & 0 \end{array}$$

is commutative, i.e., $-3\sigma_2 \mathcal{C} = 4\mathcal{C}\tilde{\sigma}$, where $\tilde{\sigma} : \odot^{k-1} \mathbb{C}^4 \otimes \wedge^3 \mathbb{C}^4 \longrightarrow \odot^{k-2} \mathbb{C}^4 \otimes \wedge^4 \mathbb{C}^4$ is given by

$$(5.9) \quad (\tilde{\sigma}\tilde{\Xi})_{B_2 \dots B_{k-1}}^{A_1 \dots A_4} = \sum_{B_1} M^{B_1 [A_1 \tilde{\Xi}_{B_1 \dots B_{k-1}}^{A_2 \dots A_4}]}.$$

This is because

$$\begin{aligned}
-3(\sigma_2 \mathcal{C} \tilde{\Xi})_{B_1 \dots B_{k-3}}^{A_1 A_2 A_3} &= -3 \sum_{E, F} M^{E[A_1 \tilde{\Xi} | F] A_2 A_3}_{B_1 \dots B_{k-3} E F} \\
&= - \sum_{E, F} \left(M^{E A_1 \tilde{\Xi} F A_2 A_3}_{B_1 \dots B_{k-3} E F} - M^{E A_2 \tilde{\Xi} F A_1 A_3}_{B_1 \dots B_{k-3} E F} - M^{E A_3 \tilde{\Xi} F A_2 A_1}_{B_1 \dots B_{k-3} E F} \right) \\
&= 4 \sum_{E, F} M^{E[F \tilde{\Xi} A_1 A_2 A_3] F E} = 4(\mathcal{C} \tilde{\sigma} \tilde{\Xi})_{B_1 \dots B_{k-3}}^{A_1 A_2 A_3}
\end{aligned}$$

by $\sum_{E, F} M^{E F \tilde{\Xi} A_1 A_2 A_3}_{B_1 \dots B_{k-3} F E} = 0$ since ξ is symmetric in E, F while M is antisymmetric in E, F .

Now we construct an inverse image of σ_2 by an inverse image of $\tilde{\sigma}$. Suppose that A_1, \dots, A_4 are different. There must be at least one of A_1, A_2, A_3, A_4 equal to one of B_1, \dots, B_{k-2} . Without loss of generality, we assume $A_1 = B_{k-2}$. For $\xi \in \mathcal{V}_3$, we construct a lifting $\tilde{\xi} \in \odot^{k-2} \mathbb{C}^4 \otimes \wedge^4 \mathbb{C}^4$ as follows

$$(5.10) \quad \tilde{\xi}_{B_1 \dots B_{k-2}}^{A_1 A_2 A_3 A_4} = \xi_{B_1 \dots B_{k-3}}^{A_2 A_3 A_4},$$

when $A_1 = B_{k-2}$. $\tilde{\xi}$ is well defined because if there also exists $A_2 = B_{k-3}$, we must have $\tilde{\xi}_{B_1 \dots B_{k-2}}^{A_1 A_2 A_3 A_4} = -\xi_{B_1 \dots B_{k-4} B_{k-2}}^{A_1 A_3 A_4}$ by $\xi_{B_1 \dots B_{k-3}}^{A_2 A_3 A_4} = -\xi_{B_1 \dots B_{k-4} B_{k-2}}^{A_1 A_3 A_4}$. The latter identity follows from

$$0 = \sum_E \xi_{B_1 \dots B_{k-4} E}^{E A_3 A_4} = \sum_{E=A_1, A_2} \xi_{B_1 \dots B_{k-4} E}^{E A_3 A_4},$$

by $\mathcal{C} \xi = 0$ for $\xi \in \mathcal{V}_3 = \ker \sigma_3$. We have

$$(5.11) \quad \mathcal{C}(\tilde{\xi})_{B_1 \dots B_{k-3}}^{A_1 A_2 A_3} = \sum_C \tilde{\xi}_{B_1 \dots B_{k-3} C}^{C A_1 A_2 A_3} = \xi_{B_1 \dots B_{k-3}}^{A_1 A_2 A_3}$$

for any fixed $A_1, A_2, A_3, B_1, \dots, B_{k-3}$. Now define $\tilde{\Xi} \in \odot^{k-1} \mathbb{C}^4 \otimes \wedge^3 \mathbb{C}^4$ by

$$\tilde{\Xi}_{B_1 \dots B_{k-1}}^{E_2 A_1 A_2} := \sum_{E_1} M_{E_1(B_1)}^{-1} \tilde{\xi}_{B_2 \dots B_{k-1}}^{E_2 A_1 A_2 E_1},$$

and $\Xi := \mathcal{C} \tilde{\Xi}$. Then

$$\Xi_{B_1 \dots B_{k-2}}^{A_1 A_2} = \sum_{E_1, E_2} M_{E_1(B_1)}^{-1} \tilde{\xi}_{B_2 \dots B_{k-2} E_2}^{E_2 A_1 A_2 E_1}.$$

and $\Xi \in \mathcal{V}_2$, since $\mathcal{C} \circ \mathcal{C} \tilde{\Xi} = 0$. Now we show $\sigma_2 \Xi = C \xi$ for some constant $C \neq 0$.

(5.12)

$$\begin{aligned}
(k-1)(\sigma_2 \Xi)_{B_2 \dots B_{k-2}}^{A_1 A_2 A_3} &= (k-1) \sum_{B_1} M^{B_1[A_1 \Xi | A_2 A_3] E_1}_{B_1 \dots B_{k-2}} \\
&= \sum_{B_1, E_1, E_2} \left[M^{B_1[A_1 M_{E_1 E_2}^{-1} \tilde{\xi}_{B_1 \dots B_{k-2}}^{A_2 A_3} E_2 E_1]} + \sum_{s=1}^{k-2} M^{B_1[A_1 M_{E_1 B_s}^{-1} \tilde{\xi}_{B_1 \dots \hat{B}_s \dots B_{k-2} E_2}]} \right] \\
&= \sum_{B_1, E_1, E_2} M_{E_1 E_2}^{-1} M^{B_1[A_1 \tilde{\xi}_{B_1 \dots B_{k-2}}^{A_2 A_3} E_2 E_1]} - \sum_{B_1, E_1} M_{E_1 B_1}^{-1} M^{B_1[A_1 \xi_{B_2 \dots B_{k-2}}^{A_2 A_3} E_1]} \\
&\quad - \sum_{s=2}^{k-2} \sum_{B_1, E_1} M_{E_1 B_s}^{-1} M^{B_1[A_1 \xi_{B_1 \dots \hat{B}_s \dots B_{k-2}}^{A_2 A_3} E_1]} := \Sigma_1 + \Sigma_2 + \Sigma_3,
\end{aligned}$$

by expanding symmetrization and using (5.11). It is easy to see that

$$(5.13) \quad \Sigma_2 = - \sum_{E_1} \delta_{E_1}^{[A_1 \xi_{B_2 \dots B_{k-2}}^{A_2 A_3} E_1]} = -\xi_{B_2 \dots B_{k-2}}^{A_1 A_2 A_3}.$$

On the other hand, it follows from $\xi \in \mathcal{V}_3 = \ker \sigma_3$, i.e. $4(\sigma_3 \xi)_{B_2 \dots \widehat{B}_s \dots B_{k-2}}^{A_1 E_1 A_2 A_3} = 0$, that

$$0 = 4 \sum_{B_1} M^{B_1 [A_1 \xi_{B_1 \dots \widehat{B}_s \dots B_{k-2}}^{E_1 A_2 A_3}]} = 3 \sum_{B_1} M^{B_1 [A_1 \xi_{B_1 \dots \widehat{B}_s \dots B_{k-2}}^{A_2 A_3] E_1} - \sum_{B_1} M^{B_1 E_1 \xi_{B_1 \dots \widehat{B}_s \dots B_{k-2}}^{A_1 A_2 A_3}}.$$

Apply this identity to Σ_3 in (5.12) to get

$$(5.14) \quad \begin{aligned} \Sigma_3 &= -\frac{1}{3} \sum_{s=2}^{k-2} \sum_{B_1, E_1} M_{E_1 B_s}^{-1} M^{B_1 E_1} \xi_{B_1 \dots \widehat{B}_s \dots B_{k-2}}^{A_1 A_2 A_3} = -\frac{1}{3} \sum_{s=2}^{k-2} \sum_{B_1} \delta_{B_s}^{B_1} \xi_{B_1 \dots \widehat{B}_s \dots B_{k-2}}^{A_1 A_2 A_3} \\ &= \frac{3-k}{3} \xi_{B_2 \dots B_{k-2}}^{A_1 A_2 A_3}, \end{aligned}$$

by M^{-1} inverse to M again. Note that $M^{B_1 [A_1 \widetilde{\xi}_{B_1 \dots B_{k-2}}^{A_2 A_3 E_1 E_2}]} = 0$ by $\wedge^5 \mathbb{C}^4 = \{0\}$, which implies

$$(5.15) \quad M^{B_1 E_1 \widetilde{\xi}_{B_1 \dots B_{k-2}}^{A_2 A_3 A_1 E_2}} + M^{B_1 E_2 \widetilde{\xi}_{B_1 \dots B_{k-2}}^{A_2 A_3 E_1 A_1}} = 3M^{B_1 [A_1 \widetilde{\xi}_{B_1 \dots B_{k-2}}^{A_2 A_3] E_1 E_2}.$$

Then apply (5.15) to Σ_1 in (5.12) to get

$$(5.16) \quad \begin{aligned} \Sigma_1 &= \frac{1}{3} \sum_{B_1, E_1, E_2} M_{E_1 E_2}^{-1} \left(M^{B_1 E_1 \widetilde{\xi}_{B_1 \dots B_{k-2}}^{A_2 A_3 A_1 E_2}} + M^{B_1 E_2 \widetilde{\xi}_{B_1 \dots B_{k-2}}^{A_2 A_3 E_1 A_1}} \right) \\ &= \frac{1}{3} \sum_{B_1, E_2} \delta_{E_2}^{B_1} \widetilde{\xi}_{B_1 \dots B_{k-2}}^{A_2 A_3 A_1 E_2} - \frac{1}{3} \sum_{B_1, E_1} \delta_{E_1}^{B_1} \widetilde{\xi}_{B_1 \dots B_{k-2}}^{A_2 A_3 E_1 A_1} = -\frac{2}{3} \xi_{B_2 \dots B_{k-2}}^{A_1 A_2 A_3}, \end{aligned}$$

by (5.11). Now apply (5.13), (5.14) and (5.16) to (5.12) to get

$$(k-1)(\sigma_2 \Xi)_{B_2 \dots B_{k-2}}^{A_1 A_2 A_3} = -\frac{k+2}{3} \xi_{B_2 \dots B_{k-2}}^{A_1 A_2 A_3}.$$

Hence, σ_2 is surjective. Proposition 1.1 is proved. \square

Data availability statement: This manuscript has no associated data.

REFERENCES

- [1] R. Baston, Quaternionic complexes, *J. Geom. Phys.* **8** (1992), 29-52.
- [2] J. Bureš, A. Damiano and I. Sabadini, Explicit resolutions for the complex of several Fueter operators, *J. Geom. Phys.* **57**(3) (2007), 765-775.
- [3] F. Colombo, V. Souček and D. Struppa, Invariant resolutions for several Fueter operators, *J. Geom. Phys.* **56** (7) (2006), 1175-1191.
- [4] D. C. Chang, I. Markina and W. Wang, On the Hodge-type decomposition and cohomology groups of k -Cauchy-Fueter complexes over domains in the quaternionic space, *J. Geom. Phys.* **107** (2016), 15-34.
- [5] S.-C. Chen and M.-C. Shaw, *Partial Differential Equations in Several Complex Variables*, AMS/IP Stud. Adv. Math. Vol. 19, American Mathematical Society/ International Press, Providence, RI/Boston, MA, 2001.
- [6] R. Delanghe, F. Sommen and V. Souček, *Clifford algebra and spinor-valued functions: a function theory for the Dirac operator*, Mathematics and Its Applications **53**, Kluwer academic publishers, Dordrecht, 1992.
- [7] M. Eastwood, R. Penrose and R. Wells, Cohomology and massless fields, *Comm. Math. Phys.* **78** (3) (1980) 305-351.
- [8] W. Fulton and J. Harris, *Representation theory*, a first course, Graduate Text in Mathematics **129**, Springer-Verlag, New York, 1991.
- [9] M. Gaffney, Hilbert space methods in the theory of harmonic integrals, *Trans. Amer. Math. Soc.*, **78** (1955), 426-444.
- [10] L. Hörmander, L^2 estimates and existence theorems for the $\bar{\partial}$ operator, *Acta Math.* **113**(1)(1965) 89-152.
- [11] L. Hörmander, A history of existence theorems for the Cauchy-Riemann complex in L^2 spaces, *J. Geom. Anal.* **13**(2)(2003) 329-357.
- [12] Q.-Q Kang and W. Wang, On Radon-Penrose transformation and k -Cauchy-Fueter operator, *Sci. China Math.* **55**(9) (2012) 1921-1936.
- [13] Q.-Q Kang and W. Wang, On Penrose integral formula and series expansion of k -regular functions on the quaternionic space \mathbb{H}^n , *J. Geom. Phys.* **64** (2013) 192-208.

- [14] Q.-Q Kang and W. Wang, k -monogenic functions over 6-dimensional Euclidean space, *Complex Anly. Oper. Theory* **12**(5) (2018) 1219-1235.
- [15] C. Morrey, *Multiple integrals in the calculus of variations*, in: Die Grundlehren der mathematischen Wissenschaften, vol. 130, Springer-Verlag New York, Inc., New York, 1966.
- [16] L. Mason, R. Reid-Edwards and A. Taghavi-Chabert, Conformal field theories in six-dimensional twistor space, *J. Geom. Phys.* **62** (2012) 2353-2375.
- [17] R. Penrose and W. Rindler, *Spinors and space-time*, Vol. 1, Two-spinor calculus and relativistic fields, Cambridge University Press, Cambridge, 1984.
- [18] R. Penrose and W. Rindler, *Spinors and space-time*, Vol. 2, Spinor and twistor methods in space-time geometry, Cambridge University Press, Cambridge, 1986.
- [19] C. Sämann and M. Wolf, On twistors and conformal field theories from six dimensions, *J. Math. Phys.* **54**(1)(2013) 345-200.
- [20] V. Souček, Clifford analysis for higher spins, in F. Brackx, R. Delanghe, H. Serras (Eds.) *Clifford algebras and their applications in mathematical physics*, Kluwer academic publishers, Dordrecht (1993) 223-232.
- [21] V. Souček, Generalized Cauchy-Riemann equations on manifolds, in J. Chisholm, A. Common and D. Reidel (Eds.) *Proceedings of the workshop Clifford algebra and their applications in mathematical physics*, Publ. Comp. (1986) 219-227.
- [22] W. Wang, The k -Cauchy-Fueter complex, Penrose transformation and Hartogs' phenomenon for quaternionic k -regular functions, *J. Geom. Phys.* **60** (2010) 513-530.
- [23] W. Wang, On the weighted L^2 estimate for the k -Cauchy-Fueter operator and the weighted k -Bergman kernel, *J. Math. Anal. Appl.* **452**(6) (2017) 85-707.
- [24] W. Wang, The Neumann problem for the k -Cauchy-Fueter complex over k -Pseudoconvex domains in \mathbb{R}^4 and the L^2 estimate, *J. Geom. Anal.* (2019) 1233-1258.

(Qianqian Kang) DEPARTMENT OF MATHEMATICS, ZHEJIANG INTERNATIONAL STUDIES UNIVERSITY, HANGZHOU 310012, PR CHINA

Email address: qqkang@zisu.edu.cn

(Wei Wang) DEPARTMENT OF MATHEMATICS, ZHEJIANG UNIVERSITY, ZHEJIANG 310027, PR CHINA

Email address: wwang@zju.edu.cn

(Yuchen Zhang) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI 230026, PR CHINA

Email address: yuchen95@mail.ustc.edu.cn