

BOUNDED COHOMOLOGY OF CLASSIFYING SPACES FOR FAMILIES OF SUBGROUPS

KEVIN LI

ABSTRACT. We introduce a bounded version of Bredon cohomology for groups relative to a family of subgroups. Our theory generalizes bounded cohomology and differs from Mineyev–Yaman’s relative bounded cohomology for pairs. We obtain cohomological characterizations of relative amenability and relative hyperbolicity, analogous to the results of Johnson and Mineyev for bounded cohomology.

1. INTRODUCTION

Bounded cohomology is a homotopy invariant of topological spaces with deep connections to Riemannian geometry via the simplicial volume of manifolds [Gro82]. An astonishing phenomenon known as Gromov’s Mapping Theorem is that for any CW-complex X , the classifying map $X \rightarrow B\pi_1(X)$ induces an isometric isomorphism on bounded cohomology. This emphasizes the importance of the corresponding theory of bounded cohomology for groups, which is also of independent interest due to its plentiful applications in geometric group theory [Mon01, Mon06, Fri17]. The bounded cohomology $H_b^n(G; V)$ of a (discrete) group G with coefficients in a normed G -module V is the cohomology of the cochain complex of bounded G -maps $G^{n+1} \rightarrow V$. The inclusion of bounded G -maps into (not necessarily bounded) G -maps induces the so called comparison map $H_b^n(G; V) \rightarrow H^n(G; V)$. On the one hand, the bounded cohomology groups are very difficult to compute in general. On the other hand, they characterize interesting group-theoretic properties such as amenability [Joh72] and hyperbolicity [Min01, Min02].

Theorem 1.1 (Johnson). *Let G be a group. The following are equivalent:*

- (i) G is amenable;
- (ii) $H_b^n(G; V^\#) = 0$ for any dual normed $\mathbb{R}G$ -module $V^\#$ and all $n \geq 1$;
- (iii) $H_b^1(G; V^\#) = 0$ for any dual normed $\mathbb{R}G$ -module $V^\#$.

Theorem 1.2 (Mineyev). *Let G be a finitely presented group. The following are equivalent:*

- (i) G is hyperbolic;
- (ii) The comparison map $H_b^n(G; V) \rightarrow H^n(G; V)$ is surjective for any normed $\mathbb{Q}G$ -module V and all $n \geq 2$;
- (iii) The comparison map $H_b^2(G; V) \rightarrow H^2(G; V)$ is surjective for any normed $\mathbb{R}G$ -module V .

Date: May 11, 2021.

2020 Mathematics Subject Classification. 55N35, 20J06, 43A07, 20F67.

Key words and phrases. Bounded cohomology, Bredon cohomology, relative amenability, relative hyperbolicity.

There are well-studied notions of relative amenability and relative hyperbolicity in the literature [JOR12, Hru10]. In the present article we introduce a new “relative bounded cohomology theory” characterizing these relative group-theoretic properties as a bounded version of Bredon cohomology. For a group G , a family of subgroups \mathcal{F} is a non-empty set of subgroups which is closed under conjugation and taking subgroups. For a set of subgroup \mathcal{H} of G , we denote by $\mathcal{F}\langle\mathcal{H}\rangle$ the smallest family containing \mathcal{H} . The Bredon cohomology $H_{\mathcal{F}}^n(G; V)$ with coefficients in a G -module V (or more general coefficient systems) is a generalization of group cohomology, which is recovered when \mathcal{F} consists only of the trivial subgroup. A fundamental feature of Bredon cohomology is that for a normal subgroup N of G there is an isomorphism $H_{\mathcal{F}\langle N\rangle}^n(G; V) \cong H^n(G/N; V^N)$. From a topological point of view, the Bredon cohomology of G can be identified with the equivariant cohomology of the classifying space $E_{\mathcal{F}}G$ for the family \mathcal{F} . Especially the classifying spaces $E_{\mathcal{F}LN}G$ and $E_{\mathcal{V}CY}G$ for the family of finite groups and virtually cyclic groups have received a lot of attention in recent years due to their prominent role in the Isomorphism Conjectures of Baum–Connes and Farrell–Jones, respectively.

We introduce the *bounded Bredon cohomology* $H_{\mathcal{F},b}^n(G; V)$ of G with coefficients in a normed G -module V , which generalizes bounded cohomology (Definition 3.1). Our theory still is well-behaved with respect to normal subgroups (Corollary 3.11) and admits a topological interpretation in terms of classifying spaces for families (Theorem 3.10). We obtain the following generalizations of Theorems 1.1 and 1.2. A group G is called *amenable relative to a set of subgroups \mathcal{H}* if there exists a G -invariant mean on the G -set $\coprod_{H \in \mathcal{H}} G/H$.

Theorem 1.3 (Theorem 4.4). *Let G be a group and \mathcal{H} be a set of subgroups. The following are equivalent:*

- (i) G is amenable relative to \mathcal{H} ;
- (ii) $H_{\mathcal{F}\langle\mathcal{H}\rangle,b}^n(G; V^\#) = 0$ for any dual normed $\mathbb{R}G$ -module $V^\#$ and all $n \geq 1$;
- (iii) $H_{\mathcal{F}\langle\mathcal{H}\rangle,b}^1(G; V^\#) = 0$ for any dual normed $\mathbb{R}G$ -module $V^\#$.

We also provide a characterization of relative amenability in terms of relatively injective modules (Proposition 4.6) and of families containing a finite index subgroup (Theorem 4.5). Recall that a finite set of subgroups \mathcal{H} is called a *malnormal* (resp. *almost malnormal*) collection if for all $H_i, H_j \in \mathcal{H}$ and $g \in G$ we have $H_i \cap gH_jg^{-1}$ is trivial (resp. finite) or $i = j$ and $g \in H_i$. A group G is said to be of type $F_{n,\mathcal{F}}$ for a family of subgroups \mathcal{F} , if there exists a model for $E_{\mathcal{F}}G$ with cocompact n -skeleton.

Theorem 1.4 (Theorem 5.4). *Let G be a finitely generated torsionfree group and \mathcal{H} be a finite malnormal collection of subgroups. Suppose that G is of type $F_{2,\mathcal{F}\langle\mathcal{H}\rangle}$ (e.g. G and all subgroups in \mathcal{H} are finitely presented). Then the following are equivalent:*

- (i) G is hyperbolic relative to \mathcal{H} ;
- (ii) The comparison map $H_{\mathcal{F}\langle\mathcal{H}\rangle,b}^n(G; V) \rightarrow H_{\mathcal{F}\langle\mathcal{H}\rangle}^n(G; V)$ is surjective for any normed $\mathbb{Q}G$ -module V and all $n \geq 2$;
- (iii) The comparison map $H_{\mathcal{F}\langle\mathcal{H}\rangle,b}^2(G; V) \rightarrow H_{\mathcal{F}\langle\mathcal{H}\rangle}^2(G; V)$ is surjective for any normed $\mathbb{R}G$ -module V .

In Theorem 1.4 the equivalence of (i) and (iii) still holds if the group G contains torsion and \mathcal{H} is almost malnormal, see Remark 5.5. Note that condition (iii) is trivially satisfied for groups of Bredon cohomological dimension $\text{cd}_{\mathcal{F}\langle\mathcal{H}\rangle}$ equal to 1.

The topological interpretation of bounded Bredon cohomology via classifying spaces for families was used by Löh–Sauer [LS20] to give a new proof of the Nerve Theorem and Vanishing Theorem for amenable covers. We prove a converse of [LS20, Proposition 5.2], generalizing a recent result of [MR] where the case of a normal subgroup is treated.

Theorem 1.5 (Theorem 6.3). *Let G be a group and \mathcal{F} be a family of subgroups. The following are equivalent:*

- (i) *All subgroups in \mathcal{F} are amenable;*
- (ii) *The canonical map $H_{\mathcal{F},b}^n(G; V^\#) \rightarrow H_b^n(G; V^\#)$ is an isomorphism for any dual normed $\mathbb{R}G$ -module $V^\#$ and all $n \geq 1$;*
- (iii) *The canonical map $H_{\mathcal{F},b}^1(G; V^\#) \rightarrow H_b^1(G; V^\#)$ is an isomorphism for any dual normed $\mathbb{R}G$ -module $V^\#$.*

As an application of Theorem 1.5, the comparison map vanishes for groups which admit a “small” model for $E_{\mathcal{F}}G$, where \mathcal{F} is any family consisting of amenable subgroups (Corollary 6.4). Examples are graph products of amenable groups (e.g. right-angled Artin groups) and fundamental groups of graphs of amenable groups.

There is another natural relative cohomology theory given by the relative cohomology of a pair of spaces. For a set of subgroups \mathcal{H} , it gives rise to the cohomology $H^n(G, \mathcal{H}; V)$ of the group pair (G, \mathcal{H}) introduced by Bieri–Eckmann [BE78]. A bounded version $H_b^n(G, \mathcal{H}; V)$ was defined by Mineyev–Yaman [MY] to give a characterization of relative hyperbolicity (see also [Fra18]). A characterization of relative amenability in terms of this relative theory was obtained in [JOR12]. There is a canonical map $H_{\mathcal{F}\langle\mathcal{H}\rangle}^n(G; V) \rightarrow H^n(G, \mathcal{H}; V)$ for $n \geq 2$ which is an isomorphism if \mathcal{H} is malnormal (see Remark 2.1). Similarly, there is a map for the bounded versions but we do not know when it is an isomorphism due to the failure of the excision axiom for bounded cohomology (see Remark 3.12). We also mention that Mineyev–Yaman’s relative bounded cohomology was extended to pairs of groupoids in [Bla16].

Acknowledgements. The present work is part of the author’s PhD project. He wishes to thank his supervisors Nansen Petrosyan for suggesting this topic as well as for numerous discussions and Ian Leary for his support. We are grateful to the organizers of the “Virtual workshop: Simplicial Volumes and Bounded Cohomology” held in September 2020 during which parts of this work were discussed. We thank Clara Löh for several interesting conversations, Sam Hughes and Eduardo Martínez-Pedroza for helpful comments.

CONTENTS

1. Introduction	1
2. Preliminaries on Bredon cohomology and classifying spaces	4
3. Bounded Bredon cohomology	6
4. Characterization of relative amenability	10
5. Characterization of relative hyperbolicity	12
6. Mapping Theorem for families of subgroups	17
References	19

2. PRELIMINARIES ON BREDON COHOMOLOGY AND CLASSIFYING SPACES

In this section we briefly recall the notion of Bredon cohomology for groups and its topological interpretation as the equivariant cohomology of classifying spaces for families of subgroups. For an introduction to Bredon cohomology we refer to [Flu] and for a survey on classifying spaces to [Lüc05].

Let G be a group, which shall always mean a discrete group. A *family of subgroups* \mathcal{F} is a non-empty set of subgroups of G that is closed under conjugation by elements of G and under taking subgroups. Typical examples are $\mathcal{TR} = \{1\}$, $\mathcal{FLN} = \{\text{finite subgroups}\}$, $\mathcal{VCY} = \{\text{virtually cyclic subgroups}\}$, and $\mathcal{ALL} = \{\text{all subgroups}\}$. We will moreover be interested in $\mathcal{AME} = \{\text{amenable subgroups}\}$. For a set of subgroups \mathcal{H} , one can consider the smallest family containing \mathcal{H} which is $\mathcal{F}(\mathcal{H}) = \{\text{conjugates of elements in } \mathcal{H} \text{ and their subgroups}\}$ and called the *family generated by* \mathcal{H} . When \mathcal{H} consists of a single subgroup H , we denote $\mathcal{F}(\mathcal{H})$ instead by $\mathcal{F}(H)$ and call it the *family generated by* H . We denote by G/\mathcal{H} the G -set $\coprod_{H \in \mathcal{H}} G/H$.

Let R be a ring and \mathbf{Mod}_R denote the category of R -modules. We will often suppress the ring R , so that G -modules are understood to be RG -modules. The (\mathcal{F} -restricted) *orbit category* $\mathcal{O}_{\mathcal{F}}G$ has as objects G -sets of the form G/H with $H \in \mathcal{F}$ and as morphisms G -maps. An $\mathcal{O}_{\mathcal{F}}G$ -module is a contravariant functor $M: \mathcal{O}_{\mathcal{F}}G \rightarrow \mathbf{Mod}_R$, the category of which is denoted by $\mathcal{O}_{\mathcal{F}}G\text{-Mod}_R$. Note that $\mathcal{O}_{\mathcal{TR}}G\text{-Mod}_R$ can be identified with the category of G -modules. For a G -module V , there is a coinduced $\mathcal{O}_{\mathcal{F}}G$ -module $V^?$ given by $V^?(G/H) = V^H$. (In the literature this is sometimes called a *fixed-point functor*.) Observe that $(-)^?$ is right-adjoint to the restriction $\mathcal{O}_{\mathcal{F}}G\text{-Mod}_R \rightarrow \mathcal{O}_{\mathcal{TR}}G\text{-Mod}_R$, $M \mapsto M(G/1)$. For a G -space X and a G -CW-complex Y with stabilizers in \mathcal{F} , there are singular and cellular $\mathcal{O}_{\mathcal{F}}G$ -chain complexes $C_*(X^?)(G/H) = C_*(X^H)$ and $C_*^{\text{cell}}(Y^?)(G/H) = C_*^{\text{cell}}(Y^H)$, where $C_*(X^H)$ and $C_*^{\text{cell}}(Y^H)$ denote the usual singular and cellular chain complexes, respectively.

The *Bredon cohomology* of G with coefficients in an $\mathcal{O}_{\mathcal{F}}G$ -module M is defined as

$$H_{\mathcal{F}}^n(G; M) := \text{Ext}_{\mathcal{O}_{\mathcal{F}}G\text{-Mod}_R}^n(R, M)$$

for $n \geq 0$, where R is regarded as a constant $\mathcal{O}_{\mathcal{F}}G$ -module. It can be computed as the cohomology of the cochain complex $\text{Hom}_{\mathcal{O}_{\mathcal{F}}G\text{-Mod}_R}(R[(G/\mathcal{F})^{*+1}]^?, M)$. We define the G -chain complex $C_*^{\mathcal{F}}(G)$ given by G -modules

$$C_n^{\mathcal{F}}(G) := R[(G/\mathcal{F})^{n+1}]$$

with the diagonal G -action and differentials $\partial_n: C_n^{\mathcal{F}}(G) \rightarrow C_{n-1}^{\mathcal{F}}(G)$,

$$\partial_n(g_0H_0, \dots, g_nH_n) = \sum_{i=0}^n (-1)^i (g_0H_0, \dots, \widehat{g_iH_i}, \dots, g_nH_n).$$

For a G -module V , the G -cochain complex $C_{\mathcal{F}}^*(G; V)$ is given by

$$C_{\mathcal{F}}^n(G; V) := \text{Hom}_R(C_n^{\mathcal{F}}(G), V)$$

so that

$$H_{\mathcal{F}}^n(G; V) := H_{\mathcal{F}}^n(G; V^?) \cong H^n(C_{\mathcal{F}}^*(G; V)^G).$$

For a G -space X with stabilizers in \mathcal{F} , the *Bredon cohomology* of X with coefficients in an $\mathcal{O}_{\mathcal{F}}G$ -module M is defined as

$$H_G^n(X; M) := H^n(\mathrm{Hom}_{\mathcal{O}_{\mathcal{F}}G\text{-Mod}_R}(C_*(X^?), M))$$

for $n \geq 0$. If X is a G -CW-complex, then $H_G^n(X; M)$ can be computed using $C_*^{\mathrm{cell}}(X^?)$ instead of $C_*(X^?)$.

A *classifying space* $E_{\mathcal{F}}G$ for the family \mathcal{F} is a terminal object in the G -homotopy category of G -CW-complexes with stabilizers in \mathcal{F} . It can be shown that a G -CW-complex X is a model for $E_{\mathcal{F}}G$ if and only if the fixed-point set X^H is contractible for $H \in \mathcal{F}$ and empty otherwise. An explicit model is given by the geometric realization Y of the semi-simplicial set $\{(G/\mathcal{F})^{n+1} \mid n \geq 0\}$ with the usual face maps, generalizing Segal's construction of EG . Then Y has (non-equivariant) n -cells corresponding to $(G/\mathcal{F})^{n+1}$ and we refer to it as the *simplicial model* for $E_{\mathcal{F}}G$. The cellular $\mathcal{O}_{\mathcal{F}}G$ -chain complex of any model for $E_{\mathcal{F}}G$ is a projective resolution of the constant $\mathcal{O}_{\mathcal{F}}G$ -module R and thus we have

$$(1) \quad H_{\mathcal{F}}^n(G; M) \cong H_G^n(E_{\mathcal{F}}G; M)$$

for any $\mathcal{O}_{\mathcal{F}}G$ -module M . If N is a normal subgroup of G , then a model for $E_{\mathcal{F}\langle N \rangle}G$ is given by $E(G/N)$ regarded as a G -CW-complex and we find

$$(2) \quad H_{\mathcal{F}\langle N \rangle}^n(G; M) \cong H^n(G/N; M(G/N)).$$

Remark 2.1 (Bieri–Eckmann's relative cohomology). For a group G and a set of subgroups \mathcal{H} , Bieri–Eckmann [BE78] have introduced the *relative cohomology* $H^n(G, \mathcal{H}; V)$ of the pair (G, \mathcal{H}) with coefficients in a G -module V . It can be identified with the relative cohomology $H_G^n(EG, \coprod_{H \in \mathcal{H}} G \times_H EH; V)$ of the pair of G -spaces $(EG, \coprod_{H \in \mathcal{H}} G \times_H EH)$. Here a model for EG is chosen that contains $\coprod_{H \in \mathcal{H}} G \times_H EH$ as a subcomplex by taking mapping cylinders. Hence there is a long exact sequence

$$\cdots H^n(G, \mathcal{H}; V) \rightarrow H^n(G; V) \rightarrow \prod_{H \in \mathcal{H}} H^n(H; V) \rightarrow \cdots,$$

which is one of the main features of the relative cohomology groups.

There is a relation between Bredon cohomology and Bieri–Eckmann's relative cohomology as follows. Consider the G -space X obtained as the G -pushout

$$\begin{array}{ccc} \coprod_{H \in \mathcal{H}} G \times_H EH & \longrightarrow & EG \\ \downarrow & & \downarrow \\ \coprod_{H \in \mathcal{H}} G/H & \longrightarrow & X, \end{array}$$

where the left vertical map is induced by collapsing each EH to a point. Then the G -space X has stabilizers in $\mathcal{F}\langle \mathcal{H} \rangle$ and hence admits a G -map $X \rightarrow E_{\mathcal{F}\langle \mathcal{H} \rangle}G$. For an $\mathcal{O}_{\mathcal{F}}G$ -module M , we have maps

$$\begin{array}{ccc} H_G^n(X; M) & \longleftarrow & H_G^n(X, \coprod_{H \in \mathcal{H}} G/H; M) \\ \uparrow & & \downarrow \cong \\ H_G^n(E_{\mathcal{F}\langle \mathcal{H} \rangle}G; M) & & H_G^n(EG, \coprod_{H \in \mathcal{H}} G \times_H EH; M), \end{array}$$

where the right vertical map is an isomorphism by excision. Now, if \mathcal{H} is a malnormal collection, then X is a model for $E_{\mathcal{F}(\mathcal{H})}G$ and we have

$$H_{\mathcal{F}(\mathcal{H})}^n(G; M) \cong H^n(G, \mathcal{H}; M(G/1))$$

for $n \geq 2$. This was shown in [ANCMSS, Theorem 4.16] for the special case when \mathcal{H} consists of a single subgroup.

3. BOUNDED BREDON COHOMOLOGY

In this section we introduce a bounded version of Bredon cohomology and develop some of its basic properties. We follow the exposition in [Fri17] for bounded cohomology. Throughout, let G be a group and \mathcal{F} be a family of subgroups.

From now on, let the ring R be one of \mathbb{Z} , \mathbb{Q} or \mathbb{R} . A *normed G -module* V is a G -module equipped with a G -invariant norm $\|\cdot\|: V \rightarrow \mathbb{R}$. (That is, for all $v, u \in V$, $r \in R$, and $g \in G$ we have $\|v\| = 0$ if and only if $v = 0$, $\|rv\| \leq |r| \cdot \|v\|$, $\|v + u\| \leq \|v\| + \|u\|$, and $\|g \cdot v\| = \|v\|$.) A morphism $f: V \rightarrow W$ of normed G -modules is a morphism of G -modules with finite operator-norm $\|f\|_\infty$. By $\text{bHom}_R(V, W)$ we denote the G -module of R -linear maps $f: V \rightarrow W$ with finite operator-norm, where the G -action is given by $(g \cdot f)(v) = g \cdot f(g^{-1}v)$. We denote the topological dual $\text{bHom}_R(V, \mathbb{R})$ of V by $V^\#$. For a set S and a normed module V , we denote by $\text{bMap}(S, V)$ the module of functions $S \rightarrow V$ with bounded image. Instead of $\text{bMap}(S, \mathbb{R})$ we also write $\ell^\infty(S)$.

The following is our key definition. Recall the notation $G/\mathcal{F} = \coprod_{H \in \mathcal{F}} G/H$ and consider $C_n^{\mathcal{F}}(G) = R[(G/\mathcal{F})^{n+1}]$ as a normed G -module equipped with the ℓ^1 -norm with respect to the generators. For a normed G -module V , we define the cochain complex $C_{\mathcal{F}, b}^*(G; V)$ of normed G -modules by

$$C_{\mathcal{F}, b}^n(G; V) := \text{bHom}_R(C_n^{\mathcal{F}}(G), V)$$

together with the differentials $\delta^n: C_{\mathcal{F}, b}^n(G; V) \rightarrow C_{\mathcal{F}, b}^{n+1}(G; V)$,

$$\delta^n(f)(g_0 H_0, \dots, g_{n+1} H_{n+1}) = \sum_{i=0}^{n+1} (-1)^i f(g_0 H_0, \dots, \widehat{g_i H_i}, \dots, g_{n+1} H_{n+1}).$$

Definition 3.1 (Bounded Bredon cohomology of groups). The *bounded Bredon cohomology* of G with coefficients in a normed G -module V is defined as

$$H_{\mathcal{F}, b}^n(G; V) := H^n(C_{\mathcal{F}, b}^*(G; V)^G)$$

for $n \geq 0$. The inclusion $C_{\mathcal{F}, b}^n(G; V) \subset C_{\mathcal{F}}^n(G; V)$ induces a map

$$c_{\mathcal{F}}^n: H_{\mathcal{F}, b}^n(G; V) \rightarrow H_{\mathcal{F}}^n(G; V)$$

called the *comparison map*.

Note that for $\mathcal{F} = \mathcal{TR}$, Definition 3.1 recovers the usual definition of bounded cohomology.

Remark 3.2 (Coefficient modules). We only consider normed G -modules as coefficients, rather than more general $\mathcal{O}_{\mathcal{F}}G$ -modules equipped with a “compatible norm”. Hence strictly speaking our theory is a bounded version of Nucinkis’ cohomology relative to the G -set G/\mathcal{F} [Nuc99], rather than a bounded version of Bredon cohomology.

Remark 3.3 (Canonical semi-norm). The ℓ^∞ -norm on $C_{\mathcal{F},b}^n(G;V)$ descends to a *canonical semi-norm* on $H_{\mathcal{F},b}^n(G;V)$. However, we do not consider semi-norms anywhere in this article and regard $H_{\mathcal{F},b}^n(G;V)$ merely as an R -module.

Bounded Bredon cohomology satisfies the following basic properties.

Lemma 3.4. *The following hold:*

- (i) *Let $0 \rightarrow V_0 \rightarrow V_1 \rightarrow V_2 \rightarrow 0$ be a short exact sequence of normed G -modules such that $0 \rightarrow V_0^H \rightarrow V_1^H \rightarrow V_2^H \rightarrow 0$ is exact for each $H \in \mathcal{F}$. Then there exists a long exact sequence*

$$0 \rightarrow H_{\mathcal{F},b}^0(G;V_0) \rightarrow H_{\mathcal{F},b}^0(G;V_1) \rightarrow H_{\mathcal{F},b}^0(G;V_2) \rightarrow H_{\mathcal{F},b}^1(G;V_0) \rightarrow \cdots;$$

- (ii) $H_{\mathcal{F},b}^0(G;V) \cong V^G$ for any normed G -module V ;
 (iii) $H_{\mathcal{F},b}^1(G;\mathbb{R}) = 0$.

Proof. (i) Observe that the sequence of cochain complexes

$$0 \rightarrow C_{\mathcal{F},b}^*(G;V_0)^G \rightarrow C_{\mathcal{F},b}^*(G;V_1)^G \rightarrow C_{\mathcal{F},b}^*(G;V_2)^G \rightarrow 0$$

is exact. Then the associated long exact sequence on cohomology is as desired.

- (ii) We have $H_{\mathcal{F},b}^0(G;V) = \ker(\delta^0)$, where

$$\delta^0: \text{bHom}_{RG}(R[G/\mathcal{F}], V) \rightarrow \text{bHom}_{RG}(R[(G/\mathcal{F})^2], V)$$

is given by $\delta^0(f)(g_0H_0, g_1H_1) = f(g_1H_1) - f(g_0H_0)$. Hence $\ker(\delta^0)$ consists precisely of the constant G -maps $G/\mathcal{F} \rightarrow V$, which are in correspondence to V^G .

- (iii) We identify

$$C_{\mathcal{F},b}^n(G;\mathbb{R})^G \cong \text{bMap}\left(\coprod_{H_0, \dots, H_n \in \mathcal{F}} H_0 \setminus (G/H_1 \times \cdots \times G/H_n), \mathbb{R}\right)$$

for $n \geq 1$ and $C_{\mathcal{F},b}^0(G;\mathbb{R})^G \cong \text{bMap}\left(\coprod_{H_0 \in \mathcal{F}} *_{H_0}, \mathbb{R}\right)$. The differentials of this “inhomogeneous” complex in low degrees are given by

$$\delta^0(f)(H_0g_1H_1) = f(*_{H_1}) - f(*_{H_0})$$

$$\delta^1(\varphi)(H_0(g_1H_1, g_2H_2)) = \varphi(H_1g_1^{-1}g_2H_2) - \varphi(H_0g_2H_2) + \varphi(H_0g_1H_1).$$

Then it is not difficult to check that $\ker(\delta^1) = \text{im}(\delta^0)$. \square

We also define the bounded cohomology of a G -space X as follows. Denote by $S_n(X)$ the set of singular n -simplices in X and consider $C_n(X) = R[S_n(X)]$ equipped with the ℓ^1 -norm as a normed G -module. For a normed G -module V , we define the cochain complex $C_b^*(X;V)$ of normed G -modules by

$$C_b^n(X;V) := \text{bHom}_R(C_n(X), V)$$

together with the usual differentials.

Definition 3.5 (Bounded cohomology of G -spaces). The (G -equivariant) *bounded cohomology* of a G -space X with coefficients in a normed G -module V is defined as

$$H_{G,b}^n(X;V) := H^n(C_b^*(X;V)^G)$$

for $n \geq 0$. The inclusion $C_b^n(X;V) \rightarrow C^n(X;V)$ induces a map

$$c_X^n: H_{G,b}^n(X;V) \rightarrow H_G^n(X;V)$$

called the *comparison map*.

Note that the functors $H_{G,b}^*$ are G -homotopy invariant and that $H_{G,b}^n(G/H; V)$ is isomorphic to V^H for $n = 0$ and trivial otherwise. However, beware that $H_{G,b}^*$ is neither a G -cohomology theory, nor can it be computed cellularly for G -CW-complexes, as is the case already when G is the trivial group.

Relative homological algebra. We develop the relative homological algebra that will allow us to compute bounded Bredon cohomology via resolutions, analogous to Ivanov's approach for bounded cohomology [Iva85].

A map $p: A \rightarrow B$ of G -modules is called \mathcal{F} -strongly surjective if for each $H \in \mathcal{F}$ there exists a map $\tau_H: B \rightarrow A$ of H -modules such that $p \circ \tau_H = \text{id}_B$. A G -module P is called *relatively \mathcal{F} -projective* if for every \mathcal{F} -strongly surjective G -map $p: A \rightarrow B$ and every G -map $\phi: P \rightarrow B$, there exists a G -map $\Phi: P \rightarrow A$ such that $p \circ \Phi = \phi$. A chain complex of G -modules is called *relatively \mathcal{F} -projective* if each chain module is relatively \mathcal{F} -projective. A resolution (C_*, ∂_*) of G -modules is called \mathcal{F} -strong if it is contractible as a resolution of H -modules for each $H \in \mathcal{F}$. (That is, there exist H -maps $k_*^H: C_* \rightarrow C_{*+1}$ such that $\partial_{n+1} \circ k_n^H + k_{n-1}^H \circ \partial_n = \text{id}_{C_n}$.)

Lemma 3.6. *The following hold:*

- (i) *If S is a G -set with stabilizers in \mathcal{F} , then the G -module $R[S]$ is relatively \mathcal{F} -projective;*
- (ii) *The resolution $C_*^{\mathcal{F}}(G) \rightarrow R$ of G -modules is \mathcal{F} -strong;*
- (iii) *If X is a G -space with contractible fixed-point set X^H for each $H \in \mathcal{F}$, then the resolution $C_*(X) \rightarrow R$ of G -modules is \mathcal{F} -strong.*

Proof. (i) Given a lifting problem as in the definition of relative \mathcal{F} -projectivity, we construct a lift as follows. Let T be a set of representatives of $G \backslash S$ and denote the stabilizer of an element $t \in T$ by G_t . Then for every $s \in S$ there exist unique elements $t_s \in T$ and $g_s G_{t_s} \in G/G_{t_s}$ such that $g_s^{-1}s = t_s$. Define $\Phi: R[S] \rightarrow A$ on generators by

$$\Phi(s) = g_s \cdot \tau_{G_{t_s}}(\phi(g_s^{-1}s))$$

which is a well-defined G -equivariant lift of ϕ .

- (ii) For $H \in \mathcal{F}$, define $k_*^H: C_*^{\mathcal{F}}(G) \rightarrow C_{*+1}^{\mathcal{F}}(G)$ on generators by

$$k_n^H(g_0 H_0, \dots, g_n H_n) = (eH, g_0 H_0, \dots, g_n H_n).$$

Then k_*^H is an H -equivariant contraction.

- (iii) For $H \in \mathcal{F}$, fix a point $x_H \in X^H$ and define a contraction $k_*^H: C_*(X) \rightarrow C_{*+1}(X)$ of H -chain complexes inductively as follows. Starting with $k_{-1}^H: R \rightarrow C_0(X)$, $r \mapsto r \cdot x_H$, we may assume that k_{n-1}^H has been constructed. Let s be a singular n -simplex in X and denote its stabilizer by H_s . Then there exists a singular $(n+1)$ -simplex s' with 0-th vertex x_H and opposite face s , satisfying $\partial_{n+1}(s') + k_{n-1}^H(\partial_n(s)) = s$. Moreover, since X^{H_s} is contractible we may choose s' such that its image is contained in X^{H_s} . Now, for each H -orbit of singular n -simplices in X choose a representative s , define $k_n^H(s)$ to be s' and then extend H -equivariantly. \square

The proof of the following proposition is standard and omitted.

Proposition 3.7. *Let $f: V \rightarrow W$ be a map of G -modules, $P_* \rightarrow V$ be a G -chain complex with P_n relatively \mathcal{F} -projective for all $n \geq 0$, and $C_* \rightarrow W$ be an \mathcal{F} -strong*

G-resolution. Then there exists a G -chain map $f_: P_* \rightarrow C_*$ extending f , which is unique up to G -chain homotopy.*

While relatively \mathcal{F} -projective \mathcal{F} -strong resolutions are useful to compute Bredon homology, the following dual approach will compute bounded Bredon cohomology.

A map $i: A \rightarrow B$ of normed G -modules is called *\mathcal{F} -strongly injective* if for each $H \in \mathcal{F}$ there exists a map $\sigma_H: B \rightarrow A$ of normed H -modules with $\|\sigma_H\|_\infty \leq K$ such that $\sigma_H \circ i = \text{id}_A$, for a uniform constant $K \geq 0$. A normed G -module I is called *relatively \mathcal{F} -injective* if for every \mathcal{F} -strongly injective G -map $i: A \rightarrow B$ and every map $\psi: A \rightarrow I$ of normed G -modules, there exists a map $\Psi: B \rightarrow I$ of normed G -modules such that $\Psi \circ i = \psi$. A chain complex of normed G -modules is called *relatively \mathcal{F} -injective* if each chain module is relatively \mathcal{F} -injective. A resolution of normed G -modules is called *\mathcal{F} -strong* if it is contractible as a resolution of normed H -modules for each $H \in \mathcal{F}$.

Dual to Lemma 3.6 and Proposition 3.7 we obtain the following.

Lemma 3.8. *Let V be a normed G -module. The following hold:*

- (i) *If S is a G -set with stabilizers in \mathcal{F} , then $\text{bHom}_R(R[S], V)$ is a relatively \mathcal{F} -injective normed G -module;*
- (ii) *The resolution $V \rightarrow C_{\mathcal{F},b}^*(G; V)$ of normed G -modules is \mathcal{F} -strong;*
- (iii) *If X is a G -space with contractible fixed-point set X^H for each $H \in \mathcal{F}$, then the resolution $V \rightarrow C_b^*(X; V)$ of normed G -modules is \mathcal{F} -strong.*

Proposition 3.9. *Let $f: V \rightarrow W$ be a map of normed G -modules, $V \rightarrow C^*$ be an \mathcal{F} -strong resolution, and $W \rightarrow I^*$ be a G -chain complex with I^n relatively \mathcal{F} -injective for all $n \geq 0$. Then there exists a G -chain map $f^*: C^* \rightarrow I^*$ extending f , which is unique up to G -chain homotopy.*

As a consequence of Proposition 3.9, we may use any relatively \mathcal{F} -injective \mathcal{F} -strong resolution to compute bounded Bredon cohomology. We obtain the isomorphisms analogous to (1) and (2) for Bredon cohomology.

Theorem 3.10. *Let G be a group, \mathcal{F} be a family of subgroups, and V be a normed G -module. For all $n \geq 0$ there is an isomorphism*

$$H_{\mathcal{F},b}^n(G; V) \cong H_{G,b}^n(E_{\mathcal{F}}G; V).$$

Proof. Both $C_{\mathcal{F},b}^*(G; V)$ and $C_b^*(E_{\mathcal{F}}G; V)$ are relatively \mathcal{F} -injective \mathcal{F} -strong resolutions of V by Lemma 3.8 and hence G -homotopy equivalent by Proposition 3.9. \square

Corollary 3.11. *Let G be a group, N be a normal subgroup of G , and V be a normed G -module. For all $n \geq 0$ there is an isomorphism*

$$H_{\mathcal{F}\langle N \rangle, b}^n(G; V) \cong H_b^n(G/N; V^N).$$

Proof. As a model for $E_{\mathcal{F}\langle N \rangle}G$ we take $E(G/N)$ regarded as a G -space. Then it suffices to observe that

$$\text{bHom}_{RG}(R[S_n(E(G/N))], V) \cong \text{bHom}_{R[G/N]}(R[S_n(E(G/N))], V^N)$$

and to apply Theorem 3.10 twice. \square

Remark 3.12 (Mineyev–Yaman’s relative bounded cohomology). Mineyev–Yaman have introduced the bounded analogue of Bieri–Eckmann’s relative cohomology for

pairs (Remark 2.1) in [MY]. For a group G , a set of subgroups \mathcal{H} , and a normed G -module V , their relative bounded cohomology groups $H_b^n(G, \mathcal{H}; V)$ can be identified with $H_{G,b}^n(EG, \coprod_{H \in \mathcal{H}} G \times_H EH; V)$ and therefore fit in a long exact sequence

$$\cdots \rightarrow H_b^n(G, \mathcal{H}; V) \rightarrow H_b^n(G; V) \rightarrow \prod_{H \in \mathcal{H}} H_b^n(H; V) \rightarrow \cdots$$

As in Remark 2.1, we denote by X the G -space obtained as a G -pushout from EG by collapsing $G \times_H EH$ to G/H for each $H \in \mathcal{H}$. Then we have maps

$$\begin{array}{ccc} H_{G,b}^n(X; V) & \longleftarrow & H_{G,b}^n(X, \coprod_{H \in \mathcal{H}} G/H; V) \\ \uparrow & & \downarrow \\ H_{G,b}^n(E_{\mathcal{F}(\mathcal{H})}G; V) & & H_{G,b}^n(EG, \coprod_{H \in \mathcal{H}} G \times_H EH; V). \end{array}$$

For $n \geq 2$, the horizontal map is an isomorphism and hence we obtain a map

$$H_{\mathcal{F}(\mathcal{H}),b}^n(G; V) \rightarrow H_b^n(G, \mathcal{H}; V).$$

However, even if \mathcal{H} is a malnormal collection in which case X is a model for $E_{\mathcal{F}(\mathcal{H})}G$, this map need not be an isomorphism due to the failure of the excision axiom for bounded cohomology.

4. CHARACTERIZATION OF RELATIVE AMENABILITY

In this section we prove a characterization of relatively amenable groups in terms of bounded Bredon cohomology analogous to Theorem 1.1.

Recall that a G -invariant mean on a G -set S is an \mathbb{R} -linear map $m: \ell^\infty(S) \rightarrow \mathbb{R}$ which is normalized, non-negative, and G -invariant. (That is, for the constant function $1 \in \ell^\infty(S)$, $f \in \ell^\infty(S)$, and $g \in G$ we have $m(1) = 1$, $m(f) \geq 0$ if $f \geq 0$, and $m(g \cdot f) = m(f)$.) Note that for a G -map $S_1 \rightarrow S_2$ of G -sets, a G -invariant mean on S_1 is pushed forward to a G -invariant mean on S_2 .

Definition 4.1 (Relative amenability). A group G is *amenable relative* to a set of subgroups \mathcal{H} if the G -set G/\mathcal{H} admits a G -invariant mean. When G is amenable relative to \mathcal{H} consisting of a single subgroup H , we also say that H is *co-amenable* in G .

When \mathcal{H} is a finite set of subgroups, we recover the notion of relative amenability studied in [JOR12] (see also [MP03]).

Example 4.2. Let G be a group, H be a subgroup, and \mathcal{H} be a set of subgroups.

- (i) If G is amenable, then G is amenable relative to \mathcal{H} ;
- (ii) If H is a normal subgroup, then H is co-amenable in G if and only if the quotient group G/H is amenable;
- (iii) If H has finite index in G or contains the commutator subgroup $[G, G]$, then H is co-amenable in G ;
- (iv) If \mathcal{H} is finite and G is amenable relative to \mathcal{H} , then \mathcal{H} contains an element that is co-amenable in G ;
- (v) G is amenable relative to \mathcal{H} if and only if G is amenable relative to $\mathcal{F}(\mathcal{H})$.

The following Lemma is proved analogously to [Fri17, Lemma 3.2].

Lemma 4.3. *Let G be a group and \mathcal{H} be a set of subgroups. The following are equivalent:*

- (i) G is amenable relative to \mathcal{H} ;
- (ii) There exists a non-trivial G -invariant functional $\varphi \in \ell^\infty(G/\mathcal{H})^\#$;
- (iii) G/\mathcal{H} admits a G -invariant finitely additive probability measure.

For a family of subgroups \mathcal{F} , consider the short exact sequence of normed $\mathbb{R}G$ -modules

$$0 \rightarrow \mathbb{R} \rightarrow \ell^\infty(G/\mathcal{F}) \rightarrow \ell^\infty(G/\mathcal{F})/\mathbb{R} \rightarrow 0,$$

where \mathbb{R} is regarded as the constant functions, and the exact sequence of topological duals

$$(3) \quad 0 \rightarrow (\ell^\infty(G/\mathcal{F})/\mathbb{R})^\# \rightarrow \ell^\infty(G/\mathcal{F})^\# \rightarrow \mathbb{R} \rightarrow 0.$$

We define the *relative Johnson class* $[J_{\mathcal{F}}] \in H_{\mathcal{F},b}^1(G; (\ell^\infty(G/\mathcal{F})/\mathbb{R})^\#)$ as the cohomology class of the 1-cocycle $J_{\mathcal{F}} \in C_{\mathcal{F},b}^1(G; (\ell^\infty(G/\mathcal{F})/\mathbb{R})^\#)$ given by

$$J_{\mathcal{F}}(g_0H_0, g_1H_1) = \epsilon_{g_1H_1} - \epsilon_{g_0H_0},$$

where $\epsilon_{g_iH_i}$ is the evaluation map at g_iH_i for $i = 0, 1$.

Theorem 4.4. *Let G be a group and \mathcal{F} be a family of subgroups. The following are equivalent:*

- (i) G is amenable relative to \mathcal{F} ;
- (ii) $H_{\mathcal{F},b}^n(G; V^\#) = 0$ for any dual normed $\mathbb{R}G$ -module $V^\#$ and all $n \geq 1$;
- (iii) $H_{\mathcal{F},b}^1(G; V^\#) = 0$ for any dual normed $\mathbb{R}G$ -module $V^\#$;
- (iv) The relative Johnson class $[J_{\mathcal{F}}] \in H_{\mathcal{F},b}^1(G; (\ell^\infty(G/\mathcal{F})/\mathbb{R})^\#)$ vanishes.

Proof. Suppose that G is amenable relative to \mathcal{F} and let $m_{\mathcal{F}}$ be a G -invariant mean on G/\mathcal{F} . Then a G -cochain contraction $k^*: C_{\mathcal{F},b}^*(G; V^\#) \rightarrow C_{\mathcal{F},b}^{*-1}(G; V^\#)$ is given by

$$k^n(f)(g_0H_0, \dots, g_{n-1}H_{n-1})(v) = m_{\mathcal{F}}(gH \mapsto f(gH, g_0H_0, \dots, g_{n-1}H_{n-1})(v))$$

for $f \in C_{\mathcal{F},b}^n(G; V^\#)$, $(g_0H_0, \dots, g_{n-1}H_{n-1}) \in (G/\mathcal{F})^n$, and $v \in V$.

The implications (ii) \Rightarrow (iii) \Rightarrow (iv) are obvious. Suppose that the relative Johnson class $[J_{\mathcal{F}}]$ vanishes. Note that the short exact sequence (3) of normed $\mathbb{R}G$ -modules stays exact when restricting to H -fixed-points for each $H \in \mathcal{H}$. Therefore we obtain an associated long exact sequence on bounded cohomology

$$0 \rightarrow ((\ell^\infty(G/\mathcal{F})/\mathbb{R})^\#)^G \rightarrow (\ell^\infty(G/\mathcal{F})^\#)^G \rightarrow \mathbb{R} \xrightarrow{\partial^0} H_{\mathcal{F},b}^1(G; (\ell^\infty(G/\mathcal{F})/\mathbb{R})^\#) \rightarrow \dots$$

by Lemma 3.4 (i) and (ii). Observe that the image of the connecting homomorphism ∂^0 is precisely $\mathbb{R} \cdot [J_{\mathcal{F}}]$ and hence trivial by assumption. Thus there exists a non-trivial G -invariant element in $\ell^\infty(G/\mathcal{F})^\#$ and we conclude by Lemma 4.3. \square

By a proof similar to that of Theorem 4.4, the vanishing of the first bounded Bredon cohomology with coefficients in any (not necessarily dual) normed $\mathbb{R}G$ -module characterizes that the family \mathcal{F} contains a finite index subgroup of G . Let $\ell^1(G/\mathcal{F})$ denote the normed $\mathbb{R}G$ -module of summable functions $f: G/\mathcal{F} \rightarrow \mathbb{R}$ with norm $\|f\|_1 = \sum_{gH \in G/\mathcal{F}} f(gH)$ and let $\ell_0^1(G/\mathcal{F})$ be the kernel of $\|\cdot\|_1$. We define the class $[K_{\mathcal{F}}] \in H_{\mathcal{F},b}^1(G; \ell_0^1(G/\mathcal{F}))$ as the cohomology class of the 1-cocycle $K_{\mathcal{F}} \in C_{\mathcal{F},b}^1(G; \ell_0^1(G/\mathcal{F}))$ given by

$$K_{\mathcal{F}}(g_0H_0, g_1H_1) = \chi_{g_1H_1} - \chi_{g_0H_0},$$

where $\chi_{g_iH_i}$ is the characteristic function supported at g_iH_i for $i = 0, 1$.

Theorem 4.5. *Let G be a group and \mathcal{F} be a family of subgroups. The following are equivalent:*

- (i) \mathcal{F} contains a finite index subgroup of G ;
- (ii) $H_{\mathcal{F},b}^n(G; V) = 0$ for any normed $\mathbb{R}G$ -module V and all $n \geq 1$;
- (iii) $H_{\mathcal{F},b}^1(G; V) = 0$ for any normed $\mathbb{R}G$ -module V ;
- (iv) The class $[K_{\mathcal{F}}] \in H_{\mathcal{F},b}^1(G; \ell_0^1(G/\mathcal{F}))$ vanishes.

Note that for $\mathcal{F} = \mathcal{TR}$, Theorem 4.5 recovers the characterization of finite groups via bounded cohomology [Fri17, Theorem 3.12]. Furthermore, we obtain a characterization of relative amenability via relatively \mathcal{F} -injective modules, analogous to [Fri17, Proposition 4.18].

Proposition 4.6. *Let G be a group and \mathcal{F} be a family of subgroups. The following are equivalent:*

- (1) G is amenable relative to \mathcal{F} ;
- (2) Any dual normed $\mathbb{R}G$ -module $V^\#$ is relatively \mathcal{F} -injective;
- (3) The trivial normed $\mathbb{R}G$ -module \mathbb{R} is relatively \mathcal{F} -injective.

Proof. Suppose that G is amenable relative to \mathcal{F} and let $m_{\mathcal{F}}$ be a G -invariant mean on G/\mathcal{F} . The inclusion $V^\# \rightarrow C_{\mathcal{F},b}^0(G; V^\#)$ of normed G -modules admits a right inverse r given by

$$r(f)(v) = m_{\mathcal{F}}(gH \mapsto f(gH)(v))$$

for $f \in C_{\mathcal{F},b}^0(G; V^\#)$ and $v \in V$. Then the relative \mathcal{F} -injectivity of $V^\#$ follows from the relative \mathcal{F} -injectivity of $C_{\mathcal{F},b}^0(G; V^\#)$.

Clearly, condition (ii) implies (iii). Suppose that \mathbb{R} is relatively \mathcal{F} -injective. Consider the strongly \mathcal{F} -injective map $i: \mathbb{R} \rightarrow \ell^\infty(G/\mathcal{F})$ of normed G -modules that has an H -section τ_H given by $\tau_H(f) = f(eH)$ for each $H \in \mathcal{F}$. Then the identity $\text{id}_{\mathbb{R}}$ admits an extension along i which yields a non-trivial G -invariant element in $\ell^\infty(G/\mathcal{F})$. By Lemma 4.3 this finishes the proof. \square

5. CHARACTERIZATION OF RELATIVE HYPERBOLICITY

In this section we prove a characterization of relatively hyperbolic groups in terms of bounded Bredon cohomology analogous to Theorem 1.2.

Let G be a finitely generated group and \mathcal{H} be a finite set of subgroups. Recall that G is *hyperbolic relative to \mathcal{H}* if the coned-off Cayley graph is hyperbolic and fine. For example, hyperbolic groups are hyperbolic relative to the trivial subgroup, free products $G_1 * G_2$ are hyperbolic relative to $\{G_1, G_2\}$, and fundamental groups of finite volume hyperbolic manifolds are hyperbolic relative to the cusp subgroups. If G is hyperbolic relative to \mathcal{H} , then \mathcal{H} is almost malnormal and hence malnormal if G is torsionfree.

From now on, let the ring R be either \mathbb{Q} or \mathbb{R} . A map $f: C \rightarrow B$ of normed RG -modules is called *undistorted* if there exists a constant $K \geq 0$ such that for all $b \in \text{im}(f)$ there exists $c \in C$ with $f(c) = b$ such that $\|c\|_C \leq K \cdot \|b\|_B$. A normed RG -module P is called *boundedly projective* if for every undistorted epimorphism $f: C \rightarrow B$ and every map $\phi: P \rightarrow B$ of normed RG -modules, there exists a map $\Phi: P \rightarrow C$ of normed RG -modules such that $f \circ \Phi = \phi$.

The following lemma [MY, Lemma 52] is useful to construct G -equivariant maps.

Lemma 5.1 (Mineyev–Yaman). *Let G be a group and S be a G -set with finite stabilizers. Then $\mathbb{Q}[S]$ is projective as a $\mathbb{Q}G$ -module and boundedly projective as a normed $\mathbb{Q}G$ -module when equipped with the ℓ^1 -norm.*

Let X be a G -CW-complex with cocompact $(n+1)$ -skeleton and consider for $k \geq 0$ the cellular chains $C_k^{\text{cell}}(X; R)$ as a normed RG -module equipped with the ℓ^1 -norm. We say that X satisfies a *linear homological isoperimetric inequality over R in degree n* if the boundary map

$$\partial_{n+1} : C_{n+1}^{\text{cell}}(X; R) \rightarrow C_n^{\text{cell}}(X; R)$$

is undistorted. Equivalently, there exists a constant $K \geq 0$ such that for every cellular n -boundary $b \in B_n^{\text{cell}}(X; R)$ we have $\|b\|_{\partial} \leq K \cdot \|b\|_1$, where

$$\|b\|_{\partial} := \inf\{\|c\|_1 \mid c \in C_{n+1}^{\text{cell}}(X; R), \partial_{n+1}(c) = b\}$$

(which is sometimes called the *filling norm*).

If G is hyperbolic relative to \mathcal{H} , Mineyev–Yaman [MY, Theorem 41] have constructed the so called “ideal complex” X . It is in particular a cocompact G -CW-complex with precisely one equivariant 0-cell G/H for each $H \in \mathcal{H}$ and finite edge-stabilizers. Moreover, X is (non-equivariantly) contractible and hence a model for $E_{\mathcal{F}(\mathcal{H})}G$ provided that G is torsionfree. We summarize some of its properties [MY, Theorem 47 and 51].

Theorem 5.2 (Mineyev–Yaman). *Let G be a finitely generated torsionfree group and \mathcal{H} be a finite set of subgroups. If G is hyperbolic relative to \mathcal{H} , then there exists a cocompact model X for $E_{\mathcal{F}(\mathcal{H})}G$ satisfying the following:*

- (i) X satisfies linear homological isoperimetric inequalities over \mathbb{Q} in degree n for all $n \geq 1$;
- (ii) There exists a map $q : X^{(0)} \times X^{(0)} \rightarrow C_1^{\text{cell}}(X; \mathbb{Q})$ with $\partial_1(q(a, b)) = b - a$, called a homological \mathbb{Q} -bicombing, that is G -equivariant and satisfies

$$\|q(a, b) + q(b, c) - q(a, c)\|_1 \leq K$$

for all $a, b, c \in X^{(0)}$ and a uniform constant $K \geq 0$.

The following criterion for relative hyperbolicity is a combination of [Fra18, Proposition 8.3 and Theorem 8.5] (see also [MP16, Theorems 1.6 and 1.10]).

Theorem 5.3 (Franceschini, Martínez-Pedroza). *Let G be a group and \mathcal{H} be a finite set of subgroups. Then G is hyperbolic relative to \mathcal{H} if there exists a G -CW-complex Z satisfying the following:*

- (i) Z is simply-connected;
- (ii) The 2-skeleton $Z^{(2)}$ is cocompact;
- (iii) \mathcal{H} is a set of representatives of distinct conjugacy classes of vertex-stabilizers such that each infinite stabilizer is represented;
- (iv) The edge-stabilizers of Z are finite;
- (v) Z satisfies a linear homological isoperimetric inequality over \mathbb{R} in degree 1.

We prove the following characterization of relative hyperbolicity closely following Mineyev’s original proof [Min01, Theorem 11] and [Min02, Theorem 9].

Theorem 5.4. *Let G be a finitely generated torsionfree group and \mathcal{H} be a finite malnormal collection of subgroups. Let \mathcal{F} be the family $\mathcal{F}(\mathcal{H})$ and suppose that G is of type $F_{2, \mathcal{F}}$. Then the following are equivalent:*

- (i) G is hyperbolic relative to \mathcal{H} ;
- (ii) The comparison map $H_{\mathcal{F},b}^n(G; V) \rightarrow H_{\mathcal{F}}^n(G; V)$ is surjective for any normed $\mathbb{Q}G$ -module V and all $n \geq 2$;
- (iii) The comparison map $H_{\mathcal{F},b}^2(G; V) \rightarrow H_{\mathcal{F}}^2(G; V)$ is surjective for any normed $\mathbb{R}G$ -module V .

Proof. Suppose that G is hyperbolic relative to \mathcal{H} . Let X be the model for $E_{\mathcal{F}}G$ that is given by Mineyev–Yaman’s ideal complex (Theorem 5.2) and Y be the simplicial model for $E_{\mathcal{F}}G$ with (non-equivariant) n -cells corresponding to $(G/\mathcal{F})^{n+1}$ for all $n \geq 0$. We claim that there is a G -chain map

$$\varphi_*: C_*^{\text{cell}}(Y; \mathbb{Q}) \rightarrow C_*^{\text{cell}}(X; \mathbb{Q})$$

with φ_n bounded for all $n \geq 2$, admitting a G -homotopy left inverse. We construct φ_* inductively as follows. In degree 0, we define

$$\varphi_0: C_0^{\text{cell}}(Y; \mathbb{Q}) = \mathbb{Q}[G/\mathcal{F}] \rightarrow C_0^{\text{cell}}(X; \mathbb{Q})$$

to map a generator of the form eH to the vertex of X with stabilizer containing H . Then extend G -equivariantly and \mathbb{Q} -linearly to all of $\mathbb{Q}[G/\mathcal{F}]$. In degree 1, we define $\varphi_1: C_1^{\text{cell}}(Y; \mathbb{Q}) \rightarrow C_1^{\text{cell}}(X; \mathbb{Q})$ on generators by

$$\varphi_1(g_0H_0, g_1H_1) = q(\varphi_0(g_0H_0), \varphi_0(g_1H_1)),$$

where q is the homological \mathbb{Q} -bicombing on X from Theorem 5.2 (ii). Since both q and φ_0 are G -equivariant, so is φ_1 . In degree 2, we consider the maps

$$(4) \quad \begin{array}{ccc} C_2^{\text{cell}}(Y; \mathbb{Q}) & \xrightarrow{\partial_2^Y} & C_1^{\text{cell}}(Y; \mathbb{Q}) \\ & & \downarrow \varphi_1 \\ C_2^{\text{cell}}(X; \mathbb{Q}) & \xrightarrow{\partial_2^X} & C_1^{\text{cell}}(X; \mathbb{Q}) \end{array}$$

and observe that the composition $\varphi_1 \circ \partial_2^Y$ is bounded by properties of q and that ∂_2^X is undistorted by Theorem 5.2 (i). There is a G -invariant decomposition

$$C_2^{\text{cell}}(Y; \mathbb{Q}) \cong \mathbb{Q}[S_1] \oplus \mathbb{Q}[S_2],$$

where S_1 and S_2 denote the sets of 2-cells of Y with trivial resp. non-trivial stabilizers. We obtain a bounded G -map $\varphi_2: C_2^{\text{cell}}(Y; \mathbb{Q}) \rightarrow C_2^{\text{cell}}(X; \mathbb{Q})$ by using the bounded projectivity of $\mathbb{Q}[S_1]$ (Lemma 5.1) and by setting φ_2 to be zero on $\mathbb{Q}[S_2]$. This renders the square (4) commutative because the edge-stabilizers of X are trivial.

Assuming that φ_n has been constructed, one analogously defines a bounded G -map φ_{n+1} using that ∂_{n+1}^X is undistorted by Theorem 5.2 (i). Thus one obtains a G -chain map φ_* with φ_n bounded for $n \geq 2$. To conclude the claim, we note that $C_*^{\text{cell}}(Y; \mathbb{Q})$ is a relatively \mathcal{F} -projective \mathcal{F} -strong resolutions of \mathbb{Q} by Lemma 3.6. Hence by Proposition 3.7 any G -chain map $\psi_*: C_*^{\text{cell}}(X; \mathbb{Q}) \rightarrow C_*^{\text{cell}}(Y; \mathbb{Q})$ extending $\text{id}_{\mathbb{Q}}$ is a G -homotopy left inverse of φ_* .

Now, let V be a normed $\mathbb{Q}G$ -module. Applying $\text{Hom}_{\mathbb{Q}G}(-, V)$ yields a cochain map

$$\varphi^*: C_{\text{cell}}^*(X; V)^G \rightarrow C_{\text{cell}}^*(Y; V)^G$$

with homotopy right inverse ψ^* . In particular, the composition $\varphi^* \circ \psi^*$ induces the identity on $H^*(C_{\text{cell}}^*(Y; V)^G) \cong H_{\mathcal{F}}^*(G; V)$. Finally, for $n \geq 2$ let $c \in C_{\text{cell}}^n(Y; V)^G$

be a cocycle. Then $\varphi^n(\psi^n(c))$ and c represent the same cohomology class in $H_{\mathcal{F}}^n(G; V)$. We have

$$\|\varphi^n(\psi^n(c))\|_{\infty} = \|\psi^n(c) \circ \varphi_n\|_{\infty} \leq \|\psi^n(c)\|_{\infty} \cdot \|\varphi_n\|_{\infty},$$

where φ_n is bounded by construction and so is $\psi^n(c) \in C_{\text{cell}}^n(X; V)^G$ because X has only finitely many orbits of n -cells. Thus we have shown that for $n \geq 2$ every cohomology class in $H_{\mathcal{F}}^n(G; V)$ admits a bounded representative.

Obviously condition (ii) implies (iii). Suppose that the comparison map is surjective in degree 2 for coefficients in any normed $\mathbb{R}G$ -module. Let Z be a model for $E_{\mathcal{F}}G$ with cocompact 2-skeleton. Since \mathcal{H} is malnormal, by collapsing fixed-point sets of Z we may assume that for every non-trivial subgroup $H \in \mathcal{F}$ the fixed-point set Z^H consists of precisely one point.¹ In order to apply Theorem 5.3 and conclude that G is hyperbolic relative to \mathcal{H} , it remains to verify that Z satisfies a linear homological isoperimetric inequality over \mathbb{R} in degree 1.

We take as coefficients the cellular 1-boundaries $V := B_1^{\text{cell}}(Z; \mathbb{R})$ equipped with the norm $\|\cdot\|_{\partial}$. Let Y be the simplicial model for $E_{\mathcal{F}}G$. Then there is a G -chain homotopy equivalence

$$\psi_* : C_*^{\text{cell}}(Z; \mathbb{R}) \rightarrow C_*^{\text{cell}}(Y; \mathbb{R})$$

with G -homotopy inverse φ_* . Applying $\text{Hom}_{\mathbb{R}G}(-, V)$ yields a cochain homotopy equivalence

$$\psi^* : C_{\text{cell}}^*(Y; V)^G \rightarrow C_{\text{cell}}^*(Z; V)^G$$

with homotopy inverse φ^* . In particular, the composition $\psi^* \circ \varphi^*$ induces the identity on $H^*(C_{\text{cell}}^*(Z; V)^G) \cong H_{\mathcal{F}}^*(G; V)$. Consider the 2-cocycle $u \in C_{\text{cell}}^2(Z; V)^G$ given by the boundary map

$$u = \partial_2 : C_2^{\text{cell}}(Z; \mathbb{R}) \rightarrow B_1^{\text{cell}}(Z; \mathbb{R}) = V.$$

Then we can write

$$(5) \quad u = (\psi^2 \circ \varphi^2)(u) + \delta_Z^1(v)$$

for some $v \in C_{\text{cell}}^1(Z; V)^G$. Since the comparison map $H_{\mathcal{F}, b}^2(G; V) \rightarrow H_{\mathcal{F}}^2(G; V)$ is surjective by hypothesis, we can write

$$(6) \quad \varphi^2(u) = u' + \delta_Y^1(v')$$

for a *bounded* 2-cocycle $u' \in C_{\text{cell}}^2(Y; V)^G$ and some $v' \in C_{\text{cell}}^1(Y; V)^G$. For a fixed vertex $y \in Y^{(0)} = G/\mathcal{F}$, let $\text{Cone}_y : C_1^{\text{cell}}(Y; \mathbb{R}) \rightarrow C_2^{\text{cell}}(Y; \mathbb{R})$ be defined on generators by

$$\text{Cone}_y((g_0 H_0, g_1 H_1)) = (y, g_0 H_0, g_1, H_1).$$

Obviously Cone_y preserves the ℓ^1 -norms. For a G -CW-complex W , we denote the evaluation pairing by

$$\langle \cdot, \cdot \rangle_W : C_{\text{cell}}^*(W; V)^G \times C_*^{\text{cell}}(W; \mathbb{R}) \rightarrow V.$$

¹A detailed proof of this fact was communicated to us by Sam Hughes. It will appear in a forthcoming paper of his, joint with Martínez-Pedroza and Sánchez Saldaña.

Now, for $b \in C_1^{\text{cell}}(Z; \mathbb{R})$ and $c \in C_2^{\text{cell}}(Z; \mathbb{R})$ with $\partial_2(c) = b$, we find by (5) that

$$\begin{aligned} b &= \langle u, c \rangle_Z = \langle (\psi^2 \circ \varphi^2)(u) + \delta_Z^1(v), c \rangle_Z \\ &= \langle (\psi^2 \circ \varphi^2)(u), c \rangle_Z + \langle v, \partial_2^Y(c) \rangle_Z \\ &= \langle \varphi^2(u), \psi_2(c) \rangle_Y + \langle v, b \rangle_Z. \end{aligned}$$

Since $\varphi^2(u)$ is a cocycle and $\psi_2(c) - \text{Cone}_y(\partial_2^Y(\psi_2(c)))$ is a cycle and hence a boundary, we have

$$\begin{aligned} \langle \varphi^2(u), \psi_2(c) \rangle_Y &= \langle \varphi^2(u), \text{Cone}_y(\partial_2^Y(\psi_2(c))) \rangle_Y \\ &= \langle \varphi^2(u), \text{Cone}_y(\psi_1(b)) \rangle_Y \\ &= \langle u' + \delta_Y^1(v'), \text{Cone}_y(\psi_1(b)) \rangle_Y \\ &= \langle u', \text{Cone}_y(\psi_1(b)) \rangle_Y + \langle v', \partial_2^Y(\text{Cone}_y(\psi_1(b))) \rangle_Y \\ &= \langle u', \text{Cone}_y(\psi_1(b)) \rangle_Y + \langle v', \psi_1(b) \rangle_Y \\ &= \langle u', \text{Cone}_y(\psi_1(b)) \rangle_Y + \langle \psi^1(v'), b \rangle_Z, \end{aligned}$$

where we used (6). Together, we have

$$b = \langle u', \text{Cone}_y(\psi_1(b)) \rangle_Y + \langle \psi^1(v') + v, b \rangle_Z$$

and it follows that

$$\begin{aligned} \|b\|_{\partial} &\leq \|\langle u', \text{Cone}_y(\psi_1(b)) \rangle_Y\|_{\partial} + \|\langle \psi^1(v') + v, b \rangle_Z\|_{\partial} \\ &\leq \|u'\|_{\infty} \cdot \|\text{Cone}_y(\psi_1(b))\|_1 + \|\psi^1(v') + v\|_{\infty} \cdot \|b\|_1 \\ &= \|u'\|_{\infty} \cdot \|\psi_1(b)\|_1 + \|\psi^1(v') + v\|_{\infty} \cdot \|b\|_1 \\ &\leq (\|u'\|_{\infty} \cdot \|\psi_1\|_{\infty} + \|\psi^1(v') + v\|_{\infty}) \cdot \|b\|_1. \end{aligned}$$

Finally, u' is bounded by construction and so are ψ_1 and $\psi^1(v') + v$ because they are G -maps with domain $C_1^{\text{cell}}(Z; \mathbb{R})$ and Z has only finitely many orbits of 1-cells. Thus we have shown that Z satisfies a linear homological isoperimetric inequality over \mathbb{R} in degree 1. This finishes the proof. \square

Remark 5.5 (Groups with torsion). In Theorem 5.4, if the group G is not assumed to be torsionfree and \mathcal{H} is instead assumed to be almost malnormal, one can still prove the equivalence of (i) and (iii). However, a few modifications are necessary which we shall only outline.

Assuming that G is hyperbolic relative to \mathcal{H} , Mineyev–Yaman’s ideal complex has to be replaced by a Rips type construction X due to Martínez-Pedroza–Przytycki that is a model for $E_{\mathcal{F} \cup \mathcal{FIN}}G$. This complex X satisfies a linear homological isoperimetric inequality over \mathbb{Z} in degree 1 ([MPP19, Corollary 1.5]). It is part of a hyperbolic tuple in the sense of [MY, Definition 38] and hence admits a homological \mathbb{Q} -bicombing by [MY, Theorem 47]. Then one can construct a G -chain map φ_* with φ_2 bounded similarly as before and conclude surjectivity of the comparison map in degree 2 for the family $\mathcal{F} \cup \mathcal{FIN}$. This implies the same for the family \mathcal{F} over the ring \mathbb{R} .

For the converse implication, since \mathcal{H} is almost malnormal, there exists a model Z for $E_{\mathcal{F}}G$ such that for every infinite subgroup $H \in \mathcal{F}$ the fixed-point set Z^H consists of precisely one point. Then one shows as before that Z satisfies a linear homological isoperimetric inequality over \mathbb{R} in degree 1.

We do not know whether condition (ii) is equivalent to (i) and (iii) in this case.

6. MAPPING THEOREM FOR FAMILIES OF SUBGROUPS

In this section we prove the converse of a result by Löh–Sauer [LS20, Proposition 5.2] and give an application to vanishing results for the comparison map.

Recall the Johnson class $[J] := [J_{\mathcal{T}\mathcal{R}}] \in H_b^1(G; (\ell^\infty(G)/\mathbb{R})^\#)$ from Section 4, whose vanishing is equivalent to the amenability of G . The following lemma is part of the proof of [MR, Theorem 3.1.3].

Lemma 6.1 (Moraschini–Raptis). *Let H be a subgroup of a group G . If the image of the Johnson class under the restriction map*

$$\text{res}_H^G: H_b^1(G; (\ell^\infty(G)/\mathbb{R})^\#) \rightarrow H_b^1(H; (\ell^\infty(G)/\mathbb{R})^\#)$$

is trivial, then H is amenable.

For a family of subgroups \mathcal{F} , the up to G -homotopy unique G -map $EG \rightarrow E_{\mathcal{F}}G$ induces a canonical map on bounded cohomology

$$\text{can}_{\mathcal{F}}: H_{\mathcal{F},b}^n(G; V) \rightarrow H_b^n(G; V)$$

for any normed G -module V and all $n \geq 0$. Here and in the following we use Theorem 3.10 implicitly.

Lemma 6.2. *Let G be a group, \mathcal{F} be a family of subgroups, and V be a normed G -module. For each $H \in \mathcal{F}$, the composition*

$$\text{res}_H^G \circ \text{can}_{\mathcal{F}}: H_{\mathcal{F},b}^n(G; V) \rightarrow H_b^n(H; V)$$

vanishes for all $n \geq 1$.

Proof. Consider the following commutative diagram

$$\begin{array}{ccccc} H_{G,b}^n(E_{\mathcal{F}}G; V) & \xrightarrow{\text{can}_{\mathcal{F}}} & H_{G,b}^n(EG; V) & & \\ \downarrow & & \downarrow & \searrow^{\text{res}_H^G} & \\ H_{H,b}^n(E_{\mathcal{F}}G; V) & \longrightarrow & H_{H,b}^n(EG; V) & \xrightarrow{\cong} & H_{H,b}^n(EH; V), \end{array}$$

where the vertical maps are induced by viewing a G -space as an H -space. Observe that the lower left corner $H_{H,b}^n(E_{\mathcal{F}}G; V)$ vanishes for $n \geq 1$, since when viewed as an H -space $E_{\mathcal{F}}G$ is a model for $E_{\mathcal{A}\mathcal{L}\mathcal{L}}H$ and hence H -equivariantly contractible. \square

Note that Lemma 6.2 holds analogously for ordinary Bredon cohomology. We obtain a group-theoretic Mapping Theorem for families of subgroups.

Theorem 6.3. *Let G be a group and \mathcal{F} be a family of subgroups. The following are equivalent:*

- (i) *All subgroups in \mathcal{F} are amenable;*
- (ii) *The canonical map $H_{\mathcal{F},b}^n(G; V^\#) \rightarrow H_b^n(G; V^\#)$ is an isomorphism for any dual normed $\mathbb{R}G$ -module $V^\#$ and all $n \geq 1$;*
- (iii) *The canonical map $H_{\mathcal{F},b}^1(G; V^\#) \rightarrow H_b^1(G; V^\#)$ is an isomorphism for any dual normed $\mathbb{R}G$ -module $V^\#$;*
- (iv) *The Johnson class $[J] \in H_b^1(G; (\ell^\infty(G)/\mathbb{R})^\#)$ lies in the image of the canonical map $\text{can}_{\mathcal{F}}$.*

Proof. The fact that condition (i) implies (ii) was proved in [LS20, Proposition 5.2] for $\mathcal{F} = \mathcal{AM}\mathcal{E}$ and $V^\# = \mathbb{R}$. The same proof works for any subfamily $\mathcal{F} \subset \mathcal{AM}\mathcal{E}$ and any dual normed $\mathbb{R}G$ -module $V^\#$.

The implications (ii) \Rightarrow (iii) \Rightarrow (iv) are clear. Suppose that the Johnson class lies in the image of the canonical map $\text{can}_{\mathcal{F}}$. Then for each $H \in \mathcal{F}$, the image of the Johnson class under the restriction map res_H^G is trivial by Lemma 6.2 and hence H is amenable by Lemma 6.1. \square

Note that for $\mathcal{F} = \mathcal{F}\langle N \rangle$, where N is a normal subgroup of G , we recover [MR, Theorem 3.1.3] by Corollary 3.11. By considering $\mathcal{F} = \mathcal{AL}\mathcal{L}$, Theorem 6.3 can also be viewed as a generalization of Theorem 1.1.

Corollary 6.4. *Let X be a CW-complex with fundamental group G and \mathcal{F} be a family consisting of amenable subgroups of G . Suppose that there exists a model for $E_{\mathcal{F}}G$ whose orbit space $G \backslash E_{\mathcal{F}}G$ is homotopy equivalent to a k -dimensional CW-complex. Then the comparison map $c_X^n: H_b^n(X; \mathbb{R}) \rightarrow H^n(X; \mathbb{R})$ vanishes for all $n > k$.*

Proof. By Gromov's Mapping Theorem (see e.g. [Fri17, Theorem 5.9]), the comparison map c_X^n vanishes if the comparison map $c_{EG}^n: H_{G,b}^n(EG; \mathbb{R}) \rightarrow H_G^n(EG; \mathbb{R})$ vanishes. The G -map $EG \rightarrow E_{\mathcal{F}}G$ induces a commutative square

$$\begin{array}{ccc} H_{G,b}^n(EG; \mathbb{R}) & \xrightarrow{c_{EG}^n} & H_G^n(EG; \mathbb{R}) \\ \text{can}_{\mathcal{F}} \uparrow \cong & & \uparrow \\ H_{G,b}^n(E_{\mathcal{F}}G; \mathbb{R}) & \xrightarrow{c_{E_{\mathcal{F}}G}^n} & H_G^n(E_{\mathcal{F}}G; \mathbb{R}), \end{array}$$

where the canonical map $\text{can}_{\mathcal{F}}$ is an isomorphism by Theorem 6.3. Since we are considering trivial coefficients, the lower right corner can be identified with the (non-equivariant) cohomology of the orbit space

$$H_G^n(E_{\mathcal{F}}G; \mathbb{R}) \cong H^n(G \backslash E_{\mathcal{F}}G; \mathbb{R})$$

(see e.g. [Flu, Theorem 4.2]). This concludes the proof. \square

As an application of Corollary 6.4 we obtain the following well-known examples.

Example 6.5. The comparison map vanishes in all positive degrees for CW-complexes with the following fundamental groups:

- (i) Graph products of amenable groups (e.g. right-angled Artin groups);
- (ii) Fundamental groups of graphs of groups with amenable vertex groups.

Indeed, if G_Γ is a graph product of amenable groups, we consider the family \mathcal{F} generated by the vertex groups and direct products of vertex groups whenever the corresponding vertices form a clique in the underlying graph Γ . Then there exists a model for $E_{\mathcal{F}}(G_\Gamma)$ with contractible orbit space, which can be constructed by induction on the number of vertices of Γ .

If G is the fundamental group of a graph of groups with amenable vertex groups, we consider the family \mathcal{F} generated by the vertex groups. Then the Bass–Serre tree is a 1-dimensional model for $E_{\mathcal{F}}G$. Recall that the comparison map always vanishes in degree 1, since $H_b^1(G; \mathbb{R})$ is trivial for any group G .

Similar to Theorem 6.3, when considering all (not necessarily dual) normed $\mathbb{R}G$ -modules one obtains the following generalization of [MR, Theorem B] to families of subgroups. Recall the class $[K] := [K_{\mathcal{T}\mathcal{R}}] \in H_b^1(G, \ell_0^1(G))$ from Section 4, whose vanishing is equivalent to the finiteness of G .

Theorem 6.6. *Let G be a group and \mathcal{F} be a family of subgroups. The following are equivalent:*

- (i) *All subgroups in \mathcal{F} are finite;*
- (ii) *The canonical map $H_{\mathcal{F},b}^n(G; V) \rightarrow H_b^n(G; V)$ is an isomorphism for any normed $\mathbb{R}G$ -module V and all $n \geq 1$;*
- (iii) *The canonical map $H_{\mathcal{F},b}^1(G; V) \rightarrow H_b^1(G; V)$ is an isomorphism for any normed $\mathbb{R}G$ -module V ;*
- (iv) *The class $[K] \in H_b^1(G; (\ell_0^1(G)))$ lies in the image of the canonical map $\text{can}_{\mathcal{F}}$.*

Note that Theorem 6.6 can also be interpreted as a generalization of the characterization of finite groups [Fri17, Theorem 3.12] by considering $\mathcal{F} = \mathcal{ALL}$.

REFERENCES

- [ANCMSS] José Antonio Arciniega-Nevárez, José Luis Cisneros-Molina, and Luis Jorge Sánchez Saldaña. Relative group (co)homology theories with coefficients and the comparison homomorphism. Preprint, arXiv:1809.01209, 2018.
- [BE78] Robert Bieri and Beno Eckmann. Relative homology and Poincaré duality for group pairs. *J. Pure Appl. Algebra*, 13(3):277–319, 1978.
- [Bla16] Matthias Blank. Relative bounded cohomology for groupoids. *Geom. Dedicata*, 184:27–66, 2016.
- [Flu] Martin Fluch. On Bredon (co-)homological dimension of groups. PhD thesis, arXiv:1009.4633, 2010.
- [Fra18] Federico Franceschini. A characterization of relatively hyperbolic groups via bounded cohomology. *Groups Geom. Dyn.*, 12(3):919–960, 2018.
- [Fri17] Roberto Frigerio. *Bounded cohomology of discrete groups*, volume 227 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2017.
- [Gro82] Michael Gromov. Volume and bounded cohomology. *Inst. Hautes Études Sci. Publ. Math.*, (56):5–99, 1982.
- [Hru10] G. Christopher Hruska. Relative hyperbolicity and relative quasiconvexity for countable groups. *Algebr. Geom. Topol.*, 10(3):1807–1856, 2010.
- [Iva85] N. V. Ivanov. Foundations of the theory of bounded cohomology. In *Studies in topology*, V, volume 143, pages 69–109, 177–178. 1985.
- [Joh72] Barry Edward Johnson. *Cohomology in Banach algebras*. American Mathematical Society, Providence, R.I., 1972.
- [JOR12] Ronghui Ji, Crichton Ogle, and Bobby W. Ramsey. Relative property A and relative amenability for countable groups. *Adv. Math.*, 231(5):2734–2754, 2012.
- [LS20] Clara Löh and Roman Sauer. Bounded cohomology of amenable covers via classifying spaces. *Enseign. Math.*, 66(1-2):151–172, 2020.
- [Lüc05] Wolfgang Lück. Survey on classifying spaces for families of subgroups. In *Infinite groups: geometric, combinatorial and dynamical aspects*, volume 248 of *Progr. Math.*, pages 269–322. Birkhäuser, Basel, 2005.
- [Min01] Igor Mineyev. Straightening and bounded cohomology of hyperbolic groups. *Geom. Funct. Anal.*, 11(4):807–839, 2001.
- [Min02] Igor Mineyev. Bounded cohomology characterizes hyperbolic groups. *Q. J. Math.*, 53(1):59–73, 2002.
- [Mon01] Nicolas Monod. *Continuous bounded cohomology of locally compact groups*, volume 1758 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2001.
- [Mon06] Nicolas Monod. An invitation to bounded cohomology. In *International Congress of Mathematicians. Vol. II*, pages 1183–1211. Eur. Math. Soc., Zürich, 2006.
- [MP03] Nicolas Monod and Sorin Popa. On co-amenability for groups and von Neumann algebras. *C. R. Math. Acad. Sci. Soc. R. Can.*, 25(3):82–87, 2003.

- [MP16] Eduardo Martínez-Pedroza. A note on fine graphs and homological isoperimetric inequalities. *Canad. Math. Bull.*, 59(1):170–181, 2016.
- [MPP19] Eduardo Martínez-Pedroza and Piotr Przytycki. Dismantlable classifying space for the family of parabolic subgroups of a relatively hyperbolic group. *J. Inst. Math. Jussieu*, 18(2):329–345, 2019.
- [MR] Marco Moraschini and George Raptis. Amenability and acyclicity in bounded cohomology theory. Preprint, arXiv:2105.02821, 2021.
- [MY] Igor Mineyev and Asli Yaman. Relative hyperbolicity and bounded cohomology. Preprint, <https://faculty.math.illinois.edu/~mineyev/math/art/rel-hyp.pdf>, 2007.
- [Nuc99] Brita E. A. Nucinkis. Cohomology relative to a G -set and finiteness conditions. *Topology Appl.*, 92(2):153–171, 1999.

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SOUTHAMPTON, SOUTHAMPTON SO17 1BJ, UNITED KINGDOM

Email address: `kevin.li@soton.ac.uk`