

Normalized Laplacian spectra of central vertex join and central edge join of graphs

Jahfar T K *and Chithra A V †

Department of Mathematics, National Institute of Technology, Calicut, Kerala, India-673601

Abstract

In this paper, we compute normalized Laplacian spectra of central graph of a regular graph, central vertex join and central edge join of two regular graphs. Also, we determine their Kemeny's constant and degree Kirchhoff index.

AMS classification: 05C50

Keywords: *Central graph, normalized Laplacian spectrum, central vertex join, central edge join, Kirchhoff index, Kemeny's constant.*

1 Introduction

Let $G = (V(G), E(G))$ be a simple undirected graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and its edge set $E(G) = \{e_1, e_2, \dots, e_m\}$. The degree d_i of the vertex $v_i \in V(G)$ is the number of vertices adjacent to v_i . The adjacency matrix $A(G) = [a_{ij}]$ of the graph G is a square symmetric matrix of order n whose $(i, j)^{th}$ entry is defined as,

$$a_{ij} = \begin{cases} 1, & \text{if } v_i \text{ and } v_j \text{ are adjacent in } G, \\ 0, & \text{otherwise.} \end{cases}$$

The Laplacian matrix of G , denoted by $L(G)$ is defined as $L(G) = D(G) - A(G)$, where $D(G)$ is the diagonal matrix with vertex degrees. The characteristic polynomial of the $n \times n$ matrix M of G is defined as $f(M, x) = |xI_n - M|$, where I_n is the identity matrix of order n . Eigenvalues of $A(G)$ are denoted by $\lambda_1(G) \geq \lambda_2(G) \geq \dots \geq \lambda_n(G)$. Eigenvalues of $A(G)$ together with their multiplicities are called the spectrum of G . The normalized Laplacian matrix $\mathcal{L}(G) = (\mathcal{L}_{ij})$ of G is a square symmetric matrix of order n whose $(i, j)^{th}$ entry is defined as,

$$\mathcal{L}_{ij} = \begin{cases} 1, & \text{if } v_i = v_j \text{ and } d_i \neq 0, \\ -\frac{1}{\sqrt{d_i d_j}}, & \text{if } v_i \text{ and } v_j \text{ are adjacent in } G, \\ 0, & \text{otherwise.} \end{cases}$$

*jahfartk@gmail.com

†chithra@nitc.ac.in

The normalized Laplacian matrix $\mathcal{L}(G)$ of G is defined as

$$\mathcal{L}(G) = I_n - D^{-\frac{1}{2}}(G)A(G)D^{-\frac{1}{2}}(G).$$

The eigenvalues of the matrix $\mathcal{L}(G)$ are called the normalized Laplacian eigenvalues of G , and will be denoted by $0 = \tilde{\mu}_1(G) < \tilde{\mu}_2(G) \leq \dots \leq \tilde{\mu}_n(G)$. The eigenvalues of $\mathcal{L}(G)$ together with their multiplicities are called the normalized Laplacian spectrum of G . If G is an r -regular graph, then the adjacency and the normalized Laplacian eigenvalues of G are related by $\tilde{\mu}_i(G) = 1 - \frac{\lambda_i(G)}{r}$ [1, 2]. Two graphs are said to be \mathcal{L} -cospectral, if they have the same \mathcal{L} -spectrum. Otherwise, they are non \mathcal{L} -cospectral graphs. In 1993, Klein and Randić [3] introduced the resistance distance between two vertices v_i and v_j in G , denoted by $r_{i,j}^*$, is defined as the effective resistance between them when unit resistors are distributed on every edge of G . The degree Kirchoff index of a connected graph G is defined by Chen et al. in [4] as

$$Kf^*(G) = \sum_{i < j} d_i r_{i,j}^* d_j,$$

where $r_{i,j}^*$ denotes the resistance distance between vertices v_i and v_j in G . In [4], the authors proved that

$$Kf^*(G) = 2m \sum_{i=2}^n \frac{1}{\tilde{\mu}_i(G)}$$

The Kemeny's constant $K(G)$ of a graph G [5, 6] is defined as the expected number of steps from a randomly chosen starting point in the graph to a randomly chosen destination point in the graph. It is defined in terms of the normalized Laplacian eigenvalues of G as

$$K(G) = \sum_{i=2}^n \frac{1}{\tilde{\mu}_i(G)}.$$

The various application of Kemeny's constant to perturbed Markov chains, random walks on directed graphs are studied in [6]. We recall that for two matrices A and B , of the same size, the Hadamard product $A \circ B$ of A and B is a matrix of the same size with entries given by $(A \circ B)_{ij} = (A)_{ij} \cdot (B)_{ij}$ (that is entry wise multiplication) [7]. The incidence matrix of G , $I(G)$ is the $n \times m$ matrix whose $(i, j)^{th}$ entry is 1 if v_i is incident to e_j and 0 otherwise. It is known [8] that, $I(G)I(G)^T = A(G) + D(G)$. In particular, G is an r -regular graph, then $I(G)I(G)^T = A(G) + rI_n$. Also, $I(G)I(G)^T = 2rI_n - r\mathcal{L}(G)$ [9]. The adjacency matrix of the complement of a graph G is $A(\bar{G}) = J_n - I_n - A(G)$. Throughout the paper for any positive integer m , I_m denotes the $m \times m$ identity matrix, J_m denotes the $m \times m$ matrix whose all entries equal to one and $J_{m \times 1}$ denotes the column vector of size m with all entries equal to one. For any positive integers p and q , $O_{p \times q}$ denotes the zero matrix of size $p \times q$ and $K_{p,p}$ is the complete bipartite graph with $2p$ vertices.

Definition 1.1. [10] *Let G be a simple graph with n vertices and m edges. The central graph of G , denoted by $C(G)$ is obtained by subdividing each edge of G exactly once and joining all the nonadjacent vertices in G .*

The number of vertices and edges in $C(G)$ are $m + n$ and $m + \frac{n(n-1)}{2}$ respectively.

The rest of the paper is organized as follows. In Section 2, we present necessary definitions and known results, which will be used later. In Section 3, we calculate the normalized Laplacian spectrum of central graph of regular graph. In Section 4, normalized Laplacian spectra of central vertex join and central edge join of two regular graphs are obtained. In Section 5, some \mathcal{L} -cospectral non-regular graphs are constructed. Also, we investigate some graph invariants like the degree Kirchoff index and the Kemeny's constant of central vertex join and central edge join of two regular graphs.

2 Preliminaries

In this section, some definitions and results are given which will be useful to prove our main results.

Lemma 2.1. [8] *Let U, V, W and X be matrices with U invertible. Let*

$$S = \begin{pmatrix} U & V \\ W & X \end{pmatrix}.$$

Then $\det(S) = \det(U)\det(X - WU^{-1}V)$.

If X is invertible, then $\det(S) = \det(X)\det(U - VX^{-1}W)$. If U and W are commutes, then $\det(S) = \det(UX - WV)$.

Lemma 2.2. [8] *Let G be a connected r -regular graph on n vertices with adjacency matrix A having t distinct eigenvalues $r = \lambda_1, \lambda_2, \dots, \lambda_t$. Then there exists a polynomial*

$$P(x) = n \frac{(x - \lambda_2)(x - \lambda_3) \dots (x - \lambda_t)}{(r - \lambda_2)(r - \lambda_3) \dots (r - \lambda_t)}.$$

such that $P(A) = J_n, P(r) = n$ and $P(\lambda_i) = 0$ for $\lambda_i \neq r$.

Definition 2.1. [11] *The M -coronal $\chi_M(x)$ of $n \times n$ matrix M is defined as the sum of the entries of the matrix $(xI_n - M)^{-1}$ (if exists), that is,*

$$\chi_M(x) = J_{n \times 1}^T (xI_n - M)^{-1} J_{n \times 1}.$$

Throughout this, we use the notation $\chi_M(x, \alpha) = J_{n \times 1}^T (xI_n - M \circ (\alpha J_n + (1 - \alpha)I_n))^{-1} J_{n \times 1}$, as in [12], if $\alpha = 1$, then $\chi_M(x, 1)$ is the usual M -coronal $\chi_M(x)$.

Let G be an r -regular graph of order n . Then $\chi_{\mathcal{L}(G)}(x, \alpha) = \frac{n}{x + \alpha - 1}$ [12].

Lemma 2.3. [13] *Let G be an r -regular graph on n vertices, then $\chi_A(x) = \frac{n}{x - r}$.*

For Laplacian matrix each row sum is zero, so $\chi_L(x) = \frac{n}{x}$.
Note that $\chi_{kL}(x) = \chi_L(x)$ for any real number k .

Lemma 2.4. [14] *Let A be an $n \times n$ real matrix. Then $\det(A + \alpha J_n) = \det(A) + \alpha J_{n \times 1}^T \text{adj}(A) J_{n \times 1}$, where α is a real number and $\text{adj}(A)$ is the adjoint of A .*

Corollary 2.5. [14] *Let A be an $n \times n$ real matrix. Then*

$$\det(xI_n - A - \alpha J_n) = (1 - \alpha \chi_A(x)) \det(xI_n - A).$$

Lemma 2.6. [11] For any real numbers $c, d > 0$, $(cI_n - dJ_n)^{-1} = \frac{1}{c}I_n + \frac{d}{c(c-nd)}J_n$.

Definition 2.2. [15] Let G_1 and G_2 be any two graphs on n_1, n_2 vertices and m_1, m_2 edges respectively. The central vertex join of G_1 and G_2 is the graph $G_1 \dot{\vee} G_2$ is obtained from $C(G_1)$ and G_2 by joining each vertex of G_1 with every vertex of G_2 .

The graph $G_1 \dot{\vee} G_2$ has $m_1 + n_1 + n_2$ vertices and $m_1 + m_2 + n_1n_2 + \frac{n_1(n_1-1)}{2}$ edges.

Definition 2.3. [15] Let G_i be any two graphs with n_i vertices and m_i edges respectively for $i = 1, 2$. Then the central edge join of two graphs G_1 and G_2 is the graph $G_1 \vee G_2$ is obtained from $C(G_1)$ and G_2 by joining each vertex corresponding to edges of G_1 with every vertex of G_2 .

The graph $G_1 \vee G_2$ has $m_1 + n_1 + n_2$ vertices and $m_1 + m_2 + m_1n_2 + \frac{n_1(n_1-1)}{2}$ edges.

3 Normalized Laplacian spectra of central graphs

In this section, we determine the normalized Laplacian spectrum of central graph of regular graphs.

Theorem 3.1. Let G be an r -regular graph with n vertices. Then the normalized Laplacian characteristic polynomial of central graph of G is

$$f(\mathcal{L}(C(G)), x) = (x-1)^{m-n} \prod_{i=1}^n \left((x-1) \left((x-1) - \frac{P(\lambda_i) - 1 - \lambda_i}{n-1} \right) - \frac{(\lambda_i + r)}{2(n-1)} \right).$$

Proof. Let $I(G)$ be the incidence matrix of G and $A(\bar{G})$ be the adjacency matrix of complement of graph G . Then the adjacency matrix of $C(G)$ is

$$A(C(G)) = \begin{pmatrix} A(\bar{G}) & I(G) \\ I(G)^T & O_{m \times m} \end{pmatrix}.$$

The degree matrix of $C(G)$ is

$$D(C(G)) = \begin{pmatrix} (n-1)I_n & O_{n \times m} \\ O_{m \times n} & 2I_m \end{pmatrix}.$$

Then the normalized Laplacian matrix of $C(G)$ is

$$\mathcal{L}(C(G)) = I_{m+n} - D^{-\frac{1}{2}}(C(G))A(C(G))D^{-\frac{1}{2}}(C(G)) = \begin{pmatrix} I_n - \frac{A(\bar{G})}{n-1} & -\frac{I(G)}{\sqrt{2(n-1)}} \\ -\frac{I(G)^T}{\sqrt{2(n-1)}} & I_m \end{pmatrix}.$$

Using Lemmas 2.1 and 2.2, we have the normalized Laplacian characteristic polynomial

of $C(G)$ is

$$\begin{aligned}
f(\mathcal{L}(C(G)), x) &= \det \begin{pmatrix} (x-1)I_n + \frac{J_n - I_n - A(G)}{n-1} & \frac{I(G)}{\sqrt{2(n-1)}} \\ \frac{I(G)^T}{\sqrt{2(n-1)}} & (x-1)I_m \end{pmatrix} \\
&= (x-1)^m \det \left((x-1)I_n + \frac{J_n - I_n - A(G)}{n-1} - \frac{I(G)I(G)^T}{2(x-1)(n-1)} \right) \\
&= (x-1)^{m-n} \det \left((x-1) \left((x-1)I_n + \frac{J_n - I_n - A(G)}{n-1} \right) - \frac{I(G)I(G)^T}{2(n-1)} \right) \\
&= (x-1)^{m-n} \det \left((x-1) \left((x-1)I_n + \frac{J_n - I_n - A(G)}{n-1} \right) - \frac{(A(G) + rI_n)}{2(n-1)} \right) \\
f(\mathcal{L}(C(G)), x) &= (x-1)^{m-n} \prod_{i=1}^n \left((x-1) \left((x-1) + \frac{P(\lambda_i) - 1 - \lambda_i}{n-1} \right) - \frac{(\lambda_i + r)}{2(n-1)} \right).
\end{aligned}$$

□

The following corollary describes the complete normalized Laplacian spectrum of $C(G)$ when G is regular graph.

Corollary 3.2. *Let G be an r -regular graph on n vertices. Then the normalized Laplacian spectrum of $C(G)$ consists of*

1. 1 with multiplicity $m - n$.
2. Two roots of the equation $(x-1) \left((x-1) + \frac{n-1-r}{n-1} \right) - \frac{r}{(n-1)} = 0$.
3. Two roots of the equation $(x-1) \left((x-1) + \frac{-1-\lambda_i}{n-1} \right) - \frac{(\lambda_i+r)}{2(n-1)} = 0$ for $i = 2, \dots, n$.

Example 3.1. Let $G = K_{p,p}$. Then the adjacency eigenvalues of G are 0 (multiplicity $2p - 2$) and $\pm p$. Therefore, the normalized Laplacian eigenvalues of $C(G)$ are 0, $\frac{3p-1}{2p-1}, 1$ (multiplicity $p^2 - 2p$), roots of the equation $(2p-1)x^2 + (1-3p)x + p = 0$ and roots of the equation $(4p-2)x^2 - (8p-2)x - 3p = 0$ (each root with multiplicity $2p - 2$).

In particular, $G = K_{3,3}$. The normalized Laplacian eigenvalues of $C(G)$ are 0, $\frac{8}{5}, 1$ (multiplicity 3), roots of the equation $5x^2 - 8x + 3 = 0$ and roots of the equation $10x^2 - 22x + 9 = 0$ (each root with multiplicity 4).

4 Normalized Laplacian spectra of central vertex join and central edge join of Graphs

In this section, we determine the normalized Laplacian spectrum of central vertex join and central edge join of two regular graphs.

Theorem 4.1. *Let G_i be an r_i -regular graph with n_i vertices and m_i edges for $i=1,2$.*

Then the normalized Laplacian characteristic polynomial of $G_1 \dot{\vee} G_2$ is given by

$$\begin{aligned}
& f(\mathcal{L}(G_1 \dot{\vee} G_2), x) \\
&= (x-1)^{m_1-n_1} \prod_{i=2}^{n_2} \left(\left(x - \frac{n_1}{r_2+n_1} \right) I_{n_2} - \frac{r_2}{r_2+n_1} \tilde{\mu}_i(G_2) \right) \\
& \times \left[\left(x - \frac{n_1}{r_2+n_1} \right) \left((x-1)^2 - \frac{r_1}{(n_1+n_2-1)} - \frac{(1+r_1)(x-1)}{n_1+n_2-1} \right) \right. \\
& \quad \left. - \frac{n_1 n_2 (x-1) - n_1 (x-1)(x n_1 + x r_2 - n_1)}{(n_1+n_2-1)(n_1+r_2)} \right] \\
& \times \prod_{i=2}^{n_1} \left((x-1)^2 - \frac{r_1}{(n_1+n_2-1)} - \frac{(1+r_1)(x-1)}{n_1+n_2-1} + \frac{r_1 \tilde{\mu}_i(G_1)}{2(n_1+n_2-1)} + \frac{r_1(x-1) \tilde{\mu}_i(G_1)}{n_1+n_2-1} \right).
\end{aligned}$$

Proof. Let $I(G_1)$ be the incidence matrix of G_1 . Then by a proper labeling of vertices, the adjacency matrix of $G_1 \dot{\vee} G_2$ can be written as

$$A(G_1 \dot{\vee} G_2) = \begin{pmatrix} A(\bar{G}_1) & I(G_1) & J_{n_1 \times n_2} \\ I(G_1)^T & O_{m_1 \times m_1} & O_{m_1 \times n_2} \\ J_{n_2 \times n_1} & O_{n_2 \times m_1} & A(G_2) \end{pmatrix}.$$

The degree matrix of $G_1 \dot{\vee} G_2$ is

$$D(G_1 \dot{\vee} G_2) = \begin{pmatrix} (n_1+n_2-1)I_{n_1} & O_{n_1 \times m_1} & O_{n_1 \times n_2} \\ O_{m_1 \times n_1} & 2I_{m_1} & O_{m_1 \times n_2} \\ O_{n_2 \times n_1} & O_{n_2 \times m_1} & (r_2+n_1)I_{n_2} \end{pmatrix}.$$

$$D^{-\frac{1}{2}}(G_1 \dot{\vee} G_2) = \begin{pmatrix} \frac{I_{n_1}}{\sqrt{(n_1+n_2-1)}} & O_{n_1 \times m_1} & O_{n_1 \times n_2} \\ O_{m_1 \times n_1} & \frac{I_{m_1}}{\sqrt{2}} & O_{m_1 \times n_2} \\ O_{n_2 \times n_1} & O_{n_2 \times m_1} & \frac{I_{n_2}}{\sqrt{(r_2+n_1)}} \end{pmatrix}.$$

Then the normalized Laplacian matrix of $G_1 \dot{\vee} G_2$ is

$$\mathcal{L}(G_1 \dot{\vee} G_2) = \begin{pmatrix} I_{n_1} - \frac{A(\bar{G}_1)}{n_1+n_2-1} & -\frac{I(G_1)}{\sqrt{2(n_1+n_2-1)}} & -\frac{J_{n_1 \times n_2}}{\sqrt{(n_1+n_2-1)(n_1+r_2)}} \\ -\frac{I(G_1)^T}{\sqrt{2(n_1+n_2-1)}} & I_{m_1} & O_{m_1 \times n_2} \\ -\frac{J_{n_2 \times n_1}}{\sqrt{(n_1+n_2-1)(n_1+r_2)}} & O_{n_2 \times m_1} & I_{n_2} - \frac{A(G_2)}{(n_1+r_2)} \end{pmatrix}$$

$$\mathcal{L}(G_1 \dot{\vee} G_2) = \begin{pmatrix} I_{n_1} - \frac{A(\bar{G}_1)}{n_1+n_2-1} & -\frac{I(G_1)}{\sqrt{2(n_1+n_2-1)}} & -\frac{J_{n_1 \times n_2}}{\sqrt{(n_1+n_2-1)(n_1+r_2)}} \\ -\frac{I(G_1)^T}{\sqrt{2(n_1+n_2-1)}} & I_{m_1} & O_{m_1 \times n_2} \\ -\frac{J_{n_2 \times n_1}}{\sqrt{(n_1+n_2-1)(n_1+r_2)}} & O_{n_2 \times m_1} & \mathcal{L}(G_2) \circ B \end{pmatrix},$$

where $B = \frac{r_2}{r_2+n_1}J_{n_2} + \frac{n_1}{r_2+n_1}I_{n_2}$.

Using Definition 2.1, Lemmas 2.1, 2.2, 2.3 and Corollary 2.5, we have the normalized Laplacian characteristic polynomial of $G_1 \dot{\vee} G_2$ is

$$\begin{aligned} f(\mathcal{L}(G_1 \dot{\vee} G_2), x) &= \det \begin{pmatrix} (x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1+n_2-1} & \frac{I(G_1)}{\sqrt{2(n_1+n_2-1)}} & \frac{J_{n_1 \times n_2}}{\sqrt{(n_1+n_2-1)(n_1+r_2)}} \\ \frac{I(G_1)^T}{\sqrt{2(n_1+n_2-1)}} & (x-1)I_{m_1} & O_{m_1 \times n_2} \\ \frac{J_{n_2 \times n_1}}{\sqrt{(n_1+n_2-1)(n_1+r_2)}} & O_{n_2 \times m_1} & xI_{n_2} - (\mathcal{L}(G_2) \circ B) \end{pmatrix} \\ &= \det \left(xI_{n_2} - (\mathcal{L}(G_2) \circ B) \right) \det S, \end{aligned}$$

$$\begin{aligned} \text{where } S &= \begin{pmatrix} (x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1+n_2-1} & \frac{I(G_1)}{\sqrt{2(n_1+n_2-1)}} \\ \frac{I(G_1)^T}{\sqrt{2(n_1+n_2-1)}} & (x-1)I_{m_1} \end{pmatrix} - \left[\begin{pmatrix} \frac{J_{n_1 \times n_2}}{\sqrt{(n_1+n_2-1)(n_1+r_2)}} \\ O_{m_1 \times n_2} \end{pmatrix} (xI_{n_2} - (\mathcal{L}(G_2) \circ B))^{-1} \right. \\ &\quad \left. \times \begin{pmatrix} \frac{J_{n_2 \times n_1}}{\sqrt{(n_1+n_2-1)(n_1+r_2)}} & O_{n_2 \times m_1} \end{pmatrix} \right] \\ &= \begin{pmatrix} (x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1+n_2-1} - \frac{J_{n_1 \times n_2}}{\sqrt{(n_1+n_2-1)(n_1+r_2)}} (xI_{n_2} - (\mathcal{L}(G_2) \circ B))^{-1} \frac{J_{n_2 \times n_1}}{\sqrt{(n_1+n_2-1)(n_1+r_2)}} & \frac{I(G_1)}{\sqrt{2(n_1+n_2-1)}} \\ \frac{I(G_1)^T}{\sqrt{2(n_1+n_2-1)}} & (x-1)I_{m_1} \end{pmatrix} \\ &= \begin{pmatrix} (x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1+n_2-1} - \frac{\chi_{\mathcal{L}(G_2)}(x, \frac{r_2}{r_2+n_1})}{(n_1+n_2-1)(n_1+r_2)} J_{n_1} & \frac{I(G_1)}{\sqrt{2(n_1+n_2-1)}} \\ \frac{I(G_1)^T}{\sqrt{2(n_1+n_2-1)}} & (x-1)I_{m_1} \end{pmatrix}. \end{aligned}$$

Therefore,

$$\begin{aligned} \det(S) &= \det \begin{pmatrix} (x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1+n_2-1} - \frac{\chi_{\mathcal{L}(G_2)}(x, \frac{r_2}{r_2+n_1})}{(n_1+n_2-1)(n_1+r_2)} J_{n_1} & \frac{I(G_1)}{\sqrt{2(n_1+n_2-1)}} \\ \frac{I(G_1)^T}{\sqrt{2(n_1+n_2-1)}} & (x-1)I_{m_1} \end{pmatrix} \\ &= (x-1)^{m_1} \det \left((x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1+n_2-1} - \frac{\chi_{\mathcal{L}(G_2)}(x, \frac{r_2}{r_2+n_1})}{(n_1+n_2-1)(n_1+r_2)} J_{n_1} - \frac{I(G_1)I(G_1)^T}{2(n_1+n_2-1)(x-1)} \right) \\ &= (x-1)^{m_1} \det \left((x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1+n_2-1} - \frac{\chi_{\mathcal{L}(G_2)}(x, \frac{r_2}{r_2+n_1})}{(n_1+n_2-1)(n_1+r_2)} J_{n_1} \right. \\ &\quad \left. - \frac{r_1}{(n_1+n_2-1)(x-1)} I_{n_1} + \frac{r_1 \mathcal{L}(G_1)}{2(n_1+n_2-1)(x-1)} \right) \end{aligned}$$

$$\begin{aligned}
&= (x-1)^{m_1} \det \left((x-1)I_{n_1} + \frac{J_{n_1} - I_{n_1} - A(G_1)}{n_1 + n_2 - 1} - \frac{\chi_{\mathcal{L}(G_2)}\left(x, \frac{r_2}{r_2+n_1}\right)}{(n_1 + n_2 - 1)(n_1 + r_2)} J_{n_1} \right. \\
&\quad \left. - \frac{r_1}{(n_1 + n_2 - 1)(x-1)} I_{n_1} + \frac{r_1 \mathcal{L}(G_1)}{2(n_1 + n_2 - 1)(x-1)} \right) \\
&= (x-1)^{m_1} \det \left((x-1)I_{n_1} + \frac{J_{n_1} - I_{n_1} - r_1 I_{n_1} + r_1 \mathcal{L}(G_1)}{n_1 + n_2 - 1} - \frac{\chi_{\mathcal{L}(G_2)}\left(x, \frac{r_2}{r_2+n_1}\right)}{(n_1 + n_2 - 1)(n_1 + r_2)} J_{n_1} \right. \\
&\quad \left. - \frac{r_1}{(n_1 + n_2 - 1)(x-1)} I_{n_1} + \frac{r_1 \mathcal{L}(G_1)}{2(n_1 + n_2 - 1)(x-1)} \right) \\
&= (x-1)^{m_1} \left(1 - \left(\frac{\chi_{\mathcal{L}(G_2)}\left(x, \frac{r_2}{r_2+n_1}\right)}{(n_1 + n_2 - 1)(n_1 + r_2)} - \frac{1}{n_1 + n_2 - 1} \right) \right. \\
&\quad \left. \chi_{k\mathcal{L}(G_1)} \left(x - 1 - \frac{r_1}{(n_1 + n_2 - 1)(x-1)} - \frac{1+r_1}{n_1 + n_2 - 1} \right) \right) \\
&\quad \times \det \left(\left(x - 1 - \frac{r_1}{(n_1 + n_2 - 1)(x-1)} - \frac{1+r_1}{n_1 + n_2 - 1} \right) I_{n_1} + \frac{r_1 \mathcal{L}(G_1)}{2(n_1 + n_2 - 1)(x-1)} + \frac{r_1 \mathcal{L}(G_1)}{n_1 + n_2 - 1} \right)
\end{aligned}$$

where $k = -\frac{r_1}{2(n_1+n_2-1)(x-1)} - \frac{r_1}{n_1+n_2-1}$.

It is easy to see that

$$\begin{aligned}
\det \left(xI_{n_2} - (\mathcal{L}(G_2) \circ B) \right) &= \det \left(\left(x - \frac{n_1}{r_2 + n_1} \right) I_{n_2} - \frac{r_2}{r_2 + n_1} \mathcal{L}(G_2) \right). \\
\chi_{\mathcal{L}(G_2)} \left(x, \frac{r_2}{r_2 + n_1} \right) &= \frac{n_2}{x - 1 + \frac{r_2}{r_2+n_1}}. \\
\chi_{k\mathcal{L}(G_1)} \left(x - 1 - \frac{r_1}{(n_1 + n_2 - 1)(x-1)} - \frac{1+r_1}{n_1 + n_2 - 1} \right) &= \frac{n_1}{x - 1 - \frac{r_1}{(n_1+n_2-1)(x-1)} - \frac{1+r_1}{n_1+n_2-1}}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
\det(S) &= (x-1)^{m_1-n_1} \left(1 - \left(\frac{\chi_{\mathcal{L}(G_2)}(x, \frac{r_2}{r_2+n_1})}{(n_1+n_2-1)(n_1+r_2)} - \frac{1}{n_1+n_2-1} \right) \right. \\
&\quad \times \left. \left(\frac{n_1}{x-1 - \frac{r_1}{(n_1+n_2-1)(x-1)} - \frac{1+r_1}{n_1+n_2-1}} \right) \right) \\
&\quad \times \det \left(\left((x-1)^2 - \frac{r_1}{(n_1+n_2-1)} - \frac{(1+r_1)(x-1)}{n_1+n_2-1} \right) I_{n_1} + \frac{r_1 \mathcal{L}(G_1)}{2(n_1+n_2-1)} + \frac{r_1(x-1) \mathcal{L}(G_1)}{n_1+n_2-1} \right). \\
f(\mathcal{L}(G_1 \dot{\vee} G_2), x) &= \det \left(x I_{n_2} - (\mathcal{L}(G_2) \circ B) \right) \det S \\
&= \det \left(\left(x - \frac{n_1}{r_2+n_1} \right) I_{n_2} - \frac{r_2}{r_2+n_1} \mathcal{L}(G_2) \right) \\
&\quad \times (x-1)^{m_1-n_1} \left[1 - \left(\frac{\frac{n_2}{x-1 + \frac{r_2}{r_2+n_1}}}{(n_1+n_2-1)(n_1+r_2)} - \frac{1}{n_1+n_2-1} \right) \right. \\
&\quad \quad \quad \times \left. \left(\frac{n_1}{x-1 - \frac{r_1}{(n_1+n_2-1)(x-1)} - \frac{1+r_1}{n_1+n_2-1}} \right) \right. \\
&\quad \times \left. \det \left((x-1)^2 - \frac{r_1}{(n_1+n_2-1)} - \frac{(1+r_1)(x-1)}{n_1+n_2-1} + \frac{r_1 \mathcal{L}(G_1)}{2(n_1+n_2-1)} + \frac{r_1(x-1) \mathcal{L}(G_1)}{n_1+n_2-1} \right) \right] \\
&= \prod_{i=1}^{n_2} \left(\left(x - \frac{n_1}{r_2+n_1} \right) I_{n_2} - \frac{r_2}{r_2+n_1} \tilde{\mu}_i(G_1) \right) \\
&\quad \times (x-1)^{m_1-n_1} \left[1 - \left(\frac{n_1 n_2 (x-1) - n_1 (x-1) (x n_1 + x r_2 - n_1)}{\left(x-1 + \frac{r_2}{r_2+n_1} \right) (n_1+n_2-1) (n_1+r_2) \left((x-1)^2 - \frac{r_1}{(n_1+n_2-1)} - \frac{(1+r_1)(x-1)}{n_1+n_2-1} \right)} \right) \right] \\
&\quad \prod_{i=1}^{n_1} \left((x-1)^2 - \frac{r_1}{(n_1+n_2-1)} - \frac{(1+r_1)(x-1)}{n_1+n_2-1} + \frac{r_1 \tilde{\mu}_i(G_1)}{2(n_1+n_2-1)} + \frac{r_1(x-1) \tilde{\mu}_i(G_1)}{n_1+n_2-1} \right) \\
&= (x-1)^{m_1-n_1} \prod_{i=2}^{n_2} \left(\left(x - \frac{n_1}{r_2+n_1} \right) I_{n_2} - \frac{r_2}{r_2+n_1} \tilde{\mu}_i(G_2) \right) \\
&\quad \times \left[\left(x-1 + \frac{r_2}{r_2+n_1} \right) \left((x-1)^2 - \frac{r_1}{(n_1+n_2-1)} - \frac{(1+r_1)(x-1)}{n_1+n_2-1} \right) \right. \\
&\quad \quad \quad \left. - \frac{n_1 n_2 (x-1) - n_1 (x-1) (x n_1 + x r_2 - n_1)}{(n_1+n_2-1)(n_1+r_2)} \right] \\
&\quad \prod_{i=2}^{n_1} \left((x-1)^2 - \frac{r_1}{(n_1+n_2-1)} - \frac{(1+r_1)(x-1)}{n_1+n_2-1} + \frac{r_1 \tilde{\mu}_i(G_1)}{2(n_1+n_2-1)} + \frac{r_1(x-1) \tilde{\mu}_i(G_1)}{n_1+n_2-1} \right)
\end{aligned}$$

$$\begin{aligned}
&= (x-1)^{m_1-n_1} \prod_{i=2}^{n_2} \left(\left(x - \frac{n_1}{r_2+n_1} \right) I_{n_2} - \frac{r_2}{r_2+n_1} \tilde{\mu}_i(G_2) \right) \\
&\quad \times \left[\left(x - \frac{n_1}{r_2+n_1} \right) \left((x-1)^2 - \frac{r_1}{(n_1+n_2-1)} - \frac{(1+r_1)(x-1)}{n_1+n_2-1} \right) \right. \\
&\quad \quad \left. - \frac{n_1 n_2 (x-1) - n_1 (x-1)(x n_1 + x r_2 - n_1)}{(n_1+n_2-1)(n_1+r_2)} \right] \\
&\quad \prod_{i=2}^{n_1} \left((x-1)^2 - \frac{r_1}{(n_1+n_2-1)} - \frac{(1+r_1)(x-1)}{n_1+n_2-1} + \frac{r_1 \tilde{\mu}_i(G_1)}{2(n_1+n_2-1)} + \frac{r_1(x-1) \tilde{\mu}_i(G_1)}{n_1+n_2-1} \right).
\end{aligned}$$

□

The following corollary describes the complete normalized Laplacian spectrum of $G_1 \dot{\vee} G_2$ when G_1 and G_2 are regular graphs.

Corollary 4.2. *Let G_i be an r_i -regular graph with n_i vertices and m_i edges for $i=1,2$. Then the normalized Laplacian spectrum of $G_1 \dot{\vee} G_2$ consists of*

1. 1 repeated $m_1 - n_1$ times.
2. Roots of the equation,

$$\left(x - \frac{n_1}{r_2+n_1} \right) - \frac{r_2}{r_2+n_1} \tilde{\mu}_i(G_2) = 0 \text{ for } i = 2, \dots, n_2.$$

3. Three roots of the equation,

$$\begin{aligned}
&\left(x - \frac{n_1}{r_2+n_1} \right) \left((x-1)^2 - \frac{r_1}{n_1+n_2-1} - \frac{(1+r_1)(x-1)}{n_1+n_2-1} \right) \\
&- \left(\frac{n_1 n_2 (x-1) - n_1 (x-1)(x n_1 + x r_2 - n_1)}{(n_1+n_2-1)(n_1+r_2)} \right) = 0.
\end{aligned}$$

4. Two roots of the equation,

$$(x-1)^2 - \frac{r_1}{(n_1+n_2-1)} - \frac{(1+r_1)(x-1)}{n_1+n_2-1} + \frac{r_1 \tilde{\mu}_i(G_1)}{2(n_1+n_2-1)} + \frac{r_1(x-1) \tilde{\mu}_i(G_1)}{n_1+n_2-1} = 0 \text{ for } i = 2, \dots, n_1.$$

Example 4.1. Let $G_1 = K_{p,p}$ and $G_2 = K_2$. Then the normalized Laplacian eigenvalues of G_1 are 0, 1(multiplicity $2p-2$) and 2. The normalized Laplacian eigenvalues of G_2 are 0 and 2. Therefore, the normalized Laplacian eigenvalues of $G_1 \dot{\vee} G_2$ are 0, $\frac{2p+2}{2p+1}, 1$ (multiplicity $p^2 - 2p$), roots of the equation $(4p^2 + 4p + 1)x^2 - (10p^2 + 11p + 3)x + (6p^2 + 6p + 2) = 0$, roots of the equation $(4p+2)x^2 - (6p+6)x + 2p+4 = 0$ and roots of the equation $(4p+2)x^2 - (8p+6)x + 3p+4 = 0$ (each root with multiplicity $2p-2$).

In particular, $G_1 = K_{3,3}$ and $G_2 = K_2$. Then the normalized Laplacian eigenvalues of G_1 are 0, 1(multiplicity 4) and 2. The normalized Laplacian eigenvalues of G_2 are 0 and 2. Therefore, the normalized Laplacian eigenvalues of $G_1 \dot{\vee} G_2$ are 0, $\frac{8}{7}, 1$ (multiplicity 3), roots of the equation $49x^2 - 126x + 74 = 0$, roots of the equation $14x^2 - 24x + 10 = 0$ and roots of the equation $14x^2 - 30x + 13 = 0$ (each root with multiplicity 4).

Next we consider the normalized Laplacian spectrum of $G_1 \vee G_2$.

Theorem 4.3. *Let G_i be an r_i -regular graph with n_i vertices and m_i edges for $i=1,2$. Then the normalized Laplacian characteristic polynomial of $G_1 \vee G_2$ is given by*

$$\begin{aligned} f(\mathcal{L}(G_1 \vee G_2), x) &= (x-1)^{m_1-n_1-1} \prod_{i=2}^{n_2} \left((x - \frac{m_1}{r_2+m_1}) I_{n_2} - \frac{r_2}{r_2+m_1} \tilde{\mu}_i(G_2) \right) \\ &\prod_{i=2}^{n_1} \left[(x-1)^2 + \frac{(-1-r_1+r_1\tilde{\mu}_i(G_1))(x-1)}{n_1-1} - \frac{(2r_1-r_1\tilde{\mu}_i(G_1))}{(n_1-1)(n_2+2)} \right] \\ &\times \left[\left((x-1)^2 + \frac{(n_1-1-r_1)(x-1)}{n_1-1} - \frac{2r_1}{(n_1-1)(n_2+2)} \right) \right. \\ &\left. \left((x - \frac{m_1}{r_2+m_1})(x-1) - \frac{n_2 m_1}{(n_2+2)(r_2+m_1)} \right) - \frac{n_2 r_1^2 n_1}{(n_1-1)(n_2+2)^2(r_2+m_1)} \right]. \end{aligned}$$

Proof. Let $I(G_1)$ be the incidence matrix of G_1 . Then by a proper labeling of vertices, adjacency matrix of $G_1 \vee G_2$ can be written as

$$A(G_1 \vee G_2) = \begin{pmatrix} A(\bar{G}_1) & I(G_1) & O_{n_1 \times n_2} \\ I(G_1)^T & O_{m_1 \times m_1} & J_{m_1 \times n_2} \\ O_{n_2 \times n_1} & J_{n_2 \times m_1} & A(G_2) \end{pmatrix}.$$

The degree matrix of $G_1 \vee G_2$ is

$$D(G_1 \vee G_2) = \begin{pmatrix} (n_1-1)I_{n_1} & O_{n_1 \times m_1} & O_{n_1 \times n_2} \\ O_{m_1 \times n_1} & (n_2+2)I_{m_1} & O_{m_1 \times n_2} \\ O_{n_2 \times n_1} & O_{n_2 \times m_1} & (r_2+m_1)I_{n_2} \end{pmatrix}.$$

$$D^{-\frac{1}{2}}(G_1 \vee G_2) = \begin{pmatrix} \frac{I_{n_1}}{\sqrt{(n_1-1)}} & O_{n_1 \times m_1} & O_{n_1 \times n_2} \\ O_{m_1 \times n_1} & \frac{I_{m_1}}{\sqrt{n_2+2}} & O_{m_1 \times n_2} \\ O_{n_2 \times n_1} & O_{n_2 \times m_1} & \frac{I_{n_2}}{\sqrt{(r_2+m_1)}} \end{pmatrix}.$$

Then the normalized Laplacian matrix of $G_1 \vee G_2$ is

$$\mathcal{L}(G_1 \vee G_2) = \begin{pmatrix} I_{n_1} - \frac{A(\bar{G}_1)}{n_1-1} & -\frac{I(G_1)}{\sqrt{(n_1-1)(n_2+2)}} & O_{n_1 \times n_2} \\ -\frac{I(G_1)^T}{\sqrt{(n_1-1)(n_2+2)}} & I_{m_1} & -\frac{J_{m_1 \times n_2}}{\sqrt{(n_2+2)(r_2+m_1)}} \\ O_{n_2 \times n_1} & -\frac{J_{n_2 \times m_1}}{\sqrt{(n_2+2)(r_2+m_1)}} & I_{n_2} - \frac{A(G_2)}{(r_2+m_1)} \end{pmatrix}$$

$$\mathcal{L}(G_1 \vee G_2) = \begin{pmatrix} I_{n_1} - \frac{A(\bar{G}_1)}{n_1-1} & -\frac{I(G_1)}{\sqrt{(n_1-1)(n_2+2)}} & O_{n_1 \times n_2} \\ -\frac{I(G_1)^T}{\sqrt{(n_1-1)(n_2+2)}} & I_{m_1} & -\frac{J_{m_1 \times n_2}}{\sqrt{(n_2+2)(r_2+m_1)}} \\ O_{n_2 \times m_1} & -\frac{J_{n_2 \times m_1}}{\sqrt{(n_2+2)(r_2+m_1)}} & \mathcal{L}(G_2) \circ C \end{pmatrix}$$

where $C = \frac{r_2}{r_2+m_1} J_{n_2} + \frac{m_1}{r_2+m_1} I_{n_2}$.

Using Definition 2.1, Lemmas 2.1, 2.2, 2.3, 2.6 and Corollary 2.5, we have the normalized Laplacian characteristic polynomial of $G_1 \vee G_2$ is

$$f(\mathcal{L}(G_1 \vee G_2), x) = \det \begin{pmatrix} (x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1-1} & \frac{I(G_1)}{\sqrt{(n_1-1)(n_2+2)}} & O_{n_1 \times n_2} \\ \frac{I(G_1)^T}{\sqrt{(n_1-1)(n_2+2)}} & (x-1)I_{m_1} & \frac{J_{m_1 \times n_2}}{\sqrt{(n_2+2)(r_2+m_1)}} \\ O_{n_2 \times m_1} & \frac{J_{n_2 \times m_1}}{\sqrt{(n_2+2)(r_2+m_1)}} & xI_{n_2} - (\mathcal{L}(G_2) \circ C) \end{pmatrix}$$

$$= \det \left(xI_{n_2} - (\mathcal{L}(G_2) \circ C) \right) \det S,$$

$$\text{where } S = \begin{pmatrix} (x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1-1} & \frac{I(G_1)}{\sqrt{(n_1-1)(n_2+2)}} \\ \frac{I(G_1)^T}{\sqrt{(n_1-1)(n_2+2)}} & (x-1)I_{m_1} \end{pmatrix} - \left[\begin{pmatrix} O_{n_1 \times n_2} \\ \frac{J_{m_1 \times n_2}}{\sqrt{(n_2+2)(r_2+m_1)}} \end{pmatrix} \right. \\ \left. \times (xI_{n_2} - (\mathcal{L}(G_2) \circ C))^{-1} \begin{pmatrix} O_{n_2 \times m_1} & \frac{J_{n_2 \times m_1}}{\sqrt{(n_2+2)(r_2+m_1)}} \end{pmatrix} \right]$$

$$S = \begin{pmatrix} (x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1-1} & \frac{I(G_1)}{\sqrt{(n_1-1)(n_2+2)}} \\ \frac{I(G_1)^T}{\sqrt{(n_1-1)(n_2+2)}} & (x-1)I_{m_1} - \frac{J_{m_1 \times n_2}}{\sqrt{(n_2+2)(r_2+m_1)}} (xI_{n_2} - (\mathcal{L}(G_2) \circ C))^{-1} \frac{J_{n_2 \times m_1}}{\sqrt{(n_2+2)(r_2+m_1)}} \end{pmatrix} \\ = \begin{pmatrix} (x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1-1} & \frac{I(G_1)}{\sqrt{(n_1-1)(n_2+2)}} \\ \frac{I(G_1)^T}{\sqrt{(n_1-1)(n_2+2)}} & (x-1)I_{m_1} - \frac{\chi_{\mathcal{L}(G_2)}(x, \frac{r_2}{r_2+m_1})}{(n_2+2)(r_2+m_1)} J_{m_1} \end{pmatrix}.$$

Therefore,

$$\det(S) = \det \begin{pmatrix} (x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1-1} & \frac{I(G_1)}{\sqrt{(n_1-1)(n_2+2)}} \\ \frac{I(G_1)^T}{\sqrt{(n_1-1)(n_2+2)}} & (x-1)I_{m_1} - \frac{\chi_{\mathcal{L}(G_2)}(x, \frac{r_2}{r_2+m_1})}{(n_2+2)(r_2+m_1)} J_{m_1} \end{pmatrix} \\ = \det \left((x-1)I_{m_1} - \frac{\chi_{\mathcal{L}(G_2)}(x, \frac{r_2}{r_2+m_1})}{(n_2+2)(r_2+m_1)} J_{m_1} \right) \det \left[(x-1)I_{n_1} + \frac{A(\bar{G}_1)}{n_1-1} \right. \\ \left. - \frac{I(G_1)}{\sqrt{(n_1-1)(n_2+2)}} \left((x-1)I_{m_1} - \frac{\chi_{\mathcal{L}(G_2)}(x, \frac{r_2}{r_2+m_1})}{(n_2+2)(r_2+m_1)} J_{m_1} \right)^{-1} \frac{I(G_1)^T}{\sqrt{(n_1-1)(n_2+2)}} \right] \\ = (x-1)^{m_1} \left(1 - \frac{\chi_{\mathcal{L}(G_2)}(x, \frac{r_2}{r_2+m_1})}{(n_2+2)(r_2+m_1)} \cdot \frac{m_1}{x-1} \right) \det \left[(x-1)I_{n_1} + \frac{J_{n_1} - I_{n_1} - A(G_1)}{n_1-1} \right. \\ \left. - \frac{I(G_1)}{\sqrt{(n_1-1)(n_2+2)}} \left(\frac{1}{(x-1)} I_{m_1} + \frac{\beta}{(x-1)(x-1-m_1\beta)} J_{m_1} \right) \frac{I(G_1)^T}{\sqrt{(n_1-1)(n_2+2)}} \right],$$

where $\beta = \frac{\chi_{\mathcal{L}(G_2)}(x, \frac{r_2}{r_2+m_1})}{(n_2+2)(r_2+m_1)}$

$$\begin{aligned}
\det S &= (x-1)^{m_1} \left(1 - \beta \frac{m_1}{x-1}\right) \det \left[(x-1)I_{n_1} + \frac{J_{n_1} - I_{n_1} - A(G_1)}{n_1-1} - \frac{I(G_1)I(G_1)^T}{(x-1)(n_1-1)(n_2+2)} \right. \\
&\quad \left. - \frac{I(G_1)}{\sqrt{(n_1-1)(n_2+2)}} \left(\frac{\beta}{(x-1)(x-1-m_1\beta)} J_{m_1} \right) \frac{I(G_1)^T}{\sqrt{(n_1-1)(n_2+2)}} \right] \\
&= (x-1)^{m_1} \left(1 - \beta \frac{m_1}{x-1}\right) \det \left[(x-1)I_{n_1} + \frac{J_{n_1} - I_{n_1} - A(G_1)}{n_1-1} \right. \\
&\quad \left. - \frac{I(G_1)I(G_1)^T}{(x-1)(n_1-1)(n_2+2)} - \frac{\beta}{(x-1)(x-1-m_1\beta)} \frac{I(G_1)J_{m_1}I(G_1)^T}{(n_1-1)(n_2+2)} \right] \\
&= (x-1)^{m_1} \left(1 - \beta \frac{m_1}{x-1}\right) \det \left[(x-1)I_{n_1} + \frac{J_{n_1} - I_{n_1} - A(G_1)}{n_1-1} \right. \\
&\quad \left. - \frac{I(G_1)I(G_1)^T}{(x-1)(n_1-1)(n_2+2)} - \frac{\beta r_1^2 J_{n_1}}{(x-1)(x-1-m_1\beta)(n_1-1)(n_2+2)} \right] \\
&= (x-1)^{m_1} \left(1 - \beta \frac{m_1}{x-1}\right) \det \left[(x-1)I_{n_1} + \frac{J_{n_1} - I_{n_1} - A(G_1)}{n_1-1} - \frac{I(G_1)I(G_1)^T}{(x-1)(n_1-1)(n_2+2)} \right] \\
&\quad \left[1 - \frac{\beta r_1^2 \chi_{\frac{J_{n_1}-I_{n_1}-A(G_1)}{n_1-1} + \frac{I(G_1)I(G_1)^T}{(x-1)(n_1-1)(n_2+2)}}(x-1)}{(x-1)(x-1-m_1\beta)(n_1-1)(n_2+2)} \right].
\end{aligned}$$

Thus,

$$\begin{aligned}
\det(S) &= (x-1)^{m_1} \left(1 - \frac{n_2 m_1}{(x-1)\left(x - \frac{m_1}{r_2+m_1}\right)(n_2+2)(r_2+m_1)}\right) \\
&\quad \times \det \left[(x-1)I_{n_1} + \frac{J_{n_1} - I_{n_1} - r_1 I_{n_1} + r_1 \mathcal{L}(G_1)}{n_1-1} - \frac{(2r_1 I_{n_1} - r_1 \mathcal{L}(G_1))}{(x-1)(n_1-1)(n_2+2)} \right] \\
&\quad \times \left[1 - \frac{\frac{n_2 r_1^2 n_1}{\left(x - \frac{m_1}{r_2+m_1}\right)(n_2+2)(r_2+m_1)\left(x-1 + \frac{n_1-1-r_1}{n_1-1} - \frac{2r_1}{(x-1)(n_1-1)(n_2+2)}\right)}}{(x-1)(x-1-m_1\beta)(n_1-1)(n_2+2)} \right].
\end{aligned}$$

$$\begin{aligned}
& f(\mathcal{L}(G_1 \vee G_2), x) = \det \left(xI_{n_2} - (\mathcal{L}(G_2) \circ C) \right) \det S \\
& = \det \left(\left(x - \frac{m_1}{r_2 + m_1} \right) I_{n_2} - \frac{r_2}{r_2 + m_1} \mathcal{L}(G_2) \right) \\
& \quad \times (x-1)^{m_1} \left(1 - \frac{n_2 m_1}{(x-1) \left(x - \frac{m_1}{r_2 + m_1} \right) (n_2 + 2)(r_2 + m_1)} \right) \\
& \quad \times \det \left[(x-1)I_{n_1} + \frac{J_{n_1} - I_{n_1} - r_1 I_{n_1} + r_1 \mathcal{L}(G_1)}{n_1 - 1} - \frac{(2r_1 I_{n_1} - r_1 \mathcal{L}(G_1))}{(x-1)(n_1 - 1)(n_2 + 2)} \right] \\
& \quad \times \left[1 - \frac{\frac{n_2 r_1^2 n_1}{\left(x - \frac{m_1}{r_2 + m_1} \right) (n_2 + 2)(r_2 + m_1) \left(x - 1 + \frac{n_1 - 1 - r_1}{n_1 - 1} - \frac{2r_1}{(x-1)(n_1 - 1)(n_2 + 2)} \right)}}{(x-1) \left(x - 1 - \frac{m_1 n_2}{\left(x - \frac{m_1}{r_2 + m_1} \right) (n_2 + 2)(r_2 + m_1)} \right)} (n_1 - 1)(n_2 + 2) \right] \\
& = \prod_{i=2}^{n_2} \left(\left(x - \frac{m_1}{r_2 + m_1} \right) I_{n_2} - \frac{r_2}{r_2 + m_1} \tilde{\mu}_i(G_2) \right) \\
& \quad \times (x-1)^{m_1 - n_1} \left[\left(x - \frac{m_1}{r_2 + m_1} \right) - \frac{n_2 m_1}{(x-1)(n_2 + 2)(r_2 + m_1)} \right] \\
& \quad \prod_{i=1}^{n_1} \left[(x-1)^2 + \frac{(P(\lambda_i) - 1 - r_1 + r_1 \tilde{\mu}_i(G_1))(x-1)}{n_1 - 1} - \frac{(2r_1 - r_1 \tilde{\mu}_i(G_1))}{(n_1 - 1)(n_2 + 2)} \right] \\
& \quad \times \left[1 - \frac{\frac{n_2 r_1^2 n_1}{\left(x - \frac{m_1}{r_2 + m_1} \right) (n_2 + 2)(r_2 + m_1) \left(x - 1 + \frac{n_1 - 1 - r_1}{n_1 - 1} - \frac{2r_1}{(x-1)(n_1 - 1)(n_2 + 2)} \right)}}{(x-1) \left(x - 1 - \frac{m_1 n_2}{\left(x - \frac{m_1}{r_2 + m_1} \right) (n_2 + 2)(r_2 + m_1)} \right)} (n_1 - 1)(n_2 + 2) \right] \\
& = \prod_{i=2}^{n_2} \left(\left(x - \frac{m_1}{r_2 + m_1} \right) I_{n_2} - \frac{r_2}{r_2 + m_1} \tilde{\mu}_i(G_2) \right) \\
& \quad \times (x-1)^{m_1 - n_1} \left[\left(x - \frac{m_1}{r_2 + m_1} \right) - \frac{n_2 m_1}{(x-1)(n_2 + 2)(r_2 + m_1)} \right] \\
& \quad \prod_{i=2}^{n_1} \left[(x-1)^2 + \frac{(-1 - r_1 + r_1 \tilde{\mu}_i(G_1))(x-1)}{n_1 - 1} - \frac{(2r_1 - r_1 \tilde{\mu}_i(G_1))}{(n_1 - 1)(n_2 + 2)} \right] \\
& \quad \times \left[(x-1)^2 + \frac{(n_1 - 1 - r_1)(x-1)}{n_1 - 1} - \frac{2r_1}{(n_1 - 1)(n_2 + 2)} \right. \\
& \quad \left. - \frac{n_2 r_1^2 n_1}{\left(x - \frac{m_1}{r_2 + m_1} \right) (n_2 + 2)(r_2 + m_1) \left(x - 1 - \frac{m_1 n_2}{\left(x - \frac{m_1}{r_2 + m_1} \right) (n_2 + 2)(r_2 + m_1)} \right)} (n_1 - 1)(n_2 + 2) \right]
\end{aligned}$$

$$\begin{aligned}
&= (x-1)^{m_1-n_1-1} \prod_{i=2}^{n_2} \left(\left(x - \frac{m_1}{r_2+m_1} \right) I_{n_2} - \frac{r_2}{r_2+m_1} \tilde{\mu}_i(G_2) \right) \\
&\quad \prod_{i=2}^{n_1} \left[(x-1)^2 + \frac{(-1-r_1+r_1\tilde{\mu}_i(G_1))(x-1)}{n_1-1} - \frac{(2r_1-r_1\tilde{\mu}_i(G_1))}{(n_1-1)(n_2+2)} \right] \\
&\quad \times \left[\left((x-1)^2 + \frac{(n_1-1-r_1)(x-1)}{n_1-1} - \frac{2r_1}{(n_1-1)(n_2+2)} \right) \right. \\
&\quad \left. \left(\left(x - \frac{m_1}{r_2+m_1} \right) (x-1) - \frac{n_2 m_1}{(n_2+2)(r_2+m_1)} \right) - \frac{n_2 r_1^2 n_1}{(n_1-1)(n_2+2)^2(r_2+m_1)} \right].
\end{aligned}$$

□

The following corollary describes the complete normalized Laplacian spectrum of $G_1 \vee G_2$ when G_1 and G_2 are regular graphs.

Corollary 4.4. *Let G_i be an r_i -regular graph with n_i vertices and m_i edges for $i=1,2$. Then the normalized Laplacian spectrum of $G_1 \vee G_2$ consists of*

1. 1 repeated $m_1 - n_1 - 1$ times.
2. Roots of the equation,

$$\left(x - \frac{m_1}{r_2+m_1} \right) - \frac{r_2}{r_2+m_1} \tilde{\mu}_i(G_2) = 0 \text{ for } i = 2, \dots, n_2.$$

3. Four roots of the equation,

$$\begin{aligned}
&\left((x-1)^2 + \frac{(n_1-1-r_1)(x-1)}{n_1-1} - \frac{2r_1}{(n_1-1)(n_2+2)} \right) \\
&\left(\left(x - \frac{m_1}{r_2+m_1} \right) (x-1) - \frac{n_2 m_1}{(n_2+2)(r_2+m_1)} \right) - \frac{n_2 r_1^2 n_1}{(n_1-1)(n_2+2)^2(r_2+m_1)} = 0.
\end{aligned}$$

4. Two roots of the equation,

$$\left(x-1 \right)^2 + \frac{(-1-r_1+r_1\tilde{\mu}_i(G_1))(x-1)}{n_1-1} - \frac{(2r_1-r_1\tilde{\mu}_i(G_1))}{(n_1-1)(n_2+2)} = 0 \text{ for } i = 2, \dots, n_1.$$

Example 4.2. Let $G_1 = K_{p,p}$ and $G_2 = K_2$. Then the normalized Laplacian eigenvalues of G_1 are 0, 1(multiplicity $2p-2$) and 2. The normalized Laplacian eigenvalues of G_2 are 0 and 2. Therefore, the normalized Laplacian eigenvalues of $G_1 \vee G_2$ are 0, $\frac{p^2+2}{p^2+1}, 1$ (multiplicity p^2-2p-1), roots of the equation $(32p^3-16p^2+32p-16)x^3+(-112p^3+48p^2-80p+32)x^2+(120p^3-40p^2+56p-16)x+8p^2-40p^3-8p=0$, roots of the equation $(2p-1)x^2-(3p-1)x+p=0$ and roots of the equation $(8p-4)x^2-(16p-4)x+7p=0$ (each root with multiplicity $2p-2$).

In particular, $G_1 = K_{3,3}$ and $G_2 = K_2$. Then the normalized Laplacian eigenvalues of G_1 are 0, 1(multiplicity 4) and 2. The normalized Laplacian eigenvalues of G_2 are 0 and 2. Therefore, the normalized Laplacian eigenvalues of $G_1 \vee G_2$ are 0, $\frac{11}{10}, 1$ (multiplicity 2), roots of the equation $800x^3-2800x^2+3032x-1032=0$, roots of the equation $5x^2-8x+3=0$ and roots of the equation $20x^2-44x+21=0$ (each root with multiplicity 4).

5 Applications

In this section, we construct a family of non-regular \mathcal{L} -cospectral graphs using central joins. Also, we compute the degree Kirchhoff index and the Kemeny's constant of central graph, central vertex join and central edge join of two regular graphs.

In the following theorem we construct non-regular \mathcal{L} cospectral graphs using central vertex join and central edge join of two regular graphs.

Theorem 5.1. *Let G_1 and G_2 (not necessarily distinct) be \mathcal{L} -cospectral regular graphs, H_1 and H_2 (not necessarily distinct) are another \mathcal{L} -cospectral regular graphs. Then $\mathcal{L}(G_1 \dot{\vee} H_1)$ and $\mathcal{L}(G_2 \dot{\vee} H_2)$ (respectively, $\mathcal{L}(G_1 \vee H_1)$ and $\mathcal{L}(G_2 \vee H_2)$) are \mathcal{L} -cospectral non-regular graphs.*

Proof. If G_1 and G_2 are two \mathcal{L} -cospectral regular graphs, then they have the same regularity with same number of vertices and same number of edges. It is also true for H_1 and H_2 . By applying Theorem 4.1 on the concerned graphs and comparing their normalized Laplacian characteristic polynomial we get the required result. \square

Example 5.1. Consider two non isomorphic regular cospectral graphs G and H as in [16]. Graphs $C(G)$ and $C(H)$ as shown in Figure:1. Also, $G \dot{\vee} K_2$ and $H \dot{\vee} K_2$ are non isomorphic. By Theorem 5.1, we obtain $G \dot{\vee} K_2$ and $H \dot{\vee} K_2$ are \mathcal{L} -cospectral non-regular graphs.





Figure:1 Non-regular non isomorphic \mathcal{L} cospectral graphs



$G \dot{\vee} K_2$

$H \dot{\vee} K_2$

Figure:2 Non-regular non isomorphic \mathcal{L} cospectral graphs

By Corollary 3.2, we can readily obtain the following theorem.

Theorem 5.2. *Let G be an r -regular graph of order n and size m . Then*

$$K(C(G)) = \left[m - n + \frac{n-1}{n-1+r} + \sum_{i=2}^n \frac{2(2n + \lambda_i - 1)}{2n + \lambda_i - r} \right].$$

The following result is obtained from Theorem 5.2.

Let G be an r -regular graph of order n and size m . Then

$$Kf^*(C(G)) = (2m + n(n-1)) \left[m - n + \frac{n-1}{n-1+r} + \sum_{i=2}^n \frac{2(2n + \lambda_i - 1)}{2n + \lambda_i - r} \right].$$

By Corollary 4.2, we can readily obtain the following theorem.

Theorem 5.3. *Let G_i be an r_i -regular graph with n_i vertices and m_i edges for $i=1,2$. Then*

$$K(G_1 \dot{\vee} G_2) = \left[m_1 - n_1 + \frac{n_1(2n_1 + 3n_2 + r_1 + r_2) + 2n_2r_2 - 2n_1 - r_2 - r_1r_2}{n_1(n_1 + 2n_2 - 2) + n_2r_2} + \sum_{i=2}^{n_2} \frac{r_2 + n_1}{n_1 + r_2\tilde{\mu}_i(G_2)} + \sum_{i=2}^{n_1} \frac{2(-r_1\tilde{\mu}_i(G_1) + 2n_1 + 2n_2 + r_1 - 1)}{2n_1 + 2n_2 - r_1\tilde{\mu}_i(G_1)} \right].$$

The following result is obtained from Theorem 5.3.

Let G_i be an r_i -regular graph with n_i vertices and m_i edges for $i = 1, 2$.

Then

$$Kf^*(G_1 \dot{\vee} G_2) = 2m_1 + 2m_2 + 2n_1n_2 + n_1(n_1 - 1) \left[m_1 - n_1 + \frac{n_1(2n_1 + 3n_2 + r_1 + r_2) + 2n_2r_2 - 2n_1 - r_2 - r_1r_2}{n_1(n_1 + 2n_2 - 2) + n_2r_2} + \sum_{i=2}^{n_2} \frac{r_2 + n_1}{n_1 + r_2\tilde{\mu}_i(G_2)} + \sum_{i=2}^{n_1} \frac{2(-r_1\tilde{\mu}_i(G_1) + 2n_1 + 2n_2 + r_1 - 1)}{2n_1 + 2n_2 - r_1\tilde{\mu}_i(G_1)} \right].$$

By Corollary 4.4, we get the following theorem.

Theorem 5.4. Let G_i be an r_i -regular graph with n_i vertices and m_i edges for $i=1,2$. Then

$$K(G_1 \vee G_2) = \left[m_1 - n_1 - 1 + \frac{r_2(4n_1n_2 - n_2^2 + 6n_2r_1 + 4n_1 - 4n_2 + 4r_1 - 4)}{m_1(2n_2^2r_1 + 2n_1n_2 + 6n_2r_1 + 4n_1 + 2n_2) + 2n_2r_1 + n_2^2r_1r_2} \right. \\ \left. \frac{n_2^2(n_1r_2 + 2n_1m_1 + 2r_1r_2 + 3m_1r_1) + m_1(10n_1n_2 - 2n_2^2 + 10n_2r_1 + 12n_1 - 10n_2 + 8r_1 - 12)}{m_1(2n_2^2r_1 + 2n_1n_2 + 6n_2r_1 + 4n_1 + 2n_2) + 2n_2r_1 + n_2^2r_1r_2} + \right. \\ \left. \sum_{i=2}^{n_2} \frac{r_2 + m_1}{m_1 + r_2\tilde{\mu}_i(G_2)} + \sum_{i=2}^{n_1} \frac{-n_2r_1\tilde{\mu}_i(G_1) - 2r_1\tilde{\mu}_i(G_1) + 2n_1n_2 + n_2r_1 + 4n_1 - n_2 + 2r_1 - 2}{n_1n_2 + n_2r_1 + 2n_1 - n_2r_1\tilde{\mu}_i(G_1) - 3r_1\tilde{\mu}_i(G_1)} \right].$$

The following result is obtained from Theorem 5.4.

Let G_i be an r_i -regular graph with n_i vertices and m_i edges for $i = 1, 2$. Then

$$Kf^*(G_1 \vee G_2) = 2m_1 + 2m_2 + 2m_1m_2 + n_1(n_1 - 1) \\ \left[m_1 - n_1 - 1 + \frac{r_2(4n_1n_2 - n_2^2 + 6n_2r_1 + 4n_1 - 4n_2 + 4r_1 - 4)}{m_1(2n_2^2r_1 + 2n_1n_2 + 6n_2r_1 + 4n_1 + 2n_2) + 2n_2r_1 + n_2^2r_1r_2} \right. \\ \left. \frac{n_2^2(n_1r_2 + 2n_1m_1 + 2r_1r_2 + 3m_1r_1) + m_1(10n_1n_2 - 2n_2^2 + 10n_2r_1 + 12n_1 - 10n_2 + 8r_1 - 12)}{m_1(2n_2^2r_1 + 2n_1n_2 + 6n_2r_1 + 4n_1 + 2n_2) + 2n_2r_1 + n_2^2r_1r_2} + \right. \\ \left. \sum_{i=2}^{n_2} \frac{r_2 + m_1}{m_1 + r_2\tilde{\mu}_i(G_2)} + \sum_{i=2}^{n_1} \frac{-n_2r_1\tilde{\mu}_i(G_1) - 2r_1\tilde{\mu}_i(G_1) + 2n_1n_2 + n_2r_1 + 4n_1 - n_2 + 2r_1 - 2}{n_1n_2 + n_2r_1 + 2n_1 - n_2r_1\tilde{\mu}_i(G_1) - 3r_1\tilde{\mu}_i(G_1)} \right].$$

6 Conclusion

In this paper, the normalized Laplacian spectrum of central graph, central vertex join and central edge join of two regular graphs are calculated. Also, we construct non-regular \mathcal{L} -cospectral graphs. Moreover, we investigate some graph invariants like the degree Kirchhoff index and the Kemeny's constant of central graph, central vertex join and central edge join of two regular graphs.

References

- [1] Ivan Gutman, Boris Furtula, and Ş Burcu Bozkurt. On Randić energy. *Linear Algebra and its Applications*, 442:50–57, 2014.
- [2] S Burcu Bozkurt, A Dilek Güngör, Ivan Gutman, and A Sinan Cevik. Randić matrix and Randić energy. *MATCH Commun. Math. Comput. Chem*, 64(1):239–250, 2010.
- [3] Douglas J Klein and Milan Randić. Resistance distance. *Journal of mathematical chemistry*, 12(1):81–95, 1993.
- [4] Haiyan Chen and Fuji Zhang. Resistance distance and the normalized Laplacian spectrum. *Discrete Applied Mathematics*, 155(5):654–661, 2007.
- [5] Steve Butler. Algebraic aspects of the normalized Laplacian. In *Recent Trends in Combinatorics*, pages 295–315. Springer, 2016.

- [6] Jeffrey J Hunter. The role of Kemeny's constant in properties of Markov chains. *Communications in Statistics-Theory and Methods*, 43(7):1309–1321, 2014.
- [7] Roger A Horn and Charles R Johnson. Topics in matrix analysis. Cambridge university press. *Cambridge, UK*, 1991.
- [8] Dragoš M Cvetković, Michael Doob, and Horst Sachs. *Spectra of graphs, theory and applications*, volume 10. Academic Press, New York, 1980.
- [9] Dragos M Cvetković, Peter Rowlinson, and Slobodan Simic. *An introduction to the theory of graph spectra*. Cambridge University Press Cambridge, 2010.
- [10] J Vernold Vivin, MM Akbar Ali, and K Thilagavathi. On harmonious coloring of central graphs. *Advances and applications in discrete mathematics*, 2(1):17–33, 2008.
- [11] Arpita Das and Pratima Panigrahi. Spectra of R-vertex join and R-edge join of two graphs. *Discussiones Mathematicae-General Algebra and Applications*, 38(1):19–32, 2018.
- [12] Gui-xian Tian and Shu-Yu Cui. On the normalized Laplacian spectra of some subdivision joins of two graphs. *arXiv:1703.06595*, 2017.
- [13] Cam McLeman and Erin McNicholas. Spectra of coroneae. *Linear algebra and its applications*, 435(5):998–1007, 2011.
- [14] Xiaogang Liu and Zuhe Zhang. Spectra of subdivision-vertex join and subdivision-edge join of two graphs. *Bulletin of the Malaysian Mathematical Sciences Society*, 42(1):15–31, 2019.
- [15] TK Jahfar and AV Chithra. Central vertex join and central edge join of two graphs. *AIMS Mathematics*, 5(6):7214–7233, 2020.
- [16] Edwin R Van Dam and Willem H Haemers. Which graphs are determined by their spectrum? *Linear Algebra and its applications*, 373:241–272, 2003.