

On (Coalitional) Exchange-Stable Matching

Jiehua Chen, Adrian Chmurovic, Fabian Jogl, and Manuel Sorge
TU Wien, Austria
{jiehua.chen, manuel.sorge}@tuwien.ac.at

Abstract

We study (*coalitional*) *exchange stability*, which Alcalde [Economic Design, 1995] introduced as an alternative solution concept for matching markets involving property rights, such as assigning persons to two-bed rooms. Here, a matching of a given STABLE MARRIAGE or STABLE ROOMMATES instance is called *coalitional exchange-stable* if it does not admit any *exchange-blocking coalition*, that is, a subset S of agents in which everyone prefers the partner of some other agent in S . The matching is *exchange-stable* if it does not admit any *exchange-blocking pair*, that is, an exchange-blocking coalition of size two.

We investigate the computational and parameterized complexity of the COALITIONAL EXCHANGE-STABLE MARRIAGE (resp. COALITIONAL EXCHANGE ROOMMATES) problem, which is to decide whether a STABLE MARRIAGE (resp. STABLE ROOMMATES) instance admits a coalitional exchange-stable matching. Our findings resolve an open question and confirm the conjecture of Cechlárová and Manlove [Discrete Applied Mathematics, 2005] that COALITIONAL EXCHANGE-STABLE MARRIAGE is NP-hard even for complete preferences without ties. We also study bounded-length preference lists and a local-search variant of deciding whether a given matching can reach an exchange-stable one after at most k *swaps*, where a swap is defined as exchanging the partners of the two agents in an exchange-blocking pair.

1 Introduction

An instance in a matching market consists of a set of agents that each have preferences over other agents with whom they want to be matched with. The goal is to find a matching, i.e., a subset of disjoint pairs of agents, which is *fair*. A classical notion of fairness is *stability* [13], meaning that no two agents can form a *blocking pair*, i.e., they would prefer to be matched with each other rather than with the partner assigned by the matching. This means that a matching is fair if the agents cannot take local action to improve their outcome. If we assign property rights via the matching, however, then the notion of blocking pairs may not be actionable, as Alcalde [3] observed: For example, if the matching represents an assignment of persons to two-bed rooms, then two persons in a blocking pair may not be able to deviate from the assignment because they may not find a new room that they could share. Instead, we may consider the matching to be *fair* if no two agents form an *exchange-blocking pair*, that is, they would prefer to have each other's partner rather than to have the partner given by the matching [3]. In other words, they would like to *exchange* their partners. Note that such an exchange would be straightforward in the room-assignment problem mentioned before. We refer to Alcalde [3], Cechlárová [9], and Cechlárová and Manlove [10] for more discussion and examples of markets involving property rights.

If a matching does not admit an exchange-blocking pair, then we say the matching is *exchange-stable*. If we also want to exclude the possibility that several agents may collude to favorably exchange partners, then we arrive at *coalitional exchange-stability* [3], a concept that is more stringent than the exchange-stability. In contrast to classical stability and exchange-stability for perfect matchings (i.e., everyone is matched), it is not hard to observe that coalitional exchange-stability implies *Pareto-optimality*, another fairness concept

which asserts that no other matching can make at least one agent better-off without making some other agent worse-off (see also Abraham and Manlove [2]). Note that, in contrast, the classical Gale/Shapley stability and Pareto-optimality are incompatible to each other.

Cechlárová and Manlove [10] showed that the problem of deciding whether an exchange-stable matching exists is NP-hard, even for the marriage case (where the agents are partitioned into two subsets of equal size such that each agent of either subset has preferences over the agents of the other subset) with complete preferences but without ties. They left open whether the NP-hardness transfers to the case with coalitional exchange-stability, but observed NP-containment.

Our contributions. We study the algorithmic complexity of problems revolving around (coalitional) exchange-stability. In particular, we establish a first NP-hardness result for deciding coalitional exchange-stability, confirming a conjecture of Cechlárová and Manlove [10]. The NP-hardness reduction is based on a novel *switch-gadget* wherein each preference list contains at most three agents. Utilizing this, we can carefully complete the preferences so as to prove the desired NP-hardness. We then investigate the impact of the maximum length d of a preference list. We find that NP-hardness for both exchange-stability and coalitional exchange-stability starts already when $d = 3$, while it is fairly easy to see that the problem becomes polynomial-time solvable for $d = 2$. For $d = 3$, we obtain a fixed-parameter algorithm for exchange-stability regarding a parameter which is related to the number of switch-gadgets.

Finally, we look at a problem variant, called PATH TO EXCHANGE-STABLE MARRIAGE (P-ESM), for uncoordinated (or decentralized) matching markets. Starting from an initial matching, in each iteration the two agents in an exchange-blocking pair may exchange their partners. An interesting question regarding the behavior of the agents in uncoordinated markets is whether such iterative exchange actions can reach a stable state, i.e., exchange-stability, and how hard is it to decide. It is fairly straight-forward to verify that if the number k of exchanges is bounded by a constant, then we can decide in polynomial-time whether an exchange-stable matching is reachable since there are only polynomially many possible sequences of exchanges to be checked. From the parameterized complexity point of view, we obtain an XP algorithm for k , i.e., the exponent in the polynomial running time depends on k . We further show that this dependency is unlikely to be removed by showing W[1]-hardness with respect to k .

Related work. Alcalde [3] introduced (coalitional) exchange stability and discussed restricted preference domains where (coalitional) exchange stability is guaranteed to exist. Abizada [1] showed a weaker condition (on the preference domain) to guarantee the existence of exchange stability. Cechlárová and Manlove [10] proved that it is NP-complete to decide whether an exchange-stable matching exists, even for the marriage case with complete preferences without ties. Aziz and Goldwasser [4] introduced several relaxed notions of coalitional exchange-stability and discussed their relations.

The P-ESM problem is inspired by the PATH-TO-STABILITY VIA DIVORCES (PSD) problem, originally introduced by Knuth [17], see also Biró and Norman [5] for more background. Very recently, Chen [11] showed that PSD is NP-hard and W[1]-hard when parameterized by the number of divorces. P-ESM can also be considered as a local search problem and is a special case of the LOCAL SEARCH EXCHANGE-STABLE SEAT ARRANGEMENT (LOCAL-STA) problem, introduced by Bodlaender et al. [6]: Given a set of agents, each having cardinal preferences (i.e., real values) over the other agents, an undirected graph G with the same number of vertices as agents, and an initial assignment (bijection) of the agents to the vertices in G , is it possible to swap two agents' assignments iteratively so as to reach an exchange-stable assignment? Herein an assignment is called *exchange-stable* if no

two agents can each have a higher sum of cardinal preferences over the other’s neighboring agents. P-ESM is a restricted variant of LOCAL-STA, where G consists of disjoint edges and the agents have ordinal preferences. Bodlaender et al. [7] showed that LOCAL-STA is W[1]-hard wrt. the number k of swaps. Their reduction relies on the fact that the given graph contains cliques and stars, and the preferences of the agents may contain ties. Our results for P-ESM that LOCAL-STA is W[1]-hard even if the given graph consists of disjoint edges and the preferences do not have ties.

Finally, we mention that Irving [16] and McDermid et al. [18] studied the complexity of computing stable matchings in the marriage setting with preference lists, requiring additionally that the matching be man-exchange stable, that is, no two men form an exchange-blocking pair. The resulting problem is NP-hard, even if the men’s preference lists are of length at most c for each $c \geq 4$ [18], but it becomes polynomial-time solvable when the men’s preference lists have length at most three [16]. This contrasts with COALITIONAL EXCHANGE-STABLE MARRIAGE which, as we show, remains NP-hard even when all preference lists have length at most three.

Organization. In Section 2, we introduce relevant concepts and notation, and define our central problems. In Section 3, we investigate the complexity of deciding (coalitional) exchange-stability, both when the preferences are complete and when the preferences length are bounded. In Section 4, we provide algorithms for profiles with preference length bounded by three. In Section 5, we turn to the local search variant of reaching exchange-stability.

2 Basic Definitions and Observations

For each natural number t , we denote the set $\{1, 2, \dots, t\}$ by $[t]$.

Let $V = \{1, 2, \dots, 2n\}$ be a set of $2n$ agents. Each agent $i \in V$ has a nonempty subset of agents $V_i \subseteq V$ which he finds *acceptable* as a partner and has a *strict preference list* \succ_i on V_i (i.e., a linear order on V_i). The *length* of preference list \succ_i is defined as the number of acceptable agents of i , i.e., $|V_i|$. Here, $x \succ_i y$ means that i *prefers* x to y .

We assume that the acceptability relation among the agents is *symmetric*, i.e., for each two agents x and y it holds that x is acceptable to y if and only if y is acceptable to x . For two agents x and y , we call x *most acceptable* to y if x is a maximal element in the preference list of y . For notational convenience, we write $X \succ Y$ to indicate that for each pair of agents $x \in X$ and $y \in Y$ it holds that $x \succ y$.

A *preference profile* \mathcal{P} is a tuple $(V, (\succ_i)_{i \in V})$ consisting of an agent set V and a collection $(\succ_i)_{i \in V}$ of preference lists for all agents $i \in V$. For a graph G , by $V(G)$ and $E(G)$ we refer to its vertex set and edge set, respectively. Given a vertex $v \in V(G)$, by $N_G(v)$ and $d_G(v)$ we refer to the neighborhood of v and degree of v in G , respectively. To a preference profile \mathcal{P} with agent set V we assign an *acceptability graph* $G(\mathcal{P})$ which has V as its vertex set and two agents are connected by an edge if they find each other acceptable. A preference profile \mathcal{P} may have the following properties:

- It is *bipartite*, if the agent set V can be partitioned into two agent sets U and W of size n each, such that each agent from one set has a preference list over a subset of the agents from the other set.
- It has *complete* preferences if the underlying acceptability graph $G(\mathcal{P})$ is a complete graph or a complete bipartite graph on two disjoint sets of vertices of equal size; otherwise it has *incomplete* preferences.

Profile \mathcal{P} has *bounded length* d if each preference list in \mathcal{P} has length at most d .

(Coalitional) exchange-stable matchings. A *matching* M for a given profile \mathcal{P} is a subset of disjoint edges from the underlying acceptability graph $G(\mathcal{P})$. Given a matching M for \mathcal{P} , and two agents x and y , if it holds that $\{x, y\} \in M$, then we use $M(x)$ (resp. $M(y)$) to refer to y (resp. x), and we say that x and y are their respective assigned *partners* under M and that they are *matched* to each other; otherwise we say that $\{x, y\}$ is an *unmatched pair* under M . If an agent x is *not* assigned any partner by M , then we say that x is *unmatched by* M and we put $M(x) = x$. We assume that each agent x prefers to be matched than remaining unmatched. To formalize this, we will always say that x prefers all acceptable agents from V_x to himself x .

A matching M is *perfect* if every agent is assigned a partner. It is *maximal* if for each unmatched pair $\{x, y\} \in E(G(\mathcal{P})) \setminus M$ it holds that x or y is matched under M . For two agents x, y , we say that x *envies* y under M if x prefers the partner of y , i.e., $M(y)$, to his partner $M(x)$. We omit the “under M ” if it is clear from the context.

We say that matching M admits an *exchange-blocking coalition* (in short *ebc*) if there exists a sequence $\rho = (x_0, x_1, \dots, x_{r-1})$ of r agents, $r \geq 2$, such that each agent x_i envies her successor x_{i+1} in ρ (index $i+1$ taken modulo r). The *size* of an ebc is defined as the number of agents in the sequence. An *exchange-blocking pair* (in short *ebp*) is an ebc of size two. A matching M of \mathcal{P} is *exchange-stable* (resp. *coalitional exchange-stable*) if it does not admit any ebp (resp. ebc.) Note that an coalitional exchange-stable matching is exchange-stable.

For an illustration, let us consider the following example.

Example 1. The following bipartite preference profile \mathcal{P} with agent sets $U = \{x, y, z\}$ and $W = \{a, b, c\}$ admits 2 (coalitional) exchange-stable matchings M_1 and M_2 with $M_1 = \{\{x, c\}, \{y, b\}, \{z, a\}\}$ (marked in red boxes) and $M_2 = \{\{x, b\}, \{y, c\}, \{z, a\}\}$ (marked in blue boxes). Matching M_3 with $M_3 = \{\{x, c\}, \{y, a\}, \{z, b\}\}$ is not exchange-stable and hence not coalitional exchange-stable since for instance (y, z) is an exchange-blocking pair of M_3 .

$$\begin{array}{l} x: a \succ \boxed{b} \succ \boxed{c}, \quad a: y \succ x \succ \boxed{z}, \\ y: \boxed{b} \succ a \succ \boxed{c}, \quad b: \boxed{x} \succ \boxed{y} \succ z, \\ z: \boxed{a} \succ c \succ b, \quad c: \boxed{x} \succ \boxed{y} \succ z. \end{array}$$

As already observed by Cechlárová and Manlove [10], exchange-stable (or coalitional exchange-stable) matchings may not exist, even for bipartite profiles with complete preferences.

Example 2. The following bipartite preference profile \mathcal{P} with agent sets $U = \{x, y, z\}$ and $W = \{a, b, c\}$ admits one exchange-stable matching M with $M = \{\{x, b\}, \{y, c\}, \{z, a\}\}$. However, it is not coalitional exchange-stable as for instance (x, y, z) is an ebc.

$$\begin{array}{l} x: a \succ b \succ c, \quad a: y \succ z \succ x, \\ y: b \succ c \succ a, \quad b: z \succ x \succ y, \\ z: c \succ a \succ b, \quad c: x \succ y \succ z. \end{array}$$

By assumption, if a matching M is coalitional exchange-stable, then it is maximal.

Lemma 1. Every exchange-stable matching is maximal.

Proof. Let \mathcal{P} be a preference profile and M be an exchange-stable matching of \mathcal{P} .

Suppose, for the sake of contradiction, that there exists an unmatched pair $\{x, y\} \in E(G(\mathcal{P})) \setminus M$ with $M(x) = x$ and $M(y) = y$. Since x and y are acceptable to each other, by our assumption that each agent prefers to be matched than remaining unmatched, we obtain that x prefers y to $M(x)$, and y prefers x to $M(y)$. In other words, (x, y) is an exchange-blocking pair of M , a contradiction. \square

Moreover, ebc can only occur in cycles:

Lemma 2. Let \mathcal{P} be a preference profile and let M be a maximal matching of \mathcal{P} . Then, for each ebc ρ of M with $\rho = (x_0, \dots, x_{r-1})$ it holds that $\{x_0, M(x_0), \dots, x_{r-1}, M(x_{r-1})\}$ forms a (not necessarily induced) cycle in $G(\mathcal{P})$.

Proof. Let \mathcal{P}, M, ρ be as defined in the statement with $\rho = (x_0, \dots, x_{r-1})$. By definition each agent x_i , $0 \leq i \leq r-1$, envies his successor x_{i+1} in ρ ($i+1$ taken modulo r). In other words, $\{x_i, M(x_{i+1})\} \in E(G(\mathcal{P}))$ holds for each $i \in \{0, \dots, r-1\}$. Since M is maximal, for each $x_i \in \rho$ it must also hold that agent x_i or agent $M(x_{i+1})$ must be matched, i.e., $x_i \neq M(x_i)$ or $x_{i+1} \neq M(x_{i+1})$.

Define the following pieces for all $i \in \{0, \dots, r-1\}$: $(s_i) := (M(x_i), x_i)$ if $\{x_i, M(x_i)\} \in M$; $(s_i) = (x_i)$ otherwise. Then, it is straightforward to verify that $(s_0, s_1, \dots, s_{r-1}, M(x_0))$ is a cycle in $G(\mathcal{P})$. \square

Central problem definitions. We are interested in the computational complexity of deciding whether a given profile admits a coalitional exchange-stable matching.

COALITIONAL EXCHANGE-STABLE ROOMMATES (CESR)

Input: A preference profile \mathcal{P} .

Question: Does \mathcal{P} admit a coalitional exchange-stable matching?

The bipartite restriction of CESR, called COALITIONAL EXCHANGE-STABLE MARRIAGE (CESM), has as input a *bipartite* preference profile. EXCHANGE-STABLE ROOMMATES (ESR) and EXCHANGE-STABLE MARRIAGE (ESM) are defined analogously.

We are also interested in the case when the preferences have bounded length. In this case, not every coalitional exchange-stable (or exchange-stable) matching is perfect. In keeping with the literature [9, 10], we focus on the perfect case.

d -COALITIONAL EXCHANGE-STABLE ROOMMATES (d -CESR)

Input: A preference profile \mathcal{P} with preferences of bounded length d .

Question: Does \mathcal{P} admit a coalitional exchange-stable and *perfect* matching?

We analogously define the bipartite restriction d -COALITIONAL EXCHANGE-STABLE MARRIAGE (d -CESM), and the exchange-stable variants d -EXCHANGE-STABLE ROOMMATES (d -ESR) and d -EXCHANGE-STABLE MARRIAGE (d -ESM). Note that the above problems are contained in NP [10].

Finally, we investigate a local search variant regarding exchange-stability. To this end, given two matchings M and N of the same profile \mathcal{P} , we say that M is *one-swap reachable* from N if there exists an exchange-blocking pair (x, y) of N such that $M = (N \setminus \{\{x, N(x)\}, \{y, N(y)\}\}) \cup \{\{x, y\}, \{N(x), N(y)\}\}$. Accordingly, we say that M is *k -swaps reachable* from N if there exists a sequence $(M_0, M_1, \dots, M_{k'})$ of k' matchings of profile \mathcal{P} such that (a) $k' \leq k$, $M_0 = N$, $M_{k'} = M$, and (b) for each $i \in [k']$, M_i is one-swap reachable from M_{i-1} .

For an illustration, let us consider Example 1 again.

Example 3. Consider matching M_3 from Example 1, which admits two ebps (x, z) and (y, z) . If we let y and z exchange their partners, then we reach M_1 , which is (coalitional) exchange-stable. If we let x and z exchange their partners, however, then we reach matching $M_4 = \{\{x, b\}, \{y, a\}, \{z, c\}\}$, which will never reach an exchange-stable matching due to the following: (x, y) is a unique exchange-blocking pair of M_4 . If we let x and y exchange partners under M_4 , then we obtain matching M_5 with $M_5 = \{\{x, a\}, \{y, b\}, \{z, c\}\}$, and it admits a unique exchange-blocking pair (a, b) . If we let a and b exchange partners under M_5 , then we obtain matching M_4 , again.

The local search problem variant that we are interested in is defined as follows:

PATH TO EXCHANGE-STABLE MARRIAGE (P-ESM)

Input: A bipartite preference profile \mathcal{P} , a matching M_0 of \mathcal{P} , and an integer k .

Question: Does \mathcal{P} admit an exchange-stable matching M which is k -swap reachable from M_0 ?




Figure 1: The three possible exchange-stable matchings discussed in Lemma 4.

3 Deciding (Coalitional) Exchange-Stability is NP-complete

Cechlárová and Manlove [10] proved NP-completeness for deciding whether a profile with complete and strict preferences admits an exchange-stable matching, by reducing from the NP-complete R-3SAT problem, where each clause has at most three literals and each literal appears at most two times [14]. It is, however, not immediate how to adapt Cechlárová and Manlove’s proof to show hardness for coalitional exchange-stability since their constructed exchange-stable matching is not always coalitional exchange-stable. To obtain a hardness reduction for coalitional exchange-stable, we first study the case when the preferences have length bounded by three, and show that 3-COALITIONAL EXCHANGE-STABLE MARRIAGE is NP-hard, even for strict preferences. The idea is different than that by Cechlárová and Manlove. To simplify the reduction, we will reduce from an NP-complete variant of R-3SAT.

(2,2)-3SAT

Input: A Boolean formula $\phi(X)$ with variable set X in 3CNF (i.e., a set of clauses each containing at most 3 literals) such that no clause contains both the positive and the negated literal of the same variable, and each literal appears *exactly* two times.

Question: Is ϕ satisfiable?

Lemma 3. (2,2)-3SAT is NP-complete.

Proof. Clearly, the problem belongs to NP. We provide a reduction from the NP-complete R-3SAT, where each clause has at most three literals and each literal appears at most two times [14].


First, we assume that no variable appears only negatively or only positively: if this were the case for some variable, then we could set it to true or false and simplify the formula. Second, we assume that no clause contains both the positive and negated literal of the same variable more as otherwise it is always satisfiable and we can delete it from the formula.

Now, for each literal lit_i (either x_i or \bar{x}_i) which appears only once, add two new variables a_i and b_i , and four new clauses $(\text{lit}_i, a_i, \bar{b}_i)$, (a_i, \bar{b}_i) , (\bar{a}_i, b_i) , and (\bar{a}_i, b_i) .

It is straight-forward to see that in the constructed instance no clause contains the positive and negated literal of the same variable and each literal appears exactly two times. Moreover, the original instance is a yes-instance if and only if the newly constructed instance is a yes-instance for (2,2)-3SAT since the newly added variables only affect the newly added clauses which can be satisfied by setting all newly added variables to true. \square

A crucial ingredient for our reduction is the following *switch-gadget* which enforces that each exchange-stable matching results in a valid truth assignment. The gadget and its properties are summarized in the following lemma.

Lemma 4. Let \mathcal{P} be a bipartite preference profile on agent sets U and W . Let $A = \{a^z \mid z \in \{0, 1, \dots, 6\}\}$ and $B = \{b^z \mid z \in \{0, 1, \dots, 6\}\}$ be two disjoint sets of agents, and let $Q = \{\alpha, \beta, \gamma, \delta\}$ be four further distinct agents with $A \cup \{\alpha, \gamma\} \subseteq U$ and $B \cup \{\beta, \delta\} \subseteq W$. The preferences of the agents from A and B are as follows; the preferences of the other agents are arbitrary but fixed.



Define the following matchings:

$$N^1 := \{\{\alpha, b^0\}, \{a^6, \delta\}\} \cup \{\{a^{z-1}, b^z\} \mid z \in [6]\},$$

$$N^2 := \{\{a^0, \beta\}, \{\gamma, b^6\}\} \cup \{\{a^z, b^{z-1}\} \mid z \in [6]\},$$

$$N^D := \{\{\alpha, b^0\}, \{a^0, \beta\}, \{a^6, \delta\}, \{\gamma, b^6\}, \{a^1, b^2\}, \{a^2, b^1\}, \{a^3, b^3\}, \{a^4, b^5\}, \{a^5, b^4\}\}.$$

See Figure 1 for an illustration of the gadget and the matchings above. Every perfect matching M of \mathcal{P} satisfies the following,

- (1) If M is exchange-stable, then (i) either $N^1 \subseteq M$, or (ii) $N^2 \subseteq M$, or (iii) $N^D \subseteq M$.
- (2) If $N^1 \subseteq M$, then every ebc of M which involves an agent from A (resp. B) also involves α (resp. δ).
- (3) If $N^2 \subseteq M$, then every ebc of M which involves an agent from A (resp. B) also involves γ (resp. β).
- (4) If $N^D \subseteq M$, then every ebc of M which involves an agent from A (resp. B) also involves an agent from $\{\alpha, \gamma\}$ (resp. $\{\beta, \delta\}$).

Proof. For statement (1), since M is perfect, every agent is matched. In particular, $M(a^3) \in \{b^2, b^3, b^4\}$. We show that if M is exchange-stable, then the partner $M(a^3)$ decides which of the three cases in the first statement holds.

If $M(a^3) = b^2$, then $M(a^2) \neq b^3$ as otherwise (b^2, b^3) is an exchange-blocking pair. Hence, $M(a^2) = b^1$ and $M(b^3) = a^4$ since b^2 , a^2 , and a^3 are already matched to other agents. By the acceptability relations, $M(a^1) = b^0$, $M(a^0) = \beta$, $M(b^4) = a^5$, $M(b^5) = a^6$, and $M(b^6) = \gamma$. This implies that M satisfies the condition given in case (1ii).

If $M(a^3) = b^3$, then $M(a^2) \neq b^2$ and $M(a^4) \neq b^4$ as otherwise (a^2, a^3) or (b^3, b^4) is an exchange-blocking pair. Hence, $M(a^2) = b^1$ and $M(a^4) = b^5$ since b^3 is already matched to a^3 . By the acceptability relations, $M(b^2) = a^1$, $M(b^4) = a^5$, $M(a^0) = \beta$, $M(a^6) = \delta$, $M(b^0) = \alpha$, $M(b^6) = \gamma$. This implies that M satisfies the condition given in case (1iii).

If $M(a^3) = b^4$, then $M(a^4) \neq b^3$ as otherwise (a^3, a^4) is an exchange-blocking pair. Hence, $M(a^4) = b^5$ and $M(b^3) = a^2$ since b^5 and a^2 are the only agents available to a^4 and b^3 , respectively. By the acceptability relations, $M(a^5) = b^6$, $M(a^6) = \delta$, $M(b^2) = a^1$, $M(b^1) = a^0$, and $M(b^0) = \alpha$. This implies M satisfies the condition given in case (1i). Together, this completes the proof for Statement (1).

To show Statements (2)–(4), we show their contra-positives, by means of a straightforward verification. For Statement (2), assume that $N^1 \subseteq M$ and let ρ denote an arbitrary ebc of M which does not involve α .

We aim to show that ρ does not involve any agent from A . Suppose, for the sake contradiction, that ρ involves an agent x with $x \in A$.

Certainly, ρ cannot involve any agent from $S = \{a^0, a^2, a^5\}$ since each of the agents from S already receives his most-preferred agent. Moreover, $x \neq a^1$ since a^1 only envies the partner of b^0 , which is α and $\alpha \notin \rho$. Consequently, $x \neq a^3$, because a^3 envies only $a^1, a^2 \in S \cup \{a^1\}$. Thus, $x \neq a^4$ because a^4 envies only a^3 and a^2 . Finally, $x \neq a^6$ because a^6 envies only a^4 , a contradiction.

Analogously, assume that $N^1 \subseteq M$ and let ρ denote an arbitrary ebc of M which does not involve δ . Certainly, ρ cannot involve any agent from $S = \{b^1, b^4, b^6\}$ since each of the agents from S already receives his most-preferred agent. Moreover, $x \neq b^5$ since b^5 only envies the partner of a^6 , which is δ , but $\delta \notin \rho$. Consequently, $x \neq b^3$. It follows that, $x \neq b^2$, because b^2 envies only b^3 and b^4 . Finally, $x \neq b^0$, because b^0 envies only b^2 , a contradiction. This completes the proof for Statement (2).

For Statement (3), assume that $N^2 \subseteq M$ and let ρ denote an arbitrary ebc of M which does not involve γ . Suppose, for the sake contradiction, that ρ involves an agent x with $x \in A$. Certainly, $x \notin \{a^1, a^3, a^6\}$ since each of these agents is matched with his most preferred partner. Moreover, $x \neq a^5$ since $\gamma \notin \rho$. Consequently, $x \notin \{a^4, a^2, a^0\}$ by an analogous reasoning, a contradiction.

Analogously, assume that $N^2 \subseteq M$ and let ρ denote an arbitrary ebc of M which does not involve β . Suppose, for the sake contradiction, that ρ involves an agent x with $x \in B$. Certainly, $x \notin \{b^0, b^3, b^5\}$ since each of these agents is matched with his most preferred partner. Moreover, $x \neq b^1$ since $\beta \notin \rho$. Consequently, $x \notin \{b^2, b^4, b^6\}$ by an analogous reasoning, a contradiction. This completes the proof for Statement (3).

For Statement (4), assume that $N^D \subseteq M$ and let ρ denote an arbitrary ebc of M which involves neither α nor γ . Suppose, for the sake contradiction, that ρ involves an agent x with $x \in A$. Certainly, $x \notin \{a^1, a^5\}$ since a^1 only envies α and a^5 only envies γ , but neither α nor γ is included in ρ . Consequently, $x \neq a^3$ since a^3 only envies a^1 . Using a similar reasoning, we infer that $x \notin \{a^2, a^4, a^6, a^0\}$. Altogether, no agent from A is involved in ρ .

Analogously, assume that $N^D \subseteq M$ and let ρ denote an arbitrary ebc of M which involves neither β nor δ . Certainly, $x \notin \{b^1, b^5\}$ since b^1 only envies β and b^5 only envies δ . Consequently, $x \neq b^3$ since b^3 only envies b^5 . Using a similar reasoning, we infer that $x \notin \{b^2, b^4, b^6, b^0\}$. Altogether, no agent from B is involved in ρ . This completes the proof for Statement (4). \square

Using Lemma 4, we can show NP-completeness for bounded preference length.

Theorem 1. *3-CESM, 3-ESM, 3-CESR, and 3-ESR are NP-complete.*

Proof. As already mentioned [10], by checking for cycles in the envy graph all discussed problems are in NP. For the sake of completeness, we show the NP-containment here. Given a preference profile \mathcal{P} and a matching M , we can check in polynomial time whether M is coalitional exchange-stable for \mathcal{P} as follows. Compute an auxiliary directed graph H that has the agent set V as vertices and an arc (u, v) if agent u envies v under M . Observe that M admits an exchange-blocking coalition if and only if there is a directed cycle in H . Thus, it suffices to check for a directed cycle in H . It follows that all problems are in NP.

For the NP-hardness, it is enough to show that 3-CESM and 3-ESM are NP-hard. We use the same reduction from (2,2)-3SAT for both. Let (X, C) be an instance of (2,2)-3SAT where $X = \{x_1, x_2, \dots, x_n\}$ is the set of variables and $\phi = \{C_1, C_2, \dots, C_m\}$ the set of clauses.

$$\begin{array}{l}
\forall i \in [\hat{n}]: \\
\bullet v_i: [y_i] \succ [\bar{y}_i], \\
\bullet x_i: [w_i] \succ [b_{i,o_1(x_i)}^0], \\
\bullet \bar{x}_i: [w_i] \succ [b_{i,o_1(\bar{x}_i)}^0], \\
\forall j \in [\hat{m}]: \\
\bullet c_j: [E_j], \\
\bullet f_j^i: d_j \succ b_{i,j}^6, \\
\bullet a_{i,j}^0: [b_{i,j}^1] \succ [e_j^i], \\
\bullet a_{i,j}^1: [b_{i,j}^0] \succ [b_{i,j}^2] \succ b_{i,j}^1, \\
\bullet a_{i,j}^3: [b_{i,j}^2] \succ [b_{i,j}^3] \succ [b_{i,j}^4], \\
\bullet a_{i,j}^4: [b_{i,j}^4] \succ [b_{i,j}^3] \succ [b_{i,j}^5], \\
\bullet a_{i,j}^5: [b_{i,j}^6] \succ [b_{i,j}^4] \succ b_{i,j}^5, \\
\bullet a_{i,j}^6: [b_{i,j}^5] \succ [\delta_{i,j}], \\
\bullet w_i: [x_i] \succ [\bar{x}_i], \\
\bullet y_i: [v_i] \succ [a_{i,o_2(x_i)}^6], \\
\bullet \bar{y}_i: [v_i] \succ [a_{i,o_2(\bar{x}_i)}^0], \\
\bullet d_j: [F_j], \\
\bullet e_j^i: c_j \succ a_{i,j}^0, \\
\bullet b_{i,j}^0: [a_{i,j}^1] \succ [\alpha_{i,j}], \\
\bullet b_{i,j}^1: [a_{i,j}^0] \succ [a_{i,j}^2] \succ a_{i,j}^1, \\
\bullet b_{i,j}^3: [a_{i,j}^4] \succ [a_{i,j}^3] \succ [a_{i,j}^2], \\
\bullet b_{i,j}^4: [a_{i,j}^3] \succ [a_{i,j}^5] \succ a_{i,j}^4, \\
\bullet b_{i,j}^5: [a_{i,j}^6] \succ [a_{i,j}^4] \succ a_{i,j}^5, \\
\bullet b_{i,j}^6: [a_{i,j}^5] \succ [f_j^i].
\end{array}$$

$\forall i \in [\hat{n}], \forall j \in [\hat{m}]$ with $\text{lit}_i \in C_j$:

Figure 2: The preferences constructed in the proof for Theorem 1. Recall that for each literal $\text{lit}_i \in X \cup \bar{X}$, the indices $o_1(\text{lit}_i)$ and $o_2(\text{lit}_i)$ denote the two indices $j < j'$ of the clauses that contain lit_i . For each clause $C_j \in \phi$, the expression $[E_j]$ (resp. $[F_j]$) denotes an arbitrary but fixed order of the agents in E_j (resp. F_j).

We construct a bipartite preference profile on two disjoint agent sets U and W . The set U (resp. W) will be partitioned into three different agent-groups: the variable-agents, the switch-agents, and the clause-agents. The general idea is to use the variable-agents and the clause-agents to determine a truth assignment and satisfying literals, respectively. Then, we use the switch-agents from Lemma 4 to make sure that the selected truth assignment is consistent with the selected satisfying literals. For each $\text{lit}_i \in X \cup \bar{X}$ that appears in two different clauses C_j and C_k with $j < k$, we use $o_1(\text{lit}_i)$ and $o_2(\text{lit}_i)$ to refer to the indices j and k ; recall that in ϕ each literal appears exactly two times. For example, if literal x_i appears in C_3 and C_5 , then $o_1(x_i) = 3$ and $o_2(x_i) = 5$.

For illustration of the construction below, refer to Figure 2.

The variable-agents. For each variable $x_i \in X$, introduce 6 *variable-agents* $v_i, w_i, x_i, \bar{x}_i, y_i, \bar{y}_i$. Add v_i, x_i, \bar{x}_i to U , and w_i, y_i, \bar{y}_i to W . For each literal $\text{lit}_i \in X \cup \bar{X}$ let $y(\text{lit}_i)$ denote the corresponding Y -variable-agent, that is, $y(x_i) = y_i$ and $y(\bar{x}_i) = \bar{y}_i$. Define $\bar{X} := \{\bar{x}_i \mid i \in [\hat{n}]\}$, and $\bar{Y} := \{\bar{y}_i \mid i \in [\hat{n}]\}$.

The clause-agents. For each clause $C_j \in C$, introduce two *clause-agents* c_j, d_j . Further, for each literal $\text{lit}_i \in C_j$ with $\text{lit} \in \{x, \bar{x}\}$, introduce two more *clause-agents* e_j^i, f_j^i . Add c_j, f_j^i to U , and d_j, e_j^i to W .

For each clause $C_j \in \phi$, define $E_j := \{e_j^i \mid \text{lit}_i \in C_j\}$, and $F_j := \{f_j^i \mid \text{lit}_i \in C_j\}$. Moreover, define $E := \bigcup_{C_j \in \phi} E_j$ and $F := \bigcup_{C_j \in \phi} F_j$.

The switch-agents. For each clause $C_j \in C$, and each literal $\text{lit}_i \in C_j$ introduce fourteen *switch-agents* $a_{i,j}^z, b_{i,j}^z$, $z \in \{0, 1, \dots, 6\}$. Define $A_{i,j} = \{a_{i,j}^z \mid z \in \{0, 1, \dots, 6\}\}$ and $B_{i,j} = \{b_{i,j}^z \mid z \in \{0, 1, \dots, 6\}\}$. Add $A_{i,j}$ to U and $B_{i,j}$ to W .




Figure 3: Relevant part of the acceptability graph for variable x_4 and clauses C_3 , and C_5 discussed in the proof of Theorem 1.

In total, we have the following agent sets:

$$U := \{v_i \mid i \in [\hat{n}]\} \cup X \cup \bar{X} \cup \{c_j \mid j \in [\hat{m}]\} \cup F \cup \bigcup_{C_j \in \phi \wedge \text{lit}_i \in C_j} A_{i,j},$$

$$W := \{w_i \mid i \in [\hat{n}]\} \cup Y \cup \bar{Y} \cup \{d_j \mid j \in [\hat{m}]\} \cup E \cup \bigcup_{C_j \in \phi \wedge \text{lit}_i \in C_j} B_{i,j}.$$

Note that we use the same symbol x_i for both the variable and the variable-agent to strengthen the connection. The meaning will, however, be clear from the context.

The preference lists. The preference lists of the agents are shown in Figure 2. Herein, the preferences of the switch-agents of each occurrence of the literal correspond to those given in Lemma 4. Note that all preferences are specified up to defining the agents $\alpha_{i,j}$ and $\delta_{i,j}$, which we do now. Defining them in an appropriate way will connect the two groups of switch-agents that correspond to the same literal as well as literals to clauses. For each literal $\text{lit}_i \in X \cup \bar{X}$, recall that $\text{o}_1(i)$ and $\text{o}_2(i)$ are the indices of the clauses which contain lit_i with $\text{o}_1(i) < \text{o}_2(i)$. Define the agents $\alpha_{i,\text{o}_1(\text{lit}_i)}$, $\delta_{i,\text{o}_1(\text{lit}_i)}$, $\alpha_{i,\text{o}_2(\text{lit}_i)}$, and $\delta_{i,\text{o}_2(\text{lit}_i)}$ as follows:

$$\alpha_{i,\text{o}_1(\text{lit}_i)} := \text{lit}_i, \delta_{i,\text{o}_1(\text{lit}_i)} := b_{i,\text{o}_2(\text{lit}_i)}^0, \alpha_{i,\text{o}_2(\text{lit}_i)} := a_{i,\text{o}_1(\text{lit}_i)}^6, \delta_{i,\text{o}_2(\text{lit}_i)} := y(\text{lit}_i). \quad (1)$$

For an illustration, assume that literal x_4 appears in C_3 and C_5 , and literal \bar{x}_4 only appears in C_2 and C_6 with

$$C_2 = (x_2 \vee \bar{x}_4 \vee \bar{x}_5), \quad C_3 = (x_1 \vee x_4 \vee x_6), \quad C_5 = (x_3 \vee x_4), \quad C_6 = (\bar{x}_4 \vee x_5).$$

Then, $\alpha_{4,3} = x_4$, $\delta_{4,3} = b_{4,5}^0$, $\alpha_{4,5} = a_{4,3}^6$, $\delta_{4,5} = y_4$, $\alpha_{4,2} = \bar{x}_4$, $\delta_{4,2} = b_{4,6}^0$, $\alpha_{4,6} = a_{4,5}^6$, and $\delta_{4,6} = \bar{y}_4$. The relevant part of the acceptability graph for the variable-agents, switch-agents, and clause-agents which correspond to literal x_4 and clauses C_3, C_5 are depicted in Figure 3.

This completes the construction of the instance for 3-CESM, which can clearly be done in polynomial-time. Let \mathcal{P} denote the constructed instance with $\mathcal{P} = (U \uplus W, (\succ_x)_{x \in U \cup W})$. It is straight-forward to verify that \mathcal{P} is bipartite and contains no ties and that each preference list \succ_x has length bounded by three. Before we give the correctness proof, for each literal $\text{lit}_i \in X \cup \bar{X}$ and each clause C_j with $\text{lit}_i \in C_j$ we define the following three matchings:

$$\begin{aligned} N_{i,j}^1 &:= \{\{\alpha_{i,j}, b_{i,j}^0\}, \{a_{i,j}^6, \delta_{i,j}\}\} \cup \{\{a_{i,j}^{z-1}, b_{i,j}^z\} \mid z \in [6]\}, \\ N_{i,j}^2 &:= \{\{a_{i,j}^0, e_j^i\}, \{b_{i,j}^6, f_j^i\}\} \cup \{\{a_{i,j}^z, b_{i,j}^{z-1}\} \mid z \in [6]\}, \\ N_{i,j}^D &:= \{\{\alpha_{i,j}, b_{i,j}^0\}, \{a_{i,j}^0, e_j^i\}, \{a_{i,j}^6, \delta_{i,j}\}, \{f_j^i, b_{i,j}^6\}, \\ &\quad \{a_{i,j}^1, b_{i,j}^2\}, \{a_{i,j}^2, b_{i,j}^1\}, \{a_{i,j}^3, b_{i,j}^3\}, \{a_{i,j}^4, b_{i,j}^5\}, \{a_{i,j}^5, b_{i,j}^4\}\}. \end{aligned} \quad (2)$$

Now we show the correctness, i.e., ϕ admits a satisfying assignment if and only if \mathcal{P} admits a perfect and coalitional exchange-stable (resp. exchange-stable) matching. For the “only if” direction, assume that $\sigma: X \rightarrow \{t, f\}$ is a satisfying assignment for ϕ . Then, we define a perfect matching M as follows.

- For each variable $x_i \in X$, let $M(\bar{x}_i) := w_i$ and $M(v_i) := \bar{y}_i$ if $\sigma(x_i) = t$; otherwise, let $M(x_i) := w_i$ and $M(v_i) := y_i$.
- For each clause $C_j \in \phi$, fix an arbitrary literal whose truth value satisfies C_j and denote the index of this literal as $s(j)$. Then, let $M(c_j) := e_j^{s(j)}$ and $M(f_j^{s(j)}) := d_j$.
- Further, for each literal $\text{lit}_i \in X \cup \bar{X}$ and each clause C_j with $\text{lit}_i \in C_j$, do the following:
 - (a) If $s(j) = i$, then add to M all pairs from $N_{i,j}^1$.
 - (b) If $s(j) \neq i$ and lit_i is set true under σ (i.e., $\sigma(x_i) = t$ iff. $\text{lit}_i = x_i$), then add to M all pairs from $N_{i,j}^D$.
 - (c) If $s(j) \neq i$ and lit_i is set to false under σ (i.e., $\sigma(x_i) = t$ iff. $\text{lit}_i = \bar{x}_i$), then add to M all pairs from $N_{i,j}^2$.

One can verify that M is perfect. Hence, it remains to show that M is coalitional exchange-stable. Note that this would also imply that M is exchange-stable.

Suppose, for the sake of contradiction, that M admits an ebc ρ . First, observe that for each variable-agent $z \in X \cup \bar{X} \cup Y \cup \bar{Y}$ it holds that $M(z)$ either is matched with his most-preferred partner (i.e., either v_i or w_i) or only envies someone who is matched with his most-preferred partner. Hence, no agent from $X \cup \bar{X} \cup Y \cup \bar{Y}$ is involved in ρ . Analogously, no agent from $E \cup F$ is involved in ρ .

Next, we claim the following.

Claim 1. *For each literal $\text{lit}_i \in X \cup \bar{X}$ and each clause C_j with $\text{lit}_i \in C_j$, it holds that neither $\alpha_{i,j}$ nor $\delta_{i,j}$ is involved in ρ .*

Proof. Suppose, for the sake of contradiction, that there exists some $\text{lit}_i \in X \cup \bar{X}$ with $\text{lit}_i \in C_j$ such that $\alpha_{i,j} \in \rho$ or $\delta_{i,j} \in \rho$.

If $\alpha_{i,j} \in \rho$, then by the definition of $\alpha_{i,j}$, it follows that $j = o_2(\text{lit}_i)$ as otherwise $\alpha_{i,j} = \text{lit}_i$ which would be a contradiction because we have just shown that no agent from $X \cup \bar{X}$ is involved in ρ . This implies that $\alpha_{i,j} = a_{i,o_1(\text{lit}_i)}^6 \in \rho$. We infer from the preferences of $a_{i,o_1(\text{lit}_i)}^6$ that $M(a_{i,o_1(\text{lit}_i)}^6) = \delta_{i,o_1(\text{lit}_i)}$. By the definition of M , we have that $N_{i,o_1(\text{lit}_i)}^1 \subseteq M$ or $N_{i,o_1(\text{lit}_i)}^D \subseteq M$. From Lemma 4(2) and Lemma 4(4) (setting $\alpha = \alpha_{i,o_1(\text{lit}_i)}$, $\beta = e_{o_1(\text{lit}_i)}^i$, $\gamma = f_{o_1(\text{lit}_i)}^i$, and $\delta = \delta_{i,o_1(\text{lit}_i)}$) we infer that ρ involves $f_{o_1(\text{lit}_i)}^i \in F$ or ρ involves $\alpha_{i,o_1(\text{lit}_i)}$; observe that $\alpha_{i,o_1(\text{lit}_i)} = \text{lit}_i \in X$. This is a contradiction since we have already proved that no agent from $X \cup \bar{X} \cup F$ is involved in ρ .

If $\delta_{i,j} \in \rho$, then by the definition of $\delta_{i,j}$, we have $j = o_1(\text{lit}_i)$ as otherwise $\delta_{i,j} = y(\text{lit}_i)$ which would be a contradiction because we have just shown that no agent from $Y \cup \bar{Y}$ is involved in ρ . This implies that $\delta_{i,j} = b_{i,o_2(\text{lit}_i)}^0 \in \rho$. By the preferences of $b_{i,o_2(\text{lit}_i)}^0$ we infer that $M(b_{i,o_2(\text{lit}_i)}^0) = \alpha_{i,o_2(\text{lit}_i)}$. By the definition of M , we have that $N_{i,o_2(\text{lit}_i)}^1 \subseteq M$ or $N_{i,o_2(\text{lit}_i)}^D \subseteq M$. By Lemma 4(2) and Lemma 4(4) (setting $\alpha = \alpha_{i,o_2(\text{lit}_i)}$, $\beta = e_{o_2(\text{lit}_i)}^i$, $\gamma = f_{o_2(\text{lit}_i)}^i$, and $\delta = \delta_{i,o_2(\text{lit}_i)}$), we infer that ρ involves $e_{o_2(\text{lit}_i)}^i \in E$ or $\delta_{i,o_2(\text{lit}_i)}$; observe that $\delta_{i,o_2(\text{lit}_i)} = y(\text{lit}_i)$. This is a contradiction since we have already proved that no agent from $Y \cup \bar{Y} \cup E$ is involved in ρ . (of Claim 1) \diamond

Now, using the above observations and claim, we continue with the proof. We successively prove that no agent is involved in ρ , starting with the agents in U .

- If v_i is involved in ρ for some $i \in [\hat{n}]$, then he only envies someone who is matched with y_i . By the preferences of y_i , this means that $M(y_i) = a_{i,o_2(x_i)}^6$ and that v_i envies $a_{i,o_2(x_i)}^6$. Hence, $a_{i,o_2(x_i)}^6$ is also involved in ρ . Moreover, since $M(a_{i,o_2(x_i)}^6) = y_i$, we have $N_{i,o_2(x_i)}^1 \subseteq$

M or $N_{i,o_2(x_i)}^D \subseteq M$. By Lemma 4(2) and Lemma 4(4) (setting $\alpha = \alpha_{i,o_2(x_i)}$, $\beta = e_{o_2(x_i)}^i$, $\gamma = f_{o_2(x_i)}^i$, and $\delta = \delta_{i,o_2(x_i)}$), ρ involves an agent from $\{\alpha_{i,o_2(x_i)}, f_{o_2(x_i)}^i\}$. Since no agent from F is involved in ρ , it follows that ρ involves $\alpha_{i,o_2(x_i)}$, a contradiction to Claim 1.

- Analogously, if $c_j \in \rho$ for some $j \in [\hat{m}]$, then this means that E_j contains two agents e_j^i and e_j^t such that $M(c_j) = e_j^t$ but c_j prefers e_j^i to e_j^t , and $M(e_j^t) \in \rho$. Since M is perfect and c_j is not available, it follows that $M(e_j^t) = a_{i,j}^0$, implying that $a_{i,j}^0 \in \rho$. Moreover, by the definition of M we have that $N_{i,j}^2 \subseteq M$ or $N_{i,j}^D \subseteq M$. By Lemmas 4(3)–(4) (setting $\alpha = \alpha_{i,j}$, $\beta = e_j^i$, $\gamma = f_j^i$, and $\delta = \delta_{i,j}$), ρ involves an agent from $\{\alpha_{i,j}, f_j^i\}$, a contradiction since no agent from F_j is involved in ρ and by Claim 1 $\alpha_{i,j}$ is not in ρ .
- Analogously, we can obtain a contradiction if w_i with $i \in [\hat{n}]$ is in ρ : By the definition of M , if $w_i \in \rho$, then $M(x_i) = b_{i,o_1(x_i)}^0$ and w_i envies $b_{i,o_1(x_i)}^0$. Hence, $b_{i,o_1(x_i)}^0$ is also involved in ρ . Moreover, since $M(b_{i,o_1(x_i)}^0) = x_i$, it follows that $N_{i,o_1(x_i)}^1 \subseteq M$ or $N_{i,o_1(x_i)}^D \subseteq M$. By Lemmas 4(2) and (4) (setting $\alpha = \alpha_{i,o_1(x_i)}$, $\beta = e_{o_1(x_i)}^i$, $\gamma = f_{o_1(x_i)}^i$, and $\delta = \delta_{i,o_1(x_i)}$), ρ involves an agent from $\{e_{o_1(x_i)}^i, \delta_{i,o_1(x_i)}\}$. Since no agent from E is involved in ρ , it follows that ρ involves $\delta_{i,o_1(x_i)}$, a contradiction to Claim 1.
- Again, analogously, if $d_j \in \rho$ for some $j \in [\hat{m}]$, then we obtain that $\delta_{i,j}$ is involved in ρ , which is a contradiction to Claim 1.
- Finally, if ρ involves an agent from $A_{i,j}$ (resp. $B_{i,j}$), then by Lemma 4(2) and (4) (setting $\alpha = \alpha_{i,j}$, $\beta = e_j^i$, $\gamma = f_j^i$, and $\delta = \delta_{i,j}$), it follows that ρ involves an agent from $\{\alpha_{i,j}, f_j^i\}$ (resp. $\{\beta_{i,j}, e_j^i\}$), a contradiction to our observation and to Claim 1.

Summarizing, we have showed that M is coalitional exchange-stable and exchange-stable.

For the “if” direction, assume that M is a perfect and exchange-stable matching for \mathcal{P} . We show that there is a satisfying assignment for ϕ . Note that this then also implies that, if M is perfect and coalitional exchange-stable, then there is a satisfying assignment for ϕ .

We claim that the selection of the partner of w_i defines a satisfying truth assignment for ϕ . More specifically, define a truth assignment $\sigma: X \rightarrow \{\mathbf{t}, \mathbf{f}\}$ with $\sigma(x_i) = \mathbf{t}$ if $M(w_i) = \bar{x}_i$, and $\sigma(x_i) = \mathbf{f}$ otherwise. We claim that σ satisfies ϕ . To this end, consider an arbitrary clause C_j and the corresponding clause-agent. Since M is perfect, it follows that $M(c_j) = e_j^i$ for some $\text{lit}_i \in C_j$. Since e_j^i is not available, it also follows that $M(a_{i,j}^0) = b_{i,j}^1$. By Lemma 4(1) (setting $\alpha = \alpha_{i,j}$, $\beta = e_j^i$, $\gamma = f_j^i$, and $\delta = \delta_{i,j}$), it follows that $N_{i,j}^1 \subseteq M$. In particular, $M(\alpha_{i,j}) = b_{i,j}^0$ so that $\alpha_{i,j}$ is not available to other agents anymore.

Now, if we can show that lit_i is matched to $b_{i,o_1(\text{lit}_i)}^0$, then since M is perfect, we have $M(w_i) = \bar{x}_i$ if $\text{lit}_i = x_i$, and $M(w_i) = \bar{x}_i$ otherwise. By definition, we have $\sigma(x_i) = \mathbf{t}$ if $\text{lit}_i = x_i$ and $\sigma(x_i) = \mathbf{f}$ otherwise. Thus, C_j is satisfied under σ , implying that σ is a satisfying assignment. It remains to show that lit_i is matched to $b_{i,o_1(\text{lit}_i)}^0$. We distinguish between two cases;

- If $j = o_1(\text{lit}_i)$, then by assumption, $\alpha_{i,o_1(\text{lit}_i)}$ is matched to $b_{i,o_1(\text{lit}_i)}^0$, as required.
- If $j = o_2(\text{lit}_i)$, then by definition, it holds that $\alpha_{i,j} = a_{i,o_1(\text{lit}_i)}^6$ and $\delta_{i,o_1(\text{lit}_i)} = b_{i,j}^0$. In other words, $M(a_{i,o_1(\text{lit}_i)}^6) = \delta_{i,o_1(\text{lit}_i)}$. By Lemma 4(1) (setting $\alpha = \alpha_{i,o_1(\text{lit}_i)}$, $\beta = e_{o_1(\text{lit}_i)}^i$, $\gamma = f_{o_1(\text{lit}_i)}^i$, and $\delta = \delta_{i,o_1(\text{lit}_i)}$), it follows that $N_{i,j}^1 \subseteq M$ or $N_{i,j}^D \subseteq M$. In both cases, it follows that $\alpha_{i,o_1(i)}$ is matched to $b_{i,o_1(i)}^0$.

We have thus proved that there is a satisfying assignment for ϕ if and only if there is a perfect coalitional exchange-stable (resp. exchange-stable) matching for \mathcal{P} . \square

Next, we show how to complete the preferences of the agents constructed in the proof of Theorem 1 to show hardness for complete and strict preferences.

Theorem 2. *CESM and CESR are NP-complete even for complete and strict preferences.*

Proof. CESM is contained in NP by the same argument as in Theorem 1. For NP-hardness we adapt the proof of Theorem 1. Recall that in that proof, for a given (2,2)-3SAT

instance (X, ϕ) with $X = \{x_1, x_2, \dots, x_{\hat{n}}\}$ and $\phi = \{C_1, C_2, \dots, C_{\hat{m}}\}$, we construct two disjoint agent sets U and W with

$$U := \{v_i \mid i \in [\hat{n}]\} \cup X \cup \overline{X} \cup \{c_j \mid j \in [\hat{m}]\} \cup F \cup \bigcup_{C_j \in \phi \wedge \text{lit}_i \in C_j} A_{i,j},$$

$$W := \{w_i \mid i \in [\hat{n}]\} \cup Y \cup \overline{Y} \cup \{d_j \mid j \in [\hat{m}]\} \cup E \cup \bigcup_{C_j \in \phi \wedge \text{lit}_i \in C_j} B_{i,j}, \text{ where}$$

$\overline{X} := \{\overline{x}_i \mid i \in [\hat{n}]\}$, $\overline{Y} := \{\overline{y}_i \mid i \in [\hat{n}]\}$, $A_{i,j} = \{a_{i,j}^z \mid z \in \{0, 1, \dots, 6\}\}$, $B_{i,j} = \{b_{i,j}^z \mid z \in \{0, 1, \dots, 6\}\}$, $E_j := \{e_j^i \mid \text{lit}_i \in C_j\}$, $F_j := \{f_j^i \mid \text{lit}_i \in C_j\}$, and $E := \bigcup_{C_j \in \phi} E_j$ and $F := \bigcup_{C_j \in \phi} F_j$. For each agent $z \in U \cup W$ let L_z denote the preference list of z constructed in the proof. The basic idea is to extend the preference list L_z by appending to it the remaining agents appropriately.

We introduce some more notations. Let \triangleright_U and \triangleright_W denote two arbitrary but fixed linear orders of the agents in U and W , respectively. Now, for each subset of agents $S \subseteq U$ (resp. $S \subseteq W$), let $[S]_{\triangleright}$ denote the fixed order of the agents in S induced by \triangleright_U (resp. \triangleright_W), and let $S \setminus L_z$ denote the subset $\{t \in S \mid t \notin L_z\}$, where $z \in W$ (resp. $z \in U$). Finally, for each agent $z \in U$ (resp. $z \in W$), let R_z denote the subset of agents which do not appear in L_z or in $Y \cup \overline{Y} \cup E$ (resp. $X \cup \overline{X} \cup F$). That is, $R_z := (W \setminus (Y \cup \overline{Y} \cup E)) \setminus L_z$ (resp. $R_z := (U \setminus (X \cup \overline{X} \cup F)) \setminus L_z$).

Now, we define the preferences of the agents as follows.

$$\forall z \in U, z: L_z \succ [Y \cup \overline{Y} \cup E \setminus L_z]_{\triangleright} \succ [R_z]_{\triangleright}, \text{ and } \forall z \in W, z: L_z \succ [X \cup \overline{X} \cup F \setminus L_z]_{\triangleright} \succ [R_z]_{\triangleright}.$$

For instance, the complete preference list of an agent called $a_{i,j}^0$ (corresponding to the literal lit_i which appears in clause C_j) is

$$a_{i,j}^0: b_{i,j}^1 \succ e_j^i \succ [X \cup \overline{X} \cup F \setminus \{e_j^i\}]_{\triangleright} \succ \left[\bigcup_{k=0}^6 B^k \cup C \cup V \setminus \{b_{i,j}^1\} \right]_{\triangleright}.$$

Let \mathcal{P}' denote the newly constructed preference profile. Clearly, the constructed preferences are complete and strict. It remains to show the correctness. Assume that ϕ admits a satisfying assignment $\sigma: X \rightarrow \{t, f\}$. First of all, observe that the following claim holds.

Claim 2. *Besides the agents of $X \cup \overline{X} \cup F \cup Y \cup \overline{Y} \cup E$ every other agent appears as the most-preferred agent of some other agent.*

Using this, we claim the following for each coalitional exchange-stable matching of \mathcal{P}' .

Claim 3. *If M is a coalitional exchange-stable matching for \mathcal{P}' , then*

- (i) *for each agent $z \in U \cup W$ it holds that $M(z) \notin R_z$, and*
- (ii) *for each agent $z \in U \cup W \setminus (X \cup \overline{X} \cup F \cup Y \cup \overline{Y} \cup E)$ it holds that $M(z) \in L_z$.*

Proof. Let M be a coalitional exchange-stable matching of \mathcal{P}' . By Lemma 1, M is maximal. Since $G(\mathcal{P}')$ is complete it follows that M is perfect. For Statement (i), suppose, for the sake of contradiction, that there exists an agent $z_0 \in U \cup W$ such that $M(z_0) \in R_{z_0}$. This means that $M(z_0) \notin X \cup \overline{X} \cup F \cup Y \cup \overline{Y} \cup E$. Now consider the following inductive definition of agents z_i for $i = 1, 2, \dots$ such that

- (a) z_i prefers $M(z_{i-1})$ to $M(z_i)$,
- (b) $M(z_i) \notin X \cup \overline{X} \cup F \cup Y \cup \overline{Y} \cup E$, and
- (c) $z_i \notin \{z_0, z_1, \dots, z_{i-1}\}$.

Let z_{i-1} already be defined. Since $M(z_{i-1}) \notin X \cup \overline{X} \cup F \cup Y \cup \overline{Y} \cup E$, by Claim 2, there exists an agent who considers $M(z_{i-1})$ as the most-preferred agent. Define z_i to be an arbitrary agent who considers $M(z_{i-1})$ as the most-preferred agent. Clearly, (a) holds. Observe that z_0 prefers $M(z_0)$ to $M(z_i)$ since otherwise (z_0, z_1, \dots, z_i) would form an exchange-blocking coalition. Thus, $M(z_i) \in R_{z_0}$, which implies (b). Finally, for the sake of contradiction assume that (c) does not hold, that is, there is an agent z_j with $j < i$ such that $z_j = z_i$. By definition, z_j and z_i consider $M(z_{j-1}) = M(z_{i-1})$ as the most-preferred agent. Since M is a matching, $z_{j-1} = z_{i-1}$, a contradiction to the fact that $z_{i-1} \notin \{z_0, z_1, \dots, z_{i-2}\}$. Hence, we have found an infinite sequence of pairwise distinct agents, a contradiction, proving Statement (i).

For Statement (ii), suppose, for the sake of contradiction, that there exists an agent $z \in U \cup W \setminus (X \cup \overline{X} \cup F \cup Y \cup \overline{Y} \cup E)$ such that $M(z) \notin L_z$. By symmetry, it follows that $z = M(M(z)) \notin L_{M(z)}$. By the first statement, $z \notin R_z$. Hence, $z \in (X \cup \overline{X} \cup F \cup Y \cup \overline{Y} \cup E) \setminus L_z$, a contradiction. (of Claim 3) \diamond

Now, we are ready to show the correctness, i.e., ϕ admits a satisfying assignment if and only if \mathcal{P}' admits a coalitional exchange-stable matching.

For the “only if” direction, assume that ϕ admits a satisfying assignment, say $\sigma: X \rightarrow \{t, f\}$. We claim that the coalitional exchange-stable matching M for \mathcal{P} that we defined in the “only if” direction of the proof for Theorem 1 is a coalitional exchange-stable matching for \mathcal{P}' . Clearly, M is a perfect matching for \mathcal{P}' since $G(\mathcal{P}')$ is a supergraph of $G(\mathcal{P})$. Since each agent $z \in U \cup W$ has $M(z) \in L_z$, for each two agents $z, z' \in U$ (resp. W), it holds that z envies z' only if $M(z') \in L_z$. In other words, if M would admit an exchange-blocking coalition $\rho = (z_0, z_1, \dots, z_{r-1})$ ($r \geq 2$) for \mathcal{P}' , then for each $i \in \{0, 1, \dots, r-1\}$ it must hold that $M(z_i) \in L_{z_{i-1}}$ (z_{i-1} taken modulo r). But then, ρ is also an exchange-blocking coalition for \mathcal{P} , a contradiction to our “only if” part of the proof for Theorem 1.

For the “if” direction, let M be a coalitional exchange-stable matching for \mathcal{P}' . Note that in the “if” part of the proof of Theorem 1 we heavily utilize the properties given in Lemma 4(1). Now, to construct a satisfying assignment for ϕ from M , we will prove that the lemma also holds for profile \mathcal{P}' . To this end, for each literal $\text{lit}_i \in X \cup \overline{X}$ and each clause C_j with $\text{lit}_i \in C_j$, recall the three matchings we have defined in equation (2) (on Page 10):

$$\begin{aligned} N_{i,j}^1 &:= \{\{\alpha_{i,j}, b_{i,j}^0\}, \{a_{i,j}^6, \delta_{i,j}\}\} \cup \{\{a_{i,j}^{z-1}, b_{i,j}^z\} \mid z \in [6]\}, \\ N_{i,j}^2 &:= \{\{a_{i,j}^0, e_j^i\}, \{b_{i,j}^6, f_j^i\}\} \cup \{\{a_{i,j}^z, b_{i,j}^{z-1}\} \mid z \in [6]\}, \\ N_{i,j}^D &:= \{\{\alpha_{i,j}, b_{i,j}^0\}, \{a_{i,j}^0, e_j^i\}, \{a_{i,j}^6, \delta_{i,j}\}, \{f_j^i, b_{i,j}^6\}, \\ &\quad \{a_{i,j}^1, b_{i,j}^2\}, \{a_{i,j}^2, b_{i,j}^1\}, \{a_{i,j}^3, b_{i,j}^3\}, \{a_{i,j}^4, b_{i,j}^5\}, \{a_{i,j}^5, b_{i,j}^4\}\}. \end{aligned}$$

The agents $\alpha_{i,j}$ and $\delta_{i,j}$ were defined according to equation (1); for the sake of completeness, we repeat here:

$$\alpha_{i, \text{lit}_i} := \text{lit}_i, \delta_{i, \text{lit}_i} := b_{i, \text{lit}_i}^0, \alpha_{i, \text{lit}_i} := a_{i, \text{lit}_i}^6, \delta_{i, \text{lit}_i} := y(\text{lit}_i).$$

Claim 4. *Matching M satisfies for each literal $\text{lit}_i \in X \cup \overline{X}$ and each clause $C_j \in \phi$ with $\text{lit}_i \in C_j$, either (i) $N_{i,j}^1 \subseteq M$, or (ii) $N_{i,j}^2 \subseteq M$, or (iii) $N_{i,j}^D \subseteq M$.*

Proof. The proof is almost the same as the one given for Lemma 4(1). We repeat for the sake of completeness.

Since M is coalitional exchange-stable for \mathcal{P}' , by Claim 3(ii) it follows that

$$\forall \text{lit}_i \in X \cup \overline{X}, \forall C_j \in \phi \text{ with } \text{lit}_i \in C_j, \forall z \in \{0, 1, \dots, 6\}: M(a_{i,j}^z) \in L_{a_{i,j}^z}. \quad (3)$$

We distinguish between three subcases.

If $M(a_{i,j}^3) = b_{i,j}^2$, then $M(a_{i,j}^2) \neq b_{i,j}^3$ as otherwise $(b_{i,j}^2, b_{i,j}^3)$ is an exchange-blocking pair. Hence, $M(a_{i,j}^2) = b_{i,j}^1$ and $M(b_{i,j}^3) = a_{i,j}^4$ since $b_{i,j}^2$, $a_{i,j}^2$, and $a_{i,j}^3$ are already matched to others. By (3), $M(a_{i,j}^1) = b_{i,j}^0$, $M(a_{i,j}^0) = e_j^i$, $M(b_{i,j}^4) = a_{i,j}^5$, $M(b_{i,j}^5) = a_{i,j}^6$, and $M(b_{i,j}^6) = f_j^i$. This implies that M contains $N_{i,j}^2$.

If $M(a_{i,j}^3) = b_{i,j}^3$, then $M(a_{i,j}^2) \neq b_{i,j}^2$ and $M(a_{i,j}^4) \neq b_{i,j}^4$ as otherwise $(a_{i,j}^2, a_{i,j}^3)$ or $(b_{i,j}^3, b_{i,j}^4)$ is an exchange-blocking pair. By (3), $M(a_{i,j}^2) = b_{i,j}^1$ and $M(a_{i,j}^4) = b_{i,j}^5$ since $b_{i,j}^3$ is already matched to $a_{i,j}^3$. Again, by (3), $M(b_{i,j}^2) = a_{i,j}^1$, $M(b_{i,j}^4) = a_{i,j}^5$, $M(a_{i,j}^0) = e_j^i$, $M(a_{i,j}^6) = \delta_{i,j}$, $M(b_{i,j}^0) = \alpha_{i,j}$, $M(b_{i,j}^6) = f_j^i$. This implies that M contains $N_{i,j}^D$.

Finally, if $M(a_{i,j}^3) = b_{i,j}^4$, then $M(a_{i,j}^4) \neq b_{i,j}^3$ as otherwise $(a_{i,j}^3, a_{i,j}^4)$ is an exchange-blocking pair. Hence, by (3), $M(a_{i,j}^4) = b_{i,j}^5$ and $M(b_{i,j}^3) = a_{i,j}^2$ since $b_{i,j}^5$ and $a_{i,j}^2$ are the only agents available to $a_{i,j}^4$ and $b_{i,j}^3$, respectively. By the acceptability relations, $M(a_{i,j}^5) = b_{i,j}^6$, $M(a_{i,j}^6) = \delta_{i,j}$, $M(b_{i,j}^2) = a_{i,j}^1$, $M(b_{i,j}^1) = a_{i,j}^0$, and $M(b_{i,j}^0) = \alpha_{i,j}$. This implies that M contains $N_{i,j}^1$. Together, this completes the proof for Statement (4). (of Claim 4) \diamond

Now we show that the function $\sigma: X \rightarrow \{\mathbf{t}, \mathbf{f}\}$ with $\sigma(x_i) = \mathbf{t}$ if $M(w_i) = \bar{x}_i$, and $\sigma(x_i) = \mathbf{f}$ otherwise is a satisfying truth assignment for ϕ . Clearly, ϕ is a valid truth assignment since by Claim 3(ii) every variable agent w_i is matched to either x_i or \bar{x}_i . We claim that σ satisfies ϕ . To this end, consider an arbitrary clause C_j and the corresponding clause-agent c_j . By Claim 3(ii), we know that $M(c_j) = e_j^i$ for some $\text{lit}_i \in C_j$. Since e_j^i is not available, by Claim 3(ii), it also follows that $M(a_{i,j}^0) = b_{i,j}^1$. By Claim 4, it follows that $N_{i,j}^1 \subseteq M$. In particular, $M(\alpha_{i,j}) = b_{i,j}^0$ so that $\alpha_{i,j}$ is not available to other agents anymore.

We aim to show that $\alpha_{i,\mathbf{o}_1(\text{lit}_i)}$ is matched to $b_{i,\mathbf{o}_1(\text{lit}_i)}^0$ by M , which implies that lit_i is not available to w_i since $\alpha_{i,\mathbf{o}_1(\text{lit}_i)} = \text{lit}_i$ by the definition of $\alpha_{i,\mathbf{o}_1(\text{lit}_i)}$. We distinguish two cases:

- If $j = \mathbf{o}_1(\text{lit}_i)$, then by the definition of $\alpha_{i,j}$, it follows that $\alpha_{i,\mathbf{o}_1(\text{lit}_i)}$ is matched to $b_{i,\mathbf{o}_1(\text{lit}_i)}^0$, as required.
- If $j = \mathbf{o}_2(\text{lit}_i)$, then by the definition of $\alpha_{i,j}$, we have $\alpha_{i,j} = a_{i,\mathbf{o}_1(\text{lit}_i)}^6$ and by the definition of $\delta_{i,\mathbf{o}_1(\text{lit}_i)}$ we have $\delta_{i,\mathbf{o}_1(\text{lit}_i)} = b_{i,\mathbf{o}_2(\text{lit}_i)}^0 = b_{i,j}^0$. In particular, since $M(\alpha_{i,j}) = b_{i,j}^0$ we have $M(a_{i,\mathbf{o}_1(\text{lit}_i)}^6) = \delta_{i,\mathbf{o}_1(\text{lit}_i)}$. By Claim 4, it follows that $N_{i,\mathbf{o}_1(\text{lit}_i)}^1 \subseteq M$ or $N_{i,\mathbf{o}_1(\text{lit}_i)}^D \subseteq M$. In both cases, it follows that $\alpha_{i,\mathbf{o}_1(\text{lit}_i)}$ is matched to $b_{i,\mathbf{o}_1(\text{lit}_i)}^0$.

We have just shown that lit_i is *not* available to w_i . Hence, by Claim 3(ii), $M(w_i) = \bar{x}_i$ if $\text{lit}_i = x_i$, and $M(w_i) = \bar{x}_i$ otherwise. By definition, we have that $\sigma(x_i) = \mathbf{t}$ if $\text{lit}_i = x_i$ and $\sigma(x_i) = \mathbf{f}$ otherwise. This implies that C_j is satisfied under σ , implying that σ is a satisfying assignment. \square

4 Algorithms for Bounded Preferences Length

In this section we give algorithms for the profiles with bounded preference length. In Subsection 4.1 we give linear-time algorithms for length at most two, and in Subsection 4.2 we give a fixed-parameter algorithm for length at most three.

4.1 Preference list length at most two

Bounding the length of the preference lists by two allows us to decide whether a coalitional exchange-stable and perfect matching exists in linear time. The reason is that for preferences of length at most two, the underlying acceptability graph consists of disjoint paths and cycles. Together with Lemma 2, we infer that only agents from a cycle may induce exchange-blocking coalitions, and hence can be checked for coalitional exchange-stability in linear time.

ALGORITHM 1: Algorithm for checking coalitional exchange-stability when each preference list has length bounded by two.

Input: A preference profile \mathcal{P} with preferences length at most two.
Output: A coalitional exchange-stable matching for \mathcal{P} or **no** if none exists.

```

1  $M \leftarrow \emptyset$ 
2 if  $G(\mathcal{P})$  contains a connected component of odd size then return no;
3 foreach connected component  $C$  in  $G(\mathcal{P})$  do
4   if  $G(\mathcal{P})[C]$  is a path then
5     Let  $E(G(\mathcal{P})[C]) = \{\{x_0, x_1\}, \{x_1, x_2\}, \dots, \{x_{2t-2}, x_{2t-1}\}\}$ 
6      $M \leftarrow M \cup \{\{x_{2i-2}, x_{2i-1}\} \mid i \in [t]\}$ 
7   if  $G(\mathcal{P})[C]$  is a cycle then
8     Let  $E(G(\mathcal{P})[C]) = \{\{x_0, x_1\}, \{x_1, x_2\}, \dots, \{x_{2t-2}, x_{2t-1}\}, \{x_{2t-1}, x_0\}\}$ 
9      $M_1 \leftarrow \{\{x_{2i-2}, x_{2i-1}\} \mid i \in [t]\}$ 
10     $M_2 \leftarrow \{\{x_{2i-1}, x_{2i}\} \mid i \in [t-1]\} \cup \{\{x_{2t-1}, x_0\}\}$ 
11    if  $M_1$  is coalitional exchange-stable for  $\mathcal{P}$  restricted to  $C$  then  $M \leftarrow M \cup M_1$ ;
12    else if  $M_2$  is coalitional exchange-stable for  $\mathcal{P}$  restricted to  $C$  then  $M \leftarrow M \cup M_2$ ;
13    else return no;
14 return  $M$ 

```

Theorem 3. 2-ESM, 2-ESR, 2-CESM, and 2-CESR can be solved in linear time.

Proof. Let \mathcal{P} be a preference profile with preference-list length bounded by two, meaning that $G(\mathcal{P})$ consists of solely paths and cycles. The algorithm works as follows and we give pseudo code for coalitional exchange-stability in Algorithm 1. For exchange-stability the pseudo code is analogous; we only need to exchange coalitional exchange-stable with exchange-stable in lines 11–12.

We consider each path and cycle independently and accept if and only if each one admits a perfect (coalitional) exchange-stable matching. Since we are aiming for a perfect matching that is exchange-stable or coalitional exchange-stable, by Lemma 2, it suffices to only consider cycles of even length as paths do not induce any exchange-blocking coalitions and cycles of odd length cannot result in a perfect matching.

For cycles of even length, since there are only two possible perfect matchings, it suffices to check whether one of them avoids exchange-blocking coalitions or exchange-blocking pairs. Since constructing all perfect matchings for a path or a cycle (there are at most two) and checking each of them for coalitional exchange-stability or exchange-stability can be done in linear time, the described algorithms run in linear time. \square

4.2 Fixed-parameter algorithm for 3-ESR

We now turn to preference length at most three. In Theorem 1 we have seen that even this case remains NP-hard, even for bipartite preference profiles. Moreover, the proof suggests that a main obstacle that one has to deal with when solving 3-ESM (and hence 3-ESR) are the switch gadgets. Here we essentially show that they are indeed the *only* obstacles, that is, if there are few of them present in the input, then we can solve the problem efficiently. We capture the essence of the switch gadgets with the following structure that we call hourglasses.

Definition 1. Let \mathcal{P} be a preference profile and $V_H \subseteq V$ a subset of $2h$ agents with $V_H = \{u_i, w_i \mid 0 \leq i \leq h-1\}$. We call the subgraph $G(\mathcal{P})[V_H]$ induced by V_H an *hourglass* of height h if it satisfies the following:

- For each $i \in \{0, h-1\}$ the vertex degree of u_i and w_i are both at least two in $G(\mathcal{P})[V_H]$;
- For each $i \in [h-2]$, the vertex degree of u_i and w_i are exactly three in $G(\mathcal{P})[V_H]$;




Figure 4: An hourglass of height three.

- For each $i \in \{0, 1, \dots, h-1\}$ we have $\{u_i, w_i\} \in E(G(\mathcal{P})[V_H])$;
- For each $i \in \{0, 1, \dots, h-2\}$ we have $\{u_i, w_{i+1}\}, \{u_{i+1}, w_i\} \in E(G(\mathcal{P})[V_H])$.

We refer to the agents u_i and w_i from V_H as *layer- i* agents. We call an hourglass H *maximal* if no larger agent subset $V' \supsetneq V(H)$ exists that induces an hourglass.

Given an hourglass H in $G(\mathcal{P})$, we call a matching M for \mathcal{P} *perfect for H* if for each agent $v \in V(H)$ we have $M(v) \in V(H) \setminus \{v\}$. Further, M is *exchange-stable for H* if no two agents from $V(H)$ can form an exchange-blocking pair.

Notice that the smallest hourglass has height two and is a four-cycle. Figure 4 shows a maximal hourglass of height three which contains two different non-maximal hourglasses of height two. Let us mention that it is possible that the vertex sets of two distinct maximal hourglasses have nonempty intersection.

We are ready to show the following fixed-parameter tractability result.

Theorem 4. *An instance of 3-ESR with $2n$ agents and ℓ maximal hourglasses can be solved in $O(6^\ell \cdot n\sqrt{n})$ time.*

The main ideas are as follows. The first observation is that a matching for a maximal hourglass can interact with the rest of the graph in only six different ways: The only agents in an hourglass H of height h that may have neighbors outside H are the layer-0 and layer- $(h-1)$ agents; let us call them *connecting agents* of H . A matching M may match these agents either to agents inside or outside H . Requiring M to be perfect means that an even number of the connecting agents has to be matched inside H . This then gives a bound of at most six different possibilities of the matching M with respect to whether the connecting agents are matched inside or outside H (see Figure 5). Let us call this the *signature* of M with respect to H . Hence, we may try all 6^ℓ possible combinations of signatures for all hourglasses and check whether one of them leads to a solution (i.e., an exchange-stable matching).

The second crucial observation is that each exchange-blocking pair of a perfect matching yields a four-cycle and hence, is contained in some maximal hourglass. Thus, the task of checking whether a combination of signatures leads to a solution decomposes into (a) checking whether each maximal hourglass H allows for an exchange-stable matching adhering to the signature we have chosen for H and (b) checking whether the remaining acceptability graph after deleting all agents that are in hourglasses or matched by the chosen signatures admits a perfect matching.

Task (b) can clearly be done in $O(n \cdot \sqrt{n})$ time by performing any maximum-cardinality matching algorithm (note that the graph $G(\mathcal{P})$ has $O(n)$ edges). We then prove that task (a) for all six signatures can be reduced to checking whether a given hourglass admits a perfect and exchange-stable matching. This, in turn, we show to be linear-time solvable by giving a dynamic program that fills a table, maintaining some limited but crucial facts about the structure of partial matchings for the hourglass.

In the remainder of this section we prove Theorem 4. To this end, let \mathcal{P} denote an input instance of 3-ESR. We observe that the hourglasses are the only parts where an

exchange-blocking pair can form:

Observation 1. *For each exchange-blocking pair (x, y) of a perfect matching M of \mathcal{P} there exists an hourglass in $G(\mathcal{P})$ which contains both x, y and their partners under M .*

Proof. Let M be an arbitrary perfect matching of \mathcal{P} which admits an exchange-blocking pair, say $\{x, y\}$. Then, it must hold that $\{x, M(x)\}, \{x, M(y)\}, \{y, M(y)\}, \{y, M(x)\} \in E(G(\mathcal{P}))$. This implies that $\{x, y, M(x), M(y)\}$ induces an hourglass of height two. \square

4.2.1 Perfect and exchange-stable matchings for a maximal hourglass

We can check in linear time whether a perfect and exchange-stable matching exists for a given hourglass.

Lemma 5. *Let H be a height- h hourglass with agents $V(H) = \{u_i, w_i \mid i \in \{0, 1, \dots, h-1\}\}$. In $O(h)$ time, one can decide whether there exists a perfect and exchange-stable matching for H .*

Proof. Clearly, if $h \leq 4$, we can check for existence of a perfect and exchange-stable matching in constant time. Otherwise, we distinguish between three cases.

Case 1: $\{u_0, w_{h-1}\}, \{u_{h-1}, w_0\} \in E(H)$. Since the maximum degree is three and by the definition of hourglass, this means that H is bipartite. We claim that the matching M with $M := \{u_{i-1}, w_i \mid i \in [h]\}$ (indices taken modulo h) is perfect and exchange-stable for H . Clearly, M is perfect for H . Since $h \geq 5$, any alternating cycle wrt. M has length at least six. By Lemma 2, an exchange-blocking coalition implies that there is a cycle of length at most four and thus we obtain that M is exchange-stable for H . Hence, in this case, we immediately return **yes** by returning M .

Case 2: $\{u_0, u_{h-1}\}, \{w_0, w_{h-1}\} \in E(H)$. By the definition of hourglasses, there are two possible types of perfect matchings M : Either (a) both $\{u_0, u_{h-1}\}$ and $\{w_0, w_{h-1}\}$ are in M or (b) neither is. Consider a matching M of type (a). Observe that no agent u_i , $1 < i < h-1$, envies u_0 or u_{h-1} and analogous for agents w_i , $1 < i < h-1$. If M is not exchange-stable then either (i) (u_0, w_0) or (u_{h-1}, w_{h-1}) form an exchange-blocking pair or (ii) no exchange-blocking pair involves an agent from $\{u_0, u_{h-1}, w_0, w_{h-1}\}$. This is because, since $h \geq 5$, there is no alternating cycle of length at most 4 involving exactly one agent from $\{u_0, u_{h-1}, w_0, w_{h-1}\}$. Hence, there is a perfect exchange-stable matching of type (a) for H if and only neither (u_0, w_0) nor (u_{h-1}, w_{h-1}) are an exchange-blocking pair and the sub-hourglass H' with $H' = H[\{u_i, w_i \mid i \in [h-2]\}]$ has a perfect exchange-stable matching. Notice that in H' the layer-0 agents both have degree two. Thus we may use the algorithm for Case 3 below.

Analogously, there is perfect exchange-stable matching M of type (b) if and only if the hourglass obtained from H by removing both $\{u_0, u_{h-1}\}$ and $\{w_0, w_{h-1}\}$ has a perfect exchange-stable matching. Hence, again we may use the algorithm for Case 3 below.

Case 3: $d_H(u_0) = 2$ or $d_H(w_0) = 2$. This means that all perfect and exchange-stable matchings for H must match u_0 with w_0 or match u_0 and w_0 with the agents from layer 1. In either case, such a matching must match the first layer of unmatched agents within this layer or to agents in a larger layer. This pattern will continue until the last but one layer. We use the following notions to describe the pattern formally. For each $i \in \{0, \dots, h-1\}$, we say that layer i is *horizontally matched* in M if $\{u_i, w_i\} \in M$. For each $i \in [h-1]$, we say that layers $i-1$ and i are *cross-matched* in M if $\{u_{i-1}, w_i\}, \{u_i, w_{i-1}\} \in M$. Then, by induction, we observe that for each $i \in \{0, 1, \dots, h-1\}$, layer- i is either horizontally matched or cross-matched. To check which match yields a correct answer, we can use dynamic programming.

The dynamic program aims to fill two tables D_h, D_c , each of size $O(h)$, which specify for each layer whether there exists a perfect matching that ends with a cross-match or a horizontal match. Formally, $D_h[i] = \mathbf{true}$ if there exists a perfect and exchange-stable matching for $H[\{u_z, w_z \mid 0 \leq z \leq i\}]$ where layer i is horizontally matched, and $D_h[i] = \mathbf{false}$ otherwise. Analogously, $D_c[i] = \mathbf{true}$ if there exists a perfect and exchange-stable matching for $H[\{u_z, w_z \mid 0 \leq z \leq i\}]$ where layers $i-1$ and i are cross-matched, and $D_c[i] = \mathbf{false}$ otherwise. Clearly, by definition, there exists a perfect and exchange-stable matching for the entire hourglass H that matches the agents from H among themselves if and only if $D_h[h-1]$ or $D_c[h-1]$ is **true**. Hence, to show our desired statement it suffices to show that we can compute $D_h[h-1]$ and $D_c[h-1]$ correctly in $O(h)$ time.

Next, we describe how to fill the tables. First, we fill the table for layer $i = 0$, i.e., $D_h[0]$ and $D_c[0]$. The only possibility for a perfect and exchange-stable matching for layer 0 is a horizontal match, i.e., $D_c[0] := \mathbf{false}$ and $D_h[0] := \mathbf{true}$. For $i = 1$, we have two possible perfect matchings such that agents from layer up to 1 are matched among themselves, and thus can determine $D_h[1]$ and $D_c[1]$ in constant time.

Now, to compute the remaining entries of tables D_h and D_c , we claim that for all $i \in \{2, \dots, h-1\}$, the following holds:

$$D_h[i] := D_c[i-1] \vee (D_h[i-1] \wedge \tau_h^{i-1,i}), \quad (4)$$

$$D_c[i] := (D_c[i-2] \vee D_h[i-2]) \wedge \tau_c^{i-1,i}. \quad (5)$$

Herein, $\tau_c^{i-1,i}$ is set **true** if and only if cross-matching layers $i-1$ and i is exchange-stable for $H[\{u_{i-1}, w_{i-1}, u_i, w_i\}]$. Analogously, $\tau_h^{i-1,i}$ is set **true** if and only if horizontally matching layers $i-1$ and i respectively is exchange-stable for $H[\{u_{i-1}, w_{i-1}, u_i, w_i\}]$.

It remains to prove that the equations (4)–(5) are correct.

We first prove equation 4. “ $D_h[i] \Rightarrow D_c[i-1] \vee (D_h[i-1] \wedge \tau_h^{i-1,i})$ ”: Let M be a matching witnessing that $D_h[i] = \mathbf{true}$. If M cross-matches layers $i-2$ and $i-1$, then $D_c[i-1] = \mathbf{true}$. Otherwise, M matches layer $i-1$ horizontally. Thus, $D_h[i-1] \wedge \tau_h^{i-1,i} = \mathbf{true}$.

“ $D_h[i] \Leftarrow D_c[i-1] \vee (D_h[i-1] \wedge \tau_h^{i-1,i})$ ”: If $D_c[i-1] = \mathbf{true}$, then there is a perfect and exchange-stable matching M for layers 0 to $i-1$ that cross-matches layers $i-2$ and $i-1$. Extending M to layer i by matching layer i horizontally yields a perfect and exchange-stable matching for layers up to i because of the following:

- u_i (resp. w_i) does not envy u_{i-1} (resp. w_{i-1}).
- u_{i-2} (resp. w_{i-2}) does not envy u_i (resp. w_i).

If $D_c[i-1] = \mathbf{false}$, $D_h[i-1] \wedge \tau_h^{i-1,i} = \mathbf{true}$. That is, there is a perfect matching M for layers 0 to $i-1$ which matches layer $i-1$ horizontally and, since $\tau_h^{i-1,i} = \mathbf{true}$, extending M to layer i by matching layer i horizontally does not incur an exchange-blocking pair.

Now we show equation (5). “ $D_c[i] \Rightarrow (D_c[i-2] \vee D_h[i-2]) \wedge \tau_c^{i-1,i}$ ”: Consider a matching M for layers 0 to i witnessing $D_c[i] = \mathbf{true}$. Since M is exchange-stable and layers i and $i-1$ are cross-matched by M , we have $\tau_c^{i-1,i} = \mathbf{true}$. Clearly, either layer $i-2$ is horizontally matched or layers $i-3$ and $i-2$ are cross-matched by M and hence, $D_c[i-2] = \mathbf{true}$ or $D_h[i-2] = \mathbf{true}$.

“ $D_c[i] \Leftarrow (D_c[i-2] \vee D_h[i-2]) \wedge \tau_c^{i-1,i}$ ”: Since $D_c[i-2] \vee D_h[i-2] = \mathbf{true}$, there is a perfect and exchange-stable matching M for layers up to $i-2$. Since $\tau_c^{i-1,i}$ is true, cross-matching layers $i-1$ and i is exchange-stable for $H[\{u_{i-1}, w_{i-1}, u_i, w_i\}]$. Observe that adding to M the two pairs $\{u_{i-1}, w_i\}$ and $\{u_i, w_{i-1}\}$ yields an exchange-stable matching for layers 0 to i because neither u_{i-2} nor w_{i-2} envy u_{i-1} or w_{i-1} , respectively, and neither u_i nor w_i envies u_{i-2} or w_{i-2} , respectively.

Finally, it is straight-forward to see that the table entries can be computed in $O(h)$ time by using the just proved equations (4)–(5). \square




Figure 5: Six categories of perfect matchings for an hourglass (see Lemma 6). Notice that the edges which are incident to some vertices without labels may not belong to the hourglass.

4.2.2 Overview and solution structure

We now give an overview of how the FPT algorithm works. Exchange-blocking pairs form only inside hourglasses (Observation 1). Therefore, any perfect matching which is exchange-stable for all hourglasses is exchange-stable. The bound on the preference list length implies that a maximal hourglass H of height h has at most four agents (u_0, w_0, u_{h-1} , and w_{h-1}) that might have a partner outside of H . Each of these four vertices can either be matched to a partner inside or outside the hourglass. Lemma 6 below shows that there are only six different ways a perfect matching can assign partners outside the hourglass to these four vertices. Note that these six possibilities also include the situation that a matching might not assign partners outside the hourglass to these four vertices. We check each of these possibilities to see if they can be part of an exchange-stable matching on the hourglass. Finally, we consider all combinations of possible matchings for different hourglasses. If one of these combination is an exchange-stable matching for all hourglasses and can be extended to form a perfect matching for the rest of the graph, then we return YES. Otherwise, we return NO.

We now observe that there are only six categories for a perfect matching to match the agents of an hourglass. We use the following terminology. Let H be a hourglass in $G(\mathcal{P})$ with $V(H) = \{u_i, w_i \mid 0 \leq i \leq h-1\}$, and let M be a matching for \mathcal{P} . We say that an agent $w_i \in V(H)$ (resp. $u_i \in V(H)$) is *matched straight* if (i) $1 \leq i \leq h-2$ and $M(w_i) \in \{u_{i-1}, u_i, u_{i+1}\}$, or (ii) $i = 0$ and $M(w_i) \in \{u_i, u_{i+1}\}$, or (iii) $i = h-1$ and $M(w_i) \in \{u_{i-1}, u_i\}$ (resp. $i > 0$ and $M(u_i) \in \{w_{i-1}, w_i, w_{i+1}\}$, or $i = 0$ and $M(u_i) \in \{w_i, w_{i+1}\}$, or $i = h-1$ and $M(u_i) \in \{w_{i-1}, w_i\}$). Otherwise we say that $w_i \in V(H)$ (resp. $u_i \in V(H)$) is *matched askew*. Note that if an agent in $V(H)$ is matched askew, then it is either u_0, u_{h-1}, w_0 , or w_{h-1} and the partner is either outside of $V(H)$ or on the other side of the hourglass. Furthermore, if an agent is matched askew, then its partner is uniquely determined.

Lemma 6. *Let H be a height- h hourglass with $V(H) = \{u_i, w_i \mid 0 \leq i \leq h-1\}$. Then, every perfect matching M for \mathcal{P} falls into exactly one of the following six categories (see Figure 5):*

- (I) *Vertices u_0 and w_{h-1} are matched askew and all other vertices of H are matched straight.*
- (II) *Vertices w_0 and u_{h-1} are matched askew and all other vertices of H are matched*

straight.

- (III) Vertices u_0 and w_0 are matched askew and all other vertices of H are matched straight.
- (IV) Vertices u_{h-1} and w_{h-1} are matched askew and all other vertices of H are matched straight.
- (V) Vertices u_0 , w_0 , u_{h-1} , and w_{h-1} are matched askew and all other vertices of H are matched straight.
- (VI) All vertices in H are matched straight.

Below we also say that M belongs to category $C \in \{\text{(I)}, \dots, \text{(VI)}\}$ with respect to H if it satisfies point C above.

Lemma 6. Consider a specific subgraph $H' \subseteq H$ in which the agents from layer-0 and layer- $h-1$ have degree two and the remaining layers has degree three. Formally, $V(H') = V(H)$ and $E(H') = \{\{u_i, w_i\} \mid 0 \leq i \leq h-1\} \cup \{\{u_i, w_{i+1}\}, \{u_{i+1}, w_i\} \mid 0 \leq i \leq h-2\}$. Notice that H' is a bipartite graph on two equal-size subsets. Hence, under M an even number, ν , of agents from $V(H)$ are matched with edges not contained in H' . Note that each such agent is matched askew (and each agent that is matched askew is counted in ν). Furthermore, each such agent is either u_0 , w_0 , u_{h-1} , or w_{h-1} , hence $\nu \in \{0, 2, 4\}$.

If $\nu = 0$, then M corresponds to category (VI).

If $\nu = 4$, then M corresponds to category (V).

If $\nu = 2$, observe that it is not possible that both u_0 and u_{h-1} are matched with edges outside of H' as otherwise it would not be possible to match all remaining agents with edges inside H' . Similarly, it is not possible that both w_0 and w_{h-1} are matched with edges outside of H' . Hence, the agents matched outside of H' are either u_0 and w_{h-1} (category (I)), u_{h-1} and w_0 (category (II)), u_0 and w_0 (category (III)), or u_{h-1} and w_{h-1} (category (IV)). \square

4.2.3 Checking for eligible matchings for hourglasses

Next, we show that, given a height- h hourglass H and a category $C \in \{\text{(I)}, \dots, \text{(VI)}\}$, we can check in $O(h)$ time whether there exists an exchange-stable matching for H which belongs to category i (see Lemma 6 and Figure 5).

Lemma 7. Let H be a maximal hourglass in $G(\mathcal{P})$ of height h with $h \geq 5$ and let $C \in \{\text{(I)}, \dots, \text{(VI)}\}$. In time $O(h)$ it is possible to decide whether there is a matching for \mathcal{P} that matches all agents in H , is exchange-stable for H (i.e. does not contain an exchange-blocking pair in $V(H)$), and belongs to category C .

Proof. We consider each category individually. Consider first the case $C = \text{(I)}$. Let M be a matching that matches all agents in H . Observe that $M(w_0) = u_1$. Hence, $M(w_1) = u_2$, $M(w_2) = u_3$, and so forth. Since M is perfect, by Lemma 2, each exchange-blocking pair induces a length-4 alternating cycle wrt. M . However, since $h \geq 5$, every alternating cycle wrt. M containing an edge $\{x, M(x)\}$ with $x \in V(H)$ has length at least six. So no two agents from $V(H)$ form an exchange-blocking pair, that is, M is exchange-stable for H . Thus, there is always a matching that matches all agents in H and is exchange-stable for H .

The case $C = \text{(II)}$ is analogous to $C = \text{(I)}$ and omitted.

The case $C = \text{(VI)}$ follows from Lemma 5.

For $C = \text{(III)}, \text{(IV)}, \text{(V)}$, we show that checking for an exchange-stable matching in category C can be reduced to checking whether there exists a perfect and exchange-stable matching (i.e. in category (VI)) for the smaller hourglasses H' obtained by ignoring the agents from layer-0 (category (III)), layer- $(h-1)$ (category (IV)), or both (category (V)). The statement then follows from Lemma 5.

Consider $C = \text{(III)}$. Define the induced subgraph $H' := H[\{u_i, w_i \mid i \in [h-1]\}]$. We show that there is an exchange-stable matching that matches all agents of H and belongs to category (III) if and only if there is a perfect exchange-stable matching for H' .

For the forward direction, assume that M is an exchange-stable matching for H and falls in category (III) . Let M' be the matching obtained from M by removing the pairs matching the agents from $\{v_0, w_0\}$. By the definition of category (III) , M' is perfect for H' . Since $M' \subseteq M$ and M is exchange-stable for H , matching M' is also exchange-stable for H' , as required.

For the backward direction, assume that M' is a perfect and exchange-stable matching for H' . Without loss of generality, assume that both u_0 and w_0 have degree three in $G(\mathcal{P})$ as otherwise no exchange-stable matching in category (III) exists. Let x and y denote the neighbors of u_0 and w_0 , respectively, outside of H . Notice that such agents must exist and $x \neq y$ as otherwise no matching in category (III) exists. Now, we claim that matching M derived from M' by defining $M := M' \cup \{\{u_0, x\}, \{w_0, y\}\}$ is exchange-stable for H . Suppose, for the sake of contradiction, that M is not exchange-stable for H , and let (a, b) denote an exchange-blocking pair with $a, b \in V(H)$. Clearly, at least one of $\{a, b\}$, say a , is from layer-0 of H as otherwise (a, b) would also be exchange-blocking H' . By symmetry, suppose that $a = u_0$ without loss of generality. By Observation 1, there must be an hourglass which contains (u_0, b) , a contradiction to H being maximal.

The case $C = \text{(IV)}$ is analogous to $C = \text{(III)}$ and omitted.

Consider $C = \text{(V)}$. If all edges incident with the agents u_0, u_{h-1}, w_0 , and w_{h-1} are contained in $V(H)$, then H is a connected component in $G(\mathcal{P})$ and thus, checking for an exchange-stable matching that matches all agents of H is equivalent to checking for a perfect exchange-stable matching for H and can be done in the required time using Lemma 5. If not all edges incident with the agents u_0, u_{h-1}, w_0 , and w_{h-1} are contained in $V(H)$, then observe that if two agents u_0, u_{h-1}, w_0 , and w_{h-1} share a neighbor (in $G(\mathcal{P})$) outside of H , then there is no matching that matches all agents of H and belongs to category (V) . We thus from now on assume that no two agents u_0, u_{h-1}, w_0 , and w_{h-1} share a neighbor outside of H . Define the induced subgraph $H' := H[\{u_i, w_i \mid i \in [h-2]\}]$. We claim that there is exchange-stable matching that matches all agents of H and belongs to category (III) if and only if there is a perfect exchange-stable matching for H' . If this is true, then the proof is finished in this case by using the algorithm for Lemma 5.

The forward direction is analogous to the one for $C = \text{(III)}$ and omitted.

For the backward direction, assume that M' is a perfect and exchange-stable matching for H' . Let M be the matching that is obtained from M' by matching each of the agents u_0, u_{h-1}, w_0 , and w_{h-1} such that they are matched askew (note that there is a unique way to do so). Clearly, M matches all agents of H and belongs to category (V) . We claim that M is exchange-stable for H . If none of the agents u_0, u_{h-1}, w_0 , and w_{h-1} are matched inside H then M is exchange-stable for H by the same argument as for $C = \text{(III)}$. Hence, assume that two of these agents are matched inside H . To get a contradiction, assume that (a, b) is an exchange-blocking pair in H . Since M' is exchange-stable for H' , at least one agent, say a , is in layer 0 or layer $h-1$ of H . Note that a is not matched with a partner x outside of H , since no agent in H finds x acceptable (recall that no two agents u_0, u_{h-1}, w_0 , and w_{h-1} share a neighbor outside of H). Hence, $M(a) \in V(H)$ and one of $a, M(a)$ is in layer 0 and the other in layer $h-1$. However, since $h \geq 5$, no neighbor of a neighbor a finds $M(a)$ acceptable except for a itself. This is a contradiction to (a, b) being an exchange-blocking pair. Thus, M is exchange-stable for H , as required.

Consider $C = \text{(VI)}$. To check for an hourglass in this category, we may employ the algorithm from Lemma 5 for the hourglass obtained from H by removing all edges with one endpoint in $\{u_0, w_0\}$ and one endpoint in $\{u_{h-1}, w_{h-1}\}$. \square

4.2.4 Overlapping hourglasses

The above results allow us to deal with hourglasses that do not overlap with any other hourglass. However, it is not always the case that hourglasses do not overlap. We now show that if two hourglasses overlap, then their size is bounded. This then allows us to specify the interaction of sets of overlapping hourglasses with the rest of the graph in a similar way as Lemma 6.

We first observe the interaction between two maximal hourglasses.

Lemma 8. *Let H_1 and H_2 be two different maximal hourglasses of profile \mathcal{P} such that H_1 has height h_1 and H_2 has height h_2 . Then, the following holds.*

- (i) *If $h_1 \geq 4$, then for each two agents u and w in the same layer of H_1 , it holds that $u \in V(H_2)$ if and only if $w \in V(H_2)$.*
- (ii) *If $V(H_1) \cap V(H_2) \neq \emptyset$, then $h_1 = h_2 = 2$ or $h_1 = h_2 = 3$.*
- (iii) *If $V(H_1) \cap V(H_2) \neq \emptyset$ and $h_1 = 2$, then $h_2 = 2$ and $|V(H_1) \cap V(H_2)| = 3$.*
- (iv) *If $V(H_1) \cap V(H_2) \neq \emptyset$ and $h_1 = 3$, then $h_2 = 3$ and $|V(H_1) \cap V(H_2)| = 5$.*

Proof. Let \mathcal{P} be a preference profile and H_1 and H_2 two maximal hourglasses in $G(\mathcal{P})$ with heights h_1 and h_2 , respectively. Let $V(H_1) = \{u_i, w_i \mid 0 \leq i \leq h_1 - 1\}$.

For Statement (i), assume that $h_1 \geq 4$. Let u_i and w_i be two agents which are in layer i in H_1 , $i \in \{0, 1, \dots, h_1 - 1\}$. By symmetry, it suffices to show that if $u_i \notin V(H_2)$ then $w_i \notin V(H_2)$. Suppose, for the sake of contradiction, that $u_i \notin V(H_2)$ but $w_i \in V(H_2)$. By the definition of hourglasses, it follows that w_i is in either the first or the last layer of H_2 (since w has degree two in H_2). Without loss of generality, assume that w_i is in layer 0 in H_2 . We distinguish between two cases.

Case 1: $i = 0$ or $i = h_1 - 1$. We only consider the case when $i = 0$ since the other case is symmetric (by reversing the indices). Since $i = 0$, meaning that $w_i = w_0$ is in layer-0 in H_2 , we have that w_0 has three neighbors u_0 , u_1 , and x for some agent $x \in V(G(\mathcal{P})) \setminus \{u_j, w_j \mid 0 \leq j \leq 2\}$ (note that $h \geq 4$) such that $u_1, x \in V(H_2)$ (recall that $u_0 \notin V(H_2)$). Moreover, since w_0 is in layer-0 in H_2 , it follows that w_0 is with either u_1 or x in the same layer in H_2 .

If w_0 and u_1 are in the same layer in H_2 , then by the maximality of H_2 , it follows that $w_1 \in V(H_2)$ (as otherwise we could enlarge H_2 by adding to it u_0 and w_1). Notice that w_1 and u_0 are adjacent in $G(\mathcal{P})$ but $u_0 \notin V(H_2)$. Since $h_1 \geq 4$, by the definition of hourglasses, this means that the third neighbor of w_1 , namely u_2 , is also in H_2 such that w_1 and u_2 are in the last layer in H_2 (recall that w_0 and u_1 are in layer-0 in H_2 , and w_1 and u_1 are adjacent). This implies that H_2 has two layers such that $\{w_0, u_2\} \in E(H_2)$ (note that $\{u_1, u_2\} \notin E(G(\mathcal{P}))$ since u_1 is adjacent to w_0, w_1, w_2 already). However, this also means that $x = u_2$ such that $V(H_2) = \{w_0, u_1, w_1, u_2\} \subsetneq V(H_1)$, a contradiction to H_2 being maximal.

If w_0 and x are in the same layer in H_2 , then u_1 is with either w_1 or w_2 in the same layer in H_2 . If u_1 and w_1 are in the same layer in H_2 , then since $u_0 \notin V(H_2)$, this further implies that w_1 and hence u_1 are in the last layer in H_2 . Hence, $h_2 = 2$ and $\{w_1, x\} \in E(H_2)$. This implies that $x = u_2$ since w_1 is adjacent to u_0, u_1 , and u_2 . In other words, $\{w_0, u_2, u_1, w_1\} \subsetneq V(H_1)$, a contradiction to H_2 being maximal.

If u_1 and w_2 are in the same layer in H_2 , then u_1 and w_2 are not in the last layer in H_2 as otherwise we could add $\{w_1, u_2\}$ to H_2 to extend the height by one, a contradiction. Since u_1 and w_2 are not in the last layer in H_2 , this means that u_1 has degree three in H_2 . Moreover, u_1 and w_2 are both in layer-1 in H_2 ; recall that $\{u_1, w_0\} \in E(H_2)$ and w_0 is in layer-0 in H_2 . This implies that $\{w_0, x\} \in E(H_2)$ such that $x \in \{u_2, u_3\}$. Since $h_1 \geq 4$ we infer that $x \neq u_2$ and hence $x = u_3$. Again, since u_1 and w_2 are in layer-1 but not the last layer in H_2 , we also infer that u_2 and w_1 are in layer-2 in H_2 . Since $u_0 \notin V(H_2)$, it follows that w_1 is in the last layer in H_2 and hence $h_2 = 3$. This means that $V(H_2) = \{u_3, w_0, u_1, w_2, u_2, w_1\} \subsetneq V(H_1)$, a contradiction to H_2 being maximal.

Case 2: $i \in [h_1 - 2]$. This means that w_i is adjacent to u_{i-1}, u_i , and u_{i+1} in H_1 such that u_{i-1} and u_{i+1} are both in $V(H_2)$. Furthermore, w_i is with either u_{i-1} or u_{i+1} in layer-0 or the last layer in H_2 . We only consider the case that that w_i and u_{i-1} are in layer-0 in H_2 as the other three cases are symmetric and analogous.

If w_i and u_{i-1} are in layer-0 in H_2 , then w_{i-1} must also be in $V(H_2)$ as otherwise we could enlarge H_2 by one more layer by adding to it u_i and w_{i-1} , a contradiction. Now, since $w_{i-1} \in V(H_2)$ but $v_i \notin V(H_2)$, it follows that w_{i-1} is in the last layer in H_2 . Similarly as above, we infer that $h_2 = 2$ and hence, $V(H_2) = \{u_{i-1}, w_i, u_{i+1}, w_{i-1}\} \subsetneq V(H_1)$, a contradiction to H_2 being maximal.

Overall, we have shown that u_i and w_i are either both in H_2 or both not in H_2 .

For Statement (ii) assume that $V(H_1) \cap V(H_2) \neq \emptyset$. We first show that $h_1, h_2 \leq 3$. To achieve this, it suffices to show that $h_1 \leq 3$ since H_1 and H_2 are symmetric. Suppose, for the sake of contradiction, that $h_1 \geq 4$. Since H_1 and H_2 are two different hourglasses, by Statement (i), there are two layers i and j such that u_i and w_i are in $V(H_2)$ while u_j and w_j are not. Without loss of generality, assume that $i < j$ (the case that $j < i$ can be shown by reversing the indices). Then, there must also be a layer i such that u_i and w_i are in $V(H_2)$ while u_{i+1} and w_{i+1} are not. This means that u_i and w_i are in layer-0 or layer- $(h-1)$ in H_2 . If u_i and w_i are in different layers in H_2 , then it must hold that $h_2 = 2$ since both u_i and w_i have degree two in H_2 . However, we could enlarge the height of H_2 by one by adding to it the two agents u_{i+1} and w_{i+1} , a contradiction. Similarly, if u_i and w_i are in the same layer in H_2 , then we could as well enlarge the height of H_2 by one by adding to it the two agents u_{i+1} and w_{i+1} , a contradiction.

Now, to show the statement, it remains to show that $h_1 = h_2$. By our above reasoning, suppose, for the sake of contradiction, that $h_1 = 2$ while $h_2 = 3$. Then, by the maximality of H_1 and by the definition of hourglasses, either 2 or 3 agents from $V(H_1)$ are contained in $V(H_2)$. If $|V(H_1) \cap V(H_2)| = 2$, then without loss of generality, assume that $V(H_1) \cap V(H_2) = \{u_0, w_0\}$. Then, this also means that u_0 (resp. w_0) is in the first or the last layer in H_2 . It cannot be the case that u_0 and w_0 are in the same layer in H_2 as otherwise we could enlarge H_2 by adding to it the pair $\{u_1, w_1\}$, a contradiction to H_2 being maximal. If u_0 and w_0 are in different layers, then H_2 has only two layers, a contradiction too.

Now, it remains to consider the case with $|V(H_1) \cap V(H_2)| = 3$. Without loss of generality, assume that $V(H_1) \cap V(H_2) = \{u_0, w_0, u_1\}$. This means that u_0 (resp. u_1) is in the first or the last layer in H_2 since they both have degree two in H_2 . Agents u_0 and u_1 cannot be in the same layer in H_2 as otherwise $\{u_0, u_1\} \in E(H_2)$ such that u_0 or u_1 would have degree four in $G(\mathcal{P})$ (since u_0 or u_1 would need to be adjacent to the agent which is in the same layer as w_0). Hence, without loss of generality, assume that u_0 and u_1 are in layer-0 and layer-2 in H_2 . Moreover, w_0 must be in layer-1 in H_2 as otherwise either u_0 or w_0 would have degree at least four in $G(\mathcal{P})$, a contradiction. This means that there are two other agents $x, y \notin V(H_1)$ such that u_0 and x are layer-0 while w_0 and y are in layer-2 in H_2 , respectively. However, this implies that we could enlarge H_1 by adding to it agents x and y , a contradiction. This completes the proof for Statement (ii).

For Statement (iii), assume that $V(H_1) \cap V(H_2) \neq \emptyset$ and $V(H_1) = \{u_0, w_0, u_1, w_1\}$. By Statement (ii), we know that $h_2 = 2$ such that $|V(H_1) \cap V(H_2)| \in \{2, 3\}$. Using a reasoning similar to the one given for Statement (ii) we can infer that $|V(H_1) \cap V(H_2)| = 3$.

Finally, for Statement (iv), assume that $V(H_1) \cap V(H_2) \neq \emptyset$ and $V(H_1) = \{u_0, w_0, u_1, w_1, u_2, w_2\}$. By Statement (ii), we know that $h_2 = 3$.

Since H_1 and H_2 are two different maximal hourglasses, let u_i , $0 \leq i \leq 2$, be an agent which does not appear in $V(H_2)$. Notice that we can assume that $i \in \{0, 1\}$ as the case with $i = 2$ is symmetrical to the case $i = 0$. We distinguish between four subcases.

If $i = 0$ and $w_0 \in V(H_2)$, then there exists another neighboring agent $x \in V(H_2) \setminus \{u_1, w_0\}$ such that $\{x, w_0\} \in E(H_2)$, $u_1 \in V(H_2)$, and w_0 is in layer-0 or layer-2 in H_2 .




Figure 6: Possible subgraph induced by overlapping maximal hourglasses of height two.

This also means that w_0 is with either x or u_1 in the same layer in H_2 . If w_0 and x are in the same layer in H_2 , then u_1 is in layer-1 in H_2 so that w_1 and w_2 are also in $V(H_2)$. Since $u_0 \notin V(H_2)$ it follows that w_1 is in layer-0 or layer-2 in H_2 . Hence, w_2 and u_1 are in layer-2 in H_2 . Moreover, $u_2 \in V(H_2)$ since w_1 is only adjacent to u_0, u_1 , and u_2 but $u_0 \notin V(H_2)$. Altogether, we infer that $V(H_2) = \{u_1, u_2, w_0, w_1, w_2, x\}$, meaning that $|V(H_2) \cap V(H_1)| = 5$.

If $i = 0$ and $w_0 \notin V(H_2)$, then either both u_1 and w_1 are in $V(H_2)$ or both are not. If both u_1 and w_1 are in $V(H_2)$, then they must be in the same first or last layer as otherwise $h_2 = 2$. But then, we could enlarge H_2 by adding to it $\{u_0, w_0\}$, a contradiction. If both u_1 and w_1 are not in $V(H_2)$, then analogously, both u_2 and w_2 are in $V(H_2)$ or both are not. In either case, we could obtain the contradiction that we could enlarge H_2 .

If $i = 1$ and $w_1 \in V(H_2)$, then it must hold that $u_0, u_2 \in V(H_2)$ such that w_1 is with either u_0 or u_2 in the same first or last layer in H_2 . Without loss of generality, assume that w_1 and u_0 are in the same layer in H_2 (the other case is symmetric). Then, u_2 must be in layer-1 in H_2 , meaning that u_2 has degree three in H_2 . This implies that there exists an agent $x \in V(H_2) \setminus \{w_2\}$ such that $\{x, u_2\} \in E(H_2)$. Furthermore, we also infer that $w_2 \in V(H_2)$. Since $u_1 \notin V(H_2)$, it follows that w_2 in the first layer or the last layer, and hence not with u_2 in layer-1 in H_2 . Thus, there exists an agent $y \in V(H_2) \setminus \{u_2\}$ such that y and w_2 are in the same layer in H_1 . Now, if $\{x, y\} \cap V(H_1) = \emptyset$, then we could enlarge H_1 by adding to it $\{x, y\}$ since $\{x, u_2\}, \{y, w_2\}$ are two edges in $E(G(\mathcal{P}))$, a contradiction. Hence, $|\{x, y\} \cap V(H_1)| \geq 1$. Since $\{u_0, w_1, u_2, w_2\} \subseteq V(H_2)$ and $u_1 \notin V(H_2)$. We obtain that exactly one of $\{x, y\}$ is in $V(H_1)$, i.e., $|V(H_1) \cap V(H_2)| = 5$, as required.

If $i = 1$ and $w_1 \notin V(H_2)$, then either both u_0 and w_0 are in $V(H_2)$ or both are not. If both u_0 and w_0 are in $V(H_2)$, then they must be in the same first or last layer as otherwise $h_2 = 2$. But then, we could enlarge H_2 by adding to it $\{u_1, w_1\}$, a contradiction. If both u_0 and w_0 are not in $V(H_2)$, then analogously, both u_2 and w_2 are in $V(H_2)$ or both are not. In either case, we could obtain the contradiction that we could enlarge H_2 . \square

By Lemma 8 we know that hourglasses of height at least four do not overlap with other hourglasses. We now show that the number of possible matchings for sets of overlapping hourglasses of height two or three is a small constant.

Lemma 9. *Let C be a collection of at least two maximal hourglasses of height two in $G(\mathcal{P})$ such that the union of their vertex sets induces a connected subgraph of $G(\mathcal{P})$ and no maximal hourglass of height two in $G(\mathcal{P})$ can be added to C maintaining this connectedness property. Then, there are at most three matchings that match all vertices in hourglasses in C .*

Proof. We observe that each subgraph induced by the vertex sets of hourglasses in C is isomorphic to a supergraph of the graph G shown in Figure 6. Let C contain two hourglasses H_1 and H_2 . Let H_1 contain the cycle (a, b, c, d) . Two hourglasses of height two overlap in exactly three vertices by Lemma 8(iii). Without loss of generality, assume $V(H_1) \cap V(H_2) = \{a, b, c\}$. Since $G(\mathcal{P})$ has maximum degree 3 and H_2 contains a cycle with four vertices, d




Figure 7: Possible subgraph induced by overlapping maximal hourglasses of height three.

has two neighbors in $V(H_1) \cap V(H_2)$ that have a common neighbor in $V(H_1) \cap V(H_2)$. In other words, d 's neighbors are a and c . Thus, the subgraph induced by $V(H_1) \cup V(H_2)$ is isomorphic to a supergraph of G .

Now consider another hourglass, H_3 such that $V(H_3) \cap V(H_1) \neq \emptyset$. By the above, the subgraph induced by $V(H_1) \cup V(H_2)$ is isomorphic to a supergraph of G . By symmetry, assume without loss of generality that $V(H_1) = \{a, b, c, d\}$ and $V(H_2) = \{a, b, c, e\}$. Since hourglasses overlap in exactly three vertices, Lemma 8(iii) we have $V(H_2) \cap V(H_1) \neq \emptyset$. We claim that $V(H_3) \subset V(H_1) \cup V(H_2)$. For the sake of a contradiction, assume that $f \in V(H_3) \setminus (V(H_1) \cup V(H_2))$. Since f is part of a cycle in H_3 , it has two neighbors among $V(H_1) \cup V(H_2)$. Since a and c have degree three, these two neighbors are among b, e, d . By symmetry assume $b, d \in V(H_3)$ without loss of generality. Notice that $e \notin V(H_3)$ as it would have to be a common neighbor of b and d and hence would have to have degree four. Thus, exactly one of $a, c \in V(H_3)$, a contradiction to the fact that $|V(H_1) \cap V(H_3)| = |V(H_2) \cap V(H_3)| = 3$. Thus, $V(H_3) \subset V(H_1) \cup V(H_2)$ and thus if C contains at least three hourglasses, its induced subgraph is isomorphic to a supergraph of G , as claimed.

Consider a matching that matches all agents in C . Since C induces a subgraph isomorphic to G , matching M matches a and c , and these two have degree three in G , two of b, e, d are matched to a and c , respectively, and the remaining agent is matched to a uniquely determined agent outside of G . There are hence at most three possible matchings that match all agents in the hourglasses in C . \square

Now we turn to hourglasses of height three.

Lemma 10. *Let C be a collection of at least two maximal hourglasses of height three in $G(\mathcal{P})$ such that the union of their vertex sets induces a connected subgraph of $G(\mathcal{P})$ and no maximal hourglass of height three in $G(\mathcal{P})$ can be added to C maintaining this connectedness property. Then, there is a constant number of matchings that match all vertices in hourglasses in C . Moreover, there are at most three different sets S of vertices that are not contained in hourglasses in C and such that there exists a matching M that matches all vertices in hourglasses in C and such that S is exactly the set of vertices that are not contained in hourglasses in C and matched to vertices in hourglasses in C by M .*

Proof. We observe that each subgraph induced by the vertex sets of hourglasses in C is isomorphic to a supergraph of the graph G shown in Figure 7. Let C contain two hourglasses H_1 and H_2 . Let $V(H_1) = \{a, b, c, d, e, f\}$ and let the edges mandated by the definition of hourglasses be as shown in Figure 7. Let thus a, e be in layer 0 and f, c in layer 2 of H_1 . By Lemma 8(iv) we have $|V(H_1) \cap V(H_2)| = 5$, that is, there is exactly one vertex $g \in V(H_2) \setminus V(H_1)$. Since the minimum degree is two in H_2 , vertex g has at least two neighbors in $V(H_1)$. Moreover, these vertices must have degree two in $V(H_1)$, that is, g is neighbors with two of a, e, f, c . Since g is part of a cycle with four vertices in H_2 , it has two neighbors among these four vertices that have a common neighbor. Moreover, that common neighbor must be in $V(H_1)$ because $|V(H_1) \cap V(H_2)| = 5$. That is, g is neighbors with

either a, c or e, f , say e, f . Thus, $V(H_1) \cup V(H_2)$ induces a subgraph that is isomorphic to a supergraph of G . Now observe that because $|V(H_1) \cap V(H_2)| = 5$ and maximum degree three we have either $a, b \in V(H_2)$ or $b, c \in V(H_2)$. By symmetry, assume without loss of generality that $a, b \in V(H_2)$, that is $V(H_2) = \{a, b, e, d, f, g\}$.

Now let H_3 be a maximal hourglass of height three such that $V(H_3) \cap V(H_1) \neq \emptyset$. By Lemma 8(iv) we have $|V(H_3) \cap V(H_1)| = 5$ and thus $V(H_3) \cap V(H_2) \neq \emptyset$, meaning that $|V(H_3) \cap V(H_2)| = 5$. We claim that $V(H_3) \subseteq V(H_2) \cup V(H_1)$. For a contradiction assume that there exists $h \in V(H_3) \setminus (V(H_2) \cup V(H_1))$. By the same argument as for g above and since e, f have degree three already, we get that h is neighbors with a and c and that $a, b, c \in V(H_3)$. Since $|V(H_3) \cap V(H_1)| = 5$ exactly two of e, d, f are in $V(H_3)$. This is a contradiction to $|V(H_3) \cap V(H_2)| = 5$. Thus, indeed $V(H_3) \subseteq V(H_2) \cup V(H_1)$. It follows that each subgraph induced by the vertex sets of hourglasses in C is isomorphic to a supergraph of the graph G shown in Figure 7.

Now let M be a matching that matches all vertices of hourglasses in C . It is clear that, because there is a constant number of vertices in C and by maximum degree three, there is a constant number of matchings matching vertices in C . We claim that exactly one of a, c, g is matched to vertices not in hourglasses in C and all other vertices of C are matched among themselves. Note that this implies the statement of the lemma because each of a, c, g has two neighbors in C and so the partner outside (if any) is uniquely determined. Indeed, since d has degree three, it is matched to either e, f , or b . The remaining two are matched to a, b , or c . Thus the statement holds. \square

4.2.5 Computing hourglasses

Our FPT algorithm starts by computing the collection of maximal hourglasses. We now gather the crucial tool to compute this collection.

Lemma 11. *There is an algorithm \mathcal{A} such that for each maximal hourglass H in $G(\mathcal{P})$ there is an edge $\{u, w\}$ such that on input of u, w algorithm \mathcal{A} computes H if the height of H is at least three, or computes a set of hourglasses containing H if the height of H is two. Algorithm \mathcal{A} runs in $O(1)$ time if there is no maximal hourglass containing u, w and in $O(h)$ time if there is such an hourglass, where h is the hourglass' height.*

Proof. Algorithm \mathcal{A} works as follows. First, it computes all maximal hourglasses of height two containing the input edge $\{u, w\}$. Observe that this can be done in constant time by enumerating all cycles on four vertices containing $\{u, w\}$, then checking whether they are contained in an hourglass of size three that contains $\{u, w\}$, and if not, reporting them as output. (All hourglasses of height three that contain $\{u, w\}$ can be enumerated in constant time since the maximum degree is three.) If no hourglasses are found in this step, it runs Algorithm 2 with input u, w .

Let $V(H) = \{u_i, w_i \mid i \in \{0, 1, \dots, h-1\}\}$. If the height of H is two, then it is clear that \mathcal{A} finds H on input of one of its edges. If the height of H is at least three, then we claim that Algorithm 2 finds H on input of u_1, w_1 . Indeed, in this case a, b in Line 9 is either u_0, w_0 or u_2, w_2 and a, b in Line 14 is the other pair. By a similar argument, MaxHG will compute the remaining vertices of H in the two calls (note that in each call, the next pair u_i, w_i to add is unique).

For the running time it only remains to observe that Algorithm 2 runs in $O(h)$ time. This follows from the fact that checking for Line 9, Line 14, and the if-condition in the calls of MaxHG can be done in constant time since the maximum degree is three. \square

ALGORITHM 2: Algorithm to find a maximal hourglass

Input: A preference profile \mathcal{P} with agent set V and with preferences length at most three, and an edge $\{u, w\}$ in $G(\mathcal{P})$.

Output: A maximal hourglass in $G(\mathcal{P})$ that contains $\{u, w\}$ or **no** if none exists.

▷ **Recursively extend an hourglass $G'[U \cup W]$ to one more layer**

```
1 MaxHG( $u, w, U, W, G'$ ):
2   if  $\exists x \in N_{G'}(u) \setminus \{w\}$  and  $y \in N_{G'}(w) \setminus \{u\}$  s.t.  $\{x, y\} \in E(G')$  then
3      $U \leftarrow U \cup \{y\}$ 
4      $W \leftarrow W \cup \{x\}$ 
5      $G' \leftarrow G' - x - y$ 
6     MaxHG( $x, y, U, W, G'$ )
7 Main( $u, w$ ):
8    $G_1 \leftarrow G$ ;
9    $U_1 \leftarrow \emptyset$ ;  $W_1 \leftarrow \emptyset$ ;
10  ▷ Try to find a four-cycle and build the first half of a maximal hourglass
11  if  $\exists a \in N_{G_1}(u) \setminus \{w\}$  and  $b \in N_{G_1}(w) \setminus \{u\}$  s.t.  $\{a, b\} \in E(G')$  then
12     $U_1 \leftarrow U_1 \cup \{u, b\}$ 
13     $W_1 \leftarrow W_1 \cup \{w, a\}$ 
14     $G_1 \leftarrow G_1 - \{u, a\} - \{w, b\}$ 
15    MaxHG( $a, b, U_1, W_1, G_1$ )
16  ▷ Try to extend to build the second half of the maximal hourglass
17  if  $\exists a \in N_{G_1}(u) \setminus \{w\}$  and  $b \in N_{G_1}(w) \setminus \{u\}$  s.t.  $\{a, b\} \in E(G')$  then
18     $U_1 \leftarrow U_1 \cup \{b\}$ 
19     $W_1 \leftarrow W_1 \cup \{a\}$ 
20    MaxHG( $a, b, U_1, W_1, G_1$ )
21  if  $|U_1| \geq 2$  then return  $(U_1, W_1)$ ;
22  return no
```

4.2.6 FPT algorithm

We now describe an algorithm which solves 3-ESM in $f(6^\ell \cdot n \cdot \sqrt{n})$ time, where ℓ denotes the number of maximal hourglases. We are given an instance \mathcal{P} of 3-ESR. We first find a collection of all maximal hourglases in $G(\mathcal{P})$ as follows. We maintain for each agent v a list $L(v)$ of hourglases of height at most three that contain v , we initialize these as empty lists. Next, for each edge $\{u, w\}$ in $G(\mathcal{P})$ we run Algorithm 2 with input u, w . If the result is an hourglass of height at least four, we remove its agents from the graph and continue with the next edge. If the result is an hourglass of height at most three, we save a pointer to this hourglass in $L(v)$ for each vertex v in the hourglass. After we have done this for each edge, we have a collection of maximal hourglases $H_1, \dots, H_{\ell'}$ by Lemma 11. Since no two hourglases of height at least four overlap by Lemma 8, each hourglass in $G(\mathcal{P})$ occurs among the hourglases we have computed, but some may occur multiple times. To remove duplicates we proceed as follows. First, since we have removed the agents of hourglases of height at least four, no such hourglass occurs twice. We iterate over all agents in V and for each $v \in V$ we check all pairs of hourglases $L(v)$ and if they contain the same vertices, we keep only the first one.

Let us now tally the running time of computing the collection of hourglases. When iterating over the edges of $G(\mathcal{P})$, we can compute a maximal hourglass containing the edge in question in $O(h)$ time where h is the height of the hourglass by Lemma 11 (or in $O(1)$ time determine that there is no hourglass containing the edge). Since the vertices of hourglases of height at least four are removed, the time spent in loop iterations for such hourglases is $O(n)$. Since the other hourglases have constant size, they are discovered a constant number of times, each of which takes constant time, and hence the time spent in

loop iterations for hourglasses of size at most three is $O(n)$ as well. The deduplication step takes $O(n)$ time because the length of $L(v)$ is constant for each vertex v : Each hourglass occurring in $L(v)$ is contained in the set $N_6(v)$ of vertices that have distance at most 6 from v . Since the maximum degree is constant, $|N_6(v)|$ is constant, and the number of edges in $N_6(v)$ is constant. Hence, each hourglass in $L(v)$ is discovered at most a constant number of times. Thus the deduplication step takes constant time for each vertex and $O(n)$ time overall. After the deduplication step, the number of maximal hourglasses we have found is ℓ .

We now compute an auxiliary graph that contains all maximal hourglasses of height two or three and an edge between two hourglasses if they overlap. Note that this auxiliary graph can be computed in $O(n)$ time by using the lists $L(v)$. We group the hourglasses into collections $C_1, \dots, C_{\ell'}$ given by the connected components of the auxiliary graph. Note that, by Lemma 8 each collection C_i contains either only hourglasses of height two or only of height three.

Next, we iterate over all possible sets of the following combinations:

1. Each combination of a collection C_i of hourglasses of height two and one of the three possible matchings that match all vertices in C_i (see Lemma 9) if that matching is exchange-stable for C_i .
2. Each combination of a collection C_i of hourglasses of height three and one of the three possible vertex sets S that are matched to vertices in C_i by a matching M that matches all vertices of C_i (see Lemma 10) if that matching is exchange-stable for C_i . (Note that this can be checked in constant time because there is a constant number of matchings that match all vertices of C_i by Lemma 10.)
3. Each combination of an hourglass H_i of height at least four and the six categories described in Lemma 6 if there is a matching of that category that matches all vertices of H_i and is exchange-stable for H_i . (Whether such a matching exists can be checked in overall $O(n)$ time by Lemma 7.)

For each combination, we check if any agent outside an hourglass is assigned more than one partner (note that the partners are in each combination uniquely determined). If this is the case then we go to the next combination. Otherwise, we delete all maximal hourglasses and all agents already matched by this combination and compute a perfect matching on the remaining instance. If such a perfect matching exists then return the corresponding perfect matching, otherwise try the next combination. Should no combination result in a perfect matching, return **no**.

Let us investigate the running time of this algorithm. By Lemmas 6, 9 and 10 there are in total 6^ℓ possible combinations, so checking the hourglass-category combinations takes overall $O(6^\ell \cdot n)$ time. Using a maximum-cardinality matching algorithm (see e.g., Hopcroft and Karp [15], running time proven by Micali and Vazirani [19]) we check for a perfect matching in $O(n \cdot \sqrt{n})$ time (recall that the acceptability graph has $O(n)$ edges). In total this gives us a running time of $O(6^\ell \cdot (n + n \cdot \sqrt{n})) = O(6^\ell \cdot (n + n \cdot \sqrt{n}))$ which, if correct, shows that the problem is fixed-parameter tractable with respect to the number of maximal hourglasses.

It remains to show that the algorithm is correct. If \mathcal{P} admits a perfect and exchange-stable matching M , then M is exchange-stable for each maximal hourglass H_i , $i \in [\ell]$, such that each agent in $V(H_i)$ is matched and analogously for each collection C_i . By Lemma 8 two hourglasses overlap only if both have height two or both have height three. Thus, matching M decomposes into disjoint matchings that match all agents in (1) each hourglass of height at least four, (2) each collection C_i of hourglasses of height two, (3) each collection C_i of hourglasses of height three, and (4) remaining agents that are not

matched by the matchings corresponding to (1), (2), or (3). By Lemma 6, for each maximal hourglass H_i of height at least four, M corresponds to one of the (up to) six possible categories. And by Lemma 7 we will correctly obtain a matching of the right category in one of the combinations. By Lemma 9 we correctly determine a matching for collections C_i hourglasses of height two in one of the combinations.. By Lemma 10 we correctly determine a matching for collections C_i of hourglasses of height three in one of the combinations.. Finally, the Hopcroft-Karp algorithm will find a perfect matching for the remaining agents. Thus, in one of the combinations we have obtained a perfect matching. By Observation 1 and since we have checked for exchange-stability by direct checks for collections C_i and via Lemma 7 for hourglasses of height at least four, the found perfect matching is exchange-stable for \mathcal{P} .

Hence, we have proved Theorem 4.

5 Paths to Exchange-Stability

We now study the parameterized complexity of P-ESM with respect to the number of swaps. Observe that it is straightforward to decide an instance of P-ESM with $2n$ agents in $O((2n)^{2k+2})$ time by trying k times all of the $O(n^2)$ possibilities for the next swap and then checking whether the resulting matching is exchange-stable. The next theorem shows that the dependency of the exponent in the running time cannot be removed unless the unlikely collapse $\text{FPT} = \text{W}[1]$ occurs.

Theorem 5. *PATH TO EXCHANGE-STABLE MARRIAGE is $\text{W}[1]$ -hard with respect to the number k of swaps.*

Proof. We provide a parameterized reduction from the $\text{W}[1]$ -complete INDEPENDENT SET problem, parameterized by the size of the independent set [12]. Therein, we are given a graph H and an integer h and want to decide whether there is an h -vertex *independent* set, i.e., a subset of h pairwise nonadjacent vertices.

Let $I = (H, h)$ be an instance of INDEPENDENT SET with vertex set $V(H) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(H)$. We construct an instance $I' = (\mathcal{P}, M_0, 2h)$ of P-ESM where \mathcal{P} has two disjoint agent sets U and W , each of size $2n + h$.

Both U and W consist of h *selector-agents* and $2n$ *vertex-agents* with preferences which encode the adjacency of the vertices in $V(H)$. More precisely, for each $j \in [h]$, we create two selector-agents, called s_j and t_j , and add them to U and W , respectively. For each $i \in [n]$, we create four vertex-agents, called x_i, u_i, y_i, w_i , add x_i and u_i to U , and add y_i and w_i to W . Altogether, we have $U = \{s_j \mid j \in [h]\} \cup \{u_i, x_i \mid i \in [n]\}$ and $W = \{t_j \mid j \in [h]\} \cup \{w_i, y_i \mid i \in [n]\}$.

Now, we define the preferences of the agents from $U \cup W$. For notational convenience, we define two subsets of agents which shall encode the neighborhood of a vertex: For each vertex $v_i \in V(H)$, define $Y(v_i) := \{y_z \mid \{v_i, v_z\} \in E(H)\}$ and $U(v_i) := \{u_z \mid \{v_i, v_z\} \in E(H)\}$.

$$\begin{aligned} \forall j \in [h]: & \quad s_j : w_1 \succ \dots \succ w_n \succ t_j, & t_j : u_1 \succ \dots \succ u_n \succ x_1 \succ \dots \succ x_n \succ s_j, \\ \forall i \in [n]: & \quad x_i : t_1 \succ \dots \succ t_h \succ y_i, & y_i : u_i \succ x_i \succ [U(v_i)], \\ \forall i \in [n]: & \quad u_i : w_i \succ [Y(v_i)] \succ y_i \succ t_1 \succ \dots \succ t_h, & w_i : s_1 \succ \dots \succ s_h \succ u_i. \end{aligned}$$

Figure 8 illustrates the underlying acceptability graph for a graph consisting of a single edge. Herein, $[Y(v_i)]$ (resp. $[U(v_i)]$) denotes the unique preference list where the agents in $Y(v_i)$ (resp. $U(v_i)$) are ordered ascendingly according to their indices.

Observe that the acceptability graph $G(\mathcal{P})$ includes the following edges:

- For all $i \in [h]$ and $j \in [n]$, the edges $\{s_i, t_i\}$, $\{s_i, w_j\}$, $\{t_i, x_j\}$, $\{t_i, u_j\}$, $\{w_j, u_j\}$, $\{y_j, x_j\}$, $\{y_j, u_j\}$ are in $E(G(\mathcal{P}))$.




Figure 8: The left part shows graph $H = (\{v_1, v_2\}, \{\{v_1, v_2\}\})$. Then $(H, 1)$ is an instance of INDEPENDENT SET. We reduce this instance to the P-ESM instance of which the acceptability graph is shown on the right.

– For all edges $\{v_i, v_{i'}\} \in E(G)$, the edges $\{u_i, y_{i'}\}$ and $\{u_{i'}, y_i\}$ are in $E(G(\mathcal{P}))$.

We define an initial matching M_0 on $G(\mathcal{P})$ as $M_0 = \{\{s_j, t_j\} \mid j \in [h]\} \cup \{\{w_i, u_i\}, \{y_i, x_i\} \mid i \in [n]\}$. This completes the construction of I' , which can clearly be done in polynomial time. It is straight-forward to check that that \mathcal{P} is bipartite and the construction can be done in linear time.

Next, we show the correctness of the reduction, i.e., I is a positive instance of INDEPENDENT SET if and only if I' is a positive instance of P-ESM. For notational convenience, define $S = \{s_1, \dots, s_h\}$, $T = \{t_1, \dots, t_h\}$, $X = \{x_1, \dots, x_n\}$, $Y = \{y_1, \dots, y_n\}$, and $W = \{w_1, \dots, w_n\}$:

For the forward direction, assume that (H, h) is a yes-instance of INDEPENDENT SET. We need to show that we can reach an exchange-stable matching for the constructed instance of P-ESM in $2h$ swaps from M_0 . Let $V' = \{v_{j_1}, \dots, v_{j_h}\}$ denote an h -vertex independent set. For each $z \in [h]$, one can verify that t_z and w_{j_z} envy each other, implying that (t_z, w_{j_z}) is exchange-blocking pair of M_0 . Therefore, we swap their partners and eventually obtain M_h with $M_h(t_z) = u_{j_z}$ and $M_h(w_{j_z}) = s_z$ for all $z \in [h]$. The partners of the remaining agents remain unchanged, i.e., they are the same as in M_0 . Now, for each $z \in [h]$, (x_{j_z}, u_{j_z}) is an exchange-blocking pair of M_h , since x_{j_z} prefers t_z to his partner y_{j_z} whereas u_{j_z} prefers y_{j_z} to his partner t_z . We swap their partners and obtain the matching M_{2h} where $M_{2h}(x_{j_z}) = t_z$ and $M_{2h}(u_{j_z}) = y_{j_z}$ for all $z \in [h]$. The partners of the remaining agents remain unchanged, i.e., are the same as in M_h . It remains to be shown that M_{2h} is exchange-stable, i.e., no agent is involved in an exchange-blocking pair of M_{2h} :

- Each selector-agent $s_z \in S$ is matched to w_{j_z} . Since all selector agents from S have the same preferences over $\{w_i \mid i \in [n]\}$, no two agents from S can form an exchange-blocking pair. Neither can an agent from S and an agent from $\{u_i \mid i \in [n]\}$ jointly form an exchange-blocking pair since u_i is either matched with its most-preferred agent or matched with y_i who is not acceptable to s_j . Finally, no agent from S can form with an agent from X an exchange-blocking pair since every agent from S either prefers its own partner or does not find the partner of the agents from X acceptable.
- Each vertex-agent $x_i \in X$ is either matched to some selector-agent t_j or to y_i . If x_i is matched to some t_j . Then x_i prefers all selector-agents t_z with $z < j$. Notice that $M_{2h}(t_z) = x_{i'}$ for some vertex-agent $x_{i'} \in X \setminus \{x_i\}$ other than x_i . Since all vertex-agents from X have the same preferences over all selector-agents from T , $x_{i'}$ does not envy x_i , and will not form with x_i an exchange-blocking pair of M_{2h} .

If x_i is matched to y_i , then x_i envies every other agent $x_{i'} \in X \setminus \{x_i\}$ which is matched to some selector-agent from T . However, no agent from $X \setminus \{x_i\}$ finds y_i acceptable and

hence, will not form with x_i an exchange-blocking pair of M_{2h} .

- For each $i \in [n]$, vertex-agent u_i is either matched to w_i or to y_i . In the former case, u_i does not prefer anyone over its current partner. In the latter case, we know that w_i is matched to some selector-agent s_z . Agent u_i envies s_z , however s_z is not involved in any exchange-blocking pair (see above).

Agent u_i also prefers any $y_j \in Y(v_i)$ to his partner y_i . Suppose, for the sake of contradiction, that u_i and $M_{2h}(y_j)$ form an exchange-blocking pair of M_{2h} with $y_j \in Y(v_i)$. By our construction, either $M(y_j) = x_j$ or $M(y_j) = u_j$ holds. In the former case, x_j does not find y_i acceptable and will not form with u_i an exchange-blocking pair, a contradiction. In the latter case, we infer that $v_j \in V'$, implying that v_j and v_i are non-adjacent, a contradiction too.

- Each vertex-agent $y_i \in Y$ is either matched to u_i or to x_i . In the former case, u_i is most acceptable to agent y_i . Therefore, y_i does not envy any other agent. In the latter case, y_i prefers only u_i to x_i . We know that u_i must be matched to w_i . However, w_i does not find x_i acceptable. Therefore, y_i is not involved in any exchange-blocking pair.
- Each selector-agent $t_z \in T$ is matched to some vertex-agent x_i . Since all selector agents from T have the same preferences over X , no two selector-agents will form an exchange-blocking pair of M_{2h} . Neither will any agent $y_{i'}$ from Y form an exchange-blocking pair with t_z by the previous reasoning. Neither will any agent w_i form an exchange-blocking pair with t_z since he does not find x_i acceptable.
- For each $i \in [n]$, vertex-agent w_i is either matched to some selector-agent $s_z \in S$ or to u_i . In the former case, w_i envies $M_{2h}(s_j)$ for all $j < z$. However, for each $j \in [h] \setminus \{z\}$, we know that the partner of s_j under M_{2h} is a vertex-agent $w_{i'}$ for some $i' \in [n] \setminus \{i\}$ who also prefers s_j to s_z , and will not form with w_i an exchange-blocking pair.
In the latter case, agent w_i prefers all agents from S to his partner u_i . However, no partner of the agents from S finds u_i acceptable, and will not form with w_i an exchange-blocking pair.

For the backwards direction, assume that $(\mathcal{P}, M_0, 2h)$ is a positive instance, meaning that there exists an exchange-stable matching M_ℓ that can be reached in at most $2h$ swaps from M_0 (and $\ell \leq 2h$). We show that the set $V' = \{v_i \mid M(w_i) \in S\}$ is an independent set of size h , implying that (H, h) is a positive instance of INDEPENDENT SET.

First of all, we observe that M_ℓ is a perfect matching since it is reachable from M_0 which is perfect.

Claim 5. *Each selector-agent $s_j \in S$ has $M_\ell(s_j) \in W$.*

Proof. Towards a contradiction, suppose that there is a selector-agent s_j with $M_\ell(s_j) = t_j$. Without loss of generality, assume that $h < n$ (as otherwise the instance is solvable in polynomial-time). This means there remains a vertex-agent w_i with $M(w_i) = u_i$. It is straightforward to see that w_i and t_j form an exchange-blocking pair, a contradiction. (of Claim 5) \diamond

Claim 6. *No vertex-agent $y_i \in Y$ is matched to some agent from $U(v_i)$ in M_ℓ .*

Proof. Suppose, for the sake of contradiction, that $M_\ell(y_i) \in U(v_i)$. Then, by the preferences of x_i we infer that $M_\ell(x_i) = t_j$ with $t_j \in T$ (since M_ℓ is perfect). However, we also infer that t_i and y_i envy each other, a contradiction. (of Claim 6) \diamond

Claim 7. *No vertex-agent u_i has $M_\ell(u_i) \in T$.*

Proof. Suppose, for the sake of contradiction, that $M_\ell(u_i) \in T$. By Claim 6, we know that $M_\ell(y_i) \in U(v_i)$. Hence, $M_\ell(y_i) = x_i$. Now, x_i and u_i envy each other, a contradiction. (of Claim 7) \diamond

Claim 8. For each $i \in [n]$, if $M_\ell(w_i) \in S$, then $M_\ell(y_i) = u_i$.

Proof. If $M_\ell(w_i) \in S$, then u_i must be matched to y_i since it cannot be matched to any agent from T (Claim 7) nor to any agent from $Y(u_i)$ (Claim 6). (of Claim 8) \diamond

Claim 9. V' contains exactly h vertices.

Proof. We know that every selector-agent $s_j \in S$ is matched to some vertex-agent w_i (Claim 5). A vertex v_i is selected into V' if w_i is matched to some s_j . Since $|S| = h$, we will select exactly h vertices into V' . (of Claim 9) \diamond

Claim 10. No two vertices in V' are adjacent in G .

Proof. Suppose we have selected two adjacent vertices $v_i, v_{i'}$ with $\{v_i, v_{i'}\} \in$. This means that $M_\ell(w_i), M_\ell(w_{i'}) \in S$. By Claim 8, we infer that $M_\ell(u_i) = y_i$ and $M_\ell(u_{i'}) = y_{i'}$. Since v_i and $v_{i'}$ are adjacent in G , it follows $y_i \in Y(u_{i'})$ and $y_{i'} \in Y(u_i)$, meaning that u_i and $u_{i'}$ envy each other, a contradiction. (of Claim 10) \diamond

Claims 9 and 10 immediately imply that V' is an h -vertex independent set. Since the reduction is correct, it is a parameterized reduction. (of Claim 6) \diamond

6 Conclusion

We conclude by mentioning that all obtained results transfer to the case when the preferences may contain ties (i.e., two agents are considered equally good).

Regarding preference restrictions [8], it would be interesting to know whether deciding (coalitional) exchange-stability for complete preferences would become tractable for restricted preferences domains, such as single-peakedness or single-crossingness.

Moreover, as it is unclear whether the number of exchange steps is polynomially bounded, the NP-containment of the problem of checking whether a given matching may reach an exchange-stable matching is open.

Acknowledgments. JC was supported by the WWTF, research project (VRG18-012). MS was supported by the Alexander von Humboldt Foundation.

References

- [1] A. Abizada. Exchange-stability in roommate problems. *Review of Economic Design*, 23:3–12, 2019. 2
- [2] D. J. Abraham and D. F. Manlove. Pareto optimality in the roommates problem. Technical report, University of Glasgow, Department of Computing Science, 2004. TR-2004-182. 2
- [3] J. Alcalde. Exchange-proofness or divorce-proofness? Stability in one-sided matching markets. *Economic Design*, 1:275–287, 1995. 1, 2

- [4] H. Aziz and A. Goldwaser. Coalitional exchange stable matchings in marriage and roommate market. In *Proceedings of the 16th International Conference on Autonomous Agents and Multiagent Systems (AAMAS '17)*, pages 1475–1477, 2017. Extended Abstract. **2**
- [5] P. Biró and G. Norman. Analysis of stochastic matching markets. *International Journal of Game Theory*, 42(4):1021–1040, 2013. **2**
- [6] H. L. Bodlaender, T. Hanaka, L. Jaffke, H. Ono, Y. Otachi, and T. C. van der Zanden. Hedonic seat arrangement problems. In A. E. F. Seghrouchni, G. Sukthankar, B. An, and N. Yorke-Smith, editors, *Proceedings of the 19th International Conference on Autonomous Agents and Multiagent Systems (AAMAS '20)*, pages 1777–1779, 2020. Extended Abstract. **2**
- [7] H. L. Bodlaender, T. Hanaka, L. Jaffke, H. Ono, Y. Otachi, and T. C. van der Zanden. Hedonic seat arrangement problems. Technical report, arXiv:2002.10898 (cs.GT), 2020. **3**
- [8] R. Bredereck, J. Chen, U. P. Finnendahl, and R. Niedermeier. Stable roommate with narcissistic, single-peaked, and single-crossing preferences. *Autonomous Agents and Multi-Agent Systems*, 34(53):1–29, 2020. **33**
- [9] K. Cechlárová. On the complexity of exchange-stable roommates. *Discrete Applied Mathematics*, 116(3):279–287, 2002. **1, 5**
- [10] K. Cechlárová and D. F. Manlove. The exchange-stable marriage problem. *Discrete Applied Mathematics*, 152(1-3):109–122, 2005. **1, 2, 4, 5, 6, 8**
- [11] J. Chen. Reaching stable marriage via divorces is hard. Technical report, arXiv:1906.12274(cs.GT), 2020. **2**
- [12] M. Cygan, F. V. Fomin, L. Kowalik, D. Lokshtanov, D. Marx, M. Pilipczuk, M. Pilipczuk, and S. Saurabh. *Parameterized Algorithms*. Springer, 2015. **30**
- [13] D. Gale and L. S. Shapley. College admissions and the stability of marriage. *The American Mathematical Monthly*, 120(5):386–391, 1962. **1**
- [14] M. R. Garey and D. S. Johnson. *Computers and Intractability—A Guide to the Theory of NP-Completeness*. W. H. Freeman and Company, 1979. **6**
- [15] J. E. Hopcroft and R. M. Karp. An $n^{5/2}$ algorithm for maximum matchings in bipartite graphs. *SIAM Journal on Computing*, 2(4):225–231, 1973. **29**
- [16] R. W. Irving. Stable matching problems with exchange restrictions. *Journal of Combinatorial Optimization*, 16(4):344–360, 2008. **3**
- [17] D. Knuth. *Mariages Stables*. Les Presses de L’Université de Montréal, 1976. **2**
- [18] E. McDermid, C. T. Cheng, and I. Suzuki. Hardness results on the man-exchange stable marriage problem with short preference lists. *Information Processing Letters*, 101(1):13–19, 2007. **3**
- [19] S. Micali and V. V. Vazirani. An $o(\sqrt{|v|}|e|)$ algorithm for finding maximum matching in general graphs. In *Proceedings of the 21st Annual IEEE Symposium on Foundations of Computer Science (FOCS '80)*, pages 17–27, 1980. **29**