GEOMETRIC DEDUCTION OF THE SOLUTIONS TO MODULAR EQUATIONS

MD. SHAFIUL ALAM AND TOSHIYUKI SUGAWA

ABSTRACT. In his notebooks, Ramanujan presented without proof many remarkable formulae for the solutions to generalized modular equations. Much later, proofs of the formulae were provided by making use of highly nontrivial identities for theta series and hypergeometric functions. We offer a geometric approach to the proof of those formulae. We emphasize that our proofs are geometric and independent of such identities.

1. INTRODUCTION

For given integers $p \ge 2$, Ramanujan, an Indian mathematical genius, considered the equation

(1.1)
$$\frac{{}_{2}F_{1}(a,1-a;1;1-\beta)}{{}_{2}F_{1}(a,1-a;1;\beta)} = p \frac{{}_{2}F_{1}(a,1-a;1;1-\alpha)}{{}_{2}F_{1}(a,1-a;1;\alpha)},$$

known as the generalized modular equation of degree p and signature 1/a. He left many formulae describing relations between α and β in his unpublished notebooks but he did not record any proof of those formulae (see [9] and [23]). Here, $_2F_1(a,b;c;z)$ denotes the hypergeometric function whose definition will be given in Section 2. The case when 1/a = 2 corresponds to the classical modular equation. Indeed, the complete elliptic integral of the first kind is described by

$$\mathbf{K}(r) = \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-r^2x^2)}} = \frac{\pi}{2} {}_2F_1(\frac{1}{2}, \frac{1}{2}; 1; r^2)$$

and the function

$$\mu(r) = \frac{\pi}{2} \cdot \frac{\mathbf{K}(\sqrt{1-r^2})}{\mathbf{K}(r)} = \frac{\pi}{2} \cdot \frac{{}_2F_1(\frac{1}{2}, \frac{1}{2}; 1; 1-r^2)}{{}_2F_1(\frac{1}{2}, \frac{1}{2}; 1; r^2)}$$

is known to be the modulus of the Grötzsch ring $\{z \in \mathbb{C} : |z| < 1\} \setminus [0, r]$ for 0 < r < 1 and plays an important role in the theory of plane quasiconformal mappings (see, for instance,

File: main.tex, printed: 2021-5-13, 0.46

²⁰²⁰ Mathematics Subject Classification. Primary 30F35; Secondary 33C05, 33E05, 30F10.

Key words and phrases. modular equation, hypergeometric function, Fuchsian group, generalized complete elliptic integral.

The first author was supported by Data Sciences Program II of Graduate School of Information Sciences, Tohoku University. The authors were supported in part by JSPS KAKENHI Grant Number JP17H02847.

[21] or [5]). The modular equation (1.1) for 1/a = 2 now takes the form $\mu(s) = p\mu(r)$ with $\alpha = r^2$ and $\beta = s^2$. When p = 2, the solution to this modular equation is given by

(1.2)
$$s = \frac{1 - \sqrt{1 - r^2}}{1 + \sqrt{1 - r^2}} = \left(\frac{1 - \sqrt{1 - r^2}}{r}\right)^2$$

(see [21, (2.4) in p. 60] or [5, (5.4)]).

Ramanujan mainly considered the case when the signature 1/a is 3, 4 or 6. See [7], [8], [9], [10], [12] and [22] for his work in relation with the modular equations. Berndt, Bhargava and Garvan [11] gave proofs for those formulae of Ramanujan (see also [3]). Their proofs make use of highly nontrivial identities for Jacobi's theta functions and hypergeometric functions in addition to a number of ingeneous ideas. For example, they gave rigorous proofs for the following results.

Theorem 1.3 ([11, Theorem 7.1]). When p = 2 and q = 3, the solutions α and β to the equation (1.1) are related by

(1.4)
$$(\alpha\beta)^{1/3} + \left\{ (1-\alpha)(1-\beta) \right\}^{1/3} = 1.$$

Theorem 1.5 ([11, Lemma 7.4]). When p = 3 and q = 3, the solutions α and β to the equation (1.1) are related by

(1.6)
$$(1-\alpha)^{1/3} = \frac{1-\beta^{1/3}}{1+2\beta^{1/3}}.$$

We note that the above relations can be transformed to polynomial equations. For instance, (1.4) may be transformed to

(1.7)
$$(2\alpha - 1)^3 \beta^3 - 3\alpha (4\alpha^2 - 13\alpha + 10)\beta^2 + 3\alpha (2\alpha^2 - 10\alpha + 9)\beta - \alpha^3 = 0.$$

In particular, we observe that there are at most three values of β satisfying the modular equation (1.4) for each α . We can say that α and β satisfy a polynomial equation of degree 3 in this case. It is rather surprising that α and β are related algebraically, because the hypergeometric function is transcendental for the corresponding parameters. For instance, we do not have a complete answer to the question for which p and a the solutions to the modular equation (1.1) is algebraic. In this note, we propose a geometric approach to this problem. In particular, the geometric observation suggests that it is more natural to look at q = 1/(1-2a) rather than the signature 1/a. We will call q the order of the modular equation (1.1). For instance, the signatures 1/a = 2, 3, 4, 6 correspond to $q = \infty, 3, 2, 3/2$, respectively. Though our approach does not cover the case 1/a = 6, it may allow us to approach other cases when q are integers > 3. Let $G_q = \langle T, W \rangle$ be the Fuchsian group generated by $T\tau = \tau + \lambda$ and $W\tau = \tau/(1+\lambda\tau)$, where $\lambda = 2\cos\frac{\pi}{2a}$ and let M_p be the Möbius transformation defined by $M_p \tau = p \tau$. To speak about the solutions to (1.1), we have to clarify the range of the solutions. Originally, the range of α and β was the interval [0, 1]. However, in our setting, it is natural to choose the quotient Riemann surface $X = G_q \setminus \mathbb{H}$ as the range of the solutions. Here, we take $X = \widehat{\mathbb{C}} \setminus \{0, 1\}$ for finite q and $X = \mathbb{C} \setminus \{0, 1\}$ for $q = \infty$. Then we have the following result.

Theorem 1.8. Let n, p and q be integers with $n, p, q \ge 2$ (possibly $q = \infty$) and set a = (q-1)/(2q). Then, the solutions (α, β) to the equation (1.1) in $X = G_q \setminus \mathbb{H}$ satisfy the equation $P(\alpha, \beta) = 0$ for an irreducible polynomial P(x, y) of degree n if and only if $G_q \cap G_q^{M_p}$ is a subgroup of G_q of index n, where $G_q^{M_p} = M_p^{-1}G_qM_p$.

As we will see in the proof, the solutions α and β are parametrized as $\alpha = \varphi(z)$ and $\beta = \psi(z)$ on the Riemann surface $Z = (G_q \cap G_q^{M_p}) \setminus \mathbb{H}$ and they satisfy the polynomial equation $P(\alpha, \beta) = 0$ of order n. More precisely, P(x, y) has the forms

$$P(x,y) = a_0(x)y^n + a_1(x)y^{n-1} + \dots + a_{n-1}(x)y + a_n(x)$$

= $b_0(y)x^n + b_1(y)x^{n-1} + \dots + b_{n-1}(y)x + b_n(y),$

where $a_j(x) \in \mathbb{C}[x], b_j(y) \in \mathbb{C}[y]$ (j = 0, 1, ..., n), such that P(x, y) is irreducible as an element of $\mathbb{C}(x)[y]$ and $\mathbb{C}(y)[x]$, respectively. (Recall that $\mathbb{C}[x]$ and $\mathbb{C}(x)$ stand for the \mathbb{C} -algebra of polynomials in x and the field of rational functions of x with coefficients in \mathbb{C} , respectively.)

As applications of it, we will show (1.2) and Theorems 1.3 and 1.5 in Sections 4 and 5. In Section 2, we recall basic facts about hypergeometric functions and Hecke groups and its subgroup of index 2. Section 3 is devoted to the statement of our main lemma and its proof. Theorem 1.8 is also proved in Section 3 with the help of the main lemma. We emphasize that our approach does not need ingenious ideas as in [11]. Without knowledge about Ramanujan's formulae, we can deduce those formulae possibly with laborious computations only. In this paper, we treat only the case when the Riemann surface Z is planar. When Z is non-planar, it is technically difficult to find an explicit form of the polynomial P(x, y). Let \hat{Z} be the compactification of Z. Since the parametrizations $\varphi, \psi : \hat{Z} \to \hat{\mathbb{C}}$ are *n*-sheeted covering maps with critical values contained in $\{0, 1, \infty\}$ if $|G_q : G_q \cap G_q^{M_p}| = n$, Belyi's theorem implies that the compact Riemann surface \hat{Z} is an algebraic curve defined over $\overline{\mathbb{Q}}$ (see [19]). Therefore, in principle, we could determine the surface Z and maps φ, ψ by the combinatorial information about the coverings. We hope to give further examples when Z is non-planar in the future work.

Acknowledgement. The authors would like to thank Professor Hiroshige Shiga for suggestions which simplified the proof of Lemma 3.2 in the early version of the manuscript. They also thank Professor Matti Vuorinen for many suggestions.

2. Hypergeometric functions and Hecke groups

The hypergeometric function $_2F_1(a, b; c; z)$ is defined by

$$_{2}F_{1}(a,b;c;z) = \sum_{n=0}^{\infty} \frac{(a)_{n}(b)_{n}}{(c)_{n}n!} z^{n}, \quad |z| < 1,$$

for parameters $a, b, c \in \mathbb{C}$ with $c \neq 0, -1, -2, \ldots$ By Euler's integral representation formula, we know that ${}_2F_1(a, b; c; z)$ analytically extends to the slit domain $\mathbb{C} \setminus [1, +\infty)$. Let $0 < a \leq 1/2$. Then it is classically known [13] that the function

$$\tau = f_a(z) = i \cdot \frac{{}_2F_1(a, 1-a; 1; 1-z)}{{}_2F_1(a, 1-a; 1; z)}$$

maps the upper half-plane $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$ onto the domain $\Delta_a = \{\tau \in \mathbb{H} : 0 < \text{Re } \tau < \sin \pi a, |2\tau \sin \pi a - 1| > 1\}$ and that Δ_a is the hyperbolic triangle with vertices at $0, \infty$ and $ie^{-\pi ai}$ of interior angles 0, 0 and $(1 - 2a)\pi$, respectively (see also [4, Lemma 4.1]). Note also that f_a extends homeomorphically to the boundary and

(2.1)
$$f_a(0) = \infty, \quad f_a(1) = 0, \quad f_a(\infty) = ie^{-\pi a i} = e^{\pi i (1-2a)/2}.$$

Suppose now that q = 1/(1-2a) is an integer or ∞ . Let $\pi_q : \Delta_a \to \mathbb{H}$ be the inverse map $z = f_a^{-1}(\tau)$ of $\tau = f_a(z)$. Then, by repeated applications of the Schwarz reflection principle, π_q extends to a holomorphic map from \mathbb{H} into $\widehat{\mathbb{C}} \setminus \{0, 1\}$. Here, we note that π_q is locally q to 1 at the point $ie^{-\pi ai}$. By construction, the covering group $G_q = \{\gamma \in \operatorname{Aut}(\mathbb{H}) :$ $\pi_q \circ \gamma = \pi_q\}$ is the triangle group of signature (q, ∞, ∞) arising from the hyperbolic triangle Δ_a , q = 1/(1-2a). Here, the group $\operatorname{Aut}(\mathbb{H})$ of analytic automorphism of \mathbb{H} is identified with $\operatorname{PSL}(2,\mathbb{R}) = \operatorname{SL}(2,\mathbb{R})/\{\pm I\}$. Since G_q is a subgroup of the Hecke group H_{2q} of index 2, we recall basics of Hecke groups. See [14] for details about the Hecke groups. Moreover, we refer to [6] and [20] as general references for Fuchsian groups.

By its form, the function f_a satisfies the relation $f_a(1-z) = -1/f_a(z)$. Since $\pi_q = f_a^{-1}$ on Δ_a , the following result follows.

Lemma 2.2. The covering map π_q satisfies the functional equation

$$\pi_q(\tau) + \pi_q(-1/\tau) = 1, \quad \tau \in \mathbb{H}.$$

For an integer $k \geq 3$, the Hecke group H_k is defined as the discrete subgroup of $PSL(2, \mathbb{R})$ generated by the two elements $\pm S$ and $\pm T_k$, where

$$S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad T_k = \begin{pmatrix} 1 & \lambda_k \\ 0 & 1 \end{pmatrix} \text{ and } \lambda_k = 2\cos\frac{\pi}{k}.$$

See [6] or [14]. Here and in what follows, we often identify a matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with the Möbius transformation $\tau \mapsto A\tau = (a\tau + b)/(c\tau + d)$ so that S, T_k are regarded as elements of Aut(\mathbb{H}) = PSL(2, \mathbb{R}). Note that H_3 is the classical modular group PSL(2, \mathbb{Z}). We put

$$U_k = T_k S = \begin{pmatrix} \lambda_k & -1 \\ 1 & 0 \end{pmatrix}.$$

Then U_k is an elliptic element of order k with fixed point at $e^{\pi i/k}$. Let F_k be the set of points $\tau \in \mathbb{H}$ with $|\operatorname{Re} \tau| \leq \cos \frac{\pi}{k}, |\tau| \geq 1$. Then F_k is a fundamental domain for H_k with side pairing transformations S, T_k . (Note that S has a fixed point at $\tau = i$.) In particular, we see that H_k is a triangle group of signature $(2, k, \infty)$.

From now on, we restrict ourselves on the case when k = 2q is an even number with $q \ge 2$. Then $\hat{F}_q = F_{2q} \cup S(F_{2q})$ is a fundamental domain for the (normal) subgroup G of H_{2q} of index 2 generated by

$$T_{2q} = \begin{pmatrix} 1 & \lambda_{2q} \\ 0 & 1 \end{pmatrix}$$
 and $W_{2q} = S^{-1}T_{2q}^{-1}S = \begin{pmatrix} 1 & 0 \\ \lambda_{2q} & 1 \end{pmatrix}$.

To adapt with our aim, we modify the fundamental domain as follows. For a = (q - 1)/(2q), let $\tilde{F}_q = \Delta_a \cup \Delta'_a$, where Δ'_a is the reflection of Δ_a across the line $\operatorname{Re} \tau = \sin \pi a$. Then, \tilde{F}_q serves as a fundamental domain of G, which is the same as G_q introduced in Introduction and the above-defined covering group of π_q . (This appears in Example 1.3 of [14].) Note that the element

$$V_q := T_{2q} W_{2q}^{-1} = U_{2q}^2 = \begin{pmatrix} \lambda_{2q}^2 - 1 & -\lambda_{2q} \\ \lambda_{2q} & -1 \end{pmatrix} = \begin{pmatrix} \lambda_q + 1 & -\lambda_{2q} \\ \lambda_{2q} & -1 \end{pmatrix}$$

is elliptic of order q and that G_q is generated by T_{2q} and V_q (see Figure 1 and Figure 2). Here, we note the following property.



FIGURE 1. Fundamental domains of $H_{2q} = \langle T_{2q}, S \rangle$ and $G = \langle T_{2q}, W_{2q} \rangle$.

Lemma 2.3. Let p, q be integers with $p, q \ge 2$ (where q may be ∞) and let $K = G_q \cap G_q^{M_p}$, where $M_p \tau = p\tau$. Then $|G_q: K| = |G_q^{M_p}: K|$.

Proof. If both of the indices are infinite, there is nothing to show. Assume $n = |G_q: K| < \infty$. Then the hyperbolic area Area(Z) of the Riemann orbifold $Z = K \setminus \mathbb{H}$ is computed as (see [6, p. 150] or [20, p. 13])

$$\operatorname{Area}(Z) = n \operatorname{Area}(G_q \setminus \mathbb{H}) = n \operatorname{Area}(F_q) = 2n(\pi - \pi/q).$$



FIGURE 2. Fundamental domain \tilde{F}_q of $G_q = \langle T_{2q}, V_q \rangle$.

Since $G_q^{M_p} \setminus \mathbb{H}$ has the same hyperbolic area as $G_q \setminus \mathbb{H}$, we have

$$|G_q^{M_p}:K| = \frac{\operatorname{Area}(K \setminus \mathbb{H})}{\operatorname{Area}(G_q^{M_p} \setminus \mathbb{H})} = \frac{\operatorname{Area}(Z)}{\operatorname{Area}(G_q \setminus \mathbb{H})} = n$$

as required. When $|G_q^{M_p}:K| < \infty$, the same argument works for the proof.

Remark 2.4. In view of the proof, one can also show that the formula $|G : G \cap G^M| = |G^M : G \cap G^M|$ holds for a cofinite Fuchsian group G; namely, if the surface $G \setminus \mathbb{H}$ has finite hyperbolic area. However, this formula is not true in general. For instance, Jørgensen, Marden and Pommerenke [18] constructed a Fuchsian group Γ and its subgroup G of index 2 such that $G^M = M^{-1}GM$ is a proper subgroup of G for some $M \in \Gamma$ so that $G \cap G^M = G^M$. In particular, $|G : G \cap G^M| > |G^M : G \cap G^M| = 1$.

The index $|G_q : K|$ can be computed explicitly when $q = 3, 4, \infty$ (see [2]). In the present paper, our main concerns are about the limiting case as $q \to \infty$ and the case of q = 3. Those cases will be discussed in Sections 4 and 5, respectively.

3. A result on Fuchsian groups

Let Γ be a Fuchsian group acting on the upper half-plane \mathbb{H} and X be the quotient Riemann surface $\Gamma \setminus \mathbb{H}$. We denote by π the canonical projection $\pi : \mathbb{H} \to X$. When $\gamma \in \Gamma$ is a parabolic element with fixed point at $\tau_0 \in \partial \mathbb{H}$, it is known that $\pi(\tau)$ tends to a puncture, say P, of X as $\tau \to \tau_0$ nontangentially. As a convention, we will write $\pi(\tau_0) = P$ in the sequel. A point $\tau_0 \in \mathbb{H}$ is called a fixed point of an element γ of PSL(2, \mathbb{R}) if $\gamma \tau_0 = \tau_0$ and the set of fixed points of γ in \mathbb{H} is denoted by Fix(γ). A non-identity element γ is called elliptic if Fix($\gamma \neq \emptyset$. Let $M \in PSL(2, \mathbb{R}) = Aut(\mathbb{H})$ such that $M \notin \Gamma$. Then we consider

the number (possibly ∞) defined by

(3.1)
$$N_{X_0}(M,\Gamma) = \sup_{x \in X_0} \#\{\pi(M\tau) : \tau \in \pi^{-1}(x)\}.$$

for $X_0 \subset X$. The next result is our main lemma and it gives a criterion for finiteness of $N_{X_0}(M, \Gamma)$.

Lemma 3.2. Let Γ be a Fuchsian group and $M \in PSL(2, \mathbb{R}) \setminus \Gamma$. Suppose that X_0 is an uncountable subset of the quotient Riemann surface $X = \Gamma \setminus \mathbb{H}$. Then $N_{X_0}(M, \Gamma) < \infty$ if and only if $\Gamma^M := M^{-1}\Gamma M$ is commensurable with Γ in the sense that $|\Gamma : G| < \infty$ for $G = \Gamma \cap \Gamma^M$. Moreover, in this case, $N_{X_0}(M, \Gamma) = |\Gamma : G|$.

Proof. The "if" part is almost trivial. Let $N := |\Gamma : G| < \infty$. Then we take $\gamma_1, \ldots, \gamma_N \in \Gamma$ so that $\Gamma = G\gamma_1 \cup \cdots \cup G\gamma_N$. For $x \in X$ and $\tau_0 \in \pi^{-1}(x)$, we observe

$$M(\pi^{-1}(x)) = M\Gamma\tau_0 = \bigcup_{j=1}^{N} MG\gamma_j\tau_0 = \bigcup_{j=1}^{N} MGM^{-1} \cdot M\gamma_j\tau_0.$$

Since $MGM^{-1} \subset \Gamma$, the set $MGM^{-1} \cdot M\gamma_j\tau_0$ is projected to the point $\pi(M\gamma_j\tau_0)$ by π . Therefore, $\#(\pi(M(\pi^{-1}(x)))) \leq N$. Hence, we conclude that

(3.3)
$$N_{X_0}(M,\Gamma) \le |\Gamma:G|.$$

To show the "only if" part, we assume that $N := N_{X_0}(M, \Gamma)$ is finite. Let

$$E_1 = \bigcup_{\gamma,\delta} \operatorname{Fix}(\gamma^{-1}\delta),$$

where γ and δ range over $\gamma \in \Gamma$ and $\delta \in \Gamma^M$ with $\gamma \neq \delta$. Since each $\operatorname{Fix}(\gamma^{-1}\delta)$ contains at most one point, the set E_1 is at most countable. Moreover, the set $E = \Gamma \cdot E_1 = \{\gamma \tau : \gamma \in \Gamma, \tau \in E_1\}$ is also at most countable. Take a point τ_0 from the uncountable subset $\pi^{-1}(X_0) \setminus E$ of \mathbb{H} and fix it. We regard $G \setminus \Gamma$ as the set of right cosets $\{G\gamma : \gamma \in \Gamma\}$. As we saw above, each set $G\gamma \cdot \tau_0$ projects to one point $\pi(M\gamma\tau_0)$ under the mapping $\pi \circ M :$ $\mathbb{H} \to X$. We now show that the mapping $\phi : G \setminus \Gamma \to X$ defined by $\phi(G\gamma) = \pi(M\gamma\tau_0)$ is injective. To this end, we suppose that $\phi(G\gamma_1) = \phi(G\gamma_2)$ for some $\gamma_1, \gamma_2 \in \Gamma$. Then $M\gamma_2\tau_0 = \gamma M\gamma_1\tau_0$ for some $\gamma \in \Gamma$. It says that $\gamma_1\tau_0$ is a fixed point of $(\gamma_2\gamma_1^{-1})^{-1}\delta$, where $\delta = M^{-1}\gamma M \in \Gamma^M$. Since $\gamma_1\tau_0 \notin E$, the element $(\gamma_2\gamma_1^{-1})^{-1}\delta$ must be the identity, which implies $\gamma_2\gamma_1^{-1} = \delta \in \Gamma \cap \Gamma^M = G$. Hence $G\gamma_1 = G\gamma_2$.

We have seen that $\phi: G \setminus \Gamma \to X$ is injective. On the other hand, the image $\phi(G \setminus \Gamma) = \pi(M\Gamma\tau_0)$ consists of at most N points. Therefore, we obtain

$$|\Gamma:G| = \#(G \setminus \Gamma) \le N.$$

Combining with (3.3), we obtain $|\Gamma : G| = N_{X_0}(M, \Gamma)$ as required.

In particular, we see that $N_{X_0}(M, \Gamma)$ does not depend on the uncountable set $X_0 \subset X$. Thus we denote by $N(M, \Gamma)$ the common number $N_{X_0}(M, \Gamma)$.

We are now ready to prove Theorem 1.8.

Proof of Theorem 1.8. Recall that a = (q-1)/(2q) and $M_p\tau = p\tau$. In this proof, we will use the notation introduced in Section 2. By definition, the generalized modular equation (1.1) may be expressed by $f_a(\beta) = pf_a(\alpha)$. Therefore, α and β in $\widehat{\mathbb{C}} \setminus \{0, 1\}$ satisfy (1.1) if and only if $\alpha = \pi_q(\tau), \beta = \pi_q(p\tau)$ for some $\tau \in \mathbb{H}$.

First we assume that α and β in (1.1) satisfy the algebraic equation $P(\alpha, \beta) = 0$, where

$$P(x,y) = \sum_{j=0}^{n} a_j(x)y^j, \quad a_j(x) \in \mathbb{C}[x] \ (j = 0, 1, \dots, n),$$

is an irreducible polynomial of degree n in $\mathbb{C}(x)[y]$. Let X_0 be the set of those points $x \in X$ for which $a_k(x) \neq 0$ for some k. Then for a fixed $x_0 \in X_0$, the algebraic equation $P(x_0, y) = 0$ in y has at most n solutions. Thus we conclude that $N_{X_0}(M_p, G_q) \leq n$. The previous lemma now implies that $|G_q : G_q \cap G_q^{M_p}| \leq n$. By the irreducibility of P(x, y), we see that equality holds.

Conversely, we assume that the equality $|G_q: K| = n$ holds, where $K = G_q \cap G_q^{M_p}$. Note that $|G_q^{M_p}: K| = n$ by Lemma 2.3. We denote by Z the quotient Riemann surface $K \setminus \mathbb{H}$. Let $\rho: \mathbb{H} \to Z$ be the canonical projection and let $\varphi: Z \to X$ and $\psi: Z \to X$ be the induced mappings satisfying the relations $\pi_q = \varphi \circ \rho$ and $\pi_q \circ M_p = \psi \circ \rho$, respectively. Thus we have the following commutative diagram:



Note that the solution (α, β) to the modular equation (1.1) is now parametrized by $\alpha = \varphi(z)$ and $\beta = \psi(z)$ for $z \in Z$. We denote by \hat{X} and \hat{Z} the compact Riemann surfaces obtained by filling in the punctures of X and Z, respectively. Note that $\widehat{\mathbb{C}}$ can be taken as \hat{X} . Then φ and ψ extend to n-sheeted branched covering maps of \hat{Z} onto $\hat{X} = \widehat{\mathbb{C}}$. In particular, ψ may be regarded as a meromorphic function on \hat{Z} . Let X_0 denote X minus the set of critical values of $\varphi : Z \to X$; indeed $X_0 = \widehat{\mathbb{C}} \setminus \{0, 1, \infty\}$ in this case. Similarly, X_1 is defined for $\psi : Z \to X$. For a point $x_0 \in X_0$ we choose a small disk $U = U(x_0)$ with $x_0 \in U \subset X_0$. Let s_j be the elementary symmetric functions of $\psi \circ \eta_1, \ldots, \psi \circ \eta_n$ of degree j, where η_1, \ldots, η_n are local inverses of φ of U. Then for each $j = 1, 2, \ldots, n$, all $s_j : U(x_0) \to X$ piece together to one function and it extends meromorphically to $\hat{X} = \widehat{\mathbb{C}}$ and ψ satisfies the equation

$$\psi^n - s_1 \circ \varphi \cdot \psi^{n-1} + \dots + (-1)^{n-1} s_{n-1} \circ \varphi \cdot \psi + (-1)^n s_n \circ \varphi = 0$$

(see [16, Theorem 8.3] for details). Note that each $s_j : \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ is a rational function. By writing $(-1)^j s_j(x) = a_j(x)/a_0(x)$ for polynomials $a_0(x), \ldots, a_n(x) \in \mathbb{C}[x]$ without non-trivial common factor, we define

$$P(x,y) = a_0(x)y^n + a_1(x)y^{n-1} + \dots + a_{n-1}(x)y + a_n(x).$$

Then $P(\alpha, \beta) = P(\varphi(z), \psi(z)) = 0$ for $z \in Z$. Conversely, suppose that $x_0 \in X_0$ with $a_0(x_0) \neq 0$ and $y_0 \in X_1$ satisfy $P(x_0, y_0) = 0$. Let $\varphi^{-1}(x_0) = \{z_1, \ldots, z_n\}$ and let $\sigma_1, \ldots, \sigma_n$ be the elementary symmetric functions of $\psi(z_1), \ldots, \psi(z_n)$. Note that $\sigma_j = s_j(x_0)$ for $j = 1, \ldots, n$. Then,

$$a_0(x_0)(y_0 - \psi(z_1)) \cdots (y_0 - \psi(z_n)) = a_0(x_0)(y_0^n - \sigma_1 y_0^{n-1} + \dots + (-1)^n \sigma_n)$$
$$= \sum_{j=0}^n a_j(x_0) y_0^{n-j} = P(x_0, y_0) = 0.$$

Hence, $y_0 = \psi(z_k)$ for some k. Since $x_0 = \varphi(z_k)$, $y_0 = \psi(z_k)$, we have shown that the converse is true. We next show that the polynomial P(x, y) is irreducible in y with coefficients in $\mathbb{C}[x]$. Suppose, on the contrary, that P(x, y) reduces to the product $P_1(x, y)P_2(x, y)$ of nonconstant polynomials P_1, P_2 . Let $Y_l = \{(x, y) \in X_0 \times X_1 : P_l(x, y) = 0\}$ for l = 1, 2. We claim that $Y_1 \cap Y_2 = \emptyset$. Indeed, if $(x_0, y_0) \in Y_1 \cap Y_2$, then the polynomial $P(x_0, y) = P_1(x_0, y)P_2(x_0, y)$ has a multiple zero at $y = y_0$, which is impossible because the set $\psi^{-1}(y_0)$ consists of n points for $y_0 \in X_1$. Since $\varphi \times \psi : Z_0 \to X_0 \times X_1$ is contained in either Y_1 or Y_2 . However, this is impossible. The claim has been shown.

Finally, consider the polynomial P(x, y) in $\mathbb{C}(y)[x]$. Then, P(x, y) is a polynomial in x with coefficients in $\mathbb{C}[y]$, and we may write

$$P(x,y) = b_0(y)x^m + b_1(y)x^{m-1} + \dots + b_{m-1}(y)x + b_m(y), \quad b_j[y] \in \mathbb{C}[y],$$

where $m = \max_{0 \le j \le n} \deg a_j(x)$. Note that $b_0 \ne 0$. Since $\psi : Z \rightarrow X$ is *n*-sheeted, as in the case of $\varphi : Z \rightarrow X$, we can see that m = n and P(x, y) is irreducible in $\mathbb{C}(y)[x]$. \Box

Let us summarize the above observations for later use. Suppose that $K = G_q \cap G_q^{M_p}$ is a subgroup of G_q of finite index n. The intermediate Riemann surface $Z = K \setminus \mathbb{H}$ may be used as a parameter space of the solutions (α, β) to (1.1). Indeed, the solutions are given by $\alpha = \varphi(z), \beta = \psi(z)$ for $z \in Z$, where $\varphi, \psi : Z \to X = G_q \setminus \mathbb{H}$ are (possibly branched) covering maps satisfying the relations $\varphi(\rho(\tau)) = \pi_q(\tau)$ and $\psi(\rho(\tau)) = \pi_q(p\tau)$ for $\tau \in \mathbb{H}$. Note that φ and ψ extend to the compactifications \hat{Z} to $\hat{X} = \widehat{\mathbb{C}}$ as *n*-sheeted branched (analytic) covering maps. The polynomial P(x, y) whose zero set describes the solutions (α, β) can be computed as in the above proof. The following lemma helps us to compute ψ when we know about φ .

Lemma 3.4. Under the assumption of Theorem 1.8, the Möbius transformation $SM_p\tau = -1/(p\tau)$ induces an analytic involution $\omega : Z \to Z$ which satisfies the relation $\rho \circ SM_p = \omega \circ \rho$ and the functional equation

$$\psi = 1 - \varphi \circ \omega.$$

Proof. Let $K = G_q \cap G_q^{M_p}$. We first note that the Möbius transformations S and M_p satisfy the relation $SM_p = M_p^{-1}S$ and thus $M_pSM_p = S$. Recall that S normalizes G_q ; namely, $G_q^S := S^{-1}G_qS = G_q$. Therefore,

$$K^{SM_p} = G_q^{SM_p} \cap (G_q^{M_p})^{SM_p} = G_q^{M_p} \cap G_q^{M_p SM_p} = G_q^{M_p} \cap G_q^S = K,$$

which means that $SM_p : \mathbb{H} \to \mathbb{H}$ descends to an automorphism $\omega : Z \to Z$ such that $\rho \circ SM_p = \omega \circ \rho$. Since $(SM_p)^2 = I$, we see that $\omega \circ \omega = \text{id}$. We recall Lemma 2.2 which says that $\pi_q \circ S = 1 - \pi_q$. Then we compute

$$\psi \circ \omega \circ \rho = \psi \circ \rho \circ SM_p = (\pi_q \circ M_p) \circ SM_p = \pi_q \circ S = 1 - \pi_q = 1 - \varphi \circ \rho$$

and therefore we have $\psi \circ \omega = 1 - \varphi$. Since ω is an involution, we obtain the required relation.

Remark 3.5. Since ω comes from the normalizer of K, ω is indeed an automorphism of the Riemann *orbifold* $K \setminus \mathbb{H}$. In particular, ω maps a cone point of angle $2\pi/m$ to another (possibly the same) cone point of the same angle for an integer $m \geq 2$.

4. Applications to the classical case $q = \infty$

As $q \to \infty$, the Hecke group H_{2q} converges to the group $H_{\infty} = \langle S, T_{\infty} \rangle$, which is a subgroup of the modular group $PSL(2,\mathbb{Z})$ of index 3, where $T_{\infty}z = z + 2$. At the same time, the subgroup G_q converges to the principal congruence subgroup $\Gamma(2)$ of level 2, which will be denoted by G_{∞} in this section. That is to say,

$$G_{\infty} = \{A \in \mathrm{SL}(2,\mathbb{Z}) : A \equiv I \mod 2\} / \{\pm I\}.$$

This corresponds to the case when a = 1/2 in the modular equation (1.1). The fundamental domain \tilde{F}_{∞} is described by $0 \leq \text{Re } \tau \leq 2, |\tau - 1/2| \geq 1/2, |\tau - 3/2| \geq 1/2$ for $\tau \in \mathbb{H}$. A pair of generators of G_{∞} are given by

$$T = T_{\infty} = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$$
 and $V = \begin{pmatrix} 3 & -2 \\ 2 & -1 \end{pmatrix}$.

Note that V is a parabolic element with fixed point at $\tau = 1$ and that V(0) = 2. Also, let

$$W = -V^{-1}T = \begin{pmatrix} 1 & 0\\ 2 & 1 \end{pmatrix}.$$

It is well known that $\Gamma(2) = G_{\infty}$ is a Fuchsian group uniformizing the thrice punctured sphere $\widehat{\mathbb{C}} \setminus \{0, 1, \infty\}$ (see §§3.4-5 in Chapter 7 of [1] and §§4.3 in Chapter 1 of [15], where the symbol λ is used for π_{∞}). The canonical projection $\pi_{\infty} : \mathbb{H} \to X$ may be considered to be $\pi_{\infty} : \mathbb{H} \to \widehat{\mathbb{C}} \setminus \{0, 1, \infty\} = \mathbb{C} \setminus \{0, 1\}$ with

(4.1)
$$\pi_{\infty}(0) = 1, \quad \pi_{\infty}(\infty) = 0 \quad \text{and} \quad \pi_{\infty}(1) = \infty$$

by (2.1). Let $\rho : \mathbb{H} \to Z, \varphi, \psi : Z \to X$ be as in the proof of Theorem 1.8. Here is a simple but useful observation. The conjugation of a matrix by the Möbius transformation M_p is computed by

$$M_p^{-1} \begin{pmatrix} a & b \\ c & d \end{pmatrix} M_p = \begin{pmatrix} a & b/p \\ pc & d \end{pmatrix}$$

Hence, an element $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ of G_{∞} is a member of $G_{\infty}^{M_p}$ precisely if $c \equiv 0 \pmod{2p}$.

4.1. Case p = 2. We first look at the most classical case when p = 2. Let $K = G_{\infty} \cap G_{\infty}^{M_2}$. Then

$$A_1 := T = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}, \quad A_2 := V^2 = \begin{pmatrix} 5 & -4 \\ 4 & -3 \end{pmatrix}, \quad A_3 := V^{-1}TV = \begin{pmatrix} -3 & 2 \\ -8 & 5 \end{pmatrix}$$

are side-paring transformations of the hyperbolic polygon $F = \tilde{F}_{\infty} \cup V^{-1}(\tilde{F}_{\infty})$ (see Figure 3).



FIGURE 3. Fundamental domain F of $K = G_{\infty} \cap G_{\infty}^{M_2}$.

Therefore, the elements A_1, A_2, A_3 generate a subgroup of G_{∞} of index 2. In view of the forms of A_j , we see that $A_j \in K$ for j = 1, 2, 3 and thus we have $K = \langle A_1, A_2, A_3 \rangle$ and $|G_{\infty}: K| = 2$. It is easy to observe that the quotient surface $Z = K \setminus \mathbb{H}$ is conformally equivalent to a four-times punctured sphere. Thus we can assume that $Z = \widehat{\mathbb{C}} \setminus \{0, \infty, 1, b\}$, where $\rho(0) = 0, \rho(1) = \infty, \rho(\infty) = 1, \rho(1/2) = b$. In view of (4.1), we have the conditions $\varphi(0) = 1, \varphi(\infty) = \infty, \varphi(1) = \varphi(b) = 0$. Here we used the fact that 1/2 is conjugate to ∞ under the action of G_{∞} . We note that the extension $\varphi: \widehat{Z} = \widehat{\mathbb{C}} \to \widehat{X} = \widehat{\mathbb{C}}$ is a rational map of degree 2. Since $\varphi^{-1}(1) = \{0\}, \varphi^{-1}(\infty) = \{\infty\}$, we see that φ takes the values 0 and ∞ with multiplicity 2. Therefore, φ should have the form $\varphi(z) = cz^2 + 1$ for a constant $c \neq 0$. Since $\varphi(1) = 0$, we conclude that c = -1 and b = -1. Hence, $\varphi(z) = 1 - z^2$ under the above normalization.

Next we determine the form of $\psi : Z \to X$. By the relation $\pi_{\infty}(2\tau) = \psi(\rho(\tau))$, we have the necessary conditions $\psi(0) = \pi_{\infty}(0) = 1, \psi(\infty) = \pi_{\infty}(2) = \pi_{\infty}(0) = 1, \psi(1) = \pi_{\infty}(\infty) = 0$ and $\psi(-1) = \pi_{\infty}(1) = \infty$. In particular, $\psi^{-1}(0) = \{1\}, \psi^{-1}(\infty) = \{-1\}$. Since $\psi : \hat{Z} = \widehat{\mathbb{C}} \to \hat{X} = \widehat{\mathbb{C}}$ is a rational map of degree 2, ψ has the form

$$\psi(z) = \frac{c(z-1)^2}{(z+1)^2}$$

for a constant $c \neq 0$. Since $\psi(0) = 1$, we have c = 1. We note that $\psi(\infty) = 1$ is also satisfied. In this way, the solution (α, β) of the modular equation (1.1) with a = 1/2, p = 2 is parametrized by

$$\alpha = \varphi(z) = 1 - z^2$$
 and $\beta = \psi(z) = \frac{(z-1)^2}{(z+1)^2}$

Eliminating the variable z, we see that

$$\beta = \left(\frac{1 - \sqrt{1 - \alpha}}{1 + \sqrt{1 - \alpha}}\right)^2,$$

which is equivalent to (1.2) with $\alpha = r^2$ and $\beta = s^2$. The polynomial in Theorem 1.8 is given by

$$P(x,y) = x^2y^2 - 2(x^2 - 8x + 8)y + x^2.$$

In this case, the polynomial is not symmetric in x and y.

4.2. Case p = 3. Next we consider the case when p = 3. We now consider the hyperbolic 10-gon $F = \tilde{F}_{\infty} \cup V(\tilde{F}_{\infty}) \cup V^{-1}(\tilde{F}_{\infty}) \cup W(\tilde{F}_{\infty})$ with vertices at $0, 1/3, 2/5, 1/2, 2/3, 1, 4/3, 3/2, 2, \infty$ in the counterclockwise order. Then the elements

$$A_{1} := T, \quad A_{2} := V^{-1}T^{2}V^{-1} = \begin{pmatrix} 5 & -8\\ 12 & -19 \end{pmatrix}, \quad A_{3} := V^{-1}T^{-1}V^{-1} = \begin{pmatrix} -7 & 10\\ -12 & 17 \end{pmatrix},$$
$$A_{4} := V^{-3} = \begin{pmatrix} -5 & 6\\ -6 & 7 \end{pmatrix}, \quad A_{5} := V^{-1}TV^{-1}T^{-1}V = \begin{pmatrix} -5 & 2\\ -18 & 7 \end{pmatrix}$$

form side-pairing transformations of F (see Figure 4).

Therefore, $K = \langle A_1, A_2, A_3, A_4, A_5 \rangle$ is a subgroup of G_{∞} for which F is a fundamental domain. In view of the forms of A_j , we observe that $K \subset G_{\infty} \cap G_{\infty}^{M_3}$. On the other hand, the elements V, V^{-1}, W are not contained in $G_{\infty}^{M_3}$, therefore $K = G_{\infty} \cap G_{\infty}^{M_3}$ and $|G_{\infty}:K| = 4$. It is a simple task to see that the quotient surface $Z = K \setminus \mathbb{H}$ is a six-times punctured sphere. Recall that $\varphi, \psi: Z \to X$ extend to rational functions of degree 4. We may normalize the punctures so that $\rho(\infty) = 0, \rho(0) = \rho(2/5) = \rho(2) = 1, \rho(1/3) = \infty, \rho(1/2) = \rho(3/2) = b, \rho(1) = c, \rho(2/3) = \rho(4/3) = d$. Thus $\varphi(0) = \pi_{\infty}(\infty) = 0$. Similarly, we obtain $\varphi(1) = 1, \varphi(\infty) = \infty, \varphi(b) = 0, \varphi(c) = \infty, \varphi(d) = 1$. We have to



FIGURE 4. Fundamental domain F of $K = G_{\infty} \cap G_{\infty}^{M_3}$.

compute multiplicities of these values. For instance, we have $\varphi^{-1}(\infty) = \{\infty, c\}$. The part of the basic fundamental domain \tilde{F}_{∞} corresponding to ∞ under π_{∞} is the cusp neighbourhood $D = \tilde{F}_{\infty} \cap \{\tau \in \mathbb{H} : |\tau - 1| < \varepsilon\}$ for a small enough $\varepsilon > 0$. Since V and V^{-1} fix 1 while W sends 1 to 1/3, the multiplicity of φ at $\rho(1) = c$ is 3 and that is 1 at $\rho(1/3) = \infty$. We write $(\varphi)_{\infty} = 3 \cdot c + 1 \cdot \infty$ as a divisor for short¹. In the same way, we have $(\varphi)_0 = 3 \cdot b + 1 \cdot 0$ and $(\varphi)_1 = 1 \cdot d + 3 \cdot 1$. In particular, φ may be expressed by

$$\varphi(z) = e \frac{z(z-b)^3}{(z-c)^3}$$
 and $\varphi(z) - 1 = e' \frac{(z-d)(z-1)^3}{(z-c)^3}$

for constants e and e'. Since b, c, d are different from 1, we have the unique solution b = -2, c = -1/2, d = -1 and e = e' = 1/8. Hence,

$$\alpha = \varphi(z) = \frac{z(z+2)^3}{(2z+1)^3}.$$

Next we determine the form of ω given in Lemma 3.4. Since SM_3 swaps 0 and ∞ (respectively 1/3 and -1), ω swaps $\rho(0) = 1$ and $\rho(\infty) = 0$ (respectively, $\rho(1/3) = \infty$ and $\rho(-1) = \rho(1) = -1/2$). Hence, $\omega(z) = (1-z)/(1+2z)$. By Lemma 3.4, we obtain

$$\beta = \psi(z) = 1 - \varphi(\omega(z)) = \frac{z^3(z+2)}{2z+1}.$$

¹When the equation $\varphi(z) = w$ has solutions z_j with multiplicities m_j for j = 1, 2, ..., N, we write $(\varphi)_w = m_1 \cdot z_1 + m_2 \cdot z_2 + \cdots + m_N \cdot z_N$ as a divisor on Z.

We compute

$$\alpha\beta = \frac{z^4(z+2)^4}{(2z+1)^4}$$
 and $(1-\alpha)(1-\beta) = \frac{(1-z^2)^4}{(2z+1)^4}.$

Note that φ and ψ both map the interval [0, 1] onto itself homeomorphically. Hence, for $\alpha, \beta \in [0, 1]$, we obtain the relation

(4.2)
$$(\alpha\beta)^{1/4} + \left\{ (1-\alpha)(1-\beta) \right\}^{1/4} = 1.$$

This is known as Legendre's modular equation (see (5.37) in [5] and (4.1.16) in [12]). Formula (4.2) may be transformed to the polynomial equation $P(\alpha, \beta) = 0$ as in Theorem 1.8, where

$$P(x,y) = y^4 + 2x^3y^3 - 2xy - x^4.$$

This is known as the modular equation of third order in Jacobi's form (see (3.42) in [24]).

5. Case of degree 3 in the theory of signature 3

When q = 3, the standard generators of G_3 are

$$T := T_6 = \begin{pmatrix} 1 & \sqrt{3} \\ 0 & 1 \end{pmatrix}$$
 and $V := V_3 = \begin{pmatrix} 2 & -\sqrt{3} \\ \sqrt{3} & -1 \end{pmatrix}$.

Recall that V is an elliptic element of order 3 with fixed point at $\tau_0 := e^{\pi i/6} = (\sqrt{3} + i)/2$. Recall also that the canonical projection $\pi_3 : \mathbb{H} \to X = \widehat{\mathbb{C}} \setminus \{0, 1\}$ satisfies

(5.1)
$$\pi_3(0) = 1, \quad \pi_3(\infty) = 0 \quad \text{and} \quad \pi_3(\tau_0) = \infty$$

by (2.1). The following result is useful.

Lemma 5.2. $A \in PSL(2, \mathbb{R})$ belongs to G_3 precisely when A is represented by a matrix of the form

(5.3)
$$\begin{pmatrix} a & b\sqrt{3} \\ c\sqrt{3} & d \end{pmatrix}, \quad a, b, c, d \in \mathbb{Z}, \ ad - 3bc = 1.$$

Proof. We denote by G the Möbius transformations represented by the matrices in (5.3). Then it is well known that G is a proper subgroup of H_6 (see [17]). On the other hand, it is obvious that G_3 is contained in G. Since $|H_6:G_3| = 2$, we have $G = G_3$ as required. \Box



FIGURE 5. Fundamental domain F of $K = G_3 \cap G_3^{M_3}$.

5.1. Case p = 3. Since the case of p = 3 is simpler than that of p = 2, we start with this case. Let $F = \tilde{F}_3 \cup V(\tilde{F}_3) \cup V^2(\tilde{F}_3)$. Then F is a hyperbolic polygon with six vertices at $0, \sqrt{3}/3, \sqrt{3}/2, 2\sqrt{3}/3, \sqrt{3}, \infty$. Let K be the group generated by the side pairing transformations

$$A_1 = T, \quad A_2 = VTV^{-1} = \begin{pmatrix} -5 & 4\sqrt{3} \\ -3\sqrt{3} & 7 \end{pmatrix}, \quad A_3 = V^{-1}TV = \begin{pmatrix} -2 & \sqrt{3} \\ -3\sqrt{3} & 4 \end{pmatrix}$$

of F (see Figure 5).

It is easy to see that $K = G_3 \cap G_3^{M_3}$ by Lemma 5.2 and that K is torsion-free and satisfies $|G_3:K| = 3$. Since the quotient surface $Z = K \setminus \mathbb{H}$ is a four-times punctured sphere, we may normalize $\rho:\mathbb{H} \to Z$ so that $\rho(\infty) = 1, \rho(\sqrt{3}/3) = a, \rho(2\sqrt{3}/3) = b, \rho(0) = \rho(\sqrt{3}/2) = \rho(\sqrt{3}) = 0$ and $\rho(\tau_0) = \infty$. By the form of K, the element V normalizes K; that is, $V^{-1}KV = K$. Thus V induces an analytic automorphism $v:Z \to Z$. We recall that V is a rotation about τ_0 of angle $-2\pi/3$. It is clear that v(0) = 0 and $v(\infty) = \infty$. Since V satisfies $V(\infty) = 2\sqrt{3}/3, V(2\sqrt{3}/3) = \sqrt{3}/3$ and $V(\sqrt{3}/3) = \infty$, the map v should have the form $v(z) = e^{-2\pi i/3}z$. Hence, $b = e^{-2\pi i/3} = -(1 + i\sqrt{3})/2$ and $a = \bar{b} = (-1 + i\sqrt{3})/2$. We will determine the forms of rational maps $\varphi, \psi: \hat{Z} = \widehat{\mathbb{C}} \to \hat{X} = \widehat{\mathbb{C}}$ of degree 3. Since φ has a branch point at ∞ of order 3, we have $\varphi^{-1}(\infty) = \{\infty\}$. In particular, φ is a polynomial of degree 3. We also have $\varphi^{-1}(1) = \{0\}$. Therefore, φ should be of the form $1 + cz^3$ for a constant c. Since SM_3 swaps 0 and ∞ (respectively $\sqrt{3}/3$ and $-\sqrt{3}/3 \equiv 2\sqrt{3}/3$ (mod K), ω swaps 1 and 0 (respectively, a and b), Hence, after some

computations, we get the form $\omega(z) = (1-z)/(1+2z)$. We now apply Lemma 3.4 to obtain $\psi(z) = 1 - \varphi(\omega(z)) = \omega(z)^3 = (1-z)^3/(1+2z)^3$. In summary, we have

$$\alpha = \varphi(z) = 1 - z^3$$
 and $\beta = \psi(z) = \frac{(1-z)^3}{(1+2z)^3}$

We are now able to show Theorem 1.5 easily. The polynomial P(x, y) in Theorem 1.8 is computed as

$$P(x,y) = (8x - 9)^3 y^3 + 3 (64x^3 + 504x^2 - 1053x + 486) y^2 + 3 (8x^3 - 171x^2 + 405x - 243) y + x^3.$$

5.2. Case p = 2. Let F be the same as in the previous subsection. We choose side-pairing transformations A_1, A_2, A_3 of F at this time as follows:

$$A_1 = T, \quad A_2 = V^2 T V^2 = \begin{pmatrix} 1 & -\sqrt{3} \\ 2\sqrt{3} & -5 \end{pmatrix}, \quad A_3 = (SV)^{-1} T S V = \begin{pmatrix} 5 & -3\sqrt{3} \\ 4\sqrt{3} & -7 \end{pmatrix}.$$

Then A_1, A_2 and A_3 generate the torsion-free group $K = G_3 \cap G_3^{M_2}$ and the relation $|G_3:K| = 3$ follows (see Figure 6).



FIGURE 6. Fundamental domain F of $K = G_3 \cap G_3^{M_2}$.

We also see that $Z = K \setminus \mathbb{H}$ is a four-times punctured sphere. We normalize the map $\rho : \mathbb{H} \to Z$ so that $\rho(\infty) = 0, \rho(0) = \rho(\sqrt{3}) = 1$ and $\rho(\tau_0) = -1$. Let $b = \rho(\sqrt{3}/3) = \rho(2\sqrt{3}/3)$ and $c = \rho(\sqrt{3}/2)$. Since $\pi_3(\tau) = \varphi(\rho(\tau))$ has a branch point of order 3 at $\tau = \tau_0$, the map $\varphi(z)$ has a branch point of order 3 at z = -1; that is, $\varphi^{-1}(\infty) = \{-1\}$. Since

 $V(\infty) = 2\sqrt{3}/3, V^2(\infty) = \sqrt{3}/3$, we have $(\varphi)_0 = 1 \cdot 0 + 2 \cdot b$. Therefore, the rational map φ of degree 3 should have the form

$$\varphi(z) = \frac{mz(z-b)^2}{(z+1)^3}$$

for a constant $m \neq 0$. Observe that $V(\tilde{F}_3)$ and $V^2(\tilde{F}_3)$ share the cusps at $\sqrt{3}$ and 0, respectively, with \tilde{F}_3 . Hence, we see that $(\varphi)_1 = 2 \cdot 1 + 1 \cdot c$ and

$$\varphi(z) - 1 = \frac{m'(z-c)(z-1)^2}{(z+1)^3}$$

for a constant $m' \neq 0$. Then we have b = -3, c = -2, m = 1/2 and m' = -1/2 so that $\varphi(z) = z(z+3)^2/[2(z+1)^3]$. Next we will determine ω in Lemma 3.4. Since SM_2 swaps 0 and ∞ (respectively, $\sqrt{3}/3$ and $-\sqrt{3}/2 \equiv \sqrt{3}/2 \mod K$), ω swaps 1 and 0 (respectively, b = -3 and c = -2), the form of ω is (1-z)/(1+z). Now Lemma 3.4 yields $\psi(z) = 1 - \varphi(\omega(z)) = z^2(z+3)/4$. We summarize the results as

$$\alpha = \varphi(z) = \frac{z(z+3)^2}{2(z+1)^3}$$
 and $\beta = \psi(z) = \frac{z^2(z+3)}{4}$.

We remark that these expressions of α and β are found in Ramanujan's notebook (see [11, Theorem 6.1]). Since

$$\alpha\beta = \frac{z^3(z+3)^3}{8(z+1)^3}$$
 and $(1-\alpha)(1-\beta) = \frac{(1-z)^3(z+2)^3}{8(z+1)^3}$,

it is now easy to obtain the formula in Theorem 1.3. We also obtain the polynomial P(x, y) in Theorem 1.8 as

$$P(x,y) = (2x-1)^3 y^3 - 3x(4x^2 - 13x + 10)y^2 + 3x(2x^2 - 10x + 9)y - x^3,$$

which is equivalent to (1.4).

References

- 1. L. V. Ahlfors, Complex Analysis, 3rd ed., McGraw Hill, New York, 1979.
- 2. Md. S. Alam, On Ramanujan's modular equations and Hecke groups, Preprint (2021).
- G. D. Anderson, S.-L. Qiu, M. K. Vamanamurthy, and M. Vuorinen, Generalized elliptic integrals and modular equations, Pacific J. Math. 192 (2000), 1–37.
- 4. G. D. Anderson, T. Sugawa, M. K. Vamanamurthy, and M. Vuorinen, *Hyperbolic metric on the twice punctured sphere with one conic singularity*, Math. Z. **266** (2010), 181–191.
- G. D. Anderson, M. K. Vamanamurthy, and M. K. Vuorinen, Conformal Invariants, Inequalities, and Quasiconformal Maps, Wiley-Interscience, 1997.
- A. F. Beardon, *The Geometry of Discrete Groups*, Graduate Texts in Mathematics 91, Springer-Verlag, New York, 1983.
- 7. B. C. Berndt, Ramanujan's Notebooks, Part I, Springer-Verlag, New York, 1985.
- 8. B. C. Berndt, Ramanujan's Notebooks, Part II, Springer-Verlag, New York, 1989.
- 9. B. C. Berndt, Ramanujan's Notebooks, Part III, Springer-Verlag, New York, 1991.
- 10. B. C. Berndt, Ramanujan's Notebooks, Part V, Springer-Verlag, New York, 1998.

- B. C. Berndt, S. Bhargava, and F. G. Garvan, Ramanujan's theories of elliptic functions to alternative bases, Trans. Amer. Math. Soc. 347 (1995), 4163–4244.
- 12. J. M. Borwein and P. B. Borwein, Pi and the AGM, Wiley, New York, 1987.
- 13. C. Carathéodory, Funktionentheorie. Band II, Birkhäuser Verlag, Basel, 1950.
- I. N. Cangül and D. Singerman, Normal subgroups of Hecke groups and regular maps, Math. Proc. Camb. Phil. Soc. 123 (1998), 59–74.
- 15. H. M. Farkas and I. Kra, *Theta Constants, Riemann Surfaces and the Modular Group*, Graduate Studies in Mathematics, vol. 37, American Mathematical Society, 2001.
- 16. O. Forster, Lectures on Riemann Surfaces, Springer-Verlag, New York, 1981.
- 17. J. I. Hutchinson, On a class of automorphic functions, Trans. Amer. Math. Soc. 3 (1902), 1–11.
- T. Jørgensen, A. Marden, and C. Pommerenke, *Two examples of covering surfaces*, Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978), Ann. of Math. Stud., vol. 97, Princeton Univ. Press, Princeton, N.J., 1981, pp. 305–317.
- 19. G. A. Jones and J. Wolfart, Dessins d'Enfants on Riemann Surfaces, Springer, New York, 2016.
- 20. S. Katok, Fuchsian Groups, The University of Chicago Press, Chicago and London, 1992.
- 21. O. Lehto and K. I. Virtanen, Quasiconformal Mappings in the Plane, 2nd Ed., Springer-Verlag, 1973.
- 22. S. Ramanujan, Notebooks (2 volumes), Tata Institute of Fundamental Research, Bombay, 1957.
- 23. S. Ramanujan, The lost notebook and other unpublished papers, Narosa, New Delhi, 1988.
- 24. R. Roy, *Elliptic and Modular Functions from Gauss to Dedekind to Hecke*, Cambridge University Press, 2017.

GRADUATE SCHOOL OF INFORMATION SCIENCES, TOHOKU UNIVERSITY, AOBA-KU, SENDAI 980-8579, JAPAN – AND – DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BARISHAL, BARISHAL-8254, BANGLADESH

Email address: shafiul@dc.tohoku.ac.jp, msalam@bu.ac.bd

GRADUATE SCHOOL OF INFORMATION SCIENCES, TOHOKU UNIVERSITY, AOBA-KU, SENDAI 980-8579, JAPAN

Email address: sugawa@math.is.tohoku.ac.jp