# ON THE MATROIDAL PATH IDEALS

#### MEHRDAD NASERNEJAD, KAZEM KHASHYARMANESH, AYESHA ASLOOB QURESHI

ABSTRACT. We prove that the set of all paths of a fixed length in a complete multipartite graph is the bases of a matroid. Moreover, we discuss the Cohen-Macaulayness and depth of powers of *t*-path ideals of a complete multipartite graph.

#### INTRODUCTION

Matroid theory is one of the most attractive areas in combinatorics and is deeply rooted in graph theory and linear algebra. The well-known examples of matroids include uniform matroids, graphical matroids and linear matroids. We refer to [23], [20] and [14] for basic definitions and notions in matroid theory. The polymatroids originated in [9] and the *discrete polymatroids* appeared in [14] as a multiset analogue of matroids. In commutative algebra, the matroidal ideals as the squarefree version of polymatroidal ideals, hold a very special place due to their nice algebraic and homological properties. A monomial ideal  $I \subset K[x_1, \ldots, x_n]$  is called *polymatroidal*, if the set of exponent vectors of the minimal generating set of I corresponds to the set of bases of a discrete polymatroid. In particular, a squarefree polymatroidal ideal is simply referred to as a *matroidal* ideal, because in this case, the set of exponent vectors of the minimal generating set of I corresponds to the set of bases of a matroid. The algebraic and homological properties of polymatroidal ideals have been studied by many authors, for example, see [17], [16], [12], [2], [15]. It is known from [7] and [17] that the product of polymatroidal ideals is again polymatroidal. Moreover, polymatroidal ideals have linear quotients, and hence linear resolutions, see [7]. In particular, all powers of a polymatroidal ideal have linear resolutions. If Iis a polymatroidal ideal, then the Rees algebra  $\mathcal{R}(I)$  is normal, see [22, Proposition 3.11], [16, Theorem 3.4]. Furthermore, polymatroidal ideals are known to have the persistence property, that is,  $\operatorname{Ass}(I^k) \subset \operatorname{Ass}(I^{k+1})$  for all k, see [16, Proposition 3.3], and they even have strong persistence property, that is,  $I^{k+1}: I = I^k$  for all k, see [15, Proposition 2.4].

In this paper, we introduce another class of matroidal ideals, namely, the class of the *t*-path ideals of complete multipartite graphs. Let G be a finite simple graph with n vertices. We refer to a path of length t - 1 in G as a *t*-path. Let S be a polynomial ring over a field K in n variables. To simplify the notation, throughout

<sup>2020</sup> Mathematics Subject Classification. Primary 05B35, 05E40; Secondary 13C13.

Key words and phrases. Veronese ideals, depth of powers of ideals, matroids, polymatroidal ideals, complete multipartite graphs.

The third author was funded by the project 118F321 under the program 2509 of the Scientific and Technological Research Council of Turkey(TUBITAK)..

this text, we identify the vertices of G with the variables in S. The *t*-path ideal  $I_t(G) \subset S = K[x_1, \ldots, x_n]$  is generated by those monomials  $x_{i_1} \cdots x_{i_t}$  such that  $x_{i_1}, \ldots, x_{i_t}$  is a *t*-path in G. The *t*-path ideals of graphs were introduced by Conca and De Negri in [8] and later on discussed by several authors for different classes of graphs, in particular for directed trees, trees and cycles, see [10], [3], [5]. When t = 2, then  $I_2(G)$  is the *edge ideal* of G which is usually denoted as I(G). Let  $K_{n_1,\ldots,n_r}$  be the complete *r*-partite graph with r partition sets of sizes  $n_1,\ldots,n_r$ . The main result of this paper is

# **Theorem 2.5.** Let $t \ge 2$ . If $I_t(K_{n_1,\dots,n_r}) \ne 0$ , then it is a matroidal ideal.

In other words, let  $t \geq 2$  and  $\mathcal{M}_t(G)$  be a collection of subsets of V(G) such that  $A \in \mathcal{M}_t(G)$  if and only if there exists a *t*-path in *G* with vertex set *A*. Then it follows from Theorem 2.5 that if  $\mathcal{M}_t(K_{n_1,\dots,n_r}) \neq \emptyset$ , then it is the set of bases of a matroid. Up to our knowledge there is not much known about *t*-path ideals. In our case, after establishing Theorem 2.5, the *t*-path ideals of a complete *r*-partite graph inherit all the nice properties of matroidal ideals.

The contents of this paper are distributed as following: in Section 1, the definitions and basic facts about t-paths and t-path ideals of complete r-partite graphs are provided. The Section 2 is devoted to establish the matroidal property of  $I_t(K_{n_1,\ldots,n_r})$ in Theorem 2.5. It follows from [6, Theorem 3.2] and [1, Theorem 2.2] that an edge ideal I(G) is matroidal if and only if G is a complete r-partite graph. In Theorem 2.3, we give another proof of this fact. It is natural to ask that if the t-path ideal of a graph is matroidal for some t, then does it follow that its non-zero k-path ideal is also matroidal for  $k \ge t$ ? In Example 2.6, we provide a graph whose 3-path ideal is matroidal but 4-path ideal is not matroidal.

The identification of  $I_t(K_{n_1,\ldots,n_r})$  as a matroidal ideal, helps tremendously to study its algebraic properties. In Section 3, the Cohen-Macaulay property of  $I_t(K_{n_1,\ldots,n_r})$ is discussed. It is known from [12, Theorem 4.2] that a matroidal ideal is Cohen-Macaulay if and only if it is a principal ideal or a squarefree Veronese ideal. It remains to investigate for which t and under which conditions on  $n_1, \ldots, n_r$ , the ideal  $I_t(K_{n_1,\ldots,n_r})$  is squarefree Veronese. As a main result of Section 3, we prove the following:

**Theorem 3.3.** The *t*-path ideal  $I_t(K_{n_1,\ldots,n_r}) \neq 0$  is Cohen-Macaulay if and only if  $n_i \leq \lfloor t/2 \rfloor$ , for each  $i = 1, \ldots, r$ .

The stability indices related to associated primes and depth of powers of polymatroidal ideals are of particular interest and have been discussed in several papers, for example, see [16, 15, 18]. The limit depth and index of depth stability is computed for *t*-path ideals of complete bipartite graphs in Theorem 4.3, and for 3-path ideals of complete *r*-partite graphs in Theorem 4.7. All of these concepts are defined and discussed in Section 4. From Theorem 4.3 and Theorem 4.7, it is evident that there is no uniform description for the depth of the powers of  $I_t(K_{n_1,\dots,n_r})$ . However, in Theorem 4.8, it is observed that if  $r \ge 3$  and all of the  $n_i$ 's are bigger than or equal to  $\lfloor t/2 \rfloor$ , then the limit depth of  $I_t(K_{n_1,\dots,n_r})$  is 0.

## 1. Preliminaries

First we recall some basis definition and notion related to matroids. Let  $[n] = \{1, \ldots, n\}$  and P([n]) denote the set of all subsets of [n]. For any  $S \subseteq [n]$ , the cardinality of S is denoted by |S|. A matroid  $\mathcal{M}$  on the ground set [n] is a non-empty collection of subsets of [n] satisfying the following properties:

- (1) if  $A \in \mathcal{M}$  and  $B \subset A$ , then  $B \in \mathcal{M}$ ;
- (2) if  $A, B \in \mathcal{M}$  and  $|B| \leq |A|$ , then there exists  $a \in A \setminus B$  such that  $B \cup \{a\} \in \mathcal{M}$ .

A maximal element (with respect to inclusion) in  $\mathcal{M}$  is called a *base*. Condition (2) can be used to show that all bases of  $\mathcal{M}$  have the same cardinalities. Let  $\mathcal{B}$  be the set containing all the bases of  $\mathcal{M}$ , then  $\mathcal{B}$  is distinguished by the following "exchange property".

(EP) For any  $A, B \in \mathcal{B}$ , if  $a \in A \setminus B$ , then there exists  $b \in B \setminus A$  such that  $(A \setminus \{a\}) \cup \{b\} \in \mathcal{B}$ .

Given any subset C of P([n]), there exists a matroid on [n] with C as its sets of bases if and only if C satisfies the exchange property.

Let K be a field and  $S = K[x_1, \ldots, x_n]$  be a polynomial ring over K. Let I be a monomial ideal in S. The set  $\mathcal{G}(I)$  denotes the unique minimal set of monomial generators of I. For a given  $\mathbf{a} = (a_1, \ldots, a_n)$  with non-negative entries,  $\mathbf{x}^{\mathbf{a}}$  denotes the monomial  $x_1^{a_1} \cdots x_n^{a_n}$  in S. A monomial ideal  $I \subset S$  generated in a single degree is called *polymatroidal* if for all monomials  $\mathbf{x}^{\mathbf{a}}, \mathbf{x}^{\mathbf{b}} \in \mathcal{G}(I)$  with  $a_i > b_i$ , there exists j with  $a_j < b_j$  such that  $x_j(\mathbf{x}^{\mathbf{a}}/x_i) \in \mathcal{G}(I)$ . A squarefree polymatroidal ideal is simply referred to as a *matroidal* ideal. In other words, a monomial ideal I in S is matroidal, if  $\mathcal{G}(I)$  can be identified as a set of bases of a matroid.

Next we recall some basic definitions and notions from graph theory. All graphs considered in this paper will be simple, undirected and finite. Let G be a graph with the vertex set V(G) and the edge set E(G). A path in G is a sequence of distinct vertices  $x_{i_1}, \ldots, x_{i_t}$  such that  $\{x_{i_j}, x_{i_{j+1}}\} \in E(G)$  for  $j = 1, \ldots, t - 1$ . The length of a path is the number of edges in it. We will refer to a path of length t - 1 as a t-path. In other words, a t-path in G is a path with t vertices. A subset  $A \subseteq V(G)$ is called *independent* if no vertices in A are adjacent in G.

A graph G is called *complete r-partite*, if V(G) can be partitioned into r independent sets  $V_1, \ldots, V_r$  such that a and b are adjacent for all  $a \in V_i$  and all  $b \in V_j$  with  $1 \leq i \neq j \leq r$ . Such a partition is called an r-partition of G. Moreover, if  $|V_i| = n_i$  for all  $i = 1, \ldots, r$ , then the complete r-partite graph is denoted by  $K_{n_1,\ldots,n_r}$ .

Let G be a graph with |V(G)| = n. As mentioned in the introduction, we identify the vertices of the graph G with the variables in the polynomial ring  $S = K[x_1, \ldots, x_n]$ .

**Definition 1.1.** The *t*-path ideal of G, denoted by  $I_t(G)$ , is

$$I_t(G) := (x_{i_1} \cdots x_{i_t} : x_{i_1}, \dots, x_{i_t} \text{ is a } t\text{-path in } G) \subset S.$$

When t = 2, then  $I_2(G)$  is simply the *edge ideal* I(G) of G. If there is no t-path in G, then we set  $I_t(G) = 0$ .

The following example illustrates the definitions that have been stated above.

**Example 1.2.** Let t = 4 and  $K_{1,2,3}$  be as shown in Figure 1.2 with partition sets  $\{x_1\}, \{x_2, x_3\}$  and  $\{x_4, x_5, x_6\}$ . Then  $I_4(K_{1,2,3})$  has the following generators:




FIGURE 1. The graph  $K_{1,2,3}$ 

 $x_1x_2x_3x_4, x_1x_2x_3x_5, x_1x_2x_3x_6, x_1x_2x_4x_5, x_1x_2x_4x_6, x_1x_2x_5x_6, x_1x_5x_7, x_1x_2x_5x_7, x_1x_7, x_1x_$ 

 $x_1x_3x_4x_5, x_1x_3x_4x_6, x_1x_3x_5x_6, x_2x_3x_4x_5, x_2x_3x_4x_6, x_2x_3x_5x_6.$ 

Recall that if  $u = x_1^{a_1} \cdots x_n^{a_n}$  is a monomial in  $S = K[x_1, \ldots, x_n]$ , then the support of u is given by  $\operatorname{supp}(u) = \{x_i : a_i > 0\}$ . We set  $\deg_{x_i} u = a_i$ , for all  $i = 1, \ldots, n$ . For a monomial ideal  $I \subset S$ , the support of I is  $\operatorname{supp}(I) = \bigcup_{u \in \mathcal{G}(I)} \operatorname{supp}(u)$ . If  $\operatorname{supp}(I) = \{x_1, \ldots, x_n\}$ , then I is said to be fully supported. If  $\mathcal{G}(I) = \{u_1, \ldots, u_m\}$ , then  $\operatorname{gcd}(I) = \operatorname{gcd}(u_1, \ldots, u_m)$ .

Below we give a list of remarks that will be used frequently throughout the paper. For any integer a, the notations  $\lceil a \rceil$  and  $\lfloor a \rfloor$  denote the ceiling and floor functions of a, respectively. Let  $G = K_{n_1,\ldots,n_r}$  and  $V_1,\ldots,V_r$  be the r-partition of G. Then we have the following.

**Remarks 1.3.** Let  $P : x_{i_1}, x_{i_2}, \ldots, x_{i_t}$  be a *t*-path in G, and  $u = x_{i_1}x_{i_2}\cdots x_{i_t} \in \mathcal{G}(I_t(G))$ . If  $x_{i_j} \in V_s$  for some  $1 \leq s \leq r$  and  $2 \leq j \leq t - 1$ , then  $x_{i_{j-1}}, x_{i_{j+1}} \notin V_s$ . This leads to the following conclusions:

(i) A *t*-path in G can have at most  $\lfloor t/2 \rfloor$  vertices in  $V_s$ .

- (ii) Let  $x_{i_j} \in V_s$  for some  $1 \leq j \leq t$ . If there exists  $x_k \in V_s$  such that  $x_k \notin V(P)$ , then  $x_{i_j}$  can be replaced by  $x_k$  in P to obtain a new t-path P' in G. In particular,  $(u/x_{i_j})x_k \in \mathcal{G}(I_t(G))$ .
- (iii) If there exists some  $V_s$  such that  $V(P) \cap V_s = \emptyset$ , then any  $x_{i_j} \in V(P)$  can be replaced in P by any  $x_k \in V_s$ , to obtain a new path P' in G. In particular,  $(u/x_{i_j})x_k \in \mathcal{G}(I_t(G))$ , for all  $x_k \in V_s$  and for each  $j = 1, \ldots, t$ .
- (iv) If  $I_t(G) \neq 0$ , then  $I_t(G)$  is fully supported. This can be easily seen due to statements (ii) and (iii).
- (v) Let  $I_t(G) \neq 0$ . Then  $I_t(G)$  is a principal ideal if and only if  $t = n_1 + \cdots + n_r$ . This also follows from statements (ii) and (iii).

#### 2. The Matroidal path ideals of graphs

It is proved in [2, Theorem 1.1] that a monomial ideal  $I \subset S = K[x_1, \ldots, x_n]$  is polymatroidal if and only if I: u is polymatroidal for all monomials u in S. In particular,  $I: x_i$  is polymatroidal if I is polymatroidal. The following theorem gives further information about  $I: x_i$ .

**Theorem 2.1.** Let  $I \subset S = K[x_1, \ldots, x_n]$  be a polymatroidal ideal and  $1 \leq i \leq n$ . Let  $I = \sum_{j=0}^{d} I_j x_i^j$ , where  $d = \max\{\deg_{x_i} u : u \in \mathcal{G}(I)\}$  and  $I_j = (u/x_i^j : u \in \mathcal{G}(I), \deg_{x_i} u = j)$  for each  $j = 0, \ldots, d$ . Then  $I_0 \subseteq I_1 \subseteq \cdots \subseteq I_d$ .

Proof. If d = 0, then  $I = I_0$  and there is nothing to prove. Let  $d \ge 1$  and set  $J_1 := \sum_{j=1}^d I_j x_i^{j-1}$ . It is easy to see that  $I = I_0 + x_i J_1$  and  $(I : x_i) = I_0 + J_1$ . Following the proof of [2, Theorem 1.1], one can deduce that  $I_0 \subseteq J_1$ , and hence  $(I : x_i) = J_1$  is a polymatroidal ideal. Now, on account of  $I_0 = (u : u \in \mathcal{G}(I), \deg_{x_i} u = 0)$  and  $I_1 = (u/x_i : u \in \mathcal{G}(I), \deg_{x_i} u = 1)$ , we obtain  $I_0 \subseteq I_1$ . If d = 1, then we are done. Otherwise, if  $d \ge 2$ , we set  $J_2 := \sum_{j=2}^d I_j x_i^{j-2}$ . Thus,  $J_1 = I_1 + x_i J_2$ . Once again, from the proof of [2, Theorem 1.1] we obtain that  $I_1 \subseteq J_2$ , and thus  $I_1 \subseteq I_2$ . Therefore,  $I_0 \subseteq I_1 \subseteq I_2$ . Now, set  $J_k = \sum_{j=k}^d I_j x_i^{j-k}$ , where  $1 \le k \le d$ . By continuing this procedure, we can conclude that  $I_0 \subseteq I_1 \subseteq \cdots \subseteq I_d$ , as required.  $\Box$ 

If  $I \subset K[x_1, \ldots, x_n]$  is a matroidal ideal, then  $\max\{\deg_{x_i} u : u \in \mathcal{G}(I)\}$  is either 0 or 1. For each variable  $x_i$ , we set  $I_{0,i} = (u : x_i \notin \operatorname{supp}(u))$  and  $I_{1,i} = (u/x_i : x_i \in \operatorname{supp}(u))$ . Then, we have  $I = I_{0,i} + x_i I_{1,i}$ , for all  $i = 1, \ldots, n$ . With this notation, we have the following specialization of Theorem 2.1.

**Proposition 2.2.** Let  $I \subset K[x_1, \ldots, x_n]$  be a fully supported matroidal ideal generated in degree d with gcd(I) = 1. We have the following:

- (i) for each i = 1, ..., n we have  $\mathcal{G}(I_{1,i}) \subseteq \bigcup_{t=1, t \neq i}^{n} \mathcal{G}(I_{1,t})$ .
- (ii) if  $I_{1,i} \subseteq I_{1,j}$  for some  $1 \le i, j \le n$  with  $i \ne j$ . Then  $I_{1,i} = I_{1,j}$ .

Proof. (i) The assumption that I is fully supported gives that  $I_{1,i} \neq 0$ , for each i = 1, ..., n. Moreover, gcd(I) = 1 gives that  $I_{0,i} \neq 0$ , for each i = 1, ..., n. Let  $u' \in \mathcal{G}(I_{1,i})$  and  $v \in \mathcal{G}(I_{0,i})$ . Then it follows from the definition of  $I_{0,i}$  and  $I_{1,i}$  that  $v \in \mathcal{G}(I)$  and  $u := u'x_i \in \mathcal{G}(I)$ . Since  $\deg_{x_i} u > \deg_{x_i} v$ , one can conclude from the exchange property that there exists some s with  $\deg_{x_s} u < \deg_{x_s} v$  such that  $u'x_s = x_s(u/x_i) \in \mathcal{G}(I)$ . Then,  $u' \in \mathcal{G}(I_{1,s})$ . We therefore have  $\mathcal{G}(I_{1,i}) \subseteq \bigcup_{t=1, t\neq i}^n \mathcal{G}(I_{1,t})$ , as claimed.

(ii) Suppose, on the contrary, that there exists some monomial  $u \in \mathcal{G}(I_{1,j}) \setminus \mathcal{G}(I_{1,i})$ . Thus,  $x_j u \in \mathcal{G}(I)$ . If  $x_i \in \operatorname{supp}(u)$ , then  $(x_j u)/x_i \in \mathcal{G}(I_{1,i})$ . Using the assumption  $I_{1,i} \subseteq I_{1,j}$ , we obtain  $(x_j u)/x_i \in I_{1,j}$  which gives  $(x_j^2 u)/x_i \in I$ . Since I is matroidal,  $(x_j u)/x_i \in I$ . This yields a contradiction to  $x_j u \in \mathcal{G}(I)$ . Hence,  $x_i \notin \operatorname{supp}(u)$ . Since  $x_j u \in \mathcal{G}(I)$ , we conclude that  $x_j u \in \mathcal{G}(I_{0,i})$ . From Theorem 2.1, we know that  $I_{0,i} \subset I_{1,i}$ . Then there exists some monomial  $w \in \mathcal{G}(I_{1,i})$  such that w divides  $x_j u$ . Again, by using the assumption,  $I_{1,i} \subseteq I_{1,j}$ , we have  $x_j \notin \operatorname{supp}(w)$  and therefore, w|u. Since both w and u are monomials of degree d-1, we conclude that u = w, as required. It follows from [6, Theorem 3.2] that if I(G) is a matroidal ideal, then G is a complete multipartite graph. The converse follows as a corollary of [1, Theorem 2.2]. We give another straightforward proof of this fact in the language of edge ideals of complete multipartite graphs.

# **Theorem 2.3.** Let G be a graph without isolated vertices. The edge ideal I(G) is a matroidal ideal if and only if G is a complete r-partite graph for some $r \ge 2$ .

Proof. We first assume that G is a complete r-partite graph, that is,  $G = K_{n_1,..,n_r}$ with vertex partition  $V_1, \ldots, V_r$  and  $|V_i| = n_i$  for all  $i = 1, \ldots, r$ . Let  $u := x_k x_l \in I(G)$  and  $v := x_i x_j \in I(G)$  such that  $x_k$  does not divide v. We need to show that either  $x_i(u/x_k) = x_i x_l \in I(G)$  or  $x_j(u/x_k) = x_j x_l \in I(G)$ . Let  $x_l \in V_t$  for some  $1 \le t \le r$ . The graph G is complete r-partite and  $\{x_i, x_j\} \in E(G)$ , hence it follows that at least one of the vertices between  $x_i$  and  $x_j$  does not belong to  $V_t$ . If  $x_i \notin V_t$ , then  $x_i x_l \in I(G)$  and similarly if  $x_j \notin V_t$ , then  $x_j x_l \in I(G)$ , as required.

Conversely, let I(G) be a matroidal ideal and |V(G)| = n. We can assume that G does not contain any isolated vertices, and hence I(G) is fully supported. If I(G) is a principal ideal, then there is nothing to prove and the assertion holds trivially. Let I(G) be not a principal ideal. If  $gcd(I(G)) = x_i$ , for some variable  $x_i$ , then G is a complete bipartite graph with r-partition  $V_1 = \{x_i\}$  and  $V_2 = V(G) \setminus \{x_i\}$ , and again the assertion holds in this case.

Finally, we assume that gcd(I) = 1. Note that for each i = 1, ..., n, we have  $I(G)_{1,i} = (x_q : x_q \in N_G(x_i))$ , where  $N_G(x_i)$  denotes the neighbourhood of  $x_i$  in G. Then from Proposition 2.2(ii), we conclude that if  $I(G)_{1,i} \subseteq I(G)_{1,j}$ , for some  $i \neq j$ then  $I(G)_{1,i} = I(G)_{1,j}$ ; equivalently, if  $N_G(x_i) \subseteq N_G(x_j)$ , then  $N_G(x_i) = N_G(x_j)$ . We can partition vertices of G into disjoint sets, say  $V_1, \ldots, V_r$  such that for all  $k = 1, \ldots, r$ , the vertices in  $V_k$  have same neighbourhood. Since G is a simple graph, it is clear that each of the  $V_k$  is an independent set of G. We claim that  $G = K_{n_1,\ldots,n_r}$  with r-partition  $V_1, \ldots, V_r$ . To prove our claim, it only remains to verify the following: if  $x_i \in V_k$  and  $x_j \in V_\ell$  with  $k \neq \ell$ , then  $\{x_i, x_j\} \in E(G)$ . By virtue of Proposition 2.2(ii), we have  $N_G(x_i) \not\subseteq N_G(x_j)$  and  $N_G(x_j) \not\subseteq N_G(x_i)$ . Let  $x_k \in N_G(x_i) \setminus N_G(x_j)$  and  $x_\ell \in N_G(x_j) \setminus N_G(x_i)$ . Then  $\{x_i, x_k\}$  and  $\{x_j, x_\ell\}$  are disjoint edges in G. Since I(G) is matroidal, then by applying exchange property on  $x_i x_k$  and  $x_j x_\ell$ , we conclude that  $\{x_i, x_j\} \in E(G)$ , as claimed. This completes the proof.

To prove the main result of this section, we first need the following:

**Lemma 2.4.** Let  $G = K_{n_1,\ldots,n_r}$  and  $t \geq 3$ . Furthermore, let  $P_1 : x_{i_1},\ldots,x_{i_{t-1}}$  be a (t-1)-path in G and  $P_2 : x_{j_1},\ldots,x_{j_t}$  be a t-path in G. Then there exists  $x_{j_k} \in V(P_2) \setminus V(P_1)$  such that one of the following statements is satisfied.

- (1)  $x_{j_k}, x_{i_1}, \ldots, x_{i_{t-1}}$  is a t-path in G.
- (2)  $x_{i_1}, \ldots, x_{i_{t-1}}, x_{j_k}$  is a *t*-path in *G*.
- (3)  $t \ge 4$  and there exists  $2 \le p \le t-2$  such that  $x_{i_1}, \ldots, x_{i_p}, x_{j_k}, x_{i_{p+1}}, \ldots, x_{i_{t-1}}$ is a t-path in G.

*Proof.* Let  $V_1, \ldots, V_r$  be the *r*-partition of  $G = K_{n_1,\ldots,n_r}$ . We may assume that  $x_{i_1} \in V_1$ . If there exists  $x_{j_k} \in V(P_2) \setminus V(P_1)$  such that  $x_{j_k} \notin V_1$ , then  $\{x_{j_k}, x_{i_1}\} \in E(G)$ , and  $P : x_{j_k}, x_{i_1}, \ldots, x_{i_{t-1}}$  is a *t*-path in *G*, as given in (1).

If (1) is not true, then every  $x_{j_k} \in V(P_2) \setminus V(P_1)$  is such that  $x_{j_k} \in V_1$ . If  $x_{i_{t-1}} \notin V_1$ , then for any  $x_{j_k} \in V(P_2) \setminus V(P_1)$ , we have  $\{x_{i_{t-1}}, x_{j_k}\} \in E(G)$ , and hence we obtain a *t*-path  $P : x_{i_1}, \ldots, x_{i_{t-1}}, x_{j_k}$  in *G*, as given in (2). Note that if t = 3, then in  $P_1 : x_{i_1}, x_{i_2}$ , both vertices cannot be in  $G_1$  because *G* is a complete *r*-partite graphs. Hence either (1) or (2) must be true for t = 3.

If both (1) and (2) do not hold, then  $t \ge 4$ . In this case, we try to construct a path as given in (3). The negation of (1) and (2) gives us that  $x_{i_1}, x_{i_{t-1}} \in V_1$ and every  $x_{j_k} \in V(P_2) \setminus V(P_1)$  is such that  $x_{j_k} \in V_1$ . If there exists some  $2 \le p \le t-2$  such that  $x_{i_p}, x_{i_{p+1}} \notin V_1$ , then for any  $x_{j_k} \in V(P_2) \setminus V(P_1)$ , we have  $\{x_{i_p}, x_{j_k}\}, \{x_{j_k}, x_{i_{p+1}}\} \in E(G)$ , and  $P: x_{i_2}, \ldots, x_{i_p}, x_{j_k}, x_{i_{p+1}}, \ldots, x_{i_t}$  is a *t*-path in *G* as given in (3). Finally, suppose that each of (1), (2), and (3) do not hold. Then we are in the following situation:

- (i)  $x_{i_1}, x_{i_{t-1}} \in V_1;$
- (ii) every  $x_{j_k} \in V(P_2) \setminus V(P_1)$  is such that  $x_{j_k} \in V_1$ . This also gives that for every  $x_{j_k} \in V(P_2) \setminus V_1$ , we have  $x_{j_k} \in V(P_1)$ ;
- (iii) for every  $2 \le p \le t 2$  either  $x_{i_p} \in V_1$  or  $x_{i_{p+1}} \in V_1$ .

Since  $x_{i_1} \in V_1$  and  $\{x_{i_1}, x_{i_2}\} \in E(G)$ , this implies that  $x_{i_2} \notin V_1$ . It follows now from (iii) that  $x_{i_3} \in V_1$ . By continuing this process, we obtain from (iii) that  $x_{i_{t-2}} \notin V_1$  since  $x_{i_{t-1}} \in V_1$  and  $\{x_{i_{t-2}}, x_{i_{t-1}}\} \in E(G)$ . This shows that  $P_1$  starts and ends at vertices in  $V_1$ , and all vertices in  $P_1$  with odd indices are also in  $V_1$ . Furthermore, all vertices in  $V(P_1)$  with even indices do not belong to  $V_1$ . Hence, t-1must be an odd integer, and in  $V(P_1)$ , there are t/2-1 vertices that do not belong to  $V_1$ , that is,  $|V(P_1) \setminus V_1| = t/2 - 1$ . Moreover, it follows from (ii) that every vertex in  $V(P_2)$  which is outside of  $V_1$  must belong to  $V(P_1)$ , that is,  $V(P_2) \setminus V_1 \subseteq V(P_1) \setminus V_1$ . Since  $P_2$  is a t-path with even number of vertices in a complete r-partite graph G, this gives that at least half of its vertices are not in  $V_1$ , that is,  $|V(P_2) \setminus V_1| \ge t/2$ . This contradicts  $|V(P_2) \setminus V_1| \le |V(P_1) \setminus V_1| = t/2 - 1$ . This yields that at least one of the statements in (1), (2), or (3) must hold.

Now we state the main result of this section.

# **Theorem 2.5.** Let $t \ge 2$ . If $I_t(K_{n_1,\ldots,n_r}) \ne 0$ , then it is a matroidal ideal.

Proof. We set  $G := K_{n_1,\ldots,n_r}$  and let  $V_1,\ldots,V_r$  be the *r*-partition of *G*. The assertion is true for  $I_2(G)$ , as shown in Theorem 2.3. Let  $t \ge 3$  and  $I_t(G) \ne 0$ . If  $I_t(G)$  is a principal ideal, then there is nothing to prove and the assertion holds trivially. Let  $|\mathcal{G}(I_t(G))| \ge 2$ . To prove that  $I_t(G)$  is matroidal, we need to show that if  $P_1 : x_{i_1},\ldots,x_{i_t}$  and  $P_2 : x_{j_1},\ldots,x_{j_t}$  are two *t*-paths with  $x_{i_l} \notin V(P_2)$  for some  $1 \le l \le t$ , then there exists some  $x_{j_k} \in V(P_2) \setminus V(P_1)$ , such that one obtains a new path by removing  $x_{i_l}$  from  $P_1$  and inserting  $x_{j_k}$  in an appropriate position in  $P_1$ .

We may assume that  $x_{i_l} \in V_1$ . If  $x_{i_1}, \ldots, x_{i_{l-1}}, x_{i_{l+1}}, \ldots, x_{i_t}$  is (t-1)-path in G then we obtain the desired conclusion by using Lemma 2.4. In particular, if l = 1 or l = t, then removing  $x_{i_l}$  from  $P_1$  gives a (t-1)-path in G. Now assume

that  $x_{i_1}, \ldots, x_{i_{l-1}}, x_{i_{l+1}} \ldots, x_{i_t}$  is not a (t-1)-path in G. This is possible only if  $\{x_{i_{l-1}}, x_{i_{l+1}}\} \notin E(G)$ , that is,  $x_{i_{l-1}}, x_{i_{l+1}}$  belong to  $V_a$  for some  $2 \le a \le r$ . If there exists some  $x_{j_k} \in V(P_2) \setminus V(P_1)$  such that  $x_{j_k} \notin V_a$ , then

$$x_{i_1}, \ldots, x_{i_{l-1}}, x_{j_k}, x_{i_{l+1}}, \ldots, x_{i_t}$$

is a *t*-path in *G*, and the proof is complete. Otherwise, every  $x_{j_k} \in V(P_2) \setminus V(P_1)$  is such that  $x_{j_k} \in V_a$ . If  $x_{i_1} \notin V_a$ , then  $\{x_{i_1}, x_{i_{l+1}}\} \in E(G)$ , and we get  $x_{i_{l-1}}, x_{i_{l-2}}, \ldots, x_{i_1}, x_{i_{l+1}}, \ldots, x_{i_t}$  as a (t-1)-path in *G*. We again use Lemma 2.4 to obtain the desired conclusion. A similar argument gives us the desired conclusion when  $x_{i_t} \notin V_a$ . Otherwise, we are in the following situation:

- (1)  $x_{i_{l-1}}, x_{i_{l+1}}, x_{i_1}, x_{i_t} \in V_a;$
- (2) every  $x_{j_k} \in V(P_2) \setminus V(P_1)$  is such that  $x_{j_k} \in V_a$ . This implies that if  $x_{j_k} \in V(P_2) \setminus V_a$ , then  $x_{j_k} \in V(P_1)$ .

We claim that there must exist some p with either  $2 \le p \le l-3$  or  $l+2 \le p \le t-2$ such that  $x_{i_p}, x_{i_{p+1}} \notin V_a$ . Indeed, if our claim is true and we can find such p with  $2 \le p \le l-3$ , then

$$x_{i_1}, \ldots, x_{i_p}, x_{i_{l-1}}, x_{i_{l-2}}, \ldots, x_{i_{p+1}}, x_{i_{l+1}}, \ldots, x_{i_t}$$

is a (t-1)-path in G, and the desired result can be deduced from Lemma 2.4. Similarly, if we can find such p with  $l+2 \le p \le t-2$ , then

$$x_{i_1}, \ldots x_{i_{l-1}}, x_{i_p}, x_{i_{p-1}}, \ldots, x_{i_{l+1}}, x_{i_{p+1}}, \ldots, x_{i_t}$$

is a (t-1)-path in G, and one can conclude the desired result from Lemma 2.4.

To establish our claim, suppose, on the contrary, that for every p with  $2 \le p \le l-3$ and  $l+2 \le p \le t-2$ , we have either  $x_{i_p} \in V_a$  or  $x_{i_{p+1}} \in V_a$ . Since  $x_{i_1} \in V_a$  and  $\{x_{i_1}, x_{i_2}\} \in E(G)$ , this yields that  $x_{i_2} \notin V_a$ . It follows now from the assumption that  $x_{i_3} \in V_a$ . Moreover, from (1) it follows that  $x_{i_{l-2}} \notin V_a$  since  $x_{i_{l-1}} \in V_a$  and  $\{x_{i_{l-2}}, x_{i_{l-1}}\} \in E(G)$ . This forces l-1 to be an odd integer. Similarly, since  $x_{i_{l+1}}, x_{i_\ell} \in V_a$ , we derive that the number of vertices in the path  $x_{i_{l+1}}, \ldots, x_{i_\ell}$  is odd as well. Collectively, we obtain that t itself must be odd. This shows that  $V(P_1)$  contains (t+1)/2 vertices from  $V_a$  and the other (t-1)/2 vertices are not from  $V_a$ . Note that these (t-1)/2 vertices that are not from  $V_a$  include  $x_{i_l}$  as well. Considering that, by (2), every vertex in  $V(P_2) \setminus V(P_1)$  belongs to  $V_a$  and  $x_{i_l} \notin V(P_2)$ , this implies that in  $V(P_2)$  there are at most (t-1)/2 - 1 vertices that are not from  $V_a$ , which is impossible by Remark 1.3(i). Hence the claim holds.  $\Box$ 

It is known from [16, Proposition 3.11] that polymatroidal ideals are normal. In [21, Corollary 2.11], it is shown that the t-path ideals of complete bipartite graphs are normal. This result can also be seen now as a corollary of the above theorem.

In general, for a given  $t \ge 3$ , one can find graphs that are not complete r-partite but their t-path ideal  $I_t(G)$  is matroidal. As a very simple example, let G itself be a path on t vertices. Then  $I_t(G)$  is matroidal because it is a principal ideal, while G is not complete r-partite.

Theorems 2.3 and 2.5 together show that if  $I_2(G)$  is matroidal then  $I_t(G)$  is also matroidal for all  $t \geq 2$ . Therefore, it is natural to ask the following question: If

 $I_t(G)$  is matroidal for some t, then is it true that  $I_k(G)$  is also matroidal for all  $k \ge t$ ? The following example shows that it is not true in general.

**Example 2.6.** Let G be the graph as shown in the following figure. Then




FIGURE 2. The graph G

$$I_3(G) = (x_1 x_2 x_4, x_1 x_3 x_4, x_1 x_4 x_5, x_1 x_4 x_6, x_2 x_3 x_4, x_2 x_4 x_5, x_2 x_4 x_6)$$

 $x_3x_4x_5, x_3x_4x_6, x_4x_5x_6).$ 

Note that  $I_3(G) = x_4 J$ , where

$$J = (x_1x_2, x_1x_3, x_1x_5, x_1x_6, x_2x_3, x_2x_5, x_2x_6, x_3x_5, x_3x_6, x_5x_6).$$

The ideal J is indeed the edge ideal of the complete graph on the vertex set  $\{x_1, x_2, x_3, x_5, x_6\}$ , and by Theorem 2.3, J is matroidal. This shows that  $I_3(G)$  is also matroidal because it is the product of two matroidal ideals, namely  $(x_4)$  and J. Set  $u := x_1 x_2 x_3 x_4$  and  $v := x_3 x_4 x_5 x_6$ . Then  $u, v \in I_4(G)$ , but both  $x_5(u/x_1)$  and  $x_6(u/x_1)$  do not belong to G. Hence,  $I_4(G)$  is not matroidal.

In the view of Theorem 2.5 and Example 2.6, we close this section with the following questions.

- (1) For a given t > 2, characterize the graphs for which  $I_t(G)$  is matroidal.
- (2) For which graphs it is true that if  $I_t(G)$  is matroidal for some t, then  $I_k(G)$  is also matroidal for all  $k \ge t$ ?

3. Cohen-Macaulay property of  $I_t(K_{n_1,\ldots,n_r})$ 

Now we investigate the Cohen-Macaulay property for  $I_t(K_{n_1,\ldots,n_r})$ . Let I be a monomial ideal in the polynomial ring  $S = K[x_1,\ldots,x_n]$ . The ideal I is called the *Veronese* ideal of degree d in S if I is generated by all monomial of degree d in S. Similarly, one defines the squarefree counterpart of a Veronese ideal. The monomial ideal I is called *squarefree Veronese* ideal of degree d in S if I is generated by all squarefree monomials of degree d in S. It follows from [12, Theorem 4.2] that a polymatroidal ideal I is Cohen-Macaulay if and only if I is

- (i) a principal ideal, or
- (ii) a Veronese ideal, or
- (iii) a squarefree Veronese ideal.

In particular, if I is a matroidal ideal, then it is Cohen-Macaulay if and only if it is a principal ideal or squarefree Veronese ideal. From Remark 1.3 (v) it follows that  $I_t(K_{n_1,\dots,n_r})$  is a principal ideal if and only if  $t = n_1 + \cdots + n_r$ , and hence in this case it can be viewed as the squarefree Veronese ideal of degree t = n in S. Therefore, for a given t, to be able to characterize  $n_1, \ldots, n_r$  such that  $I_t(K_{n_1,\ldots,n_r})$  is Cohen-Macaulay, one only needs to check that when  $I_t(K_{n_1,\ldots,n_r})$  is squarefree Veronese of degree t in S. More precisely, one needs to check that for which  $n_1, \ldots, n_r$  with  $n_1 + \cdots + n_r = n$ , every subset of [n] of size t can be viewed as a path in  $K_{n_1,\ldots,n_r}$ . In the case of edge ideal of  $K_{n_1,\ldots,n_r}$ , the answer immediately follows from [12, Theorem 4.2].

**Proposition 3.1.** Let G be a graph such that I(G) is matroidal. Then I(G) is Cohen-Macaulay if and only if G is a complete graph.

*Proof.* Let |V(G)| = n. From [12, Theorem 4.2], it follows that I(G) is Cohen-Macaulay if and only if I(G) is squarefree Veronese of degree 2 in  $S = K[x_1, \ldots, x_n]$ , that is, for every  $1 \le i \ne j \le n$ , we have  $x_i x_j \in I(G)$ . Moreover,  $x_i x_j \in I(G)$  for every  $1 \le i \ne j \le n$  if and only if G is a complete graph.

By using Theorem 2.3, the above proposition can be rephrased as follows:

**Corollary 3.2.** The edge ideal  $I(K_{n_1,...,n_r})$  is Cohen-Macaulay if and only if  $n_1 = \cdots = n_r = 1$ .

Now we state the main result of this section in which we discuss the general case when  $t \geq 2$ .

**Theorem 3.3.** The t-path ideal  $I_t(K_{n_1,\ldots,n_r}) \neq 0$  is Cohen-Macaulay if and only if  $n_i \leq \lfloor t/2 \rfloor$ , for each  $i = 1, \ldots, r$ .

Proof. Let  $V_1, \ldots, V_r$  be the r-partition of  $G = K_{n_1,\ldots,n_r}$  with  $|V_i| = n_i$ , for each  $i = 1, \ldots, r$ . Let  $n = n_1 + \cdots + n_r$ . From [12, Theorem 4.2] and Remark 1.3 (v), it follows that if  $I_t(G)$  is Cohen-Macaulay, then it is a squarefree Veronese ideal of degree t in  $S = K[x_1, \ldots, x_n]$ , and hence  $x_{i_1} \cdots x_{i_t} \in I_t(G)$ , for every  $1 \le i_1 < \cdots < i_t \le n$ . Moreover, from Remark 1.3(i), it follows that if there exists some  $V_i$  with  $n_i > \lceil t/2 \rceil$ , then for any subset T of  $V_i$  with  $|T| = \lceil t/2 \rceil + 1$ , there does not exist any t-path in G that contains all the elements of T. Hence, if there exists some  $V_i$  of G with  $n_i > \lceil t/2 \rceil$ , then I(G) is not squarefree Veronese, and hence not Cohen-Macaulay.

Now assume that  $n_i \leq \lfloor t/2 \rfloor$ , for each  $i = 1, \ldots, r$ . Let  $u = x_{i_1} \cdots x_{i_t}$  be a squarefree monomial in S. We need to show that the elements in  $A := \operatorname{supp}(u)$  can be interpreted as a t-path in G. Let  $A_i = A \cap V_i$  and  $a_i = |A_i|$ . Since  $n_i \leq \lfloor t/2 \rfloor$ , we get  $a_i \leq \lfloor t/2 \rfloor$ . Without loss of generality, we may assume that the elements in  $\{x_{i_1}, \ldots, x_{i_t}\}$  are arranged such that the first  $a_1$  elements lie in  $V_1$ , the next  $a_2$  elements lie in  $V_2$  and so on. If t is odd, then

$$x_1, x_{\frac{t+1}{2}+1}, x_2, x_{\frac{t+1}{2}+2}, \dots, x_{\frac{t-1}{2}}, x_t, x_{\frac{t+1}{2}}$$

is a t-path in G, since consecutive vertices in above sequence belong to different  $V_i$ 's. Similarly, if t is even, then

$$x_1, x_{\frac{t}{2}+1}, x_2, x_{\frac{t}{2}+2}, \dots, x_{\frac{t}{2}}, x_t$$

is a t-path in G. This completes the proof.

4. LIMIT DEPTH OF  $I_t(K_{n_1,\ldots,n_r})$ 

Let  $(R, \mathfrak{m})$  be a Noetherian local ring or a standard graded algebra over a field Kwith graded maximal ideal  $\mathfrak{m}$ . Let  $I \subset R$  be an ideal, which is graded if R is standard graded K-algebra. It is known from Brodmann [4] that depth $(R/I^k)$  stabilizes for large k, that is, depth $(R/I^k)$  is constant for  $k \gg 0$ . The smallest t > 0, for which depth $(R/I^t) = depth(R/I^k)$  for all  $k \geq t$  is called the *index of depth stability* of Iand is denoted by dstab(I), as defined in [16]. Moreover, depth $(I^{dstab(I)})$  is called the *limit depth* of I and is denoted by  $\lim_{k\to\infty} depth(R/I^k)$ , see [11].

In this section we will study the dstab $(S/I_t(K_{n_1,\ldots,n_r}))$  and  $\lim_{k\to\infty} \operatorname{depth}(S/I_t(K_{n_1,\ldots,n_r})^k)$ . In [15], the linear relation graphs of monomial ideals were introduced to help in understanding their analytic spreads. Let  $I \subset S$  be a monomial ideal with  $\mathcal{G}(I) = \{u_1,\ldots,u_m\}$ . The *linear relation graph*  $\Gamma$  of I is the graph with the edge set

$$E(\Gamma) = \{\{x_i, x_j\}: \text{ there exists } u_k, u_l \in \mathcal{G}(I) \text{ such that } x_i u_k = x_j u_l\},\$$

and  $V(\Gamma) = \bigcup_{\{x_i, x_j\} \in E(\Gamma)} \{x_i, x_j\}$ . It is known from [15, Lemma 4.2 and Theorem 4.1] that if the linear relation graph of a matroidal ideal I has m vertices and s connected components, then

(1) 
$$\operatorname{dstab}(I) < \ell(I) = m - s + 1$$

where  $\ell(I)$  denotes the analytic spread of I, that is, the Krull dimension of the fiber ring  $\mathcal{R}(I)/\mathfrak{m}\mathcal{R}(I)$ . Combining this with [16, Corollary 3.5], one can deduce that

(2) 
$$\lim_{k \to \infty} \operatorname{depth}(S/I^k) = n - (m - s + 1).$$

Therefore, to compute the limit depth of t-path ideals of complete r-partite graphs, it is enough to compute the number of vertices and the number of connected components in their linear relation graphs.

**Remark 4.1.** In [11, Proposition 2.1], it is shown that for any graded ideal I, the depth $(S/I^k)$  is a non-increasing function of k if all powers of I have a linear resolution. This is indeed the case for matroidal ideals. Therefore, one concludes the following: if limit depth of a matroidal ideal is s and the k is the smallest integer for which depth $(S/I^k) = s$ , then dstab(I) = k.

**Remark 4.2.** It follows from [7, Theorem 2.5] that if I is a fully supported matroidal ideal generated in degree d, then depth(I) = d - 1. It is shown in Remark 1.3 (iv) that if  $I_t(K_{n_1,\ldots,n_r}) \neq 0$ , then it is a fully supported ideal. This shows that whenever  $I_t(K_{n_1,\ldots,n_r}) \neq 0$ , we have depth $(I_t(K_{n_1,\ldots,n_r})) = t - 1$ .

At first, we will compute limit depth of *t*-path ideals of complete bipartite graphs.

**Theorem 4.3.** Let  $G = K_{p,q}$  with n = p + q and  $t \ge 2$ . Then we have the following:

- (i) if  $p = \lfloor t/2 \rfloor$  and  $q = \lceil t/2 \rceil$ , then  $\lim_{k \to \infty} \operatorname{depth}(S/I_t(G)^k) = n 1$  and  $\operatorname{dstab}(I_t(G)) = 1$ .
- (ii) if  $p = \lfloor t/2 \rfloor$  and  $q > \lceil t/2 \rceil$ , then  $\lim_{k \to \infty} \operatorname{depth}(S/I_t(G)^k) = \lfloor t/2 \rfloor$  and  $\operatorname{dstab}(I_t(G)) = \lceil (q-1)/(q-\lceil t/2 \rceil) \rceil$ .

(iii) if  $p > \lfloor t/2 \rfloor$  and  $q > \lceil t/2 \rceil$ , then  $\lim_{k\to\infty} \operatorname{depth}(S/I_t(G)^k) = 0$  and  $1 < \operatorname{dstab}(I_t(G)) < n$  if t is odd, and  $\lim_{k\to\infty} \operatorname{depth}(S/I_t(G)^k) = 1$  and  $1 \leq \operatorname{dstab}(I_t(G)) < n-1$ , if t is even.

*Proof.* Let  $V_1 = \{x_1, \ldots, x_p\}$  and  $V_2 = \{y_1, \ldots, y_q\}$  be the 2-partition of G.

(i) In this case n = p + q = t and  $I_t(G)$  is a principal ideal and the assertion follows trivially.

(ii) Is  $p = \lfloor t/2 \rfloor$  and  $q > \lceil t/2 \rceil$ , then it is evident from Remark 1.3(i) that all vertices of  $V_1$  appear in every t-path in G. Then, it follows from the definition of  $\Gamma$  that  $V_1 \cap V(\Gamma) = \emptyset$ . It follows immediately from Remark 1.3(ii) that  $\{y_i, y_j\} \in E(\Gamma)$  for all  $1 \leq i \neq j \leq q$ . Therefore  $V(\Gamma) = V_2$  and  $\Gamma$  is a complete graph on q vertices, and has only one connected component. We conclude  $\ell(I) = q$  and use the equality in (2) to compute  $\lim_{k\to\infty} \operatorname{depth}(S/I_t(G)^k) = n - q = p = \lfloor t/2 \rfloor$ .

To prove the assertion about dstab $(I_t(G))$ , we first observe that  $I_t(G) = JL$ , where  $J = (x_1 \cdots x_p)$  and L is generated by all monomials of degree  $\lceil t/2 \rceil$  in qvariables. Indeed, as discussed before, all vertices of  $V_1$  appear in every t-path in G, equivalently,  $x_1 \cdots x_p$  divides every generator of  $I_t(G)$ . Moreover, for every subset  $\{y_{i_1}, \ldots, y_{i_m}\}$  of  $V_2$  of size  $m = \lceil t/2 \rceil$ , there is a t-path P in G with V(P) = $\{y_{i_1}, \ldots, y_{i_m}\} \cup V_1$ , equivalently,  $y_{i_1} \cdots y_{i_m} \in L$ . Hence, J is a principal ideal and L is the squarefree Veronese ideal of degree  $\lceil t/2 \rceil$  in variables  $y_1, \ldots, y_q$ . Let  $S_1 =$  $K[x_1, \ldots, x_p]$  and  $S_2 = K[y_1, \ldots, y_q]$ . Then depth $(S_1/J^k) = p - 1$  and it follows from [16, Corollary 5.7] that depth $(S_2/L^k) = \max\{0, k(\lceil t/2 \rceil - q) + q - 1\}$  for all kand  $\lim_{k\to\infty} depth(S_2/L^k) = 0$  with dstab $(L) = \lceil (q-1)/(q-\lceil t/2 \rceil)\rceil$ .

The fact that the ideals J and L are generated in different sets of variables facilitates to conclude that  $(I_t(G))^k = J^k L^k$ , and depth $(S/I_t(G)^k) = depth(S_1/J^k) + depth(S_2/L^k) + 1$ , for all k, for example, see [19, Lemma 2.2]. Therefore,

$$depth(S/I_t(G)^k) = p + \max\{0, k(\lceil t/2 \rceil - q) + q - 1\}.$$

Since  $\lim_{k\to\infty} \operatorname{depth}(S/I_t(G)^k) = p$  and  $\operatorname{dstab}(L) = \lceil (q-1)/(q-\lceil t/2\rceil) \rceil$ , it follows that  $\operatorname{dstab}(I_t(G)) = \lceil (q-1)/(q-\lceil t/2\rceil) \rceil$ .

(iii) Let  $p > \lceil t/2 \rceil$  and  $q > \lfloor t/2 \rfloor$ . Then it follows from Remark 1.3(ii), that  $\{x_i, x_j\} \in E(\Gamma)$ , for all  $1 \leq i \neq j \leq p$  and  $\{y_i, y_j\} \in E(\Gamma)$ , for all  $1 \leq i \neq j \leq q$ . If t is even, then every path in G contains half variables from  $V_1$  and the other half from  $V_2$ . Consequently, given any t-path in G, we cannot replace any vertex from  $V_1$  with any vertex from  $V_2$ . This shows that  $\{x_i, y_j\} \notin E(\Gamma)$ . Therefore,  $\Gamma$  has exactly two connected components, and  $\ell(I) = p + q - 2 + 1 = n - 1$ . By using equality in (2), we obtain  $\lim_{k\to\infty} \operatorname{depth}(S/I_t(G)^k) = n - (n-1) = 1$  and from (1), we obtain  $\operatorname{dstab}(I_t(G)) < n - 1$ . Remark 4.2 gives that  $\operatorname{depth}(I_t(G)) = t - 1$ . Hence  $\operatorname{dstab}(I_t(G) \geq 1$  and the equality holds only if t = 2 due to Remark 4.1.

If t is odd, then set  $m = \lfloor t/2 \rfloor$ . The sequence  $x_1, y_1, x_2, y_2, \ldots, x_m, y_m$  is a (t-1)-path in G. It can be extended to a t-path by either joining the edge  $\{y_{m+1}, x_1\}$ , or the edge  $\{y_m, x_{m+1}\}$ . Hence, for  $u = y_{m+1}x_1y_1x_2y_2\ldots x_my_m$  and  $v = x_1y_1x_2y_2\ldots x_my_mx_{m+1}$ , we have  $x_{m+1}u = y_{m+1}v$ , which gives  $\{x_{m+1}, y_{m+1}\} \in E(\Gamma)$ . This shows that  $\Gamma$  is connected. In fact, by repeating the same argument as above, one can show that  $\Gamma$  is a complete graph on n vertices. Therefore,  $\ell(I) = n-1+1 = n$ 

and by using the equality in (2), we obtain  $\lim_{k\to\infty} \operatorname{depth}(S/I_t(G)^k) = 0$  and from (1), we obtain  $\operatorname{dstab}(I_t(G)) < n$ . Again from Remark 4.2 we have  $\operatorname{depth}(I_t(G)) = t - 1 > 0$ . Hence  $\operatorname{dstab}(I_t(G) > 1$ . This finishes the proof.

Now we consider the case in which G is an r-partite graph with  $r \geq 3$ . Let  $d, a_1, \ldots, a_n$  be positive integers with  $\sum_{i=1}^n a_i \geq d$ . Then the ideal generated by all monomials  $x_1^{b_1} \cdots x_1^{b_n}$  of degree d with  $b_i \leq a_i$ , for all  $i = 1, \ldots, n$  is called the ideal of *Veronese type*, and is denoted by  $I_{d;a_1,\ldots,a_n}$ . In [16], the index of depth stability of Veronese type ideals is discussed in detail. Note that if  $a_1 = \cdots = a_n = 1$ , then  $I_{d;a_1,\ldots,a_n}$  is squarefree Veronese ideal in n variables. From Theorem 3.3 and results obtained in [16], we conclude the following

**Proposition 4.4.** Let  $t \ge 2$  and  $G = K_{n_1,\dots,n_r}$  with |V(G)| = n and  $n_i \le \lceil t/2 \rceil$ , for all  $1 \le i \le r$ . Then  $\lim_{k\to\infty} \operatorname{depth}(S/I_t(G)^k) = 0$  and  $\operatorname{dstab}(I_t(G)) = \lceil (n-1)/(n-t) \rceil$ .

In the view of Proposition 4.4, it is enough to consider those cases in which at least one of the  $n_i$ 's is greater than  $\lceil t/2 \rceil$ . From Theorem 4.3, it is clear that limit depth of  $I_t(G)$  depends on t and how vertices are distributed among the partition sets. We see this behaviour in subsequent results too. Now we will describe limit depth and dstab for  $I_3(K_{n_1,\ldots,n_r})$  when  $r \ge 3$ . Note that since  $r \ge 3$ , we must have  $3 \le n = n_1 + \cdots + n_r$ . First we discuss the cases when  $n \in \{3, 4\}$ . The following lemma is a consequence of Proposition 4.4.

**Lemma 4.5.** Let  $G = K_{n_1,\ldots,n_r}$  with  $r \ge 3$ . Then we have the following:

- (i) If |V(G)| = 3, then  $I_3(G)$  is a principal ideal and  $\lim_{k\to\infty} \operatorname{depth}(S/I_3(G)^k) = 2$  and  $\operatorname{dstab}(I_3(G)) = 1$ .
- (ii) If |V(G)| = 4, then G is either a complete graph on four vertices or isomorphic to  $K_{2,1,1}$ . Moreover,  $\lim_{k\to\infty} \operatorname{depth}(S/I_3(G)^k) = 0$  and  $\operatorname{dstab}(I_3(G)) = 3$ .

In the following text,  $\mathfrak{m}$  denotes the unique graded maximal ideal in the polynomial ring S whose variables correspond to the vertices of the graph G.

**Proposition 4.6.** Let  $G = K_{n_1,\dots,n_r}$  with  $r \ge 3$ . Then  $\lim_{k\to\infty} \operatorname{depth}(I(G)^k) = 0$ and  $\operatorname{dstab}(I(G)) = 2$ .

Proof. From Remark 4.2, we know that depth(I(G)) = 1. Then by following Remark 4.1, it is enough to show that depth $(I(G)^2) = 0$ . Given a graph H, in [13, Theorem 2.1], the equivalent condition for depth $(I(H)^2) = 0$  is described as follows: depth $(I(H)^2) = 0$  if and only if H contains a cycle of length 3 and every other vertex of G has a neighbour in this cycle. It can be easily seen that G satisfies this condition. Let  $V_1, \ldots, V_r$  be the r-partition of G and since  $r \geq 3$ , we can choose  $x_1 \in V_1, x_2 \in V_2$ , and  $x_3 \in V_3$ . Then the subgraph induced by  $x_1, x_2$  and  $x_3$  is cycle of length 3 in G. Moreover, it follows from the definition of G that any  $x_j \in V(G)$  is adjacent to at least one of the vertices in  $\{x_1, x_2, x_3\}$ . Hence depth $(I(G)^2) = 0$ , as required.

**Theorem 4.7.** Let  $G = K_{n_1,\ldots,n_r}$  with  $r \ge 3$  and  $|V(G)| \ge 5$ . Then  $\lim_{k\to\infty} \operatorname{depth}(I_3(G)^k) = 0$  and  $\operatorname{dstab}(I_3(G)) = 2$ .

*Proof.* From Remark 4.2, we know that depth $(I_3(G)) = 2$ , and then by following Remark 4.1, it is enough to show that depth $(I_3(G)^2) = 0$ . Let  $V_1, \ldots, V_r$  be the *r*-partition of *G*.

First, assume that  $|V_1| = \cdots = |V_{r-1}| = 1$ . If  $|V_r| = 1$  or  $|V_r| = 2$ , then the assertion follows from Proposition 4.4. More precisely, If  $|V_r| = 1$ , then |V(G)| = r, and if  $|V_r| = 2$ , then |V(G)| = r + 1. Since  $r \ge 5$ , in both cases, we have  $dstab(I_3(G)) = \lceil (s-1)/(s-3) \rceil = 2$  for s = r, r+1. Finally, if  $|V_r| > 2$ , then take  $x_1 \in V_1, x_2 \in V_2$  and  $x_3, x_4, x_5 \in V_r$ , and set  $u := x_1 x_2 x_3 x_4 x_5$ . Then for any  $x_i \in V_r$ , we have  $ux_i = (x_i x_2 x_3)(x_4 x_1 x_5) \in I_3(G)^2$  because  $x_i, x_2, x_3$  and  $x_4, x_1, x_5$  are 3-paths in G. Furthermore, for any  $x_i \in V(G) \setminus V_r$ , we have  $ux_i = (x_1 x_3 x_2)(x_4 x_i x_5) \in I_3(G)^2$  because  $x_1, x_3, x_2$  and  $x_4, x_i, x_5$  are 3-paths in G. Therefore,  $I_3(G)^2 : u = \mathfrak{m}$ , and hence depth $(I_3(G)^2) = 0$ , as required.

Now, assume that  $|V_i|, |V_j| \ge 2$ , for some  $1 \le i \ne j \le r$ . Take  $x_1, x_2 \in V_i, x_3, x_4 \in V_j$ , and  $x_5 \in V_k$ . Set  $u := x_1 x_2 x_3 x_4 x_5$ ; then  $u \notin I_3(G)^2$ . Then  $I_3(G)^2 : u = \mathfrak{m}$ . Indeed, for any  $x_m \in V_i$ , we have  $ux_m \in I_3(G)^2$  because  $x_1, x_5, x_2$  and  $x_3, x_m, x_4$  are 3-paths in G. Similarly, for any  $x_m \in V_j$ , we have  $ux_m \in I_3(G)^2$ . Finally, if  $x_m \in V(G) \setminus (V_i \cup V_j)$ , then again  $ux_m \in I_3(G)^2$  because  $x_1, x_m, x_2$  and  $x_3, x_5, x_4$  are 3-paths in G. This shows that depth $(I_3(G)^2) = 0$ , as claimed.

Now we give the final result of this section which shows that in the case of  $r \ge 3$  and  $t \ge 3$ , if the number of vertices in each partition set is big enough, then limit depth of  $I_t(G)$  is 0.

**Theorem 4.8.** Let  $G = K_{n_1,\dots,n_r}$  with  $r \ge 3$  and  $n_i \ge \lceil t/2 \rceil$  for all  $1 \le i \le r$ . Then  $\lim_{k\to\infty} \operatorname{depth}(I_t(G)^k) = 0$  and  $1 < \operatorname{dstab}(I_t(G)) < n$  for  $t \ge 3$ .

Proof. From Remark 4.2, we conclude that depth $(I_t(G)) = t - 1 > 0$ . Let |V(G)| = n. Due to (1) and (2), it is enough to prove that the linear relation graph  $\Gamma$  of  $I_t(G)$  has one connected component and  $|V(\Gamma)| = n$ . To prove this, we will show that  $\Gamma$  is a complete graph with  $V(\Gamma) = V(G)$ . Let  $V_1, \ldots, V_r$  be the *r*-partition of *G*. Let  $x_i, x_j \in V(G)$ , for some  $i \neq j$ . We can choose  $A \subset V(G)$  such that  $|A| = t, x_i \in A$ ,  $x_j \notin A$  and  $|A \cap V_k| < \lceil t/2 \rceil$ , for all  $k = 1, \ldots, r$ . Such a choice of *A* is possible because  $n_i \geq \lceil t/2 \rceil$  and  $r \geq 3$ . Take  $B = (A \setminus \{x_i\}) \cup \{x_j\}$ . Then |B| = t and  $|B \cap V_k| \leq \lceil t/2 \rceil$ , for all  $k = 1, \ldots, r$ . Then by following the similar construction of the path at the end of the proof of Theorem 3.3, we obtain a *t*-path *P* in *G* with vertices in *A* and a *t*-path *P'* in *G* with vertices in *B*. Let *u* be the monomial in  $\mathcal{G}(I_t(G))$  that corresponds to the *t*-path *P* and *v* be the monomial in  $\mathcal{G}(I_t(G))$  that corresponds to the *t*-path *P'*. Then  $x_iv = x_ju$  and  $\{x_i, x_j\} \in E(\Gamma)$ , as required.

#### Acknowledgement

The authors are grateful to the referee for careful reading of the paper and the valuable comments and suggestions.

### References

- R. R. Asghar, S. Yassemi, On the weakly polymatroidal property of the edge ideals of hypergraphs, Comm. in Algebra, 42 (2014), 1011–1021.
- S. Bandari, J. Herzog, Monomial localizations and polymatroidal ideals, European J. Combin. 34 (2013), 752–763.
- [3] R. Bouchat, H. T. Hà, A. O. Keefe, Path ideals of rooted trees and their grad Betti numbers, J. Combin. Theory Ser. A, 118 (8) (2011), 2411–2425.
- [4] M. Brodmann, The asymptotic nature of the analytic spread, Math. Proc. Cambridge Philos. Soc. 86 (1979), 35–39.
- [5] D. Campos, R. Gunderson, S. Morey, C. Paulsen, T. Polstra, Depths and Cohen-Macaulay properties of path ideals, J. Pure Appl. Algebra, 218 (8) (2014), 1537–1543.
- [6] H. Chiang-Hsieh, Some arithmetic properties of matroidal ideals, Comm. in Algebra, 38 (2010), 944–952.
- [7] A. Conca, J. Herzog, Castelnuovo-Mumford regularity of products of ideals, Collect. Math. 54(2), (2003), 137–152.
- [8] A. Conca, E. D. Negri, *M-sequences, graph ideals, and ladder ideals of linear type*, J. Algebra, 211(2)(1999), 599–624.
- [9] J. Edmonds, Submodular functions, matroids, and certain polyhedra. In: R. Guy, H. Hanani, N. Sauer, J. Schonheim, (eds.) Combinatorial Structures and Their Applications. Gordon and Breach, New York (1970).
- [10] J. He, A. V. Tuyl, Algebraic Properties of the Path Ideal of a Tree, Comm. in Algebra, 38(5) (2010), 1725-1742.
- [11] J. Herzog, T. Hibi, The depth of powers of an ideal, J. Alg. **291** (2005), 534–550.
- [12] J. Herzog, T. Hibi, Cohen-Macaulay polymatroidal ideals, European J. Combin. 27(4) (2006), 513–517.
- [13] J. Herzog, T. Hibi, Bounding the socles of powers of squarefree monomial ideals, In: Commutative algebra and noncommutative algebraic geometry. II, (2015), 223–229.
- [14] J. Herzog, T. Hibi, Discrete polymatroids, J. Algebr. Comb., 16 (2002), 239–268.
- [15] J. Herzog, A. A. Qureshi, Persistence and stability properties of power of ideals, J. Pure Appl. Algebra, 219 (3) (2015), 530–542.
- [16] J. Herzog, A. Rauf, M. Vladoiu, The stable set of associated prime ideals of a polymatroidal ideal, J. Algebr. Comb. 37 (2) (2013), 289–312,
- [17] J. Herzog, Y. Takayama, Resolutions by mapping cones, The Roos Festschrift Vol. 2. Homology Homotopy Appl. 4 (2002), 277–294.
- [18] J. Herzog, M. Vladoiu, Squarefree monomial ideals with constant depth function, J. Pure Appl. Algebra, 217 (2013), 1764–1772.
- [19] L. T. Hoa, N. D. Tam, On some invariants of a mixed product of ideals, Arch. Math. 94(4) (2010), 327–337.
- [20] J. G. Oxley, Matroid Theory, Oxford University Press, Oxford, New York, (1992).
- [21] G. Restuccia, R. H. Villarreal, On the normality of monomial ideals of mixed products, Comm. in Algebra, 29(8) (2001), 3571–3580,
- [22] R. H. Villarreal, Rees cones and monomial rings of matroids, Linear Algebra Appl., 428 (2008), 2933–2940.
- [23] D. J. A. Welsh, *Matroid Theory*, Academic Press, London, (1976).

KAZEM KHASHYARMANESH, MEHRDAD NASERNEJAD, DEPARTMENT OF PURE MATHEMATICS, FERDOWSI UNIVERSITY OF MASHHAD, P.O.BOX 1159-91775, MASHHAD, IRAN

Email address: m\_nasernejad@yahoo.com, khashyar@ipm.ir

AYESHA ASLOOB QURESHI, SABANCI UNIVERSITY, FACULTY OF ENGINEERING AND NATURAL SCIENCES, ORTA MAHALLE, TUZLA 34956, ISTANBUL, TURKEY

Email address: aqureshi@sabanciuniv.edu