# UPPER LEVEL SETS OF LELONG NUMBERS ON $\mathbb{P}^2$ AND CUBIC CURVES

ALİ ULAŞ ÖZGÜR KİŞİSEL AND OZCAN YAZICI

ABSTRACT. Let T be a positive closed current of bidimension (1, 1) with unit mass on  $\mathbb{P}^2$  and  $V_{\alpha}(T)$  be the upper level sets of Lelong numbers  $\nu(T, x)$  of T. For any  $\alpha \geq \frac{1}{3}$ , we show that  $|V_{\alpha}(T) \setminus C| \leq 2$  for some cubic curve C.

**2010** Mathematics Subject Classification: 32U05, 32U25, 32U35, 32U40. Keywords: Positive closed currents, Lelong numbers, Cubic curves.

### 1. INTRODUCTION

Suppose that T is a positive closed current of bidimension (1,1) on the complex projective plane  $\mathbb{P}^2$  such that

$$||T|| = \int_{\mathbb{P}^2} T \wedge \omega = 1$$

where  $\omega$  is the Fubini-Study form on  $\mathbb{P}^2$ . The Lelong number of T at a point  $x \in \mathbb{P}^2$  is defined by

$$\nu(T,x) = \lim_{r \to 0} \frac{1}{\pi r^2} \int_{B(x,r)} T \wedge \omega$$

where B(x, r) is the Euclidean ball with center x and radius r > 0 (See [4] for equivalent definitions and properties of Lelong numbers).

For  $0 < \alpha < 1$ , upper level sets of Lelong numbers  $\nu(T, .)$  are defined by

$$E_{\alpha}(T) := \{ x \in \mathbb{P}^2 | \nu(T, x) \ge \alpha \}$$
$$V_{\alpha}(T) := \{ x \in \mathbb{P}^2 | \nu(T, x) > \alpha \}.$$

It is a remarkable result of Siu ([7]) that  $E_{\alpha}(T)$  is an analytic subvariety of  $\mathbb{P}^2$  of dimension at most 1 for any  $\alpha > 0$ . It follows from Chow's theorem that  $E_{\alpha}(T)$  is an algebraic variety.

In [1], Coman proved that if  $\alpha \geq \frac{1}{2}$ ,  $|V_{\alpha}(T) \setminus L| \leq 1$  for some complex line L and if  $\alpha \geq \frac{2}{5}$ ,  $|V_{\alpha}(T) \setminus C| \leq 1$  for some conic. In [1, 3, 5, 6], further geometric results are obtained for the sets  $V_{\alpha}(T)$  where T is a positive closed current of bidimension (p, p) in  $\mathbb{P}^n$  and  $p \geq 1$ . Our goal here is to understand the geometry of level sets  $V_{\alpha}(T)$  for smaller  $\alpha > 0$ . In this direction, we show that for any  $\alpha \geq \frac{1}{3}$ ,  $|V_{\alpha}(T) \setminus C| \leq 2$  for some cubic curve C. More precisely, our main result is the following:

**Theorem 1.1.** If  $\alpha \ge 1/3$  then either

- $|V_{\alpha}(T) \setminus C| \leq 1$  for some (possibly reducible) conic,
- or  $V_{\alpha}(T)$  is a finite set and  $|V_{\alpha}(T) \setminus C| \leq 2$  for some cubic C.

The proof of our main result is based on the construction of entire plurisubharmonic functions of logarithmic growth and with weighted logarithmic poles at some points of  $\mathbb{C}^2$  (See Lemma 3.2). In doing so, we followed the ideas in [1], where such plurisubharmonic functions with smaller number of poles were constructed.

Date: September 21, 2021.

#### **2.** Preliminaries

We say that a plurisubharmonic function  $u \in PSH(\mathbb{C}^n)$  has logarithmic growth if

$$\gamma_u = \lim_{||z|| \to \infty} \sup \frac{u(z)}{\log ||z||} < \infty.$$

A plurisubharmonic function  $u\in PSH(\mathbb{C}^n)$  is said to have logarithmic pole of weight  $\alpha>0$  at  $p\in\mathbb{C}^n$  if

$$u(z) = \alpha \log ||z - p|| + \mathcal{O}(1)$$

in an open neighborhood of *p*.

In the following example, we show that the number 1/3 in the theorem is optimal.

**Example 2.1.** Suppose that  $0 < \alpha < 1/3$ . Consider 6 generic lines  $L_1, \ldots, L_6$  in  $\mathbb{P}^2$  and consider the set S of 15 points of pairwise intersections of these lines. Let

$$T := \frac{1}{6} \sum_{j=1}^{6} [L_j]$$

Then, clearly ||T|| = 1. For each  $p \in S$ , we have  $\nu(T, p) = 1/3$ . Therefore, since  $\alpha < 1/3$ , we must have  $p \in V_{\alpha}(T)$  and consequently  $S \subseteq V_{\alpha}(T)$ . Let us now show that  $V_{\alpha}(T)$  cannot be contained in the union of a cubic and two points: Let  $S = \{x_1, \ldots, x_{15}\}$  and  $\Gamma$  be a cubic containing S except two points, say  $x_1, x_2$ . We may assume that  $L_1$  and  $L_2$  do not contain any of the points  $x_1$  and  $x_2$ . Then Bezout's theorem shows that  $C = L_1 \cup L_2 \cup L$  for some line L. But this implies that the 4 points in  $S \setminus (L_1 \cup L_2 \cup \{x_1, x_2\})$  are collinear, which contradicts to the configuration of points of S.

The following proposition plays an important role in our proof of the main theorem.

**Proposition 2.2** (Proposition 2.1 in [1]). Let  $S = \{x_1, \ldots, x_k\} \subset \mathbb{C}^n$  and T be a positive closed current on  $\mathbb{P}^n$  of bidimension (1, 1) with unit mass. If  $u \in PSH(\mathbb{C}^n)$  has logarithmic growth, it is locally bounded outside a finite set, and  $u(z) \leq \alpha_j \log ||z - x_j|| + O(1)$  for z near  $x_j$  where  $\alpha_j > 0$ ,  $1 \leq j \leq k$ , then

$$\sum_{j}^{k} \alpha_{j} \nu(T, x_{j}) \le \gamma_{u}.$$

In order to construct a plurisubharmonic function u as in Proposition 2.2, we will use the following result of Coman-Nivoche.

**Theorem 2.3** (Theorem 4.1 in [2]). Let r be a positive integer and P,Q be polynomials such that  $S = \{z \in \mathbb{C}^2 : P(z) = Q(z) = 0\}$  and for any  $x \in S$ ,  $ord(P, x) \ge r$ ,  $ord(Q, x) \ge r$  and the intersection number  $(P.Q)_x = r^2$ . Then

$$u = \frac{1}{2r} \log(|P|^2 + |Q|^2)$$

is a plurisubharmonic function in  $\mathbb{C}^2$  with logarithmic growth, which is locally bounded outside of S and has logarithmic pole of weight 1 at the points of S.

## 3. PROOF OF THE MAIN RESULT

Assume from now on that S is a subset of  $\mathbb{P}^2$  such that |S| = 12 where |S| denotes the number of points in S. Say  $S = \{x_1, \ldots, x_{12}\}$ . Let  $\mathcal{X}$  be the subspace of  $\mathbb{C}[X, Y, Z]$  containing all homogeneous polynomials P(X, Y, Z) of degree 6 such that P vanishes on all points of S, and the first partial derivatives of P vanish on  $x_1, \ldots, x_6$ . Each of these vanishings imposes a linear condition on the

coefficients of *P*; the totality of these conditions may or may not be linearly independent. We can then estimate the dimension of  $\mathcal{X}$  from below:

$$dim(\mathcal{X}) \ge \binom{6+2}{2} - (12+2\times 6) = 4$$

Recall that the space of all homogeneous cubic polynomials in  $\mathbb{C}[X, Y, Z]$  has dimension 10, therefore for any given 9 points in  $\mathbb{P}^2$ , there exists a (possibly reducible) cubic curve passing through these points. Through the paper, by abuse of notation, for an algebraic curve  $C \subset \mathbb{P}^2$ , the homogenous defining polynomial will also be denoted by C.

**Lemma 3.1.** Suppose that there exist irreducible cubics  $C_1$  and  $C_2$  such that

- $C_1$  contains  $x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9$
- C<sub>2</sub> contains  $x_1, x_2, x_3, x_4, x_5, x_6, x_{10}, x_{11}, x_{12}$

such that  $C_1$  does not contain  $x_{10}, x_{11}, x_{12}$  and  $C_2$  does not contain  $x_7, x_8, x_9$ . Then,  $\mathcal{X}$  contains two polynomials P, Q such that the common zero set of P and Q is discrete and contains S.

*Proof.* Let  $P_1 = C_1C_2$ . Then, it is clear that  $P_1$  belongs to  $\mathcal{X}$ . Since  $dim(\mathcal{X}) \ge 4$ , we can choose  $P_2, P_3, P_4 \in \mathcal{X}$  such that the set  $\{P_1, P_2, P_3, P_4\}$  is linearly independent. If for some  $i \in \{2, 3, 4\}$ , neither  $C_1$  nor  $C_2$  divide  $P_i$ , then we can take  $P = P_1$  and  $Q = P_i$ : In virtue of the irreducibility of  $C_1$  and  $C_2$ , it is clear that in this case P and Q will have no common components, so their common zero locus will be discrete; it will contain S as well by construction. Also, if both  $C_1$  and  $C_2$  divide  $P_i$  for some  $i \in \{2, 3, 4\}$  then we obtain the contradiction  $P_1 = P_i$ , so from now on we may assume that this does not happen. Suppose now that  $C_1$  divides  $P_i$  and  $C_2$  divides  $P_j$  where  $i \neq j$  and  $i, j \in \{2, 3, 4\}$ . Then  $P_i + P_j \in \mathcal{X}$  and neither  $C_1$  nor  $C_2$  divides  $P_i + P_j$ . Therefore, we can take  $P = P_1, Q = P_i + P_j$  in this case, and we are done.

The remaining case is when one of  $C_1$  or  $C_2$  divides  $P_2$ ,  $P_3$ ,  $P_4$ , and the other doesn't divide any of them. Suppose without loss of generality that  $C_1$  divides  $P_2$ ,  $P_3$  and  $P_4$ . We can write  $P_2 = C_1D_2$ ,  $P_3 = C_1D_3$  and  $P_4 = C_1D_4$  where  $D_2$ ,  $D_3$ ,  $D_4$  are (possibly reducible) cubics. Since  $C_1$  does not contain  $x_{10}$ ,  $x_{11}$ ,  $x_{12}$ , each  $D_i$  must contain these three points. For  $j \in \{1, 2, 3, 4, 5, 6\}$ , if  $x_j$  is a smooth point of  $C_1$ , then in order for  $P_2$ ,  $P_3$  and  $P_4$  to have order 2 at  $x_j$ , the cubics  $D_2$ ,  $D_3$  and  $D_4$ must contain  $x_j$  (this cannot be inferred if  $x_j$  is a singular point of  $C_1$ ). Since  $C_1$  is an irreducible cubic by assumption, it can have at most one singular point. Therefore, without loss of generality we may assume that  $x_2$ ,  $x_3$ ,  $x_4$ ,  $x_5$ ,  $x_6$  are smooth points of  $C_1$ , hence they belong to  $D_2$ ,  $D_3$  and  $D_4$ .

We now have 3 cubics  $D_2, D_3, D_4$ , each of which intersects the irreducible cubic  $C_2$  at a set of points containing the 8 distinct points  $x_2, x_3, x_4, x_5, x_6, x_{10}, x_{11}, x_{12}$ . By Bezout's theorem, the total intersection number of  $D_2$  and  $C_2$  is 9. If there is a 9th intersection point  $x^*$  of  $D_2$  and  $C_2$ distinct from the previous 8 points, then by Cayley-Bacharach theorem,  $D_3$  must also contain  $x^*$ . But then, by Noether's AF+BG theorem,  $D_3$  must lie on the pencil through  $C_2$  and  $D_2$ , namely there exists  $[\alpha : \beta] \in \mathbb{CP}^1$  such that  $D_3 = \alpha C_2 + \beta D_2$ . This would imply that  $P_3 = \alpha P_1 + \beta P_2$ , contradicting the linear independence of  $\{P_1, P_2, P_3, P_4\}$ . The only other possibility is that  $C_2$  and  $D_2$  have intersection number 2 at one of the 8 intersection points, without loss of generality at  $x_2$ . Then we can assume that the intersection numbers  $\mu_{x_2}(C_2, D_3)$  and  $\mu_{x_2}(C_2, D_4)$  are also equal to 2; else a generic member of the pencil through  $D_2$  and one of  $D_3$  or  $D_4$  will intersect  $C_2$  in 9 distinct points, and we can repeat the argument in the beginning of this paragraph by using this generic member instead of  $D_2$ .

To resolve this final case, note that  $\mu_{x_2}(C_2, D_2) = 2$  is the dimension of the  $\mathbb{C}$ -vector space  $\mathcal{O}_{\mathbb{P}^2, x_2}/(C_2, D_2)$  where  $\mathcal{O}_{\mathbb{P}^2, x_2}$  denotes the local ring of the projective plane at  $x_2$ . Denoting by  $m_{\mathbb{P}^2, x_2}$  its maximal ideal, we observe that  $m_{\mathbb{P}^2, x_2}/(C_2, D_2)$  is 1-dimensional. But now, both  $D_3$  and  $D_4$  contain  $x_2$ , hence their representatives in this local ring both belong to  $m_{\mathbb{P}^2, x_2}$ . Hence, their equivalence classes in  $m_{\mathbb{P}^2, x_2}/(C_2, D_2)$  will be linearly dependent. This implies that there exists

 $[\alpha:\beta] \in \mathbb{CP}^1$  such that the equivalence class of  $\alpha D_3 + \beta D_4$  is zero in all  $\mathcal{O}_{\mathbb{P}^2,p}/(C_2, D_2)$ , where p ranges over all intersection points of  $C_2$  and  $D_2$ . By Noether's AF+BG theorem again, we deduce that  $\alpha D_3 + \beta D_4$  lies on the pencil through  $C_2$  and  $D_2$ . This contradicts the linear independence of  $\{P_1, P_2, P_3, P_4\}$  in this final case as well, finishing the proof.

By  $m_j(S)$ , we denote the maximum number of points of S which are contained in an algebraic curve of degree j. For a finite generic subset  $S \subset \mathbb{C}^2$  which contains at least 9 points,  $m_1(S) = 2$ ,  $m_2(S) = 5$  and  $m_3(S) = 9$ .

**Lemma 3.2.** Let  $S \subset \mathbb{C}^2$  such that |S| = 12 and  $m_3(S) = 9$ . Then there exists a  $u \in PSH(\mathbb{C}^2)$  of logarithmic growth which is locally bounded outside a finite set and satisfies one of the following:

- $\gamma_u = 6$  and u has logarithmic poles at the points of S of total weight 18.
- $\gamma_u = 5$  and u has logarithmic poles at the points of S of total weight 15.
- $\gamma_u = 4$  and u has logarithmic poles at the points of S of total weight 12.
- $\gamma_u = 3$  and u has logarithmic poles at the points of S of total weight 9.

*Proof.* We will consider the different cases for the configurations of points of S. Since  $m_3(S) = 9$ , we have  $5 \le m_2(S) \le 7$ . Also note that  $m_1(S) \le m_2(S) - 2$ .

**Case 1.**  $m_2(S) = 5$ . Consider the cubics  $C_1$  and  $C_2$  passing through

$$x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9$$

and

$$x_1, x_2, x_3, x_4, x_5, x_6, x_{10}, x_{11}, x_{12}$$

respectively. Since  $m_2(S) = 5$ , we have  $m_1(S) \leq 3$ . This implies that  $C_1$  and  $C_2$  are irreducible cubics. Since  $m_3(S) = 9$ ,  $C_1$  and  $C_2$  does not contain any other point of S. Then we choose P, Q as in Lemma 3.1. By Theorem 2.3,  $u = \frac{1}{2} \log(|P|^2 + |Q|^2)$  is a plurisubharmonic function which has logarithmic poles at the points of S of total weight 18, locally bounded outside a finite set and  $\gamma_u = 6$ .

**Case 2.**  $m_2(S) = 6, m_1(S) \leq 3$ . Consider the cubics  $C_1$  and  $C_2$  passing through

$$x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9$$

and

$$x_1, x_2, x_3, x_4, x_5, x_6, x_{10}, x_{11}, x_{12}$$

respectively. If  $m_1(S) = 2$  then  $C_1$  and  $C_2$  are irreducible and we return Case 1. Otherwise, we may assume that  $m_1(S) = 3$  and  $C_1$  is reducible. Then  $C_1 = l_1 \cup C$  where  $l_1$  is a line through 3 points of  $\{x_1, \ldots, x_9\}$  and C is a conic (possibly reducible) through the remaining 6 points of  $\{x_1, \ldots, x_9\}$ . Let  $\mathcal{X}$  be the subspace of  $\mathbb{C}[X, Y, Z]$  containing all homogeneous polynomials P(X, Y, Z) of degree 6 such that P vanishes on all points of S, and vanishes to second order at the points  $x_1, \ldots, x_6$ . Then we take  $P_1 = C_1C_2 \in \mathcal{X}$  and by counting the dimension of  $\mathcal{X}$ , we find another  $P_2 \in \mathcal{X}$  such that  $P_1$ and  $P_2$  are linearly independent. If  $P_1$  and  $P_2$  have no common factor then  $u = \frac{1}{2} \log(|P_1|^2 + |P_2|^2)$ is a plurisubharmonic function which has logarithmic poles at the points of S of total weight 18, locally bounded outside a finite set and  $\gamma_u = 6$ .

Otherwise, we first assume that  $P_2$  is divisible by  $l_1$ . Then  $P_2 = l_1\gamma$  where  $\gamma$  is a curve of degree 5 which passes through the 15 points (counting multiplicity) of  $S \setminus l_1$ . If  $\gamma$  and  $CC_2$  have no common factor then  $u = \frac{1}{2} \log(|\gamma|^2 + |CC_2|^2)$  is a plurisubharmonic function which has logarithmic poles at the points of S of total weight 15, locally bounded outside a finite set and  $\gamma_u = 5$ . Otherwise,  $\gamma$  is divisible by C or  $C_2$  or an irreducible factor of C or  $C_2$ . If C divides  $\gamma$  then  $\gamma = C\gamma'$  where  $\gamma'$  is a cubic which shares the same 9 points of S with  $C_2$ . If  $\gamma'$  and  $C_2$  have no common factor then  $u = \frac{1}{2} \log(|\gamma'|^2 + |C_2|^2)$  is a plurisubharmonic function which has logarithmic poles at the points of S of total weight 9, locally bounded outside a finite set and  $\gamma_u = 3$ . Otherwise,  $\gamma'$  and  $C_2$  either

have a degree 1 or degree 2 common factor. If they have a degree 2 common factor, then it must contain 6 points of S, so the remaining lines share the same 3 points, so they coincide, showing that  $\gamma' = C_2$ . If they have a degree 1 common factor, then it must contain 3 points of S, so the remaining conics share the same 6 points; these conics then coincide by applying Bezout's theorem (regardless of whether the conic is irreducible or reducible). In all cases,  $P_1$  and  $P_2$  become linearly dependent, causing a contradiction. A similar argument works if a component of C or  $C_2$  divides  $\gamma$ .

In the case that  $P_2$  is divisible by C or a component of C or  $C_2$  or a component of  $C_2$ , the same discussion follows as above.

**Case 3.**  $m_2(S) = 6$  and  $m_1(S) = 4$ . Let *l* be the unique line containing 4 points for *S*, say  $x_9, x_{10}, x_{11}, x_{12}$ . We consider the cubics  $C_1$  and  $C_2$  passing through

$$x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_9, x_{10},$$

and

$$x_1, x_2, x_3, x_4, x_5, x_6, x_8, x_{11}, x_{12}$$

respectively. If both  $C_1$  and  $C_2$  are irreducible then we return Case 1. Otherwise, we may assume that  $C_1$  is reducible, say  $C_1 = l_1 \cup C$  where  $l_1$  is a line and C is a conic. Since  $m_2(S) = 6$ , the line  $l_1$  must contain at least 3 points out of the 9 points of S that  $C_1$  contains, consequently  $l_1 \neq l$  since otherwise  $l_1$  would also have to contain  $x_{11}$  and  $x_{12}$ , contradicting  $m_1(S) = 4$ . Since l is the unique line containing 4 points,  $l_1$  contains exactly 3 points of  $x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_9, x_{10}$  and C contains the remaining 6 points. Then the result follows from the same argument as in Case 2.

**Case 4.**  $m_2(S) = 7$ . By rearranging the points of *S*, we may assume that there is a conic *C* through  $x_1, \ldots, x_7$ .

First we assume that C is irreducible. Let  $\tilde{C}$  be the conic passing through the points

$$x_8, x_9, x_{10}, x_{11}, x_{12}.$$

Since  $m_3(S) = 9$ ,  $\tilde{C}$  is irreducible, otherwise the line component of  $\tilde{C}$  containing at least 3 points together with C would give us a cubic containing at least 10 points of S. Let  $\mathcal{X}$  be the vector space of homogeneous polynomials of degree 4, passing through the points of S. Then  $P_1 = C\tilde{C} \in \mathcal{X}$ . The dimension of  $\mathcal{X}$  is at least 15 - 12 = 3, therefore we can choose  $P_2 \in \mathcal{X}$  such that  $P_1$  and  $P_2$  are linearly independent. It follows from Bezout's theorem that  $P_1$  and  $P_2$  have no common factor. Thus  $u = \frac{1}{2} \log(|P_1|^2 + |P_2|^2)$  is a plurisubharmonic function which has logarithmic poles at the points of S of total weight 12, locally bounded outside a finite set and  $\gamma_u = 4$ .

Now we consider the case where C is reducible. We note that there cannot be any line containing 5 or more points of S, since otherwise this line together with a conic through 5 other points of S would contradict  $m_3(S) = 9$ . Therefore C must split into two lines at least one of which has 4 points. By rearranging the points, we may assume that l is the line containing 4 points,  $x_1, x_2, x_3, x_4$  of C. Let us list all possible lines which contain 4 points of S. Any such two lines must intersect at a point of S since  $m_2(S) = 7$ . At most two such lines intersect at any point of S since  $m_3(S) = 9$ . Considering these restrictions, by rearranging the points of S, we may assume that there are at most 5 lines, each containing 4 points of S, (see Figure 1). The proof below will be presented as if all 5 lines are actually present, but the argument works without any change if there are less than 5 such lines.

Let  $C_1$  and  $C_2$  be the cubics through the points

 $x_4, x_5, x_6, x_8, x_{11}, x_{12}, x_3, x_7, x_{10}$ 

and

$$x_4, x_5, x_6, x_8, x_{11}, x_{12}, x_1, x_2, x_9$$



FIGURE 1.

respectively. Then  $C_1$  and  $C_2$  do not contain any 4 collinear points of S. If  $C_1$  and  $C_2$  are irreducible then the result follows from Lemma 3.1. Otherwise, we may assume that  $C_1$  is reducible. Let  $C_1 = C \cup l$  where C is a conic which contains 6 or 7 points of S and l is a line. If C contains 7 points of S, then C must be an irreducible conic and the result follows from the above argument. Otherwise, C contains 6 points of S and we conclude the result by a similar argument to the one in Case 2.

Before we prove our main result, we will eliminate some degenerate configurations of points in  $V_{1/3}(T)$ .

**Proposition 3.3.** Say *T* is a positive closed (1, 1) current with unit mass such that the set  $V_{1/3}(T)$  is finite and  $V_{1/3}(T) \setminus C$  contains at least 2 points for every cubic *C*. Then, (a)  $m_1(V_{1/3}(T)) \leq 4$ , (b)  $m_2(V_{1/3}(T)) \leq 7$ , (c) An irreducible cubic can contain at most 9 points of  $V_{1/3}(T)$ .

*Proof.* (a) Assume that  $V_{1/3}(T)$  contains at least 5 collinear points. Say  $x_1, \ldots, x_5$  are contained in L. Then by Siu's decomposition theorem, [7], T = a[L] + R for some current R of bidegree (1,1) having 0 Lelong number generically on L. Using [4], for any  $\epsilon > 0$ , there exists a current R' with ||R'|| = ||R|| = 1 - a,  $\nu(R', x) = \nu(R, x) - \epsilon$  at any  $x \in \mathbb{P}^2$  and R' is smooth generically on L. Thus the intersection  $R' \wedge [L]$  is well-defined. It follows that

$$1 - a = \int R' \wedge [L] \ge \sum_{i=1}^{5} \nu(R' \wedge [L], x_i) > \sum_{i=1}^{5} \nu(R', x_i) \nu([L], x_i) > 5\left(\frac{1}{3} - a - \epsilon\right).$$

The second inequality follows from [4, Corollary 5.10] and the last inequality follows from the fact that  $\nu([L], x_i) \ge 1$ . Since  $\epsilon > 0$  is arbitrary,  $a \ge \frac{1}{6}$ . For any  $x \in V_{1/3}(T) \setminus L$ ,

$$\nu\left(\frac{R}{1-a},x\right) > \frac{2}{5}.$$

By [1], all points (except possibly one point) in  $V_{1/3}(T) \setminus L$  are contained in a conic *C*. Then  $|V_{1/3}(T) \setminus (C \cup L)| \leq 1$ , contradicting the assumption. Hence,  $m_1(V_{1/3}(T)) \leq 4$ .

(b) Assume that  $V_{1/3}(T)$  contains at least 8 points on an irreducible conic *C*. Say  $x_1, \ldots, x_8$  are contained in *C*. Then by Siu's decomposition theorem, [7], T = a[C] + R for some current *R* of bidegree (1,1) having 0 Lelong number generically on *C*. Using [4], for any  $\epsilon > 0$ , there exists a current *R'* with ||R'|| = ||R|| = 1 - 2a,  $\nu(R', x) = \nu(R, x) - \epsilon$  at any  $x \in \mathbb{P}^2$  and *R'* is smooth generically on *C*. It follows that

$$2(1-2a) = \int R' \wedge [C] \ge \sum_{i=1}^{8} \nu(R' \wedge [C], x_i) > \sum_{i=1}^{8} \nu(R', x_i) \nu([C], x_i) > 8\left(\frac{1}{3} - a - \epsilon\right)$$

Since  $\epsilon > 0$  is arbitrary,  $a \ge \frac{1}{6}$ . For any  $x \in V_{1/3}(T) \setminus C$ ,

$$\nu\left(\frac{R}{1-2a},x\right) > \frac{1}{2}$$

By [1], all points (except possibly one point) in  $V_{1/3}(T) \setminus C$  are contained in a line *L*. Hence  $|V_{1/3}(T) \setminus (C \cup L)| \le 1$ , contradicting the assumption.

Similar argument as above works if C is a union of 2 lines, each of which contains 4 points of S. Let  $C = L_1 \cup L_2$  where  $L_1$  and  $L_2$  are the lines containing  $x_1, \ldots, x_4$  and  $x_5, \ldots, x_8$ , respectively. Then  $T = a[L_1] + b[L_2] + R$  for some current R of bidegree (1, 1) having 0 Lelong number generically on C. Using [4], for any  $\epsilon > 0$ , there exists a current R' with ||R'|| = ||R|| = 1 - a - b,  $\nu(R', x) = \nu(R, x) - \epsilon$ at any  $x \in \mathbb{P}^2$  and R' is smooth generically on C. It follows that

$$2(1-a-b) = \int R' \wedge ([L_1] + [L_2]) \ge \sum_{i=1}^{8} \nu(R' \wedge ([L_1] + [L_2]), x_i)$$
  
> 
$$\sum_{i=1}^{4} \nu(R', x_i) \nu([L_1], x_i) + \sum_{i=5}^{8} \nu(R', x_i) \nu([L_2], x_i) > 4\left(\frac{1}{3} - a - \epsilon + \frac{1}{3} - b - \epsilon\right).$$

Since  $\epsilon > 0$  is arbitrary,  $a + b \ge \frac{1}{3}$ . For any  $x \in V_{1/3}(T) \setminus C$ ,

$$\nu\left(\frac{R}{1-a-b},x\right) > \frac{1}{2}.$$

By [1], all points (except possibly one point) in  $V_{1/3}(T) \setminus C$  are contained in a line *L*. Hence  $|V_{1/3}(T) \setminus (C \cup L)| \le 1$ , contradicting the assumption. Hence  $m_2(V_{1/3}(T)) \le 7$ .

(c) Assume that  $V_{1/3}(T)$  contains at least 10 points,  $x_1, \ldots, x_{10}$  on an irreducible cubic C. Then by Siu's decomposition theorem, [7], T = a[C] + R for some current R of bidegree (1, 1) having 0 Lelong number generically on C and  $a \leq \frac{1}{3}$ . Using [4], for any  $\epsilon > 0$ , there exists a current R' with ||R'|| = ||R|| = 1 - 3a,  $\nu(R', x) = \nu(R, x) - \epsilon$  at any  $x \in \mathbb{P}^2$  and R' is smooth generically on C. It follows that

$$3(1-3a) = \int R' \wedge [C] \ge \sum_{i=1}^{10} \nu(R' \wedge [C], x_i) > \sum_{i=1}^{10} \nu(R', x_i) \nu([C], x_i) > 10 \left(\frac{1}{3} - a - \epsilon\right).$$

Since  $\epsilon > 0$  is arbitrary,  $a \ge \frac{1}{3}$  and hence  $T = \frac{1}{3}[C]$ . This implies that  $V_{1/3}(T)$  may contain only the singular point of C and hence we conclude that at most 9 points of  $V_{1/3}(T)$  may lie on an irreducible cubic.

*Proof.* (of Theorem 1.1 ) We first assume that  $E_{\beta}(T)$  contains an algebraic curve C for some  $\beta > \alpha$ . By Siu's decomposition theorem, there exists a positive closed (1,1) current R on  $\mathbb{P}^2$  with 0 Lelong number generically on C such that  $T = \beta[C] + R$ . Then  $\deg(C) \leq \frac{1}{\beta} < 3$ . If C is a conic then  $||R|| = 1 - 2\beta < \frac{1}{3}$ . This implies that  $V_{\alpha}(T) = C$ . If C is a line, then  $||R|| = 1 - \beta < \frac{2}{3}$ . It follows that  $V_{\alpha}(T) = C \cup V_{\alpha}(R)$ . By Theorem 1.1 in [1], there exist a line L and a point p such that

$$V_{\alpha}(R) \subset V_{\frac{1}{2}}\left(\frac{R}{||R||}\right) \subset L \cup \{p\}$$

Then  $|V_{\frac{1}{2}}(T) \setminus (L \cup C)| \leq 1$ .

Now we assume that  $E_{\beta}(T)$  is 0 dimensional for all  $\beta > \alpha$ . Then  $V_{\alpha}(T)$  is at most countable. If  $V_{\alpha}(T)$  is infinite then  $E_{\frac{1}{3}}(T)$  contains an algebraic curve C of degree at most 3 and  $T = \frac{1}{3}[C] + R$ . If C is a cubic then R = 0. This implies that  $\nu(T, p) > \frac{1}{3}$  only at the singular points of C and hence  $V_{\frac{1}{3}}(T)$  is finite. Hence C must be a conic or line. If C is a conic then  $||R|| = \frac{1}{3}$  and  $V_{\alpha}(T) \subset C$ . If C is a line then  $||R|| = \frac{2}{3}$ . Since  $V_{\alpha}(R) \subset V_{\frac{1}{2}}\left(\frac{R}{||R||}\right)$ , Theorem 1.1 in [1] implies that there exists a line L such that  $|V_{\alpha}(R) \setminus L| \leq 1$ . Thus  $|V_{\alpha}(T) \setminus (L \cup C)| \leq 1$ .

Finally we assume that  $V_{\alpha}(T)$  is finite and  $|V_{\alpha}(T) \setminus C| > 2$  for any cubic C. Then  $V_{\alpha}(T)$  has a subset S with |S| = 12.

By Proposition 3.3,  $m_1(S) \le 4$  and  $m_2(S) \le 7$ . We also note that  $m_3(S) \le 11$ : If a cubic contains all the points of S, then it must be reducible by Proposition 3.3. But this contradicts to the fact that  $m_1(S) \le 4$  and  $m_2(S) \le 7$ .

**Case 1.**  $m_3(S) = 9$ . By applying Proposition 2.2 with one of the plurisubharmonic functions u in Lemma 3.2, we obtain that  $\alpha < \frac{1}{3}$  which contradicts to the choice of  $\alpha$ .

Now we move to the cases where  $m_3(S) = 10$ . Let  $\Gamma$  be a cubic containing 10 points of S. By above discussion,  $\Gamma$  must be reducible,  $m_2(S) \leq 7$  and  $m_1(S) \leq 4$ . Thus  $m_2(S) = 6$  or 7. First we assume that  $m_2(S) = 6$ .

**Case 2.**  $m_3(S) = 10$ ,  $m_2(S) = 6$ . Then  $m_1(S) = 4$  and there is a unique line containing 4 points of S. Let L be the line containing the points  $x_1, x_2, x_3, x_4$ ,  $C_1$  and  $C_2$  be the cubics through

$$x_1, x_2, x_5, x_7, x_8, x_9, x_{10}, x_{11}, x_{12}$$

and

$$x_3, x_4, x_6, x_7, x_8, x_9, x_{10}, x_{11}, x_{12}$$

respectively. If  $C_1$  and  $C_2$  are irreducible, then by Proposition 3.3, they can not contain any other point of S. In this case, we choose the polynomials  $P, Q \in \mathcal{X}$  as in Lemma 3.1 and define  $u = \frac{1}{2}\log(|P|^2 + |Q|^2)$ . Then u satisfies the first weight condition in Lemma 3.2 and result follows as in Case 1 above.

Hence we may assume that  $C_1 = l_1 \cup C'$  where  $l_1$  is a line and C' is a conic. Since  $m_3(S) = 10$ ,  $l_1 \neq L$ . Then  $l_1$  contains 3 points of S and C' contains the remaining 6 points of  $C_1$ . We have a similar decomposition for  $C_2$ . Then by a similar argument as in Case 2 of Lemma 3.2, we obtain a plurisubharmonic function u which satisfies one of the weight conditions in Lemma 3.2. It follows from Proposition 2.2 with u that  $\alpha < \frac{1}{3}$  which contradicts to the choice of  $\alpha$ .

**Case 3.**  $m_3(S) = 10$ ,  $m_2(S) = 7$ . Then  $m_1(S) = 3$  or 4. Let C be a conic with 7 points,  $x_1, \ldots, x_7$  and  $\tilde{C}$  be the conic through  $x_8, \ldots, x_{12}$ . First we assume that C is irreducible. If  $\tilde{C}$  is

also irreducible, then we take  $P_1 = CC$  and  $P_2$  a polynomial of degree 4 which passes through the points of S and linearly independent with  $P_1$ . It follows from Bezout's theorem that  $P_1$  and  $P_2$  have no common factor. Then  $u = \frac{1}{2} \log(|P_1|^2 + |P_2|^2)$  is a plurisubharmonic function with total weight 12 at the points of S,  $\gamma_u = 4$  and locally bounded outside S. Proposition 2.2 with u implies that  $\alpha < \frac{1}{3}$  which contradicts to the choice of  $\alpha$ .

Thus, by rearranging the points of S, we may assume that  $\tilde{C}$  is reducible and  $x_8, x_9, x_{12}$  are collinear. We consider the conics  $C_i$  passing through the points  $x_8, x_9, x_{10}, x_{11}, x_i$  for i = 1, ..., 7. If  $C_i$  is irreducible for some i, then we take  $P_1 = CC_i$  and a polynomial  $P_2$  of degree 4 which vanishes at the points of  $S \setminus \{x_{12}\}$  and vanishes to second order at  $x_i$ . Then it follows from Bezout's theorem that  $P_1$  and  $P_2$  have no common factor. Hence  $u = \frac{1}{2} \log(|P_1|^2 + |P_2|^2)$  is a plurisubharmonic function with total weight 12 at the points of  $S \setminus \{x_{12}\}$ ,  $\gamma_u = 4$  and locally bounded outside S. This implies with Proposition 2.2 that  $\alpha < \frac{1}{3}$ .

Now we assume that C is irreducible and  $C_i$  is reducible for all i : 1, ..., 7. By  $L_{i,j}$  we denote the line through  $x_i$  and  $x_j$ . Since  $m_3(S) = 10$  and  $x_8, x_9, x_{12}$  are collinear, the line  $L_{8,9}$  does not contain either of the points  $x_{10}$  or  $x_{11}$ . For the moment, we also assume that  $L_{10,11}$  does not contain  $x_8, x_9, x_{12}$ . Then  $m_1\{x_8, x_9, x_{10}, x_{11}\} = 2$ , each  $x_i$ , i = 1, ..., 7, belongs to  $L_{jk}$  for some  $8 \le j, k \le 11$ .

By (P1), we refer to the property that  $L_{8,9}$  intersects C only at  $x_7$  and  $L_{10,11}$  possibly intersects C only at  $x_7$ . Then we take  $P_1 = CL_{8,9}L_{10,11}$  and  $P_2$  is a polynomial of degree 4 linearly independent with  $P_1$ , which vanishes at the points of  $S \setminus \{x_{12}\}$  and vanishes to second order at  $x_7$ . Then  $P_1$  and  $P_2$  have no common factor. Hence  $u = \frac{1}{2} \log(|P_1|^2 + |P_2|^2)$  is a plurisubharmonic function with total weight 12 at the points of  $S \setminus \{x_{12}\}$ ,  $\gamma_u = 4$ , u is locally bounded outside S and we are done.

By (P2), we refer to the property that  $L_{8,9}$  intersects C only at  $x_7$  and  $L_{10,11}$  intersects C only at  $x_6$ . Then we take  $P_1 = CL_{8,9}L_{10,11}$  and  $P_2$  is a polynomial of degree 4 linearly independent with  $P_1$ , which vanishes at the points of  $S \setminus \{x_{12}\}$  and vanishes to second order at  $x_7$ . Then  $P_1$  and  $P_2$  have no common factor and the result follows as above.

By (P3), we refer to the property that  $L_{8,10}$  intersects C at  $x_6, x_7$  and  $L_{9,11}$  intersects C only at  $x_5$ . Then we take  $P_1 = CL_{8,10}L_{9,11}$  and  $P_2$  is a polynomial of degree 4 linearly independent with  $P_1$ , which vanishes at the points of  $S \setminus \{x_{12}\}$  and vanishes to second order at  $x_5$ . Then  $P_1$  and  $P_2$  have no common factor and the result follows as above.

Note that in (P1), (P2), (P3), we may replace the point  $x_9$  with  $x_{12}$  and the same conclusion holds. Thus we may assume that the configurations (P1), (P2) and (P3) do not hold for the lines  $L_{jk}$  where  $j, k \in \{8, 9, 10, 11\}$  and  $j, k \in \{8, 10, 11, 12\}$ .

Consider the pairs of lines  $(L_{8,9}, L_{10,11}), (L_{8,10}, L_{9,11}), (L_{8,11}, L_{9,10})$ . Then each pair contains at most 3 points of  $\{x_1, \ldots, x_7\} \subset C$  since  $m_2(S) = 7$ . Since  $m_1\{x_8, x_9, x_{10}, x_{11}\} = 2$ , each point  $x_i$ ,  $i = 1, \ldots, 7$ , belongs to a unique pair of lines. If the pair  $(L_{8,9}, L_{10,11})$  contains only one point, say  $x_7$  of  $C \cap S$  then property (P1) holds. Otherwise, since  $m_2(S) = 7$ , we may assume that the pair  $(L_{8,9}, L_{10,11})$  contains 2 points, say  $x_1, x_2$ . If  $x_1$  and  $x_2$  lie on different lines, then property (P2) holds. Thus we may assume that  $x_1, x_2 \in L_{10,11}$ . Note that one of these points may belong to  $L_{8,9}$  as well. One of the remaining pairs of lines, say  $(L_{8,10}, L_{9,11})$  must contain 3 points of  $C \cap S$ . Assuming that (P3) does not hold, we may assume that  $x_6, x_7 \in L_{8,10}$  and  $x_5, x_7 \in L_{9,11}$ . We consider the similar arrangement for the lines  $L_{jk}$  where  $j, k \in \{8, 10, 11, 12\}$ . Then either the pair  $(L_{8,11}, L_{10,12})$  or  $(L_{8,10}, L_{11,12})$  must contain 3 points of  $C \cap S$ .

First we assume that the first pair  $(L_{8,11}, L_{10,12})$  contains 3 points of  $C \cap S$ . As (P3) does not hold, we may assume that  $x_3, x_4 \in L_{8,11}$  and  $x_4, x_5 \in L_{10,12}$ . Then each line  $L_{9,10}$  and  $L_{11,12}$  may contain only one point of  $C \cap S$ , say  $x_3$  and  $x_6$  respectively. But then (P2) holds for the lines  $L_{9,10}$ and  $L_{11,12}$ .



FIGURE 2.

Similarly, we assume now that the other pair  $(L_{8,10}, L_{11,12})$  contains 3 points of  $C \cap S$ . Say  $x_6, x_7 \in L_{8,10}$  and  $x_4, x_6 \in L_{11,12}$ . Since (P1) and (P2) do not hold for the pair  $(L_{8,11}, L_{10,12})$ ,  $x_3 \in L_{10,12}$  and  $x_5 \in L_{10,12} \cap L_{9,11}$ . As  $x_3, x_9, x_{10}, x_{12}$  are not collinear,  $x_3 \notin L_{9,10}$ . Then  $x_4 \in L_{9,10}$  and (P1) or (P2) (if  $x_3 \in L_{8,11}$ ) holds for the pair  $(L_{9,10}, L_{8,11})$ . Hence we are done in the case that C is irreducible and  $L_{10,11}$  does not contain any of the points  $x_8, x_9, x_{12}$ .

Now we assume that the line  $L_{10,11}$  contains  $x_{12}$ . Since  $m_1(S) \leq 4$ , each line  $L_{8,9}$  and  $L_{10,11}$  contains at most 1 point of  $C \cap S$  and hence (P1) or (P2) holds for the pair  $(L_{8,9}, L_{10,11})$ .

Finally, we may assume that there does not exist any irreducible conic containing 7 points of S. Then  $m_1(S) = 4$ . We will list all possible lines containing 4 points of S. Since  $m_2(S) = 7$ , any such two lines must intersect at a point of S. Since |S| = 12, there are at most 3 such lines passing through at any point  $x_i \in S$ .

First we consider the case that there does not exist any point in S through which 3 such lines pass. Then considering the above restrictions, there are at most 5 lines containing 4 points of S (see Figure 2).

We consider the cubics  $C_1$  and  $C_2$  passing through the points of  $S \setminus \{x_2, x_6, x_{10}\}$  and  $S \setminus \{x_5, x_8, x_9\}$ . Then the result follows by a similar argument to the one in Case 4 of Lemma 3.2.

Now we consider the case that there is only one point, say  $x_1$ , through which 3 lines containing 4 points of S pass. We may assume that the points in the sets  $\{x_1, x_2, x_3, x_4\}$ ,  $\{x_1, x_5, x_6, x_7\}$  and  $\{x_1, x_8, x_9, x_{10}\}$  are collinear. Then there are at most 5 lines containing 4 points of S (see Figure 3). We consider the cubics  $C_1$  and  $C_2$  passing through the points of  $S \setminus \{x_1, x_4, x_{11}\}$  and  $S \setminus \{x_2, x_5, x_{10}\}$ . Then the result follows by a similar argument to the one in Case 4 of Lemma 3.2.

Finally, we consider the case that there is more than one point through which 3 lines containing 4 points of S pass. Then, other than  $x_1$ , there is a point of S through which 3 lines containing 4 points of S pass. This point can be a point on a line through  $x_1$  or a point outside of the 3 lines through  $x_1$ .



FIGURE 3.



FIGURE 4.

In the first case, we may take this point as  $x_2$  (see Figure 4). It is easy to check that there are no 4 collinear points in the sets  $S \setminus \{x_2, x_7, x_{10}\}$  and  $S \setminus \{x_1, x_{11}, x_{12}\}$ . Consider the cubics  $C_1$  and  $C_2$  passing through the points of  $S \setminus \{x_2, x_7, x_{10}\}$  and  $S \setminus \{x_1, x_{11}, x_{12}\}$ . Then the result follows by a similar argument to the one in Case 4 of Lemma 3.2.

In the second case, we may assume that 3 lines containing 4 points of S pass through  $x_{11}$  (see Figure 5). If there does not exist any other line containing 4 points of S, then we take the cubics  $C_1$  and  $C_2$  passing through the points of  $S \setminus \{x_4, x_5, x_9\}$  and  $S \setminus \{x_1, x_{11}, x_{12}\}$ . Then the result follows by a similar argument to the one in Case 4 of Lemma 3.2. Otherwise, we may assume that  $x_2, x_6, x_{10}, x_{12}$  are collinear and return to the first case.

**Case 4.**  $m_3(S) = 11$ . In this case,  $m_1(S) = 4$  and  $m_2(S) = 7$ . Let  $\Gamma$  be the cubic with 11 points of S. Then by Proposition 3.3,  $\Gamma$  must be reducible, say  $\Gamma = C \cup L$ . C must be irreducible,



FIGURE 5.

otherwise one of the components of C together with L would contain 8 points of S which contradicts to  $m_2(S) = 7$ . We may assume that the points  $x_1, \ldots, x_7$  are contained in C and  $x_8, \ldots, x_{11}$  are contained in L. Since we assumed that  $|V_{1/3}(T) \setminus \gamma| > 1$  for any cubic  $\gamma$ , there is a point  $p \in V_{1/3}(T) \setminus (\Gamma \cup x_{12})$ . Let  $L_{p,12}$  denote the line which passes through p and  $x_{12}$ . First we assume that  $L_{p,12} \cap L \cap S = \emptyset$ . We take  $P_1 = CL_{p,12}L$  and  $P_2$  a polynomial of degree 4, linearly independent with  $P_1$ , through the points of  $S \cup \{p\}$ . It follows from Bezout's theorem that  $P_1$  and  $P_2$  have no common factor. Thus  $u = \frac{1}{2} \log(|P_1|^2 + |P_2|^2)$  is a plurisubharmonic function with total weight 13 at the points of  $S \cup \{p\}$  locally bounded outside S and  $\gamma_u = 4$ . Proposition 2.2 with u implies that  $\alpha < \frac{4}{13}$  which contradicts to the choice of  $\alpha$ .

Now we consider the case that  $L_{p,12}$  intersects L at a point of S and  $L_{p,12} \cap C \cap S = \emptyset$ . Then the same polynomials  $P_1$  and  $P_2$  above work.

Finally, we consider the case that  $L_{p,12} \cap L \cap S \neq \emptyset$  and  $L_{p,12} \cap C \cap S \neq \emptyset$ . Say  $x_1 \in L_{p,12} \cap C$  and  $x_{11} \in L_{p,12} \cap L$ . Then take  $P_1 = CLL_{p,12}$  and  $P_2$  a polynomial of degree 4, linearly independent with  $P_1$ , which vanishes at the points of  $(S \cup \{p\}) \setminus \{x_{10}, x_{11}\}$  and vanishes to the second order at  $x_1$ . Then  $P_1$  and  $P_2$  have no common factor and the result follows as above.

**Remark 3.4.** We should note that a slightly better conclusion follows from the proof of Theorem 1.1. To get a contradiction in the proof, we assumed that  $V_{1/3}(T)$  contains a subset S with 12 points and  $|V_{1/3}(T) \setminus C| > 1$  for any cubic C. Hence if  $V_{1/3}(T)$  is finite then  $|V_{1/3}(T)| = 11$  or  $|V_{1/3}(T) \setminus C| \le 1$  for some cubic C.

Acknowledgments. We would like to thank to the referee for his/her careful reading and suggestions. The second author is supported by TÜBİTAK 3501 Proj. No 120F084.

### REFERENCES

- [1] Coman, D. Entire Pluricomplex Green Functions and Lelong Numbers of Projective Currents. *Proc. Amer. Math. Soc.* 134 (2005), 1927-1935. 1, 1, 2.2, 3
- [2] Coman, Dan; Nivoche, S., Plurisubharmonic functions with singularities and affine invariants for finite sets in  $\mathbb{C}^n$ . *Math. Ann.* 322 (2002), no. 2, 317-332 2.3
- [3] Coman,D.,Truong,T.T.:Geometric properties of upper level sets of Lelong numbers on projective spaces. *Math. Ann.* 361, 981-994 (2015). 1

- [4] Demailly, J.P., Monge-Ampére operators, Lelong numbers and intersection theory, in *Complex analysis and geometry*, pp. 115-193. Plenum, New York (1993) 1, 3
- [5] Heffers, J.J., A property of upper level sets of Lelong numbers of currents on P<sup>2</sup>. Int. J. Math. 28(14), 1750110, 18 (2017). 1
- [6] Heffers, J.J., On Lelong numbers of positive closed currents on  $\mathbb{P}^n$ . Complex Var. Elliptic Equ. 64, 352-360 (2019). 1
- [7] Siu,Y.T., Analyticity of sets associated to Lelong numbers and the extension of closed positive currents. *Invent. Math.* 27, 53-156 (1974). 1, 3

DEPARTMENT OF MATHEMATICS, MIDDLE EAST TECHNICAL UNIVERSITY, 06800 ANKARA, TURKEY *Email address*: akisisel@metu.edu.tr

DEPARTMENT OF MATHEMATICS, MIDDLE EAST TECHNICAL UNIVERSITY, 06800 ANKARA, TURKEY *Email address:* oyazici@metu.edu.tr