PRINCIPAL SPECIALIZATION OF DUAL CHARACTERS OF FLAGGED WEYL MODULES

KAROLA MÉSZÁROS, AVERY ST. DIZIER, AND ARTHUR TANJAYA

ABSTRACT. Schur polynomials are special cases of Schubert polynomials, which in turn are special cases of dual characters of flagged Weyl modules. The principal specialization of Schur and Schubert polynomials has a long history, with Macdonald famously expressing the principal specialization of any Schubert polynomial in terms of reduced words. We study the principal specialization of dual characters of flagged Weyl modules. Our result yields an alternative proof of a conjecture of Stanley about the principal specialization of Schubert polynomials, originally proved by Weigandt.

1. INTRODUCTION

Schubert polynomials \mathfrak{S}_w were introduced by Lascoux and Schützenberger in [14] as distinguished polynomial representatives for the cohomology classes of Schubert cycles in the flag variety. Schubert polynomials generalize Schur polynomials, a classical basis of the ring of symmetric functions.

The principal specialization of Schur polynomials has a long history: $s_{\lambda}(1,\ldots,1)$ counts the number of semistandard Young tableaux of shape λ , a number famously enumerated by the hook-content formula, see for instance [21]. Macdonald [16, Eq. 6.11] famously expressed the principal specialization $\mathfrak{S}_w(1,\ldots,1)$ of the Schubert polynomial \mathfrak{S}_w in terms of the reduced words of the permutation w. Fomin and Kirillov [9] placed this expression in the context of plane partitions for dominant permutations, while after two decades Billey et al. [2] provided a combinatorial proof. Principal specialization of Schubert polynomials has inspired a flurry of recent interest [10, 18, 20, 22, 23]. A major catalyst for the current line of study into $\mathfrak{S}_w(1,\ldots,1)$ is the following result of Weigandt, which generalizes a conjecture of Stanley ([22, Conjecture 4.1]).

Theorem 1.1 ([23, Theorem 1.1]). For any permutation $w \in S_n$,

$$\mathfrak{S}_w(1,\ldots,1) \ge 1 + p_{132}(w),$$

where $p_{132}(w)$ is the number of 132-patterns in w.

Weigandt's proof of Theorem 1.1 works by exploiting the structure of pipe dreams, one of the earliest combinatorial models for Schubert polynomials [1, 8]. We give an alternative proof of Theorem 1.1 by generalizing its statement to the setting of dual characters of flagged Weyl modules of diagrams:

Theorem 1.2. For any diagram D, the dual character χ_D of the flagged Weyl module of D satisfies

$$\chi_D(1,\ldots,1) \ge \operatorname{rank}(D) + 1.$$

We show in Corollary 3.4 that Theorem 1.2 specializes to Theorem 1.1. Additionally, Theorem 1.2 implies an analogous result for key polynomials (Corollary 3.5).

Outline of this paper. In Section 2 we define dual characters of flagged Weyl modules of diagrams, and we provide necessary background. In Section 3, we define the rank of a diagram and prove Theorem 1.2. We characterize the case of equality in Theorem 1.2 and connect to zero-one polynomials. We conclude in Section 4 by describing a simple upper bound version of Theorem 1.2, and conjecturing a characterization for when equality holds.

Karola Mészáros is partially supported by CAREER NSF Grant DMS-1847284. Avery St. Dizier received support from NSF Grant DMS-2002079.

2. Background

We first define flagged Weyl modules and their dual characters. We then recall the definition of Schubert polynomials and the connection between Schubert polynomials and dual characters. The exposition of this section follows that of [7].

By a *diagram*, we mean a sequence $D = (C_1, C_2, \ldots, C_n)$ of finite subsets of [n], called the *columns* of D. We interchangeably think of $D \subseteq [n] \times [n]$ as a collection of boxes (i, j) in a grid, viewing an element $i \in C_j$ as a box in row i and column j of the grid. When we draw diagrams, we read the indices as in a matrix: i increases top-to-bottom and j increases left-to-right.

Let $G = \operatorname{GL}(n, \mathbb{C})$ be the group of $n \times n$ invertible matrices over \mathbb{C} and B be the subgroup of G consisting of the $n \times n$ upper-triangular matrices. The flagged Weyl module is a representation of B associated to a diagram D. The flagged Weyl module of D will be denoted by \mathcal{M}_D . We will use the construction of \mathcal{M}_D in terms of determinants given in [17].

Denote by Y the $n \times n$ matrix with indeterminates y_{ij} in the upper-triangular positions and zeros elsewhere. Let $\mathbb{C}[Y]$ be the polynomial ring in the indeterminates $\{y_{ij}\}_{i \leq j}$. Note that B acts on $\mathbb{C}[Y]$ on the right via left translation: if $f(Y) \in \mathbb{C}[Y]$, then a matrix $b \in B$ acts on f by $f(Y) \cdot b = f(b^{-1}Y)$. For any $R, S \subseteq [n]$, let Y_S^R be the submatrix of Y obtained by restricting to rows R and columns S.

For $R, S \subseteq [n]$, we say $R \leq S$ if #R = #S and the *k*th least element of R does not exceed the *k*th least element of S for each k. For any diagrams $C = (C_1, \ldots, C_n)$ and $D = (D_1, \ldots, D_n)$, we say $C \leq D$ if $C_j \leq D_j$ for all $j \in [n]$.

Definition 2.1. For a diagram $D = (D_1, \ldots, D_n)$, the *flagged Weyl module* \mathcal{M}_D is defined by

$$\mathcal{M}_D = \operatorname{Span}_{\mathbb{C}} \left\{ \prod_{j=1}^n \det \left(Y_{D_j}^{C_j} \right) \middle| C \le D \right\}$$

 \mathcal{M}_D is a *B*-module with the action inherited from the action of *B* on $\mathbb{C}[Y]$.

Note that since Y is upper-triangular, the condition $C \leq D$ is technically unnecessary since det $\left(Y_{D_j}^{C_j}\right) = 0$ unless $C_j \leq D_j$. Conversely, if $C_j \leq D_j$, then det $\left(Y_{D_j}^{C_j}\right) \neq 0$. For any B-module N, the character of N is defined by char $(N)(x_1, \ldots, x_n) = \operatorname{tr}(X : N \to N)$ where X

For any *B*-module *N*, the character of *N* is defined by $char(N)(x_1, \ldots, x_n) = tr(X : N \to N)$ where *X* is the diagonal matrix with diagonal entries x_1, \ldots, x_n , and *X* is viewed as a linear map from *N* to *N* via the *B*-action. Define the *dual character* of *N* to be the character of the dual module N^* :

$$\operatorname{char}^{*}(N)(x_{1},\ldots,x_{n}) = \operatorname{tr}(X:N^{*} \to N^{*})$$

= $\operatorname{char}(N)(x_{1}^{-1},\ldots,x_{n}^{-1}).$

Definition 2.2. For a diagram $D \subseteq [n] \times [n]$, let $\chi_D = \chi_D(x_1, \ldots, x_n)$ be the dual character

$$\chi_D = \operatorname{char}^* \mathcal{M}_D$$

Example 2.3. Let *D* be the diagram

$$D = (\{1,3\},\{2,3\}) =$$

Then the diagrams C with $C \leq D$ are



The corresponding determinants are

 $y_{11}y_{22}y_{33}^2$, $y_{11}y_{12}y_{23}y_{33} - y_{11}y_{13}y_{22}y_{33}$ $y_{11}y_{12}y_{33}^2$ $y_{11}y_{22}y_{23}y_{33}$ $y_{11}y_{12}y_{23}^2 - y_{11}y_{13}y_{22}y_{23}$ $y_{11}y_{12}y_{23}y_{33}$ These determinants are all linearly independent eigenvectors of X, so

$$\chi_D(x_1, x_2, x_3) = x_1 x_2 x_3^2 + 2x_1^2 x_2 x_3 + x_1^2 x_3^2 + x_1 x_2^2 x_3 + x_1^2 x_2^2.$$

Definition 2.4. For any diagram $D \subseteq [n] \times [n]$ with columns D_1, \ldots, D_n , we write x^D for the monomial

$$x^D = \prod_{j=1}^n \prod_{i \in D_j} x_i$$

The following two easy results describe the supports and coefficients of dual characters of diagrams.

Theorem 2.5 (cf. [6, Theorem 7]). For any diagram $D \subseteq [n] \times [n]$, the monomials appearing in χ_D are exactly

$$\left\{ x^D \mid C \le D \right\}$$

Corollary 2.6. Let $D \subseteq [n] \times [n]$ be a diagram. Fix any diagram $C^{(1)} \leq D$ and set $\mathbf{m} = x^{C^{(1)}}$. Suppose $C^{(1)}, \ldots, C^{(r)}$ are all the diagrams $C \leq D$ such that $x^{C} = \mathbf{m}$. Then, the coefficient of \mathbf{m} in χ_{D} is equal to

the number of linearly independent polynomials over
$$\mathbb{C}$$
 among $\left\{ \prod_{j=1}^{n} \det \left(Y_{D_j}^{C_j^{(i)}} \right) \middle| i \in [r] \right\}$

2.1. Schubert Polynomials. Recall the *divided difference operators* ∂_j for $j \in [n-1]$ are operators on the polynomial ring $\mathbb{C}[x_1, \ldots, x_n]$ defined by

$$\partial_j(f) = \frac{f - (s_j \cdot f)}{x_j - x_{j+1}} = \frac{f(x_1, \dots, x_n) - f(x_1, \dots, x_{j+1}, x_j, \dots, x_n)}{x_j - x_{j+1}}$$

Definition 2.7. The Schubert polynomial \mathfrak{S}_w of $w \in S_n$ is defined recursively on the weak Bruhat order. Let $w_0 = n n - 1 \cdots 2 1 \in S_n$, the longest permutation in S_n . If $w \neq w_0$ then there is $j \in [n-1]$ with w(j) < w(j+1) (called an *ascent* of w). The polynomial \mathfrak{S}_w is defined by

$$\mathfrak{S}_{w} = \begin{cases} x_{1}^{n-1} x_{2}^{n-2} \cdots x_{n-1} & \text{if } w = w_{0}, \\ \partial_{j} \mathfrak{S}_{ws_{j}} & \text{if } w(j) < w(j+1). \end{cases}$$

Definition 2.8. The *Rothe diagram* D(w) of a permutation $w \in S_n$ is the diagram

$$D(w) = \{(i,j) \in [n] \times [n] \mid i < w^{-1}(j) \text{ and } j < w(i)\}.$$

The diagram D(w) can be visualized as the set of boxes left in the $n \times n$ grid after you cross out all boxes weakly below (i, w(i)) in the same column, or weakly right of (i, w(i)) in the same row for each $i \in [n]$.

Example 2.9. If w = 31542, then



The Schubert polynomial of w is computed by

$$\mathfrak{S}_w = \partial_2 \partial_1 \partial_3 \partial_2 \partial_4 (x_1^4 x_2^3 x_3^2 x_4)$$

Via Rothe diagrams, Schubert polynomials occur as special cases of dual characters of flagged Weyl modules:

Theorem 2.10 ([13]). Let w be a permutation with Rothe diagram D(w). Then,

$$\mathfrak{S}_w = \chi_{D(w)}.$$

2.2. Key Polynomials. Key polynomials were first introduced by Demazure for Weyl groups [3], and studied in the context of the symmetric group by Lascoux and Schützenberger in [14, 15]. Recall the key polynomial κ_{α} of a composition $\alpha = (\alpha_1, \alpha_2, ...)$ is defined as follows. When α is a partition, $\kappa_{\alpha} = x^{\alpha}$. Otherwise, suppose $\alpha_i < \alpha_{i+1}$ for some *i*. Then

$$\kappa_{\alpha} = \partial_i(x_i \kappa_{\beta}), \text{ where } \beta = (\alpha_1, \dots, \alpha_{i+1}, \alpha_i, \dots).$$

Definition 2.11 ([11,19]). Fix a composition α , and set

$$l = \max\{i : \alpha_i \neq 0\}$$
 and $n = \max\{l, \alpha_1, \dots, \alpha_l\}.$

The skyline diagram of α is the diagram $D(\alpha) \subseteq [n] \times [n]$ containing the leftmost α_i boxes in row *i* for each $i \in [n]$.

Example 2.12. If $\alpha = (3, 2, 0, 1, 1)$, then



The key polynomial of α is computed by

$$\mathbf{x}_{\alpha} = \partial_3(x_3\partial_4(x_4(x_1^3x_2^2x_3x_4))).$$

Theorem 2.13 ([4]). Let α be a composition with skyline diagram $D(\alpha)$. Then

$$\kappa_{\alpha} = \chi_{D(\alpha)}$$

3. A Lower Bound for $\chi_D(1,\ldots,1)$

In this section, we prove a lower bound for the principal specialization of the dual character of any diagram. We then specialize this bound to Schubert and key polynomials.

Definition 3.1. Fix a diagram D. For each box $(i, j) \in D$, the rank of that box is

 $\operatorname{rank}_{D}(i,j) = \#\{k \colon 1 \le k \le i \text{ and } (k,j) \notin D\}.$

The rank of D is

$$\operatorname{rank}(D) = \sum_{(i,j)\in D} \operatorname{rank}_D(i,j).$$

Lemma 3.2. Let D be a diagram and let $r = \operatorname{rank}(D)$. Then there are diagrams $C^0, C^1, \ldots, C^{r-1}$ such that $C^0 < C^1 < \cdots < C^{r-1} < D$ and $\operatorname{rank}(C^k) = k$.

Proof. If r = 0 then the chain consists of just D and there is nothing to prove. Assume r > 0. We begin with the case that D has a single nonempty column. Without loss of generality, we may write $D = (D_1) = (\{a_1, \ldots, a_m\})$. Since rank(D) > 0, $D_1 \neq \{1, \ldots, m\}$. Let k be the largest integer less than a_m such that $k \notin D_1$. Choose i so that $a_i = k + 1$ (which must exist by definition of k). Define

$$C_1 = (D_1 \setminus \{a_i\}) \cup \{k\}$$

Then $C_1 < D_1$, and rank $(C_1) = \operatorname{rank}(D_1) - 1$. By induction, the result follows whenever D has a single nonempty column. Since

$$\operatorname{rank}((D_1,\ldots,D_n)) = \sum_{j \in [n]} \operatorname{rank}((D_j)),$$

the general case follows from the single column case by performing the above construction to one column at a time. $\hfill \Box$

Recall the *inverse lexicographic order* on monomials: $x^a <_{invlex} x^b$ if there exists $1 \le i \le n$ such that $a_j = b_j$ for $i + 1 \le j \le n$, and $a_i < b_i$.

Lemma 3.3. If C < D, then $x^C \neq x^D$.

Proof. Let $C = (C_1, \ldots, C_n)$ and $D = (D_1, \ldots, D_n)$, so $C_j \leq D_j$ for all $j \in [n]$. Fix a column j. Then $C_j \leq D_j$ means we can write $C_j = \{a_1, \ldots, a_m\}$ and $D_j = \{b_1, \ldots, b_m\}$ with $a_i \leq b_i$ for all $i \in [m]$. Consequently,

$$\prod_{i \in C_j} x_i \leq_{\text{invlex}} \prod_{i \in D_j} x_i.$$

Since C < D, we know $C_j < D_j$ for at least one $j \in [n]$. For any such j, we have

$$\prod_{i \in C_j} x_i <_{\text{invlex}} \prod_{i \in D_j} x_i,$$

 \mathbf{SO}

$$x^{C} = \prod_{j=1}^{n} \prod_{i \in C_{j}} x_{i} <_{\text{invlex}} \prod_{j=1}^{n} \prod_{i \in D_{j}} x_{i} = x^{D}$$

In particular $x^C \neq x^D$.

Theorem 1.2. For any diagram D,

$$\chi_D(1,\ldots,1) \ge \operatorname{rank}(D) + 1$$

Proof. By Theorem 2.5,

$$\chi_D(1,\ldots,1) \ge \# \left\{ x^C \mid C \le D \right\}$$

By Lemma 3.2, there exists a chain of $r = \operatorname{rank}(D) + 1$ diagrams $C^0 < C^1 < \cdots < C^{r-1} < D$. Thus, by Lemma 3.3,

$$\# \left\{ x^C \mid C \le D \right\} \ge \# \left\{ x^C \mid C \in \{C^0, C^1, \dots, C^{r-1}, D\} \right\} = r + 1 = \operatorname{rank}(D) + 1.$$

By specializing Theorem 1.2 to Rothe diagrams, we obtain a new proof of Theorem 1.1:

Corollary 3.4 ([23, Theorem 1.1]). For any permutation $w \in S_n$,

$$\mathfrak{S}_w(1,\ldots,1) \ge 1 + p_{132}(w),$$

where $p_{132}(w)$ is the number of 132-patterns in w.

Proof. It is enough to show that $p_{132}(w) = \operatorname{rank}(D(w))$. By viewing 132-patterns of w graphically in D(w), one easily observes that 132-patterns are in transparent bijection with tuples (i, j, k) such that $(i, j) \in D(w)$, $1 \le k < i$, and $(k, j) \notin D(w)$. The quantity $\operatorname{rank}(D)$ exactly counts these tuples.

By specializing Theorem 1.2 to skyline diagrams, we obtain an analogous result for key polynomials. For a composition α , let rinv(α) denote the set of *right inversions* of α , the pairs i < j such that $\alpha_i < \alpha_j$.

Corollary 3.5. For any composition α ,

$$\kappa_{\alpha}(1,\ldots,1) \ge 1 + \sum_{(i,j)\in \operatorname{rinv}(\alpha)} (\alpha_j - \alpha_i).$$

We now characterize the case of equality in Theorem 1.2.

Definition 3.6. Let D be any diagram. A pair of boxes $(i, j), (i', j') \in D$ is called an *unstable pair* if

- rank_D $(i, j) \ge 1;$
- $\operatorname{rank}_D(i', j') \ge 1;$
- If i = i' or j = j', then $\operatorname{rank}_D(i, j) + \operatorname{rank}_D(i', j') \ge 3$.

Proposition 3.7. A diagram D satisfies $\chi_D(1, \ldots, 1) = \operatorname{rank}(D) + 1$ if and only if D does not contain an unstable pair.

Proof. Suppose D contains an unstable pair $\{(i, j), (i', j')\}$. A simple case analysis shows one can move boxes in D upwards to create diagrams $C, C' \leq D$ of the same rank with $x^C \neq x^{C'}$. This implies $\chi_D(1, \ldots, 1) \neq \operatorname{rank}(D) + 1$.

Assume D contains no unstable pair. If $\operatorname{rank}(D) = 0$, then the result follows. Pick $(i, j) \in D$ with $\operatorname{rank}_D(i, j) \geq 1$. If $\operatorname{rank}_D(i, j) > 1$, then any other positive rank box would form an unstable pair with (i, j). Hence (i, j) is the only positive rank box of D, and the result follows easily.

Suppose rank_D(i, j) = 1. To avoid unstable pairs, all other positive rank boxes of D either lie in row i, or they all lie in column j. In either case, they must all have rank exactly 1. If all positive rank boxes of D lie in column i, then one observes there is a unique diagram $C \leq D$ with rank k for each k = 0, 1, ..., rank(D), implying the result.

If all positive rank boxes of D lie in row j, then one observes that all diagrams $C \leq D$ of a fixed rank have the same monomial x^{C} , and their determinants span an eigenspace of dimension one in the flagged Weyl module.

We now relate equality in Theorem 1.2 with the question of when χ_D is zero-one. Recall a polynomial f is called *zero-one* if all nonzero coefficients in f equal 1.

Proposition 3.8. If a diagram D satisfies $\chi_D(1, \ldots, 1) = \operatorname{rank}(D) + 1$, then χ_D is zero-one.

Proof. In order for $\chi_D(1, \ldots, 1) = \operatorname{rank}(D) + 1$, it must happen that all diagrams $C \leq D$ with a fixed rank induce the same monomial x^C and have dependent determinants in the flagged Weyl module. Since all diagrams $C, C' \leq D$ with $x^C = x^{C'}$ must have the same rank, it follows that all eigenspaces in the flagged Weyl module of D have dimension one.

We now provide a conjectural characterization of diagrams D such that χ_D is zero-one. Consider the six box configurations shown in Figure 1. In each configuration, an \times (red) indicates the absence of a box; a shaded square (gray) indicates the presence of a box; and an unshaded square (white) indicates no restriction on the presence of a box.

Definition 3.9. Let *D* be any diagram. We say *D* contains a *multiplicitous configuration* if there are $r_1 < r_2 < r_3 < r_4$ and $c_1 < c_2$ so that *D* restricted to rows $\{r_1, r_2, r_3, r_4\}$ and columns $\{c_1, c_2\}$ equals one of the configurations shown in Figure 1, up to possibly swapping the order of the columns.



FIGURE 1. The six multiplicitous configurations.

Example 3.10. Consider the diagrams D and D' shown in Figure 2. The diagram D does not contain instances of any multiplicitous configurations. The diagram D' contains instances of each of the multiplicitous configurations.

Proposition 3.11. If a diagram D contains a multiplicitous configuration, then χ_D is not zero-one.

Proof. It follows immediately from [7, Theorem 5.8] that if the restriction of a diagram D to rows $\{i_1, \ldots, i_p\}$ and columns $\{j_1, \ldots, j_q\}$ equals a diagram D', then the largest coefficient appearing in χ_D is bounded below by the largest coefficient appearing in $\chi_{D'}$. One easily checks that the dual characters of each of the multiplicitous configurations are not zero-one.

Conjecture 3.12. If D is a diagram such that χ_D is not zero-one, then D contains a multiplicitous configuration.



FIGURE 2.

Together, Proposition 3.11 and Conjecture 3.12 specialize to known results for Schubert and key polynomials: [7, Theorem 1] when D is the Rothe diagram of a permutation, and [12, Theorem 1.1] when D is the skyline diagram of a composition.

4. An Upper Bound for $\chi_D(1,\ldots,1)$

In this final section, we recall a trivial upper bound for the principal specialization of the dual character of any diagram. We make a conjecture for the case of equality. From Corollary 2.6, it follows immediately that if c_{α} is the coefficient of x^{α} in χ_D , then

$$c_{\alpha} \le \#\{C \le D \mid x^C = x^{\alpha}\}.$$

In particular,

$$\chi_D(1,...,1) \le \#\{C \mid C \le D\}$$

Fan and Guo gave the following characterization for equality when the diagram D is northwest. Recall a diagram D is northwest if whenever $(i, j), (i', j') \in D$ with i > i' and j < j', one has $(i', j) \in D$.

Theorem 4.1 ([5]). For any northwest diagram D,

$$\chi_D(1,...,1) = \#\{C \mid C \le D\}$$

if and only if D contains no instance of the configuration shown in Figure 3.



FIGURE 3.

We conjecturally extend Theorem 4.1 to all diagrams.

Conjecture 4.2. Let D be any diagram. Then $\chi_D(1, \ldots, 1) = \#\{C \mid C \leq D\}$ if and only if D contains no instance of the configuration shown in Figure 4.



FIGURE 4.

References

- [1] N. Bergeron and S. Billey. RC-graphs and Schubert polynomials. Experiment. Math., 2(4):257-269, 1993.
- S. Billey, A. Holroyd, and B. Young. A bijective proof of Macdonald's reduced word formula. Algebr. Comb., 2(2):217–248, 2019.
- [3] M. Demazure. Key polynomials and a flagged Littlewood–Richardson rule. Bull. Sci. Math., 98:163–172, 1974.
- [4] M. Demazure. Une nouvelle formule des caracteres. J. Combin. Theory Ser. A, 70(1):107–143, 1995.
- [5] N. Fan and P.L. Guo. Upper bounds of Schubert polynomials. Sci. China Math., 1(4):415–423, 2020.
- [6] A. Fink, K. Mészáros, and A. St. Dizier. Schubert polynomials as integer point transforms of generalized permutahedra. Adv. Math., 332:465–475, 2018.
- [7] A. Fink, K. Mészáros, and A. St. Dizier. Zero-one Schubert polynomials. Math. Z., 2020.
- [8] S. Fomin and A. N. Kirillov. The Yang-Baxter equation, symmetric functions, and Schubert polynomials. Discrete Math., 153(1):123–143, 1996. Proceedings of the 5th Conference on Formal Power Series and Algebraic Combinatorics.
- [9] S. Fomin and A.N. Kirillov. Reduced words and plane partitions. J. Algebraic Combin., 6(4):311–319, 1997.
- [10] Y. Gao. Principal specializations of Schubert polynomials and pattern containment. European J. Combin., 94:103291, 2021.
- [11] J. Haglund, M. Haiman, and N. Loehr. A combinatorial formula for nonsymmetric Macdonald polynomials. Amer. J. Math., 130(2):359–383, 2008.
- [12] R. Hodges and A. Yong. Multiplicity-free key polynomials, 2020. arXiv:2007.09229.
- [13] W. Kraśkiewicz and P. Pragacz. Foncteurs de Schubert. C. R. Acad. Sci. Paris Sér. I Math., 304(9):209–211, 1987.
- [14] A. Lascoux and M.-P. Schützenberger. Polynômes de Schubert. C. R. Acad. Sci. Paris Sér. I Math., 294(13):447–450, 1982.
- [15] A. Lascoux and M.-P Schützenberger. Structure de Hopf de l'anneau de cohomologie et de l'anneau de Grothendieck d'une variété de drapeaux. C. R. Acad. Sci. Paris Sér. I Math., 295(11):629–633, 1982.
- [16] I.G. Macdonald. Notes on Schubert polynomials. Publ. LaCIM, UQAM, Montrèal, 1991.
- [17] P. Magyar. Schubert polynomials and Bott-Samelson varieties. Comment. Math. Helv., 73(4):603-636, 1998.
- [18] K. Mészáros and A. Tanjaya. Inclusion-exclusion on Schubert polynomials, 2021. arXiv:2102.11179.
- [19] C. Monical, N. Tokcan, and A. Yong. Newton polytopes in algebraic combinatorics. Selecta Math. (N.S.), 25(66), 2019.
- [20] A. Morales, I. Pak, and G. Panova. Asymptotics of principal evaluations of Schubert polynomials for layered permutations. Proc. Amer. Math. Soc., 147(4):1377–1389, 2019.
- [21] R.P. Stanley. Enumerative Combinatorics, volume 2 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1999.
- [22] R.P. Stanley. Some Schubert shenanigans, 2017. arXiv:1704.00851.
- [23] A. Weigandt. Schubert polynomials, 132-patterns, and Stanley's conjecture. Algebr. Comb., 1(4):415-423, 2018.

KAROLA MÉSZÁROS, DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NY 14853. karola@math.cornell.edu

AVERY ST. DIZIER, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801. stdizie2@illinois.edu

ARTHUR TANJAYA, DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA NY 14853. amt333@cornell.edu