

AN UPPERBOUND FOR THE NUMBER OF CRITICAL POINTS OF THE SYSTOLE FUNCTION ON THE SURFACE MODULI SPACE

YUE GAO

ABSTRACT. We obtain an upper bound for the number of critical points of the systole function on \mathcal{M}_g . Besides, we obtain an upper bound for the number of those critical points whose systole is smaller than a constant.

CONTENTS

1.	Introduction	1
2.	Preliminaries	3
2.1.	Topological Morse function	3
2.2.	Teichmüller space and length function	3
2.3.	Filling curves	4
3.	Upper bound	4
4.	Further questions	9
	References	9

1. INTRODUCTION

The systole of a closed hyperbolic surface is the length of the shortest geodesic on the surface. It can also be treated as a function on the Teichmüller space \mathcal{T}_g , denoted as $\text{sys}_g : \mathcal{T}_g \rightarrow \mathbb{R}_+$. The systole function is also well-defined on the moduli space \mathcal{M}_g .

The systole function on \mathcal{T}_g or \mathcal{M}_g has been shown to be a topological Morse function (for its definition, see Section 2) by Akrouf [Akr03]. Similarly to a Morse function, a topological Morse function has regular and critical points, and critical points of a topological Morse function on a space provide information of the space's topology. For example, let $f : M \rightarrow \mathbb{R}$ be a Morse function on a differential manifold M (or a topological Morse function $f : M \rightarrow \mathbb{R}$ on a topological manifold M), then the Euler characteristic $\chi(M)$ of M is given by

$$\chi(M) = \sum_k (-1)^k \# \{\text{critical points of } f \text{ with index } k\}.$$

Remark 1. For the moduli space \mathcal{M}_g , Harer and Zagier obtained its orbifold Euler characteristic:

$$\chi(\mathcal{M}_g) = \frac{B_{2g}}{4g(g-1)}$$

in [HZ86]. Here B_{2g} is the $2g$ -th Bernoulli number.

The study on critical points of systole function initiated in a series of work by Schmutz [Sch93], [Sch94], [Sch98], [Sch99]. He proved that systole is a topological Morse function on the space of cusped surfaces before Akrouf and classified critical points of systole function on genus 2 moduli space \mathcal{M}_2 . Besides he provided several series of critical points, including locally maximal points.

Recently, Fortier-Bourque and Rafi obtained new examples of critical points (including locally maximal points) of systole function [FB19] [FBR20]. Part of their examples have relatively large systole and part of theirs have small indices. Critical points with such properties had not been known before.

Examples of critical points and the Euler characteristic $\chi(\mathcal{M}_g)$ provide lower bounds for the number of critical points of systole function. The largest lower bound is given by $\chi(\mathcal{M}_g)$ and asymptotic to $\frac{\sqrt{\pi}}{\sqrt{g(g-1)}} \left(\frac{g}{\pi e}\right)^{2g}$. Some examples offer lower bound for number of critical points with some restrictions.

The main result of this paper is an upper bound of this number. We let sys_g be the systole function and $\text{Crit}(\text{sys}_g)$ be the set of its critical points. For $L > 0$ we let

$$\text{Crit}(\text{sys}_g)^{\leq L} = \{\Sigma \in \text{Crit}(\text{sys}_g) \subset \mathcal{M}_g \mid \text{sys}(\Sigma) \leq L\}.$$

Then we get an upper bound for $|\text{Crit}(\text{sys}_g)^{\leq L}|$ when $g \geq 2$ (Proposition 3.4). When g is sufficiently large, we have

Theorem 3.5. *If g is large enough then the number of points in $\text{Crit}(\text{sys}_g)$ on \mathcal{M}_g is bounded from above by*

$$|\text{Crit}(\text{sys}_g)| \leq (6g)^{Cg^{\frac{7}{2}}}.$$

Here $C = 6054$.

Also we have a theorem on how the critical points distribute on \mathcal{M}_g .

Theorem 3.6. *When g is sufficiently large, then*

$$|\text{Crit}(\text{sys}_g)^{\leq L}| \leq \left(2^{C_1 g e^L} \cdot 2^{C_2 g e^{\frac{5L}{4}}}\right) \cdot \left((e^L)^{C_3 g e^L} \cdot (e^L)^{C_4 g e^{\frac{5L}{4}}}\right) \cdot g^{C_3 g e^L}.$$

Here $C_1 = 200000$, $C_2 = 5040$, $C_3 = 15400$, and $C_4 = 1300$.

Here is a comparison of our upper bounds with some known lower bounds. Our upper bound for $|\text{Crit}(\text{sys}_g)^{\leq L}|$ is $(Cg)^{C'g}$ as g grows for some $C = C(L) > 0$ and $C' = C'(L) > 0$. On the other hand, there is a series $\{g_k\}$, $g_k \rightarrow \infty$ as $k \rightarrow \infty$, and there is a known lower bound for $|\text{Crit}(\text{sys}_{g_k})^{\leq L}|$, also is $(Cg_k)^{C'g_k}$ as k grows for some other $C, C' > 0$ (see [FBR20, Theorem B], Theorem 3.7 and Corollary 3.8). As for $|\text{Crit}(\text{sys}_g)|$, the gap between our upper bound $((Cg)^{C'g^{\frac{7}{2}}})$ and the lower bound implied by $\chi(\mathcal{M}_g)$ is not that small. Currently we still do not have much knowledge on critical points with large systole to obtain a sharper estimate to $|\text{Crit}(\text{sys}_g)|$.

In an unpublished manuscript [Thu86], Thurston claimed that let $X_g \subset \mathcal{M}_g$ be the space consisting of surfaces with filling shortest geodesics, then X_g is a spine (deformation retract) of \mathcal{M}_g (See also [Ji14] [FB19], [FBR20]). Here by filling we mean that the union of the shortest geodesics cut the surfaces into polygonal disks. Unfortunately, recently his proof was shown incomplete [Ji14]. Thurston's claim is still unsolved. Critical points of the systole function are contained in X_g [Sch99, Corollary 20]. and our result characterizes critical points of the systole function.

We use two combinatorial properties in [Sch99] (Corollary 20, 21) to obtain our result. These two properties tell us that by bounding number of filling graph (with restrictions to number of vertices, edges and degree at each vertex) from above, we can bound $|\text{Crit}(\text{sys}_g)|$ from above.

The method to bound number of filling graphs on surfaces from above is similar to Kahn-Markovic's method to bound number of triangulations on surfaces from above in [KM12].

In Section 2, some preliminaries we need in our proof are offered. Then we prove Theorem 3.5 and Theorem 3.6 in Section 3. In Section 4, we give some further questions to consider.

Acknowledgement: The author would like to acknowledge Prof. Yi Liu for many helpful discussions.

2. PRELIMINARIES

2.1. Topological Morse function. Let M^n be a n -dimensional topological manifold.

Definition 2.1. *A function $f : M^n \rightarrow \mathbb{R}$ is a topological Morse function if at each point $p \in M$, there is a neighborhood U of p and a map $\psi : U \rightarrow \mathbb{R}^n$. Here ψ is a homeomorphism between U and its image, such that $f \circ \psi^{-1}$ is either a linear function or*

$$f \circ \psi^{-1}((x_1, x_2, \dots, x_n)) = f(p) - x_1^2 - x_2^2 - \dots - x_j^2 + x_{j+1}^2 + \dots + x_n^2.$$

In the former case, the point p is called a regular point of f , while in the latter case the point p is called a singular point with index j .

2.2. Teichmüller space and length function. We denote by \mathcal{T}_g the Teichmüller space consisting of marked hyperbolic surface with genus g , and \mathcal{M}_g to be the moduli space consisting of hyperbolic surface with genus g . It is known that

$$\mathcal{M}_g \cong \mathcal{T}_g / \Gamma_g.$$

Here Γ_g is the mapping class group.

For $\Sigma \in \mathcal{T}_g$, the set of all the shortest geodesics is denoted by $S(\Sigma)$. The length of an essential curve α on Σ is denoted by $l_\alpha(\Sigma)$. For $\alpha \in S(\Sigma)$,

$$l_\alpha(\Sigma) \leq l_\beta(\Sigma), \forall \text{ simple closed geodesic } \beta \subset \Sigma.$$

Length of the shortest geodesics can be treated as a function on \mathcal{T}_g , we denote it as sys_g or shortly sys . Obviously for $\alpha \in S(\Sigma)$

$$\text{sys}(\Sigma) = l_\alpha(\Sigma) = \inf_{\forall \text{ simple closed geodesic } \beta \subset \Sigma} l_\beta(\Sigma).$$

Since systole is an invariant function on \mathcal{T}_g under the action of the mapping class group, systole function can be also defined as a function on \mathcal{M}_g .

$$\begin{aligned} \text{sys} : \mathcal{M}_g &\rightarrow \mathbb{R}^+ \\ \Sigma &\mapsto \text{sys}(\Sigma). \end{aligned}$$

Akrout showed that

Corollary 2.2 ([Akr03], Corollary). *The systole function is a topological Morse function on \mathcal{T}_g .*

Therefore systole function is also a topological Morse function on \mathcal{M}_g .

The set of all the critical points of sys_g in \mathcal{M}_g is denoted by $\text{Crit}(\text{sys}_g)$. The number of the critical points in \mathcal{M}_g , is denoted by $|\text{Crit}(\text{sys}_g)|$.

2.3. Filling curves.

Definition 2.3. For a closed surface Σ , we assume $F = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$ is a finite set of essential simple closed curves. F is said to be filling if and only if each component of $\Sigma \setminus F$ is a disk.

We need criteria regarding critical points of the systole function provided by the following two corollaries in [Sch99].

Corollary 2.4 ([Sch99] Corollary 20). If $\Sigma \in \mathcal{T}_g$ is a critical point of the systole function, then $S(\Sigma)$ is a filling set.

Definition 2.5. If F and F' are filling sets on Σ and Σ' respectively, then (Σ, F) is equivalent to (Σ', F') if and only if there is a homeomorphism from Σ to Σ' that maps the elements of F to those of F' .

Corollary 2.6 ([Sch99] Corollary 21). For $\Sigma, \Sigma' \in \mathcal{T}_g$, if $(\Sigma, S(\Sigma)), (\Sigma', S(\Sigma'))$ are equivalent then Σ and Σ' cannot both be critical points of the systole function.

3. UPPER BOUND

On a $\Sigma \in \text{Crit}(\text{sys})$, the union of shortest geodesics can be treated as a graph. Vertices of this graph are the intersection points of shortest geodesics. Edges of the graph consist of subarcs of shortest geodesics connecting two intersection points of shortest geodesics. This graph is denoted by Γ .

To obtain the upper bound of $|\text{Crit}(\text{sys}_g)|$ on \mathcal{M}_g , first we need to estimate the number of vertices, edges and degree at each vertex of the graph Γ defined in the previous paragraph. Then we count the number of graphs satisfying this restriction by the method of Kahn-Markovic in [KM12]. This number bounds $|\text{Crit}(\text{sys}_g)|$ from above because of Corollary 2.4 and 2.6.

It is known that two shortest geodesics intersect at most once, see for example [Par14].

By this fact, we have the following estimate:

Lemma 3.1. For any $\Sigma \in \text{Crit}(\text{sys}_g)$, the graph $\Gamma = (V, E)$ satisfies

- (1) $|V| \leq 2545ge^{\text{sys}(\Sigma)}$,
- (2) $|E| \leq 3|V| + 6g - 6 \leq 7700ge^{\text{sys}(\Sigma)}$,
- (3) The genus¹ of Γ (denoted by g_Γ) is bounded by $\pi \frac{\sqrt{g(g-1)}}{\log(4g-2)} \leq g_\Gamma \leq 3|V| + 6g - 6 \leq 7700ge^{\text{sys}(\Sigma)}$.
- (4) For all $v \in V$, $\deg(v) \leq 2e^{\text{sys}(\Sigma)/4}$.

Proof. (1) In [Par13], for a surface $\Sigma \in \mathcal{M}_g$, Parlier defined $F(\Sigma)$ to be the minimum number of radius- r -disks needed for covering Σ and $G(\Sigma)$ to be supremum number of shortest geodesics of Σ that pass through a radius- r -disk, where $r := \text{arcsinh}\left(\frac{1}{2\sinh(\text{sys}(\Sigma)/4)}\right)$. These two numbers are bounded from above by

$$F(\Sigma) < 16(g-1)e^{\text{sys}(\Sigma)/2} \text{ and } G(\Sigma) < 17.83e^{\text{sys}(\Sigma)/4}.$$

¹For a connected graph $\Gamma = (V, E)$, the genus of Γ is the number of circles of Γ , more precisely, is $\min_{E_0 \in \mathcal{E}} |E_0|$. Here $\mathcal{E} = \{E_0 \subset E | \Gamma \setminus E_0 \text{ is a tree.}\}$.

In each radius- r -disk, there are at most $\binom{G(\Sigma)}{2}$ intersection points since it is known that two shortest geodesics intersect at most once. Then on the surface Σ , number of intersection points of shortest geodesics is bounded from above by

$$\frac{F(\Sigma)G^2(\Sigma)}{2}.$$

Then (1) follows by direct calculation.

(2) We can add some edges to Γ to get a triangulation of Σ . Therefore we can bound the number of edges of Γ from above by bounding the number of edges of the triangulations containing Γ from above.

If Γ is a triangulation of Σ , then the number of edges and faces of this triangulation satisfies the following relation:

$$(3.1) \quad 2E = 3F.$$

The Euler characteristic formula is

$$(3.2) \quad V - E + F = 2 - 2g.$$

Then by (3.1) and (3.2), (2) follows.

(3) The genus of a graph is $g_\Gamma = 1 - \chi(\Gamma)$. Here $\chi(\Gamma)$ is the Euler characteristic of Γ . Then the upper bound follows from (1) and (2).

By [APP11, Theorem 3], if $\Sigma \in \mathcal{M}_g$ has filling shortest geodesics then $|S(\Sigma)| \geq \pi \frac{\sqrt{g(g-1)}}{\log(4g-2)}$. Therefore $\pi \frac{\sqrt{g(g-1)}}{\log(4g-2)}$ also lower bounds g_Γ .

(4) This follows directly from [Par13, Lemma 2.4], for details, see [Par13]. \square

We may add some vertices to Γ such that (1) every edge of the resulting graph connects two different vertices, (2) number of vertices, edges, and degree of the resulting graph is still bounded by the bounds in Lemma 3.1. We still denote the resulting graph by Γ .

By Corollary 2.4, each component of $\Sigma \setminus \Gamma$ is a polygonal disk. We add some edges to Γ so that we get a triangulation of Σ and the number of vertices, edges and degree at each vertex of the triangulation are bounded by the bounds in Lemma 3.1. (See Figure 1.)




FIGURE 1. A triangulation of the polygon

We recall a definition in [KM12]:

Definition 3.2 ([KM12], Definition 2.1). *For $k > 0$, $g \geq 2$, a triangulation τ of genus g surface is in the set $T(k, g)$ if and only if*

- each vertex of τ has the degree at most k ,
- the graph τ has at most kg vertices and at most kg edges.

By the construction above, we have

Lemma 3.3. *There is a constant $0 < C < 7700$, for all $\Sigma \in \text{Crit}(\text{sys}_g)$, Γ , the graph consisting of shortest geodesics of Σ is a subgraph of a triangulation $\tau \in T(Ce^{\text{sys}(\Sigma)}, g)$.*

By Corollary 2.6, shortest geodesics on two different critical points of sys_g must be non-equivalent. Therefore number of critical points of sys_g whose systole length is smaller than $L > 0$ is bounded from above by number of filling subgraphs of triangulations in $T(Ce^L, g)$ that satisfies the restriction of Lemma 3.1.

Proposition 3.4. *For $L > 0$ we let*

$$\text{Crit}(\text{sys}_g)^{\leq L} = \{\Sigma \in \text{Crit}(\text{sys}_g) \subset \mathcal{M}_g \mid \text{sys}(\Sigma) \leq L\},$$

and let $|V_0| = 2545ge^L$, $|g_{\Gamma_0}| = 7700ge^L$, and $\text{deg}_0 = 2e^{L/4}$.

Then

$$(3.3) \quad |\text{Crit}(\text{sys}_g)^{\leq L}| \leq \text{Cat}(|V_0| - 1) \frac{(|V_0|)^{2g_{\Gamma_0}}}{\left[\pi \frac{\sqrt{g(g-1)}}{\log(4g-2)} \right]!} ([\text{deg}_0]!)^{|V_0|}.$$

Here $\text{Cat}(|V| - 1)$ is the $(|V| - 1)$ -th Catalan number.

Proof. The proof here is similar to the proof of [KM12, Lemma 2.2].

Fix some genus g surface, we assume that

$$\mathcal{G} = \{\Gamma \mid \exists \tau \in T(Ce^L, g), \text{ s.t. } \Gamma \subset \tau, \Gamma \text{ is filling, } \Gamma \text{ satisfies the restriction in Lemma 3.1}\}.$$

Any $\Gamma \in \mathcal{G}$ can be constructed in this way. We pick a $\tau \in T(Ce^L, g)$, and then choose a spanning tree T of τ . Since $\tau \in T(Ce^L, g)$, T has at most $2545ge^L$ vertices. After that, we add g_{Γ} edges $e_1, e_2, \dots, e_{g_{\Gamma}}$ to the tree T in an arbitrary way. Thus $T \cup \{e_1, \dots, e_{g_{\Gamma}}\}$ is a graph with genus at most g_{Γ} . Next we give each vertex of $T \cup \{e_1, \dots, e_{g_{\Gamma}}\}$ a cyclic ordering, thus $T \cup \{e_1, \dots, e_{g_{\Gamma}}\}$ is a *ribbon graph*². We thicken the edges of the ribbon graph and get a surface with boundary. If the genus of this surface is not g , we discard it. Finally, we fill each boundary component of the surface by a disk, thus get a closed surface S_g . Every graph in \mathcal{G} can be constructed this way.

By this construction, $|\mathcal{G}|$ can be bounded from above by $|\mathcal{G}| \leq abc$. Here

$$\begin{aligned} a &= \{\text{number of unlabelled trees } T \text{ with } n \leq 2545ge^L \text{ vertices}\} \\ b &= \{\text{number of ways of adding } g_{\Gamma} \text{ many unlabelled edges } e_1, \dots, e_{g_{\Gamma}} \text{ to } T\} \\ c &= \{\text{number of cyclic orderings of edges of } T \cup \{e_1, \dots, e_{g_{\Gamma}}\}\}. \end{aligned}$$

Because Catalan number $\text{Cat}(n)$ enumerates number of ordered planar trees with $n + 1$ vertices (see for example [Sta15]) and by proof of [KM12, Lemma 2.2] and

²A ribbon graph is a graph with every vertex of the graph assigned a cyclic ordering.

Lemma 3.1,

$$\begin{aligned} a &\leq \text{Cat}(|V_0| - 1) \\ b &\leq \frac{(|V_0|)^{2g_{\Gamma_0}}}{\left[\pi \frac{\sqrt{g(g-1)}}{\log(4g-2)} \right]!} \\ c &\leq (\lceil \text{deg}_0 \rceil!)^{|V_0|}. \end{aligned}$$

Then this proposition follows. \square

By area comparison, for any $\Sigma \in \mathcal{M}_g$, $\text{sys}(\Sigma) \leq \log(2g^2)$. Therefore if we let $L = \log(2g^2)$, then $\text{Crit}(\text{sys}_g) = \text{Crit}(\text{sys}_g)^{\leq L}$. We let $g \rightarrow \infty$, then we have the following theorem:

Theorem 3.5. *If g is large enough then*

$$|\text{Crit}(\text{sys}_g)| \leq (6g)^{Cg^{\frac{7}{2}}}.$$

Here $C = 6054$.

Proof. This theorem follows from direct calculation. First we let $L = \log(2g^2)$ and then we obtain formulae of $|V_0|$, g_{Γ_0} and deg_0 in Proposition 3.4 in g :

$$\begin{aligned} |V_0| &= 2545ge^L = 5090g^3 \\ g_{\Gamma_0} &= 7700ge^L = 15400g^3 \\ \text{deg}_0 &= 2e^{\frac{L}{4}} = 2^{\frac{5}{4}}g^{\frac{1}{2}}. \end{aligned}$$

When n is sufficiently large, $\text{Cat}(n) \sim \frac{4^n}{\sqrt{\pi n^{\frac{3}{2}}}}$, see for example [Sta15]. Thus when g is large enough,

$$(3.4) \quad \text{Cat}(|V_0| - 1) \leq \frac{4^{|V_0|-1}}{\sqrt{\pi}(|V_0| - 1)^{\frac{3}{2}}} \leq 4^{|V_0|-1} \leq 4^{5090g^3}.$$

Besides

$$(3.5) \quad \frac{(|V_0|)^{2g_{\Gamma_0}}}{\left[\pi \frac{\sqrt{g(g-1)}}{\log(4g-2)} \right]!} \leq (|V_0|)^{2g_{\Gamma_0}} = (5090g^3)^{15400g^3},$$

and

$$(3.6) \quad \begin{aligned} (\lceil \text{deg}_0 \rceil!)^{|V_0|} &\leq \text{deg}_0^{\text{deg}_0 |V_0|} \\ &= \left(2^{\frac{5}{4}}g^{\frac{1}{2}} \right)^{2^{\frac{5}{4}}g^{\frac{1}{2}} \cdot 5090g^3} \\ &\leq (5.6g)^{6053.1g^{\frac{7}{2}}}. \end{aligned}$$

Then by (3.3), (3.4), (3.5) and (3.6), when g is large enough,

$$|\text{Crit}(\text{sys}_g)| \leq (6g)^{6054g^{\frac{7}{2}}}.$$

\square

If $L > 0$ is a constant, we have

Theorem 3.6. *If $L > 0$ is a constant, g is sufficiently large, then*

$$|\text{Crit}(\text{sys}_g)^{\leq L}| \leq \left(2^{C_1 g e^L} \cdot 2^{C_2 g e^{\frac{5L}{4}}}\right) \cdot \left((e^L)^{C_3 g e^L} \cdot (e^L)^{C_4 g e^{\frac{5L}{4}}}\right) \cdot g^{C_3 g e^L}.$$

Here $C_1 = 200000$, $C_2 = 5040$, $C_3 = 15400$, and $C_4 = 1300$.

Proof. If L is a constant, g is sufficiently large, then similarly to Theorem 3.5, we have

$$(3.7) \quad \begin{aligned} \text{Cat}(|V_0| - 1) &\leq 4^{|V_0| - 1} \\ &= 4^{2545 g e^L - 1} \end{aligned}$$

$$(3.8) \quad \begin{aligned} \frac{(|V_0|)^{2g_{r_0}}}{\left[\pi \frac{\sqrt{g(g-1)}}{\log(4g-2)}\right]!} &\leq (|V_0|)^{2g_{r_0}} \leq (2545 g e^L)^{7700 g e^L \cdot 2} \\ &= (2545 g e^L)^{15400 g e^L} \end{aligned}$$

$$(3.9) \quad \begin{aligned} ([\deg_0]!)^{|V_0|} &\leq \deg_0^{\deg_0 |V_0|} \leq \left(2e^{\frac{L}{4}}\right)^{2e^{\frac{L}{4}} \cdot 2545 g e^L} \\ &= \left(2e^{\frac{L}{4}}\right)^{5040 g e^{\frac{5L}{4}}}. \end{aligned}$$

Then by (3.3), (3.7), (3.8) and (3.9), when g is large enough,

$$|\text{Crit}(\text{sys}_g)^{\leq L}| \leq (2546 g e^L)^{15401 g e^L}.$$

$$\begin{aligned} |\text{Crit}(\text{sys}_g)^{\leq L}| &\leq 4^{2545 g e^L - 1} \cdot (2545 g e^L)^{15400 g e^L} \cdot \left(2e^{\frac{L}{4}}\right)^{5040 g e^{\frac{5L}{4}}} \\ &\leq \left(2^{5090 g e^L} \cdot (2^{12})^{15400 g e^L} \cdot 2^{5040 g e^{\frac{5L}{4}}}\right) \\ &\quad \left((e^L)^{15400 g e^L} \cdot (e^{\frac{L}{4}})^{5040 g e^{\frac{5L}{4}}}\right) \cdot g^{15400 g e^L} \\ &\leq \left(2^{189890 g e^L} \cdot 2^{5040 g e^{\frac{5L}{4}}}\right) \cdot \left((e^L)^{15400 g e^L} \cdot (e^L)^{1273 g e^{\frac{5L}{4}}}\right) \cdot g^{15400 g e^L}. \end{aligned}$$

□

Fortier-Bouque and Rafi [FBR20] obtained a series of local maxima of the systole function. Letting $n = 3$ in [FBR20, Theorem B], we have

Theorem 3.7 ([FBR20], Theorem B). *There is $\beta > 0$ and a sequence g_k , $g_k \rightarrow \infty$ as $k \rightarrow \infty$ such that*

$$|\text{Crit}(\text{sys}_{g_k})^{\leq 10}| \geq (\beta g_k)^{\frac{1}{3} g_k}.$$

This lower bound is not far from the upperbound we obtain. Letting $L = 10$ in Theorem 3.6, we immediately get the following corollary:

Corollary 3.8. *There are constants $\beta, \beta' > 0$ and a sequence g_k , $g_k \rightarrow \infty$ as $k \rightarrow \infty$ such that*

$$(\beta g_k)^{\frac{1}{3} g_k} \leq |\text{Crit}(\text{sys}_{g_k})^{\leq 10}| \leq (\beta' g_k)^{4 \times 10^8 g_k}.$$

4. FURTHER QUESTIONS

We have some further questions to consider:

With the restriction $\text{sys} \leq C$, the upper bound is quite close to the lower bound.

Then

Question 1. *Could we obtain a sharper estimate to critical points with large systoles?*

Moreover,

Question 2. *For all $C > 0$, does*

$$\frac{|\text{Crit}^{<C}(\text{sys}_g)|}{|\text{Crit}(\text{sys}_g)|}$$

tend to 0 as $g \rightarrow \infty$?

The lower bound in [FBR20, Theorem B] is provided merely by locally maximal points. Therefore,

Question 3. *Could we estimate the number of critical points of the systole function with a specific index?*

On the distribution of the systole function:

Question 4. *Let $l_g := \inf_{\Sigma \in \text{Crit}(\text{sys}_g)} \text{sys}_g(\Sigma)$, by [Sch99, Corollary 20] and Mumford compactness criterion, l_g has a positive universal lower bound. Can we estimate l_g ?*

REFERENCES

- [Akr03] Hugo Akrouit. Singularités topologiques des systoles généralisées. *Topology*, 42(2):291–308, 2003.
- [APP11] James W Anderson, Hugo Parlier, and Alexandra Pettet. Small filling sets of curves on a surface. *Topology and its Applications*, 158(1):84–92, 2011.
- [FB19] Maxime Fortier Bourque. Hyperbolic surfaces with sublinearly many systoles that fill. *Commentarii Mathematici Helvetici*, 2019.
- [FBR20] Maxime Fortier Bourque and Kasra Rafi. Local maxima of the systole function. *Journal of the European Mathematical Society*, 2020.
- [HZ86] John Harer and Don Zagier. The euler characteristic of the moduli space of curves. *Inventiones mathematicae*, 85(3):457–485, 1986.
- [Ji14] Lizhen Ji. Well-rounded equivariant deformation retracts of teichmüller spaces. *L'Enseignement Mathématique*, 60(1):109–129, 2014.
- [KM12] Jeremy Kahn and Vladimir Marković. Counting essential surfaces in a closed hyperbolic three-manifold. *Geometry and Topology*, 16(1):601–624, 2012.
- [Par13] Hugo Parlier. Kissing numbers for surfaces. *Journal of Topology*, 6(3):777–791, 2013.
- [Par14] Hugo Parlier. Simple closed geodesics and the study of teichmüller spaces. *Handbook of Teichmüller Theory, Volume IV*, pages 113–134, 2014.
- [Sch93] Paul Schmutz. Reimann surfaces with shortest geodesic of maximal length. *Geometric & Functional Analysis GAFA*, 3(6):564–631, 1993.
- [Sch94] Paul Schmutz. Systoles on riemann surfaces. *manuscripta mathematica*, 85(1):429–447, 1994.
- [Sch98] Paul Schaller. Geometry of riemann surfaces based on closed geodesics. *Bulletin of the American Mathematical Society*, 35(3):193–214, 1998.
- [Sch99] Paul Schmutz Schaller. Systoles and topological morse functions for riemann surfaces. *Journal of Differential Geometry*, 52(3):407–452, 1999.
- [Sta15] Richard P Stanley. *Catalan numbers*. Cambridge University Press, 2015.
- [Thu86] William Thurston. A spine for teichmüller space. *preprint*, 1986.

BICMR, PEKING UNIVERSITY, BEIJING 100871, CHINA
Email address: yue-gao@pku.edu.cn