

# UPPER BOUND FOR THE NUMBER OF SPANNING FORESTS OF REGULAR GRAPHS

FERENC BENCs AND PÉTER CSIKVÁRI

ABSTRACT. We show that if  $G$  is a  $d$ -regular graph on  $n$  vertices, then the number of spanning forests  $F(G)$  satisfies  $F(G) \leq d^n$ . The previous best bound due to Kahale and Schulman gave  $(d + 1/2 + O(1/d))^n$ . We also have the more precise conjecture that

$$F(G)^{1/n} \leq \frac{(d-1)^{d-1}}{(d^2 - 2d - 1)^{d/2-1}}.$$

If this conjecture is true, then the expression on the right hand side is the best possible.

## 1. INTRODUCTION

In this paper, we study the number of spanning forests of regular graphs. Let  $\tau(G)$  and  $F(G)$  denote the number of spanning trees and spanning forests, respectively. By a spanning forest  $F$  we mean an acyclic set of edges. The word spanning simply refers to the fact that we always consider  $V(G)$  to be the vertex set of the forest  $F$ .

One expects similar theorems to be true for  $\tau(G)$  and  $F(G)$ , still, it turns out that they are very different in nature. It is known that  $\tau(G)$  can be computed in polynomial time by the matrix-tree theorem [15]. On the other hand,  $F(G)$  is #P-hard to compute [13]. In fact, for studying  $\tau(G)$  one can use the whole machinery of Laplacian matrices and eigenvalues, whereas for  $F(G)$  there is no such toolkit. Nevertheless one expects similar or at least analogous behaviours in certain problems even if we cannot prove them for spanning forests. For instance, it is known [9, 15] that if  $e, f \in E(G)$  are fixed edges of a graph  $G$ , and  $\mathbf{T}$  is a random spanning tree chosen uniformly at random, then

$$\mathbb{P}(e, f \in \mathbf{T}) \leq \mathbb{P}(e \in \mathbf{T})\mathbb{P}(f \in \mathbf{T}).$$

For a uniformly chosen random spanning forest  $\mathbf{F}$  the similar statement

$$\mathbb{P}(e, f \in \mathbf{F}) \leq \mathbb{P}(e \in \mathbf{F})\mathbb{P}(f \in \mathbf{F})$$

is a well-known open problem [11, 18].

A classical result due to McKay gives an upper bound on the number of spanning trees of regular graphs.

**Theorem 1.1** (McKay [17]). *Let  $d \geq 3$  and let  $G$  be a  $d$ -regular graph on  $n$  vertices. There exists an absolute constant  $c_d$  such that*

$$\tau(G) \leq \frac{c_d \ln(n)}{n} \left( \frac{(d-1)^{d-1}}{(d^2 - 2d)^{d/2-1}} \right)^n.$$

In this paper we conjecture the following analogue of this theorem.

---

2010 *Mathematics Subject Classification.* Primary: 05C30. Secondary: 05C31, 05C70.

*Key words and phrases.* forests, Tutte polynomial.

The first author is supported by the NKFIH (National Research, Development and Innovation Office, Hungary) grant KKP-133921. The second author is partially supported by the Counting in Sparse Graphs Lendület Research Group.

**Conjecture 1.2.** Let  $d \geq 3$  and let  $G$  be a  $d$ -regular graph on  $n$  vertices. Then

$$F(G) \leq \left( \frac{(d-1)^{d-1}}{(d^2 - 2d - 1)^{d/2-1}} \right)^n.$$

This conjecture is motivated by the following theorem of the authors.

**Theorem 1.3** (Bencs and Csikvári [2]). *Let  $v(H)$  denote the number of vertices of a graph  $H$ . Let  $F(H)$  denote the number of spanning forests of the graph  $H$ . Furthermore, let  $g(H)$  denote the girth of the graph  $H$ , that is, the length of the shortest cycle. Let  $(G_n)_n$  be a sequence of  $d$ -regular graphs, where  $d \geq 3$ , such that  $\lim_{n \rightarrow \infty} g(G_n) = \infty$ . Then*

$$\lim_{n \rightarrow \infty} F(G_n)^{1/v(G_n)} = \frac{(d-1)^{d-1}}{(d^2 - 2d - 1)^{d/2-1}}.$$

*If  $(G_n)_n$  is a sequence of random  $d$ -regular graphs with growing number of vertices, then the same statement holds true asymptotically almost surely.*

Although we are not able to prove Conjecture 1.2, we can prove the following result.

**Theorem 1.4.** *Let  $d \geq 2$  be an integer. Let  $G$  be a  $d$ -regular graph on  $n$  vertices. Then  $F(G) \leq d^n$ .*

To compare Conjecture 1.2 and Theorem 1.4 we note that

$$\frac{(d-1)^{d-1}}{(d^2 - 2d - 1)^{d/2-1}} = d - \frac{1}{2d} - \frac{1}{3d^2} - \frac{1}{8d^3} + O\left(\frac{1}{d^4}\right).$$

We also note that the previous best bound on  $F(G)$  is due to Kahale and Schulman [14], and they proved that  $F(G) \leq C_d^n$ , where

$$C_d = d + \frac{1}{2} + \frac{1}{8d} + \frac{13}{48d^2} + O\left(\frac{1}{d^3}\right).$$

Thus, the actual improvement in the constant is more than  $1/2$ . Note that Borbényi, Csikvári and Luo [3] were able to improve on the results of Kahale and Schulman for certain values of  $d$ , but even those bounds are far from the result of Theorem 1.4. Let us also mention that we will actually prove a slightly stronger result, namely,

$$F(G)^{1/v(G)} \leq \sqrt{2} \mu_{K_{d+1}} \left( \frac{d+1}{\sqrt{2}} \right)^{1/(d+1)},$$

where  $\mu_G(x)$  is the matching polynomial of a graph  $G$  (see Section 2.1), and  $K_{d+1}$  is the complete graph on  $d+1$  vertices. A considerable work is actually needed to prove that the right hand side is at most  $d$  if  $d \geq 4$ .

One may wish to compare our results to various specific graph classes. One notable special case concerns the number of forests of Archimedean lattices. In this case we recommend the short survey of Chang and Shrock [5].

**Notation.** Throughout the paper  $G = (V, E)$  denotes a graph with vertex set  $V$  and edge set  $E$ . We will denote  $|E(G)|$  by  $e(G)$ , and  $|V(G)|$  by  $v(G)$ . If we fix a graph  $G$ , then we use  $v(G) = n$ . The degree of a vertex  $v$  is the number of neighbors of  $v$ , and is denoted by  $d_v$ .  $K_n$  denotes the complete graph on  $n$  vertices.

**Organization of this paper.** In the next section we study a polynomial that we call  $R_G(z)$ . This polynomial was introduced in paper [2]. It turns out that  $F(G) \leq R_G(2)$ . Since  $R_G(z)$  is very strongly connected to the so-called matching polynomial we also

collect all relevant results about the matching polynomial here. In Section 3 we prove Theorem 1.4. In Section 4 we revisit Conjecture 1.2.

## 2. THE POLYNOMIAL $R_G(z)$

In this section we study a polynomial that is related to the matching polynomial of a graph  $G$ . Recall that a set of edges  $M$  is a matching if no two edges of  $M$  have a common end vertex. A  $k$ -matching is simply a matching of size  $k$ .

**Definition 2.1.** Let

$$R_G(z) = \sum_{M \in \mathcal{M}(G)} (-z)^{|M|} \prod_{v \notin V(M)} (z + d_v - 1),$$

where  $d_v$  is the degree of the vertex  $v$ , and  $\mathcal{M}(G)$  is the set of matchings of  $G$ , including the empty one.

**Remark 2.2.** This polynomial was introduced by the authors in paper [2] in order to study the Tutte polynomial. In this paper we strongly build on [2].

Next we need the concept of pseudo-forests for a better understanding of  $R_G(z)$ .

**Definition 2.3.** We say that an  $A \subseteq E(G)$  is a pseudo-forest of  $G$  if each of its connected components are forests or unicyclic graphs. Let  $\mathcal{PF}(G)$  be the set of pseudo-forests of  $G$ . For a pseudo-forest  $A$  let  $c(A)$  be the number of cycles in  $A$ .

The main result about  $R_G(z)$  is the following characterization in terms of pseudo-forests.

**Theorem 2.4** (Bencs and Csikvári [2]). *For a graph  $G$  we have*

$$R_G(z + 1) = \sum_{k=0}^n \left( \sum_{\substack{A \in \mathcal{PF}(G) \\ |A|=k}} 2^{c(A)} \right) z^{n-k}.$$

Let us introduce the polynomial

$$F_G(z) = \sum_{k=0}^n f_k(G) z^{n-k},$$

where  $f_k(G)$  denotes the number of forests with exactly  $k$  edges. It turns out that  $F_G(z)$  is a specialization of the Tutte polynomial [20]. The Tutte polynomial of a graph  $G$  is defined as follows:

$$T_G(x, y) = \sum_{A \subseteq E(G)} (x - 1)^{k(A) - k(G)} (y - 1)^{k(A) + |A| - |V|},$$

where  $k(A)$  is the number of connected components of the graph with vertex set  $V = V(G)$  and edge set  $A$ , and  $k(G)$  is the number of connected components of the graph  $G$ . For a comprehensive introduction to the Tutte polynomial see the surveys [4, 6, 8, 21].

It is easy to see that

$$F_G(z) = z^{k(G)} T_G(z + 1, 1).$$

Since a forest is a pseudo-forest without any cycle we immediately have

$$F_G(z) \leq R_G(z + 1)$$

for positive  $z$ . In particular,

$$(2.1) \quad F(G) = F_G(1) \leq R_G(2)$$




FIGURE 1. A path-tree from vertex 1.

**2.1. Matching polynomial.** In this section, we collect a few facts about the so-called matching polynomial. The matching polynomial of a graph  $G$  is defined as follows. Let

$$\mu_G(z) = \sum_{k=0}^{n/2} (-1)^k m_k(G) z^{n-2k},$$

where  $m_k(G)$  denotes the number of matchings of size  $k$ . Note that for a  $d$ -regular graph  $G$  we have

$$(2.2) \quad R_G(z) = \sum_{M \in \mathcal{M}(G)} (-z)^{|M|} (d+z-1)^{n-2|M|} = z^{n/2} \mu_G \left( \frac{d-1+z}{\sqrt{z}} \right).$$

The following theorem about the matching polynomial is fundamental for us.

**Theorem 2.5** (Heilmann and Lieb [12]). *All zeros of the matching polynomial  $\mu_G(z)$  are real. Furthermore, if the largest degree  $\Delta$  satisfies  $\Delta \geq 2$ , then all zeros lie in the interval  $(-2\sqrt{\Delta-1}, 2\sqrt{\Delta-1})$ .*

Let  $\mu_G(z) = \prod_{i=1}^n (z - \alpha_i)$ , and  $s_k(G) = \sum_{i=1}^n \alpha_i^k$ . The quantities  $s_k(G)$  have a combinatorial meaning. They count the so-called tree-like walks. Before introducing the concept of tree-like walks, we first require the definition of a path-tree.

**Definition 2.6.** Let  $G$  be graph with a given vertex  $u$ . The *path-tree*  $T(G, u)$  is defined as follows. The vertices of  $T(G, u)$  are the paths in  $G$  which start at the vertex  $u$  and two paths as vertices are joined by an edge if one of them is a one-step extension of the other. See Figure 1.

**Definition 2.7.** A closed tree-like walk of length  $\ell$  is a closed walk on  $T(G, u)$  of length  $\ell$  starting and ending at the vertex  $u$ .

Note that a priori the tree-like walk is on the path-tree, although one can make a correspondence with certain walks on the graph itself. Indeed, a walk in the tree  $T(G, u)$  from  $u$  can be imagined as follows. Suppose that in the graph  $G$  a worm is sitting at vertex  $u$  at the beginning. Then at each step the worm can either grow or pull back its head. When it grows it can move its head to a neighboring unoccupied vertex while keeping its tail at vertex  $u$ . At each step the worm occupies a path in the graph  $G$ . A closed walk in the tree  $T(G, u)$  from  $u$  to  $u$  corresponds to the case when at the final

step the worm occupies only vertex  $u$ . Let  $s_k(G, u)$  be the number of tree-like closed walks of length  $k$  starting and ending at vertex  $u$ . Note that for different vertices  $u$  and  $v$  the quantities  $s_k(G, u)$  and  $s_k(G, v)$  may be different.

**Lemma 2.8** (Godsil [10]). *The number of all closed tree-like walks of length  $\ell$  is exactly  $s_\ell(G)$ , that is,  $s_\ell(G) = \sum_{u \in V(G)} s_\ell(G, u)$ .*

The following inequalities will be useful when we prove Theorem 1.4.

**Lemma 2.9.** *For every  $d$ -regular graph  $G$  and  $k \geq 1$  we have*

$$\frac{s_k(G)}{v(G)} \geq \frac{s_k(K_{d+1})}{v(K_{d+1})}.$$

*Proof.* The path-tree of the complete graph on  $d+1$  vertices is a subtree of the path-tree of any regular graph. Indeed, if we have a path  $P$  in  $G$  consisting of  $t$  vertices, then we can extend it to at least  $d+1-t$  ways to a path on  $t+1$  vertices. In  $K_{d+1}$  we have equality here.

From this it follows that if  $u \in V(G)$ , then  $s_k(G, u) \geq s_k(K_{d+1}, r)$ , where  $r$  is any vertex of  $K_{d+1}$ . Then

$$\frac{s_k(G)}{v(G)} = \frac{1}{v(G)} \sum_{u \in V(G)} s_k(G, u) \geq s_k(K_{d+1}, r) = \frac{s_k(K_{d+1})}{v(K_{d+1})}.$$

□

**Lemma 2.10.** *Let  $G$  be a  $d$ -regular graph with  $v(G)$  vertices, and let  $a > 2\sqrt{d-1}$ . Then we have*

$$\frac{1}{v(G)} \log \mu_G(a) \leq \frac{1}{v(K_{d+1})} \log \mu_{K_{d+1}}(a).$$

*Proof.* For a graph  $G$  we can write the matching polynomial as  $\mu_G(z) = \prod_{i=1}^{v(G)} (z - \alpha_i)$ , where  $|\alpha_i| < 2\sqrt{d-1} < a$  by Theorem 2.5. Then we have

$$\begin{aligned} \frac{1}{v(G)} \log \mu_G(a) &= \frac{1}{v(G)} \sum_{i=1}^{v(G)} \log(a - \alpha_i) = \log(a) + \frac{1}{v(G)} \sum_{i=1}^{v(G)} \log\left(1 - \frac{\alpha_i}{a}\right) = \\ &= \log(a) - \frac{1}{v(G)} \sum_{k=1}^{\infty} \frac{1}{ka^k} \sum_{i=1}^n \alpha_i^k = \log(a) - \sum_{k=1}^{\infty} \frac{1}{ka^k} \frac{s_k(G)}{v(G)}. \end{aligned}$$

The upper bound now follows using that  $\frac{s_k(G)}{v(G)} \geq \frac{s_k(K_{d+1})}{v(K_{d+1})}$ . □

### 3. PROOF OF THEOREM 1.4

In this section we prove Theorem 1.4. Note that the cases  $d = 2$  and  $d = 3$  are trivial since for  $d = 2$

$$F(G) \leq 2^{e(G)} = 2^{dn/2} = 2^n,$$

and for  $d = 3$  we have

$$F(G) \leq 2^{e(G)} = 2^{3n/2} = (2\sqrt{2})^n < 3^n.$$

From now on we can assume that  $d \geq 4$ . By inequality 2.1 and identity 2.2 we have that if  $G$  is a  $d$ -regular graph on  $n$  vertices, then

$$F(G) \leq R_G(2) = 2^{n/2} \mu_G\left(\frac{d+1}{\sqrt{2}}\right).$$

$d$	$\frac{(d-1)^{d-1}}{(d^2-2d-1)^{d/2-1}}$	$\sqrt{2} \cdot \mu_{K_{d+1}} \left( \frac{d+1}{\sqrt{2}} \right)^{1/(d+1)}$	$d - \frac{1}{2d}$
4	3.85714285714286	3.91947904192452	3.87500000000000
5	4.88706270925576	4.91723641784679	4.90000000000000
6	5.90737240075614	5.92330714974640	5.91666666666667
7	6.92165915952326	6.93081266948571	6.92857142857143
8	7.93218265702205	7.93782732697178	7.93750000000000
9	8.94023598867791	8.94392512794576	8.94444444444444
10	9.94659007255980	9.94911825176517	9.95000000000000
11	10.9517282616543	10.9535296122445	10.9545454545455
12	11.9559676088814	11.9572931718296	11.9583333333333
13	12.9595242435679	12.9605264078823	12.9615384615385
14	13.9625503782448	13.9633255787092	13.9642857142857
15	14.9651562263964	14.9657677145412	14.9666666666667
16	15.9674235136022	15.9679140832706	15.9687500000000
17	16.9694141030687	16.9698135007115	16.9705882352941
18	17.9711756732928	17.9715050723952	17.9722222222222
19	18.9727455571083	18.9730203479578	18.9736842105263
20	19.9741533998487	19.9743849792194	19.9750000000000

TABLE 1. Numerical values of the constants appearing in this paper for  $4 \leq d \leq 20$ .

Hence it is enough to prove that if  $d \geq 4$ , then

$$2^{n/2} \mu_G \left( \frac{d+1}{\sqrt{2}} \right) < d^n.$$

First we show that the quantity  $\sqrt{2} \mu_G \left( \frac{d+1}{\sqrt{2}} \right)^{1/v(G)}$  is maximised by the graph  $K_{d+1}$  among  $d$ -regular graphs. Then we show that

$$\sqrt{2} \mu_{K_{d+1}} \left( \frac{d+1}{\sqrt{2}} \right)^{1/(d+1)} < d$$

if  $d \geq 4$ . Let us start with the first inequality.

**Proposition 3.1.** *Let  $d \geq 4$ . For a  $d$ -regular graph  $G$  on  $n$  vertices we have*

$$\sqrt{2} \mu_G \left( \frac{d+1}{\sqrt{2}} \right)^{1/n} \leq \sqrt{2} \mu_{K_{d+1}} \left( \frac{d+1}{\sqrt{2}} \right)^{1/(d+1)}.$$

*Proof.* This follows from Lemma 2.10 since  $\frac{d+1}{\sqrt{2}} > 2\sqrt{d-1}$  for every  $d \geq 4$ .  $\square$

So we have

$$F(G) < \left( \sqrt{2} \mu_{K_{d+1}} \left( \frac{d+1}{\sqrt{2}} \right)^{1/(d+1)} \right)^n$$

for every  $d$ -regular graph  $G$  on  $n$  vertices.

As Table 1 suggests this bound is actually better than  $d - \frac{1}{2d}$  from  $d \geq 9$ , and for all  $d$  it is less than  $d$ . We only prove this latter result.

**Lemma 3.2.** *For  $d \geq 4$  we have*

$$\sqrt{2}\mu_{K_{d+1}} \left( \frac{d+1}{\sqrt{2}} \right)^{1/(d+1)} < d.$$

The following proof is quite long and technical, but completely standard analysis.

*Proof.* It will be convenient to introduce the notation  $n = d + 1$ . Then the required inequality is equivalent with

$$2^{n/2}\mu_{K_n} \left( \frac{n}{\sqrt{2}} \right) < (n-1)^n$$

for  $n \geq 5$ . The plan is the following: we will show that if we divide by  $n^n$ , then

$$\frac{2^{n/2}}{n^n}\mu_{K_n} \left( \frac{n}{\sqrt{2}} \right) = \frac{1}{e} - \frac{1}{en} + O\left(\frac{1}{n^2}\right),$$

and

$$\frac{(n-1)^n}{n^n} = \frac{1}{e} - \frac{1}{2en} + O\left(\frac{1}{n^2}\right),$$

where  $e = 2.71..$  is the base of the natural logarithm. We will prove these statements with quantitative error bounds to determine those numbers  $n$  for which the error term is less than the second main term. For smaller  $n$  one may check the statement with computer.

Note that  $m_k(K_n) = \frac{n!}{2^k k!(n-2k)!}$ , and so

$$\begin{aligned} \frac{2^{n/2}}{n^n}\mu_{K_n} \left( \frac{n}{\sqrt{2}} \right) &= \frac{2^{n/2}}{n^n} \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k m_k(K_n) \left( \frac{n}{\sqrt{2}} \right)^{n-2k} \\ &= \frac{2^{n/2}}{n^n} \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \frac{n!}{2^k k!(n-2k)!} \left( \frac{n}{\sqrt{2}} \right)^{n-2k} \\ &= \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{(-1)^k}{k!} \frac{n(n-1)\dots(n-2k+1)}{n^{2k}} \\ &= \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{(-1)^k}{k!} \prod_{i=0}^{2k-1} \left( 1 - \frac{i}{n} \right). \end{aligned}$$

Similarly,

$$\frac{(n-1)^n}{n^n} = \left( 1 - \frac{1}{n} \right)^n = \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{1}{n^k} = \sum_{k=0}^n \frac{(-1)^k}{k!} \prod_{i=0}^{k-1} \left( 1 - \frac{i}{n} \right).$$

In what follows we will often use the following two observations. If  $b_1 > b_2 > \dots > b_r > 0$  and  $r > m$ , then

$$\left| \sum_{j=m+1}^r (-1)^j b_j \right| \leq b_{m+1}.$$

We also need that if  $a_1, \dots, a_r > 0$  such that  $\sum_{i=1}^r a_i < 1$ , then

$$1 - \sum_{i=1}^r a_i \leq \prod_{i=1}^r (1 - a_i) < 1 - \sum_{i=1}^r a_i + \frac{1}{2} \left( \sum_{i=1}^r a_i \right)^2.$$

Here the lower bound can be proved by induction on  $r$ , while the upper bound follows from

$$\prod_{i=1}^r (1 - a_i) < \prod_{i=1}^r e^{-a_i} = \exp\left(-\sum_{i=1}^r a_i\right) < 1 - \sum_{i=1}^r a_i + \frac{1}{2} \left(\sum_{i=1}^r a_i\right)^2$$

since  $e^{-x} = 1 - x + \frac{x^2}{2} - \frac{x^3}{3!} + \dots$  and  $\sum_{j=2}^{\infty} \left(-\frac{x^{2j-1}}{(2j-1)!} + \frac{x^{2j}}{(2j)!}\right) < 0$  if  $x < 1$ . The sequences  $b_k = \frac{1}{k!} \prod_{i=0}^{2k-1} \left(1 - \frac{i}{n}\right)$  and  $b'_k = \frac{1}{k!} \prod_{i=0}^{k-1} \left(1 - \frac{i}{n}\right)$  are monotone decreasing, so for  $m < \lfloor n/2 \rfloor$  we have

$$\sum_{k=0}^{\lfloor n/2 \rfloor} \frac{(-1)^k}{k!} \prod_{i=0}^{2k-1} \left(1 - \frac{i}{n}\right) = \sum_{k=0}^m \frac{(-1)^k}{k!} \prod_{i=0}^{2k-1} \left(1 - \frac{i}{n}\right) + R_m^{(1)}$$

and

$$\sum_{k=0}^n \frac{(-1)^k}{k!} \prod_{i=0}^{k-1} \left(1 - \frac{i}{n}\right) = \sum_{k=0}^m \frac{(-1)^k}{k!} \prod_{i=0}^{k-1} \left(1 - \frac{i}{n}\right) + r_m^{(1)},$$

where  $|R_m^{(1)}|, |r_m^{(1)}| < \frac{1}{(m+1)!}$ . We will choose  $m$  in such a way that  $\sum_{j=0}^{2m-1} \frac{j}{n} = \frac{\binom{2m}{2}}{n} < 1$ . Then for  $k \leq m$  we have

$$\prod_{i=0}^{2k-1} \left(1 - \frac{i}{n}\right) = 1 - \frac{\binom{2k}{2}}{n} + R_k \quad \text{and} \quad \prod_{i=0}^{k-1} \left(1 - \frac{i}{n}\right) = 1 - \frac{\binom{k}{2}}{n} + r_k,$$

where  $|R_k| \leq \frac{1}{2} \left(\frac{\binom{2k}{2}}{n}\right)^2$  and  $|r_k| \leq \frac{1}{2} \left(\frac{\binom{k}{2}}{n}\right)^2$ . Hence

$$\begin{aligned} \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{(-1)^k}{k!} \prod_{i=0}^{2k-1} \left(1 - \frac{i}{n}\right) &= \sum_{k=0}^m \frac{(-1)^k}{k!} \prod_{i=0}^{2k-1} \left(1 - \frac{i}{n}\right) + R_m^{(1)} \\ &= \sum_{k=0}^m \frac{(-1)^k}{k!} \left(1 - \frac{\binom{2k}{2}}{n} + R_k\right) + R_m^{(1)} \\ &= \sum_{k=0}^m \frac{(-1)^k}{k!} + \frac{1}{n} \sum_{k=0}^m (-1)^k \frac{\binom{2k}{2}}{k!} + \sum_{k=0}^m \frac{(-1)^k}{k!} R_k + R_m^{(1)} \\ &= \sum_{k=0}^m \frac{(-1)^k}{k!} + \frac{1}{n} \left(-\frac{1}{e} + R_m^{(2)}\right) + \sum_{k=0}^m \frac{(-1)^k}{k!} R_k + R_m^{(1)}, \end{aligned}$$

where  $|R_m^{(2)}| \leq \frac{\binom{2(m+1)}{2}}{(m+1)!}$ . Similarly,

$$\begin{aligned} \sum_{k=0}^n \frac{(-1)^k}{k!} \prod_{i=0}^{k-1} \left(1 - \frac{i}{n}\right) &= \sum_{k=0}^m \frac{(-1)^k}{k!} \prod_{i=0}^{k-1} \left(1 - \frac{i}{n}\right) + r_m^{(1)} \\ &= \sum_{k=0}^m \frac{(-1)^k}{k!} \left(1 - \frac{\binom{k}{2}}{n} + r_k\right) + r_m^{(1)} \\ &= \sum_{k=0}^m \frac{(-1)^k}{k!} + \frac{1}{n} \sum_{k=0}^m (-1)^k \frac{\binom{k}{2}}{k!} + \sum_{k=0}^m \frac{(-1)^k}{k!} r_k + r_m^{(1)} \\ &= \sum_{k=0}^m \frac{(-1)^k}{k!} + \frac{1}{n} \left(-\frac{1}{2e} + r_m^{(2)}\right) + \sum_{k=0}^m \frac{(-1)^k}{k!} r_k + r_m^{(1)}, \end{aligned}$$



where  $|r_m^{(2)}| \leq \frac{\binom{m+1}{2}}{(m+1)!}$ . Note that

$$\begin{aligned} & \left| \left( \frac{R_m^{(2)}}{n} + \sum_{k=0}^m \frac{(-1)^k}{k!} R_k + R_m^{(1)} \right) - \left( \frac{r_m^{(2)}}{n} + \sum_{k=0}^m \frac{(-1)^k}{k!} r_k + r_m^{(1)} \right) \right| \\ & \leq \frac{|R_m^{(2)}|}{n} + \sum_{k=0}^m \frac{|R_k|}{k!} + |R_m^{(1)}| + \frac{|r_m^{(2)}|}{n} + \sum_{k=0}^m \frac{r_k}{k!} + |r_m^{(1)}| \\ & \leq \frac{\binom{2(m+1)}{2}}{n \cdot (m+1)!} + \frac{1}{2n^2} \sum_{k=0}^m \frac{\binom{2k}{2}^2}{k!} + \frac{1}{(m+1)!} + \frac{\binom{m+1}{2}}{n \cdot (m+1)!} + \frac{1}{2n^2} \sum_{k=0}^m \frac{\binom{k}{2}^2}{k!} + \frac{1}{(m+1)!} \\ & \leq \frac{5/2 \cdot m^2 + 7/2 \cdot m + 1}{n \cdot (m+1)!} + \frac{175e}{8n^2} + \frac{2}{(m+1)!} \end{aligned}$$

since

$$\sum_{k=0}^{\infty} \frac{\binom{2k}{2}^2}{k!} = 42e \quad \text{and} \quad \sum_{k=0}^m \frac{\binom{k}{2}^2}{k!} = \frac{7e}{4}.$$

In general, if we have a polynomial  $P(x) = \sum_{r=0}^t a_r x(x-1)\dots(x-r+1)$ , then

$$\sum_{k=0}^{\infty} \frac{P(k)}{k!} = \sum_{r=0}^t a_r \sum_{k=r}^{\infty} \frac{1}{(k-r)!} = e \sum_{r=0}^t a_r,$$

and in our case

$$\binom{2k}{2}^2 = 4k(k-1)(k-2)(k-3) + 20k(k-1)(k-2) + 17k(k-1) + k,$$

and

$$\binom{k}{2}^2 = \frac{1}{4}k(k-1)(k-2)(k-3) + k(k-1)(k-2) + \frac{1}{2}k(k-1).$$

Hence

$$\begin{aligned} & \sum_{k=0}^n \frac{(-1)^k}{k!} \prod_{i=0}^{k-1} \left(1 - \frac{i}{n}\right) - \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{(-1)^k}{k!} \prod_{i=0}^{2k-1} \left(1 - \frac{i}{n}\right) \\ & \geq \frac{1}{2e} \cdot \frac{1}{n} - \left( \frac{5/2 \cdot m^2 + 7/2 \cdot m + 1}{n \cdot (m+1)!} + \frac{175e}{8n^2} + \frac{2}{(m+1)!} \right). \end{aligned}$$

Let  $m = \lfloor \sqrt{n/2} \rfloor$ , then  $\binom{2m}{2} < 2m^2 \leq n$ . If  $n \geq 392$ , then  $m \geq 14$ , and we have

$$\frac{5/2 \cdot m^2 + 7/2 \cdot m + 1}{n \cdot (m+1)!} \leq \frac{7m^2}{(m+1)! \cdot n} < \frac{7}{13! \cdot n} < \frac{1}{1000n},$$

and

$$\frac{175e}{8n^2} \leq \frac{175e}{8 \cdot 392 \cdot n} < \frac{1}{6n},$$

and

$$\frac{2}{(m+1)!} \leq \frac{2}{13! \cdot m \cdot (m+1)} \leq \frac{2}{6000 \cdot n/3} = \frac{1}{1000n}.$$

Hence

$$\frac{1}{2e} \cdot \frac{1}{n} - \left( \frac{5/2 \cdot m^2 + 7/2 \cdot m + 1}{n \cdot (m+1)!} + \frac{7e}{4n^2} + \frac{2}{(m+1)!} \right) \geq \left( \frac{1}{2e} - \frac{1}{1000} - \frac{1}{6} - \frac{1}{1000} \right) \frac{1}{n} > 0.$$

So for  $n \geq 392$  we have

$$\frac{2^{n/2}}{n^n} \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k m_k(K_n) \left( \frac{n}{\sqrt{2}} \right)^{n-2k} < \frac{(n-1)^n}{n^n}.$$

The authors checked the cases  $5 \leq n \leq 392$  with a computer. For courtesy we made available the Sagemath code of the computation at [1].  $\square$

#### 4. CONJECTURE 1.2 REVISITED

In this section we revisit Conjecture 1.2. Here we review a result of Borbényi, Csikvári and Luo [3]. They showed that a well-known conjecture mentioned already in the introduction actually implies Conjecture 1.2. This well-known conjecture is about a negative correlation inequality. The precise statement is the following.

**Conjecture 4.1** ([11, 18]). Let  $G$  be a graph and let  $\mathbf{F}$  be a random forest chosen uniformly from all the forests of  $G$ . Let  $e, f \in E(G)$ , then

$$\mathbb{P}(e, f \in \mathbf{F}) \leq \mathbb{P}(e \in \mathbf{F})\mathbb{P}(f \in \mathbf{F}).$$

Assuming Conjecture 4.1 Borbényi, Csikvári and Luo proved a result on forests of 2-covers which then implies Conjecture 1.2.

Recall that a  $k$ -cover (or  $k$ -lift)  $H$  of a graph  $G$  is defined as follows. The vertex set of  $H$  is  $V(H) = V(G) \times \{0, 1, \dots, k-1\}$ , and the edge set of  $H$  is  $E(H) = \cup_{(u,v) \in E(G)} M_{u,v}$ , where  $M_{u,v}$  is a perfect matching between the vertex set  $L_u = \{(u, i) \mid 0 \leq i \leq k-1\}$  and  $L_v = \{(v, i) \mid 0 \leq i \leq k-1\}$ .




FIGURE 2. A 2-lift.

When  $k = 2$  one can encode a 2-lift  $H$  by putting signs on the edges of the graph  $G$ : the  $+$  sign means that we use the matching  $((u, 0), (v, 0)), ((u, 1), (v, 1))$  at the edge  $(u, v)$ , the  $-$  sign means that we use the matching  $((u, 0), (v, 1)), ((u, 1), (v, 0))$  at the edge  $(u, v)$ . For instance, if we put  $+$  signs to every edge, then we simply get the disjoint union  $G \cup G$ , and if we put  $-$  signs everywhere, then the obtained 2-cover  $H$  is simply the tensor product  $G \times K_2$ . Observe that if  $G$  is bipartite, then  $G \cup G = G \times K_2$ , but other 2-covers might differ from  $G \cup G$ .

**Theorem 4.2** (Borbényi, Csikvári and Luo [3]). *Let  $G$  be a graph, and let  $H$  be a 2-cover of  $G$ . If Conjecture 4.1 is true, then we have*

$$F(G \cup G) \leq F(H).$$

Thus,  $F(G)^{1/v(G)} \leq F(H)^{1/v(H)}$ .

There is a nice property of covers that is related to the girth. For every graph  $G$ , there is a sequence of graphs  $(G_n)_n$  such that  $G_0 = G$ ,  $G_k$  is a 2-cover of  $G_{k-1}$ , and  $g(G_k) \rightarrow \infty$ . This is an observation due to Linial [16], his proof is also given in [7]. This observation, Theorem 1.3 and Theorem 4.2 (assuming Conjecture 4.1) together implies that

$$\sup_{G \in \mathcal{G}_d} F(G)^{1/v(G)} = \frac{(d-1)^{d-1}}{(d^2 - 2d - 1)^{d/2-1}},$$

where  $\mathcal{G}_d$  denotes the family of  $d$ -regular graphs.

One might wonder whether an analogue of Theorem 1.4 is true for non-regular graphs too. The most natural guess

$$F(G) \leq \prod_{v \in V(G)} d_v$$

can only be true if we require some minimum degree condition. We speculate that this inequality can be true if the minimum degree is at least 3. Minimum degree 2 is not enough as gluing together  $r$  cycles of length  $k$  at a fixed vertex shows that for the obtained graph  $G$  we have  $F(G) > \prod_{v \in V(G)} d_v$  if  $(2^k - 1)^r > (2r)2^{r(k-1)}$  which is true for every fixed  $k$  and sufficiently large  $r$ . Note that the inequality  $F(G) \leq \prod_{v \in V(G)} (d_v + 1)$  is easily seen to be true for every graph  $G$  (cf. [19]), but gives an exponentially weaker bound for sparse graphs.

**Acknowledgment.** The authors are very grateful to the reviewers for the comments and suggestions, especially for pointing out a computational error in the original manuscript.

#### REFERENCES

- [1] Ferenc Bencs and Péter Csikvári. Supplementary computational results, [https://github.com/bencsf/upper\\_bound\\_spanning\\_forests](https://github.com/bencsf/upper_bound_spanning_forests). 2022.
- [2] Ferenc Bencs and Péter Csikvári. Evaluations of Tutte polynomials of regular graphs. *Journal of Combinatorial Theory, Series B*, 157:500–523, 2022.
- [3] Márton Borbényi, Péter Csikvári, and Haoran Luo. On the number of forests and connected spanning subgraphs. *Graphs and Combinatorics*, 37(6):2655–2678, 2021.
- [4] Thomas Brylawski and James Oxley. The Tutte polynomial and its applications. *Matroid applications*, 40:123–225, 1992.
- [5] Shu-Chiuan Chang and Robert Shrock. Exponential growth constants for spanning forests on Archimedean lattices: Values and comparisons of upper bounds. *International Journal of Modern Physics B*, 35(06):2150085, 2021.
- [6] Henry H. Crapo. The Tutte polynomial. *Aequationes Mathematicae*, 3(3):211–229, 1969.
- [7] Péter Csikvári. Lower matching conjecture, and a new proof of Schrijver’s and Gurvits’s theorems. *Journal of the European Mathematical Society*, 19(6):1811–1844, 2017.
- [8] Joanna A. Ellis-Monaghan and Criel Merino. Graph polynomials and their applications I: The Tutte polynomial. In *Structural analysis of complex networks*, pages 219–255. Springer, 2011.
- [9] Tomás Feder and Milena Mihail. Balanced matroids. In *Proceedings of the twenty-fourth annual ACM symposium on Theory of computing*, pages 26–38, 1992.
- [10] Chris Godsil. *Algebraic combinatorics*, volume 6. CRC Press, 1993.
- [11] Geoffrey R. Grimmett and Stephan N. Winkler. Negative association in uniform forests and connected graphs. *Random Structures & Algorithms*, 24(4):444–460, 2004.
- [12] Ole J. Heilmann and Elliott H. Lieb. Theory of monomer-dimer systems. In *Statistical Mechanics*, pages 45–87. Springer, 1972.
- [13] François Jaeger, Dirk L. Vertigan, and Dominic J. A. Welsh. On the computational complexity of the Jones and Tutte polynomials. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 108, pages 35–53. Cambridge University Press, 1990.
- [14] Nabil Kahale and Leonard J Schulman. Bounds on the chromatic polynomial and on the number of acyclic orientations of a graph. *Combinatorica*, 16(3):383–397, 1996.

- [15] Gustav Kirchhoff. Ueber die Auflösung der Gleichungen, auf welche man bei der Untersuchung der linearen Vertheilung galvanischer Ströme geführt wird. *Annalen der Physik*, 148(12):497–508, 1847.
- [16] Nathan Linial. Lifts of graphs, (talk slides), [http://www.cs.huji.ac.il/~nati/PAPERS/lifts\\_talk.pdf](http://www.cs.huji.ac.il/~nati/PAPERS/lifts_talk.pdf).
- [17] Brendan D. McKay. Spanning trees in regular graphs. *European Journal of Combinatorics*, 4(2):149–160, 1983.
- [18] Robin Pemantle. Towards a theory of negative dependence. *Journal of Mathematical Physics*, 41(3):1371–1390, 2000.
- [19] Carsten Thomassen. Spanning trees and orientations of graphs. *Journal of Combinatorics*, 1(2):101–111, 2010.
- [20] William Thomas Tutte. A contribution to the theory of chromatic polynomials. *Canadian Journal of Mathematics*, 6:80–91, 1954.
- [21] Dominic Welsh. The Tutte polynomial. *Random Structures & Algorithms*, 15(3-4):210–228, 1999.

KORTEWEG DE VRIES INSTITUTE FOR MATHEMATICS, UNIVERSITY OF AMSTERDAM. P.O. BOX 94248 1090 GE AMSTERDAM THE NETHERLANDS.

*Email address:* [ferenc.bencs@gmail.com](mailto:ferenc.bencs@gmail.com)

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, H-1053 BUDAPEST, REÁLTANODA UTCA 13-15

*Email address:* [peter.csikvari@gmail.com](mailto:peter.csikvari@gmail.com)