

Constraint tableaux for two-dimensional fuzzy logics^{*}

Marta Bílková¹[0000-0002-3490-2083], Sabine Frittella²[0000-0003-4736-8614], and Daniil Kozhemiachenko²[0000-0002-1533-8034]

¹ The Czech Academy of Sciences, Institute of Computer Science, Prague
bilkova@cs.cas.cz

² INSA Centre Val de Loire, Univ. Orléans, LIFO EA 4022, France
sabine.frittella@insa-cvl.fr, daniil.kozhemiachenko@insa-cvl.fr

Abstract. We introduce two-dimensional logics based on Łukasiewicz and Gödel logics to formalize paraconsistent fuzzy reasoning. The logics are interpreted on matrices, where the common underlying structure is the bi-lattice (twisted) product of the $[0, 1]$ interval. The first (resp. second) coordinate encodes the positive (resp. negative) information one has about a statement. We propose constraint tableaux that provide a modular framework to address their completeness and complexity.

Keywords: Constraint tableaux · Łukasiewicz logic · Gödel logic · Two-dimensional logics · Paraconsistent logics.

1 Introduction

A two-dimensional treatment of uncertainty. Belnap-Dunn four-valued logic BD [5,10,26], also referred to as First Degree Entailment FDE, provides a logical framework to reason with both incomplete and inconsistent information. In BD, formulas are evaluated on the Belnap-Dunn square (Figure 1, left) where the four values encode the information available about the formula: $\{t, f, b, n\}$ (true, false, both, neither). Hence, b and n correspond to inconsistent and incomplete information respectively. The shift in perspective lies in the the values encoding the information available about the formula rather than the intrinsic truth or falsity of the formula which may never be known. This idea was generalized by introducing the algebraic notion of bilattices by Ginsberg [14] in the context of AI, and studied further in [28,20]. Bilattices contain two lattice orders simultaneously: a truth order, and a knowledge (or an information) order. Belnap-Dunn square, the smallest interlaced bilattice, can be seen as the product bilattice of the two-element lattice where the four values are seen as pairs of classical values which can be naturally interpreted as representing two independent dimensions of information – the positive and the negative one. We can understand them as providing positive and negative support for statements independently.

^{*} The research of Marta Bílková was supported by RVO: 67985807. The research of Sabine Frittella and Daniil Kozhemiachenko was funded by the grant ANR JCJC 2019, project PRELAP (ANR-19-CE48-0006).

Non-standard probabilities [11,21] generalise the notion of independent positive and negative support of a statement in presence of uncertainty. They encode the positive and negative probabilistic information about a statement φ with a couple $p(\varphi) = (p^+(\varphi), p^-(\varphi))$. The maps are such that $p^-(\varphi) = p^+(\neg\varphi)$, p^+ is a monotone map w.r.t. BD entailment relation, and satisfies the import-export axiom $p^+(\varphi \wedge \psi) + p^+(\varphi \vee \psi) = p^+(\varphi) + p^+(\psi)$. Since formulas are interpreted in BD, one cannot prove that $p(\neg\varphi) = 1 - p(\varphi)$, and $p(\varphi) = p(\neg\varphi) = 1$ can be the case when one has contradictory information about φ . We can diagrammatically represent non-standard probabilities on a continuous extension of Belnap-Dunn square (Figure 1, center), which we see as the product bilattice of the unit real interval $[0, 1] \odot [0, 1]$ in subsection 2.1³. We employ expansions of this algebra in subsection 2.2 to provide semantics to two-dimensional fuzzy logics.

A broader motivation. This paper is a part of the project introduced in [6] aiming to develop a modular logical framework for reasoning based on uncertain, incomplete and inconsistent information. We model an agent who builds her beliefs based on probabilistic information aggregated from multiple sources. A convenient framework to formalize such probabilistic reasoning is that of two-layer modal logics, first introduced in [12,19] and further developed in [7,4]. Roughly speaking, the lower layer of events or evidence encodes the information given by the sources, while the upper layer encodes reasoning with the agent's belief based on this information, and the belief modalities connect the two layers. In [6], we have proposed two-layer modal logics to formalise such probabilistic reasoning in a potentially paraconsistent context.

In this article, we study two families of logics suitable for the upper layer, allowing us to reason with graded, probabilistic or not, incomplete and inconsistent information. We aim at a two-dimensional formalism that separates the positive and negative dimensions of information or support not only on the level of evidence, but also on the level of beliefs.

The logics. In some scenarios, it is reasonable to represent agents beliefs as probabilities (e.g. a company reasoning with information based on statistical data). For such cases, we propose logics derived from Łukasiewicz logic [9, Chapter VI]. We choose Łukasiewicz logic as a starting point because it can express arithmetical operation and therefore the (non-standard) probability axioms [19,6].

In other cases, the agent's aggregated belief is not a probability. Indeed, agents may be able to compare their belief on two different statements while not being necessarily able to say exactly to what extent they believe. Just as Łukasiewicz logic can be seen as a logic of measure or quantity, Gödel logic [9, Chapter VII] can be considered a logic of order. In this context it is therefore natural to consider Gödel logic as the starting point.

To comply with the two-dimensionality aim, we define the logics semantically, using expansions of the product bilattice $[0, 1] \odot [0, 1]$ with connectives derived from standard semantics of Łukasiewicz logic or Gödel logic. Two-dimensional

³ In the context of Nelson's paraconsistent logics such product construction has been called twisted product of algebras [29], or twist structures [25, Chapter 8].

treatment of implication is of a particular interest. We consider two possibilities: the first dualizes implication by co-implication, the second understands negative support of an implication as a conjunction of the positive support of the antecedent and the negative support of the consequent. The first option connects to one of Wansing’s logic of [30], namely I_4C_4 , and goes back to bi-intuitionistic logic [15,27], the second option connects to Nelson’s logic $N4$ [24].

Depending on the choice of connectives, and the choice of the set of designated values on the resulting algebra, we encounter both logics which are paraconsistent and logics which are not. Before proceeding further, we need to clarify the notion of paraconsistency. Unless specified otherwise, we construe ‘logic’ as a set of valid formulas, not as sets of valid entailments. Hence, while not all logics considered in the paper lack explosion w.r.t. their entailment \vdash , $p, \neg p \vDash q$, in none of them $(p \wedge \neg p) \rightarrow q$ is valid. It is in this sense that we call the logics presented here ‘paraconsistent’.

Proof theory. Proof theory for many-valued logics is mostly presented in either of the following three forms. Hilbert style axiomatic calculi (cf., e.g. [19,22]); different versions of sequent and hypersequent calculi [17,22]; tableaux and decomposition calculi (cf., e.g. [18] for Lukasiewicz logic and [3] for Gödel logic).

Each of these proof formalisms has its own advantages: Hilbert calculi provide an explicit list of postulates which facilitates the establishing relations between different logics (e.g. whether one logic is an extension of another). The rules of (hyper)sequent calculi provide structural insights into the algebraic properties of the connectives of the given logic. On the other hand, tableaux and decomposition systems are easily automatisable and can be readily used to determine an upper bound on the complexity of the validity and satisfiability problems for the logic in question. Another advantage of the tableaux is that their semantical nature allows for a straightforward formalisation of different entailment relations defined on the same algebra. Since the logics we are going to introduce are hybrids between FDE and Lukasiewicz or Gödel logic, we opt for combining the constraint tableaux framework with the FDE-tableaux by D’Agostino [8].

Structure of the paper. Section 2 presents preliminaries on bilattices and matrices and introduces the logics for a two dimensional treatment of uncertainty and their properties (proofs are in the Appendix). Section 3 presents the constraint tableaux for these logics and discusses their soundness and completeness, and the complexity of the proof search. Section 4 presents further lines of research.

2 The logics for a two-dimensional treatment of uncertainty.

2.1 Preliminaries

First, we describe the algebras we are going to use to interpret the logics. Their construction relies on the standard MV-algebra, and the standard Gödel algebra, which provide the standard semantics of Lukasiewicz and Gödel logic

respectively (we refer the reader to [9, Chapters VI, VII] for a basic exposure to Gödel and Łukasiewicz logics and their standard semantics). In what follows, $[0, 1]$ denotes the real unit interval with its natural order, and $[0, 1]^{\text{op}}$ denotes the interval with the reversed order.

The standard MV-algebra $[0, 1]_{\mathbf{L}} = ([0, 1], 0, \wedge, \vee, \&, \rightarrow_{\mathbf{L}})$ is defined as follows: for all $a, b \in [0, 1]$ the standard operations are given by

$$\begin{aligned} a \wedge b &:= \min(a, b) & a \&b &:= \max(0, a + b - 1) \\ a \vee b &:= \max(a, b) & a \rightarrow_{\mathbf{L}} b &:= \min(1, 1 - a + b) \end{aligned}$$

Moreover, we define the negation $\sim_{\mathbf{L}} a := a \rightarrow_{\mathbf{L}} 0$, the constant $1 := \sim_{\mathbf{L}} 0$, the truncated sum $a \oplus b := \sim_{\mathbf{L}} a \rightarrow_{\mathbf{L}} b$, and the truncated subtraction $a \ominus b := a \& \sim b$.

The MV-algebra $[0, 1]_{\mathbf{L}}^{\text{op}} = ([0, 1]^{\text{op}}, 1, \vee, \wedge, \oplus, \ominus)$ arises turning the standard MV-algebra upside down, and is isomorphic to it. Here, we have $\sim_{\mathbf{L}} a := 1 \ominus a$.

The standard Gödel algebra $[0, 1]_{\mathbf{G}} = ([0, 1], 0, \wedge, \vee, \rightarrow_{\mathbf{G}})$ is defined as follows: for all $a, b \in [0, 1]$, the standard operations are given by $a \wedge b := \min(a, b)$, $a \vee b := \max(a, b)$, and the implication is defined as follows. We at the same time spell out a definition of a co-implication we shall need later on:

$$a \rightarrow_{\mathbf{G}} b = \begin{cases} 1, & \text{if } a \leq b \\ b & \text{else} \end{cases} \quad b \prec_{\mathbf{G}} a = \begin{cases} 0, & \text{if } b \leq a \\ b & \text{else} \end{cases}$$

We define a negation $\sim_{\mathbf{G}} a := a \rightarrow_{\mathbf{G}} 0$, and $1 := \sim_{\mathbf{G}} 0$.

The algebra $[0, 1]_{\mathbf{G}}^{\text{op}} = ([0, 1]^{\text{op}}, 1, \vee, \wedge, \prec_{\mathbf{G}})$ arises by dualizing the standard Gödel algebra (in particular, similarly as $\rightarrow_{\mathbf{G}}$ is the residuum of \wedge , $\prec_{\mathbf{G}}$ is the residuum of \vee). A negation can be defined on this algebra as $\sim_{\mathbf{G}} a := 1 \prec_{\mathbf{G}} a$.

Product bilattices Given an arbitrary lattice $\mathbf{L} = (L, \wedge_L, \vee_L)$, we can construct the *product bilattice* $\mathbf{L} \odot \mathbf{L} = (L \times L, \wedge, \vee, \sqcap, \sqcup, \neg)$ [1,2]. In what follows, we essentially use the product bilattice $[0, 1] \odot [0, 1]$, constructed from the lattice $([0, 1], \min, \max)$. We only consider the $\{\wedge, \vee, \neg\}$ reduct of this structure in this paper, and not to complicate notation denote it by $[0, 1] \odot [0, 1]$. It is defined as follows: for all $(a_1, a_2), (b_1, b_2) \in [0, 1] \times [0, 1]$,

$$\begin{aligned} (a_1, a_2) \leq (b_1, b_2) &:= a_1 \leq b_1 \wedge a_2 \leq b_2 \\ \neg(a_1, a_2) &:= (a_2, a_1) \\ (a_1, a_2) \wedge (b_1, b_2) &:= (\min(a_1, b_1), \max(a_2, b_2)) \\ (a_1, a_2) \vee (b_1, b_2) &:= (\max(a_1, b_1), \min(a_2, b_2)). \end{aligned}$$

We use expansions of $[0, 1] \odot [0, 1]$ by implication connectives derived from the Łukasiewicz or Gödel implication described above. There are two possibilities we consider. The first dualizes the implication by the co-implication, the second results in negating implication by the conjunction of the positive part of the antecedent and the negative part of the consequent. For Łukasiewicz logics these result in:

$$(a_1, a_2) \rightarrow (b_1, b_2) := (a_1 \rightarrow_{\mathbf{L}} b_1, b_2 \ominus a_2) \quad (a_1, a_2) \rightarrow (b_1, b_2) := (a_1 \rightarrow_{\mathbf{L}} b_1, a_1 \& b_2)$$

For Gödel logics we obtain:

$$(a_1, a_2) \rightarrow (b_1, b_2) := (a_1 \rightarrow_{\mathbf{G}} b_1, b_2 \prec_{\mathbf{G}} a_2) \quad (a_1, a_2) \rightarrow (b_1, b_2) := (a_1 \rightarrow_{\mathbf{G}} b_1, a_1 \wedge b_2)$$

In the first option, the interpretation arises as the one on the product algebra $[0, 1]_{\mathbf{L}} \times [0, 1]_{\mathbf{L}}^{\text{op}}$ or $[0, 1]_{\mathbf{G}} \times [0, 1]_{\mathbf{G}}^{\text{op}}$. In the Gödel case, it relates how the implication is interpreted in Wansing's logic I_4C_4 [30]. In the second options, \rightarrow is not congruential, and a strong congruential implication can be defined as $(a \rightarrow b) \wedge (\neg b \rightarrow \neg a)$. The second option corresponds to how implication is interpreted in product residuated bilattices of [20]. In the Gödel case, it relates to how the implication is interpreted in Nelson's logic N4 [24].

- We denote by $[0, 1]_{\mathbf{L}} \odot [0, 1]_{\mathbf{L}}(\rightarrow)$ and $[0, 1]_{\mathbf{L}} \odot [0, 1]_{\mathbf{L}}(\rightarrow)$ the corresponding expansions of $[0, 1] \odot [0, 1]$ defined using the Łukasiewicz connectives.
- We denote by $[0, 1]_{\mathbf{G}} \odot [0, 1]_{\mathbf{G}}(\rightarrow)$ and $[0, 1]_{\mathbf{G}} \odot [0, 1]_{\mathbf{G}}(\rightarrow)$ the corresponding expansions of $[0, 1] \odot [0, 1]$ defined using the Gödel connectives.

2.2 The logics

The logics considered in this paper are defined through matrix semantics [13]. We consider logical matrices of the form (\mathbf{A}, D) where \mathbf{A} is one of the four algebras described above, and $D \subseteq A$ is a set of designated values. As sets of designated values, we use various lattice filters of the form $(x, y)^\dagger := \{(x', y') \mid x \leq x' \text{ and } y' \leq y\}$ (see Figure 1, center). The motivation is the following: x represents the threshold of having enough evidence to say there is reasonable evidence supporting the truth of the formula, while y represents the threshold below which one considers not to have enough evidence to say that there is reasonable evidence supporting the falsity of the formula. Of particular interest are filters $(1, 0)^\dagger$ ('exactly certain') and $(1, 1)^\dagger$ ('positively certain').




Fig. 1. Belnap-Dunn square $\mathbf{4}$ (left), its continuous probabilistic extension with the filter $(x, y)^\dagger$ (center) and the geometric interpretation of \neg , \sim and $\neg\sim$ for \mathbf{L}^2 (right).

Each logical matrix determines the set of valid formulas in a given language (formulas, which are, for each valuation, designated), and a consequence relation (an entailment) between sets of formulas and formulas, defined as preservation of designated values. Regarding the two types of implication introduced in the previous subsection, the standard semantics of the logics is set as follows:

- Logics with the \rightarrow implication in the language (i.e. the logics $\mathbf{L}_{(x,y)}^2(\rightarrow)$ and $\mathbf{G}_{(x,y)}^2(\rightarrow)$ below) are given by the matrices $([0, 1]_{\mathbf{L}} \odot [0, 1]_{\mathbf{L}}(\rightarrow), (x, y)^\dagger)$ and $([0, 1]_{\mathbf{G}} \odot [0, 1]_{\mathbf{G}}(\rightarrow), (x, y)^\dagger)$ respectively.

- Logics with the \rightarrow implication in the language (i.e. the logics $\mathbb{L}_{(x,y)}^2(\rightarrow)$ and $\mathbb{G}_{(x,y)}^2(\rightarrow)$ below) are given by the matrices $([0, 1]_{\mathbb{L}} \odot [0, 1]_{\mathbb{L}}(\rightarrow), (x, y)^\uparrow)$ and $([0, 1]_{\mathbb{G}} \odot [0, 1]_{\mathbb{G}}(\rightarrow), (x, y)^\uparrow)$ respectively.

We however need to treat Łukasiewicz and Gödel logics separately. Therefore it is practical to define the language and semantics for them separately in a compact way as follows. We refer by \mathbb{L}^2 to Łukasiewicz logics, and by \mathbb{G}^2 to Gödel logics, specifying the filter in the subscript.

Definition 1 (Language and semantics of \mathbb{L}^2). *We fix a countable set Prop of propositional letters and consider the following language:*

$$\phi := \mathbf{0} \mid p \mid \neg\phi \mid (\phi \wedge \phi) \mid (\phi \vee \phi) \mid (\phi \rightarrow \phi) \mid (\phi \rightarrow \phi)$$

where $p \in \text{Prop}$. We define $\sim\phi := \phi \rightarrow \mathbf{0}$, $\sim_w\phi := \phi \rightarrow \mathbf{0}$, $\phi_1 \odot \phi_2 := \sim(\phi_1 \rightarrow \sim\phi_2)$, and $\phi_1 \leftrightarrow \phi_2 := (\phi_1 \rightarrow \phi_2) \odot (\phi_2 \rightarrow \phi_1)$.

Let $v : \text{Prop} \rightarrow [0, 1] \times [0, 1]$, and denote v_1 and v_2 its left and right coordinates, respectively. We extend v as follows.

$$\begin{aligned} v(\mathbf{0}) &= (0, 1) & v(\phi_1 \rightarrow \phi_2) &= (v_1(\phi_1) \rightarrow_{\mathbb{L}} v_1(\phi_2), v_2(\phi_2) \ominus v_2(\phi_1)) \\ v(\neg\phi) &= (v_2(\phi), v_1(\phi)) & v(\phi_1 \rightarrow \phi_2) &= (v_1(\phi_1) \rightarrow_{\mathbb{L}} v_1(\phi_2), v_1(\phi_1) \& v_2(\phi_2)) \\ & & v(\phi_1 \wedge \phi_2) &= (v_1(\phi_1) \wedge v_1(\phi_2), v_2(\phi_1) \vee v_2(\phi_2)) \\ & & v(\phi_1 \vee \phi_2) &= (v_1(\phi_1) \vee v_1(\phi_2), v_2(\phi_1) \wedge v_2(\phi_2)) \end{aligned}$$

Notice that

$$\begin{aligned} v(\sim\phi) &= (1 - v_1(\phi), 1 - v_2(\phi)) \\ v(\phi_1 \odot \phi_2) &= (v_1(\phi_1) \& v_1(\phi_2), v_2(\phi_1) \oplus v_2(\phi_2)) \\ v(\phi_1 \leftrightarrow \phi_2) &= (1 - |v_1(\phi_1) - v_1(\phi_2)|, |v_2(\phi_1) - v_2(\phi_2)|) \end{aligned}$$

Remark 1. In section 2.3, we use the fact that \neg corresponds to a symmetry w.r.t. the horizontal axis, \sim to a symmetry w.r.t. the point $(0.5, 0.5)$, $\sim\neg$ and $\neg\sim$ are correspond to a symmetry w.r.t. the horizontal axis (see Figure 1, right).

Definition 2 (Language and semantics of \mathbb{G}^2). *We fix a countable set Prop of propositional letters and consider the following language:*

$$\phi := \mathbf{0} \mid \mathbf{1} \mid p \mid \neg\phi \mid (\phi \wedge \phi) \mid (\phi \vee \phi) \mid (\phi \rightarrow \phi) \mid (\phi \prec \phi) \mid (\phi \rightarrow \phi)$$

where $p \in \text{Prop}$. We define $\sim\phi := \phi \rightarrow \mathbf{0}$, and $\sim_w\phi := \phi \rightarrow \mathbf{0}$.

Let $v : \text{Prop} \rightarrow [0, 1] \times [0, 1]$, and denote v_1 and v_2 its left and right coordinates, respectively. We extend v as follows.

$$\begin{aligned} v(\mathbf{0}) &= (0, 1) & v(\phi_1 \wedge \phi_2) &= (v_1(\phi_1) \wedge v_1(\phi_2), v_2(\phi_1) \vee v_2(\phi_2)) \\ v(\mathbf{1}) &= (1, 0) & v(\phi_1 \vee \phi_2) &= (v_1(\phi_1) \vee v_1(\phi_2), v_2(\phi_1) \wedge v_2(\phi_2)) \\ v(\neg\phi) &= (v_2(\phi), v_1(\phi)) & v(\phi_1 \rightarrow \phi_2) &= (v_1(\phi_1) \rightarrow_{\mathbb{G}} v_1(\phi_2), v_2(\phi_2) \prec_{\mathbb{G}} v_2(\phi_1)) \\ & & v(\phi_1 \rightarrow \phi_2) &= (v_1(\phi_1) \rightarrow_{\mathbb{G}} v_1(\phi_2), v_1(\phi_1) \wedge v_2(\phi_2)) \end{aligned}$$

Definition 3 (Validity and consequence). *Let ϕ be a formula and Γ a set of formulas of \mathbb{L}^2 (resp. \mathbb{G}^2) and $v[\Gamma] := \{v(\gamma) \mid \gamma \in \Gamma\}$.*

- ϕ is $\mathbb{L}_{(x,y)}^2$ -valid (resp. $\mathbb{G}_{(x,y)}^2$ -valid) iff $\forall v, v(\phi) \geq (x, y)$.
- $\Gamma \models_{\mathbb{L}_{(x,y)}^2} \phi$ (resp. $\Gamma \models_{\mathbb{G}_{(x,y)}^2} \phi$) iff $\forall v, (v[\Gamma] \subseteq (x, y)^\uparrow)$ implies $v(\phi) \geq (x, y)$.

Convention 1 *We introduce the following notation.*

- $\mathbf{L}_{(x,y)}^2(\rightarrow)$ stands for the $\mathbf{L}_{(x,y)}^2$ logics over $\{\mathbf{0}, \neg, \wedge, \vee, \rightarrow\}$.
- $\mathbf{G}_{(x,y)}^2(\rightarrow)$ stands for the $\mathbf{G}_{(x,y)}^2$ logics over $\{\mathbf{0}, \mathbf{1}, \neg, \wedge, \vee, \rightarrow, \prec\}$.
- $\mathbf{L}_{(x,y)}^2(\rightarrow)$ stands for the $\mathbf{L}_{(x,y)}^2$ logics over $\{\mathbf{0}, \neg, \wedge, \vee, \rightarrow\}$.
- $\mathbf{G}_{(x,y)}^2(\rightarrow)$ stands for the $\mathbf{G}_{(x,y)}^2$ logics over $\{\mathbf{0}, \mathbf{1}, \neg, \wedge, \vee, \rightarrow\}$.

We note that $\mathbf{0}$ of $\mathbf{L}_{(x,y)}^2(\rightarrow)$ can be defined as $\neg(p \rightarrow p)$. However, since there is no definition of $\mathbf{0}$ using \rightarrow , we leave it in both languages for the sake of preserving the same tableau rules for all logics. Likewise, although $\mathbf{0}$ and $\mathbf{1}$ are definable in $\mathbf{G}_{(x,y)}^2(\rightarrow)$, their presence in the language simplifies the proofs of their semantical properties (cf. Propositions 6 and 7).

Remark 2. Let ϕ be a formula over $\{0, \wedge, \vee, \supset\}$. Denote ϕ^\bullet the formula obtained from it by substituting \supset for \rightarrow , and ϕ° by substituting \supset for \rightarrow . Since v_1 's behave precisely like the valuations in Lukasiewicz (Gödel) logic, one can see that ϕ is L-valid (G-valid) iff ϕ^\bullet is $\mathbf{L}_{(1,0)}^2(\rightarrow)$ -valid ($\mathbf{G}_{(1,0)}^2(\rightarrow)$ -valid). Furthermore, ϕ is L-valid (G-valid) iff ϕ° is $\mathbf{L}_{(1,1)}^2(\rightarrow)$ -valid ($\mathbf{G}_{(1,1)}^2(\rightarrow)$ -valid). Thus, $\mathbf{L}_{(1,1)}^2(\rightarrow)$ and $\mathbf{L}_{(1,1)}^2(\rightarrow)$ are conservative extensions of L while $\mathbf{G}_{(1,1)}^2(\rightarrow)$ and $\mathbf{G}_{(1,1)}^2(\rightarrow)$ are conservative extensions of G.

Proposition 1. *Let ϕ be a $\mathbf{L}_{(x,y)}^2(\rightarrow)$ (resp. $\mathbf{G}_{(x,y)}^2(\rightarrow)$) formula without constants and $(1, 1) \notin (x, y)^\uparrow$. Then ϕ is not $\mathbf{L}_{(x,y)}^2(\rightarrow)$ (resp. $\mathbf{G}_{(x,y)}^2(\rightarrow)$) valid.*

Because of this, we are going to consider only $\mathbf{L}_{(x,y)}^2(\rightarrow)$'s and $\mathbf{G}_{(x,y)}^2(\rightarrow)$'s whose sets of designated values extend $(1, 1)^\uparrow$, that is $\mathbf{L}_{(x,1)}^2(\rightarrow)$ and $\mathbf{G}_{(x,1)}^2(\rightarrow)$. In the remainder of the article $\phi, \varphi, \chi, \psi$ denote formulas. Unless there is some ambiguity, we do not specify to which language they belong.

2.3 Semantical properties of $\mathbf{L}_{(x,y)}^2(\rightarrow)$

In this section, we are going to explore how the choice of $(x, y)^\uparrow$ affects the set of $\mathbf{L}_{(x,y)}^2(\rightarrow)$ -valid formulas. In particular, we are providing families of formulas differentiating different $\mathbf{L}_{(x,y)}^2(\rightarrow)$ -validities.

Definition 4 (Closure under conflation). *We say that a filter D of $[0, 1] \odot [0, 1]$ is closed under conflation if for any $(x, y) \in D$, we have $(1 - y, 1 - x) \in D$.*

In bilattices, the negation \neg corresponds to a symmetry w.r.t. the horizontal axis and the conflation corresponds to a symmetry w.r.t. the vertical axis. Notice that a filter $(x, y)^\uparrow$ is closed under conflation iff $y = 1 - x$. In $\mathbf{L}_{(x,y)}^2(\rightarrow)$, conflation can be defined as $\neg\sim$ or equivalently $\sim\neg$ (cf. Figure 1, right).

Proposition 2.

- Let $y \geq 1 - x$. Then ϕ is $\mathbf{L}_{(x,y)}^2(\rightarrow)$ -valid iff ϕ is $\mathbf{L}_{(x,1-x)}^2(\rightarrow)$ -valid.
- Let $y < 1 - x$. Then ϕ is $\mathbf{L}_{(x,y)}^2(\rightarrow)$ -valid iff ϕ is $\mathbf{L}_{(1-y,y)}^2(\rightarrow)$ -valid.

The following statements show that by choosing different sets of designated values, we can alter the sets of tautologies.

Proposition 3. *Let $m, n \in \{2, 3, \dots\}$. Then $L^2_{(\frac{m-1}{m}, \frac{1}{m})} \subsetneq L^2_{(\frac{n-1}{n}, \frac{1}{n})}$ iff $m > n$.*

Note, however, that while $L_{(\frac{1}{2}, \frac{1}{2})}$ validates $p \vee \sim p$, it does not collapse into classical logic as the following propositions show.

Proposition 4. *Let $m, n \in \{3, 4, \dots\}$. Then $L^2_{(\frac{m-2}{2m}, \frac{m+2}{2m})} \subsetneq L^2_{(\frac{n-2}{2n}, \frac{n+2}{2n})}$ iff $m > n$.*

We end this subsection by noting that all $L^2_{(x,y)}(\rightarrow)$'s where $(x, y)^\uparrow$ is prime are paraconsistent in the following sense: $p, \neg p \not\vdash_{L^2_{(x,y)}(\rightarrow)} q$. Furthermore, if $(\frac{1}{2}, \frac{1}{2})^\uparrow \in (x, y)^\uparrow$, the logic is paraconsistent even w.r.t. \sim since $p, \sim p \not\vdash_{L^2_{(x,y)}(\rightarrow)} q$. Last but not least, most $L^2_{(x,y)}(\rightarrow)$'s are not closed under modus ponens.

Proposition 5. *Let $L^2_{(1,0)}(\rightarrow) \subsetneq L^2_{(x,y)}(\rightarrow)$. Then $L^2_{(x,y)}(\rightarrow)$ is not closed under modus ponens.*

2.4 Semantical properties of $G^2(\rightarrow)$

In this section, we show that all $G^2_{(x,y)}(\rightarrow)$ logics have the same set of valid formulas. This means that just as the original Gödel logic, $G^2(\rightarrow)$ can be seen as the logic of comparative truth. Furthermore, the presence of the second dimension allows to interpret $G^2(\rightarrow)$ as the logic of comparative truth and falsehood.

Proposition 6. *Let ϕ be a formula over $\{\mathbf{0}, \mathbf{1}, \neg, \wedge, \vee, \rightarrow, \prec\}$. For any $v(p) = (x, y)$ let $v^*(p) = (1 - y, 1 - x)$. Then $v(\phi) = (x, y)$ iff $v^*(\phi) = (1 - y, 1 - x)$.*

Proposition 7. *Let ϕ be a formula over $\{\mathbf{0}, \mathbf{1}, \neg, \wedge, \vee, \rightarrow, \prec\}$ such that $v(\phi) \geq (x, y)$ for any v and some fixed $(x, y) \neq (0, 1)$. Then $v'(\phi) = (1, 0)$ for any v' .*

The last three propositions show that in contrast to $L^2(\rightarrow)$, the changing of the set of designated values does not change the set of valid formulas as long as the set remains a filter⁴ on $[0, 1] \odot [0, 1]$ generated by a single point. However, while the sets of tautologies remain the same, the entailment relation can be made paraconsistent. Indeed, it suffices to choose any prime $(x, y)^\uparrow$ and the entailment ceases to be explosive in the following sense: $p, \neg p \not\vdash_{G^2_{(x,y)}} q$.

Furthermore, the propositions have an important corollary which simplifies the construction of the tableaux proofs.

Corollary 1. *$v(\phi) = (1, 0)$ for any v iff $v'_1(\phi) = 1$ for any v' .*

3 Tableaux

First, we give a general definition of a constraint tableaux, then in sections 3.1 and 3.2, we introduce tableaux for L^2 's and G^2 's.

⁴ Notice that $p \vee \neg p$ would be valid for $D = [0, 1] \odot [0, 1] \setminus \{(0, 1)\}$. But D is not a filter.

Definition 5 (Constraint tableaux). Let Label be a set of labels and \mathcal{L} a set of formulas. A constraint is one of these three expressions:

- Labelled formulas of the form $L : \phi$ with $L \in \text{Label}$ and $\phi \in \mathcal{L}$,
- Numerical constraints of the form $c \leq d$ or $c < d$ with $c, d \in [0, 1]$,
- Formulaic constraints of the form $L : \phi \leq L' : \phi'$ or $L : \phi < L' : \phi'$ with $L, L' \in \text{Label}$ and $\phi, \phi' \in \mathcal{L}$.

A constraint tableau is a downward branching tree each branch of which is a non-empty set of constraints. Each branch \mathcal{B} can be extended by applications of a given set of rules. If no rule application adds new entries to \mathcal{B} , it is called complete.

Henceforth, we only state the rules and the closure conditions for the branches. In what follows we identify a branch with the set of constraints that appear at some point on the branch.

Since our logics are hybrids between FDE and L (or G), we can combine the constraint tableaux framework with the FDE-tableaux by D'Agostino [8]. In particular, it means that we use two kinds of labelled formulas and formulaic constraints: those that concern the left coordinate (evidence for the statement) and those that concern the right coordinate (evidence against the statement).

3.1 Constraint tableaux for L^2

Definition 6 (Constraint tableau for L^2 — $\mathcal{T}(L^2_{(x,y)})$). Branches contain labelled formulas of the form $\phi \leq_1 i$, $\phi \leq_2 i$, $\phi \geq_1 i$, or $\phi \geq_2 i$, and numerical constraints of the form $i \leq j$ with $i, j \in [0, 1]$. We call atomic labelled formulas labelled formulas where $\phi \in \text{Prop}$.

Each branch can be extended by an application of one of the rules in Figure 2 where $i, j \in [0, 1]$. Let i 's be in $[0, 1]$ and x 's be variables ranging over the real interval $[0, 1]$. We define the translation τ from labelled formulas to linear inequalities as follows:

$$\tau(\phi \leq_1 i) = x_\phi^L \leq i, \quad \tau(\phi \geq_1 i) = x_\phi^L \geq i, \quad \tau(\phi \leq_2 i) = x_\phi^R \leq i, \quad \tau(\phi \geq_2 i) = x_\phi^R \geq i.$$

A tableau branch

$$\mathcal{B} = \{\phi_1 \circ i_1, \dots, \phi_m \circ i_m, \phi'_1 \bullet j_1, \dots, \phi'_n \bullet j_n, k_1 \leq l_1, \dots, k_q \leq l_q\}$$

is closed if the system of inequalities

$$\tau(\phi_1 \circ i_1), \dots, \tau(\phi_m \circ i_m), \tau(\phi'_1 \bullet j_1), \dots, \tau(\phi'_n \bullet j_n), k_1 \leq l_1, \dots, k_q \leq l_q$$

does not have solutions. Otherwise, \mathcal{B} is open. A tableau is closed if all its branches are closed.

ϕ has a $\mathcal{T}(L^2_{(x,y)})$ proof if the tableaux beginning with $\{\phi \leq_1 c, c < x\}$ and $\{\phi \geq_2 d, d > y\}$ are both closed.

Remark 3 (How to interpret the rules of $\mathcal{T}(L^2_{(x,y)})$?). Consider for instance the rule $\rightarrow \leq_2$. It's meaning is: $v_2(\phi_1 \rightarrow \phi_2) \leq i$ iff there is $j \in [0, 1]$ s.t. $v_2(\phi_1) \geq j$ and $v_2(\phi_2) \leq i + j$. While rule $\wedge \leq_1$ means $v_1(\phi_1 \wedge \phi_2) \leq i$ iff either $v_1(\phi_1) \leq i$ or $v_1(\phi_2) \leq i$.

$$\begin{array}{cccc}
\mathbf{0} \leq_1 \frac{\mathbf{0} \leq_1 i}{0 \leq i} & \mathbf{0} \leq_2 \frac{\mathbf{0} \leq_2 i}{1 \leq i} & \mathbf{0} \geq_1 \frac{\mathbf{0} \geq_1 i}{0 \geq i} & \mathbf{0} \geq_2 \frac{\mathbf{0} \geq_2 i}{1 \geq i} \\
\neg \leq_1 \frac{\neg \phi \leq_1 i}{\phi \leq_2 i} & \neg \leq_2 \frac{\neg \phi \leq_2 i}{\phi \leq_1 i} & \neg \geq_1 \frac{\neg \phi \geq_1 i}{\phi \geq_2 i} & \neg \geq_2 \frac{\neg \phi \geq_2 i}{\phi \geq_1 i} \\
\rightarrow \leq_1 \frac{\phi_1 \rightarrow \phi_2 \leq_1 i}{i \geq 1 \left| \begin{array}{l} \phi_1 \geq_1 1 - i + j \\ \phi_2 \leq_1 j \\ j \leq i \end{array} \right.} & \rightarrow \leq_2 \frac{\phi_1 \rightarrow \phi_2 \leq_2 i}{\phi_1 \geq_2 j \\ \phi_2 \leq_2 i + j} & \rightarrow \geq_1 \frac{\phi_1 \rightarrow \phi_2 \geq_1 i}{\phi_1 \leq_1 1 - i + j \\ \phi_2 \geq_1 j} & \rightarrow \geq_2 \frac{\phi_1 \rightarrow \phi_2 \geq_2 i}{i \leq 0 \left| \begin{array}{l} \phi_1 \leq_2 j \\ \phi_2 \geq_2 i + j \\ j \leq 1 - i \end{array} \right.} \\
\rightarrow \leq_1 \frac{\phi_1 \rightarrow \phi_2 \leq_1 i}{i \geq 1 \left| \begin{array}{l} \phi_1 \geq_1 1 - i + j \\ \phi_2 \leq_1 j \\ j \leq i \end{array} \right.} & \rightarrow \leq_2 \frac{\phi_1 \rightarrow \phi_2 \leq_2 i}{\phi_1 \leq_2 i + j \\ \phi_2 \leq_1 1 - j} & \rightarrow \geq_1 \frac{\phi_1 \rightarrow \phi_2 \geq_1 i}{\phi_1 \leq_1 1 - i + j \\ \phi_2 \geq_1 j} & \rightarrow \geq_2 \frac{\phi_1 \rightarrow \phi_2 \geq_2 i}{i \leq 0 \left| \begin{array}{l} \phi_1 \geq_2 i + j \\ \phi_2 \geq_1 1 - j \\ j \leq 1 - i \end{array} \right.} \\
\wedge \leq_1 \frac{\phi_1 \wedge \phi_2 \leq_1 i}{\phi_1 \leq_1 i \mid \phi_2 \leq_1 i} & \wedge \leq_2 \frac{\phi_1 \wedge \phi_2 \leq_2 i}{\phi_1 \leq_2 i \\ \phi_2 \leq_2 i} & \wedge \geq_1 \frac{\phi_1 \wedge \phi_2 \geq_1 i}{\phi_1 \geq_1 i \\ \phi_2 \geq_1 i} & \wedge \geq_2 \frac{\phi_1 \wedge \phi_2 \geq_2 i}{\phi_1 \geq_2 i \mid \phi_2 \geq_2 i} \\
\vee \leq_1 \frac{\phi_1 \vee \phi_2 \leq_1 i}{\phi_1 \leq_1 i \\ \phi_2 \leq_1 i} & \vee \leq_2 \frac{\phi_1 \vee \phi_2 \leq_2 i}{\phi_1 \leq_2 i \mid \phi_2 \leq_2 i} & \vee \geq_1 \frac{\phi_1 \vee \phi_2 \geq_1 i}{\phi_1 \geq_1 i \mid \phi_2 \geq_1 i} & \vee \geq_2 \frac{\phi_1 \vee \phi_2 \geq_2 i}{\phi_1 \geq_2 i \\ \phi_2 \geq_2 i}
\end{array}$$

Fig. 2. Rules of $\mathcal{T}(\mathbf{I}_{(x,y)}^2)$. Vertical bars denote splitting of the branch.

To prove completeness and soundness, we need the following definitions.

Definition 7 (Satisfying valuation of a branch). *Let v be a valuation and $k \in \{1, 2\}$. v satisfies a labelled formula $\phi \leq_k i$ (resp. $\phi \geq_k i$) iff $v_k(\phi) \leq i$ (resp. $v_k(\phi) \geq i$). v satisfies a branch \mathcal{B} iff v satisfies any labelled formula \mathcal{B} . A branch \mathcal{B} is satisfiable iff there is a valuation which satisfies it.*

Theorem 1 (Soundness and completeness). *ϕ is $L_{(x,y)}^2(\rightarrow)$ -valid (resp. $L_{(x,y)}^2(\rightarrow)$ -valid) iff there is a $\mathcal{T}(L_{(x,y)}^2)$ proof for it.*

Proof. The soundness follows from the fact that no closed branch is realisable and that if a premise of the rule is realisable, then all labelled formulas are satisfied in at least one of the conclusions.

To show completeness, we proceed by contraposition. We need to show that complete open branches are satisfiable.

Assume that \mathcal{B} is a complete open branch. We construct the satisfying valuation as follows. Let $*$ $\in \{\leq_1, \geq_1, \leq_2, \geq_2\}$ and p_1, \dots, p_m be the propositional variables appearing in the atomic labelled formulas in \mathcal{B} . Let $\{p_1 * i_1, \dots, p_m * i_n\}$ and $\{k_1 \leq l_1, \dots, k_q \leq l_q\}$ be the sets of all atomic labelled formulas and all numerical constraints in \mathcal{B} . Notice that one variable might appear in many atomic labelled formulas, hence we might have $m \neq n$. Since \mathcal{B} is complete and open, the following system of linear inequalities over the set of variables $\{x_{p_1}^L, x_{p_1}^R, \dots, x_{p_m}^L, x_{p_m}^R\}$ must have at least one solution under the constraints listed:

$$\tau(p_1 * i_1), \dots, \tau(p_m * i_n), k_1 \leq l_1, \dots, k_q \leq l_q.$$

Let $c = (c_1^L, c_1^R, \dots, c_m^L, c_m^R)$ be a solution to the above system of inequalities such that c_j^L (resp. c_j^R) is the value of $x_{p_j}^L$ (resp. $x_{p_j}^R$). Define the valuation v as follows: $v(p_j) = (c_j^L, c_j^R)$.

It remains to show by induction on ϕ that all formulas present at \mathcal{B} are satisfied by v . The basis case of variables holds by construction of v . We consider only the most instructive case of $\phi_1 \rightarrow \phi_2 \geq_2 i$ as the other cases are straightforward.

Assume that $\phi_1 \rightarrow \phi_2 \geq_2 i \in \mathcal{B}$. Then, by completeness of \mathcal{B} , either $i \leq 0 \in \mathcal{B}$, in which case, $\phi_1 \rightarrow \phi_2 \geq_2 i$ is trivially satisfied, or $\phi_1 \leq_2 j, \phi_2 \geq_2 i + j \in \mathcal{B}$. Furthermore, by the induction hypothesis, v satisfies $\phi_1 \leq_2 j$ and $\phi_2 \geq_2 i + j$, and we also have that $j \leq 1 - i$. Now, to show that v satisfies $\phi_1 \rightarrow \phi_2 \geq_2 i$, recall from semantics that $v_2(\phi_1 \rightarrow \phi_2) = \max(0, v_2(\phi_2) - v_2(\phi_1))$.

Now, we have

$$\max(0, v_2(\phi_2) - v_2(\phi_1)) \geq \max(0, i + j - j) = \max(0, i) = i$$

as desired.

The cases of other connectives can be tackled in a similar fashion. \square

3.2 Constraint tableaux for G^2

Definition 8 (Constraint tableaux for G^2 — $\mathcal{T}(G^2)$). *Let $\lesssim \in \{<, \leq\}$ and $\gtrsim \in \{<, \leq\}$. Branches contain:*

- formulaic constraints of the form $\mathbf{x} : \phi \lesssim \mathbf{x}' : \phi'$ with $\mathbf{x} \in \{1, 2\}$;

- numerical constraints of the form $c \lesssim c'$ with $c, c' \in \{1, 0\}$;
- labelled formulas of the form $\mathbf{x} : \phi * c$ with $* \in \{\lesssim, \gtrsim\}$.

We abbreviate all these types of entries with $\mathfrak{X} \lesssim \mathfrak{X}'$. Each branch can be extended by an application of one of the rules in Figure 2 where $\mathbf{c} \neq \mathbf{c}'$, $c \neq c'$, $\mathbf{c}, \mathbf{c}' \in \{\mathbf{0}, \mathbf{1}\}$ and $c, c' \in \{0, 1\}$. A tableau's branch \mathcal{B} is closed iff at least one of the following conditions applies:

- the transitive closure of \mathcal{B} under \lesssim contains $\mathfrak{X} < \mathfrak{X}$,
- $0 \geq 1 \in \mathcal{B}$ or $\mathfrak{X} > 1 \in \mathcal{B}$ or $\mathfrak{X} < 0 \in \mathcal{B}$.

A tableau is closed iff all its branches are closed. We say that there is a tableau proof of ϕ iff there is a closed tableau starting from $1 : \phi < 1$.

Remark 4 (Interpretation of the constraints). Formulaic constraint $\mathbf{x} : \phi \leq \mathbf{x}' : \phi'$ encodes the fact that $v_{\mathbf{x}}(\phi) \leq v_{\mathbf{x}'}(\phi')$, similarly labelled formula $\mathbf{x} : \phi \leq c$ encodes the fact that $v_{\mathbf{x}}(\phi) \leq c$.

Definition 9 (Satisfying valuation of a branch). Let $\mathbf{x}, \mathbf{x}' \in \{1, 2\}$. A branch \mathcal{B} is satisfied by a valuation v iff

- $v_{\mathbf{x}}(\phi) \leq v_{\mathbf{x}'}(\phi')$ for any $\mathbf{x} : \phi \leq \mathbf{x}' : \phi' \in \mathcal{B}$ and
- $v_{\mathbf{x}}(\phi) \leq c$ for any $\mathbf{x} : \phi \leq c \in \mathcal{B}$ s.t. $c \in \{0, 1\}$.

Theorem 2 (Soundness and completeness). ϕ is G^2 -valid iff it has a $\mathcal{T}(G^2)$ proof.

Proof. For soundness, we check that if the premise of the rule is satisfied, then so is at least one of its conclusions.

For completeness, we show that every complete open branch \mathcal{B} is satisfiable. We construct the satisfying valuation as follows. If $\mathbf{x} : p \geq 1 \in \mathcal{B}$, we set $v_1(p) = 1$. If $1 : p \leq 0 \in \mathcal{B}$, we set $v_1(p) = 0$. We do likewise for $2 : p \leq 0$ and $2 : p \geq 1$. To set the values of the remaining variables q_1, \dots, q_n , we proceed as follows. Denote \mathcal{B}^+ the transitive closure of \mathcal{B} under \lesssim and let

$$[\mathbf{x} : q_i] = \left\{ \mathbf{x}' : q_j \left| \begin{array}{l} \mathbf{x} : q_i \leq \mathbf{x}' : q_j \in \mathcal{B}^+ \text{ or } \mathbf{x} : q_i \geq \mathbf{x}' : q_j \in \mathcal{B}^+ \\ \text{but} \\ \mathbf{x} : q_i < \mathbf{x}' : q_j, \mathbf{x} : q_i > \mathbf{x}' : q_j \notin \mathcal{B}^+ \end{array} \right. \right\}$$

It is clear that there are at most $2n$ $[\mathbf{x} : q_i]$'s since the only possible loop in \mathcal{B}^+ is $\mathbf{x} : r \leq \dots \leq \mathbf{x} : r$, but in such a loop all elements belong to $[\mathbf{x} : r]$. We put $[\mathbf{x} : q_i] \preceq [\mathbf{x}' : q_j]$ iff there are $\mathbf{x} : r \in [\mathbf{x} : q_i]$ and $\mathbf{x}' : r' \in [\mathbf{x}' : q_j]$ s.t. $\mathbf{x} : r \leq \mathbf{x}' : r' \in \mathcal{B}^+$.

We now set the valuation of these variables as follows

$$v_{\mathbf{x}}(q_i) = \frac{|\{[\mathbf{x}' : q'] \mid [\mathbf{x}' : q'] \preceq [\mathbf{x} : q_i]\}|}{2n} \quad (*)$$

Thus, all constraints containing only variables are satisfied.

It remains to show that all other constraints are satisfied. For that, we prove that if at least one conclusion of the rule is satisfied, then so is the premise. We consider only the case of $\rightarrow_2 \lesssim$. Let $1 : \phi_1 \lesssim \mathfrak{X}$ be satisfied. W.l.o.g., assume that $\mathfrak{X} = 2 : \psi$ and $\lesssim = <$. Thus, $v_1(\phi_1) < v_2(\psi)$. Recall that $v_2(\phi_1 \rightarrow \phi_2) = \min(v_1(\phi_1), v_2(\phi_2))$. Hence, $v_2(\phi_1 \rightarrow \phi_2) < v_2(\psi)$, and $2 : \phi_1 \rightarrow \phi_2 < 2 : \psi$ is

$$\begin{array}{c}
 \mathbf{c}_1 \gtrsim \frac{1: \mathbf{c} \lesssim \mathfrak{X}}{c \lesssim \mathfrak{X}} \quad \mathbf{c}_2 \lesssim \frac{2: \mathbf{c} \lesssim \mathfrak{X}}{c' \lesssim \mathfrak{X}} \quad \mathbf{c}_1 \gtrsim \frac{1: \mathbf{c} \gtrsim \mathfrak{X}}{c \gtrsim \mathfrak{X}} \quad \mathbf{c}_2 \gtrsim \frac{2: \mathbf{c} \gtrsim \mathfrak{X}}{c' \gtrsim \mathfrak{X}} \\
 \neg_1 \lesssim \frac{1: \neg \phi \lesssim \mathfrak{X}}{2: \phi \lesssim \mathfrak{X}} \quad \neg_2 \lesssim \frac{2: \neg \phi \lesssim \mathfrak{X}}{1: \phi \lesssim \mathfrak{X}} \quad \neg_1 \gtrsim \frac{1: \neg \phi \gtrsim \mathfrak{X}}{2: \phi \gtrsim \mathfrak{X}} \quad \neg_2 \gtrsim \frac{2: \neg \phi \gtrsim \mathfrak{X}}{1: \phi \gtrsim \mathfrak{X}} \\
 \wedge_1 \gtrsim \frac{1: \phi_1 \wedge \phi_2 \gtrsim \mathfrak{X}}{1: \phi_1 \gtrsim \mathfrak{X} \quad 1: \phi_2 \gtrsim \mathfrak{X}} \quad \wedge_2 \lesssim \frac{2: \phi_1 \wedge \phi_2 \lesssim \mathfrak{X}}{2: \phi_1 \lesssim \mathfrak{X} \quad 2: \phi_2 \lesssim \mathfrak{X}} \quad \vee_1 \lesssim \frac{1: \phi_1 \vee \phi_2 \lesssim \mathfrak{X}}{1: \phi_1 \lesssim \mathfrak{X} \quad 1: \phi_2 \lesssim \mathfrak{X}} \quad \vee_2 \gtrsim \frac{2: \phi_1 \vee \phi_2 \gtrsim \mathfrak{X}}{2: \phi_1 \gtrsim \mathfrak{X} \quad 2: \phi_2 \gtrsim \mathfrak{X}} \\
 \wedge_1 \lesssim \frac{1: \phi_1 \wedge \phi_2 \lesssim \mathfrak{X}}{1: \phi_1 \lesssim \mathfrak{X} \mid 1: \phi_2 \lesssim \mathfrak{X}} \quad \wedge_2 \gtrsim \frac{2: \phi_1 \wedge \phi_2 \gtrsim \mathfrak{X}}{2: \phi_1 \gtrsim \mathfrak{X} \mid 2: \phi_2 \gtrsim \mathfrak{X}} \\
 \vee_1 \gtrsim \frac{1: \phi_1 \vee \phi_2 \gtrsim \mathfrak{X}}{1: \phi_1 \gtrsim \mathfrak{X} \mid 1: \phi_2 \gtrsim \mathfrak{X}} \quad \vee_2 \lesssim \frac{2: \phi_1 \vee \phi_2 \lesssim \mathfrak{X}}{2: \phi_1 \lesssim \mathfrak{X} \mid 2: \phi_2 \lesssim \mathfrak{X}} \\
 \rightarrow_1 \leq \frac{1: \phi_1 \rightarrow \phi_2 \leq \mathfrak{X}}{\mathfrak{X} \geq 1 \mid \begin{array}{l} \mathfrak{X} < 1 \\ 1: \phi_2 \leq \mathfrak{X} \\ 1: \phi_1 > 1: \phi_2 \end{array}} \quad \rightarrow_1 \gtrsim \frac{1: \phi_1 \rightarrow \phi_2 \gtrsim \mathfrak{X}}{1: \phi_1 \leq 1: \phi_2 \mid 1: \phi_2 \gtrsim \mathfrak{X}} \quad \rightarrow_1 < \frac{1: \phi_1 \rightarrow \phi_2 < \mathfrak{X}}{1: \phi_2 < \mathfrak{X} \quad 1: \phi_1 > 1: \phi_2} \\
 \rightarrow_2 \lesssim \frac{2: \phi_1 \rightarrow \phi_2 \lesssim \mathfrak{X}}{2: \phi_2 \leq 2: \phi_1 \mid 2: \phi_2 \lesssim \mathfrak{X}} \quad \rightarrow_2 \geq \frac{2: \phi_1 \rightarrow \phi_2 \geq \mathfrak{X}}{\mathfrak{X} \leq 0 \mid \begin{array}{l} \mathfrak{X} > 0 \\ 2: \phi_2 \geq \mathfrak{X} \\ 2: \phi_2 > 2: \phi_1 \end{array}} \quad \rightarrow_2 > \frac{2: \phi_1 \rightarrow \phi_2 > \mathfrak{X}}{2: \phi_2 > \mathfrak{X} \quad 2: \phi_2 > 2: \phi_1} \\
 \prec_1 \lesssim \frac{1: \phi_1 \prec \phi_2 \lesssim \mathfrak{X}}{1: \phi_1 \leq 1: \phi_2 \mid 1: \phi_1 \lesssim \mathfrak{X}} \quad \prec_1 > \frac{1: \phi_1 \prec \phi_2 > \mathfrak{X}}{1: \phi_1 > \mathfrak{X} \quad 1: \phi_1 > 1: \phi_2} \quad \prec_1 \geq \frac{1: \phi_1 \prec \phi_2 \geq \mathfrak{X}}{\mathfrak{X} \leq 0 \mid \begin{array}{l} \mathfrak{X} > 0 \\ 1: \phi_1 \geq \mathfrak{X} \\ 1: \phi_1 > 1: \phi_2 \end{array}} \\
 \prec_2 \gtrsim \frac{2: \phi_1 \prec \phi_2 \gtrsim \mathfrak{X}}{2: \phi_1 \gtrsim \mathfrak{X} \mid 2: \phi_2 \leq 2: \phi_1} \quad \prec_2 \leq \frac{1: \phi_1 \prec \phi_2 \leq \mathfrak{X}}{\mathfrak{X} \geq 1 \mid \begin{array}{l} \mathfrak{X} < 1 \\ 2: \phi_1 \leq \mathfrak{X} \\ 2: \phi_2 > 2: \phi_1 \end{array}} \quad \prec_2 < \frac{2: \phi_1 \prec \phi_2 < \mathfrak{X}}{2: \phi_1 < \mathfrak{X} \quad 2: \phi_1 < 2: \phi_2} \\
 \rightarrow_1 \leq \frac{1: \phi_1 \rightarrow \phi_2 \leq \mathfrak{X}}{\mathfrak{X} \geq 1 \mid \begin{array}{l} \mathfrak{X} < 1 \\ 1: \phi_2 \leq \mathfrak{X} \\ 1: \phi_1 > 1: \phi_2 \end{array}} \quad \rightarrow_1 \gtrsim \frac{1: \phi_1 \rightarrow \phi_2 \gtrsim \mathfrak{X}}{1: \phi_1 \leq 1: \phi_2 \mid 1: \phi_2 \gtrsim \mathfrak{X}} \quad \rightarrow_1 < \frac{1: \phi_1 \rightarrow \phi_2 < \mathfrak{X}}{1: \phi_2 < \mathfrak{X} \quad 1: \phi_1 > 1: \phi_2} \\
 \rightarrow_2 \lesssim \frac{2: \phi_1 \rightarrow \phi_2 \lesssim \mathfrak{X}}{1: \phi_1 \lesssim \mathfrak{X} \mid 2: \phi_2 \lesssim \mathfrak{X}} \quad \rightarrow_2 \gtrsim \frac{2: \phi_1 \rightarrow \phi_2 \gtrsim \mathfrak{X}}{1: \phi_1 \gtrsim \mathfrak{X} \quad 2: \phi_2 \gtrsim \mathfrak{X}}
 \end{array}$$

Fig. 3. Rules of $\mathcal{T}(\mathbb{G}^2)$. Vertical bars denote branching; $\mathbf{c} \neq \mathbf{c}'$, $c \neq c'$, $\mathbf{c}, \mathbf{c}' \in \{0, 1\}$, $c, c' \in \{0, 1\}$.

satisfied as desired. By the same reasoning, we have that if $2 : \phi_2 \lesssim \mathfrak{X}$ is satisfied, then so is $2 : \phi_1 \rightarrow \phi_2 \lesssim \mathfrak{X}$.

The cases of other rules can be showed in the same fashion. \square

3.3 Applications

Corollary 2. *Satisfiability for any $L_{(x,y)}^2(\rightarrow)$ and $L_{(x,1)}^2(\rightarrow)$ is \mathcal{NP} -complete while their validities are $\text{co}\mathcal{NP}$ -complete.*

Proof. Let $|\phi|$ be the number of symbols in ϕ . Observe, from the proof of Theorem 1, that each tableau branch gives rise to two bounded mixed-integer programming problems (bMIP) — each of the length $O(\rho(|\phi|))$ for some polynomial ρ . Recall that bMIP is \mathcal{NP} -complete (cf. [16]). Thus we can non-deterministically guess an open branch and then solve its two bMIPs (one arising from inequalities with \leq_1 , and the other from those with \leq_2). This yields the \mathcal{NP} - and $\text{co}\mathcal{NP}$ -easiness for satisfiability and validity, respectively.

To obtain the \mathcal{NP} -hardness, we use the same method as in [23,16]. For each classical formula ϕ one can construct a formula $\text{two}(\phi)$ (cf. the detailed definition in [23, Lemmas 3.1–3.3]). Then by [23, Lemma 3.2], ϕ is classically valid iff $\phi^C := \text{two}(\phi) \supset \phi$ is L -valid. Furthermore, if ϕ^C is not valid, there is an L -valuation v such that $v(\phi^C) = 0$. Recall that ϕ^\bullet (resp. ϕ°) denotes the formula obtained by substituting \supset for \rightarrow (resp. \rightarrow) in the formula ϕ (cf. Remark 2). Thus, ϕ is classically valid iff $(\phi^C)^\bullet$ is $L_{(1,0)}^2(\rightarrow)$ -valid and $(\phi^C)^\circ$ is $L_{(1,1)}^2(\rightarrow)$ -valid. Furthermore, if $(\phi^C)^\bullet$ is not valid, there is v' such that $v'((\phi^C)^\bullet) = (0, 1)$, and $(\phi^C)^\circ$ is not valid, there is v' such that $v'((\phi^C)^\circ) = (0, x)$ for some x . Since $(0, 1)$ is not included in any non-trivial filter on $[0, 1] \odot [0, 1]$, and since no non-trivial filter can include $(1, 1)$ and some $(0, x)$ simultaneously, we obtain

$$\models_{CPL} \phi \text{ iff } (\phi^C)^\bullet \text{ is } L_{(x,y)}^2(\rightarrow)\text{-valid iff } (\phi^C)^\circ \text{ is } L_{(x,y)}^2(\rightarrow)\text{-valid}$$

as desired. \square

Remark 5 (Removing the branching). We have introduced branching rules in our tableaux in order to make them more intuitive. It is possible, however, to make all rules *linear* just as it was done originally in [16]. For example, the linear versions of $\rightarrow \leq_1$ and $\rightarrow \geq_2$ look as follows ($y \in \{0, 1\}$):

$$\frac{\phi_1 \rightarrow \phi_2 \leq_1 i}{\phi_1 \geq_1 1 - i + j - y \quad y \leq i \quad \phi_2 \leq_1 j + y \quad j \leq i} \quad \frac{\phi_1 \rightarrow \phi_2 \geq_2 i}{\phi_1 \leq_2 j + y \quad y \leq 1 - i \quad \phi_2 \geq_2 i + j - y \quad j \leq 1 - i}$$

Other rules can be easily acquired since \vee and \wedge can be defined via \rightarrow and $\mathbf{0}$ in the language of L^2 . Rules without branching improve efficiency of the proof search by removing the need to guess the branch whose bMIP we should solve.

Corollary 3. *Satisfiability for $G^2(\rightarrow)$ and $G^2(\rightarrow)$ is \mathcal{NP} -complete.*

Proof. It follows from the proof of theorem 2 that the satisfiability of $G^2(\rightarrow)$ and $G^2(\rightarrow)$ is in \mathcal{NP} : we take the valuation from $(*)$, and it takes polynomial time to check that it indeed satisfies the formula.

The \mathcal{NP} -hardness follows since \mathbf{G}^2 's are conservative extensions of \mathbf{G} whose satisfiability and validity are \mathcal{NP} - and $\text{co}\mathcal{NP}$ -complete respectively. \square

We can also use the tableaux to check whether a set Γ of assumptions entails a formula ϕ in the logics we consider. This yields the finite strong completeness for \mathbf{G}^2 's and \mathbf{L}^2 's by means of tableaux, and extends the complexity results to the finitary entailment.

Corollary 4. *Let Γ be a finite set of formulas. Then $\Gamma \models_{\mathbf{L}^2_{(x,y)}} \phi$ iff the left tableau closes, $\Gamma \models_{\mathbf{G}^2(\rightarrow)} \phi$ iff the central tableau closes, and $\Gamma \models_{\mathbf{G}^2(\rightarrow)} \phi$ iff the left tableau closes.*

$$\begin{array}{c}
 \bigcup_{\phi' \in \Delta} \{\phi' \geq_1 x, \phi' \leq_2 y\} \\
 \begin{array}{ccc}
 / & \backslash & \\
 \phi \geq_1 c & \phi \geq_2 d & \\
 c < x & d > y &
 \end{array}
 \end{array}
 \quad
 \begin{array}{c}
 \{1 : \phi' \geq 1, \phi' \leq 0 \mid \phi' \in \Gamma\} \\
 \begin{array}{ccc}
 / & \backslash & \\
 1 : \phi < 1 & 2 : \phi > 0 &
 \end{array}
 \end{array}
 \quad
 \begin{array}{c}
 \{1 : \phi' \geq 1 \mid \phi' \in \Gamma\} \\
 1 : \phi < 1
 \end{array}$$

Thus, the finitary entailment for any of these logics is $\text{co}\mathcal{NP}$ -complete.

4 Conclusions and further research

Using constraint tableaux, we have provided a modular treatment of the Lukasiewicz and Gödel based two-dimensional logics. Our next steps are: (1) to study the structural proof theory of these logics, and of the two layer logics introduced in [6]; (2) to study and compare the logics in terms of consequence relations, to provide a Hilbert style axiomatization (for those where modus ponens is sound), and to prove standard completeness — cases we understand so far are the following four: $\mathbf{L}^2_{(1,1)}(\rightarrow)$, $\mathbf{L}^2_{(1,0)}(\rightarrow)$ which is the logic $\mathbf{L}_{(-)}$ of [6], $\mathbf{G}^2_{(1,0)}(\rightarrow)$ whose validities coincide with the axiomatic extension of Wansing's I_4C_4 [30] with the prelinearity axiom, and $\mathbf{G}^2_{(1,1)}(\rightarrow)$ whose consequence coincides with the axiomatic extension of Nelson's $N4^\perp$ [24,25] with the prelinearity axiom.

In a broader sense we naturally aim to provide a general treatment of two-dimensional graded logics. Indeed, within the research project introduced in [6], we want to develop a modular logical framework for reasoning based on heterogeneous information (such as crisp or fuzzy data, personal beliefs, etc.) that can be both incomplete and inconsistent. In addition, we do not wish to commit to a specific logic to model the reasoning of the agent(s), because different situations may call for different logics — modeling the reasoning of a group of experts is different from modeling the reasoning of the crowd. Doing so requires to be able to manipulate and combine logics for these different situations in a modular way.

References

1. Avron, A.: The structure of interlaced bilattices. *Mathematical Structures in Computer Science* **6**(3), 287–299 (1996)
2. Avron, A., Arieli, O.: Reasoning with logical bilattices. *Journal of Logic, Language and Information* **5**, 25–63 (1996)
3. Avron, A., Konikowska, B.: Decomposition proof systems for Gödel-Dummett logics. *Studia Logica* **69**(2), 197–219 (2001)
4. Baldi, P., Cintula, P., Noguera, C.: On two-layered modal logics for uncertainty (2020), manuscript
5. Belnap, N.: How a computer should think. In: Omori, H., Wansing, H. (eds.) *New Essays on Belnap-Dunn Logic*, Synthese Library (Studies in Epistemology, Logic, Methodology, and Philosophy of Science), vol. 418. Springer, Cham (2019)
6. Bílková, M., Frittella, S., Majer, O., Nazari, S.: Belief based on inconsistent information. In: Martins, M.A., Sedlár, I. (eds.) *Dynamic Logic. New Trends and Applications*. pp. 68–86. Springer International Publishing, Cham (2020)
7. Cintula, P., Noguera, C.: Modal logics of uncertainty with two-layer syntax: A general completeness theorem. In: *Proceedings of WoLLIC 2014*. pp. 124–136 (2014)
8. D’Agostino, M.: *Investigations into the complexity of some propositional calculi*. Oxford University Computing Laboratory, Oxford (1990)
9. Di Nola, A., Leustean, I.: Łukasiewicz logic and MV-algebras. In: Cintula, P., Hájek, P., Noguera, C. (eds.) *Handbook of Mathematical Fuzzy Logic Vol. 2*. College Publications (2011)
10. Dunn, J.M.: Intuitive semantics for first-degree entailments and ‘coupled trees’. *Philosophical Studies* **29**(3), 149–168 (1976)
11. Dunn, J.M.: Contradictory information: Too much of a good thing. *Journal of Philosophical Logic* **39**, 425–452 (2010)
12. Fagin, R., Halpern, J.Y., Megiddo, N.: A logic for reasoning about probabilities. *Information and Computation* **87**, 78–128 (1990)
13. Font, J.: *Abstract Algebraic Logic-An Introductory Textbook*, Studies in Logic, vol. 60. College Publications, London (2016)
14. Ginsberg, M.: Multivalued logics: A uniform approach to reasoning in AI. *Computer Intelligence* **4**, 256–316 (1988)
15. Goré, R.: Dual intuitionistic logic revisited. In: Dyckhoff, R. (ed.) *Automated Reasoning with Analytic Tableaux and Related Methods*. pp. 252–267. Springer Berlin Heidelberg, Berlin, Heidelberg (2000)
16. Hähnle, R.: Many-valued logic and mixed integer programming. *Annals of mathematics and Artificial Intelligence* **12**(3-4), 231–263 (1994)
17. Hähnle, R.: Advanced many-valued logics. In: Gabbay, D., Guentner, F. (eds.) *Handbook of Philosophical Logic*. vol. 2nd, pp. 297–395. Springer Netherlands, Dordrecht (2001). https://doi.org/10.1007/978-94-017-0452-6_5
18. Hähnle, R.: Tableaux and related methods. In: Robinson, J.A., Voronkov, A. (eds.) *Handbook of Automated Reasoning*, vol. 2, pp. 100–178. Elsevier and MIT Press (2001). <https://doi.org/10.1016/b978-044450813-3/50005-9>
19. Hájek, P.: *Metamathematics of Fuzzy Logic*. Trends in Logic 4, Springer, Dordrecht (1998)
20. Jansana, R., Riviaccio, U.: Residuated bilattices. *Soft Comput.* **16**(3), 493–504 (2012)
21. Klein, D., Majer, O., Raffie-Rad, S.: Probabilities with gaps and gluts (2020), <http://arxiv.org/abs/2003.07408>, to appear in *Journal of Philosophical Logic*

22. Metcalfe, G., Olivetti, N., Gabbay, D.: Proof Theory for Fuzzy Logics. Applied Logic Series 36, Springer (2008)
23. Mundici, D.: Satisfiability in many-valued sentential logic is NP-complete. Theoretical Computer Science **52**(1), 145–153 (1987). [https://doi.org/10.1016/0304-3975\(87\)90083-1](https://doi.org/10.1016/0304-3975(87)90083-1)
24. Nelson, D.: Constructible falsity. The Journal of Symbolic Logic **14**(1), 16–26 (1949), <http://www.jstor.org/stable/2268973>
25. Odintsov, S.: Constructive negations and paraconsistency, Trends in logic, vol. 26. Springer (2008)
26. Omori, H., Wansing, H.: 40 years of FDE: An Introductory Overview. Studia Logica **105**(6), 1021–1049 (Dec 2017). <https://doi.org/10.1007/s11225-017-9748-6>
27. Rauszer, C.: An algebraic and Kripke-style approach to a certain extension of intuitionistic logic. dissertation, Institute of Mathematics, Polish Academy of Sciences (1980)
28. Riviaccio, U.: An Algebraic Study of Bilattice-based Logics. Ph.D. thesis, University of Barcelona - University of Genoa (2010)
29. Vakarelov, D.: Notes on N-lattices and constructive logic with strong negation. Studia logica **36**(1–2), 109–125 (1977)
30. Wansing, H.: Constructive negation, implication, and co-implication. Journal of Applied Non-Classical Logics **18**(2–3), 341–364 (2008). <https://doi.org/10.3166/jancl.18.341-364>

5 Appendix

Proposition 1. Let ϕ be a $L_{(x,y)}^2(\rightarrow)$ (resp. $G_{(x,y)}^2(\rightarrow)$) formula without constants and $(1, 1) \notin (x, y)^\uparrow$. Then ϕ is not $L_{(x,y)}^2(\rightarrow)$ (resp. $G_{(x,y)}^2(\rightarrow)$) valid.

Proof. Observe that all connectives preserve $(1, 1)$. The result follows. \square

5.1 Proofs of Section 2.3. Semantical properties of $L_{(x,y)}^2(\rightarrow)$

Proposition 2

- Let $y \geq 1 - x$. Then ϕ is $L_{(x,y)}^2(\rightarrow)$ -valid iff ϕ is $L_{(x,1-x)}^2(\rightarrow)$ -valid.
- Let $y < 1 - x$. Then ϕ is $L_{(x,y)}^2(\rightarrow)$ -valid iff ϕ is $L_{(1-y,y)}^2(\rightarrow)$ -valid.

Proof. We prove only the first case, as the second one can be tackled in the same manner. Assume that ϕ is $L_{(x,1-x)}^2(\rightarrow)$ -valid. Since $y \geq 1 - x$, $(x, 1 - x)^\uparrow \subseteq (x, y)^\uparrow$, then ϕ is $L_{(x,y)}^2(\rightarrow)$ -valid as well. Now let us show that if ϕ is $L_{(x,y)}^2(\rightarrow)$ -valid, then ϕ is $L_{(x,1-x)}^2(\rightarrow)$ -valid as well.

For any valuation v , let us define the valuation v^* as follows: $v^*(p) := (1 - v_2(p), 1 - v_1(p))$. We now show that the extension of v^* satisfies the following property: for every formula ϕ , we have $v^*(\phi) = (1 - v_2(\phi), 1 - v_1(\phi))$.

Notice that $v(\phi) = (x', y') \Leftrightarrow v(\neg \sim \phi) = (1 - y', 1 - x')$ and $v(\psi_1 \leftrightarrow \psi_2) = (1, 0) \Leftrightarrow v(\psi_1) = v(\psi_2)$. Since $(1, 0) \in (x, y)^\uparrow$ for any x and y , a formula always

evaluated at $(1, 0)$ is necessarily $L_{(x,y)}^2(\rightarrow)$ -valid. Notice that

$$\begin{aligned} v(\neg\neg\neg\psi_1 \leftrightarrow \neg\neg\neg\psi_1) &= (1, 0), \\ v(\neg\neg(\psi_1 \wedge \psi_2) \leftrightarrow (\neg\neg\psi_1 \wedge \neg\neg\psi_2)) &= (1, 0), \\ v(\neg\neg(\psi_1 \vee \psi_2) \leftrightarrow (\neg\neg\psi_1 \vee \neg\neg\psi_2)) &= (1, 0), \\ v(\neg\neg(\psi_1 \rightarrow \psi_2) \leftrightarrow (\neg\neg\psi_1 \rightarrow \neg\neg\psi_2)) &= (1, 0). \end{aligned}$$

Indeed, it follows from the fact that these formulas are derivable in the Hilbert calculus for $L_{(-)}$ from [6, Lemma 2] which is sound and complete w.r.t. $[0, 1]_{\mathbb{L}} \odot [0, 1]_{\mathbb{L}}(\rightarrow)_{(1,0)}$. Using the four equalities above, one can prove by induction on ϕ that $v^*(\phi) = (1 - v_2(\phi), 1 - v_1(\phi))$ for any formula ϕ . Thus for every valuation v we have defined its counterpart v^* such that $v^*(\phi)$ is the reflection of $v(\phi)$ along the vertical axis of the lattice.

Let ϕ be $L_{(x,y)}^2(\rightarrow)$ -valid but not $L_{(x,1-x)}^2(\rightarrow)$ -valid. Then, there is a v s.t. $v(\phi) \in (x, y)^\uparrow$ but $v(\phi) \notin (x, 1-x)^\uparrow$. But then $v^*(\phi) \notin (x, y)^\uparrow$ since only $(x, 1-x)^\uparrow$'s are closed under conflation. Contradiction. \square

Proposition 3 Let $m, n \in \{2, 3, \dots\}$. Then, $L_{(\frac{m-1}{m}, \frac{1}{m})}^2 \subsetneq L_{(\frac{n-1}{n}, \frac{1}{n})}^2$ iff $m > n$.

Proof. The inclusion follows immediately from the fact that

$$\left(\frac{m-1}{m}, \frac{1}{m}\right)^\uparrow \subsetneq \left(\frac{n-1}{n}, \frac{1}{n}\right)^\uparrow \text{ iff } m > n$$

To show the strictness of the inclusion, consider the following family of formulas.

$$F_n := \bigvee_{1 \leq i < j \leq n+1} (p_i \leftrightarrow p_j)$$

Notice that $v(\phi_1 \leftrightarrow \phi_2) = (1 - |v_1(\phi_1) - v_1(\phi_2)|, |v_2(\phi_1) - v_2(\phi_2)|)$. In addition, we have that for $n \geq 2$:

$$v(F_n) \geq \left(\frac{n-1}{n}, \frac{1}{n}\right)$$

Indeed, we can observe that $v_1(\psi_1 \leftrightarrow \psi_2)$ is the *complement to the distance* between $v_1(\psi_1)$ and $v_1(\psi_2)$ on $[0, 1]$, while $v_2(\psi_1 \leftrightarrow \psi_2)$ is the distance between $v_2(\psi_1)$ and $v_2(\psi_2)$. Thus, $v_1(F_n)$ is the *maximal complement to the distance between any two points out of n on $[0, 1]$* , while $v_2(F_n)$ is the *minimal such distance*. The lower bound on $v(F_n)$ is produced when we place points on $[0, 1]$ with *equal intervals* between them. Hence, in each $L_{(\frac{k-1}{k}, \frac{1}{k})}^2(\rightarrow)$, only F_n 's with $n \geq k$ are valid. Furthermore, none of the F_n 's are valid in $L_{(1,0)}$. This gives us the desired strictness of the inclusion. \square

Proposition 4 Let $m, n \in \{3, 4, \dots\}$. Then $L_{(\frac{m-2}{2m}, \frac{m+2}{2m})}^2 \subsetneq L_{(\frac{n-2}{2n}, \frac{n+2}{2n})}^2$ iff $m > n$.

Proof. As in Proposition 3, inclusion follows from the fact that

$$\left(\frac{m-2}{2m}, \frac{m+2}{2m}\right)^\uparrow \subsetneq \left(\frac{n-2}{2n}, \frac{n+2}{2n}\right)^\uparrow \text{ iff } m > n$$

Again, the non-trivial part is the strictness. Recall that $\psi_1 \odot \psi_2 = \sim(\psi_1 \rightarrow \sim\psi_2)$. Now, consider the family of formulas $F_2 \odot F_n$ for $n \geq 3$. We have that $v(F_2 \odot F_n) \geq \left(\max\left(0, \frac{1}{2} + \frac{n-1}{n} - 1\right), \min\left(1, \frac{1}{2} + \frac{1}{n}\right) \right) \geq \left(\frac{n-2}{2n}, \frac{n+2}{2n}\right)$.

As in Proposition 3, we can see that $F_2 \odot F_k$ is $L_{(\frac{n-2}{2n}, \frac{n+2}{2n})}$ -valid iff $k \geq n$. \square

Proposition 5 Let $L_{(1,0)}^2(\rightarrow) \subsetneq L_{(x,y)}^2(\rightarrow)$. Then $L_{(x,y)}^2(\rightarrow)$ is not closed under modus ponens.

Proof. Observe that by proposition 2, $L_{(1,0)}^2(\rightarrow) \subsetneq L_{(x,y)}^2(\rightarrow)$ iff $x < 1$ and $y > 0$. Furthermore, by proposition 2, it suffices to consider only the $L_{(x',1-x')}^2(\rightarrow)$'s. Since for any ψ and ψ' , $\psi \rightarrow (\psi' \rightarrow (\psi \odot \psi'))$ is $L_{(1,0)}^2$ -valid, it is enough to find $L_{(x',1-x')}^2(\rightarrow)$ -valid formulas ψ and ψ' such that $\psi \odot \psi'$ is not $L_{(x',1-x')}^2(\rightarrow)$ -valid.

We consider two cases. Either (1) $(x', 1-x')^\uparrow$ is not an extension of $(\frac{1}{2}, \frac{1}{2})^\uparrow$ or (2) it is. In the first case, we have $\frac{1}{2} < x' \leq 1$. Hence, there is a $k \in \mathbb{N}$ s.t. $k \geq 3$ and $\frac{k-2}{k-1} < x' \leq \frac{k-1}{k}$. Hence, F_k is $L_{(x',1-x')}^2(\rightarrow)$ -valid. By Proposition 3, there is v such that $v(F_{k-1}) = (\frac{k-2}{k-1}, \frac{1}{k-1}) \notin (x', 1-x')^\uparrow$. Hence, F_{k-1} is not $L_{(x',1-x')}^2(\rightarrow)$ -valid. Observe that in such case $F_k \odot F_k$ is not $L_{(x',1-x')}^2(\rightarrow)$ -valid. Indeed, consider the valuation v' such that $v'(p_1) = (1, 0)$, $v'(p_{k+1}) = (0, 1)$, and $v'(p_i) = \frac{i-1}{k}$ for every $2 \leq i \leq k$. We get that

$$v'(F_k \odot F_k) = \left(\frac{k-2}{k}, \frac{2}{k}\right) < (x', 1-x'). \quad (\text{because } \frac{k-2}{k-1} < x')$$

Hence, $F_k \odot F_k$ is not $L_{(x',1-x')}^2(\rightarrow)$ -valid as desired.

For the second case, we have $0 \leq x \leq \frac{1}{2}$. Note that $v'(F_2 \odot F_2) = (0, 1)$ and thus not valid in any $L_{(x,y)}^2$. This proves the second case as F_2 is $L_{(x,y)}^2$ -valid iff $(x, y)^\uparrow \supseteq (\frac{1}{2}, \frac{1}{2})^\uparrow$. \square

5.2 Proofs of Section 2.4. Semantical properties of $\mathbf{G}^2(\rightarrow)$

Proposition 6 Let ϕ be a formula over $\{\mathbf{0}, \mathbf{1}, \neg, \wedge, \vee, \rightarrow, \prec\}$. For any $v(p) = (x, y)$, let $v^*(p) = (1-y, 1-x)$. Then $v(\phi) = (x, y)$ iff $v^*(\phi) = (1-y, 1-x)$.

Proof. Let χ and χ' be formulas. Let us show that, for all $\mathbf{x}, \mathbf{x}', \mathbf{y}, \mathbf{y}' \in \{1, 2\}$ such that $\mathbf{x} \neq \mathbf{y}$ and $\mathbf{x}' \neq \mathbf{y}'$, we have

$$v_{\mathbf{x}}(\chi) \geq v_{\mathbf{x}'}(\chi') \Leftrightarrow v_{\mathbf{y}'}^*(\chi') \geq v_{\mathbf{y}}^*(\chi) \quad (*)$$

We proceed by the induction on the number of unary and binary connectives in both χ and χ' . The only non-trivial case is that of \rightarrow . For \rightarrow , we have

$$\begin{aligned} v_1(\psi_1 \rightarrow \psi_2) < v_{\mathbf{x}}(\chi) &\Leftrightarrow v_1(\psi_1) > v_1(\psi_2) \text{ and } v_1(\psi_2) < v_{\mathbf{x}}(\chi) \\ &\Leftrightarrow v_2^*(\psi_1) < v_2^*(\psi_2) \text{ and } v_2^*(\psi_2) > v_{\mathbf{y}}^*(\chi) \quad (\text{by IH}) \\ &\Leftrightarrow v_2^*(\psi_1 \rightarrow \psi_2) > v_{\mathbf{y}}^*(\chi) \end{aligned}$$

and

$$\begin{aligned}
& v_1(\psi_1 \rightarrow \psi_2) = v_{\mathbf{x}}(\chi) \\
\Leftrightarrow & v_1(\psi_1) > v_1(\psi_2) = v_{\mathbf{x}}(\chi) \text{ or } (v_1(\psi_1) \leq v_1(\psi_2) \text{ and } v_{\mathbf{x}}(\chi) = v_1(\mathbf{1})) \\
\Leftrightarrow & v_2^*(\psi_1) < v_2^*(\psi_2) = v_{\mathbf{y}}^*(\chi) \text{ or } (v_2^*(\psi_1) \geq v_2^*(\psi_2) \text{ and } v_{\mathbf{y}}^*(\chi) = v_2^*(\mathbf{1})) \quad (\text{by IH}) \\
\Leftrightarrow & v_2^*(\psi_1 \rightarrow \psi_2) = v_{\mathbf{y}}^*(\chi)
\end{aligned}$$

Now we can prove the statement by induction on ϕ . The basis cases of variables and constants hold by the construction of v^* . We only present the case of \rightarrow . We consider two cases: $(x, y) \neq (1, 0)$ and $(x, y) = (1, 0)$.

In the first and in the second cases, we have

$$\begin{aligned}
& v(\psi_1 \rightarrow \psi_2) = (x, y) \\
\Leftrightarrow & v_1(\psi_1) > v_1(\psi_2) = x \text{ and } v_2(\psi_1) < v_2(\psi_2) = y \\
\Leftrightarrow & v_2^*(\psi_1) < v_2^*(\psi_2) = 1 - x \text{ and } v_1^*(\psi_1) > v_1^*(\psi_2) = 1 - y \quad (\text{by IH and } (*)) \\
\Leftrightarrow & v^*(\psi_1 \rightarrow \psi_2) = (1 - y, 1 - x)
\end{aligned}$$

and

$$\begin{aligned}
v(\psi_1 \rightarrow \psi_2) = (1, 0) & \Leftrightarrow v_1(\psi_1) \leq v_1(\psi_2) \text{ and } v_2(\psi_1) \geq v_2(\psi_2) \\
& \Leftrightarrow v_2^*(\psi_1) \geq v_2^*(\psi_2) \text{ and } v_1^*(\psi_1) \leq v_1^*(\psi_2) \quad (\text{by IH and } (*)) \\
& \Leftrightarrow v^*(\psi_1 \rightarrow \psi_2) = (1, 0).
\end{aligned}$$

□

In order to prove Proposition 7, we need the following claims.

Claim 1. (NNF) Each formula over $\{\mathbf{0}, \mathbf{1}, \neg, \wedge, \vee, \rightarrow, \prec\}$ is equivalent to a formula in a \neg -negation normal form (NNF) in $\mathbf{G}^2(\rightarrow)$.

Proof. We create the NNF by repeating applications of instances of the following formulas, which are always designated (as they are always evaluated at $(1, 0)$).

$$\begin{array}{ll}
\neg\neg\phi \leftrightarrow \phi & \neg(\phi \wedge \psi) \leftrightarrow (\neg\phi \vee \neg\psi) \\
\neg\mathbf{0} \leftrightarrow \mathbf{1} & \neg(\phi \vee \psi) \leftrightarrow (\neg\phi \wedge \neg\psi) \\
\neg\mathbf{1} \leftrightarrow \mathbf{0} & \neg(\phi \rightarrow \psi) \leftrightarrow (\neg\psi \prec \neg\phi) \\
& \neg(\phi \prec \psi) \leftrightarrow (\neg\psi \rightarrow \neg\phi)
\end{array}$$

A similar claim was shown in [6] for the logic $\mathbf{L}_{(\neg)}$ which coincides with $\mathbf{L}_{(1,0)}^2(\rightarrow)$, and can be shown for all the logics we consider in this paper, using the corresponding implication and varying the way we negate it (the one but last formula on the right). For logics with \rightarrow we only get weakly equivalent negation normal form. □

Claim 2. For every Gödel formula ϕ over $\{\mathbf{0}, \mathbf{1}, \wedge, \vee, \rightarrow, \prec\}$, for every valuation v such that $v(\phi) < 1$ and for every $0 < x \leq 1$, we have that

1. $v(\phi) \leq \max\{v(p) \mid p \in \text{Var}(\phi)\}$ and
2. there exists a valuation v' such that $v'(\phi) \leq x$.

Proof. 1) We proceed by induction. The only non-trivial parts are $\phi = \phi_1 \rightarrow \phi_2$ and $\phi = \phi_1 \prec \phi_2$.

If $v(\phi_1 \rightarrow \phi_2) < 1$, then $v(\phi_2) < v(\phi_1)$ and $v(\phi_1 \rightarrow \phi_2) = v(\phi_2) < 1$. Let $q \in \mathbf{Var}(\phi)$ be such that $v(q) = \max\{v(p) \mid p \in \mathbf{Var}(\phi)\}$. We consider two cases: either $q \in \mathbf{Var}(\phi_1)$ or $q \in \mathbf{Var}(\phi_2)$. In both cases we obtain the desired result by the induction hypothesis.

If $v(\phi_1 \prec \phi_2) < 1$, then $v(\phi_1) < 1$ and $v(\phi_1 \prec \phi_2) = v(\phi_1)$ or $v(\phi_1 \prec \phi_2) = 0$ and $v(\phi_1) \leq v(\phi_2)$. Let $q \in \mathbf{Var}(\phi)$ be such that $v(q) = \max\{v(p) \mid p \in \mathbf{Var}(\phi)\}$. We consider two cases: either $q \in \mathbf{Var}(\phi_1)$ or $q \in \mathbf{Var}(\phi_2)$. In both cases we obtain the desired result by the induction hypothesis.

2) Let $v(\phi) < 1$, $n > 0$ and $\frac{1}{n} \leq x$. We construct $v'(\phi)$ as follows: $v'(p) = \frac{v(p)}{n}$. We can prove by induction on ϕ that $v'(\phi) < 1$. Hence, by the previous item, we have that

$$v'(\phi) \leq \max\{v'(p) \mid p \in \mathbf{Var}(\phi)\} = \frac{1}{n} \max\{v(p) \mid p \in \mathbf{Var}(\phi)\} \leq x$$

as required. \square

Claim 3. Let ϕ be a formula over $\{\mathbf{0}, \mathbf{1}, \neg, \wedge, \vee, \rightarrow, \prec\}$, for every valuation v such that $v_1(\phi) \neq 1$ and for every $0 < x \leq 1$, we have that

1. $v_1(\phi) \leq \max(\{v_1(p) \mid p \in \mathbf{Var}(\phi)\} \cup \{v_2(p) \mid p \in \mathbf{Var}(\phi)\})$ and
2. there exists a valuation v' such that $v'_1(\phi) \leq x$.

Proof. W.l.o.g. we can assume that ϕ is in NNF.

1) Let v be a valuation such that $v_1(\phi) \neq 1$. ϕ being a formula in NNF in the language $\{\mathbf{0}, \mathbf{1}, \neg, \wedge, \vee, \rightarrow, \prec\}$ over the set \mathbf{Prop} , it can be perceived as a formula ϕ' in the language $\{\mathbf{0}, \mathbf{1}, \wedge, \vee, \rightarrow, \prec\}$ over the set of literals \mathbf{Lit} in place of propositional atoms. Let w be the new valuation over the set of literals defined as follows: $w(p) = v_1(p)$ and $w(\neg p) = v_1(\neg p) = v_2(p)$. By applying Claim 2 to ϕ' and w , we get that

$$w(\phi') \leq \max\{w(l) \mid l \in \mathbf{Lit}(\phi')\} = \max(\{v_1(p) \mid p \in \mathbf{Var}(\phi)\} \cup \{v_2(p) \mid p \in \mathbf{Var}(\phi)\})$$

as required.

2) By Claim 2, there exists w' on the set of literals such that $w'(\phi') \leq x$. Let v' be defined as follows: $v'(p) = (w(p), w(\neg p))$ for every $p \in \mathbf{Prop}$. We get that $v'(\phi) \leq x$ as required. \square

Proposition 7 Let ϕ be a formula over $\{\mathbf{0}, \mathbf{1}, \neg, \wedge, \vee, \rightarrow, \prec\}$ such that $v(\phi) \geq (x, y)$ for any v and some fixed $(x, y) \neq (0, 1)$. Then $v'(\phi) = (1, 0)$ for any v' .

Proof. Assume that ϕ is $\mathbf{G}_{(x,y)}^2$ -valid and w.l.o.g. that ϕ is in NNF. Observe that (x, x) points are not affected by \neg . Now recall that by proposition 6, if $v(\phi) \neq (1, 0)$, there is a v' such that $v'_1(\phi) \neq 1$. Furthermore, notice that if $\{(0, 0), (1, 1)\} \subseteq (x, y)^\uparrow$ then $(x, y)^\uparrow = (0, 1)$. Hence, we have that $\{(0, 0), (1, 1)\} \not\subseteq (x, y)^\uparrow$, which implies that

$$\begin{aligned} & \exists(z, z) \forall(x', y') \in (x, y)^\uparrow : (z, z) \neq (1, 1) \text{ and } z \geq y' \\ & \text{or} \\ & \exists(z, z) \forall(x', y') \in (x, y)^\uparrow : (z, z) \neq (0, 0) \text{ and } z \leq x'. \end{aligned}$$

By proposition 6, we know that $v(\phi) = (0, 0)$ iff $v^*(\phi) = (1, 1)$, hence we can state w.l.o.g. that $(0, 0) \notin (x, y)^\uparrow$ and that $v(\phi) \neq (0, 0)$ for every v . Thus, there is a (z, z) such that $(z, z) \neq (0, 0)$ and for any $(x', y') \in (x, y)^\uparrow$ $z \leq x'$.

Assume, for contradiction, that $v'(\phi) \neq (1, 0)$. There are two cases.

Case 1: $v'_1(\phi) \neq 1$. Since $x > 0$, by Claim 3, we have that there exists a valuation v' such that $v'_1(\phi) \leq \frac{x}{2} < x$. Hence, $v'(\psi) \notin (x, y)^\uparrow$, which contradicts the fact that ψ is $G_{(x,y)}^2$ -valid.

Case 2: $v'_1(\phi) = 1$. Then $v'_2(\phi) \neq 0$ and $v_1^*(\phi) = 1 - v'_2(\phi) \neq 1$ (see Proposition 6 for definition of v^*). We proceed as in the previous case. □

Corollary 1 $v(\phi) = (1, 0)$ for any v iff $v'_1(\phi) = 1$ for any v' .

Proof. Left to right is obvious. We prove right to left by contraposition. Assume that $v(\phi) \neq (1, 0)$. Then, either $v_1(\phi) \neq 1$ or $v_2(\phi) = c \neq 0$. In the first case, we get the result. In the second case we obtain by proposition 6 that $1 \neq v_1^*(\phi) = 1 - c$ as required. □