

Yet another eigenvalue algorithm for solving polynomial systems

Matías R. Bender* Simon Telen†

Abstract

In latest years, several advancements have been made in symbolic-numerical eigenvalue techniques for solving polynomial systems. In this article, we add to this list. We design an algorithm which solves systems with isolated solutions reliably and efficiently. In overdetermined cases, it reduces the task to an eigenvalue problem in a simpler and considerably faster way than in previous methods, and it can outperform the homotopy continuation approach. We provide many examples and an implementation in the proof-of-concept Julia package `EigenvalueSolver.jl`.

Key words — polynomial systems, eigenvalue theorem, symbolic-numerical algorithm
AMS subject classifications — 65H04, 65H10

1 Introduction

Polynomial systems arise in many areas of applied science [42, 20]. This paper is concerned with solving such systems of equations using numerical computations, that is, using finite precision, floating point arithmetic. Two important classes of numerical algorithms are algebraic algorithms [30, 36] and homotopy continuation methods [16, 42]. See [20, Ch. 2] for an overview. In this work, we focus on algorithms of the former type.

Algebraic algorithms are also called eigenvalue algorithms. They consist of two steps. Step (A) uses linear algebra operations to reduce the problem to an eigenvalue problem or univariate polynomial root finding problem. Step (B) is to solve the eigenvalue or univariate root finding problem using numerical tools. Classical examples include Gröbner basis and resultant algorithms, see [23, Ch. 2] or [6]. These use symbolic manipulations for step (A), pushing the numerical linear algebra back to the eigenvalue computation in step (B). The reason for this is that, when performed in finite precision arithmetic, these approaches are numerically unstable for step (A), see for instance [35]. Border basis methods have been developed to remedy this unstable behaviour [39, 43] and variants based on nullspace computations were introduced in [25]. Methods for performing step (B) are based on linear algebra [19] or, recently, on multilinear algebra [49].

Two special types of structured matrices play a central role in algebraic algorithms: Macaulay (or Sylvester) matrices and multiplication matrices. Macaulay matrices have a sparse, quasi-Toeplitz structure. They contain the coefficients of the equations and are manipulated in step (A). The result of these manipulations is a set of multiplication matrices. These are structured

*Department of Mathematics, Technische Universität Berlin, mbender@math.tu-berlin.de

†Max Planck Institute for Mathematics in the Sciences, Leipzig, simon.telen@mis.mpg.de

in the sense that they commute. Multiplication matrices represent multiplication operators in the coordinate ring of the solution set [23, Ch. 5] and their eigenstructure reveals the coordinates of the solutions [22, Ch. 2]. As the Macaulay matrices are typically much larger than multiplication matrices, step (A) determines the running time of the algorithm. This motivates the efforts in active research, including the present paper, to design algorithms which use smaller Macaulay matrices.

In practice, to construct multiplication matrices from Macaulay matrices, we need to choose a basis for the aforementioned coordinate ring. The numerical stability of step (A) strongly depends on this choice [47]. Gröbner and border basis methods use bases corresponding to special sets of monomials. For instance, they require these monomials to come from a monomial ordering [23, Ch. 2, §2] or to be ‘connected-to-1’ [39]. Recent developments showed that numerical linear algebra heuristics can be applied to choose bases that improve the accuracy substantially [47]. This has led to the development of truncated normal forms [46], which use much more general bases of monomials coming from QR factorizations with optimal column pivoting, or non-monomial bases coming from singular value decompositions or Chebyshev representations [40].

Other than making a good choice of basis, in order to stabilize algebraic algorithms it is necessary to take solutions at infinity into account. Loosely speaking, a polynomial system has solutions at infinity if the slightest random perturbation of the nonzero coefficients introduces new solutions with large coordinates. This is best understood in the language of toric geometry [24]. Situations in which there are finitely many solutions at infinity (see Assumption 1) can be handled by introducing an extra randomization in the algorithm, which was first used in [44, 11]. Where classically the multiplication matrices represent ‘multiplication with a polynomial g ’, the multiplication matrices in these papers represent ‘multiplication with a rational function g/f_0 ’, where f_0 is a random polynomial that does not vanish at any of the solutions to the system. For details and a geometric interpretation, we refer to [44, 11]. We will use a similar approach in this paper. In terms of our results, choosing the denominator f_0 randomly is essential in cases where the conditions in Lemma 2.1 are not satisfied for $f_0 = 1$, while they are for a generic f_0 (Example 3).

We summarize the contributions of the present paper. First, we adapt the eigenvalue theorem [22, Ch. 2, Thm. 4.5] to reduce the problem of solving polynomial systems to the computation of eigenvalues. Our new version allows to compute solutions from matrices that need not represent classical multiplication operators; see Example 5. We propose an easy-to-state and easy-to-verify criterion for Macaulay matrices to be ‘large enough’ for constructing such matrices (Lemma 2.1). Moreover, we identify a broad class of overdetermined polynomial systems, namely semi-regular unmixed systems, for which these Macaulay matrices are much smaller than those in classical algorithms, e.g. [27, 38]. We distil these new insights, together with the recent advances in numerical eigenvalue algorithms explained above, into an algorithm (Algorithm 2). We introduce the notion of admissible tuples (Definition 2.1), which parametrize Macaulay matrices satisfying our criterion from Lemma 2.1 and show how to construct such tuples for structured systems of equations. Additionally, we adapt [40, Sec. 4] to obtain an algorithm for computing smaller admissible tuples for overdetermined, unmixed systems (Algorithm 3). We provide a Julia implementation of our algorithms, available online at <https://github.com/simontelen/JuliaEigenvalueSolver>. Our experiments show the efficiency and accuracy of this package. They contain a comparison with the state-of-the-art Julia package `HomotopyContinuation.jl` [14]. We show that our eigenvalue methods are competitive, and in strongly overdetermined cases, they are considerably faster.

To make the paper accessible to a wide audience, we state most of our results and proofs using only terminology from linear algebra. For results that require more background in algebraic (and in particular toric) geometry, we sketch proofs and provide full references.

The paper is organized as follows. In Section 2, we introduce our adapted eigenvalue theorem, admissible tuples and our algorithm. In Section 3, we present constructions for admissible tuples for different families of polynomial systems. Finally, in Section 4, we demonstrate the effectiveness of our algorithms through extensive numerical experimentation. Our computations are done using the Julia package `EigenvalueSolver.jl`.

2 The algorithm

In this section, we present a symbolic-numerical algorithm to solve polynomial systems (Algorithm 2). We show that the solutions of the system can be obtained from the eigenvalues of certain matrices M_g defined in (2.2). For some choice of input for Algorithm 2, these matrices represent *multiplication operators*, see Remark 2.3. In this case, the results of this section are well-known, e.g. [37]. However, in general, our matrices M_g may not have this interpretation. This is illustrated in Example 5. The upshot in these cases is that they can be computed more efficiently.

Consider the polynomial ring $R := \mathbb{C}[x_1, \dots, x_n]$ and a tuple of s polynomials $\mathcal{F} := (f_1, \dots, f_s) \in R^s$, with $s \geq n$. Our aim in this section is to present an algorithm for solving the system of equations $\mathcal{F}(x) = 0$, where we use the short notation x for (x_1, \dots, x_n) . A point $\zeta \in \mathbb{C}^n$ is called a *solution* of \mathcal{F} if $\mathcal{F}(\zeta) = 0$, that is, $f_i(\zeta) = 0$ for every $i \in \{1, \dots, s\}$. For a vector $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$, we denote by x^α the monomial $\prod_{i=1}^n x_i^{\alpha_i} \in R$. We say that α is the *exponent* of the monomial x^α . In what follows, we write each polynomial f_i as

$$f_i := \sum_{\alpha \in \mathbb{N}^n} c_{i,\alpha} x^\alpha.$$

where $c_{i,\alpha} \in \mathbb{C}$ are the *coefficients* of f_i and finitely many of them are nonzero. We define the *support* A_i of f_i as the set of exponents $\alpha \in \mathbb{N}^n$ corresponding to non-zero coefficients $c_{i,\alpha} \in \mathbb{C}$,

$$A_i := \{\alpha \in \mathbb{N}^n : c_{i,\alpha} \neq 0\}.$$

Given two subsets $E_1, E_2 \subset \mathbb{N}^n$, we denote by $E_1 + E_2$ the *Minkowski sum* of E_1, E_2 , that is,

$$E_1 + E_2 := \{\alpha + \beta : \alpha \in E_1, \beta \in E_2\}.$$

For a finite set of exponents $E \subset \mathbb{N}^n$, we write R_E for the subvector space of R spanned by the monomials with exponent in E . That is,

$$R_E := \bigoplus_{\alpha \in E} \mathbb{C} \cdot x^\alpha.$$

Observe that, given $g_1 \in R_{E_1}$ and $g_2 \in R_{E_2}$, we have that $g_1 g_2 \in R_{E_1 + E_2}$.

Consider a tuple of s finite sets of exponents $\mathbf{E} := (E_1, \dots, E_s)$, where $E_i \subset \mathbb{N}^n$, and another finite set of exponents $D \subset \mathbb{N}^n$ such that for every $i \in \{1, \dots, s\}$, D contains the exponents in $A_i + E_i$. An essential ingredient for our eigenvalue algorithm is the *Sylvester map*

$$\begin{aligned} \text{Sylv}_{(\mathcal{F}, \mathbf{E}; D)} : R_{E_1} \times \dots \times R_{E_s} &\rightarrow R_D \\ (g_1, \dots, g_s) &\mapsto \sum_i g_i f_i. \end{aligned}$$

This is a linear map between finite dimensional vector spaces, so we can represent it by a matrix

$$M(\mathcal{F}, \mathbf{E}; D) \in \mathbb{C}^{\#D \times (\sum_i \#E_i)}.$$

Matrices obtained by using the standard monomial bases for the vector spaces R_{E_i} and R_D in this representation are often called *Macaulay matrices*. We index the rows of the matrix with the exponents belonging to D and the columns with pairs $\{(i, \beta_i) : i \in \{1, \dots, s\}, \beta_i \in E_i\}$. The $(\alpha, (i, \beta_i))$ -entry of $M(\mathcal{F}, \mathbf{E}; D)$ contains the coefficient $c_{i, \alpha - \beta_i}$ of f_i , that is,

$$M(\mathcal{F}, \mathbf{E}; D)_{(\alpha, (i, \beta_i))} := c_{i, \alpha - \beta_i}.$$

Observe that this coefficient might be zero. The ordering of the exponents is of no importance in the scope of this work. We will therefore not specify it and assume that some ordering is fixed for all tuples A_i, E_i, D throughout the paper.

Example 1. To avoid subscripts, we replace the variables x_1 and x_2 by x and y , respectively. Consider the sets of exponents A_1, \dots, A_3 and the system $\mathcal{F} := (f_1, f_2, f_3)$ given by

$$\begin{aligned} A_1 &:= \{(0, 0), (1, 0), (0, 1), (0, 2)\}, & f_1 &:= -1 + 2x + 2y + y^2 \in R_{A_1}, \\ A_2 &:= \{(0, 0), (1, 0), (2, 0), (0, 1)\}, & f_2 &:= -1 + x + x^2 + y \in R_{A_2}, \\ A_3 &:= \{(0, 0), (1, 0), (2, 0), (0, 1)\}, & f_3 &:= -1 + 2x + 2x^2 + y \in R_{A_3}. \end{aligned}$$

We construct the Macaulay matrix $M(\mathcal{F}, \mathbf{E}; D)$, where $\mathbf{E} := (E_1, E_2, E_3)$ and

$$\begin{aligned} E_1 &:= \{(0, 0), (1, 0)\}, & D &:= \{(0, 0), (1, 0), (2, 0), (0, 1), \\ E_2 = E_3 &:= \{(0, 0), (0, 1)\}, & & (1, 1), (2, 1), (0, 2), (1, 2)\}. \end{aligned}$$

$$M(\mathcal{F}, \mathbf{E}; D) = \begin{array}{c|cccccc} & f_1 & x f_1 & f_2 & y f_2 & f_3 & y f_3 \\ \hline 1 & -1 & & -1 & & -1 & \\ y & 2 & & 1 & -1 & 1 & -1 \\ y^2 & 1 & & & 1 & & 1 \\ x & 2 & -1 & 1 & & 2 & \\ xy & & 2 & & 1 & & 2 \\ xy^2 & & 1 & & & & \\ x^2 & & 2 & 1 & & 2 & \\ x^2 y & & & & 1 & & 2 \end{array}$$

△

Remark 2.1. Given $\zeta \in \mathbb{C}^n$ and a finite subset $E \subset \mathbb{N}^n$, we denote by ζ^E the row vector

$$(\zeta^\alpha : \alpha \in E).$$

The vector obtained by the product $\zeta^D \cdot M(\mathcal{F}, \mathbf{E}; D) \in \mathbb{C}^{\sum_i \#E_i}$ is indexed by the tuples $\{(i, \beta_i) : i \in \{1, \dots, s\}, \beta_i \in E_i\}$ and the (i, β_i) -entry is given by $f_i(\zeta) \zeta^{\beta_i}$. If ζ is a solution of \mathcal{F} , then ζ^D belongs to the cokernel of $M(\mathcal{F}, \mathbf{E}; D)$. Moreover, if $\zeta \in \mathbb{C}^n$ is such that $\zeta^{E_i} \neq 0$ for all i , the opposite implication also holds. This is the case, for instance, for any solution $\zeta \in (\mathbb{C} \setminus \{0\})^n$.

We define the value $\text{HF}(\mathcal{F}, \mathbf{E}; D)$ as the corank of $M(\mathcal{F}, \mathbf{E}; D)$:

$$\text{HF}(\mathcal{F}, \mathbf{E}; D) := \#D - \text{Rank}(M(\mathcal{F}, \mathbf{E}; D)).$$

Let $\text{Coker}(\mathcal{F}, \mathbf{E}; D) \in \mathbb{C}^{\text{HF}(\mathcal{F}, \mathbf{E}; D) \times \#D}$ be a cokernel matrix (or left null space matrix) of $M(\mathcal{F}, \mathbf{E}; D)$. That is, $\text{Coker}(\mathcal{F}, \mathbf{E}; D)$ has rank $\text{HF}(\mathcal{F}, \mathbf{E}; D)$ and

$$\text{Coker}(\mathcal{F}, \mathbf{E}; D) \cdot M(\mathcal{F}, \mathbf{E}; D) = 0$$

We will index the columns of $\text{Coker}(\mathcal{F}, \mathbf{E}; D)$ with the exponents in D .

Example 2 (Cont.). The system \mathcal{F} has one solution $(-1, 1) \in \mathbb{C}^2$. The vector

$$(-1, 1)^D = (1, 1, 1, -1, -1, -1, 1, 1)$$

belongs to the cokernel of $M(\mathcal{F}, \mathbf{E}; D)$. Moreover, we have that $\text{HF}(\mathcal{F}, \mathbf{E}; D) = 2$ and

$$\text{Coker}(\mathcal{F}, \mathbf{E}; D) = \begin{bmatrix} 1 & y & y^2 & x & xy & xy^2 & x^2 & x^2y \\ 1 & 1 & 1 & -1 & -1 & -1 & 1 & 1 \\ 0 & 0 & 0 & 0 & -1 & 2 & 0 & 1 \end{bmatrix}. \quad \triangle$$

Consider two finite sets of exponents $A_0, E_0 \in \mathbb{N}^n$ such that $A_0 + E_0 \subset D$. For each polynomial $f_0 \in R_{A_0}$, we define the matrix N_{f_0} as

$$N_{f_0} := \text{Coker}(\mathcal{F}, \mathbf{E}; D) \cdot M(f_0, E_0; D). \quad (2.1)$$

Observe that $N_{f_0} \in \mathbb{C}^{\text{HF}(\mathcal{F}, \mathbf{E}; D) \times \#E_0}$ and the columns of N_{f_0} are indexed by the exponents in E_0 (more precisely, by the pairs $(0, \alpha)$ for each $\alpha \in E_0$).

Lemma 2.1. *For any $f_0 \in R_{A_0}$ we have that*

$$\text{HF}((f_0, \mathcal{F}), (E_0, \mathbf{E}); D) = 0 \iff N_{f_0} \text{ has rank } \text{HF}(\mathcal{F}, \mathbf{E}; D),$$

where $(f_0, \mathcal{F}) = (f_0, f_1, \dots, f_s)$ and $(E_0, \mathbf{E}) = (E_0, E_1, \dots, E_s)$. Moreover, in that case, for every solution $\zeta \in \mathbb{C}^n$ of \mathcal{F} such that the vector ζ^D is non-zero, we have $f_0(\zeta) \neq 0$.

Proof. The \Rightarrow direction of the first statement follows directly from

$$\begin{pmatrix} N_{f_0} & 0 \end{pmatrix} = \text{Coker}(\mathcal{F}, \mathbf{E}; D) \cdot M((f_0, \mathcal{F}), (E_0, \mathbf{E}); D).$$

For the \Leftarrow direction, suppose that N_{f_0} has rank $\text{HF}(\mathcal{F}, \mathbf{E}; D)$. Then $\text{Coker}(f_0, E_0; D) \cap \text{Coker}(\mathcal{F}, \mathbf{E}; D) = \{0\}$, which implies that the cokernel of $M((f_0, \mathcal{F}), (E_0, \mathbf{E}); D)$ is trivial, and hence it has rank $\#D$. The second statement follows from the fact that $M((f_0, \mathcal{F}), (E_0, \mathbf{E}); D)$ has trivial cokernel; as ζ^D is a non-zero vector, if $f_0(\zeta) = 0$, by Remark 2.1, ζ^D belongs to the cokernel of $M((f_0, \mathcal{F}), (E_0, \mathbf{E}); D)$. \square

Example 3 (Cont.). We consider the sets of exponents A_0, E_0 and the polynomial f_0 given by

$$A_0 = \{(0, 0), (1, 0), (0, 1)\}, \quad E_0 = \{(0, 0), (1, 0), (0, 1)\}, \quad f_0 := 1 + 3x + y \in R_{A_0}.$$

In this case, we have

$$N_{f_0} = \begin{bmatrix} 1 & x & y \\ -1 & 1 & -1 \\ 0 & -1 & -3 \end{bmatrix}.$$

Observe that, even though $1 \in R_{A_0}$, the matrix N_1 is not full-rank. \triangle

In what follows, we say that a property holds for *generic* points of a vector space if it holds for all points not contained in a subset of Lebesgue measure zero. Note that N_{f_0} is a matrix whose entries depend linearly on the coefficients of f_0 . This means that if there exists $f_0 \in R_{A_0}$ such that N_{f_0} has rank $\text{HF}(\mathcal{F}, \mathbf{E}; D)$, then $\text{rank}(N_h) = \text{HF}(\mathcal{F}, \mathbf{E}; D)$ for generic elements $h \in R_{A_0}$. Below, we assume that there is $f_0 \in R_{A_0}$ such that $\text{rank}(N_{f_0}) = \text{HF}(\mathcal{F}, \mathbf{E}; D)$ and we fix such an $f_0 \in R_{A_0}$. This assumption is very mild and given $\mathcal{F}, A_0, \mathbf{E}, D$, it is easy to check if it holds.

For ease of notation, we will write $\gamma := \text{HF}(\mathcal{F}, \mathbf{E}; D)$. Given a set of exponents $B \subset E_0$, we define the submatrix $N_{f_0, B} = \text{Coker}(\mathcal{F}, \mathbf{E}; D) \cdot \mathbf{M}(f_0, B; D) \in \mathbb{C}^{\gamma \times \#B}$ of N_{f_0} consisting of its columns indexed by B . We fix $B \subset E_0$ of cardinality γ such that $N_{f_0, B} \in \mathbb{C}^{\gamma \times \gamma}$ is invertible. For each $g \in R_{A_0}$, we define the matrix $M_g \in \mathbb{C}^{\gamma \times \gamma}$, defined as

$$M_g := N_{g, B} \cdot N_{f_0, B}^{-1}. \quad (2.2)$$

Example 4 (Cont.). We fix the basis $B = \{1, x\}$ and the matrix $N_{f_0, B} = \begin{bmatrix} -1 & 1 \\ 0 & -1 \end{bmatrix}$. Then, for $g = -1 + 3x + 2y$, we have

$$M_g = \begin{bmatrix} -2 & 2 \\ 0 & -2 \end{bmatrix} \cdot \begin{bmatrix} -1 & -1 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \quad \text{and} \quad M_x = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \quad \triangle$$

Remark 2.2. The map $g \in R_{A_0} \mapsto M_g \in \mathbb{C}^{\gamma \times \gamma}$ is a linear map. That is, for $\lambda \in \mathbb{C}$, $g_1, g_2 \in R_{A_0}$,

$$M_{g_1 + \lambda g_2} = M_{g_1} + \lambda M_{g_2}.$$

Moreover, M_{f_0} is the identity matrix.

A key observation is that we can solve the system of equations $\mathcal{F}(x) = 0$ by computing the eigenstructure of these matrices M_g , for $g \in R_{A_0}$. For that, we adapt the classical eigenvalue theorem from computational algebraic geometry (see Remark 2.3). We say that a non-zero row vector v is a *left eigenvector* of a matrix M with corresponding eigenvalue λ if it satisfies $v \cdot M = \lambda v$.

Theorem 2.1 (Eigenvalue theorem). *Using the notation introduced above, consider a polynomial system \mathcal{F} and a polynomial f_0 such that N_{f_0} has full-rank; see Equation (2.1). For each solution $\zeta \in \mathbb{C}^n$ of \mathcal{F} such that $\zeta^D \neq 0$, M_g from (2.2) has a left eigenvector v_ζ such that $v_\zeta \cdot \text{Coker}(\mathcal{F}, \mathbf{E}; D) = \zeta^D$. The corresponding eigenvalue is $\frac{g}{f_0}(\zeta)$. Conversely, if v is a left eigenvector of M_g such that $v \cdot \text{Coker}(\mathcal{F}, \mathbf{E}; D)$ is proportional to $\zeta^D \neq 0$ for some $\zeta \in \mathbb{C}^n$ such that $\zeta^{E_i} \neq 0$ for all i , then ζ is a solution of \mathcal{F} . Moreover, the corresponding eigenvalue of M_g is $\frac{g}{f_0}(\zeta)$.*

Remark 2.3. In some cases, the previous theorem can be derived from the classical eigenvalue theorem from computational algebraic geometry, where M_g represents the multiplication map

$$M_g : R / \langle \mathcal{F} \rangle \rightarrow R / \langle \mathcal{F} \rangle, h \mapsto M_g(h) = hg, \quad \text{for } g \in R.$$

Here $\langle \mathcal{F} \rangle \subset R$ is the ideal generated by f_1, \dots, f_s . As we will see (Example 5), this is not always the case. In the context of computer algebra, the eigenvalue theorem was introduced in [37] (eigenvalues) and in [1] (eigenvectors). For a historic overview and a proof in terms of matrices, see [21] and [27] respectively.

To prove Theorem 2.1, we need two auxiliary lemmas.

Lemma 2.2. Let $\zeta \in \mathbb{C}^n$ be a solution of \mathcal{F} such that $\zeta^D \neq 0$ and let $g \in R_{A_0}$ be such that $g(\zeta) = 0$. Then, the matrix M_g is singular.

Proof. By Remark 2.1, ζ^D belongs to the cokernel of $M(\mathcal{F}, \mathbf{E}; D)$, hence there is a row vector $v_\zeta \in \mathbb{C}^\gamma \setminus \{0\}$ such that $v_\zeta \cdot \text{Coker}(\mathcal{F}, \mathbf{E}; D) = \zeta^D$. Moreover, ζ^D belongs to the cokernel of $M(g, E_0; D)$. Hence, $v_\zeta \cdot N_g = 0$ and so $v_\zeta \cdot N_{g,B} = 0$. \square

Lemma 2.3. Let $\zeta \in \mathbb{C}^n$ be a solution of \mathcal{F} . If $\zeta^B = 0$, then $\zeta^D = 0$.

Proof. Since ζ is a solution of \mathcal{F} , there is a row vector v_ζ such that $v_\zeta \cdot \text{Coker}(\mathcal{F}, \mathbf{E}; D) = \zeta^D$. By observing that $N_{f_0,B} = \text{Coker}(\mathcal{F}, \mathbf{E}; D) \cdot M(f_0, B; D)$, the lemma follows from

$$\begin{aligned} v_\zeta \cdot N_{f_0,B} &= (v_\zeta \cdot \text{Coker}(\mathcal{F}, \mathbf{E}; D)) \cdot M(f_0, B; D) \\ &= \zeta^D \cdot M(f_0, B; D) \\ &= (\zeta^\alpha f_0(\zeta) : \alpha \in B) = 0, \end{aligned}$$

where the last line uses $\zeta^B = 0$. Since $N_{f_0,B}$ is invertible, this implies $v_\zeta = 0$, and thus $\zeta^D = 0$. \square

Proof of Theorem 2.1. The proof is based on the following observations. By Remark 2.2, the eigenvalues of M_g correspond to the values $\lambda \in \mathbb{C}$ such that $M_g - \lambda \text{id} = M_{g-\lambda f_0}$ is singular. By Lemma 2.1, we have $f_0(\zeta) \neq 0$ for each solution ζ of \mathcal{F} . Let ζ be a solution of \mathcal{F} . If $\lambda = \frac{g}{f_0}(\zeta)$, the polynomial $g - \lambda f_0 \in R_{A_0}$ vanishes at ζ . As by assumption $\zeta^D \neq 0$, from Lemma 2.2 we deduce that $M_{g-\lambda f_0}$ is singular. Therefore, $\frac{g}{f_0}(\zeta)$ is an eigenvalue of M_g . For the associated left eigenvector, let v_ζ be as in the proof of Lemma 2.2. We have $v_\zeta \cdot N_{g-\lambda f_0,B} = v_\zeta \cdot (N_{g,B} - \lambda N_{f_0,B}) = 0$ and multiplying from the right by $N_{f_0,B}^{-1}$ gives $v_\zeta \cdot (M_g - \lambda \text{id}) = 0$.

Conversely, suppose that v is a left eigenvector of M_g such that $v \cdot \text{Coker}(\mathcal{F}, \mathbf{E}; D) = \zeta^D \neq 0$ for some $\zeta \in \mathbb{C}^n$ (we may assume equality after scaling). By Remark 2.1, under the assumption $\zeta^{E_i} \neq 0$ for all i , ζ is a solution of \mathcal{F} , see Remark 2.1. We now compute the corresponding eigenvalue. By definition, $v \cdot (M_g - \lambda \text{id}) = 0$ for some λ . Multiplying from the right by $N_{f_0,B}$ we see that $v \cdot N_{g-\lambda f_0,B} = (v \cdot \text{Coker}(\mathcal{F}, \mathbf{E}; D)) \cdot M(g - \lambda f_0, B, D) = \zeta^D \cdot M(g - \lambda f_0, B, D) = 0$. By Lemma 2.3, $\zeta^B \neq 0$ and since $f_0(\zeta) \neq 0$ (Lemma 2.1) we conclude $\lambda = \frac{g}{f_0}(\zeta)$. \square

Example 5 (Cont.). The unique eigenvalue of $M_g = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$ is $2 = \frac{g}{f_0}(-1, 1)$. Moreover, we have that $1 = \frac{x}{f_0}(-1, 1)$ is an eigenvalue of $M_x = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ whose associated eigenvector $(1, 0)$ satisfies $(1, 0) \cdot \text{Coker}(\mathcal{F}, \mathbf{E}; D) = (-1, 1)^D$. Observe that, as the system \mathcal{F} has only one solution, namely $(-1, 1)$, with multiplicity one, the matrices M_x and M_g are not multiplication operators. \triangle

We now characterize row vectors v that are an eigenvector of all matrices in $\mathcal{M} := \{M_h : h \in R_{A_0}\}$. Observe that, by Remark 2.2, \mathcal{M} is a vector space. For any nonzero row vector $v \in \mathbb{C}^\gamma \setminus \{0\}$, we define the subspace

$$\mathcal{M}(v) = \{M_h \in \mathcal{M} : v \text{ is a left eigenvector of } M_h\}.$$

One can check that $\mathcal{M}(v)$ is a vector space. We say that v is an eigenvector of \mathcal{M} if $\mathcal{M}(v) = \mathcal{M}$.

Example 6 (Cont.). Any vector in \mathbb{C}^2 is an eigenvector of $M_g = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$, but for $h = 1 + x + y$, we obtain $M_h = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$, which has left eigenvectors $(1, 0)$ and $(0, 1)$. These vectors are also eigenvectors of $M_x = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$. It is straightforward to check that $\{M_g, M_h, M_x\}$ generate \mathcal{M} as a vector space, so that $(1, 0)$ and $(0, 1)$ are eigenvectors of \mathcal{M} . This means that there might be eigenvectors of \mathcal{M} which are not associated to solutions of \mathcal{F} . Also, observe that the matrices in \mathcal{M} commute. \triangle

Proposition 2.1. *Fix a non-zero vector $v \in \mathbb{C}^\gamma$. We have that v is an eigenvector of \mathcal{M} if and only if it is a left eigenvector of M_h for a generic $h \in R_{A_0}$.*

Proof. The ‘only if’ direction is clear. For the ‘if’ direction, let $U \subsetneq R_{A_0}$ be the subset of polynomials h for which v is an eigenvector of M_h . It is easy to see that U is a linear subspace, so that it is closed in the Zariski topology. It is also dense by assumption, so we conclude that $U = R_{A_0}$. \square

Proposition 2.2. *Consider eigenvectors v_1, \dots, v_m of \mathcal{M} . We have that v_1, \dots, v_m correspond to the same eigenvalue of M_h for all $M_h \in \mathcal{M}$ if and only if they correspond to the same eigenvalue of M_h for a generic $h \in R_{A_0}$.*

Proof. Again, the ‘only if’ direction is clear. For the opposite implication, note that

$$W = \{h \in R_{A_0} : v_1, \dots, v_m \text{ correspond to the same eigenvalue of } M_h\}$$

is a vector subspace of R_{A_0} . Since it is also dense by hypothesis, we conclude that $W = R_{A_0}$ as in the proof of Proposition 2.1. \square

Example 7 (Cont.). Instead of the basis B fixed in Example 4, in what follows we consider $B = \{x, y\}$ and so $N_{f_0, B'} = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$. This way, we obtain the following matrices:

$$M'_g = N_{g, B'} N_{f_0, B'}^{-1} = \begin{bmatrix} 2 & 0 \\ -\frac{3}{4} & \frac{5}{4} \end{bmatrix} \quad \text{and} \quad M'_x = \begin{bmatrix} 1 & 0 \\ \frac{1}{4} & \frac{1}{4} \end{bmatrix}$$

While $(1, 0)$ is a common left eigenvector of M'_g and M'_x corresponding to the unique solution of the system, the vector $(1, 1)$ is a left eigenvector of M'_g , but it is not of M'_x . Comparing this example with Example 6, we observe that, depending on the choice of the basis B , there may be spurious common eigenvectors of \mathcal{M} . \triangle

Propositions 2.1 and 2.2 give a simple procedure for computing, for a given eigenvalue λ_g of M_g , the intersection of the corresponding left eigenspace of M_g with the eigenvectors of \mathcal{M} . Suppose that this eigenspace is spanned by the rows of the matrix V_{λ_g} . By Proposition 2.1, we simply need to check which elements in the row span of V_{λ_g} are also eigenvectors of M_h , for a random element $h \in R_{A_0}$. Proposition 2.2 guarantees that these eigenvectors, if they exist, belong to a unique eigenvalue of M_h . This is summarized in Algorithm 1. In line 8 of the algorithm, we solve the generalized eigenvalue problem (GEP) given by the pencil $(B_1, B_2) := (V_{\lambda_g} \cdot M_h \cdot O, V_{\lambda_g} \cdot O)$, that is, we compute all eigenvalues μ_i and a basis for the left eigenspace $C_i = \{c_i \in \mathbb{C}^m \mid c_i B_1 = \mu_i c_i B_2\}$. In line 9, we select (if possible) the unique eigenvalue μ_i whose corresponding eigenspace C_i gives the desired intersection $\mathcal{V} = C_i \cdot V_{\lambda_g}$. Proposition 2.2 also has the following direct corollary.

Corollary 2.1. *Let λ_g be an eigenvalue of M_g and let $\mathcal{V} \in \mathbb{C}^{m \times \gamma}$ be a matrix whose rows are a basis for the left eigenspace of M_g corresponding to λ_g , intersected with the eigenvectors of \mathcal{M} . If g is generic, there is exactly one tuple $(\lambda_\alpha)_{\alpha \in A_0}$ such that*

$$\mathcal{V} M_{x^\alpha} = \lambda_\alpha \mathcal{V}, \quad \text{for all } \alpha \in A_0.$$

Algorithm 1 GETEIGENSPACE

Input: An eigenvalue λ_g of M_g for generic $g \in R_{A_0}$, a matrix V_{λ_g} of size $m \times \gamma$ whose rows contain a basis for the corresponding eigenspace and the matrix M_h for a generic $h \in R_{A_0}$

Output: A matrix \mathcal{V} whose rows are a basis for the intersection of the row span of V_{λ_g} with the eigenvectors of \mathcal{M} .

```

1: if  $m = 1$  then
2:   if  $\text{rank} \begin{bmatrix} V_{\lambda_g} \\ V_{\lambda_g} \cdot M_h \end{bmatrix} = 1$  then
3:      $\mathcal{V} \leftarrow V_{\lambda_g}$ 
4:   else
5:      $\mathcal{V} \leftarrow \{0\}$ 
6:   else
7:      $O \leftarrow$  random matrix of size  $\gamma \times m$ 
8:      $\{(\mu_i, C_i)\} \leftarrow$  solve the GEP  $c_i \cdot (V_{\lambda_g} \cdot M_h \cdot O) = \mu_i c_i \cdot (V_{\lambda_g} \cdot O)$ 
9:      $C \leftarrow C_i$  such that  $C_i \cdot V_{\lambda_g} \cdot M_h = \mu_i \cdot C_i \cdot V_{\lambda_g}$ 
10:    if  $C$  is empty then
11:       $\mathcal{V} \leftarrow \{0\}$ 
12:    else
13:       $\mathcal{V} \leftarrow C \cdot V_{\lambda_g}$ 
14:  return  $\mathcal{V}$ 

```

Remark 2.4. This has the practical implication that $\tilde{M} = \mathcal{V}M_{x^\alpha}T(\mathcal{V}T)^{-1} = \text{diag}(\lambda_\alpha, \dots, \lambda_\alpha)$ for a random matrix $T \in \mathbb{C}^{\gamma \times m}$ has only one eigenvalue λ_α , equal to $\text{trace}(\tilde{M})/m$. If \mathcal{V} has only one row we obtain λ_α from the Rayleigh quotient $\lambda_\alpha = \mathcal{V}M_{x^\alpha}\mathcal{V}^*/(\mathcal{V}\mathcal{V}^*)$, where $*$ is the conjugate transpose.

Proposition 2.3 (Criterion for eigenvalues). *Let λ_g be an eigenvalue of M_g . If $\lambda_g = \frac{g}{f_0}(\zeta)$ for some solution $\zeta \in \mathbb{C}^n$ of \mathcal{F} satisfying $\zeta^D \neq 0$, then the tuple $(\lambda_\alpha)_{\alpha \in A_0}$ from Corollary 2.1 satisfies*

$$C \cdot (\lambda_\alpha)_{\alpha \in A_0} = \zeta^{A_0} \text{ for some } C \in \mathbb{C} \setminus \{0\}.$$

Proof. Let $\mathcal{V} \in \mathbb{C}^{m \times \gamma}$ be a matrix whose rows are a basis for the left eigenspace of M_g corresponding to λ_g intersected with the eigenvectors of \mathcal{M} . If $\lambda_g = \frac{g}{f_0}(\zeta)$ for some solution $\zeta \in \mathbb{C}^n$ of \mathcal{F} satisfying $\zeta^D \neq 0$, then by Theorem 2.1 we know that v_ζ is a corresponding eigenvector of \mathcal{M} . Therefore, there exists $c_\zeta \in \mathbb{C}^m \setminus \{0\}$ such that $c_\zeta \mathcal{V} = v_\zeta$. Another consequence of Theorem 2.1 is

$$c_\zeta \mathcal{V}M_{x^\alpha} = \frac{x^\alpha}{f_0}(\zeta)c_\zeta \mathcal{V}. \quad \square$$

The results discussed above suggest several ways of extracting the coordinates of a solution $\zeta \in \mathbb{C}^n$ of $\mathcal{F}(x) = 0$ from the eigenstructure of the matrices M_g . Both the eigenvectors (Theorem 2.1) and the eigenvalues (Proposition 2.3) reveal vectors of the form ζ^A for some set of exponents $A \subset \mathbb{N}^n$. We now recall how to compute the coordinates of ζ from the vector ζ^A and discuss the assumptions that we need on A in order to be able to do this.

For any subset $A \subset \mathbb{N}^n$, we write

$$\mathbb{N}A := \left\{ \sum_{\alpha \in A} n_\alpha \cdot \alpha : n_\alpha \in \mathbb{N} \right\} \subset \mathbb{N}^n, \quad \mathbb{Z}A := \left\{ \sum_{\alpha \in A} m_\alpha \cdot \alpha : m_\alpha \in \mathbb{Z} \right\} \subset \mathbb{Z}^n.$$

If $A = \{\alpha_1, \dots, \alpha_k\}$ with $\alpha_1 = 0$ and the condition $\mathbb{Z}A = \mathbb{Z}^n$ is satisfied, then for $\ell = 1, \dots, n$, there exist integers $m_{2,\ell}, \dots, m_{k,\ell}$ such that $m_{2,\ell}\alpha_2 + \dots + m_{k,\ell}\alpha_k = e_\ell$, where e_ℓ is the ℓ -th standard basis vector of \mathbb{Z}^n . These integers $m_{j,\ell}$ can be computed, for instance, using the Smith normal form of an integer matrix whose columns are the elements of A . If this is the case, from $\zeta^A = (\zeta^{\alpha_1}, \dots, \zeta^{\alpha_k})$ we can compute the ℓ -th coordinate ζ_ℓ of ζ as

$$\zeta_\ell = \prod_{j=2}^k (\zeta^{\alpha_j})^{m_{j,\ell}}, \quad \ell = 1, \dots, n. \quad (2.3)$$

This approach can be used to compute the coordinates of $\zeta \in (\mathbb{C} \setminus \{0\})^n$ from ζ^A , i.e. all points with all non-zero coordinates. Note that some of the $m_{j,\ell}$ may be negative, which may be problematic in the case where ζ has zero coordinates. If the stronger condition $\mathbb{N}A_0 = \mathbb{N}^n$ is satisfied (this implies $e_\ell \in A$, $\ell = 1, \dots, n$), then the integers $m_{j,\ell}$ can be taken non-negative and we can obtain the coordinates of all points ζ in \mathbb{C}^n from ζ^A . We will continue under the assumption that we are mostly interested in computing points in $(\mathbb{C} \setminus \{0\})^n$, as this is commonly assumed in a sparse setting. However, solutions in \mathbb{C}^n can be computed by replacing $\mathbb{Z}A = \mathbb{Z}^n$ in what follows by the stronger assumption $\mathbb{N}A = \mathbb{N}^n$. Note that if $\mathbb{Z}A = \mathbb{Z}^n$, the outlined approach suggests a way of checking whether or not a vector $q \in \mathbb{C}^{\#A}$ with $q_{\alpha_1} = 1$ is of the form ζ^A for some $\zeta \in (\mathbb{C} \setminus \{0\})^n$. Indeed, one computes the coordinates $\zeta_\ell = \prod_{j=2}^k (q_{\alpha_j})^{m_{j,\ell}}$ and checks whether $\zeta^A = q$.

We turn to the eigenvalue method for extracting the roots ζ from the matrices M_g . Let $\{\zeta_1, \dots, \zeta_\delta\} \subset \mathbb{C}^n$ be a set of solutions of \mathcal{F} such that $\zeta_i^D \neq 0$ for all i . By Theorem 2.1, for each of these solutions there is an eigenvalue λ_g of the matrix M_g and a space of dimension $m \geq 1$, spanned by the rows of a matrix \mathcal{V} , of eigenvectors of \mathcal{M} . Suppose we have computed this matrix \mathcal{V} (for instance, using Algorithm 1). We write $A_0 = \{\alpha_1, \dots, \alpha_k\} \subset \mathbb{N}^n$ and assume that $\alpha_1 = 0$. The unique eigenvalue (Corollary 2.1) of $M_{x^{\alpha_j}}$ corresponding to \mathcal{V} is denoted by λ_{ij} and can be computed using Remark 2.4. As $\zeta_i^D \neq 0$, by Theorem 2.1, there is $v \in \mathcal{V}$ such that $v \cdot \text{COKER}(F, E, D) = C \cdot \zeta_i^D$, for non-zero $C \in \mathbb{C}$. Therefore, by Theorem 2.1,

$$\lambda_{ij} = \frac{\zeta_i^{\alpha_j}}{f_0(\zeta_i)} \quad \text{and hence} \quad \frac{\lambda_{ij}}{\lambda_{i1}} = \zeta_i^{\alpha_j}, \quad i = 1, \dots, \delta, j = 1, \dots, k. \quad (2.4)$$

We would like to recover the coordinates of ζ_i from the tuple $(\zeta_i^{\alpha_2}, \dots, \zeta_i^{\alpha_k}) \in \mathbb{C}^{k-1}$. Assuming $\mathbb{Z}A_0 = \mathbb{Z}^n$ and applying (2.3), we find

$$\zeta_{i,\ell} = \prod_{j=2}^k \left(\frac{\lambda_{ij}}{\lambda_{i1}} \right)^{m_{j,\ell}}, \quad \ell = 1, \dots, n. \quad (2.5)$$

Remark 2.5. In many cases, one can take $A_0 = \{0, e_1, \dots, e_n\}$, in which case $m_{j,\ell} = 1$ if $j = \ell + 1$ and $m_{j,\ell} = 0$ otherwise.

Motivated by this discussion, we make the following definition.

Definition 2.1. We say that a tuple $(\mathcal{F} = (f_1, \dots, f_s), A_0, (E_0, \mathbf{E}) = (E_0, \dots, E_s), D)$ is *admissible* if it satisfies the following three conditions,

- **Compatibility condition:** For $i = 0, \dots, s$, $A_i + E_i \subset D$.

- **Rank condition:** There exists $f_0 \in R_{A_0}$ such that $\text{rank}(N_{f_0}) = \text{HF}(\mathcal{F}, \mathbf{E}; D)$.
- **Lattice condition:** The set A_0 satisfies $0 \in A_0$ and $\mathbb{Z}A_0 = \mathbb{Z}^n$.

The results in this section lead to Algorithm 2 for solving $\mathcal{F}(x) = 0$, given an admissible tuple $(\mathcal{F}, A_0, (E_0, \mathbf{E}), D)$. This algorithm computes a candidate set of solutions containing every solution in $(\mathbb{C} \setminus \{0\})^n$. It might contain spurious points, since there might be eigenvalues that do not correspond to solutions but do come from a common eigenvector of \mathcal{M} , see Example 5. One can identify these points, for instance, by evaluating the *relative backward error*, see Equation (4.1).

In what follows, we discuss some aspects of the algorithm in more detail. In practice, the number of columns $\sum_{i=1}^s \#E_i$ of the Macaulay matrix $M(\mathcal{F}, \mathbf{E}; D)$ is often much larger than the number $\#D$ of rows. Multiplying from the right by a random matrix of size $(\sum_{i=1}^s \#E_i) \times \#D$ does not affect the left nullspace, but reduces the complexity of computing it. This is what happens in line 2. See [40, Sec. 4.2] for details. If $(\sum_{i=1}^s \#E_i) \lesssim \#D$, that is, the number of columns is not much larger than the number of rows, this step can be skipped.

Remark 2.6. *By the lattice condition, we have that $1 \in R_{A_0}$. However, the rank condition might not be satisfied for $f_0 = 1$. That is, it might happen that $\text{rank}(N_1) < \text{Coker}(\mathcal{F}, \mathbf{E}; D)$. This is the case, for instance, in Example 3. To overcome this issue, we choose f_0 randomly in R_{A_0} .*

Numerical considerations. In theory, we may pick B arbitrary such that $N_{f_0, B}$ is an invertible matrix. In practice, *it is crucial to pick B such that $N_{f_0, B}$ is well-conditioned*. This was shown in [46, 47]. For that, we select a random f_0 and, in line 6, we use a standard numerical linear algebra procedure for selecting a well-conditioned submatrix from N_{f_0} : *QR factorization with optimal column pivoting*. This computes matrices Q_0, R_0 and a permutation $p = (p_1, \dots, p_{\#E_0})$ of the columns of N_{f_0} such that $N_{f_0}[:, p] = Q_0 R_0$, where $Q_0 \in \mathbb{C}^{\gamma \times \gamma}$ is a unitary matrix, $R_0 \in \mathbb{C}^{\gamma \times \#E_0}$ is upper triangular and $N_{f_0}[:, p]$ is N_{f_0} with its columns permuted according to p . The leftmost γ columns of R_0 form the square, upper triangular matrix \hat{R}_0 . The column permutation p is such that columns p_1, \dots, p_γ form a well-conditioned submatrix of N_{f_0} . In line 8, these columns are selected to form the matrix $N_{f_0, B}$. Using the identities $N_{f_0, B}^* M_g^* = N_{g, B}^*$, where $*$ is the conjugate transpose, and $N_{f_0, B} = Q_0 \hat{R}_0$, we see that the solution $Q_0^* M_g^* Q_0$ to the linear system $\hat{R}_0^* X = N_{g, B}^* Q_0$ is similar to the matrix M_g^* in this section, and it can be obtained by back substitution since R_0^* is lower triangular. Since we extract the coordinates of the roots from the eigenvalues, not the eigenvectors, we may work with $Q_0^* M_g^* Q_0$ as well. This is exploited in line 11. In line 17, we invoke Algorithm 1. Lines 18-25 are a straightforward implementation of Remark 2.4. As pointed out, in the case $m = 1$, λ_{ij} can alternatively be computed as a Rayleigh quotient.

Remark 2.7. *Alternatively, by Theorem 2.1, when \mathcal{V} is one-dimensional, we may check if, for a vector $v \in \mathcal{V}$, there is a non-zero constant C and $\zeta \in \mathbb{C}^n$ such that $C \cdot \zeta^D = v_\zeta \cdot \text{Coker}(\mathcal{F}, \mathbf{E}; D)$. If $0 \in D$ and $\mathbb{Z}D = \mathbb{Z}^n$, we scale v such that $C = 1$ and find ζ from ζ^D as above. When the matrices M_g are multiplication operators, this approach is usually referred as the eigenvector criterion [1]. This idea can be extended to the case where \mathcal{V} has dimension > 1 . Extracting vectors of the form ζ^D from a vector space can be viewed as a harmonic retrieval problem, see [49, Sec. 3.3].*

Theorem 2.2 (Correctness). *Algorithm 2 computes a set of points containing every solution of \mathcal{F} in the algebraic torus $(\mathbb{C} \setminus \{0\})^n$.*

Algorithm 2 SOLVE

Input: An admissible tuple $(\mathcal{F}, A_0, (E_0, \mathbf{E}), D)$ with $A_0 = \{\alpha_1 = 0, \alpha_2, \dots, \alpha_k\}$

Output: A candidate set of solutions of \mathcal{F} , containing all solutions in $(\mathbb{C} \setminus \{0\})^n$

- 1: $O \leftarrow$ random matrix of size $(\sum_{i=1}^s \#E_i) \times \#D$
 - 2: $MO \leftarrow M(\mathcal{F}, \mathbf{E}; D) \cdot O$
 - 3: Compute $\text{Coker}(\mathcal{F}, \mathbf{E}; D)$ via the SVD of MO
 - 4: $f_0 \leftarrow$ random element in R_{A_0}
 - 5: $N_{f_0} \leftarrow$ matrix of size $\gamma \times (\#E_0)$ given by $\text{Coker}(\mathcal{F}, \mathbf{E}; D) \cdot M(f_0, E_0; D)$.
 - 6: $Q_0, R_0, p \leftarrow$ apply QR decomposition with optimal pivoting to N_{f_0}
 - 7: $\hat{R}_0 \leftarrow$ square, upper triangular matrix given by the first γ columns of R_0
 - 8: $B \leftarrow$ exponents in E_0 corresponding to columns p_1, \dots, p_γ of N_{f_0}
 - 9: **for** $j = 1, \dots, k$ **do**
 - 10: $N_{x^{\alpha_j}, B} \leftarrow \text{Coker}(\mathcal{F}, \mathbf{E}; D) \cdot M(x^{\alpha_j}, B; D)$
 - 11: $M_{x^{\alpha_j}}^* \leftarrow$ solve $\hat{R}_0^* X = N_{x^{\alpha_j}, B}^* Q_0$ for X by back substitution
 - 12: $M_g \leftarrow$ random linear combination of $M_{x^\alpha}, \alpha \in A_0$
 - 13: $\{(\mu_1, V_{\mu_1}), \dots, (\mu_\delta, V_{\mu_\delta})\} \leftarrow$ distinct eigenvalues of M_g and corresponding left eigenspaces
 - 14: $M_h \leftarrow$ a different random linear combination of $M_{x^\alpha}, \alpha \in A_0$
 - 15: $Z \leftarrow \{\}$
 - 16: **for** $i = 1, \dots, \delta$ **do**
 - 17: $\mathcal{V} \leftarrow \text{GETEIGENSPACE}(\mu_i, V_{\mu_i}, M_h)$
 - 18: **if** $\mathcal{V} \neq \{0\}$ **then**
 - 19: $m \leftarrow$ number of rows of \mathcal{V}
 - 20: $T \leftarrow$ random matrix of size $\gamma \times m$
 - 21: **for** $j = 1, \dots, k$ **do**
 - 22: $\tilde{M} \leftarrow \mathcal{V} M_{x^{\alpha_j}} T (\mathcal{V} T)^{-1}$
 - 23: $\lambda_{i,j} \leftarrow \text{trace}(\tilde{M})/m$ (when $m = 1$, use Rayleigh quotient, see Remark 2.4)
 - 24: $\zeta_i \leftarrow$ if possible, compute the coordinates of ζ_i via (2.3) and (2.4)
 - 25: $Z \leftarrow Z \cup \{\zeta_i\}$
 - 26: **return** Z
-

Proof. As our input is an admissible tuple, the compatibility condition implies that the matrix N_{f_0} is well-defined. By the rank condition and the fact that f_0 is generic, N_{f_0} has full rank. See the discussion below Lemma 2.1. Hence, the matrices M_g and M_h are well-defined and agree with the ones defined in (2.2). Let ζ_1 be a solution of \mathcal{F} such that $\zeta_1 \in (\mathbb{C} \setminus \{0\})^n$. As $\zeta_1^D \neq 0$, by Theorem 2.1, we can assume with no loss of generality that $\mu_1 = \frac{g}{f_0}(\zeta_1)$. Let $\mathcal{V} := \text{GETEIGENSPACE}(\mu_1, V_{\mu_1}, M_h)$, for generic $h \in R_{A_0}$. As h is generic, by Proposition 2.2, all vectors in \mathcal{V} belong to the same eigenvalue of $M_{x^{\alpha_j}}$, for $j = 1, \dots, k$. Hence, by Proposition 2.3, there is a non-zero constant $C \in \mathbb{C}$ such that the element $\lambda_{1,j}$ computed in line 23 agrees with $C \zeta_1^{\alpha_j}$, for $\alpha_j \in A_0$. Observe that, as $\zeta_1 \in (\mathbb{C} \setminus \{0\})^n$, $\lambda_{1,j} \neq 0$. Therefore, as the admissible tuple satisfies the lattice condition $\mathbb{Z}A_0 = \mathbb{Z}^n$, we can recover the coordinates of ζ_1 using (2.4) and $\zeta_1 \in Z$. \square

Remark 2.8. By adapting the previous proof, we observe that algorithm 2 computes every solution ζ of \mathcal{F} such that $\zeta^D \neq 0$ and the formula given in (2.4) is well-defined. That is, for every $j \in \{1, \dots, k\}$ and $\ell \in \{1, \dots, n\}$, $m_{j,\ell} \geq 0$ or $\zeta^{\alpha_j} \neq 0$.

It is clear that the size of the matrices in Algorithm 2 depends on the cardinality of the exponent sets in the admissible tuple. Constructing admissible tuples for certain families of polynomial systems is an active field of research, strongly related to the study of *regularity* of ideals in polynomial rings, in the sense of commutative algebra [26, Sec. 20.5]. Recent progress in this area, for the case where $n = s$, was made in [11]. In the next section, we will summarize some of these results by explicitly describing some admissible tuples for systems with important types of structures.

As mentioned above, the matrices M_g considered in this section play the role of *multiplication operators* in the algebra R/I , where I is the ideal generated by the polynomials in \mathcal{F} [22, Ch. 2]. In the very general setting we consider here, assuming only that $(\mathcal{F}, A_0, (E_0, \mathbf{E}), D)$ is an admissible tuple, the matrices M_g do not necessarily represent such multiplication operators. However, under some extra assumptions, they do commute. In this case, we can simplify Algorithm 2 by computing the simultaneous Schur factorization of $(M_{x^\alpha})_{\alpha \in A_0}$ as in [11, Sec. 3.3].

Theorem 2.3 (Criterion for commutativity). *Let $(\mathcal{F}, A_0, (E_0, \mathbf{E}), D)$ be an admissible tuple and $\gamma = \text{HF}(\mathcal{F}, \mathbf{E}; D)$. Let f_0 be such that N_{f_0} satisfies the Rank condition (Definition 2.1). If*

$$\text{HF}((f_0^2, \mathcal{F}), (E_0, E_1 + A_0, \dots, E_s + A_0); D + A_0) - \text{HF}(\mathcal{F}, (E_1 + A_0, \dots, E_s + A_0); D + A_0) = \gamma,$$

then for every $g_1, g_2 \in R_{A_0}$ and every valid choice of $B \subset A_0$, we have that $M_{g_1} M_{g_2} = M_{g_2} M_{g_1}$.

Proof. In what follows, we fix two vector spaces $I_D := \text{Im}(\text{Sylv}_{(\mathcal{F}, (E_1, \dots, E_s); D)})$ and $I_{D+A_0} := \text{Im}(\text{Sylv}_{(\mathcal{F}, (E_1+A_0, \dots, E_s+A_0); D+A_0)})$. Observe that, for every $g \in R_{A_0}$ and $f \in I_D$, $gf \in I_{D+A_0}$. We write $B = \{b_1, \dots, b_\gamma\}$ and given $v \in \mathbb{C}^\gamma$, we set $v \cdot B := \sum_i v_i x^{b_i}$.

In this proof, for each $g \in R_{A_0}$, we consider the map $\tilde{M}_g := N_{f_0, B}^{-1} \cdot M_g \cdot N_{f_0, B}$. The maps M_g and \tilde{M}_g are similar, so it is enough to prove that $\tilde{M}_{g_1} \tilde{M}_{g_2} = \tilde{M}_{g_2} \tilde{M}_{g_1}$. It is not hard to show that for $v, w \in \mathbb{C}^\gamma$, such that $\tilde{M}_g(v) = w \in \mathbb{C}^\gamma$, we have $g(v \cdot B) \equiv f_0(w \cdot B)$ modulo I_D .

First, observe that, for every $v \in \mathbb{C}^\gamma$, $g_1 g_2(v \cdot B) \equiv f_0^2((\tilde{M}_{g_1} \tilde{M}_{g_2} v) \cdot B)$ modulo I_{D+A_0} . Indeed, $g_1(v \cdot B) = f_0((\tilde{M}_{g_1} v) \cdot B) + h_1$, for $h_1 \in I_D$ and for $w = \tilde{M}_{g_1} v$, we have that $g_2(w \cdot B) = f_0((\tilde{M}_{g_2} w) \cdot B) + h_2$, for $h_2 \in I_D$. Hence, $g_1 g_2(v \cdot B) = g_2 f_0((\tilde{M}_{g_1} v) \cdot B) + g_2 h_1 = f_0^2((\tilde{M}_{g_2} \tilde{M}_{g_1} v) \cdot B) + f_0 h_2 + g_2 h_1$. As $f_0 h_2 + g_2 h_1 \in I_{D+A_0}$, the claim follows. Since $g_1 g_2 = g_2 g_1$, it also holds that $g_1 g_2(v \cdot B) \equiv f_0^2((\tilde{M}_{g_2} \tilde{M}_{g_1} v) \cdot B)$ modulo I_{D+A_0} .

Second, we show that $\{f_0^2 x^{b_i} : b_i \in B\}$ is a basis of the vector space V spanned by $\{f_0^2 x^e : e \in E_0\}$ modulo I_{D+A_0} . By construction of $B \subset E_0$, $\{f_0 x^{b_i} : b_i \in B\}$ is a basis of the vector space spanned by $\{f_0 x^e : e \in E_0\}$ modulo I_D , so $\{f_0^2 x^{b_i} : b_i \in B\}$ generates V . Moreover, by the assumption on the difference of coranks, the dimension of the vector space V is $\gamma = \#\{f_0^2 x^{b_i} : b_i \in B\}$.

By the first observation, we have that $f_0^2(((\tilde{M}_{g_1} \tilde{M}_{g_2} - \tilde{M}_{g_2} \tilde{M}_{g_1}) v) \cdot B) \equiv 0$ modulo I_{D+A_0} for every $v \in \mathbb{C}^\gamma$. By the second observation, the elements in $\{f_0^2 x^{b_i} : b_i \in B\}$ are linearly independent modulo I_{D+A_0} . Therefore, we have that $\tilde{M}_{g_1} \tilde{M}_{g_2} = \tilde{M}_{g_2} \tilde{M}_{g_1}$. \square

Remark 2.9. *This criterion is similar to Bayer and Stillman's criterion to compute the Castelnuovo-Mumford regularity of ideals defining a zero dimensional projective scheme [5, Thm. 1.10]. Under further assumptions on A_0, E_0, D , and B , the commutativity of the matrices implies that γ is the number of isolated solutions of the system \mathcal{F} , see [39, Thm. 3.1].*

Example 8 (Cont.). Theorem 2.3 is independent of the chosen basis B . Its hypotheses are not satisfied by the admissible tuple of our running example, as

$$\text{HF}((f_0^2, \mathcal{F}), (E_0, E_1 + A_0, \dots, E_s + A_0); D + A_0) - \text{HF}(\mathcal{F}, (E_1 + A_0, \dots, E_s + A_0); D + A_0) = 1 < 2.$$

However, as we showed in Example 6 for $B = \{1, x\}$, the matrices in \mathcal{M} do commute. \triangle

3 Construction of admissible tuples

In this section, we fix an s -tuple of sets of exponents $\mathbf{A} := (A_1, \dots, A_s)$, where $A_i \subset \mathbb{N}^n$, and consider a polynomial system $\mathcal{F} = (f_1, \dots, f_s) \in R_{A_1} \times \dots \times R_{A_s}$. We construct tuples that are admissible under mild assumptions on \mathcal{F} (Assumption 1). This allows us to compute the solutions of the system \mathcal{F} using Algorithm 2. Section 3.1 states explicit formulas for admissible tuples that in practice are near-optimal in the case where $s = n$. In the overdetermined case ($s > n$), we can obtain admissible tuples leading to smaller matrices by using incremental constructions. These are the topic of Subsection 3.2.

The section uses the following notation. The *convex hull* of a finite subset $E \subset \mathbb{R}^n$ is the polytope $\text{Conv}(E) \subset \mathbb{R}^n$ defined as,

$$\text{Conv}(E) := \left\{ \sum_{e \in E} \lambda_e e : \sum_{e \in E} \lambda_e = 1, \lambda_e \geq 0 \text{ for all } e \in E \right\}.$$

By a *lattice polytope* we mean a convex polytope $P \subset \mathbb{R}^n$ that arises as $\text{Conv}(E)$, where $E \subset \mathbb{N}^n$. Such a lattice polytope is called *full-dimensional* if it has a positive Euclidean volume in \mathbb{R}^n . Given two polytopes $P_1, P_2 \subset \mathbb{R}^n$ and $c \in \mathbb{N}$, we denote by $P_1 + P_2$ the *Minkowski sum* of P_1, P_2 and by $c \cdot P_1$ the *c-dilation* of P_1 , that is,

$$P_1 + P_2 := \{\alpha + \beta : \alpha \in P_1, \beta \in P_2\}, \quad c \cdot P_1 := \{c\alpha : \alpha \in P_1\}.$$

We denote the Cartesian product of two subsets $P_1 \subset \mathbb{R}^{n_1}$ and $P_2 \subset \mathbb{R}^{n_2}$ by $P_1 \times P_2 := \{(\alpha, \beta) : \alpha \in P_1, \beta \in P_2\} \subset \mathbb{R}^{n_1} \times \mathbb{R}^{n_2} = \mathbb{R}^{n_1+n_2}$. Throughout, we use the notation $\Delta_n = \text{Conv}(\{0, e_1, \dots, e_n\}) \subset \mathbb{R}^n$ for the standard simplex in \mathbb{R}^n .

Example 9. Consider the sets of exponents $E_1 = \{(0,0), (1,0), (1,1), (2,0), (0,1)\}$ and $E_2 = \{(0,0), (1,0), (0,1)\}$. In Figure 1, the polytopes $P_1 := \text{Conv}(E_1)$, $P_2 := \text{Conv}(E_2)$, and $P_1 + P_2 \subset \mathbb{R}^2$ are displayed. Observe that P_2 is the two-dimensional standard simplex Δ_2 . \triangle




Figure 1: Polytopes from Example 9.

3.1 Explicit constructions

We present explicit constructions of admissible tuples for the following types of polynomial systems, listed in (more or less) increasing order of generality.

1. **Dense systems.** These are systems for which f_i may involve all monomials of degree at most d_i , where $(d_1, \dots, d_s) \in \mathbb{N}_{>0}^s$ is an s -tuple of positive natural numbers. For dense systems, we have $A_i = \{\alpha \in \mathbb{N}^n : \alpha_1 + \dots + \alpha_n \leq d_i\} = (d_i \cdot \Delta_n) \cap \mathbb{N}^n$.
2. **Unmixed systems.** We say that the polynomial system \mathcal{F} is *unmixed* if there is a full-dimensional lattice polytope P and integers d_1, \dots, d_s such that $d_i \cdot P = \text{Conv}(A_i)$. The *codegree* of P is the smallest $t \in \mathbb{N}_{>0}$ such that $t \cdot P$ contains a point with integer coordinates in its interior. Note that dense systems can be viewed as unmixed systems with $P = \Delta_n$.
3. **Multi-graded dense systems.** A different, natural generalization of the dense case allows different degrees for different subgroups of the variables x_1, \dots, x_n . Let $\{\mathcal{I}_1, \dots, \mathcal{I}_r\}$ be a partition of $\{1, \dots, n\}$, i.e. $\mathcal{I}_j \subset \{1, \dots, n\}$, $\mathcal{I}_j \cap \mathcal{I}_k = \emptyset$ and $\bigcup_{j=1}^r \mathcal{I}_j = \{1, \dots, n\}$. This way we obtain subsets $x_{\mathcal{I}_1}, \dots, x_{\mathcal{I}_r} \subset \{x_1, \dots, x_n\}$ of the variables, indexed by the \mathcal{I}_j . In a multi-graded dense system, f_i may contain all monomials of degree at most $d_{i,j}$ in the variables $x_{\mathcal{I}_j}$. If the variables are ordered such that the first n_1 variables are indexed by \mathcal{I}_1 , the next n_2 variables by \mathcal{I}_2 and so on, this means $A_i = ((d_{i,1} \cdot \Delta_{n_1}) \times \dots \times (d_{i,r} \cdot \Delta_{n_r})) \cap \mathbb{N}^n$. Necessarily we have $n_1 + \dots + n_r = n$. A dense system is a multi-graded dense system with $r = 1$.
4. **Multi-unmixed systems.** This is a generalization of the unmixed and the multi-graded dense case, where there are full-dimensional lattice polytopes $P_1 \subset \mathbb{R}^{n_1}, \dots, P_r \subset \mathbb{R}^{n_r}$ such that $0 \in P_i$ and $n = \sum_i n_i$ and for each $i \in \{1, \dots, s\}$, an r -tuple $(d_{i,1}, \dots, d_{i,r}) \in \mathbb{N}^r$ such that $\text{Conv}(A_i) = (d_{i,1} \cdot P_1) \times \dots \times (d_{i,r} \cdot P_r)$. That is, the convex hull of A_i is the product of dilations of the polytopes P_1, \dots, P_r . Note that a multi-graded dense system is a multi-unmixed system with $P_i = \Delta_{n_i}$, and an unmixed system is a multi-unmixed system with $r = 1$.
5. **Mixed systems.** This is the most general case, our only assumption on each A_i is that the lattice polytope $\sum_{i=1}^s \text{Conv}(A_i) \subset \mathbb{R}^n$ is full-dimensional.

If the full-dimensionality requirements in the previous list are not fulfilled, one can reformulate the system using fewer variables. For polynomial systems from these nested families, admissible tuples are presented in Table 1. In what follows, we discuss them in more detail.

The tuples presented in Table 1 are admissible under a *zero-dimensionality assumption* on the system \mathcal{F} . Unfortunately, it is not enough to require that $\mathcal{F}(x) = 0$ has finitely many solutions in \mathbb{C}^n or $(\mathbb{C} \setminus \{0\})^n$. Loosely speaking, we need that the lifting of \mathcal{F} to a certain larger solution space has finitely many solutions. This is best understood in the context of toric geometry. We refer the reader to [44, Sec. 3] or [11, Sec. 2] for a description of the zero-dimensionality assumption in this language. Here, we omit terminology from toric geometry and state the assumption in terms of *face systems*, following [12]. We will use the notation

$$\mathcal{F} = (f_1, \dots, f_s) \in R_{A_1} \times \dots \times R_{A_s}, \quad f_i = \sum_{\alpha \in A_i} c_{i,\alpha} x^\alpha. \quad (3.1)$$

For any vector $v \in \mathbb{R}^n$, we define

$$A_{i,v} := \{\alpha \in A_i : \langle v, \alpha \rangle = \min_{\beta \in A_i} \langle v, \beta \rangle\},$$

where $\langle v, \alpha \rangle = v_1 \alpha_1 + \dots + v_n \alpha_n \in \mathbb{R}$. For $i = 1, \dots, s$, fix any $\beta_{i,v} \in A_{i,v}$. This gives a new system

$$\mathcal{F}_v = (f_{1,v}, \dots, f_{s,v}) \in R_{A_{1,v}} \times \dots \times R_{A_{s,v}}, \quad f_{i,v} = \sum_{\alpha \in A_{i,v}} c_{i,\alpha} x^{\alpha - \beta_{i,v}},$$

called the *face system* associated to v . The exponents $\alpha - \beta_{i,v}$, $\alpha \in A_i$ occurring in the polynomials $f_{i,v}$ lie in a lattice of rank $< n$ when $v \neq 0$. We denote this lattice by

$$M_v = \mathbb{Z} \cdot \left\{ \bigcup_{i=1}^s (A_{i,v} - \beta_{i,v}) \right\}.$$

Let r_v be the rank of M_v . Applying a change of coordinates, \mathcal{F}_v is a system of Laurent polynomials in r_v variables on the torus $(\mathbb{C} \setminus \{0\})^{r_v}$. Its solutions are independent of the choice of $\beta_{i,v} \in A_{i,v}$.

Assumption 1 (Zero-dimensionality assumption). For every $v \in \mathbb{R}^n$, the *face system* $\mathcal{F}_v(x) = 0$ has finitely many (possibly zero) solutions in $(\mathbb{C} \setminus \{0\})^{r_v}$.

Setting $v = 0$, Assumption 1 implies that $\mathcal{F}(x) = 0$ has finitely many solutions in $(\mathbb{C} \setminus \{0\})^n$.

Remark 3.1. Assumption 1 holds for a generic element $\mathcal{F} \in R_{A_1} \times \cdots \times R_{A_s}$, in the sense of Section 2. In fact, for a generic system \mathcal{F} all face systems \mathcal{F}_v for $v \neq 0$ have no solutions in $(\mathbb{C} \setminus \{0\})^{r_v}$, and the condition for this to hold only depends on the coefficients associated to some vertices of the polytopes $\text{Conv}(A_i)$, see [17]. The fact that we can allow finitely many solutions for all face systems comes from the recent contributions [44, 11]. In practice, this means that our algorithm is robust in the presence of isolated solutions at or near infinity (where this is understood in the appropriate toric sense).

Theorem 3.1. Consider a polynomial system $\mathcal{F} = (f_1, \dots, f_s)$ with supports A_1, \dots, A_s satisfying Assumption 1. Consider $(A_0, (E_0, \dots, E_s), D)$ as defined in Table 1. Then, we have that $(\mathcal{F}, A_0, (E_0, \dots, E_s), D)$ is an admissible tuple.

Proof. We sketch the proof. We need to show that the three conditions in Definition 2.1 are satisfied. Observe that, by construction, the elements from the tuple satisfy the Compatibility condition and A_0 satisfies the Lattice condition. By Assumption 1, for generic $f_0 \in R_{A_0}$, the system (f_0, \dots, f_s) has no solutions on the toric variety associated to the lattice polytope $\text{Conv}(A_1) + \cdots + \text{Conv}(A_s)$ and we can adapt [11, Thm. 4.3] straightforwardly to the case of no solutions (the Koszul complex of sheaves in that proof is exact by [33, Ch. 2.B, Prop.1.4.a], see also [38, Thm. 3.C]). Hence, following the same procedure as in [11, Sec. 4], we can show that $\text{HF}((f_0, \mathcal{F}), (E_0, \mathbf{E}); D) = 0$. Therefore, by Lemma 2.1, the Rank condition holds. \square

Remark 3.2 (The number γ and the number of solutions). Consider \mathcal{F} satisfying Assumption 1. We fix an admissible tuple $(\mathcal{F}, A_0, (E_0, \dots, E_s), D)$ from Table 1. If $n = s$, the dimension $\gamma = \text{HF}(\mathcal{F}, \mathbf{E}; D)$ is the number of solutions defined by \mathcal{F} on the compact toric variety $X \supset (\mathbb{C} \setminus \{0\})^n$ from [11, Thm. 4.4], counted with multiplicities. For generic \mathcal{F} , all solutions have multiplicity 1 and lie in $(\mathbb{C} \setminus \{0\})^n$, which means that γ is the mixed volume of the polytopes $\text{Conv}(A_1), \dots, \text{Conv}(A_n)$ [12, Thm. A]. Additionally, in these cases, we have a complete characterization of the invariant subspaces of M_g as it represents a multiplication operator, see [11, Sec. 3.2]. It was pointed out to us by Laurent Busé that [18, Lem. 6.2] should imply that the same holds for $s > n$, see the proof of [15, Prop.3] for an example of how to prove such a result in the multihomogeneous case.

1. Dense case, $\text{Conv}(A_i) = d_i \cdot \Delta_n$		([11, Cor. 4.3])
A_0	$\Delta_n \cap \mathbb{N}^n$	($d_0 = 1$)
E_i	$((\sum_{j \neq i} d_j - n) \cdot \Delta_n) \cap \mathbb{N}^n$	
D	$((\sum_j d_j - n) \cdot \Delta_n) \cap \mathbb{N}^n$	
2. Unmixed case, $\text{Conv}(A_i) = d_i \cdot P$		([11, Thm. 4.5])
A_0	$P \cap \mathbb{N}^n$	($d_0 = 1$)
E_i	$((\sum_{j \neq i} d_j - \text{CODEGREE}(P) + 1) \cdot P) \cap \mathbb{N}^n$	
D	$((\sum_j d_j - \text{CODEGREE}(P) + 1) \cdot P) \cap \mathbb{N}^n$	
3. Multi-graded dense case, $\text{Conv}(A_i) = (d_{i,1} \cdot \Delta_{n_1} \times \cdots \times d_{i,r} \cdot \Delta_{n_r})$		([11, Ex. 10])
A_0	$(\Delta_{n_1} \times \cdots \times \Delta_{n_r}) \cap \mathbb{N}^n$	($d_{0,k} = 1$)
E_i	$\prod_k ((\sum_{j \neq i} d_{j,k} - n_k) \cdot \Delta_{n_k}) \cap \mathbb{N}^n$	
D	$\prod_k ((\sum_j d_{j,k} - n_k) \cdot \Delta_{n_k}) \cap \mathbb{N}^n$	
4. Multi-unmixed case, $\text{Conv}(A_i) = (d_{i,1} P_1 \times \cdots \times d_{i,r} P_r)$		([11, Ex. 10])
A_0	$(P_1 \times \cdots \times P_r) \cap \mathbb{N}^n$	($d_{0,k} = 1$)
E_i	$\prod_k ((\sum_{j \neq i} d_{j,k} - \text{CODEGREE}(P_k) + 1) \cdot P_k) \cap \mathbb{N}^n$	
D	$\prod_k ((\sum_j d_{j,k} - \text{CODEGREE}(P_k) + 1) \cdot P_k) \cap \mathbb{N}^n$	
5. Mixed case, $\text{Conv}(A_i) = P_i$		([11, Thm. 4.4])
A_0	$\Delta_n \cap \mathbb{N}^n$	($P_0 = \Delta_n$)
E_i	$(\sum_{j \neq i} P_j) \cap \mathbb{N}^n$	
D	$(\sum_j P_j) \cap \mathbb{N}^n$	

Table 1: Admissible tuples for five families of structured polynomial systems. In the table, we assume that all $d_i > 0$, $d_{i,j} \geq 0$ and $P \subset \mathbb{R}^n, P_i \subset \mathbb{R}^{n_i}$ are full dimensional lattice polytopes.

Macaulay matrices defined by the tuples from Table 1 have been used in different algorithms for solving sparse polynomial systems, e.g. sparse resultants [30], truncated normal forms [47], Gröbner bases [8, 9], and others [38]. When restricted to Macaulay matrices, these constructions are often near-optimal when $s = n$. However, there exist other kind of smaller matrices which can be also used to solve the system [7, 10]. When $s > n$, we can often work with much smaller Macaulay matrices. This is the topic of the next subsection.

3.2 Incremental constructions

Even though the tuples from Theorem 3.1 are admissible, they might lead to the construction of unnecessarily big matrices in Algorithm 2. To avoid this, we present an incremental approach which leads to the construction of potentially smaller matrices. For ease of exposition, we consider

only the unmixed case. The ideas can be extended to the other cases.

In what follows, we fix a polytope P such that $0 \in P$ and integers $d_0, \dots, d_s \in \mathbb{N}_{>0}$. We consider sets of exponents $A_0, A_1, \dots, A_s \subset \mathbb{N}^n$ such that, for each $i \in \{0, \dots, s\}$, we have $\text{Conv}(A_i) = d_i \cdot P$. For each $\lambda \in \mathbb{N}$, we define $\mathbf{E}^\lambda = (E_1^\lambda, \dots, E_s^\lambda)$ with

$$E_i^\lambda := \begin{cases} ((\lambda - d_i) \cdot P) \cap \mathbb{N}^n & \text{if } \lambda \geq d_i \\ \emptyset & \text{otherwise} \end{cases} \quad \text{for } i = 0, \dots, s, \quad \text{and} \quad D^\lambda := (\lambda \cdot P) \cap \mathbb{N}^n. \quad (3.2)$$

Theorem 3.2. *With the above notation, consider an unmixed polynomial system $\mathcal{F} \in R_{A_1} \times \dots \times R_{A_s}$, with $\text{Conv}(A_i) = d_i \cdot P$, satisfying Assumption 1. For any $\lambda \in \mathbb{N}$ such that there is $f_0 \in R_{A_0}$ satisfying $\text{rank}(N_{f_0}) = \text{HF}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda)$, we have that the tuple $(\mathcal{F}, A_0, (E_0^\lambda, \mathbf{E}^\lambda), D^\lambda)$ is admissible. Moreover, for any $\lambda \geq \sum_i d_i - \text{CODEGREE}(P) + 1$ and for generic $f_0 \in R_{A_0}$, we have that $\text{rank}(N_{f_0}) = \text{HF}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda)$.*

Proof. By construction, the tuple satisfies the Compatibility and Lattice conditions. By assumption, it satisfies the Rank condition, so it is admissible. The proof follows as in Theorem 3.1. \square

The bound upper bound on λ obtained in Theorem 3.2 is not tight for overdetermined systems. Below, we will present a broad class of overdetermined unmixed systems, namely semi-regular* sequences, for which we can improve it.

Remark 3.3 (The number γ and the number of solutions). *In contrast to Remark 3.2, the condition $\text{rank}(N_{f_0}) = \text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda)$ does not imply that $\gamma = \text{HF}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda)$ agrees with the number of solutions of \mathcal{F} on some toric compactification. In fact, in Section 4, we will present examples of semi-regular* sequences (Definition 3.1) where γ is strictly larger than the number of solutions. In these cases, the matrices M_g from Equation (2.2) are not multiplication operators. For readers familiar with the concept of Castelnuovo-Mumford regularity, we note that this happens because the degree D^λ belongs to the regularity of $\{f_0, \mathcal{F}\}$, but not necessarily to that of \mathcal{F} .*

Theorem 3.2 suggests an algorithm for finding an admissible tuple for an unmixed system \mathcal{F} : we simply check, for a random element $f_0 \in R_{A_0}$ and increasing values of λ , whether $\text{rank}(N_{f_0}) = \text{HF}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda)$ with $N_{f_0} = \text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda) \cdot \mathbf{M}(\mathcal{F}, E_0^\lambda; D^\lambda)$. In order to do this efficiently, instead of computing $\text{Coker}(\mathcal{F}, \mathbf{E}^{\lambda+1}; D^{\lambda+1})$ directly as the left nullspace of the large matrix $\mathbf{M}(\mathcal{F}, \mathbf{E}^{\lambda+1}; D^{\lambda+1})$, we will obtain it from the previously computed $\text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda)$ and a smaller Macaulay matrix. This technique was applied in the dense setting ($P = \Delta_n$) in [4, 40], where it is also called ‘degree-by-degree’ approach. See also [41] for a recent complexity analysis.

Note that, by construction, $E_i^\lambda \subset E_i^{\lambda+1}$ and $D^\lambda \subset D^{\lambda+1}$. The first step is to construct the following 2×2 block matrix

$$(\text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda) \times \text{id}) := \begin{bmatrix} D^\lambda & D^{\lambda+1} \setminus D^\lambda \\ \text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda) & 0 \\ 0 & \text{id} \end{bmatrix}. \quad (3.3)$$

Here id denotes the identity matrix of size $\#(D^{\lambda+1} \setminus D^\lambda)$. Note that the columns of the matrix $(\text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda) \times \text{id})$ are indexed by $D^{\lambda+1}$, where the first block column is indexed by $D^\lambda \subset D^{\lambda+1}$. Next, we set $\mathbf{E}^{\lambda+1} \setminus \mathbf{E}^\lambda := (E_1^{\lambda+1} \setminus E_1^\lambda, \dots, E_s^{\lambda+1} \setminus E_s^\lambda)$ and construct the Macaulay matrix

$M(\mathcal{F}, \mathbf{E}^{\lambda+1} \setminus \mathbf{E}^\lambda; D^{\lambda+1})$. Here we require that the ordering of the rows is compatible with the ordering of the columns in (3.3). Let $L^{\lambda+1}$ be a left nullspace matrix of the matrix product

$$(\text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda) \times \text{id}) \cdot M(\mathcal{F}, \mathbf{E}^{\lambda+1} \setminus \mathbf{E}^\lambda; D^{\lambda+1}). \quad (3.4)$$

Then $\text{Coker}(\mathcal{F}, \mathbf{E}^{\lambda+1}; D^{\lambda+1}) = L^{\lambda+1} \cdot (\text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda) \times \text{id})$ is a left nullspace matrix for the Macaulay matrix $M(\mathcal{F}, \mathbf{E}^{\lambda+1}; D^{\lambda+1})$. The power of this approach lies in the fact that (3.4) is much smaller than $M(\mathcal{F}, \mathbf{E}^{\lambda+1}; D^{\lambda+1})$, which leads to a much cheaper left nullspace computation.

This gives an iterative algorithm for *updating* the left nullspace matrix $\text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda)$. We start our iteration by considering $\lambda = \max_i d_i$, as we want to take into account all of the equations. This discussion is summarized in Algorithm 3. Note that the algorithm computes the cokernel $\text{Coker}(\mathcal{F}, \mathbf{E}; D)$ for the admissible tuple as a by-product, as well as the matrix N_{f_0} . This allows us to skip the steps before line 6 in Algorithm 2.

Algorithm 3 GETADMISSIBLETUPLEUNMIXED

Input: An unmixed system \mathcal{F} satisfying Assumption 1, the polytope $P \ni 0$, the degrees (d_1, \dots, d_s) .

Output: An admissible tuple $(\mathcal{F}, A_0, (E_0, \mathbf{E}), D)$, a left nullspace matrix $\text{Coker}(\mathcal{F}, (E_0, \mathbf{E}); D)$ and a corresponding matrix N_{f_0}

- 1: $d_0 \leftarrow 1$
 - 2: $A_0 \leftarrow P \cap \mathbb{N}^n$
 - 3: $f_0 \leftarrow$ a random element of R_{A_0}
 - 4: $\lambda \leftarrow \max_i d_i$
 - 5: $\text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda) \leftarrow$ left nullspace of $M(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda)$, for the sets of exponents in (3.2)
 - 6: $r \leftarrow$ number of rows of $\text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda)$
 - 7: $N_{f_0} = \text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda) \cdot M(f_0, E_0^\lambda; D^\lambda)$
 - 8: **while** $\text{rank}(N_{f_0}) \neq r$ **do**
 - 9: $(\text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda) \times \text{id}) \leftarrow$ the matrix from (3.3)
 - 10: $L^{\lambda+1} \leftarrow$ a left nullspace matrix for the matrix in (3.4)
 - 11: $\text{Coker}(\mathcal{F}, \mathbf{E}^{\lambda+1}; D^{\lambda+1}) \leftarrow L^{\lambda+1} \cdot (\text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda) \times \text{id})$
 - 12: $\lambda \leftarrow \lambda + 1$
 - 13: $r \leftarrow$ number of rows of $\text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda)$
 - 14: $N_{f_0} = \text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda) \cdot M(f_0, E_0^\lambda; D^\lambda)$
 - 15: **return** $(\mathcal{F}, A_0, (E_0^\lambda, \mathbf{E}^\lambda), D^\lambda), \text{Coker}(\mathcal{F}, \mathbf{E}^\lambda; D^\lambda), N_{f_0}$
-

Remark 3.4 (Other incremental constructions). *There are alternative incremental constructions for the matrices N_{f_0} which also reuse information from previous steps to speed up the computations. An example is the F5 criterion in the context of Gröbner bases [31]. These ideas extend naturally to the mixed setting, see [9]. However, these approaches based on monomial orderings lead to bad numerical behaviour. In the context of sparse resultants for mixed systems, Canny and Emiris [28] proposed an alternative incremental algorithm to construct admissible tuples leading to smaller Macaulay matrices. Their procedure can be enhanced with the approach followed in this section.*

In the rest of this subsection, we identify a broad class of overdetermined unmixed systems for which we can obtain smaller admissible tuples than the ones in Theorem 3.1. We will need some

more notation. The *Ehrhart series* of a polytope P is the series

$$\text{ES}_P(t) = \sum_{\lambda \geq 0} \#((\lambda \cdot P) \cap \mathbb{Z}^n) t^\lambda.$$

The *Hilbert series* of a polynomial system $\mathcal{F}_0 := (f_0, \dots, f_s) \in R_{A_0} \times \dots \times R_{A_s}$, is

$$\text{HS}_{\mathcal{F}_0}(t) := \sum_{\lambda \geq 0} \text{HF}(\mathcal{F}_0, \mathbf{E}^\lambda; D^\lambda) t^\lambda.$$

Definition 3.1 (Semi-regularity*). We say that \mathcal{F}_0 is a semi-regular* sequence if

$$\text{HS}_{\mathcal{F}_0}(t) = \left[\text{ES}_P(t) \prod_{i=0}^s (1 - t^{d_i}) \right]_+,$$

where $[\cdot]_+$ means that we truncate the series in its first negative coefficient.

Observe that we write semi-regular* sequence with an asterisk as the usual definition of *semi-regular sequence* asks for this condition on the Hilbert series to hold for every subsystem (f_0, \dots, f_i) , $i \leq s$. However, semi-regular sequences are too restrictive for our purposes.

Even in the case where P is a standard simplex, semi-regular* sequences are not understood as well as regular sequences. For example, Fröberg's conjecture states that being a semi-regular* sequence is a generic condition [32]. This conjecture, supported by a lot of empirical evidence, was extended to the unmixed case [31].

Theorem 3.3. Consider an unmixed polynomial system $\mathcal{F} \in R_{A_1} \times \dots \times R_{A_s}$ and a polynomial $f_0 \in R_{A_0}$, with $\text{Conv}(A_i) = d_i \cdot P$. Let λ_{\min} be the smallest integer among the degrees of the monomials in $\text{ES}_P(t) \prod_{i=0}^s (1 - t^{d_i})$ standing with a non-positive coefficient. We have that, if (f_0, \mathcal{F}) is a semi-regular* sequence, then the tuple $((f_0, \mathcal{F}), A_0, (E_0^{\lambda_{\min}}, \mathbf{E}^{\lambda_{\min}}); D^{\lambda_{\min}})$ is admissible.

Proof. The proof follows from the fact that $\text{HF}((f_0, \mathcal{F}), E_0^{\lambda_{\min}}, \mathbf{E}^{\lambda_{\min}}; D^{\lambda_{\min}}) = 0$ as the sequence is semi-regular*. \square

It follows directly from Theorem 3.2 that, whenever (f_0, \mathcal{F}) is semi-regular*, $\lambda_{\min} \leq \sum_i d_i - \text{CODEGREE}(P) + 1$. In Section 4.2, we present generic families of zero-dimensional overdetermined systems \mathcal{F} such that (f_0, \mathcal{F}) is semi-regular*. For these systems, we show that the previous inequality can be strict.

Semi-regular* sequences give us an inexpensive heuristic to discover values for λ for which we can obtain admissible tuples. It was observed in practice [2, 31] that for many systems \mathcal{F} not having much solutions outside the torus (see Remark 3.1), they can be extended to semi-regular* sequences. Moreover, there are asymptotic estimates for the expected value of λ [2].

4 Experiments

In this section we illustrate several aspects of the methods presented in this paper via numerical experiments. We implemented these algorithms in the new Julia package `EigenvalueSolver.jl`,

which is freely available at <https://github.com/simontelen/JuliaEigenvalueSolver>. For all computations involving polytopes, we use `Polymake.jl` (version 0.5.3), which is a Julia interface to Polymake [34]. We compare our results with the package `HomotopyContinuation.jl` (version 2.3.1), which is state-of-the-art software for solving systems of polynomial equations using homotopy continuation [14]. All computations were run on a 16 GB MacBook Pro with an Intel Core i7 processor working at 2.6 GHz.

To evaluate the quality of a numerical approximation $\zeta \in \mathbb{C}^n$ of a solution for a polynomial system \mathcal{F} given by (3.1). We define the *backward error* $\text{BWE}(\zeta)$ of ζ as

$$\text{BWE}(\zeta) = \frac{1}{s} \sum_{i=1}^s \frac{|f_i(\zeta)|}{\sum_{\alpha \in A_i} |c_{i,\alpha} \zeta^\alpha| + 1}. \quad (4.1)$$

This error can be interpreted as a measure for the relative distance of \mathcal{F} to a system \mathcal{F}' for which $\mathcal{F}'(\zeta) = 0$, see [45, App. C].

Additionally, we validate our computed solutions via *certification*. For that, we use the certification procedure implemented in the function `certify` of `HomotopyContinuation.jl`, which is based on interval arithmetic, as described in [13]. This function takes as an input a list of approximate solutions to \mathcal{F} and tries to compute a list of small boxes in \mathbb{C}^n , each of them containing an approximate input solution and exactly one *actual* solution to \mathcal{F} . The total number of connected components in the union of these boxes is denoted by `crt` in what follows. Each of these connected components contains exactly one solution of \mathcal{F} , and one or more approximate input solutions. This means that `crt` is a lower bound on the number of solutions to \mathcal{F} . If `crt` equals the number of solutions, the solutions of \mathcal{F} are in one-to-one correspondence with the approximate input solutions. In this case, we say that all solutions are *certified*. The function `certify` assumes that \mathcal{F} is *square*, i.e. \mathcal{F} should have as many equations as variables ($s = n$). If this is not the case ($s > n$), we use `certify` on a system obtained by taking n random \mathbb{C} -linear combinations of f_1, \dots, f_s .

The main function of our package `EigenvalueSolver.jl` is `solve_EV`, which implements Algorithm 2. It takes as an input an admissible tuple (see Definition 2.1). This tuple can be computed using the auxiliary functions provided in our implementation, which are tailored to take into account the specific structure of the systems. These functions use the explicit and incremental constructions from Section 3.

It is common in applications that we have to solve many different generic systems \mathcal{F} with the same supports A_1, \dots, A_s . In this case, the computation of the admissible tuple can be seen as an *offline* computation that needs to happen only once. We will therefore report both the *offline* and the *online* computation time. The offline computation time is the time needed for computing an admissible tuple *and* executing `solve_EV`. The *online* computation re-uses a previously computed admissible tuple to execute `solve_EV`.

Table 2 summarizes the notation that we use to describe our experiments.

The section is organized as follows. In Section 4.1, we consider square systems ($s = n$) and show how to use `EigenvalueSolver.jl` to solve them. In Section 4.2, we solve overdetermined systems ($s > n$) using our incremental algorithm. We perform several experiments summarized in Table 4 and Table 5. In Section 4.3, we consider systems for which one solutions drifts off to ‘infinity’. In Section 4.4, we compare our algorithm with homotopy continuation methods.

n	number of variables
δ	number of solutions
crt	number of connected components computed by certify
t_{on}	online computation time in seconds
t_{off}	offline computation time in seconds
$\overline{\text{BWE}}$	maximum backward error of all computed approximate solutions
$\overline{\text{BWE}}$	geometric mean of the backward errors of all computed solutions
γ	the number of rows of $\text{Coker}(\mathcal{F}, \mathbf{E}; D)$
$\#D$	cardinality of D , i.e. the number of columns of $\text{Coker}(\mathcal{F}, \mathbf{E}; D)$

Table 2: Notation in the experiments in Section 4.

4.1 Square systems

In this subsection, we demonstrate some of the functionalities of `EigenvalueSolver.jl` by solving square systems, that is $s = n$, for each of the families in Table 1. The code used for the examples can be found at <https://github.com/simontelen/JuliaEigenvalueSolver> in the Jupyter notebook `/example/demo_EigenvalueSolver.ipynb`. We fix the parameters of Table 1 and consider specific supports A_i as described below. We construct random polynomial systems by assigning random real coefficients to each of the monomials, which we draw from a standard normal distribution. By Remark 3.2, the number γ equals the number of solutions δ for all examples in this subsection.

For our first example, we intersect two degree 20 curves in the plane. That is, we consider a square, dense system \mathcal{F}_1 with $n = 2$ and $d_1 = d_2 = 20$. The equations are generated by the following simple commands:

```
@polyvar x[1:2]; ds = [20;20];
f = EigenvalueSolver.getRandomSystem_dense(x, ds)
```

By Bézout’s theorem, this system has $\delta = 400$ different solutions, which we can compute via

```
sol = EigenvalueSolver.solve_CI_dense(f, x; DBD = false)
```

In the previous line, the option `DBD = false` indicates that we do not want to use the ‘degree-by-degree’ approach for solving this system, that is, the incremental approach described in Section 3.2. Experiments show that this strategy is only beneficial for square systems with $n \geq 3$. The letters CI in the name of the function stand for *complete intersection*, which indicates that a zero-dimensional square system is expected as its input. In this example, we have $\#D = 820$ and the computation took $t_{\text{off}} = 0.83$ seconds. To validate the solutions, we compute their backward errors.

```
BWEs = EigenvalueSolver.get_residual(f, sol, x)
```

The maximal value, computed using the command `maximum(BWEs)`, is $\text{BWE} \approx 10^{-12}$. The function `certify` from `HomotopyContinuation.jl` certifies `crt = 400` distinct solutions. If we perform the same computation with parameters $n = 3$, $(d_1, d_2, d_3) = (4, 8, 12)$, we obtain $\delta = \gamma = \text{crt} = 384$, $\#D = 2300$, $t_{\text{off}} = 3.10$, $\text{BWE} \approx 10^{-11}$.

```
@polyvar x[1:3]; ds = [4;8;12];
f = EigenvalueSolver.getRandomSystem_dense(x, ds)
sol = EigenvalueSolver.solve_CI_dense(f, x)
```

For our next example, we consider an unmixed system \mathcal{F}_2 with parameters

$$n = 2, \quad P = \text{Conv}(A), \quad A = \{(0,0), (1,0), (1,1), (0,1), (2,2)\}, \quad (d_1, d_2) = (5, 12). \quad (4.2)$$

The following code executes this example,

```
@polyvar x[1:2]; A = [0 0; 1 0; 1 1; 0 1; 2 2]; d = [5;12];
f = EigenvalueSolver.getRandomSystem_unmixed(x, A, d)
sol, A0, E, D = EigenvalueSolver.solve_CI_unmixed(f, x, A, d)
```

In this case, we obtain $\delta = \gamma = \text{crt} = 240$, $\#D = 685$, $t_{\text{off}} = 0.94$, $\text{BWE} \approx 10^{-11}$. We remark that the function `solve_CI_unmixed` also returns the admissible tuple (A_0, \mathbf{E}, D) , so that it can be used to solve another generic unmixed system with the same parameters, without redoing the polyhedral computations to generate this tuple. This can be done in the following way,

```
sol = EigenvalueSolver.solve_EV(f, x, A0, E, D; check_criterion = false)
```

The option `check_criterion = false` in the previous line indicates that the input tuple is admissible, so we do not need to spend time on checking whether the criterion in Lemma 2.1 is satisfied. Using this option, the *online* computation is faster and takes $t_{\text{on}} = 0.41$ seconds, yet the parameters $\delta, \gamma, \text{crt}, \text{BWE}$ are comparable to the offline case. To illustrate how the unmixed function exploits the structure of the equations, in Figure 2, we plot the exponents in D for this example, together with the exponents in D for our dense example \mathcal{F}_1 . In both plots, we have highlighted the exponents in the set B that were selected using QR factorization with optimal column pivoting. These monomial bases clearly do not correspond to any standard (Gröbner) or border basis. Figure 2 should be compared to, for instance, Figure 2 in [47].




Figure 2: Exponents in D , constructed as in Table 1, for the dense systems \mathcal{F}_1 (left) and the unmixed system \mathcal{F}_2 (right). The exponents are shown as lattice points. Dark coloured dots correspond to the set $B \subset D$ chosen by the QR factorization in line 6 in Algorithm 2.

family	parameters	results
multi-graded, dense	$n_1 = n_2 = 2, (d_{j,k}) = \begin{pmatrix} 1 & 6 \\ 2 & 1 \\ 3 & 2 \\ 4 & 1 \end{pmatrix},$	$\delta = \gamma = \text{crt} = 219, \#D = 3025,$ $t_{\text{off}} = 15.46, \text{BWE} \approx 10^{-11}$
multi-unmixed	$n_1 = n_2 = 2, (d_{j,k}) = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}, P_1 = \text{Conv}(A)$ $P_2 = 2 \cdot \Delta_2$	$\delta = \gamma = \text{crt} = 96, \#D = 2745,$ $t_{\text{off}} = 12.84, t_{\text{on}} = 11.92, \text{BWE} \approx 10^{-9}$

Table 3: Computational data and results for multi-graded and multi-unmixed examples. Here A is as in (4.2).

We can solve multi-graded dense and multi-unmixed systems using the implemented functions `solve_CI_multi_dense` and `solve_CI_multi_unmixed`, respectively. Table 3 summarizes our choice of parameters and the results of our experiments for these systems.

To conclude this subsection, we present a classical example of a square mixed system in $n = 3$ variables coming from molecular biology [29, Sec. 3.3]. The following code generates and solves these equations:

```
@polyvar t[1:3]
b = [-13 -1 -1 24 -1; -13 -1 -1 24 -1; -13 -1 -1 24 -1]
mons1 = [1 t[2]^2 t[3]^2 t[2]*t[3] t[2]^2*t[3]^2]
mons2 = [1 t[3]^2 t[1]^2 t[3]*t[1] t[3]^2*t[1]^2]
mons3 = [1 t[1]^2 t[2]^2 t[1]*t[2] t[1]^2*t[2]^2]
f = [b[1,:] '*mons1'; b[2,:] '*mons2'; b[3,:] '*mons3'] [:]
sol, A0, E, D = EigenvalueSolver.solve_CI_mixed(f,t)
```

In this case, we obtain $\delta = \gamma = \text{crt} = 16, \#D = 200, t_{\text{off}} = 0.53, t_{\text{on}} = 0.02, \text{BWE} \approx 10^{-13}$. The function `certify` tells us that all 16 solutions are real, confirming the observation made in [29].

4.2 Overdetermined systems

We now consider examples of *overdetermined systems*, by which we mean cases where $s > n$. We will limit ourselves to unmixed systems and use Algorithm 3 to find admissible tuples leading to small Macaulay matrices. These systems arise, for instance, in tensor decomposition problems [48]. We present examples where γ is significantly larger than δ and show that, nevertheless, our algorithms successfully extract $\delta < \gamma$ relevant eigenvalues and consistently return all solutions of the input systems.

We observe that the Macaulay matrices constructed in this section are smaller than the ones obtained using other symbolic-numerical techniques as (sparse) resultants [27] or its generalization [38]. The admissible tuples used in those symbolic-numerical algorithms lead to multiplication operators, for which $\gamma = \delta$. As observed in Remark 3.3, our matrices M_g are too large to be multiplication operators. The extra time needed for computing the eigenvalues of these larger matrices is negligible compared to the time won by computing M_g from a smaller Macaulay matrix.

The overdetermined systems considered in this section are constructed as follows. For a fixed number of variables n , number of solutions δ and set of exponents A , we generate δ random points $\zeta_1, \dots, \zeta_\delta$ in \mathbb{C}^n by drawing their coordinates from a complex standard normal distribution.

We construct a Vandermonde type matrix V_{dm} whose rows consist of the vectors $\zeta_i^A / \|\zeta_i^A\|_2, i = 1, \dots, \delta$. The nullspace of V_{dm} is computed using SVD and its columns represent $s = \#A - \delta$ polynomials f_1, \dots, f_s with support A . If we do not pick too many points, we have that $s > n$ and the solutions of $\mathcal{F} = (f_1, \dots, f_s)$ are exactly the points $\zeta_1, \dots, \zeta_\delta$.

4.2.1 Dense, overdetermined systems

In this subsection, we consider dense overdetermined systems, i.e. $A_0 = \Delta_n \cap \mathbb{N}^n$ and $A = (d \cdot \Delta_n) \cap \mathbb{N}^n$ for some degree $d \in \mathbb{N}_{>0}$. The *offline* computation uses Algorithm 3 to find an admissible tuple, as well as a left nullspace, and then execute Algorithm 2 from line 6 on. The *online* computation uses this admissible tuple to execute Algorithm 2 directly. This means that the offline version uses an incremental strategy for computing the left nullspace, while the online version works directly with the large Macaulay matrix. The online version can be adapted to work incrementally as well. We have chosen not to do this in order to illustrate that, depending on n, s , the incremental approach may be less or more efficient than the direct approach. In cases where the incremental approach is more efficient, this may cause $t_{off} < t_{on}$. In the square case ($s = n$), this happens for $n \geq 3$ [40, 41], but our results show that in the overdetermined case this might not happen. Further research is necessary to make an automated choice. Table 4 gives an overview of the computational results. The column indexed by $\widetilde{\#D}$ represents the size of the matrix that would be used in classical approaches. This is discussed in the final paragraph of this subsection.

The first 10 rows in Table 4 correspond to systems of 6 equations in 3 variables of increasing degree $d = 2, 4, \dots, 20$. Note that $\gamma > \delta$ for $d > 4$. In all cases, δ distinct solutions were computed using our algorithms and $crt = \delta$. This means that exactly δ out of γ eigenvalues were selected and correctly processed to compute solution coordinates. The maximum backward error grows faster with the degree of the equations than for square systems [47]. This can be remedied by using larger admissible tuples to bring γ closer to δ , at the cost of computing cokernels of larger matrices. However, our experiment shows that we can find certified approximations for all 1765 intersection points of 6 threefolds of degree 20 within less than 10 minutes. All of these are within two Newton refinement steps from having a backward error of machine precision.

The next 5 rows of Table 4 contain results for 18 dense equations in 6 variables of increasing degree $d = 2, 3, \dots, 6$. Note that $t_{on} > t_{off}$ for $d > 2$. This is due to the incremental approach for the offline phase, as mentioned above.

In the following 7 rows of Table 4, we illustrate the effect of increasing the number of variables when we fix the degree $d = 3$. We work with overdetermined systems for which $s = 2n$. Although the complexity of eigenvalue methods usually scales badly with the number of variables, these results show that when the system is ‘sufficiently overdetermined’, our algorithms can find feasible admissible tuples to solve cubic equations in 8 variables in no more than 20 seconds.

Finally, the last rows of Table 4 correspond to systems of cubic equations in 15 variables with an increasing number $\delta = 200, 300, \dots, 600$ of solutions. Note that the computation time *decreases* with the number of solutions. The reason is that for all these values of δ , we can work with the same support D for the Macaulay matrix. This means that the matrix has the same number of rows for each system. The number of columns, however, depends on the number of equations, which increases with decreasing δ by construction. For $\delta = 700$, we need a larger set of exponents D , causing memory issues.

All systems (f_0, \mathcal{F}) appearing in Table 4 are semi-regular*. By Theorem 3.3, the minimal value of λ such that $((f_0, \mathcal{F}), A_0, (E_0^\lambda, E^\lambda); D^\lambda)$ is an admissible tuple is the degree λ_{min} of the

n	s	d	δ	crt	γ	$\#D$	$\widetilde{\#D}$	BWE	$\overline{\text{BWE}}$	t_{off}	t_{on}
3	6	2	4	4	4	10	10	5.75e-16	3.20e-16	1.25e-03	1.34e-03
3	6	4	29	29	29	84	84	1.70e-14	2.54e-15	9.41e-03	6.33e-03
3	6	6	78	78	100	220	286	7.07e-12	2.23e-14	7.00e-02	5.23e-02
3	6	8	159	159	224	560	816	1.21e-12	4.67e-14	4.47e-01	2.90e-01
3	6	10	280	280	465	969	1540	6.32e-10	6.63e-13	1.99e+00	1.32e+00
3	6	12	449	449	820	1540	2600	5.09e-09	7.90e-12	8.76e+00	6.04e+00
3	6	14	674	674	1280	2600	4495	1.51e-08	7.78e-12	3.88e+01	2.21e+01
3	6	16	963	963	1938	3654	6545	3.57e-07	3.98e-11	1.26e+02	7.36e+01
3	6	18	1324	1324	2776	4960	9139	1.83e-06	5.77e-10	3.54e+02	2.08e+02
3	6	20	1765	1765	3780	7140	12341	1.11e-05	9.96e-10	9.85e+02	5.38e+02
6	18	2	10	10	10	84	84	1.53e-14	2.96e-15	1.45e-02	8.82e-03
6	18	3	66	66	66	462	462	4.51e-14	5.59e-15	1.69e-01	1.74e-01
6	18	4	192	192	204	1716	3003	2.95e-12	6.36e-14	2.11e+00	3.79e+00
6	18	5	444	444	1225	5005	8008	7.52e-12	1.76e-13	5.18e+01	7.86e+01
6	18	6	906	906	4060	12376	27132	5.28e-10	2.37e-12	1.01e+03	1.33e+03
2	4	3	6	6	6	10	10	3.02e-15	1.12e-15	1.15e-03	1.23e-03
3	6	3	14	14	14	35	35	5.95e-15	1.49e-15	2.92e-03	2.35e-03
4	8	3	27	27	27	126	126	3.85e-14	2.27e-15	1.56e-02	1.27e-02
5	10	3	46	46	46	252	252	6.59e-14	8.89e-15	4.19e-02	2.04e-01
6	12	3	72	72	126	462	924	3.70e-12	1.46e-13	1.61e-01	1.51e-01
7	14	3	106	106	127	1716	3432	6.20e-12	3.96e-14	2.29e+00	4.26e+00
8	16	3	149	149	483	3003	6435	8.31e-12	1.05e-13	1.16e+01	1.92e+01
15	616	3	200	200	200	3876	3876	1.45e-13	1.04e-14	9.78e+01	5.80e+01
15	516	3	300	300	300	3876	3876	3.66e-13	9.37e-15	8.25e+01	5.64e+01
15	416	3	400	400	400	3876	3876	5.46e-13	1.44e-14	8.50e+01	5.42e+01
15	316	3	500	500	500	3876	3876	4.25e-13	1.26e-14	6.38e+01	5.81e+01
15	216	3	600	600	600	3876	3876	4.86e-13	1.41e-14	4.91e+01	4.65e+01

Table 4: Computational results for overdetermined, dense systems. See Table 2 for the notation.

lowest-degree monomial with a non-positive coefficient in

$$\text{ES}_{\Delta_n}(t) \prod_{i=0}^s (1 - t^{d_i}) = \frac{(1-t)(1-t^d)^s}{(1-t^n)}.$$

To illustrate the gain of using such a minimal λ_{\min} , we included the number $\widetilde{\#D}$ which corresponds to the number of monomials in the D^λ for the smallest λ which gives $\gamma = \delta$. That is, the smallest λ for which the matrices M_g in our algorithm represent multiplication matrices. For $n = 3, s = d = 6$, λ_{\min} is 9, and the admissible tuple has $\#D = \#(9\Delta_2 \cap \mathbb{Z}^2) = 220$ lattice points. Multiplication matrices are obtained from $\#D = \#(10\Delta_2 \cap \mathbb{Z}^2) = 286$. To see the benefit of our incremental construction over the bounds from Table 1, note that case 1 gives $\#D = \#(34\Delta_2 \cap \mathbb{Z}^2) = 7770$, and the Minkowski sum of the Newton polytopes (Table 1, case 5) gives $\#D = \#(37\Delta_2 \cap \mathbb{Z}^2) = 9880$.

4.2.2 Unmixed, overdetermined systems

We now use our algorithms to solve overdetermined unmixed systems. The results are summarized in Table 5. First, we set $n = 3$ and choose δ such that $s = 6$. We define A_0 as the columns of

$$\begin{pmatrix} 2 & 2 & 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & 1 & 0 & 2 & 1 & 2 & 0 \end{pmatrix}.$$

The support A is obtained as $A = (d \cdot \text{Conv}(A_0)) \cap \mathbb{N}^3$ for increasing values of d . The conclusions are similar to those for the $n = 3$ experiments in the previous subsection. Note that $d = 8$ is the only reported case for which one solution could not be certified.

Next, we set $n = 15$, $\delta = 100$ and we define $A_0 = \{0, e_1, e_2, \dots, e_{13}, e_{13} + e_{14}, e_{14} + e_{15}\}$, where e_i is the i -th standard basis vector of \mathbb{Z}^{15} . We set $A = (2 \cdot \text{Conv}(A_0)) \cap \mathbb{N}^{15}$. There are 136 exponents in A , of degree at most 4.

n	s	d	δ	crt	γ	# D	BWE	$\overline{\text{BWE}}$	t_{off}	t_{on}
3	6	1	3	3	3	33	8.91e-16	5.36e-16	1.25e+00	7.59e-01
3	6	2	27	27	27	165	2.71e-13	1.99e-14	1.96e+00	2.30e-02
3	6	3	76	76	93	291	3.94e-12	8.52e-14	2.07e+00	9.36e-02
3	6	4	159	159	216	708	8.53e-11	6.62e-13	3.27e+00	5.06e-01
3	6	5	285	285	415	1405	1.99e-08	6.42e-12	6.25e+00	2.78e+00
3	6	6	463	463	891	1881	2.00e-06	6.15e-11	1.56e+01	1.06e+01
3	6	7	702	702	1387	3133	4.56e-05	7.06e-10	5.66e+01	4.51e+01
3	6	8	1011	1010	2031	4845	9.29e-05	3.78e-10	1.86e+02	1.61e+02
15	36	2	100	100	100	3876	2.88e-13	7.42e-15	8.07e+01	4.59e+01

Table 5: Computational results for overdetermined, unmixed systems. See Table 2 for the notation.

Remark 4.1 (Noisy coefficients). *As an important direction for future research, we note that our eigenvalue algorithms can be used to compute ‘solutions’ to overdetermined systems with noisy coefficients. For instance, the noise level needs to be taken into account when setting the relative tolerance for computing the left nullspace in line 3 of Algorithm 2. This is expected to work especially well for strongly overdetermined problems with only a few solutions.*

4.3 Solutions at infinity

An important feature of our algorithms is that they can deal with systems having isolated solutions *at or near infinity*. To illustrate this, we work with the same set-up as in Section 4.2.1 with parameters $n = 7$, $d = 3$ and $s = 14$, implying $\delta = 106$. We generate 106 random complex points $\zeta_1, \dots, \zeta_{106}$ as before, and then multiply the coordinates of ζ_{106} by a factor 10^e for increasing values of e . That is, we let one of 106 solutions drift off to infinity. Figure 3 shows the maximal 2-norm of the computed solutions as well as the maximal backward error BWE for $e = 0, \dots, 14$. The results clearly show that the accuracy is not affected by the ‘outlier’ solution. As e grows larger, the solution ζ_{106} corresponds to an isolated solution of the face system \mathcal{F}_v with $v = (1, 1, 1, 1, 1, 1)$, see Remark 3.1. For all considered values of e , our algorithm computed $\text{crt} = \delta = 106$ distinct certified approximate solutions.




Figure 3: Max. backward error BWE (—●—) and norm of the largest solution (—●—) for the experiments in Sec. 4.3.

4.4 Comparison with homotopy continuation methods

Homotopy continuation algorithms form another important class of numerical methods for solving polynomial systems [42]. These methods transform a *start system* with known solutions continuously into the *target system*, which is the system we want to solve, and track the solutions along the way. This process can usually only be set up for square systems, i.e. $s = n$. In these cases, especially when $n = s$ is large (≥ 4), homotopy continuation methods often outperform eigenvalue methods. When the system \mathcal{F} is overdetermined ($s > n$), homotopy methods solve a square system $\mathcal{F}_{\text{square}}$ obtained by taking n random \mathbb{C} -linear combinations of the s input polynomials. The set of solutions of \mathcal{F} is contained in the set of solutions of $\mathcal{F}_{\text{square}}$, so that the solutions of \mathcal{F} can be extracted by an additional ‘filtering’ step. Often $\mathcal{F}_{\text{square}}$ has *many more* solutions than \mathcal{F} , so that many of the tracked paths do not end at a solution of \mathcal{F} . Below, we use the notation δ_{square} for the number of solutions of $\mathcal{F}_{\text{square}}$.

Several implementations of homotopy methods exist, including Bertini [3] and PHCpack [50]. Here, we choose to compare our computational results with the relatively recent Julia implementation `HomotopyContinuation.jl` [14]. The motivation is twofold: it is implemented in the same programming language as `EigenvalueSolver.jl`, and it is considered the state of the art for the functionalities we are interested in. We point out that due to the extremely efficient implementation of numerical path tracking in `HomotopyContinuation.jl`, the package can outperform our eigenvalue solver even when δ_{square} is significantly larger than δ . For instance, in the case $n = 3, d = 20$ from Table 4, we have $\delta_{\text{square}} = 8000 > \delta = 1765$, but `HomotopyContinuation.jl` tracks all these 8000 paths in no more than 40 seconds. The performance is comparable for the row $n = 6, d = 5$ in Table 4, where `HomotopyContinuation.jl` tracks $\delta_{\text{square}} = 15625$ paths in about 45 seconds. For all the above computations, we used the option `start_system = :total_degree`, which is optimal for dense systems and avoids polyhedral computations to generate start systems.

However, for strongly overdetermined systems, our algorithm outperforms the homotopy approach. For example, for all the cases $n = 15, d = 3$, Table 4 shows that our algorithms take no more than 2 minutes for $\delta \leq 600$. On the other hand, the number δ_{square} equals $3^{15} = 14348907$, for which `HomotopyContinuation.jl` shows an estimated duration of more than 2 days. Additionally, for the case $n = 15, d = 2$ in Table 5, we have $\delta_{\text{square}} = 32765$ and the path tracking takes over 10 minutes, as compared to 48 seconds for the online version of our algorithm and 65 seconds for the offline version. In this last case we use the default `start_system = :polyhedral`.

We conclude that for strongly overdetermined systems ($s \gg n$), `EigenvalueSolver.jl` outperforms `HomotopyContinuation.jl`, which suggests that eigenvalue methods are more suitable to deal with this kind of systems.

Acknowledgments Part of this work was done during the visit of the second author to TU Berlin for the occasion of the MATH+ Thematic Einstein Semester on Algebraic Geometry, Varieties, Polyhedra, Computation. We thank the organizers of this nice semester for making this collaboration possible. We thank the anonymous reviewer for their useful comments and constructive remarks. The first author was funded by the ERC under the European’s Horizon 2020 research and innovation programme (grant agreement No 787840).

References

- [1] W. Auzinger and H. J. Stetter. An Elimination Algorithm for the Computation of All Zeros of a System of Multivariate Polynomial Equations. In *Numerical Mathematics Singapore 1988: Proceedings of the International Conference on Numerical Mathematics*, pages 11–30, Basel, 1988. Birkhäuser Basel.
- [2] M. Bardet, J.-C. Faugère, B. Salvy, and B.-Y. Yang. Asymptotic behaviour of the degree of regularity of semi-regular polynomial systems. In *Proceedings of MEGA*, volume 5, 2005.
- [3] D. J. Bates, A. J. Sommese, J. D. Hauenstein, and C. W. Wampler. *Numerically solving polynomial systems with Bertini*. SIAM, 2013.
- [4] K. Batselier, P. Dreesen, and B. De Moor. A fast recursive orthogonalization scheme for the Macaulay matrix. *Journal of Computational and Applied Mathematics*, 267:20–32, 2014.
- [5] D. Bayer and M. Stillman. A criterion for detecting m-regularity. *Inventiones mathematicae*, 87(1):1–11, 1987.
- [6] M. R. Bender. *Algorithms for sparse polynomial systems: Groebner basis and resultants*. PhD thesis, Sorbonne Université, June 2019.
- [7] M. R. Bender, J.-C. Faugère, A. Mantzaflaris, and E. Tsigaridas. Koszul-type determinantal formulas for families of mixed multilinear systems. *SIAM Journal on Applied Algebra and Geometry*, in press.
- [8] M. R. Bender, J.-C. Faugère, and E. Tsigaridas. Towards mixed gröbner basis algorithms: The multihomogeneous and sparse case. In *Proceedings of the 2018 ACM International Symposium on Symbolic and Algebraic Computation*, ISSAC ’18, pages 71–78. ACM, 2018.
- [9] M. R. Bender, J.-C. Faugère, and E. Tsigaridas. Gröbner basis over semigroup algebras: Algorithms and applications for sparse polynomial systems. *Proceedings of the 44th International Symposium on Symbolic and Algebraic Computation*, 2019.
- [10] M. R. Bender, J.-C. Faugère, A. Mantzaflaris, and E. Tsigaridas. Bilinear Systems with Two Supports: Koszul Resultant Matrices, Eigenvalues, and Eigenvectors. In *Proceedings of the 2018 ACM International Symposium on Symbolic and Algebraic Computation*, ISSAC ’18, pages 63–70, New York, NY, USA, 2018. ACM.
- [11] M. R. Bender and S. Telen. Toric Eigenvalue Methods for Solving Sparse Polynomial Systems. *arXiv:2006.10654v2 [cs, math]*, June 2020. preprint, arXiv: 2006.10654v2.
- [12] D. N. Bernshtein. The number of roots of a system of equations. *Functional Analysis and Its Applications*, 9(3):183–185, July 1975.
- [13] P. Breiding, K. Rose, and S. Timme. Certifying zeros of polynomial systems using interval arithmetic. *arXiv preprint arXiv:2011.05000*, 2020.
- [14] P. Breiding and S. Timme. Homotopycontinuation.jl: A package for homotopy continuation in Julia. In *International Congress on Mathematical Software*, pages 458–465. Springer, 2018.
- [15] L. Busé, M. Chardin, and N. Nemati. Multigraded Sylvester forms, Duality and Elimination Matrices. *arXiv:2104.08941 [cs, math]*, Apr. 2021. arXiv: 2104.08941.
- [16] P. Bürgisser and F. Cucker. *Condition: The Geometry of Numerical Algorithms*. Grundlehren der mathematischen Wissenschaften. Springer-Verlag, Berlin, 2013.
- [17] J. Canny and J. M. Rojas. An Optimal Condition for Determining the Exact Number of Roots of a Polynomial System. In *Proceedings of the 1991 International Symposium on Symbolic and Algebraic Computation*, ISSAC ’91, pages 96–102, New York, NY, USA, 1991. ACM.
- [18] M. Chardin. Powers of ideals and the cohomology of stalks and fibers of morphisms. *Algebra & Number Theory*, 7(1):1–18, Jan. 2013. Publisher: MSP.
- [19] R. M. Corless, P. M. Gianni, and B. M. Trager. A reordered schur factorization method for zero-dimensional polynomial systems with multiple roots. In *Proceedings of the 1997 international symposium on Symbolic and algebraic computation*, pages 133–140, 1997.
- [20] D. A. Cox. *Applications of Polynomial Systems*. CBMS Regional Conference Series in Mathematics. Conference Board of the Mathematical Sciences, 2020.

- [21] D. A. Cox. Stickelberger and the eigenvalue theorem. *arXiv preprint arXiv:2007.12573*, 2020.
- [22] D. A. Cox, J. Little, and D. O’Shea. *Using Algebraic Geometry*. Graduate Texts in Mathematics. Springer-Verlag, New York, 2 edition, 2005.
- [23] D. A. Cox, J. Little, and D. O’Shea. *Ideals, varieties, and algorithms: an introduction to computational algebraic geometry and commutative algebra*. Springer Science & Business Media, 2013.
- [24] D. A. Cox, J. Little, and H. K. Schenck. *Toric Varieties*. American Mathematical Soc., 2011.
- [25] P. Dreesen, K. Batselier, and B. De Moor. Back to the roots: Polynomial system solving, linear algebra, systems theory. *IFAC Proceedings Volumes*, 45(16):1203–1208, 2012.
- [26] D. Eisenbud. *Commutative Algebra: with a View Toward Algebraic Geometry*. Graduate Texts in Mathematics. Springer-Verlag, New York, 2004.
- [27] I. Z. Emiris. On the Complexity of Sparse Elimination. *Journal of Complexity*, 12(2):134–166, June 1996.
- [28] I. Z. Emiris and J. F. Canny. Efficient Incremental Algorithms for the Sparse Resultant and the Mixed Volume. *Journal of Symbolic Computation*, 20(2):117–149, Aug. 1995.
- [29] I. Z. Emiris and B. Mourrain. Computer algebra methods for studying and computing molecular conformations. *Algorithmica*, 25(2):372–402, 1999.
- [30] I. Z. Emiris and B. Mourrain. Matrices in elimination theory. *Journal of Symbolic Computation*, 28(1-2):3–44, 1999.
- [31] J.-C. Faugère, P.-J. Spaenlehauer, and J. Svartz. Sparse gröbner bases: the unmixed case. In *Proceedings of the 39th International Symposium on Symbolic and Algebraic Computation*, pages 178–185, 2014.
- [32] R. Fröberg. An inequality for hilbert series of graded algebras. *Mathematica Scandinavica*, 56(2):117–144, 1985. Publisher: Mathematica Scandinavica.
- [33] I. M. Gelfand, M. M. Kapranov, and A. V. Zelevinsky. *Discriminants, Resultants, and Multidimensional Determinants*. Birkhäuser Boston, Boston, MA, 1994.
- [34] M. Kaluba, B. Lorenz, and S. Timme. Polymake.jl: A new interface to polymake. In *International Congress on Mathematical Software*, pages 377–385. Springer, 2020.
- [35] A. Kondratyev, H. J. Stetter, and F. Winkler. Numerical computation of Gröbner bases. *Proceedings of CASC2004 (Computer Algebra in Scientific Computing)*, pages 295–306, 2004.
- [36] M. Kreuzer and L. Robbiano. *Computational Commutative Algebra 1*. Springer-Verlag, Berlin, 2000.
- [37] D. Lazard. Résolution des systèmes d’équations algébriques. *Theoretical Computer Science*, 15(1):77–110, Jan. 1981.
- [38] C. Massri. Solving a sparse system using linear algebra. *Journal of Symbolic Computation*, 73:157–174, 2016.
- [39] B. Mourrain. A New Criterion for Normal Form Algorithms. In G. Goos, J. Hartmanis, J. van Leeuwen, M. Fossorier, H. Imai, S. Lin, and A. Poli, editors, *Applied Algebra, Algebraic Algorithms and Error-Correcting Codes*, volume 1719, pages 430–442. Springer Berlin Heidelberg, Berlin, Heidelberg, 1999.
- [40] B. Mourrain, S. Telen, and M. Van Barel. Truncated normal forms for solving polynomial systems: Generalized and efficient algorithms. *Journal of Symbolic Computation*, 2019.
- [41] S. Parkinson, H. Ringer, K. Wall, E. Parkinson, L. Erikson, D. Christensen, and T. J. Jarvis. Analysis of normal-form algorithms for solving systems of polynomial equations. *arXiv preprint arXiv:2104.03526*, 2021.
- [42] A. Sommese and C. Wampler. *The numerical solution of systems of polynomials arising in engineering and science*. World Scientific, Jan. 2005.
- [43] H. J. Stetter. *Numerical polynomial algebra*, volume 85. Siam, 2004.
- [44] S. Telen. Numerical root finding via Cox rings. *Journal of Pure and Applied Algebra*, 224(9), 2020.
- [45] S. Telen. *Solving Systems of Polynomial Equations*. PhD thesis, KU Leuven, Leuven, Belgium, 2020.
- [46] S. Telen, B. Mourrain, and M. Van Barel. Solving polynomial systems via truncated normal forms. *SIAM Journal on Matrix Analysis and Applications*, 39(3):1421–1447, 2018.
- [47] S. Telen and M. Van Barel. A stabilized normal form algorithm for generic systems of polynomial equations. *Journal of Computational and Applied Mathematics*, 342:119–132, 2018.
- [48] S. Telen and N. Vannieuwenhoven. Normal forms for tensor rank decomposition. *arXiv preprint arXiv:2103.07411*, 2021.
- [49] J. Vanderstukken, A. Stegeman, and L. De Lathauwer. Systems of polynomial equations, higher-order tensor decompositions and multidimensional harmonic retrieval: a unifying framework—part i: The canonical polyadic decomposition. *Technical Report 17-133, KU Leuven — ESAT/STADIUS*, 2017.
- [50] J. Verschelde. Algorithm 795: PHCpack: A general-purpose solver for polynomial systems by homotopy continuation. *ACM Transactions on Mathematical Software (TOMS)*, 25(2):251–276, 1999.