

THE EREMENKO-LYUBICH CONSTANT

LASSE REMPE

ABSTRACT. Eremenko and Lyubich proved that an entire function f whose set of singular values is bounded is expanding at points where $|f(z)|$ is large. These expansion properties have been at the centre of the subsequent study of this class of functions, now called the *Eremenko–Lyubich class*. We improve the estimate of Eremenko and Lyubich, and show that the new estimate is asymptotically optimal. As a corollary, we obtain an elementary proof that functions in the Eremenko–Lyubich class have lower order at least $1/2$.

1. INTRODUCTION

Eremenko and Lyubich [EL92] introduced the class \mathcal{B} of transcendental entire functions whose set of critical and asymptotic values is bounded. The dynamics of functions in this class, now called the *Eremenko–Lyubich class*, has been studied intensively in recent years; see e.g. [Rem09, RRRS11, Six14, Bis15a, Bis15b, MPS20] and the survey [Six18]. The key property of functions in this class is that they satisfy strong expansion properties near infinity. More precisely, let $R > 0$ be so large that $|f(0)| \leq R$ and such that all critical and asymptotic values of f have modulus at most R . Then there is a holomorphic function

$$F: \mathcal{T} \rightarrow \mathbb{H}_{\ln R} := \{x + iy: x > \ln R\}$$

with $\exp \circ F = f \circ \exp$, where

$$\mathcal{T} = \{\zeta \in \mathbb{C}: |f(\exp(\zeta))| > R\} = \{\zeta \in \mathbb{C}: \operatorname{Re} F(\zeta) > \ln R\}.$$

The map F is a conformal isomorphism when restricted to any connected component of \mathcal{T} ; these components are also called the (*logarithmic*) *tracts* of F .

Eremenko and Lyubich [EL92, Lemma 1] use the Koebe quarter theorem to show that

$$(1.1) \quad |F'(\zeta)| \geq \frac{\operatorname{Re} F(\zeta) - \ln R}{4\pi}$$

for all $\zeta \in \mathcal{T}$; in other words,

$$(1.2) \quad |f'(z)| \geq \frac{(\ln|f(z)| - \ln R) \cdot |f(z)|}{4\pi|z|}$$

whenever $|f(z)| > R$.

The inequality (1.1) has played a fundamental role in the study of the class \mathcal{B} . It therefore seems natural to ask whether the above estimates can be improved. The purpose of this note is to establish the optimal constant in (1.1) and (1.2).

Date: June 27, 2022.

2020 Mathematics Subject Classification. Primary 30D15, Secondary 30D05, 37F10.

1.1. Theorem (Optimal expansion estimates). *Let f and F be as above. Then*

$$(1.3) \quad |F'(\zeta)| \geq \frac{\operatorname{Re} F(\zeta) - \ln R}{2}$$

for all $\zeta \in \mathcal{T}$, and hence

$$(1.4) \quad |f'(z)| \geq \frac{(\ln|f(z)| - \ln R) \cdot |f(z)|}{2|z|}$$

whenever $|f(z)| > R$.

Moreover, the constant 2 is optimal. Indeed, for $f(z) = \cos \sqrt{z}$,

$$(1.5) \quad \lim_{r \rightarrow +\infty} \frac{|f'(-r)| \cdot r}{(\ln|f(-r)|) \cdot |f(-r)|} = \frac{1}{2}.$$

Every component of $\mathcal{V} := \exp(\mathcal{T})$ is simply-connected, and f is bounded on $\partial\mathcal{V}$. By an old result of Wiman [Wim03], it follows that f must have order $\rho(f) \geq 1/2$, where

$$\rho(f) = \limsup_{r \rightarrow \infty} \frac{\ln \ln M(r, f)}{\ln r}.$$

Here $M(r, f) = \max_{|z|=r} |f(z)|$. That $\rho(f) \geq 1/2$ also follows from the famous $\cos \pi\rho$ theorem, proved later by Wiman [Wim15]. See also [Hay64, p. 119], and compare [Lan95, p. 1788] and [BE95, Proof of Corollary 2]. In fact, f must even have *lower order* at least $1/2$, i.e.

$$(1.6) \quad \liminf_{r \rightarrow \infty} \frac{\ln \ln M(r, f)}{\ln r} \geq \frac{1}{2}.$$

According to Heins [Hei48], this result is also due to Wiman; see [RS05, Lemma 3.5], where (1.6) is proved using a version of the Ahlfors distortion theorem. As an application of Theorem 1.1, we obtain a new elementary proof of (1.6).

1.2. Corollary (Lower order of functions in \mathcal{B}). *Let $f \in \mathcal{B}$. Then f has lower order at least $1/2$. More precisely, there are constants $c > 0$ and $r_0 > 1$ such that*

$$\ln M(r, f) \geq c \cdot \sqrt{r}$$

for $r \geq r_0$.

Acknowledgements. I am grateful to Walter Bergweiler and Alex Eremenko for helpful comments on a draft of this paper, and in particular for discussing the history of Corollary 1.2. I also thank Chris Bishop, Tania Grisel Benítez López, Weiwei Cui, Dave Sixsmith and James Waterman for interesting discussions and comments concerning the paper. I am particularly grateful to David Martí-Pete for a careful reading of an earlier manuscript; his corrections and thoughtful comments improved the presentation of the article.

2. PROOF OF THE ESTIMATE

2.1. Lemma (Expansion in logarithmic coordinates). *Suppose that $T \subset \mathbb{C}$ is a simply-connected domain such that $\exp|_T$ is injective, let $\rho \in \mathbb{R}$ and suppose that $\varphi: T \rightarrow \mathbb{H}_\rho$ is a conformal isomorphism. Then*

$$|\varphi'(\zeta)| \geq \frac{\operatorname{Re} \varphi(\zeta) - \rho}{2}$$

for all $\zeta \in T$.

Proof. Post-composing with a translation, we may assume that $\rho = 0$. Let \mathbb{D} denote the unit disc, and consider the Möbius transformation

$$M: \mathbb{D} \rightarrow \mathbb{H}_0; \quad z \mapsto (1 - z)/(1 + z).$$

Let $\zeta \in T$. The map

$$g: \mathbb{D} \rightarrow \exp(T - \zeta); \quad z \mapsto \exp(\varphi^{-1}(\operatorname{Re}(\varphi(\zeta)) \cdot M(z) + i \cdot \operatorname{Im} \varphi(\zeta)) - \zeta)$$

is a conformal isomorphism with $g(0) = 1$. Since $0 \notin \exp(T - \zeta)$, the Koebe quarter theorem implies that

$$4 \geq |g'(0)| = \frac{\operatorname{Re}(\varphi(\zeta)) \cdot |M'(0)|}{|\varphi'(\zeta)|} = \frac{2 \operatorname{Re}(\varphi(\zeta))}{|\varphi'(\zeta)|}.$$

So $|\varphi'(\zeta)| \geq \operatorname{Re} \varphi(\zeta)/2$, as claimed. \square

Proof of Theorem 1.1. If T is a connected component of \mathcal{T} , then T and $\varphi := F|_T$ satisfy the hypotheses of Lemma 2.1. This proves (1.3). To deduce (1.4) from (1.3), differentiate the functional equation $\exp \circ F = f \circ \exp$.

For $f(z) = \cos \sqrt{z}$ and $r > 0$, we have

$$|f'(-r)| = \frac{|\sin i\sqrt{r}|}{2\sqrt{r}}.$$

As $r \rightarrow \infty$,

$$|\sin i\sqrt{r}| = \frac{e^{\sqrt{r}}}{2} + O(e^{-\sqrt{r}}) = |f(-r)| + O(e^{-\sqrt{r}}).$$

Moreover,

$$\ln|f(-r)| = \sqrt{r} - \ln 2 + O(1/|f(x)|^2).$$

Hence

$$|f'(-r)| - \frac{(\ln|f(-r)| + \ln 2) \cdot |f(-r)|}{2r} = O(e^{-\sqrt{r}}/\sqrt{r})$$

as $r \rightarrow \infty$. We may rewrite this as

$$\frac{|f'(-r)| \cdot r}{(\ln|f(-r)|) \cdot |f(-r)|} - \frac{1}{2} = \frac{\ln 2}{\ln|f(-r)|} + O\left(\frac{e^{-\sqrt{r}} \cdot r}{(\ln|f(-r)|) \cdot |f(-r)| \cdot \sqrt{r}}\right) = O(e^{-\sqrt{r}}),$$

which proves (1.5). \square

For future reference, we also record the following reformulation of Lemma 2.1 in terms of the hyperbolic density of T ; see e.g. [BM07] for background on hyperbolic geometry.

2.2. Corollary (Hyperbolic density of logarithmic tracts). *Suppose that $T \subset \mathbb{C}$ is a simply-connected domain such that $\exp|_T$ is injective, and let ρ_T denote the density of the hyperbolic metric of T . Then*

$$(2.1) \quad \rho_T(\zeta) \geq \frac{1}{2} \quad \text{for all } \zeta \in T.$$

Proof. Let $\varphi: T \rightarrow \mathbb{H}_0$ be a conformal isomorphism. The hyperbolic density of \mathbb{H}_0 is given by $\rho_{\mathbb{H}_0}(\omega) = 1/\operatorname{Re}\omega$. The hyperbolic density ρ_T satisfies

$$\rho_T(\zeta) = \rho_{\mathbb{H}_0}(\omega) \cdot |\varphi'(\zeta)| = \frac{|\varphi'(\zeta)|}{\operatorname{Re}\varphi(\zeta)} \geq \frac{1}{2}$$

by Lemma 2.1. □

The following implies Corollary 1.2.

2.3. Corollary (Lower order in logarithmic coordinates). *Suppose that $T \subset \mathbb{C}$ is a simply-connected domain such that $\exp|_T$ is injective, let $\rho \in \mathbb{R}$ and suppose that*

$$\varphi: T \rightarrow \mathbb{H}_\rho$$

is a conformal isomorphism with $\varphi^{-1}(t) \rightarrow \infty$ as $t \rightarrow +\infty$, $t > 0$. Then there are constants $\gamma > 0$ and $\sigma_0 > 0$ such that

$$\sup_{\operatorname{Re}\zeta=\sigma} \ln \operatorname{Re}\varphi(\zeta) \geq \frac{\sigma}{2} - \gamma$$

for $\sigma \geq \sigma_0$.

Proof. Without loss of generality, we may suppose that $\rho = 0$. Set $\zeta_0 := \varphi^{-1}(1)$ and $\gamma := \operatorname{Re}\zeta_0/2$ and $\sigma_0 := \operatorname{Re}\zeta_0$. Let $\sigma > \sigma_0$; by the assumption on φ and the intermediate value theorem, there is $t > 1$ such that $\operatorname{Re}\varphi^{-1}(t) = \sigma$. Set $\zeta := \varphi^{-1}(t)$. By Lemma 2.1,

$$|\zeta - \zeta_0| \leq \int_1^t |(\varphi^{-1})'(x)| dx \leq \int_1^t \frac{2dx}{x} = 2 \ln t = 2 \ln \operatorname{Re}\varphi(\zeta).$$

So


$$\ln \operatorname{Re}\varphi(\zeta) \geq \frac{|\zeta - \zeta_0|}{2} = \frac{\sigma}{2} - \gamma. \quad \square$$

REFERENCES

- [BE95] Walter Bergweiler and Alexandre Eremenko, *On the singularities of the inverse to a meromorphic function of finite order*, Rev. Mat. Iberoamericana **11** (1995), no. 2, 355–373.
- [Bis15a] Christopher J. Bishop, *Constructing entire functions by quasiconformal folding*, Acta Math. **214** (2015), no. 1, 1–60.
- [Bis15b] ———, *Models for the Eremenko-Lyubich class*, J. Lond. Math. Soc. (2) **92** (2015), no. 1, 202–221.
- [BM07] A. F. Beardon and D. Minda, *The hyperbolic metric and geometric function theory*, Quasiconformal mappings and their applications, Narosa, New Delhi, 2007, pp. 9–56. MR 2492498 (2011c:30108)
- [EL92] A. È. Eremenko and M. Yu. Lyubich, *Dynamical properties of some classes of entire functions*, Ann. Inst. Fourier (Grenoble) **42** (1992), no. 4, 989–1020.
- [Hay64] W. K. Hayman, *Meromorphic functions*, Oxford Mathematical Monographs, Clarendon Press, Oxford, 1964.

- [Hei48] Maurice Heins, *Entire functions with bounded minimum modulus; subharmonic function analogues*, Ann. of Math. (2) **49** (1948), 200–213.
- [Lan95] J. K. Langley, *On the multiple points of certain meromorphic functions*, Proc. Amer. Math. Soc. **123** (1995), no. 6, 1787–1795.
- [MPS20] David Martí-Pete and Mitsuhiro Shishikura, *Wandering domains for entire functions of finite order in the Eremenko-Lyubich class*, Proc. Lond. Math. Soc. (3) **120** (2020), no. 2, 155–191.
- [Rem09] Lasse Rempe, *Rigidity of escaping dynamics for transcendental entire functions*, Acta Math. **203** (2009), no. 2, 235–267.
- [RRRS11] Günter Rottenfuß, Johannes Rückert, Lasse Rempe, and Dierk Schleicher, *Dynamic rays of bounded-type entire functions*, Ann. of Math. (2) **173** (2011), no. 1, 77–125.
- [RS05] P. J. Rippon and G. M. Stallard, *Dimensions of Julia sets of meromorphic functions*, J. London Math. Soc. (2) **71** (2005), no. 3, 669–683.
- [Six14] D. J. Sixsmith, *A new characterisation of the Eremenko-Lyubich class*, J. Anal. Math. **123** (2014), 95–105.
- [Six18] David J. Sixsmith, *Dynamics in the Eremenko-Lyubich class*, Conform. Geom. Dyn. **22** (2018), 185–224.
- [Wim03] A. Wiman, *Über die angenäherte Darstellung von ganzen Funktionen*, Ark. Mat. Astron. Fys. **1** (1903), 105–111 (Swedish).
- [Wim15] ———, *Über eine Eigenschaft der ganzen Funktionen von Höhe Null*, Math. Ann. **76**, 197–211 (1915)., 1915.

DEPT. OF MATHEMATICAL SCIENCES, UNIVERSITY OF LIVERPOOL, LIVERPOOL L69 7ZL, UK,

 <https://orcid.org/0000-0001-8032-8580>

Email address: `l.rempe@liverpool.ac.uk`