

# Bicomplex Version of Lebesgue's Dominated Convergence Theorem and Hyperbolic Invariant Measure

Chinmay Ghosh<sup>1</sup>, Soumen Mondal<sup>2</sup>

<sup>1</sup>Department of Mathematics

Kazi Nazrul University

Nazrul Road, P.O.- Kalla C.H.

Asansol-713340, West Bengal, India

chinmayarp@gmail.com

<sup>2</sup>28, Dolua Dakshinpara Haridas Primary School

Beldanga, Murshidabad

Pin-742133

West Bengal, India

mondalsoumen79@gmail.com

## Abstract

In this article we have studied bicomplex valued measurable functions on an arbitrary measurable space. We have established the bicomplex version of Lebesgue's dominated convergence theorem and some other results related to this theorem. Also we have proved the bicomplex version of Lebesgue-Radon-Nikodym theorem. Finally we have introduced the idea of hyperbolic version of invariant measure.

**AMS Subject Classification** (2010) : 28E99, 30G35.

**Keywords and Phrases:** Bicomplex measurable function; Bicomplex Lebesgue integrable function; Hyperbolic invariant measure.

## 1 Introduction

In 1882 Corrado Segre [10] introduced a new number system called bicomplex numbers. Unlike quaternions this number system is a commutative generalization of complex numbers by four reals. The book of G. B. Price [8] is

a good resource of the analysis of bicomplex numbers. Many works have been done on bicomplex functional analysis. Few researchers have worked on bicomplex dynamics, bicomplex topological modules. In this article we have studied bicomplex valued measurable functions on an arbitrary measurable space. We have established the bicomplex version of Lebesgue's dominated convergence theorem and some other results related to this theorem. Also we have proved the bicomplex version of Lebesgue-Radon-Nikodym theorem. Finally we have introduced the idea of hyperbolic version of invariant measure. To prove the results in the first two subsections in our main results we have used the ideas of the book of W. Rudin [9] and for the results in the last subsection we have used [7] and [11].

## 2 Basis definitions

We denote the set of real and complex numbers by  $\mathbb{R}$  and  $\mathbb{C}$  respectively. We may think three imaginary numbers  $\mathbf{i}_1, \mathbf{i}_2$  and  $\mathbf{j}$  governed by the rules

$$\mathbf{i}_1^2 = -1, \mathbf{i}_2^2 = -1, \mathbf{j}^2 = 1$$

$$\begin{aligned} \mathbf{i}_1 \mathbf{i}_2 &= \mathbf{i}_2 \mathbf{i}_1 = \mathbf{j} \\ \mathbf{i}_1 \mathbf{j} &= \mathbf{j} \mathbf{i}_1 = -\mathbf{i}_2 \\ \mathbf{i}_2 \mathbf{j} &= \mathbf{j} \mathbf{i}_2 = -\mathbf{i}_1. \end{aligned}$$

Then we have two complex planes  $\mathbb{C}(\mathbf{i}_1) = \{x + \mathbf{i}_1 y : x, y \in \mathbb{R}\}$  and  $\mathbb{C}(\mathbf{i}_2) = \{x + \mathbf{i}_2 y : x, y \in \mathbb{R}\}$ , both of which are identical to  $\mathbb{C}$ . Bicomplex numbers are defined as  $\zeta = z_1 + \mathbf{i}_2 z_2$  for  $z_1, z_2 \in \mathbb{C}(\mathbf{i}_1)$ . The set of all bicomplex numbers is denoted by  $\mathbb{T}$ . In particular if  $z_1 = x, z_2 = \mathbf{i}_1 y$  where  $x, y \in \mathbb{R}$  we get  $\zeta = x + \mathbf{j}y$  and these type of numbers are called hyperbolic numbers or duplex numbers. The set of all hyperbolic numbers is denoted by  $\mathbb{D}$ . For  $(z_1 + \mathbf{i}_2 z_2), (w_1 + \mathbf{i}_2 w_2) \in \mathbb{T}$ , the addition and multiplication are defined as

$$\begin{aligned} (z_1 + \mathbf{i}_2 z_2) + (w_1 + \mathbf{i}_2 w_2) &= (z_1 + w_1) + \mathbf{i}_2 (z_2 + w_2) \\ (z_1 + \mathbf{i}_2 z_2)(w_1 + \mathbf{i}_2 w_2) &= (z_1 w_1 - z_2 w_2) + \mathbf{i}_2 (z_1 w_2 + z_2 w_1). \end{aligned}$$

With these operations  $\mathbb{T}$  forms a commutative ring with zero divisors. The elements  $z_1 + \mathbf{i}_2 z_2 \in \mathbb{T}$  such that  $z_1^2 + z_2^2 = 0$  are the zero divisors. The interesting property of a bicomplex number is its idempotent representation. Setting  $\mathbf{e}_1 = \frac{1+\mathbf{j}}{2}$  and  $\mathbf{e}_2 = \frac{1-\mathbf{j}}{2}$ , we get

$$z_1 + \mathbf{i}_2 z_2 = (z_1 - \mathbf{i}_1 z_2) \mathbf{e}_1 + (z_1 + \mathbf{i}_1 z_2) \mathbf{e}_2.$$

Many calculations become easier for this representation.

Throughout this article we will consider  $\mathfrak{M}$  to be a  $\sigma$ -algebra in a set  $X$ , unless stated otherwise.

## 2.1 Partial order on $\mathbb{D}$

The set of nonnegative hyperbolic numbers is

$$\mathbb{D}^+ = \{\nu_1 \mathbf{e}_1 + \nu_2 \mathbf{e}_2 : \nu_1, \nu_2 \geq 0\}.$$

A hyperbolic number  $\zeta$  is said to be (strictly) positive if  $\zeta \in \mathbb{D}^+ \setminus \{0\}$ . The set of nonnegative hyperbolic numbers is also defined as

$$\mathbb{D}^+ = \{x + y\mathbf{k} : x^2 - y^2 \geq 0, x \geq 0\}.$$

On the realization of  $\mathbb{D}^+$ , M.E. Luna-Elizarraras et.al.[6] defined a partial order relation on  $\mathbb{D}$ . For two hyperbolic numbers  $\zeta_1, \zeta_2$  the relation  $\preceq_{\mathbb{D}}$  is defined as

$$\zeta_1 \preceq_{\mathbb{D}} \zeta_2 \text{ if and only if } \zeta_2 - \zeta_1 \in \mathbb{D}^+.$$

One can check that this relation is reflexive, transitive and antisymmetric. Therefore  $\preceq_{\mathbb{D}}$  is a partial order relation on  $\mathbb{D}$ . This partial order relation  $\preceq_{\mathbb{D}}$  on  $\mathbb{D}$  is an extension of the total order relation  $\leq$  on  $\mathbb{R}$ . We say  $\zeta_1 \prec_{\mathbb{D}} \zeta_2$  if  $\zeta_1 \preceq_{\mathbb{D}} \zeta_2$  but  $\zeta_1 \neq \zeta_2$ . Also we say  $\zeta_2 \succeq_{\mathbb{D}} \zeta_1$  if  $\zeta_1 \preceq_{\mathbb{D}} \zeta_2$  and  $\zeta_2 \succ_{\mathbb{D}} \zeta_1$  if  $\zeta_1 \prec_{\mathbb{D}} \zeta_2$ .

**Definition 1** For any hyperbolic number  $\zeta = \nu_1 \mathbf{e}_1 + \nu_2 \mathbf{e}_2$ , the  $\mathbb{D}$ -modulus of  $\zeta$  is defined by

$$|\zeta|_{\mathbb{D}} = |\nu_1 \mathbf{e}_1 + \nu_2 \mathbf{e}_2|_{\mathbb{D}} = |\nu_1| \mathbf{e}_1 + |\nu_2| \mathbf{e}_2 \in \mathbb{D}^+$$

where  $|\nu_1|$  and  $|\nu_2|$  are the usual modulus of real numbers.

**Definition 2** A subset  $A$  of  $\mathbb{D}$  is said to be  $\mathbb{D}$ -bounded if there exists  $M \in \mathbb{D}^+$  such that  $|\zeta|_{\mathbb{D}} \preceq_{\mathbb{D}} M$  for any  $\zeta \in A$ .

Set

$$\begin{aligned} A_1 &= \{x \in \mathbb{R} : \exists y \in \mathbb{R}, x\mathbf{e}_1 + y\mathbf{e}_2 \in A\}, \\ A_2 &= \{y \in \mathbb{R} : \exists x \in \mathbb{R}, x\mathbf{e}_1 + y\mathbf{e}_2 \in A\}. \end{aligned}$$

If  $A$  is  $\mathbb{D}$ -bounded then  $A_1$  and  $A_2$  are bounded subset of  $\mathbb{R}$ .

**Definition 3** For a  $\mathbb{D}$ -bounded subset  $A$  of  $\mathbb{D}$ , the **supremum** of  $A$  with respect to the  $\mathbb{D}$ -modulus is defined by

$$\sup_{\mathbb{D}} A = \sup A_1 \mathbf{e}_1 + \sup A_2 \mathbf{e}_2.$$

**Definition 4** A sequence of hyperbolic numbers  $\{\zeta_n\}_{n \geq 1}$  is said to be **convergent to**  $\zeta \in \mathbb{D}$  if for  $\varepsilon \in \mathbb{D}^+ \setminus \{0\}$  there exists  $k \in \mathbb{N}$  such that

$$|\zeta_n - \zeta|_{\mathbb{D}} \prec_{\mathbb{D}} \varepsilon.$$

Then we write

$$\lim_{n \rightarrow \infty} \zeta_n = \zeta.$$

**Definition 5** A sequence of hyperbolic numbers  $\{\zeta_n\}_{n \geq 1}$  is said to be  **$\mathbb{D}$ -Cauchy sequence**  $\zeta \in \mathbb{D}$  if for  $\varepsilon \in \mathbb{D}^+ \setminus \{0\} \exists N \in \mathbb{N}$  such that

$$|\zeta_{N+m} - \zeta_N|_{\mathbb{D}} \prec_{\mathbb{D}} \varepsilon$$

for all  $m = 1, 2, 3, \dots$ .

Note that a sequence of hyperbolic numbers  $\{\zeta_n\}_{n \geq 1}$  is **convergent** if and only if it is a  **$\mathbb{D}$ -Cauchy sequence**.

**Definition 6** A hyperbolic series  $\sum_{n=1}^{\infty} \zeta_n$  is **convergent** if and only if its partial sums is a  **$\mathbb{D}$ -Cauchy sequence**, i.e., for any  $\varepsilon \in \mathbb{D}^+ \setminus \{0\} \exists N \in \mathbb{N}$  such that

$$\left| \sum_{k=1}^m \zeta_{N+k} \right|_{\mathbb{D}} \prec_{\mathbb{D}} \varepsilon$$

for any  $m \in \mathbb{N}$ .

**Definition 7** A hyperbolic series  $\sum_{n=1}^{\infty} \zeta_n$  is  **$\mathbb{D}$ -absolutely convergent** if the series  $\sum_{n=1}^{\infty} |\zeta_n|_{\mathbb{D}}$  is convergent.

Every  **$\mathbb{D}$ -absolutely convergent** series is convergent.

### 3 Main Results

In this section we have established our main results. We have arranged these in three subsections. In the first subsection we have proved the bicomplex version of Lebesgue's dominated convergence theorem. In the second subsection we have focussed on the bicomplex version of Lebesgue-Radon-Nikodym theorem and also we have established bicomplex version of Hahn decomposition theorem. Finally in the last subsection we have introduced the idea of hyperbolic version of invariant measure.

### 3.1 Bicomplex Version of Lebesgue's Dominated Convergence Theorem

**Definition 8** [2] Let  $X$  be a measurable space then the bicomplex valued function  $f = f_1\mathbf{e}_1 + f_2\mathbf{e}_2$  is called  $\mathbb{T}$ -measurable on  $X$  if  $f_1$  and  $f_2$  are complex measurable functions on  $X$ . In particular if  $f_1$  and  $f_2$  are real measurable functions on  $X$  then  $f = f_1\mathbf{e}_1 + f_2\mathbf{e}_2$  is called  $\mathbb{D}$ -measurable function on  $X$ .

For a bicomplex measurable function  $f = f_1\mathbf{e}_1 + f_2\mathbf{e}_2$  one can easily check that  $|f| = |f_1|\mathbf{e}_1 + |f_2|\mathbf{e}_2$  is  $\mathbb{D}$ -measurable. Also for two  $\mathbb{T}$ -measurable functions  $f$  and  $g$  it is routine check up that  $f + g$  and  $fg$  are also  $\mathbb{T}$ -measurable functions.

**Theorem 1** If  $f$  is a  $\mathbb{T}$ -measurable function on a measurable space  $X$  then there is a  $\mathbb{T}$ -measurable function  $\alpha = \alpha_1\mathbf{e}_1 + \alpha_2\mathbf{e}_2$  such that  $|\alpha_1| = 1, |\alpha_2| = 1$  and  $f = \alpha|f|$ .

**Proof.** Let  $f = f_1\mathbf{e}_1 + f_2\mathbf{e}_2$ .

Since  $f$  is a  $\mathbb{T}$ -measurable function on a measurable space  $X$ ,  $f_1$  and  $f_2$  are complex measurable functions on  $X$ .

So there exist complex measurable functions  $\alpha_1, \alpha_2$  such that  $|\alpha_1| = 1, |\alpha_2| = 1$  and  $f_1 = \alpha_1|f_1|, f_2 = \alpha_2|f_2|$ .

Set  $\alpha = \alpha_1\mathbf{e}_1 + \alpha_2\mathbf{e}_2$ . Obviously  $\alpha$  is a  $\mathbb{T}$ -measurable function on  $X$  and the result follows.

**Definition 9** Let  $\mathfrak{M}$  be a  $\sigma$ -algebra in a set  $X$ . A bicomplex function  $\mu = \mu_1\mathbf{e}_1 + \mu_2\mathbf{e}_2$  defined on  $X$  is called a  $\mathbb{T}$ -measure on  $\mathfrak{M}$  if  $\mu_1, \mu_2$  are complex measures on  $\mathfrak{M}$ . In particular if  $\mu_1, \mu_2$  are positive measures on  $\mathfrak{M}$  i.e range of both  $\mu_1, \mu_2$  are  $[0, \infty]$  then  $\mu$  is called a  $\mathbb{D}$ -measure on  $\mathfrak{M}$  and if  $\mu_1, \mu_2$  are real measures on  $\mathfrak{M}$  i.e range of both  $\mu_1, \mu_2$  are  $[0, \infty)$  then  $\mu$  is called a  $\mathbb{D}^+$ -measure on  $\mathfrak{M}$ .

**Definition 10** Let  $\mu = \mu_1\mathbf{e}_1 + \mu_2\mathbf{e}_2$  be a  $\mathbb{D}$ -measure on an arbitrary measurable space  $X$ . We say  $f = f_1\mathbf{e}_1 + f_2\mathbf{e}_2$  to be bicomplex Lebesgue integrable function on  $X$  if

$$\int_X |f_1| d\mu_1 < \infty,$$

$$\int_X |f_2| d\mu_2 < \infty$$

i.e.,  $f_i \in L^1(\mu_i)$ , the set of all complex Lebesgue integrable functions with respect to  $\mu_i$  for  $i = 1, 2$ .

In that case we write

$$\int_X f d\mu = \left( \int_X f_1 d\mu_1 \right) \mathbf{e}_1 + \left( \int_X f_2 d\mu_2 \right) \mathbf{e}_2.$$

We define  $L_{\mathbb{T}}^1(\mu)$  to be the collection of all bicomplex Lebesgue integrable functions on  $X$ .

**Theorem 2** *Let  $f, g \in L_{\mathbb{T}}^1(\mu)$  and  $\alpha, \beta \in \mathbb{C}$ . Then  $\alpha f + \beta g \in L_{\mathbb{T}}^1(\mu)$ , and*

$$\int_X (\alpha f + \beta g) d\mu = \alpha \int_X f d\mu + \beta \int_X g d\mu.$$

**Proof.** Let  $f = f_1 \mathbf{e}_1 + f_2 \mathbf{e}_2, g = g_1 \mathbf{e}_1 + g_2 \mathbf{e}_2, \mu = \mu_1 \mathbf{e}_1 + \mu_2 \mathbf{e}_2, \alpha = \alpha_1 \mathbf{e}_1 + \alpha_2 \mathbf{e}_2, \beta = \beta_1 \mathbf{e}_1 + \beta_2 \mathbf{e}_2$ .

Then

$$\alpha f + \beta g = (\alpha_1 f_1 + \beta_1 g_1) \mathbf{e}_1 + (\alpha_2 f_2 + \beta_2 g_2) \mathbf{e}_2.$$

Since  $f, g \in L_{\mathbb{T}}^1(\mu)$  we have  $\alpha_1 f_1 + \beta_1 g_1 \in L^1(\mu_1)$  and  $\alpha_2 f_2 + \beta_2 g_2 \in L^1(\mu_2)$  and therefore  $\alpha f + \beta g \in L_{\mathbb{T}}^1(\mu)$ .

The last part follows from the facts

$$\begin{aligned} \int_X (\alpha_1 f_1 + \beta_1 g_1) d\mu_1 &= \alpha_1 \int_X f_1 d\mu_1 + \beta_1 \int_X g_1 d\mu_1, \\ \int_X (\alpha_2 f_2 + \beta_2 g_2) d\mu_2 &= \alpha_2 \int_X f_2 d\mu_2 + \beta_2 \int_X g_2 d\mu_2, \end{aligned}$$

and

$$\int_X (\alpha f + \beta g) d\mu = \left( \int_X (\alpha_1 f_1 + \beta_1 g_1) d\mu_1 \right) \mathbf{e}_1 + \left( \int_X (\alpha_2 f_2 + \beta_2 g_2) d\mu_2 \right) \mathbf{e}_2.$$

**Theorem 3** *If  $f \in L_{\mathbb{T}}^1(\mu)$ , then*

$$\left| \int_X f d\mu \right|_{\mathbb{D}} \preceq_{\mathbb{D}} \int_X |f|_{\mathbb{D}} d\mu.$$

**Proof.** Let  $f = f_1 \mathbf{e}_1 + f_2 \mathbf{e}_2, \mu = \mu_1 \mathbf{e}_1 + \mu_2 \mathbf{e}_2$ .

Since  $f \in L^1_{\mathbb{T}}(\mu)$ , we have  $f_1 \in L^1(\mu_1)$  and  $f_2 \in L^1(\mu_2)$ .

Therefore,

$$\begin{aligned} \left| \int_X f d\mu \right|_{\mathbb{D}} &= \left| \left( \int_X f_1 d\mu_1 \right) \mathbf{e}_1 + \left( \int_X f_2 d\mu_2 \right) \mathbf{e}_2 \right|_{\mathbb{D}} \\ &= \left| \int_X f_1 d\mu_1 \right| \mathbf{e}_1 + \left| \int_X f_2 d\mu_2 \right| \mathbf{e}_2 \preceq_{\mathbb{D}} \int_X |f_1| d\mu_1 \mathbf{e}_1 + \int_X |f_2| d\mu_2 \mathbf{e}_2 = \int_X |f|_{\mathbb{D}} d\mu. \end{aligned}$$

**Theorem 4 (Lebesgue's Dominated Convergence Theorem)** Let  $\{f_n = f_{n1} \mathbf{e}_1 + f_{n2} \mathbf{e}_2\}$  be a sequence of  $\mathbb{T}$ -measurable functions on  $X$  such that

$$\lim_{n \rightarrow \infty} f_n(x) = f(x)$$

exists for all  $x \in X$ . If there exists  $g = g_1 \mathbf{e}_1 + g_2 \mathbf{e}_2 \in L^1_{\mathbb{T}}(\mu)$  such that

$$|f_{ni}(x)| \leq g_i(x)$$

for all  $n = 1, 2, 3, \dots$ ;  $i = 1, 2$ ;  $x \in X$ , then  $f \in L^1_{\mathbb{T}}(\mu)$ ,

$$\lim_{n \rightarrow \infty} \int_X |f_n(x) - f(x)| d\mu = 0,$$

and

$$\lim_{n \rightarrow \infty} \int_X f_n(x) d\mu = f(x).$$

**Proof.** Since  $\{f_n\}$  is a sequence of  $\mathbb{T}$ -measurable functions on  $X$ , both  $\{f_{n1}\}$  and  $\{f_{n2}\}$  are sequences of complex measurable functions on  $X$ .

Let  $f = f_1 \mathbf{e}_1 + f_2 \mathbf{e}_2, \mu = \mu_1 \mathbf{e}_1 + \mu_2 \mathbf{e}_2$ .

Thus

$$\lim_{n \rightarrow \infty} f_{ni}(x) = f_i(x)$$

for  $i = 1, 2$ .

Now since

$$|f_{ni}(x)| \leq g_i(x)$$

for all  $n = 1, 2, 3, \dots$ ;  $i = 1, 2$ ;  $x \in X$ , we get  $f_i \in L^1(\mu_i)$  for  $i = 1, 2$  and therefore  $f \in L^1_{\mathbb{T}}(\mu)$ .

Also,

$$\lim_{n \rightarrow \infty} \int_X |f_{n_i}(x) - f_i(x)| d\mu_i = 0,$$

and

$$\lim_{n \rightarrow \infty} \int_X f_{n_i}(x) d\mu_i = f_i(x)$$

for  $i = 1, 2$ . Hence

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_X |f_n(x) - f(x)|_{\mathbb{D}} d\mu \\ &= \lim_{n \rightarrow \infty} \left( \int_X |f_{n_1}(x) - f_1(x)| d\mu_1 \right) \mathbf{e}_1 + \lim_{n \rightarrow \infty} \left( \int_X |f_{n_2}(x) - f_2(x)| d\mu_2 \right) \mathbf{e}_2 \\ &= 0 \end{aligned}$$

and

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_X f_n(x) d\mu \\ &= \lim_{n \rightarrow \infty} \left( \int_X f_{n_1}(x) d\mu_1 \right) \mathbf{e}_1 + \lim_{n \rightarrow \infty} \left( \int_X f_{n_2}(x) d\mu_2 \right) \mathbf{e}_2 \\ &= f_1(x) \mathbf{e}_1 + f_2(x) \mathbf{e}_2 \\ &= f(x). \end{aligned}$$

### 3.2 Bicomplex Version of Lebesgue-Radon-Nikodym Theorem

Let  $\mathfrak{M}$  be a measure space and  $E \in \mathfrak{M}$ . Let  $P = \{E_k\}$  be a partition of  $E$ . Then for all  $E \in \mathfrak{M}$  the  $\mathbb{T}$ -measure  $\mu = \mu_1 \mathbf{e}_1 + \mu_2 \mathbf{e}_2$  on  $\mathfrak{M}$  satisfies

$$\mu(E) = \sum_{k=1}^{\infty} \mu(E_k)$$

for every partition  $\{E_k\}$  of  $E$ .



Let  $\lambda = \lambda_1 \mathbf{e}_1 + \lambda_2 \mathbf{e}_2$  be a  $\mathbb{D}$ -measure on  $\mathfrak{M}$ . We say that  $\lambda$  dominates  $\mu$  on  $\mathfrak{M}$  if  $|\mu_i(E)| \leq \lambda_i(E)$  for all  $E \in \mathfrak{M}$  and for  $i = 1, 2$ . The  $\mathbb{D}$ -modulus of  $\mu$ , denoted by  $|\mu|_{\mathbb{D}}$ , is defined on  $\mathfrak{M}$  by

$$|\mu|_{\mathbb{D}}(E) = \sup_P \sum_{k=1}^{\infty} |\mu(E_k)|_{\mathbb{D}}$$

for all  $E \in \mathfrak{M}$ .

**Theorem 5** For a  $\mathbb{T}$ -measure  $\mu$  on  $\mathfrak{M}$ ,  $|\mu|_{\mathbb{D}}$  is a  $\mathbb{D}$ -measure on  $\mathfrak{M}$ .

**Proof.** Let  $\mu = \mu_1 \mathbf{e}_1 + \mu_2 \mathbf{e}_2$ .

Since  $\mu_1, \mu_2$  being complex measures on  $\mathfrak{M}$ ,  $|\mu_1|$  and  $|\mu_2|$  are positive measures on  $\mathfrak{M}$ .

Hence  $|\mu|_{\mathbb{D}} = |\mu_1| \mathbf{e}_1 + |\mu_2| \mathbf{e}_2$  is a  $\mathbb{D}$ -measure on  $\mathfrak{M}$ .

**Theorem 6** For a  $\mathbb{T}$ -measure  $\mu$  on  $X$ ,

$$|\mu|_{\mathbb{D}}(X) \prec_{\mathbb{D}} \infty_{\mathbb{D}}.$$

**Proof.** Let  $\mu = \mu_1 \mathbf{e}_1 + \mu_2 \mathbf{e}_2$ .

Since  $\mu_1, \mu_2$  being complex measures on  $\mathfrak{M}$ ,

$$|\mu_i|(X) < \infty$$

for  $i = 1, 2$ .

Hence

$$|\mu|_{\mathbb{D}}(X) = |\mu_1|(X) \mathbf{e}_1 + |\mu_2|(X) \mathbf{e}_2 \prec_{\mathbb{D}} \infty_{\mathbb{D}}.$$

Let  $\mu, \lambda$  be two  $\mathbb{T}$ -measures on  $\mathfrak{M}$  and  $c \in \mathbb{D}$ . For all  $E \in \mathfrak{M}$  define

$$\begin{aligned} (\mu + \lambda)(E) &= \mu(E) + \lambda(E), \\ (c\mu)(E) &= c\mu(E). \end{aligned}$$

One can easily check that  $\mu + \lambda$  and  $c\mu$  are also  $\mathbb{T}$ -measures on  $\mathfrak{M}$ . The collection of all  $\mathbb{T}$ -measures on  $\mathfrak{M}$  forms a module space over  $\mathbb{D}$ .

Slight modifying the definition of hyperbolic valued signed measure from [3], we now define it to be a  $\mathbb{T}$ -measure on  $\mathfrak{M}$  having range in  $\mathbb{D}^+ \cup \mathbb{D}^-$ . Let  $\mu$  be a signed  $\mathbb{D}$ -measure on  $\mathfrak{M}$ . Then both  $\mu^+ = \frac{1}{2}(|\mu|_{\mathbb{D}} + \mu)$  and  $\mu^- = \frac{1}{2}(|\mu|_{\mathbb{D}} - \mu)$  are  $\mathbb{D}$ -measures on  $\mathfrak{M}$ . Obviously  $\mu^+$  and  $\mu^-$  are  $\mathbb{D}$ -bounded. Also the Jordan decomposition of a signed  $\mathbb{D}$ -measure is given by

$$\begin{aligned} \mu &= \mu^+ - \mu^-, \\ |\mu|_{\mathbb{D}} &= \mu^+ + \mu^-. \end{aligned}$$

**Definition 11** Let  $\mu = \mu_1 \mathbf{e}_1 + \mu_2 \mathbf{e}_2$  be a  $\mathbb{D}$ -measure and  $\lambda = \lambda_1 \mathbf{e}_1 + \lambda_2 \mathbf{e}_2$  be a  $\mathbb{T}$ -measure on  $\mathfrak{M}$ . Then  $\lambda$  is said to be absolutely  $\mathbb{T}$ -continuous with respect to  $\mu$  if  $\lambda_i$  is absolutely continuous with respect to  $\mu_i$  for  $i = 1, 2$ . We denote this by  $\lambda \ll_{\mathbb{T}} \mu$ .

If for  $A \in \mathfrak{M}$ ,  $\lambda_i$  is concentrated on  $A$  for  $i = 1, 2$ , then  $\lambda$  is said to be  $\mathbb{T}$ -concentrated on  $A$ .

Two  $\mathbb{T}$ -measures  $\lambda' = \lambda'_1 \mathbf{e}_1 + \lambda'_2 \mathbf{e}_2, \lambda'' = \lambda''_1 \mathbf{e}_1 + \lambda''_2 \mathbf{e}_2$  on  $\mathfrak{M}$  are called mutually  $\mathbb{T}$ -singular if  $\lambda'_i$  and  $\lambda''_i$  are mutually singular for  $i = 1, 2$ . We denote this by  $\lambda' \perp_{\mathbb{T}} \lambda''$ .

**Theorem 7** Let  $\lambda, \lambda'$  and  $\lambda''$  be  $\mathbb{T}$ -measures on  $\mathfrak{M}$ . Also let  $\mu$  be a  $\mathbb{D}$ -measure on  $\mathfrak{M}$ . Then the following hold:

- a) If  $\lambda$  is  $\mathbb{T}$ -concentrated on  $A$ , then  $|\lambda|_{\mathbb{D}}$  is also so.
- b) If  $\lambda' \perp_{\mathbb{T}} \lambda''$ , then  $|\lambda'|_{\mathbb{D}} \perp_{\mathbb{T}} |\lambda''|_{\mathbb{D}}$ .
- c) If  $\lambda' \perp_{\mathbb{T}} \mu, \lambda'' \perp_{\mathbb{T}} \mu$ , then  $\lambda' + \lambda'' \perp_{\mathbb{T}} \mu$ .
- d) If  $\lambda' \ll_{\mathbb{T}} \mu, \lambda'' \ll_{\mathbb{T}} \mu$ , then  $\lambda' + \lambda'' \ll_{\mathbb{T}} \mu$ .
- e) If  $\lambda \ll_{\mathbb{T}} \mu$ , then  $|\lambda|_{\mathbb{D}} \ll_{\mathbb{T}} \mu$ .
- f) If  $\lambda' \ll_{\mathbb{T}} \mu, \lambda'' \perp_{\mathbb{T}} \mu$ , then  $\lambda' \perp_{\mathbb{T}} \lambda''$ .
- g) If  $\lambda \ll_{\mathbb{T}} \mu$  and  $\lambda \perp_{\mathbb{T}} \mu$  then  $\lambda = 0$ .

**Proof.** Let  $\lambda = \lambda_1 \mathbf{e}_1 + \lambda_2 \mathbf{e}_2, \lambda' = \lambda'_1 \mathbf{e}_1 + \lambda'_2 \mathbf{e}_2, \lambda'' = \lambda''_1 \mathbf{e}_1 + \lambda''_2 \mathbf{e}_2$  and  $\mu = \mu_1 \mathbf{e}_1 + \mu_2 \mathbf{e}_2$ .

a)  $\lambda$  is  $\mathbb{T}$ -concentrated on  $A$  implies  $\lambda_i$  is concentrated on  $A$  for  $i = 1, 2$ .

Thus  $|\lambda_i|$  is concentrated on  $A$  for  $i = 1, 2$ .

Therefore  $|\lambda|_{\mathbb{D}} = |\lambda_1| \mathbf{e}_1 + |\lambda_2| \mathbf{e}_2$  is  $\mathbb{T}$ -concentrated on  $A$ .

b)  $\lambda' \perp_{\mathbb{T}} \lambda''$  implies  $\lambda'_i$  and  $\lambda''_i$  are mutually singular for  $i = 1, 2$ .

Thus  $|\lambda'_i|$  and  $|\lambda''_i|$  are mutually singular for  $i = 1, 2$ .

Therefore  $|\lambda'|_{\mathbb{D}} = |\lambda'_1| \mathbf{e}_1 + |\lambda'_2| \mathbf{e}_2$  and  $|\lambda''|_{\mathbb{D}} = |\lambda''_1| \mathbf{e}_1 + |\lambda''_2| \mathbf{e}_2$  are mutually  $\mathbb{T}$ -singular.

c)  $\lambda' \perp_{\mathbb{T}} \mu, \lambda'' \perp_{\mathbb{T}} \mu$  implies  $\lambda'_i$  and  $\mu_i$  are mutually singular for  $i = 1, 2$  and  $\lambda''_i$  and  $\mu_i$  are mutually singular for  $i = 1, 2$ .

Thus  $\lambda'_i + \lambda''_i$  and  $\mu_i$  are mutually singular for  $i = 1, 2$ .

Therefore  $\lambda' + \lambda'' = (\lambda'_1 + \lambda''_1) \mathbf{e}_1 + (\lambda'_2 + \lambda''_2) \mathbf{e}_2$  and  $\mu$  are mutually  $\mathbb{T}$ -singular.

d)  $\lambda' \ll_{\mathbb{T}} \mu, \lambda'' \ll_{\mathbb{T}} \mu$  implies  $\lambda'_i$  is absolutely continuous with respect to  $\mu_i$  for  $i = 1, 2$  and  $\lambda''_i$  is absolutely continuous with respect to  $\mu_i$  for  $i = 1, 2$ .

Thus  $\lambda'_i + \lambda''_i$  is absolutely continuous with respect to  $\mu_i$  for  $i = 1, 2$ .

Therefore  $\lambda' + \lambda'' = (\lambda'_1 + \lambda''_1) \mathbf{e}_1 + (\lambda'_2 + \lambda''_2) \mathbf{e}_2$  is absolutely  $\mathbb{T}$ -continuous with respect to  $\mu$ .

e)  $\lambda \ll_{\mathbb{T}} \mu$  implies  $\lambda_i$  is absolutely continuous with respect to  $\mu_i$  for  $i = 1, 2$ .

Thus  $|\lambda_i|$  is absolutely continuous with respect to  $\mu_i$  for  $i = 1, 2$ .

Therefore  $|\lambda|_{\mathbb{D}} = |\lambda_1| \mathbf{e}_1 + |\lambda_2| \mathbf{e}_2$  is absolutely  $\mathbb{T}$ -continuous with respect to  $\mu$ .

f)  $\lambda' \ll_{\mathbb{T}} \mu$  implies  $\lambda'_i$  is absolutely continuous with respect to  $\mu_i$  for  $i = 1, 2$  and  $\lambda'' \perp_{\mathbb{T}} \mu$  implies  $\lambda''_i$  and  $\mu_i$  are mutually singular for  $i = 1, 2$ .

Thus  $\lambda'_i$  and  $\lambda''_i$  are mutually singular for  $i = 1, 2$ .

Therefore  $\lambda' = \lambda'_1 \mathbf{e}_1 + \lambda'_2 \mathbf{e}_2$  and  $\lambda'' = \lambda''_1 \mathbf{e}_1 + \lambda''_2 \mathbf{e}_2$  are mutually  $\mathbb{T}$ -singular.

g)  $\lambda \ll_{\mathbb{T}} \mu$  implies  $\lambda_i$  is absolutely continuous with respect to  $\mu_i$  for  $i = 1, 2$  and  $\lambda \perp_{\mathbb{T}} \mu$  implies  $\lambda_i$  and  $\mu_i$  are mutually singular for  $i = 1, 2$ .

Thus  $\lambda_i = 0$  for  $i = 1, 2$ .

Therefore  $\lambda = \lambda_1 \mathbf{e}_1 + \lambda_2 \mathbf{e}_2 = 0$ .

**Theorem 8 (Lebesgue-Radon-Nikodym Theorem)** *Let  $\mu$  be a  $\sigma$ -finite  $\mathbb{D}$ -measure on  $\mathfrak{M}$ , and let  $\lambda$  be  $\mathbb{T}$ -measure on  $\mathfrak{M}$ .*

a) *There is a unique pair of  $\mathbb{T}$ -measures  $\lambda', \lambda''$  on  $\mathfrak{M}$  such that*

$$\lambda = \lambda' + \lambda''$$

where  $\lambda' \ll_{\mathbb{T}} \mu, \lambda'' \perp_{\mathbb{T}} \mu$ . If  $\lambda$  is  $\mathbb{D}$ -finite measure on  $\mathfrak{M}$  then  $\lambda', \lambda''$  are also so.

b) *For all  $E \in \mathfrak{M}$  there is a unique  $h \in L^1_{\mathbb{T}}(\mu)$  such that*

$$\lambda'(E) = \int_E h d\mu.$$

**Proof.** Let  $\mu = \mu_1 \mathbf{e}_1 + \mu_2 \mathbf{e}_2, \lambda = \lambda_1 \mathbf{e}_1 + \lambda_2 \mathbf{e}_2$ .

Since  $\mu$  is a  $\sigma$ -finite  $\mathbb{D}$ -measure on  $\mathfrak{M}$ ,  $\mu_1$  and  $\mu_2$  are positive  $\sigma$ -finite measures on  $\mathfrak{M}$ . Also  $\lambda_1, \lambda_2$  are complex measures on  $\mathfrak{M}$ .

a) Then for each  $i = 1, 2$  there is a unique pair of complex measures  $\lambda'_i, \lambda''_i$  on  $\mathfrak{M}$  such that

$$\lambda_i = \lambda'_i + \lambda''_i$$

where  $\lambda'_i$  is absolutely continuous with respect to  $\mu_i$  and  $\lambda''_i, \mu_i$  are mutually singular. If  $\lambda$  is positive and finite measure on  $\mathfrak{M}$  then  $\lambda'_i, \lambda''_i$  are also so.

Hence the result follows from these facts.

b) For each  $i = 1, 2$  and for all  $E \in \mathfrak{M}$  there is a unique  $h_i \in L^1(\mu_i)$  such that

$$\lambda'_i(E) = \int_E h_i d\mu_i.$$

Therefore for all  $E \in \mathfrak{M}$  there is a unique  $h = h_1 \mathbf{e}_1 + h_2 \mathbf{e}_2 \in L_{\mathbb{T}}^1(\mu)$  such that

$$\begin{aligned} \lambda'(E) &= \lambda'_1(E) \mathbf{e}_1 + \lambda'_2(E) \mathbf{e}_2 \\ &= \left( \int_E h_1 d\mu_1 \right) \mathbf{e}_1 + \left( \int_E h_2 d\mu_2 \right) \mathbf{e}_2 \\ &= \int_E h d\mu. \end{aligned}$$

**Theorem 9** *Let  $\lambda$  be  $\mathbb{T}$ -measure on  $\mathfrak{M}$  and  $\mu$  be  $\mathbb{D}$ -measure on  $\mathfrak{M}$ . Then the following are equivalent:*

- a)  $\lambda \ll_{\mathbb{T}} \mu$ .
- b) For every  $\varepsilon \in \mathbb{D}^+ / \{0\}$  there exists  $\delta \in \mathbb{D}^+ / \{0\}$  such that  $|\lambda(E)|_{\mathbb{D}} \prec_{\mathbb{D}} \varepsilon$  for all  $E \in \mathfrak{M}$  with  $|\mu(E)|_{\mathbb{D}} \prec_{\mathbb{D}} \delta$ .

**Proof.** Let  $\lambda = \lambda_1 \mathbf{e}_1 + \lambda_2 \mathbf{e}_2, \mu = \mu_1 \mathbf{e}_1 + \mu_2 \mathbf{e}_2$  where  $\lambda_1, \lambda_2$  are complex measures and  $\mu_1, \mu_2$  are positive measures on  $\mathfrak{M}$ .

Then for  $i = 1, 2$  the following two statements are equivalent

- i)  $\lambda_i$  is absolutely continuous with respect to  $\mu_i$ .
- ii) For every  $\varepsilon_i > 0$  there exists  $\delta_i > 0$  such that  $|\lambda_i(E)| < \varepsilon_i$  for all  $E \in \mathfrak{M}$  with  $\mu_i(E) < \delta_i$ .

If for  $i = 1, 2, \lambda_i$  is absolutely continuous with respect to  $\mu_i$  we get  $\lambda \ll_{\mathbb{T}} \mu$ .

Also if for every  $\varepsilon_i > 0$  there exists  $\delta_i > 0$  such that  $|\lambda_i(E)| < \varepsilon_i$  for all  $E \in \mathfrak{M}$  with  $\mu_i(E) < \delta_i$  we can say for every  $\varepsilon = \varepsilon_1 \mathbf{e}_1 + \varepsilon_2 \mathbf{e}_2 \in \mathbb{D}^+ / \{0\}$  there exists  $\delta = \delta_1 \mathbf{e}_1 + \delta_2 \mathbf{e}_2 \in \mathbb{D}^+ / \{0\}$  such that  $|\lambda(E)|_{\mathbb{D}} = |\lambda_1(E)| \mathbf{e}_1 + |\lambda_2(E)| \mathbf{e}_2 \prec_{\mathbb{D}} \varepsilon$  for all  $E \in \mathfrak{M}$  with  $|\mu(E)|_{\mathbb{D}} = |\mu_1(E)| \mathbf{e}_1 + |\mu_2(E)| \mathbf{e}_2 \prec_{\mathbb{D}} \delta$ .

**Theorem 10** *Let  $\mathfrak{M}$  be a  $\sigma$ -algebra on  $X$ . Let  $\mu$  be  $\mathbb{T}$ -measure on  $\mathfrak{M}$ . Then there exists a  $\mathbb{T}$ -measurable function  $h$  such that  $|h(x)|_{\mathbb{D}} = \mathbf{e}_1 + \mathbf{e}_2$  for all  $x \in X$  and such that*

$$d\mu = h d|\mu|_{\mathbb{D}}.$$

**Proof.** Let  $\mu = \mu_1 \mathbf{e}_1 + \mu_2 \mathbf{e}_2$ .

Since for  $i = 1, 2, \mu_i$  is a complex measure on the  $\sigma$ -algebra  $\mathfrak{M}$  in  $X$ , there exists measurable functions  $h_i$  such that  $|h_i(x)| = 1$  for all  $x \in X$  and such that

$$d\mu_i = h_i d|\mu_i|.$$

Setting  $h = h_1\mathbf{e}_1 + h_2\mathbf{e}_2$ , we get

$$\begin{aligned} d\mu &= d\mu_1\mathbf{e}_1 + d\mu_2\mathbf{e}_2 \\ &= h_1d|\mu_1|\mathbf{e}_1 + h_2d|\mu_2|\mathbf{e}_2 \\ &= hd|\mu|_{\mathbb{D}}. \end{aligned}$$

**Theorem 11** Let  $\mu$  be a  $\mathbb{D}$ -measure on  $\mathfrak{M}$ ,  $g \in L^1_{\mathbb{T}}(\mu)$ , and for all  $E \in \mathfrak{M}$

$$\lambda(E) = \int_E g d\mu.$$

Then for all  $E \in \mathfrak{M}$ ,

$$|\lambda|_{\mathbb{D}}(E) = \int_E |g|_{\mathbb{D}} d\mu.$$

**Proof.** Let  $\mu = \mu_1\mathbf{e}_1 + \mu_2\mathbf{e}_2$ ,  $\lambda = \lambda_1\mathbf{e}_1 + \lambda_2\mathbf{e}_2$  and  $g = g_1\mathbf{e}_1 + g_2\mathbf{e}_2$ . Then  $\mu_1, \mu_2$  are positive measures on  $\mathfrak{M}$ .

Since  $g \in L^1_{\mathbb{T}}(\mu)$ , then  $g_1 \in L^1(\mu_1)$  and  $g_2 \in L^1(\mu_2)$ .

Also  $\lambda(E) = \int_E g d\mu \Rightarrow \lambda_1(E) = \int_E g_1 d\mu_1$  and  $\lambda_2(E) = \int_E g_2 d\mu_2$ .

Then by Consequences of the Radon-Nikodym Theorem [9], we have

$$|\lambda_1|(E) = \int_E |g_1| d\mu_1 \text{ and } |\lambda_2|(E) = \int_E |g_2| d\mu_2.$$

Therefore

$$\begin{aligned} |\lambda|_{\mathbb{D}}(E) &= |\lambda_1|(E)\mathbf{e}_1 + |\lambda_2|(E)\mathbf{e}_2 \\ &= \left( \int_E |g_1| d\mu_1 \right) \mathbf{e}_1 + \left( \int_E |g_2| d\mu_2 \right) \mathbf{e}_2 \\ &= \int_E (|g_1|\mathbf{e}_1 + |g_2|\mathbf{e}_2) d(\mu_1\mathbf{e}_1 + \mu_2\mathbf{e}_2) \\ &= \int_E |g|_{\mathbb{D}} d\mu. \end{aligned}$$

**Theorem 12 (Hahn Decomposition Theorem)** *Let  $\mathfrak{M}$  be a  $\sigma$ -algebra on  $X$ . Let  $\mu$  be a  $\mathbb{D}^+$ -measure on  $\mathfrak{M}$ . Then there exists a partition  $\{A, B, C, D\} \subset \mathfrak{M}$  of  $X$  such that for all  $E \in \mathfrak{M}$ ,*

$$\mu^+(E) = \mu(E \cap A) + \mathbf{e}_1 |\mu(E \cap C)|_{\mathbb{D}} + \mathbf{e}_2 |\mu(E \cap D)|_{\mathbb{D}},$$

$$\mu^-(E) = -\mu(E \cap B) - \mu(E \cap C) - \mu(E \cap D) + \mathbf{e}_1 |\mu(E \cap C)|_{\mathbb{D}} + \mathbf{e}_2 |\mu(E \cap D)|_{\mathbb{D}}. \quad (1)$$

**Proof.** By Theorem 10,  $d\mu = hd|\mu|_{\mathbb{D}}$ , where  $|h(x)|_{\mathbb{D}} = \mathbf{e}_1 + \mathbf{e}_2$  for all  $x \in X$ . Since  $\mu$  is hyperbolic, it follows that  $h$  is hyperbolic, hence  $h(x) = \pm \mathbf{e}_1 \pm \mathbf{e}_2$  for all  $x \in X$ . Put

$$A = \{x : h(x) = \mathbf{e}_1 + \mathbf{e}_2\},$$

$$B = \{x : h(x) = -\mathbf{e}_1 - \mathbf{e}_2\},$$

$$C = \{x : h(x) = \mathbf{e}_1 - \mathbf{e}_2\},$$

$$D = \{x : h(x) = -\mathbf{e}_1 + \mathbf{e}_2\}.$$

Since  $\mu^+ = \frac{1}{2}(|\mu|_{\mathbb{D}} + \mu)$ , and since

$$\frac{1}{2}(1 + h) = \begin{cases} h & \text{on } A, \\ 0 & \text{on } B, \\ \mathbf{e}_1 & \text{on } C, \\ \mathbf{e}_2 & \text{on } D, \end{cases}$$

we have, for any  $E \in \mathfrak{M}$ ,

$$\begin{aligned} \mu^+(E) &= \frac{1}{2} \int_E (1 + h) d|\mu|_{\mathbb{D}} \\ &= \int_{E \cap A} h d|\mu|_{\mathbb{D}} + \int_{E \cap C} \mathbf{e}_1 d|\mu|_{\mathbb{D}} + \int_{E \cap D} \mathbf{e}_2 d|\mu|_{\mathbb{D}} \\ &= \mu(E \cap A) + \mathbf{e}_1 |\mu(E \cap C)|_{\mathbb{D}} + \mathbf{e}_2 |\mu(E \cap D)|_{\mathbb{D}}. \end{aligned}$$

Since  $\mu(E) = \mu(E \cap A) + \mu(E \cap B) + \mu(E \cap C) + \mu(E \cap D)$  and since  $\mu = \mu^+ - \mu^-$ , (1) follows.

### 3.3 Hyperbolic Invariant Measure

**Definition 12** *Let  $\mathfrak{B}$  be a Borel  $\sigma$ -algebra on a metric space  $X$ . A  $\mathbb{D}$ -measure  $\mu$  on  $X$  is  $\mathbb{D}$ -finite if  $\mu(X) \prec_{\mathbb{D}} \infty_{\mathbb{D}}$  and  $\mu$  is called a Borel  $\mathbb{D}$ -probability measure if  $\mu(X) = \mathbf{e}_1 + \mathbf{e}_2$  or  $\mathbf{e}_1$  or  $\mathbf{e}_2$ .*

For any  $\mathbb{D}$ -finite, nonzero,  $\mathbb{D}$ -measure  $\widehat{\mu}$  on  $X$  we may define a Borel  $\mathbb{D}$ -probability measure as

$$\mu(A) = \frac{\widehat{\mu}(A)}{\widehat{\mu}(X)}$$

for all  $A \in \mathfrak{B}$ .

Throughout this section we will need a set  $X$  equipped with a Borel  $\sigma$ -algebra  $\mathfrak{B}$  and a measurable function  $f : X \rightarrow X$ . We denote by  $\mathcal{M}_{\mathbb{D}}$ , the space of all Borel  $\mathbb{D}$ -probability measures on  $X$ .

**Definition 13** *A Borel  $\mathbb{D}$ -probability measure  $\mu = \mu_1 \mathbf{e}_1 + \mu_2 \mathbf{e}_2$  on  $X$  is said to be  $\mathbb{D}$ -invariant with respect to a measurable function  $f : X \rightarrow X$  if  $\mu_1$  and  $\mu_2$  are invariant i.e.,  $\mu_1(f^{-1}(A)) = \mu_1(A)$  and  $\mu_2(f^{-1}(A)) = \mu_2(A)$  for all  $A \in \mathfrak{B}$ .*

**Definition 14 (Push-forward of measures)** *Let  $f_* : \mathcal{M}_{\mathbb{D}} \rightarrow \mathcal{M}_{\mathbb{D}}$  be the map from the space of  $\mathbb{D}$ -probability measures to itself, defined by*

$$f_*\mu(A) := \mu(f^{-1}(A)) = \mu_1(f^{-1}(A)) \mathbf{e}_1 + \mu_2(f^{-1}(A)) \mathbf{e}_2 = f_*\mu_1(A) \mathbf{e}_1 + f_*\mu_2(A) \mathbf{e}_2.$$

We call  $f_*\mu$  the push forward of  $\mu$  by  $f$ .

The mapping is well defined and  $\mu$  is invariant iff  $f_*\mu = \mu$ . For any  $\mu \in \mathcal{M}_{\mathbb{D}}$  and any  $i \geq 1$  we also let

$$f_*^i \mu(A) := \mu(f^{-i}(A)).$$

We now prove some properties of the map  $f_*$ .

**Lemma 1** *For all  $\varphi \in L^1_{\mathbb{T}}(\mu)$  we have  $\int \varphi d(f_*\mu) = \int \varphi \circ f d\mu$ .*

**Proof.** *First let  $\varphi = \chi_A$  be the characteristic function of some  $A \subseteq X$ . Then*

$$\begin{aligned} \int \chi_A d(f_*\mu) &= f_*\mu(A) \\ &= \mu(f^{-1}(A)) \\ &= \mu_1(f^{-1}(A)) \mathbf{e}_1 + \mu_2(f^{-1}(A)) \mathbf{e}_2 \\ &= \left( \int \chi_{f^{-1}(A)} d\mu_1 \right) \mathbf{e}_1 + \left( \int \chi_{f^{-1}(A)} d\mu_2 \right) \mathbf{e}_2 \\ &= \int (\chi_A \circ f) d\mu. \end{aligned}$$

*So, the statement is true for characteristic functions and thus for simple functions. Hence the statement is true for general integrable functions by standard approximation arguments.*

**Corollary 1**  $f_* : \mathcal{M}_{\mathbb{D}} \rightarrow \mathcal{M}_{\mathbb{D}}$  is continuous.

**Proof.** Consider a sequence  $\mu_n \rightarrow \mu$  in  $\mathcal{M}_{\mathbb{D}}$ . Then, by Lemma 1, for any continuous function  $\varphi : X \rightarrow \mathbb{D}$  we have

$$\int \varphi d(f_*\mu_n) = \int \varphi \circ f d\mu_n \rightarrow \int \varphi \circ f d\mu = \int \varphi d(f_*\mu)$$

which implies  $f_*\mu_n \rightarrow f_*\mu$ . Hence  $f_*$  is continuous.

**Corollary 2**  $\mu$  is  $\mathbb{D}$ -invariant with respect to a measurable function  $f : X \rightarrow X$  if and only if  $\int \varphi \circ f d\mu = \int \varphi d\mu$  for all continuous function  $\varphi : X \rightarrow \mathbb{D}$ .

**Proof.**  $\varphi = \varphi_1 \mathbf{e}_1 + \varphi_2 \mathbf{e}_2$  is continuous  $\iff \varphi_1$  and  $\varphi_2$  are continuous.

Suppose first that  $\mu$  is  $\mathbb{D}$ -invariant. Then  $f_*\mu = \mu$ . Now using Lemma 1, we have

$$\int \varphi \circ f d\mu = \int \varphi df_*\mu = \int \varphi d\mu.$$

For the converse, we have

$$\int \varphi d\mu = \int \varphi \circ f d\mu = \int \varphi d(f_*\mu).$$

which implies

$$\left( \int \varphi_1 d\mu_1 \right) \mathbf{e}_1 + \left( \int \varphi_2 d\mu_2 \right) \mathbf{e}_2 = \left( \int \varphi_1 df_*\mu_1 \right) \mathbf{e}_1 + \left( \int \varphi_2 df_*\mu_2 \right) \mathbf{e}_2.$$

So,

$$\int \varphi_1 d\mu_1 = \int \varphi_1 df_*\mu_1$$

and

$$\int \varphi_2 d\mu_2 = \int \varphi_2 df_*\mu_2$$

for every continuous function  $\varphi_1, \varphi_2 : X \rightarrow \mathbb{R}$ .

By the Riesz Representation Theorem, measures correspond to linear functionals and therefore this can be restated as saying that

$$\mu_1(\varphi_1) = f_*\mu_1(\varphi_1)$$

and

$$\mu_2(\varphi_2) = f_*\mu_2(\varphi_2)$$

for all continuous function  $\varphi_1, \varphi_2 : X \rightarrow \mathbb{R}$ .



Hence

$$\mu_1 = f_*\mu_1 \text{ and } \mu_2 = f_*\mu_2$$

and so

$$f_*\mu = \mu.$$

Therefore  $\mu$  is  $\mathbb{D}$ -invariant.

We now prove a general result which gives conditions to guarantee that atleast some  $\mathbb{D}$ -invariant  $\mathbb{D}$ -probability measure exists.

Let

$$\mathcal{M}_{\mathbb{D}}(f) = \{\mu \in \mathcal{M}_{\mathbb{D}} : \mu \text{ is } \mathbb{D}\text{-invariant with respect to } f\}.$$

**Theorem 13** *Suppose  $M$  is compact metric space and  $f : M \rightarrow M$  is continuous. Then  $\mathcal{M}_{\mathbb{D}}(f)$  is non-empty, convex, compact.*

**Proof.** *Krylov-Boguliobov Theorem [7] states that if  $M$  is compact metric space and  $f : M \rightarrow M$  is continuous, then the set  $\{\mu^* \in \mathcal{M}^* : \mu^* \text{ is } f\text{-invariant}\}$  is non-empty, where  $\mu^*$  is probability measure and  $\mathcal{M}^*$  is the space of probability measures.*

*It is clear that*

$$\{\mu^* \in \mathcal{M}^* : \mu^* \text{ is } f\text{-invariant}\} \subseteq \mathcal{M}_{\mathbb{D}}(f).$$

*Hence  $\mathcal{M}_{\mathbb{D}}(f)$  is non-empty.*

*Now let  $\mu^1, \mu^2 \in \mathcal{M}_{\mathbb{D}}(f)$ . Then  $\mu^1(f^{-1}(A)) = \mu^1(A)$  and  $\mu^2(f^{-1}(A)) = \mu^2(A)$ , for all  $A \in \mathfrak{B}$ .*

*Now for  $t \in [0, 1]$ , let  $\mu = t\mu^1 + (1 - t)\mu^2$ .*

*Then*

$$\begin{aligned} \mu(f^{-1}(A)) &= t\mu^1(f^{-1}(A)) + (1 - t)\mu^2(f^{-1}(A)) \\ &= t\mu^1(A) + (1 - t)\mu^2(A) \\ &= \mu(A). \end{aligned}$$

*Hence  $\mathcal{M}_{\mathbb{D}}(f)$  is convex.*

*To show compactness, suppose that  $\mu_n$  is a sequence in  $\mathcal{M}_{\mathbb{D}}(f)$  converging to some  $\mu \in \mathcal{M}_{\mathbb{D}}$ . Then by Lemma 1 we have, for any continuous function  $\varphi$ , that  $\int \varphi \circ f d\mu = \lim_{n \rightarrow \infty} \int \varphi \circ f d\mu_n = \lim_{n \rightarrow \infty} \int \varphi d\mu_n = \int \varphi d\mu$ . Therefore by Corollary 2,  $\mu$  is  $\mathbb{D}$ -invariant and so  $\mu \in \mathcal{M}_{\mathbb{D}}(f)$ .*

## References

- [1] D. Alpay, M. E. Luna-Elizarraras, M. Shapiro: Kolmogorov's axioms for probabilities with values in hyperbolic numbers, *Adv. Appl. Clifford Algebras*, 27, 913-929 (2017).
- [2] S. Dubey, R. Kumar and K. Sharma (2014), A note on bicomplex Orlicz spaces, arXiv:1401.7112v1[math.FA], 1-12.
- [3] C. Ghosh, S. Biswas and T. Yasin, Hyperbolic valued signed measures. *Int. J. Math. Trends Technol.* 55(7), 515–522 (2018).
- [4] M. P. Goswami, *Bicomplex Integral Transforms and Applications*, Ph.D. Thesis, MNIT, Jaipur, (October, 2017).
- [5] R. Kumar, K. Sharma, R. Tundup and S. Wazir (2017), Orlicz Spaces with Bicomplex Scalars, arXiv:1401.7112v2[math.FA], 1-33.
- [6] M.E. Luna-Elizarraras, M. Shapiro, D. C. Struppa, A. Vajiac: Bicomplex holomorphic functions: The algebra, geometry and analysis of bicomplex numbers, *Frontiers in Mathematics*, Birkhäuser Basel, 2015.
- [7] S. Luzzatto, *Introduction to Smooth Ergodic Theory*, Abdus Salam International Centre for Theoretical Physics, <https://www.ictp.it/~luzzatto>, (May 15, 2020).
- [8] G. B. Price, *An Introduction to Multicomplex Spaces and Functions*, Marcel Dekker Inc., New York, 1991.
- [9] W. Rudin, *Real and Complex Analysis*, 3rd Edition, Tata McGraw-Hill, New Delhi, 2006.
- [10] C. Segre, Le rappresentazioni reali delle forme complesse Gli Enti Iperalgebrici, *Math. Ann.* 40 (1892), 413–467.
- [11] M. Viana, K. Oliveira, *Foundations of Ergodic Theory*, Cambridge University Press, 2016.