ON COMPACT CLASSES OF SOLUTIONS OF DIRICHLET PROBLEM IN SIMPLY CONNECTED DOMAINS

O. DOVHOPIATYI, E. SEVOST'YANOV

May 21, 2021

Abstract

The article is devoted to questions concerning the problems of compactness of solutions of the Dirichlet problem for the Beltrami equation in some simply connected domain. In terms of prime ends, we have proved results of a detailed form for the case when the maximal dilations of these solutions satisfy certain integral constraints. In addition, in this article we have proved theorems on the local and global behavior of plane and spatial mappings with direct and inverse modulus conditions.

1 Introduction

In our recent joint publication [SD], we proved the compactness theorem of the classes of solutions of the Dirichlet problem for the Beltrami equation in a simply connected Jordan domain whose characteristics satisfy the constraints of the integral type. In this article we are talking about solutions defined in an arbitrary simply connected domain. Since such domains do not have to be Jordanian, this is a slight relaxation of conditions compared to [SD]. Note the existence of solutions of the Dirichlet problem under certain assumptions (see, e.g., [P]). Note that such solutions, generally speaking, do not have a homeomorphic extension to the boundary of the domain in the usual sense, since simply connected domains do not have to be locally connected on their boundary. However, such a continuation usually takes place in terms of the so-called prime ends.

Let D be a domain in \mathbb{C} . In what follows, a mapping $f: D \to \mathbb{C}$ is assumed to be sensepreserving, moreover, we assume that f has partial derivatives almost everywhere. Put $f_{\overline{z}} = (f_x + if_y)/2$ and $f_z = (f_x - if_y)/2$. The complex dilatation of f at $z \in D$ is defined as follows: $\mu(z) = \mu_f(z) = f_{\overline{z}}/f_z$ for $f_z \neq 0$ and $\mu(z) = 0$ otherwise. The maximal dilatation of f at z is the following function:

$$K_{\mu}(z) = K_{\mu_f}(z) = \frac{1 + |\mu(z)|}{1 - |\mu(z)|}.$$
(1.1)

Note that the Jacobian of f at $z \in D$ may be calculated according to the relation

$$J(z,f) = |f_z|^2 - |f_{\overline{z}}|^2.$$

Since we assume that the map f is sense preserving, the Jacobian of this map is positive at all points of its differentiability. Let $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$, and let $\mu : D \to \mathbb{D}$ be a Lebesgue measurable function. Without reference to some mapping f, we define the *maximal dilatation* corresponding to its complex dilatation μ by (1.1). It is easy to see that

$$K_{\mu_f}(z) = \frac{|f_z| + |f_{\overline{z}}|}{|f_z| - |f_{\overline{z}}|}$$

whenever partial derivatives of f exist at $z \in D$ and, in addition, $J(z, f) \neq 0$.

Let D be a domain in \mathbb{R}^n , $n \ge 2$. Recall some definitions (see, for example, [KR₁], [KR₂], [IS] or [SSI]). Let ω be an open set in \mathbb{R}^k , $k = 1, \ldots, n-1$. A continuous mapping $\sigma \colon \omega \to \mathbb{R}^n$ is called a *k*-dimensional surface in \mathbb{R}^n . A surface is an arbitrary (n-1)-dimensional surface σ in \mathbb{R}^n . A surface σ is called a Jordan surface, if $\sigma(x) \neq \sigma(y)$ for $x \neq y$. In the following, we will use σ instead of $\sigma(\omega) \subset \mathbb{R}^n$, $\overline{\sigma}$ instead of $\overline{\sigma(\omega)}$ and $\partial \sigma$ instead of $\overline{\sigma(\omega)} \setminus \sigma(\omega)$. A Jordan surface $\sigma \colon \omega \to D$ is called a *cut* of D, if σ separates D, that is $D \setminus \sigma$ has more than one component, $\partial \sigma \cap D = \emptyset$ and $\partial \sigma \cap \partial D \neq \emptyset$.

A sequence of cuts $\sigma_1, \sigma_2, \ldots, \sigma_m, \ldots$ in D is called a *chain*, if:

(i) the set σ_{m+1} is contained in exactly one component d_m of the set $D \setminus \sigma_m$, wherein $\sigma_{m-1} \subset D \setminus (\sigma_m \cup d_m)$; (ii) $\bigcap_{m=1}^{\infty} d_m = \emptyset$.

Two chains of cuts $\{\sigma_m\}$ and $\{\sigma'_k\}$ are called *equivalent*, if for each m = 1, 2, ... the domain d_m contains all the domains d'_k , except for a finite number, and for each k = 1, 2, ... the domain d'_k also contains all domains d_m , except for a finite number.

The end of the domain D is the class of equivalent chains of cuts in D. Let K be the end of D in \mathbb{R}^n , then the set $I(K) = \bigcap_{m=1}^{\infty} \overline{d_m}$ is called the impression of the end K. Throughout the paper, $\Gamma(E, F, D)$ denotes the family of all paths $\gamma : [a, b] \to \overline{\mathbb{R}^n}$ such that $\gamma(a) \in E$, $\gamma(b) \in F$ and $\gamma(t) \in D$ for every $t \in [a, b]$. In what follows, M denotes the modulus of a family of paths, and the element dm(x) corresponds to the Lebesgue measure in \mathbb{R}^n , $n \ge 2$, see [Va]. Following [Na₂], we say that the end K is a prime end, if K contains a chain of cuts $\{\sigma_m\}$ such that $\lim_{m\to\infty} M(\Gamma(C, \sigma_m, D)) = 0$ for some continuum C in D. In the following, the following notation is used: the set of prime ends corresponding to the domain D, is denoted by E_D , and the completion of the domain D by its prime ends is denoted \overline{D}_P .

Consider the following definition, which goes back to Näkki [Na₂], see also [KR₁]–[KR₂]. We say that the boundary of the domain D in \mathbb{R}^n is *locally quasiconformal*, if each point $x_0 \in \partial D$ has a neighborhood U in \mathbb{R}^n , which can be mapped by a quasiconformal mapping φ onto the unit ball $\mathbb{B}^n \subset \mathbb{R}^n$ so that $\varphi(\partial D \cap U)$ is the intersection of \mathbb{B}^n with the coordinate hyperplane.

For the sets $A, B \subset \mathbb{R}^n$ we set, as usual,

diam
$$A = \sup_{x,y \in A} |x - y|$$
, dist $(A, B) = \inf_{x \in A, y \in B} |x - y|$.

Sometimes we also write d(A) instead of diam A and d(A, B) instead of dist (A, B), if no misunderstanding is possible. The sequence of cuts σ_m , $m = 1, 2, \ldots$, is called *regular*, if $\overline{\sigma_m} \cap \overline{\sigma_{m+1}} = \emptyset$ for $m \in \mathbb{N}$ and, in addition, $d(\sigma_m) \to 0$ as $m \to \infty$. If the end K contains at least one regular chain, then K will be called *regular*. We say that a bounded domain D in \mathbb{R}^n is *regular*, if D can be quasiconformally mapped to a domain with a locally quasiconformal boundary whose closure is a compact in \mathbb{R}^n , and, besides that, every prime end in D is regular. Note that space $\overline{D}_P = D \cup E_D$ is metric, which can be demonstrated as follows. If $g: D_0 \to D$ is a quasiconformal mapping of a domain D_0 with a locally quasiconformal boundary onto some domain D, then for $x, y \in \overline{D}_P$ we put:

$$\rho(x,y) := |g^{-1}(x) - g^{-1}(y)|, \qquad (1.2)$$

where the element $g^{-1}(x)$, $x \in E_D$, is to be understood as some (single) boundary point of the domain D_0 . The specified boundary point is unique and well-defined by [IS, Theorem 2.1, Remark 2.1], cf. [Na₂, Theorem 4.1]. It is easy to verify that ρ in (1.2) is a metric on \overline{D}_P , and that the topology on \overline{D}_P , defined by such a method, does not depend on the choice of the map g with the indicated property.

We say that a sequence $x_m \in D$, m = 1, 2, ..., converges to a prime end of $P \in E_D$ as $m \to \infty$, write $x_m \to P$ as $m \to \infty$, if for any $k \in \mathbb{N}$ all elements x_m belong to d_k except for a finite number. Here d_k denotes a sequence of nested domains corresponding to the definition of the prime end P. Note that for a homeomorphism of a domain D onto D', the end of the domain D uniquely corresponds to some sequence of nested domains in the image under the mapping.

Consider the following Cauchy problem:

$$f_{\overline{z}} = \mu(z) \cdot f_z \,, \tag{1.3}$$

$$\lim_{\zeta \to P} \operatorname{Re} f(\zeta) = \varphi(P) \qquad \forall \ P \in E_D , \qquad (1.4)$$

where $\varphi : E_D \to \mathbb{R}$ is a predefined continuous function. In what follows, we assume that D is some simply connected domain in \mathbb{C} . The solution of the problem (1.3)–(1.4) is called *regular*, if one of two is fulfilled: or f(z) = const in D, or f is an open discrete $W^{1,1}_{loc}(D)$ -mapping such that $J(z, f) \neq 0$ for almost any $z \in D$. Given $z_0 \in D$, a function $\varphi : E_D \to \mathbb{R}$, a function $\Phi : \overline{\mathbb{R}^+} \to \overline{\mathbb{R}^+}$ and a function $\mathcal{M}(\Omega)$ of open sets $\Omega \subset D$, we denote by $\mathfrak{F}^{\mathcal{M}}_{\varphi,\Phi,z_0}(D)$ the class of all regular solutions $f : D \to \mathbb{C}$ of the Cauchy problem (1.3)–(1.4) that satisfy the condition $\operatorname{Im} f(z_0) = 0$ and, in addition,

$$\int_{\Omega} \Phi(K_{\mu}(z)) \cdot \frac{dm(z)}{(1+|z|^2)^2} \leqslant \mathcal{M}(\Omega)$$
(1.5)

for any open set $\Omega \subset D$. The following statement generalizes [Dyb, Theorem 2] to the case of arbitrary simply connected domains.

Theorem 1.1. Let D be some simply connected domain in \mathbb{C} , and let $\Phi : \overline{\mathbb{R}^+} \to \overline{\mathbb{R}^+}$ be a continuous increasing convex function, which satisfies the condition

$$\int_{\delta}^{\infty} \frac{d\tau}{\tau \Phi^{-1}(\tau)} = \infty$$

for some $\delta > \Phi(0)$. Assume that the function \mathcal{M} is bounded, and the function φ in (1.4) is continuous. Then the family $\mathfrak{F}_{\varphi,\Phi,z_0}^{\mathcal{M}}(D)$ is compact in \overline{D}_P .

2 Convergence theorems for mappings with upper estimates for modulus distortion

The proof of the main result is based on the theorems on the global behavior of mappings satisfying the weight Poletsky inequality. Results of a similar type in some other situations have been obtained by us before, see, for example, $[Sev_1]$, $[SevSkv_1]$ and $[SevSkv_3]$. The case we will consider below concerns regular domains and mappings with one normalization condition. This case is considered for the first time in this degree of generality.

Given $p \ge 1$, M_p denotes the *p*-modulus of a family of paths, and the element dm(x) corresponds to a Lebesgue measure in \mathbb{R}^n , $n \ge 2$, see [Va]. In what follows, we usually write $M(\Gamma)$ instead of $M_n(\Gamma)$. Everywhere below, unless otherwise stated, the boundary and the closure of a set are understood in the sense of an extended Euclidean space \mathbb{R}^n . Let $x_0 \in \overline{D}$, $x_0 \neq \infty$,

$$S(x_0, r) = \{x \in \mathbb{R}^n : |x - x_0| = r\}, S_i = S(x_0, r_i), \quad i = 1, 2,$$

$$A = A(x_0, r_1, r_2) = \{x \in \mathbb{R}^n : r_1 < |x - x_0| < r_2\}.$$
(2.1)

Everywhere below, unless otherwise stated, the closure \overline{A} and the boundary ∂A of the set A are understood in the topology of the space $\overline{\mathbb{R}^n} = \mathbb{R}^n \cup \{\infty\}$. Let $Q : \mathbb{R}^n \to \mathbb{R}^n$ be a Lebesgue measurable function satisfying the condition $Q(x) \equiv 0$ for $x \in \mathbb{R}^n \setminus D$, and let $p \ge 1$. Given sets E and F and a given domain D in $\overline{\mathbb{R}^n} = \mathbb{R}^n \cup \{\infty\}$, we denote by $\Gamma(E, F, D)$ the family of all paths $\gamma : [0, 1] \to \overline{\mathbb{R}^n}$ joining E and F in D, that is, $\gamma(0) \in E$, $\gamma(1) \in F$ and $\gamma(t) \in D$

for all $t \in (0, 1)$. According to [MRSY, Chap. 7.6], a mapping $f : D \to \overline{\mathbb{R}^n}$ is called a *ring* Q-mapping at the point $x_0 \in \overline{D} \setminus \{\infty\}$ with respect to p-modulus, if the condition

$$M_p(f(\Gamma(S_1, S_2, D))) \leqslant \int_{A \cap D} Q(x) \cdot \eta^p(|x - x_0|) \, dm(x)$$

$$(2.2)$$

holds for all $0 < r_1 < r_2 < d_0 := \sup_{x \in D} |x - x_0|$ and all Lebesgue measurable functions $\eta : (r_1, r_2) \to [0, \infty]$ such that

$$\int_{r_1}^{r_2} \eta(r) \, dr \ge 1 \,. \tag{2.3}$$

The mapping $f: D \to \overline{\mathbb{R}^n}$ is called a ring *Q*-mapping in $\overline{D} \setminus \{\infty\}$ with respect to *p*-modulus if (2.2) holds for any $x_0 \in \overline{D} \setminus \{\infty\}$. This definition can also be applied to the point $x_0 = \infty$ by inversion: $\varphi(x) = \frac{x}{|x|^2}, \infty \mapsto 0$. In what follows, *h* denotes the so-called chordal metric defined by the equalities

$$h(x,y) = \frac{|x-y|}{\sqrt{1+|x|^2}\sqrt{1+|y|^2}}, \quad x \neq \infty \neq y, \quad h(x,\infty) = \frac{1}{\sqrt{1+|x|^2}}.$$
 (2.4)

For a given set $E \subset \overline{\mathbb{R}^n}$, we set

$$h(E) := \sup_{x,y \in E} h(x,y),$$
 (2.5)

The quantity h(E) in (2.5) is called the *chordal diameter* of the set E. For given sets $A, B \subset \mathbb{R}^n$, we put $h(A, B) = \inf_{x \in A, y \in B} h(x, y)$, where h is a chordal metric defined in (2.4).

Let I be a fixed set of indices and let D_i , $i \in I$, be some sequence of domains. Following [NP, Sect. 2.4], we say that a family of domains $\{D_i\}_{i\in I}$ is equi-uniform with respect to p-modulus if for any r > 0 there exists a number $\delta > 0$ such that the inequality

$$M_p(\Gamma(F^*, F, D_i)) \ge \delta \tag{2.6}$$

holds for any $i \in I$ and any continua $F, F^* \subset D$ such that $h(F) \ge r$ and $h(F^*) \ge r$.

Given a Lebesgue measurable function $Q: \mathbb{R}^n \to [0, \infty]$ and a point $x_0 \in \mathbb{R}^n$ we set

$$q_{x_0}(t) = \frac{1}{\omega_{n-1}r^{n-1}} \int_{S(x_0,t)} Q(x) \, d\mathcal{H}^{n-1} \,, \tag{2.7}$$

where \mathcal{H}^{n-1} denotes (n-1)-dimensional Hausdorff measure. The following lemma was formulated and proved in [Sev₅]; however, for completeness of presentation, we present it in full.

Lemma 2.1. Let $1 \leq p \leq n$, and let $\Phi : [0, \infty] \to [0, \infty]$ be a strictly increasing convex function such that the relation

$$\int_{\delta_0}^{\infty} \frac{d\tau}{\tau \left[\Phi^{-1}(\tau)\right]^{\frac{1}{p-1}}} = \infty$$
(2.8)

holds for some $\delta_0 > \tau_0 := \Phi(0)$. Let \mathfrak{Q} be a family of functions $Q : \mathbb{R}^n \to [0, \infty]$ such that

$$\int_{D} \Phi(Q(x)) \frac{dm(x)}{(1+|x|^2)^n} \leqslant M_0 < \infty$$
(2.9)

for some $0 < M_0 < \infty$. Now, for any $0 < r_0 < 1$ and for every $\sigma > 0$ there exists $0 < r_* = r_*(\sigma, r_0, \Phi) < r_0$ such that

$$\int_{\varepsilon}^{r_0} \frac{dt}{t^{\frac{n-1}{p-1}} q_{x_0}^{\frac{1}{p-1}}(t)} \ge \sigma, \qquad \varepsilon \in (0, r_*),$$

for any $Q \in \mathfrak{Q}$.

Proof. Using the substitution of variables $t = r/r_0$, for any $\varepsilon \in (0, r_0)$ we obtain that

$$\int_{\varepsilon}^{r_0} \frac{dr}{r^{\frac{n-1}{p-1}} q_{x_0}^{\frac{1}{p-1}}(r)} \ge \int_{\varepsilon}^{r_0} \frac{dr}{r q_{x_0}^{\frac{1}{p-1}}(r)} = \int_{\varepsilon/r_0}^{1} \frac{dt}{t q_{x_0}^{\frac{1}{p-1}}(tr_0)} = \int_{\varepsilon/r_0}^{1} \frac{dt}{t \widetilde{q}_0^{\frac{1}{p-1}}(t)}, \quad (2.10)$$

where $\tilde{q}_0(t)$ is the average integral value of the function $\tilde{Q}(x) := Q(r_0 x + x_0)$ over the sphere |x| = t, see the ratio (2.7). Then, according to [RS, Lemma 3.1],

$$\int_{\varepsilon/r_0}^{1} \frac{dt}{t\widetilde{q}_0^{\frac{1}{p-1}}(t)} \ge \frac{1}{n} \int_{eM_*(\varepsilon/r_0)}^{\frac{M_*(\varepsilon/r_0)r_0^n}{\varepsilon^n}} \frac{d\tau}{\tau \left[\Phi^{-1}(\tau)\right]^{\frac{1}{p-1}}},$$
(2.11)

where

$$M_*(\varepsilon/r_0) = \frac{1}{\Omega_n \left(1 - (\varepsilon/r_0)^n\right)} \int_{A(0,\varepsilon/r_0,1)} \Phi\left(Q(r_0x + x_0)\right) \, dm(x) =$$
$$= \frac{1}{\Omega_n \left(r_0^n - \varepsilon^n\right)} \int_{A(x_0,\varepsilon,r_0)} \Phi\left(Q(x)\right) \, dm(x)$$

and $A(x_0, \varepsilon, r_0)$ is defined in (2.1) for $r_1 := \varepsilon$ and $r_2 := r_0$. Observe that $|x| \leq |x - x_0| + |x_0| \leq r_0 + |x_0|$ for any $x \in A(x_0, \varepsilon, r_0)$. Thus

$$M_*\left(\varepsilon/r_0\right) \leqslant \frac{\beta(x_0)}{\Omega_n\left(r_0^n - \varepsilon^n\right)} \int_{A(x_0,\varepsilon,r_0)} \Phi(Q(x)) \frac{dm(x)}{\left(1 + |x|^2\right)^n},$$

where $\beta(x_0) = (1 + (r_0 + |x_0|)^2)^n$. Therefore,

$$M_*\left(\varepsilon/r_0\right) \leqslant \frac{2\beta(x_0)}{\Omega_n r_0^n} M_0$$

for $\varepsilon \leq r_0 / \sqrt[n]{2}$, where M_0 is a constant in (2.9). Observe that

$$M_*\left(\varepsilon/r_0\right) > \Phi(0) > 0\,,$$

because Φ is increasing. Now, by (2.10) and (2.11) we obtain that

$$\int_{\varepsilon}^{r_0} \frac{dr}{r^{\frac{n-1}{p-1}} q_{x_0}^{\frac{1}{p-1}}(r)} \ge \frac{1}{n} \int_{\frac{2\beta(x_0)M_0e}{\Omega_n r_0^n}}^{\frac{\Phi(0)r_0^{-}}{\varepsilon^n}} \frac{d\tau}{\tau \left[\Phi^{-1}(\tau)\right]^{\frac{1}{p-1}}}.$$
(2.12)

The desired conclusion follows from (2.12) and (2.8).

Definitions of a condenser and its capacity may be found, for example, in [Ri, Sect. 10, Ch. II]. The following statement holds (see, e.g., [Ma, item (8.9)]).

Proposition 2.1. Let E = (A, C) be a condenser and let 1 . Then

$$\operatorname{cap}_{p} E \ge n\Omega_{n}^{\frac{p}{n}} \left(\frac{n-p}{p-1}\right)^{p-1} \left[m(C)\right]^{\frac{n-p}{n}}$$

where m(C) is the Lebesgue measure of C.

Consider another auxiliary family of mappings. For $p \ge 1$, given numbers $\delta > 0$, $0 < M_0 < \infty$, a domain $D \subset \mathbb{R}^n$, $n \ge 2$, and a given strictly increasing convex function $\Phi \colon \overline{\mathbb{R}^+} \to \overline{\mathbb{R}^+}$ denote by $\mathfrak{A}_{\Phi,p,\delta,M_0}(D)$ the family all open discrete mappings $f : D \to \mathbb{R}^n$ satisfying relations (2.2)–(2.3) with some $Q = Q_f$ in \overline{D} with respect to *p*-modulus. The following statement is true.

Lemma 2.2. Let $p \in (n-1, n)$, and let $\delta_0 > \tau_0 := \Phi(0)$ be such that the condition (2.9) holds. Now the family $\mathfrak{A}_{\Phi,p,\delta,M_0}(D)$ is equicontinuous in D.

In the lemma 2.2 the equicontinuity of the family of mappings $\mathfrak{A}_{\Phi,p,\delta,M_0}(D)$ should be understood as between the spaces (D,d) and (\mathbb{R}^n,d) , where d is the Euclidean metric.

Proof. Let us fix $x_0 \in D$. Let us use the approach used to prove Lemma 2.4 in [GSS₂]. Let $0 < r_0 < \text{dist}(x_0, \partial D)$. Consider the condenser E = (A, C), where $A = B(x_0, r_0)$, $C = \overline{B(x_0, \varepsilon)}$. In this case, according to [SalSev, Lemma 1]

$$\operatorname{cap}_p f(E) \leqslant \frac{\omega_{n-1}}{I^{p-1}},$$

where

$$I = I(x_0, \varepsilon, r_0) = \int_{\varepsilon}^{r_0} \frac{dr}{r^{\frac{n-1}{p-1}} q_{x_0}^{\frac{1}{p-1}}(r)}$$

and $f(E) = \left(f(B(x_0, r_0)), f(\overline{B(x_0, \varepsilon)}) \right)$. By Lemma 2.1 and due to the condition (2.8) there is a function $\alpha = \alpha(\varepsilon)$ and a number $0 < \varepsilon'_0 < r_0$ such that $\alpha(\varepsilon) \to 0$ as $\varepsilon \to 0$ and, in addition,

$$\operatorname{cap}_p f(E) \leqslant \alpha(\varepsilon)$$

for any $\varepsilon \in (0, \varepsilon'_0)$ and $f \in \mathfrak{A}_{\Phi, p, \delta, M_0}(D)$. By Proposition 2.1 we obtain that

$$\alpha(\varepsilon) \geqslant \operatorname{cap}_p f(E) \geqslant n\Omega_n^{\frac{p}{n}} \left(\frac{n-p}{p-1}\right)^{p-1} \left[m(f(C))\right]^{\frac{n-p}{n}},$$

where m(C) denotes the Lebesgue measure of C. In other words,

$$m(f(C)) \leq \alpha_1(\varepsilon)$$
,

where $\alpha_1(\varepsilon) \to 0$ as $\varepsilon \to 0$. From the last relation it follows that there exists a number $\varepsilon_1 \in (0, 1)$, such that

$$m(f(C)) \leqslant 1 \,, \tag{2.13}$$

where $C = \overline{B(x_0, \varepsilon_1)}$.

Let $E_1 = (A_1, C_{\varepsilon})$, $A_1 = B(x_0, \varepsilon_1)$, and $C_{\varepsilon} = \overline{B(x_0, \varepsilon)}$, $\varepsilon \in (0, \varepsilon_1)$. By Lemma 2.1 there exists a function $\alpha_2(\varepsilon)$ and a number $0 < \varepsilon'_0 < \varepsilon_1$ such that

$$\operatorname{cap}_p f(E_1) \leqslant \alpha_2(\varepsilon)$$

for any $\varepsilon \in (0, \varepsilon'_0)$, where $\alpha_2(\varepsilon) \to 0$ as $\varepsilon \to 0$. On the other hand, by [Kr, Proposition 6], cf. [MRV, Lemma 5.9]

$$\left(c_1 \frac{\left(d(f(\overline{B(x_0,\varepsilon)}))\right)^p}{\left(m(f(B(x_0,\varepsilon_1)))\right)^{1-n+p}}\right)^{\frac{1}{n-1}} \leqslant \operatorname{cap}_p f(E_1) \leqslant \alpha_2(\varepsilon).$$
(2.14)

By (2.13) and (2.14), we obtain that

$$d(f(\overline{B(x_0,\varepsilon)})) \leqslant \alpha_3(\varepsilon), \qquad (2.15)$$

where $\alpha_3(\varepsilon) \to 0$ as $\varepsilon \to 0$. The proof of Lemma 2.2 is complete because the mapping $f \in \mathfrak{A}_{\Phi,p,\delta,M_0}(D)$ is arbitrary. \Box

Given $p \ge 1$, numbers $\delta > 0$, $0 < M_0 < \infty$, a domain $D \subset \mathbb{R}^n$, $n \ge 2$, a point $a \in D$ and a strictly increasing convex function $\Phi \colon \overline{\mathbb{R}^+} \to \overline{\mathbb{R}^+}$ denote by $\mathfrak{F}_{\Phi,a,p,\delta,M_0}(D)$ the family of all homeomorphisms $f : D \to \overline{\mathbb{R}^n}$ satisfying (2.2)–(2.3) in \overline{D} for some $Q = Q_f$ such that $h(f(a), \partial f(D)) \ge \delta$, $h(\overline{\mathbb{R}^n} \setminus f(D)) \ge \delta$ and, in addition, (2.9) holds.

Theorem 2.1. Let $p \in (n-1,n]$, let D be regular, and let $D'_f = f(D)$ be bounded domains with a locally quasiconformal boundary which are equi-uniform with respect to p-modulus over all $f \in \mathfrak{F}_{\Phi,a,p,\delta,M_0}(D)$. If there is $\delta_0 > \tau_0 := \Phi(0)$ such that (2.8) holds, then any $f \in \mathfrak{F}_{\Phi,a,p,\delta,M_0}(D)$ has a continuous extension $\overline{f} : \overline{D}_P \to \mathbb{R}^n$ and, in addition, the family $\mathfrak{F}_{\Phi,a,p,\delta,M_0}(\overline{D})$ of all extended mappings $\overline{f} : \overline{D}_P \to \mathbb{R}^n$ is equicontinuous in \overline{D}_P .

Remark 2.1. In Theorem 2.1, the equicontinuity should be understood in the sense of mappings acting between the spaces (X, d) and (X', d'), where $X = \overline{D}_P$ is the complement of the domain D by its prime ends, and d is one of the possible metrics that correspond to the topological space \overline{D}_P in (1.2). In addition, $X' = \overline{\mathbb{R}^n}$ and d' = h is a chordal (spherical) metric.

An example of a family of plane mappings $f_n(z) = z^n$, $n = 1, 2, ..., z \in \mathbb{D}$, indicates the inaccuracy of the analogue of Theorem 2.1 for mappings with branching, in particular, this theorem is not true under the normalization condition $f_n(0) = 0$, n = 1, 2, ...

Proof. Put $f \in \mathfrak{F}_{\Phi,A,p,\delta,M_0}(D)$ and $Q = Q_f(x)$. Given $x \in \mathbb{R}^n$ we set

$$Q'(x) = \begin{cases} Q(x), & x \in D, Q(x) \ge 1\\ 1, & x \in D, Q(x) < 1\\ 1, & x \notin D \end{cases}$$

Observe that the function Q'(x) satisfies the relation (2.9) up to some constant. Indeed,

$$\int_{D} \Phi(Q'(x)) \frac{dm(x)}{(1+|x|^2)^n} = \int_{\{x \in D: Q(x) < 1\}} \Phi(Q'(x)) \frac{dm(x)}{(1+|x|^2)^n} + \int_{\{x \in D: Q(x) \ge 1\}} \Phi(Q'(x)) \frac{dm(x)}{(1+|x|^2)^n} \le \delta + \Phi(1) \int_{\mathbb{R}^n} \frac{dm(x)}{(1+|x|^2)^n} = M'_0 < \infty.$$

In this case, by Lemma 2.1

$$\int_{\varepsilon}^{\tau_0} \frac{dt}{t^{\frac{n-1}{p-1}} q_{x_0}^{\prime \frac{1}{p-1}}(t)} \to \infty$$

for any $0 < r_0 < 1$ and $\varepsilon \to 0$, where $q'_{x_0}(t) = \frac{1}{\omega_{n-1}r^{n-1}} \int_{S(x_0,t)} Q'(x) d\mathcal{H}^{n-1}$. Besides that,

 $\int_{\varepsilon}^{r_0} \frac{dt}{t^{\frac{n-1}{p-1}} q_{x_0}^{\prime \frac{1}{p-1}}(t)} < \infty \text{ for any } \varepsilon \in (0, r_0), \text{ because } q_{x_0}^{\prime}(t) \ge 1 \text{ for almost any } t \in (0, r_0). \text{ Set } \psi(t) = \frac{1}{t^{\frac{n-1}{p-1}} q_{x_0}^{\frac{1}{p-1}}(t)} \text{ for } t \in (0, r_0). \text{ Now, by the Fubini theorem, we obtain that }$

$$\frac{\omega_{n-1}}{I^{p-1}} = \frac{1}{I^p} \int_{\varepsilon < |x-x_0| < r_0} Q(x) \cdot \psi^p(|x-x_0|) \ dm(x) \to 0, \quad \varepsilon \to 0, \quad (2.16)$$

where $I := I(\varepsilon, r_0) = \int_{\varepsilon}^{r_0} \frac{dt}{t^{\frac{n-1}{p-1}}q_{x_0}^{\prime \frac{1}{p-1}}(r)}$. In this case, by [Sev₅, Lemma 3] a mapping $f \in \mathfrak{F}_{\Phi,A,p,\delta,M_0}(D)$ has a continuous extension to \overline{D}_P . In particular, the strong accessibility of the boundary of $D'_f = f(D)$ with respect to *p*-modulus follows from [SevSkv₁, Remark]. Observe that $f \in \mathfrak{F}_{\Phi,a,p,\delta,M_0}(\overline{D})$ does not take the point infinity for $p \neq n$ (see, e.g., [GSS₁, Lemmas 2.6 and 3.1]). Now, the equicontinuity of $\mathfrak{F}_{\Phi,a,p,\delta,M_0}(\overline{D})$ inside *D* follows by Theorem 4.1 in [RS] for p = n and Lemma 2.2 for p < n.

We prove the equicontinuity of the family $\mathfrak{F}_{\Phi,a,p,\delta,M_0}(\overline{D})$ in $E_D := \overline{D}_P \setminus D$. Let us assume the opposite, namely that there are $\varepsilon_* > 0$, $P_0 \in E_D$, a sequence $x_m \in \overline{D}_P$, $x_m \to P_0$ as $m \to \infty$, and a mapping $f_m \in \mathfrak{F}_{\Phi,a,p,\delta,M_0}(\overline{D})$ such that

$$h(f_m(x_m), f_m(P_0)) \ge \varepsilon_*, \quad m = 1, 2, \dots$$

$$(2.17)$$

Since f_m as a continuous extension at P_0 , we may assume that $x_m \in D$ and, in addition, there is a sequence $x'_m \in \overline{D}_P, x'_m \to P_0$ as $m \to \infty$, such that

$$h(f_m(x_m), f_m(x'_m)) \ge \varepsilon_*/2, \quad m = 1, 2, \dots$$
(2.18)

Let d_m be a sequence of cuts σ_m corresponding to P_0 such that $\sigma_m \subset S(x_0, r_m)$, $x_0 \in \partial D$ and $r_m \to 0$ as $m \to \infty$ (see [KR₁, Lemma 2]). Without loss of a generality, we may assume that $x_m, x'_m \in d_m$. Let γ_m be a path joining x_m and x'_m in d_m .

Since the domain D is regular, the space \overline{D}_P contains at least two prime ends P_1 and $P_2 \in E_D$. Let $P_1 \subset E_D$ be a prime end that does not coincide with P_0 . Suppose that $G_m, m = 1, 2, \ldots$, is a sequence of domains that corresponds to a prime end P_1 . Since the mapping f_m has a continuous extension on \overline{D}_P for any $m = 1, 2, \ldots$, we may choose a sequence $\zeta_m \in G_m, \zeta_m \to P_1$ as $m \to \infty$, such that $h(f_m(\zeta_m), f_m(P_1)) \to 0$ as $m \to \infty$. Note that

$$h(f_m(a), f_m(\zeta_m)) \ge h(f_m(a), f_m(P_1)) - h(f_m(\zeta_m), f_m(P_1)) \ge \delta/2,$$
 (2.19)

for any $m \ge m_0$ and some $m_0 \in \mathbb{N}$. We construct a sequence of continua K_m , m = 1, 2, ...as follows. We join the points ζ_1 and a by an arbitrary path in D, which we denote by K_1 . Next, we join the points ζ_2 and ζ_1 by a path K'_1 , in G_1 . Combining the paths K_1 and K'_1 , we obtain a path K_2 , joining the points a and ζ_2 . And so on. Suppose that at some step we have a path K_m , that join the points ζ_m and a. Join the points ζ_{m+1} and ζ_m with a path K'_m , which lies in G_m . Combining the paths K_m and K'_m , we obtain a path K_{m+1} . And so on (see Figure 1). We show that there is a number $m_1 \in \mathbb{N}$ such that




Figure 1: To proof of Theorem 2.1

$$d_m \cap K_m = \emptyset \quad \forall \quad m \ge m_1 \,. \tag{2.20}$$

We prove this from the opposite, namely, suppose that (2.20) does not hold. Then there is an increasing sequence of numbers $m_k \to \infty$, $k \to \infty$, and points $\xi_k \in K_{m_k} \cap d_{m_k}$, $m = 1, 2, \ldots, .$ Then $\xi_k \to P_0$ as $k \to \infty$.

Note that two cases are possible: either all elements ξ_k belong to $D \setminus G_1$ for $k = 1, 2, \ldots$, or there is a number k_1 such that $\xi_{k_1} \in G_1$. In the second case, consider the sequence ξ_k , $k > k_1$. Note that two cases are possible: or ξ_k for $k > k_1$ belong to $D \setminus G_2$, or there is $k_2 > k_1$ such that $\xi_{k_2} \in G_2$. In the second case, consider the sequence ξ_k , $k > k_2$, and so on. Assume that the element $\xi_{k_{l-1}} \in G_{l-1}$ is already constructed. Note that two cases are possible: either ξ_k belong to $D \setminus G_l$ for $k > k_{l-1}$, or there is a number $k_l > k_{l-1}$ such that $\xi_{k_l} \in G_l$, and etc. This procedure can be both finite or infinite, depending on which we have two possible situations:

1) or there are numbers $n_0 \in \mathbb{N}$ and $l_0 \in \mathbb{N}$ such that that $\xi_k \in D \setminus G_{n_0}$ for all $k > l_0$;

2) or for each there is an element ξ_{k_l} such that $\xi_{k_l} \in G_l$, and the sequence k_l is increasing by $l \in \mathbb{N}$.

Consider each of these cases separately and show that in both of them we come to a contradiction. Let situation 1) holds. Observe that all elements of the sequence ξ_k belong to K_{n_0} , hence the existence of the subsequence ξ_{k_r} , $r = 1, 2, \ldots$, convergent as $r \to \infty$ to some point $\xi_0 \in D$. However, $\xi_k \in d_{m_k}$ and therefore $\xi_0 \in \bigcap_{m=1}^{\infty} \overline{d_m} \subset \partial D$ (see [KR₂, Proposition 1]). The obtained contradiction indicates the impossibility of the case 1). Suppose that case 2) holds, then simultaneously $\xi_k \to P_0$ and $\xi_k \to P_1$ as $k \to \infty$. Since space \overline{D}_P is metric with a metric ρ in (1.2), by the triangle inequality it follows that $P_1 = P_0$, which contradicts the choice of P_1 . The obtained contradiction indicates the validity of the relation (2.20).

By the relation (2.20) and by the definition of cuts $\sigma_m \subset S(x_0, r_m)$, we obtain that

$$\Gamma\left(|\gamma_m|, K_m, D\right) > \Gamma(S(x_0, r_m), S(x_0, \widetilde{\varepsilon_0}), D).$$

Thus

$$f_m(\Gamma(|\gamma_m|, K_m, D)) > f_m(\Gamma(S(x_0, r_m), S(x_0, \widetilde{\varepsilon_0}), D))$$

whence, by the definition of the class $\mathfrak{F}_{\Phi,a,p,\delta,M_0}(\overline{D})$

$$M_p(f_m(\Gamma(|\gamma_m|, K_m, D))) \leqslant$$

$$\leqslant M_p(f_m(\Gamma(S(x_0, r_m), S(x_0, \widetilde{\varepsilon_0}), D)) \leqslant \int_{A \cap D} Q_m(x) \cdot \eta^p(|x - x_0|) \, dm(x) \,, \qquad (2.21)$$

where η is any Lebesgue measurable function satisfying (2.3) for $r_1 \mapsto r_m$ and $r_2 \mapsto \tilde{\varepsilon_0}$, in addition, $Q_m := Q_{f_m}$ corresponds to the function Q in (2.2). Let us to prove the inequality

$$M_p(f_m(\Gamma(S(x_0, r_m), S(x_0, \widetilde{\varepsilon_0}), D)) \leqslant \frac{\omega_{n-1}}{I_m^{p-1}}, \qquad (2.22)$$

where $I_m = \int_{r_m}^{\tilde{\varepsilon}_0} \frac{dr}{r^{\frac{n-1}{p-1}}q^{\frac{1}{p-1}}_{mx_0}(r)}$, $q_{mx_0}(t) = \frac{1}{\omega_{n-1}r^{n-1}} \int_{S(x_0,t)} Q_m(x) d\mathcal{H}^{n-1}$ and $Q_m := Q_{f_m}$ (we set $Q_m(x) \equiv 0$ for $x \notin D$). To do this, we will reason similarly to the proof of Lemma 2.1 in [SalSev]. We may assume that $I \neq 0$, since (2.22) is obviously in this case. We may also assume that $I \neq \infty$, because otherwise we may consider $Q(x) + \delta$ instead of Q(x) in (2.22), and then go to the limit as $\delta \to 0$. Let $0 \neq I \neq \infty$. Then $q_{x_0}(r) \neq 0$ for $r \in (r_m, \tilde{\varepsilon_0})$. Put

$$\psi(t) = \begin{cases} 1/[t^{\frac{n-1}{p-1}}q_{x_0}^{\frac{1}{p-1}}(t)], & t \in (r_m, \widetilde{\varepsilon_0}) \\ 0, & t \notin (r_m, \widetilde{\varepsilon_0}) \end{cases}$$

By the Fubini theorem

$$\int_{A} Q_m(x) \cdot \psi^p(|x - x_0|) \, dm(x) = \omega_{n-1} I_m \,, \tag{2.23}$$

where $A = A(r_m, \tilde{\varepsilon_0}, x_0)$ is defined in (2.1). Observe that the function $\eta_1(t) = \psi(t)/I$, $t \in (r_m, \tilde{\varepsilon_0})$, satisfies (2.3). Now, by (2.2) and (2.23) we obtain the relation (2.22), as required.

Finally, from (2.21), (2.22) and from Lemma 2.1 it follows that

$$M_p(f_m(\Gamma(|\gamma_m|, K_m, D))) \leqslant \frac{\omega_{n-1}}{\left(\int\limits_{r_m}^{\tilde{\varepsilon}_0} \frac{dr}{r^{\frac{n-1}{p-1}}q^{\frac{1}{p-1}}(r)}\right)^{p-1}} \to 0, \quad m \to \infty,$$
(2.24)

where $q_{mx_0}(t) = \frac{1}{\omega_{n-1}r^{n-1}} \int_{S(x_0,t)} Q_m(x) d\mathcal{H}^{n-1}$ and $Q_m := Q_{f_m}$ corresponds to the function Q in (2.7). The relation (2.21) contradicts the equi-uniformity of the sequence of domains $D'_m := f_m(D)$. Indeed, $h(f_m(K_m)) \ge \delta/2$ according to (2.19), and $h(f_m(|\gamma_m|)) \ge \varepsilon_*/2$ by the relation (2.18). Hence, since the sequence of domains $D'_m := f_m(D)$ is equi-uniform, we obtain that

$$M_p(f_m(\Gamma(|\gamma_m|, K_m, D))) = M_p(\Gamma(f_m(|\gamma_m|), f_m(K_m), f_m(D))) \ge \delta_* > 0$$

for some $\delta_* > 0$ and any $m = 1, 2, \ldots$, which contradicts the relation (2.24). The obtained contradiction indicates the incorrectness of the assumption in (2.17). Theorem is proved. \Box

3 Equicontinuity of families of mappings with inverse Poletsky inequality

In this section we deal with mappings $f: D \to \mathbb{R}^n$ області $D \subset \mathbb{R}^n$, $n \ge 2$. The main purpose is to summarize the results of our previous article [SSD]. In particular, it is necessary for proving the key theorems of compactness of mapping classes from the next section. Let $y_0 \in \mathbb{R}^n$, $0 < r_1 < r_2 < \infty$ and

$$A = A(y_0, r_1, r_2) = \{ y \in \mathbb{R}^n : r_1 < |y - y_0| < r_2 \} .$$

Let, as above, $M(\Gamma)$ be a conformal modulus of families of paths Γ in \mathbb{R}^n (see e.g. [Va, rn. 6]). Let $f: D \to \mathbb{R}^n$, $n \ge 2$, and let $Q: \mathbb{R}^n \to [0, \infty]$ be a Lebesgue measurable function such that $Q(x) \equiv 0$ for $x \in \mathbb{R}^n \setminus f(D)$. Let $A = A(y_0, r_1, r_2)$. Let $\Gamma_f(y_0, r_1, r_2)$ denotes the family of all paths $\gamma: [a, b] \to D$ such that $f(\gamma) \in \Gamma(S(y_0, r_1), S(y_0, r_2), A(y_0, r_1, r_2))$, i.e., $f(\gamma(a)) \in S(y_0, r_1), f(\gamma(b)) \in S(y_0, r_2)$, and $\gamma(t) \in A(y_0, r_1, r_2)$ for any a < t < b. We say that f satisfies the inverse Poletsky inequality at $y_0 \in f(D)$ if the relation

$$M(\Gamma_f(y_0, r_1, r_2)) \leqslant \int_{A(y_0, r_1, r_2) \cap f(D)} Q(y) \cdot \eta^n(|y - y_0|) \, dm(y)$$
(3.1)

holds for any Lebesgue measurable function $\eta: (r_1, r_2) \to [0, \infty]$ such that

$$\int_{r_1}^{r_2} \eta(r) \, dr \ge 1 \,. \tag{3.2}$$

The following statement holds.

Theorem 3.1. Let $D \subset \mathbb{R}^n$, $n \ge 2$, be a domain that has a weakly flat boundary, and let $D' \subset \mathbb{R}^n$ be a regular domain. Suppose that f is an open discrete and closed mapping of D onto D' that satisfies the relation (3.1) at each point $y_0 \in D'$. Suppose that for each point $y_0 \in D'$ and $0 < r_1 < r_2 < r_0 := \sup_{y \in D'} |y - y_0|$ there is a set $E \subset [r_1, r_2]$ of a positive linear Lebesgue measure such that the function Q is integrable on $S(y_0, r)$ for almost all $r \in E$. Then the mapping f has a continuous extension $\overline{f}: \overline{D} \to \overline{D'}_P$, moreover $\overline{f}(\overline{D}) = \overline{D'}_P$.

Proof. Fix $x_0 \in \partial D$. It is necessary to show the possibility of continuous extension of the mapping of f to the point x_0 . Using, if necessary, the Möbius transformation $\varphi : \infty \mapsto 0$ and taking into account the invariance of the modulus M in left part of the relation (3.1) (see [Va, Theorem 8.1]), we may assume that $x_0 \neq \infty$.

Assume that the conclusion about the continuous extension of the mapping of f to the point x_0 is not correct. Then any prime end $P_0 \in E_{D'}$ is not a limit of f at the point x_0 , that is, there is a sequence $x_k \to x_0$ as $k \to \infty$ and a number $\varepsilon_0 > 0$ such that $\rho(f(x_k), P_0) \ge \varepsilon_0$ for any $k \in \mathbb{N}$, where ρ is one of metric in (1.2). Since by condition the domain D' is regular, it can be mapped to a bounded domain D_* with a locally quasi-conformal boundary using some quasi-conformal mapping $g: D' \to D_*$. Note that the points of the boundary of the domain D' and the prime ends of the domain D_* are in one-to-one correspondence, see, for example, (see [IS, Theorem 2.1], cf. [Na₂, Theorem 4.1]). In this case, since \overline{D}_* is a compactum in \mathbb{R}^n , the metric space $(\overline{D'}_P, \rho)$ is compact, as well. Now, we may assume that $f(x_k)$ converges to some element $P_1 \neq P_0$ as $k \to \infty$. Since f has no a limit at x_0 , there is at least one more sequence y_k , $k = 1, 2..., y_k \to x_0$ as $k \to \infty$, such that $\rho(f(y_k), P_1) \ge \varepsilon_1$ for any $k \in \mathbb{N}$ and some $\varepsilon_1 > 0$. Again, since the metric space $(\overline{D'}_P, \rho)$ is compact, we may assume that $f(y_k) \to P_2$ as $k \to \infty$, $P_1 \neq P_2$, $P_2 \in \overline{D'}_P$. Since the mapping f is closed, it preserves the boundary of the domain, see [Vu, Theorem 3.3]. So, $P_1, P_2 \in E_{D'}$.

Let σ_m and σ'_m , $m = 0, 1, 2, \ldots$, are sequences of cuts which correspond to prime ends P_1 and P_2 , respectively. Assume that cuts σ_m , $m = 0, 1, 2, \ldots$, belong to spheres $S(z_0, r_m)$ centered at some point $z_0 \in \partial D'$, where $r_m \to 0$ as $m \to \infty$ (such a sequence σ_m exists by [IS, Lemma 3.1], cf. [KR₂, Lemma 1]). Let d_m and g_m , $m = 0, 1, 2, \ldots$, be sequences of domains in D', corresponding to cuts σ_m and σ'_m , respectively. Since the space $(\overline{D'}_P, \rho)$ is metric, we may consider that d_m and g_m are disjoint for any $m = 0, 1, 2, \ldots$, in particular,

$$d_0 \cap g_0 = \varnothing \,. \tag{3.3}$$

Since $f(x_k)$ converges to P_1 as $k \to \infty$, for any $m \in \mathbb{N}$ there is k = k(m) such that $f(x_k) \in d_m$ for $k \ge k = k(m)$. By renumbering the sequence x_k , we may ensure that $f(x_k) \in d_k$ for any natural k. Similarly, we may assume that $f(y_k) \in g_k$ for any $k \in \mathbb{N}$. Fix points $f(x_1)$ and $f(y_1)$. Since by the definition of the prime end $\bigcap_{k=1}^{\infty} d_k = \bigcap_{l=1}^{\infty} g_l = \emptyset$, there are numbers k_1 and $k_2 \in \mathbb{N}$ such that $f(x_1) \notin d_{k_1}$ and $f(y_1) \notin g_{k_2}$. By the definition of d_k and g_k , we have that $d_k \subset d_{k_0}$ for any $k \ge k_1$ and $g_k \subset g_{k_2}$ for any $k \ge k_2$. Now, we obtain that

$$f(x_1) \notin d_k$$
, $f(y_1) \notin g_k$, $k \ge \max\{k_1, k_2\}$.

Let γ_k be a path joining $f(x_1)$ and $f(x_k)$ in d_1 , and let γ'_k be a path joining $f(y_1)$ and $f(y_k)$ in g_1 . In addition, let α_k and β_k are total f-liftings of γ_k and γ'_k in D starting at x_k and y_k , respectively (they exists by [Vu, Lemma 3.7]), see Figure 2). Observe that the points $f(x_1)$




Figure 2: To proof of Theorem 3.1

and $f(y_1)$ cannot have more than a finite number of pre-images in D under the mapping f, see [Vu, Lemma 3.2]. Then there is $R_0 > 0$ such that $\alpha_k(1), \beta_k(1) \in D \setminus B(x_0, R_0)$ for any

 $k = 1, 2, \dots$ Since the boundary of D is weakly flat, for every P > 0 there is $k = k_P \ge 1$ such that

$$M(\Gamma(|\alpha_k|, |\beta_k|, D)) > P \qquad \forall \ k \ge k_P.$$

$$(3.4)$$

Let us to show that the condition (3.4) contradicts to (3.1). Indeed, let $\gamma \in \Gamma(|\alpha_k|, |\beta_k|, D)$. Then $\gamma : [0,1] \to D$, $\gamma(0) \in |\alpha_k|$ and $\gamma(1) \in |\beta_k|$. In particular, $f(\gamma(0)) \in |\gamma_k|$ and $f(\gamma(1)) \in |\gamma'_k|$. In this case, by (3.3) and (3.4) we obtain that $|f(\gamma)| \cap d_1 \neq \emptyset \neq |f(\gamma)| \cap (D' \setminus d_1)$ for $k \ge \max\{k_1, k_2\}$. By [Ku, Theorem 1.I.5.46] $|f(\gamma)| \cap \partial d_1 \neq \emptyset$, i.e., $|f(\gamma)| \cap S(z_0, r_1) \neq \emptyset$, because $\partial d_1 \cap D' \subset \sigma_1 \subset S(z_0, r_1)$ by the definition of the cut σ_1 . Let $t_1 \in (0, 1)$ be such that $f(\gamma(t_1)) \in S(z_0, r_1)$ and $f(\gamma)|_1 := f(\gamma)|_{[t_1, 1]}$. Without loss of generality, we may assume that $f(\gamma(t_2)) \in S(z_0, r_0)$. Set $f(\gamma)|_2 := f(\gamma)|_{[t_1, t_2]}$. Now, the path $f(\gamma)|_2$ is a subpath of $f(\gamma)$ and, moreover, $f(\gamma)|_2 \in \Gamma(S(z_0, r_1), S(z_0, r_0), D')$. Without loss of generality, we may assume that $f(\gamma)|_2 \subset B(z_0, r_0)$. Therefore, $\Gamma(|\alpha_k|, |\beta_k|, D) > \Gamma_f(z_0, r_1, r_0)$. From the last relation, by the minorization of the modulus of families of paths (see, e.g., [Fu, Theorem 1(c)]) we obtain that

$$M(\Gamma(|\alpha_k|, |\beta_k|, D)) \leqslant$$

$$\leqslant M(\Gamma_f(z_0, r_1, r_0)) \leqslant \int_A Q(y) \cdot \eta^n(|y - z_0|) \, dm(y) \,, \qquad (3.5)$$

where $A = A(z_0, r_1, r_0)$ and η be an arbitrary Lebesgue measurable function with (3.2). Bellow we set, as usual, $a/\infty = 0$ for $a \neq \infty$, $a/0 = \infty$ for a > 0 and $0 \cdot \infty = 0$ (see, e.g., [Sa, 3.I]). Set

$$I = \int_{r_1}^{r_0} \frac{dt}{tq_{z_0}^{1/(n-1)}(t)},$$
(3.6)

where

$$q_{z_0}(r) = \frac{1}{\omega_{n-1}r^{n-1}} \int_{S(z_0,r)} Q(y) \, d\mathcal{H}^{n-1}(y)$$

and ω_{n-1} demotes the area of the unit sphere \mathbb{S}^{n-1} in \mathbb{R}^n . By the assumption, there is a set $E \subset [r_1, r_0]$ of a positive linear Lebesgue measure such that $q_{z_0}(t)$ is finite for almost all $t \in E$. Thus, $I \neq 0$ in (3.6). In this case, the function $\eta_0(t) = \frac{1}{Itq_{z_0}^{1/(n-1)}(t)}$ satisfies the relation (3.2) for $r_2 := r_0$. Substituting this function into the right-hand side of the inequality (3.5) and applying Fubini's theorem, we obtain that

$$M(\Gamma(|\alpha_k|, |\beta_k|, D)) \leqslant \frac{\omega_{n-1}}{I^{n-1}} < \infty.$$
(3.7)

The relation (3.7) contradicts the condition (3.4). The obtained contradiction refutes the assumption that there is no limit of the mapping f at the point x_0 .

It remains to check the equality $\overline{f}(\overline{D}) = \overline{D'}_P$. Obviously, $\overline{f}(\overline{D}) \subset \overline{D'}_P$. We show that $\overline{D'}_P \subset \overline{f}(\overline{D})$. Indeed, let $y_0 \in \overline{D'}_P$. Then either $y_0 \in D'$ or $y_0 \in E_{D'}$. If $y_0 \in D'$, then $y_0 = f(x_0)$ and $y_0 \in \overline{f}(\overline{D})$, because f maps D onto D'. Finally, let $y_0 \in E_{D'}$, then by the

regularity of the domain D' there is a sequence $y_k \in D'$ such that $\rho(y_k, y_0) \to 0$ as $k \to \infty$, $y_k = f(x_k)$ and $x_k \in D$, where ρ is one of possible metrics in $\overline{D'}_P$. Since \mathbb{R}^n is a compact space, we may assume that $x_k \to x_0$, where $x_0 \in \overline{D}$. Observe that $x_0 \in \partial D$, because f is open. Now $f(x_0) = y_0 \in \overline{f}(\partial D) \subset \overline{f}(\overline{D})$. The theorem is completely proved. \Box

Corollary 3.1. The statement of Theorem 3.1 remains valid if the condition for the existence of the set $E \subset [r_1, r_2]$ with the specified property is replaced by the condition $Q \in L^1(D')$.

Proof. Indeed, by the Fubini theorem (see, e.g., [Sa, Theorem 8.1.III])

$$\int_{B(y_0,r_0)} Q(x) \, dm(x) = \int_0^1 \int_{S(y_0,r)} Q(x) \, d\mathcal{H}^{n-1} dr < \infty \,,$$

because the function Q is integrable in D' (here we set $Q(y) \equiv 0$ for $y \notin D'$). It follows that Q is integrable on almost all spheres $S(y_0, r)$, $0 < r_0 < \leq r_0$. In this case, the desired statement follows from Theorem 3.1 for $E = [r_1, r_2]$. \Box

The following lemma was proved in [SSI, Lemma 2.1].

Lemma 3.1. Let $D \subset \mathbb{R}^n$, $n \ge 2$, be a regular domain, and let $x_m \to P_1$, $y_m \to P_2$ as $m \to \infty$, $P_1, P_2 \in \overline{D}_P$, $P_1 \ne P_2$. Suppose that $d_m, g_m, m = 1, 2, \ldots$, are sequences of descending domains, corresponding to P_1 and $P_2, d_1 \cap g_1 = \emptyset$, and $x_0, y_0 \in D \setminus (d_1 \cup g_1)$. Then there are arbitrarily large $k_0 \in \mathbb{N}$, $M_0 = M_0(k_0) \in \mathbb{N}$ and $0 < t_1 = t_1(k_0), t_2 = t_2(k_0) < 1$ for which the following condition is fulfilled: for each $m \ge M_0$ there are non-intersecting paths

$$\gamma_{1,m}(t) = \begin{cases} \widetilde{\alpha}(t), & t \in [0, t_1], \\ \widetilde{\alpha_m}(t), & t \in [t_1, 1] \end{cases}, \quad \gamma_{2,m}(t) = \begin{cases} \widetilde{\beta}(t), & t \in [0, t_2], \\ \widetilde{\beta_m}(t), & t \in [t_2, 1] \end{cases}$$

such that:

1) $\gamma_{1,m}(0) = x_0, \ \gamma_{1,m}(1) = x_m, \ \gamma_{2,m}(0) = y_0 \text{ and } \gamma_{2,m}(1) = y_m;$ 2) $|\gamma_{1,m}| \cap \overline{g_{k_0}} = \emptyset = |\gamma_{2,m}| \cap \overline{d_{k_0}};$ 3) $\widetilde{\alpha_m}(t) \in d_{k_0+1} \text{ for } t \in [t_1, 1] \text{ and } \widetilde{\beta_m}(t) \in g_{k_0+1} \text{ for } t \in [t_2, 1] \text{ (see Figure 3).}$

Let $\gamma_{1,m}$ and $\gamma_{2,m}$ be the paths from Lemma 3.1, and let $|\gamma_{1,m}|$ and $|\gamma_{2,m}|$ be their images in \mathbb{R}^n , respectively. The following statement is proved in [SSI, Lemma 2.2] for the case of homeomorphisms. Its proof and content are similar to [Sev₄, Lemma 2], however, for completeness of presentation, we give it in full in the text.

Lemma 3.2. Let D and D' be domains in \mathbb{R}^n , $n \ge 2$, let D' be a regular domain, and let f be an open discrete and closed mapping of D onto D' satisfying the condition (3.1) at any point $y_0 \in \overline{D'}$ with some function $Q \in L^1(D')$. Let d_m be a sequence of decreasing domains which correspond to cuts σ_m , $m = 1, 2, \ldots$, lying on the spheres $S(\overline{x_0}, r_m)$ and such that $\overline{x_0} \in \partial D'$, wherein $r_m \to 0$ as $m \to \infty$. Then in the conditions and notation of Lemma 3.1




Figure 3: To the statement of Lemma 3.1

we may choose the number $k_0 \in \mathbb{N}$ for which there is $0 < N = N(k_0, ||Q||_1, D') < \infty$, independent on m and f, such that

$$M(\Gamma_m) \leqslant N, \qquad m \ge M_0 = M_0(k_0),$$

where Γ_m is a family of paths $\gamma : [0,1] \to D$ in D such that $f(\gamma) \in \Gamma(|\gamma_{1,m}|, |\gamma_{2,m}|, D')$. *Proof.* Let k_0 be an arbitrary number for which the statement of Lemma 3.1 holds. By the definition of $\gamma_{1,m}$ and of the family Γ_m we may write that

$$\Gamma_m = \Gamma_m^1 \cup \Gamma_m^2 \,, \tag{3.8}$$

where Γ_m^1 is a family of paths $\gamma \in \Gamma_m$ such that $f(\gamma) \in \Gamma(|\widetilde{\alpha}|, |\gamma_{2,m}|, D')$ and Γ_m^2 is a family of paths $\gamma \in \Gamma_m$ such that $f(\gamma) \in \Gamma(|\widetilde{\alpha}_m|, |\gamma_{2,m}|, D')$.

Taking into account the notation of Lemma 3.1, we put

 $\varepsilon_0 := \min\{ \operatorname{dist}(|\widetilde{\alpha}|, \overline{g_{k_0}}), \operatorname{dist}(|\widetilde{\alpha}|, |\widetilde{\beta}|) \} > 0 \,.$

Let us consider the covering $\bigcup_{x \in |\widetilde{\alpha}|} B(x, \varepsilon_0/4)$ of $|\widetilde{\alpha}|$. Since $|\widetilde{\alpha}|$ is a compactum in D', there are numbers i_1, \ldots, i_{N_0} such that $|\widetilde{\alpha}| \subset \bigcup_{i=1}^{N_0} B(z_i, \varepsilon_0/4)$, where $z_i \in |\widetilde{\alpha}|$ for $1 \leq i \leq N_0$. By [Ku, Theorem 1.I.5.46]

$$\Gamma(|\widetilde{\alpha}|, |\gamma_{2,m}|, D') > \bigcup_{i=1}^{N_0} \Gamma(S(z_i, \varepsilon_0/4), S(z_i, \varepsilon_0/2), A(z_i, \varepsilon_0/4, \varepsilon_0/2)).$$
(3.9)

Fix $\gamma \in \Gamma_m^1$, $\gamma : [0,1] \to D$, $\gamma(0) \in |\widetilde{\alpha}|, \gamma(1) \in |\gamma_{2,m}|$. It follows from (3.9) that $f(\gamma)$ has a subpath $f(\gamma)_1 := f(\gamma)|_{[p_1,p_2]}$ such that

$$f(\gamma)_1 \in \Gamma(S(z_i, \varepsilon_0/4), S(z_i, \varepsilon_0/2), A(z_i, \varepsilon_0/4, \varepsilon_0/2))$$

for some $1 \leq i \leq N_0$. Then $\gamma|_{[p_1,p_2]}$ is a subpath of γ and belongs to $\Gamma_f(z_i, \varepsilon_0/4, \varepsilon_0/2)$, because

$$f(\gamma|_{[p_1,p_2]}) = f(\gamma)|_{[p_1,p_2]} \in \Gamma(S(z_i,\varepsilon_0/4), S(z_i,\varepsilon_0/2), A(z_i,\varepsilon_0/4,\varepsilon_0/2)).$$

Thus

$$\Gamma_m^1 > \bigcup_{i=1}^{N_0} \Gamma_f(z_i, \varepsilon_0/4, \varepsilon_0/2) \,. \tag{3.10}$$

Put

$$\eta(t) = \begin{cases} 4/\varepsilon_0, & t \in [\varepsilon_0/4, \varepsilon_0/2], \\ 0, & t \notin [\varepsilon_0/4, \varepsilon_0/2] \end{cases}$$

Observe that the function η satisfies the relation (3.2). Then, by the definition of f in (3.1), by the relation (3.10) and due to the subadditivity of the modulus of families of paths (see [Va, Theorem 6.2]), we obtain that

$$M(\Gamma_m^1) \leqslant \sum_{i=1}^{N_0} M(\Gamma_f(z_i, \varepsilon_0/4, \varepsilon_0/2)) \leqslant \sum_{i=1}^{N_0} \frac{N_0 4^n \|Q\|_1}{\varepsilon_0^n}, \qquad m \geqslant M_0, \qquad (3.11)$$

where $||Q||_1 = \int_{D'} Q(x) dm(x)$. In addition, by [Ku, Theorem 1.I.5.46], $\Gamma_m^2 > \Gamma_f(\overline{x_0}, r_{k_0+1}, r_{k_0})$. Arguing similarly as above, we put

$$\eta(t) = \begin{cases} 1/(r_{k_0} - r_{k_0+1}), & t \in [r_{k_0+1}, r_{k_0}], \\ 0, & t \notin [r_{k_0+1}, r_{k_0}] \end{cases}$$

Now, by the last relation we obtain that

$$M(\Gamma_m^2) \leqslant \frac{\|Q\|_1}{(r_{k_0} - r_{k_0+1})^n}, \quad m \geqslant M_0.$$
 (3.12)

Thus, by (3.8), (3.11) and (3.12), due to the subadditivity of the modulus of families of paths (see [Va, Theorem 6.2]), we obtain that

$$M(\Gamma_m) \leq \left(\frac{N_0 4^n}{\varepsilon_0^n} + \frac{1}{(r_{k_0} - r_{k_0+1})^n}\right) \|Q\|_1, \quad m \geq M_0.$$

The right part of the last relation does not depend on m, so we may put

$$N := \left(\frac{N_0 4^n}{\varepsilon_0^n} + \frac{1}{(r_{k_0} - r_{k_0+1})^n}\right) \|Q\|_1$$

Lemma 3.2 is proved. \Box

Given $\delta > 0, M > 0$ domains $D, D' \subset \mathbb{R}^n$, $n \ge 2$, and a continuum $A \subset D'$ denote by $\mathfrak{S}_{\delta,A,M}(D,D')$ the family of all open discrete and closed mappings f of D onto D' such that the condition (3.1) holds for any $y_0 \in D'$ and such that $h(f^{-1}(A), \partial D) \ge \delta$ and $\|Q_f\|_{L^1(D')} \le M$. The following theorem with certain differences in its formulation is given in [Sev₄, Theorem 2], and its proof is completely similar to proof of this statement. However, for completeness, we present it in full in the text.

Theorem 3.2. Let D be a domain with a weakly flat boundary, and let D' be a regular domain. Now, any $f \in \mathfrak{S}_{\delta,A,M}(D,D')$ has a continuous extension $\overline{f}: \overline{D} \to \overline{D'}_P$, wherein

 $\overline{f}(\overline{D}) = \overline{D'}_P$ and, in addition, the family $\mathfrak{S}_{\delta,A,M}(\overline{D},\overline{D'})$ of all extended mappings $\overline{f}: \overline{D} \to \overline{D'}_P$ is equicontinuous in \overline{D} .

Proof. The possibility of continuous extension of the mapping $f \in \mathfrak{S}_{\delta,A,M}(D,D')$ to the boundary of D follows by Theorem 3.1 and Corollary 3.1. The equicontinuity of $\mathfrak{S}_{\delta,A,M}(D,D')$ at inner points of D is proved in [SSD, Theorem 1.1].

Let us to show the equicontinuity of $\mathfrak{S}_{\delta,A,\mathrm{M}}(\overline{D},\overline{D'})$ on ∂D . Assume the contrary. Now, there is a point $z_0 \in \partial D$, a number $\varepsilon_0 > 0$, a sequence $z_m \in \overline{D}$ and a mapping $\overline{f}_m \in \mathfrak{S}_{\delta,A,M}(\overline{D},\overline{D'})$ such that $z_m \to z_0$ as $m \to \infty$ and

$$\rho(\overline{f}_m(z_m), \overline{f}_m(z_0)) \geqslant \varepsilon_0, \quad m = 1, 2, \dots,$$
(3.13)

where ρ is some of possible metrics in $\overline{D'}_P$ defined in (1.4). Since $f_m = \overline{f}_m|_D$ has a continuous extension to \overline{D} , we may assume that $z_m \in D$ and, in addition, there is one more sequence $z'_m \in D, z'_m \to z_0$ as $m \to \infty$ such that $\rho(f_m(z'_m), \overline{f}_m(z_0)) \to 0$ as $m \to \infty$. In this case, it follows from (3.13) that

$$\rho(f_m(z_m), f_m(z'_m)) \ge \varepsilon_0/2, \quad m \ge m_0.$$

Since D' is regular, the space $\overline{D'}_P$ is compact. Thus, we may assume that $f_m(z_m)$ and $f_m(z'_m)$ converge to some $P_1, P_2 \in \overline{D'}_P$, $P_1 \neq P_2$, as $m \to \infty$. Let d_m and g_m be sequences of decreasing domains corresponding to prime ends P_1 and P_2 , respectively. Due to [IS, Lemma 3.1], cf. [KR₂, Lemma 1], we may assume that the sequence of cuts σ_m which corresponds to $d_m, m = 1, 2, \ldots$, belongs to spheres $S(\overline{x_0}, r_m)$, where $\overline{x_0} \in \partial D'$ and $r_m \to 0$ as $m \to \infty$. Put $x_0, y_0 \in A$ such that $x_0 \neq y_0$ and $x_0 \neq P_1 \neq y_0$, where the continuum $A \subset D'$ is taken from the conditions of Theorem 3.2. Without loss of generality, we may assume that $d_1 \cap g_1 = \emptyset$ and $x_0, y_0 \notin d_1 \cup g_1$.

By Lemmas 3.1 and 3.2, we may find disjoint paths $\gamma_{1,m} : [0,1] \to D'$ and $\gamma_{2,m} : [0,1] \to D'$ and a number N > 0 such that $\gamma_{1,m}(0) = x_0$, $\gamma_{1,m}(1) = f_m(z_m)$, $\gamma_{2,m}(0) = y_0$, $\gamma_{2,m}(0) = f_m(z'_m)$, wherein

$$M(\Gamma_m) \leqslant N, \quad m \geqslant M_0,$$
 (3.14)

where Γ_m consists of those and only those paths $\gamma \ge D$ for which $f_m(\gamma) \in \Gamma(|\gamma_{1,m}|, |\gamma_{2,m}|, D')$ (see Figure 4). On the other hand, let $\gamma_{1,m}^*$ and $\gamma_{2,m}^*$ be the total f_m -lifting of the paths $\gamma_{1,m}$ and $\gamma_{2,m}$ starting at the points z_m and z'_m , respectively (such lifting exist by [Vu, Lemma 3.7]). Now, $\gamma_{1,m}^*(1) \in f_m^{-1}(A)$ and $\gamma_{2,m}^*(1) \in f_m^{-1}(A)$. Since by the condition $h(f_m^{-1}(A), \partial D) > \delta > 0$, $m = 1, 2, \ldots$, we obtain that

$$h(|\gamma_{1,m}^{*}|) \ge h(z_{m}, \gamma_{1,m}^{*}(1)) \ge (1/2) \cdot h(f_{m}^{-1}(A), \partial D) > \delta/2,$$

$$h(|\gamma_{2,m}^{*}|) \ge h(z'_{m}, \gamma_{2,m}^{*}(1)) \ge (1/2) \cdot h(f_{m}^{-1}(A), \partial D) > \delta/2$$
(3.15)

for sufficiently large $m \in \mathbb{N}$. Choose the ball $U := B_h(z_0, r_0) = \{z \in \mathbb{R}^n : h(z, z_0) < r_0\}$, where $r_0 > 0$ and $r_0 < \delta/4$. Observe that $|\gamma_{1,m}^*| \cap U \neq \emptyset \neq |\gamma_{1,m}^*| \cap (D \setminus U)$ for sufficiently




Figure 4: To proof of Theorem 3.2.

large $m \in \mathbb{N}$, because $h(f_m(|\gamma_{1,m}|)) \ge \delta/2$ and $z_m \in |\gamma_{1,m}^*|$, $z_m \to z_0$ as $m \to \infty$. Arguing similarly, we may conclude that $|\gamma_{2,m}^*| \cap U \ne \emptyset \ne |\gamma_{2,m}^*| \cap (D \setminus U)$. Since $|\gamma_{1,m}^*|$ and $|\gamma_{2,m}^*|$ are continua, by [Ku, Theorem 1.I.5.46]

$$|\gamma_{1,m}^*| \cap \partial U \neq \emptyset, \quad |\gamma_{2,m}^*| \cap \partial U \neq \emptyset.$$
(3.16)

Put P := N > 0, where N is a number from the relation (3.14). Since D has a weakly flat boundary, we may find a neighborhood $V \subset U$ of z_0 such that

$$M(\Gamma(E, F, D)) > N \tag{3.17}$$

for any continua $E, F \subset D$ with $E \cap \partial U \neq \emptyset \neq E \cap \partial V$ and $F \cap \partial U \neq \emptyset \neq F \cap \partial V$. Observe that

$$|\gamma_{1,m}^*| \cap \partial V \neq \emptyset, \quad |\gamma_{2,m}^*| \cap \partial V \neq \emptyset.$$
(3.18)

for sufficiently large $m \in \mathbb{N}$. Indeed, $z_m \in |\gamma_{1,m}^*|$ and $z'_m \in |\gamma_{2,m}^*|$, where $z_m, z'_m \to z_0 \in V$ as $m \to \infty$. Thus, $|\gamma_{1,m}^*| \cap V \neq \emptyset \neq |\gamma_{2,m}^*| \cap V$ for sufficiently large $m \in \mathbb{N}$. Besides that, $h(V) \leq h(U) = 2r_0 < \delta/2$ and $|\gamma_{1,m}^*| \cap (D \setminus V) \neq \emptyset$ because $h(|\gamma_{1,m}^*|) > \delta/2$ by (3.15). Then $|\gamma_{1,m}^*| \cap \partial V \neq \emptyset$ (see [Ku, Theorem 1.I.5.46]). Similarly, $h(V) \leq h(U) = 2r_0 < \delta/2$. Now, since by (3.15) $h(|\gamma_{2,m}^*|) > \delta/2$, we obtain that $|\gamma_{2,m}^*| \cap (D \setminus V) \neq \emptyset$. By [Ku, Theorem 1.I.5.46], $|\gamma_{1,m}^*| \cap \partial V \neq \emptyset$. Thus, (3.18) is proved. By (3.17), (3.16) and (3.18), we obtain that

$$M(\Gamma(|\gamma_{1,m}^*|, |\gamma_{2,m}^*|, D)) > N.$$
(3.19)

The inequality (3.19) contradicts to (3.14), since $\Gamma(|\gamma_{1,m}^*|, |\gamma_{2,m}^*|, D) \subset \Gamma_m$ and thus

$$M(\Gamma(|\gamma_{1,m}^*|, |\gamma_{2,m}^*|, D)) \leqslant M(\Gamma_m) \leqslant N$$

The obtained contradiction indicates the incorrectness of the assumption in (3.13). The theorem is proved. \Box

One of the versions of the following statement is established in $[SevSkv_1, item v, Lemma 2]$ for homeomorphisms and "good" boundaries, see also $[SevSkv_2, Lemma 4.1]$. Let us also point out the case relating to mappings with branching and bad boundaries, see [SSD, Lemma 6.1],

as well as the case of bad boundaries and homeomorphisms, see [SSI, Lemma 2.13]. We are interested in the "most general" case when the mapping is only open and discrete, and the mapped domain is regular.

Lemma 3.3. Let $n \ge 2$, and let D and D' be domains in \mathbb{R}^n . Assume that D has a weakly flat boundary, none of the components of which degenerates into a point, and D' is regular. Let A be a non-degenerate continuum in D' and $\delta > 0$. Assume that f_m is a sequence of open discrete and closed mappings of D onto D' satisfying the following condition: for any $m = 1, 2, \ldots$ there is a continuum $A_m \subset D$, $m = 1, 2, \ldots$, such that $f_m(A_m) = A$ and $h(A_m) \ge \delta > 0$. If there is $0 < M_1 < \infty$ such that f_m satisfies (3.1) at any $y_0 \in D'$ and $m = 1, 2, \ldots$ with some $Q = Q_m(y)$ for which $||Q_m||_{L^1(D')} \le M_1$, then there exists $\delta_1 > 0$ such that

$$h(A_m, \partial D) > \delta_1 > 0 \quad \forall \ m \in \mathbb{N}.$$

Proof. Due to the compactness of the space $\overline{\mathbb{R}^n}$ the boundary of the domain D is not empty and is compact, so that the distance $\overline{\mathbb{R}^n}h(A_m, \partial D)$ is well-defined.

We will prove from the opposite. Suppose that the conclusion of the lemma is not true. Then for each $k \in \mathbb{N}$ there is a number $m = m_k$ such that $h(A_{m_k}, \partial D) < 1/k$. We may assume that the sequence m_k is increasing by k. Since A_{m_k} is compact, there are $x_k \in A_{m_k}$ i $y_k \in \partial D$ such that $h(A_{m_k}, \partial D) = h(x_k, y_k) < 1/k$ (see Figure 5). Since ∂D is a compact set,




Figure 5: To proof of Lemma 3.3

we may consider that $y_k \to y_0 \in \partial D$ as $k \to \infty$. Now we also have that $x_k \to y_0 \in \partial D$ as $k \to \infty$. Let K_0 be a component of ∂D containing y_0 . Obviously, K_0 is a continuum in $\overline{\mathbb{R}^n}$. Since ∂D is weakly flat, by Theorem 3.1 f_{m_k} has a continuous extension $\overline{f}_{m_k} : \overline{D} \to \overline{D'_P}$. Moreover, the mapping \overline{f}_{m_k} is uniformly continuous in \overline{D} for any fixed k, because \overline{f}_{m_k} is continuous on the compact set \overline{D} . Now, for any $\varepsilon > 0$ there is $\delta_k = \delta_k(\varepsilon) < 1/k$ such that

$$\rho(\overline{f}_{m_k}(x), \overline{f}_{m_k}(x_0)) < \varepsilon \tag{3.20}$$

 $\forall x, x_0 \in \overline{D}, \quad h(x, x_0) < \delta_k, \quad \delta_k < 1/k,$

where ρ is one of the possible metric in $\overline{D'}_P$,

$$\rho(x,y) := |g^{-1}(x) - g^{-1}(y)|, \qquad (3.21)$$

where the element $g^{-1}(x)$ is understood as some (single) point of the boundary D_0 for $x \in E_{D'}$, which is well-defined due to [IS, Theorem 2.1]; see also [Na₂, теорема 4.1]. Let $\varepsilon > 0$ be some number such that

$$\varepsilon < (1/2) \cdot \operatorname{dist} \left(\partial D_0, g^{-1}(A) \right), \tag{3.22}$$

where A is a continuum from the conditions of lemma and $g: D_0 \to D'$ is a quasiconformal mapping of D_0 onto D, while D is a domain with a quasiconformal boundary corresponding to the definition of the metric ρ . Put $B_h(x_0, r) = \{x \in \mathbb{R}^n : h(x, x_0) < r\}$. Given $k \in \mathbb{N}$, we set

$$B_k := \bigcup_{x_0 \in K_0} B_h(x_0, \delta_k), \quad k \in \mathbb{N}.$$

Since B_k is a neighborhood of a continuum K_0 , by [HK, Lemma 2.2] there is a neighborhood U_k of K_0 such that $U_k \subset B_k$ and $U_k \cap D$ is connected. We may consider that U_k is open, so that $U_k \cap D$ is linearly path connected (see [MRSY, Proposition 13.1]). Let $h(K_0) = m_0$. Then we may find $z_0, w_0 \in K_0$ such that $h(K_0) = h(z_0, w_0) = m_0$. Thus, there are sequences $\overline{y_k} \in U_k \cap D$, $z_k \in U_k \cap D$ and $w_k \in U_k \cap D$ such that $z_k \to z_0$, $\overline{y_k} \to y_0$ and $w_k \to w_0$ as $k \to \infty$. We may consider that

$$h(z_k, w_k) > m_0/2 \quad \forall \ k \in \mathbb{N} \,. \tag{3.23}$$

Since the set $U_k \cap D$ is linearly connected, we may joint the points z_k , $\overline{y_k}$ and w_k using some path $\gamma_k \in U_k \cap D$. As usually, we denote by $|\gamma_k|$ the locus of the path γ_k in D. Then $f_{m_k}(|\gamma_k|)$ is a compact set in D'. If $x \in |\gamma_k|$, then we may find $x_0 \in K_0$ such that $x \in B(x_0, \delta_k)$. Fix $\omega \in A \subset D$. Since $x \in |\gamma_k|$ and, in addition, x is an inner point of D, we may use the notation $f_{m_k}(x)$ instead $\overline{f}_{m_k}(x)$. By (3.20) and (3.22), and by the triangle inequality, we obtain that

$$\rho(f_{m_k}(x),\omega) \ge \rho(\omega,\overline{f}_{m_k}(x_0)) - \rho(\overline{f}_{m_k}(x_0),f_{m_k}(x)) \ge$$

 $\geq \operatorname{dist}(\partial D_0, g^{-1}(A)) - (1/2) \cdot \operatorname{dist}(\partial D_0, g^{-1}(A)) = (1/2) \cdot \operatorname{dist}(\partial D_0, g^{-1}(A)) > \varepsilon \quad (3.24)$

for sufficiently large $k \in \mathbb{N}$. Passing to inf in (3.24) over all $x \in |\gamma_k|$ and $\omega \in A$, we obtain that

$$\rho(f_{m_k}(|\gamma_k|), A) > \varepsilon, \quad k = 1, 2, \dots$$
(3.25)

We now show that there is $\varepsilon_1 > 0$ such that

dist
$$(f_{m_k}(|\gamma_k|), A) > \varepsilon_1, \quad \forall \ k = 1, 2, \dots$$
 (3.26)

Indeed, let (3.26) be violated. Then for the number $\varepsilon_l = 1/l$, l = 1, 2, ... there are $\xi_l \in |\gamma_{k_l}|$ and $\zeta_l \in A$ such that

$$|f_{m_{k_l}}(\xi_l) - \zeta_l| < 1/l, \quad l = 1, 2, \dots$$
 (3.27)

Without loss of generality, we may assume that the sequence of numbers k_l , l = 1, 2, ..., is increasing. Since A is a compact set, we may consider that the sequence ζ_l converge to some $\zeta_0 \in A$ as $l \to \infty$. By the triangle inequality and from (3.27) it follows that

$$|f_{m_{k_l}}(\xi_l) - \zeta_0| \to 0, \quad l \to \infty.$$
(3.28)

On the other hand, recall that $\rho(f_{m_k}(x), \omega) = |g^{-1}(f_{m_k}(x)) - g^{-1}(\omega)|$, where $g: D_0 \to D'$ is a quasiconformal mapping of D_0 onto D' (see (3.21)). In particular, g^{-1} is a continuous mapping in D'. Thus, by the triangle inequality and (3.28) we obtain that

$$|g^{-1}(f_{m_{k_l}}(\xi_l)) - g^{-1}(\zeta_l)| \leq$$

$$\leq |g^{-1}(f_{m_{k_l}}(\xi_l)) - g^{-1}(\zeta_0)| + |g^{-1}(\zeta_0) - g^{-1}(\zeta_l)| \to 0, \quad l \to \infty.$$
(3.29)

However, by the definition of ρ and by (3.29) we obtain that

$$\rho(f_{m_{k_l}}(|\gamma_{k_l}|), A) \leqslant \rho(f_{m_{k_l}}(\xi_l), \zeta_l) = |g^{-1}(f_{m_{k_l}}(\xi_l)) - g^{-1}(\zeta_l)| \to 0, \quad l \to \infty,$$

that contradicts to (3.25). The resulting contradiction indicates the validity of relation (3.26).

We cover the set A with balls $B(x, \varepsilon/4), x \in A$. Since A is compact, we may assume that $A \subset \bigcup_{i=1}^{M_0} B(x_i, \varepsilon/4), x_i \in A, i = 1, 2, ..., M_0, 1 \leq M_0 < \infty$. By the definition, M_0 depends only on A, in particular, M_0 does not depend on k. Put

$$\Gamma_k := \Gamma(A_{m_k}, |\gamma_k|, D) \,.$$

Let $\Gamma_{ki} := \Gamma_{f_{m_k}}(x_i, \varepsilon/4, \varepsilon/2)$, in other words, Γ_{ki} consists of all paths $\gamma : [0, 1] \to D$ such that $f_{m_k}(\gamma(0)) \in S(x_i, \varepsilon/4)$, $f_{m_k}(\gamma(1)) \in S(x_i, \varepsilon/2)$ i $\gamma(t) \in A(x_i, \varepsilon/4, \varepsilon/2)$ for 0 < t < 1. We show that

$$\Gamma_k > \bigcup_{i=1}^{M_0} \Gamma_{ki} \,. \tag{3.30}$$

Indeed, let $\tilde{\gamma} \in \Gamma_k$, in other words, $\tilde{\gamma} : [0,1] \to D$, $\tilde{\gamma}(0) \in A_{m_k}$, $\tilde{\gamma}(1) \in |\gamma_k|$ and $\tilde{\gamma}(t) \in D$ for $0 \leq t \leq 1$. Then $\gamma^* := f_{m_k}(\tilde{\gamma}) \in \Gamma(A, f_{m_k}(|\gamma_k|), D')$. Since the balls $B(x_i, \varepsilon/4), 1 \leq i \leq M_0$, form the coverage of the compact set A, we may find $i \in \mathbb{N}$ such that $\gamma^*(0) \in B(x_i, \varepsilon/4)$ and $\gamma^*(1) \in f_{m_k}(|\gamma_k|)$. By the relation (3.26), $|\gamma^*| \cap B(x_i, \varepsilon/4) \neq \emptyset \neq |\gamma^*| \cap (D' \setminus B(x_i, \varepsilon/4))$. Thus, by [Ku, Theorem 1.I.5.46] there is $0 < t_1 < 1$ such that $\gamma^*(t_1) \in S(x_i, \varepsilon/4)$. We may assume that $\gamma^*(t) \notin B(x_i, \varepsilon/4)$ for $t > t_1$. Set $\gamma_1 := \gamma^*|_{[t_1, 1]}$. By (3.26) it follows that $|\gamma_1| \cap B(x_i, \varepsilon/2) \neq \emptyset \neq |\gamma_1| \cap (D \setminus B(x_i, \varepsilon/2))$. Thus, by [Ku, Theorem 1.I.5.46] there is $t_1 < t_2 < 1$ such that $\gamma^*(t_2) \in S(x_i, \varepsilon/2)$. We may assume that $\gamma^*(t) \in B(x_i, \varepsilon/2)$ for any $t < t_2$. Putting $\gamma_2 := \gamma^*|_{[t_1, t_2]}$, we observe that a path γ_2 is a subpath of γ^* , which belongs to $\Gamma(S(x_i, \varepsilon/4), S(x_i, \varepsilon/2), A(x_i, \varepsilon/4, \varepsilon/2))$.

Finally, $\widetilde{\gamma}$ has a subpath $\widetilde{\gamma_2} := \widetilde{\gamma}|_{[t_1, t_2]}$ such that $f_{m_k} \circ \widetilde{\gamma_2} = \gamma_2$, while

$$\gamma_2 \in \Gamma(S(x_i, \varepsilon/4), S(x_i, \varepsilon/2), A(x_i, \varepsilon/4, \varepsilon/2)).$$

Thus, the relation (3.30) is proved. Set

$$\eta(t) = \begin{cases} 4/\varepsilon, & t \in [\varepsilon/4, \varepsilon/2], \\ 0, & t \notin [\varepsilon/4, \varepsilon/2]. \end{cases}$$

Observe that η satisfies the relation (3.2) for $r_1 = \varepsilon/4$ and $r_2 = \varepsilon/2$. Since f_{m_k} satisfies the relation (3.1), we obtain that

$$M(\Gamma_{f_{m_k}}(x_i, \varepsilon/4, \varepsilon/2)) \leqslant (4/\varepsilon)^n \cdot \|Q\|_1 < M_0 < \infty.$$
(3.31)

By (3.30) and (3.31) and due to the subadditivity of the modulus of families of paths, we obtain that

$$M(\Gamma_k) \leqslant \frac{4^n M_0}{\varepsilon^n} \int_{D'} Q(y) \, dm(y) \leqslant M_1 \cdot M_0 < \infty \,. \tag{3.32}$$

Arguing similarly to the proof of relations (3.15) and using the condition (3.23), we obtain that $M(\Gamma_k) \to \infty$ as $k \to \infty$, which contradicts to (3.32). The resulting contradiction proves the lemma.

Given domains $D, D' \subset \mathbb{R}^n$, points $a \in D$, $b \in D'$ and a number $M_0 > 0$ denote by $\mathfrak{S}_{a,b,M_0}(D,D')$ the family of open discrete and closed mappings f of D onto D' satisfying the relation (3.1) for some $Q = Q_f$, $||Q||_{L^1(D')} \leq M_0$ for any $y_0 \in f(D)$, such that f(a) = b. The following statement was proved in [SSD, Theorem 7.1] in the case of a fixed function Q (cf. Theorem 4.2 in [SD]).

Theorem 3.3. Assume that D has a weakly flat boundary, none of the components of which degenerates into a point, and D' is regular. Then any $f \in \mathfrak{S}_{a,b,M_0}(D,D')$ has a continuous extension $\overline{f}: \overline{D} \to \overline{D'}_P$, while $\overline{f}(\overline{D}) = \overline{D'}_P$ and, in addition, the family $\mathfrak{S}_{a,b,M_0}(\overline{D},\overline{D'})$ of all extended mappings $\overline{f}: \overline{D} \to \overline{D'}_P$ is equicontinuous in \overline{D} .

Proof. The possibility of continuous extension of $f \in \mathfrak{S}_{a,b,M_0}(D,D')$ to a continuous mapping $\overline{f}: \overline{D} \to \overline{D'}_P$ is a statement of Theorem 3.1, as well as the equality $\overline{f}(\overline{D}) = \overline{D'}_P$. The equicontinuity of $\mathfrak{S}_{a,b,M_0}(D,D')$ at inner points of D is a result of [SSD, Theorem 1.1].

It remains to establish the equicontinuity of the family of extended mappings $\overline{f}: \overline{D} \to \overline{D'}_P$ at the boundary points of the domain D.

We prove this statement from the opposite. Assume that the family $\mathfrak{S}_{a,b,M_0}(\overline{D},\overline{D'})$ is not equicontinuous at some point $x_0 \in \partial D$. Then there are points $x_m \in D$ and mappings $f_m \in \mathfrak{S}_{a,b,M_0}(\overline{D},\overline{D'}), m = 1, 2, \ldots$, such that $x_m \to x_0$ as $m \to \infty$, moreover,

$$\rho(f_m(x_m), f_m(x_0)) \geqslant \varepsilon_0, \quad m = 1, 2, \dots$$
(3.33)

for some $\varepsilon_0 > 0$, where ρ is one of the metrics in $\overline{D'}_P$ (see, e.g., [KR₁, Remark 2]). We choose in an arbitrary way the point $y_0 \in D'$, $y_0 \neq b$, and join it to the point b by some path in D', which we denote by α . Let $A := |\alpha|$ and let A_m be a total f_m -lifting of α starting at a (it exists by [Vu, Lemma 3.7]). Observe that $h(A_m, \partial D) > 0$ due to the closeness of f_m . Now, the following two cases are possible: either $h(A_m) \to 0$ as $m \to \infty$, or $h(A_{m_k}) \geq \delta_0 > 0$ as $k \to \infty$ for some increasing sequence of numbers m_k and some $\delta_0 > 0$.

In the first of these cases, obviously, $h(A_m, \partial D) \ge \delta > 0$ for some $\delta > 0$. Then, by Theorem 3.2, the family $\{f_m\}_{m=1}^{\infty}$ is equicontinuous at the point x_0 , however, this contradicts the condition (3.33).

In the second case, if $h(f(A_{m_k})) \ge \delta_0 > 0$ for sufficiently large k, we also have that $f(A_{m_k}, \partial D) \ge \delta_1 > 0$ for some $\delta_1 > 0$ by Lemma 3.3. Again, by Theorem 3.2, the family $\{f_{m_k}\}_{k=1}^{\infty}$ is equicontinuous at the point x_0 , and this contradicts the condition (3.33).

Thus, in both of the two possible cases, we came to a contradiction (3.33), and this indicates the incorrect assumption of the absence of the equicontinuity of the family $\mathfrak{S}_{a,b,M_0}(D,D')$ in \overline{D} . The theorem is proved. \Box

4 Compactness of families of solutions of the Dirichlet problem

Proof of Theorem 1.1. In general, we will use the scheme of proving Theorem 1.2 in [SD].

I. Let $f_m \in \mathfrak{F}^{\mathcal{M}}_{\varphi,\Phi,z_0}(D)$, $m = 1, 2, \ldots$ By Stoilow's factorization theorem (see, e.g., [St, 5(III).V]) a mapping f_m has a representation

$$f_m = \varphi_m \circ g_m \,, \tag{4.1}$$

where g_m is some homeomorphism, and φ_m is some analytic function. By Lemma 1 in [Sev₂], the mapping g_m belongs to the Sobolev class $W_{\text{loc}}^{1,1}(D)$ and has a finite distortion. Moreover, by [A, (1).C, Ch. I]

$$f_{m_z} = \varphi_{m_z}(g_m(z))g_{m_z}, \qquad f_{m_{\overline{z}}} = \varphi_{m_z}(g_m(z))g_{m_{\overline{z}}}$$
(4.2)

for almost all $z \in D$. Therefore, by the relation (4.2), $J(z, g_m) \neq 0$ for almost all $z \in D$, in addition, $K_{\mu_{f_m}}(z) = K_{\mu_{g_m}}(z)$.

II. We prove that $\partial g_m(D)$ contains at least two points. Suppose the contrary. Then either $g_m(D) = \mathbb{C}$, or $g_m(D) = \mathbb{C} \setminus \{a\}$, where $a \in \mathbb{C}$. Consider first the case $g_m(D) = \mathbb{C}$. By Picard's theorem $\varphi_m(g_m(D))$ is the whole plane, except perhaps one point $\omega_0 \in \mathbb{C}$. On the other hand, for every $m = 1, 2, \ldots$ the function $u_m(z) := \operatorname{Re} f_m(z) = \operatorname{Re} (\varphi_m(g_m(z)))$ is continuous on the compact set \overline{D} under the condition (1.4) by the continuity of φ . Therefore, there exists $C_m > 0$ such that $|\operatorname{Re} f_m(z)| \leq C_m$ for any $z \in D$, but this contradicts the fact that $\varphi_m(g_m(D))$ contains all points of the complex plane except, perhaps, one. The situation $g_m(D) = \mathbb{C} \setminus \{a\}, a \in \mathbb{C}$, is also impossible, since the domain $g_m(D)$ must be simply connected in \mathbb{C} as a homeomorphic image of the simply connected domain D.

Therefore, the boundary of the domain $g_m(D)$ contains at least two points. Then, according to Riemann's mapping theorem, we may transform the domain $g_m(D)$ onto the unit disk \mathbb{D} using the conformal mapping ψ_m . Let $z_0 \in D$ be a point from the condition of the theorem. By using an auxiliary conformal mapping

$$\widetilde{\psi_m}(z) = \frac{z - (\psi_m \circ g_m)(z_0)}{1 - z(\psi_m \circ g_m)(z_0)}$$

of the unit disk onto itself we may consider that $(\psi_m \circ g_m)(z_0) = 0$. Now, by (4.1) we obtain that

$$f_m = \varphi_m \circ g_m = \varphi_m \circ \psi_m^{-1} \circ \psi_m \circ g_m = F_m \circ G_m, \quad m = 1, 2, \dots$$

where $F_m := \varphi_m \circ \psi_m^{-1}$, $F_m : \mathbb{D} \to \mathbb{C}$, and $G_m = \psi_m \circ g_m$. Obviously, a function F_m is analytic, and G_m is a regular Sobolev homeomorphism in D. In particular, $\operatorname{Im} F_m(0) = 0$ for any $m \in \mathbb{N}$.

III. We prove that the L^1 -norms of the functions $K_{\mu_{G_m}}(z)$ are bounded from above by some universal positive constant C > 0 over all $m = 1, 2, \ldots$ Indeed, by the convexity of the function Φ in (1.5) and by [Bou, Proposition 5, I.4.3], the slope $[\Phi(t) - \Phi(0)]/t$ is a non-decreasing function. Hence there exist constants $t_0 > 0$ and $C_1 > 0$ such that

$$\Phi(t) \ge C_1 \cdot t \qquad \forall \ t \in [t_0, \infty) \,. \tag{4.3}$$

Fix $m \in \mathbb{N}$. By (1.5) and (4.3), we obtain that

$$\int_{D} K_{\mu_{G_m}}(z) \, dm(z) = \int_{\{z \in D: K_{\mu_{G_m}}(z) < t_0\}} K_{\mu_{G_m}}(z) \, dm(z) + \int_{\{z \in D: K_{\mu_{G_m}}(z) \ge t_0\}} K_{\mu_{G_m}}(z) \, dm(z) \leqslant \\ \leqslant t_0 \cdot m(D) + \frac{1}{C_1} \int_{D} \Phi(K_{\mu_{G_m}}(z)) \, dm(z) \leqslant \\ \leqslant t_0 \cdot m(D) + \frac{\sup(1+|z|^2)^2}{C_1} \int_{D} \Phi(K_{\mu_{G_m}}(z)) \cdot \frac{1}{(1+|z|^2)^2} \, dm(z) \leqslant \\ \leqslant t_0 \cdot m(D) + \frac{\sup(1+|z|^2)^2}{C_1} \mathcal{M}(D) < \infty \,,$$

because $\mathcal{M}(D) < \infty$ by the assumption of the theorem.

IV. We prove that each map G_m , m = 1, 2, ..., has a continuous extension to E_D , in addition, the family of extended maps \overline{G}_m , m = 1, 2, ..., is equicontinuous in \overline{D}_P . Indeed, as proved in item **III**, $K_{\mu_{G_m}} \in L^1(D)$. By [KPRS, Theorem 3] (see also [LSS, Theorem 3.1])

each G_m , m = 1, 2, ..., is a ring *Q*-homeomorphism in \overline{D} for $Q = K_{\mu G_m}(z)$, where μ is defined in (1.3), and K_{μ} my be calculated by the formula (1.1). Note that the unit disk \mathbb{D} is a uniform domain as a finitely connected flat domain at its boundary with a finite number of boundary components (see, for example, [Na₁, Theorem 6.2 and Corollary 6.8]). Then it is desirable the conclusion is a statement of Theorem 2.1.

V. Let us prove that the inverse homeomorphisms G_m^{-1} , $m = 1, 2, \ldots$, have a continuous extension \overline{G}_m^{-1} to $\partial \mathbb{D}$ in terms of prime ends in D, and $\{\overline{G}_m^{-1}\}_{m=1}^{\infty}$ is equicontinuous in $\overline{\mathbb{D}}$ as a family of mappings from $\overline{\mathbb{D}}$ to \overline{D}_P . Since by the item IV mappings G_m , $m = 1, 2, \ldots$, are ring $K_{\mu_{G_m}}(z)$ -homeomorphisms in D, the corresponding inverse mappings G_m^{-1} satisfy (3.1) (in this case, D corresponds the unit disk \mathbb{D} in (3.2), $f \mapsto G_m$, $Q \mapsto K_{\mu_{G_m}}(z)$, and $f(D) \mapsto D$). Since $G_m^{-1}(0) = z_0$ for any $m = 1, 2, \ldots$, the possibility of a continuous extension of G_m^{-1} to $\partial \mathbb{D}$, and the equicontinuity of $\{\overline{G}_m^{-1}\}_{m=1}^{\infty}$ in terms of $G_m^{-1}: \overline{\mathbb{D}} \to \overline{D}_P$ follows by Theorem 3.3.

VI. Since, as proved above the family $\{G_m\}_{m=1}^{\infty}$ is equicontinuous in D, according to Arzela-Ascoli criterion there exists an increasing subsequence of numbers m_k , $k = 1, 2, \ldots$, such that G_{m_k} converges locally uniformly in D to some continuous mapping $G: D \to \overline{\mathbb{C}}$ as $k \to \infty$ (see, e.g., [Va, Theorem 20.4]). By [SD, Lemma 2.1], either G is a homeomorphism with values in \mathbb{R}^n , or a constant in $\overline{\mathbb{R}^n}$. Let us prove that the second case is impossible. Let us apply the approach used in proof of the second part of Theorem 21.9 in [Va]. Suppose the contrary: let $G_{m_k}(x) \to c = const$ as $k \to \infty$. Since $G_{m_k}(z_0) = 0$ for all $k = 1, 2, \ldots$, we have that c = 0. By item \mathbf{V} , the family of mappings G_m^{-1} , $m = 1, 2, \ldots$, is equicontinuous in \mathbb{D} . Then

$$h(z, G_{m_k}^{-1}(0)) = h(G_{m_k}^{-1}(G_{m_k}(z)), G_{m_k}^{-1}(0)) \to 0$$

as $k \to \infty$, which is impossible because z is an arbitrary point of the domain D. The obtained contradiction refutes the assumption made above. Thus, $G: D \to \mathbb{C}$ is a homeomorphism.

VII. According to **V**, the family of mappings $\{\overline{G}_m^{-1}\}_{m=1}^{\infty}$ is equicontinuous in $\overline{\mathbb{D}}$. By the Arzela-Ascoli criterion (see, e.g., [Va, Theorem 20.4]) we may consider that $\overline{G}_{m_k}^{-1}(y)$, $k = 1, 2, \ldots$, converges to some mapping $\widetilde{F} : \overline{\mathbb{D}} \to \overline{D}$ as $k \to \infty$ uniformly in \overline{D} . Let us to prove that $\widetilde{F} = \overline{G}^{-1}$. For this purpose, we show that $G(D) = \mathbb{D}$. Fix $y \in \mathbb{D}$. Since $G_{m_k}(D) = \mathbb{D}$ for every $k = 1, 2, \ldots$, we obtain that $G_{m_k}(x_k) = y$ for some $x_k \in D$. Since Dis regular, the metric space (\overline{D}_P, ρ) is compact. Thus, we may assume that $\rho(x_k, x_0) \to 0$ as $k \to \infty$, where $x_0 \in \overline{D}_P$. By the triangle inequality and the equicontinuity of $\{\overline{G}_m\}_{m=1}^{\infty}$ onto \overline{D}_P , see **IV**, we obtain that

$$|\overline{G}(x_0) - y| = |\overline{G}(x_0) - \overline{G}_{m_k}(x_k)| \leq |\overline{G}(x_0) - \overline{G}_{m_k}(x_0)| + |\overline{G}_{m_k}(x_0) - \overline{G}_{m_k}(x_k)| \to 0$$

as $k \to \infty$. Thus, $\overline{G}(x_0) = y$. Observe that $x_0 \in D$, because G is a homeomorphism. Since $y \in \mathbb{D}$ is arbitrary, the equality $G(D) = \mathbb{D}$ is proved. In this case, $G_{m_k}^{-1} \to G^{-1}$ locally uniformly in \mathbb{D} as $k \to \infty$ (see, e.g., [RSS, Lemma 3.1]). Thus, $\widetilde{F}(y) = G^{-1}(y)$ for every $y \in \mathbb{D}$.

Finally, since $\widetilde{F}(y) = G^{-1}(y)$ for any $y \in \mathbb{D}$ and, in addition, \widetilde{F} has a continuous extension on $\partial \mathbb{D}$, due to the uniqueness of the limit at the boundary points we obtain that $\widetilde{F}(y) = \overline{G}^{-1}(y)$ for $y \in \overline{\mathbb{D}}$. Therefore, we have proved that $\overline{G}_{m_k}^{-1} \to \overline{G}^{-1}$ uniformly in $\overline{\mathbb{D}}$ with as $k \to \infty$ with respect to the metrics ρ in \overline{D}_P .

VIII. By **VII**, for $y = e^{i\theta} \in \partial \mathbb{D}$

$$\operatorname{Re} F_{m_k}(e^{i\theta}) = \varphi\left(\overline{G}_{m_k}^{-1}(e^{i\theta})\right) \to \varphi\left(\overline{G}^{-1}(e^{i\theta})\right)$$

$$(4.4)$$

as $k \to \infty$ uniformly on $\theta \in [0, 2\pi)$. Since by the construction $\text{Im } F_{m_k}(0) = 0$ for any $k = 1, 2, \ldots$, by the Schwartz formula (see, e.g., [GK, section 8.III.3]) the analytic function F_{m_k} is uniquely restored by its real part, namely,

$$F_{m_k}(y) = \frac{1}{2\pi i} \int\limits_{S(0,1)} \varphi\left(\overline{G}_{m_k}^{-1}(t)\right) \frac{t+y}{t-y} \cdot \frac{dt}{t} \,. \tag{4.5}$$

Set

$$F(y) := \frac{1}{2\pi i} \int_{S(0,1)} \varphi\left(\overline{G}^{-1}(t)\right) \frac{t+y}{t-y} \cdot \frac{dt}{t}.$$
(4.6)

Let $K \subset \mathbb{D}$ be an arbitrary compact set, and let $y \in K$. By (4.5) and (2.9) we obtain that

$$|F_{m_k}(y) - F(y)| \leq \frac{1}{2\pi} \int_{S(0,1)} \left| \varphi(\overline{G}_{m_k}^{-1}(t)) - \varphi(\overline{G}^{-1}(t)) \right| \left| \frac{t+y}{t-y} \right| |dt|.$$

$$(4.7)$$

Since K is compact, there is $0 < R_0 = R_0(K) < \infty$ such that $K \subset B(0, R_0)$. By the triangle inequality $|t + y| \leq 1 + R_0$ and $|t - y| \geq |t| - |y| \geq 1 - R_0$ for $y \in K$ and any $t \in \mathbb{S}^1$. Thus

$$\left|\frac{t+y}{t-y}\right| \leqslant \frac{1+R_0}{1-R_0} := M = M(K) \,.$$

Put $\varepsilon > 0$. By (4.4), for a number $\varepsilon' := \frac{\varepsilon}{M}$ there is $N = N(\varepsilon, K) \in \mathbb{N}$ such that $\left| \varphi \left(\overline{G}_{m_k}^{-1}(t) \right) - \varphi \left(\overline{G}^{-1}(t) \right) \right| < \varepsilon'$ for any $k \ge N(\varepsilon)$ and $t \in \mathbb{S}^1$. Now, by (4.7)

$$|F_{m_k}(y) - F(y)| < \varepsilon \quad \forall \ k \ge N.$$
(4.8)

It follows from (4.8) that the sequence F_{m_k} converges to F as $k \to \infty$ in the unit disk locally uniformly. In particular, we obtain that Im F(0) = 0. Note that F is analytic function in \mathbb{D} (see remarks made at the end of item 8.III in [GK]), and

$$\operatorname{Re} F(re^{i\psi}) = \frac{1}{2\pi} \int_{0}^{2\pi} \varphi\left(\overline{G}^{-1}(e^{i\theta})\right) \frac{1-r^2}{1-2r\cos(\theta-\psi)+r^2} d\theta$$

for $z = re^{i\psi}$. By [GK, Theorem 2.10.III.3]

$$\lim_{\zeta \to z} \operatorname{Re} F(\zeta) = \varphi(\overline{G}^{-1}(z)) \quad \forall \ z \in \partial \mathbb{D}.$$
(4.9)

Observe that F either is a constant or open and discrete (see, e.g., [St, Ch. V, I.6 and II.5]). Thus, $f_{m_k} = F_{m_k} \circ G_{m_k}$ converges to $f = F \circ G$ locally uniformly as $k \to \infty$, where $f = F \circ G$ either is a constant or open and discrete. Moreover, by (4.9)

$$\lim_{\zeta \to P} \operatorname{Re} f(\zeta) = \lim_{\zeta \to P} \operatorname{Re} F(G(\zeta)) = \varphi(G^{-1}(G(P))) = \varphi(P) \,.$$

IX. Since by VI G is a homeomorphism, by [L, Lemma 1 and Theorem 1] G is a regular solution of the equation (1.3) for some function $\mu : \mathbb{C} \to \mathbb{D}$. Since the set of points of the function F, where its Jacobian is zero, consist only of isolated points (see [St, Ch. V, 5.II and 6.II]), f is regular solution of the Dirichlet problem (1.3)–(1.4) whenever $F \not\equiv const$. Note that the relation (1.5) holds for the corresponding function $K_{\mu} = K_{\mu f}$ (see e.g. [L, Lemma 1]). Therefore, $f \in \mathfrak{F}_{\varphi,\Phi,z_0}^{\mathcal{M}}(D)$. \Box

References

- [A] AHLFORS, L.V.: Lectures on Quasiconformal Mappings. Van Nostrand, Toronto, 1966.
- [Bou] BOURBAKI, N.: Functions of Real Variable. Springer, Berlin, 2004.
- [Dyb] DYBOV, YU.P.: Compactness of classes of solutions of the Dirichlet problem for the Beltrami equations // Proc. Inst. Appl. Math. and Mech. of NAS of Ukraine 19, 2009, 81-89 (in Russian).
- [Fu] FUGLEDE, B.: Extremal length and functional completion. Acta Math. 98, 1957, 171–219.
- [GSS₁] GOLBERG, A., R. SALIMOV AND E. SEVOST'YANOV: Singularities of discrete open mappings with controlled *p*-module. - J. Anal. Math. 127, 2015, 303–328.
- [GSS₂] GOLBERG A., R. SALIMOV AND E. SEVOST'YANOV: Normal Families of Discrete Open Mappings with Controlled *p*-Module. - Contemporary Mathematics 667, 2016, 83–103.
- [GK] HURWITZ, A. AND R. COURANT: The Function Theory. Nauka, Moscow, 1968 (in Russian).
- [IS] ILYUTKO, D.P. AND E.A. SEVOST'YANOV: On prime ends on Riemannian manifolds. J. Math. Sci. 241:1, 2019, 47–63.
- [HK] HERRON, J. AND P. KOSKELA: Quasiextremal distance domains and conformal mappings onto circle domains. - Compl. Var. Theor. Appl. 15, 1990, 167–179.
- [KPRS] KOVTONYUK, D.A., I.V. PETKOV, V.I. RYAZANOV, R.R. SALIMOV: The boundary behavior and the Dirichlet problem for the Beltrami equations. - St. Petersburg Math. J. 25:4, 2014, 587-603.
- [KR₁] KOVTONYUK, D.A., V.I. RYAZANOV: On the theory of prime ends for space mappings. -Ukrainian Math. J. 67:4, 2015, 528-541.
- [KR₂] KOVTONYUK, D.A., V.I. RYAZANOV: Prime ends and Orlicz-Sobolev classes. St. Petersburg Math. J. 27:5, 2016, 765–788.

- [Kr] KRUGLIKOV, V.I.: Capacity of condensers and spatial mappings quasiconformal in the mean (in Russian). - Mat. Sb. (N.S.) 130(172):2(6), 1986, 185–206.
- [Ku] KURATOWSKI, K.: Topology, v. 2. Academic Press, New York–London, 1968.
- [LSS] LOMAKO, T., R. SALIMOV R. AND E. SEVOST'YANOV: On equicontinuity of solutions to the Beltrami equations. - Ann. Univ. Bucharest (math. series) V. LIX:2, 2010, 261-271.
- [L] LOMAKO, T.: On the theory of convergence and compactness for Beltrami equations. -Ukrainian Mathematical Journal 63:3, 2011, 393–402.
- [MRV] MARTIO, O., S. RICKMAN, AND J. VÄISÄLÄ: Definitions for quasiregular mappings. Ann. Acad. Sci. Fenn. Ser. A1 448, 1969, 1–40.
- [MRSY] MARTIO, O., V. RYAZANOV, U. SREBRO, AND E. YAKUBOV: Moduli in modern mapping theory. - Springer Science + Business Media, LLC, New York, 2009.
- [Ma] MAZ'YA, V.: Lectures on isoperimetric and isocapacitary inequalities in the theory of Sobolev spaces. - Contemp. Math. 338, 2003, 307—340.
- [Na₁] NÄKKI, R.: Extension of Loewner's capacity theorem. Trans. Amer. Math. Soc. 180, 1973, 229–236.
- [Na₂] NÄKKI, R.: Prime ends and quasiconformal mappings. J. Anal. Math. 35, 1979, 13–40.
- [NP] NÄKKI, R. AND B. PALKA: Uniform equicontinuity of quasiconformal mappings. Proc. Amer. Math. Soc. 37:2, 1973, 427–433.
- [P] PETKOV, I.V.: Dirichlet problem for Beltrami equation in simply connected domains. Dopovidi NAN Ukrainy 11, 2015, 12–17 (in Russian).
- [Ri] RICKMAN, S.: Quasiregular mappings. Results in Mathematic and Related Areas (3), 26, Springer-Verlag, Berlin, 1993.
- [RSS] RYAZANOV, V., R. SALIMOV AND E. SEVOST'YANOV: On Convergence Analysis of Space Homeomorphisms. - Siberian Advances in Mathematics 23:4, 2013, 263-293.
- [RS] RYAZANOV, V., E. SEVOST'YANOV: Equicontinuity of mappings quasiconformal in the mean.Ann. Acad. Sci. Fenn. 36, 2011, 231-244.
- [Sa] SAKS, S.: Theory of the Integral. Dover, New York, 1964.
- [SalSev] SALIMOV, R.R. AND E.A. SEVOST'YANOV: Analogs of the Ikoma-Schwartz lemma and Liouville theorem for mappings with unbounded characteristic. - Ukrainian Math. J. 63:10, 2012, 1551–1565.
- [Sev1] SEVOST'YANOV, E.A.: Equicontinuity of homeomorphisms with unbounded characteristic.
 Siberian Advances in Mathematics 23:2, 2013, 106–122.

- [Sev₂] SEVOST'YANOV, E.A.: Analog of the Montel Theorem for Mappings of the Sobolev Class with Finite Distortion. - Ukrainian Math. J. 67:6, 2015, 938–947.
- [Sev₃] SEVOST'YANOV E.A.: Boundary behavior and equicontinuity for families of mappings in terms of prime ends. - St. Petersburg Math. J. 30:6, 2019, 973–1005.
- [Sev₄] SEVOST'YANOV, E.A.: Boundary extension of mappings with the inverse Poletsky inequality by prime ends. https://arxiv.org/abs/2004.04800.
- [Sev₅] SEVOST'YANOV, E.A.: On global behavior of mappings with integral constraints. https://arxiv.org/abs/2101.08859
- [SD] SEVOST'YANOV, E.A., O.P. DOVHOPIATYI: On compact classes of solutions of the Dirichlet problem with integral restrictions. - https://arxiv.org/abs/2009.12705.
- [SevSkv₁] SEVOST'YANOV, E.A., S.A. SKVORTSOV: On the equicontinuity of families of mappings in the case of variable domains. - Ukrainian Mathematical Journal 71:7, 2019, 1071–1086.
- [SevSkv₂] SEVOST'YANOV, E.A., S.A. SKVORTSOV: On mappings whose inverse satisfy the Poletsky inequality. - Ann. Acad. Scie. Fenn. Math. 45, 2020, 259-277.
- [SevSkv₃] SEVOST'YANOV, E.A., S.A. SKVORTSOV: On equicontinuity of families of mappings with one normalization condition. - Mat. Zametki 109:4, 2021, 597–607 (in Russian).
- [SSD] SEVOST'YANOV, E.A., S.A. SKVORTSOV AND O.P. DOVHOPIATYI: On non-homeomorphic mappings with inverse Poletsky inequality. - Journal of Mathematical Sciences 252:4, 2021, 541–557.
- [SSI] SEVOST'YANOV, E.A., S.A. SKVORTSOV AND N.S. ILKEVYCH: On the global behavior of inverse mappings in terms of prime ends. Ann. Acad. Sci. Fenn. Math. (accepted for print).
- [St] STOILOW, S.: Principes Topologiques de la Théorie des Fonctions Analytiques. Gauthier-Villars, Paris, 1956.
- [Va] VÄISÄLÄ, J.: Lectures on n-dimensional quasiconformal mappings. Lecture Notes in Math. 229, Springer-Verlag, Berlin etc., 1971.
- [Vu] VUORINEN, M.: Exceptional sets and boundary behavior of quasiregular mappings in n-space.
 Ann. Acad. Sci. Fenn. Ser. A 1. Math. Dissertationes, 11, 1976, 1–44.

Oleksandr Dovhopiatyi

 Zhytomyr Ivan Franko State University,
 Bol'shaya Berdichevskaya Str., 10 008 Zhytomyr, UKRAINE alexdov1111111@gmail.com

Evgeny Sevost'yanov

1. Zhytomyr Ivan Franko State University,

40 Bol'shaya Berdichevskaya Str., 10 008 Zhytomyr, UKRAINE

2. Institute of Applied Mathematics and Mechanics

of NAS of Ukraine,

1 Dobrovol'skogo Str., 84 100 Slavyansk, UKRAINE

esevos tyanov 2009 @gmail.com