# **Binomial Fibonacci Power Sums**

Kunle Adegoke Department of Physics and Engineering Physics Obafemi Awolowo University 220005 Ile-Ife, Nigeria adegoke00@gmail.com

2010 Mathematics Subject Classification: Primary 11B39; Secondary 11B37. Keywords: Fibonacci number, Lucas number, summation identity, series, binomial coefficient.

#### Abstract

We evaluate various binomial sums involving the powers of Fibonacci and Lucas numbers.

# 1 Introduction

Our main goal in this paper is to evaluate the following sums of powers of Fibonacci and Lucas numbers involving the binomial coefficients:

$$\sum_{k=0}^{n} (\pm 1)^{k} \binom{n}{k} F_{j(rk+s)}^{2m}, \qquad \sum_{k=0}^{n} (\pm 1)^{k} \binom{n}{k} L_{j(rk+s)}^{2m},$$
$$\sum_{k=0}^{n} (\pm 1)^{k} \binom{n}{k} F_{j(2rk+s)}^{2m+1}, \qquad \sum_{k=0}^{n} (\pm 1)^{k} \binom{n}{k} L_{j(2rk+s)}^{2m+1};$$

thereby extending the work of Wessner [11], Hoggatt and Bicknell [3, 4], Long [9], Kiliç et al [6] and several previous researchers. Here n is any non-negative integer, j, m, r and s are any integers and  $F_t$  and  $L_t$  are the Fibonacci and Lucas numbers.

There is a dearth of binomial cubic Fibonacci and Lucas identities in existing literature. We will show that, for any non-negative integer n and any integer s,

$$\sum_{k=0}^{n} \binom{n}{k} F_{k+s}^{3} = \frac{1}{5} (2^{n} F_{2n+3s} + 3F_{n-s}),$$
$$\sum_{k=0}^{n} \binom{n}{k} L_{k+s}^{3} = 2^{n} L_{2n+3s} + 3L_{n-s},$$

$$\begin{split} \sum_{k=0}^{n} \binom{n}{k} (-1)^{k} F_{k+s}^{3} &= \frac{1}{5} ((-1)^{n} 2^{n} F_{n+3s} - (-1)^{s} 3F_{2n+s}), \\ \sum_{k=0}^{n} (-1)^{k} \binom{n}{k} L_{k+s}^{3} &= (-1)^{n} 2^{n} L_{n+3s} + (-1)^{s} 3L_{2n+s}, \\ \sum_{k=0}^{n} \binom{n}{k} 2^{k} F_{k+s}^{3} &= \begin{cases} 5^{n/2-1} (F_{3n+3s} - (-1)^{s} 3F_{s}), & n \text{ even}; \\ 5^{(n-3)/2} (L_{3n+3s} + (-1)^{s} 3L_{s}) & n \text{ odd}, \end{cases} \\ \sum_{k=0}^{n} \binom{n}{k} 2^{k} L_{k+s}^{3} &= \begin{cases} 5^{n/2} (L_{3n+3s} + (-1)^{s} 3L_{s}), & n \text{ even}; \\ 5^{(n+1)/2} (F_{3n+3s} - (-1)^{s} 3F_{s}), & n \text{ odd}. \end{cases} \end{split}$$

We will also derive the following binomial summation identities which we believe are new:

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} F_{2jr+p}^{n-k} F_{p}^{k} F_{j(rk+s)}^{2} = \frac{1}{5} (F_{2jr}^{n} L_{pn-2js} - (-1)^{js} 2F_{jr}^{n} L_{jr+p}^{n}),$$

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} F_{2jr+p}^{n-k} F_{p}^{k} L_{j(rk+s)}^{2} = F_{2jr}^{n} L_{pn-2js} + (-1)^{js} 2F_{jr}^{n} L_{jr+p}^{n},$$

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} L_{2jr+p}^{n-k} L_{p}^{k} F_{j(rk+s)}^{2} = \begin{cases} 5^{n/2-1} F_{2jr}^{n} L_{pn-2js} - (-1)^{js} 5^{n-1} 2F_{jr}^{n} F_{jr+p}^{n}, & n \text{ even}; \\ 5^{(n-1)/2} F_{2jr}^{n} F_{pn-2js} - (-1)^{js} 5^{n-1} 2F_{jr}^{n} F_{jr+p}^{n}, & n \text{ odd}, \end{cases}$$

and

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} L_{2jr+p}^{n-k} L_{p}^{k} L_{j(rk+s)}^{2} = \begin{cases} 5^{n/2} F_{2jr}^{n} L_{pn-2js} + (-1)^{js} 5^{n} 2F_{jr}^{n} F_{jr+p}^{n}, & n \text{ even}; \\ 5^{(n+1)/2} F_{2jr}^{n} F_{pn-2js} + (-1)^{js} 5^{n} 2F_{jr}^{n} F_{jr+p}^{n}, & n \text{ odd}. \end{cases}$$

The Fibonacci numbers,  $F_n$ , and the Lucas numbers,  $L_n$ , are defined, for  $n \in \mathbb{Z}$ , through the recurrence relations

$$F_n = F_{n-1} + F_{n-2}, (n \ge 2), \quad F_0 = 0, F_1 = 1;$$
 (1)

and

$$L_n = L_{n-1} + L_{n-2}, (n \ge 2), \quad L_0 = 2, L_1 = 1;$$
 (2)

with

$$F_{-n} = (-1)^{n-1} F_n, \quad L_{-n} = (-1)^n L_n.$$
 (3)

Throughout this paper, we denote the golden ratio,  $(1 + \sqrt{5})/2$ , by  $\alpha$  and write  $\beta = (1 - \sqrt{5})/2 = -1/\alpha$ , so that  $\alpha\beta = -1$  and  $\alpha + \beta = 1$ .

Explicit formulas (Binet formulas) for the Fibonacci and Lucas numbers are

$$F_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \quad L_n = \alpha^n + \beta^n, \quad n \in \mathbb{Z}.$$
 (4)

Koshy [7] and Vajda [10] have written excellent books dealing with Fibonacci and Lucas numbers.

Our results emanate from the following general Fibonacci and Lucas summation identities (Lemma 2):

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^{k} F_{j(rk+s)}^{m} = \frac{1}{(\sqrt{5})^{m}} \sum_{i=0}^{m} (-1)^{i(js+1)} \binom{m}{i} \alpha^{(m-2i)js} \left(x + (-1)^{ijr} \alpha^{(m-2i)jr} z\right)^{n},$$
(BF)

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^{k} L_{j(rk+s)}^{m} = \sum_{i=0}^{m} (-1)^{ijs} \binom{m}{i} \alpha^{(m-2i)js} \left( x + (-1)^{ijr} \alpha^{(m-2i)jr} z \right)^{n}.$$
(BL)

For low m, the identities (BF) and (BL) are more useful in the equivalent form

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^{k} F_{j(rk+s)}^{m} = \frac{1}{(\sqrt{5})^{m}} \sum_{i=0}^{m} (-1)^{i} \binom{m}{i} \beta^{ijs} \alpha^{(m-i)js} \left(x + \beta^{ijr} \alpha^{(m-i)jr} z\right)^{n}, \quad (BF')$$

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^k L^m_{j(rk+s)} = \sum_{i=0}^{m} \binom{m}{i} \beta^{ijs} \alpha^{(m-i)js} \left(x + \beta^{ijr} \alpha^{(m-i)jr} z\right)^n.$$
(BL')

When m = 1, we have the weighted linear binomial identities:

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^k F_{j(rk+s)} = \frac{1}{\sqrt{5}} \left( \alpha^{js} (x + \alpha^{jr} z)^n - \beta^{js} (x + \beta^{jr} z)^n \right),$$
(F1)

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^k L_{j(rk+s)} = \alpha^{js} (x + \alpha^{jr} z)^n + \beta^{js} (x + \beta^{jr} z)^n, \tag{L1}$$

which are valid for n a non-negative integer, j, r, s integers and real or complex x and z. Most linear binomial Fibonacci identities can be obtained from identities (F1) and (L1) by substituting appropriate choices of n, j, r, s, x and z. For example, if we write 2r for r and set  $x = (-1)^{jr}$ , z = 1 in (F1) and (L1), we obtain

$$\sum_{k=0}^{n} (-1)^{jrk} \binom{n}{k} F_{j(2rk+s)} = (-1)^{jrn} L_{jr}^{n} F_{j(rn+s)},$$
(5)

$$\sum_{k=0}^{n} (-1)^{jrk} \binom{n}{k} L_{j(2rk+s)} = (-1)^{jrn} L_{jr}^{n} L_{j(rn+s)};$$
(6)

which are valid for n a non-negative integer and integers r, s and j. The special case (s = 0) of identity (6) was also derived by Layman [8]. As another example of linear binomial Fibonacci identities that may be derived from (F1) and (L1), write 2r for r and set  $x = (-1)^{jr}$ , z = -1. This gives

$$\sum_{k=0}^{n} (-1)^{(jr+1)k} \binom{n}{k} F_{j(2rk+s)} = \begin{cases} 5^{n/2} F_{jr}^{n} F_{j(rn+s)}, & n \text{ even}; \\ (-1)^{jr+1} 5^{(n-1)/2} F_{jr}^{n} L_{j(rn+s)}, & n \text{ odd}; \end{cases}$$
(7)

$$\sum_{k=0}^{n} (-1)^{(jr+1)k} \binom{n}{k} L_{j(2rk+s)} = \begin{cases} 5^{n/2} F_{jr}^{n} L_{j(rn+s)}, & n \text{ even}; \\ (-1)^{jr+1} 5^{(n+1)/2} F_{jr}^{n} F_{j(rn+s)}, & n \text{ odd.} \end{cases}$$
(8)

Setting  $(x = F_{p+jr}, z = -F_p)$  and also  $(x = L_{p+jr}, z = -L_p)$  and making use of the identities of Hoggat et al, (see Lemma 3), where p is any integer, we find

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} F_{p+jr}^{n-k} F_{p}^{k} F_{j(rk+s)} = (-1)^{js+1} F_{jr}^{n} F_{pn-js};$$
(9)

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} F_{p+jr}^{n-k} F_{p}^{k} L_{j(rk+s)} = (-1)^{js+1} F_{jr}^{n} L_{pn-js};$$
(10)

and

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} L_{p+jr}^{n-k} L_{p}^{k} F_{j(rk+s)} = \begin{cases} (-1)^{js+1} 5^{\frac{n}{2}} F_{jr}^{n} F_{pn-js}, & n \text{ even}; \\ (-1)^{js+1} 5^{\frac{n-1}{2}} F_{jr}^{n} L_{pn-js}, & n \text{ odd}; \end{cases}$$

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} L_{p+jr}^{n-k} L_{p}^{k} L_{j(rk+s)} = \begin{cases} (-1)^{js} 5^{\frac{n}{2}} F_{jr}^{n} L_{pn-js}, & n \text{ even}; \\ (-1)^{js} 5^{\frac{n+1}{2}} F_{jr}^{n} F_{pn-js}, & n \text{ odd}. \end{cases}$$
(11)

Identities (9), (10) were obtained by Carlitz [1] while these and (11) and (12) may be found in Dresel [2]. The special case (s = 0) of (9) was also derived by Layman [8].

### 2 Required identities and preliminary results

**Lemma 1.** For real or complex z, let a given well-behaved function h(z) have, in its domain, the representation  $h(z) = \sum_{k=c_1}^{c_2} g(k) z^{f(k)}$  where f(k) and g(k) are given real sequences and  $c_1, c_2 \in [-\infty, \infty]$ . Let j be an integer. Then,

$$\sum_{k=c_1}^{c_2} g(k) z^{f(k)} F_{jf(k)}^m = \frac{1}{(\sqrt{5})^m} \sum_{i=0}^m (-1)^i \binom{m}{i} h\left(\beta^{ij} \alpha^{(m-i)j} z\right),\tag{F}$$

$$\sum_{k=c_1}^{c_2} g(k) z^{f(k)} L^m_{jf(k)} = \sum_{i=0}^m \binom{m}{i} h\left(\beta^{ij} \alpha^{(m-i)j} z\right).$$
(L)

*Proof.* We have

$$\begin{split} \sum_{k=c_1}^{c_2} g(k) z^{f(k)} F_{jf(k)}^m &= \sum_{k=c_1}^{c_2} g(k) z^{f(k)} \frac{\left(\alpha^{jf(k)} - \beta^{jf(k)}\right)^m}{(\sqrt{5})^m} \\ &= \frac{1}{(\sqrt{5})^m} \sum_{k=c_1}^{c_2} g(k) z^{f(k)} \sum_{i=0}^m (-1)^i \binom{m}{i} \beta^{ijf(k)} \alpha^{(m-i)jf(k)} \\ &= \frac{1}{(\sqrt{5})^m} \sum_{i=0}^m (-1)^i \binom{m}{i} \sum_{k=c_1}^{c_2} g(k) \left(\beta^{ij} \alpha^{(m-i)j} z\right)^{f(k)} \\ &= \frac{1}{(\sqrt{5})^m} \sum_{i=0}^m (-1)^i \binom{m}{i} h \left(\beta^{ij} \alpha^{(m-i)j} z\right). \end{split}$$

The proof of (L) is similar.

Since  $\beta^i \alpha^{m-i} = (-1)^i \alpha^{m-2i}$ , identities (F) and (L) can also be written as

$$\sum_{k=c_1}^{c_2} g(k) z^{f(k)} F_{jf(k)}^m = \frac{1}{(\sqrt{5})^m} \sum_{i=0}^m (-1)^i \binom{m}{i} h\left((-1)^{ij} \alpha^{(m-2i)j} z\right), \tag{F'}$$

$$\sum_{k=c_1}^{c_2} g(k) z^{f(k)} L^m_{jf(k)} = \sum_{i=0}^m \binom{m}{i} h\left( (-1)^{ij} \alpha^{(m-2i)j} z \right).$$
(L')

**Lemma 2.** For non-negative integers m and n, integers j, r and s and real or complex x and z,

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^k F_{j(rk+s)}^m = \frac{1}{(\sqrt{5})^m} \sum_{i=0}^{m} (-1)^{i(js+1)} \binom{m}{i} \alpha^{(m-2i)js} \left(x + (-1)^{ijr} \alpha^{(m-2i)jr} z\right)^n,$$
(BF)

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^{k} L_{j(rk+s)}^{m} = \sum_{i=0}^{m} (-1)^{ijs} \binom{m}{i} \alpha^{(m-2i)js} \left( x + (-1)^{ijr} \alpha^{(m-2i)jr} z \right)^{n}.$$
(BL)

*Proof.* Consider the binomial identity

$$h(z) = \sum_{k=0}^{n} g(k) z^{f(k)} = z^{s} (x + z^{r})^{n},$$
(13)

where

$$f(k) = rk + s, \quad g(k) = \binom{n}{k} x^{n-k}.$$
(14)

Thus,

$$h\left((-1)^{ij}\alpha^{(m-2i)j}z\right) = (-1)^{ijs}\alpha^{(m-2i)js}z^s(x+(-1)^{ijr}\alpha^{(m-2i)jr}z^r)^n.$$
 (15)

Use of (14) and (15) in identity  $(\mathbf{F}')$ , with  $c_1 = 0$ ,  $c_2 = n$  gives

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^{rk} F_{j(rk+s)}^{m} = \frac{1}{(\sqrt{5})^{m}} \sum_{i=0}^{m} (-1)^{i(js+1)} \binom{m}{i} \alpha^{(m-2i)js} \left(x + (-1)^{ijr} \alpha^{(m-2i)jr} z^{r}\right)^{n},$$

from which identity (BF) follows when we write  $z^{1/r}$  for z. To prove (BL), use (14) and (15) in identity (L').

It is sometimes convenient to use the ( $\alpha$  vs  $\beta$ ) version of identities (BF) and (BL):

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^k F_{j(rk+s)}^m = \frac{1}{(\sqrt{5})^m} \sum_{i=0}^{m} (-1)^i \binom{m}{i} \beta^{ijs} \alpha^{(m-i)js} \left(x + \beta^{ijr} \alpha^{(m-i)jr} z\right)^n, \quad (BF')$$

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^{k} L_{j(rk+s)}^{m} = \sum_{i=0}^{m} \binom{m}{i} \beta^{ijs} \alpha^{(m-i)js} \left( x + \beta^{ijr} \alpha^{(m-i)jr} z \right)^{n}.$$
(BL')

Lemma 3 (Hoggatt et al [5]). For p and q integers,

$$L_{p+q} - L_p \alpha^q = -\beta^p F_q \sqrt{5},\tag{16}$$

$$L_{p+q} - L_p \beta^q = \alpha^p F_q \sqrt{5},\tag{17}$$

$$F_{p+q} - F_p \alpha^q = \beta^p F_q, \tag{18}$$

$$F_{p+q} - F_p \beta^q = \alpha^p F_q. \tag{19}$$

Quadratic binomial Fibonacci identities may be obtained from m = 2 in (BF') and (BL'):

$$5\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^{k} F_{j(rk+s)}^{2} = \alpha^{2js} (x + \alpha^{2jr} z)^{n} + \beta^{2js} (x + \beta^{2jr} z)^{n} - 2(-1)^{js} (x + (-1)^{jr} z)^{n},$$
(F2)

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^k L_{j(rk+s)}^2 = \alpha^{2js} (x + \alpha^{2jr} z)^n + \beta^{2js} (x + \beta^{2jr} z)^n + 2(-1)^{js} (x + (-1)^{jr} z)^n.$$
(L2)

**Theorem 1.** For non-negative integer n and integers j, r, s, p,

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} F_{2jr+p}^{n-k} F_{p}^{k} F_{j(rk+s)}^{2} = \frac{1}{5} (F_{2jr}^{n} L_{pn-2js} - (-1)^{js} 2F_{jr}^{n} L_{jr+p}^{n}), \quad p \neq 0,$$
(20)

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} F_{2jr+p}^{n-k} F_{p}^{k} L_{j(rk+s)}^{2} = F_{2jr}^{n} L_{pn-2js} + (-1)^{js} 2F_{jr}^{n} L_{jr+p}^{n}, \quad p \neq 0, \qquad (21)$$

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} L_{2jr+p}^{n-k} L_{p}^{k} F_{j(rk+s)}^{2} = \begin{cases} 5^{n/2-1} F_{2jr}^{n} L_{pn-2js} - (-1)^{js} 5^{n-1} 2F_{jr}^{n} F_{jr+p}^{n}, & n \text{ even}; \\ 5^{(n-1)/2} F_{2jr}^{n} F_{pn-2js} - (-1)^{js} 5^{n-1} 2F_{jr}^{n} F_{jr+p}^{n}, & n \text{ odd}, \end{cases}$$

$$(22)$$

and

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} L_{2jr+p}^{n-k} L_{p}^{k} L_{j(rk+s)}^{2} = \begin{cases} 5^{n/2} F_{2jr}^{n} L_{pn-2js} + (-1)^{js} 5^{n} 2F_{jr}^{n} F_{jr+p}^{n}, & n \text{ even}; \\ 5^{(n+1)/2} F_{2jr}^{n} F_{pn-2js} + (-1)^{js} 5^{n} 2F_{jr}^{n} F_{jr+p}^{n}, & n \text{ odd.} \end{cases}$$

$$(23)$$

 $\textit{Proof.}\,$  Choose  $x=F_{2jr+p},\,z=-F_p$  in (F2), noting Lemma 3, to obtain

$$5\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} F_{2jr+p}^{n-k} F_{p}^{k} F_{j(rk+s)}^{2} = F_{2jr}^{n} (\alpha^{2js} \beta^{pn} + \alpha^{pn} \beta^{2js}) - 2(-1)^{js} (F_{2jr+p} - (-1)^{jr} F_{p})^{n},$$

from which identity (20) follows. The same (x, z) choice in (L2) produces identity (21).

Set  $x = L_{2jr+p}$ ,  $z = -L_p$  in (F2), utilizing Lemma 3. This gives

$$5\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} L_{2jr+p}^{n-k} L_{p}^{k} F_{j(rk+s)}^{2} = F_{2jr}^{n} (\sqrt{5})^{n} (\alpha^{pn-2js} + (-1)^{n} \beta^{pn-2js}) - 2(-1)^{js} (L_{2jr+p} - (-1)^{jr} L_{p})^{n};$$

and hence identity (22). The same (x, z) choice in (L2) produces identity (23).

Cubic binomial Fibonacci identities may be obtained from m = 3 in (BF') and (BL'):

$$5\sqrt{5}\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^{k} F_{j(rk+s)}^{3} = \alpha^{3js} (x + \alpha^{3jr} z)^{n} - \beta^{3js} (x + \beta^{3jr} z)^{n} - (-1)^{js} 3\alpha^{js} (x + (-1)^{jr} \alpha^{jr} z)^{n} + (-1)^{js} 3\beta^{js} (x + (-1)^{jr} \beta^{jr} z)^{n},$$
(F3)

$$\sum_{k=0}^{n} \binom{n}{k} x^{n-k} z^{k} L_{j(rk+s)}^{3} = \alpha^{3js} (x + \alpha^{3jr} z)^{n} + \beta^{3js} (x + \beta^{3jr} z)^{n} + (-1)^{js} 3\alpha^{js} (x + (-1)^{jr} \alpha^{jr} z)^{n} + (-1)^{js} 3\beta^{js} (x + (-1)^{jr} \beta^{jr} z)^{n}.$$
(L3)

**Theorem 2.** For non-negative integer n and any integer s,

$$\sum_{k=0}^{n} \binom{n}{k} F_{k+s}^{3} = \frac{1}{5} (2^{n} F_{2n+3s} + 3F_{n-s}), \qquad (24)$$

$$\sum_{k=0}^{n} \binom{n}{k} L_{k+s}^{3} = 2^{n} L_{2n+3s} + 3L_{n-s}, \qquad (25)$$

$$\sum_{k=0}^{n} \binom{n}{k} (-1)^{k} F_{k+s}^{3} = \frac{1}{5} ((-1)^{n} 2^{n} F_{n+3s} - (-1)^{s} 3F_{2n+s}),$$
(26)

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} L_{k+s}^{3} = (-1)^{n} 2^{n} L_{n+3s} + (-1)^{s} 3 L_{2n+s},$$
(27)

$$\sum_{k=0}^{n} \binom{n}{k} 2^{k} F_{k+s}^{3} = \begin{cases} 5^{n/2-1} (F_{3n+3s} - (-1)^{s} 3F_{s}), & n \text{ even}; \\ 5^{(n-3)/2} (L_{3n+3s} + (-1)^{s} 3L_{s}) & n \text{ odd}, \end{cases}$$
(28)

$$\sum_{k=0}^{n} \binom{n}{k} 2^{k} L_{k+s}^{3} = \begin{cases} 5^{n/2} (L_{3n+3s} + (-1)^{s} 3L_{s}), & n \text{ even}; \\ 5^{(n+1)/2} (F_{3n+3s} - (-1)^{s} 3F_{s}) & n \text{ odd.} \end{cases}$$
(29)

Proof. Set x = 1, z = 1, j = 1, r = 1 in (F3) to obtain

$$5\sqrt{5}\sum_{k=0}^{n} \binom{n}{k} F_{k+s}^{3} = 2^{n}(\alpha^{3s+2n} - \beta^{3s+2n}) + 3(\alpha^{n-s} - \beta^{n-s});$$

and hence identity (24). To prove identity (25), use these (x, z, j, ...) values in (L3). To prove identity (26), set x = 1, z = -1, j = 1, r = 1 in (F3) to get

$$5\sqrt{5}\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} F_{k+s}^{3} = (-1)^{n} 2^{n} (\alpha^{n+3s} - \beta^{n+3s}) - 3(-1)^{s} (\alpha^{2n+s} - \beta^{2n+s}),$$

from which the identity follows. The proof of (27) is similar. Use these values in (L3). The proof of (28) proceeds with the choice j = 1, r = 1, x = 1, z = 2 in (F3), giving

$$5\sqrt{5}\sum_{k=0}^{n} 2^{k} \binom{n}{k} F_{k+s}^{3} = (\sqrt{5})^{n} (\alpha^{3n+3s} - (-1)^{n} \beta^{3n+3s}) - 3(-1)^{n+s} (\sqrt{5})^{n} (\alpha^{s} - (-1)^{n} \beta^{s}),$$

from which the identity follows in accordance with the parity of n. The proof of (29) is similar. Use these (x, z, j, ...) values in (L3).

**Lemma 4.** Let a, b, c and d be rational numbers and  $\lambda$  an irrational number. Then,

$$a + \lambda b = c + \lambda d \iff a = c, \quad b = d$$

**Lemma 5.** If m is an integer and (f(i)) a real sequence, then,

$$\sum_{i=0}^{2m} f(i) = f(m) + \sum_{i=0}^{m-1} \left( f(i) + f(2m-i) \right), \tag{30}$$

$$\sum_{i=0}^{2m+1} f(i) = f(2m+1) - f(m) + \sum_{i=0}^{m} (f(i) + f(2m-i)).$$
(31)

In particular, if f(2m - i) = f(i), then,

$$\sum_{i=0}^{2m} f(i) = f(m) + 2\sum_{i=0}^{m-1} f(i),$$
(32)

$$\sum_{i=0}^{2m+1} f(i) = f(2m+1) - f(m) + 2\sum_{i=0}^{m} f(i).$$
(33)

Lemma 6. For p and q integers,

$$1 + (-1)^{p} \alpha^{2q} = \begin{cases} (-1)^{p} \alpha^{q} F_{q} \sqrt{5}, & p \text{ and } q \text{ have different parity;} \\ (-1)^{p} \alpha^{q} L_{q}, & p \text{ and } q \text{ have the same parity.} \end{cases}$$
(34)

$$1 - (-1)^{p} \alpha^{2q} = \begin{cases} (-1)^{p-1} \alpha^{q} L_{q}, & p \text{ and } q \text{ have different parity;} \\ (-1)^{p-1} \alpha^{q} F_{q} \sqrt{5}, & p \text{ and } q \text{ have the same parity.} \end{cases}$$
(35)

*Proof.* We have

$$(-1)^{p+q} + (-1)^p \alpha^{2q} = \alpha^{p+q} \beta^{p+q} + \alpha^{p+2q} \beta^p$$
  
=  $\alpha^{p+q} \beta^p (\alpha^q + \beta^q)$   
=  $(-1)^p \alpha^q L_q.$  (36)

Similarly,

$$(-1)^{p+q} - (-1)^p \alpha^{2q} = (-1)^{p-1} \alpha^q F_q \sqrt{5}.$$
(37)

Corresponding to (36) and (37) we have

$$(-1)^{p+q} + (-1)^p \beta^{2q} = (-1)^p \beta^q L_q \tag{38}$$

and

$$(-1)^{p+q} - (-1)^p \beta^{2q} = (-1)^p \beta^q F_q \sqrt{5}.$$
(39)

Identities (36), (37), (38) and (39) imply

$$(-1)^q + \alpha^{2q} = \alpha^q L_q,\tag{40}$$

$$(-1)^q - \alpha^{2q} = -\alpha^q F_q \sqrt{5},\tag{41}$$

$$(-1)^q + \beta^{2q} = \beta^q L_q, \tag{42}$$

$$(-1)^q - \beta^{2q} = \beta^q F_q \sqrt{5}.$$
 (43)

# 3 Main results

**Theorem 3.** Let m and n be non-negative integers and let j, r and s be any integers. Then,

$$\begin{split} &\sum_{k=0}^{n} \binom{n}{k} F_{j(rk+s)}^{2m} \\ &= \begin{cases} 5^{-m} \sum_{i=0}^{m-1} (-1)^{i(js+jrn+1)} \binom{2m}{i} L_{(m-i)jr}^{n} L_{(m-i)(jrn+2js)} + (-1)^{m(js+1)} \binom{2m}{m} 5^{-m} 2^{n}, & jmr \ even; \\ 5^{n/2-m} \sum_{i=0}^{m-1} (-1)^{i(s+1)} \binom{2m}{i} F_{(m-i)jr}^{n} L_{(m-i)(jrn+2js)}, & jmr \ odd, \ n \ even; \\ 5^{(n+1)/2-m} \sum_{i=0}^{m-1} (-1)^{is} \binom{2m}{i} F_{(m-i)jr}^{n} F_{(m-i)(jrn+2js)}, & jmr \ odd, \ n \ odd; \end{cases}$$

$$(44)$$

$$\sum_{k=0}^{n} \binom{n}{k} L_{j(rk+s)}^{2m} = \begin{cases} \sum_{i=0}^{m-1} (-1)^{i(js+jrn)} \binom{2m}{i} L_{(m-i)jr}^{n} L_{(m-i)(jrn+2js)} + (-1)^{mjs} \binom{2m}{m} 2^{n}, & jmr \ even; \\ 5^{n/2} \sum_{i=0}^{m-1} (-1)^{is} \binom{2m}{i} F_{(m-i)jr}^{n} L_{(m-i)(jrn+2js)}, & jmr \ odd, \ n \ even; \\ 5^{(n+1)/2} \sum_{i=0}^{m-1} (-1)^{i(s+1)} \binom{2m}{i} F_{(m-i)jr}^{n} F_{(m-i)(jrn+2js)}, & jmr \ odd, \ n \ odd. \end{cases}$$
(45)

*Proof.* In (BF) write 2m for m and set x = 1 and z = 1. This gives

$$5^{m} \sum_{k=0}^{n} \binom{n}{k} F_{j(rk+s)}^{2m} = \sum_{i=0}^{2m} (-1)^{i(js+1)} \binom{2m}{i} \alpha^{(m-i)2js} \left(1 + (-1)^{ijr} \alpha^{(m-i)2jr}\right)^{n}.$$
 (46)

Now, on account of Lemma 6, identity (34), we have

$$1 + (-1)^{ijr} \alpha^{(m-i)2jr} = \begin{cases} (-1)^{ijr} \alpha^{(m-i)jr} L_{(m-i)jr}, & jrm \text{ even};\\ (-1)^i \alpha^{(m-i)jr} F_{(m-i)jr} \sqrt{5}, & jrm \text{ odd.} \end{cases}$$
(47)

Thus, using (47) in (46), we have

$$5^{m} \sum_{k=0}^{n} \binom{n}{k} F_{j(rk+s)}^{2m} = \begin{cases} \sum_{i=0}^{2m} (-1)^{i(js+jrn+1)} \binom{2m}{i} \alpha^{(m-i)(jrn+2js)} L_{(m-i)jr}^{n}, \quad jrm \text{ even};\\ (\sqrt{5})^{n} \sum_{i=0}^{2m} (-1)^{i(js+n+1)} \binom{2m}{i} \alpha^{(m-i)(jrn+2js)} F_{(m-i)jr}^{n}, \quad jrm \text{ odd.} \end{cases}$$

$$\tag{48}$$

Observe that the left side of (48) evaluates to a rational number since it is the finite sum of rational numbers. Since,

$$2\alpha^{(m-i)(jrn+2js)} = L_{(m-i)(jrn+2js)} + F_{(m-i)(jrn+2js)}\sqrt{5},$$
(49)

identity (44) now follows by comparing both sides of identity (48) in each case of jmr even or jmr odd, invoking Lemma 4 with  $\lambda = \sqrt{5}$ . Note the use of Lemma 5, identity (30) to re-write the (i = 0 to 2m) sum. The proof of identity (45) is similar; set x = 1 and z = 1 in (BL) and write 2m for m.

**Theorem 4.** Let m and n be non-negative integers and let j, r and s be any integers. Then,

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} F_{j(rk+s)}^{2m} = \begin{cases} 5^{-m} (-1)^{n} \sum_{i=0}^{m-1} (-1)^{i(s+n+1)} \binom{2m}{i} L_{(m-i)jr}^{n} L_{(m-i)(jrn+2js)} + (-1)^{s+1} \binom{2m}{m} 5^{-m} 2^{n}, & jmr \ odd; \\ 5^{n/2-m} \sum_{i=0}^{m-1} (-1)^{i(js+1)} \binom{2m}{i} F_{(m-i)jr}^{n} L_{(m-i)(jrn+2js)}, & jmr \ even, \ n \ even; \\ -5^{(n+1)/2-m} \sum_{i=0}^{m-1} (-1)^{i(js+jr+1)} \binom{2m}{i} F_{(m-i)jr}^{n} F_{(m-i)(jrn+2js)}, & jmr \ even, \ n \ odd; \end{cases}$$

$$(50)$$

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} L_{j(rk+s)}^{2m} = \begin{cases} (-1)^{n} \sum_{i=0}^{m-1} (-1)^{i(s+n)} \binom{2m}{i} L_{(m-i)jr} L_{(m-i)(jrn+2js)} + (-1)^{s} \binom{2m}{m} 2^{n}, & jmr \ odd; \\ 5^{n/2} \sum_{i=0}^{m-1} (-1)^{ijs} \binom{2m}{i} F_{(m-i)jr}^{n} L_{(m-i)(jrn+2js)}, & jmr \ even, \ n \ even; \\ -5^{(n+1)/2} \sum_{i=0}^{m-1} (-1)^{i(js+jr)} \binom{2m}{i} F_{(m-i)jr}^{n} F_{(m-i)(jrn+2js)}, & jmr \ even, \ n \ odd. \end{cases}$$
(51)

*Proof.* In (BF) write 2m for m and set x = 1 and z = -1. This gives

$$5^{m} \sum_{k=0}^{n} (-1)^{k} \binom{n}{k} F_{j(rk+s)}^{2m} = \sum_{i=0}^{2m} (-1)^{i(js+1)} \binom{2m}{i} \alpha^{(m-i)2js} \left(1 - (-1)^{ijr} \alpha^{(m-i)2jr}\right)^{n}.$$
 (52)

Now, on account of Lemma 6, identity (35), we have

$$1 - (-1)^{ijr} \alpha^{(m-i)2jr} = \begin{cases} (-1)^{ijr-1} \alpha^{(m-i)jr} F_{(m-i)jr} \sqrt{5}, & jrm \text{ even}; \\ (-1)^{i-1} \alpha^{(m-i)jr} L_{(m-i)jr}, & jrm \text{ odd.} \end{cases}$$
(53)

Thus, using (53) in (52), we have

$$5^{m} \sum_{k=0}^{n} (-1)^{k} \binom{n}{k} F_{j(rk+s)}^{2m} = \begin{cases} \sum_{i=0}^{2m} (-1)^{in+is+i-n} \binom{2m}{i} \alpha^{(m-i)(jrn+2js)} L_{(m-i)jr}^{n}, \quad jrm \text{ odd}; \\ (\sqrt{5})^{n} \sum_{i=0}^{2m} (-1)^{ijnr+ijs+i-n} \binom{2m}{i} \alpha^{(m-i)(jrn+2js)} F_{(m-i)jr}^{n}, \quad jrm \text{ even}. \end{cases}$$
(54)

The left side of (54) evaluates to a rational number since it is the finite sum of rational numbers. Making use of identity (49), identity (50) follows by comparing both sides of identity (54) in each case of *jmr* even or *jmr* odd, invoking Lemma 4 with  $\lambda = \sqrt{5}$ . The proof of identity (51) is similar; put x = 1 and z = -1 in (BL) and write 2m for m.

The proofs of Theorems 5 and 6 are similar to those of Theorems 3 and 4. We therefore omit the details and indicate only the appropriate choices of x, z, m and r to be made in identities (BF) and (BL) in each case.

**Theorem 5.** Let m and n be non-negative integers and let j, r and s be any integers. Then,

$$\sum_{k=0}^{n} \binom{n}{k} F_{j(2rk+s)}^{2m+1} = \begin{cases} 5^{-m} \sum_{i=0}^{m} (-1)^{i(js+1)} \binom{2m+1}{i} L_{(2m+1-2i)jr}^{n} F_{(2m+1-2i)(jrn+js)}, & jr \ even; \\ 5^{n/2-m} \sum_{i=0}^{m} (-1)^{i(js+1)} \binom{2m+1}{i} F_{(2m+1-2i)jr}^{n} F_{(2m+1-2i)(jrn+js)}, & jr \ odd, \ n \ even; \\ 5^{(n-1)/2-m} \sum_{i=0}^{m} (-1)^{i(js+1)} \binom{2m+1}{i} F_{(2m+1-2i)jr}^{n} L_{(2m+1-2i)(jrn+js)}, & jr \ odd, \ n \ odd; \end{cases}$$

$$(55)$$

$$\sum_{k=0}^{n} \binom{n}{k} L_{j(2rk+s)}^{2m+1} = \begin{cases} \sum_{i=0}^{m} (-1)^{ijs} \binom{2m+1}{i} L_{(2m+1-2i)jr}^{n} L_{(2m+1-2i)(jrn+js)}, & jr \ even; \\ 5^{n/2} \sum_{i=0}^{m} (-1)^{ijs} \binom{2m+1}{i} F_{(2m+1-2i)jr}^{n} L_{(2m+1-2i)(jrn+js)}, & jr \ odd, \ n \ even; \\ 5^{(n+1)/2} \sum_{i=0}^{m} (-1)^{ijs} \binom{2m+1}{i} F_{(2m+1-2i)jr}^{n} F_{(2m+1-2i)(jrn+js)}, & jr \ odd, \ n \ odd. \end{cases}$$
(56)

*Proof.* Set x = 1, z = 1 and write 2m + 1 for m and 2r for r in identities (BF) and (BL). Note that

$$1 + \alpha^{(2m+1-2i)2jr} = \begin{cases} \alpha^{(2m+1-2i)jr} L_{(2m+1-2i)jr}, & jr \text{ even}; \\ \alpha^{(2m+1-2i)jr} F_{(2m+1-2i)jr} \sqrt{5}, & jr \text{ odd.} \end{cases}$$

**Theorem 6.** Let m and n be non-negative integers and let j, r and s be any integers. Then,

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} F_{j(2rk+s)}^{2m+1} = \begin{cases} (-1)^{n} 5^{-m} \sum_{i=0}^{m} (-1)^{i(js+1)} \binom{2m+1}{i} L_{(2m+1-2i)jr}^{n} F_{(2m+1-2i)(jrn+js)}, & jr \ odd; \\ 5^{n/2-m} \sum_{i=0}^{m} (-1)^{i(js+1)} \binom{2m+1}{i} F_{(2m+1-2i)jr}^{n} F_{(2m+1-2i)(jrn+js)}, & jr \ even, \ n \ even; \\ -5^{(n-1)/2-m} \sum_{i=0}^{m} (-1)^{i(js+1)} \binom{2m+1}{i} F_{(2m+1-2i)jr}^{n} L_{(2m+1-2i)(jrn+js)}, & jr \ even, \ n \ odd; \end{cases}$$
(57)

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} L_{j(2rk+s)}^{2m+1} = \begin{cases} (-1)^{n} \sum_{i=0}^{m} (-1)^{ijs} \binom{2m+1}{i} L_{(2m+1-2i)jr}^{n} L_{(2m+1-2i)(jrn+js)}, & jr \ odd; \\ 5^{n/2} \sum_{i=0}^{m} (-1)^{ijs} \binom{2m+1}{i} F_{(2m+1-2i)jr}^{n} L_{(2m+1-2i)(jrn+js)}, & jr \ even, \ n \ even; \\ -5^{(n+1)/2} \sum_{i=0}^{m} (-1)^{ijs} \binom{2m+1}{i} F_{(2m+1-2i)jr}^{n} F_{(2m+1-2i)(jrn+js)}, & jr \ even, \ n \ odd. \end{cases}$$
(58)

*Proof.* Put x = 1, z = -1 and write 2m + 1 for m and 2r for r in identities (BF) and (BL). Note that

$$1 - \alpha^{(2m+1-2i)2jr} = \begin{cases} -\alpha^{(2m+1-2i)jr} F_{(2m+1-2i)jr} \sqrt{5}, & jr \text{ even}; \\ -\alpha^{(2m+1-2i)jr} L_{(2m+1-2i)jr}, & jr \text{ odd.} \end{cases}$$

# References

- L. Carlitz, Some classes of Fibonacci sums, The Fibonacci Quarterly 16:5 (1978), 411– 426.
- [2] L. A. G. Dresel, Transformations of Fibonacci-Lucas identities, in Applications of Fibonacci Numbers, Vol. 5, Dordrecht: Kluwer, 1993, pp. 169–184.
- [3] V. E. Hoggatt Jr and M. Bicknell, Some new Fibonacci identities, The Fibonacci Quarterly 2:1 (1964), 29–32.
- [4] V. E. Hoggatt Jr and M. Bicknell, Fourth power identities from Pascal's triangle, The Fibonacci Quarterly 2:4 (1964), 261–266.
- [5] V. E. Hoggatt, Jr., J. W. Phillips and H. T. Leonard, Jr., Twenty-four master identities, *The Fibonacci Quarterly* 9:1 (1971), 1–17.
- [6] E. Kiliç and I. Akkus and N. Omür and Y. T. Ulutaş, Formulas for binomial sums including powers of Fibonacci and Lucas numbers, UPB Scientific Bulletin, Series A 77:4 (2015), 69–78.
- [7] T. Koshy, Fibonacci and Lucas Numbers with Applications, Wiley-Interscience, 2001.
- [8] J. W. Layman, Certain general binomial-Fibonacci sums, The Fibonacci Quarterly 15:3 (1977), 362–366.
- C. T. Long, Some binomial Fibonacci identities, in Applications of Fibonacci Numbers, Vol. 3, Dordrecht: Kluwer, 1990, pp. 241–254.
- [10] S. Vajda, Fibonacci and Lucas Numbers, and the Golden Section: Theory and Applications, Dover Press, 2008.
- [11] J. Wessner, Binomial sums of Fibonacci powers, The Fibonacci Quarterly 4:4 (1966), 355–358.

Concerned with sequences: <u>A000032</u>, <u>A000045</u>.