Nicola Bastianello, Liam Madden, Ruggero Carli, Emiliano Dall'Anese

Abstract—This paper proposes a framework to study the convergence of stochastic optimization and learning algorithms. The framework is modeled over the different challenges that these algorithms pose, such as (i) the presence of random additive errors (e.g. due to stochastic gradients), and (ii) random coordinate updates (e.g. due to asynchrony in distributed set-ups). The paper covers both convex and strongly convex problems, and it also analyzes online scenarios, involving changes in the data and costs. The paper relies on interpreting stochastic algorithms as the iterated application of stochastic operators, thus allowing us to use the powerful tools of operator theory. In particular, we consider operators characterized by additive errors with sub-Weibull distribution (which parameterize a broad class of errors by their tail probability), and random updates. In this framework we derive convergence results in mean and in high probability, by providing bounds to the distance of the current iteration from a solution of the optimization or learning problem. The contributions are discussed in light of federated learning applications.

*Index Terms*—Stochastic operators, inexact optimization, online optimization, high probability convergence, federated learning

## I. INTRODUCTION

Recent technological advances in a range of disciplines – from machine learning, to data-driven optimization and control, with their applications to smart power grids, traffic networks, healthcare, etc. – introduced a wide set of challenges to the implementation and analysis of optimization and learning algorithms. As a motivating example, we elaborate on these challenges in the context of federated learning [1], [2], which was designed to allow a set of agents to cooperatively train a model without the need to directly share data. Due to the distributed set-up, algorithms designed in this framework need to deal with *asynchrony, and limited or unreliable communications*. This is similarly a challenge in any application where multi-agent systems are deployed, such as distributed optimization and parallel computing [3]–[6]. Additionally, due to the size of available data-sets, the agents may need to resort

R. Carli is with the Department of Information Engineering (DEI), University of Padova, Italy. carlirug@dei.unipd.it. to the use of *stochastic gradients*, which are computed on a sub-set of the available data [7]–[9]. Using stochastic gradients reduces the computational cost but may lead to less accuracy. Alternatively, when gradients are not directly accessible, for example due to sparse users' feedback [10], agents may need to approximate them from functional evaluations (0-th order gradients) [11]–[13]. Finally, the data on which the agents train their model may change over time, due to changes in the phenomenon being observed or new data arriving in real time [14]–[17]. This turns the problem into an *online learning* problem, with the agents now tracking the optimal model as it changes in response to changes in the data.

Abstracting away from this example, in many applications of optimization and learning we need to design and analyze algorithms that are provably robust to different sources of stochasticity and changes in the data. In particular, in this paper we focus on the analysis side of this challenge by leveraging the formalism of *operator theory*. Operator theory has been shown to offer a valuable tool for the analysis of algorithms, both in traditional static settings [18]–[21] and in online settings [14], [16], [22], [23]. The key insight is that algorithmic steps can be interpreted as operators (or maps when working in an Euclidean space), with fixed points of the operators coinciding with optimal solutions of the optimization or learning problem [19]-[21]. This link gives access to powerful tools to characterize the convergence of existing algorithms [18], [20], [21], and it further inspires the design of new ones, e.g. [24]-[26].

However, as discussed above, in many applications of interest the algorithms we apply are stochastic. To analyze their convergence, we will therefore make use of the formalism of stochastic operators, which we review in the following. One source of stochasticity is that of random coordinate updates, in which only some part of the operator is applied at any given iteration [4], [6], [27]-[30]. This class of operators model distributed algorithms in which only some agents, e.g. due to asynchrony, perform an update and hence apply the corresponding part of the algorithm/operator. Operators subject to additive errors, e.g. due to the use of stochastic gradients, have also been studied, see [7], [27]-[29], [31]. It is typical to handle stochastic errors in the algorithmic map or operator by either assuming that their norm vanishes asymptotically, or by multiplying them by a vanishing parameter. While this choice allows one to prove almost sure convergence, in many practical applications the additive error is not (or cannot be made) vanishing, especially in an online scenario.

N. Bastianello is with the School of Electrical Engineering and Computer Science, and Digital Futures, KTH Royal Institute of Technology, Sweden. nicolba@kth.se.

L. Madden is with the Department of Electrical and Computer Engineering, University of British Columbia, Vancouver, BC, CA. liam@ece.ubc.ca

E. Dall'Anese is with the Department of Electrical, Computer and Energy Engineering, and affiliate faculty of the Department of Applied Mathematics, University of Colorado Boulder, Boulder, CO, US. emiliano.dallanese@colorado.edu

To delineate our contribution in the context of the discussion above, we analyze algorithms that can be characterized by the stochastic iteration (formalized in section III):

$$x_i^{\ell+1} = \begin{cases} \mathcal{T}_i \boldsymbol{x}^{\ell} + e_i^{\ell} & \text{w.p. } p_i \\ x_i^{\ell} & \text{w.p. } 1 - p_i \end{cases}, \quad \ell \in \mathbb{N}$$
(1)

where  $\mathcal{T} \boldsymbol{x} = (\mathcal{T}_1 \boldsymbol{x}, \dots, \mathcal{T}_n \boldsymbol{x})$  is an operator,  $\boldsymbol{x} = (x_1, \dots, x_n)$ , and  $e_i^{\ell}$  is an additive error. The stochasticity in (1) has two sources: (i) each "coordinate" of the operator is updated with probability  $p_i$  (e.g. due to asynchrony), and (ii) the update is performed using an inexact version of the operator (e.g. stochastic gradients). Additionally, we will be interested in the (iii) online scenario in which the operator changes over time to model applications in online optimization and learning; that is, at time  $\ell + 1$  we apply  $\mathcal{T}_i^{\ell+1}$ . The iteration (1) will be studied under the assumption that the operator is either contractive or averaged (in which case (1) can be seen as a stochastic Krasnosel'skii-Mann).

As in *e.g.*, [27], [28], we model inexact operators as operators subject to the additive errors  $e_i$ . However, differently from previous results, in this paper we perform our analysis for errors with *sub-Weibull distributions*. That is, the norm of  $e_i$  satisfies

$$\mathbb{P}\left[\|e_i\| \ge \epsilon\right] \le 2\exp\left(-\left(\frac{\epsilon}{\nu}\right)^{1/\theta}\right), \quad \forall \ \epsilon > 0 \qquad (2)$$

for some parameters  $\theta \ge 0$ ,  $\nu > 0$ . The class of sub-Weibull r.v.s allows us to consider distributions for the additive errors that may have heavy tails [32]–[35]; this class is also general and it includes the sub-Gaussian and sub-exponential classes as sub-cases, as well as random variable whose distribution has a finite support [36], [37]. As explained shortly, the sub-Weibull model will allow us to provide high probability bounds for the convergence of (1) by deriving pertinent concentration results. As defined in (2), sub-Weibull random variables are characterized by a tail parameter  $\theta$ , that defines the decay rate of the tails. Besides the convenient properties of the sub-Weibull class (e.g. closure under scaling, sum, and product) that allow for a unified theoretical analysis, there is a growing body of work showing that heavy-tailed distributions arise in machine learning applications, see [32], [38]-[44]. This is especially relevant while training (deep) neural networks, and this phenomenon can also arise in decentralized learning set-ups [45].

**Contributions.** Overall, this paper offers the following contributions:

We study the convergence of (1), and provide convergence results in mean and in high probability when the errors follow a sub-Weibull distribution. In particular, bounds are offered for the distance from the fixed point (if the operator is contractive) or for the *cumulative fixed point residual* (if the operator is averaged). These error bounds hold for any iteration *l* ∈ N with a given, arbitrary probability. We also show that the high-probability bounds – in the form P[||x|| ≤ ε(δ)] ≥ 1 − δ for the random variable x – scale with a factor ~ log(1/δ), as

opposed to a scaling  $\sim 1/\delta$  that one would obtain via Markov's inequality.

- 2) The framework we propose leverages a sub-Weibull model [32]–[35] for the norm of the additive errors. To the best of our knowledge, this is the first time that sub-Weibull models are employed in combination with operator-theoretic tools. By using sub-Weibulls, we are able to model a very broad class of random variables, fitting different practical applications.
- 3) As mentioned above, the convergence results proposed in this paper hold with high probability, as opposed to the almost sure convergence results of *e.g.* [27]. We discuss the difference between the two viewpoints, and also further characterize the convergence of (1) in almost sure terms (under some additional assumptions, such as vanishing errors).
- 4) The analysis provided also holds in online scenarios, in which the operator characterizing (1) changes over time.

**Organization.** In section II we review some preliminaries in operator theory, and introduce the sub-Weibull formalism. In section III, building on the motivating example of federated learning, we formalize and discuss the stochastic framework. In sections IV and V we present and discuss, respectively, mean and high probability convergence results derived in the proposed framework. Finally, section VI presents some illustrative numerical results.

#### II. PRELIMINARIES

## A. Operators

Let  $\mathbb{H}_i$ ,  $i \in \{1, ..., n\}$  be (possibly infinite-dimensional) Hilbert spaces with inner product  $\langle \cdot, \cdot \rangle_i$ , induced norm  $\|\cdot\|_i$ and identity  $\mathcal{I}_i : \mathbb{H}_i \to \mathbb{H}_i$ . We consider the direct sum space  $\mathbb{H} = \mathbb{H}_1 \oplus \cdots \oplus \mathbb{H}_n$ , whose elements are  $\boldsymbol{x} = (x_1, \ldots, x_n)$ with  $x_i \in \mathbb{H}_i$  for all  $i \in \{1, \ldots, n\}$ . For  $\boldsymbol{x}, \boldsymbol{y} \in \mathbb{H}$ , we define the inner product as  $\langle \boldsymbol{x}, \boldsymbol{y} \rangle = \sum_{i=1}^n \langle x_i, y_i \rangle_i$ , and denote the induced norm and identity of  $\mathbb{H}$  as  $\|\cdot\|$  and  $\mathcal{I} : \mathbb{H} \to \mathbb{H}$ , respectively. We consider operators  $\mathcal{T} : \mathbb{H} \to \mathbb{H}$  defined as

$$\mathcal{T}\boldsymbol{x} = (\mathcal{T}_1\boldsymbol{x},\ldots,\mathcal{T}_n\boldsymbol{x})$$
 (3)

where  $\mathcal{T}_i : \mathbb{H} \to \mathbb{H}_i$  for all  $i \in \{1, \ldots, n\}$ .

A central theme of the paper is to compute fixed points of a given operator via iterative algorithms. To this end, in the following we introduce pertinent definitions and results. For a background on operator theory we refer to *e.g.* [20], [46].

**Definition 1** (Fixed points). Let  $\mathcal{T} : \mathbb{H} \to \mathbb{H} : \mathbf{x} \mapsto \mathcal{T}\mathbf{x} = (\mathcal{T}_1\mathbf{x}, \ldots, \mathcal{T}_n\mathbf{x})$ . The point  $\bar{\mathbf{x}} \in \mathbb{H}$  is a fixed point of  $\mathcal{T}$  if  $\bar{\mathbf{x}} = \mathcal{T}\bar{\mathbf{x}}$ . We denote the fixed set of  $\mathcal{T}$  as fix $(\mathcal{T}) = \{\mathbf{x} \in \mathbb{H} | \mathbf{x} = \mathcal{T}\mathbf{x}\}$ .

Notice that a fixed point  $\bar{x} = (\bar{x}_1, \dots, \bar{x}_n)$  by definition is such that  $\bar{x}_i = \mathcal{T}_i \bar{x}$  for all  $i \in \{1, \dots, n\}$ . In the following, we review well-known definitions and properties of operators (see, e.g., [20]).

**Definition 2** (Non-expansive, contractive). *The operator*  $\mathcal{T}$  *is*  $\zeta$ -Lipschitz continuous,  $\zeta \geq 0$ , *if:* 

$$\|\mathcal{T}\boldsymbol{x} - \mathcal{T}\boldsymbol{y}\| \leq \zeta \|\boldsymbol{x} - \boldsymbol{y}\|, \quad \forall \boldsymbol{x}, \boldsymbol{y} \in \mathbb{H}.$$
 (4)

The operator is non-expansive if  $\zeta = 1$ , and contractive if  $\zeta \in (0, 1)$ .

**Definition 3** (Averaged). The operator  $\mathcal{T}$  is averaged if and only if there exist  $\alpha \in (0,1)$  and a non-expansive operator  $\mathcal{R} : \mathbb{H} \to \mathbb{H}$  such that  $\mathcal{T} = (1 - \alpha)\mathcal{I} + \alpha \mathcal{R}$ . Equivalently,  $\mathcal{T}$ is  $\alpha$ -averaged if

$$\left\|\mathcal{T}\boldsymbol{x}-\mathcal{T}\boldsymbol{y}\right\|^{2} \leq \left\|\boldsymbol{x}-\boldsymbol{y}\right\|^{2} - \frac{1-\alpha}{\alpha}\left\|(\mathcal{I}-\mathcal{T})\boldsymbol{x}-(\mathcal{I}-\mathcal{T})\boldsymbol{y}\right\|^{2}$$

*for any*  $x, y \in \mathbb{H}$ *.* 

We discuss now the existence of fixed points of nonexpansive operators.

**Lemma 1** (Browder's theorem). Let  $\mathbb{D} \subset \mathbb{H}$  be a non-empty, convex, compact subset, and let  $\mathcal{T} : \mathbb{D} \to \mathbb{D}$  be non-expansive, then  $\operatorname{fix}(\mathcal{T}) \neq \emptyset$  [20, Theorem 4.29].

**Lemma 2** (Banach-Picard theorem). Let  $\mathcal{T} : \mathbb{H} \to \mathbb{H}$ be a contractive operator, then fix( $\mathcal{T}$ ) is a singleton [20, Theorem 1.50]. The unique fixed point is the limit of the sequence generated by:

$$x^{\ell+1} = \mathcal{T}x^{\ell}, \quad \ell \in \mathbb{N}.$$

If we want to compute a fixed point of the non-expansive operator  $\mathcal{T}$ , the following results can be used.

**Lemma 3.** Let  $\mathcal{T} : \mathbb{H} \to \mathbb{H}$  be a non-expansive operator with  $\operatorname{fix}(\mathcal{T}) \neq \emptyset$ . Then for any  $\alpha \in (0, 1)$ , the  $\alpha$ -averaged operator  $\mathcal{T}_{\alpha} = (1 - \alpha)\mathcal{I} + \alpha\mathcal{T}$  is such that  $\operatorname{fix}(\mathcal{T}_{\alpha}) = \operatorname{fix}(\mathcal{T})$ .

*Proof.* Let  $\bar{x} \in \text{fix}(\mathcal{T})$ , by definition  $\bar{x} = \mathcal{T}\bar{x}$ . Therefore given  $\alpha \in (0, 1)$  we have  $(1 - \alpha)\bar{x} + \alpha \mathcal{T}\bar{x} = \bar{x} - \alpha \bar{x} + \alpha \bar{x} = \bar{x}$ .  $\Box$ 

**Lemma 4** (Krasnosel'skiĭ-Mann theorem). Let  $\mathbb{D} \subset \mathbb{H}$  be a non-empty closed convex subset, and let  $\mathcal{T} : \mathbb{D} \to \mathbb{D}$  be a non-expansive operator (notice that by Lemma 1 fix $(\mathcal{T}) \neq \emptyset$ ). Let  $\alpha \in (0, 1)$ , then the Krasnosel'skiĭ-Mann iteration

$$\boldsymbol{x}^{\ell+1} = (1-\alpha)\boldsymbol{x}^{\ell} + \alpha \mathcal{T} \boldsymbol{x}^{\ell}, \quad \ell \in \mathbb{N},$$

guarantees that  $\|(\mathcal{I} - \mathcal{T})\mathbf{x}^{\ell}\| \to 0$  as  $\ell \to \infty$  [20, Theorem 5.15].

In the remainder of the paper we will thus focus on studying the convergence of the Banach-Picard in two cases: i) when the operator is contractive, and ii) when it is averaged.

### B. Probability

In this section, we provide some definitions and results in probability theory that will be used in the paper to derive convergence results in high-probability. Throughout the paper, the underlying probability space will be  $(\Omega, \mathcal{F}, \mathbb{P})$ . We start by introducing the definition of *sub-Weibull random variables* [32]–[35].

**Definition 4** (Sub-Weibull random variable). A random variable x is said to be sub-Weibull if there exist  $\theta \ge 0$ ,  $\nu > 0$  such that

$$\left\|x\right\|_{k} = \mathbb{E}\left[\left|x\right|^{k}\right]^{1/k} \le \nu k^{\theta}, \quad \forall k \ge 1,$$

and we denote it by  $x \sim \text{subW}(\theta, \nu)$ .

The following equivalence result relates the moments growth condition of Definition 4 with a bound for the tail probability [35, Lemma 5].

**Lemma 5** (Sub-Weibull tail probability). Let  $x \sim \text{subW}(\theta, \nu)$ , then the tail probability verifies the bound

$$\mathbb{P}\left[|x| \ge \epsilon\right] \le 2\exp\left(-\left(\frac{\epsilon}{c(\theta)\nu}\right)^{1/\theta}\right), \quad \forall \ \epsilon > 0, \qquad (5)$$

where  $c(\theta) := (2e/\theta)^{\theta}$ .

Lemma 5 shows that the tails of a sub-Weibull r.v. become heavier as the parameter  $\theta$  grows larger. Moreover, setting  $\theta = 1/2$  and  $\theta = 1$  yields the class of sub-Gaussians and sub-exponential random variables, respectively; see, *e.g.*, [36], [37]. The following lemma shows how the tail probability equation (5) can be used to give high probability bounds.

**Lemma 6** (High probability bound). Let  $x \sim \text{subW}(\theta, \nu)$ , then for any  $\delta \in (0, 1)$ , w.p.  $1 - \delta$  we have the bound:

$$|x| \le \log^{\theta}(2/\delta)c(\theta) \ \nu$$

*Proof.* By Lemma 5 we have, for any  $\epsilon > 0$ :

$$\mathbb{P}\left[|x| \ge \epsilon\right] \le 2 \exp\left(-\left(\epsilon/(c(\theta)\nu)\right)^{1/\theta}\right).$$

Setting the right-hand side equal to  $\delta$  and solving for  $\epsilon$  we get  $\epsilon = c(\theta)\nu \log^{\theta}(2/\delta)$  which implies that, w.p.  $1 - \delta$  we have  $|x| \le \epsilon = c(\theta)\nu \log^{\theta}(2/\delta)$ .

We characterize now the properties of the sub-Weibull class of random variables.

**Lemma 7** (Inclusion). Let  $x \sim \text{subW}(\theta, \nu)$ , then  $x \sim \text{subW}(\theta', \nu')$  for any  $\theta' \geq \theta$ ,  $\nu' \geq \nu$ .

*Proof.* By assumption we have  $||x||_k \leq \nu k^{\theta}$ . Using the fact that  $\nu k^{\theta} \leq \nu' k^{\theta'}$  (which holds since  $k \geq 1$ ) yields the thesis; cf. [32, Proposition 1].

**Lemma 8** (Closure). The class of sub-Weibull random variables is closed w.r.t. product by a scalar, sum, product, and exponentiation, according to the following rules.

- 1) Product by scalar: let  $x \sim \text{subW}(\theta, \nu)$  and  $a \in \mathbb{R}$ , then  $ax \sim \text{subW}(\theta, |a|\nu)$ ;
- 2) Sum: let  $x_i \sim \text{subW}(\theta_i, \nu_i)$ ,  $i \in \{1, 2\}$ , possibly dependent, then  $x_1 + x_2 \sim \text{subW}(\max\{\theta_1, \theta_2\}, \nu_1 + \nu_2)$ ;
- 3) Product: let  $x_i \sim \text{subW}(\theta_i, \nu_i)$ ,  $i \in \{1, 2\}$ , and independent, then  $x_1 x_2 \sim \text{subW}(\theta_1 + \theta_2, \nu_1 \nu_2)$ .
- 4) Power: let  $x \sim \text{subW}(\theta, \nu)$  and a > 0, then  $x^a \sim \text{subW}(a\theta, \nu^a \max\{1, a^{a\theta}\})$ .

We conclude with some remarks.

**Remark 1** (Square of sub-Weibull). A consequence of 4) in Lemma 8 is that the square of  $x \sim \text{subW}(\theta, \nu)$  is itself a sub-Weibull, characterized by  $x^2 \sim \text{subW}(2\theta, 4^{\theta}\nu^2)$ . A particular case is the well known fact that the square of a sub-Gaussian ( $\theta = 1/2$ ) is sub-exponential ( $\theta = 1$ ), see e.g. [37, Lemma 2.7.6]. **Remark 2** (Mean of sub-Weibulls). Notice that the definition of sub-Weibulls and their properties does not require that their mean be zero. Moreover, if  $x \sim \text{subW}(\theta, \nu)$  and  $x \ge 0$  almost surely, then  $\mathbb{E}[x] \le \nu$ , since  $||x||_1 = \mathbb{E}[x]$ .

**Remark 3** (Bounded r.v.s). We can see that bounded r.v.s are sub-Weibull with  $\theta = 0$ ; indeed, let x be a r.v. such that a.s.  $|x| \leq b$ , then  $||x||_k \leq b = bk^0$ ,  $k \geq 1$ . This characterization is "optimal" in terms of  $\theta$ , since  $\theta = 0$  corresponds to the lightest possible tail. However, it is sub-optimal in the other parameter,  $\nu$ , which does not reflect the overall distribution of x, only its maximum absolute value. Alternatively, we know by [37, p. 24] that the class of sub-Gaussian random variables includes that of bounded r.v.s; this implies that bounded r.v.s are sub-Weibull with  $\theta = 1/2$ .

#### **III. MOTIVATION AND FRAMEWORK DEVELOPMENT**

We start this section by discussing a motivating example in the context of *federated learning* [1], [2], around which we will formalize our stochastic operator framework.

#### A. Motivating example: federated learning



Fig. 1. The architecture in federated learning.

In *federated learning*, a set of n agents, aided by a coordinator, aim to cooperatively train a model without directly sharing the data they store [1], [2]. In order to keep the data private, the agents share the results of local training with the coordinator, which aggregates them into a more accurate model. Formally, the goal is to solve the optimization problem

$$\min_{x_i \in \mathbb{R}^q} \sum_{i=1}^n f_i(x_i) \quad \text{s.t. } x_1 = x_2 = \ldots = x_n \tag{6}$$

where  $f_i : \mathbb{R}^q \to \mathbb{R} \cup \{+\infty\}$  is the loss function of agent *i*, defined over the local dataset. Let  $\{d_i^h\}_{h=1}^{m_i}$  be the  $m_i$  data points stored by *i*, then usually the local loss is of the form

$$f_i(x_i) = \sum_{h=1}^{m_i} \lambda(x_i; d_i^h) \tag{7}$$

with  $\lambda$  being a training loss (e.g. quadratic or logistic).

In principle, (6) could be solved via projected gradient descent by applying,  $i \in \{1, ..., n\}$ :

$$x_i^{\ell+1} = \mathcal{T}_i x^{\ell} := \frac{1}{n} \sum_{j=1}^n \left( x_j^{\ell} - \gamma \nabla f_j(x_j^{\ell}) \right), \quad \ell \in \mathbb{N}, \quad (8)$$

where the agents compute local gradient descent steps, and the coordinator computes their average (the projection onto  $x_1 = \ldots x_n$ ) and transmits it to the agents. In the following, using the notation defined in section II-A we write (8) as  $x^{\ell+1} = \mathcal{T}x^{\ell}$ . However, when solving learning problems there are several practical constraints that make implementing (8) unrealistic. In the following we discuss some of these practical challenges, and how they motivate the theoretical developments of subsequent sections. The section concludes by presenting a modified version of (8) that accounts for these challenges, and which can be interpreted as a stochastic operator, fitting in the framework of section III-B.

Challenge 1: additive errors: Learning problems are often high dimensional, both in the size of the unknown  $x \in \mathbb{R}^{q}$ (especially when training neural networks), and in the size of the local data sets. The size of the unknown poses a first challenge, since algorithm (8) requires sharing of qdimensional vectors, and the larger q the more expensive these communications become. In practice, then, quantization or compression is applied before the agents communicate with the coordinator [47], lessening the burden at the cost of introducing some error.

A second challenge arises when computing the gradient of local losses (7), since the larger  $m_i$  is the more computationally expensive gradient computations are. To reduce the cost of gradient evaluations, a standard strategy in learning is to use *stochastic gradients* [48], which approximate the true gradient using only a random subset of the local data points. But, similarly to communications reduction techniques, approximating local gradients introduces some error. In the following, we denote by  $\hat{\nabla} f_i(x_i)$  a stochastic local gradient.

We are therefore interested in analyzing the modified version of (8) given by  $x^{\ell+1} = \mathcal{T}x^{\ell} + e^{\ell}$ , where  $e^{\ell}$  is a random vector modeling the errors introduced by the use of inexact communications and stochastic gradients.

**Remark 4** (Local training). In practice, federated learning algorithm use local training, that is, the agents perform multiple local gradient steps in between communication rounds. While this reduces the ratio of communications to gradient evaluations, it introduces client drift [49]; interpreting it as an additive error, one may then apply the results of the next sections to analyze the convergence of algorithms employing local training.

*Challenge 2: asynchrony:* The agents cooperating in the learning procedure may be *highly heterogeneous*, in that they have different computational resources [2]. In (8) all the agents compute a local gradient at the same time – however, heterogeneous agents will have different computation times. Therefore, requiring the agents to work synchronously implies that the faster agents will sit idle while waiting for the slower ones to conclude the local computations [3].

One way to solve this problem is to allow *asynchronous* activation of the agents, so that each agent is free to perform local training at its own pace. More formally, let  $\mathbb{I}^{\ell} \subset \{1, \ldots, n\}$ ,  $|\mathbb{I}^{\ell}| \leq n$ , be the subset of agents that concluded a local computation at time  $\ell \in \mathbb{N}$ . Then we are interested in algorithms in which the coordinator only aggregates information from  $\mathbb{I}^{\ell}$ , instead of all agents. Algorithm 1 presents a modified

version of (8) that allows for asynchronous activations (see Appendix B for the derivation).

We also remark that asynchrony could be enforced by design as an additional measure to reduce the amount of communications required at each time.

Challenge 3: online problems: The problem (6) discussed so far is static, in the sense that the data sets defining the local losses do not change over time. However, in many learning applications the agents may be continuously collecting new data and consequently modifying their loss. This results in an online learning problem, characterized by [14]-[16]

$$\min_{x_i \in \mathbb{R}^q} \sum_{i=1}^n f_i^{\ell}(x_i) \quad \text{s.t. } x_1 = x_2 = \ldots = x_n.$$

In this set-up the solution(s) to the problem at time  $\ell \in \mathbb{N}$ may not coincide with those at time  $\ell + 1$ , and our analysis of Algorithm 1's performance needs to account for this fact.

# Algorithm 1 Federated gradient descent.

**Input:** Initial conditions  $z^0$ ,  $x_i^0$ , and step-size  $\gamma$ .

1: for  $\ell = 0, 1, 2, \ldots$ 

// compressed communications

- the coordinator compresses and transmits  $z^{\ell}$  to the 2. active agents  $\mathbb{I}^{\ell}$
- for  $i \in \{1, ..., n\}$ 3:

// asynchrony

- if  $i \in \mathbb{I}^{\ell}$  then
  - // inexact gradients
- the agent performs a local (stochastic) gradient 5: step

$$x_i^{\ell+1} = z^\ell - \gamma \hat{\nabla} f_i(z^\ell)$$

#### // compressed communications

and compresses and transmits  $x_i^{\ell+1}$  to the co-6: ordinator

7: else

4:

- otherwise sets  $x_i^{\ell+1} = x_i^{\ell}$ 8:
- end if 9:
- 10: end for
- 11: the coordinator computes

$$z^{\ell+1} = \frac{1}{n} \left( \sum_{i \in \mathbb{I}^\ell} x_i^{\ell+1} + \sum_{i \not\in \mathbb{I}^\ell} x_i^\ell \right)$$

12: end for

#### B. Stochastic operator framework

Motivated by the example discussed so far, we are now ready to formalize the stochastic operator framework of interest in this paper. Consider an operator  $\mathcal{T} \boldsymbol{x} = (\mathcal{T}_1 \boldsymbol{x}, \dots, \mathcal{T}_n \boldsymbol{x})$ , in the following we analyze the convergence of the update, for  $i \in \{1, ..., n\}$ :

$$x_{i}^{\ell+1} = \begin{cases} \mathcal{T}_{i}^{\ell+1} \boldsymbol{x}^{\ell} + e_{i}^{\ell} & \text{w.p. } p_{i} \\ x_{i}^{\ell} & \text{w.p. } 1 - p_{i} \end{cases}$$
(9)  
$$= (1 - u_{i}^{\ell}) x_{i}^{\ell} + u_{i}^{\ell} \left( \mathcal{T}_{i}^{\ell+1} \boldsymbol{x}^{\ell} + e_{i}^{\ell} \right) \\=: \hat{\mathcal{T}}_{i}^{\ell+1} \boldsymbol{x}^{\ell} + u_{i}^{\ell} e_{i}^{\ell} \end{cases}$$

where  $\ell \in \mathbb{N}$  is the iteration index,  $u_i^{\ell}$  are Bernoulli random variables that indicate whether a coordinate is updated or not at iteration  $\ell$ , and  $e^{\ell} = (e_1^{\ell}, \dots, e_n^{\ell})$  is a random vector of additive noise. As discussed in section III-A, in federated learning only the coordinates corresponding to active nodes (with  $u_i^{\ell}$ ) are updated, and additive noise may be due to inexact communications or stochastic gradients (hence  $e_i^{\ell}$ ). The operator  $\mathcal{T}^{\ell}$  is allowed to change over time to account for online learning problems defined by streaming sources of data.

In the following we introduce and discuss some assumptions to formalize the framework. We remark that throughout the paper, the initial condition  $x^0 \in \mathbb{H}$  will be assumed to be deterministic.

Assumption 1 (Stochastic framework 1). The following is assumed.

- (i)  $u^{\ell} = (u_1^{\ell}, \dots, u_n^{\ell})$  is a  $\{0, 1\}^n$ -valued random vector with  $p_i := \mathbb{P}[u_i = 1] > 0, i \in \{1, \dots, n\}.$
- (ii) The additive error e<sup>ℓ</sup> ∈ H is an H-valued random vector such that E [||e<sup>ℓ</sup>||<sup>2</sup>] ≤ ν<sup>2</sup>.
  (iii) The random processes {u<sup>ℓ</sup>}<sub>ℓ∈N</sub> and {e<sup>ℓ</sup>}<sub>ℓ∈N</sub> are i.i.d.
- and independent of each other.

We provide now two assumptions on the operator defining (9), which will be used to provide two different sets of results. Notice that these assumptions apply to the underlying *deterministic* operator  $\mathcal{T}^{\ell+1}$  only.

Assumption 2 (Contractive set-up). Consider an operator  $\mathcal{T}^{\ell}: \mathbb{D} \to \mathbb{D}$ . The following is assumed.

- (i) The operator  $\mathcal{T}^{\ell} : \mathbb{H} \to \mathbb{H}$  is  $\zeta$ -contractive for all  $\ell \in \mathbb{N}$ , and we denote by  $\bar{x}^{\ell}$  its unique fixed point.
- (ii) There exists  $\sigma \geq 0$  such that the fixed points of consecutive operators have a bounded distance

$$\|\bar{\boldsymbol{x}}^{\ell} - \bar{\boldsymbol{x}}^{\ell-1}\| \leq \sigma, \quad \forall \ell \in \mathbb{N}.$$

**Assumption 3** (Averaged set-up). Consider an operator  $\mathcal{T}^{\ell}$ :  $\mathbb{D} \to \mathbb{D}$ . The following is assumed.

- (i) The operator  $\mathcal{T}^{\ell} : \mathbb{D} \to \mathbb{D}$  is  $\alpha$ -averaged for all  $\ell \in \mathbb{N}$ , with  $\mathbb{D} \subset \mathbb{H}$  being convex and compact.
- (ii) There exists  $\sigma \geq 0$  such that given  $\bar{x}^{\ell} \in fix(\mathcal{T}^{\ell}), \ \ell \in \mathbb{N}$ , then

$$\inf_{\bar{\boldsymbol{x}}^{\ell-1}\in \operatorname{fix}(\mathcal{T}^{\ell-1})} \left\| \bar{\boldsymbol{x}}^{\ell} - \bar{\boldsymbol{x}}^{\ell-1} \right\| \leq \sigma.$$

In the following we denote by  $diam(\mathbb{D})$  the diameter of  $\mathbb{D}$ . Before moving on to the convergence analysis, we discuss the framework and the assumptions that characterize it.

1) Update model: We note that Assumption 1(i) does not require independence among the components of  $u^{\ell}$ . Independence is only assumed between  $u^{\ell}$  and  $u^{h}$ , for any pair  $\ell, h \in \mathbb{N}, \ \ell \neq h.$ 

Under Assumption 1(i) there may exist times during which *none* of the coordinates are updated, because  $u_i^{\ell} = 0$  for all  $i \in \{1, \ldots, n\}$ . This is different from the framework used in *e.g.* [27], [28], where at least one  $u_i^{\ell}$  must always be different from zero. Accounting for such events is important in online scenarios, where the problem changes (e.g. due to changes in the environment) even when updates are not performed.

2) Additive error model: The error model characterized by Assumption 1(ii) implies that the variance of the additive error is bounded. This is a common assumption in learning applications, where the error  $e^{\ell}$  quantifies the difference between the gradient and a stochastic approximation thereof [48], [50]:

$$e^{\ell} = \nabla f(\boldsymbol{x}^{\ell}) - \hat{\nabla} f(\boldsymbol{x}^{\ell}).$$

Additionally, some quantization or compression errors also verify this assumption. This is the case for uniform or random quantization [51]; see also [47] for a comprehensive overview.

We remark that this model implies that errors are persistently present over time. This sets the paper apart from previous works on stochastic operators in which the additive error is assumed to decay to zero (or, equivalently, that the error is multiplied by a decaying parameter) [27]–[29]. Finally, we note that Assumptions 1 and 4 do not require independence of the error components at a fixed time  $\ell$ , or between the errors drawn at different times  $\ell$  and h. Finally, the errors could also be biased, that is, they could have mean different from zero.

3) Operators: In Assumption 3, the operators are defined on a compact  $\mathbb{D}$ , which is verified when for example  $\mathcal{T}^{\ell}$ contain a projection onto  $\mathbb{D}$ . On the one hand, this implies that Browder's theorem (cf. Lemma 1) holds, and hence that the operators have a non-empty fixed set. On the other, it allows us to guarantee that additive errors do not lead to divergence. This assumption is not required in *e.g.* [27] since the additive errors converge to zero asymptotically, and thus cannot lead to divergence.

In the context of convex optimization, we can relate Assumptions 2 and 3 to the properties of the cost function on which the operator  $\mathcal{T}^{\ell}$  is defined. For example, in the federated learning set-up of section III-A convex loss functions yield averaged operators, while strongly convex ones yield contractive operators [21] (cf. also the discussion in Appendix B).

Finally, a remark about the dynamic nature of the operators. As discussed in section III-A, in many applications we may be interested in solving optimization problems that change over time, as new data come in [16], [52], which motivates our choice to analyze time-varying operators. In this context, we can interpret (9) as an online algorithm, in which the output  $(x^{\ell})$  computed at time  $\ell$  is used to warm-start the computation at time  $\ell + 1$ . For this reason, it is necessary to provide bounds on the difference between consecutive operators (or equivalently, consecutive problems) to ensure that they are "close enough" for the warm-starting to provide an improvement in performance. This guarantee is provided by Assumption 2(ii) and Assumption 3(ii), which can be obtained for example bounding the rate of change of the operators.

## **IV. MEAN CONVERGENCE ANALYSIS**

We start by characterizing in section IV-A the mean convergence of (9) for contractive and averaged operators. The results and their implications will be discussed in section IV-B, while section IV-C will present some corollaries on the asymptotic convergence.

## A. Main results

Proposition 1 (Mean - contractive operators). Let Assumptions 1 and 2 hold, and let  $\{x^{\ell}\}_{\ell \in \mathbb{N}}$  be the sequence generated by (9). Then we have the following bound, for any  $\ell \in \mathbb{N}$ 

$$\mathbb{E}\left[\left\|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}\right\|\right] \leq \sqrt{\frac{\bar{p}}{\bar{p}}} \left(\chi^{\ell} \left\|\boldsymbol{x}^{0} - \bar{\boldsymbol{x}}^{0}\right\| + \frac{1 - \chi^{\ell}}{1 - \chi}(\chi\sigma + \nu)\right)$$
  
where  $\bar{p} := \max_{i} p_{i}, \ \underline{p} := \min_{i} p_{i}, \ and \ \chi := \sqrt{1 - \underline{p} + \underline{p}\zeta^{2}} \in \mathbb{E}$ 

(0, 1).

*Proof.* See Appendix C-A. 
$$\Box$$

**Proposition 2** (Mean – averaged operators). Let Assumptions 1 and 3 hold, and let  $\{x^{\ell}\}_{\ell \in \mathbb{N}}$  be the sequence generated by (9). Then we have the following bound, for any  $\ell \in \mathbb{N}$ 

$$\mathbb{E}\left[\frac{1}{\ell+1}\sum_{h=0}^{\ell}\left\|(\mathcal{I}-\mathcal{T}^{h+1})\boldsymbol{x}^{h}\right\|^{2}\right] \leq \frac{\alpha}{\underline{p}(1-\alpha)} \times \\ \times \left(\frac{1}{\ell+1}\left\|\boldsymbol{x}^{0}-\bar{\boldsymbol{x}}^{0}\right\|^{2}+\sigma^{2}+\nu^{2}+2\operatorname{diam}(\mathbb{D})(\sigma+\nu)\right)$$
  
*roof.* See Appendix C-B.

*Proof.* See Appendix C-B.

#### B. Discussion

1) Difference from deterministic convergence: In the following we discuss the difference between Propositions 1 and 2 and the convergence of the static and deterministic update  $\boldsymbol{x}^{\ell+1} = \mathcal{T}\boldsymbol{x}^{\ell}, \ \ell \in \mathbb{N}.$ 

a) Contractive case: By contractiveness we know that  $x^{\ell+1} = \mathcal{T}x^{\ell}$  converges linearly to fix $(\mathcal{T}) = \{\bar{x}\}$ , that is  $\|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}\| \leq \zeta^{\ell} \|\boldsymbol{x}^{0} - \bar{\boldsymbol{x}}\|$  [20, Theorem 1.50]. Comparing this bound with that of Proposition 1 we notice first of all that the introduction of random coordinate updates degrades the *convergence rate* – in mean – from  $\zeta$  to  $\chi \geq \zeta$ , in accordance with the results of [28].

Secondly, the presence of additive errors implies that we do not reach exact convergence to the fixed point, but rather to a neighborhood thereof. Moreover, at any given time  $\ell \in \mathbb{N}$ , the additive error may cause the overall stochastic operator to be expansive  $\|x^{\ell+1} - \bar{x}\| \geq \|x^{\ell} - \bar{x}\|$ , but the underlying contractiveness of  $\mathcal{T}$  ensure that this lead to inexact convergence rather than divergence.

b) Averaged case: The convergence of (9) for averaged operators can be proved only in terms of the *cumulative fixed* point residual  $\frac{1}{\ell+1} \sum_{h=0}^{\ell} ||(\mathcal{I} - \mathcal{T}^{h+1}) \boldsymbol{x}^h||^2$ , due to averaged-ness being weaker than contractiveness. On the other hand, for the deterministic update  $x^{\ell+1} = \mathcal{T}x^{\ell}$  it is possible to prove that  $\{\|(\mathcal{I} - \mathcal{T})\boldsymbol{x}^{\ell}\|\}_{\ell \in \mathbb{N}}$  is a monotonically decreasing sequence that converges to zero [20, Theorem 5.15]. Similarly to the contractive case, this is no longer the case in the presence of additive errors, and the metric used to analyze convergence needs to account for the overall evolution of the fixed point residual.

We remark that the concept of cumulative fixed point residual is similar to that of *regret* in convex optimization, and in particular to that of dynamic regret in online optimization [53]. Moreover, it includes as a particular case the different concept of regret based on the residual of the proximal gradient method proposed in [54].

2) *Time-variability:* As mentioned in section III-A, the stochastic framework we analyze allows for the operator to change over time, as this models *online optimization algorithms* [16], [52]. But the time-variability of the operators, and hence of their fixed point(s), can be seen as a second source of additive errors besides  $e^{\ell}$ . This error is always present, *also when no update is performed*, to account for the fact that the environment changes at every iteration  $\ell \in \mathbb{N}$ . This is the case in online optimization when the problem depends on *e.g.* measurements of a system, and the system will evolve even when no measurement is performed.

a) Path length: In online optimization, a widely used concept is that of *path-length*, that is, the cumulative distance between consecutive optimizers. This concept can be straightforwardly extended to the operator theoretical set-up, by defining it as the cumulative distance between consecutive fixed points:  $\sum_{h=0}^{\ell-1} \|\bar{x}^{h+1} - \bar{x}^h\| \le \ell\sigma$ . The path-length often appears in regret bounds, see *e.g.* [53], but notice that in the worst case it *grows linearly with*  $\ell$ , and to carry out the convergence analysis the additional assumption that it grows sub-linearly is required. What instead appears in our bound is a *weighted* path-length

$$\sum_{h=0}^{\ell-1} \chi^{\ell-h-1} \left\| \bar{\boldsymbol{x}}^{h+1} - \bar{\boldsymbol{x}}^{h} \right\| \leq \frac{1-\chi^{\ell}}{1-\chi} \sigma$$

which asymptotically reaches a fixed value, independent of  $\ell$ . This is due to the fact that we use the contractiveness of the operator, allowing to reach a tighter bound.

3) Convergence without random coordinate updates: We remark that the results of Propositions 1 and 2 can be adapted in a straightforward manner to a scenario where no random coordinate updates are performed. Indeed, if  $p_i = 1$ ,  $i \in \{1, \ldots, n\}$  then the two results provide bounds to the convergence of *inexact operators* modeled by

$$oldsymbol{x}^{\ell+1} = \mathcal{T}oldsymbol{x}^\ell + oldsymbol{e}^\ell, \quad \ell \in \mathbb{N}.$$

This class of stochastic updates includes several optimization methods such as *stochastic gradient descent* and 0-*th order methods* [7]–[9], [11]–[13], which are widely used in machine learning applications.

Section V-C will instead provide some convergence results when only random coordinate updates are present.

#### C. Asymptotic convergence results

The mean convergence results of section IV-A can be further used to characterize the *asymptotic, almost sure* convergence of (9), as proved in the following. The results are presented in the contractive case (cf. Assumption 2), but the same argument can be applied in the averaged case as well.

1) Convergence to a neighborhood: We start by showing that the output of (9) asymptotically and almost surely converges to a bounded neighborhood of the fixed point trajectory  $\{\bar{x}^{\ell}\}_{\ell \in \mathbb{N}}$ .

**Corollary 1** (Asymptotic a.s. convergence to neighborhood). Let Assumptions 1 and 2 hold, and let  $\{x^{\ell}\}_{\ell \in \mathbb{N}}$  be the sequence generated by (9). Then it holds that:

$$\limsup_{\ell \to \infty} \| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell} \| \le \sqrt{\frac{\bar{p}}{p}} \frac{\chi \sigma + \nu}{1 - \chi} \qquad a.s..$$
  
See Appendix C-C.

Proof. See Appendix C-C.

2) Exact asymptotic convergence: Corollary 1 shows that in general we can only prove asymptotic convergence to a neighborhood of the fixed point trajectory. The following result proves that a zero asymptotic error can be achieved under the assumptions that the operator is static and the additive error  $e^{\ell}$  vanishing.

**Corollary 2** (Exact a.s. convergence). Let Assumptions 1 and 2 hold, with the difference that

- (i) the errors are vanishing, with  $\mathbb{E}\left[\left\|\boldsymbol{e}^{\ell}\right\|^{2}\right] \leq (\nu^{\ell})^{2}$  and  $\lim_{\ell \to \infty} \nu^{\ell} = 0;$
- (ii) for all  $\ell \in \mathbb{N}$ ,  $\mathcal{T}^{\ell} = \mathcal{T}$ ; we denote by  $\bar{x}$  the unique fixed point of  $\mathcal{T}$ .
- Let  $\{x^{\ell}\}_{\ell \in \mathbb{N}}$  be the sequence generated by (9), then

$$\limsup_{\ell \to \infty} \left\| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}} \right\| = 0 \qquad a.s.$$

Proof. See Appendix C-D.

The result of Corollary 2 is similar to [27], [28], in which the additive errors are assumed to be vanishing (or to be multiplied by a vanishing parameter).

Corollaries 1 and 2 leveraged Markov's inequality to characterize the almost sure behavior of (9) – but these results hold only asymptotically, and do not characterize the transient. The question then is: *can we provide high probability bounds that hold for any*  $\ell \in \mathbb{N}$ ? A first observation is that almost sure convergence cannot be guaranteed during the transient, owing to the ever-present disturbance of the additive errors  $e^{\ell}$ . The next section will instead provide transient bounds that hold with assigned probability.

#### V. HIGH PROBABILITY CONVERGENCE ANALYSIS

In order to derive the high probability bounds for the transient, we introduce the following assumption, which models additive errors as sub-Weibull r.v.s.

**Assumption 4** (Stochastic framework 2). Assumption 1 (i) and (iii) hold, and moreover:

(ii) the additive error  $e^{\ell} \in \mathbb{H}$  is an  $\mathbb{H}$ -valued random vector such that its norm is sub-Weibull, that is  $\|e^{\ell}\|^2 \sim \mathrm{subW}(\theta, \nu^2)$ .

Sub-Weibull r.v.s can be used to model a broad range of additive errors that arise *e.g.* in federated learning applications (cf. section III-A). For example, quantization and compression techniques that result in bounded additive errors can be modeled as sub-Weibulls as discussed in Remark 3.

More importantly, a series of recent papers have highlighted how stochastic gradients present heavy tails in many learning applications [39]–[44], especially involving the training of deep neural networks. The same issues can therefore arise in a federated learning set-up, where each agent computes a local stochastic gradient [45].

## A. Contractive set-up

In the following we will make use of the scaled norm  $|||\boldsymbol{x}|||^2 := \sum_{i=1}^n \frac{1}{p_i} ||x_i||_i^2$  introduced in Appendix C-A.

**Proposition 3** (High prob. – contractive operators). Let Assumptions 2 and 4 hold, and assume that the random coordinate update operators  $\hat{T}^{\ell}$  defined in (9) satisfy

$$\left| \left| \left| \hat{\mathcal{T}}^{\ell} \boldsymbol{x} - \bar{\boldsymbol{x}}^{\ell} \right| \right| \leq \eta(\ell) \left| \left| \left| \boldsymbol{x} - \bar{\boldsymbol{x}}^{\ell} \right| \right| \right|$$
(10)

with  $\eta(\ell)$  i.i.d., and there exist  $\gamma, \chi \in (0,1)$  such that

$$\prod_{h=\ell_1}^{\ell_2} \eta(h) \sim \text{subW}\left(\gamma^{\ell_2-\ell_1+1}, \chi^{\ell_2-\ell_1+1}\right).$$
(11)

Let  $\{x^{\ell}\}_{\ell \in \mathbb{N}}$  be the sequence generated by (9), then for any  $\ell \in \mathbb{N}$ , given  $\delta \in (0, 1)$ , with probability  $1 - \delta$  the following bound holds:

$$\begin{aligned} \left\| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell} \right\| &\leq \sqrt{\frac{\bar{p}}{\underline{p}}} \log^{\theta+1}(2/\delta) c(\theta+1) \times \\ & \times \left( \chi^{\ell} \left\| \boldsymbol{x}^{0} - \bar{\boldsymbol{x}} \right\| + \frac{1-\chi^{\ell}}{1-\chi} (\chi \sigma + \nu) \right). \end{aligned}$$

Proof. See Appendix D-A.

The following sections discuss the assumptions required in Proposition 3, as well as the implications of such convergence result.

1) Assumption (10): In this section we discuss and motivate the use of assumption (10) in Proposition 3. We start by remarking that the assumption is stated in terms of the random coordinate updates operators  $\hat{T}^{\ell}$  defined in (9), and thus depends exclusively on  $\{u_i^{\ell}\}_{i=1}^n$ . In our setup, the underlying



Fig. 2. Random coordinate updates can lead to expansion.

operators  $\mathcal{T}^{\ell}$  are assumed to be contractive, but these random counterparts may not be, as exemplified in Figure 2 where updating only the second coordinate leads to expansion. Assumption (10) thus allows for the whole gamut of possibilities – contraction, non-expansion, expansion – since  $\eta(\ell) \leq 1$ .

While an expansion is possible at any given time  $\ell$ , the results of section IV motivate the second part of the assumption, (11). First, the random operators  $\hat{\mathcal{T}}^{\ell}$  are contractive in mean (w.r.t. the scaled norm  $||| \cdot |||$ ), since inspecting the proof of Proposition 1 in Appendix C-A we see that (cf. (16))

$$\mathbb{E}\left[\left|\left|\left|\hat{\mathcal{T}}^{\ell+1}\boldsymbol{x}^{\ell}-\bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|\right] \leq \chi \mathbb{E}\left[\left|\left|\left|\boldsymbol{x}^{\ell}-\bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|\right]$$

with  $\chi \in (0,1)$ . Thus by the discussion in Remark 2 we expect the second sub-Weibull parameter of  $\prod_{h=\ell_1}^{\ell_2} \eta(h)$  to be in (0,1).

Secondly, Corollary 2 shows that the iterated application of a random coordinate update operator leads to almost sure convergence to the fixed point. This observation thus motivates the choice of a tail parameter for  $\prod_{h=\ell_1}^{\ell_2} \eta(h)$  which decays as the number of iterations increases.

2) Interpretation of high probability convergence results: The high probability bound of Proposition 3 states that, w.p.  $1 - \delta$ , the error satisfies:

$$\begin{aligned} \left\| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell} \right\| &\leq \sqrt{\frac{\bar{p}}{\underline{p}}} \log^{\theta + 1}(2/\delta) c(\theta + 1) \times \\ & \times \left( \chi^{\ell} \left\| \boldsymbol{x}^{0} - \bar{\boldsymbol{x}} \right\| + \frac{1 - \chi^{\ell}}{1 - \chi} (\chi \sigma + \nu) \right). \end{aligned}$$

First of all, we can see that the right-hand side is multiplied by  $\log^{\theta+1}(2/\delta)$ , which grows as  $\delta$  decreases. This implies that the more confidence we want in the bound (which requires  $\delta$  to be smaller), the looser the bound becomes. Intuitively, smaller  $\delta$  requires that we enlarge the bound to include more trajectory realizations  $\{x^{\ell}\}_{\ell \in \mathbb{N}}$ . Similarly, if  $\theta$  grows larger, then the bound becomes looser. This is a consequence of the fact that the larger  $\theta$  is, the heavier the tails of the additive noise are.

We can further observe that the high probability bound – multiplicative factors notwithstanding – shares the structure of the mean bound of Proposition 1. Indeed, both bounds have a first term that depends on the initial condition and decays to zero as  $\ell \to \infty$ , and a second term that bounds the asymptotic distance from the fixed point, and which depends on the additive error and the time-variability of the operators.

3) Sub-Weibull v. Markov's inequality: The fact that the dependence on  $1/\delta$  appears through its logarithm in the bound of Proposition 3 is an important motivation for the use of the proposed sub-Weibull framework. This feature of the bounds ensures that the right-hand side grows relatively slowly when we ask for increasing confidence (that is,  $\delta \rightarrow 0$ ).

Consider instead the following alternative high probability bound, which is based on Markov's inequality: with this approach, the dependence on the right-hand side is with  $1/\delta$ and not its logarithm. Although Markov's inequality holds for a more general class of random variables than sub-Weibull, this result makes it clear that using the sub-Weibull framework allows to derive sharper bounds, while still considering a number of relevant distributions as sub-cases.

**Lemma 9.** Let Assumptions 2 and 4 hold, and let  $\{x^{\ell}\}_{\ell \in \mathbb{N}}$  be the sequence generated by (9). Then with probability  $1 - \delta$ ,  $\delta \in (0, 1)$ , we have that

$$\left\|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}\right\| \leq \frac{1}{\delta} \sqrt{\frac{\bar{p}}{\underline{p}}} \left( \chi^{\ell} \left\|\boldsymbol{x}^{0} - \bar{\boldsymbol{x}}^{0}\right\| + \frac{1 - \chi^{\ell}}{1 - \chi} (\chi \sigma + \nu) \right).$$

*Proof.* By Markov's inequality we know that  $\mathbb{P}\left[\left\|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}\right\| \ge \mathbb{E}\left[\left\|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}\right\| / \delta\right]\right] \le \delta$ . Using the bound of Proposition 1 then yields the thesis.

## B. Averaged set-up

In this section, we provide a convergence analysis for averaged operators. The treatment of this case requires a somewhat more restrictive assumption than (10), owing to the fact that averagedness is a weaker property than contractiveness. Section V-B1 will discuss this assumption in detail.

**Proposition 4** (High prob. – averaged operators). Let Assumptions 3 and 4 hold, and assume that the random coordinate update operators  $\hat{\mathcal{T}}^{\ell+1}$  defined in (9) satisfy

$$\begin{aligned} \left| \left| \left| \hat{\mathcal{T}}^{\ell+1} \boldsymbol{x} - \bar{\boldsymbol{x}}^{\ell+1} \right| \right| \right|^2 \\ \leq \left| \left| \left| \boldsymbol{x} - \bar{\boldsymbol{x}}^{\ell+1} \right| \right| \right|^2 - \frac{1 - \alpha(\ell)}{\alpha(\ell)} \right| \left| \left| (\mathcal{I} - \mathcal{T}^{\ell+1}) \boldsymbol{x} \right| \right| \right|^2 \end{aligned}$$
(12)

where  $\alpha(\ell) \in (0,1)$  almost surely.

0

Let  $\{x^{\ell}\}_{\ell \in \mathbb{N}}$  be the sequence generated by (9), then for any  $\ell \in \mathbb{N}$ , given  $\delta \in (0, 1)$ , with probability  $1 - \delta$  the following bound holds:

$$\frac{1}{\ell+1} \sum_{h=0}^{\ell} \left\| (\mathcal{I} - \mathcal{T}^{h+1}) \boldsymbol{x}^{h} \right\|^{2} \leq \log^{\theta} (2/\delta) c(\theta) \frac{\bar{p}}{\underline{p}} \times \frac{\bar{\alpha}}{1-\bar{\alpha}} \left( \frac{\left\| \boldsymbol{x}^{0} - \bar{\boldsymbol{x}}^{0} \right\|^{2}}{\ell+1} + \sigma^{2} + \nu^{2} + 2\sqrt{\underline{p}} \operatorname{diam}(\mathbb{D})(\sigma+\nu) \right)$$

where  $\bar{\alpha} = \max_{h \in \{0,\dots,\ell\}} \alpha(h)$ .

*Proof.* See Appendix D-B. 
$$\Box$$

1) Assumption (12): As discussed in section V-A1, the presence of random coordinate updates may cause expansion, and assumption (12) is introduced to exclude such behavior. This is needed due to the weaker nature of averagedness. Indeed, we are currently not able to provide high probability bounds for the metric  $\frac{1}{\ell+1} \sum_{h=0}^{\ell} ||(\mathcal{I} - \mathcal{T}^{h+1}) \boldsymbol{x}^h||^2$  when expansion may occur.

Let us now discuss how assumption (12) can be guaranteed in practice. If expansion may occur, then we can model the operators  $\hat{T}^{\ell}$  as being  $\beta(\ell)$ -conically averaged (in the scaled norm  $|||\cdot|||)$  [55]

$$\begin{split} & \left| \left| \left| \hat{\mathcal{T}}^{\ell+1} \boldsymbol{x} - \bar{\boldsymbol{x}}^{\ell+1} \right| \right| \right|^2 \\ & \leq \left| \left| \left| \boldsymbol{x} - \bar{\boldsymbol{x}}^{\ell+1} \right| \right| \right|^2 - \frac{1 - \beta(\ell)}{\beta(\ell)} \right| \left| \left| (\mathcal{I} - \mathcal{T}^{\ell+1}) \boldsymbol{x} \right| \right| \right|^2 \end{split}$$

where  $\beta(\ell) > 0$  almost surely. The case of  $\beta(\ell) > 1$  then models expansion. Inspecting Figure 2, it is clear that expansion – when it occurs – is bounded, which implies that  $\beta(\ell) < \bar{\beta}$  a.s.

Therefore, following [55, Proposition 2.9], we can guarantee averagedness by applying a *relaxed version* of the operators  $\hat{T}^{\ell}$ :

$$(1-\gamma)\mathcal{I} + \gamma \hat{\mathcal{T}}^{\ell}$$

for  $\gamma \in (0, 1/\overline{\beta})$ . This shows that modifying the operator being applied can indeed lead to assumption (12) being guaranteed.

## C. Convergence without additive errors

We conclude this section with two high probability convergence results derived in the absence of additive errors ( $e^{\ell} = 0$ a.s.) and when the operator is static ( $\mathcal{T}^{\ell} = \mathcal{T}, \ell \in \mathbb{N}$ ). We remark that Propositions 3 and 4 do hold in this scenario by setting  $\nu = 0$  and  $\sigma = 0$ ; however, the following results present more refined bounds.

**Proposition 5** (Without additive noise). Let Assumptions 2 and 4 hold, with  $e^{\ell} = 0$  a.s. and  $\mathcal{T}^{\ell} = \mathcal{T}$ , for all  $\ell \in \mathbb{N}$ . Let  $\{x^{\ell}\}_{\ell \in \mathbb{N}}$  be the trajectory generated by (9).

Let  $\epsilon \in (0, p]$  with  $p = \prod_{i=1}^{n} p_i$ , then with probability  $1 - \delta(\epsilon, \ell)$  we have:

$$\left\|oldsymbol{x}^\ell - oldsymbol{ar{x}}
ight\| \leq \zeta^{\ell(p-\epsilon)} \left\|oldsymbol{x}^0 - oldsymbol{ar{x}}
ight\|$$

where  $fix(\mathcal{T}) = \{\bar{x}\}$  and

$$\delta(\epsilon, \ell) = \exp\left(-\ell D(p - \epsilon ||p)\right),$$

$$D(p - \epsilon ||p) = (p - \epsilon) \log\left(1 - \frac{\epsilon}{p}\right) + (1 - p + \epsilon) \log\left(1 + \frac{\epsilon}{1 - p}\right).$$
(13)

*Proof.* Setting  $\sigma = 0$  and  $e^{\ell} = 0$  in (19) we have

$$\left\|oldsymbol{x}^\ell - oldsymbol{ar{x}}
ight\| \leq \zeta^{eta(\ell)} \left\|oldsymbol{x}^0 - oldsymbol{ar{x}}
ight\|$$

where  $\beta(\ell) \sim \mathcal{B}(\ell, p)$ . Using Sanov's theorem [56, Theorem D.3] (in particular its symmetric form in [56, eq. (D.7)]) we know that

$$\mathbb{P}\left[\beta(\ell) \le \ell(p-\epsilon)\right] \le \exp\left(-\ell D(p-\epsilon||p)\right)$$

and, since  $\mathbb{P}\left[\zeta^{\beta(\ell)} \geq \zeta^{\ell(p-\epsilon)}\right] = \mathbb{P}\left[\beta(\ell) \leq \ell(p-\epsilon)\right]$ , the thesis follows.

We observe that in Proposition 5 the convergence rate is characterized by  $\zeta^{p-\epsilon}$ , which is larger than the convergence rate  $\zeta$  achieved in the deterministic case (cf. the discussion in section IV-B1). We further notice that  $\delta(\epsilon, \ell)$  is a decreasing function of  $\ell$ , with  $\delta(\epsilon, \ell) \to 0$  as  $\ell \to \infty$ . This implies that asymptotically, the bound holds with probability  $\lim_{\ell\to\infty} 1 - \delta(\epsilon, \ell) = 1$ . As a consequence, Proposition 5 yields the known fact that *asymptotic almost sure convergence is achieved* [4], [27] – but with the difference that a probabilistic bound is also provided for any  $\ell \in \mathbb{N}$ .

A similar result can also be derived for the averaged case.

**Proposition 6** (Without additive noise – averaged). Let Assumptions 3 and 4 hold, with  $e_i = 0$  a.s. for all  $i \in \{1, ..., n\}$ ; let  $\{x^{\ell}\}_{\ell \in \mathbb{N}}$  be the trajectory generated by (9).

Let  $\epsilon \in (0,p]$  with  $p = \prod_{i=1}^{n}$ , then with probability  $1 - \delta(\epsilon, \ell+1)$  we have:

$$\left\| (\mathcal{I} - \mathcal{T}) \boldsymbol{x}^{\ell} \right\|^{2} \leq \frac{1}{(\ell+1)(p-\epsilon)} \frac{\alpha}{1-\alpha} \left\| \boldsymbol{x}^{0} - \bar{\boldsymbol{x}} \right\|^{2}$$

where  $\delta(\epsilon, \ell + 1)$  and  $D(p - \epsilon || p)$  are defined as in (13). *Proof.* See Appendix D-C.

Recalling again the discussion in section IV-B1, in the deterministic case we know that  $\{\|(\mathcal{I} - \mathcal{T})\boldsymbol{x}^{\ell}\|\}_{\ell \in \mathbb{N}}$  is monotonically decreasing, and such that [21]

$$\left\| (\mathcal{I} - \mathcal{T}) \boldsymbol{x}^{\ell} \right\|^{2} \leq \frac{1}{\ell + 1} \frac{\alpha}{1 - \alpha} \left\| \boldsymbol{x}^{0} - \bar{\boldsymbol{x}} \right\|^{2}$$

Therefore we see that the introduction of random coordinate updates degrades the convergence rate, namely from  $1/(\ell+1)$  to the larger  $1/((\ell+1)(p-\epsilon))$ .

## VI. ILLUSTRATIVE NUMERICAL RESULTS

In this section we illustrate the accordance of the theoretical results provided in this paper with numerical results derived when applying the federated learning Algorithm 1 to solve a logistic regression problem. In particular, we consider problem (6),  $\min_{x_i \in \mathbb{R}^q} \sum_{i=1}^n f_i(x_i)$  s.t.  $x_1 = x_2 = \ldots = x_n$ , where the local data  $\{(a_i^h, b_i^h) \in \mathbb{R}^{1 \times q} \times \{-1, 1\}\}_{h=1}^{m_i}$  define the loss

$$f_i(x) = \sum_{h=1}^{m_i} \log \left( 1 + \exp(-b_i^h a_i^h x) \right) + \frac{\epsilon}{2} \|x\|^2.$$

In our experiments we have n = 10 agents with  $m_i = 150$  data-points each, and the problem size is q = 10. The data are randomly generated using the make\_classification utility of sklearn [57]. The regularization weight is set to  $\epsilon = 0.075$ . Owing to the regularization, the problem is strongly convex, and thus the federated algorithm we apply is contractive (cf. Appendix B).

In the following sections we apply Algorithm 1 to this problem, subject to the challenges of *asynchrony* and *additive errors*. The simulations are implemented using tvopt [58], and all results are averaged over 100 Monte Carlo iterations.

## A. Asynchrony

We start by considering Algorithm 1 in which each agent has a probability p of updating at iteration  $\ell \in \mathbb{N}$ ; therefore we choose  $\{u_i\}_{i=1}^n$  to be i.i.d. Bernoulli of mean p. Table I reports the empirical convergence rate attained for the different update probabilities. Such rate is estimated by computing the slope of the mean error curves  $\{\mathbb{E} [|| \mathbf{x}^{\ell} - \bar{\mathbf{x}} ||]\}_{\ell \in \mathbb{N}}$ . The results show

 TABLE I

 Empirical convergence rate for different update probabilities.

Update probability	Convergence rate
0.1	0.9985
0.2	0.9972
0.3	0.9958
0.4	0.9945
0.5	0.9931
0.6	0.9917
0.7	0.9904
0.8	0.9890
0.9	0.9876
1.0	0.9862

how the larger the probability of each agent performing an update, the smaller the convergence rate. This is in accordance with the results of Proposition 1. We remark that the algorithm still converges to the optimal solution, even if at different rates.

## B. Synthetic additive errors

We evaluate now the effect of additive errors on the performance of Algorithm 1, in a fully synchronous case  $(p_i = 1)$ . In particular, each local update to  $x_i^{\ell}$  (line 5 in the pseudocode) is subject to an additive error  $e_i^{\ell}$  whose components are drawn from the Weibull distribution <sup>1</sup>. With this choice, by Lemma 10 we know that the norm of  $e_i^{\ell}$  is sub-Weibull and thus satisfies Assumption 1(ii).

**Lemma 10** (Norm of sub-Weibull vectors). Let  $e = [e_1, \ldots, e_q]^\top$  be a random vector in  $\mathbb{R}^q$ ,  $q < +\infty$ , such that  $e_i \sim \operatorname{subW}(\theta, \nu)$ ,  $i \in \{1, \ldots, q\}$ . Then the Euclidean norm of e is sub-Weibull with

$$||e||^2 \sim \operatorname{subW}(2\theta, \max\{1, 2^{2\theta}\}q\nu).$$

*Proof.* We want to characterize  $||e||^2 = \sum_{i=1}^{q} e_i^2$  as a sub-Weibull. By Lemma 8 we know that  $e_i^2 \sim \text{subW}(2\theta, \max\{1, 2^{2\theta}\}\nu^2)$ , and it follows that  $\sum_{i=1}^{q} e_i^2 \sim \text{subW}(2\theta, q \max\{1, 2^{2\theta}\}\nu^2)$ .

In Table II we report the mean asymptotic error (computed as the maximum mean error in the last 4/5 of the simulation) for different values of  $\theta$  and  $\nu$ . Additionally, we report the accuracy of the trained model on the training and test sets, computed as the percentage of correctly classified data points. The test set consists of 150 data points generated from the same distribution as the training points. In accordance with

TABLE II Asymptotic error, training and test accuracies for different sub-Weibull distributions of the additive noise.

( heta, u)	As. err.	Training acc. [%]	Test acc. [%]
(1/2, 0.01)	1.305	90.167	86.653
(1/2, 0.1)	17.842	72.757	76.727
(1, 0.01)	1.482	89.720	85.653
(1, 0.1)	20.406	71.939	76.627
(2, 0.01)	3.162	84.945	84.527
(2, 0.1)	43.330	67.924	72.060

the theoretical results (cf. Proposition 3), the heavier the tail (*i.e.* the larger  $\theta$ ) or the larger the scaling parameter  $\nu$ , the larger the error attained by the algorithm. As expected, the accuracy also degrades the larger the errors are. However, it is interesting to notice that in moving from  $\nu = 0.01$  to  $\nu = 0.1$ , the test accuracy becomes better than the training accuracy. This may be due to the fact that larger additive errors improve the generalization capabilities of the model.

#### C. Stochastic gradients

We conclude this section by discussing the performance of Algorithm 1 when the agents use stochastic gradients during the local updates. In particular, the gradients are approximated by using a random subset of B data points. We notice that the larger the batch size, the smaller the asymptotic error, since the additive error we introduce is smaller. This is also illustrated by the training accuracy which roughly increases with B. On

<sup>&</sup>lt;sup>1</sup>Notice that usually the Weibull distribution is characterized by the CDF  $1 - \exp(-(x/\nu)^c)$  [59], and hence is indeed a sub-Weibull, with parameters  $\theta = 1/c$  and  $\nu$  (cf. [32]).

TABLE III Asymptotic error, training and test accuracies for different batch sizes of the stochastic gradients.

B	As. err.	Training acc. [%]	Test acc. [%]
1	1.298	92.251	89.680
5	1.117	92.248	89.753
10	0.970	92.289	89.727
50	0.468	92.619	88.673
100	0.186	92.803	88.667
Full grad.	0	92.733	88.667

the other hand, the test accuracy degrades as the batch size grows larger than 10, signaling that the smaller the errors introduced by the stochastic gradients, the more the trained model is over-fitting the training data.

# APPENDIX A Proof of Lemma 8

*Proof of 1)* The result follows by  $||ax||_k = |a| ||x||_k \le |a|\nu k^{\theta}$ .

*Proof of 2)* For completeness we report the proof provided in [32, Proposition 3]. Using the triangle inequality we write

$$\begin{aligned} \|x_1 + x_2\|_k &\leq \|x_1\|_k + \|x_2\|_k \overset{(i)}{\leq} \nu_1 k^{\theta_1} + \nu_2 k^{\theta_2} \\ &\overset{(ii)}{\leq} (\nu_1 + \nu_2) k^{\max\{\theta_1, \theta_2\}}, \end{aligned}$$

where (i) holds by the assumption that  $x_i$  are sub-Weibull, and (ii) holds since  $k \ge 1$ .

*Proof of 3*) By definition of  $\|\cdot\|_k$  we can write

$$\begin{split} \|xy\|_k &\leq \mathbb{E}\left[|x|^k|y|^k\right]^{1/k} \stackrel{\text{(i)}}{=} \mathbb{E}\left[|x|^k\right]^{1/k} \mathbb{E}\left[|y|^k\right]^{1/k} \\ &\stackrel{\text{(ii)}}{\leq} \nu_1 k^{\theta_1} \nu_2 k^{\theta_2} = \nu_1 \nu_2 k^{\theta_1 + \theta_2} \end{split}$$

where (i) holds by independence and (ii) by sub-Weibull assumption.

*Proof of 4)* By definition of  $\|\cdot\|_k$  we have  $\|x^a\|_k = \mathbb{E}\left[|x^a|^k\right]^{1/k} = \mathbb{E}\left[|x|^{ak}\right]^{1/k}$ . Now, we distinguish two cases: if 0 < a < 1 then by Jensen's inequality we have

$$\mathbb{E}\left[|x|^{ak}\right]^{1/k} \le \left(\mathbb{E}\left[|x|^k\right]^{1/k}\right)^a \le (\nu k^{\theta})^a;$$

instead if  $a \ge 1$  then we can write

$$\mathbb{E}\left[|x|^{ak}\right]^{1/k} = \left(\mathbb{E}\left[|x|^{ak}\right]^{1/ak}\right)^a \stackrel{\text{(i)}}{\leq} \left(\nu(ak)^\theta\right)^a = \nu^a a^{a\theta} k^\theta$$

where (i) holds by the fact that  $ak \ge 1$  and that x is sub-Weibull.

# Appendix B

# Algorithm 1

A. Derivation of the algorithm

Consider (8), and redefine,  $i \in \{1, \ldots, n\}$ 

$$x_i^{\ell+1} = z^\ell - \gamma \nabla f_i(z^\ell) = \mathcal{G}_i z^\ell$$

where  $z^{\ell} = \frac{1}{n} \sum_{i=1}^{n} x_i^{\ell}$  represents the average computed by the coordinator.

Consider now the asynchronous set-up described in section III-A, in which only the subset  $\mathbb{I}^{\ell}$  of the agents is active. In this case, we need to modify the local update as follows

$$x_i^{\ell+1} = \begin{cases} \mathcal{G}_i z^\ell & \text{if } i \in \mathbb{I}^\ell \\ x_i^\ell & \text{otherwise,} \end{cases}$$

so that only the active agents update their state. Since the coordinator receives new information only from the active agents, then we can modify its update as

$$z^{\ell+1} = \frac{1}{n} \left( \sum_{i \in \mathbb{I}^{\ell}} x_i^{\ell+1} + \sum_{i \notin \mathbb{I}^{\ell}} x_i^{\ell} \right).$$

With this update, the coordinator aggregates the new information received from  $\mathbb{I}^{\ell}$  with the most recent information received from the inactive agents (which may have been transmitted several iterations ago, *e.g.* if  $x_i^{\ell+1} = x_i^{\ell} = x_i^{\ell-1}$ ).

#### B. Interpretation as stochastic operator

The goal now is to show that Algorithm 1 derived in the previous section can be interpreted as a stochastic operator that fits into the framework of the paper.

We start with the deterministic algorithm (all agents are always active). Using  $z^{\ell} = \frac{1}{n} \sum_{i=1}^{n} x_i^{\ell}$ , the algorithm is described as

$$x_i^{\ell+1} = \mathcal{G}_i\left(\frac{1}{n}\sum_{j=1}^n x_j^\ell\right).$$

Letting  $\boldsymbol{x} = (x_1, \ldots, x_n)$  the overall algorithm therefore is characterized by the update  $\boldsymbol{x}^{\ell+1} = \mathcal{T} \boldsymbol{x}^{\ell} = (\mathcal{G} \circ \mathcal{A}) \boldsymbol{x}^{\ell}$ with  $\circ$  denoting composition of operators, and where  $\mathcal{G} \boldsymbol{x} = (\mathcal{G}_1 x_1, \ldots, \mathcal{G}_n x_n)$  and

$$\mathcal{A}\boldsymbol{x} = \left(\frac{1}{n}\sum_{i=1}^{n}x_{i}, \dots, \frac{1}{n}\sum_{i=1}^{n}x_{i}\right)$$

Let us now turn to the asynchronous set-up in which only the active agents  $\mathbb{I}^{\ell}$  perform an update. Letting  $u_i^{\ell}$  be a Bernoulli r.v. which is 1 if agent *i* updates at time  $\ell$ , then we can write

$$x_i^{\ell+1} = (1 - u_i^\ell) x_i^\ell + \mathcal{T}_i \boldsymbol{x}^\ell,$$

which fits exactly into the framework of section III-B.

We conclude this section by discussing the properties of  $\mathcal{T}$  as derived from the properties of the problem

$$\min_{x_i \in \mathbb{R}^q} \sum_{i=1}^n f_i(x_i) \quad \text{s.t. } x_1 = x_2 = \ldots = x_n.$$

We remark first that  $\mathcal{A}$  is non-expansive, and we need to characterize the properties of  $\mathcal{G}$ . Assume that the local costs have  $\overline{\lambda}$ -Lipschitz continuous gradients, and that they are  $\underline{\lambda}$ -strongly convex, where we allow  $\underline{\lambda} = 0$  to signify that the costs are convex. Then we have the following cases:

- convex costs (λ = 0): if γ < 2/λ then G is averaged, and hence so is T [21, section 3.3];
- strongly convex costs (λ > 0): if γ < 2/λ then G is contractive and by [60, Lemma 4.11] so is T.</li>

Finally, notice that the fixed point(s) of  $\mathcal{T}$  do not coincide with solutions of the optimization problem, but rather given  $\bar{x} \in \text{fix}(\mathcal{T})$  then  $\bar{z} = \frac{1}{n} \sum_{i=1}^{n} \bar{x}_i$  is a solution to the problem.

# APPENDIX C PROOFS OF SECTION IV

#### A. Proof of Proposition 1

Similarly to e.g. [4], [27], the first step is to define the norm

$$|||\boldsymbol{x}||| := \sqrt{\sum_{i=1}^{n} \frac{1}{p_i} \|x_i\|_i^2}$$

for which, letting  $\bar{p} := \max_i p_i$  and  $p = \min_i p_i$ , it holds that

$$\underline{p}|||\boldsymbol{x}|||^2 \le ||\boldsymbol{x}||^2 \le \bar{p}|||\boldsymbol{x}|||^2.$$
(14)

By the triangle inequality and the fact that  $|||(u_1^{\ell}e_1^{\ell},\ldots,u_n^{\ell}e_n^{\ell})||| \le |||e^{\ell}|||$  we have

$$|||oldsymbol{x}^{\ell+1}-oldsymbol{ar{x}}^{\ell+1}|||\leq \left|\left|\left|\hat{\mathcal{T}}^{\ell+1}oldsymbol{x}^{\ell}-oldsymbol{ar{x}}^{\ell+1}
ight|
ight|+|||oldsymbol{e}^{\ell}|||$$

where  $\hat{\mathcal{T}}^{\ell+1}$  is defined in (9). Taking the expected value, by (14) we have  $\mathbb{E}\left[\left|\left|\left|e^{\ell}\right|\right|\right|\right] \leq \mathbb{E}\left[\left|\left|e^{\ell}\right|\right|\right]/\sqrt{p}$ , and by Assumption 1(ii) and Jensen's inequality we have  $\mathbb{E}\left[\left\|\boldsymbol{e}^{\ell}\right\|\right] \leq \nu$ , hence  $\mathbb{E}\left[\left|\left|\left|e^{\ell}\right|\right|\right|\right] \leq \nu/\sqrt{p}$ . We focus now on the first term. By the law of total expectation and Jensen's inequality we have

$$\begin{split} \mathbb{E}\left[\left|\left|\left|\hat{\mathcal{T}}^{\ell+1}\boldsymbol{x}^{\ell}-\bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|\right] &= \mathbb{E}\left[\mathbb{E}_{\ell}\left[\left|\left|\left|\hat{\mathcal{T}}^{\ell+1}\boldsymbol{x}^{\ell}-\bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|\right]\right]\right] \\ &\leq \mathbb{E}\left[\sqrt{\mathbb{E}_{\ell}\left[\left|\left|\left|\hat{\mathcal{T}}^{\ell+1}\boldsymbol{x}^{\ell}-\bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|^{2}\right]}\right] \end{split}$$

where  $\mathbb{E}_{\ell}[\cdot]$  denotes the expectation conditioned on  $x^{\ell}$ . By linearity of the expected value we can write

$$\mathbb{E}_{\ell} \left[ \left| \left| \left| \hat{\mathcal{T}}^{\ell+1} \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1} \right| \right| \right|^{2} \right] = \\
= \sum_{i=1}^{n} \mathbb{E}_{\ell} \left[ \frac{1}{p_{i}} \left\| \hat{\mathcal{T}}_{i}^{\ell+1} \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}_{i}^{\ell+1} \right\|_{i}^{2} \right] \\
\stackrel{(a)}{=} \sum_{i=1}^{n} \mathbb{E}_{\ell} \left[ \frac{u_{i}^{\ell}}{p_{i}} \left\| \mathcal{T}_{i}^{\ell+1} \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}_{i}^{\ell+1} \right\|_{i}^{2} + \frac{1 - u_{i}^{\ell}}{p_{i}} \left\| \boldsymbol{x}_{i}^{\ell} - \bar{\boldsymbol{x}}_{i}^{\ell+1} \right\|_{i}^{2} \right] \\
\stackrel{(b)}{=} \left\| \mathcal{T}^{\ell+1} \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1} \right\|^{2} + \sum_{i=1}^{n} \frac{1 - p_{i}}{p_{i}} \left\| \boldsymbol{x}_{i}^{\ell} - \bar{\boldsymbol{x}}_{i}^{\ell+1} \right\|_{i}^{2} \quad (15) \\
\stackrel{(c)}{\leq} \zeta^{2} \left\| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1} \right\|^{2} + \sum_{i=1}^{n} \frac{1 - p_{i}}{p_{i}} \left\| \boldsymbol{x}_{i}^{\ell} - \bar{\boldsymbol{x}}_{i}^{\ell+1} \right\|_{i}^{2} \\
= \sum_{i=1}^{n} \frac{1}{p_{i}} (1 - p_{i} + p_{i} \zeta^{2}) \left\| \boldsymbol{x}_{i}^{\ell} - \bar{\boldsymbol{x}}_{i}^{\ell+1} \right\|_{i}^{2} \\
\stackrel{(d)}{\leq} \chi^{2} \left\| \left| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1} \right| \right\|^{2} \quad (16)$$

where (a) holds by definition of  $\hat{\mathcal{T}}^{\ell+1}$ , (b) by the fact that  $u_i^{\ell} \sim Ber(p_i)$ , (c) holds by the contractiveness in Assumption 2(i), and (d) by defining  $\chi = \sqrt{\max_i 1 - p_i + p_i \zeta^2} =$  $\sqrt{1-p+p\zeta^2}.$ 

Putting this bound together with that for  $\mathbb{E} \left[ ||| e^{\ell} ||| \right]$ :

$$\mathbb{E}\left[\left|\left|\left|\boldsymbol{x}^{\ell+1} - \bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|\right] \leq \chi \mathbb{E}\left[\left|\left|\left|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|\right] + \nu/\sqrt{\underline{p}} \\ \leq \chi \mathbb{E}\left[\left|\left|\left|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}\right|\right|\right|\right] + \chi \sigma + \nu/\sqrt{\underline{p}}$$
(17)

where the last inequality holds by triangle inequality and Assumption 2(ii). Iterating and using the geometric sum

$$\mathbb{E}\left[\left|\left|\left|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}\right|\right|\right|\right] \leq \chi^{\ell}\left|\left|\left|\boldsymbol{x}^{0} - \bar{\boldsymbol{x}}^{0}\right|\right|\right| + \frac{1 - \chi^{\ell}}{1 - \chi}(\chi\sigma + \frac{\nu}{\sqrt{\underline{p}}}),$$
  
and the thesis follows by (14).

and the thesis follows by (14).

# B. Proof of Proposition 2

Let  $\langle\!\langle \cdot \rangle\!\rangle$  be the inner product that induces  $|||\cdot|||$ , then we can write

$$\begin{split} &|||\boldsymbol{x}^{\ell+1} - \bar{\boldsymbol{x}}^{\ell+1}|||^{2} = \\ &= \left|||\hat{\mathcal{T}}^{\ell+1}\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1}|||^{2} + \left|||(u_{1}^{\ell}e_{1}^{\ell}, \dots, u_{n}^{\ell}e_{n}^{\ell})|||^{2} + \\ &+ 2\langle\!\langle \hat{\mathcal{T}}^{\ell+1}\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1}, (u_{1}^{\ell}e_{1}^{\ell}, \dots, u_{n}^{\ell}e_{n}^{\ell})\rangle\!\rangle \\ &\leq \left|||\hat{\mathcal{T}}^{\ell+1}\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1}|||^{2} + \left|||\boldsymbol{e}^{\ell}|||^{2} + 2\operatorname{diam}(\mathbb{D})\sqrt{\underline{p}}|||\boldsymbol{e}^{\ell}||| \\ & \text{where the last inequality follows by } \left|||(u_{n}^{\ell}e_{n}^{\ell} - u_{n}^{\ell}e_{n}^{\ell})\rangle\right|| \leq \\ \end{aligned}$$

where the last inequality follows by  $|||(u_1^{\ell}e_1^{\ell},\ldots,u_n^{\ell}e_n^{\ell})||| \leq$  $|||e^{\ell}|||$  and by Cauchy-Schwarz inequality. Using the law of total expectation we can write  $\mathbb{E}\left[\left|\left|\left|\hat{\mathcal{T}}^{\ell+1}\boldsymbol{x}^{\ell}-\bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|^{2}\right]=$  $\mathbb{E}\left[\mathbb{E}_{\ell}\left[\left|\left|\left|\hat{\mathcal{T}}^{\ell+1}\boldsymbol{x}^{\ell}-\bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|^{2}\right]\right] \text{ where recall that } \mathbb{E}_{\ell}\left[\cdot\right] \text{ is }$ conditioned on  $x^{\ell}$ . Following the steps leading to (15) we have

$$\mathbb{E}_{\ell} \left[ \left| \left| \left| \hat{\mathcal{T}}^{\ell+1} \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1} \right| \right| \right|^{2} \right] = \\ = \left\| \mathcal{T}^{\ell+1} \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1} \right\|^{2} + \sum_{i=1}^{n} \frac{1-p_{i}}{p_{i}} \left\| \boldsymbol{x}_{i}^{\ell} - \bar{\boldsymbol{x}}_{i}^{\ell+1} \right\|_{i}^{2} \\ \stackrel{(a)}{\leq} \left\| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1} \right\|^{2} - \frac{1-\alpha}{\alpha} \left\| (\mathcal{I} - \mathcal{T}^{\ell+1}) \boldsymbol{x}^{\ell} \right\|^{2} + \\ + \sum_{i=1}^{n} \frac{1-p_{i}}{p_{i}} \left\| \boldsymbol{x}_{i}^{\ell} - \bar{\boldsymbol{x}}_{i}^{\ell+1} \right\|_{i}^{2} \\ = \left\| \left\| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1} \right\| \right\|^{2} - \frac{1-\alpha}{\alpha} \left\| (\mathcal{I} - \mathcal{T}^{\ell+1}) \boldsymbol{x}^{\ell} \right\|^{2}$$

where (a) follows by averagedness in Assumption 3(i). Let  $\bar{x}^{\ell} \in \operatorname{fix}(\mathcal{T}^{\ell})$ , then

$$\begin{aligned} \left| \left| \left| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1} \right| \right| \right|^2 &= \left| \left| \left| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell} \right| \right| \right|^2 + \left| \left| \left| \bar{\boldsymbol{x}}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1} \right| \right| \right|^2 + \\ &+ 2 \langle \left\langle \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}, \bar{\boldsymbol{x}}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1} \right\rangle \rangle \\ &\leq \left| \left| \left| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell} \right| \right| \right|^2 + \sigma^2 / \underline{p} + 2 \operatorname{diam}(\mathbb{D}) \sigma / \sqrt{\underline{p}} \end{aligned} \end{aligned}$$

where we used Assumption 3(ii), (14), and the fact that  $\mathbb{D}$  is bounded. Putting all these results together yields

$$\mathbb{E}\left[\left|\left|\left|\boldsymbol{x}^{\ell+1} - \bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|^{2}\right] \leq \mathbb{E}\left[\left|\left|\left|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}\right|\right|\right|^{2}\right] + \sigma^{2}/\underline{p} + 2\operatorname{diam}(\mathbb{D})\sigma/\sqrt{\underline{p}} + \mathbb{E}\left[\left|\left|\left|\boldsymbol{e}^{\ell}\right|\right|\right|^{2} + 2\operatorname{diam}(\mathbb{D})\right|\right|\left|\boldsymbol{e}^{\ell}\right|\right|\right|\right] + -\frac{1-\alpha}{\alpha}\left\|(\mathcal{I} - \mathcal{T}^{\ell+1})\boldsymbol{x}^{\ell}\right\|^{2} \leq \mathbb{E}\left[\left|\left|\left|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}\right|\right|\right|^{2}\right] + d - \frac{1-\alpha}{\alpha}\left\|(\mathcal{I} - \mathcal{T}^{\ell+1})\boldsymbol{x}^{\ell}\right\|^{2}$$
(18)

where the bound

$$d := \frac{1}{\underline{p}} \left( \sigma^2 + \nu^2 \right) + \frac{2 \operatorname{diam}(\mathbb{D})}{\sqrt{\underline{p}}} \left( \sigma + \nu \right)$$

was derived using Assumption 1 (ii) and Jensen's inequality.

Reordering (18) and averaging over time yields

$$egin{aligned} &rac{1}{\ell+1}\sum_{h=0}^{\ell}\mathbb{E}\left[\left\|(\mathcal{I}-\mathcal{T}^{h+1})oldsymbol{x}^{h}
ight\|^{2}
ight]\leq\ &\leqrac{lpha}{1-lpha}\left(rac{1}{\ell+1}|||oldsymbol{x}^{0}-oldsymbol{ar{x}}^{0}|||^{2}+d
ight) \end{aligned}$$

where we used the telescopic sum and removed the negative term  $-\mathbb{E}\left[\left|\left|\left|\boldsymbol{x}^{\ell+1}-\bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|^2\right]$ . The thesis follows by (14) and  $\sqrt{\underline{p}} \leq 1$ .

## C. Proof of Corollary 1

By Proposition 1 we know that

$$\mathbb{E}\left[\left\|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}\right\|\right] \leq \sqrt{\frac{\bar{p}}{\underline{p}}} \left(\chi^{\ell} \left\|\boldsymbol{x}^{0} - \bar{\boldsymbol{x}}^{0}\right\| + \frac{1 - \chi^{\ell}}{1 - \chi} (\chi \sigma + \nu)\right)$$

and, defining the random variable

$$y^{\ell} := \max\left\{0, \sqrt{\frac{\underline{p}}{\bar{p}}} \left\| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell} \right\| - \frac{1 - \chi^{\ell}}{1 - \chi} (\chi \sigma + \nu) 
ight\},$$

this fact implies  $\mathbb{E}\left[y^{\ell}\right] \leq \chi^{\ell} \left\|\boldsymbol{x}^{0} - \bar{\boldsymbol{x}}^{0}\right\|.$ 

By Markov's inequality then we have that, for any  $\epsilon > 0$ :  $\mathbb{P}\left[y^{\ell} \ge \epsilon\right] \le \mathbb{E}\left[y^{\ell}\right]/\epsilon \le \chi^{\ell} \left\|\boldsymbol{x}^{0} - \bar{\boldsymbol{x}}^{0}\right\|/\epsilon$ , and summing over time yields

$$\sum_{\ell=0}^{\infty} \mathbb{P}\left[y^{\ell} \geq \epsilon\right] \leq \frac{1}{1-\chi} \frac{\left\|\boldsymbol{x}^0 - \bar{\boldsymbol{x}}^0\right\|}{\epsilon} < \infty.$$

By the Borel-Cantelli lemma, this fact implies that almost surely  $\limsup_{\ell \to \infty} y^{\ell} \leq \epsilon$ , and, since the inequality holds for any  $\epsilon > 0$  the thesis is proved.

## D. Proof of Corollary 2

By assumption (ii) the operator is static, which implies that  $\sigma = 0$ ; hereafter  $\bar{x}$  denotes the unique fixed point of  $\mathcal{T}$ .

Following the same derivation leading to (17) yields

$$\mathbb{E}\left[\left|\left|\left|\boldsymbol{x}^{\ell+1} - \bar{\boldsymbol{x}}\right|\right|\right|\right] \leq \chi \mathbb{E}\left[\left|\left|\left|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}\right|\right|\right|\right] + \nu^{\ell}/\sqrt{\underline{p}}$$

with the difference that now the right-most term is a function of  $\ell$  as well. Iterating and using (14) we have

$$\mathbb{E}\left[\left\|\boldsymbol{x}^{\ell}-\bar{\boldsymbol{x}}\right\|\right] \leq \sqrt{\frac{\bar{p}}{\bar{p}}}\left(\chi^{\ell}\left\|\boldsymbol{x}^{0}-\bar{\boldsymbol{x}}\right\| + \sum_{h=0}^{\ell-1}\chi^{\ell-h-1}\nu^{h}\right).$$

Similarly to Corollary 1, by Markov's inequality we have

$$\lim_{\ell \to \infty} \left\| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}} \right\| \le \sqrt{\frac{\bar{p}}{\underline{p}}} \lim_{\ell \to \infty} \sum_{h=0}^{\ell-1} \chi^{\ell-h-1} \nu^h = 0$$

where the right-hand side is equal to zero since  $\{\nu^{\ell}\}_{\ell \in \mathbb{N}}$  is summable and we can apply [61, Lemma 3.1(a)] because  $\chi \in (0, 1)$ .

## APPENDIX D Proofs of Section V

## A. Proof of Proposition 3

We start by deriving the following chain of inequalitites:

$$\begin{split} |||\boldsymbol{x}^{\ell+1} - \bar{\boldsymbol{x}}^{\ell+1}||| &\leq ||| \hat{\mathcal{T}}^{\ell+1} \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1}||| + |||\boldsymbol{e}^{\ell}||| \\ &\leq \eta(\ell) |||\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1}||| + |||\boldsymbol{e}^{\ell}||| \\ &\leq \eta(\ell) |||\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}||| + \eta(\ell) |||\bar{\boldsymbol{x}}^{\ell+1} - \bar{\boldsymbol{x}}^{\ell}||| + |||\boldsymbol{e}^{\ell}||| \\ &\leq \eta(\ell) |||\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}||| + \eta(\ell) \frac{\sigma}{\sqrt{p}} + \frac{||\boldsymbol{e}^{\ell}||}{\sqrt{p}} \end{split}$$

where (i) follows by the triangle inequality and the fact that  $|||(\ldots, u_i^{\ell} e_i^{\ell}, \ldots)||| \leq |||e^{\ell}|||$ ; (ii) by assumption (10); (iii) by triangle inequality, and (iv) by (14) and Assumption 2(ii). Iterating this inequality and using (14) again, we derive

$$|||\boldsymbol{x}^{\ell+1} - \bar{\boldsymbol{x}}^{\ell+1}||| \leq \frac{1}{\sqrt{\underline{p}}} \left( \prod_{h=0}^{\ell} \eta(h) \| \boldsymbol{x}^{0} - \bar{\boldsymbol{x}}^{0} \| + \sigma \sum_{h=0}^{\ell} \prod_{j=h}^{\ell} \eta(j) + \sum_{h=0}^{\ell} \| \boldsymbol{e}^{h} \| \prod_{j=h+1}^{\ell} \eta(j) \right).$$
(19)

The goal now is to show that the right hand side of (19) is sub-Weibull. First of all, by (10), Lemma 8, and simplifying, we get that

$$\sum_{h=0}^{\ell} \prod_{j=h}^{\ell} \eta(j) \sim \operatorname{subW}\left(\gamma, \chi \frac{1-\chi^{\ell+1}}{1-\chi}\right).$$

Similarly, using the fact that  $\|\boldsymbol{e}^{\ell}\|^2 \sim \operatorname{subW}(\theta, \nu)$  and hence  $\|\boldsymbol{e}^{\ell}\| \sim \operatorname{subW}(\theta/2, \nu)$  by Lemma 8, we get

$$\sum_{h=0}^{\ell} \left\| \boldsymbol{e}^h \right\| \prod_{j=h+1}^{\ell} \eta(j) \sim \operatorname{subW} \left( \theta + 1, \nu \frac{1 - \chi^{\ell+1}}{1 - \chi} \right).$$

Combining these results, and using Lemma 8 again, yields

(19) 
$$\sim \frac{1}{\sqrt{\underline{p}}} \operatorname{subW} \left( \theta + 1, \chi^{\ell+1} \| \boldsymbol{x}^0 - \bar{\boldsymbol{x}}^0 \| + \frac{1 - \chi^{\ell+1}}{1 - \chi} (\chi \sigma + \nu) \right).$$

Using (14) and Lemma 6 then yields the thesis.

## B. Proof of Proposition 4

Using the fact that  $|||(\ldots, u_i^{\ell} e_i^{\ell}, \ldots)||| \leq |||e^{\ell}|||$ , and Cauchy-Schwarz inequality we can write

$$\begin{split} &|||\boldsymbol{x}^{\ell+1} - \bar{\boldsymbol{x}}^{\ell+1}|||^2 \leq \\ &\leq \left|\left|\left|\hat{\mathcal{T}}^{\ell+1}\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|^2 + \left|\left|\left|\boldsymbol{e}^{\ell}\right|\right|\right|^2 + 2\operatorname{diam}(\mathbb{D})|||\boldsymbol{e}^{\ell}|||. \end{split}$$

But by assumption, the random coordinate update operator  $\hat{\mathcal{T}}^{\ell+1}$  is  $\alpha(\ell)$ -averaged, hence

$$||\boldsymbol{x}^{\ell+1} - \bar{\boldsymbol{x}}^{\ell+1}|||^{2} \leq |||\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1}|||^{2} + (20) - \frac{1 - \alpha(\ell)}{\alpha(\ell)}|||(\mathcal{I} - \mathcal{T}^{\ell+1})\boldsymbol{x}^{\ell}|||^{2} + |||\boldsymbol{e}^{\ell}|||^{2} + 2\operatorname{diam}(\mathbb{D})|||\boldsymbol{e}^{\ell}|||.$$

Now, by Assumption 3, Cauchy-Schwarz inequality, and (14) we have that

$$\left|\left|\left|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell+1}\right|\right|\right|^{2} \leq \left|\left|\left|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell}\right|\right|\right|^{2} + \frac{\sigma^{2}}{\underline{p}} + \frac{2\operatorname{diam}(\mathbb{D})\sigma}{\sqrt{\underline{p}}}$$

and using this fact into (20) yields

$$\begin{aligned} \left| \left| \left| \boldsymbol{x}^{\ell+1} - \bar{\boldsymbol{x}}^{\ell+1} \right| \right| \right|^2 &\leq \left| \left| \left| \boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}^{\ell} \right| \right| \right|^2 + \\ &- \frac{1 - \alpha(\ell)}{\alpha(\ell)} \left| \left| \left| \left( \mathcal{I} - \mathcal{T}^{\ell+1} \right) \boldsymbol{x}^{\ell} \right| \right| \right|^2 + \\ &+ \frac{\left\| \boldsymbol{e}^{\ell} \right\|^2 + \sigma^2}{\underline{p}} + 2 \operatorname{diam}(\mathbb{D}) \frac{\left\| \boldsymbol{e}^{\ell} \right\| + \sigma}{\sqrt{\underline{p}}}. \end{aligned}$$

Rearranging (21), averaging over time, and using the telescopic sum we have

$$\begin{aligned} &\frac{1}{\ell+1}\sum_{h=0}^{\ell}\frac{1-\alpha(h)}{\alpha(h)}|||(\mathcal{I}-\mathcal{T}^{h+1})\boldsymbol{x}^{h}|||^{2} \leq \\ &\leq \frac{1}{\ell+1}\left(\frac{\|\boldsymbol{x}^{0}-\bar{\boldsymbol{x}}^{0}\|^{2}}{\underline{p}} + \right. \\ &+ \sum_{h=0}^{\ell}\frac{\|\boldsymbol{e}^{h}\|^{2}+\sigma^{2}}{\underline{p}} + 2\operatorname{diam}(\mathbb{D})\frac{\|\boldsymbol{e}^{h}\|+\sigma}{\sqrt{\underline{p}}}\right). \end{aligned}$$

Now defining  $\bar{\alpha} = \max_{h \in \{0,...,\ell\}} \alpha(h)$ , and using (14) we have that

$$\frac{1}{\ell+1} \sum_{h=0}^{\ell} \left\| (\mathcal{I} - \mathcal{T}^{h+1}) \boldsymbol{x}^{h} \right\|^{2} \leq (22)$$

$$\leq \frac{\bar{p}}{\bar{p}} \frac{1}{\ell+1} \frac{\bar{\alpha}}{1-\bar{\alpha}} \left( \left\| \boldsymbol{x}^{0} - \bar{\boldsymbol{x}}^{0} \right\|^{2} + \sum_{h=0}^{\ell} \left\| \boldsymbol{e}^{h} \right\|^{2} + \sigma^{2} + 2 \operatorname{diam}(\mathbb{D}) \sqrt{\bar{p}} \left( \left\| \boldsymbol{e}^{h} \right\| + \sigma \right) \right).$$

Finally, by Assumption 4 and the properties of sub-Weibull r.v.s (cf. Lemma 8) we know that the right hand side of (22) is a sub-Weibull with parameters  $\theta$  and

$$\sigma^2 + \nu^2 + 2 \operatorname{diam}(\mathbb{D}) \sqrt{\underline{p}}(\sigma + \nu)$$

and by Lemma 6 this yields the thesis.

### C. Proof of Proposition 6

Setting  $\sigma = 0$  and  $\|e^{\ell}\| = 0$  in (20) yields

$$\left\|\boldsymbol{x}^{\ell+1} - \bar{\boldsymbol{x}}\right\|^{2} \leq \left\|\boldsymbol{x}^{\ell} - \bar{\boldsymbol{x}}\right\|^{2} - u^{\ell} \frac{1-\alpha}{\alpha} \left\| (\boldsymbol{\mathcal{I}} - \boldsymbol{\mathcal{T}}) \boldsymbol{x}^{\ell} \right\|^{2}, \quad (23)$$

which implies  $\|\boldsymbol{x}^{\ell+1} - \bar{\boldsymbol{x}}\|^2 \leq \|\boldsymbol{x}^\ell - \bar{\boldsymbol{x}}\|^2$ , that is, the operator is *stochastic Fejér monotone* [27], [28]. This means that the fixed point residual  $\{\|(\mathcal{I} - \mathcal{T})\boldsymbol{x}^\ell\|\}_{\ell \in \mathbb{N}|u^\ell=1}$  is a.s. monotonically decreasing [27, Theorem 3.2] (cf. in particular [27, (3.3)]).

Therefore, summing (23) over time and using Fejér monotonicity we can write

$$egin{split} eta(\ell) \left\| (\mathcal{I} - \mathcal{T}) oldsymbol{x}^\ell 
ight\|^2 &\leq \sum_{h=0}^\ell u^h \left\| (\mathcal{I} - \mathcal{T}) oldsymbol{x}^h 
ight\|^2 \ &\leq rac{lpha}{1-lpha} \left\| oldsymbol{x}^0 - oldsymbol{ar{x}} 
ight\|^2 \end{split}$$

where we used the fact that the sequence  $\{\|\boldsymbol{x}^{\ell+1} - \boldsymbol{x}^{\ell}\|\}_{\ell \in \mathbb{N}|u^{\ell}=1}$  has  $\beta(\ell)$  non-zero terms. Dividing by  $\beta(\ell)$  on both sides we get

$$\left\| \boldsymbol{x}^{\ell+1} - \boldsymbol{x}^{\ell} \right\|^2 \leq rac{1}{eta(\ell)} rac{lpha}{1-lpha} \left\| \boldsymbol{x}^0 - ar{\boldsymbol{x}} \right\|^2.$$

We have now the following fact

$$\mathbb{P}\left[\frac{1}{\beta(\ell)} \left\|\boldsymbol{x}^{0} - \bar{\boldsymbol{x}}\right\|^{2} \ge \frac{1}{(p-\epsilon)(\ell+1)} \left\|\boldsymbol{x}^{0} - \bar{\boldsymbol{x}}\right\|^{2}\right] = \mathbb{P}\left[\beta(\ell) \le (\ell+1)(p-\epsilon)\right] \stackrel{(i)}{\le} \exp\left(-(\ell+1)D(p-\epsilon)\right]$$

where (i) holds by Sanov's theorem [56, Theorem D.3] (cf. [56, eq. (D.7)]), and the thesis follows.

#### REFERENCES

- T. Li, A. K. Sahu, A. Talwalkar, and V. Smith, "Federated Learning: Challenges, Methods, and Future Directions," *IEEE Signal Processing Magazine*, vol. 37, no. 3, pp. 50–60, May 2020.
- [2] T. Gafni, N. Shlezinger, K. Cohen, Y. C. Eldar, and H. V. Poor, "Federated Learning: A signal processing perspective," *IEEE Signal Processing Magazine*, vol. 39, no. 3, pp. 14–41, May 2022.
- [3] Z. Peng, T. Wu, Y. Xu, M. Yan, and W. Yin, "Coordinate Friendly Structures, Algorithms and Applications," *Annals of Mathematical Sciences* and Applications, vol. 1, no. 1, pp. 57–119, 2016.
- [4] P. Bianchi, W. Hachem, and F. Iutzeler, "A Coordinate Descent Primal-Dual Algorithm and Application to Distributed Asynchronous Optimization," *IEEE Transactions on Automatic Control*, vol. 61, no. 10, pp. 2947–2957, 2016.
- [5] N. Bastianello, R. Carli, L. Schenato, and M. Todescato, "Asynchronous distributed optimization over lossy networks via relaxed admm: Stability and linear convergence," *IEEE Transactions on Automatic Control*, vol. 66, no. 6, pp. 2620–2635, 2021.
- [6] S. Salzo and S. Villa, "Parallel random block-coordinate forward-backward algorithm: a unified convergence analysis," *Mathematical Programming*, Apr. 2021.
- [7] R. Dixit, A. S. Bedi, R. Tripathi, and K. Rajawat, "Online Learning with Inexact Proximal Online Gradient Descent Algorithms," *IEEE Transactions on Signal Processing*, vol. 67, no. 5, pp. 1338 – 1352, 2019.
- [8] S. Liu, P.-Y. Chen, B. Kailkhura, G. Zhang, A. O. Hero III, and P. K. Varshney, "A Primer on Zeroth-Order Optimization in Signal Processing and Machine Learning: Principals, Recent Advances, and Applications," *IEEE Signal Processing Magazine*, vol. 37, no. 5, pp. 43–54, Sep. 2020.
- [9] L. Bottou, F. E. Curtis, and J. Nocedal, "Optimization methods for largescale machine learning," *Siam Review*, vol. 60, no. 2, pp. 223–311, 2018.
- [10] A. Simonetto, E. Dall'Anese, J. Monteil, and A. Bernstein, "Personalized optimization with user's feedback," *Automatica*, vol. 131, p. 109767, Sep. 2021.
- [11] J. C. Duchi, M. I. Jordan, M. J. Wainwright, and A. Wibisono, "Optimal Rates for Zero-Order Convex Optimization: The Power of Two Function Evaluations," *IEEE Transactions on Information Theory*, vol. 61, no. 5, pp. 2788–2806, May 2015.
- [12] Y. Nesterov and V. Spokoiny, "Random Gradient-Free Minimization of Convex Functions," *Foundations of Computational Mathematics*, vol. 17, no. 2, pp. 527–566, Apr. 2017.
- [13] A. S. Berahas, L. Cao, K. Choromanski, and K. Scheinberg, "A Theoretical and Empirical Comparison of Gradient Approximations in Derivative-Free Optimization," *Foundations of Computational Mathematics*, vol. 22, no. 2, pp. 507–560, Apr. 2022.
- [14] S. Shalev-Shwartz, "Online Learning and Online Convex Optimization," *Foundations and Trends in Machine Learning*, vol. 4, no. 2, pp. 107– 194, 2011.
- [15] E. Dall'Anese, A. Simonetto, and A. Bernstein, "On the Convergence of the Inexact Running Krasnosel'skii–Mann Method," *IEEE Control Systems Letters*, vol. 3, no. 3, pp. 613–618, Jul. 2019.
- [16] A. Simonetto, E. Dall'Anese, S. Paternain, G. Leus, and G. B. Giannakis, "Time-Varying Convex Optimization: Time-Structured Algorithms and Applications," *Proceedings of the IEEE*, vol. 108, no. 11, pp. 2032–2048, Nov. 2020.

- [17] A. Hauswirth, S. Bolognani, G. Hug, and F. Dorfler, "Timescale Separation in Autonomous Optimization," *IEEE Transactions on Automatic Control*, vol. 66, no. 2, pp. 611–624, 2021.
- [18] E. K. Ryu and S. Boyd, "A primer on monotone operator methods," *Applied and Computational Mathematics*, vol. 15, no. 1, pp. 3–43, 2016.
- [19] P. L. Combettes, "Monotone operator theory in convex optimization," *Mathematical Programming*, vol. 170, no. 1, pp. 177–206, Jul. 2018.
- [20] H. H. Bauschke and P. L. Combettes, *Convex analysis and monotone operator theory in Hilbert spaces*, 2nd ed., ser. CMS books in mathematics. Cham: Springer, 2017.
- [21] D. Davis and W. Yin, "Convergence Rate Analysis of Several Splitting Schemes," in *Splitting Methods in Communication, Imaging, Science, and Engineering*, R. Glowinski, S. J. Osher, and W. Yin, Eds. Cham: Springer International Publishing, 2016, pp. 115–163.
- [22] A. Simonetto, "Time-Varying Convex Optimization via Time-Varying Averaged Operators," arXiv:1704.07338 [math], Apr. 2017. [Online]. Available: http://arxiv.org/abs/1704.07338
- [23] A. Jadbabaie, A. Rakhlin, S. Shahrampour, and K. Sridharan, "Online Optimization: Competing with Dynamic Comparators," in *PMLR*, no. 38, 2015, pp. 398 – 406.
- [24] A. Beck and M. Teboulle, "A Fast Iterative Shrinkage-Thresholding Algorithm for Linear Inverse Problems," *SIAM Journal on Imaging Sciences*, vol. 2, no. 1, pp. 183–202, Jan. 2009.
- [25] D. Davis and W. Yin, "A Three-Operator Splitting Scheme and its Optimization Applications," *Set-Valued and Variational Analysis*, vol. 25, no. 4, pp. 829–858, Dec. 2017.
- [26] A. Themelis and P. Patrinos, "SuperMann: A Superlinearly Convergent Algorithm for Finding Fixed Points of Nonexpansive Operators," *IEEE Transactions on Automatic Control*, vol. 64, no. 12, pp. 4875–4890, Dec. 2019.
- [27] P. L. Combettes and J.-C. Pesquet, "Stochastic Quasi-Fejér Block-Coordinate Fixed Point Iterations with Random Sweeping," *SIAM Journal on Optimization*, vol. 25, no. 2, pp. 1221–1248, 2015.
- [28] —, "Stochastic quasi-Fejér block-coordinate fixed point iterations with random sweeping II: mean-square and linear convergence," *Mathematical Programming*, vol. 174, no. 1-2, pp. 433–451, 2019.
- [29] V. Berinde, *Iterative approximation of fixed points*, 2nd ed., ser. Lecture notes in mathematics. Berlin; New York: Springer, 2007, no. 1912.
- [30] Z. Peng, Y. Xu, M. Yan, and W. Yin, "ARock: an Algorithmic Framework for Asynchronous Parallel Coordinate Updates," *SIAM Journal on Scientific Computing*, vol. 38, no. 5, pp. A2851–A2879, Jan. 2016.
- [31] V. S. Borkar, "A concentration bound for contractive stochastic approximation," Systems & Control Letters, vol. 153, p. 104947, 2021.
- [32] M. Vladimirova, S. Girard, H. Nguyen, and J. Arbel, "Sub-Weibull distributions: Generalizing sub-Gaussian and sub-Exponential properties to heavier tailed distributions," *Stat*, vol. 9, no. 1, Jan. 2020.
- [33] H. Zhang and Song Xi Chen, "Concentration Inequalities for Statistical Inference," *Communications in Mathematical Research*, vol. 37, no. 1, pp. 1–85, Jun. 2021.
- [34] A. K. Kuchibhotla and A. Chakrabortty, "Moving beyond sub-Gaussianity in high-dimensional statistics: Applications in covariance estimation and linear regression," *Information and Inference: A Journal* of the IMA, vol. 11, no. 4, pp. 1389–1456, 2022.
- [35] K. C. Wong, Z. Li, and A. Tewari, "Lasso guarantees for β-mixing heavy-tailed time series," *Annals of Statistics*, vol. 48, no. 2, pp. 1124– 1142, 2020.
- [36] S. Boucheron, G. Lugosi, and P. Massart, *Concentration inequalities:* a nonasymptotic theory of independence, 1st ed. Oxford: Oxford University Press, 2013.
- [37] R. Vershynin, High-Dimensional Probability: An Introduction with Applications in Data Science, 1st ed. Cambridge University Press, Sep. 2018.
- [38] M. Vladimirova, J. Verbeek, P. Mesejo, and J. Arbel, "Understanding Priors in Bayesian Neural Networks at the Unit Level," in *Proceedings* of the 36th International Conference on Machine Learning, ser. Proceedings of Machine Learning Research, K. Chaudhuri and R. Salakhutdinov, Eds., vol. 97. PMLR, Jun. 2019, pp. 6458–6467.
- [39] M. Gürbüzbalaban, U. Simsekli, and L. Zhu, "The Heavy-Tail Phenomenon in SGD," arXiv:2006.04740 [cs, math, stat], Feb. 2021. [Online]. Available: http://arxiv.org/abs/2006.04740
- [40] W. Zhu, Z. Lou, and W. B. Wu, "Beyond Sub-Gaussian Noises: Sharp Concentration Analysis for Stochastic Gradient Descent," *Journal of Machine Learning Research*, vol. 23, no. 46, pp. 1–22, 2022.
- [41] E. Gorbunov, M. Danilova, and A. Gasnikov, "Stochastic optimization with heavy-tailed noise via accelerated gradient clipping," in *Advances* in *Neural Information Processing Systems*, H. Larochelle, M. Ranzato,

R. Hadsell, M. Balcan, and H. Lin, Eds., vol. 33. Curran Associates, Inc., 2020, pp. 15 042–15 053.

- [42] A. Cutkosky and H. Mehta, "High-probability bounds for non-convex stochastic optimization with heavy tails," in *Advances in Neural Information Processing Systems*, M. Ranzato, A. Beygelzimer, Y. Dauphin, P. Liang, and J. W. Vaughan, Eds., vol. 34. Curran Associates, Inc., 2021, pp. 4883–4895.
- [43] T. D. Nguyen, T. H. Nguyen, A. Ene, and H. L. Nguyen, "High Probability Convergence of Clipped-SGD Under Heavy-tailed Noise," Feb. 2023, arXiv:2302.05437 [math]. [Online]. Available: http://arxiv.org/abs/2302.05437
- [44] D. Jakovetić, D. Bajović, A. K. Sahu, S. Kar, N. Milosevic, and D. Stamenković, "Nonlinear Gradient Mappings and Stochastic Optimization: A General Framework with Applications to Heavy-Tail Noise," *SIAM Journal on Optimization*, pp. 394–423, Jun. 2023.
- [45] M. Gurbuzbalaban, Y. Hu, U. Simsekli, K. Yuan, and L. Zhu, "Heavy-Tail Phenomenon in Decentralized SGD," arXiv:2205.06689, May 2022.
- [46] A. Cegielski, Iterative methods for fixed point problems in Hilbert spaces, ser. Lecture notes in mathematics. New York: Springer Verlag, 2012, no. 2057.
- [47] Z. Zhao, Y. Mao, Y. Liu, L. Song, Y. Ouyang, X. Chen, and W. Ding, "Towards efficient communications in federated learning: A contemporary survey," *Journal of the Franklin Institute*, Jan. 2023.
- [48] H. Yuan and T. Ma, "Federated Accelerated Stochastic Gradient Descent," in Advances in Neural Information Processing Systems, H. Larochelle, M. Ranzato, R. Hadsell, M. F. Balcan, and H. Lin, Eds., vol. 33. Curran Associates, Inc., 2020, pp. 5332–5344.
- [49] S. P. Karimireddy, S. Kale, M. Mohri, S. Reddi, S. Stich, and A. T. Suresh, "SCAFFOLD: Stochastic Controlled Averaging for Federated Learning," in *Proceedings of the 37th International Conference on Machine Learning*, ser. Proceedings of Machine Learning Research, H. D. III and A. Singh, Eds., vol. 119. PMLR, Jul. 2020, pp. 5132–5143.
- [50] S. Ghadimi and G. Lan, "Stochastic first-and zeroth-order methods for nonconvex stochastic programming," *SIAM Journal on Optimization*, vol. 23, no. 4, pp. 2341–2368, 2013.
- [51] A. Reisizadeh, A. Mokhtari, H. Hassani, and R. Pedarsani, "An Exact Quantized Decentralized Gradient Descent Algorithm," *IEEE Transactions on Signal Processing*, vol. 67, no. 19, pp. 4934–4947, Oct. 2019.
- [52] E. Dall'Anese, A. Simonetto, S. Becker, and L. Madden, "Optimization and Learning With Information Streams: Time-varying algorithms and applications," *IEEE Signal Processing Magazine*, vol. 37, no. 3, pp. 71– 83, May 2020.
- [53] A. Mokhtari, S. Shahrampour, A. Jadbabaie, and A. Ribeiro, "Online optimization in dynamic environments: Improved regret rates for strongly convex problems," in 2016 IEEE 55th Conference on Decision and Control (CDC), Dec. 2016, pp. 7195–7201.
- [54] N. Hallak, P. Mertikopoulos, and V. Cevher, "Regret Minimization in Stochastic Non-Convex Learning via a Proximal-Gradient Approach," in *Proceedings of the 38th International Conference on Machine Learning*, ser. Proceedings of Machine Learning Research, M. Meila and T. Zhang, Eds., vol. 139. PMLR, Jul. 2021, pp. 4008–4017.
- [55] S. Bartz, M. N. Dao, and H. M. Phan, "Conical averagedness and convergence analysis of fixed point algorithms," *Journal of Global Optimization*, vol. 82, no. 2, pp. 351–373, Feb. 2022.
- [56] M. Mohri, A. Rostamizadeh, and A. Talwalkar, *Foundations of machine learning*, 2nd ed., ser. Adaptive computation and machine learning. Cambridge, Massachusetts: The MIT Press, 2018.
- [57] F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau, M. Brucher, M. Perrot, and E. Duchesnay, "Scikit-learn: Machine learning in Python," *Journal of Machine Learning Research*, vol. 12, pp. 2825–2830, 2011.
- [58] N. Bastianello, "tvopt: A Python Framework for Time-Varying Optimization," in 2021 60th IEEE Conference on Decision and Control (CDC), 2021, pp. 227–232.
- [59] H. Rinne, *The Weibull distribution: a handbook*. Boca Raton: CRC Press, 2009.
- [60] H. H. Bauschke, S. M. Moffat, and X. Wang, "Firmly Nonexpansive Mappings and Maximally Monotone Operators: Correspondence and Duality," *Set-Valued and Variational Analysis*, vol. 20, no. 1, pp. 131– 153, Mar. 2012.
- [61] S. Sundhar Ram, A. Nedić, and V. V. Veeravalli, "Distributed Stochastic Subgradient Projection Algorithms for Convex Optimization," *Journal* of Optimization Theory and Applications, vol. 147, no. 3, pp. 516–545, Dec. 2010.