

When is Assortment Optimization Optimal?

Will Ma

Graduate School of Business, Columbia University, New York, NY 10027, wm2428@gsb.columbia.edu

Given many different items in a product category, each with its own fixed price point, which subset should a retailer offer to its customers? Assortment optimization describes the process of finding a subset that maximizes average revenue, based on a model for how customers choose between items in that product category. In this paper we ask whether offering an assortment is actually optimal, given the emergence of more sophisticated selling practices, such as offering certain items only through lotteries.

To formalize this question, we introduce a mechanism design problem where the items have fixed prices and the seller optimizes over (randomized) allocations, given a distribution for how a buyer ranks the items. Under our formulation, deterministic mechanisms correspond to assortments, while randomized mechanisms correspond to lotteries for selling items with fixed prices. We derive a sufficient condition, based purely on the buyer’s preference distribution, that guarantees assortments to be optimal within the larger class of randomized mechanisms. Our sufficient condition captures many preference distributions commonly studied in assortment optimization, including Multi-Nomial Logit (MNL), Markov Chain, a mixture of MNL with an Independent Demand model, and simple cases of Nested Logit. When our condition does not hold, we also bound the suboptimality of assortments compared to lotteries. Finally, from our paper emerges two results of independent interest: an example showing that Nested Logit is not captured by Markov Chain choice models, and a tighter Linear Programming relaxation for assortment optimization.

1. Introduction

A ubiquitous question faced by online and brick-and-mortar retailers alike is which *assortment* of products to carry. On one hand, carrying a wide variety of products allows the retailer to satisfy more customers’ needs, capturing greater market share. On the other hand, carrying cheap brands can cannibalize the sales of luxury brands, hurting the retailer’s margins. Correctly designing the assortment of products has a tremendous impact on a retailer’s bottom line (Kök et al. 2008).

This well-studied problem can be formalized as follows. There is a universe of substitute products, and each product j has a fixed price r_j which has been pre-determined by the brand or the competition. The buyer has an a-priori *ranking* of the products in the universe, including the “no-purchase” option represented by $j = 0$ with $r_j = 0$. After seeing a subset, or assortment, of products S , the buyer chooses their most-preferred product among $S \cup \{0\}$, paying the fixed price of that product. The ranking can be interpreted as the products sorted in decreasing order of *utility* $v_j - r_j$,

where v_j denotes some underlying cardinal valuation for product j . The *assortment optimization* problem is then: given exogenous prices r_j and a distribution of rankings (or valuations) for the buyer, compute an S to offer which maximizes expected revenue.

Our paper asks—can the seller *go beyond* the revenue of the optimal assortment? In practice, some designer fashion brands have begun using randomized lotteries¹ to sell certain products. These lotteries occur for new products being launched, and users who sign up agree that if they “win” a product, they will be charged the fixed price of that product. We emphasize that these lotteries do not change the price; they merely make some products more difficult to obtain. Every product still has a fixed price, which is important for protecting the brand value of high-end products.

We begin by providing a concrete example showing that such a lottery can earn greater expected revenue than any assortment. We note that due to the fixed prices, our example differs from the lotteries that have been extensively studied in the Bayesian mechanism design literature, in which the pricing of each product and lottery option is a decision variable (see Subsection 2.1).

EXAMPLE 1 (GENERAL SUBOPTIMALITY OF ASSORTMENTS). There are four items A, B, C, D with fixed prices $r_A = 2, r_B = r_C = r_D = 1$. The ranking is uniformly drawn from $\{(BA), (CA), (DA), (CB), (DB), (DC)\}$, where we have expressed each ranking as a *list* of items sorted in decreasing order of preference, omitting the items ranked below 0. On this example, it is easy to check that one cannot do better with an assortment than offering $\{A, B\}$ or $\{A, B, C\}$, both of which have expected revenue **7/6**. On the other hand, consider the *top- k lottery*, in which the buyer submits up to k distinct items, and then randomly receives one of them with probability $1/k$ each (if the buyer reports s items where $s < k$, then they receive option 0 with probability $1 - s/k$). It can be checked that a top-2 lottery on this example has expected revenue

$$\frac{1.5 + 1.5 + 1.5 + 1 + 1 + 1}{6} = \mathbf{7.5/6}, \quad (1)$$

greater than that of any assortment. To explain (1), note that with any of the possible lists the buyer would submit 2 items, making a purchase w.p. 1 and paying an average of 1.5 if the list contains A and an average of 1 otherwise. In essence, the top-2 lottery has allowed the retailer to capture full market share while reducing the cannibalized sales of the high-priced item A . \square

Example 1 describes a specific “top- k ” lottery that can outperform any assortment. In practice, the seller can design many variants of lotteries, including lotteries that restrict to a subset of items S (which resembles assortment optimization), generalized top- k lotteries where items can be submitted multiple times, etc. In this paper, we formalize the space of all allowable lotteries using mechanism design, in which the buyer must be incentivized to submit their true preferences and

¹ An example is End Clothing; see their weekly launches on endclothing.com.

participate in the lottery. We then derive conditions under which such lotteries are no better than assortments, i.e. conditions under which the phenomenon observed in Example 1 cannot occur. We show that these conditions are actually satisfied by many *choice models* from the assortment optimization literature, which impose different structures on the distribution of rankings. Finally, we analyze the suboptimality of assortments when these conditions are not met.

1.1. Formulation of Mechanism Design Problem

We formulate a mechanism design problem in which a revenue-maximizing seller can optimize over allocations, but not prices. Although there is a large literature on mechanism design over allocations (reviewed in Subsection 2.2), our problem has two major differences. First, most mechanisms in this literature are prior-free. By contrast, we are optimizing with respect to a given Bayesian prior on the buyer’s preferences. Second, the tension in this literature is traditionally caused by resource scarcity, where there are *multiple* agents who want the same item and hence cannot all be satisfied. By contrast, the tension in our problem exists even for a *single* buyer with no resource limitations, because there is a mismatch between the seller’s preferences (given by the product revenues r_j) and the buyer’s preferences (given by the ranking, which includes a no-purchase option). This mismatch corresponds exactly to the tension in the assortment optimization problem.

Formally, in our mechanism design problem the buyer reports a list $\ell = (\ell_1, \dots, \ell_s)$ containing the s items they rank above 0 in decreasing order of preference. The buyer is then assigned, possibly randomly, an item j to purchase for price r_j (where j could be 0). The buyer must prefer the random assignment from truthful reporting over any other assignment, including the option of being assigned item 0 w.p. 1 (which is obtained by not participating in the mechanism).

Note that there is ambiguity² in how an ordinal ranking’s preferences over randomized assignments should be defined. To resolve this ambiguity, we adopt the definition that truthfully reporting to the mechanism and agreeing to purchase any item assigned must maximize the buyer’s probability of receiving one of their k most-preferred items, *simultaneously* for all $k = 1, \dots, s + 1$. Since the buyer’s $s + 1$ ’st most-preferred item is 0, under this definition, the mechanism must w.p. 1 assign the buyer either an item on their list or item 0. In other words, the mechanism is *ex-post individually rational*—the buyer does not regret participating in the mechanism *under any realization*, since they can only be assigned an item they are willing to purchase. This also explains why the buyer need not report their ranking below 0.

The randomized truthfulness we are imposing on the mechanism can be seen as fairly strong, in that it equivalently says for *any* set of cardinal utilities consistent with the reported list,

² For example, would the buyer prefer to surely receive their 2nd choice, or have a 50-50 between their 1st choice and 3rd choice?

truthful participation must maximize the buyer’s expected utility. This constraint was pioneered by Gibbard (1977), and is standard for an ordinal ranking facing a randomized outcome (see Chakrabarty and Swamy 2014). Hereafter we proceed with our strong randomized truthfulness constraint and refer to mechanisms satisfying it, along with the implied ex-post individual rationality, as “IC” (incentive-compatible). This constraint is formalized along with our problem in **Section 3**, and further discussion of other IC constraints and models can be found in **Subsection 3.2**.

We note that top- k lotteries, or more generally top- k lotteries restricted to items from a subset S , clearly satisfy this definition of IC—the buyer is no worse off in any way by truthfully reporting their up-to- k most-preferred items from S . When $k = 1$, this top- k lottery has an extremely simple implementation—let the buyer choose from assortment S . In analyzing when assortments are optimal, we are analyzing when this simple implementation is as powerful as eliciting preferences.

1.2. Description of Results

First we show that assortments are optimal within the larger class of IC mechanisms whenever the list distribution follows a *Markov Chain choice model*, given any fixed item prices. Markov Chain already represents a fairly broad family of list distributions, capturing the most common *Multi-Nomial Logit (MNL)* choice model, as we explain in **Section 4**.

We also extend this result to hold under a more general sufficient condition, motivated as follows. One way to view a Markov Chain choice model is—given any two potential prefixes for the buyer’s list that end with the *same* item, the conditional probabilities of the buyer choosing any item from any assortment disjoint from these prefixes are *equal*. As a concrete example, if I take the customers whose lists begin with (CB) , and compare them to the customers whose lists begin with (B) , then these two groups have the same choice probabilities on all assortments not containing items B or C . Our sufficient condition relaxes this property satisfied by Markov Chain choice models in the following way—if one of the two prefixes *contains the other*, then it is allowed to have *higher* conditional choice probabilities. On this example, since prefix (CB) contains (B) , the probability of the customer purchasing e.g. item A from assortment $\{A\}$ is allowed to be higher conditional on prefix (CB) rather than (B) . Note that Example 1, in which assortments were not optimal, goes directly against this condition—there the conditional probability of the customer purchasing item A was higher starting from (B) than (CB) , being 100% instead of 0%.

Our sufficient condition is formally defined in **Subsection 5.1**, with our main result that assortments are optimal under this condition proved in **Subsection 5.2**. There we also provide further intuition on why this condition causes assortments to be optimal, and discuss our proof techniques.

Knowing that assortments are optimal under this sufficient condition allows us to capture additional distributions, such as a mixture of MNL with singleton lists (a choice model of interest


as demonstrated by Cao et al. (2020)). Indeed, mixing with singleton lists would not cause our sufficient condition to be violated, since it can only *lower* the conditional choice probabilities for singleton prefixes, which *are contained in* all other prefixes that end with the same item. Our sufficient condition also allows us to capture Tversky’s Elimination by Aspects model, and special cases of Nested Logit with a small number of items, two other models not captured by a Markov Chain alone. These results can be found in **Subsection 5.3**. In fact, to our knowledge our paper is the first to show that a Nested Logit choice model cannot be captured by any Markov Chain. Given this fact, our sufficient condition could also be of general interest, since it unifies (at least for a small number of items) these two choice models that have been studied frequently but disparately.

Finally, we investigate the power that lotteries can have over assortments when our sufficient condition does not hold. First, we show in **Subsection 6.1** that a simple subclass of lotteries dubbed *budget-additive mechanisms*, which includes both assortments and top- k lotteries, can outperform assortments by a factor of at most $e \approx 2.71$. In other words, assortments are *approximately* optimal if we only compare against simple lotteries. We show in **Subsection 6.2** that assortments are also approximately optimal if customers are “picky”, having *short lists* like in the model of Feldman et al. (2019). In stark contrast though, we show in **Subsection 6.3** that assortments can be suboptimal by an unbounded factor without any restrictions on the lottery or list length. In passing, we also show in **Subsection 6.4** that our mechanism design formulation provides a Linear Programming (LP) relaxation for assortment optimization that is tighter than the state-of-the-art from Bertsimas and Mišić (2019). This has implications for computing the optimal assortment.

We conclude in **Section 7**. In the appendix, we briefly discuss two further extensions that we find interesting. First, although we focus on having a single buyer in this paper, our problem can be easily extended to multiple buyers, who compete for a limited inventory of the items. A formulation of this generalized problem, as well as a simple example on which a *top-trading cycle* mechanism (with optimized initial assignments) is revenue-optimal, is provided in **Appendix A**. This connects our work to the extensive literature on mechanism design over allocations (see Subsection 2.2). In **Appendix B**, we show that our problem provides a feasible, but not necessarily optimal, solution in a setting where the seller knows the induced ordinal distribution but not the buyer’s true distribution over cardinal utilities. This connects our work to the growing literature on *robustness* in mechanism design (see e.g. Carroll 2017).

1.3. Relevance of Results

Our result about assortments being optimal has the following practical implications. As discussed earlier, a seller who is constrained by fixed item prices can still design many variants of lotteries, each with different implementations. Our paper characterizes the space of all possible lotteries via mechanism design, and asserts that using these lotteries cannot increase expected revenue.

Figure 1 Hierarchy of choice models for which assortments are optimal

*See Subsection 5.3 for the detailed specification.

Abbreviated references: TvR04 (Talluri and Van Ryzin 2004), RSS10 (Rusmevichientong et al. 2010), BGG16 (Blanchet et al. 2016), FT17 (Feldman and Topaloglu 2017), DGSY20 (Désir et al. 2020), M20 (Ma 2020).

Of course, this result is predicated on having a particular model for the distribution of customer preferences, such as MNL or Markov Chain. Nonetheless, we emphasize that these simple choice models, specifically MNL, represents the extent to which choice models have been deployed in practice (see Feldman et al. 2021)—even if true customer behavior is believed to be more complicated, MNL provides an approximation that is simple enough to explain to managers. Our main result tells a company that once they have settled on using MNL to make decisions, assortments suffice to solve the revenue maximization problem. This also has computational implications—optimization over randomized mechanisms is intractable³ for the MNL choice model, but once we know the optimal mechanism to be an assortment, assortment optimization for MNL is easy to solve.

We provide a hierarchy of the choice models captured by our sufficient condition, along with references to assortment optimization on them, in Figure 1. We note that although our sufficient condition captures the mixture of a Markov Chain choice model with singletons, which is an open problem for assortment optimization on the frontier of Figure 1, we do not actually solve this open problem, since our result showing assortments to be optimal is non-constructive. Nonetheless, as discussed in Subsection 1.2, our results have many implications for choice modeling and assortment optimization, which are relevant even to a company uninterested in randomized lotteries.

³This is because under MNL, any list has a positive probability of being realized, and hence the LP describing the optimal mechanism has exponentially many variables and constraints. See Subsection 3.1 for the LP formulation.

2. Literature Review

Our paper unifies elements from three areas with extensive work: assortment optimization, Bayesian mechanism design, and mechanism design over allocations. We believe our problem arises quite naturally from the intersection of these areas. We now review the latter two areas in mechanism design and some further related work (assortment optimization was already discussed in Section 1).

2.1. Bayesian Mechanism Design

This area studies the design of allocation and pricing rules to maximize the expected revenue from selling a set of items, given Bayesian priors on the valuations of a set of buyers. Landmark papers (Myerson 1981, Riley and Zeckhauser 1983) showed that when selling a *single* item, a *deterministic* mechanism is always optimal, corresponding to a posted price in the special case of one buyer. Since then, understanding the power of randomized lotteries for selling multiple items has been a central goal of Bayesian mechanism design. Common questions (with representative papers) are:

1. Identifying conditions under which deterministic mechanisms are as powerful as randomized ones (Armstrong 1996, Thanassoulis 2004, Alaei et al. 2013, Haghpanah and Hartline 2015, 2019);
2. Bounding the suboptimality of simple deterministic mechanisms for a buyer with *unit-demand* valuations (Chawla et al. 2007, 2010, Briest et al. 2015, Chawla et al. 2015);
3. Bounding the suboptimality of simple deterministic mechanisms for a buyer with *additive, independent* valuations (Hart and Reny 2015, Hart and Nisan 2017, Babaioff et al. 2020).

Two features present throughout these papers are: (i) pricing is a decision variable; and consequently (ii) the buyer’s preference is modeled using a cardinal valuation distribution. Our paper, motivated by assortment optimization and the fixed-price lotteries emerging in designer fashion, adds new twists to this literature because: (i) pricing is exogenously-determined; and (ii) the buyer’s preference is modeled using an ordinal ranking distribution. Under these new twists, our paper focuses exactly on question 1 above, asking “when is assortment optimization optimal?”, following a sequence of similar titles in this area (Haghpanah and Hartline 2019, Bergemann et al. 2021). We also make progress on questions 2 and 3 of bounding the suboptimality of deterministic mechanisms, in Section 6.

We now elaborate on the aforementioned papers on unit demand, in which randomized lotteries allocate at most one item to the buyer, closely resembling our problem. The existing papers show that allowing randomized allocations in the mechanism, *assuming that each option can be priced discriminatively*, increases revenue by a factor of ∞ for arbitrary distributions over item valuations (Briest et al. 2015), at most 4 if these valuations are independent, and at most 3.16 if these valuations are IID (Chawla et al. 2015). Our paper shows that allowing randomized allocations, *assuming that prices of options must be tied to the arbitrarily-given fixed prices for items*, does **not**

increase revenue for common ordinal distributions, and also particular cardinal distributions⁴. The buyer in our mechanism requires a stronger IC constraint than expected utility maximization as in the existing papers, though. Nonetheless, we wanted to highlight this stark contrast in the power of randomization depending on whether pricing is a decision variable.

2.2. Mechanism Design over Allocations

We study mechanism design for agents who report ordinal preferences over a set of alternatives and incur a (randomized) outcome from that set. This is a vast research area that includes problems in social choice (voting, facility location) and resource allocation (house allocation, matching). The main difference between our problem and these is that our designer is selfish and trying to maximize their own exogenously-defined utility (in the form of the product revenues r_j), instead of selecting a socially desirable outcome based on the preferences of agents. The tension in our problem comes from a *mismatch between the preferences of the designer and one agent*, whereas the tension in these problems comes from having *multiple agents*, who either disagree on the best outcome (in the case of social choice) or want the same limited resource (in the case of resource allocation). Consequently, our problem is interesting even with a single agent, whereas all of these problems are non-trivial only with multiple agents. We now discuss the social choice and resource allocation problems in more detail, highlighting these differences and any shared elements with our problem.

In social choice, the designer’s goal is to aggregate the rankings (“votes”) of multiple agents and select an outcome that reflects the majority opinion. Results for this problem in its full generality are mostly negative (Gibbard 1973, Satterthwaite 1975), even for randomized mechanisms that are IC under the same definition as we study (Gibbard 1977). However, positive results can be derived after imposing structural assumptions on the rankings of agents, such as “single-peakedness”, which results in the facility location problem (see Schummer and Vohra 2007, Sec 10.2). As mentioned earlier, our problem differs from these because the designer has an exogenous utility r_j for each option j . The designer cannot trivially force the outcome $\arg \max_j r_j$ because the agent has an *outside option* which forces the mechanism to satisfy individual rationality. By contrast, the social choice problem would be trivial with a single agent—the designer would simply let the agent dictatorially choose their favorite outcome from the set.

Meanwhile, outside options have already been studied in a different stream of literature—on the house allocation problem, in which multiple agents each bring a house to the market to trade. For this problem, the celebrated top-trading-cycles (TTC) mechanism of David Gale guarantees that no individual ends up with a house less-preferred to the outside option they started with—in fact, no *coalition* is worse off by bringing their houses to the market instead of doing outside

⁴ Our sufficient condition captures MNL, which can be described using a cardinal-utility model in which the valuation of every item (including the no-purchase option) incurs an independent Gumbel shock with CDF $F(x) = e^{-e^{-x}}$.

trading among themselves (Shapley and Scarf 1974). Moreover, this mechanism is IC (Roth 1982), and is the only pareto-optimal mechanism to satisfy both incentive compatibility and individual rationality (Ma 1994). Despite sharing the element of outside options with our problem, the house allocation problem is substantially different, because once again the designer is interested in social outcomes (e.g. pareto-optimality) instead of maximizing r_j .

The other resource allocation problems have the same difference with our problem. If n items are to be allocated among n agents with preferences over those items (but no endowed item as in the house allocation problem), then under mild conditions, the only feasible mechanisms are serial dictatorships (SD), in which the agents sequentially pick their most-preferred remaining item in some order (Svensson 1999). More recent papers have studied the desirability of SD when the agent order is chosen at random (Bhalgat et al. 2011), or additional mechanisms for allocating the items under a relaxed notion of randomized ordinal IC (Chakrabarty and Swamy 2014). If the items also have preferences over the agents, then this becomes the stable matching problem, which has the widely studied and applied deferred acceptance (DA) mechanism (see Schummer and Vohra 2007, Sec 10.4). In any case, these problems all differ from ours because the designer does not have exogenous utilities r_j that could conflict with the agent preferences.

Finally, our work also differs from this mechanism design over allocations literature because we consider a *Bayesian* setting, with a well-defined objective (in terms of r_j 's) to *optimize* over a prior. By contrast, the aforementioned mechanisms TTC, SD, DA are *prior-free*, and designed with *multiple objectives* (in terms of which agent's preferences to prioritize) in mind. Consequently, the aforementioned works either establish prior-free properties for these mechanisms (e.g. no coalitions, pareto-optimality), or justify them as the *unique* mechanisms that satisfy a given combination of properties. In the next subsection we mention some literature that optimizes over the parameters of DA mechanisms. Meanwhile, we note that (prior-free) optimization formulations of problems such as facility location can also be made using the framework of Procaccia and Tennenholtz (2013).

2.3. Further Related Work

Optimization of allocation mechanisms has been studied extensively in papers about school choice (Ashlagi and Shi 2014, 2016, Bodoh-Creed 2020, Feigenbaum et al. 2020, Shi 2021). Aside from the single vs. multiple agent difference emphasized in Subsection 2.2, these papers also differ from ours in that they focus on optimizing over policy levers available in the school choice application, instead of the mechanism itself (which is fixed to a student-proposing variant of DA). We should mention that Shi (2021) also connects assortment optimization with mechanism design, using it as a *subroutine* for optimizing student priorities. Another relevant point is that he proposes a definition of cardinal utility for Markov Chain choice models, which could be used to formulate alternate IC constraints in our problem.

Connections between assortment optimization and mechanism design have also been previously made in Ma (2020). However, he focuses solely on deterministic mechanisms, in which case the single-buyer problem is equivalent to assortment optimization, and the non-triviality comes from having multiple buyers in a *service-constrained* environment (as introduced in Alaei et al. (2013)). By contrast, we focus on a single buyer and randomized mechanisms, which to our surprise, has not been studied until now. We note that mechanisms for the supplier side, to procure an assortment of products, have meanwhile been studied in Saban and Weintraub (2020).

Finally, we mention that some of the mechanisms discussed here, e.g. top- k lotteries, can be presented in a way that is similar to opaque products in practice, as described in Elmachtoub et al. (2015), Elmachtoub and Hamilton (2021).

3. Problem Definition and Sequence of Relaxations

There is a seller with n items, denoted by $N = \{1, \dots, n\}$. Each item $j \in N$ must be sold at its commonly-known, exogenously-given price $r_j \geq 0$.

There is a single buyer who is willing to purchase at most one item from N . The private type of the buyer is given by an ordered subset of N , indicating the items they are willing to purchase, in decreasing order of preference. For example, if the type is $(1, 3)$, then the buyer's first choice is to pay r_1 for item 1, second choice is to pay r_3 for item 3, and third choice is to purchase nothing at all; the buyer is not willing to consider other items. We will refer to the type as a *list* ℓ , which is assumed to be a strictly ordered subset of N .

The buyer's type is drawn from a commonly-known *list distribution*, given by a support \mathcal{L} of ordered subsets and a probability $p(\ell)$ for each list $\ell \in \mathcal{L}$. We will refer to the list distribution using p , which satisfies $\sum_{\ell \in \mathcal{L}} p(\ell) = 1$.

DEFINITION 1 (LIST NOTATION). We define the following notation on lists ℓ :

- Let $|\ell|$ denote the length of a list ℓ . For example, if $\ell = (1, 3)$, then $|\ell| = 2$.
- For $k = 1, \dots, |\ell|$, let ℓ_k denote the k 'th item on ℓ , and $\ell_{1:k}$ denote the sublist formed by the first k items given in order. For example, if $\ell = (1, 3, 2)$, then $\ell_1 = 1$, and $\ell_{1:2} = (1, 3)$.
- If we apply set-specific operations to a list or sublist, then we are referring to the set of items on that list or sublist. For example, if $\ell = (1, 3, 2)$, then $\ell \cap \{2, 3\} = \{2, 3\}$ and $\ell_{1:2} \cap \{2, 3\} = \{3\}$.

DEFINITION 2 (ASSORTMENT S , $\text{REV}[S]$, AND OPT^S). An *assortment* S is a subset of items which is offered to the buyer. For each $j \in S$, the buyer chooses to purchase item j if and only if j appears on their randomly-drawn list and is earliest (i.e. most preferred) among items in S on their list. We let $\Pr[j \succeq S_0]$ denote the probability of this event, suppressing the dependence on p

but adding a subscript “0” as a reminder that j has to be preferred over the “no-purchase” option. The expected revenue collected by an assortment S can then be expressed as

$$\text{REV}[S] := \sum_{j \in S} r_j \Pr[j \succeq S_0]. \quad (2)$$

Given item prices r_1, \dots, r_n and list distribution p , the *assortment optimization* problem is to find a subset $S \subseteq N$ which maximizes (2). We let OPT^S denote its optimal objective value.

3.1. Mechanism Design Generalization of Assortment Optimization

A (direct revelation) mechanism takes in a reported list, and then possibly using randomness, assigns up to one item for the buyer to purchase. As discussed in Subsection 1.1, we impose the incentive constraint that for any k , truthfully participating in the mechanism must maximize the buyer’s probability of receiving one of their k most-preferred items. This implies that the mechanism cannot assign an item less-preferred than the “no-purchase” option. We hereafter make no distinction between the buyer’s reported list vs. true list and assume that they will pay the price of any item they are assigned.

DEFINITION 3 (MECHANISM x , $\text{REV}[x]$, AND OPT^x). For any list ℓ and item $j \in \ell$, let $x_j(\ell)$ specify the probability of the mechanism assigning j given report ℓ . Our mechanism design problem is then formulated by the following LP:

$$\begin{aligned} \max \quad & \sum_{\ell \in \mathcal{L}} p(\ell) \sum_{j \in \ell} r_j x_j(\ell) \\ \text{s.t.} \quad & \sum_{j \in \ell_{1:k}} x_j(\ell) \geq \sum_{j \in \ell_{1:k} \cap \ell'} x_j(\ell') \quad \forall \ell \in \mathcal{L}, k \leq |\ell|, \ell' \in \mathcal{L} \quad (3) \\ & \sum_{j \in \ell} x_j(\ell) \leq 1 \quad \forall \ell \in \mathcal{L} \quad (4) \\ & x_j(\ell) \geq 0 \quad \forall \ell \in \mathcal{L}, j \in \ell \quad (5) \end{aligned}$$

We refer to a *mechanism* using x and say that it is *IC* if it feasibly satisfies (3)–(5). We let $\text{REV}[x]$ denote the objective at a given x and use OPT^x to refer to the optimal objective value of the LP.

Note that the mechanism only needs to be defined on lists ℓ in the buyer’s support \mathcal{L} , and the feasible region depends only on \mathcal{L} , with the probabilities $p(\ell)$ appearing in the objective. Constraints (4) ensure that the buyer is assigned at most one item, with $1 - \sum_{j \in \ell} x_j(\ell)$ understood as the probability of the buyer receiving nothing after reporting ℓ .

PROPOSITION 1 (Relating Assortments S to Mechanisms x). For any assortment $S \subseteq N$,

$$x_j(\ell) = \begin{cases} 1, & \text{if } j \text{ appears before any other item in } S \text{ on } \ell \\ 0, & \text{otherwise} \end{cases} \quad \forall \ell \in \mathcal{L}, j \in \ell \quad (6)$$

defines an *IC* mechanism with $\text{REV}[x] = \text{REV}[S]$. Therefore, $\text{OPT}^S \leq \text{OPT}^x$. On the other hand, any deterministic *IC* mechanism can be described by (6) for some corresponding assortment S .

Proposition 1 is proved in Appendix C, but we only need the forward direction, which is straightforward, to proceed. The statement as a whole is our analogue of the fact that in classical single-buyer auctions, a mechanism is deterministic if and only if it corresponds to posted pricing. Here, a deterministic mechanism corresponds to an assortment.

3.2. Discussion of Modeling Assumptions

Before proceeding, we discuss three modeling assumptions made in our problem formulation.

First, we have implicitly assumed the buyer to have an intrinsic, privately-known ranking (or equivalently, utility) for every item in the universe N . In practice, the buyer may need to search to discover their own ranking, or their utilities could be influenced by the assortment shown. Consequently, there are limits to using rankings/utilities to model customer behavior (Jagabathula and Rusmevichientong 2018), and there exist models out there that capture search (Wang and Sahin 2018) or inconsistent utilities (Chen and Mišić 2020, Chen et al. 2019). Nonetheless, for simplicity we assume the buyer’s behavior to be fully governed by some ranking/utility, all throughout this paper. This is also a pervasive assumption in mechanism design.

Second, having assumed the ranking/utility model, we impose on the mechanism a fairly strong IC constraint. We did not study an alternate setup where the buyer reports cardinal utilities and the mechanism only needs to maximize the buyer’s expectation, for the following reasons.

1. Mechanisms in the less-constrained setup may not be ex-post individually rational, which we believe to be necessary for practical implementation, since otherwise the buyer could just not pay for their assigned item after seeing an unfavorable lottery realization.

2. It is cognitively much easier for buyers to report an ordinal ranking than decide on cardinal utilities⁵. This also allows us to make a characterization based purely on the ranking distribution, which is important for capturing choice models that are defined purely combinatorially (e.g. Markov Chain) for which the cardinal utilities are ambiguous.

3. It appears difficult⁶ for assortments to be competitive with the less-constrained mechanisms.

Additionally, in Appendix B we show how our formulation provides a feasible, but not necessarily optimal, solution to a *robust* mechanism design problem in which the buyer’s true utility is cardinal but its distribution is unknown.

Finally, we note that the optimality of posting a single assortment in our model suggests that the seller cannot benefit by showing different items to different buyers. However, it is not to suggest

⁵ Although the buyer’s preference under uncertainty can be robustly described as maximizing the expectation of some underlying von Neumann-Morgenstern utilities, it is difficult for the buyer to know what these utilities are.

⁶ The less-constrained mechanisms can offer items that are *never* ranked above 0 as part of a lottery, and sell them with positive probability. Meanwhile, an assortment can never sell such items. Therefore, whenever such an item contributes the majority of the revenue, the gap between lotteries and assortments would be unbounded.

that an online retailer cannot benefit by showing *personalized assortments* (Golrezaei et al. 2014) to different users based on available information such as IP address. Crucially, our model assumes the buyers to be indistinguishable to the seller, except through preferences that they strategically disclose. If buyers can be segmented based on IP address, then our result should be applied at the segment level, determining for each indistinguishable segment whether the online retailer should show a single assortment or ask the buyers from that segment to report preferences.

3.3. Sequential Implementation of IC Mechanisms

Proposition 1 showed that $\text{OPT}^S \leq \text{OPT}^x$, and in this paper we are interested in identifying conditions under which $\text{OPT}^S = \text{OPT}^x$. To aid in upper-bounding OPT^x , our goal in this subsection is to describe every IC mechanism as a sequential exchange between the buyer and seller.

To begin, we rewrite $\text{REV}[x]$ as

$$\sum_{\ell \in \mathcal{L}} p(\ell) \sum_{k=1}^{|\ell|} r_{\ell_k} x_{\ell_k}(\ell),$$

where we are now summing over the items on list ℓ in decreasing order of preference. We aim to express each variable $x_{\ell_k}(\ell)$ as a difference in an increasing set function.

DEFINITION 4 (FUNCTION f , $\text{REV}[f]$, AND OPT^f). For an increasing set function $f : 2^N \rightarrow [0, 1]$, i.e. one which satisfies $f(S) \leq f(S')$ for all $S \subseteq S' \in 2^N$, define its “revenue” as

$$\text{REV}[f] := \sum_{\ell \in \mathcal{L}} p(\ell) \sum_{k=1}^{|\ell|} r_{\ell_k} (f(\ell_{1:k}) - f(\ell_{1:k-1})).$$

Let OPT^f denote the supremum value of $\text{REV}[f]$ over all increasing set functions $f : 2^N \rightarrow [0, 1]$.

PROPOSITION 2 (Relating Mechanisms x to Functions f). For any IC mechanism x ,

$$f(S) = \max_{\ell \in \mathcal{L}} \sum_{j \in S \cap \ell} x_j(\ell) \quad \forall S \subseteq N \quad (7)$$

defines an increasing set function $f : 2^N \rightarrow [0, 1]$ with

$$x_{\ell_k}(\ell) = f(\ell_{1:k}) - f(\ell_{1:k-1}) \quad \forall \ell \in \mathcal{L}, k = 1, \dots, |\ell| \quad (8)$$

and $\text{REV}[f] = \text{REV}[x]$. Therefore, $\text{OPT}^x \leq \text{OPT}^f$.

Proposition 2 is proved in Appendix C, but is easy to understand in the following way. Given any mechanism x , defining $f(S)$ as in (7) represents the maximum probability the buyer could have of being assigned an item in S . Clearly, this probability is increasing in S and takes values in $[0, 1]$. Finally, the IC constraints (3) can then be used to establish identity (8).

An important consequence of Proposition 2 is that every IC mechanism x can be implemented by the following sequential procedure. First, the buyer draws their most-preferred item ℓ_1 and reports

it to the mechanism. The mechanism assigns this item w.p. $f(\{\ell_1\})$, with f defined according to (7), in which case this probability equals $x_{\ell_1}(\ell)$ (since $f(\emptyset) = 0$). If ℓ_1 is not assigned, then the buyer draws and reports their second-most-preferred item ℓ_2 . The mechanism can assign this item according to an independent coin flip w.p. $\frac{f(\{\ell_1, \ell_2\}) - f(\{\ell_1\})}{1 - f(\{\ell_1\})}$, to ensure that the ex-ante probability of ℓ_2 being assigned is $f(\{\ell_1, \ell_2\}) - f(\{\ell_1\})$, equaling $x_{\ell_2}(\ell)$ by (8). This process repeats until either an item is assigned, or the buyer's list terminates when they attempt to draw their next-most-preferred item, in which case there is no recourse to sell the buyer a previously-reported item.

As emphasized by this sequential description, the probability of a list ℓ receiving its k 'th-most-preferred item under an IC mechanism can depend only on the first k items on the list, and nothing afterward. We will refer to these initial items as a *prefix*.

DEFINITION 5 (PREFIX NOTATION). A *prefix* ρ is an ordered subset of N , referring to all lists with the same (ordered) beginning as ρ . The probability of a prefix ρ occurring is

$$\Pr[\rho] := \sum_{\ell \in \mathcal{L}: \ell_{1:|\rho|} = \rho} p(\ell)$$

where we overload the $\Pr[\cdot]$ operator to indicate the probability that the randomly-drawn list begins with ρ . We say that a prefix ρ is *realizable* if $\Pr[\rho] > 0$. Any operations we defined for ordered lists ℓ in Definition 1 will also apply to prefixes ρ . We let $\rho_{\text{end}} := \rho_{|\rho|}$ denote the *endpoint* item of ρ , and $\rho_{\text{pre}} := \rho_{1:|\rho|-1}$ denote the sequence of items on ρ before its endpoint.

By exchanging sums in the original definition of $\text{REV}[f]$ and letting ρ denote $\ell_{1:k}$, we can now simplify the revenue of function f as

$$\begin{aligned} \text{REV}[f] &= \sum_{\rho} r_{\rho_{\text{end}}} (f(\rho) - f(\rho_{\text{pre}})) \left(\sum_{\ell \in \mathcal{L}: \ell_{1:|\rho|} = \rho} p(\ell) \right) \\ &= \sum_{\rho} r_{\rho_{\text{end}}} (f(\rho) - f(\rho_{\text{pre}})) \Pr[\rho]. \end{aligned} \quad (9)$$

3.4. Relaxing to Monotone Stopping Policies

Through Proposition 2, one can upper-bound OPT^x by the maximum value of expression (9) over increasing set functions $f: 2^N \rightarrow [0, 1]$. However, this relaxed optimization problem is still difficult to analyze, because the value of f on any set S has consequences for many different prefixes ρ in the sum in (9). Therefore, in this subsection we re-interpret (9) as the reward collected by an optimal stopping policy, and introduce a relaxed “cross-path” constraint under which the optimal stopping decisions can be analyzed. We now describe the optimal stopping problem, first defining a tree diagram which probabilistically generates the buyer's list.

DEFINITION 6 (TREE DIAGRAM, TRANSITION PROBABILITIES q). Consider a list distribution which is given by its realizable prefixes ρ and their positive probabilities $\Pr[\rho]$. We define its *tree*

diagram as follows. There is a *node* for every realizable prefix ρ , labeled by the endpoint ρ_{end} . There is an additional node labeled “root”, and an imaginary “terminal” node 0. For every realizable ρ , there is an arc from the node for prefix ρ_{pre} to the node for ρ , with transition probability

$$q(\rho) := \Pr[\rho] / \Pr[\rho_{\text{pre}}]$$

where ρ_{pre} is understood to be “root”, with $\Pr[\rho_{\text{pre}}] = 1$, if $|\rho| = 1$. Transition probability $q(\rho)$ can be interpreted as the likelihood of the next item being ρ_{end} conditional on a random list starting with ρ_{pre} . Note that the sum of outgoing transition probabilities from the node for any ρ is $\sum_{\rho': \Pr[\rho'] > 0, \rho'_{\text{pre}} = \rho} \Pr[\rho'] / \Pr[\rho]$, which is upper-bounded by 1.

DEFINITION 7 (OPTIMAL STOPPING PROBLEM). Given the tree diagram for a list distribution, define the following *optimal stopping problem*. Starting from “root”, the player transitions along the tree diagram according to the outgoing probabilities from the current node. Upon arriving at a node labeled j , the player can either stop to end the game with reward r_j , or proceed with the next transition. The game could also end with 0 reward upon transitioning to the “terminal” node, which occurs with probability $1 - \sum_{\rho': \Pr[\rho'] > 0, \rho'_{\text{pre}} = \rho} \Pr[\rho'] / \Pr[\rho]$ (which is non-negative) when starting from the node for prefix ρ .

An example of a tree diagram, with labeled transition probabilities, can be found at the beginning of Section 5. The optimal stopping problem from Definition 7 based on tree diagrams will be used to upper-bound OPT^x . However, we do not want to allow for any online stopping policy, i.e. the optimal one obtained from dynamic programming, because that would be too loose—an example of this can also be found at the beginning of Section 5. Therefore, we still maintain a simpler “monotonicity” constraint on the stopping decisions across sample paths, dependent on the history. Importantly though, we do allow the policy to indicate its stopping decisions based on the label j of the current node, which affects the stopping reward r_j (contrast this with the function f , which had to be defined on sets S , without being able to favor stopping on one item over another).

DEFINITION 8 (MONOTONE STOPPING POLICY ϕ , $\text{REV}[\phi]$, AND OPT^ϕ). A *monotone stopping policy* ϕ is defined by n boolean-valued functions $\phi_j : 2^{N \setminus \{j\}} \rightarrow \{0, 1\}$, one for each item j , which are *increasing*, in that $\phi_j(H) \leq \phi_j(H')$ for all $H \subseteq H' \in 2^{N \setminus \{j\}}$. For each $j \in N$ and (unordered) history $H \subseteq N \setminus \{j\}$, the value $\phi_j(H) \in \{0, 1\}$ indicates whether to deterministically stop on a node labeled j after encountering exactly the set of labels H (in any order). Define the “revenue” of ϕ

$$\text{REV}[\phi] := \sum_{\rho} r_{\rho_{\text{end}}} \Pr[\rho] \left(\phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1})) \right). \quad (10)$$

Let OPT^ϕ denote the maximum value of $\text{REV}[\phi]$ over all monotone stopping policies ϕ .

Assortments S corresponds to monotone stopping policies for which *every* ϕ_j is a *constant function*—the all-1 function if $j \in S$, and the all-0 function otherwise. For a general deterministic stopping policy ϕ , objective $\text{REV}[\phi]$ represents the expected reward collected by ϕ in the aforementioned optimal stopping problem. To see this, consider any realizable prefix ρ . Since ϕ takes values in $\{0, 1\}$, the large parentheses in (10) evaluates to 1 if and only if $\phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) = 1$ (i.e. ϕ would stop on ρ_{end} after visiting a sequence of nodes labeled $\rho_1, \dots, \rho_{k-1}$) and $\phi_{\rho_k}(\rho_{1:k-1}) = 0$ for all $k < |\rho|$ (i.e. ϕ would not have stopped earlier in this sequence). Therefore, (10) counts the contribution from all prefixes ρ which the policy would reach and stop on (collecting reward $r_{\rho_{\text{end}}}$), thereby equaling the total expected reward of ϕ .

PROPOSITION 3 (Relating Functions f to Monotone Stopping Policies ϕ). *The optimization problem for OPT^f always has an optimal solution which is an increasing boolean set function. Moreover, for any increasing boolean $f : 2^N \rightarrow \{0, 1\}$, setting*

$$\phi_j(H) = f(H \cup \{j\}) \quad \forall j \in N, H \subseteq N \setminus \{j\} \quad (11)$$

defines a monotone stopping policy ϕ with $\text{REV}[\phi] = \text{REV}[f]$. Therefore, $\text{OPT}^f \leq \text{OPT}^\phi$.

Proposition 3 is proved in Appendix C. The key observation is that the constraints enforcing f to be increasing describe a totally unimodular system, which allows us to restrict our attention *boolean-valued* functions f . To show $\text{REV}[\phi] = \text{REV}[f]$, one has to check that the linear term $f(\rho) - f(\rho_{\text{pre}})$ in (9) equals the non-linear expression $f(\rho) \prod_{k=1}^{|\rho|-1} (1 - f(\rho_{1:k}))$ when f is boolean.

4. Assortments are Optimal for Markov Chain choice models

In Section 3 we defined a sequence of relaxed optimization problems satisfying $\text{OPT}^S \leq \text{OPT}^x \leq \text{OPT}^f \leq \text{OPT}^\phi$. In this section we use the relaxation OPT^ϕ to show that assortments are optimal for Markov Chain choice models, presenting a refined proof that only works on Markov Chains. In the subsequent Section 5 we derive a different and more general proof based on a sufficient condition.

First we provide a definition of Markov Chain choice models, originally introduced in Blanchet et al. (2016). Although usually defined in terms of assortment purchase probabilities (i.e. given by $\{\Pr[j \succeq S_0] : S \subseteq N, j \in S\}$), a Markov Chain choice model has a natural interpretation as a distribution over lists. This interpretation using lists is needed when considering our mechanism design problem.

DEFINITION 9 (MARKOV CHAIN CHOICE MODEL). There is a Markov Chain with states $N \cup \{0\}$ and a probability $\sigma_{jj'}$ for transitioning from any state $j \in N$ to any state $j' \in N \cup \{0\}$, where 0 is a terminal state with no outgoing transitions. Start at each state j with probability λ_j (where $\sum_j \lambda_j = 1$) and then transition along the Markov Chain according to probabilities $\sigma_{jj'}$ (where $\sum_{j' \in N \cup \{0\}} \rho_{jj'} = 1$ for all $j \in N$). Every time a state $j \in N$ is visited for the *first* time, add j to the end of the list. The list terminates upon reaching state 0, which is assumed to occur w.p. 1.

To show that assortments are optimal for Markov Chain choice models, we do not even need the constraint that ϕ is monotone. More precisely, we show that among *all* stopping policies defined by n functions $\phi_j : 2^{N \setminus \{j\}} \rightarrow \{0, 1\}$, there always exists an optimal one corresponding to an assortment, i.e. one for which every ϕ_j is either the all-1 or all-0 function. Intuitively, this is because the state for a list generated by a Markov Chain should be fully captured by the current item j , and hence one should either always want to stop on j (in which case $\phi_j(H) = 1$ for all $H \subseteq N \setminus \{j\}$) or never want to stop on j (in which case $\phi_j(H) = 0$ for all H). However, note that the preceding argument does not immediately work, because the state does depend on the history H , in that items in H cannot appear in the future of the list. To rectify this argument, we must consider a further relaxation of stopping policies that allows for “repeat visits” and only makes sense for Markov Chain choice models. This is formalized in the theorem below, whose proof is deferred to Appendix D.

We note that even though the monotonicity constraint on ϕ is derived from incentive compatibility, a policy ϕ without the monotonicity constraint is not as powerful as a mechanism without incentive compatibility constraints. Indeed, the latter achieves the first-best revenue where the buyer’s list is revealed to the seller up-front, while the former must still decide whether to stop on each item without knowing the future of the buyer’s list.

THEOREM 1. *For a list distribution arising from a Markov Chain choice model, given any item prices r_1, \dots, r_n , assortments are optimal within the larger class of IC mechanisms.*


Although our simplified result in this section is restricted to Markov Chain choice models, it already quite expansive. In particular, Markov Chain captures the most common Multi-Nomial Logit (MNL) choice model (see Subsection 5.3), lists generated by “Outtrees” or “Intrees” (see Honhon et al. 2012), the attention-based consideration set model (see Gallego and Li 2017), and “buy-down” preferences which correspond to valuation distributions (see Ma 2020).

5. Assortments are Optimal under Sufficient Condition

In Section 4 we showed assortments to be optimal for Markov Chain choice models, by upper-bounding OPT^ϕ using a “repeat visits” optimal stopping problem that only made sense on Markov Chains. In this section we derive a different proof for assortments being optimal, for general list distributions satisfying a sufficient condition that includes Markov Chain choice models.

To motivate our sufficient condition for general list distributions, we start by identifying a property satisfied by all list distributions generated from a Markov Chain. Take any Markov Chain choice model, and consider the tree diagram (see Definition 6) describing its list distribution. Take any two prefixes ρ, ρ' with $\rho_{\text{end}} = \rho'_{\text{end}}$ that have nodes in this tree, and consider the distribution of suffixes starting from those nodes. We claim that these suffixes are “similar” in the following sense.

Figure 2 Markov Chain and tree diagram representations of the same list distribution, used to illustrate our sufficient condition.



Take any assortment S which is *disjoint from* $\rho \cup \rho'$. Then for any item $j \in S$, the probability of j being “chosen” starting from either node is identical, equal to the probability of starting from state $\rho_{\text{end}} = \rho'_{\text{end}}$ in the Markov Chain and visiting state j before any other state in S or 0.

As a concrete example, consider the Markov Chain choice model and its tree diagram in Figure 2, which describes an MNL choice model (see Subsection 5.3) with three items A, B, C whose weights equal that of the no-purchase option. Consider the prefixes $\rho = (AB)$ and $\rho' = (B)$ whose nodes are highlighted in the tree diagram. Taking assortment $S = \{C\}$, it is easy to see that the probability of encountering C in the suffix from ρ is $1/2$, while the probability of encountering C in the suffix from ρ' is $\frac{1}{3} \cdot \frac{1}{2} + \frac{1}{3} = 1/2$, both equal to the probability of visiting C before 0 when starting from state B in the Markov Chain. Note that the quantifier “disjoint from $\rho \cup \rho'$ ” on the assortment S was important. If S was allowed to be the assortment $\{A, C\}$, then the probability of C being chosen decreases from $\frac{1}{3} \cdot \frac{1}{2} + \frac{1}{3}$ to $\frac{1}{3}$ when starting from node ρ' .

Our sufficient condition can then be thought of as a more lenient version of this property satisfied by Markov Chain choice models, in that if one of the prefixes is contained within the other as a set, say $\rho' \subseteq \rho$, then the choice probabilities starting from ρ are permitted to be higher, for any assortment S disjoint from $\rho \cup \rho'$ and any item $j \in S$. This allows us to capture additional choice models, such as Markov Chain mixed with singleton lists. To illustrate this, consider the list distribution which is (ABC) or (B) each with 50% likelihood, which is a 50-50 mixture of the deterministic list (ABC) (generatable from a Markov Chain) with the singleton list (B) . This does not satisfy the equal choice probabilities property, since the probability of choosing item C from assortment $\{C\}$ is 1 when starting from prefix (AB) , and 0 when starting from prefix (B) . However, because the prefix (AB) setwise contains prefix (B) , such a difference is permitted.

We now proceed to formalize our sufficient condition, in Subsection 5.1, and prove that assortments are optimal under it, in Subsection 5.2. This proof is also based on showing that stopping policies ϕ are no better than assortments. However, unlike the case in Section 4, this is a different proof that makes use of the monotonicity constraints on ϕ . In fact, it is easy to see that non-monotone stopping policies are better than assortments on the 50-50 mixture between lists (ABC) and (B) described above—suppose $r_C = 2, r_B = 1, r_A = 0$. Then the best assortments $\{B\}$ and $\{C\}$ both earn expected revenue 1, while a non-monotone stopping policy can stop on prefix (B) without stopping on prefix (AB) , earning expected revenue $\frac{2+1}{2} = 1.5$.

5.1. Sufficient Condition

We first formalize what it means for all choice probabilities to be higher when starting from one prefix instead of another. For a realizable prefix, i.e. a prefix with non-zero probability of occurring, we refer to its conditional distribution of suffixes as its *future*, and define a notion of domination.

DEFINITION 10 (DOMINATION OF FUTURES). Given realizable prefixes ρ, ρ' , the future from ρ is said to *dominate* the future from ρ' if for all $S \subseteq N \setminus (\rho \cup \rho')$ and items $j \in S$,

$$\Pr[j \succeq S_0 | \rho] \geq \Pr[j \succeq S_0 | \rho']. \quad (12)$$

EXAMPLE 2 (DOMINATION). Consider the following futures with two items A and B :

- Future 1, which is (A) w.p. 1/2 and $()$ w.p. 1/2 (where $()$ denotes the empty future);
- Future 2, which is (A) w.p. 1/2 and (B) w.p. 1/2;
- Future 3, which is (BA) w.p. 1/2 and $()$ w.p. 1/2.

Assume that the prefixes under consideration contain neither items A nor B . It is trivial to see that Future 2 dominates Future 1. On the other hand, Future 3 does not dominate Future 1, because the probability of selling item $j = A$ under assortment $S = \{A, B\}$ is not higher for Future 3. Also note that Future 2 dominates Future 3, because it is always easier to sell items when A and B appear on different branches of the possible suffix realizations. \square

In our preceding example, Future 1 did not dominate Future 2, and Future 3 did not dominate Future 1, only because the prefixes contained neither A nor B . By contrast, if $\rho = (B)$ and $\rho' = ()$, then ρ with Future 1 would actually dominate ρ' with Future 2, because comparisons for item B (which lies in $\rho \cup \rho'$) are ignored. Similarly, $\rho = (B)$ with Future 3 would dominate $\rho' = ()$ with Future 1, because $\{A, B\}$ would no longer be a valid choice for S .

Having defined domination between futures, we now introduce our sufficient condition.

DEFINITION 11 (HISTORY-MONOTONE FUTURES). A list distribution is said to have *history-monotone futures* if for all realizable prefixes ρ, ρ' with $\rho_{\text{end}} = \rho'_{\text{end}}$, whenever $\rho_{\text{pre}} \not\subseteq \rho'_{\text{pre}}$ (i.e. ρ_{pre} viewed as a set is not contained within ρ'_{pre}), the future from ρ dominates the future from ρ' .

For every pair of prefixes with the same endpoint, our sufficient condition imposes a relationship between their futures, based on the containment between the prefixes. Note that since $\rho_{\text{end}} = \rho'_{\text{end}}$, the condition $\rho_{\text{pre}} \not\subseteq \rho'_{\text{pre}}$ is equivalent to $\rho \not\subseteq \rho'$. There are three possibilities for the containment:

1. **Neither of ρ, ρ' are contained within the other.** An example of this is $\rho = (AC), \rho' = (BC)$. In this case, Definition 11 says that the futures from ρ and ρ' must dominate each other, i.e. the choice probabilities for all S and j must be identical when conditioned on either prefix ρ or ρ' . Note that this does not⁷ impose the suffixes starting from ρ and ρ' to be identically distributed.

2. **One of ρ, ρ' is contained within the other.** An example of this is $\rho = (AC), \rho' = (C)$. In this case, since $\rho \not\subseteq \rho'$, Definition 11 says that the future from ρ must dominate the future from ρ' , i.e. the choice probabilities for all S and j must be higher when conditioned on prefix ρ .

3. **Both of ρ, ρ' are contained within the other.** An example of this is $\rho = (ABC), \rho' = (BAC)$. In this case, Definition 11 imposes *no relationship* between the futures from ρ and ρ' .

On the whole, cases 1–3 above impose that *larger* histories (in the sense of not being contained within other histories) must have *dominating* futures. This is why our sufficient condition is called *history-monotone futures*.

EXAMPLE 3 (DISTRIBUTION WITHOUT HISTORY-MONOTONE FUTURES). The simplest example of a list distribution that does *not* have history-monotone futures requires 3 items A, B, C , with the random list being (BA) or (BC) with equal probability. Consider prefixes $\rho = (CB)$ and $\rho' = (B)$, which have the same endpoint. According to history-monotone futures, the larger prefix (CB) must have the dominating future. However, letting $j = A$ and $S = \{A\}$, the choice probability $\Pr[j \succeq S_0 | (CB)]$ equals 0, while $\Pr[j \succeq S_0 | (B)]$ equals 1. Since the choice probability from the larger prefix (CB) is strictly smaller, this list distribution does not have history-monotone futures. \square

The list distribution from the introductory Example 1, which demonstrated assortments to not be optimal, can also be checked to not have history-monotone futures. In fact, we now show that assortments must be optimal for any list distribution with history-monotone futures.

5.2. Statement and Proof of Main Result

THEOREM 2. *For a list distribution with history-monotone futures, given any item prices r_1, \dots, r_n , assortments are optimal within the larger class of IC mechanisms.*

Intuition behind history-monotone futures. History-monotone futures are sufficient for assortments to be as powerful as mechanisms, by the following intuition. Recall that the revenue of mechanisms can be upper-bounded by that of monotone stopping policies ϕ , and ϕ 's that stop on

⁷ As an example, let D, E, F be three items distinct from A, B, C . Suppose one suffix distribution is (DEF) or (ED) with equal probability, while another is (DE) or (EDF) with equal probability. These suffix distributions are different, yet induce the same choice probabilities for all assortments $S \subseteq \{D, E, F\}$.

either all or none of the prefixes with the same endpoint correspond to assortments. Therefore, if a general ϕ were to do strictly better, then for some prefixes ρ, ρ' with the same endpoint $\rho_{\text{end}} = \rho'_{\text{end}}$, it must continue from ρ' while stopping on ρ , and benefit from doing so because it earns strictly greater profit in the future from ρ' than ρ . However, strictly greater profit requires the future from ρ' to *not be dominated by* ρ , which under the condition of history-monotone futures, can occur only if $\rho'_{\text{pre}} \supseteq \rho_{\text{pre}}$. Therefore, the monotonicity constraints on ϕ would impose $\phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) \leq \phi_{\rho'_{\text{end}}}(\rho'_{\text{pre}})$, i.e. ϕ can only stop on ρ (which has the worse future) if it also stops on ρ' (which has the better future). This means ϕ needs to compromise between stopping on either both or none of the prefixes ρ, ρ' , and by repeating this argument for all pairs of prefixes with the same endpoint, ϕ should indeed for each item want to stop on either all of none of its prefixes.

Proof sketch. Our goal is to formalize the intuitive argument above. The argument actually has an issue though— ϕ can earn strictly greater profit by continuing from ρ' than from ρ even if the future from ρ' is dominated by the future from ρ . To see why, consider Example 2 from Subsection 5.1, in which Future 3 (equaling (BA) w.p. $1/2$) was dominated by Future 2 (equaling (A) w.p. $1/2$, (B) w.p. $1/2$). If ϕ only stops on item A when B is in the history (which is valid under the definition of monotone stopping policies), then ϕ would collect strictly more revenue from item A by continuing from the dominated Future 3 instead of the dominating Future 2.

Therefore, we instead prove Theorem 2 using an extremal argument, where among all monotone stopping policies ϕ that are strictly better than any assortment, we take the ϕ that *maximizes* the number of 1-entries (i.e. maximizes the number of (item, history)-pairs (j, H) for which $\phi_j(H) = 1$). We modify ϕ to have one more 1-entry without decreasing its revenue, causing a contradiction and implying that such a ϕ must not have existed in the first place. To find a modification, we let S denote the set of items j for which ϕ_j is the all-1 function. We then introduce the concept of an *S -adjusted price* for a prefix ρ , which is the revenue $r_{\rho_{\text{end}}}$ minus the future revenue from assortment S conditional on starting from ρ . In light of the issue mentioned above, this concept is useful because if $\rho_{\text{end}} = \rho'_{\text{end}}$, then ρ' can only have a strictly lower S -adjusted price than ρ (i.e. it is worse to stop on) if the future from ρ' is not dominated by the future from ρ .

We show that the revenue difference $\text{REV}[\phi] - \text{REV}[S]$, which must be positive, can be expressed as a non-negative linear combination of S -adjusted prices for prefixes disjoint from S . We then use the relationship between S -adjusted prices and domination to show that under history-monotone futures, the set of S -disjoint prefixes can be partitioned into “tiers”, over which the S -adjusted prices are monotone. This monotonicity ultimately helps us find an item $j \notin S$ and history H for which changing $\phi_j(H)$ from 0 to 1 does not decrease $\text{REV}[\phi]$, completing the proof.

Intermediate concepts and results needed. We now introduce the results based on S -adjusted prices and partitioning into tiers that are needed for our proof of Theorem 2, which is

presented afterward. We note that Definition 12 extends the S -adjusted prices from Désir et al. (2020) to be defined for prefixes instead of items, and Lemma 1 extends their externality-adjustment technique to arbitrary choice models. Proofs of the intermediate results are deferred to Appendix E.

DEFINITION 12 (S -ADJUSTED PRICE FOR A PREFIX). Consider any assortment S and any realizable prefix ρ disjoint from S . Then the S -adjusted price of ρ is defined to be

$$r^S(\rho) := r_{\rho_{\text{end}}} - \sum_{j' \in S} r_{j'} \Pr[j' \succeq S_0 | \rho]. \quad (13)$$

LEMMA 1 (S -ADJUSTED REVENUE FOR A MONOTONE STOPPING POLICY). For any monotone stopping policy ϕ and assortment S ,

$$\text{REV}[\phi] - \text{REV}[S] = \sum_{\rho: \rho \cap S = \emptyset} r^S(\rho) \Pr[\rho] \phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1})). \quad (14)$$

In Lemma 1, the expression $\phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1}))$ equals 1 if and only if ϕ stops on prefix ρ after not stopping before that. Lemma 1 then intuitively says that the difference $\text{REV}[\phi] - \text{REV}[S]$ equals the expected S -adjusted price of the prefix that is stopped on over a random run of ϕ , where S -adjusted price is only defined for prefixes ρ disjoint from S .

LEMMA 2 (Partitioning Prefixes into Tiers). Suppose the list distribution has history-monotone futures, and for a fixed $S \subsetneq N$ and $j \notin S$, consider the set of realizable prefixes ρ with $\rho_{\text{end}} = j$ and $\rho \cap S = \emptyset$. These prefixes can be partitioned into T_j^S tiers, denoted by $\mathcal{T}_j^S(1), \dots, \mathcal{T}_j^S(T_j^S)$, which satisfy (after suppressing the superscript S and subscript j):

1. For any two distinct tiers t, t' satisfying $t > t'$ and any prefixes $\rho \in \mathcal{T}(t), \rho' \in \mathcal{T}(t')$, we have that $\rho \supseteq \rho'$ (when viewed as sets) and $\Pr[j' \succeq S_0 | \rho] \geq \Pr[j' \succeq S_0 | \rho']$ for all $j' \in S$;

2. Within any single tier $t \in [T]$, either:

(a) For all distinct prefixes $\rho, \rho' \in \mathcal{T}(t)$, we have that $\rho \not\subseteq \rho'$ and $\rho' \not\subseteq \rho$ (when viewed as sets) and $\Pr[j' \succeq S_0 | \rho] = \Pr[j' \succeq S_0 | \rho']$ for all $j' \in S$; or

(b) For all prefixes $\rho, \rho' \in \mathcal{T}(t)$, we have that ρ and ρ' are identical when viewed as sets.

COROLLARY 1 (Monotonicity of S -adjusted Prices). For any $j \notin S$, consider the partitioning of realizable S -disjoint prefixes with endpoint j into tiers. For tiers t, t' and prefixes $\rho \in \mathcal{T}_j^S(t), \rho' \in \mathcal{T}_j^S(t')$, the S -adjusted prices satisfy $r^S(\rho) \leq r^S(\rho')$ whenever either: $t > t'$; or $t = t'$, and $\rho \neq \rho'$ when viewed as sets.

In Subsection E.1, we present an example illustrating the intricacies in the definition of history-monotone futures and the resulting tier decomposition. Note that Corollary 1 is immediately implied by cases 1 and (2a) of Lemma 2, after applying the definition of S -adjusted prices.

Proof of Theorem 2. By Propositions 2 and 3, $\text{OPT}^x \leq \text{OPT}^f \leq \text{OPT}^\phi$, and hence the revenue of an optimal IC mechanism x is upper-bounded by the revenue of an optimal monotone stopping policy ϕ taking values in $\{0, 1\}$. We show that at least one of these optimal policies ϕ corresponds to offering an assortment. Suppose for contradiction this is not the case, and take an optimal policy ϕ which *maximizes* the number of 1-entries, i.e. maximizes $|\{j \in N, H \subseteq N \setminus \{j\} : \phi_j(H) = 1\}|$.

Let S denote the set of j for which ϕ_j is the all-1 function. Since ϕ does not correspond to offering an assortment, it must be that $S \neq N$. For items j not in S , the value of $\phi_j(H)$ is inconsequential on histories H that intersect S , because ϕ would have already stopped on any item in S . Since ϕ maximizes the number of 1-entries, we can WOLOG assume that $\phi_j(H) = 1$ for all $j \notin S$ and $H \subseteq N \setminus \{j\}$ such that $H \cap S \neq \emptyset$; note that this does not violate the monotonicity of ϕ .

We now look for an item $j \notin S$ and history $H \cap S = \emptyset$ for which we can modify $\phi_j(H)$ from 0 to 1 while preserving optimality, which would contradict ϕ having a maximum number of 1-entries. We will frequently use the identity from Lemma 1 to adjust for the externality to assortment S , which we duplicate below for convenience:

$$\text{REV}[\phi] - \text{REV}[S] = \sum_{\rho: \rho \cap S = \emptyset} r^S(\rho) \Pr[\rho] \phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1})). \quad (15)$$

To simplify notation, we let Ω^S denote the set $\{\rho \in \Omega : \rho \cap S = \emptyset\}$. For item $j \notin S$ and history $H \subseteq N \setminus S \setminus \{j\}$, let $\Omega^S(j, H)$ denote the subset of prefixes in Ω^S which first contain the elements in H in any order, followed by j . That is, $\Omega^S(j, H) = \{\rho \in \Omega^S : \rho_{1:|H|} = H, \rho_{|H|+1} = j\}$.

Recall that for items $j \notin S$, function ϕ_j is not 1 everywhere, but $\phi_j(H) = 1$ whenever $H \cap S \neq \emptyset$. Consequently, for each such j , the collection of histories $\{H \subseteq N \setminus S \setminus \{j\} : \phi_j(H) = 0\}$ is non-empty; arbitrarily choose a history H_j from this collection which is *set-wise maximal*. By this we mean that there does not exist another history H' from the same collection which contains H_j , implying that changing $\phi_j(H_j)$ from 0 to 1 would not violate the monotonicity of ϕ_j . We now note that $\phi_j(H_j)$ only appears in the RHS of (15) if $\rho \in \Omega^S(j, H_j)$: in the form of $\phi_{\rho_{\text{end}}}(\rho_{\text{pre}})$ if ρ has length exactly $|H_j| + 1$, and in the form of $(1 - \phi_{\rho_k}(\rho_{1:k-1}))$ if ρ has length greater than $|H_j| + 1$. In either case, if $\Pr[\rho] = 0$ for all $\rho \in \Omega^S(j, H_j)$, then changing $\phi_j(H_j)$ from 0 to 1 would have no impact on $\text{REV}[\phi]$, which means that we have found a feasible and optimal perturbation of ϕ which increases the number of 1-entries, leading to a contradiction.

Therefore, we can proceed assuming that for every $j \notin S$, the history H_j chosen has at least one prefix $\rho \in \Omega^S(j, H_j)$ with $\Pr[\rho] > 0$. This allows us to define

$$j^* \in \arg \max_{j \notin S} \min_{\rho \in \Omega^S(j, H_j), \Pr[\rho] > 0} r^S(\rho); \quad r^* = \min_{\rho \in \Omega^S(j^*, H_{j^*}), \Pr[\rho] > 0} r^S(\rho) \quad (16)$$

where we note that the set $j \notin S$ is also non-empty because $S \neq N$. We now argue that

$$r^* \geq r^S(\rho) \quad \forall \rho: \rho \in \Omega^S, \Pr[\rho] > 0, \phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) = 1. \quad (17)$$

To establish (17), let $j = \rho_{\text{end}}$. Since r^* was defined using a *maximin* in (16), it must be at least the minimum S -adjusted price of a realizable prefix $\rho' \in \Omega^S(j, H_j)$, i.e. $r^* \geq \min_{\rho' \in \Omega^S(j, H_j), \Pr[\rho'] > 0} r^S(\rho')$. Thus, (17) would be established by showing that for any realizable $\rho' \in \Omega^S(j, H_j)$, we have $r^S(\rho') \geq r^S(\rho)$. Since both ρ and ρ' are realizable prefixes in Ω^S which share the same endpoint j , we can apply the tier decomposition for such prefixes, where we let $\rho \in \mathcal{T}_j^S(t)$ and $\rho' \in \mathcal{T}_j^S(t')$ for tiers t, t' . We show that $r^S(\rho') \geq r^S(\rho)$ in all of the following cases.

- If $t > t'$, then $r^S(\rho) \leq r^S(\rho')$ follows immediately from Corollary 1.
- If $t < t'$, then by Lemma 2, we have $\rho' \supseteq \rho$. Since $\rho_{\text{end}} = \rho'_{\text{end}} = j$, this implies that $\rho'_{\text{pre}} \supseteq \rho_{\text{pre}}$. However, recall that ρ, ρ' were chosen so that $\phi_j(\rho_{\text{pre}}) = 1$ (by definition in (17)) and $\phi_j(\rho'_{\text{pre}}) = 0$ (because ρ'_{pre} equals H_j as a set, and H_j was chosen so that $\phi_j(H_j) = 0$). Since $\rho'_{\text{pre}} \supseteq \rho_{\text{pre}}$, the monotonicity of ϕ_j would be violated, and hence this case cannot occur.
- If $t = t'$, then as long as $\rho \neq \rho'$ when viewed as sets, Corollary 1 states that $r^S(\rho) \leq r^S(\rho')$. Otherwise, we have $\phi_j(\rho_{\text{pre}}) = \phi_j(\rho'_{\text{pre}})$, which again contradicts the facts $\phi_j(\rho_{\text{pre}}) = 1, \phi_j(\rho'_{\text{pre}}) = 0$.

Having completed the proof of (17), we now argue that r^* is also non-negative. While adjusted revenues $r^S(\rho)$ could generally be negative, if $r^* < 0$, then (17) implies that $r^S(\rho) < 0$ for all realizable prefixes ρ within Ω^S which policy ϕ stops on. However, in this case it would be impossible for the RHS of (15) to be positive, which means that $\text{REV}[\phi] \leq \text{REV}[S]$, contradicting the presumption that an assortment S could not be optimal.

Finally, equipped with (17) and the fact that $r^* \geq 0$, we argue that changing $\phi_{j^*}(H_{j^*})$ from 0 to 1, which is a feasible perturbation preserving the monotonicity of ϕ_{j^*} (since H_{j^*} was set-wise maximal), does not decrease $\text{REV}[\phi]$. To argue this, we manipulate the RHS of (15) back into a summation over lists ℓ . Let $\kappa^S(\ell)$ denote the minimum k for which $\ell_k \in S$, with $\kappa^S(\ell) = |\ell| + 1$ if $\ell \cap S = \emptyset$. Then

$$\begin{aligned} \text{REV}[\phi] - \text{REV}[S] &= \sum_{\rho: \rho \cap S = \emptyset} r^S(\rho) \left(\sum_{\ell: \ell_{1:|\rho|} = \rho} p(\ell) \right) \phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1})) \\ &= \sum_{\ell \in \mathcal{L}} p(\ell) \sum_{k < \kappa^S(\ell)} r^S(\ell_{1:k}) \phi_{\ell_k}(\ell_{1:k-1}) \prod_{k'=1}^{k-1} (1 - \phi_{\ell_{k'}}(\ell_{1:k'-1})). \end{aligned} \quad (18)$$

$\phi_{j^*}(H_{j^*})$ only has an impact on the RHS of (18) for sample paths ℓ with $p(\ell) > 0$, $\ell_{1:|H_{j^*}|} = H_{j^*}$, $\ell_{|H_{j^*}|+1} = j^*$, and $\phi_{\ell_{k'}}(\ell_{1:k'-1}) = 0$ for all $k' = 1, \dots, |H_{j^*}|$. On these sample paths, changing $\phi_{j^*}(H_{j^*})$ from 0 to 1 gains a revenue of $r_{j^*}^S(H_{j^*})$, but could lose a revenue of $r^S(\ell_{1:k})$ for some $k > |H_{j^*}| + 1$ such that $\phi_{\ell_k}(\ell_{1:k-1}) = 1$. However, the lost revenue $r^S(\ell_{1:k})$ must be no more than the gained

revenue $r_{j^*}^S(H_{j^*})$, because $r_{j^*}^S(H_{j^*}) \geq r^*$ by definition of r^* in (16) while $r^* \geq r^S(\ell_{1:k})$ since $\ell_{1:k}$ is a valid ρ in (17) (in these assertions, we have made use of the fact that $p(\ell) > 0$ and hence all of its prefixes are realizable). Alternatively, on sample paths without a $k > |H_{j^*}| + 1$ for which $\phi_{\ell_k}(\ell_{1:k-1}) = 1$, we have only gained a revenue of $r_{j^*}^S(H_{j^*})$, which is non-negative. Therefore, for all ℓ we have shown that the change on the RHS of (18) is non-negative, thereby establishing that the perturbed ϕ must also be an optimal policy, contradicting the presumption that the original ϕ had a maximum number of 1-entries among optimal policies. We have reached a contradiction in all cases, completing the proof of our main result Theorem 2. \square

5.3. Choice Models Satisfying Sufficient Condition

In this subsection we describe choice models from the literature that satisfy our sufficient condition, for which assortments are optimal. All proofs from this subsection are deferred to Appendix F.

We start with the most common MNL choice model, stated in the form of a random list.

DEFINITION 13 (MULTI-NOMIAL LOGIT (MNL)). Consider an urn which starts with balls $j = 0, \dots, n$, each with a weight $w_j > 0$. Let W denote the total weight of balls remaining in the urn. Sequentially, sample a ball *without* replacement and add its label to the end of the list, where each remaining ball j is drawn with probability w_j/W . The list terminates upon drawing the 0-ball.

The list distribution from Definition 13 is consistent with the closed-form MNL choice probabilities $\Pr[j \succeq S_0] = \frac{w_j}{\sum_{j' \in S \cup \{0\}} w_{j'}}$ denoting the likelihood of item j being chosen when an assortment S is offered (Plackett 1975, Luce 1959). As mentioned in Section 4, MNL is captured by the larger family of Markov Chain choice models (Blanchet et al. 2016). Our proposition below in conjunction with our general result based on the sufficient condition of history-monotone futures provides an alternative proof that assortments are optimal for MNL and Markov Chain choice models.

PROPOSITION 4. *A Markov Chain choice model has history-monotone futures.*

As discussed in Section 4, Markov Chain already captures a wide range of choice models from the literature. However, the following is not captured, which still satisfies our sufficient condition.

DEFINITION 14 (TVERSKY'S ELIMINATION BY ASPECTS MODEL). Consider the MNL model, where in addition the set $N \cup \{0\}$ has been partitioned into disjoint *nests* N_0, \dots, N_m , with $N_0 = \{0\}$. The list is generated as in the MNL model (where balls j are drawn with probabilities proportional to their weights w_j), except upon any ball j in nest $i \in \{1, \dots, m\}$ being drawn, the subsequent draws are constrained to balls from the same nest i (one way to implement this is to keep redrawing until the ball drawn is from nest i), all of which must be drawn before proceeding to other nests. The list terminates upon drawing the 0-ball.

The Elimination by Aspects model (Tversky 1972) is not captured by a Markov Chain choice model, since there would have to be a positive transition probability between every pair of items, which would allow nests to be traversed in a non-contiguous fashion. Despite this fact, we still show that conditional on any two prefixes ρ, ρ' with $\rho_{\text{end}} = \rho'_{\text{end}}$, the choice probabilities for assortments disjoint from $\rho \cup \rho'$ are identical, as was the case for Markov Chain.

PROPOSITION 5. *The Elimination by Aspects model has history-monotone futures.*

Next, we consider the mixture of distributions with singleton lists. As discussed at the start of Section 5, representability by a Markov Chain is not preserved under such a mixture. Nonetheless, we show that our sufficient condition *is* preserved, using the fact that if ρ contains another prefix ρ' with $\rho_{\text{end}} = \rho'_{\text{end}}$, then ρ is permitted to have higher conditional choice probabilities. Indeed, mixing with singleton lists can only reduce these choice probabilities for singleton prefixes, which are contained in all other prefixes with the same endpoint, thereby preserving our sufficient condition.

DEFINITION 15 (MIXTURE WITH SINGLETONS). The mixture of any list distribution p and singleton probabilities $\alpha_1, \dots, \alpha_n$, with $\sum_j \alpha_j \leq 1$, is defined as follows. The random list equals the singleton (j) with probability α_j , for all j . Otherwise, with probability $1 - \sum_j \alpha_j$, the random list is drawn according to p .

PROPOSITION 6. *If a list distribution has history-monotone futures, then the mixture of that list distribution with any singletons has history-monotone futures.*

Finally, we show that special cases of Nested Logit are captured by our sufficient condition.

DEFINITION 16 (GENERAL NESTED LOGIT CHOICE MODEL). Suppose the set $N \cup \{0\}$ has been partitioned into disjoint nests N_0, \dots, N_m , with $N_0 = \{0\}$. Each $j \in N \cup \{0\}$ has a weight $w_j > 0$ and each nest $i \in \{1, \dots, m\}$ has a *dissimilarity parameter* $\gamma_i \in (0, 1]$. When shown assortment S , the probability of the buyer choosing an item $j \in S$, if j lies in nest i , is given by

$$\frac{(\sum_{j' \in S \cap N_i} w_{j'})^{\gamma_i}}{w_0 + \sum_{i'=1}^n (\sum_{j' \in S \cap N_{i'}} w_{j'})^{\gamma_{i'}}} \cdot \frac{w_j}{\sum_{j' \in S \cap N_i} w_{j'}}. \quad (19)$$

The expression (19) is often referred to as the Nested Logit choice probabilities.

We note that Nested Logit corresponds to MNL if $\gamma_i = 1$ for all i , and corresponds to Tversky's Elimination by Aspects model as $\gamma_i \rightarrow 0$ for all i (Train 2009, Ch.4). We first show that for $\gamma_i \neq 1$ the Nested Logit choice probabilities are not captured by a Markov Chain choice model. To the best of our knowledge, this result has not been previously confirmed in the literature.

PROPOSITION 7. *The Nested Logit choice probabilities cannot be captured by a Markov Chain choice model, even when there are only 3 items that have equal weights and are in the same nest.*

As far as positive results, a general challenge in verifying our sufficient condition is that there is no known description of Nested Logit using a combinatorially-generated random list. In fact, multiple different list distributions could be consistent with the same set of Nested Logit choice probabilities. In light of these challenges, we now show two positive results for Nested Logit on a small number of items that construct using brute force a consistent list distribution with history-monotone futures. These results complement Proposition 7 by showing that in these special cases, although Nested Logit cannot be captured by Markov Chain, it still satisfies our sufficient condition.

PROPOSITION 8. *Any Nested Logit choice model with at most 3 items, which are in the same nest, can be represented by a list distribution with history-monotone futures.*

PROPOSITION 9. *Any Nested Logit choice model with at most 4 items, which have equal weights and are in the same nest, can be represented by a list distribution with history-monotone futures.*

All in all, our sufficient condition holds for Markov Chain choice models and the similar Elimination by Aspects model, where conditional on two different prefixes ρ, ρ' with $\rho_{\text{end}} = \rho'_{\text{end}}$, the choice probabilities for all assortments disjoint from ρ, ρ' are identical. Making use of the fact that conditional choice probabilities are permitted to be higher for “larger” prefixes, our sufficient condition is also preserved after mixing any choice model with singletons. We note that mixing with a collection of singleton lists is of broad interest in choice modeling (Cao et al. 2020), also being referred to as an *Independent Demand* model in the literature (Gallego et al. 2004). Finally, our sufficient condition is satisfied by special cases of Nested Logit, for which larger prefixes also have higher conditional choice probabilities, i.e. the longer the history, the more likely the buyer’s list is to continue into the future. To our knowledge, such intuition for Nested Logit was not known before, and could be of general interest as a property that unifies Markov Chain with Nested Logit.

We finish by making a list of choice models that, in their full generality, can be checked to not satisfy our sufficient condition: random-utility models where each item has an arbitrary independent distribution, the Mallows model (Désir et al. 2016), and the One-way Substitution and Locational Choice Models (Honhon et al. 2012). Of course, any choice model that can capture any of the bad Examples 1 and 3 also cannot satisfy our sufficient condition.

6. Revenue Gaps and Assortment Optimization for General List Distributions

Sections 4–5 considered specific list distributions for which there was no gap between the revenues of mechanisms and assortments. In this section we consider general list distributions, and establish:

- Constant-factor upper and lower bounds on the revenue gap when the mechanism is restricted to be “budget-additive”, a simple subclass that includes top- k lotteries (**Subsection 6.1**);

- The revenue gap to be bounded when lists have bounded length (**Subsection 6.2**);
- The revenue gap to be generally unbounded without any restrictions on the mechanism or the list length (**Subsection 6.3**).

Related to the question of revenue gaps, in **Subsection 6.4** we show that our new LP motivated by mechanism design provides a strictly tighter relaxation for assortment optimization than existing LP's, which are currently used to compute the optimal assortment via Integer Programming.

6.1. Bounds on the Revenue Gap for Budget-Additive Mechanisms

To define the subclass of budget-additive mechanisms, we first establish a property on the monotone-increasing set function f that was associated with every mechanism x , with $f(S)$ denoting the maximum probability that x assigns an item in S (see Proposition 2). We now show that any monotone *submodular*⁸ function f induces an IC mechanism, with the converse also being true under a mild condition.

PROPOSITION 10 (Submodularity). *For any monotone submodular $f: 2^N \rightarrow [0, 1]$, setting*

$$x_{\ell_k}(\ell) = f(\ell_{1:k}) - f(\ell_{1:k-1}) \quad \forall \ell \in \mathcal{L}, k = 1, \dots, |\ell|$$

defines an IC mechanism. Conversely, for any IC mechanism x , setting

$$f(S) = \max_{\ell \in \mathcal{L}} \sum_{j \in S \cap \ell} x_j(\ell) \quad \forall S \subseteq N$$

defines a monotone submodular function from 2^N to $[0, 1]$, under the condition that all possible lists (even those with zero probability) are included in \mathcal{L} so that the IC constraints apply to all types.

Proposition 10 is proved in Appendix G, but using its statement we can now define subclasses of mechanisms based on subclasses of monotone submodular functions, since any such function induces an IC mechanism. A well-studied subclass of monotone submodular functions are the budget-additive functions, in which f takes the form $f(S) = \min\{\sum_{j \in S} w_j, B\}$ for some non-negative weights w_j and a budget B . We assume that $B \in [0, 1]$, since we need $f(S)$ to take values in $[0, 1]$.

DEFINITION 17 (BUDGET-ADDITIVE MECHANISM). A *budget-additive mechanism* is defined by weights $w_1, \dots, w_n \geq 0$ and a budget $B \in [0, 1]$, with $f: 2^N \rightarrow [0, B]$ denoting the function $f(S) = \min\{\sum_{j \in S} w_j, B\}$. Then, the probability of the mechanism assigning a type ℓ its k 'th-most-preferred item ℓ_k is $x_{\ell_k}(\ell) = f(\ell_{1:k}) - f(\ell_{1:k-1})$, for all $\ell \in \mathcal{L}$ and $k = 1, \dots, |\ell|$.

Since the function $f(S) = \min\{\sum_{j \in S} w_j, B\}$ is easily seen to be submodular, Proposition 10 implies that budget-additive mechanisms are indeed IC. The special case of top- k lotteries is

⁸ A submodular function $f: 2^N \rightarrow \mathbb{R}$ must satisfy $f(S \cup S') + f(S \cap S') \leq f(S) + f(S')$ for all subsets $S, S' \subseteq N$.

captured by setting $w_j = 1/k$ for all j and $B = 1$. The special case of an assortment S (equivalently, a deterministic mechanism) is captured by setting $w_j = \mathbb{1}(j \in S)$ for all j and $B = 1$.

We now show that assortments can earn at least $1/e$ times the revenue of a budget-additive mechanism, without any assumption on the list distribution. Our proof entails constructing a random assortment guided by the weights w_j , and showing that this distribution of assortments has expected revenue at least $1/e$ times that of the budget-additive mechanism. This implies that at least one deterministic assortment must have the same revenue guarantee. We should note that the idea of constructing an assortment through including each item independently at random has been previously used by Aouad et al. (2018), Feldman et al. (2019) for assortment optimization.

THEOREM 3. *Given any budget-additive mechanism defined by $w_1, \dots, w_n \geq 0$ and $B \in [0, 1]$, the random assortment in which each item j is included independently with probability $1 - e^{-w_j}$ has expected revenue at least $1/e$ times that of the budget-additive mechanism.*

Proof of Theorem 3. Take any budget-additive mechanism and let f denote the function $f(S) = \min\{\sum_{j \in S} w_j, B\}$. By Proposition 2, the revenue of the mechanism can be expressed as

$$\text{REV}[f] = \sum_{\ell \in \mathcal{L}} p(\ell) \sum_{k=1}^{|\ell|} r_{\ell_k} (f(\ell_{1:k}) - f(\ell_{1:k-1})).$$

Thus, it suffices to show that for every type $\ell \in \mathcal{L}$ and position $k \in \{1, \dots, |\ell|\}$ with $f(\ell_{1:k}) - f(\ell_{1:k-1}) > 0$, the random assortment sells item ℓ_k to type ℓ with probability at least

$$\frac{1}{e} (f(\ell_{1:k}) - f(\ell_{1:k-1})). \quad (20)$$

To argue this, note that if the buyer's type realizes to ℓ , then they will choose item ℓ_k if and only if ℓ_k is included in the assortment and none of their more-preferred items $\ell_1, \dots, \ell_{k-1}$ are included. By independence, the probability of this event occurring is

$$(1 - e^{-w_{\ell_k}}) \prod_{k'=1}^{k-1} e^{-w_{\ell_{k'}}} = (1 - e^{-w_{\ell_k}}) \exp\left(-\sum_{k'=1}^{k-1} w_{\ell_{k'}}\right). \quad (21)$$

Now, by definition of f , the coefficient $f(\ell_{1:k}) - f(\ell_{1:k-1})$ can only be non-zero if $\sum_{k'=1}^{k-1} w_{\ell_{k'}} < B$ and $w_{\ell_k} > 0$, in which case it equals $\min\{w_{\ell_k}, B - \sum_{k'=1}^{k-1} w_{\ell_{k'}}\}$. Let m denote this quantity. Then, we can lower-bound the ratio of (21) to (20):

$$\begin{aligned} \frac{(1 - e^{-w_{\ell_k}}) \exp\left(-\sum_{k'=1}^{k-1} w_{\ell_{k'}}\right)}{\frac{1}{e} (f(\ell_{1:k}) - f(\ell_{1:k-1}))} &\geq \frac{(1 - e^{-m}) \exp(B - \sum_{k'=1}^{k-1} w_{\ell_{k'}} - B)}{\frac{1}{e} m} \\ &\geq \frac{(1 - e^{-m}) \exp(m - B)}{\frac{1}{e} m} \\ &\geq \frac{(1 - e^{-m}) e^{m-1}}{\frac{1}{e} m} \end{aligned}$$

where the first inequality holds because $w_{\ell_k} \geq m$, the second inequality holds because $B - \sum_{k'=1}^{k-1} w_{\ell_{k'}} \geq m$, and the third inequality holds because $B \leq 1$. The final expression can be analytically checked to be at least 1 for all values of $m \in [0, 1]$, completing the proof. \square

We note that the inclusion probabilities in Theorem 3 were optimized over all functions of the form $1 - e^{-\beta w_j}$ with $\beta > 0$. The best guarantee of $1/e$ occurred at $\beta = 1$.

Now, to complement Theorem 3, we show that assortments can earn at most $2/e$ times the revenue of budget-additive mechanisms (more specifically, top-2 lotteries), by constructing a family of instances that scale up and optimize our motivational Example 1 from the Introduction.

THEOREM 4. *Consider the following family of instances, where n denotes the number of items and M denotes a fixed large constant. Items have prices $r_j = M^j$ for all $j = 1, \dots, n$. The buyer's list always has length at most 2, and the most expensive item on it is j w.p. M^{-j} , for all j (the buyer's list is empty w.p. $1 - \sum_{j=1}^n M^{-j}$). Conditional on the most expensive item on the buyer's list being j for some $j > 1$, item j will always be the buyer's second choice, with the buyer's first choice being an item j' drawn uniformly from $\{1, \dots, j-1\}$.*

On such an instance, a top-2 lottery obtains revenue at least $n/2$. On the other hand, as $n \rightarrow \infty$, no assortment can earn revenue greater than $n/e + O(\log n)$, showing that assortments can obtain no more than $2/e$ times the revenue of the best budget-additive mechanism in the worst case.

The proof of Theorem 4 is deferred to Appendix G. The key step in upper-bounding the revenue earned by any assortment is to show that the best assortment of any size k on this instance consists of the k highest-priced items.

6.2. Revenue Gap under Bounded List Length

In Subsection 6.1 we showed that the ratio between the revenues of the best assortment and the best *budget-additive* mechanism lies in $[1/e, 2/e]$, without any further assumptions on the instance. In this subsection we show that the revenue ratio between the best assortment and the best *unrestricted* mechanism is lower-bounded by $2/(eL)$, where L denotes the maximum length of any list that can realize. Although this bound approaches 0 (and is hence meaningless) as $L \rightarrow \infty$, we show in Subsection 6.3 that this is *necessarily* the case, from a complexity-theoretic perspective.

THEOREM 5. *Define $L = \max_{\ell: p(\ell) > 0} |\ell|$. Given any mechanism x , let $f_j = \max_{\ell \ni j} x_j(\ell)$ denote the maximum probability with which x allocates item j , for all $j \in N$. Then, the random assortment in which each item j is included independently with probability $\varphi(f_j)$, where φ is the function*

$$\varphi(a) = \begin{cases} 2a/L, & a \leq 1/2; \\ 1/L, & a > 1/2; \end{cases}$$

has expected revenue at least $2/(eL)$ times that of the mechanism.

The proof of Theorem 5 is deferred to Appendix G. Based on our analysis, we designed the function $\varphi: [0, 1] \rightarrow [0, 1]$ to maximize the guarantee

$$\inf_{a \in [0, 1]} \frac{\varphi(1-a)}{1-a} (1-\varphi(a))^{L-1} \quad (22)$$

subject to the constraints that $\varphi(a)$ is non-decreasing and $\varphi(a)/a$ is non-increasing (two facts needed for our proof). It is easy to see that no higher value in (22) is possible, since when $a = 1/2$, the maximum value of $\frac{\varphi(1/2)}{1/2} (1-\varphi(1/2))^{L-1}$ over $\varphi(1/2) \in [0, 1]$ is $\frac{2}{L} (1 - \frac{1}{L})^{L-1}$. In fact, the same guarantee has been achieved by Feldman et al. (2019) based on a different function and LP, for the assortment optimization problem on lists of bounded length.

6.3. Negative Results based on Hardness of Assortment Optimization

In this subsection, we consider the computation of optimal (deterministic, budget-additive, general) mechanisms for an arbitrary list distribution that is *input explicitly*. By this, we mean that for each realizable $\ell \in \mathcal{L}$, the input contains a floating-point number for $p(\ell)$, with these probabilities summing to 1.⁹ Leveraging a strong hardness result for assortment optimization under this input model, we establish: (i) an unbounded revenue gap result for general mechanisms, and (ii) a computational hardness result for budget-additive mechanisms. To begin, we show that the optimal general mechanism under this input model can be computed in polynomial time.

PROPOSITION 11 (Computation for General Mechanisms). *On an explicitly-input list distribution, an optimal mechanism x and its revenue OPT^x can be computed in polynomial time.*

Proof of Proposition 11. The optimal mechanism and its revenue is described by the LP from Definition 3. This LP has $O(n|\mathcal{L}|)$ variables and $O(n|\mathcal{L}|^2)$ constraints, which is polynomial in the size of the input, and hence can be solved in polynomial time. \square

THEOREM 6 (from Aouad et al. (2018)). *On an explicitly-input list distribution, the optimal assortment revenue OPT^S is NP-hard to approximate within any constant factor (in fact, any factor sublinear in n).*

Theorem 6 combines with Proposition 11 to show below that there cannot be a constant (in fact, sublinear) revenue gap between mechanisms and assortments. It is also later combined with Theorem 3 to derive negative results for budget-additive mechanisms.

COROLLARY 2 (Revenue Gap for General Mechanisms). *Unless $P=NP$, on general list distributions, the revenue of mechanisms can be an unbounded factor greater than that of assortments.*

⁹ Note the contrast with the MNL and Markov Chain choice models considered in earlier sections, for which there was an implicit structural representation of the probabilities for exponentially many lists.

Proof of Corollary 2. Suppose there was a constant $B \geq 1$ such that $\text{OPT}^x \leq B \cdot \text{OPT}^S$ for all instances. By Proposition 11, the value of OPT^x can be computed in polynomial time. Since $\frac{1}{B} \cdot \text{OPT}^x \leq \text{OPT}^S \leq \text{OPT}^x$, this implies that the value of OPT^S can be approximated within a factor of B in polynomial time, which can only be the case if $P = NP$, by Theorem 6. \square

Note the contrast between Corollary 2 and the earlier Theorem 3, which showed the revenue of *budget-additive* mechanisms to be at most a factor of $e \approx 2.71$ better than assortments. These two facts together imply that general mechanisms (based on general submodular functions) can be an unbounded factor better than mechanisms based on budget-additive submodular functions.

COROLLARY 3 (Computation for Budget-additive Mechanisms). *On an explicitly-input list distribution, computing the optimal budget-additive mechanism is NP-hard.*

Proof of Corollary 3. Let OPT^w denote the revenue of an optimal budget-additive mechanism. By Theorem 3, $\frac{1}{e} \cdot \text{OPT}^w \leq \text{OPT}^S$. Meanwhile, $\text{OPT}^S \leq \text{OPT}^w$, since budget-additive mechanisms capture assortments. Therefore, computing OPT^w would allow us to approximate OPT^S within a constant factor, which is an NP-hard problem, by Theorem 6. \square

Similarly, there is a contrast between Corollary 3 and Proposition 11, highlighting that computation for the restricted subclass of budget-additive mechanisms is actually harder. In fact, for any subclass whose revenue gap with assortments is bounded, computation must be NP-hard.

6.4. A Tighter LP Relaxation for Assortment Optimization

Finally, we compare our mechanism design LP to existing LP's that are also relaxations of the assortment optimization problem on an explicitly-input list distribution. To elaborate, assortments are equivalent to deterministic mechanisms (Proposition 1), which are equivalent to integer solutions in our LP for OPT^x (Definition 3). Therefore, one could solve its corresponding Integer Program (IP), in which x is restricted to be $\{0, 1\}$ -valued, to solve the assortment optimization problem. The empirical runtime for solving such an IP, using heuristics such as Branch-and-Bound, has been observed to depend on how much larger the feasible region of the LP is than that of the IP. In particular, for two different LP's with the same integer solutions, one is *tighter* if its feasible region is contained within that of the other, in which case its IP tends to solve more quickly.

We now show that our mechanism design LP is tighter than previous LP's which have been formulated specifically to solve this explicit-input assortment optimization problem (also known as the *first-choice product line optimization* problem). It suffices to show that our LP is contained within the LP of Bertsimas and Mišić (2019), which has already been shown to be tighter than the other LP's of McBride and Zufryden (1988), Belloni et al. (2008), Feldman et al. (2019).

DEFINITION 18 (LP OF BERTSIMAS AND MIŠIĆ (2019)). In this LP, $x_j(\ell)$ denotes the probability of selling item j to list ℓ , and z_j is an auxiliary variable that can be interpreted as the probability of item j being included in the assortment. Constraints (23) enforce that the probability of selling an item cannot exceed the probability of it being in the assortment. Constraints (24) are based on the fact that a list ℓ would only purchase an item *less-preferred* to ℓ_k if ℓ_k is *not* in the assortment. Finally, constraints (25) ensure that each list is sold at most one item.

$$\begin{aligned} \max \quad & \sum_{\ell \in \mathcal{L}} p(\ell) \sum_{j \in \ell} r_j x_j(\ell) \\ & x_{\ell_k}(\ell) \leq z_{\ell_k} \quad \forall \ell \in \mathcal{L}, k \leq |\ell| \end{aligned} \quad (23)$$

$$\sum_{k' > k} x_{\ell_{k'}}(\ell) \leq 1 - z_{\ell_k} \quad \forall \ell \in \mathcal{L}, k \leq |\ell| \quad (24)$$

$$\begin{aligned} \sum_{j \in \ell} x_j(\ell) &\leq 1 \quad \forall \ell \in \mathcal{L} \\ z_j, x_j(\ell) &\geq 0 \quad \forall \ell \in \mathcal{L}, j \in \ell \end{aligned} \quad (25)$$

Bertsimas and Mišić (2019) show this to be a valid LP relaxation of the assortment optimization problem, in that its integer solutions are equivalent to assortments. We now show our LP relaxation to be tighter, in that our feasible region is contained within their feasible region (after the z -variables have been projected out).

PROPOSITION 12 (**Tighter Relaxation**). *Given any feasible solution x to the LP for OPT^x , one can define z -variables so that x is also a feasible solution to the LP from Definition 18.*

Proof of Proposition 12. Take a feasible solution x to the LP for OPT^x . Set $z_j = \max_{\ell \ni j} x_j(\ell)$ for all j , after which (23) follows. Meanwhile, (25) is implied immediately from (4). Finally, to establish (24), fix any ℓ and k . Suppose $z_{\ell_k} = x_{\ell_k}(\ell')$ for some $\ell' \ni \ell_k$. Then we have

$$z_{\ell_k} = x_{\ell_k}(\ell') \leq \sum_{j \in \ell_{1:k} \cap \ell'} x_j(\ell') \leq \sum_{j \in \ell_{1:k}} x_j(\ell) \leq 1 - \sum_{k' > k} x_{\ell_{k'}}(\ell)$$

where the first inequality holds because $x_j(\ell') \geq 0$ and the second inequality holds by the IC constraints (3). This establishes (24) and completes the proof. \square

It is also evident from the proof of Proposition 12 that this containment is strict. That is, (23)–(24) from the LP of Bertsimas and Mišić (2019) are satisfied as long as $\sum_{j \in \ell_{1:k}} x_j(\ell) \geq x_{\ell_k}(\ell')$ for $\ell' \ni \ell_k$, which is strictly weaker than our IC constraint of $\sum_{j \in \ell_{1:k}} x_j(\ell) \geq \sum_{j \in \ell_{1:k} \cap \ell'} x_j(\ell')$ for all ℓ' .

Summary of Section 6. We show that the revenue gap between mechanisms and assortments, although unbounded in the worst case (**Corollary 2**), is small for short lists (**Theorem 5**). Moreover, our LP describing the optimal mechanism can be solved in polynomial time

(**Proposition 11**), and is tighter than existing LP relaxations for the assortment optimization problem (**Proposition 12**). Therefore, our mechanism design LP should be able to speed up Integer Programming approaches to assortment optimization, an investigation we leave to future work. Additionally, the concept of budget-additive mechanisms provides a relaxation for assortment optimization with a constant-factor revenue gap (**Theorem 3**). Although this subclass of mechanisms cannot be optimized in polynomial time (**Corollary 3**), if it can be solved in special cases, then this can further speed up assortment optimization in those cases.

7. Conclusion

We summarize what we see as the three main contributions of this work.

First, on the modeling side, we introduce a Bayesian mechanism design problem where items have *fixed prices*, in which optimizing over deterministic mechanisms corresponds to assortment optimization. We present examples demonstrating that under the fixed-price restriction, lotteries can still outperform deterministic mechanisms, which to our knowledge was previously unknown.

Second, on the technical side, we derive a sufficient condition on the preference distribution under which there always exists a deterministic optimal mechanism. We establish optimality by connecting the IC constraints to *monotonicity* constraints on the stopping decisions across different sample paths in an optimal stopping problem, and deriving a new, existential local improvement argument. The techniques we use include integer programming, domination relationships in order theory, and extending the *externality-adjustment* assortment optimization framework of Désir et al. (2020) from Markov Chains to general ranking distributions. Moreover, we show that our sufficient condition is satisfied by many commonly-studied choice models from the literature, and bound the suboptimality of assortments when it is not satisfied. From this emerges two results of independent interest: an example demonstrating that Nested Logit cannot be captured by a Markov Chain choice model, and an improved Integer Programming formulation for assortment optimization.

Finally, on the managerial side, our main result suggests that for most deployments of assortment optimization in practice, usually based on the MNL choice model, assortments are in fact optimal among all fixed-price lotteries. This is in stark contrast to the traditional wisdom that discriminatively-priced lotteries can drastically increase revenue.

References

- Alaei S, Fu H, Haghpanah N, Hartline J (2013) The simple economics of approximately optimal auctions. *2013 IEEE 54th Annual Symposium on Foundations of Computer Science*, 628–637 (IEEE).
- Aouad A, Farias V, Levi R, Segev D (2018) The approximability of assortment optimization under ranking preferences. *Operations Research* 66(6):1661–1669.

-
- Armstrong M (1996) Multiproduct nonlinear pricing. *Econometrica: Journal of the Econometric Society* 51–75.
- Ashlagi I, Shi P (2014) Improving community cohesion in school choice via correlated-lottery implementation. *Operations Research* 62(6):1247–1264.
- Ashlagi I, Shi P (2016) Optimal allocation without money: An engineering approach. *Management Science* 62(4):1078–1097.
- Babaioff M, Immorlica N, Lucier B, Weinberg SM (2020) A simple and approximately optimal mechanism for an additive buyer. *Journal of the ACM (JACM)* 67(4):1–40.
- Belloni A, Freund R, Selove M, Simester D (2008) Optimizing product line designs: Efficient methods and comparisons. *Management Science* 54(9):1544–1552.
- Bergemann D, Bonatti A, Haupt A, Smolin A (2021) The optimality of upgrade pricing .
- Bertsekas D (1995) *Dynamic programming and optimal control*, volume 1 (Athena scientific Belmont, MA).
- Bertsimas D, Mišić VV (2019) Exact first-choice product line optimization. *Operations Research* 67(3):651–670.
- Bhalgat A, Chakrabarty D, Khanna S (2011) Social welfare in one-sided matching markets without money. *Approximation, randomization, and combinatorial optimization. Algorithms and techniques*, 87–98 (Springer).
- Blanchet J, Gallego G, Goyal V (2016) A markov chain approximation to choice modeling. *Operations Research* 64(4):886–905.
- Bodoh-Creed AL (2020) Optimizing for distributional goals in school choice problems. *Management Science* 66(8):3657–3676.
- Briest P, Chawla S, Kleinberg R, Weinberg SM (2015) Pricing lotteries. *Journal of Economic Theory* 156:144–174.
- Cao Y, Rusmevichientong P, Topaloglu H (2020) Revenue management under a mixture of multinomial logit and independent demand models. Technical report, Working paper.
- Carroll G (2017) Robustness and separation in multidimensional screening. *Econometrica* 85(2):453–488.
- Chakrabarty D, Swamy C (2014) Welfare maximization and truthfulness in mechanism design with ordinal preferences. *Proceedings of the 5th conference on Innovations in theoretical computer science*, 105–120 (ACM).
- Chawla S, Hartline JD, Kleinberg R (2007) Algorithmic pricing via virtual valuations. *Proceedings of the 8th ACM conference on Electronic commerce*, 243–251 (ACM).
- Chawla S, Hartline JD, Malec DL, Sivan B (2010) Multi-parameter mechanism design and sequential posted pricing. *Proceedings of the forty-second ACM symposium on Theory of computing*, 311–320 (ACM).

- Chawla S, Malec D, Sivan B (2015) The power of randomness in bayesian optimal mechanism design. *Games and Economic Behavior* 91:297–317.
- Chen N, Gallego G, Tang Z (2019) The use of binary choice forests to model and estimate discrete choices. *Available at SSRN 3430886* .
- Chen YC, Mišić V (2020) Decision forest: A nonparametric approach to modeling irrational choice. *Available at SSRN 3376273* .
- Désir A, Goyal V, Jagabathula S, Segev D (2016) Assortment optimization under the mallows model. *Advances in Neural Information Processing Systems* 29:4700–4708.
- Désir A, Goyal V, Segev D, Ye C (2020) Constrained assortment optimization under the markov chain-based choice model. *Management Science* 66(2):698–721.
- Dunning I, Huchette J, Lubin M (2017) Jump: A modeling language for mathematical optimization. *SIAM Review* 59(2):295–320, URL <http://dx.doi.org/10.1137/15M1020575>.
- Elmachtoub AN, Hamilton ML (2021) The power of opaque products in pricing. *Management Science* .
- Elmachtoub AN, Wei Y, Zhou Y (2015) Retailing with opaque products. *Available at SSRN 2659211* .
- Feigenbaum I, Kanoria Y, Lo I, Sethuraman J (2020) Dynamic matching in school choice: Efficient seat reassignment after late cancellations. *Management Science* 66(11):5341–5361.
- Feldman J, Paul A, Topaloglu H (2019) Assortment optimization with small consideration sets. *Operations Research* 67(5):1283–1299.
- Feldman J, Zhang DJ, Liu X, Zhang N (2021) Customer choice models vs. machine learning: Finding optimal product displays on alibaba. *Operations Research* .
- Feldman JB, Topaloglu H (2017) Revenue management under the markov chain choice model. *Operations Research* 65(5):1322–1342.
- Gallego G, Iyengar G, Phillips R, Dubey A (2004) Managing flexible products on a network .
- Gallego G, Li A (2017) Attention, consideration then selection choice model. *Consideration then Selection Choice Model (September 24, 2017)* .
- Gibbard A (1973) Manipulation of voting schemes: a general result. *Econometrica: journal of the Econometric Society* 587–601.
- Gibbard A (1977) Manipulation of schemes that mix voting with chance. *Econometrica: Journal of the Econometric Society* 665–681.
- Golrezaei N, Nazerzadeh H, Rusmevichientong P (2014) Real-time optimization of personalized assortments. *Management Science* 60(6):1532–1551.
- Haghpanah N, Hartline J (2015) Reverse mechanism design. *Proceedings of the Sixteenth ACM Conference on Economics and Computation*, 757–758.

-
- Haghpanah N, Hartline J (2019) When is pure bundling optimal? *Review of Economic Studies* 1.
- Hart S, Nisan N (2017) Approximate revenue maximization with multiple items. *Journal of Economic Theory* 172:313–347.
- Hart S, Reny PJ (2015) Maximal revenue with multiple goods: Nonmonotonicity and other observations. *Theoretical Economics* 10(3):893–922.
- Honhon D, Jonnalagedda S, Pan XA (2012) Optimal algorithms for assortment selection under ranking-based consumer choice models. *Manufacturing & Service Operations Management* 14(2):279–289.
- Jagabathula S, Rusmevichientong P (2018) The limit of rationality in choice modeling: Formulation, computation, and implications. *Management Science* 65(5):2196–2215.
- Kök AG, Fisher ML, Vaidyanathan R (2008) Assortment planning: Review of literature and industry practice. *Retail supply chain management* 99–153.
- Luce RD (1959) Individual choice behavior. .
- Ma J (1994) Strategy-proofness and the strict core in a market with indivisibilities. *International Journal of Game Theory* 23(1):75–83.
- Ma W (2020) Revenue-optimal deterministic auctions for multiple buyers with ordinal preferences over fixed-price items. *International Conference on Web and Internet Economics*, 162–176 (Springer).
- McBride RD, Zufryden FS (1988) An integer programming approach to the optimal product line selection problem. *Marketing Science* 7(2):126–140.
- Myerson RB (1981) Optimal auction design. *Mathematics of operations research* 6(1):58–73.
- Plackett RL (1975) The analysis of permutations. *Journal of the Royal Statistical Society: Series C (Applied Statistics)* 24(2):193–202.
- Procaccia AD, Tennenholtz M (2013) Approximate mechanism design without money. *ACM Transactions on Economics and Computation (TEAC)* 1(4):1–26.
- Riley J, Zeckhauser R (1983) Optimal selling strategies: When to haggle, when to hold firm. *The Quarterly Journal of Economics* 98(2):267–289.
- Roth AE (1982) Incentive compatibility in a market with indivisible goods. *Economics letters* 9(2):127–132.
- Rusmevichientong P, Shen ZJM, Shmoys DB (2010) Dynamic assortment optimization with a multinomial logit choice model and capacity constraint. *Operations research* 58(6):1666–1680.
- Saban D, Weintraub GY (2020) Procurement mechanisms for assortments of differentiated products .
- Satterthwaite MA (1975) Strategy-proofness and arrow’s conditions: Existence and correspondence theorems for voting procedures and social welfare functions. *Journal of economic theory* 10(2):187–217.
- Schrijver A (1998) *Theory of linear and integer programming* (John Wiley & Sons).
- Schummer J, Vohra RV (2007) Mechanism design without money. *Algorithmic Game Theory* 10:243–299.

- Shapley L, Scarf H (1974) On cores and indivisibility. *Journal of mathematical economics* 1(1):23–37.
- Shi P (2021) Optimal priority-based allocation mechanisms. *Management Science* .
- Svensson LG (1999) Strategy-proof allocation of indivisible goods. *Social Choice and Welfare* 16(4):557–567.
- Talluri K, Van Ryzin G (2004) Revenue management under a general discrete choice model of consumer behavior. *Management Science* 50(1):15–33.
- Thanassoulis J (2004) Hagglng over substitutes. *Journal of Economic theory* 117(2):217–245.
- Train KE (2009) *Discrete choice methods with simulation* (Cambridge university press).
- Tversky A (1972) Elimination by aspects: A theory of choice. *Psychological review* 79(4):281.
- Wang R, Sahin O (2018) The impact of consumer search cost on assortment planning and pricing. *Management Science* 64(8):3649–3666.

Acknowledgments

We would like to thank Joey Huchette and Juan Pablo Vielma for help with using the JuMP package (Dunning et al. 2017) to computationally check the integrality of polytopes arising from randomized ordinal IC constraints. We would also like to thank Nick Arnosti, Santiago Balseiro, Irene Lo, and Peng Shi for detailed comments on the presentation of this work, and Yilun Chen and Vineet Goyal for useful discussions.

Appendix A: Generalization to Multiple Buyers

The seller has one copy of each item in a set $N = \{1, \dots, n\}$. Each item $j \in N$ has a fixed price $r_j \geq 0$. There is a set of buyers $M = \{1, \dots, m\}$, where each buyer has a random list that is assumed to be drawn *independently*. We let \mathcal{L}_i denote the support for buyer i 's list, and $p_i(\ell_i)$ denote the probability that buyer i has list ℓ_i , with $\sum_{\ell_i \in \mathcal{L}_i} p_i(\ell_i) = 1$ for all i .

A direct mechanism maps each list *profile* $\ell := (\ell_1, \dots, \ell_m)$ lying in the set of feasible reports $\mathcal{L} := \mathcal{L}_1 \times \dots \times \mathcal{L}_m$ to an allocation matrix $X(\ell)$, where $X_{ij}(\ell)$ denotes the probability of buyer i being assigned item j . Due to only having one copy of each item, and the fact that each buyer cannot be assigned more than one item, X must satisfy the constraints

$$\sum_{i: \ell_i \ni j} X_{ij}(\ell) \leq 1 \quad \forall \ell \in \mathcal{L}, j \in N \quad (26)$$

$$\sum_{j \in \ell_i} X_{ij}(\ell) \leq 1 \quad \forall \ell \in \mathcal{L}, i \in M \quad (27)$$

$$X_{ij}(\ell) \geq 0 \quad \forall \ell \in \mathcal{L}, j \in \ell_i \quad (28)$$

Satisfying these constraints guarantees that under any profile ℓ that could be reported, the resulting matrix $X(\ell)$ will be implementable, i.e. there will be a randomized assignment of items to buyers such that no item gets assigned more than once w.p. 1, no buyer receives more than one item w.p. 1, and the probability that each buyer i receives each item j is exactly $X_{ij}(\ell)$. This is due to the integrality of the bipartite matching polytope.

Finally, with multiple buyers there are different options for the incentive-compatibility constraint, depending on what the buyers assume about each other. The most restrictive constraint is dominant-strategy incentive compatibility (DSIC), which imposes for each buyer $i \in M$, and any possibility of what the other agents report $\ell_{-i} := (\ell_{i'})_{i' \in M \setminus \{i\}}$ lying in $\mathcal{L}_{-i} := \prod_{i' \in M \setminus \{i\}} \mathcal{L}_{i'}$, that truthful reporting will satisfy the single-buyer IC constraint used throughout this paper (i.e. truthful reporting will maximize buyer i 's likelihood of receiving one of their k most-preferred items, simultaneously for all k). This can be formally expressed as

$$\sum_{j \in (\ell_i)_{1:k}} X_{ij}(\ell_i, \ell_{-i}) - \sum_{j \in (\ell_i)_{1:k} \cap \ell'_i} X_{ij}(\ell'_i, \ell_{-i}) \geq 0 \quad \forall \ell_{-i} \in \mathcal{L}_{-i}, \ell_i \in \mathcal{L}_i, k \leq |\ell_i|, \ell'_i \in \mathcal{L}_i \quad (29)$$

On the other hand, Bayesian incentive-compatibility (BIC) only imposes the single-buyer IC constraint for each $i \in M$ assuming that i knows the distributions for the other buyers and everyone buyer will report truthfully. It is more relaxed because it only imposes that (29) holds after taking an expectation over ℓ_{-i} :

$$\sum_{\ell_{-i} \in \mathcal{L}_{-i}} \left(\sum_{j \in (\ell_i)_{1:k}} X_{ij}(\ell_i, \ell_{-i}) - \sum_{j \in (\ell_i)_{1:k} \cap \ell'_i} X_{ij}(\ell'_i, \ell_{-i}) \right) \prod_{i' \in M \setminus \{i\}} p_{i'}(\ell_{i'}) \geq 0 \quad \forall \ell_i \in \mathcal{L}_i, k \leq |\ell_i|, \ell'_i \in \mathcal{L}_i \quad (30)$$

We note that under either DSIC or BIC, individual rationality is still implied assuming one uses our strong single-buyer IC constraint, which is why we only have a variable $X_{ij}(\ell)$ if $j \in \ell_i$.

The DSIC mechanism design problem can then be formulated as maximizing the expected revenue

$$\sum_{\ell \in \mathcal{L}} \left(\sum_{i \in M} \sum_{j \in \ell_i} r_j X_{ij}(\ell) \right) \prod_{i \in M} p_i(\ell_i) \quad (31)$$

subjective to constraints (26), (27), (28), and (29). The BIC mechanism design problem is identical, except that constraint (29) is replaced with (30).

We now provide a simple example with $n = m = 2$ for which the deterministic *top-trading-cycles* mechanism, using deliberate item endowments, is optimal even among the larger class of BIC mechanisms. As discussed in Subsection 2.2, the top-trading-cycles mechanism is a common one from the literature on mechanism design over allocations (more specifically, the house allocation problem), so the purpose of this example is to provide a connection to the multi-buyer generalization of our problem. We also show for this example that *serial dictatorships* are suboptimal.

EXAMPLE 4. The seller has one copy of each of two items A, B . There are two buyers, with buyer 1's list equally likely to be (A) , (B) , or (AB) , and buyer 2's list equally likely to be (A) , (B) , or (BA) . The buyers' lists are drawn independently. The fixed prices are $r_A = r_B = 1$, so the seller's objective is to maximize the expected number of items sold.

We first note that an upper bound on the expected revenue of any mechanism is $16/9$. This is because even if the seller knows the true realizations of buyer lists, they can still only sell one item if both lists realize to (A) or both lists realize to (B) , earning expected revenue $2(7/9) + 1(2/9) = 16/9$.

Now, any *serial dictatorship* mechanism can earn revenue at most $15/9$. Indeed, by symmetry, we can WOLOG assume buyer 1 to go first, choosing their favorite item from $\{A, B\}$, with buyer 2 having the option of purchasing the remaining item. Then buyer 2 has a $2/3$ chance of not purchasing an item when their list realizes to (A) (because buyer 1 takes item A with lists (A) or (AB)), and a $1/3$ chance of not purchasing when their list realizes to (B) (because buyer 1 takes item B with list (B)). Therefore, the revenue is upper-bounded by $2 - \frac{2/3 + 1/3}{3} = 15/9$.

On the other hand, consider the following mechanism. We initially give item B to buyer 1 and item A to buyer 2. Then, the buyers trade items only if *both* buyers agree. At the end, each buyer can choose to purchase the item they have. We claim that this mechanism is optimal, earning revenue $16/9$ which matches the upper bound. To see why, note that whenever a buyer's list realizes to length 2, they start with their less-preferred item, and hence a trade occurs whenever the other buyer is not willing to purchase their starting item. It can be checked that this results in both items being sold unless the lists both realize to (A) or both realize to (B) . Therefore, the expected revenue of this mechanism, which is a special case of the *top-trading-cycles* mechanism where buyer 1 is endowed with item B and buyer 2 is endowed with item A , is $16/9$. We note that if we reverse the endowments, with buyer 1 starting with A and buyer 2 starting with B , then the revenue is significantly worse, being $14/9$. \square

Appendix B: Interpretation as a Robust Mechanism

In this appendix we formulate a robust mechanism design problem in which the objective considers the *worst case* cardinal utility distribution consistent with a given ordinal distribution. We show that the formulation studied throughout this paper provides a feasible solution to the robust problem. We also construct an example where it does not provide an optimal solution.

The robust mechanism design problem is formulated as follows. We are given items $j \in N = \{1, \dots, n\}$ with fixed prices $r_j \geq 0$, and a distribution p over lists $\ell \in \mathcal{L}$, as defined throughout this paper. The robust

problem differs in that the buyer has a true cardinal utility vector $\mathbf{u} \in \mathbb{R}^{\{0\} \cup N}$, which is randomly drawn from a distribution such that the induced ordinal distribution is consistent with p . The designer commits to a cardinal mechanism, which is defined by a *menu* \mathcal{M} of *entries* $\mathbf{x} \in [0, 1]^{\{0\} \cup N}$, where each \mathbf{x} describes a randomized assignment in which the buyer purchases item j (for fixed price r_j) w.p. \mathbf{x}_j for all $j = 0, 1, \dots, n$, with $\sum_{j=0}^n \mathbf{x}_j = 1$. The 0 entry defined by $\mathbf{x}_j = \mathbb{1}(j = 0)$ for all $j = 0, \dots, n$ must lie in \mathcal{M} . Finally, the buyer chooses the option $\mathbf{x} \in \mathcal{M}$ that maximizes their own expected utility $\sum_{j=0}^n \mathbf{u}_j \mathbf{x}_j$. The seller's objective is to maximize the *worst-case* expected revenue, facing the lack of distributional information about \mathbf{u} other than it is consistent with p .

The worst-case revenue for a menu \mathcal{M} can be expressed as follows. For each $\ell \in \mathcal{L}$, let $E^{\mathcal{M}}(\ell)$ denote the *exposable* entries for list ℓ , defined as the subset of \mathbf{x} 's in \mathcal{M} for which there exist a utility vector $\mathbf{u} \in \mathbb{R}^{\{0\} \cup N}$ satisfying both consistency with ℓ , i.e.

$$\mathbf{u}_{\ell_1} > \dots > \mathbf{u}_{\ell_{\text{end}}} > \mathbf{u}_0; \mathbf{u}_0 > \mathbf{u}_j \quad \forall j \notin \ell, \quad (32)$$

and expected utility maximization for \mathbf{x} , i.e.

$$\mathbf{x} \in \arg \max_{\mathbf{x}' \in \mathcal{M}} \sum_{j=0}^n \mathbf{u}_j \mathbf{x}'_j. \quad (33)$$

Meanwhile, for each $\mathbf{x} \in \mathcal{M}$, let $r(\mathbf{x}) := \sum_{j=1}^n r_j \mathbf{x}_j$ denote the seller's expected revenue when entry \mathbf{x} is chosen. Using these two definitions, whenever the buyer's utility realizes to be consistent with a $\ell \in \mathcal{L}$, the buyer will choose the $\mathbf{x} \in E^{\mathcal{M}}(\ell)$ that *minimizes* $r(\mathbf{x})$. Therefore, the worst-case revenue of menu \mathcal{M} is

$$\sum_{\ell \in \mathcal{L}} p(\ell) \inf_{\mathbf{x} \in E^{\mathcal{M}}(\ell)} r(\mathbf{x}). \quad (34)$$

Two remarks are in order. In (32), we impose strictness, because otherwise all lists could be made to choose the 0 entry by setting $\mathbf{u}_j = 0$ for all $j = 0, \dots, n$. Given this, in (33) it does not matter whether we impose strictness, because the utility vector \mathbf{u} can always be perturbed so that the arg max in (33) is unique.

PROPOSITION 13. *Given any instance defined by $\{r_j : j \in N\}$ and $\{p(\ell) : \ell \in \mathcal{L}\}$, the optimal objective value OPT^x of the mechanism design problem from Definition 3 provides a lower bound to the robust mechanism design problem of maximizing (34) over menus \mathcal{M} containing the 0 entry.*

Proof of Proposition 13. Fix an optimal solution x to our mechanism design problem from Definition 3. Based on x , we specify a feasible menu \mathcal{M} for the robust problem whose objective value (34) is at least OPT^x . For any $\ell \in \mathcal{L}$, let $x(\ell)$ denote the vector $(1 - \sum_{j=1}^n x_j(\ell), x_1(\ell), \dots, x_n(\ell))$, which we note is a feasible entry for \mathcal{M} due to LP constraints (4) and (5). Moreover, we will let $\mathbf{0} = (1, 0, \dots, 0)$ denote the 0 entry for \mathcal{M} .

Consider the menu $\mathcal{M} = \{x(\ell) : \ell \in \mathcal{L}\} \cup \{\mathbf{0}\}$. We claim that for any $\ell \in \mathcal{L}$, the set $E^{\mathcal{M}}(\ell)$ is the singleton $\{x(\ell)\}$. This would complete the proof because $r(x(\ell)) = \sum_j r_j x_j(\ell)$, making objective function (34) identical to the objective function from Definition 3.

We first fix an $\ell \in \mathcal{L}$ and show that $x(\ell) \in E^{\mathcal{M}}(\ell)$. This is easy to see—in fact, take *any* \mathbf{u} satisfying (32). For $\mathbf{x}' \in \mathcal{M}$, there are two possibilities: $\mathbf{x}' = \mathbf{0}$, or $\mathbf{x}' = x(\ell')$ for some $\ell' \in \mathcal{L}$. If $\mathbf{x}' = \mathbf{0}$, then it is clear that

$$\sum_{j=0}^n \mathbf{u}_j (x(\ell))_j = \mathbf{u}_0 (1 - \sum_{j \in \ell} x_j(\ell)) + \sum_{j \in \ell} \mathbf{u}_j x_j(\ell) \geq \mathbf{u}_0 = \sum_{j=0}^n \mathbf{u}_j \mathbf{x}'_j \quad (35)$$

from the conditions in (32) and hence $x(\ell)$ lies in the arg max in (33). Otherwise, if $\mathbf{x}' = x(\ell')$ for some $\ell' \in \mathcal{L}$, then we can write

$$\begin{aligned}
\sum_{j=0}^n \mathbf{u}_j(x(\ell))_j &= \mathbf{u}_0 + \sum_{k=1}^{|\ell|} (\mathbf{u}_{\ell_k} - \mathbf{u}_0) x_{\ell_k}(\ell) \\
&= \mathbf{u}_0 + \sum_{k=1}^{|\ell|} (\mathbf{u}_{\ell_k} - \mathbf{u}_{\ell_{k+1}}) \sum_{k'=1}^k x_{\ell_{k'}}(\ell) \\
&\geq \mathbf{u}_0 + \sum_{k=1}^{|\ell|} (\mathbf{u}_{\ell_k} - \mathbf{u}_{\ell_{k+1}}) \sum_{j \in \ell_1, k \cap \ell'} x_j(\ell') \\
&= \mathbf{u}_0 (1 - \sum_{j \in \ell \cap \ell'} x_j(\ell')) + \sum_{j \in \ell \cap \ell'} \mathbf{u}_j x_j(\ell') \\
&\geq \sum_{j=0}^n \mathbf{u}_j(x(\ell'))_j
\end{aligned} \tag{36}$$

where in the second equality $\ell_{|\ell|+1}$ is understood to be item 0, in the first inequality we have used IC constraints (3) and the fact that the expressions in parentheses are strictly positive by (32), and in the final inequality we have used the fact that $\mathbf{u}_0 > \mathbf{u}_j$ for $j \in \ell' \setminus \ell$ by (32). This completes the proof that $x(\ell) \in E^{\mathcal{M}}(\ell)$.

We now show that $\mathbf{x}' \notin E^{\mathcal{M}}(\ell)$ for any $\mathbf{x}' \in \mathcal{M}$ distinct from $x(\ell)$. To see this, suppose for contradiction the existence of a \mathbf{u} satisfying (32) such that \mathbf{x}' lies in the arg max in (33). There are two cases: $\mathbf{x}' = \mathbf{0}$ or $\mathbf{x}' = x(\ell')$ for some $\ell' \in \mathcal{L}$, for which the deductions above resulting in either (35) or (36) can still be respectively made. In order for the inequalities to not be strict, we must have either $x_j(\ell) = 0$ for all $j \in \ell$ or $x_j(\ell) = x_j(\ell')$ for all $j = 0, \dots, n$ respectively. However, this implies that $x(\ell) = \mathbf{0}$ or $x(\ell) = x(\ell')$ respectively, contradicting the fact that \mathbf{x}' is distinct from $x(\ell)$. Therefore, $E^{\mathcal{M}}(\ell)$ cannot contain any vectors other than $x(\ell)$ and this completes the proof of Proposition 13. \square

We now show that Definition 3 is not identical to the robust mechanism design problem, by constructing an instance on which OPT^x is strictly lower than the optimal objective value of the robust problem.

EXAMPLE 5. There are five items $j = 1, \dots, 5$ with $r_1 = 2, r_2 = r_3 = r_4 = 1, r_5 = 1.5$. The buyer's ordinal distribution is uniform over the 12 lists

$$(21), (31), (41), (32), (42), (43), (5), (5), (5), (251), (351), (451).$$

It can be computed that $\text{OPT}^x = 1.3125$ for this instance, with an optimal solution being the *budget-additive mechanism* (see Subsection 6.1) with $w_1 = w_2 = w_3 = w_4 = 0.5, w_5 = 1$ and $B = 1$. Indeed, this mechanism extracts an expected revenue of 1.5 from the first three lists, an expected revenue of 1 from lists (32), (42), (43), an expected revenue of 1.5 from lists (5), (5), (5), and an expected revenue of 1.25 from the final three lists, for an overall average of $\frac{1.5+1+1.5+1.25}{4} = 1.3125$.

Meanwhile, consider the robust mechanism design problem for the same instance. A feasible solution is given by the following menu \mathcal{M} , with non-zero entries

$$(1/2, 1/2, 0, 0, 0), (1/2, 0, 1/2, 0, 0), (1/2, 0, 0, 1/2, 0), (0, 1/2, 1/2, 0, 0), (0, 1/2, 0, 1/2, 0), (0, 0, 1/2, 1/2, 0), (0, 0, 0, 0, 1).$$

The first six lists would choose the first six entries respectively, with the set of exposable entries always being a singleton. List $\ell = (5)$ would similarly have $E^{\mathcal{M}}(\ell) = \{(0, 0, 0, 0, 1)\}$. Finally, lists $\ell = (251), (351), (451)$ have $|E^{\mathcal{M}}(\ell)| = 2$, consisting of one entry that deterministically grants their 2nd choice and another entry that grants their 1st and 3rd choices with 50% probability each. However, regardless of which entry is chosen, the expected revenue would be 1.5. Therefore, the objective value (34) for this menu \mathcal{M} is $\frac{1.5+1+1.5+1.5}{4} = 1.375$, strictly greater than OPT^x . \square

Appendix C: Proofs from Section 3

Proof of Proposition 1. We first prove the forward direction, which is easy to see. To show that x is IC, note that constraints (4) and (5) are satisfied by definition. Constraints (3) are satisfied by considering two cases. First, if $\ell \in \mathcal{L}$ is such that $x_{\ell_{k'}}(\ell) = 1$ for some $k' \leq |\ell|$, then the LHS of (3) is 1 (and hence (3) is satisfied) as long as $k \geq k'$. On the other hand, if $k < k'$, then it must be the case that ℓ_1, \dots, ℓ_k are not in the assortment S , by definition of $x_j(\ell)$ in (6). Therefore, the RHS of (3) is 0 (and hence (3) is again satisfied). In the second case, ℓ is such that $x_j(\ell) = 0$ for all $j \in \ell$. This means that none of the items j on ℓ are in the assortment S . Then no matter what ℓ' the buyer reports, it is not possible for them to receive an item on their list, and thus the RHS of (3) is always 0 (and hence (3) is still satisfied). Finally, to see that $\text{REV}[x] = \text{REV}[S]$, note that $\Pr[j \succeq S_0]$ equals the sum of probabilities $p(\ell)$ over lists ℓ such that $x_j(\ell) = 1$, by definition, completing the proof of the forward direction.

To prove the converse direction, take any IC mechanism with $x_j(\ell) \in \{0, 1\}$ for all $\ell \in \mathcal{L}$ and $j \in \ell$. Define S to be the set of $j \in N$ for which there exists an ℓ' such that $x_j(\ell') = 1$, i.e. the set of items that are possible to obtain through some “lie” ℓ' . To show that (6) is satisfied, we again consider two cases. First, if $\ell \in \mathcal{L}$ is such that $x_{\ell_{k'}}(\ell) = 1$ for some $k' \leq |\ell|$, then IC constraints (3) imply that $x_{\ell_k}(\ell') = 0$ for any $k < k'$ and $\ell' \in \mathcal{L}$. In other words, none of the items lying before position k' in list ℓ are in the assortment S we defined, and hence $x_{\ell_{k'}}(\ell) = 1$ satisfies the description in (6). For the other part of description (6), the fact that $x_j(\ell) = 0$ for all $j \neq \ell_{k'}$ holds by constraints (4) and (5). In the second case, ℓ is such that $x_j(\ell) = 0$ for all $j \in \ell$. We can again use constraints (3) to see it must be the case that $j \notin S$ for any $j \in \ell$, completing the proof of the converse direction. \square

Proof of Proposition 2. It is immediate that f is non-negative and increasing, from the non-negativity constraints (5) on mechanism x . It is also immediate that f is upper-bounded by 1, from constraints (4). To establish (8), note that by the LP constraints (3), the probability of a type ℓ receiving one of their k' most-preferred items equals f evaluated on this set of items, i.e.

$$\sum_{j \in \ell_{1:k'}} x_j(\ell) = f(\ell_{1:k'}) \quad \forall \ell \in \mathcal{L}, k' = 1, \dots, |\ell|. \quad (37)$$

For any $\ell \in \mathcal{L}$ and $k = 1, \dots, |\ell|$, subtracting (37) with $k' = k - 1$ from (37) with $k' = k$ (or subtracting $f(\emptyset) = 0$ if $k = 1$), we see that the probability $x_{\ell_k}(\ell)$ of type ℓ receiving their k -th most preferred item equals $f(\ell_{1:k}) - f(\ell_{1:k-1})$, completing the proof. \square

Proof of Proposition 3. The optimization problem for OPT^f , using (9) as the expression for $\text{REV}[f]$, can be formulated by the following LP with decision variables $f(S)$:

$$\max \sum_{\rho} r_{\rho_{\text{end}}} (f(\rho) - f(\rho_{\text{pre}})) \Pr[\rho]$$

$$f(S) \leq f(S \cup \{j\}) \quad \forall S \subsetneq N, j \notin S \quad (38)$$

$$f(S) \in [0, 1] \quad \forall S \subseteq N \quad (39)$$

The matrix formed by the LHS of constraints (38) is totally unimodular, because all of its entries lie in $\{-1, 0, +1\}$, and every row contains exactly one $+1$ entry and one -1 entry (Schrijver 1998). Therefore, the feasible region defined by (38)–(39) is integral, and moreover since the objective function is linear, there must exist an integer optimal solution with $f(S) \in \{0, 1\}$ for all S . This defines an increasing boolean set function f , completing the proof of the first statement.

To show that $\text{REV}[\phi] = \text{REV}[f]$, note that when f is increasing and boolean-valued, $f(\rho) - f(\rho_{\text{pre}}) = f(\rho) \prod_{k=1}^{|\rho|-1} (1 - f(\rho_{1:k}))$. This in turn equals $\phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1}))$, by the definition of ϕ in (11). Substituting into the expression for $\text{REV}[f]$, we see that

$$\text{REV}[f] = \sum_{\rho} r_{\rho_{\text{end}}} \phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1})) \Pr[\rho],$$

completing the proof of the second statement. \square

Appendix D: Proofs from Section 4

Proof of Theorem 1. Our goal is to show that $\text{OPT}^S = \text{OPT}^x$. Since we have established in general that $\text{OPT}^S \leq \text{OPT}^x \leq \text{OPT}^{\phi}$, it suffices to show $\text{OPT}^{\phi} \leq \text{OPT}^S$. Under a Markov Chain choice model, consider the following relaxation on the optimal stopping value OPT^{ϕ} . First, instead of only adding an item j to the list when it is visited for the *first* time, we allow for stopping on an item j (ending with reward r_j) *any* time j is visited. Furthermore, we place no constraints on the stopping policy (whereas before ϕ could only make deterministic stopping decisions based on the unordered history H , and had to satisfy monotonicity constraints).

This is now an infinite-horizon optimal stopping problem which terminates in finite time w.p. 1. Since stopping is allowed on repeat visits to items j and there are no longer constraints on the stopping policy, the value-to-go under the optimal stopping policy is fully determined by the current item j , which we denote using $V[j]$. Moreover, these values-to-go must satisfy

$$V[j] = \max\{r_j, \sum_{j' \in N} \rho_{jj'} V[j']\} \quad \forall j \in N. \quad (40)$$

After solving system (40) which has $|N|$ equations and $|N|$ unknowns, a policy that stops on an item j if and only if $r_j \geq \sum_{j' \in N} \rho_{jj'} V[j']$ is optimal (see Bertsekas 1995). The expected revenue of this policy is equal to the expected revenue earned by the assortment $S = \{j \in N : r_j \geq \sum_{j' \in N} \rho_{jj'} V[j']\}$. Therefore, optimal stopping policies are no better than assortments under Markov Chain choice models, completing the proof of Theorem 1. \square

Appendix E: Proofs from Subsection 5.2

Proof of Lemma 1. We start with the definition of $\text{REV}[\phi]$ for an arbitrary monotone stopping policy ϕ , and expand $\Pr[\rho]$ as $\prod_{k=1}^{|\rho|} q(\rho_{1:k})$ (which holds by the definition of q in Definition 6):

$$\text{REV}[\phi] = \sum_{\rho} r_{\rho_{\text{end}}} \phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) q(\rho) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1})) q(\rho_{1:k}).$$

We split the sum on the RHS into the contribution from prefixes ρ that do not intersect S , and prefixes ρ' that end on S but do not intersect S before the end:

$$\begin{aligned} \text{REV}[\phi] &= \sum_{\rho: \rho \cap S = \emptyset} r_{\rho_{\text{end}}} \phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) q(\rho) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1})) q(\rho_{1:k}) \\ &\quad + \sum_{\substack{\rho': \rho'_{\text{end}} \in S, \\ \rho'_{\text{pre}} \cap S = \emptyset}} \phi_{\rho'_{\text{end}}}(\rho'_{\text{pre}}) \left(r_{\rho'_{\text{end}}} q(\rho') \prod_{k=1}^{|\rho'|-1} (1 - \phi_{\rho'_k}(\rho'_{1:k-1})) q(\rho'_{1:k}) \right). \end{aligned} \quad (41)$$

Note that we did not need to account for the contribution from prefixes that intersect S before the end, because policy ϕ would have stopped on the first intersection with S .

We now rewrite the expression inside the large parentheses in (41) as follows:

$$\begin{aligned} &r_{\rho'_{\text{end}}} q(\rho') \prod_{k=1}^{|\rho'|-1} (1 - \phi_{\rho'_k}(\rho'_{1:k-1})) q(\rho'_{1:k}) \\ &= r_{\rho'_{\text{end}}} \left(\prod_{k=1}^{|\rho'|} q(\rho'_{1:k}) \right) \left(1 - \sum_{k=1}^{|\rho'|-1} \phi_{\rho'_k}(\rho'_{1:k-1}) \prod_{k'=1}^{k-1} (1 - \phi_{\rho'_{k'}}(\rho'_{1:k'-1})) \right) \\ &= r_{\rho'_{\text{end}}} \prod_{k=1}^{|\rho'|} q(\rho'_{1:k}) - \sum_{k=1}^{|\rho'|-1} \left(r_{\rho'_{\text{end}}} \prod_{k'=k+1}^{|\rho'|} q(\rho'_{1:k'}) \right) \phi_{\rho'_k}(\rho'_{1:k-1}) q(\rho'_{1:k}) \left(\prod_{k'=1}^{k-1} (1 - \phi_{\rho'_{k'}}(\rho'_{1:k'-1})) q(\rho'_{1:k'}) \right). \end{aligned} \quad (42)$$

The first equality holds because $\prod_{k=1}^{|\rho'|-1} (1 - \phi_{\rho'_k}(\rho'_{1:k-1})) + \sum_{k=1}^{|\rho'|-1} \phi_{\rho'_k}(\rho'_{1:k-1}) \prod_{k'=1}^{k-1} (1 - \phi_{\rho'_{k'}}(\rho'_{1:k'-1}))$ sums to unity, describing the mutually exclusive and collectively exhaustive events of policy ϕ not stopping on any of the first $|\rho'| - 1$ elements of prefix ρ' , or stopping on one of them (say the k' 'th element) after not stopping before that (not stopping on elements $k' = 1, \dots, k-1$).

Substituting (42) into the expression inside large parentheses in (41), we obtain:

$$\begin{aligned} \text{REV}[\phi] &= \sum_{\rho: \rho \cap S = \emptyset} r_{\rho_{\text{end}}} \phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) q(\rho) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1})) q(\rho_{1:k}) + \sum_{\substack{\rho': \rho'_{\text{end}} \in S, \\ \rho'_{\text{pre}} \cap S = \emptyset}} r_{\rho'_{\text{end}}} \prod_{k=1}^{|\rho'|} q(\rho'_{1:k}) \\ &\quad - \sum_{\substack{\rho': \rho'_{\text{end}} \in S, \\ \rho'_{\text{pre}} \cap S = \emptyset}} \sum_{k=1}^{|\rho'|-1} \left(r_{\rho'_{\text{end}}} \prod_{k'=k+1}^{|\rho'|} q(\rho'_{1:k'}) \right) \phi_{\rho'_k}(\rho'_{1:k-1}) q(\rho'_{1:k}) \left(\prod_{k'=1}^{k-1} (1 - \phi_{\rho'_{k'}}(\rho'_{1:k'-1})) q(\rho'_{1:k'}) \right) \\ &= \sum_{\rho: \rho \cap S = \emptyset} r_{\rho_{\text{end}}} \phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) q(\rho) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1})) q(\rho_{1:k}) + \text{REV}[S] \\ &\quad - \sum_{\rho: \rho \cap S = \emptyset} \phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) q(\rho) \left(\prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1})) q(\rho_{1:k}) \right) \sum_{\substack{\rho': \rho'_{\text{end}} \in S, \\ \rho'_{\text{pre}} \cap S = \emptyset, \\ \rho'_{1:|\rho|} = \rho}} \left(r_{\rho'_{\text{end}}} \prod_{k'=|\rho|+1}^{|\rho'|} q(\rho'_{1:k'}) \right). \end{aligned}$$

In the third line above, we have used $\text{REV}[S]$ to replace a sum from the first line which described the revenue of the policy that stops on assortment S and stops nowhere else. In the fourth line above, we have let ρ denote $\rho'_{1:k}$ (a prefix disjoint from S) and exchanged the sums from the second line, pulling forward the terms which only depend on the first k elements of ρ' and rewriting them in terms of ρ . The inner sum in the fourth line is then over the ρ' which start identically as ρ , end on S , but do not intersect S before the end.

Having performed this exchange, we can now conveniently merge the sums from the third and fourth lines above to see that

$$\begin{aligned} & \text{REV}[\phi] - \text{REV}[S] \\ &= \sum_{\rho: \rho \cap S = \emptyset} \left(r_{\rho_{\text{end}}} - \sum_{\substack{\rho': \rho'_{\text{end}} \in S, \\ \rho'_{\text{pre}} \cap S = \emptyset, \\ \rho'_{1:|\rho|} = \rho}} r_{\rho'_{\text{end}}} \prod_{k=|\rho|+1}^{|\rho'|} q(\rho'_{1:k}) \right) \phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) q(\rho) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1})) q(\rho_{1:k}) \\ &= \sum_{\rho: \rho \cap S = \emptyset} \left(r_{\rho_{\text{end}}} - \sum_{j \in S} r_j \Pr[j \succeq S_0 | \rho] \right) \Pr[\rho] \phi_{\rho_{\text{end}}}(\rho_{\text{pre}}) \prod_{k=1}^{|\rho|-1} (1 - \phi_{\rho_k}(\rho_{1:k-1})). \end{aligned}$$

where the second equality uses the fact that ρ is disjoint from S to simplify the expression inside large parentheses, and also the fact that $q(\rho) \prod_{k=1}^{|\rho|-1} q(\rho_{1:k}) = \Pr[\rho]$ which holds by definition of q . Substituting in the definition of S -adjusted prices, this completes the proof of Lemma 1. \square

Proof of Lemma 2. Fix an assortment $S \subsetneq N$ along with an item $j \notin S$, and consider the realizable prefixes ρ with $\rho_{\text{end}} = j$ and $\rho \cap S = \emptyset$. Construct an undirected graph, with a vertex for each such prefix, and an edge between two prefixes ρ, ρ' if and only if $\rho_{\text{pre}} \not\supseteq \rho'_{\text{pre}}$ and $\rho'_{\text{pre}} \not\supseteq \rho_{\text{pre}}$ (neither is contained within the other when viewed as sets). Definition 11 implies that the futures from adjacent prefixes ρ, ρ' in this graph must dominate each other, and since S intersects neither ρ nor ρ' , it must be that inequality (12) holds with equality:


$$\Pr[j' \succeq S_0 | \rho] = \Pr[j' \succeq S_0 | \rho'] \quad \forall j' \in S. \quad (43)$$

Consider the connected components of this graph. By transitivity, the probability $\Pr[j' \succeq S_0 | \rho]$ must be identical across all vertices ρ in a connected component, for all $j' \in S$. Note that it was important for the graph to only include vertices for the *realizable* prefixes ρ with $\Pr[\rho] > 0$; otherwise there can only be very few¹⁰ connected components. Let C denote the number of components in this graph.

Consider ρ, ρ' in different components of this graph. Since ρ and ρ' are not adjacent, it must be that one of them contains the other (otherwise there would be an edge between ρ and ρ'), say $\rho \supseteq \rho'$. We claim that then any other prefix ϱ adjacent to ρ must also satisfy $\varrho \supseteq \rho'$. This is easy to see—since ϱ is non-adjacent to ρ' , either $\varrho \supseteq \rho'$ or $\varrho \subseteq \rho'$, but if $\varrho \subseteq \rho'$, then $\varrho \subseteq \rho$ by transitivity, contradicting the fact that ϱ is adjacent to ρ . By propagating this argument throughout the connected component containing ρ , we establish that all

¹⁰ If all prefixes were included, then there would be exactly three connected components: one for vertex (j) , one for all the prefixes ρ with $\rho_{\text{end}} = j$ and $\rho_{\text{pre}} = N \setminus S \setminus \{j\}$, and one for all other prefixes satisfying $\rho_{\text{end}} = j$ and $\rho \cap S = \emptyset$.

Figure 3 Example illustrating the details in the definition of history-monotone futures.



prefixes in this component contain ρ' . Similarly, we can argue that all prefixes in the connected component of ρ' are contained within ρ (and also contained within any ϱ in the component of ρ).

Therefore, we can take a single representative ρ_c from each of the components $c = 1, \dots, C$, and know that for any c, c' , either $\rho_c \subseteq \rho_{c'}$ or $\rho_{c'} \subseteq \rho_c$ (or both). This defines a *total preorder* on the connected components—a binary relation \subseteq which is reflexive, transitive, and any pair of elements are comparable, but the relation is not necessarily antisymmetric ($\rho \subseteq \rho'$ and $\rho' \subseteq \rho$ does not imply that $\rho = \rho'$ as ordered prefixes). Nonetheless, we can take the equivalence classes with respect to this preorder to get a total order.

Suppose that there are T equivalence classes, where $T \leq C$, and let $\mathcal{T}(t)$ denote set of all prefixes lying in connected components belonging to equivalence class t , ordered so that whenever $t > t'$, and $\rho \in \mathcal{T}(t), \rho' \in \mathcal{T}(t')$, we have that $\rho \supseteq \rho'$ and $\rho \not\subseteq \rho'$. This implies that $\rho \supseteq \rho'$ as sets. In conjunction with (43), this completes the proof of condition 1 in Lemma 2.

To establish condition 2, we claim that: an equivalence class either consists of a single component, or all of its components are isolated vertices. To show this, suppose for contradiction that an equivalence class includes two components, one which includes two vertices. Let ρ, ϱ denote adjacent prefixes in the same component and ρ' denote a prefix in another component. Since the components are in the same equivalence class, we have that $\rho \subseteq \rho' \subseteq \varrho$, but this implies that $\rho \subseteq \varrho$, contradicting the adjacency of ρ and ϱ . Having shown our claim, we can now see that equivalence classes consisting of a single component satisfy condition (2b) (where we again used (43)), while equivalence classes consisting of isolated vertices satisfy condition (2a). This completes the proof of Lemma 2. \square

E.1. Example Illustrating History-monotone Futures and Tier Decomposition

EXAMPLE 6 (HISTORY-MONOTONE FUTURES). Consider the list distribution over four items A, B, C, D whose tree diagram is shown in Figure 3. We verify that it has history-monotone futures.

There are four realizable prefixes with endpoint A — $(CBA), (CDBA), (DBA), (DCBA)$. Definition 11 is satisfied since all of their futures are empty, dominating each other.

There are five realizable prefixes with endpoint B — $(B), (CB), (DB), (CDB), (DCB)$. Since A is the only item that can appear in the future, domination is fully determined by the probabilities of visiting A from these prefixes, which are $0, 1/2, 1/2, 3/4, 1$ respectively. We verify Definition 11 using the “equivalent” condition stated at the end. (CB) and (DB) have futures which are *not dominated by* (B) , which causes no violation, since their respective histories are $\{C\}$ and $\{D\}$, which contain \emptyset . Meanwhile, (CDB) and (DCB) have futures which are not dominated by the previous ones, which also causes no violation, since for both (CDB) and (DCB) the history is $\{C, D\}$, which contains the previous ones. Finally, (DCB) has a future which is not dominated by (CDB) , which again causes no violation, since its history $\{C, D\}$ weakly contains (more specifically, is identical to) the history for (CDB) .

There are two realizable prefixes with endpoint C — $(C), (DC)$. The future from (DC) must dominate in order to satisfy Definition 11, which it does, because it is (BA) w.p. 1 while the future from (C) (ignoring item D) is (BA) w.p. $5/8$ and (B) w.p. $3/8$. Note it was important here to ignore D in the definition of domination.

There are two realizable prefixes with endpoint D — $(D), (CD)$. The future from (CD) must dominate in order to satisfy Definition 11, which it does, because ignoring item C , both futures are (BA) w.p. $3/4$ and (B) w.p. $1/4$. Note that Definition 11 would be violated if the transition probability from prefix (CDB) to A was reduced below $3/4$. \square

Example 6 can also be used to illustrate the details in the tier decomposition Lemma 2. Specifically, consider the case where $S = \{A\}, j = B$, and recall that there were five realizable prefixes with endpoint j — $(B), (CB), (DB), (CDB), (DCB)$ —none of which intersect S . The graph described in the proof of Lemma 2 will have four connected components, with an edge between $(CB), (DB)$ along with three isolated vertices. $(CDB), (DCB)$, will later be put into the same tier, resulting in

$$\mathcal{T}(1) = \{(B)\}, \quad \mathcal{T}(2) = \{(CB), (DB)\}, \quad \mathcal{T}(3) = \{(CDB), (DCB)\}.$$

Property (1) in Lemma 2 is immediately verified from the calculations in Example 6. Meanwhile, tier $t = 2$ satisfies property (2a) while tier $t = 3$ satisfies property (2b). Note that it is possible for two prefixes in a tier of type (2b) to have different futures; in this example, $\Pr[A \succeq \{A\}_0 | (CDB)] = 3/4$ which is different from $\Pr[A \succeq \{A\}_0 | (DCB)] = 1$.

Appendix F: Proofs from Subsection 5.3

Proof of Proposition 4. Take realizable prefixes ρ, ρ' with the same endpoint. We show that regardless of any containment relations between ρ_{pre} and ρ'_{pre} , as long as assortment S is disjoint from both ρ and ρ' , it will be the case that $\Pr[j \succeq S_0 | \rho] = \Pr[j \succeq S_0 | \rho']$ for any $j \in S$. To see this, consider the future from ρ . This is generated by a random path on the Markov Chain starting from state ρ_{end} , where any newly-visited items not on ρ are added to the list. The future from ρ' is generated by an identically-distributed path, except the newly-visited items on ρ' are excluded instead. However, these differences do not affect the event $j \succeq S_0$, since $j \in S$ and S is disjoint from both ρ and ρ' . Therefore, $\Pr[j \succeq S_0 | \rho] = \Pr[j \succeq S_0 | \rho']$ for all $j \in S$, satisfying the condition (12) for domination as equality and completing the proof of Proposition 4. \square

Proof of Proposition 5. Take realizable prefixes ρ, ρ' with the same endpoint j , and suppose that j is in nest i . By the rules of drawing the balls, ρ and ρ' (when viewed as sets) can only differ on items in N_i . Take any assortment S disjoint from ρ and ρ' , and we would like to show that $\Pr[j' \succeq S_0 | \rho] = \Pr[j' \succeq S_0 | \rho']$ for all $j' \in S$. Clearly if $j' \in N_i$ then this is true, since S is disjoint from any items in N_i which have already appeared in ρ or ρ' . On the other hand, if $j' \notin N_i$, then this is also true, because either both probabilities are 0 (if S contains any items in N_i), or both probabilities are calculated from an identical starting point where the items in nest N_i have all been drawn (recall that ρ and ρ' do not setwise differ outside of N_i). \square

Proof of Proposition 6. Take realizable prefixes ρ, ρ' with $\rho_{\text{end}} = \rho'_{\text{end}}$ and $\rho_{\text{pre}} \not\subseteq \rho'_{\text{pre}}$. Note that this implies $\rho_{\text{pre}} \neq \emptyset$, so ρ has length at least 2. Therefore, the future conditional on ρ did not change from before. On the other hand, the future conditional on ρ' either did not change (if $|\rho'| \geq 2$), or $\Pr[j \succeq S_0 | \rho']$ decreased for every $j \in S$ (if $|\rho'| = 1$). In either case, the future from ρ continues to dominate the future from ρ' , verifying the definition of history-monotone futures and completing the proof. \square

Proof of Proposition 7. WOLOG normalize w_0 to 1. Suppose there are 3 items, which have weight $w > 0$, in the same nest, which has dissimilarity parameter $\gamma \in (0, 1)$. Let $N = \{1, 2, 3\}$ denote the set of 3 items. For a Markov Chain choice model to capture the Nested Logit choice probabilities in (19), its parameters $\lambda_1, \lambda_2, \lambda_3, \rho_{12}, \rho_{13}, \rho_{21}, \rho_{23}, \rho_{31}, \rho_{32}$ must satisfy the following constraints.

First, λ_j must equal the probability of item j being chosen from the full assortment N , for all $j \in N$. Therefore,

$$\lambda_j = \frac{(3w)^\gamma}{1 + (3w)^\gamma} \cdot \frac{1}{3} \quad \forall j = 1, 2, 3.$$

Second, for any assortment consisting of two distinct items $j, j' \in N$, note that the probability of j being chosen from assortment $\{j, j'\}$ is $\lambda_j + \lambda_{j''} \sigma_{j'', j}$, where j'' is the item that is not j or j' . Therefore,

$$\begin{aligned} \frac{(2w)^\gamma}{1 + (2w)^\gamma} \cdot \frac{1}{2} &= \lambda_j + \lambda_{j''} \sigma_{j'', j} \\ \frac{(2w)^\gamma}{1 + (2w)^\gamma} \cdot \frac{1}{2} &= \frac{(3w)^\gamma}{1 + (3w)^\gamma} \cdot \frac{1}{3} (1 + \sigma_{j'', j}) \end{aligned}$$

which implies that

$$\sigma_{j'', j} = \frac{(3w)^{-\gamma} + 1}{(2w)^{-\gamma} + 1} \cdot \frac{3}{2} - 1 \quad \forall j \neq j'' \in \{1, 2, 3\}.$$

Finally, for a singleton assortment $\{j\}$, the Nested Logit probability of j being chosen is

$$\frac{w^\gamma}{1 + w^\gamma}. \quad (44)$$

This must equal the probability of visiting j before terminal state 0 in the Markov Chain, which can be computed via taking a sum over the probability of visiting j for the first time after transitioning exactly k times (to a non-zero state) in the Markov Chain:

$$\begin{aligned} &\frac{(3w)^\gamma}{1 + (3w)^\gamma} \cdot \frac{1}{3} + \sum_{k=1}^{\infty} 2 \frac{(3w)^\gamma}{1 + (3w)^\gamma} \cdot \frac{1}{3} \left(\frac{(3w)^{-\gamma} + 1}{(2w)^{-\gamma} + 1} \cdot \frac{3}{2} - 1 \right)^k \\ &= \frac{(3w)^\gamma}{1 + (3w)^\gamma} \cdot \frac{1}{3} \left(1 + 2 \frac{1}{\left(\frac{(3w)^{-\gamma} + 1}{(2w)^{-\gamma} + 1} \cdot \frac{3}{2} - 1 \right)^{-1} - 1} \right). \end{aligned} \quad (45)$$

However, it can be checked that this expression does not equal (44) for general values of w and γ (it can also be verified that (45) *does* equal (44) when $\gamma = 1$, which is the special case of MNL). As a concrete example, taking $w = 1$, (45) can be re-written as

$$\frac{3^{\gamma-1}}{1+3^\gamma} \left(1 + 2 \frac{1}{\left(\frac{3^{1-\gamma}+3}{2^{1-\gamma}+2} - 1\right)^{-1} - 1} \right) = \frac{3^{\gamma-1}}{1+3^\gamma} \left(1 + \frac{2}{\frac{2^{1-\gamma}+2}{3^{1-\gamma}-2^{1-\gamma}+1} - 1} \right)$$

which only equals $\frac{1}{2}$ (the value of (44)) when $\gamma = 1$. Therefore, a Nested Logit choice model with 3 items that have equal weights and are in the same nest is not captured by a Markov Chain, completing the proof.

□

Proof of Proposition 8. When there are only 3 items, the following identities are sufficient for a list distribution defined through its probabilistic tree (refer to Definition 6) to be consistent with a given set of choice probabilities. For brevity, we will omit the inner parentheses when prefixes are expressed inside the $q(\cdot)$ function.

$$q(j) = \Pr[j \succeq N_0] \quad \forall j \in N$$

$$q(j) + q(j')q(j'j) = \Pr[j \succeq (N \setminus \{j'\})_0] \quad \forall j' \neq j \in N$$

$$q(j) + q(j')q(j'j) + q(j'')q(j''j) + q(j')q(j'j'')q(j'j''j) + q(j'')q(j''j')q(j''j'j) = \Pr[j \succeq \{j\}_0] \quad \forall j \in N$$

In the third set of identities, j' and j'' refer to the two items other than j . Note that this system has $3 + 6 + 6 = 15$ variables but only $3 + 6 + 3 = 12$ equations. We will allow for a full specification of the q -probabilities by imposing that $q(j'j''j) = q(j''j'j)$ for all $j \in N$, effectively adding 3 equations. We show this always results in a list distribution with history-monotone futures.

To verify history-monotone futures for list distributions on 3 items, one only needs to check that

$$\Pr[j \succeq \{j\}_0 | (j''j')] \geq \Pr[j \succeq \{j\}_0 | (j')] \quad (46)$$

for all permutations of the items j, j', j'' . (Note that in (46) we have assumed $S = \{j\}$; if $|S| \geq 2$, then there cannot be two distinct prefixes in $N \setminus S$ with the same endpoint, and hence the condition holds vacuously.)

We now proceed to check this for an arbitrary arrangement of j, j', j'' , which for brevity we relabel as A, B, C respectively.

Note that the LHS of (46) equals $q(CBA)$, while the RHS of (46) equals $q(BA) + q(BC)q(BCA)$. The goal is to show that $q(CBA) \geq q(BA) + q(BC)q(BCA)$ when the q -values are defined through the identities above and the choice probabilities comes from a Nested Logit choice model. To solve for $q(CBA)$, which is the same as $q(BCA)$, we can write:

$$\begin{aligned} q(A) + q(B)q(BA) + q(C)q(CA) + (q(B)q(BC) + q(C)q(CB))q(CBA) &= \Pr[A \succeq \{A\}_0] \\ (q(A) + q(B)q(BA)) + (q(A) + q(C)q(CA)) + (q(B)q(BC) + q(C)q(CB))q(CBA) &= \Pr[A \succeq \{A\}_0] + q(A) \\ \Pr[A \succeq \{A, C\}_0] + \Pr[A \succeq \{A, B\}_0] + (q(B)q(BC) + q(C)q(CB))q(CBA) &= \Pr[A \succeq \{A\}_0] + \Pr[A \succeq N_0] \end{aligned}$$

This allows us to deduce:

$$q(CBA) = \frac{\Pr[A \succeq \{A\}_0] - \Pr[A \succeq \{A, B\}_0] - \Pr[A \succeq \{A, C\}_0] + \Pr[A \succeq \{A, B, C\}_0]}{\Pr[C \succeq \{A, C\}_0] - \Pr[C \succeq \{A, B, C\}_0] + \Pr[B \succeq \{A, B\}_0] - \Pr[B \succeq \{A, B, C\}_0]} \quad (47)$$

Meanwhile, we can write:

$$\frac{q(BA)}{1 - q(BC)} = \frac{q(B)q(BA)}{q(B) - q(B)q(BC)} = \frac{\Pr[A \succeq \{A, C\}_0] - \Pr[A \succeq \{A, B, C\}_0]}{\Pr[B \succeq \{A, B, C\}_0] - (\Pr[C \succeq \{A, C\}_0] - \Pr[C \succeq \{A, B, C\}_0])} \quad (48)$$

To prove (46), or equivalently $q(CBA) \geq q(BA) + q(BC)q(CBA)$, it suffices to prove that the expression on the RHS of (47) is at least the expression on the RHS of (48).

Since there is a single nest, the Nested Logit choice probability for an item j being chosen from an assortment S can be expressed as (setting $w_0 = 1$ and ignoring the nest subscript i):

$$\frac{w_j}{(\sum_{j' \in S} w_{j'})^{1-\gamma} + (\sum_{j' \in S} w_{j'})}$$

Moreover, all of the assortments in expressions (47) and (48) include item A . Therefore, we let $d(S)$ denote the denominator $(w_A + \sum_{j' \in S} w_{j'})^{1-\gamma} + (w_A + \sum_{j' \in S} w_{j'})$ where S is a subset of $\{B, C\}$. We also hereafter let a, b, c be shorthand for the weights w_A, w_B, w_C respectively. Then, the desired relationship (47) \geq (48) is equivalent to

$$\frac{\frac{a}{d(\emptyset)} - \frac{a}{d(B)} - \frac{a}{d(C)} + \frac{a}{d(BC)}}{\frac{b}{d(B)} - \frac{b}{d(BC)} + \frac{c}{d(C)} - \frac{c}{d(BC)}} \geq \frac{\frac{a}{d(C)} - \frac{a}{d(BC)}}{\frac{b}{d(BC)} + \frac{c}{d(BC)} - \frac{c}{d(C)}} \quad (49)$$

where we have omitted braces and commas in the input to the function d . Since the denominators are non-negative, we can cross-multiply and (49) is equivalent to

$$\left(\frac{1}{d(\emptyset)} - \frac{1}{d(B)} - \frac{1}{d(C)} + \frac{1}{d(BC)} \right) \left(\frac{b+c}{d(BC)} - \frac{c}{d(C)} \right) \geq \left(\frac{1}{d(C)} - \frac{1}{d(BC)} \right) \left(\frac{b}{d(B)} + \frac{c}{d(C)} - \frac{b+c}{d(BC)} \right). \quad (50)$$

After expanding (50) and canceling terms that appear on both sides, we are left with

$$\frac{b+c}{d(\emptyset)d(BC)} + \frac{c}{d(B)d(C)} \geq \frac{b}{d(B)d(C)} + \frac{c}{d(\emptyset)d(C)} + \frac{c}{d(B)d(BC)}.$$

Multiplying both sides by $d(\emptyset)d(B)d(C)d(BC)$ and substituting in the definition of d , we get

$$\begin{aligned} & (b+c)((a+b)^{1-\gamma} + a+b)((a+c)^{1-\gamma} + a+c) + c(a^{1-\gamma} + a)((a+b+c)^{1-\gamma} + a+b+c) \\ & \geq b(a^{1-\gamma} + a)((a+b+c)^{1-\gamma} + a+b+c) + c((a+b)^{1-\gamma} + a+b)((a+b+c)^{1-\gamma} + a+b+c) + c(a^{1-\gamma} + a)((a+c)^{1-\gamma} + a+c). \end{aligned}$$

Note that after expanding, any products without a $(\cdot)^{1-\gamma}$ term cancel out on both sides. Hence, it suffices to prove

$$\begin{aligned} & b(a+b)^{1-\gamma}(a+c)^{1-\gamma} + b(a+b)^{1-\gamma}(a+\phi) + b(a+b)(a+c)^{1-\gamma} \\ & + c(a+b)^{1-\gamma}(a+c)^{1-\gamma} + \cancel{c(a+b)^{1-\gamma}(a+c)} + c(\phi+b)(a+c)^{1-\gamma} + ca^{1-\gamma}(a+b+c)^{1-\gamma} + ca^{1-\gamma}(\phi+\beta+\phi) + \cancel{ca(a+b+c)^{1-\gamma}} \\ & \geq ba^{1-\gamma}(a+b+c)^{1-\gamma} + ba^{1-\gamma}(a+b+\phi) + ba(a+b+c)^{1-\gamma} \\ & + c(a+b)^{1-\gamma}(a+b+c)^{1-\gamma} + c(a+b)^{1-\gamma}(\phi+\beta+\phi) + c(\phi+b)(a+b+c)^{1-\gamma} + ca^{1-\gamma}(a+c)^{1-\gamma} + \cancel{ca^{1-\gamma}(a+c)} + \cancel{ca(a+c)^{1-\gamma}} \end{aligned}$$

which after cancellations is reduced to

$$\begin{aligned} & b(a+b)^{1-\gamma}(a+c)^{1-\gamma} + b(a+b)^{1-\gamma}(a) + b(a+b)(a+c)^{1-\gamma} + c(a+b)^{1-\gamma}(a+c)^{1-\gamma} + cb(a+c)^{1-\gamma} + ca^{1-\gamma}(a+b+c)^{1-\gamma} \\ & \geq ba^{1-\gamma}(a+b+c)^{1-\gamma} + ba^{1-\gamma}(a+b) + ba(a+b+c)^{1-\gamma} + c(a+b)^{1-\gamma}(a+b+c)^{1-\gamma} + cb(a+b+c)^{1-\gamma} + ca^{1-\gamma}(a+c)^{1-\gamma}. \end{aligned}$$

Now, we bifurcate the products on both sides depending on whether they contain one or two $(\cdot)^{1-\gamma}$ terms. We separately show that the LHS is at least the RHS for both types of terms. That is, we show

$$\begin{aligned} & b(a+b)^{1-\gamma}(a+c)^{1-\gamma} + c(a+b)^{1-\gamma}(a+c)^{1-\gamma} + ca^{1-\gamma}(a+b+c)^{1-\gamma} \\ & \geq ba^{1-\gamma}(a+b+c)^{1-\gamma} + c(a+b)^{1-\gamma}(a+b+c)^{1-\gamma} + ca^{1-\gamma}(a+c)^{1-\gamma}; \end{aligned} \quad (51)$$

$$\begin{aligned} & ba(a+b)^{1-\gamma} + b(a+b)(a+c)^{1-\gamma} + cb(a+c)^{1-\gamma} \\ & \geq b(a+b)a^{1-\gamma} + ba(a+b+c)^{1-\gamma} + cb(a+b+c)^{1-\gamma}. \end{aligned} \quad (52)$$

Doing so would suffice to establish (46) and hence complete the proof of the proposition.

Having separated the original inequality into (51) and (52), note that these new inequalities are *homogeneous*, in that the total degree of every term is the same—in (51), the total degree of every term is $3 - 2\gamma$; in (52), the total degree of every term is $3 - \gamma$. Therefore, during the proof of both (51) and (52), we can WOLOG assume that $a + b + c = 1$. Also for brevity, we will let $\delta := 1 - \gamma \in [0, 1]$.

We now rewrite (51) as

$$\begin{aligned} & (b+c)(a+b)^\delta(a+c)^\delta + ca^\delta \geq ba^\delta + c(a+b)^\delta + ca^\delta(a+c)^\delta \\ \iff & (b+c)(1+b/a)^\delta(1-b)^\delta + c \geq b+c(1+b/a)^\delta + c(1-b)^\delta \\ \iff & b((1+b/a)^\delta(1-b)^\delta - 1) \geq c((1+b/a)^\delta + (1-b)^\delta - (1+b/a)^\delta(1-b)^\delta - 1) \\ \iff & \frac{b}{c} \geq \frac{((1+b/a)^\delta - 1)(1 - (1-b)^\delta)}{((1+b/a)^\delta(1-b)^\delta - 1)} \end{aligned} \quad (53)$$

Our goal is to show that the maximum of the RHS of (53) over $\delta \in [0, 1]$ occurs at $\delta = 1$, given any values of a, b satisfying $b/a \geq b$ (which is true since $a < 1$). We will equivalently show that

$$\log((1+b/a)^\delta - 1) + \log(1 - (1-b)^\delta) - \log((1+b/a)^\delta(1-b)^\delta - 1)$$

is maximized at $\delta = 1$. Taking the derivative with respect to δ , we get

$$\frac{(1+b/a)^\delta \log(1+b/a)}{(1+b/a)^\delta - 1} - \frac{(1-b)^\delta \log(1-b)}{1 - (1-b)^\delta} - \frac{(1+b/a)^\delta(1-b)^\delta \log((1+b/a)(1-b))}{(1+b/a)^\delta(1-b)^\delta - 1}.$$

We want to show that this derivative is non-negative, i.e.

$$\begin{aligned} & \frac{\log(1+b/a)}{1 - (1+b/a)^{-\delta}} \geq \frac{\log(1-b)}{(1-b)^{-\delta} - 1} + \frac{\log(1+b/a) + \log(1-b)}{1 - (1+b/a)^{-\delta}(1-b)^{-\delta}} \\ \iff & \frac{\log(1+b/a)}{1 - (1+b/a)^{-\delta}} - \frac{\log(1+b/a)}{1 - ((1+b/a)(1-b))^{-\delta}} \geq \frac{\log(1-b)}{(1-b)^{-\delta} - 1} + \frac{\log(1-b)}{1 - ((1+b/a)(1-b))^{-\delta}} \\ \iff & \log(1+b/a) \frac{(1+b/a)^{-\delta} - ((1+b/a)(1-b))^{-\delta}}{(1 - (1+b/a)^{-\delta})(1 - ((1+b/a)(1-b))^{-\delta})} \geq \log(1-b) \frac{(1-b)^{-\delta} - ((1+b/a)(1-b))^{-\delta}}{((1-b)^{-\delta} - 1)(1 - ((1+b/a)(1-b))^{-\delta})} \end{aligned}$$

$$\begin{aligned}
&\Leftrightarrow \log(1+b/a) \frac{1-(1-b)^{-\delta}}{(1+b/a)^\delta-1} \geq \log(1-b) \frac{1-(1+b/a)^{-\delta}}{1-(1-b)^\delta} \\
&\Leftrightarrow \log\left(\frac{1}{1-b}\right) \frac{1-(1+b/a)^{-\delta}}{1-(1-b)^\delta} \geq \log(1+b/a) \frac{(1-b)^{-\delta}-1}{(1+b/a)^\delta-1} \\
&\Leftrightarrow \frac{(1-(1+b/a)^{-\delta})((1+b/a)^\delta-1)}{\log(1+b/a)} \geq \frac{(1-(\frac{1}{1-b})^{-\delta})((\frac{1}{1-b})^\delta-1)}{\log(\frac{1}{1-b})} \tag{54}
\end{aligned}$$

where we emphasize that in the third line, $1 - ((1+b/a)(1-b))^{-\delta} \geq 1 - ((1 + \frac{b}{1-b})(1-b))^{-\delta} = 0$, and in the final equivalence, all terms moved to the other side are also non-negative, and hence we never had to flip the inequality. Finally, to establish (54), note that it suffices to show the function $\frac{(1-y^{-\delta})(y^\delta-1)}{\log y} = \frac{y^\delta+y^{-\delta}-2}{\log y}$ is increasing over $y \geq 1$, since $1+b/a \geq 1 + \frac{b}{1-b} = \frac{1}{1-b} \geq 1$. To show that $\frac{y^\delta+y^{-\delta}-2}{\log y}$ is increasing in y , we take the derivative with respect to y , which can be checked to equal

$$\frac{y^\delta \log y^\delta - y^\delta - \frac{\log y^\delta}{y^\delta} - \frac{1}{y^\delta} + 2}{y(\log y)^2},$$

whose numerator can be seen to be non-negative over $[1, \infty)$ using a univariate plot of the variable y^δ . This completes the proof of (51).

To show the other inequality (52), we cancel out b from both sides, and then similarly rewrite it using $a+b+c=1$, $\delta=1-\gamma$:

$$\begin{aligned}
&a(a+b)^\delta + (a+b)(a+c)^\delta + c(a+c)^\delta \geq (a+b)a^\delta + a+c \\
&\Leftrightarrow a(a+b)^\delta + (1-b)^\delta \geq (a+b)a^\delta + 1-b \\
&\Leftrightarrow (1-b)^\delta - (1-b) \geq (a+b)a^\delta - a(a+b)^\delta \tag{55}
\end{aligned}$$

Our goal is to show that holding b and δ fixed, the RHS of (55) is increasing in a . This would complete the proof because the maximum possible value of a is $1-b$, in which case the RHS equals $(1-b)^\delta - (1-b)$, identical to the LHS. We take the derivative of the RHS of (55) with respect to a and aim to show it is non-negative:

$$\begin{aligned}
&a^\delta + (a+b)\delta a^{\delta-1} - (a+b)^\delta - a\delta(a+b)^{\delta-1} \geq 0 \\
&\Leftrightarrow 1 + (1+b/a)\delta - (1+b/a)^\delta - a \frac{\delta(1+b/a)^\delta}{a+b} \geq 0 \\
&\Leftrightarrow 1 + y\delta - y^\delta - \delta y^{\delta-1} \geq 0 \tag{56}
\end{aligned}$$

where in the final equivalence we have let $y := 1+b/a$, a quantity ranging over $[1, \infty)$. To show (56), we claim that its minimum value over $[1, \infty)$ is achieved at $y=1$, in which case it equals 0. To see why, we take the derivative of the LHS with respect to y :

$$1 + \delta - \delta y^{\delta-1} - \delta(\delta-1)y^{\delta-2} = 1 + \delta(1 - \frac{1}{y^{1-\delta}}) + \delta(1-\delta)y^{\delta-2}$$

which is clearly non-negative over $[1, \infty)$. Therefore, inequality (56) always holds, which implies (55), which in turn implies (52). Having established (51) and (52), this completes the entire proof. \square

Proof of Proposition 9. WOLOG normalize w_0 to 1. Let w denote the common weight of the items and γ denote the dissimilarity parameter. According to (19), the probability of any item being chosen when it is offered in an assortment of size k is

$$P_k := \frac{(kw)^\gamma}{1 + (kw)^\gamma} \cdot \frac{1}{k} = \frac{1}{k(1 + (kw)^{-\gamma})} \quad \forall k = 1, \dots, n. \quad (57)$$

We now define a list distribution which is consistent with the Nested Logit choice probabilities P_k and has history-monotone futures. We define this list distribution by the transition probabilities q for its tree diagram (see Definition 6). Our tree will be symmetric, and we let q_k denote the probability of transitioning to the node for any unvisited item after visiting exactly $k - 1$ previous item nodes, for all $k = 1, \dots, n$. Note that the probability of transitioning to the terminal node after visiting exactly $k - 1$ previous item nodes will be $1 - (n - k + 1)q_k$.

By symmetry, our tree is completely specified by the probabilities q_1, \dots, q_n . In order to be consistent with the choice probabilities P_k , they must satisfy the following system of equations:

$$P_k = \sum_{k'=0}^{n-k} q_{k'+1} \prod_{k''=1}^{k'} (n - k - k'' + 1)q_{k''} \quad \forall k = 1, \dots, n. \quad (58)$$

To explain the RHS, note that there are $n - k$ items not in the assortment. Any of the k items in the assortment gets chosen if and only if it appears on the list following exactly k' distinct ordered items not in the assortment, with k' ranging from 0 to $n - k$ inclusive.

Solving for the transition probabilities in the tree. We now inductively establish the identities

$$q_{m+1} \prod_{m'=1}^m q_{m'} = \frac{1}{m!} \sum_{m'=0}^m (-1)^{m-m'} \binom{m}{m'} P_{n-m'} \quad \forall m = 0, \dots, n - 1. \quad (59)$$

To do so, note that the base case $m = 0$ says $q_1 = P_n$ which follows from (58) with $k = n$. Now, suppose we have verified identity (59) for all values smaller than a particular $m \in \{1, \dots, n - 1\}$.

Using (58) with $k = n - m$, we get

$$\begin{aligned} P_{n-m} &= m!q_{m+1} \prod_{m'=1}^m q_{m'} + \sum_{k'=0}^{m-1} q_{k'+1} \prod_{k''=1}^{k'} (n - k - k'' + 1)q_{k''} \\ &= m!q_{m+1} \prod_{m'=1}^m q_{m'} + \sum_{k'=0}^{m-1} \frac{1}{k'!} \sum_{m'=0}^{k'} (-1)^{k'-m'} \binom{k'}{m'} P_{n-m'} \left(\prod_{k''=1}^{k'} (m - k'' + 1) \right) \end{aligned}$$

$$\begin{aligned}
&= m!q_{m+1} \prod_{m'=1}^m q_{m'} + \sum_{m'=0}^{m-1} P_{n-m'} \sum_{k'=m'}^{m-1} (-1)^{k'-m'} \frac{1}{k'!} \binom{k'}{m'} \left(\prod_{k''=1}^{k'} (m - k'' + 1) \right) \\
&= m!q_{m+1} \prod_{m'=1}^m q_{m'} + \sum_{m'=0}^{m-1} (-1)^{m-1-m'} P_{n-m'} \sum_{k'=m'}^{m-1} (-1)^{m-1-k'} \binom{m}{k'} \binom{k'}{m'} \\
&= m!q_{m+1} \prod_{m'=1}^m q_{m'} - \sum_{m'=0}^{m-1} (-1)^{m-m'} P_{n-m'} \binom{m}{m'}
\end{aligned}$$

where the second equality applies the induction hypothesis, the third equality switches sums, and the final equality uses the Principle of Inclusion-Exclusion on the number of ways to choose a size- m' subset from m elements by first selecting k' elements to choose from. This completes the induction and establishes (59). We can now solve for the values of q_1, \dots, q_n via taking the quotient of adjacent identities in this family.

Showing that the q -values exhibit a monotonicity property. Our goal is to show that the resulting values of q_k satisfy the following monotonicity property, which will later be useful for establishing history-monotone futures:

$$\frac{1}{q_k} \geq \frac{1}{q_{k+1}} + 1 \quad \forall k = 1, \dots, n-1. \quad (60)$$

We verify this using brute force when $n = 4$. Using (59), we can algebraically express q_1, q_2, q_3, q_4 :

$$\begin{aligned}
q_1 &= P_4 \\
q_2 q_1 &= P_3 - P_4 \\
q_3 q_2 q_1 &= \frac{P_2 - 2P_3 + P_4}{2} \\
q_4 q_3 q_2 q_1 &= \frac{P_1 - 3P_2 + 3P_3 - P_4}{6}
\end{aligned}$$

which implies:

$$\begin{aligned}
\frac{1}{q_1} &= \frac{1}{P_4} \\
\frac{1}{q_2} &= \frac{P_4}{P_3 - P_4} \\
\frac{1}{q_3} &= \frac{2(P_3 - P_4)}{P_2 - 2P_3 + P_4} \\
\frac{1}{q_4} &= \frac{3(P_2 - 2P_3 + P_4)}{P_1 - 3P_2 + 3P_3 - P_4}
\end{aligned}$$

The three inequalities to prove in (60) for $n = 4$ are:

$$\begin{aligned}
\frac{1}{P_4} &\geq \frac{P_4}{P_3 - P_4} + 1 \\
\frac{P_4}{P_3 - P_4} &\geq \frac{2(P_3 - P_4)}{P_2 - 2P_3 + P_4} + 1 \\
\frac{2(P_3 - P_4)}{P_2 - 2P_3 + P_4} &\geq \frac{3(P_2 - 2P_3 + P_4)}{P_1 - 3P_2 + 3P_3 - P_4} + 1
\end{aligned}$$

which are respectively equivalently to:

$$\frac{1}{P_4} \geq \frac{1}{P_3} + 1 \quad (61)$$

$$\frac{2}{P_3} \geq \frac{1}{P_2} + \frac{1}{P_4} \quad (62)$$

$$\frac{4}{P_1 P_3} + \frac{4}{P_2 P_4} \geq \frac{3}{P_1 P_4} + \frac{3}{P_2 P_3} + \frac{1}{P_1 P_2} + \frac{1}{P_3 P_4} \quad (63)$$

Substituting in the Nested Logit choice probabilities from (57), the first inequality (61) is equivalent to $4(1 + (4w)^{-\gamma}) \geq 3(1 + (3w)^{-\gamma}) + 1$ or $4(4w)^{-\gamma} \geq 3(3w)^{-\gamma}$ which is clearly true for all $w > 0$ and $\gamma \in (0, 1]$. Meanwhile, the second inequality (62) is equivalent to $6(1 + (3w)^{-\gamma}) \geq 4(1 + (4w)^{-\gamma}) + 2(1 + (2w)^{-\gamma})$ or $6(3w)^{-\gamma} \geq 4(4w)^{-\gamma} + 2(2w)^{-\gamma}$. Since $w > 0$, we can cancel out $w^{-\gamma}$ from both sides and the inequality is equivalent to $6(3)^{-\gamma} \geq 4(4)^{-\gamma} + 2(2)^{-\gamma}$, which can be verified to be true over $\gamma \in (0, 1]$. Finally, the third inequality (63) is equivalent to

$$\begin{aligned} & 12\left(1 + \frac{1}{w^\gamma}\right)\left(1 + \frac{1}{(3w)^\gamma}\right) + 32\left(1 + \frac{1}{(2w)^\gamma}\right)\left(1 + \frac{1}{(4w)^\gamma}\right) \\ & \geq 12\left(1 + \frac{1}{w^\gamma}\right)\left(1 + \frac{1}{(4w)^\gamma}\right) + 18\left(1 + \frac{1}{(2w)^\gamma}\right)\left(1 + \frac{1}{(3w)^\gamma}\right) + 2\left(1 + \frac{1}{w^\gamma}\right)\left(1 + \frac{1}{(2w)^\gamma}\right) + 12\left(1 + \frac{1}{(3w)^\gamma}\right)\left(1 + \frac{1}{(4w)^\gamma}\right). \end{aligned}$$

The constant terms on both sides are equal to 44. The coefficient in front of $\frac{1}{w^\gamma}$ on the LHS is $12 + \frac{32}{2^\gamma} + \frac{12}{3^\gamma} + \frac{32}{4^\gamma}$, while the same coefficient for the RHS is $14 + \frac{20}{2^\gamma} + \frac{30}{3^\gamma} + \frac{24}{4^\gamma}$. The difference $12 + \frac{32}{2^\gamma} + \frac{12}{3^\gamma} + \frac{32}{4^\gamma} - (14 + \frac{20}{2^\gamma} + \frac{30}{3^\gamma} + \frac{24}{4^\gamma})$ can be verified to be non-negative over $\gamma \in (0, 1]$. Similarly, the coefficient in front of $\frac{1}{w^{2\gamma}}$ on the LHS is $\frac{12}{3^\gamma} + \frac{32}{2^\gamma 4^\gamma}$, while the same coefficient for the RHS is $\frac{12}{4^\gamma} + \frac{18}{2^\gamma 3^\gamma} + \frac{2}{2^\gamma} + \frac{12}{3^\gamma 4^\gamma}$. The difference $\frac{12}{3^\gamma} + \frac{32}{2^\gamma 4^\gamma} - (\frac{12}{4^\gamma} + \frac{18}{2^\gamma 3^\gamma} + \frac{2}{2^\gamma} + \frac{12}{3^\gamma 4^\gamma})$ can be verified to be non-negative over $\gamma \in (0, 1]$. All in all, since the LHS has a higher coefficient for all terms $1, \frac{1}{w^\gamma}, \frac{1}{w^{2\gamma}}$, which are non-negative since $w > 0$, this completes the proof that the desired inequalities (60) indeed hold in the special case of $n = 4$.

Completing the verification of history-monotone futures. Denote the set of items using $N = \{A, B, C, D\}$. To verify history-monotone futures for list distributions on 4 items, we consider two possibilities: $|S| = 1$ and $|S| = 2$. (If $|S| \geq 3$, then there cannot be two distinct prefixes in $N \setminus S$ with the same endpoint, and hence the condition holds vacuously.) If $|S| = 1$, then WOLOG let $S = \{A\}$, $\rho_{\text{end}} = B$. The possible prefixes in $N \setminus \{A\}$ with $\rho_{\text{end}} = B$ are $(B), (CB), (DB), (CDB), (DCB)$. We will show for $j = A$ that

$$\Pr[j \succeq S_0|(B)] \leq \Pr[j \succeq S_0|(CB)] = \Pr[j \succeq S_0|(DB)] \leq \Pr[j \succeq S_0|(CDB)] = \Pr[j \succeq S_0|(DCB)]. \quad (64)$$

On the other hand, if $|S| = 2$, then WOLOG let $S = \{A, B\}$, $\rho_{\text{end}} = C$. The possible prefixes in $N \setminus S$ with $\rho_{\text{end}} = C$ are (C) and (DC) . We will show for $j = A$ (the $j = B$ case holds symmetrically) that

$$\Pr[j \succeq S_0|(C)] \leq \Pr[j \succeq S_0|(DC)]. \quad (65)$$

To show (64), substituting $j = A$ and $S = \{A\}$, note that $\Pr[A \succeq \{A\}_0 | (CDB)] = \Pr[A \succeq \{A\}_0 | (DCB)] = q_4$. Meanwhile, $\Pr[A \succeq \{A\}_0 | (CB)] = q_3 + q_3q_4$, because starting from prefix (CB) , either A can be visited right away, or it can be visited after D . By symmetry, we also have $\Pr[A \succeq \{A\}_0 | (DB)] = q_3 + q_3q_4$. Note that

$$q_3 + q_3q_4 = (1 - q_3) \frac{1}{1/q_3 - 1} + q_3 \frac{1}{1/q_4}$$

which is indeed at most q_4 since it is a convex combination of q_4 with a probability that is smaller, because $1/q_3 - 1 \geq 1/q_4$ according to (60). Similarly, $\Pr[A \succeq \{A\}_0 | (B)] = q_2 + 2q_2(q_3 + q_3q_4)$, because starting from prefix (B) , either A can be visited right away, or C or D could be visited next both of which imply the probability of event $A \succeq \{A\}_0$ is $q_3 + q_3q_4$. Note that

$$q_2 + 2q_2(q_3 + q_3q_4) = (1 - 2q_2) \frac{1}{1/q_2 - 2} + 2q_2(q_3 + q_3q_4)$$

which is indeed at most $q_3 + q_3q_4$ since it is a convex combination of $q_3 + q_3q_4$ with a probability that is smaller, because $1/q_2 - 2 \geq 1/q_3 - 1$ according to (60). This establishes all of the desired relationships in (64).

To show (65), substituting $j = A$ and $S = \{A, B\}$, note that $\Pr[A \succeq \{A, B\}_0 | (DC)] = q_3$ while $\Pr[A \succeq \{A, B\}_0 | (C)] = q_2 + 2q_2q_3 = (1 - q_2) \frac{1}{1/q_2 - 1} + 2q_2q_3$. Again we use the fact that $1/q_2 - 1 \geq 1/q_3$ according to (60), which shows $\Pr[A \succeq \{A, B\}_0 | (C)] \leq \Pr[A \succeq \{A, B\}_0 | (DC)]$. This completes the proof that the list distribution defined has history-monotone futures. \square

Appendix G: Proofs from Section 6

Proof of Proposition 10. Fix any monotone submodular function $f : 2^N \rightarrow [0, 1]$, and define x according to the statement of the proposition. We must show that constraints (3)–(5) for IC mechanisms are satisfied. The non-negativity constraints (5) follow from f being increasing, and the constraints (4) follow from f having a range of at most 1. To establish the IC constraints (3), take lists ℓ, ℓ' and index $k \leq |\ell|$. By the definition of x , the RHS of (3) can be expressed as

$$(f(\ell_{1:i_1}) - f(\ell_{1:i_1-1})) + (f(\ell_{1:i_2}) - f(\ell_{1:i_2-1})) + \cdots + (f(\ell_{1:i_{k'}}) - f(\ell_{1:i_{k'}-1})) \quad (66)$$

where $i_1 < \dots < i_{k'}$ are the indices i such that $\ell'_i \in \ell_{1:k}$, with $k' \leq k$. Applying the condition of submodularity, (66) can be upper-bounded by

$$\begin{aligned} (f(\ell_{i_1}) - f(\emptyset)) + (f(\{\ell_{i_1}, \ell_{i_2}\}) - f(\ell_{i_1})) + \cdots + (f(\{i_1, \dots, i_{k'}\}) - f(\{i_1, \dots, i_{k'-1}\})) &= f(\{i_1, \dots, i_{k'}\}) - f(\emptyset) \\ &\leq f(\ell_{1:k}) - f(\emptyset) \end{aligned}$$

where $f(\{i_1, \dots, i_{k'}\}) \leq f(\ell_{1:k})$ because the indices $i_1, \dots, i_{k'}$ are contained within $\ell_{1:k}$ and f is increasing. The final expression equals the LHS of (3), completing the proof of the forward direction.

For the converse direction, only the submodularity of f has not already been established in Proposition 2. To establish submodularity, it suffices to show that $f(S) + f(S \cup \{j, j'\}) \leq f(S \cup \{j\}) + f(S \cup \{j'\})$ for all subsets $S \subseteq N$ and distinct items $j, j' \in N \setminus S$. So take any $S \subseteq N$ and distinct items $j, j' \in N \setminus S$. Let ℓ denote the list consisting of the items in S in any fixed order, followed by j . Let ℓ' denote the list consisting of the items in S in the same fixed order, followed by j' , and then j . If ℓ reports truthfully, then their probability of receiving an item in $S \cup \{j\}$ is $f(S \cup \{j\})$, by identity (8) which was established in Proposition 2. On the other hand, if ℓ misreports ℓ' , then their probability of receiving an item in $S \cup \{j\}$ is $f(S) + x_j(\ell') = f(S) + f(S \cup \{j', j\}) - f(S \cup \{j'\})$, again making use of identity (8). Substituting $k = |S| + 1$ into the IC constraints (3), we see that the former probability must be lower-bounded by the latter, i.e. $f(S \cup \{j\}) \geq f(S) + f(S \cup \{j', j\}) - f(S \cup \{j'\})$, which suffices for establishing submodularity. \square

Proof of Theorem 4. It is immediate to see that the revenue of a top-2 lottery is

$$\sum_{j=1}^n M^{-j} \cdot \frac{O(M^{j-1}) + M^j}{2} = \sum_{j=1}^n (1/2 + O(M^{-1}))$$

which is at least $n/2$. On the other hand, consider any assortment S with $|S| = k$, for some $k \in \{1, \dots, n\}$. Conditional on the most expensive item on the buyer's list being j for some j , if $j \in S$, then the assortment will sell item j to the buyer as long as their first choice does not also lie in S , which occurs w.p. $1 - \frac{|S \cap \{1, \dots, j-1\}|}{j-1}$. Therefore, the revenue of assortment S equals

$$\begin{aligned} \sum_{j=1}^n M^{-j} \left(O(M^{j-1}) + M^j \mathbb{1}(j \in S) \left(1 - \frac{|S \cap \{1, \dots, j-1\}|}{j-1} \right) \right) &= O(1/M) + \sum_{j \in S} \left(1 - \frac{|S \cap \{1, \dots, j-1\}|}{j-1} \right) \\ &\leq O(1/M) + \sum_{k'=1}^k \left(1 - \frac{k-k'}{n-k'} \right) \\ &= O(1/M) + (n-k) \sum_{k'=1}^k \frac{1}{n-k'}. \end{aligned}$$

We explain the inequality. For all $k' = 1, \dots, k$, if j is the k' 'th-most-expensive item in S , then there must be exactly $k - k'$ elements in $|S \cap \{1, \dots, j-1\}|$. Meanwhile, the maximum possible index j could be is $n + 1 - k'$, and hence $j - 1 \leq n - k'$.

It remains to upper-bound the quantity $(n-k) \sum_{k'=1}^k 1/(n-k')$ over all $k = 1, \dots, n$. If $k < n-1$, then we upper-bound the sum by $\log \frac{n-1}{n-k-1}$, in which case the assortment revenue is at most

$$\begin{aligned} O(1/M) + (n-k) \log \frac{n-1}{n-k-1} &= O(1/M) + \left(n \left(1 - \frac{k}{n-1} \right) + \frac{k}{n-1} \right) \log \frac{n-1}{n-k-1} \\ &= O(1/M) + n \left(1 - \frac{k}{n-1} \right) \log \frac{1}{1 - \frac{k}{n-1}} + O(\log n) \\ &\leq n/e + O(\log n) \end{aligned}$$

where the inequality holds because the maximum of function $(1-a)\log\frac{1}{1-a}$ over $a \in [0, 1]$ is $1/e$, occurring at $a = 1 - 1/e$. On the other hand, if $k \geq n - 1$, then it is easy to see that the assortment revenue cannot be better than $O(\log n)$. Therefore, assortments can obtain no more than $\frac{n/e + O(\log n)}{n/2} \approx 2/e$ times the revenue of the top-2 lottery as $n \rightarrow \infty$, completing the proof. \square

Proof of Theorem 5. Given the mechanism x , define function f as in Proposition 2. Following the same reasoning as in the proof of Theorem 3, $f(\ell_{1:k}) - f(\ell_{1:k-1})$ is the probability of the mechanism selling item ℓ_k to type ℓ . Therefore, it suffices to show that the random assortment sells each type ℓ each item ℓ_k with probability at least $\frac{2}{eL}(f(\ell_{1:k}) - f(\ell_{1:k-1}))$, for $\ell \in \mathcal{L}$ and $k \leq |\ell|$.

The probability that the random assortment sells item ℓ_k to type ℓ is $\varphi(f_{\ell_k}) \prod_{k'=1}^{k-1} (1 - \varphi(f_{\ell_{k'}}))$. If $p(\ell) > 0$ and $f(\ell_{1:k}) - f(\ell_{1:k-1}) > 0$, then we can establish the inequalities

$$\begin{aligned} \frac{\varphi(f_{\ell_k}) \prod_{k'=1}^{k-1} (1 - \varphi(f_{\ell_{k'}}))}{f(\ell_{1:k}) - f(\ell_{1:k-1})} &\geq \frac{\varphi(f(\ell_{1:k}) - f(\ell_{1:k-1})) \prod_{k'=1}^{k-1} (1 - \varphi(f(\ell_{1:k-1})))}{f(\ell_{1:k}) - f(\ell_{1:k-1})} \\ &\geq \frac{\varphi(1 - f(\ell_{1:k-1}))}{1 - f(\ell_{1:k-1})} (1 - \varphi(f(\ell_{1:k-1})))^{L-1}. \end{aligned}$$

Indeed, the first inequality holds because: function φ is non-decreasing; $f_{\ell_k} = f(\ell_k) \geq f(\ell_{1:k}) - f(\ell_{1:k-1})$ since f is submodular (by Proposition 10); and $f(\ell_{k'}) \leq f(\ell_{1:k-1})$ for all $k' = 1, \dots, k-1$ since f is increasing. Subsequently, the second inequality holds because: the function $\varphi(a)/a$ is non-increasing in a ; $f(\ell_{1:k}) \leq 1$; and $k \leq |\ell| \leq L$ (since $p(\ell) > 0$).

Now, let $a = f(\ell_{1:k-1})$ and we claim that the expression $\frac{\varphi(1-a)}{1-a} (1 - \varphi(a))^{L-1}$ is at least $\frac{2}{L}(1 - \frac{1}{L})^{L-1}$ for all $a \in [0, 1]$. To verify this, one must check two cases: $a \leq 1/2$ and $a > 1/2$. In the first,

$$\frac{\varphi(1-a)}{1-a} (1 - \varphi(a))^{L-1} = \frac{1}{L \cdot (1-a)} \left(1 - \frac{2a}{L}\right)^{L-1}$$

whose RHS can be seen to be a decreasing function over $a \in [0, 1/2]$ for all integers $L \geq 2$. Therefore, the expression is lower-bounded by its value at $a = 1/2$, which is $\frac{2}{L}(1 - \frac{1}{L})^{L-1}$. In the second case, $\frac{\varphi(1-a)}{1-a} (1 - \varphi(a))^{L-1} = \frac{2(1-a)/L}{1-a} (1 - \frac{1}{L})^{L-1} = \frac{2}{L}(1 - \frac{1}{L})^{L-1}$ for all $a \in (1/2, 1)$.

We have shown that the randomized assortment sells each realizable type ℓ their k 'th most-preferred item ℓ_k with probability at least $\frac{2}{L}(1 - \frac{1}{L})^{L-1}(f(\ell_{1:k}) - f(\ell_{1:k-1}))$. Since $(1 - \frac{1}{L})^{L-1}$ is lower-bounded by $1/e$ for all values of L , this completes the proof of the theorem. \square