

A General Family of q -Hypergeometric Polynomials and Associated Generating Functions

H. M. Srivastava^{1,2,3,4,*} and Sama Arjika⁵

¹Department of Mathematics and Statistics, University of Victoria,
Victoria, British Columbia V8W 3R4, Canada

²Department of Medical Research, China Medical University Hospital,
China Medical University, Taichung 40402, Taiwan, Republic of China

³Department of Mathematics and Informatics, Azerbaijan University,
71 Jeyhun Hajibeyli Street, AZ1007 Baku, Azerbaijan

⁴Section of Mathematics, International Telematic University Uninettuno,
I-00186 Rome, Italy

E-Mail: harimsri@math.uvic.ca

***Corresponding Author**

⁵Department of Mathematics and Informatics, University of Agadez,
Post Office Box 199, Agadez, Niger

E-Mail: rjksama2008@gmail.com

Abstract

In this paper, we introduce a general family of q -hypergeometric polynomials and investigate several q -series identities such as an extended generating function and a Srivastava-Agarwal type bilinear generating function for this family of q -hypergeometric polynomials. We give a transformational identity involving generating functions for the generalized q -hypergeometric polynomials which we have introduced here. We also point out relevant connections of the various q -results, which we investigate here, with those in several related earlier works on this subject. We conclude this paper by remarking that it will be a rather trivial and inconsequential exercise to give the so-called (p, q) -variations of the q -results, which we have investigated here, because the additional parameter p is obviously redundant.

2020 Mathematics Subject Classification. Primary 05A30, 33D15, 33D45; Secondary 05A40, 11B65.

Key Words and Phrases. Basic (or q -) hypergeometric series; Homogeneous q -difference operator; q -Binomial theorem; Cauchy polynomials; Al-Salam-Carlitz q -polynomials; Rogers type formulas; Srivastava-Agarwal type generating functions.

1. Introduction, Definitions and Preliminaries

In this paper, we adopt the commonly-used conventions and notations for the basic (or q -) series and the basic (or q -) polynomials. For the convenience of the reader, we first provide a summary of the mathematical notations, basic properties and definitions to be used in this paper. We refer to the general references (see, for example, [15], [18] and [24]) for the definitions and notations.

Throughout this paper, we assume that $|q| < 1$. For complex numbers a , the q -shifted factorials are defined by

$$(a; q)_0 := 1, \quad (a; q)_n = \prod_{k=0}^{n-1} (1 - aq^k), \quad (a; q)_\infty := \prod_{k=0}^{\infty} (1 - aq^k) \quad (1.1)$$

and

$$(a_1, a_2, \dots, a_r; q)_m = (a_1; q)_m (a_2; q)_m \cdots (a_r; q)_m \quad (m \in \mathbb{N}_0 := \{0, 1, 2, \dots\} = \mathbb{N} \cup \{0\}).$$

We will frequently use the following q -identity [1]:

$$(aq^{-n}; q)_n = (q/a; q)_n (-a)^n q^{-n - \binom{n}{2}}. \quad (1.2)$$

Here, in this paper, we are mainly concerned with the Cauchy polynomials defined by (see [9]):

$$P_n(x, y) := (x - y)(x - qy) \cdots (x - q^{n-1}y) = (y/x; q)_n x^n \quad (1.3)$$

together with their Srivastava-Agarwal type generating function as given below:

$$\sum_{n=0}^{\infty} P_n(x, y) \frac{(\lambda; q)_n t^n}{(q; q)_n} = {}_2\Phi_1 \left[\begin{matrix} \lambda, y/x; \\ q; xt \end{matrix} \right]. \quad (1.4)$$

For $\lambda = 0$, we get the following generating function (see [9]):

$$\sum_{n=0}^{\infty} P_n(x, y) \frac{t^n}{(q; q)_n} = \frac{(yt; q)_\infty}{(xt; q)_\infty}. \quad (1.5)$$

The generating function (1.5) is also the homogeneous version of the Cauchy identity or the q -binomial theorem is given by (see [11])

$$\sum_{k=0}^{\infty} \frac{(a; q)_k}{(q; q)_k} z^k = {}_1\Phi_0 \left[\begin{matrix} a; \\ q; z \end{matrix} \right] = \frac{(az; q)_\infty}{(z; q)_\infty} \quad (|z| < 1). \quad (1.6)$$

Putting $a = 0$, the relation (1.6) becomes Euler's identity (see [11]):

$$\sum_{k=0}^{\infty} \frac{z^k}{(q; q)_k} = \frac{1}{(z; q)_\infty} \quad (|z| < 1) \quad (1.7)$$

and its inverse relation given below (see [11]):

$$\sum_{k=0}^{\infty} \frac{(-1)^k q^{\binom{k}{2}} z^k}{(q; q)_k} = (z; q)_\infty. \quad (1.8)$$

Saad and Sukhi [16] introduced the dual homogeneous q -difference operator Θ_{xy} as follows (see also [10, 20, 3]):

$$\Theta_{xy} \{f(x, y)\} := \frac{f(q^{-1}x, y) - f(x, qy)}{q^{-1}x - y}, \quad (1.9)$$

which, when it acts upon functions of suitably-restricted variables x and y , yields

$$\Theta_{xy}^k \{P_n(y, x)\} = (-1)^k \frac{(q; q)_n}{(q; q)_{n-k}} P_{n-k}(y, x) \quad \text{and} \quad \Theta_{xy}^k \left\{ \frac{(xt; q)_\infty}{(yt; q)_\infty} \right\} = (-t)^k \frac{(xt; q)_\infty}{(yt; q)_\infty}. \quad (1.10)$$

The Hahn polynomials (see [12], [13] and [14]) or, equivalently, the Al-Salam-Carlitz q -polynomials (see [2]) are defined as follows:

$$\phi_n^{(a)}(x|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (a; q)_k x^k \quad \text{and} \quad \psi_n^{(a)}(x|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{k(k-n)} (aq^{1-k}; q)_k x^k. \quad (1.11)$$

These polynomials are usually called the “*Al-Salam-Carlitz polynomials*” in several recent publications. Moreover, because of their considerable role in the theories of q -series and q -orthogonal polynomials, many authors investigated various extensions of the Al-Salam-Carlitz polynomials (see, for example, [6], [8] and [22]).

Recently, Cao [6, Eq. (4.7)] introduced the following two families of generalized Al-Salam-Carlitz polynomials:

$$\phi_n^{(a,b,c)}(x,y|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{(a,b;q)_k}{(c;q)_k} x^k y^{n-k} \quad (1.12)$$

and

$$\psi_n^{(a,b,c)}(x,y|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{(-1)^k q^{\binom{k+1}{2}-nk} (a,b;q)_k}{(c;q)_k} x^k y^{n-k}, \quad (1.13)$$

together with their generating functions given by (see [6, Eqs.(4.10) and (4.11)])

$$\sum_{n=0}^{\infty} \phi_n^{(a,b,c)}(x,y|q) \frac{t^n}{(q;q)_n} = \frac{1}{(xt;q)_{\infty}} {}_2\Phi_1 \left[\begin{matrix} a,b; \\ q; yt \end{matrix} \right] \quad (\max\{|yt|, |xt|\} < 1) \quad (1.14)$$

and

$$\sum_{n=0}^{\infty} \psi_n^{(a,b,c)}(x,y|q) \frac{(-1)^n q^{\binom{n}{2}} t^n}{(q;q)_n} = (xt;q)_{\infty} {}_2\Phi_1 \left[\begin{matrix} a,b; \\ q; yt \end{matrix} \right] \quad (|xt| < 1). \quad (1.15)$$

Remark 1. For $(b,c,y) = (0,0,1)$ and For $(a,b,c,y) = (1/a,0,0,1)$, (1.12) and (1.13) reduce to $\phi_n^{(a)}(x)$ and $\psi_n^{(a)}(x)$ defined in (1.11).

Motivated by the above-cited work [6], Cao *et al.* [8] introduced the extensions $\phi_n^{(a,b,c)}(x,y|q)$ and $\psi_n^{(a,b,c)}(x,y|q)$ of the Al-Salam-Carlitz polynomials, which are defined by

$$\phi_n^{(a,b,c)}(x,y|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{(a,b,c;q)_k}{(d,e;q)_k} x^{n-k} y^k \quad (1.16)$$

and

$$\psi_n^{(a,b,c)}(x,y|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{(-1)^k q^{k(k-n)} (a,b,c;q)_k}{(d,e;q)_k} x^{n-k} y^k, \quad (1.17)$$

respectively. More recently, Srivastava and Arjika [22] introduced two families $\phi_n^{(\mathbf{a},\mathbf{b})}(x,y|q)$ and $\psi_n^{(\mathbf{a},\mathbf{b})}(x,y|q)$ of the generalized Al-Salam-Carlitz q -polynomials, which are defined by

$$\phi_n^{(\mathbf{a},\mathbf{b})}(x,y|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{(a_1, a_2, \dots, a_{r+1}; q)_k}{(b_1, b_2, \dots, b_r; q)_k} x^k y^{n-k} \quad (1.18)$$

and

$$\psi_n^{(\mathbf{a},\mathbf{b})}(x,y|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{(a_1, a_2, \dots, a_{r+1}; q)_k}{(b_1, b_2, \dots, b_r; q)_k} q^{\binom{k+1}{2} - nk} x^k y^{n-k}, \quad (1.19)$$

respectively, where

$$\mathbf{a} := (a_1, a_2, \dots, a_{r+1}) \quad \text{and} \quad \mathbf{b} = (b_1, b_2, \dots, b_r).$$

On the other hand, Cao [6] introduced and studied the following family of q -polynomials:

$$V_n^{(\mathbf{a},\mathbf{c})}(x,y,z|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{(a_1, a_2, \dots, a_r; q)_k}{(c_1, c_2, \dots, c_u; q)_k} P_{n-k}(x,y) z^k, \quad (1.20)$$

where

$$\mathbf{a} := (a_1, a_2, \dots, a_r) \quad \text{and} \quad \mathbf{c} = (c_1, c_2, \dots, c_s),$$

together with their generating function given by

$$\sum_{n=0}^{\infty} V_n^{(\mathbf{a},\mathbf{c})}(x,y,z|q) \frac{t^n}{(q; q)_n} = \frac{(yt; q)_{\infty}}{(xt; q)_{\infty}} {}_r\Phi_u \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ q; zt \end{matrix} \right]. \quad (1.21)$$

Our present investigation is motivated essentially by the earlier works by Srivastava and Arjika [22] and Cao [6]. Our aim here is to introduce and study some further extensions of the above-mentioned q -polynomials.

Definition 1. In terms of q -binomial coefficient, a family of generalized q -hypergeometric polynomials $\Psi_n^{(\mathbf{a},\mathbf{b})}(x,y,z|q)$ are defined by

$$\Psi_n^{(\mathbf{a},\mathbf{b})}(x,y,z|q) := (-1)^n q^{-\binom{n}{2}} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \left[(-1)^k q^{\binom{k}{2}} \right]^{1+s-r} W_k(\mathbf{a}, \mathbf{b}) P_{n-k}(y,x) z^k, \quad (1.22)$$

where, for convenience,

$$\mathbf{a} = (a_1, a_2, \dots, a_r), \quad \mathbf{b} = (b_1, b_2, \dots, b_s) \quad \text{and} \quad W_k(\mathbf{a}, \mathbf{b}) := \frac{(a_1, a_2, \dots, a_r; q)_k}{(b_1, b_2, \dots, b_s; q)_k}.$$

The above-defined q -polynomials $\Psi_n^{(\mathbf{a},\mathbf{b})}(x,y,z|q)$ include many one-variable q -hypergeometric ${}_r\Phi_s$ series as special or limit cases. Therefore, we choose just to call them generalized q -hypergeometric polynomials.

Remark 2. The generalized q -hypergeometric polynomials $\Psi_n^{(\mathbf{a},\mathbf{b})}(x,y,z|q)$ defined in (1.22) are generalized and unified form of the Hahn polynomials and the Al-Salam-Carlitz polynomials.

- (1) Upon setting $(r,x,y) = (u+1,y,x)$ and $\mathbf{b} = \mathbf{c}$, the generalized q -hypergeometric polynomials (1.22) would reduce to the generalized Al-Salam-Carlitz q -polynomials $V_n^{(\mathbf{a},\mathbf{c})}(x,y|q)$ in (1.20) (see [7]):

$$\Psi_n^{(\mathbf{a},\mathbf{c})}(y,x,z|q) = (-1)^n q^{-\binom{n}{2}} V_n^{(\mathbf{a},\mathbf{c})}(x,y,z|q). \quad (1.23)$$

- (2) By choosing $r = s + 1$, $x = 0$ and $z = x$, the generalized q -hypergeometric polynomials (1.22) reduce to the generalized Al-Salam-Carlitz q -polynomials $\phi_n^{(\mathbf{a}, \mathbf{b})}(x, y|q)$ (see [22]):

$$\Psi_n^{(\mathbf{a}, \mathbf{b})}(0, y, x|q) = (-1)^n q^{-\binom{n}{2}} \phi_n^{(\mathbf{a}, \mathbf{b})}(x, y|q). \quad (1.24)$$

- (3) Upon setting $r = s + 1$, $y = 0$, $z = -x$ and $x = y$, the q -hypergeometric polynomials (1.22) reduce to the generalized Al-Salam-Carlitz q -polynomials $\psi_n^{(\mathbf{a}, \mathbf{b})}(x, y|q)$ (see [22]):

$$\Psi_n^{(\mathbf{a}, \mathbf{b})}(0, y, -x|q) = \psi_n^{(\mathbf{a}, \mathbf{b})}(x, y|q). \quad (1.25)$$

- (4) For $r = 3$, $s = 2$, $\mathbf{a} = (a, 0, 0)$, $\mathbf{b} = (0, 0)$ and $z = b$, the q -hypergeometric polynomials (1.22) reduce to the generalized Hahn polynomials $h_n(x, y, a, b|q)$ (see [23]):

$$\Psi_n^{(\mathbf{a}, \mathbf{c})}(y, x, z|q) = (-1)^n q^{-\binom{n}{2}} h_n(x, y, a, b|q). \quad (1.26)$$

- (5) For $r = s$ and $\mathbf{a} = \mathbf{b} = \mathbf{0}$, the q -hypergeometric polynomials $\Psi_n^{(\mathbf{a}, \mathbf{b})}(x, y, z|q)$ are the well known trivariate q -polynomials $F_n(x, y, z; q)$ (see [1]):

$$\Psi_n^{(\mathbf{0}, \mathbf{0})}(x, y, z|q) = F_n(x, y, z; q) := (-1)^n q^{-\binom{n}{2}} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (-1)^k q^{\binom{k}{2}} z^k P_{n-k}(y, x). \quad (1.27)$$

- (6) For $r = 3$, $s = 2$, $\mathbf{a} = (a, b, c)$, $\mathbf{b} = (d, e)$ and $(x, y, z) = (0, x, y)$, the generalized q -hypergeometric polynomials $\Psi_n^{(\mathbf{a}, \mathbf{b})}(y, x, z|q)$ reduce to $\phi_n^{(a, b, c; d, e)}(x, y, z|q)$ (see [8]):

$$\Psi_n^{(\mathbf{a}, \mathbf{b})}(y, x, z|q) = (-1)^n q^{-\binom{n}{2}} \phi_n^{(a, b, c; d, e)}(x, y, z|q). \quad (1.28)$$

- (7) For $r = s = 2$, $\mathbf{a} = (a, b, c)$, $\mathbf{b} = (d, e)$ and $(x, y, z) = (0, x, y)$, the q -hypergeometric polynomials (1.22) reduce to the polynomials $\psi_n^{(a, b, c; d, e)}(x, y, z|q)$ (see [8]):

$$\Psi_n^{(\mathbf{a}, \mathbf{b})}(y, x, z|q) = (-1)^n q^{-\binom{n}{2}} \psi_n^{(a, b, c; d, e)}(x, y, z|q). \quad (1.29)$$

- (8) If we let $r = 2$, $s = 1$, $\mathbf{a} = (a, 0)$, $\mathbf{b} = (0)$, $x = 0$ and $z = x$, the polynomials $\Psi_n^{(\mathbf{a}, \mathbf{b})}(x, y, z|q)$ reduce to the second Hahn polynomials $\phi_n^{(a)}(x, y|q)$ (see [5]):

$$\Psi_n^{(\mathbf{0}, \mathbf{0})}(x, ax, y|q) = (-1)^n q^{-\binom{n}{2}} \phi_n^{(a)}(x, y|q). \quad (1.30)$$

- (9) If we let $r = 2$, $s = 1$, $(\mathbf{a}, \mathbf{b}) = (\mathbf{0}, \mathbf{0})$, $y = ax$ and $z = y$, the polynomials $\Psi_n^{(\mathbf{a}, \mathbf{b})}(x, y, z|q)$ reduce to the second Hahn polynomials $\psi_n^{(a)}(x, y|q)$ (see [5]):

$$\Psi_n^{(\mathbf{0}, \mathbf{0})}(x, ax, y|q) = \psi_n^{(a)}(x, y|q). \quad (1.31)$$

- (10) For $r = 2$, $s = 1$, $(\mathbf{a}, \mathbf{b}) = (\mathbf{0}, \mathbf{0})$, $x = y = 0$ and $z = x$, the polynomials $\Psi_n^{(\mathbf{a}, \mathbf{b})}(x, y, z|q)$ reduce to the generalized Hahn polynomials $\phi_n^{(a)}(x|q)$ (see [12], [13] and [14]) or, equivalently, the Al-Salam-Carlitz q -polynomials (see [2]):

$$\Psi_n^{(\mathbf{0}, \mathbf{0})}(0, 0, x|q) = (-1)^n q^{-\binom{n}{2}} \phi_n^{(a)}(x|q). \quad (1.32)$$

- (11) Upon putting $r = s$, $(\mathbf{a}, \mathbf{b}) = (\mathbf{0}, \mathbf{0})$, $y = ax$ and $z = 1$, the generalized q -hypergeometric polynomials $\Psi_n^{(\mathbf{a}, \mathbf{b})}(x, y, z|q)$ reduce to the Hahn polynomials $\psi_n^{(a)}(x|q)$ (see [12], [13] and [14]) or, equivalently, the Al-Salam-Carlitz q -polynomials (see [2]):

$$\Psi_n^{(\mathbf{0}, \mathbf{0})}(x, ax, 1|q) = \psi_n^{(a)}(x|q). \quad (1.33)$$

This paper is organized as follows. In Section 2, we introduce the following homogeneous q -difference operator:

$${}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{array} \middle| q; -z\Theta_{xy} \right]$$

and apply it to investigate several q -series properties. In addition, we derive several extended generating functions for the generalized q -hypergeometric polynomials. In Section 3, we first state and prove the Rogers type formulas. The Srivastava-Agarwal type bilinear generating functions involving the generalized q -hypergeometric polynomials are derived in Section 4. In Section 5, we give a transformational identity involving generating functions for generalized q -hypergeometric polynomials. Finally, in our concluding Section 6, we present several remarks and observations. We also reiterate the well-documented fact that it will be a rather trivial exercise to give the so-called (p, q) -variation of the q -results, which we have investigated here, because the additional parameter p is obviously redundant.

2. Generalized q -Hypergeometric Polynomials

In this section, we begin by introducing a homogeneous q -difference hypergeometric operator as follows.

Definition 2. The homogeneous q -difference hypergeometric operator is defined by

$${}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{array} \middle| q; -z\Theta_{xy} \right] = \sum_{k=0}^{\infty} W_k(\mathbf{a}, \mathbf{b}) \frac{(-z\Theta_{xy})^k}{(q; q)_k} \left[(-1)^k q^{\binom{k}{2}} \right]^{1+s-r}. \quad (2.1)$$

We now derive the q -series identities (2.2), (2.3) and (2.4) below, which will be used later in order to derive the extended generating functions, the Rogers type formulas and the Srivastava-Agarwal type bilinear generating functions involving the generalized q -hypergeometric polynomials.

Lemma 1. Each of the following operational formulas holds true for the homogeneous q -difference operator defined by (2.1):

$${}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{array} \middle| q; -z\Theta_{xy} \right] \left\{ (-1)^n q^{-\binom{n}{2}} P_n(y, x) \right\} = \Psi_n^{(\mathbf{a}, \mathbf{b})}(x, y, z|q), \quad (2.2)$$

$${}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{array} \middle| q; -z\Theta_{xy} \right] \left\{ \frac{(xt; q)_{\infty}}{(yt; q)_{\infty}} \right\} = \frac{(xt; q)_{\infty}}{(yt; q)_{\infty}} {}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{array} \middle| q; zt \right] \quad (2.3)$$

and

$$\begin{aligned}
 {}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{array} \right. & \left. \begin{array}{c} q; -z\Theta_{xy} \\ \left\{ \frac{P_k(y, x)(xt; q)_\infty}{(xt; q)_k (yt; q)_\infty} \right\} \end{array} \right] \\
 &= t^{-k} \frac{(xt; q)_\infty}{(yt; q)_\infty} \sum_{j=0}^k \frac{(q^{-k}, yt; q)_j q^j}{(xt, q; q)_j} {}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{array} \right. \left. \begin{array}{c} q; ztq^j \\ \end{array} \right] \quad (|yt| < 1). \quad (2.4)
 \end{aligned}$$

Proof. Firstly, by applying (2.1) and (1.10), we get (2.2). Secondly, in light of (1.10), we get the desired identity (2.3). Thirdly, by making use of the q -Chu-Vandermonde formula [11, Eq. (II.6)]

$${}_2\Phi_1 \left[\begin{array}{c} q^{-n}, a; \\ c; \end{array} \right. \left. \begin{array}{c} q; q \\ \end{array} \right] = \frac{(c/a; q)_n}{(c; q)_n} a^n, \quad (2.5)$$

we find that

$$\frac{P_k(y, x)(xt; q)_\infty}{(xt; q)_k (yt; q)_\infty} = t^{-k} \frac{(xt; q)_\infty}{(yt; q)_\infty} {}_2\Phi_1 \left[\begin{array}{c} q^{-k}, yt; \\ xt; \end{array} \right. \left. \begin{array}{c} q; q \\ \end{array} \right] = t^{-k} \sum_{j=0}^{\infty} \frac{(q^{-k}; q)_j q^j}{(q; q)_j} \frac{(xtq^j; q)_\infty}{(ytq^j; q)_\infty}. \quad (2.6)$$

Therefore, by using (2.6) and (2.3) successively, we obtain

$$\begin{aligned}
 {}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{array} \right. & \left. \begin{array}{c} q; -z\Theta_{xy} \\ \left\{ \frac{P_k(y, x)(xt; q)_\infty}{(xt; q)_k (yt; q)_\infty} \right\} \end{array} \right] \\
 &= t^{-k} \sum_{j=0}^k \frac{(q^{-k}; q)_j q^j}{(q; q)_j} {}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{array} \right. \left. \begin{array}{c} q; -z\Theta_{xy} \\ \left\{ \frac{(xtq^j; q)_\infty}{(ytq^j; q)_\infty} \right\} \end{array} \right] \\
 &= t^{-k} \sum_{j=0}^k \frac{(q^{-k}; q)_j q^j}{(q; q)_j} \frac{(xtq^j; q)_\infty}{(ytq^j; q)_\infty} {}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{array} \right. \left. \begin{array}{c} q; ztq^j \\ \end{array} \right],
 \end{aligned}$$

which is asserted by Lemma 1. □

We now derive an extended generating function for the generalized q -hypergeometric polynomials $\Psi_n^{(a,b)}(x, y|q)$ by using the operator representation (2.2).

Theorem 1 (Extended generating function). *For $k \in \mathbb{N}$ and $\max\{|yt|, |zt|\} < 1$, it is asserted that*

$$\begin{aligned}
 \sum_{n=0}^{\infty} \Psi_{n+k}^{(a,b)}(x, y, z|q) (-1)^{n+k} q^{\binom{n+k}{2}} \frac{t^n}{(q; q)_n} \\
 &= \frac{(xt; q)_\infty}{t^k (yt; q)_\infty} \sum_{j=0}^k \frac{(q^{-k}, yt; q)_j q^j}{(q, xt; q)_j} {}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{array} \right. \left. \begin{array}{c} q; ztq^j \\ \end{array} \right] \quad (|yt| < 1). \quad (2.7)
 \end{aligned}$$

Proof. In light of (2.2) and (1.5), we have

$$\begin{aligned}
& \sum_{n=0}^{\infty} \Psi_n^{(\mathbf{a}, \mathbf{b})}(x, y, z|q) (-1)^{n+k} q^{\binom{n+k}{2}} \frac{t^n}{(q; q)_n} \\
&= \sum_{n=0}^{\infty} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; -z\Theta_{xy} \right] \left\{ (-1)^{n+k} q^{\binom{n+k}{2}} P_{n+k}(y, x) \right\} \frac{(-1)^{n+k} q^{\binom{n+k}{2}} t^n}{(q; q)_n} \\
&= {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; -z\Theta_{xy} \right] \left\{ P_k(y, x) \sum_{n=0}^{\infty} P_n(y, xq^k) \frac{t^n}{(q; q)_n} \right\} \\
&= {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; -z\Theta_{xy} \right] \left\{ \frac{P_k(y, x) (xt; q)_{\infty}}{(xt; q)_k (yt; q)_{\infty}} \right\}.
\end{aligned}$$

The proof of Theorem 1 is completed by using (2.4). \square

Remark 3. Setting $k = 0$ in Theorem 1, we get the following generating function for the generalized q -hypergeometric polynomials:

$$\sum_{n=0}^{\infty} \Psi_n^{(\mathbf{a}, \mathbf{b})}(x, y, z|q) (-1)^n q^{\binom{n}{2}} \frac{t^n}{(q; q)_n} = \frac{(xt; q)_{\infty}}{(yt; q)_{\infty}} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; zt \right]. \quad (2.8)$$

3. The Rogers Formula

In this section, we use the assertion (2.4) of Lemma 1 in order to derive several q -identities such as the Rogers type formula for the generalized q -hypergeometric polynomials $\Psi_n^{(\mathbf{a}, \mathbf{b})}(x, y|q)$.

Theorem 2 (The Rogers formula for $\Psi_n^{(\mathbf{a}, \mathbf{b})}(x, y, z|q)$). *For $\max\{|\frac{t}{\omega}|, |y\omega|\} < 1$, the following Rogers type formula holds:*

$$\begin{aligned}
& \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \Psi_{n+k}^{(\mathbf{a}, \mathbf{b})}(x, y, z|q) (-1)^{n+k} q^{\binom{n+k}{2}} \frac{t^n}{(q; q)_n} \frac{\omega^k}{(q; q)_k} \\
&= \frac{(x\omega; q)_{\infty}}{(t/\omega, y\omega; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(y\omega; q)_k q^k}{(q\omega/t, x\omega, q; q)_k} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; z\omega q^k \right]. \quad (3.1)
\end{aligned}$$

Proof. In light of (2.2), we have

$$\begin{aligned}
 & \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \Psi_{n+k}^{(\mathbf{a}, \mathbf{b})}(x, y, z|q) (-1)^{n+k} q^{\binom{n+k}{2}} \frac{t^n}{(q; q)_n} \frac{\omega^k}{(q; q)_k} \\
 &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; -z\Theta_{xy} \right] \{P_{n+k}(y, x)\} \frac{t^n}{(q; q)_n} \frac{\omega^k}{(q; q)_k} \\
 &= {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; -z\Theta_{xy} \right] \left\{ \sum_{n=0}^{\infty} P_n(y, x) \frac{t^n}{(q; q)_n} \sum_{k=0}^{\infty} P_k(y, q^n x) \frac{\omega^k}{(q; q)_k} \right\} \\
 &= {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; -z\Theta_{xy} \right] \left\{ \sum_{n=0}^{\infty} P_n(y, x) \frac{t^n}{(q; q)_n} \frac{(x\omega q^n; q)_{\infty}}{(y\omega; q)_{\infty}} \right\} \\
 &= \sum_{n=0}^{\infty} \frac{t^n}{(q; q)_n} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; -z\Theta_{xy} \right] \left\{ \frac{P_n(y, x) (x\omega; q)_{\infty}}{(x\omega; q)_n (y\omega; q)_{\infty}} \right\}. \tag{3.2}
 \end{aligned}$$

Now, by using (2.4) as well as (1.2), this last relation takes the following form:

$$\begin{aligned}
 & \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \Psi_{n+k}^{(\mathbf{a}, \mathbf{b})}(x, y, z|q) (-1)^{n+k} q^{\binom{n+k}{2}} \frac{t^n}{(q; q)_n} \frac{\omega^k}{(q; q)_k} \\
 &= \frac{(x\omega; q)_{\infty}}{(y\omega; q)_{\infty}} \sum_{n=0}^{\infty} \frac{(t/\omega)^n}{(q; q)_n} \sum_{k=0}^n \frac{(q^{-n}, y\omega; q)_k q^k}{(x\omega, q; q)_k} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; z\omega q^k \right] \\
 &= \frac{(x\omega; q)_{\infty}}{(y\omega; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(y\omega; q)_k q^k}{(x\omega, q; q)_k} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; z\omega q^k \right] \sum_{n=k}^{\infty} \frac{(t/\omega)^n (-1)^k q^{\binom{k}{2} - nk}}{(q; q)_{n-k}} \\
 &= \frac{(x\omega; q)_{\infty}}{(y\omega; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(y\omega; q)_k (-t/\omega)^k q^{-\binom{k}{2}}}{(x\omega, q; q)_k} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; z\omega q^k \right] \sum_{n=0}^{\infty} \frac{(t/\omega)^n q^{-nk}}{(q; q)_n} \\
 &= \frac{(x\omega; q)_{\infty}}{(y\omega; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(y\omega; q)_k (-t/\omega)^k q^{-\binom{k}{2}}}{(x\omega, q; q)_k} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; z\omega q^k \right] \frac{1}{(tq^{-k}/\omega; q)_{\infty}} \\
 &= \frac{(x\omega; q)_{\infty}}{(t/\omega, y\omega; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(y\omega; q)_k (-t/\omega)^k q^{-\binom{k}{2}}}{(x\omega, tq^{-k}/\omega, q; q)_k} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; z\omega q^k \right] \\
 &= \frac{(x\omega; q)_{\infty}}{(t/\omega, y\omega; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(y\omega; q)_k q^k}{(q\omega/t, x\omega, q; q)_k} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; z\omega q^k \right], \tag{3.3}
 \end{aligned}$$

which evidently completes the proof of Theorem 2. \square

4. The Srivastava-Agarwal Type Bilinear Generating Functions for the Generalized q -Hypergeometric Polynomials $\Psi_n^{(a,b)}(x, y, z|q)$

In this section, by applying the following homogeneous q -difference hypergeometric operator [17]:

$${}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ q, -z\Theta_{xy} \\ b_1, b_2, \dots, b_s; \end{array} \right],$$

we derive the Srivastava-Agarwal type generating functions for the generalized q -hypergeometric polynomials $\Psi_n^{(a,b)}(x, y, z|q)$ defined by (1.22). We also deduce a bilinear generating function for the Al-Salam-Carlitz polynomials $\psi_n^{(\alpha)}(x|q)$ as an application of the Srivastava-Agarwal type generating functions.

Lemma 2. (see [21, Eq. (3.20)] and [4, Eq. (5.4)]) *Each of the following generating relations holds true:*

$$\sum_{n=0}^{\infty} \phi_n^{(\alpha)}(x|q)(\lambda; q)_n \frac{t^n}{(q; q)_n} = \frac{(\lambda t; q)_{\infty}}{(t; q)_{\infty}} {}_2\Phi_1 \left[\begin{array}{c} \lambda, \alpha; \\ q; xt \\ \lambda t; \end{array} \right] \quad (\max\{|t|, |xt|\} < 1) \quad (4.1)$$

and

$$\sum_{n=0}^{\infty} \psi_n^{(\alpha)}(x|q)(1/\lambda; q)_n \frac{(\lambda t q)^n}{(q; q)_n} = \frac{(xt q; q)_{\infty}}{(\lambda x t q; q)_{\infty}} {}_2\Phi_1 \left[\begin{array}{c} 1/\lambda, 1/(\alpha x); \\ q; \alpha q \\ 1/(\lambda x t); \end{array} \right] \quad (\max\{|\lambda x t q|, |\alpha q|\} < 1). \quad (4.2)$$

For more information about the Srivastava-Agarwal type generating functions for the Al-Salam-Carlitz polynomials, one may refer to [21] and [5].

We now state and prove the Srivastava-Agarwal type bilinear generating functions asserted by Theorem 3 below.

Theorem 3. *Suppose that $\max\{|\alpha q|, |vxtq|, |xztq|\} < 1$. Then*

$$\begin{aligned} & \sum_{n=0}^{\infty} \psi_n^{(\alpha)}(x|q) \Psi_n^{(a,b)}(u, v, z|q) \frac{(-1)^n q^{\binom{n+1}{2}} t^n}{(q; q)_n} \\ &= \frac{(q/x, uxtq; q)_{\infty}}{(\alpha q, vxtq; q)_{\infty}} \sum_{n=0}^{\infty} \frac{(-1)^n q^{\binom{n}{2}} (1/(\alpha x), 1/(uxt); q)_n}{(q/x, 1/(vxt), q; q)_n} \left(\frac{\alpha u q}{v}\right)^n \\ & \quad \cdot {}_r\Phi_s \left[\begin{array}{c} a_1, a_2, \dots, a_r; \\ q, xztq^{1-n} \\ b_1, b_2, \dots, b_s; \end{array} \right]. \end{aligned} \quad (4.3)$$

Proof. In order to prove Theorem 3, we need the q -Chu-Vandermonde summation theorem given by (see [11, Eq. (II.7)])

$${}_2\Phi_1 \left[\begin{array}{c} q^{-n}, a; \\ q; \frac{cq^n}{a} \\ c; \end{array} \right] = \frac{(c/a; q)_n}{(c; q)_n} \quad (n \in \mathbb{N}_0). \quad (4.4)$$

Upon letting $(\lambda, t) = (v/u, tu)$ in the equation (4.2), we obtain

$$\begin{aligned}
 & \sum_{n=0}^{\infty} \psi_n^{(\alpha)}(x|q) p_n(v, u) \frac{(qt)^n}{(q; q)_n} \\
 &= \frac{(uxtq; q)_{\infty}}{(vxtq; q)_{\infty}} {}_2\Phi_1 \left[\begin{matrix} 1/(\alpha x), u/v; \\ q; \alpha q \end{matrix} \right] \\
 &= \frac{(uxtq; q)_{\infty}}{(vxtq; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(1/(\alpha x); q)_k (\alpha q)^k}{(q; q)_k} \frac{(u/v; q)_k}{(1/(vxt); q)_k} \text{ by (4.4)} \\
 &= \frac{(uxtq; q)_{\infty}}{(vxtq; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(1/(\alpha x); q)_k (\alpha q)^k}{(q; q)_k} {}_2\Phi_1 \left[\begin{matrix} q^{-k}, 1/(uxt); \\ q; \frac{uq^k}{v} \end{matrix} \right] \\
 &= \sum_{k=0}^{\infty} \frac{(1/(\alpha x); q)_k (\alpha q)^k}{(q; q)_k} \sum_{n=0}^{\infty} \frac{(q^{-k}; q)_n q^{nk}}{(q; q)_n} \frac{(1/(uxt); q)_n (uxtq; q)_{\infty}}{(1/(vxt); q)_n (vxtq; q)_{\infty}} \left(\frac{u}{v}\right)^n \\
 &= \sum_{k=0}^{\infty} \frac{(1/(\alpha x); q)_k (\alpha q)^k}{(q; q)_k} \sum_{n=0}^{\infty} \frac{(q^{-k}; q)_n q^{nk}}{(q; q)_n} \frac{(uxtq^{1-n}; q)_{\infty}}{(vxtq^{1-n}; q)_{\infty}}. \tag{4.5}
 \end{aligned}$$

By appealing appropriately to (4.5), the left-hand side of (4.3) becomes

$$\begin{aligned}
 & \sum_{n=0}^{\infty} \psi_n^{(\alpha)}(x|q) {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \right] q; -z\Theta_{uv} \left\{ p_n(v, u) \right\} \frac{(qt)^n}{(q; q)_n} \\
 &= {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \right] q; -z\Theta_{uv} \left\{ \sum_{n=0}^{\infty} \psi_n^{(\alpha)}(x|q) p_n(v, u) \frac{(qt)^n}{(q; q)_n} \right\} \\
 &= {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \right] q; -z\Theta_{uv} \left\{ \sum_{k=0}^{\infty} \frac{(1/(\alpha x); q)_k (\alpha q)^k}{(q; q)_k} \sum_{n=0}^{\infty} \frac{(q^{-k}; q)_n q^{nk}}{(q; q)_n} \frac{(uxtq^{1-n}; q)_{\infty}}{(vxtq^{1-n}; q)_{\infty}} \right\} \\
 &= \sum_{k=0}^{\infty} \frac{(1/(\alpha x); q)_k (\alpha q)^k}{(q; q)_k} \sum_{n=0}^{\infty} \frac{(q^{-k}; q)_n q^{nk}}{(q; q)_n} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \right] q; -z\Theta_{uv} \left\{ \frac{(uxtq^{1-n}; q)_{\infty}}{(vxtq^{1-n}; q)_{\infty}} \right\} \\
 &= \sum_{k=0}^{\infty} \frac{(1/(\alpha x); q)_k (\alpha q)^k}{(q; q)_k} \sum_{n=0}^{\infty} \frac{(q^{-k}; q)_n q^{nk}}{(q; q)_n} \frac{(uxtq^{1-n}; q)_{\infty}}{(vxtq^{1-n}; q)_{\infty}} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \right] q; xztq^{1-n} \\
 &= \frac{(uxtq; q)_{\infty}}{(vxtq; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(1/(\alpha x); q)_k (\alpha q)^k}{(q; q)_k} \sum_{n=0}^{\infty} \frac{(q^{-k}, uxtq^{1-n}, q)_n q^{nk}}{(vxtq^{1-n}, q; q)_n} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \right] q; xztq^{1-n} \\
 &= \frac{(uxtq; q)_{\infty}}{(vxtq; q)_{\infty}} \sum_{n=0}^{\infty} \frac{(-1)^n q^{\binom{n}{2}} (1/(uxt); q)_n}{(1/(vxt), q; q)_n} \left(\frac{u}{v}\right)^n {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \right] q; xztq^{1-n} \\
 & \quad \cdot \sum_{k=n}^{\infty} \frac{(1/(\alpha x); q)_k (\alpha q)^k}{(q; q)_{k-n}}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{(uxtq; q)_\infty}{(vxtq; q)_\infty} \sum_{n=0}^{\infty} \frac{(-1)^n q^{\binom{n}{2}} (1/(\alpha x), 1/(uxt); q)_n}{(1/(vxt), q; q)_n} \left(\frac{\alpha uq}{v} \right)^n {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \begin{matrix} q, xztq^{1-n} \end{matrix} \right] \\
&\quad \cdot \sum_{k=0}^{\infty} \frac{(q^n/(\alpha x); q)_k (\alpha q)^k}{(q; q)_k} \\
&= \frac{(uxtq; q)_\infty}{(vxtq; q)_\infty} \sum_{n=0}^{\infty} \frac{(-1)^n q^{\binom{n}{2}} (1/(\alpha x), 1/(uxt); q)_n}{(1/(vxt), q; q)_n} \left(\frac{\alpha uq}{v} \right)^n {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \begin{matrix} q, xztq^{1-n} \end{matrix} \right] \\
&\quad \cdot \frac{(q^{1+n}/x; q)_\infty}{(\alpha q; q)_\infty} \\
&= \frac{(q/x, uxtq; q)_\infty}{(\alpha q, vxtq; q)_\infty} \sum_{n=0}^{\infty} \frac{(-1)^n q^{\binom{n}{2}} (1/(\alpha x), 1/(uxt); q)_n}{(q/x, 1/(vxt), q; q)_n} \left(\frac{\alpha uq}{v} \right)^n {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \begin{matrix} q, xztq^{1-n} \end{matrix} \right],
\end{aligned}$$

which is precisely the right-hand side of the assertion (4.3) of Theorem 3. The proof of Theorem 3 is thus completed. \square

Remark 4. In view of the special case (1.33) of $\Psi_n^{(a,b)}(u, v, z|q)$, we deduce the bilinear generating function for the Hahn polynomials $\psi_n^{(\alpha)}(x|q)$ as asserted by Corollary 1 below.

Corollary 1. Let $\max\{|\alpha q|, |axytq|\} < 1$. Then

$$\begin{aligned}
&\sum_{n=0}^{\infty} \psi_n^{(\alpha)}(x|q) \psi_n^{(a)}(y|q) \frac{(-1)^n q^{\binom{n+1}{2}} t^n}{(q; q)_n} \\
&= \frac{(q/x, xytq, xtq; q)_\infty}{(\alpha q, axytq; q)_\infty} {}_3\Phi_2 \left[\begin{matrix} 1/(\alpha x), 1/(xyt), 1/(xt); \\ q, \frac{\alpha xtq}{a} \end{matrix} \begin{matrix} q/x, 1/(axyt); \end{matrix} \right]. \quad (4.6)
\end{aligned}$$

Remark 5. For $r = s$, $\mathbf{a} = \mathbf{b} = \mathbf{0}$, $x = y$, $y = ay$ and $z = 1$, the assertion (4.3) reduces to Corollary 1.

5. A Transformational Identity Involving Generating Functions for the Generalized q -Hypergeometric Polynomials

In this section, we derive the following transformational identity involving generating functions for the generalized q -hypergeometric polynomials. Once again, in our derivation, we apply the homogeneous q -difference operator (2.1).

Theorem 4. Let $A(n)$ and $B(n)$ satisfy the following relationship:

$$\sum_{n=0}^{\infty} A(n) P_n(v, u) = \sum_{n=0}^{\infty} B(n) \frac{(xutq^{1-n}; q)_\infty}{(xvtq^{1-n}; q)_\infty}. \quad (5.1)$$

Then

$$\sum_{n=0}^{\infty} (-1)^n q^{\binom{n}{2}} A(n) \Psi_n^{(a,b)}(u, v, z|q) = \sum_{n=0}^{\infty} B(n) \frac{(xutq^{1-n}; q)_\infty}{(xvtq^{1-n}; q)_\infty} {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \begin{matrix} q, xztq^{1-n} \end{matrix} \right], \quad (5.2)$$

provided that each of the series in (5.1) and (5.2) is absolutely convergent.

Proof. If we denote by $f(x, u, v, z)$ the right-hand side of (5.2), we find by using (5.1) and (2.2) that

$$\begin{aligned}
 f(x, u, v, z) &= \sum_{n=0}^{\infty} B(n) {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; -z\Theta_{uv} \right] \left\{ \frac{(xutq^{1-n}; q)_{\infty}}{(xvtq^{1-n}; q)_{\infty}} \right\} \\
 &= {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; -z\Theta_{uv} \right] \left\{ \sum_{n=0}^{\infty} B(n) \frac{(xutq^{1-n}; q)_{\infty}}{(xvtq^{1-n}; q)_{\infty}} \right\} \\
 &= {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; -z\Theta_{uv} \right] \left\{ \sum_{n=0}^{\infty} A(n) P_n(v, u) \right\} \\
 &= \sum_{n=0}^{\infty} A(n) {}_r\Phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r; \\ b_1, b_2, \dots, b_s; \end{matrix} \middle| q; -z\Theta_{uv} \right] \{P_n(v, u)\} \\
 &= \sum_{n=0}^{\infty} (-1)^n q^{\binom{n}{2}} A(n) \Psi_n^{(a, b)}(u, v, z|q),
 \end{aligned}$$

which is precisely the left-hand side of (5.2). Our demonstration of Theorem 4 is thus completed. \square

Remark 6. Upon setting $u = 1, v = \lambda$,

$$A(n) = \psi_n^{(\alpha)}(x|q) \frac{(qt)^n}{(q; q)_n} \quad \text{and} \quad B(n) = \sum_{k=0}^{\infty} \frac{(1/(\alpha x); q)_k (\alpha q)^k (q^{-k}; q)_n q^{nk}}{(q; q)_k (q; q)_n} \tag{5.3}$$

in (5.1), we get (4.2). Moreover, Upon specializing $A(n)$ and $B(n)$ in (5.2) as in the last equation (5.3), we get the assertion (4.3) of Theorem 3.

6. Concluding Remarks and Observations

In our present investigation, we have introduced a general family of q -hypergeometric polynomials and we have derived several q -series identities such as an extended generating function and Srivastava-Agarwal type bilinear generating functions for this family of q -hypergeometric polynomials. We have presented a transformational identity involving generating functions for the generalized q -hypergeometric polynomials which we have introduced here. We have also pointed out relevant connections of the various q -results, which we have investigated in this paper, with those in several related earlier works on this subject.

We conclude this paper by remarking that, in the recently-published survey-cum-expository review article by Srivastava [19], the so-called (p, q) -calculus was exposed to be a rather trivial and inconsequential variation of the classical q -calculus, the additional parameter p being redundant or superfluous (see, for details, [19, p. 340]). This observation by Srivastava [19] will indeed apply to any attempt to produce the rather straightforward (p, q) -variations of the results which we have presented in this paper.

Conflicts of Interest: The authors declare that they have no conflicts of interest.

REFERENCES

- [1] M. A. Abdlhusein, Two operator representations for the trivariate q -polynomials and Hahn polynomials, *Ramanujan J.* **40** (2016), 491–509.
- [2] W. A. Al-Salam and L. Carlitz, Some orthogonal q -polynomials, *Math. Nachr.* **30** (1965), 47–61.
- [3] Jian Cao and S. Arjika, A note on fractional Askey-Wilson integrals, *J. Frac. Calc. Appl.* **12** (2) (2021), 133–140.
- [4] J. Cao, Bivariate generating functions for Rogers-Szegö polynomials, *Appl. Math. Comput.* **217** (2010), 2209–2216.
- [5] J. Cao, Generalizations of certain Carlitz’s trilinear and Srivastava-Agarwal type generating functions, *J. Math. Anal. Appl.* **396** (2012), 351–362.
- [6] J. Cao, A note on q -integrals and certain generating functions, *Stud. Appl. Math.* **131** (2013), 105–118.
- [7] J. Cao, Homogeneous q -difference equations and generating functions for q -hypergeometric polynomials, *Ramanujan J.* **40** (2016), 177–192.
- [8] J. Cao, B. Xu and S. Arjika, A note on generalized q -difference equations for general Al-Salam–Carlitz polynomials, *Adv. Differ. Equ.* **2020** (2020), Article ID 668, 1–17.
- [9] W. Y. C. Chen, A. M. Fu and B. Zhang, The homogeneous q -difference operator, *Adv. Appl. Math.* **31** (2003), 659–668.
- [10] W. Y. C. Chen and Z.-G. Liu, Parameter augmenting for basic hypergeometric series. II, *J. Combin. Theory Ser. A* **80** (1997), 175–195.
- [11] G. Gasper and M. Rahman, *Basic Hypergeometric Series* (with a Foreword by Richard Askey), Second edition, Encyclopedia of Mathematics and Its Applications, Vol. **96**, Cambridge University Press, Cambridge, London and New York, 2004.
- [12] W. Hahn, Über Orthogonalpolynome, die q -Differenzgleichungen genügen, *Math. Nachr.* **2** (1949), 4–34.
- [13] W. Hahn, Über Polynome, die gleichzeitig zwei verschiedenen Orthogonalsystemen angehören, *Math. Nachr.* **2** (1949), 263–278.
- [14] W. Hahn, Beiträge zur Theorie der Heineschen Reihen. Die 24 Integrale der hypergeometrischen q -Differenzgleichung. Das q -Analogon der Laplace-Transformation, *Math. Nachr.* **2** (1949), 340–379.
- [15] R. Koekoek and R. F. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogue*, Report No. 98-17, Delft University of Technology, Delft, The Netherlands, 1998.
- [16] H. L. Saad and A. A. Sukhi, Another homogeneous q -difference operator, *Appl. Math. Comput.* **215** (2010), 4332–4339.
- [17] H. L. Saad and S. M. Khalaf, The Operator and the Polynomials, *Basrah Journal of Science* **37** (1) (2019), 25–43.
- [18] L. J. Slater, *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, London and New York, 1966.
- [19] H. M. Srivastava, Operators of basic (or q -) calculus and fractional q -calculus and their applications in geometric function theory of complex analysis, *Iran. J. Sci. Technol. Trans. A: Sci.* **44** (2020), 327–344.
- [20] H. M. Srivastava and M. A. Abdlhusein, New forms of the Cauchy operator and some of their applications, *Russian J. Math. Phys.* **23** (2016), 124–134.
- [21] H. M. Srivastava and A. K. Agarwal, Generating functions for a class of q -polynomials, *Ann. Mat. Pura Appl. (Ser. 4)* **154** (1989), 99–109.
- [22] H. M. Srivastava and S. Arjika, Generating functions for some families of the generalized Al-Salam-Carlitz q -polynomials, *Adv. Differ. Equ.* **2020** (2020), Article ID 498, 1–17.
- [23] H. M. Srivastava, S. Arjika and A. S. Kelil, Some homogeneous q -difference operators and the associated generalized Hahn polynomials, *Appl. Set-Valued Anal. Optim.* **1** (2019), 187–201.
- [24] H. M. Srivastava and P. W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Halsted Press (Ellis Horwood Limited, Chichester); John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1985.