APPROXIMATION AND LOCALIZED POLYNOMIAL FRAME ON DOUBLE HYPERBOLIC AND CONIC DOMAINS

YUAN XU

ABSTRACT. We study approximation and localized polynomial frames on a bounded double hyperbolic or conic surface and the domain bounded by such a surface and hyperplanes. The main work follows the framework developed recently in [30] for homogeneous spaces that are assumed to contain highly localized kernels constructed via a family of orthogonal polynomials. The existence of such kernels will be established with the help of closed form formulas for the reproducing kernels. The main results provide a construction of semi-discrete localized tight frame in weighted L^2 norm and a characterization of best approximation by polynomials on our domains. Several intermediate results, including the Marcinkiewicz-Zygmund inequalities, positive cubature rules, Christoeffel functions, and Bernstein type inequalities, are shown to hold for doubling weights defined via the intrinsic distance on the domain.

1. INTRODUCTION

Recently a general framework based on highly localized polynomial kernels is developed in [30] for localizable homogeneous spaces and used for studying approximation and localized polynomial frame on a finite conic surface and the domain bounded by such a surface and a hyperplane. In the present work, we establish highly localized kernels and carry out analysis on bounded double hyperbolic or conic surface and the domain bounded by such a surface and hyperplanes.

1.1. **Motivation.** Let Ω be a set in \mathbb{R}^d , either an algebraic surface or a domain with non-empty interior. A homogeneous space is a measure space $(\Omega, \varpi, \mathsf{d})$, where ϖ is a nonnegative doubling weight function with respect to the metric $\mathsf{d}(\cdot, \cdot)$ on Ω . We call a homogeneous space $(\Omega, \varpi, \mathsf{d})$ *localizable* if it contains highly localized kernels.

The kernels are constructed using orthogonal polynomials. Let dm be the Lebesgue measure on Ω . Assume that the weight function ϖ is regular so that

(1.1)
$$\langle f,g\rangle_{\varpi} := \int_{\Omega} f(x)g(x)\varpi(x)\mathrm{d}\mathsf{m}(x)$$

is a well defined inner product on the space of polynomials restricted to Ω . Let $\mathcal{V}_n(\Omega, \varpi)$ be the space of orthogonal polynomials of degree n with respect to this inner product. The projection operator $\operatorname{proj}_n : L^2(\Omega, \varpi) \mapsto \mathcal{V}_n(\Omega, \varpi)$ can be written as

(1.2)
$$\operatorname{proj}_{n}(\varpi; f, x) = \int_{\Omega} f(y) P_{n}(\varpi; x, y) \varpi(y) \mathrm{d}\mathbf{m}(y), \quad f \in L^{2}(\Omega, \varpi).$$

Date: May 25, 2021.

²⁰¹⁰ Mathematics Subject Classification. 41A10, 41A63, 42C10, 42C40.

Key words and phrases. Approximation, conic domain, Fourier orthogonal series, homogeneous spaces, localized kernel, orthogonal polynomials, polynomial frame.

where $P_n(\varpi; \cdot, \cdot)$ is the reproducing kernel of the space $\mathcal{V}_n(\Omega, \varpi)$. Let \hat{a} be a cut-off function, defined as a compactly supported function in $C^{\infty}(\mathbb{R}_+)$. Then our highly localized kernels are of the form

$$L_n(\varpi; x, y) := \sum_{j=0}^{\infty} \widehat{a}\left(\frac{j}{n}\right) P_j(\varpi; x, y).$$

The kernel is highly localized if it decays at rates faster than any inverse polynomial rate away from the main diagonal y = x in $\Omega \times \Omega$ with respect to the distance d on Ω ; see the definition in the next section. These kernels provide important tools for analysis on regular domains, such as the unit sphere and the unit ball, and are essential ingredient in recent study of approximation and localized polynomial frames; see, for example, [3, 8, 11, 14, 17, 18, 19] for some of the results on the spheres and balls and [1, 2, 12, 13, 15, 25] for various applications.

The reason that highly localized kernels are known only on a few regular domains lies in the *addition formula* for orthogonal polynomials, which are closed form formulas for the reproducing kernels of orthogonal polynomials. For spherical harmonics that are orthogonal on the unit sphere, which serve as a quintessential example for our study, the closed form formula is given by $Z_n(\langle x, y \rangle)$, where Z_n is a Gegenbauer polynomial of one variable. The addition formulas are powerful tools when they exist.

Our recent work in [30] is prompted by two aspirations. The first one is the realization that, if highly localized kernels are taken as granted, then much of the analysis can be developed within a general framework of homogeneous spaces. The second one is the possibility of establishing highly localized kernels and carrying out analysis thereafter on conic domains, which are domains largely untouched hitherto. The latter is made possible by recently discovered new addition formulas for orthogonal polynomials on conic domains [28, 29]. Altogether there are four types of of conic domains and they are standardized as:

(1) conic surface \mathbb{V}_0^{d+1} defined by

$$\mathbb{V}_0^{d+1} := \left\{ (x, t) \in \mathbb{R}^{d+1} : \|x\| = t, \, 0 \le t \le 1, \, x \in \mathbb{R}^d \right\};$$

- (2) solid cone \mathbb{V}^{d+1} bounded by \mathbb{V}_0^{d+1} and the hyperplane t = 1; (3) two-sheets hyperbolic surface ${}_{\varrho}\mathbb{X}_0^{d+1}$ defined by

$${}_{\varrho}\mathbb{X}_{0}^{d+1} := \left\{ (x,t) \in \mathbb{R}^{d+1} : \|x\|^{2} = t^{2} - \varrho^{2}, \ \varrho \leq |t| \leq 1 + \varrho, \ x \in \mathbb{R}^{d} \right\},$$

where $\rho \geq 0$, which becomes the double conic surface \mathbb{X}_0^{d+1} when $\rho = 0$; (4) solid hyperboloid ${}_{\varrho}\mathbb{X}^{d+1}$ bounded by ${}_{\varrho}\mathbb{X}_0^{d+1}$ and the hyperplanes $t = \pm 1$, which becomes solid double cone \mathbb{X}^{d+1} when $\rho = 0$.

The first two types of conic domains are studied in [30]. The present paper deals with the other two cases: double hyperbolic surfaces and hyperboloid.

1.2. Main results. We will follow the framework established in [30]. Assuming several assertions on highly localized kernels and on fast decaying polynomials, the framework provides a unified theory for two objectives. The first one leads to localized polynomial frames constructed via a semi-continuous Calderón type decomposition

$$f = \sum_{j=0}^{\infty} L_{2^j} * L_{2^j} * f, \qquad f \in L^2(\Omega, \varpi),$$

where $L_n * f$ denotes the integral operator that has $L_n(\varpi; \cdot, \cdot)$ as its kernel,

(1.3)
$$L_n * f(x) := \int_{\Omega} f(y) L_n(\varpi; x, y) \varpi(y) dy$$

Discretizing the integrals by appropriate positive cubature rules, we end up with a fame system $\{\psi_{\xi}\}_{\xi\in\Xi}$, indexed by a discrete set Ξ of well-separated points in Ω , where $\psi_{\xi}(x) = \sqrt{\lambda_{\xi}} L_{2^{j}}(\varpi; x, \xi)$ with $\lambda_{\xi} > 0$ being the coefficients in the cubature, and the frame is tight in the sense that, for all $f \in L^{2}(\Omega, \varpi)$,

$$f = \sum_{\xi \in \Xi} \langle f, \psi_{\xi} \rangle_{\varpi} \psi_{\varpi} \quad \text{and} \quad \int_{\Omega} |f(x)|^2 \varpi(x) \mathrm{d}x = \sum_{\xi \in \Xi} |\langle f, \psi_{\xi} \rangle|^2.$$

This is an extension of the extensive work on the unit sphere and the unit ball in the literature that we have described in the previous subsection. The second objective is to study the error of the best approximation by polynomials

$$E_n(f)_{L^p(\Omega,\varpi)} = \inf_{\deg g \le n} \|f - g\|_{L^p(\Omega,\varpi)}.$$

The aim is to provide a characterization via a modulus of smoothness, defined as a multiplier operator, and an equivalent K-functional, defined by the spectral differential operator that has orthogonal polynomials as eigenfunctions. Such a characterization is in line with those on the unit sphere, see [8, 20, 27] and the references therein. The characterization consists of a direct estimate, using the fast decaying of $L_n(\varpi; \cdot, \cdot)$, and an inverse estimate, using a Bernstein inequality established via highly localized kernels. It is worth mentioning that several intermediate results, such as the Marcinkiewicz-Zygmund inequalities, positive cubature rules, Christoeffel functions, and Bernstein type are of independent interests; furthermore, some of these intermediate results can be established for doubling weights, which extends the results on the interval and on the unit sphere [4, 8, 16].

In order for the framework to work on a particular domain, substantial work is needed to fulfill the assertions and assumptions, which depends heavily on the geometry of the domain and the complexity of the orthogonal structure on the domain. For each conic domain, this starts with identifying an appropriate intrinsic distance on the domain. Establishing the highly localized kernels is delicate and fairly involved, relying on the structure of addition formula and the distance function. The ε -separated set is assumed conceptually in the framework, an explicit construction is needed on each domain, which is also necessary for computational purpose. Even though the roadmap is outlined by the framework, the actual work on each domain remains challenging and amounts to a thorough understanding of the intrinsic structure on the domain.

It should also be mentioned that the structures on double hyperbolic domains \mathbb{X}_0^{d+1} and \mathbb{X}^{d+1} , which degenerate to double cones when $\rho = 0$, are distinctively different from those on the single conic domains \mathbb{V}_0^{d+1} and \mathbb{V}^{d+1} . Indeed, the distance functions for these two typos of domains are incomparable, around the conic apex, and the orthogonal structure on the double conic domains depends on the parity of the polynomials on the the variable t. For example, for the double conic surface \mathbb{X}_0^{d+1} , the closed form formula for the reproducing kernels is stablished for polynomials that are orthogonal with respect to $\mathsf{w}_{\beta,\gamma}(t) = |t|^{2\beta}(1-t^2)^{\gamma-\frac{1}{2}}$, but only for the subspace of polynomials that are even in t variable or odd in t variable. The restriction requires us to restrict to the class of functions that are even in t variable, but the entire framework can still be carried out on the domain. Using the reproducing kernels in [29], we will show that

 $(\mathbb{X}_{0}^{d+1}, \mathsf{w}_{0,\gamma}, \mathsf{d}_{\mathbb{X}_{0}})$ is a localizable homogeneous space. Similar result holds for the solid double cone \mathbb{X}^{d+1} with the weight function $W_{\beta,\gamma,\mu}(x,t) = |t|^{\beta}(1-t)^{\gamma}(t^{2}-||x||^{2})^{\mu-\frac{1}{2}}$ and its own distance function. For these homogeneous spaces we shall show that the framework in [30] on localizable tight frame and on the best approximation can be carried out completely.

1.3. Organization and convention. The general framework on the localizable homogeneous space emerges from the study on the unit sphere and the unit ball, as explained in [30], and it consists of several topics. Because of its length, we shall refer to its discussion and formulation to [30]. The present work, however, is self-contained otherwise and can be read independently.

The paper is organized as follows. We will state the assertions and just enough background to state them for the general framework in the next section. Whenever possible, we will not give definition and properties for homogeneous spaces but only on conic domains and only when they are needed. For a more extensive discussion on the background materials, we also refer to [30]. The cases of hyperbolic surface and hyperboloid will be studied in Section 3 and Section 4, respectively, and the two sections will have parallel structures, which is also the structure for the conic surface and the cone in [30], to emphasis what is required to carry out the program and for easier comparison between the cases. Each section will contain several sections and its organization will be described in the preamble of the section.

Throughout this paper, we will denote by $L^p(\Omega, \mathbf{w})$ the weighted L^p space with respect to the weight function \mathbf{w} defined on the domain Ω for $1 \leq p \leq \infty$. Its norm will be denote by $\|\cdot\|_{p,\mathbf{w}}$ for $1 \leq p \leq \infty$ with the understanding that the space is $C(\Omega)$ with the uniform norm when $p = \infty$.

Finally, we shall use c, c', c_1, c_2 etc. to denote positive constants that depend on fixed parameters and their values may change from line to line. Furthermore, we shall write $A \sim B$ if $c'A \leq B \leq cA$.

2. Preliminaries

In the first subsection we collect all assertions needed for the general framework on the localized homogeneous space. Basics of classical orthogonal polynomials that will be used latter are collected in the second subsection.

2.1. Assertions for the general framework. Let $(\Omega, \varpi, \mathsf{d})$ be a homogeneous space. For a given set $E \subset \Omega$, we define $\varpi(E) = \int_E \varpi(x) \mathrm{d}\mathfrak{m}(x)$. The weight function ϖ is a doubling weight if there exists a constant L > 0 such that

$$\varpi(B(x,2r)) \le L\varpi(B(x,r)), \qquad \forall x \in \Omega, \quad r \in (0,r_0),$$

where r_0 is the largest positive number such that $B(x,r) = \{y \in \Omega : d(x,t) < r\} \subset \Omega$. The smallest L for doubling inequality to hold is called the doubling constant of ϖ .

Let ϖ be a nonnegative weight function on Ω and let $\langle \cdot, \cdot \rangle_{\varpi}$ be the inner product defined by (1.1). For $n = 0, 1, 2, \ldots$, let $\mathcal{V}_n(\Omega, \varpi)$ be the space of orthogonal polynomials with respect to the inner product. If $\{P_{\nu,n} : 1 \leq \nu \leq \dim \mathcal{V}_n(\Omega, \varpi)\}$ is an orthogonal basis of $\mathcal{V}_n(\Omega, \varpi)$, then the reproducing kernel of $\mathcal{V}_n(\Omega, \varpi)$ satisfies

(2.1)
$$P_n(\varpi; x, y) = \sum_{\nu=1}^{\dim \mathcal{V}_n^d(\Omega, \varpi)} \frac{P_{\nu, n}(x) P_{\nu, n}(y)}{\langle P_{\nu, n}, P_{\nu, n} \rangle_{\varpi}}.$$

It is the kernel of the projection operator $\operatorname{proj}_n : L^2(\Omega, \varpi) \mapsto \mathcal{V}_n^d(\Omega, \varpi)$ stated in (1.2). The Fourier orthogonal series of $f \in L^2(\Omega, \varpi)$ is given by

(2.2)
$$f = \sum_{n=0}^{\infty} \operatorname{proj}_{n}(\varpi; x, y)$$

2.1.1. Highly localized kernels. A nonnegative function $\hat{a} \in C^{\infty}(\mathbb{R})$ is said to be admissible if it obeys either one of the conditions

(a) supp $\hat{a} \subset [0,2]$ and $\hat{a}(t) = 1, t \in [0,1]$; or (b) supp $\hat{a} \subset [1/2,2]$.

Given such a cut-off function, we define a kernel $L_n(\varpi; \cdot, \cdot)$ on $\Omega \times \Omega$ by

$$L_n(\varpi; x, y) := \sum_{j=0}^{\infty} \widehat{a}\left(\frac{j}{n}\right) P_j(\varpi; x, y)$$

Definition 2.1. The kernels $L_n(\varpi; \cdot, \cdot)$, n = 1, 2, ..., are called highly localized if they satisfy the following assertions:

Assertion 1. For $\kappa > 0$ and $x, y \in \Omega$,

$$|L_n(\varpi; x, y)| \le c_{\kappa} \frac{1}{\sqrt{\varpi(B(x, n^{-1}))} \sqrt{\varpi(B(x, n^{-1}))} (1 + n\mathsf{d}(x, y))^{\kappa}}.$$

Assertion 2. For $0 < \delta \leq \delta_0$ with some $\delta_0 < 1$ and $x_1 \in B(x_2, \frac{\delta}{n})$,

$$|L_n(\varpi; x_1, y) - L_n(\varpi; x_2, y)| \le c_{\kappa} \frac{n \mathsf{d}(x_1, x_2)}{\sqrt{\varpi(B(x_1, n^{-1}))} \sqrt{\varpi(B(x_2, n^{-1}))} (1 + n \mathsf{d}(x_2, y))^{\kappa}}$$

Assertion 3. For sufficient large $\kappa > 0$,

$$\int_{\Omega} \frac{\varpi(y)}{\varpi(B(y, n^{-1})) \left(1 + n\mathsf{d}(x, y)\right)^{\kappa}} \mathrm{d}\mathsf{m}(y) \le c.$$

The third assertion affirms the sharpness of the first two assertions, as can be seen from the following inequality [30, Lemma 2.1.4].

Lemma 2.2. Let ϖ be a doubling weight that satisfies Assertion 3 with $\kappa > 0$. For $0 , let <math>\tau = \kappa - \frac{p}{2}\alpha(\varpi)|1 - \frac{p}{2}| > 0$. Then, for $x \in \Omega$,

(2.3)
$$\int_{\Omega} \frac{\varpi(y)}{\varpi(B(y,n^{-1}))^{\frac{p}{2}} (1 + n\mathsf{d}(x,y))^{\tau}} \mathrm{d}\mathsf{m}(y) \le c \,\varpi \left(B(x,n^{-1})\right)^{1-\frac{p}{2}}.$$

In particular, the highly localized kernel satisfies

(2.4)
$$\int_{\Omega} |L_n(\varpi; x, y)|^p \, \varpi(s) \mathrm{d}\mathsf{m}(y) \le c \left[\varpi \left(B(x, n^{-1}) \right) \right]^{1-p}$$

The homogeneous space $(\Omega, \varpi, \mathsf{d})$ is called *localizable* if ϖ is a doubling weight that admits highly localized kernels.

2.1.2. Well-separated set of points. We will need well distributed points for discretization, such as in the Marcinkiewicz-Zygmund inequality and the positive cubature rules, and in our localized polynomial frame. The precise definition is given as follows.

Definition 2.3. Let Ξ be a discrete set in Ω .

- (a) A finite collection of subsets $\{S_z : z \in \Xi\}$ is called a partition of Ω if $S_z^\circ \cap S_y^\circ = \emptyset$ when $z \neq y$ and $\Omega = \bigcup_{z \in \Xi} S_z$.
- (b) Let $\varepsilon > 0$. A discrete subset Ξ of Ω is called ε -separated if $d(x, y) \ge \varepsilon$ for every two distinct points $x, y \in \Xi$.
- (c) Ξ is called maximal if there is a constant $c_d > 1$ such that

(2.5)
$$1 \le \sum_{z \in \Xi} \chi_{B(z,\varepsilon)}(x) \le c_d, \qquad \forall x \in \Omega,$$

where χ_E denotes the characteristic function of the set E.

The existence of ε -separated points is assumed in the general framework for homogeneous spaces in [30]. The construction of such points on a given domain depends on the geometry and the distance function, and it is crucial for explicit formulation and practical computation of localized frames. For conic domains \mathbb{X}_0^{d+1} and \mathbb{X}^{d+1} , we will give a construction of such points in latter sections.

2.1.3. Fast decaying polynomials and cubature rules. Let w be a doubling weight on Ω . A cubature rule is a finite sum that discretize a given integral,

$$\int_{\Omega} f(z) \mathsf{w}(x) \mathrm{d}x = \sum_{k=1}^{N} \lambda_k f(x_k),$$

where the equality holds for a given polynomials subspace. We will need positive cubature rules, for which all λ_k are positive, over ε -separated points. To quantify the coefficients λ_k , which are used for our localized polynomial frame, we need to show that the Christoffel function $\lambda_n(\mathbf{w}; \cdot)$, defined by

(2.6)
$$\lambda_n(\mathsf{w};x) := \inf_{\substack{g(x)=1\\g\in\Pi_n(\Omega)}} \int_{\Omega} |g(x)|^2 \mathsf{w}(x) \mathrm{d}\mathsf{m}(x),$$

are bounded, where $\Pi_n(\Omega)$ denotes the space of polynomials of degree *n* restricted on Ω . This is where our fourth assertion comes in, which ensures the existence of fast decaying polynomials on the domain Ω .

Assertion 4. Let Ω be compact. For each $x \in \Omega$, there is a nonnegative polynomial T_x of degree at most n that satisfies

(1) $T_x(x) = 1, T_x(y) \ge \delta > 0$ for $y \in B(x, \frac{1}{n})$ for some δ independent of n, and, for each $\gamma > 1$,

$$0 \le T_x(y) \le c_\gamma (1 + n\mathsf{d}(x, y))^{-\gamma}, \qquad y \in \Omega;$$

(2) there is a polynomial q_n such that $q_n(x)T_x(y)$ is a polynomial of degree at most rn, for some positive integer r, in x-variable and $c_1 \leq q_n(x) \leq c_2$ for $x \in \Omega$ for some positive numbers c_1 and c_2 .

For conic domains, we will establish this assertion by explicitly construction.

2.1.4. Bernstein inequality for the spectral operator. Our study of the best approximation by polynomials relies on the existence of a spectral operator for orthogonal polynomials.

Definition 2.4. Let ϖ be a weight function on Ω . We denote by \mathfrak{D}_{ϖ} the second order derivation operator that has orthogonal polynomials with respect to ϖ as eigenfunctions; more precisely,

(2.7)
$$\mathfrak{D}_{\varpi}Y = -\mu(n)Y, \quad \forall Y \in \mathcal{V}_n(\Omega, \varpi),$$

where μ is a nonnegative quadratic polynomial.

For the unit sphere, this is the Laplace-Beltrami operator. Its analogues exist on weighted sphere, ball and simplexes (cf. [9]), as well as for conic domains [28, 29]. We will need the Bernstein inequality for the power of these operators. Since $\mu(k) \geq 0$, the operator $-\mathfrak{D}_{\varpi}$ is a non-negative operator. A function $f \in L^p(\Omega; \varpi)$ belongs to the Sobolev space $W_p^r(\Omega; \varpi)$ if there is a function $g \in L^p(\Omega; \varpi)$, which we denote by $(-\mathfrak{D}_{\varpi})^{\frac{r}{2}} f$, such that

(2.8)
$$\operatorname{proj}_{n}\left(\varpi;(-\mathfrak{D}_{\varpi})^{\frac{r}{2}}f\right) = \mu(n)^{\frac{r}{2}}\operatorname{proj}_{n}(\varpi;f),$$

where we assume that $f, g \in C(\Omega)$ when $p = \infty$. The fractional differential operator $(-\mathfrak{D}_{\varpi})^{\frac{r}{2}}f$ is a linear operator on the space $W_p^r(\Omega; \varpi)$ defined by (2.8).

For r > 0, we denote by $L_n^{(r)}(\varpi; \cdot, \cdot)$ the kernel defined by

(2.9)
$$L_n^{(r)}(\varpi; x, y) = \sum_{n=0}^{\infty} \widehat{a}\left(\frac{k}{n}\right) [\mu(k)]^{r/2} P_k(\varpi; x, y),$$

which is the kernel $\mathfrak{D}_{\varpi}^{r/2}L_n(x,y)$ with $\mathfrak{D}_{\varpi}^{r/2}$ applying on x variable. Our Bernstein inequality is proved under the following assumption on the decaying of this kernel.

Assertion 5. For r > 0 and $\kappa > 0$, the kernel $L_n^{(r)}(\varpi)$ satisfies, for $x, y \in \Omega$,

$$\left|L_n^{(r)}(\varpi;x,y)\right| \le c_{\kappa} \frac{n'}{\sqrt{\varpi(B(x,n^{-1}))}\sqrt{\varpi(B(y,n^{-1}))}} (1+n\mathsf{d}(x,y))^{-\kappa}.$$

For r = 0, this reduces to the Assertion 1. We list it separately since Assertion 5 is not needed for the localized polynomial frames.

2.2. Classical orthogonal polynomials. In this subsection we collect a few results on classical orthogonal polynomials that we will need. There are three families, Jacobi polynomials, spherical harmonics and classical orthogonal polynomials on the ball. Each of three families comes from an example of a localized homogeneous space, as explained in [30], and has been extensively studied. We will recall only the basics.

2.3. Jacobi polynomials. For $\alpha, \beta > -1$, the Jacobi weight function is defined by

$$w_{\alpha,\beta}(t) := (1-t)^{\alpha}(1+t)^{\beta}, \qquad -1 < x < 1.$$

Its normalization constant $c'_{\alpha,\beta}$, defined by $c'_{\alpha,\beta} \int_{-1}^{1} w_{\alpha,\beta}(x) dx = 1$, is given by

(2.10)
$$c'_{\alpha,\beta} = \frac{1}{2^{\alpha+\beta+1}} c_{\alpha,\beta} \quad \text{with} \quad c_{\alpha,\beta} := \frac{\Gamma(\alpha+\beta+2)}{\Gamma(\alpha+1)\Gamma(\beta+1)}.$$

The Jacobi polynomials $P_n^{(\alpha,\beta)}$ satisfy the orthogonal relations

$$c_{\alpha,\beta}' \int_{-1}^{1} P_n^{(\alpha,\beta)}(x) P_m^{(\alpha,\beta)}(x) w_{\alpha,\beta}(x) \mathrm{d}x = h_n^{(\alpha,\beta)} \delta_{m,n},$$

where $h_n^{(\alpha,\beta)}$ is the square of the L^2 norm that satisfies

$$h_n^{(\alpha,\beta)} = \frac{(\alpha+1)_n(\beta+1)_n(\alpha+\beta+n+1)}{n!(\alpha+\beta+2)_n(\alpha+\beta+2n+1)}.$$

Let \widehat{a} be an admissible cut-off function. We defined a polynomial $L_n^{(\alpha,\beta)}$ by

(2.11)
$$L_n^{(\alpha,\beta)}(t) = \sum_{j=0}^{\infty} \widehat{a}\left(\frac{j}{n}\right) \frac{P_j^{(\alpha,\beta)}(t)P_j^{(\alpha,\beta)}(1)}{h_j^{(\alpha,\beta)}}$$

The estimate stated below ([3] and [8, Theorem 2.6.7]) will be used several times.

Theorem 2.5. Let ℓ be a positive integer and let η be a function that satisfy, $\eta \in C^{3\ell-1}(\mathbb{R})$, supp $\eta \subset [0,2]$ and $\eta^{(j)}(0) = 0$ for $j = 0, 1, 2, \ldots, 3\ell-2$. Then, for $\alpha \geq \beta \geq -\frac{1}{2}$, $t \in [-1,1]$ and $n \in \mathbb{N}$,

(2.12)
$$\left| \frac{d^m}{dt^m} L_n^{(\alpha,\beta)}(t) \right| \le c_{\ell,m,\alpha} \left\| \eta^{(3\ell-1)} \right\|_\infty \frac{n^{2\alpha+2m+2}}{(1+n\sqrt{1-t})^\ell}, \quad m = 0, 1, 2, \dots$$

The Jacobi polynomials with equal parameters are the Gegenbauer polynomials C_n^{λ} , which are orthogonal with respect to $w_{\lambda}(x) = (1 - x^2)^{\lambda - \frac{1}{2}}$,

(2.13)
$$c_{\lambda} \int_{-1}^{1} C_{n}^{\lambda}(x) C_{m}^{\lambda}(x) w_{\lambda}(x) \mathrm{d}x = h_{n}^{\lambda} \delta_{n,m},$$

where the normalization constant c_{λ} of w_{λ} and the norm square h_n^{λ} are given by

(2.14)
$$c_{\lambda} = \frac{\Gamma(\lambda+1)}{\Gamma(\frac{1}{2})\Gamma(\lambda+\frac{1}{2})}$$
 and $h_n^{\lambda} = \frac{\lambda}{n+\lambda}C_n^{\lambda}(1) = \frac{\lambda}{n+\lambda}\frac{(2\lambda)_n}{n!}$

2.3.1. Spherical harmonics. Spherical harmonics are homogeneous polynomials that are orthogonal on the unit sphere. Let $\mathcal{H}_n(\mathbb{S}^{d-1})$ be the space of spherical harmonics of degree n in d variables. Then $\dim \mathcal{H}_n(\mathbb{S}^{d-1}) = \binom{n+d-2}{n} + \binom{n+d-3}{n-1}$. For $n \in \mathbb{N}_0$ let $\{Y_\ell^n : 1 \leq \ell \leq \dim \mathcal{H}_n(\mathbb{S}^{d-1})\}$ be an orthonormal basis of $\mathcal{H}_n(\mathbb{S}^{d-1})$; then

$$\frac{1}{\omega_d} \int_{\mathbb{S}^{d-1}} Y_\ell^n(\xi) Y_{\ell'}^m(\xi) \mathrm{d}\sigma(\xi) = \delta_{n,m},$$

where $d\sigma$ is the surface measure of \mathbb{S}^{d-1} and ω_d denotes the surface area $\omega_d = 2\pi^{\frac{d}{2}}/\Gamma(\frac{d}{2})$ of \mathbb{S}^{d-1} . The spherical harmonics satisfy two characteristic properties. In terms of the orthonormal basis $\{Y_{\ell}^n\}$, its reproducing kernel $\mathsf{P}_n(\xi,\eta)$ satisfies the addition formula

(2.15)
$$\mathsf{P}_{n}(\xi,\eta) = \sum_{\ell=1}^{\dim \mathcal{H}_{n}(\mathbb{S}^{d-1})} Y_{\ell}^{n}(\xi) Y_{\ell}^{n}(\eta) = Z_{n}^{\frac{d-2}{2}}(\langle \xi,\eta \rangle), \quad \xi,\eta \in \mathbb{S}^{d-1},$$

where Z_n^{λ} is defined in terms of the Gegenbauer polynomial by

(2.16)
$$Z_n^{\lambda}(t) = \frac{n+\lambda}{\lambda} C_n^{\lambda}(t), \qquad \lambda = \frac{d-2}{2}.$$

Furthermore, spherical harmonics are the eigenfunctions of the Laplace-Beltrami operator Δ_0 , which is the restriction of the Laplace operator Δ on the unit sphere. More precisely (cf. [8, (1.4.9)])

(2.17)
$$\Delta_0 Y = -n(n+d-2)Y, \qquad Y \in \mathcal{H}_n^d$$

2.3.2. Orthogonal polynomials on the unit ball. The classical weight function on the unit ball \mathbb{B}^d of \mathbb{R}^d is defined by

(2.18)
$$W_{\mu}(x) = (1 - ||x||)^{\mu - \frac{1}{2}}, \quad x \in \mathbb{B}^d, \quad \mu > -\frac{1}{2}.$$

Its normalization constant is $b_{\mu}^{\mathbb{B}} = \Gamma(\mu + \frac{d+1}{2})/(\pi^{\frac{d}{2}}\Gamma(\mu + \frac{1}{2}))$. Let $\mathcal{V}_{n}^{d}(\mathbb{B}^{d}, W_{\mu})$ be the space of orthogonal polynomials of degree n with respect to W_{μ} . An orthogonal basis of $\mathcal{V}_{n}^{d}(\mathbb{B}^{d}, W_{\mu})$ can be given explicitly in terms of the Jacobi polynomials and spherical harmonics. For $0 \leq m \leq n/2$, let $\{Y_{\ell}^{n-2m} : 1 \leq \ell \leq \dim \mathcal{H}_{n-2m}(\mathbb{S}^{d-1})\}$ be an orthonormal basis of \mathcal{H}_{n-2m}^{d} . Define [9, (5.2.4)]

(2.19)
$$P_{\ell,m}^{n}(x) = P_{m}^{(\mu - \frac{1}{2}, n - 2m + \frac{d-2}{2})} \left(2\|x\|^{2} - 1\right) Y_{\ell}^{n-2m}(x).$$

Then $\{P_{\ell,m}^n : 0 \leq m \leq n/2, 1 \leq \ell \leq \dim \mathcal{H}_{n-2m}(\mathbb{S}^{d-1})\}$ is an orthogonal basis of $\mathcal{V}_n^d(W_\mu)$. Let $\mathbf{P}_n(W_\mu; \cdot, \cdot)$ denote the reproducing kernel of the space $\mathcal{V}_n(\mathbb{B}^d, W_\mu)$. This kernel satisfies an analogue of the addition formula [26]: for $x, y \in \mathbb{B}^d$,

(2.20)
$$\mathbf{P}_{n}(W_{\mu}; x, y) = c_{\mu - \frac{1}{2}} \int_{-1}^{1} Z_{n}^{\mu + \frac{d-1}{2}} \left(\langle x, y \rangle + u \sqrt{1 - \|x\|^{2}} \sqrt{1 - \|y\|^{2}} \right) \times (1 - u^{2})^{\mu - 1} \mathrm{d}u,$$

where $\mu > 0$ and it holds for $\mu = 0$ under the limit

(2.21)
$$\lim_{\mu \to 0+} c_{\mu-\frac{1}{2}} \int_{-1}^{1} f(t)(1-t^2)^{\mu-1} \mathrm{d}u = \frac{f(1)+f(-1)}{2}.$$

The orthogonal polynomials with respect to W_{μ} on the unit ball are eigenfunctions of a second order differential operator [9, (5.2.3)]: for all $u \in \mathcal{V}_n(\mathbb{B}^d, W_{\mu})$,

(2.22)
$$\left(\Delta - \langle x, \nabla \rangle^2 - (2\mu + d - 1)\langle x, \nabla \rangle\right)u = -n(n + 2\mu + d - 1)u$$

3. Homogeneous space on double conic and hyperbolic surfaces

In this chapter we work in the setting of homogeneous space on the surface

$$\mathbb{X}_{0}^{d+1} = \left\{ (x,t) : \|x\|^{2} = t^{2} - \varrho^{2}, \, x \in \mathbb{R}^{d}, \, \varrho \le |t| \le \sqrt{\varrho^{2} + 1} \right\}$$

which is a hyperbolic surface of two sheets when $\rho > 0$ and a double conic surface when $\rho = 0$. We shall verify that this domain admits a localized homogeneous space for a family of weight functions $w^{\rho}_{\beta,\gamma}$, which are related to the Gegenbauer weight functions and are even in t variable.

For $\rho = 0$, the upper part of \mathbb{X}_0^{d+1} with $t \ge 0$ is exactly the conic surface \mathbb{V}_0^{d+1} . The analysis on \mathbb{X}_0^{d+1} , however, is of a different character. As a starter, the distance on \mathbb{X}_0^{d+1} is comparable to the Euclidean distance, in contrast to the distance on \mathbb{V}_0^{d+1} . Moreover, the space of orthogonal polynomials are divided naturally into two orthogonal subspaces, consisting of polynomials even in t variable or odd in t variable. We consider only orthogonal polynomials even in t variable, since they alone satisfy the two characteristic properties: addition formula and the differential operator having orthogonal polynomials as eigenfunctions. As a consequence, we consider approximation

and localized frames for functions that are even in t variable. Many estimates and computations are easier to handle comparing with the conic surface because of simpler distance function and an addition formula that is less cumbersome. The structure of the chapter follows that of the previous two chapters, with contents arranging in the same order under similar section names.

3.1. Distance on double conic and hyperbolic surfaces. Whenever it is necessary to emphasis the dependence on ϱ , we write ${}_{\varrho}\mathbb{X}_{0}^{d+1}$ instead of \mathbb{X}_{0}^{d+1} . In most places, however, we use \mathbb{X}_{0}^{d+1} to avoid excessive notations. The surface \mathbb{X}_{0}^{d+1} can be decomposed as an upper part and a lower part,

$$\mathbb{X}_{0}^{d+1} = \mathbb{X}_{0,+}^{d+1} \cup \mathbb{X}_{0,-}^{d+1} = \{(x,t) \in \mathbb{X}_{0}^{d+1} : t \geq 0\} \cup \{(x,t) \in \mathbb{X}_{0}^{d+1} : t \leq 0\}.$$

For $\varrho = 0$, the upper part $\mathbb{X}_{0,+}^{d+1}$ is exactly the conic surface \mathbb{V}_0^{d+1} .

As \mathbb{X}_{0}^{d+1} is a bounded domain, its distance function should take into account of the boundary at the two ends, but it does not regard the apex as a boundary point because of symmetry. We first define a distance on the double conic surface; that is, when $\rho = 0$.

Definition 3.1. Let $\rho = 0$. For $(x, t), (y, s) \in \mathbb{X}_0^{d+1}$, define

$$\mathsf{d}_{\mathbb{X}_0}((x,t),(y,s)) = \arccos\left(\langle x,y\rangle + \sqrt{1-t^2}\sqrt{1-s^2}\right).$$

To see that $\mathsf{d}_{\mathbb{X}_0}(\cdot, \cdot)$ is indeed a distance on the double conic surface \mathbb{X}_0^{d+1} , we let $X = (x, \sqrt{1-t^2})$ and $Y = (y, \sqrt{1-s^2})$, so that $X, Y \in \mathbb{S}^d$ if $x, y \in \mathbb{X}_0^{d+1}$. Then

$$\mathsf{d}_{\mathbb{X}_0}((x,t),(y,s)) = \mathsf{d}_{\mathbb{S}^d}(X,Y), \quad (x,t),(y,s) \in \mathbb{X}_0^{d+1},$$

from which it follows readily that $\mathsf{d}_{\mathbb{X}_0}$ is indeed a distance on $\mathbb{X}_0^{d+1}.$ This distance is quite different from the distance $\mathsf{d}_{\mathbb{V}_0}(\cdot,\cdot)$ of \mathbb{V}_0^{d+1} defined by

$$\mathsf{d}_{\mathbb{V}_0}((x,t),(y,s)) = \arccos\left(\sqrt{\frac{ts + \langle x, y \rangle}{2}} + \sqrt{1 - t}\sqrt{1 - s}\right), \quad (x,t), (y,s) \in \mathbb{V}_0^{d+1}$$

in [30] and it resembles the distance $\mathsf{d}_{\mathbb{B}}(\cdot,\cdot)$ of the unit ball which is of the same form but with t and s replaced by ||x|| and ||y||. The distance $\mathsf{d}_{\mathbb{X}_0}$ is related to the distance on the sphere, defined by

(3.1)
$$\mathsf{d}_{\mathbb{S}}(\xi,\eta) = \arccos{\langle \xi,\eta \rangle}, \qquad \xi,\eta \in \mathbb{S}^{d-1},$$

and the distance on the interval [-1, 1], defined by

(3.2)
$$\mathsf{d}_{[-1,1]}(t,s) = \arccos\left(ts + \sqrt{1-t^2}\sqrt{1-s^2}\right), \quad t,s \in [-1,1].$$

Proposition 3.2. Let $\rho = 0$ and $d \ge 2$. For $(x, t), (y, s) \in \mathbb{X}_0^{d+1}$, write $x = t\xi$ and $y = s\eta$ with $\xi, \eta \in \mathbb{S}^{d-1}$, it holds

(3.3)
$$1 - \cos \mathsf{d}_{\mathbb{X}_0}((x,t),(y,s)) = 1 - \cos \mathsf{d}_{[-1,1]}(t,s) + ts \left(1 - \cos \mathsf{d}_{\mathbb{S}}(\xi,\eta)\right).$$

In particular,

(3.4)
$$c_1 \mathsf{d}_{\mathbb{X}_0}((x,t),(y,s)) \le \mathsf{d}_{[-1,1]}(t,s) + \sqrt{ts} \,\mathsf{d}_{\mathbb{S}}(\xi,\eta) \le c_2 \mathsf{d}_{\mathbb{X}_0}((x,t),(y,s)).$$

Proof. Using $\langle x, y \rangle$ sign $(ts) = ts \langle \xi, \eta \rangle$, we obtain

 $1 - \cos \mathsf{d}_{\mathbb{X}_0}((x,t),(y,s)) = 1 - ts \langle \xi,\eta\rangle - \sqrt{1 - t^2} \sqrt{1 - s^2}.$

Hence, (3.3) follows immediately from (3.2) and (3.1). Moreover, using (3.3), the estimate (3.4) follows from $1 - \cos \theta = 2 \sin^2 \frac{\theta}{2}$, $\frac{1}{\pi} \theta \leq \sin \frac{\theta}{2} \leq \frac{\theta}{2}$ for $0 \leq \theta \leq \pi$, and $(a+b)^2/2 \leq a^2 + b^2 \leq (a+b)^2$ for $a, b \geq 0$.

In particular, the distance on the linear segment $l_{\xi} = \{(t\xi, t) : -1 \le t \le 1\}$ for any fixed $\xi \in \mathbb{S}^{d-1}$ on the double conic surface becomes

$$\mathsf{d}_{\mathbb{X}_0}((t\xi, t), (s\xi, s)) = \mathsf{d}_{[-1,1]}(t, s).$$

We note that the line l_{ξ} passes through the origin when it passes from the upper conic surface to the lower conic surface; in fact, the line passes through $(\xi, 1)$ on the rim of the upper conic surface to $(-\xi, -1)$ on the opposite rim of the lower conic surface.

Remark 3.1. In [30, Remark 4.1], it is pointed out that the distance $\mathsf{d}_{\mathbb{V}_0}(\cdot, \cdot)$ is incompatible with the Euclidean distance at around t = 0 since, for t = s,

$$\mathsf{d}_{\mathbb{V}_0}((x,t),(y,t)) \sim \sqrt{t} \mathrm{d}_{\mathbb{S}}(\xi,\eta) \quad \text{and} \quad \|(x,t) - (y,t)\| \sim t \mathrm{d}_{\mathbb{S}}(\xi,\eta).$$

In contrast, the distance $\mathsf{d}_{\mathbb{X}_0}(\cdot, \cdot)$ on the double conic surface satisfies,

$$\mathsf{d}_{\mathbb{X}_{0}}((x,t),(y,t)) \sim |t| \mathrm{d}_{\mathbb{S}}(\xi,\eta) \sim ||(x,t) - (y,t)||$$

We will also need the following lemma in the proof of our estimates.

Lemma 3.3. Let $\varrho = 0$ and $d \ge 2$. If (x, t), (y, s) both in $\mathbb{X}_{0,+}^{d+1}$ or both in $\mathbb{X}_{0,-}^{d+1}$, then

$$|t-s| \le \mathsf{d}_{\mathbb{X}_0}((x,t),(y,s))$$
 and $|\sqrt{1-t^2} - \sqrt{1-s^2}| \le \mathsf{d}_{\mathbb{X}_0}((x,t),(y,s)).$

Proof. It suffices to consider the case $t, s \ge 0$. Let $t = \cos \theta$ and $s = \cos \phi$ with $0 \le \theta, \phi \le \pi/2$. It follows readily that $\cos \mathsf{d}_{\mathbb{X}_0}((x,t),(y,s)) \le ts + \sqrt{1-t^2}\sqrt{1-s^2} = \cos(\theta - \phi)$, which is equivalently to $|\theta - \phi| \le \mathsf{d}_{\mathbb{X}_0}((x,t),(y,s))$. Hence, the stated inequalities follows from

$$|t-s| = |\cos\theta - \cos\phi| = 2\sin\frac{\theta+\phi}{2}\sin\frac{|\theta-\phi|}{2} \le |\theta-\phi|,$$

and, similarly,

$$\left|\sqrt{1-t^2} - \sqrt{1-s^2}\right| = |\sin\theta - \sin\phi| = 2\cos\frac{\theta+\phi}{2}\sin\frac{|\theta-\phi|}{2} \le |\theta-\phi|$$

for all $t, s \in [-1, 1]$. This completes the proof.

For $\rho > 0$, the hyperbolic surface $\mathbb{X}_0^{d+1} = \mathbb{X}_{0,+}^{d+1} \cup \mathbb{X}_{0,-}^{d+1}$ consists of two disjoint parts. For all practical purpose, it is sufficient to consider the distance between points that lie in the same part. For (x, t) and (y, t) both in $\mathbb{X}_{0,+}^{d+1}$ or both in $\mathbb{X}_{0,-}^{d+1}$, we define

(3.5)
$$d^{\varrho}_{\mathbb{X}_{0}}((x,t),(y,s)) = \arccos\left(\langle x,y\rangle + \sqrt{1+\varrho^{2}-t^{2}}\sqrt{1+\varrho^{2}-s^{2}}\right)$$
$$= d_{\mathbb{X}_{0}}\left(\left(x,\sqrt{t^{2}-\varrho^{2}}\right),\left(y,\sqrt{s^{2}-\varrho^{2}}\right)\right).$$

It is easy to see that this is a distance function and, evidently, we can derive its properties as we did for the distance on the double conic surface.

3.2. A family of doubling weights. For $d \ge 2$, $\beta > -\frac{1}{2}$ and $\gamma > -\frac{1}{2}$, let $\mathsf{w}_{\beta,\gamma}^{\rho}$ be the weight function defined on the hyperbolic surface \mathbb{X}_{0}^{d+1} by

(3.6)
$$\mathsf{w}^{\varrho}_{\beta,\gamma}(t) = |t|(t^2 - \varrho^2)^{\beta - \frac{1}{2}}(\varrho^2 + 1 - t^2)^{\gamma - \frac{1}{2}}, \qquad \varrho \ge 0,$$

for $\rho \leq |t| \leq \sqrt{\rho^2 + 1}$. When $\rho = 0$, or on the double conic surface, it becomes

$$\mathsf{w}^{0}_{\beta,\gamma}(t) = |t|^{2\beta} (1-t^{2})^{\gamma-\frac{1}{2}},$$

which is integrable if $\beta > -\frac{d+1}{2}$ on \mathbb{X}_0^{d+1} . When $\beta = 0$, $w_{0,\gamma}^0$ is the classical weight function for the Gegenbauer polynomial C_n^{γ} . Evidently, the two cases are related,

$$\mathsf{w}^{\varrho}_{\beta,\gamma}(t) = |t| \mathsf{w}^{0}_{\beta-\frac{1}{2},\gamma} \left(\sqrt{t^{2} - \varrho^{2}} \right).$$

Let $d\sigma_{\varrho}$ denote the surface measure on ${}_{\varrho}X_{0}^{d+1}$. Then $d\sigma_{\varrho}(x,t) = d\omega_{\sqrt{t^{2}-\varrho^{2}}}(x)dt$, where $d\omega_{r}$ denotes the surface measure on the sphere \mathbb{S}_{r}^{d-1} of radius r. It follows then

(3.7)
$$\int_{\varrho \mathbb{X}_{0}^{d+1}} f(x,t) |t| \mathrm{d}\sigma_{\varrho}(x,t) = \int_{\varrho \le |t| \le \varrho+1} |t| \int_{\|x\| = \sqrt{t^{2} - \varrho^{2}}} f(x,t) \mathrm{d}\omega_{\sqrt{t^{2} - \varrho^{2}}}(x) \mathrm{d}t$$
$$= \int_{|s| \le 1} \int_{\|x\| = |s|} f(x,\sqrt{s^{2} + \varrho^{2}}) |s| \mathrm{d}\omega_{|s|}(x) \mathrm{d}s$$
$$= \int_{0\mathbb{X}_{0}^{d+1}} f(x,\sqrt{s^{2} + \varrho^{2}}) |s| \mathrm{d}\sigma(x,s).$$

Using this relation, it is easy to verify that the normalization constant $b^\varrho_{\beta,\gamma}$ of the weight function $w^\varrho_{\beta,\gamma}$ is given by

$$\mathbf{b}^{\varrho}_{\beta,\gamma} = \mathbf{b}_{\beta,\gamma} = \frac{\Gamma(\beta + \gamma + \frac{d+1}{2})}{\sigma_d \Gamma(\beta + \frac{d}{2}) \Gamma(\gamma + \frac{1}{2})}$$

For r > 0, (x, t) on the hyperbolic surface ${}_{\varrho} \mathbb{X}_0^{d+1}$ and $(x, t) \neq (0, 0)$, we denote the ball centered at (x, t) with radius r by

$$\mathsf{c}_{\varrho}((x,t),r) := \left\{ (y,s) \in {}_{\varrho} \mathbb{X}_0^{d+1} : \mathsf{d}_{\mathbb{X}_0}^{\varrho} \left((x,t), (y,s) \right) \leq r \right\}.$$

Lemma 3.4. For $\varrho \ge 0$ and $(x, t) \in \mathbb{X}_0^{d+1}$,

$$\mathsf{w}^{\varrho}_{\beta,\gamma}\big(\mathsf{c}_{\varrho}((x,t),r)\big) = \mathsf{w}^{0}_{\beta,\gamma}\left(\mathsf{c}_{0}\left(\left(x,\sqrt{t^{2}-\varrho^{2}}\right),r\right)\right)$$

Proof. By the definition of w(E) and (3.5),

$$\begin{split} \mathsf{w}_{\beta,\gamma}^{\varrho} \big(\mathsf{c}_{\varrho}((x,t),r) \big) &= \mathsf{b}_{\beta,\gamma} \int_{\mathsf{c}_{\varrho}((x,t),r)} \mathsf{w}_{\beta,\gamma}^{\varrho}(s) \mathrm{d}\sigma_{\varrho}(y,s) \\ &= \mathsf{b}_{\beta,\gamma} \int_{\mathsf{d}_{\mathbb{X}_{0}} \big((x,\sqrt{t^{2} - \|x\|^{2}}), (y,\sqrt{s^{2} - \|y\|^{2}}) \big) \leq r} \mathsf{w}_{\beta,\gamma}^{\varrho}(s) \mathrm{d}\sigma_{\varrho}(y,s) \\ &= \mathsf{b}_{\beta,\gamma} \int_{\mathsf{d}_{\mathbb{X}_{0}} \big((x,\sqrt{t^{2} - \|x\|^{2}}), (y,s) \big) \leq r} \mathsf{w}_{\beta,\gamma}^{0}(s) \mathrm{d}\sigma_{0}(y,s) \\ &= \mathsf{w}_{\beta,\gamma}^{0} \left(\mathsf{c}_{0} \left(\big((x,\sqrt{t^{2} - \varrho^{2}} \big), r \big) \big) \right), \end{split}$$

where we have used (3.7) in the second to last step.

Proposition 3.5. Let r > 0 and $(x,t) \in \mathbb{X}_0^{d+1}$. For $\beta > -\frac{d+1}{2}$ and $\gamma > -\frac{1}{2}$,

(3.8)
$$\mathsf{w}^{0}_{\beta,\gamma}(\mathsf{c}_{0}((x,t),r)) = \mathsf{b}^{0}_{\beta,\gamma} \int_{\mathsf{c}_{0}((x,t),r)} \mathsf{w}^{0}_{\beta,\gamma}(s) \mathrm{d}\sigma(y,s) \\ \sim r^{d} (t^{2} + r^{2})^{\beta} (1 - t^{2} + r^{2})^{\gamma}.$$

In particular, $\mathsf{w}_{\beta,\gamma}^0$ is a doubling weight on the double conic surface and the doubling index is given by $\alpha(\mathsf{w}_{\beta,\gamma}^0) = d + 2\max\{0,\beta\} + 2\max\{0,\gamma\}.$

Proof. Without loss of generality, we assume r is bounded by a small positive number, say $r \leq \frac{\pi}{12}$ and $0 \leq t \leq 1$. Since $\mathsf{w}_{\beta,\gamma}^0$ is even in t, we only need to work on $\mathbb{X}_{0,+}^{d+1} = \mathbb{V}_0^{d+1}$. Let $x = t\xi$ and $y = s\eta$ for $\xi, \eta \in \mathbb{S}^{d-1}$. By the inequality (3.3), from $\mathsf{d}_{\mathbb{X}_0}((x,t),(y,s)) \leq r$ we obtain $\mathsf{d}_{[-1,1]}(t,s) \leq r$; moreover, denote $\tau_r(t,s) = (\cos r - \sqrt{1-t^2}\sqrt{1-s^2})/(ts)$ and $\theta_r(t,s) = \arccos \tau_r(t,s)$, we also have $\mathsf{d}_{\mathbb{S}}(\xi,\eta) \leq \frac{1}{2} \arccos \tau_r(t,s) = \frac{1}{2}\theta_r(t,s)$. Then it is easy to see, since $\mathsf{d}\sigma(y,s) = s^{d-1}\mathsf{d}\sigma_{\mathbb{S}}(\eta)\mathsf{d}s$, that

$$\mathsf{w}^0_{\beta,\gamma}\big(\mathsf{c}((x,t),r)\big) = \int_{\mathsf{d}_{[-1,1]}(t,s) \leq r} s^{d-1} \int_{\mathsf{d}_{\mathbb{S}}(\xi,\eta) \leq \frac{1}{2}\theta_r(t,s)} \mathsf{w}^0_{\beta,\gamma}(s\eta,s) \mathrm{d}\sigma_{\mathbb{S}}(\eta) \mathrm{d}s.$$

By symmetry, we can choose $\xi = (1, 0, \dots, 0)$ and use the identity (cf. [8, (A.5.1)])

(3.9)
$$\int_{\mathbb{S}^{d-1}} g(\langle \xi, \eta \rangle) d\sigma(\eta) = \omega_{d-1} \int_0^\pi g(\cos \theta) (\sin \theta)^{d-2} \mathrm{d}\theta$$

with ω_{d-1} being the surface are of \mathbb{S}^{d-2} , to obtain that

$$\mathsf{w}_{\beta,\gamma}^{0}(\mathsf{c}((x,t),r)) = \omega_{d-1} \int_{\mathsf{d}_{[0,1]}(t,s) \le r} s^{d-1} \mathsf{w}_{\beta,\gamma}^{0}(s) \int_{0}^{\frac{1}{2}\theta_{r}(t,s)} (\sin\theta)^{d-2} \mathrm{d}\theta \mathrm{d}s.$$

Since $\theta \sim \sin \theta \sim \sqrt{1 - \cos \theta}$, it follows then

(3.10)
$$\mathsf{w}^{0}_{\beta,\gamma}(\mathsf{c}_{0}((x,t),r)) \sim \int_{\mathsf{d}_{[-1,1]}(t,s) \leq r} s^{d-1} \mathsf{w}^{0}_{\beta,\gamma}(s) (1-\tau_{r}(t,s))^{\frac{d-1}{2}} \mathrm{d}s.$$

We now need to consider three cases. If 3r < t < 1 - 3r, then we can use Lemma 3.3 to conclude that $s^2 \sim t^2 + r^2$ and $1 - s^2 \sim 1 - t^2 + r^2$, so that

$$\begin{split} \mathsf{w}^{0}_{\beta,\gamma}\big(\mathsf{c}_{0}((x,t),r)\big) &\sim (t^{2}+r^{2})^{\beta}(1-t^{2}+r^{2})^{\gamma} \\ &\times \int_{\mathsf{d}_{[-1,1]}(t,s) \leq r} \big(\cos(\mathsf{d}_{[-1,1]}(t,s)) - \cos r\big)^{\frac{d-1}{2}} \frac{\mathrm{d}s}{\sqrt{1-s^{2}}}. \end{split}$$

Setting $t = \cos \theta$ and $s = \cos \phi$ so that $\mathsf{d}_{[-1,1]}(t,s) = |\theta - \phi|$, then the last integral is easily seen to be

$$\int_{|\theta-\phi|\leq r} \left(\cos(\theta-\phi)-\cos r\right)^{\frac{d-1}{2}} \mathrm{d}\phi = c \int_{|\zeta|\leq r} \left(\sin\frac{\zeta-r}{2}\sin\frac{\zeta+r}{2}\right)^{\frac{d-1}{2}} \mathrm{d}\zeta \sim r^d.$$

This completes the proof of the first case. If $t \leq 3r$, then $|t-s| \leq d_{[-1,1]}(t,s) \leq r$ so that $s \leq 4r$. Evidently $1-s^2 \sim 1-t^2 \sim 1$ in this case. Furthermore, let $t = \sin \theta$ and $s = \sin \phi$; then $|\theta - \phi| = d_{[-1,1]}(t,s) \leq r$, which is easily seen to be equivalent to $|\tau_r(t,s)| \leq 1$. Hence, using $1 - \tau_r(t,s) \leq 2$ and $s \leq 4r$, we obtain by (3.10) and $\sin \phi \sim \phi$,

$$\mathsf{w}_{\beta,\gamma}(\mathsf{c}((x,t),r)) \le c \int_{\mathsf{d}_{[-1,1]}(t,s) \le r} s^{d+2\beta-1} \mathrm{d}s \sim r^{2\beta+d},$$

which proves the upper bound in (3.8). For the lower bound, we consider a subset of c((x,t),r) with $d_{[-1,1]}(t,s) \leq r/2$. Using the upper bound of s and t, we then deduce

$$1 - \tau_r(t,s) = \frac{\cos d_{[-1,1]}(t,s) - \cos r}{ts} \ge \frac{\cos \frac{r}{2} - \cos r}{12r^2} \ge \frac{1}{8\pi^2}$$

where in the last step we have used $\sin \theta \geq \frac{2}{\pi} \theta$, which shows that

$$\mathsf{w}_{\beta,\gamma}(\mathsf{c}((x,t),r)) \geq c \int_{\mathsf{d}_{[-1,1]}(t,s) \leq r} s^{d+2\beta-1} \mathrm{d}s \sim r^{2\beta+d}.$$

Finally, if $t \ge 1-3r$, then we have $s \sim t \sim 1$ for $(y,s) \in c((x,t),r)$. Since $\mathsf{d}_{[-1,1]}(t,s) = \mathsf{d}_{[-1,1]}(\sqrt{1-t^2},\sqrt{1-s^2})$, changing variable $s \mapsto \sqrt{1-s^2}$ in (3.10), we see that this case can be reduced to that of the second case. This completes the proof of (3.8).

For $\rho > 0$ and $\beta, \gamma > -\frac{1}{2}$, the weight function $w^{\rho}_{\beta,\gamma}$ is a doubling weight on the hyperbolic surface ${}_{\rho}X^{d+1}$ by Lemma 3.4.

Corollary 3.6. For $\varrho \geq 0$, $d \geq 2$, and $\beta, \gamma > -\frac{1}{2}$, the space $(\mathbb{X}_0^{d+1}, \mathsf{w}_{\beta,\gamma}^{\varrho}, \mathsf{d}_{\mathbb{X}_0}^{\varrho})$ is a homogeneous space. If $\varrho = 0$, the restriction on β can be relaxed to $\beta > -\frac{d+1}{2}$.

When $\rho = 0$, $\beta = 0$ and $\gamma = \frac{1}{2}$, the relation (3.8) is for the Lebesgue measure σ_0 on the double conic surface; in particular, if $\beta = 0$ and $\gamma = 0$, then $w_{0,0}^0(t) = (1 - t^2)^{-\frac{1}{2}}$ is the Chebyshev weight and $w_{0,0}^0(c_0((x,t),r)) \sim r^d$. The apex point t = 0 does not appear as a boundary point of \mathbb{X}_0^{d+1} .

For convenience, we will introduce the function $w_{\beta,\gamma}(n;t)$ defined by

(3.11)
$$\mathsf{w}^{\varrho}_{\beta,\gamma}(n;t) = n^{-d} \mathsf{w}^{\varrho}_{\beta,\gamma} \big(\mathbf{c}_{\varrho}((x,t), n^{-1}) \big) = \big(t^2 - \varrho^2 + n^{-2} \big)^{\beta} \big(1 - t^2 + \varrho^2 + n^{-2} \big)^{\gamma}$$

on the hyperbolic surface \mathbb{X}_0^{a+1} and use it in latter sections.

3.3. Orthogonal polynomials on hyperbolic surfaces. For $\gamma > -\frac{1}{2}$, we define the inner product on the hyperbolic surface by

$$\langle f,g\rangle_{\mathsf{w}_{\beta,\gamma}} = \mathsf{b}_{\beta,\gamma} \int_{\mathbb{X}_0^{d+1}} f(x,t) g(x,t) \mathsf{w}_{\beta,\gamma}^{\varrho}(t) \mathrm{d}\sigma_{\varrho}(x,t),$$

where $\beta > -\frac{1}{2}$ if $\varrho > 0$ and $\beta > -\frac{d+1}{2}$ if $\varrho = 0$. The orthogonal polynomials with respect to this inner product are studied in [29]. Let $\mathcal{V}_n(\mathbb{X}_0^{d+1}, \mathsf{w}_{\beta,\gamma}^{\varrho})$ be the space of these polynomials with respect to this inner product, which has the same dimension as the space of spherical harmonics \mathcal{H}_n^{d+1} . Because the weight function $\mathsf{w}_{\beta,\gamma}^{\varrho}$ is even in t, this space can be factored as

$$\mathcal{V}_n\left(\mathbb{X}_0^{d+1},\mathsf{w}_{\beta,\gamma}^{\varrho}\right) = \mathcal{V}_n^E\left(\mathbb{X}_0^{d+1},\mathsf{w}_{\beta,\gamma}^{\varrho}\right) \bigoplus \mathcal{V}_n^O\left(\mathbb{X}_0^{d+1},\mathsf{w}_{\beta,\gamma}^{\varrho}\right),$$

where the subspace $\mathcal{V}_n^E(\mathbb{X}_0^{d+1}, \mathsf{w}_{\beta,\gamma}^{\varrho})$ consists of orthogonal polynomials that are even in t variable, and the subspace $\mathcal{V}_n^O(\mathbb{X}_0^{d+1}, \mathsf{w}_{\beta,\gamma}^{\varrho})$ consists of orthogonal polynomials that are odd in t variable. Moreover,

$$\dim \mathcal{V}_n^E \left(\mathbb{X}_0^{d+1}, \mathsf{w}_{\beta, \gamma}^{\varrho} \right) = \binom{n+d-1}{n}, \qquad \dim \mathcal{V}_n^O \left(\mathbb{X}_0^{d+1}, \mathsf{w}_{\beta, \gamma}^{\varrho} \right) = \binom{n+d-2}{n-1}.$$

It turns out that an orthogonal basis can be given in terms of the spherical harmonics and the Jacobi polynomials for the subspace $\mathcal{V}_n^E(\mathbb{X}_0^{d+1}, \mathsf{w}_{\beta,\gamma}^{\varrho})$ for all $\varrho \geq 0$, but for the

subspace $\mathcal{V}_n^O(\mathbb{X}_0^{d+1}, \mathsf{w}_{\beta,\gamma}^{\varrho})$ only when $\varrho = 0$. For example, let $\{Y_\ell^m : 1 \leq \ell \leq \dim \mathcal{H}_m^d\}$ denote an orthonormal basis of \mathcal{H}_m^d . Then the polynomials

(3.12)
$$\mathsf{C}_{m,\ell}^n(x,t) = P_k^{(\gamma - \frac{1}{2}, n-2k+\beta + \frac{d-2}{2})} (2t^2 - 2\varrho^2 - 1) Y_\ell^{n-2k}(x),$$

where $1 \leq \ell \leq \dim \mathcal{H}_{n-2k}^d$ and $0 \leq k \leq n/2$, form an orthogonal basis of $\mathcal{V}_n^E(\mathbb{X}_0^{d+1}, \mathsf{w}_{\beta,\gamma}^\varrho)$. Since we will not work directly with explicit bases, we refer to [29] for further information, where these polynomials are called, when $\beta = 0$, the Gegenbauer polynomials on the hyperbolic surface or on the double conic surface when $\varrho = 0$.

Let $\mathsf{P}_n^E(\mathsf{w}_{\beta,\gamma}^\varrho;\cdot,\cdot)$ be the reproducing kernel of $\mathcal{V}_n^E(\mathbb{X}_0^{d+1},\mathsf{w}_{\beta,\gamma}^\varrho)$, which can be written in terms of the basis (3.12) as

$$\mathsf{P}_n^E\left(\mathsf{w}^{\varrho}_{\beta,\gamma};(x,t),(y,s)\right) = \sum_{m=0}^n \sum_{\ell=1}^{\dim \mathcal{H}_{n-2m}^d} \frac{\mathsf{C}_{m,\ell}^n(x,t)\mathsf{C}_{m,\ell}^n(y,s)}{\langle\mathsf{C}_{m,\ell}^n,\mathsf{C}_{m,\ell}^n\rangle_{\mathsf{w}_{\beta,\gamma}}}.$$

Let $\operatorname{proj}_{n}^{E}\left(\mathsf{w}_{\beta,\gamma}^{\varrho}\right): L^{2}\left(\mathbb{X}_{0}^{d+1}, \mathsf{w}_{\beta,\gamma}^{\varrho}\right) \mapsto \mathcal{V}_{n}^{E}\left(\mathbb{X}_{0}^{d+1}, \mathsf{w}_{\beta,\gamma}^{\varrho}\right)$ be the orthogonal projection operator. Then it can be written as

$$\operatorname{proj}_{n}^{E}(\mathsf{w}_{\beta,\gamma}^{\varrho};f) = \mathsf{b}_{\beta,\gamma} \int_{\mathbb{X}_{0}^{d+1}} f(y) \mathsf{P}_{n}^{E}(\mathsf{w}_{\beta,\gamma}^{\varrho};\cdot,(y,s)) \mathsf{w}_{\beta,\gamma}^{\varrho}(s) \mathrm{d}y \mathrm{d}s.$$

If f is a function that is even in the variable t on \mathbb{X}_0^{d+1} , then its orthogonal projection on $\mathcal{V}_n^O(\mathbb{X}_0^{d+1}, \mathsf{w}_{\beta,\gamma}^{\varrho})$ becomes zero, so that its Fourier orthogonal expansion is given by

(3.13)
$$f = \sum_{n=0}^{\infty} \operatorname{proj}_{n}^{E}(\mathsf{w}_{\beta,\gamma}^{\varrho}; f)$$

Hence, the kernel $\mathsf{P}_n^E(\mathsf{w}^{\varrho}_{\beta,\gamma};\cdot,\cdot)$ is meaningful for studying the Fourier orthogonal expansions on the hyperboloid.

If $\rho = 0$, then the upper part $\mathbb{X}_{0,+}^{d+1} = \mathbb{V}_0^{d+1}$ is the upper conic surface. The function f(x,t) that is even in t variable can be regarded as defined on \mathbb{V}_0^{d+1} or as the even extension in t variable of a function defined on the upper conic surface. Consequently, the Fourier expansion (3.13) works for the function f defined on the upper conic surface $\mathbb{X}_{0,+}^{d+1}$ when $\rho = 0$. The latter, however, is different from the Fourier expansions in the Jacobi polynomials on the conic surface \mathbb{V}_0^{d+1} discussed in [30].

The case $\beta = 0$ is the most interesting since the orthogonal polynomials for $w_{0,\gamma}$ enjoy two characteristic properties. The first one is the spectral operator that has orthogonal polynomials as eigenfunctions.

Theorem 3.7. Let $\varrho > 0$ and $\gamma > -\frac{1}{2}$. Then for $x = t\xi$, $\xi \in \mathbb{S}^{d-1}$, define the differential operator

$$\begin{aligned} \Delta_{0,\gamma}^{\varrho} &= (1+\varrho^2 - t^2) \left(1 - \frac{\varrho^2}{t^2}\right) \partial_t^2 \\ &+ \left((1+\varrho^2 - t^2) \frac{\varrho^2}{t^2} - (2\gamma + d)(t^2 - \varrho^2)\right) \frac{1}{t} \partial_t + \frac{d-1}{t} \partial_t + \frac{1}{t^2 - \varrho^2} \Delta_0^{(\xi)}. \end{aligned}$$

Then the polynomials in $\mathcal{V}_n^E(\mathbb{V}_0^{d+1}, w_{0,\gamma}^{\varrho})$ are eigenfunctions of $\Delta_{0,\gamma}^{\varrho}$,

(3.14)
$$\Delta_{0,\gamma}^{\varrho} u = -n(n+2\gamma+d-1)u, \quad \forall u \in \mathcal{V}_n^E(\mathbb{V}_0^{d+1}, w_{0,\gamma}^{\varrho})$$

The second one is the addition formula for the reproducing kernel $\mathsf{P}_n^E(\mathsf{w}_{\beta,\gamma}^\varrho;\cdot,\cdot)$, which is of the simplest form when $\beta = 0$.

Theorem 3.8. Let $d \ge 2$ and $\varrho \ge 0$. Then

(a) For $\beta, \gamma > -\frac{1}{2}$,

(3.15)
$$\mathsf{P}_{n}^{E}\left(\mathsf{w}_{\beta,\gamma}^{\varrho};(x,t),(y,s)\right) = \mathsf{P}_{n}^{E}\left(\mathsf{w}_{\beta,\gamma}^{0};\left(x,\sqrt{t^{2}-\varrho^{2}}\right),\left(y,\sqrt{s^{2}-\varrho^{2}}\right)\right).$$

(b) For $\rho = 0$, $\beta = 0$ and $\gamma \ge 0$,

(3.16)
$$\mathsf{P}_{n}^{E}\left(\mathsf{w}_{0,\gamma}^{0};(x,t),(y,s)\right) = c_{\gamma} \int_{-1}^{1} Z_{n}^{\gamma + \frac{d-1}{2}} \left(\zeta(x,t,y,s;v)\right) (1-v^{2})^{\gamma - 1} \mathrm{d}v,$$

where $c_{\gamma} = c_{\gamma-1,\gamma-1}$ and

$$\zeta(x, t, y, s; v) = \langle x, y \rangle \operatorname{sgin}(ts) + v\sqrt{1 - t^2}\sqrt{1 - s^2},$$

and the case $\gamma = 0$ holds under the limit (2.21).

The closed form formula (3.16) is essential for studying highly localized kernels.

3.4. Highly localized kernels. Let \hat{a} be a cut-off function. For $(x, t), (y, s) \in \mathbb{X}_0^{d+1}$, define the kernel $\mathsf{L}_n^E(\mathsf{w}_{0,\gamma}^{\varrho})$ by

$$\mathsf{L}_n^E(\mathsf{w}_{0,\gamma}^\varrho;(x,t),(y,s)) = \sum_{j=0}^\infty \widehat{a}\left(\frac{j}{n}\right) \mathsf{P}_j^E\big(\mathsf{w}_{0,\gamma}^\varrho;(x,t),(y,s)\big).$$

The kernel uses only orthogonal polynomials that are even in t and s variable, so that it is even in both t and s variables. We show that this kernel is highly localized when (x, t) and (y, s) are either both in $\mathbb{X}_{0,+}^{d+1}$ or both in $\mathbb{X}_{0,-}^{d+1}$. For $\gamma \geq 0$, recall by (3.11),

$$\mathsf{w}_{0,\gamma}^{\varrho}(n;t) := \left(1 + \varrho^2 - t^2 + n^{-2}\right)^{\gamma}.$$

Theorem 3.9. Let $d \ge 2$ and $\gamma \ge 0$. Let \hat{a} be an admissible cutoff function. Then, for any $\kappa > 0$, either (x, t), (y, s) both in $\mathbb{X}_{0,+}^{d+1}$ or both in $\mathbb{X}_{0,-}^{d+1}$,

$$\left|\mathsf{L}_{n}^{E}(\mathsf{w}_{0,\gamma}^{\varrho};(x,t),(y,s))\right| \leq \frac{c_{\kappa}n^{d}}{\sqrt{\mathsf{w}_{0,\gamma}^{\varrho}(n;t)}\sqrt{\mathsf{w}_{0,\gamma}^{\varrho}(n;s)}} \big(1 + n\mathsf{d}_{\mathbb{X}_{0}}^{\varrho}((x,t),(y,s))\big)^{-\kappa},$$

where we assume t and s have the same sign when $\rho > 0$.

Proof. By (3.15), it is sufficient to consider the case $\rho = 0$. The proof follows the similar procedure as in the case of the conic surface, so we shall be brief. By (3.16) we can write $L_n^E(\mathsf{w}_{0,\gamma}^0)$ in terms of the kernel for the Jacobi polynomials. Let $\lambda = \gamma + \frac{d-1}{2}$. Then

$$\mathsf{L}_{n}^{E}(\mathsf{w}_{0,\gamma}^{0};(x,t),(y,s)) = c_{\gamma} \int_{-1}^{1} L_{n}^{(\lambda - \frac{1}{2},\lambda - \frac{1}{2})} \big(\zeta(x,t,y,s;v)\big) (1 - v^{2})^{\gamma - 1} \mathrm{d}v.$$

Applying (2.12) with m = 0 and $\alpha = \beta = \lambda - 1/2$, we then obtain

$$\left|\mathsf{L}_{n}^{E}(\mathsf{w}_{0,\gamma}^{0};(x,t),(y,s))\right| \leq cn^{2\lambda+1} \int_{-1}^{1} \frac{1}{\left(1 + n\sqrt{1 - \zeta(x,t,y,s;v)^{2}}\right)^{\kappa+3\gamma+1}} (1 - v^{2})^{\gamma-1} \mathrm{d}v.$$

By the definition of $\mathsf{d}_{\mathbb{X}_0}(\cdot, \cdot)$, we have

(3.17)
$$1 - \zeta(x, t, y, s; t) = 1 - \cos \mathsf{d}_{\mathbb{X}_0}((x, t), (y, s)) + (1 - v)\sqrt{1 - t^2}\sqrt{1 - s^2}.$$

In particular, $1 - \zeta(x, t, y, s; t)$ is bounded below by either the first term or the second term in the right-hand side of (3.17), which leads to, in particular, the estimate

$$\begin{aligned} \left| \mathsf{L}_{n}^{E}(\mathsf{w}_{0,\gamma}^{0};(x,t),(y,s)) \right| &\leq c n^{2\lambda+1} \frac{1}{\left(1 + n \mathsf{d}_{\mathbb{X}_{0}}((x,t),(y,s)) \right)^{\kappa+\gamma}} \\ &\times c_{\gamma} \int_{0}^{1} \frac{(1 - v^{2})^{\gamma-1}}{\left(1 + n\sqrt{(1 - v)\sqrt{1 - t^{2}}\sqrt{1 - s^{2}}} \right)^{2\gamma+1}} \mathrm{d}v \end{aligned}$$

where we have used $1 - \cos \theta \sim \theta^2$ and the symmetry of the integral. The last integral is evidently bounded by 1 and it can be estimated by using the inequality [8, (13.5.8)]

(3.18)
$$\int_0^1 \frac{(1-t)^{a-1} dt}{(1+n\sqrt{B}+A(1-t))^b} \le c \frac{n^{-2a}}{A^a(1+n\sqrt{B})^{b-2a-1}},$$

which holds for A > 0, $B \ge 0$, a > 0 and $b \ge 2a + 1$, which leads to the estimate

$$c_{\gamma} \int_{0}^{1} \frac{(1-v^{2})^{\gamma-1}}{\left(1+n\sqrt{(1-v)\sqrt{1-t^{2}}\sqrt{1-s^{2}}}\right)^{2\gamma+1}} \mathrm{d}v \leq c \frac{n^{-2\gamma}}{\left(\sqrt{1-t^{2}}\sqrt{1-s^{2}}+n^{-1}\right)^{\gamma}}$$
$$\leq c \frac{n^{-2\gamma}}{\sqrt{\mathsf{w}_{0,\gamma}^{0}(n;t)}} (1+n\mathsf{d}_{\mathbb{X}_{0}}((x,t),(y,s)))^{\gamma},$$

where the second inequality follows from the elementary identity [8, (11.5.13)]

(3.19)
$$(a+n^{-1})(b+n^{-1}) \le 3(ab+n^{-2})(1+n|b-a|)$$

with $a = \sqrt{1 - t^2}$ and $b = \sqrt{1 - s^2}$ and Lemma 3.3. Putting the last two displayed inequalities together completes the proof.

This theorem establishes the Assertion 1 on \mathbb{X}_0^{d+1} . We now turn to Assertion 2.

Theorem 3.10. Let $d \ge 2$ and $\gamma \ge 0$. For either $(x_i, t_i), (y, s)$ all in $\mathbb{X}_{0,+}^{d+1}$ or all in $\mathbb{X}_{0,-}^{d+1}$, and $(x_1, t_1) \in \mathsf{c}_{\varrho}((x_2, t_2), c^*n^{-1})$ with c^* small and for $\kappa > 0$,

$$(3.20) \qquad \left| \mathsf{L}_{n}^{E}(\mathsf{w}_{0,\gamma}^{\varrho};(x_{1},t_{1}),(y,s)) - \mathsf{L}_{n}^{E}(\mathsf{w}_{0,\gamma}^{\varrho};(x_{2},t_{2}),(y,s)) \right| \\ \leq \frac{c_{\kappa}n^{d+1}\mathsf{d}_{\mathbb{X}_{0}}^{\varrho}((x_{1},t_{1}),(x_{2},t_{2}))}{\sqrt{\mathsf{w}_{0,\gamma}^{\varrho}(n;t_{2})}\sqrt{\mathsf{w}_{0,\gamma}^{\varrho}(n;s)} \left(1 + n\mathsf{d}_{\mathbb{X}_{0}}^{\varrho}((x_{2},t_{2}),(y,s))\right)^{\kappa}}.$$

Proof. Again, it is sufficient to consider $\rho = 0$. Denote the left-hand side of (3.20) by K. Let $\partial L(u) = L'(u)$. Using the integral expression of $\mathsf{L}_n^E(\mathsf{w}_{-1,\gamma})$, we obtain

$$K \le 2 \int_{-1}^{1} \left\| \partial L_n^{\lambda - \frac{1}{2}, \lambda - \frac{1}{2}} \right\|_{L^{\infty}(I_v)} |\zeta_1(v) - \zeta_2(v)| (1 - v^2)^{\gamma - 1} \mathrm{d}v,$$

where $\zeta_i(v) = \zeta(x_i, t_i, y, s; v)$, and I_v is the interval with end points $\zeta_1(v)$ and $\zeta_2(v)$. We claim that

(3.21)
$$|\zeta_1(v) - \zeta_2(v)| \le \mathsf{d}_{\mathbb{X}_0} \big((x_1, t_1), (x_2, t_2) \big) \big[\Sigma_1 + \Sigma_2(v) \big],$$

where

$$\Sigma_1 = \mathsf{d}_{\mathbb{X}_0}((x_2, t_2), (y, s)) + \mathsf{d}_{\mathbb{X}_0}((x_1, t_1), (x_2, t_2)),$$

$$\Sigma_2(v) = (1 - v)\sqrt{1 - s^2}.$$

To see this, we first use (3.17) to write that

$$\begin{aligned} \zeta_1(v) - \zeta_2(v) &= \cos \mathsf{d}_{\mathbb{X}_0}((x_1, t_1), (y, s)) - \cos \mathsf{d}_{\mathbb{X}_0}((x_2, t_2), (y, s)) \\ &+ (1 - v) \left(\sqrt{1 - t_1^2} - \sqrt{1 - t_2^2}\right) \sqrt{1 - s^2}. \end{aligned}$$

Denote temporarily $\alpha_i = \mathsf{d}_{\mathbb{X}_0}((x_1, t_1), (y, s))$ for i = 1, 2. The identity

$$\cos \alpha_1 - \cos \alpha_2 = 2 \sin \frac{\alpha_1 - \alpha_2}{2} \left(2 \sin \frac{\alpha_1}{2} \cos \frac{\alpha_2}{2} - \sin \frac{\alpha_2 - \alpha_1}{2} \right),$$

implies that $|\cos \alpha_1 - \cos \alpha_2| \le |\alpha_1 - \alpha_2| (|\alpha_1| + \frac{1}{2}|\alpha_1 - \alpha_2|)$, which leads to the estimate for the Σ_1 term by the triangle inequality of $\mathsf{d}_{\mathbb{X}_0}$ and Lemma 3.3. The estimate for the Σ_2 terms is trivial. Hence, (3.21) holds as claimed. Since $\max_{r \in I_v} (1 + n\sqrt{1 - r^2})^{-\kappa}$ is attained at one of the end points of the interval,

it follows from (2.12) with m = 1 and $\lambda = \gamma + \frac{d-1}{2}$ that

$$\begin{aligned} \left| \mathsf{L}_{n}^{E}(\mathsf{w}_{0,\gamma}^{0};(x_{1},t_{1}),(y,s)) - \mathsf{L}_{n}^{E}(\mathsf{w}_{-0,\gamma}^{0};(x_{2},t_{2}),(y,s)) \right| \\ &\leq c \,\mathsf{d}_{\mathbb{X}_{0}}\left((x_{1},t_{1}),(x_{2},t_{2}) \right) \int_{-1}^{1} \left[\frac{n^{2\lambda+3}}{\left(1 + n\sqrt{1-\zeta_{1}(v)^{2}} \right)^{\kappa(\gamma)}} + \frac{n^{2\lambda+3}}{\left(1 + n\sqrt{1-\zeta_{2}(v)^{2}} \right)^{\kappa(\gamma)}} \right] \\ &\times \left(\Sigma_{1} + \Sigma_{2}(v) \right) (1-v^{2})^{\gamma-1} \mathrm{d}v, \end{aligned}$$

where we choose $\kappa(\gamma) = \kappa + 3\gamma + 2$. Since $(x_1, t_1) \in \mathsf{c}((x_2, t_2), c^*n^{-1}), \Sigma_1$ is bounded by $\Sigma_1 \leq cn^{-1} (1 + n \mathsf{d}_{\mathbb{V}_0}((x_i, t_i), (y, s)))$. Hence, using the two lower bound of $1 - \zeta_i(v)$ given by the right-hand side of (3.17), we obtain

$$\begin{split} &\int_{-1}^{1} \frac{n^{2\lambda+3}}{\left(1+n\sqrt{1-\zeta_{i}(v)^{2}}\right)^{\kappa(\gamma)}} \Sigma_{1}(1-v^{2})^{\gamma-1} \mathrm{d}v \\ &\leq \frac{c \, n^{2\lambda+2}}{\left(1+n \mathsf{d}_{\mathbb{V}_{0}}((x_{i},t_{i}),(y,s))\right)^{\kappa+\gamma}} \int_{0}^{1} \frac{(1-v^{2})^{\gamma-1}}{\left(1+n\sqrt{(1-v)\sqrt{1-t_{i}^{2}}\sqrt{1-s^{2}}}\right)^{2\gamma+1}} \mathrm{d}v \\ &\leq \frac{c_{\kappa}n^{d+1}}{\sqrt{\mathsf{w}_{0,\gamma}^{0}(n;t_{i})}\sqrt{\mathsf{w}_{0,\gamma}^{0}(n;s)} \left(1+n\mathsf{d}_{\mathbb{X}_{0}}((x_{i},t_{i}),(y,s))\right)^{\kappa}}, \end{split}$$

where the second step follows from the estimate of the last integral in the proof of Theorem 3.9. Since $w_{0,\gamma}^0(n,t_1) \sim w_{0,\gamma}(n,t_2)$ and $d_{\mathbb{X}_0}((x_1,t_1),(y,s)) + n^{-1} \sim$ $d_{\mathbb{X}_0}((x_2,t_2),(y,s)) + n^{-1}$ by Lemma 3.3, we can replace (x_1,t_1) in the right-hand side by (x_2, t_2) . This shows that the integral containing Σ_1 has the desired estimate.

For the integral that contains $\Sigma_2(v) = (1-v)\sqrt{1-s^2}$, the factor 1-v increases the power of the weight to $(1 - v_1)^{\gamma}$, so that we can follow the estimate for the integral with Σ_1 but using $(1-v)^{\gamma}$, which leads to

$$\begin{split} &\int_{-1}^{1} \frac{n^{2\lambda+3}}{\left(1+n\sqrt{1-\zeta_{i}(v)^{2}}\right)^{\kappa(\gamma+\frac{1}{2})}} \Sigma_{2}(v_{1})(1-v^{2})^{\gamma-1} \mathrm{d}v \\ &\leq c \frac{n^{d+1}n^{-1}\sqrt{1-s^{2}}}{\sqrt{\mathsf{w}_{0,\gamma+\frac{1}{2}}^{0}(n;s)}\sqrt{\mathsf{w}_{0,\gamma+\frac{1}{2}}^{0}(n;t_{i})} \left(1+n\mathsf{d}_{\mathbb{V}_{0}}((y,s),(x_{i},t_{i}))\right)^{\kappa}} \\ &\leq c \frac{n^{d+1}}{\sqrt{\mathsf{w}_{\gamma,d}^{0}(n;s)}\sqrt{\mathsf{w}_{\gamma,d}^{0}(n;t_{2})} \left(1+n\mathsf{d}_{\mathbb{V}_{0}}((y,s),(x_{i},t_{i}))\right)^{\kappa}}, \end{split}$$

where the last step uses the inequality $n^{-1}\sqrt{1-s^2} \leq (\sqrt{1-t_i^2}+n^{-1})(\sqrt{1-s^2}+n^{-1})$. This takes care of the integral with $\Sigma_2(v)$ and completes the proof. \Box

The case of p = 1 of the following lemma establishes Assertion 3 for $\mathsf{w}^{\varrho}_{\beta,\gamma}$.

Lemma 3.11. Let $d \ge 2$, $\beta > -\frac{1}{2}$ and $\gamma > -\frac{1}{2}$. For $0 , assume <math>\kappa > \frac{2d+2}{p} + 2(\beta + \gamma)|\frac{1}{p} - \frac{1}{2}|$. Then for $(x, t) \in \mathbb{X}_0^{d+1}$,

$$(3.22) \quad \int_{\mathbb{X}_0^{d+1}} \frac{\mathsf{w}_{\beta,\gamma}^{\varrho}(s)}{\mathsf{w}_{\beta,\gamma}^{\varrho}(n;s)^{\frac{p}{2}} \left(1 + n\mathsf{d}_{\mathbb{X}_0}^{\varrho}((x,t),(y,s))\right)^{\kappa p}} \mathrm{d}\sigma_{\varrho}(y,s) \le cn^{-d} \mathsf{w}_{\beta,\gamma}^{\varrho}(n;t)^{1-\frac{p}{2}}.$$

Proof. We again only need to consider $\rho = 0$. Let J_p denote the left-hand side of (3.22). As shown in the proof of [30, Lemma 2.4], it is sufficient to prove the case p = 2. Furthermore, the integral over \mathbb{X}_0^{d+1} is a sum of two integrals over $\mathbb{X}_{0,+}^{d+1}$ and $\mathbb{X}_{0,-}^{d+1}$, respectively. We only need to estimate one of them. Denote the integral over $\mathbb{X}_{0,+}^{d+1} = \mathbb{V}_0^{d+1}$ by $J_{2,+}$. Then

$$J_{2,+} = \int_{\mathbb{V}_0^{d+1}} \frac{\mathsf{w}_{\beta,\gamma}^0(s)}{\mathsf{w}_{\beta,\gamma}^0(n;s)(1+n\mathsf{d}_{\mathbb{X}_0}((x,t),(y,s)))^{2\kappa}} \mathrm{d}\sigma_0(y,s).$$

Let $x = t\xi$ and $y = s\eta$. Using (3.9), we obtain

$$J_{2,+} \leq c \int_0^1 \int_{-1}^1 \frac{s^{d-1} \mathsf{w}^0_{\beta,\gamma}(s) (1-u^2)^{\frac{d-3}{2}}}{\mathsf{w}^0_{\beta,\gamma}(n;s) \left(1+n\sqrt{1-tsu-\sqrt{1-t^2}\sqrt{1-s^2}}\right)^{2\kappa}} \mathrm{d}u \mathrm{d}s.$$

Hence, using $\mathsf{w}^0_{\beta,\gamma}(s) \leq c \mathsf{w}^0_{\beta,\gamma}(n;s)(1-s^2+n^{-2})^{-\frac{1}{2}}$, making another change of variable $u \mapsto v/s$ and simplifying, it follows that

$$J_{2,+} \leq c \int_0^1 \int_{-s}^s \frac{s \left(s^2 - v^2\right)^{\frac{d-3}{2}}}{\left(1 - s^2 + n^{-2}\right)^{\frac{1}{2}} \left(1 + n\sqrt{1 - tv} - \sqrt{1 - t^2}\sqrt{1 - s^2}\right)^{2\kappa}} \mathrm{d}v \mathrm{d}s$$
$$\leq c n^d \int_{-1}^1 \int_{|v|}^1 \frac{\left(s^2 - v^2\right)^{\frac{d-3}{2}}}{\left(1 + n\sqrt{1 - tv} - \sqrt{1 - t^2}\sqrt{1 - s^2}\right)^{2\kappa}} \mathrm{d}s \mathrm{d}v,$$

where we changed the order of integration in the second step. A further change of variable $s \mapsto \sqrt{1 - \|u\|^2}$ shows then

$$J_{2,+} \le c \int_{-1}^{1} \int_{0}^{\sqrt{1-v^2}} \frac{(1-u^2-v^2)^{\frac{d-3}{2}}}{\left(1+n\sqrt{1-tv}-\sqrt{1-t^2}u\right)^{2\kappa}} \mathrm{d}u \mathrm{d}v,$$

which is an integral over the right half $\{(u, v) \in \mathbb{B}^2 : v \ge 0\}$ of the unit disk \mathbb{B}^2 . Setting $z = tv + \sqrt{1-t^2} u$ and $w = -\sqrt{1-t^2}v + tu$ in the integral, which is an orthogonal transformation, and enlarging the integral domain while taking into account that $z \ge 0$, it follows that

$$J_{2,+} \leq c \int_0^1 \frac{1}{\left(1 + n\sqrt{1-z}\right)^{2\kappa}} \int_{-\sqrt{1-z^2}}^{\sqrt{1-z^2}} (1 - z^2 - w^2)^{\frac{d-3}{2}} \mathrm{d}w \mathrm{d}z$$
$$\leq c \int_0^1 \frac{(1 - z^2)^{\frac{d-2}{2}}}{\left(1 + n\sqrt{1-z}\right)^{2\kappa}} \leq cn^{-d} \int_0^n \frac{r^{d-1}}{(1+r)^{\kappa}} \mathrm{d}r \leq cn^{-d}$$

by setting $r = n\sqrt{1-z}$ and recalling that $\kappa > d$. This completes the proof.

Proposition 3.12. For $\gamma \geq 0$ and $(x,t) \in \mathbb{X}_0^{d+1}$,

$$\int_{\mathbb{X}_0^{d+1}} \left| \mathsf{L}_n^E \big(\mathsf{w}_{0,\gamma}^\varrho; (x,t), (y,s) \big) \right|^p \mathsf{w}_{0,\gamma}^\varrho(s) \mathrm{d}\sigma_\varrho(y,s) \le \left(\frac{n^d}{\mathsf{w}_{0,\gamma}^\varrho(n;t)} \right)^{p-1}.$$

This follows by applying Lemma 3.11 on the estimate of Theorem 3.9.

We have established Assertions 1 and 3 for $\mathsf{L}_n^E(\mathsf{w}_{\beta,\gamma}^\varrho;\cdot,\cdot)$ and also Assertion 2 for $\mathsf{L}_n^E(\mathsf{w}_{0,\gamma}^\varrho;\cdot,\cdot)$. The kernel uses, however, only polynomials that are even in t and in s variable. Consequently, we have proved the following:

Corollary 3.13. For $d \ge 2$, $\varrho \ge 0$ and $\gamma \ge 0$, the space $(\mathbb{X}_0^{d+1}, \mathsf{w}_{0,\gamma}^{\varrho}, \mathsf{d}_{\mathbb{X}_0}^{\varrho})$ is a localizable homogeneous space, where its localized kernels are defined for polynomials even in t and in s variables.

3.5. Maximal ε -separated sets and MZ inequality. We provide a construction of maximal ε -separated set, as defined by Definition 2.3, on the double conic and hyperbolic surfaces.

We first consider the double conic surface; that is, $\rho = 0$. We shall need maximal ε -separated sets on the unit sphere. We adopt the following notation. For $\varepsilon > 0$, we denote by $\Xi_{\mathbb{S}}(\varepsilon)$ a maximal ε -separated set on the unit sphere \mathbb{S}^{d-1} and we let $\mathbb{S}_{\xi}(\varepsilon)$ be the subsets in \mathbb{S}^{d-1} so that the collection $\{\mathbb{S}_{\xi}(\varepsilon) : \xi \in \Xi_{\mathbb{S}}(\varepsilon)\}$ is a partition of \mathbb{S}^{d-1} , and we assume

(3.23)
$$\mathbf{c}_{\mathbb{S}}(\xi, c_1 \varepsilon) \subset \mathbb{S}_{\xi}(\varepsilon) \subset \mathbf{c}_{\mathbb{S}}(\xi, c_2 \varepsilon), \qquad \xi \in \Xi_{\mathbb{S}}(\varepsilon),$$

where $c_{\mathbb{S}}(\xi, \varepsilon)$ denotes the spherical cap centered at ξ with radius ε , c_1 and c_2 depending only on d. Such a $\Xi_{\mathbb{S}}(\varepsilon)$ exists for all $\varepsilon > 0$, see for example [8, Section 6.4], and its cardinality satisfies

(3.24)
$$c'_d \varepsilon^{-d+1} \le \# \Xi_{\mathbb{S}}(\varepsilon) \le c_d \varepsilon^{-d+1}.$$

Let $\varepsilon > 0$. We let $N = 2\lfloor \frac{\pi}{2}\varepsilon^{-1} \rfloor$, so that N is an even integer. For $1 \le j \le N$ we define

$$\theta_j := \frac{(2j-1)\pi}{2N}, \quad \theta_j^+ := \theta_j - \frac{\pi}{2N} \text{ and } \theta_j^- := \theta_j + \frac{\pi}{2N}.$$

Let $t_j = \cos \theta_j$, $t_j^- = \cos \theta_j^-$ and $t_j^+ = \cos \theta_j^+$. Thus, $t_1^+ = 1$ and $t_N^- = -1$ and

$$1 > t_1 > t_2 > \ldots > t_{\frac{N}{2}} > t_{\frac{N}{2}}^- = 0 = t_{\frac{N}{2}+1}^+ > t_{\frac{N}{2}+1}^- > \ldots > t_N > -1.$$

In particular, $t_{j-1}^+ = t_j^-$ and we can partition \mathbb{X}_0^{d+1} as the disjoint union of

$$\mathbb{X}_{0}^{(j)} := \left\{ (x,t) \in \mathbb{X}_{0}^{d+1} : t_{j}^{-} < t \leq t_{j}^{+} \right\}, \qquad 1 \leq j \leq N.$$

Furthermore, the upper and lower surfaces $\mathbb{X}_{0,+}^{d+1}$ and $\mathbb{X}_{0,-}^{d+1}$ can be partitioned by

$$\mathbb{X}_{0,+}^{d+1} = \bigcup_{j=1}^{N/2} \mathbb{X}_{0}^{(j)}$$
 and $\mathbb{X}_{0,-}^{d+1} = \bigcup_{j=N/2+1}^{N} \mathbb{X}_{0}^{(j)}$.

Let $\varepsilon_j := \pi \varepsilon/(2t_j)$. Then $\Xi_{\mathbb{S}}(\varepsilon_j)$ is the maximal ε_j -separated set of \mathbb{S}^{d-1} such that $\{\mathbb{S}_{\xi}(\varepsilon_j) : \xi \in \Xi_{\mathbb{S}}(\varepsilon_j)\}$ is a partition $\mathbb{S}^{d-1} = \bigcup_{\eta \in \Xi_{\mathbb{S}}(\varepsilon_j)} \mathbb{S}_{\eta}(\varepsilon_j)$, and $\#\Xi_{\mathbb{S}}(\varepsilon_j) \sim \varepsilon_j^{-d+1}$. For each $j = 1, \ldots, N$, we decompose $\mathbb{X}_0^{(j)}$ by

$$\mathbb{X}_{0}^{(j)} = \bigcup_{\xi \in \Xi_{\mathbb{S}}(\varepsilon_{j})} \mathbb{X}_{0}(\xi, t_{j}), \quad \text{where} \quad \mathbb{X}_{0}(\xi, t_{j}) := \left\{ (t\eta, t) : t_{j}^{-} < t \leq t_{j}^{+}, \eta \in \mathbb{S}_{\xi}(\varepsilon_{j}) \right\}.$$

Finally, we define the subset $\Xi_{\mathbb{X}_0}$ of \mathbb{X}_0^{d+1} by

$$\Xi_{\mathbb{X}_0} = \left\{ (t_j \xi, t_j) : \xi \in \Xi_{\mathbb{S}}(\varepsilon_j), \, 1 \le j \le N \right\}.$$

Proposition 3.14. Let $\varepsilon > 0$ and $N = 2\lfloor \frac{\pi}{2}\varepsilon^{-1} \rfloor$. Then $\Xi_{\mathbb{X}_0}$ is a maximal ε -separated set of \mathbb{X}_0^{d+1} and $\{\mathbb{X}_0(\xi, t_j) : (t_j\xi, t_j) \in \Xi_{\mathbb{X}_0}\}$ is a partition

$$\mathbb{X}_{0}^{d+1} = \bigcup_{(t\xi,t)\in \Xi_{\mathbb{X}_{0}}} \mathbb{X}_{0}(\xi,t) = \bigcup_{j=1}^{N} \bigcup_{\xi\in \Xi_{\mathbb{S}}(\varepsilon_{j})} \mathbb{X}_{0}(\xi,t_{j})$$

Moreover, there are positive constants c_1 and c_2 depending only on d such that

(3.25)
$$\mathbf{c}_0((t_j\xi,t_j),c_1\varepsilon) \subset \mathbb{X}_0(\xi,t_j) \subset \mathbf{c}_0((t_j\xi,t_j),c_2\varepsilon),$$

and c'_d and c_d such that

(3.26)
$$c'_d \varepsilon^{-d} \le \# \Xi_{\mathbb{X}_0} \le c_d \varepsilon^{-d}.$$

Proof. Let $(t_j\xi, t_j)$ and $(t_k\eta, t_k)$ be two distinct points in $\Xi_{\mathbb{X}_0}$. By its definition, $\mathsf{d}_{[-1,1]}(t_j, t_k) = |\theta_j - \theta_k| \geq \frac{\pi}{N} \geq \varepsilon$ if $j \neq k$. Hence,

$$\mathsf{d}_{\mathbb{X}_0}\big((t_j\xi, t_j), (t_k\eta, t_k)\big) \ge \mathsf{d}_{[0,1]}(t_j, t_k) \ge \varepsilon, \quad j \neq k.$$

If j = k, then ξ and η are both elements of $\mathbb{S}(\varepsilon_j)$, so that $\mathsf{d}_{\mathbb{S}}(\xi, \eta) \ge \varepsilon_j$. Hence, using $\frac{2}{\pi}\phi \le \sin\phi \le \phi$, we deduce from (3.3) that

$$\mathsf{d}_{\mathbb{X}_0}\big((t_j\xi,t_j),(t_j\eta,t_j)\big) \geq \frac{2}{\pi}t_j\mathsf{d}_{\mathbb{S}}(\xi,\eta) \geq \frac{2}{\pi}t_j\varepsilon_j = \varepsilon.$$

Hence, $\Xi_{\mathbb{X}_0}$ is ε -separated. Moreover, since $\#\Xi_{\mathbb{S}}(\varepsilon_j) \sim \varepsilon_j^{-d+1}$,

$$\#\Xi_{\mathbb{X}_0} = \sum_{j=1}^N \#\Xi_{\mathbb{S}}(\varepsilon_j) \sim \sum_{j=1}^N \varepsilon_j^{-d+1} \sim \varepsilon^{-d+1} \sum_{j=1}^N t_j^{d-1} \sim \varepsilon^{-d+1} N \sim \varepsilon^{-d}.$$

For the proof of (3.25), we only need to consider $X_0(\xi, t_j)$ in the upper part of the double cone, which means $1 \leq j \leq N/2$. If $\mathsf{d}_{[-1,1]}(s, t_j) \leq \delta/N$ with $\delta \leq 1/2$, then

$$|t_j - s| \le \mathsf{d}_{[-1,1]}(t_j, s) \le \frac{\delta}{N} \le \delta \sin \frac{\pi}{2N} = \delta t_{\frac{N}{2}} \le \delta t_j.$$

Hence, by $\delta \leq \frac{1}{2}$, we obtain $s \geq t_j/2$. Similarly, we see that if $s \in c_{[-1,1]}(t_j, \pi/N)$, then $s \leq c_* t_j$. By (3.23), there are constants $b_1 > 0$ and $b_2 > 0$ such that $c_{\mathbb{S}}(\xi, b_1 \varepsilon_j) \subset S_{\xi}(\varepsilon_j) \subset c_{\mathbb{S}}(\xi, b_2 \varepsilon_j)$. We claim that (3.25) holds for some $c_1 < \delta$ and some $c_2 > b_2$. Indeed, if $(y, \eta) \subset c_0((t_j \xi, t_j), c_1 \varepsilon)$, then $\mathsf{d}_{[-1,1]}(s, t_j) \leq c_1 \varepsilon \leq \delta/N$; moreover, by $s \geq t_j/2$ and $(st)^{\frac{1}{2}} \mathsf{d}_{\mathbb{S}}(\xi, \eta) \leq cc_1 \varepsilon$ by (3.4), we see that $\mathsf{d}_{\mathbb{S}}(\xi, \eta) \leq 2^{\frac{1}{4}} cc_1 \varepsilon / \sqrt{t_j} \leq b_1 \varepsilon_j$ by choosing c_1 small. This establishes the left-hand side inclusion of (3.25). The right-hand side inclusion can be similarly established. The proof is completed. \Box

For $\rho > 0$, the point set on the hyperbolic surface ${}_{\varrho}\mathbb{X}_{0}^{d+1}$ can be deduced easily from that on the double conic surface.

Proposition 3.15. For $\varepsilon > 0$, let $\Xi_{\mathbb{X}_0}$ be a maximal ε -separated set in \mathbb{X}_0^{d+1} . Define

$$\Xi_{\mathbb{X}_0}^{\varrho} := \left\{ \left(x, \sqrt{t^2 - \varrho^2} \right) : (x, t) \in \Xi_{\mathbb{X}_0} \right\}.$$

Then $\Xi_{\mathbb{X}_0}^{\varrho}$ is a maximal ε -separated set in $_{\varrho}\mathbb{X}_0^{d+1}$.

This is an immediate consequence of (3.5). In particular, if $\Xi_{\mathbb{X}_0}$ is the set given in Proposition 3.14, then we can define $\mathbb{X}_0^{\varrho}(\xi, t_j)$ accordingly so that both (3.25) and (3.26) hold.

Definition 3.16. Let $\Xi_{\mathbb{X}_0}^{\varrho}$ be a set on \mathbb{X}_0^{d+1} that does not contain (0,0). Define

$$\Xi^{\varrho}_{\mathbb{X}_0,+} = \{(x,t)\in\Xi^{\varrho}_{\mathbb{X}_0}:t>0\} \quad and \quad \Xi^{\varrho}_{\mathbb{X}_0,-} = \{(x,t)\in\Xi^{\varrho}_{\mathbb{X}_0}:t<0\}.$$

We call the set $\Xi^{\varrho}_{\mathbb{X}_0}$ evenly symmetric on \mathbb{X}_0^{d+1} if

$$\Xi^{\varrho}_{\mathbb{X}_0,-} = \left\{ (-x,-t) : (x,t) \in \Xi^{\varrho}_{\mathbb{X},+} \right\}.$$

By definition, $\Xi_{\mathbb{X}_0} = \Xi_{\mathbb{X}_0,+} \cup \Xi_{\mathbb{X}_0,-}$ and the two subsets are disjoint. For the set $\Xi_{\mathbb{X}_0}$ in Proposition 3.14, the points t_j , $1 \leq j \leq N$, are zeros of the Chebyshev polynomial of the first kind defined by $T_N(t) = \cos N \arccos(x)$ and we have

$$\begin{split} \Xi_{\mathbb{X}_{0},+} &= \left\{ (t_{j}\xi,t_{j}) \in \Xi_{\mathbb{X}_{0}} : 1 \leq j \leq \frac{N}{2} \right\}, \quad \Xi_{\mathbb{X}_{0},-} = \left\{ (t_{j}\xi,t_{j}) \in \Xi_{\mathbb{X}_{0}} : \frac{N}{2} + 1 \leq j \leq N \right\}.\\ \text{Since } t_{j} &= -t_{N-j} \text{ for } 0 \leq j \leq \frac{N}{2}, \text{ it follows that the set in Proposition 3.14 is evenly symmetric on } \mathbb{X}_{0}^{d+1}, \text{ so is its analogue on } _{\varrho}\mathbb{X}_{0}^{d+1}. \end{split}$$

We further notice that, for the set $\Xi_{\mathbb{X}_0}^{\varepsilon}$ in Proposition 3.14, with either + or -, $\Xi_{\mathbb{X}_0,\pm}$ is a maximal ε -separated sets of $\mathbb{X}_{0,\pm}^{d+1}$ and the set $\{\mathbb{X}_0(\xi,t_j):(t_j\xi,t_j)\in\Xi_{\mathbb{X}_0,\pm}\}$ is a partition of $\mathbb{X}_{0,\pm}^{d+1}$. Comparing with the maximal ε -separated set $\Xi_{\mathbb{V}_0}$ constructed on $\mathbb{V}_0^{d+1} = \mathbb{X}_{0,\pm}^{d+1}$ in [30], we see that the points in $\Xi_{\mathbb{V}_0}$ congest towards the apex, with a rate $t_j \sim N^{-2}$, whereas the points in $\Xi_{\mathbb{X}_0}$ do not. We have shown that $(\mathbb{X}_0^{d+1}, \mathbb{W}_{0,\gamma}^{\varrho}, \mathsf{d}_{\mathbb{X}_0}^{\varrho})$ is a localizable homogeneous space with the

We have shown that $(\mathbb{X}_{0}^{d+1}, \mathsf{w}_{0,\gamma}^{d}, \mathsf{d}_{\mathbb{X}_{0}}^{\mathbb{X}})$ is a localizable homogeneous space with the highly localized kernels $\mathsf{L}_{n}^{E}(\mathsf{w}_{0,\gamma}^{\varrho}; \cdot, \cdot)$. As part of the framework in [30], we can then state the Marcinkiewicz-Zygmund inequality for a doubling weight on \mathbb{X}_{0}^{d+1} , which holds under the following constrains: the weight function w need to be even in tvariable so that the integral of polynomials in $\Pi_{n}^{E}(\mathbb{X}_{0}^{d+1})$ can be written as over $\mathbb{X}_{0,+}^{d+1}$, the maximal $\frac{\delta}{n}$ -separated set $\Xi_{\mathbb{X}_{0}}$ need to be symmetric, and it works for polynomials even in t variable. Let us define, for $n = 0, 1, 2, \ldots$,

$$\Pi_n^E(\mathbb{X}_0^{d+1}) = \left\{ p \in \Pi_n(\mathbb{X}_0^{d+1}) : p(x,t) = p(x,-t), \forall (x,t) \in \mathbb{X}_0^{d+1} \right\}$$

Theorem 3.17. Let w be an doubling weight on \mathbb{X}_0^{d+1} such that w(x,t) = w(x,-t). Let $\Xi_{\mathbb{X}_0}^{\varrho}$ be a symmetric maximal $\frac{\delta}{n}$ -separated subset of \mathbb{X}_0^{d+1} and $0 < \delta \leq 1$.

(i) For all $0 and <math>f \in \Pi_m^E(\mathbb{X}^{d+1})$ with $n \le m \le cn$,

$$\sum_{z \in \Xi_{\mathbb{X}_0}} \Big(\max_{(x,t) \in \mathsf{c}_{\varrho}((z,r),\frac{\delta}{n})} |f(x,t)|^p \Big) \mathsf{w} \big(\mathsf{c}_{\varrho}((z,r),\frac{\delta}{n}) \big) \le c_{\mathsf{w},p} \|f\|_{p,\mathsf{w}}^p$$

where $c_{w,p}$ depends on p when p is close to 0 and on the doubling constant of w.

(ii) For 0 < r < 1, there is a $\delta_r > 0$ such that for $\delta \leq \delta_r$, $r \leq p < \infty$ and $f \in \prod_n^E(\mathbb{X}_0^{d+1})$,

$$\|f\|_{p,\mathsf{w}}^{p} \leq c_{\mathsf{w},r} \sum_{z \in \Xi} \Big(\min_{(x,t) \in \mathsf{c}_{\varrho}\left((z,r),\frac{\delta}{n}\right)} |f(x,t)|^{p} \Big) \mathsf{w}\big(\mathsf{c}_{\varrho}((z,r),\frac{\delta}{n})\big)$$

where $c_{w,r}$ depends only on the doubling constant of w and on r when r is close to 0.

This is a consequence of [30, Theorem 2.15]; its proof remains valid for polynomials in $\Pi_m^E(\mathbb{X}^{d+1})$ under the assumptions on symmetry.

3.6. **Positive cubature rules.** The Marcinkiewicz-Zygmund inequality is used to establish the positive cubature rule in the general framework. In order to quantify the coefficients of the cubature rule, we will need Assertion 4. This is given by the fast decaying polynomials on the hyperbolic surface given below.

Lemma 3.18. Let $d \geq 2$ and $\varrho \geq 0$. For each $(x,t) \in \mathbb{X}_0^{d+1}$, there is a polynomial $T_{(x,t)}^{\varrho}$ in $\prod_n^E(\mathbb{X}_0^{d+1})$ that satisfies

(1) $T^{\varrho}_{(x,t)}(x,t) = 1, \ T^{\varrho}_{x,t}(y,s) \ge c > 0 \ \text{if } (y,s) \in \mathsf{c}_{\varrho}((x,t),\frac{\delta}{n}), \text{ and for every } \kappa > 0,$

$$0 \leq T^{\varrho}_{(x,t)}(y,s) \leq c_{\kappa} \left(1 + \mathsf{d}^{\varrho}_{\mathbb{X}_{0}}\big((x,t),(y,s)\big)\right)^{-\kappa}, \quad (y,s) \in \mathbb{X}_{0}^{d+1}.$$

(2) there is a polynomial q(t) of degree 4n such that $q(t)T^{\varrho}_{(x,t)}$ is a polynomial of degree 5n in (x,t) variables and $1 \le q_n(t) \le c$.

Proof. For positive integer n, let $m = \lfloor \frac{n}{r} \rfloor + 1$ and define

$$S_n(\cos\theta) = \left(\frac{\sin(m+\frac{1}{2})\frac{\theta}{2}}{(m+\frac{1}{2})\sin\frac{\theta}{2}}\right)^{2r}, \qquad 0 \le \theta \le \pi.$$

Then $S_n(z)$ is an even algebraic polynomial of degree at most 2n and it satisfies

(3.27)
$$S_n(1) = 1, \qquad 0 \le S_n(\cos\theta) \le c(1+n\theta)^{-2r}, \quad 0 \le \theta \le \pi.$$

To construct polynomials on hyperbolic surface, we define, for $(x, t), (y, s) \in \mathbb{X}_0^{d+1}$,

$$S((x,t),(y,s)) = S_n \left(\langle x,y \rangle + \sqrt{1 + \varrho^2 - t^2} \sqrt{1 + \varrho^2 - s^2} \right) + S_n \left(\langle x,y \rangle - \sqrt{1 + \varrho^2 - t^2} \sqrt{1 + \varrho^2 - s^2} \right).$$

Since S_n is an even polynomial, it follows that S((x,t), (y,s)) is a polynomial of degree n in either (x,t) or (y,s) variables. Moreover, since $||y||^2 = s^2 - \rho^2$, it also follows that $S((x,t), \cdot)$ is even in s variable. Define

$$T^{\varrho}_{(x,t)}(y,s) = \frac{S((x,t),(y,s))}{1 + S_n(2t^2 - 2\varrho^2 - 1)}, \qquad (y,s) \in \mathbb{X}_0^{d+1}.$$

Then $T_{(x,t)}^{\varrho} \in \Pi_n^E(\mathbb{X}_0^{d+1})$ and $T_{(x,t)}^{\varrho}(x,t) = 1$. If $0 \leq \theta \leq \frac{2\pi}{2m+1}$, then it follows from $\sin \theta \geq \frac{2}{\pi} \theta$ for $0 \leq \theta \leq \pi/2$ that $S_n(\cos \theta) \geq \left(\frac{2}{\pi}\right)^{2r}$. Hence, since $0 \leq S_n(2t-1) \leq 1$, it follows that

$$T^{\varrho}_{(x,t)}(y,s) \ge \frac{S_n\left(\langle x,y \rangle + \sqrt{1 + \varrho^2 - t^2}\sqrt{1 + \varrho^2 - s^2}\right)}{1 + S_n(2t^2 - 2\varrho^2 - 1)} \ge \frac{1}{2}\left(\frac{2}{\pi}\right)^{2r}$$

for $(y,s) \in c((x,t), \frac{2\pi}{2m+1})$. Next, we use the estimate $S_n(z) \leq c(1 + n\sqrt{1-t^2})^{-2r}$ for $|t| \leq 1$ to obtain an upper bound for $T_{(x,t)}$. Since $1 - \langle x, y \rangle + \sqrt{1-t^2}\sqrt{1-s^2} \geq 1 - \langle x, y \rangle - \sqrt{1-t^2}\sqrt{1-s^2}$, we obtain

$$0 \le T^{\varrho}_{(x,t)}(y,s) \le c \left(1 + n\sqrt{1 - \langle x, y \rangle - \sqrt{1 - t^2}\sqrt{1 - s^2}}\right)^{-2r} = c \left(1 + n\sqrt{1 - \cos \mathsf{d}_{\mathbb{V}_0}((x,t),(y,s))}\right)^{-2r} \sim (1 + n\mathsf{d}_{\mathbb{X}_0}((x,t),(y,s)))^{-2r}$$

using $1 - \cos \theta \sim \theta^2$. This completes the proof of item (1). The item (2) follows from setting $q(t) = 1 + S_n(2t^2 - 2\varrho^2 - 1)$, which is a polynomial of degree 4n and $1 \le q(t) \le c$. This completes the proof.

The lemma establishes Assertion 4 with a polynomial in $\Pi_n^E(\mathbb{X}_0^{d+1})$. It allows us to follow the general framework to bound the Christoffel function in $\Pi_n^E(\mathbb{X}_0^{d+1})$. Let w be a doubling weight function, even in t variable, on \mathbb{X}_0^{d+1} . Let

(3.28)
$$\lambda_n^E(\mathsf{w}; x, t) := \inf_{\substack{g(x,t)=1\\g\in\Pi_n^E(\mathbb{X}_0^{d+1})}} \int_{\mathbb{X}_0^{d+1}} |g(x,t)|^2 \mathsf{w}(x,t) \mathrm{d}\sigma(x,t),$$

which is the Christoffel function for the space $\Pi_n^E(\mathbb{X}_0^{d+1})$. Let $\mathsf{K}_n^E(\mathsf{w};\cdot,\cdot)$ denote the kernel of the *n*-th partial sum operator of the the series (3.13). Then

$$\mathsf{K}_n^E(\mathsf{w};(x,t),(y,s)) = \sum_{k=0}^n \mathsf{P}_k^E(\mathsf{w};(x,t),(y,s)).$$

By the proof of [9, Theorem 4.6.6], it is related to the Christoffel function by

$$\lambda_n^E(\mathbf{w}; x, t) = \frac{1}{\mathsf{K}_n^E(\mathbf{w}; (x, t), (x, t))}, \qquad (x, t) \in \mathbb{X}_0^{d+1}.$$

Using Lemma 3.18, we can adopt [30, Propositions 2.17 and 2.18] to bound $\lambda_n^E(w)$.

Corollary 3.19. Let w be a doubling weight function on \mathbb{X}_0^{d+1} such that w(x,t) = w(x,-t) for all $(x,t) \in \mathbb{X}_0^{d+1}$. Then

$$\lambda_n^E(\mathsf{w};(x,t)) \le c \,\mathsf{w}\left(\mathsf{c}_{\varrho}\left((x,t),\frac{1}{n}\right)\right).$$

Moreover, for $\gamma \geq 0$,

$$\lambda_n^E \left(\mathsf{w}_{0,\gamma}^{\varrho}; (x,t) \right) \ge c' \, \mathsf{w}_{0,\gamma}^{\varrho} \left(\mathsf{c}_{\varrho} \left((x,t), \frac{1}{n} \right) \right) = c' n^{-d} \mathsf{w}_{0,\gamma}^{\varrho} (n;t).$$

We can now state the positive cubature rule for the hyperbolic surface, which holds for polynomials in $\Pi_n^E(\mathbb{X}_0^{d+1})$ under the assumption of symmetry for both the weight w and the set $\Xi_{\mathbb{X}_0}$.

Theorem 3.20. Let $d \ge 2$ and $\varrho \ge 0$. Let w be a doubling weight on \mathbb{X}_0^{d+1} such that w(x,t) = w(x,-t) for all $(x,t) \in \mathbb{X}_0^{d+1}$. Let Ξ^{ϱ} be a symmetric maximum $\frac{\delta}{n}$ -separated subset of \mathbb{X}_0^{d+1} . There is a $\delta_0 > 0$ such that for $0 < \delta < \delta_0$ there exist positive numbers $\lambda_{z,r}$, $(z,r) \in \Xi^{\varrho}$, so that

(3.29)
$$\int_{\mathbb{X}_0^{d+1}} f(x,t) \mathsf{w}(x,t) \mathrm{d}\sigma(x,t) = \sum_{(z,r)\in\Xi^{\varrho}} \lambda_{z,r} f(z,r), \quad \forall f \in \Pi_n^E(\mathbb{X}_0^{d+1}).$$

Moreover, $\lambda_{z,r} \sim w(c_{\varrho}((z,r), \frac{\delta}{n}))$ for all $(z,r) \in \Xi^{\varrho}$.

This is [30, Theorem 2.20] when the domain becomes \mathbb{X}_0^{d+1} and it remains valid under the assumptions on symmetry.

3.7. Localized polynomial frame. The localized polynomials are constructed using the highly localized kernel defined with a cut-off function \hat{a} of type type (b) that satisfies $\hat{a}(t) > a > 0 \qquad \text{if } t \in [2/5, 5/2]$

(3.30)
$$\begin{aligned} a(t) \ge \rho > 0, & \text{if } t \in [3/5, 5/3], \\ [\widehat{a}(t)]^2 + [\widehat{a}(2t)]^2 = 1, & \text{if } t \in [1/2, 1]. \end{aligned}$$

For the hyperbolic or double conic surface, we will also require symmetry for the weight and for the ε -separated subset. Let $\varrho \geq 0$ and let w be a doubling weight on \mathbb{X}_0^{d+1} and assume that it is even in t variable. Let $\mathsf{L}_n^E(\mathsf{w}) * f$ denote the near best approximation operator from $\Pi_{2n}^E(\mathbb{X}^{d+1})$ defined by

$$(3.31) \qquad \qquad \mathsf{L}_n^E(\mathsf{w})*f(x):=\int_{\mathbb{X}_0^{d+1}}f(y,s)\mathsf{L}_n^E(\mathsf{w};(x,t),(y,s))\mathsf{w}(y,s)\mathrm{d}\sigma_\varrho(y,s).$$

For $j = 0, 1, \ldots$, let Ξ_j^{ϱ} be a symmetric maximal $\frac{\delta}{2^j}$ -separated subset in \mathbb{X}_0^{d+1} , so that

$$\int_{\mathbb{X}_0^{d+1}} f(x,t) \mathsf{w}(x,t) \mathrm{d}\sigma_{\varrho}(x,t) = \sum_{(z,r) \in \Xi_j^{\varrho}} \lambda_{(x,r),j} f(z,r), \quad f \in \Pi_{2^j}^E(\mathbb{X}_0^{d+1}).$$

For $j = 1, 2, \ldots$, define the operator $F_j^{\varrho}(\mathsf{w})$ by

$$F_j^{\varrho}(\mathsf{w}) * f = \mathsf{L}_{2^{j-1}}^E(\mathsf{w}) * f$$

and define the frame elements $\psi_{(z,r),j}$ for $(z,r) \in \Xi_j^{\varrho}$ by

$$\psi_{(z,r),j}(x,t) := \sqrt{\lambda_{(z,r),j}} F_j^{\varrho}((x,t),(z,r)), \qquad (x,t) \in \mathbb{X}_0^{d+1}.$$

Then $\Phi = \{\psi_{(z,r),j} : (z,r) \in \Xi_j^{\varrho}, j = 1, 2, 3, ...\}$ is a tight frame. Following [30, Theorem 2.21] of the general framework, we have the following:

Theorem 3.21. Let w be a doubling weight on \mathbb{X}_0^{d+1} even in its t variable. If $f \in L^2(\mathbb{X}_0^{d+1}, w)$ and f is even in t variable, then

$$f = \sum_{j=0}^{\infty} \sum_{(z,r)\in \Xi_j^\varrho} \langle f, \psi_{(z,r),j} \rangle_{\mathsf{w}} \psi_{(z,r),j} \qquad in \ L^2(\mathbb{X}_0^{d+1}, \mathsf{w})$$

and

$$\|f\|_{2,\mathbf{w}} = \Big(\sum_{j=0}^{\infty} \sum_{(z,r)\in \Xi_j^{\varrho}} |\langle f, \psi_{(z,r),j} \rangle_{\mathbf{w}}|^2 \Big)^{1/2}.$$

Furthermore, for $\gamma \geq -\frac{1}{2}$, the frame for $w_{0,\gamma}^{\varrho}$ is highly localized in the sense that, for every $\sigma > 0$, there exists a constant $c_{\sigma} > 0$ such that

(3.32)
$$|\psi_{(z,r),j}(x,t)| \le c_{\sigma} \frac{2^{jd/2}}{\sqrt{\mathsf{w}_{0,\gamma}^{\varrho}(2^{j};t)}(1+2^{j}\mathsf{d}_{\mathbb{X}_{0}}^{\varrho}((x,t),(z,r)))^{\sigma}}}, \quad (x,t) \in \mathbb{X}_{0}^{d+1}.$$

The frame elements involve only orthogonal polynomials even in t variable and they are well defined for all doubling weight that is even in its t variable. The localization (3.32) follows from Theorem 3.9 and $\lambda_{(z,r),j} \sim 2^{-jd} \mathsf{w}^{\varrho}_{0,\gamma}(2^j;t)$, which holds for $\mathsf{w}^{\varrho}_{0,\gamma}$ by Corollary 3.19 and (3.8). It is worthwhile to point out that the localized frame is established for the Lebesgue measure on \mathbb{X}^{d+1}_0 , which is the case $\mathsf{w}_{0,\frac{1}{2}}$, and in particular for the Lebesgue measure on the upper conic surface \mathbb{V}_0^{d+1} , in contrast to the localized frame established in [30, Section 2.7].

3.8. Characterization of best approximation. For $f \in L^p(\mathbb{X}_0^{d+1}, \mathsf{w}_{0,\gamma}^{\varrho})$, we denote by $\mathsf{E}_n(f)_{p,\mathsf{w}_{0,\gamma}^{\varrho}}$ the best approximation to f from $\Pi_n(\mathbb{X}_0^{d+1})$, the space of polynomials of degree at most n restricted on the surface \mathbb{X}_0^{d+1} , in the norm $\|\cdot\|_{p,\mathsf{w}_{0,\gamma}^{\varrho}}$; that is,

$$\mathsf{E}_{n}(f)_{p,\mathsf{w}_{0,\gamma}^{\varrho}} := \inf_{g \in \Pi_{n}(\mathbb{X}_{0}^{d+1})} \|f - g\|_{p,\mathsf{w}_{0,\gamma}^{\varrho}}, \qquad 1 \le p \le \infty.$$

If f is even in t variable, then the triangle inequality and changing variable $t \mapsto -t$ shows, by the symmetry of the integrals,

$$\left\| f(x,t) - \frac{1}{2} (g(x,t) + g(x,-t)) \right\|_{p,\mathsf{w}^{\varrho}_{0,\gamma}} \le \|f - g\|_{p,\mathsf{w}^{\varrho}_{0,\gamma}}$$

Hence, we can choose the polynomial of best approximation from $\Pi_n^E(\mathbb{X}_0^{d+1})$ when f is symmetric in t variable. Following the general framework in [30], we can give a characterization of best approximation by polynomials for functions even in t variable.

We define a K-functional and a modulus of smoothness. In terms of the fractional differential operator $(-\Delta_{0,\gamma}^{\varrho})^{\frac{r}{2}}$ and a doubling weight w, even in t variable on \mathbb{X}_{0}^{d+1} , the K-functional fo $f \in L^{p}(\mathbb{X}_{0}^{d+1}, w)$ and r > 0, is defined by

$$\mathsf{K}_{r}(f,t)_{p,\mathsf{w}} := \inf_{g \in W_{p}^{r}(\mathbb{X}_{0}^{d+1},\mathsf{w})} \left\{ \|f - g\|_{p,\mathsf{w}} + t^{r} \left\| (-\Delta_{0,\gamma}^{\varrho})^{\frac{r}{2}} f \right\|_{p,\mathsf{w}} \right\},\$$

where the Sobolev space $W_p^r(\mathbb{X}_0^{d+1}, \mathbf{w})$ is the space that consists of $g \in L^p(\mathbb{X}_0^{d+1}, \mathbf{w})$, even in t variable, so that $\|(-\Delta_{0,\gamma}^{\varrho})^{\frac{r}{2}}g\|_{p,\mathbf{w}}$ is finite. The K-functional is well defined, as shown in [30, Section 3.3], where we need to require the functions being even in tvariable in the proof. Moreover, the modulus of smoothness is defined by

$$\omega_r(f;\rho)_{p,\mathsf{w}^\varrho_{0,\gamma}} = \sup_{0\le\theta\le\rho} \left\| \left(I - \mathsf{S}_{\theta,\mathsf{w}^\varrho_{0,\gamma}}\right)^{r/2} f \right\|_{p,\mathsf{w}^\varrho_{0,\gamma}}, \quad 1\le p\le\infty,$$

where the operator $\mathsf{S}_{\theta,\mathsf{w}_{0,\gamma}^{\varrho}}$ is defined by, for $n = 0, 1, 2, \ldots$ and $\lambda = \gamma + \frac{d-1}{2}$,

$$\operatorname{proj}_{n}^{E}(\mathsf{w}_{0,\gamma}^{\varrho};\mathsf{S}_{\theta,\mathsf{w}_{0,\gamma}^{\varrho}}f) = \frac{C_{n}^{\lambda-\frac{1}{2}}(\cos\theta)}{C_{n}^{\lambda-\frac{1}{2}}(1)}\operatorname{proj}_{n}^{E}(\mathsf{w}_{0,\gamma}^{\varrho};f).$$

The operator is well defined since the above relations determine it uniquely among functions even in t variable and in $L^p(\mathbb{X}_0^{d+1}, \mathsf{w}_{0,\gamma}^{\varrho})$ and the modulus of smoothness satisfies the usual properties under its name. Such a modulus of smoothness is in line with those defined on the unit sphere and the unit ball [8, 20, 27] but its structure is more complicated. There are recent results for modulus of smoothness on fairly general domains in \mathbb{R}^d [5, 6, 23, 24], defined via simple difference operators of functions, but conic domains are not included.

By [30, Theorem 3.1.2], the characterization of the best approximation holds under the Assertions 1, 3 and 5. For $w_{0,\gamma}^{\varrho}$ on \mathbb{X}_{0}^{d+1} , we have already established Assertions 1 and 3. We now establish Assertion 5, again for kernels even in t and s variables. By (3.14), the kernel $L_{n}^{(r)}(\varpi)$ in Assertion 5 becomes

$$\mathsf{L}_{n}^{(r)}\big(\mathsf{w}_{0,\gamma}^{\varrho};(x,t),(y,s)\big) = \sum_{k=0}^{\infty} \widehat{a}\left(\frac{k}{n}\right) \left(k(k+2\gamma+d-1)\right)^{\frac{r}{2}} \mathsf{P}_{k}^{E}\big(\mathsf{w}_{0,\gamma}^{\varrho};(x,t),(y,s)\big).$$

Lemma 3.22. Let $\gamma \geq -\frac{1}{2}$ and $\kappa > 0$. Then, for r > 0 and $(x,t), (y,s) \in \mathbb{X}_{0,+}^{d+1}$,

$$\left|\mathsf{L}_{n}^{(r)}\big(\mathsf{w}_{0,\gamma}^{\varrho};(x,t),(y,s)\big)\right| \leq c_{\kappa} \frac{n^{d+r}}{\sqrt{\mathsf{w}_{0,\gamma}^{\varrho}(n;t)}} \Big(1 + n\mathsf{d}_{\mathbb{X}_{0}}^{\varrho}((x,t),(y,s))\Big)^{-\kappa}.$$

Proof. By (3.15) it suffices to consider the case $\rho = 0$. By (3.16), the kernel can be written as

$$\mathsf{L}_{n}^{(r)}(\mathsf{w}_{0,\gamma}^{\varrho};(x,t),(y,s)) = c_{\gamma} \int_{-1}^{1} L_{n,r}\left(\zeta(x,t,y,s;v)\right) (1-v^{2})^{\gamma-1} \mathrm{d}v$$

where $L_{n,r}$ is defined by, with $\lambda = \gamma + \frac{d-1}{2}$,

$$L_{n,r}(t) = \sum_{k=0}^{\infty} \widehat{a}\left(\frac{k}{n}\right) \left(k(k+2\gamma+d-1)\right)^{\frac{r}{2}} \frac{P_n^{(\lambda-\frac{1}{2},\lambda-\frac{1}{2})}(1)P_n^{(\lambda-\frac{1}{2},-\lambda-\frac{1}{2})}(t)}{h^{(\lambda-\frac{1}{2},\lambda-\frac{1}{2})}}$$

Applying (2.12) with $\eta(t) = \hat{a}(t) \left(t(t+n^{-1}(2\gamma+d-1)) \right)^{\frac{r}{2}}$ and m = 0, we obtain

$$|L_{n,r}(t)| \le cn^r \frac{n^{2\lambda+1}}{(1+n\sqrt{1-t})^{\ell}}$$

Using this estimate, it is easy to see that the proof follows from the estimate already established in the proof of Theorem 3.9. $\hfill \Box$

We are now in position to state the characterization of the best approximation by polynomials for the hyperbolic surface, following [30, Theorem 3.12].

Theorem 3.23. Let $f \in L^p(\mathbb{X}_0^{d+1}, \mathsf{w})$ if $1 \leq p < \infty$ and $f \in C(\mathbb{X}_0^{d+1})$ if $p = \infty$. Assume that f satisfies f(x,t) = f(x,-t). Let $\varrho \geq 0$, r > 0 and $n = 1, 2, \ldots$. Then

(i) For $\mathbf{w} = \mathbf{w}_{0,\gamma}^{\varrho}$ with $\gamma \ge 0$,

$$\mathsf{E}_{n}(f)_{p,\mathsf{w}_{0,\gamma}^{\varrho}} \leq c \,\mathsf{K}_{r}(f;n^{-1})_{p,\mathsf{w}_{0,\gamma}^{\varrho}}$$

(ii) for every doubling weight w that satisfies w(x,t) = w(x,-t) on \mathbb{X}_0^{d+1} ,

$$\mathsf{K}_r(f; n^{-1})_{p,\mathsf{w}} \le cn^{-r} \sum_{k=0}^n (k+1)^{r-1} \mathsf{E}_k(f)_{p,\mathsf{w}}.$$

For $w = w_{0,\gamma}$, the K-functional is equivalent to the modulus of smoothness.

Theorem 3.24. Let $\gamma \geq 0$ and $f \in L_p^r(\mathbb{X}_0^{d+1}, \mathsf{w}_{0,\gamma}^\varrho)$, $1 \leq p \leq \infty$. Then for $0 < \theta \leq \pi/2$ and r > 0

$$c_1\mathsf{K}_r(f;\theta)_{p,\mathsf{w}_{0,\gamma}^\varrho} \leq \omega_r(f;\theta)_{p,\mathsf{w}_{0,\gamma}^\varrho} \leq c_2\mathsf{K}_r(f;\theta)_{p,\mathsf{w}_{0,\gamma}^\varrho}$$

Consequently, the characterization in Theorem 3.23 can be stated in terms of the modulus of smoothness in place of K-functional.

Finally, we mention that, if \hat{a} is an admissible cut-off function of type (a), then the operator $\mathsf{L}_n^E(\varpi) * f$ defined in (3.31) is the near best approximation in the sense that

$$\|\mathsf{L}_{n}^{E}(\mathsf{w}_{0,\gamma}^{\varpi}) - f\|_{p,\mathsf{w}} \le c \,\mathsf{E}_{n}(f)_{p,\mathsf{w}_{0,\gamma}^{\varrho}}, \quad 1 \le p \le \infty$$

for all $f \in L_p^r(\mathbb{X}_0^{d+1}, \mathsf{w})$, where w is a doubling weight by [30, Theorem 3.15].

4. Homogeneous space on double cone and hyperboloid

We work in the setting of homogeneous space on the solid domain defined by

$$\mathbb{X}^{d+1} = \left\{ (x,t) : \|x\|^2 \le t^2 - \varrho^2, \, x \in \mathbb{R}^d, \, \varrho \le |t| \le \sqrt{\varrho^2 + 1} \right\}$$

which is a double hyperboloid when $\rho > 0$ and a double cone when $\rho = 0$, and it is bounded by \mathbb{X}_0^{d+1} and the hyperplanes $t = \pm \sqrt{\rho^2 + 1}$ of \mathbb{R}^{d+1} . The analysis on \mathbb{X}^{d+1} differs substantially from that on the cone \mathbb{V}^{d+1} because the distance function is defined differently. We shall verify that the framework for homogeneous space is applicable on this domain for a family of weight functions related to the Gegenbauer weight and the classical weight on the unit ball, following closely the study on the hyperbolic surface. The structure of this section is parallel to that of previous section, with contents arranging in the same order and under similar section names. Furthermore, part of the proof and development follows from the counterpart on \mathbb{X}_0^{d+1} ; hence, the proof is often brief or omitted.

4.1. Distance on the solid double cone and hyperboloid. We write ${}_{\varrho} \mathbb{X}^{d+1}$ whenever it necessary to emphasis the dependence on ϱ , but will use \mathbb{X}^{d+1} most of the time. The domain \mathbb{X}_0^{d+1} can be decomposed as an upper part and a lower part,

$$\mathbb{X}^{d+1} = \mathbb{X}^{d+1}_+ \cup \mathbb{X}^{d+1}_- = \{ (x,t) \in \mathbb{X}^{d+1} : t \ge 0 \} \cup \{ (x,t) \in \mathbb{X}^{d+1} : t \le 0 \}.$$

For $\rho = 0$, the upper part is the solid cone $\mathbb{X}^{d+1}_{+} = \mathbb{V}^{d+1}$.

The distance function on the hyperboloid needs to take into account of the boundary behavior of the domain. Like the case of the hyperbolic surface, the boundary in this case are the intersection of the surface with the hyperplanes t = 1 and t = -1. In particular, the apex point is not considered a boundary point. We first define the distance function on the double cone, that is, when $\rho = 0$.

Definition 4.1. Let $\varrho = 0$. For $(x, t), (y, s) \in \mathbb{X}^{d+1}$, define

$$\mathsf{d}_{\mathbb{X}}((x,t),(y,s)) = \arccos\Big(\langle x,y \rangle + \sqrt{t^2 - \|x\|^2}\sqrt{s^2 - \|y\|^2} + \sqrt{1 - t^2}\sqrt{1 - s^2}\Big).$$

Then $d_{\mathbb{X}}(\cdot, \cdot)$ is a distance function on the double cone \mathbb{X}^{d+1} .

Let $X = (x, \sqrt{t^2 - \|x\|^2})$ and $Y = (y, \sqrt{s^2 - \|y\|^2})$, then $|t|^{-1}X$ and $|s|^{-1}Y$ belong to \mathbb{B}^{d+1} , so that $(X, t), (Y, s) \in \mathbb{X}_0^{d+2}$ and

(4.1)
$$\mathsf{d}_{\mathbb{X}^{d+1}}((x,t),(y,s)) = \mathsf{d}_{\mathbb{X}^{d+2}_0}((X,t),(Y,s)).$$

In particular, it follows that $d_{\mathbb{X}}(\cdot, \cdot)$ defines a distance on the solid double cone \mathbb{X}^{d+1} .

This distance function, however, is difference from the distance $\mathsf{d}_{\mathbb{V}}(\cdot, \cdot)$ defined in for \mathbb{V}^{d+1} in [30]. It is closely related to the distance functions $\mathsf{d}_{[-1,1]}(\cdot, \cdot)$ of [-1, 1] and the distance function on \mathbb{B}^d defined by

(4.2)
$$\mathsf{d}_{\mathbb{B}}(x',y') := \arccos\left(\langle x',y'\rangle + \sqrt{1 - \|x'\|^2}\sqrt{1 - \|y'\|^2}\right), \quad x',y' \in \mathbb{B}^d.$$

Proposition 4.2. Let $\rho = 0$ and $d \ge 2$. For $(x, t), (y, s) \in \mathbb{X}^{d+1}$, write x = tx' and y = sy' with $x', y' \in \mathbb{B}^d$. Then

(4.3)
$$1 - \cos \mathsf{d}_{\mathbb{X}}((x,t),(y,s)) = 1 - \cos \mathsf{d}_{[-1,1]}(t,s) + ts \left(1 - \cos \mathsf{d}_{\mathbb{B}}(x',y')\right).$$

In particular, if t and s have the same sign, then

(4.4)
$$c_1 \mathsf{d}_{\mathbb{X}}((x,t),(y,s)) \le \mathsf{d}_{[-1,1]}(t,s) + \sqrt{ts} \,\mathsf{d}_{\mathbb{B}}(x',t') \le c_2 \mathsf{d}_{\mathbb{X}}((x,t),(y,s)).$$

The proof is similar to that of Proposition 3.2, using (3.2) and (4.2). In particular, it follows that the distance on the line segment $l_x = \{(tx', t) : -1 \leq t \leq 1\}$, where $x' \in \mathbb{B}^d$, of the double cone becomes $\mathsf{d}_{[-1,1]}(t,s)$ as expected. We will also need the following lemma

Lemma 4.3. Let $\varrho = 0$ and $d \ge 2$. For (x,t), (y,s) either both in \mathbb{X}^{d+1}_+ or both in \mathbb{X}^{d+1}_- ,

$$\left|t-s\right| \leq \mathsf{d}_{\mathbb{X}}((x,t),(y,s)) \quad and \quad \left|\sqrt{1-t^2} - \sqrt{1-s^2}\right| \leq \mathsf{d}_{\mathbb{X}}((x,t),(y,s)).$$

and

$$\left|\sqrt{t^2 - \|x\|^2} - \sqrt{s^2 - \|y\|^2}\right| \le (\sqrt{2} + \pi) \mathsf{d}_{\mathbb{X}}((x, t), (y, s)).$$

Proof. We consider only $t, s \ge 0$. Let x = tx' and y = sy' with $x', y' \in \mathbb{B}^d$. Setting $t = \cos \theta$ and $s = \cos \phi$. Using the inequality $|\langle x', y' \rangle + \sqrt{1 - ||x'||^2} \sqrt{1 - ||y'||^2}| \le 1$, it follows readily that $\cos \mathsf{d}_{\mathbb{X}_0}((x,t),(y,s)) \le ts + \sqrt{1 - t^2} \sqrt{1 - s^2} = \cos(\theta - \phi)$, which allows us to follow the proof of Lemma 3.3 to establish the first two inequalities. For third inequality, we assume without loss of generality that $|t| \ge |s|$. Then,

$$\begin{split} \left|\sqrt{t^2 - \|x\|^2} - \sqrt{s^2 - \|y\|^2}\right| &= \left| |t|\sqrt{1 - \|x'\|^2} - |s|\sqrt{1 - \|y'\|^2} \right| \\ &\leq |t - s|\sqrt{1 - \|x'\|^2} + |s| \left|\sqrt{1 - \|x'\|^2} - \sqrt{1 - \|y'\|^2} \right| \\ &\leq \sqrt{2} \mathsf{d}_{\mathbb{X}}((x, t), (y, s)) + \sqrt{ts} \,\mathsf{d}_{\mathbb{B}}(x', y'), \end{split}$$

where the last step uses the inequality [8, (A.1.4)]

$$\left|\sqrt{1 - \|x'\|^2} - \sqrt{1 - \|y'\|^2}\right| \le \sqrt{2} \mathsf{d}_{\mathbb{B}}(x', y').$$

Hence, the third inequality follows from (4.4). This completes the proof.

For $\rho > 0$, the two parts of the solid hyperboloid, \mathbb{X}^{d+1}_+ and \mathbb{X}^{d+1}_- , are disjoint. It is sufficient to consider the distance between points that lie in the same part. We define

(4.5)
$$d_{\mathbb{X}}^{\varrho}((x,t),(y,s)) = \arg \left(\langle x,y \rangle + \sqrt{1 - \|x\|^2} \sqrt{1 - \|y\|^2} + \sqrt{1 + \varrho^2 - t^2} \sqrt{1 + \varrho^2 - s^2} \right).$$

If (x,t) and (y,t) are both in $\mathbb{X}+^{d+1}$ or both in \mathbb{X}_{-}^{d+1} , then

(4.6)
$$\mathsf{d}^{\varrho}_{\mathbb{X}}((x,t),(y,s)) = \mathsf{d}_{\mathbb{X}}\left(\left(x,\sqrt{t^2-\varrho^2}\right),\left(y,\sqrt{s^2-\varrho^2}\right)\right).$$

It is easy to see that this is a distance function and, evidently, its properties follows from those of the distance on the double cone.

4.2. A family of doubling weights. For $d \ge 2$, $\beta > -\frac{1}{2}$, $\gamma > -\frac{1}{2}$ and $\mu > -\frac{1}{2}$, let $W^{\varrho}_{\beta,\gamma,\mu}$ be the weight function defined on the hyperboloid \mathbb{X}^{d+1} by

(4.7)
$$W^{\varrho}_{\beta,\gamma,\mu}(x,t) = \mathbf{b}^{\varrho}_{\beta,\gamma,\mu} |t| (t^2 - \varrho^2)^{\beta - \frac{1}{2}} (1 + \varrho^2 - t^2)^{\gamma - \frac{1}{2}} (t^2 - \varrho^2 - ||x||^2)^{\mu - \frac{1}{2}}.$$

When $\rho = 0$ or the double cone, the weight function becomes

$$W^{0}_{\beta,\gamma,\mu}(x,t) = \mathbf{b}^{0}_{\beta,\gamma,\mu}|t|^{2\beta}(1-t^{2})^{\gamma-\frac{1}{2}}(t^{2}-\|x\|^{2})^{\mu-\frac{1}{2}},$$

which remains integrable over \mathbb{X}^{d+1} if $\beta > -\frac{d+1}{2}$. Using the identity

(4.8)
$$\int_{\varrho \mathbb{X}^{d+1}} f(x,t) |t| dx dt = \int_{\varrho \le |t| \le \varrho+1} |t| \int_{\|x\| \le \sqrt{t^2 - \varrho^2}} f(x,t) dx dt$$
$$= \int_{|s| \le 1} |s| \int_{\|x\| \le |s|} f(x, \sqrt{s^2 + \varrho^2}) dy ds$$
$$= \int_{0 \mathbb{X}^{d+1}} f(x, \sqrt{s^2 + \varrho^2}) |s| dx ds,$$

it is easy to see that the normalization constant $\mathbf{b}^{\varrho}_{\beta,\gamma,\mu}$ of $W^{\varrho}_{\beta,\gamma,\mu}(x,t)$ satisfies

$$\mathbf{b}^{\varrho}_{\beta,\gamma,\mu} = \mathbf{b}^{0}_{\beta,\gamma,\mu} = b^{\mathbb{B}}_{\mu} \frac{\Gamma(\beta + \mu + \gamma + \frac{d+1}{2})}{\Gamma(\beta + \mu \frac{d}{2})\Gamma(\gamma + \frac{1}{2})},$$

where $b^{\mathbb{B}}_{\mu}$ is the normalization constant for the weigt function ϖ_{μ} in (2.18) on \mathbb{B}^d .

For r > 0 and (x, t) on the solid hyperboloid ${}_{\varrho}\mathbb{X}^{d+1}$, we denote the ball centered at (x, t) with radius r by

$$\mathbf{c}_{\varrho}((x,t),r) := \left\{ (y,s) \in \mathbb{X}^{d+1} : \mathsf{d}_{\mathbb{X}}^{\varrho}\big((x,t),(y,s)\big) \le r \right\}$$

The following lemma can be proved similarly as that of Lemma 3.4.

Lemma 4.4. For $\rho > 0$ and $(x, t) \in \mathbb{X}^{d+1}$,

$$W^{\varrho}_{\beta,\gamma,\mu} \left(\mathbf{c}_{\varrho}((x,t),r) \right) = W^{0}_{\beta,\gamma\mu} \left(\mathbf{c}_{0} \left(\left(x, \sqrt{t^{2} - \varrho^{2}} \right), r \right) \right)$$

Proposition 4.5. Let r > 0 and $(x,t) \in \mathbb{X}^{d+1}$. Then for $\beta > -\frac{d+1}{2}$ and $\gamma > -1$ and $\mu \ge 0$,

(4.9)
$$W^{0}_{\beta,\gamma,\mu} (\mathbf{c}_{0}((x,t),r)) := \mathbf{b}^{0}_{\beta,\gamma,\mu} \int_{\mathbf{c}_{0}((x,t),r)} W^{0}_{\beta,\gamma,\mu}(y,s) \mathrm{d}y \mathrm{d}s$$
$$\sim r^{d+1} \left(t^{2} + r^{2}\right)^{\beta} \left(1 - t^{2} + r^{2}\right)^{\gamma} \left(t^{2} - \|x\|^{2} + r^{2}\right)^{\mu}$$

In particular, $W^0_{\beta,\gamma,\mu}$ is a doubling weight on the double cone and the doubling index is given by $\alpha(W^0_{\beta,\gamma,\mu}) = d + 1 + 2\mu + 2\max\{0,\beta\} + 2\max\{0,\gamma\}.$

Proof. Let $\tau_r(t,s)$ and $\theta_r(t,s) = \arccos \tau_r(t,s)$ be as in the proof of Proposition 3.5. From $\mathsf{d}_{\mathbb{X}}((x,t),(y,s)) \leq r$, we obtain $\mathsf{d}_{[-1,1]}(t,s) \leq r$ and, by (4.1)

$$\mathsf{d}_{\mathbb{B}}(x',y') \le \arccos\left(2[\tau_r(t,s)]^2 - 1\right) = \frac{1}{2}\arccos\tau_r(t,s) = \frac{1}{2}\theta_r(t,s),$$

where $\mathsf{d}_{\mathbb{B}}(\cdot, \cdot)$ is the distance on the unit ball \mathbb{B}^d . Hence, it follows that

$$W^{0}_{\beta,\gamma,\mu} (\mathbf{c}_{0}((x,t),r)) = \mathbf{b}^{0}_{\beta,\gamma,\mu} \int_{\mathsf{d}_{[-1,1]}(t,s) \leq r} s^{d} \int_{\mathsf{d}_{\mathbb{B}}(x',y') \leq \frac{1}{2}\theta_{r}(t,s)} W^{0}_{\beta,\gamma,\mu}(y,s) \mathrm{d}y \mathrm{d}s$$
$$\sim \int_{\mathsf{d}_{[-1,1]}(t,s) \leq r} s^{2\beta+2\mu+d-1} (1-s^{2})^{\gamma-\frac{1}{2}} \mathrm{d}s \int_{\mathsf{d}_{\mathbb{B}}(x',y') \leq \frac{1}{2}\theta_{r}(t,s)} (1-\|y'\|^{2})^{\mu-\frac{1}{2}} \mathrm{d}y.$$

For $\mu \ge 0$ and $0 < \rho \le 1$, it is known [19, Lemma 5.3] or [8, p. 107] that

$$\int_{\mathsf{d}_{\mathbb{B}}(x',y') \le \rho} (1 - \|y'\|^2)^{\mu - \frac{1}{2}} \mathrm{d}y' \sim (1 - \|x'\|^2 + \rho^2)^{\mu} \rho^d,$$

which implies, together with $\theta_r(t,s) \sim \sqrt{1-\tau_r, t, s}$, that

$$W^{0}_{\beta,\gamma,\mu}(\mathbf{c}_{0}((x,t),r)) \sim \int_{\mathsf{d}_{[-1,1]}(t,s) \leq r} s^{2\beta+2\mu+d-1} (1-s^{2})^{\gamma-\frac{1}{2}} \times (1-\|x'\|^{2}+1-\tau_{r}(t,s))^{\mu} (1-\tau_{r}(t,s))^{\frac{d}{2}} \mathrm{d}s.$$

If $t \ge 3r$, then $s \sim t + r$ and, by $1 - \tau_r(t, s) = (\cos \mathsf{d}_{[-1,1]}(t, s) - \cos r)/(ts)$, it follows that

$$s^{2\mu} (1 - ||x'||^2 + 1 - \tau_r(t,s))^{\mu} \sim (t^2 - ||x||^2 + r^2)^{\mu}.$$

With this term removed, the integral of the remaining integrand can be estimated by following the estimates of Case 1 and Case 3 of the proof of Proposition 3.5. If $t \leq 3r$, then $t^2 - ||x||^2 + r^2 \sim r^2$. We use $1 - ||x'||^2 + 1 - \tau_r(t,s) \leq 2$ for the upper bound and $1 - \tau_r(t,s) \geq 1/(8\pi^2)$ on the subset $\mathsf{d}_{[-1,1]}(t,s) \leq r/2$, proved in the Case 2 of the proof of Proposition 3.5, to remove the term $(1 - ||x'||^2 + 1 - \tau_r(t,s))^{\mu}$ from the integral. The rest of the proof then follows from that of the Case 2 of the proof of Proposition 3.5. This completes the proof.

Corollary 4.6. For $d \geq 2$, $\beta > -\frac{d+1}{2}$ and $\gamma > -\frac{1}{2}$, the space $(\mathbb{X}^{d+1}, W^0_{\beta,\gamma,\mu}, \mathsf{d}_{\mathbb{X}})$ is a homogeneous space.

When $\rho = 0$, $\beta = 0$, $\mu = 0$ and $\gamma = \frac{1}{2}$, the relation (4.9) is for the Lebesgue measure d**m** on the double cone,

$$\mathbf{m}(\mathbf{c}_0((x,t),r)) \sim r^{d+1}(1-t^2+r^2)^{\frac{1}{2}}(t^2-\|x\|^2+n^{-2})^{\frac{1}{2}}.$$

Furthermore, $W_{0,0,0}^0(\mathbf{c}_0((x,t),r)) \sim r^{d+1}$ and $W_{0,0,0}^0(x,y) = (1-t^2)^{-\frac{1}{2}}(t^2 - ||x||^2)^{-\frac{1}{2}}$.

4.3. Orthogonal polynomials on the hyperboloid. With respect to the weight function $W^{\varrho}_{\beta,\gamma,\mu}$, we defined the inner product

$$\langle f,g \rangle_W = \int_{\mathbb{X}^{d+1}} f(x,t)g(x,t)W^{\varrho}_{\beta,\gamma,\mu}(x,t)\mathrm{d}x\mathrm{d}t.$$

Let $\mathcal{V}_n(\mathbb{X}^{d+1}, W^{\varrho}_{\beta,\gamma,\mu})$ be the space of these orthogonal polynomials of degree n, which has the dimension dim $\mathcal{V}_n(\mathbb{X}^{d+1}, W) = \binom{n+d}{n}$. Like the decomposition on the surface \mathbb{X}_0^{d+1} , this space satisfies

$$\mathcal{V}_n\left(\mathbb{X}^{d+1}, W^{\varrho}_{\beta, \gamma, \mu}\right) = \mathcal{V}_n^E\left(\mathbb{X}^{d+1}, W^{\varrho}_{\beta, \gamma, \mu}\right) \bigoplus \mathcal{V}_n^O\left(\mathbb{X}^{d+1}, W^{\varrho}_{\beta, \gamma, \mu}\right),$$

where the subspace $\mathcal{V}_n^E(\mathbb{X}^{d+1}, W_{\beta,\gamma,\mu}^{\varrho})$ consists of orthogonal polynomials that are even in t variable, whereas the subspace $\mathcal{V}_n^O(\mathbb{X}^{d+1}, W_{\beta,\gamma,\mu}^{\varrho})$ consists of orthogonal polynomials that are odd in t variable.

An orthogonal basis can be given explicitly in terms of the Jacobi polynomials and classical orthogonal polynomials on the unit ball for the subspace $\mathcal{V}_n^E(\mathbb{X}^{d+1}, W_{\beta,\gamma,\mu}^{\varrho})$ for all $\varrho \geq 0$, but for the subspace $\mathcal{V}_n^O(\mathbb{X}^{d+1}, W_{\beta,\gamma,\mu}^{\varrho})$ only when $\varrho = 0$. For example, let $\{P_{\mathbf{k}}^{n-2k} : |\mathbf{k}| = n - 2k, \mathbf{k} \in \mathbb{N}_0^d\}$ denote an orthonormal basis of $\mathcal{V}_{n-2k}(\mathbb{B}^d, W_{\mu})$. Then the polynomials

(4.10)
$$\mathbf{C}_{n-2k,\mathbf{k}}^{n}(x,t) = P_{k}^{(\gamma-\frac{1}{2},n-2k+\beta+\mu+\frac{d-2}{2})}(2t^{2}-2\varrho^{2}-1) \times (t^{2}-\varrho^{2})^{\frac{n-2k}{2}}P_{\mathbf{k}}^{n-2k}\left(\frac{x}{\sqrt{t^{2}-\varrho^{2}}}\right)$$

with $|\mathbf{k}| = n - 2k$ and $0 \le k \le n/2$ form an orthogonal basis of $\mathcal{V}_n^E(\mathbb{V}^{d+1}, W_{\beta,\gamma,\mu}^{\varrho})$. We call these polynomials generalized Gegenbauer polynomials on the solid hyperboloid. We will not work with the basis directly; see [29] for the basis in other cases.

Let $\mathbf{P}_{n}^{E}(W_{\beta,\gamma,\mu}^{\varrho};\cdot,\cdot)$ denote the reproducing kernel of $\mathcal{V}_{n}^{E}(\mathbb{X}^{d+1}, W_{\beta,\gamma,\mu}^{\varrho})$. In terms of the basis of (4.10), we can write

$$\mathbf{P}_{n}^{E}(W_{\beta,\gamma,\mu}^{\varrho};(x,t),(y,s)) = \sum_{m=0}^{n} \sum_{|\mathbf{k}|=m} \frac{\mathbf{C}_{n-2k,\mathbf{k}}^{n}(x,t)\mathbf{C}_{n-2k,\mathbf{k}}^{n}(y,s)}{\langle \mathbf{C}_{n-2k,\mathbf{k}}^{n}, \mathbf{C}_{n-2k,\mathbf{k}}^{n} \rangle_{W_{\beta,\gamma,\mu}}}.$$

Just like on the surface \mathbb{X}_0^{d+1} , the kernel $\mathbf{P}_n^E(W_{\gamma,\mu}^{\varrho};\cdot,\cdot)$ can be used to study the Fourier orthogonal series of any function f that is even in the variable t on \mathbb{X}^{d+1} . For such a function, f(x,t) = f(x,-t), its projection on $\mathcal{V}_n^O(\mathbb{X}^{d+1}, W_{\beta,\gamma,\mu}^{\varrho})$ becomes zero, so that its Fourier orthogonal expansion is given by

(4.11)
$$f = \sum_{n=0}^{\infty} \operatorname{proj}_{n}^{E}(W_{\beta,\gamma,\mu}^{\varrho}; f),$$

where the projection $\operatorname{proj}_{n}^{E}\left(W_{\beta,\gamma,\mu}^{\varrho}\right): L^{2}\left(\mathbb{X}^{d+1}, W_{\beta,\gamma,\mu}^{\varrho}\right) \mapsto \mathcal{V}_{n}^{E}\left(\mathbb{X}^{d+1}, W_{\beta,\gamma,\mu}^{\varrho}\right)$ can be written in terms of the kernel $\mathbf{P}_{n}^{E}(W_{\beta,\gamma,\mu}^{\varrho}; \cdot, \cdot)$ as

$$\operatorname{proj}_{n}(W^{\varrho}_{\beta,\gamma,\mu};f) = c_{\mathsf{w}} \int_{\mathbb{X}^{d+1}} f(y) \mathbf{P}^{E}_{n}(W^{\varrho}_{\beta,\gamma,\mu};\cdot,(y,s)) W^{\varrho}_{\beta,\gamma,\mu}(s) \mathrm{d}y \mathrm{d}s.$$

Moreover, since f is even in t variable, we can regard it as the even extension of a function f defined on the upper hyperboloid \mathbb{X}^{d+1}_+ , which is the cone \mathbb{V}^{d+1} when $\rho = 0$. In particular, this provides a Fourier orthogonal series for functions on the cone \mathbb{V}^{d+1} , which is, however, different from the Fourier orthogonal series in the Jacobi polynomials on the cone discussed in [30].

polynomials on the cone discussed in [30]. The most interesting case on \mathbb{X}^{d+1} is $\beta = \frac{1}{2}$. To simplify the notation, we shall denote $W^{\varrho}_{\frac{1}{2},\gamma,\mu}$ by $W^{\varrho}_{\gamma,\mu}$ throughout the rest of the section; that is,

$$W^{\varrho}_{\gamma,\mu}(x,t) := |t|(1+\varrho^2-t^2)^{\gamma-\frac{1}{2}}(t^2-\varrho^2-||x||^2)^{\mu-\frac{1}{2}}.$$

The orthogonal polynomials with respect to $W_{\gamma,\mu}$ also possess two characteristic properties: the first one is the spectral operator that has orthogonal polynomials as eigenfunctions [29, Theorem 4.8].

Theorem 4.7. Let $\varrho \ge 0$, $\gamma, \mu > -\frac{1}{2}$. Define the differential operator

$$\begin{split} \mathfrak{D}^{\varrho}_{\gamma,\mu} &:= (1+\varrho^2-t^2) \left(1-\frac{\varrho^2}{t^2}\right) \partial_t^2 + \Delta_x - \langle x, \nabla_x \rangle^2 + \langle x, \nabla_x \rangle \\ &+ \frac{2}{t} (1+\varrho^2-t^2) \langle x, \nabla_x \rangle \partial_t + \left((1+\varrho^2-t^2)\frac{\varrho^2}{t^2} + 2\mu + d\right) \frac{1}{t} \partial_t \\ &- (2\gamma+2\mu+d+1) \left(\left(1-\frac{\varrho^2}{t^2}\right) t \partial_t + \langle x, \nabla_x \rangle\right) \end{split}$$

Then the polynomials in $\mathcal{V}_n^E(\mathbb{X}^{d+1}, W_{\gamma,\mu}^{\varrho})$ are eigenfunctions of $\mathfrak{D}_{\gamma,\mu}^{\varrho}$,

(4.12) $\mathfrak{D}^{\varrho}_{\gamma,\mu}u = -n(n+2\gamma+2\mu+d)u, \qquad \forall u \in \mathcal{V}^{E}_{n}\left(\mathbb{X}^{d+1}, W^{\varrho}_{\gamma,\mu}\right)$

The second one is the addition formula for the reproducing kernel $\mathbf{P}_{n}^{E}\left(\mathbf{w}_{\beta,\gamma,\mu}^{\varrho};\cdot,\cdot\right)$, which is of the simplest form when $\beta = \frac{1}{2}$ [29, Prop. 5.7 and Cor. 5.6].

Theorem 4.8. Let $d \ge 2$ and $\varrho \ge 0$. Then

(a) For
$$\beta, \gamma, \mu > -\frac{1}{2}$$
,
(4.13) $\mathbf{P}_{n}^{E} \left(W_{\beta,\gamma,\mu}^{\varrho}; (x,t), (y,s) \right) = \mathbf{P}_{n}^{E} \left(W_{\beta,\gamma,\mu}^{0}; \left(x, \sqrt{t^{2} - \varrho^{2}} \right), \left(y, \sqrt{s^{2} - \varrho^{2}} \right) \right)$.
(b) For $\varrho = 0$, and $\gamma, \mu \ge 0$,

(4.14)
$$\mathbf{P}_{n}^{E}\left(W_{\gamma,\mu}^{0};(x,t),(y,s)\right) = b_{\gamma,\mu} \int_{-1}^{1} \int_{-1}^{1} Z_{n}^{\gamma+\mu+\frac{d}{2}}\left(\xi(x,t,y,s;u,v)\right) \times (1-v^{2})^{\gamma-1}(1-u^{2})^{\mu-1} \mathrm{d}u \mathrm{d}v,$$

where
$$b_{\gamma,\mu} = c_{\gamma-1,\gamma-1}c_{\mu-1,\mu-1}$$
 with $c_{a,b}$ defined as in (2.10) and

$$\xi(x,t,y,s;u,v) = \left(\langle x,y \rangle + u\sqrt{t^2 - \|x\|^2}\sqrt{s^2 - \|y\|^2}\right)\operatorname{sign}(st) + v\sqrt{1 - s^2}\sqrt{1 - t^2},$$

and the identity (4.14) holds under the limit when $\mu = 0$ or $\gamma = 0$.

4.4. Highly localized kernels. Let \hat{a} be a cut-off function. For $(x, t), (y, s) \in \mathbb{X}^{d+1}$, the localized kernel $\mathbf{L}_{n}^{E}(W_{\gamma,\mu}^{\varrho}; \cdot, \cdot)$ is defined by

$$\mathbf{L}_{n}^{E}(W_{\gamma,\mu}^{\varrho};(x,t),(y,s)) = \sum_{j=0}^{\infty} \widehat{a}\left(\frac{j}{n}\right) \mathbf{P}_{j}^{E}(W_{\gamma,\mu}^{\varrho};(x,t),(y,s)).$$

We show that this kernel is highly localized when (x,t) and (y,s) are either both in \mathbb{X}^{d+1}_+ or both in \mathbb{X}^{d+1}_- . For $\mu, \gamma \geq 0$, define

$$W^{\varrho}_{\gamma,\mu}(n;x,t) := \left(1 + \varrho^2 - t^2 + n^{-2}\right)^{\gamma} \left(t^2 - \varrho^2 - \|x\|^2 + n^{-2}\right)^{\mu}.$$

Theorem 4.9. Let $d \ge 2$, $\mu, \gamma \ge 0$. Let \hat{a} be an admissible cutoff function. Then for any $\kappa > 0$, and (x, t), (y, s) either both in \mathbb{X}^{d+1}_+ or both in \mathbb{X}^{d+1}_- ,

$$\left|\mathbf{L}_{n}^{E}(W_{\gamma,\mu}^{\varrho};(x,t),(y,s))\right| \leq \frac{c_{\kappa}n^{d+1}}{\sqrt{W_{\gamma,\mu}^{\varrho}(n;x,t)}\sqrt{W_{\gamma,\mu}^{\varrho}(n;y,s)}} \left(1 + n\mathsf{d}_{\mathbb{X}}((x,t),(y,s))\right)^{-\kappa}.$$

Proof. Again, it is sufficient to consider the case $\rho = 0$ and we shall be brief. By (4.14) we can write $\mathbf{L}_{n}^{E}(W_{\gamma,\mu}^{0})$ in terms of the kernel for the Jacobi polynomials. Let $\lambda = \gamma + \mu + \frac{d}{2}$. Then

$$\begin{split} \mathbf{L}_{n}^{E} \big(W_{\gamma,\mu}^{0};(x,t),(y,s) \big) &= b_{\gamma,\mu} \int_{-1}^{1} \int_{-1}^{1} L_{n}^{(\lambda - \frac{1}{2}, \lambda - \frac{1}{2})} \big(\xi(x,t,y,s;u,v) \big) \\ &\times (1 - v^{2})^{\gamma - 1} (1 - u^{2})^{\mu - 1} \mathrm{d} u \mathrm{d} v. \end{split}$$

Hence, applying (2.12) with m = 1 and $\alpha = \beta = \lambda - 1/2$, we obtain

$$\begin{aligned} \left| \mathbf{L}_{n}^{E}(W_{\gamma,\mu}^{0};(x,t),(y,s)) \right| &\leq c n^{2\lambda+1} \int_{-1}^{1} \int_{-1}^{1} \frac{1}{\left(1 + n\sqrt{1 - \xi(x,t,y,s;u,v)}\right)^{\kappa+3\gamma+3\mu+2}} \\ &\times (1 - v^{2})^{\gamma-1} (1 - u^{2})^{\mu-1} \mathrm{d} u \mathrm{d} v. \end{aligned}$$

Since t and s have the same sign, it is easy to verify that

$$\begin{aligned} 1 - \xi(x, t, y, s; u, v) &= 1 - \cos \mathsf{d}_{\mathbb{X}}((x, t), (y, s)) \\ &+ (1 - u)\sqrt{t^2 - \|x\|^2}\sqrt{s^2 - \|y\|^2} + (1 - v)\sqrt{1 - t^2}\sqrt{1 - s^2}. \end{aligned}$$

The entries in both two lines in the right-hand side of the above identity are lower bounds of $1 - \xi(x, t, u, s; u, v)$. Using the first one, we obtain the estimate

$$\left|\mathbf{L}_{n}^{E}(W_{\gamma,\mu}^{0};(x,t),(y,s))\right| \leq cn^{2\lambda+1} \frac{1}{(1+n\mathsf{d}_{\mathbb{X}}((x,t),(y,s)))^{\kappa+\gamma+\mu}} I(x,t,y,s),$$

where, using the second one and the symmetry of the integral, the integral I(x, t, y, s) is given by

$$I(x,t,y,s) = c_{\gamma-\frac{1}{2},\mu-\frac{1}{2}} \int_0^1 \int_0^1 \frac{(1-u^2)^{\mu-1}(1-v^2)^{\gamma-1}}{\left(1+n\sqrt{A(1-u)}+B(1-v)\right)^{2\gamma+2\mu+2}} \mathrm{d}u \mathrm{d}v$$

with $A = \sqrt{t^2 - \|x\|^2}\sqrt{s^2 - \|y\|^2}$ and $B = \sqrt{1 - t^2}\sqrt{1 - s^2}$. The integral I(x, t, y, s) is bounded by 1 and it can also be bounded by applying (3.18) twice. Carrying out the estimates, we conclude that

$$\begin{split} I(x,t,y,s) &\leq c \frac{n^{-2\gamma-2\mu}}{(A+n^{-1})^{\mu} (B+n^{-1})^{\gamma}} \\ &\leq c \frac{n^{-2\gamma-2\mu}}{\sqrt{W_{\gamma,\mu}^{0}(n;t)} \sqrt{W_{\gamma,\mu}^{0}(n;t)}} \Big(1 + n \mathsf{d}_{\mathbb{X}}((x,t),(y,s))\Big)^{\gamma+\mu} \end{split}$$

where the second inequality follows from (3.19) and Lemma 4.3. Putting the last two displayed inequalities together, we have established (ii).

This establishes Assertion 1. The next theorem establishes Assertion 2.

Theorem 4.10. Let $d \geq 2$, $\mu, \gamma \geq -\frac{1}{2}$. Then for (x_i, t_i) and (y, s) that are either all in \mathbb{X}^{d+1}_+ or all in \mathbb{X}^{d+1}_- , $(x_1, t_1) \in \mathbf{c}_{\varrho}((x_2, t_2), c_*n^{-1})$ with c_* small and any $\kappa > 0$,

$$\begin{split} \left| \mathbf{L}_{n}^{E}(W_{\gamma,\mu}^{\varrho};(x_{1},t_{1}),(y,s)) - \mathbf{L}_{n}^{E}(W_{\gamma,\mu}^{\varrho};(x_{2},t_{2}),(y,s)) \right| \\ & \leq \frac{c_{\kappa}n^{d+1}\mathbf{d}_{\mathbb{X}}^{\varrho}((x_{1},t_{1}),(x_{2},t_{2}))}{\sqrt{W_{\gamma,\mu}^{\varrho}(n,t_{2})}\sqrt{W_{\gamma,\mu}^{\varrho}(n;s)} \big(1 + n\mathbf{d}_{\mathbb{X}}^{\varrho}((x_{2},t_{2}),(y,s))\big)^{\kappa}} \end{split}$$

Proof. Again it suffices to prove the case $\rho = 0$. Let $\xi_i(u, v) = \xi(x_i, t_i, y, s; u, v)$. From (4.3), it is easy to see that

$$\begin{split} \xi_1(u,v) - \xi_2(u,v) &= \cos \mathsf{d}_{\mathbb{X}}((x_1,t_1),(y,s)) - \cos \mathsf{d}_{\mathbb{X}}((x_2,t_2),(y,s)) \\ &+ (1-u) \left(\sqrt{t_2^2 - \|x_2\|^2} - \sqrt{t_1^2 - \|x_1\|^2} \right) \sqrt{s^2 - \|y\|^2} \\ &+ (1-v) \left(\sqrt{1-t_2^2} - \sqrt{1-t_1^2} \right) \sqrt{1-s^2}. \end{split}$$

By Lemma 4.3 and the proof of Theorem 3.10, this leads to

$$|\xi_1(u,v) - \xi_2(u,v)| \le \mathsf{d}_{\mathbb{X}}\big((x_1,t_1),(x_2,t_2)\big)\big[\Sigma_1 + \Sigma_2(u) + \Sigma_3(v)\big],$$

where

$$\begin{split} \Sigma_1 &= \mathsf{d}_{\mathbb{X}}((x_2, t_2), (y, s)) + \mathsf{d}_{\mathbb{X}}((x_1, t_1), (x_2, t_2)) \\ \Sigma_2(u) &= (1 - u)\sqrt{s^2 - \|y\|^2}, \\ \Sigma_3(v) &= (1 - v)\sqrt{1 - s^2}. \end{split}$$

Hence, following the proof of Theorem 3.10, we see that, with $\lambda = \gamma + \mu + \frac{d}{2}$,

$$\begin{aligned} \left| \mathbf{L}_{n}(W_{\gamma,\mu}^{0};(x_{1},t_{1}),(y,s)) - \mathbf{L}_{n}(W_{\gamma,\mu}^{0};(x_{2},t_{2}),(y,s)) \right| \\ &\leq c \, \mathsf{d}_{\mathbb{X}}\big((x_{1},t_{1}),(x_{2},t_{2})\big) \int_{-1}^{1} \int_{-1}^{1} \left[\frac{n^{2\lambda+3}}{\left(1 + n\sqrt{1 - \xi_{1}(u,v)^{2}}\right)^{\kappa}} + \frac{n^{2\lambda+3}}{\left(1 + n\sqrt{1 - \xi_{2}(u,v)^{2}}\right)^{\kappa}} \right] \\ &\times \big(\Sigma_{1} + \Sigma_{2}(u) + \Sigma_{3}(v)\big)(1 - u^{2})^{\mu-1}(1 - v^{2})^{\gamma-1} \mathrm{d} u \mathrm{d} v. \end{aligned}$$

The integrals that contain Σ_1 and $\Sigma_3(v)$ can be estimated exactly as in the proof of Theorem 3.10. The integral that contains $\Sigma_2(u)$ does not cause additional problem and can be handled just as the integral containing $\Sigma_3(v)$. We omit the details. \Box

The case p = 1 of the following lemma establishes Assertion 3 for $W^{\varrho}_{\gamma,\mu}$.

Lemma 4.11. Let $d \ge 2$, $\gamma > -\frac{1}{2}$ and $\mu > -\frac{1}{2}$. For $0 , assume <math>\kappa > \frac{2d+2}{p} + 2(\beta + \gamma)|\frac{1}{p} - \frac{1}{2}|$. Then for $(x, t) \in \mathbb{X}^{d+1}$,

$$\int_{\mathbb{X}^{d+1}} \frac{W_{\gamma,\mu}^{\varrho}(y,s)}{W_{\gamma,\mu}^{\varrho}(n;y,s)^{\frac{p}{2}} \left(1 + n \mathsf{d}_{\mathbb{X}}^{\varrho}((x,t),(y,s))\right)^{\kappa p}} \mathrm{d}y \mathrm{d}s \le c n^{-d-1} W_{\gamma,\mu}^{\varrho}(n;x,t)^{1-\frac{p}{2}}.$$

Proof. Again, it suffices to consider $\varrho = 0$. Let J_p denote the left-hand side the inequality to be proved. As in the case of \mathbb{X}_0 , we only need to estimate J_2 . Furthermore, following the proof of Lemma 3.11, we only need to estimate the integral in J_2 over either \mathbb{X}^{d+1}_+ or \mathbb{X}^{d+1}_- , which we choose as $\mathbb{X}^{d+1}_+ = \mathbb{V}^{d+1}$ and denote it by $J_{2,+}$. Then

$$J_{2,+} = \int_0^1 s^d \int_{\mathbb{B}^d} \frac{W^0_{\gamma,\mu}(sy',s)}{W^0_{\gamma,\mu}(n;sy',s)(1 + n\mathsf{d}_{\mathbb{X}}((x,t),(sy',s)))^{2\kappa}} \mathrm{d}y' \mathrm{d}s$$

Let x = tx' and y = sy' with $x', y' \in \mathbb{B}^d$. Using $w_{0,\gamma}^0(t) = (1-t^2)^{\gamma-\frac{1}{2}}$ and $w_{\beta,\gamma}^0(n;s) = (1-t^2+n^{-2})^{\gamma}$, we can easily verify that

$$\frac{W^0_{\gamma,\mu}(y,s)}{W^0_{\gamma,\mu}(n;y,s)} \leq c \frac{\mathsf{w}^0_{0,\gamma}(s)}{\sqrt{1 - \|y'\|^2}} \mathsf{w}^0_{0,\gamma}(n;s)$$

which leads to

$$J_{2,+} \leq \int_0^1 s^d \int_{\mathbb{B}^d} \frac{\mathsf{w}_{0,\gamma}^0(s)}{\mathsf{w}_{0,\gamma}^0(n;s)(1+n\mathsf{d}_{\mathbb{X}}((x,t),(sy',s)))^{2\kappa}} \frac{\mathrm{d}y'}{\sqrt{1-\|y'\|^2}} \mathrm{d}s.$$

Setting x = tx', $X = (x', \sqrt{1 - \|x'\|^2})$ and $Y = (y', \sqrt{1 - \|y'\|^2})$, so that $d_{\mathbb{B}^d}(x', y') = d_{\mathbb{S}^d}(X, Y)$, we use the identity

(4.15)
$$\int_{\mathbb{B}^d} g\left(y', \sqrt{1 - \|y'\|^2}\right) \frac{dy'}{\sqrt{1 - \|y'\|^2}} = \int_{\mathbb{S}^d_+} g(y) \mathrm{d}\sigma(y) dy$$

where \mathbb{S}^d_+ denotes the upper hemisphere of \mathbb{S}^d , which allows us to follow the proof of Lemma 3.11 to obtain

$$J_{2,+} \le c \int_0^1 \int_{-1}^1 \frac{s^d \mathsf{w}_{0,\gamma}^0(s)(1-u^2)^{\frac{d-2}{2}}}{\mathsf{w}_{0,\gamma}^0(n;s) \left(1+n\sqrt{1-tsu-\sqrt{1-t^2}\sqrt{1-s^2}}\right)^{2\kappa}} \mathrm{d}u \mathrm{d}s.$$

The integral in the right-hand side with d replaced by d + 1 appeared in the proof of Lemma 3.11, and it is bounded by cn^{-d-1} accordingly.

Proposition 4.12. For $\gamma \ge 0$, $\mu \ge 0$ and $(x, t) \in \mathbb{X}^{d+1}$,

$$\int_{\mathbb{X}^{d+1}} \left| \mathbf{L}_n^E \big(W_{\gamma,\mu}^{\varrho}; (x,t), (y,s) \big) \right|^p W_{\gamma,\mu}^{\varrho}(y,s) \mathrm{d}y \mathrm{d}s \le \left(\frac{n^d}{W_{\gamma,\mu}^{\varrho}(n;t)} \right)^{p-1}.$$

This follows by applying Lemma 4.11 on the estimate in Theorem 4.9.

We have established Assertions 1 - 3 for $\mathbf{L}_n^E(W_{\gamma,\mu};\cdot,\cdot)$. The kernel uses, however, only polynomials that are even in t and in s variable.

Corollary 4.13. For $d \ge 2$, $\varrho \ge 0$, $\gamma \ge 0$ and $\mu \ge 0$, the space $(\mathbb{X}^{d+1}, W^{\varrho}_{\gamma,\mu}, \mathsf{d}^{\varrho}_{\mathbb{X}})$ is a localizable homogeneous space, where its localized kernels are defined for polynomials even in t and in s variables.

4.5. Maximal ε -separated sets on the hyperbolic surface. We give a construction of maximal ε -separated on the hyperboloid and the double cone, following the definition of Definition 2.3. Our construction follows the one on \mathbb{X}_0^{d+1} in Section 3.5. We first need ε -separated set on the unit ball \mathbb{B}^d . We adopt the following notation. For $\varepsilon > 0$, we denote by $\Xi_{\mathbb{B}}(\varepsilon)$ a maximal ε -separated set on the unit ball \mathbb{B}^d and we let $\mathbb{B}_u(\varepsilon)$ be the subsets in \mathbb{B}^d so that the collection $\{\mathbb{B}_u(\varepsilon) : u \in \Xi_{\mathbb{B}}(\varepsilon)\}$ is a partition of \mathbb{B}^d , and we assume

(4.16)
$$\mathbf{c}_{\mathbb{B}}(u,c_1\varepsilon) \subset \mathbb{B}_u(\varepsilon) \subset \mathbf{c}_{\mathbb{B}}(u,c_2\varepsilon), \qquad u \in \Xi_{\mathbb{B}}(\varepsilon).$$

where $\mathbf{c}_{\mathbb{B}}(u,\varepsilon)$ denotes the ball centered at u with radius ε in \mathbb{B}^d , c_1 and c_2 depend only on d. It is known (see, for example, [19]) that such a $\Xi_{\mathbb{B}}(\varepsilon)$ exists for all $\varepsilon > 0$ and its cardinality satisfies

(4.17)
$$c'_d \varepsilon^{-d} \le \# \Xi_{\mathbb{B}}(\varepsilon) \le c_d \varepsilon^{-d}.$$

For the hyperboloid \mathbb{X}^{d+1} , we denote by $\Xi_{\mathbb{X}} = \Xi_{\mathbb{X}}(\varepsilon)$ a maximum ε -separated set and, furthermore, denote by $\{\mathbb{X}(u,t) : (tu,t) \in \Xi_{\mathbb{X}}\}$ a partition of \mathbb{X}^{d+1} . We give one construction of such sets below.

Let $\varepsilon > 0$ and let $N = 2\lfloor \frac{\pi}{2}\varepsilon^{-1} \rfloor$. We define $t_j = \cos \theta_j$ and t_j^+ and t_j^- , $1 \le j \le N$, as in Subsection 3.5. Then \mathbb{X}^{d+1} can be partitioned by

$$\mathbb{X}^{(j)} := \left\{ (x,t) \in \mathbb{X}^{d+1} : t_j^- < t \le t_j^+ \right\}, \qquad 1 \le j \le N.$$

Furthermore, the upper and lower hyperboloid \mathbb{X}^{d+1}_+ and \mathbb{X}^{d+1}_- can be partitioned by

$$\mathbb{X}^{d+1}_{+} = \bigcup_{j=1}^{N/2} \mathbb{X}^{(j)} \quad \text{and} \quad \mathbb{X}^{d+1}_{-} = \bigcup_{j=N/2+1}^{N} \mathbb{X}^{(j)}$$

Let $\varepsilon_j := \pi \varepsilon/(2t_j)$. Then $\Xi_{\mathbb{B}}(\varepsilon_j)$ is the maximal ε_j -separated set of \mathbb{B}^d such that, for each $j \ge 1$, $\{\mathbb{B}_u(\varepsilon_j) : u \in \Xi_{\mathbb{B}}(\varepsilon_j)\}$ is a partition of \mathbb{B}^d and $\#\Xi_{\mathbb{B}}(\varepsilon_j) \sim \varepsilon_j^{-d}$. For each $j = 1, \ldots, N$, we decompose $\mathbb{X}^{(j)}$ by

$$\mathbb{X}^{(j)} = \bigcup_{u \in \Xi_{\mathbb{B}}(\varepsilon_j)} \mathbb{X}(u, t_j), \quad \text{where} \quad \mathbb{X}(u, t_j) := \left\{ (tv, t) : t_j^- < t \le t_j^+, v \in \mathbb{B}_u(\varepsilon_j) \right\}.$$

Finally, we define the set $\Xi_{\mathbb{X}}$ of \mathbb{X}^{d+1} by

$$\Xi_{\mathbb{X}} = \left\{ (t_j u, t_j) : u \in \Xi_{\mathbb{B}}(\varepsilon_j), 1 \le j \le N \right\}.$$

Proposition 4.14. Let $\varepsilon > 0$ and $N = 2\lfloor \frac{\pi}{2}\varepsilon^{-1} \rfloor$. Then $\Xi_{\mathbb{X}}$ is a maximal ε -separated set of \mathbb{X}^{d+1} and $\{\mathbb{X}(t_j u, t_j) : u \in \Xi_{\mathbb{B}}(\varepsilon_j), 1 \leq j \leq N\}$ is a partition

$$\mathbb{X}^{d+1} = \bigcup_{(tu,t)\in\Xi_{\mathbb{X}}} \mathbb{X}(u,t) = \bigcup_{j=1}^{N} \bigcup_{u\in\Xi_{\mathbb{B}}(\varepsilon_j)} \mathbb{X}(u,t_j).$$

Moreover, there are positive constants c_1 and c_2 depending only on d such that

(4.18) $\mathbf{c}_0((t_ju, t_j), c_1\varepsilon) \subset \mathbb{X}(u, t_j) \subset \mathbf{c}_0((t_ju, t_j), c_2\varepsilon), \quad (t_ju, t_j) \in \Xi_{\mathbb{X}},$

and c'_d and c_d depending only on d such that

$$c_d'\varepsilon^{-d-1} \le \#\Xi_{\mathbb{X}} \le c_d\varepsilon^{-d-1}$$

Proof. The proof is parallel to that of Propositions 3.14 and follows almost verbatim. We omit the details. \Box

Proposition 4.15. For $\varepsilon > 0$, let $\Xi_{\mathbb{X}}$ be a maximal ε -separated set in \mathbb{X}^{d+1} . Define

$$\Xi_{\mathbb{X}}^{\varrho} := \left\{ \left(x, \sqrt{t^2 - \varrho^2} \right) : (x, t) \in \Xi_{\mathbb{X}} \right\}.$$

Then $\Xi^{\varrho}_{\mathbb{X}}$ is a maximal ε -separated set in the solid hyperboloid ${}_{\varrho}\mathbb{X}^{d+1}$.

This is an immediate consequence of (3.5). In particular, for $\Xi_{\mathbb{X}}$ defined in Proposition 4.14, both (3.25) and (3.26) extend to ${}_{\varrho}\mathbb{X}^{d+1}$ as well. Moreover, this set is also evenly symmetric, where the notion of evenly symmetric set on \mathbb{X}^{d+1} is defined analogously as in Definition 3.16 with \mathbb{X}_0 replaced by \mathbb{X} .

With Assertions 1–3 established for $W_{\gamma,\mu}$ for $\gamma \geq 0$ and $\mu \geq 0$, we can now apply [30, Theorem 2.15] to state the Marcinkiewicz-Zygmund inequality. However, following the consideration in the case of \mathbb{X}_0^{d+1} , our result holds under symmetry assumption on the weight W and the set $\Xi_{\mathbb{X}}$, and is restricted to the subspace of polynomials

$$\Pi_n^E(\mathbb{X}^{d+1}) = \left\{ p \in \Pi_n^{d+1} : p(x,t) = p(x,-t), \forall (x,t) \in \mathbb{X}^{d+1} \right\}.$$

Theorem 4.16. Let W be a doubling weight on \mathbb{X}^{d+1} such that W(x,t) = W(x,-t) for all $(x,t) \in \mathbb{R}^{d+1}$. Let $\Xi^{\varrho}_{\mathbb{X}}$ be a symmetric maximal $\frac{\delta}{n}$ -separated subset of \mathbb{X}^{d+1} and $0 < \delta \leq 1$.

(i) For all $0 and <math>f \in \Pi_m^E(\mathbb{X}^{d+1})$ with $n \le m \le cn$,

$$\sum_{z \in \Xi_{\mathbb{X}}^{\varrho}} \left(\max_{(x,t) \in \mathbf{c}_{\varrho}((z,r),\frac{\delta}{n})} |f(x,t)|^{p} \right) W\left(\mathbf{c}_{\varrho}((z,r),\frac{\delta}{n})\right) \le c_{W,p} \|f\|_{p,W}^{p}$$

where $c_{W,p}$ depends on p when p is close to 0 and on the doubling constant of W.

(ii) For 0 < r < 1, there is a $\delta_r > 0$ such that for $\delta \leq \delta_r$, $r \leq p < \infty$ and $f \in \prod_n^E(\mathbb{X}^{d+1})$,

$$\|f\|_{p,W}^{p} \leq c_{W,r} \sum_{z \in \Xi^{\varrho}} \left(\min_{(x,t) \in \mathbf{c}_{\varrho}\left((z,r), \frac{\delta}{n}\right)} |f(x,t)|^{p} \right) W\left(\mathbf{c}_{\varrho}((z,r), \frac{\delta}{n})\right)$$

where $c_{W,r}$ depends only on the doubling constant of W and on r when r is close to 0.

4.6. **Positive cubature rules.** We need fast decaying polynomials on the solid hyperboloid, which will verify Assertion 4.

Lemma 4.17. Let $d \ge 2$ and $\varrho \ge 0$. For each $(x,t) \in \mathbb{X}^{d+1}$, there is a polynomial $\mathcal{T}_{x,t}^{\varrho}$ in Π_n^E that satisfies

(1)
$$\mathcal{T}_{x,t}^{\varrho}(x,t) = 1, \ \mathcal{T}_{x,t}^{\varrho}(y,s) \ge c > 0 \ if \ (y,s) \in \mathbf{c}((x,t),\frac{\delta}{n}), \ and \ for \ every \ \kappa > 0,$$

$$0 \le \mathcal{T}_{x,t}^{\varrho}(y,s) \le c_{\kappa} \left(1 + \mathsf{d}_{\mathbb{X}}^{\varrho}((x,t),(y,s))\right)^{-\kappa}, \quad (y,s) \in \mathbb{X}^{d+1}.$$

(2) there is a polynomial q of degree 4n such that $q(x,t)\mathcal{T}_{x,t}^{\varrho}$ is a polynomial of degree 5n in (x,t) and $1 \leq q(x,t) \leq c$.

Proof. We construct our polynomial based on the one given in Lemma 3.18 by following the approach in Lemma 3.18. For $(x,t), (y,s) \in \mathbb{X}^{d+1}$, we introduce the notation $X = (x, \sqrt{t^2 - \varrho^2 - ||x||^2})$ and $Y = (y, \sqrt{s^2 - \varrho^2 - ||y||^2})$. Moreover, denote $X_* = (x, -\sqrt{t^2 - \varrho^2 - ||x||^2})$ and $Y_* = (y, -\sqrt{s^2 - \varrho^2 - ||y||^2})$. Then (X,t), (Y,s) and (X_*,t) and (Y_*,s) are all elements of \mathbb{X}_0^{d+2} . Let $T_{(X,t)}^{\varrho}$ denote the polynomial of degree n on \mathbb{X}_0^{d+2} defined in Lemma 3.18. We now define

$$\mathcal{T}^{\varrho}_{x,t}(y,s) := \frac{T^{\varrho}_{(X,t)}(Y,s) + T^{\varrho}_{(X,t)}(Y_*,s)}{1 + T^{\varrho}_{(X,t)}(X_*,t)}$$

Since $T^{\varrho}_{(X,t)}(X,t) = 1$, it follows that $\mathcal{T}^{\varrho}_{(x,t)}(x,t) = 1$. Moreover, since

$$T^{\varrho}_{(X,t)}(X_*,t) = \frac{S_n(2\|x\|^2 + 1 + 2\varrho^2 - 2t^2) + S_n(2\|x\|^2 - 1)}{1 + S_n(2t^2 - 2\varrho^2 - 1)}$$

and $0 \leq S_n(t) \leq c$, we see that $0 \leq T_{(X,t)}(X_*,s) \leq c$. In particular, it follows that

$$\mathcal{T}^{\varrho}_{x,t}(y,s) \ge c \, T^{\varrho}_{(X,t)}(Y,s) \ge c > 0, \qquad (y,s) \in \mathbf{c}\big((x,t), \frac{\delta}{n}\big),$$

since $\mathsf{d}_{\mathbb{X}}((x,t),(y,s)) = \mathsf{d}_{\mathbb{X}_0^{d+2}}((X,t),(Y,s))$. Furthermore, since $\cos \mathsf{d}_{\mathbb{X}_0}((X,t),(Y,s)) \ge \cos \mathsf{d}_{\mathbb{X}_0}((X,t),(Y_*,s))$, we obtain

$$\mathsf{d}_{\mathbb{X}_0}((X,t),(Y_*,s)) \ge \mathsf{d}_{\mathbb{X}_0}((X,t),(Y,s)) = \mathsf{d}_{\mathbb{X}}((x,t),(y,s)).$$

Hence, using the estimate of $T^{\varrho}_{(X,t)}$ in Lemma 3.18, we conclude that

$$0 \leq \mathcal{T}_{x,t}^{\varrho}(y,s) \leq c \left[\left(1 + n \mathsf{d}_{\mathbb{V}_0}((X,t),(Y,s)) \right)^{-\kappa} + \left(1 + n \mathsf{d}_{\mathbb{V}_0}((X,t),(Y_*,s)) \right)^{-\kappa} \right] \\ \leq c \left(1 + n \mathsf{d}_{\mathbb{V}}((x,t),(y,s)) \right)^{-\kappa}.$$

Finally, let $q(x,t) = (1 + S_n(2t^2 - 2\varrho^2 - 1))T^{\varrho}_{(X,t)}(X_*,t)$. Then q(x,t) is a polynomial of degree at most 4n, so that $q(x,t)\mathcal{T}^{\varrho}_{x,t}$ is a polynomial of degree at most 5n and $1 \leq q(x,t) \leq c$. This completes the proof.

The lemma establishes Assertion 4 with a polynomial in $\prod_{n=1}^{E} (\mathbb{X}^{d+1})$. Let W be a doubling weight function that is even in t variable on \mathbb{X}^{d+1} . We define

(4.19)
$$\lambda_n^E(W; x, t) := \inf_{\substack{g(x,t)=1\\g\in\Pi_n^E(\mathbb{X}^{d+1})}} \int_{\mathbb{X}^{d+1}} |g(x,t)|^2 W(x,t) \mathrm{d}x \mathrm{d}t,$$

which is the Christoffel function for the space $\Pi_n^E(\mathbb{X}^{d+1})$. Just like the case of \mathbb{X}_0^{d+1} , this Christoffel function is related to the kernel $\mathbf{K}_n^E(W) = \sum_{k=0}^n \mathbf{P}_k^E(W)$ by

$$\lambda_n^E(W; x, t) = \frac{1}{\mathbf{K}_n^E(W; (x, t), (x, t))}, \qquad (x, t) \in \mathbb{X}^{d+1}.$$

Moreover, [30, Propositions 2.17 and 2.18] remain valid for $\lambda_n^E(W)$ on \mathbb{X}^{d+1} by using the Lemma 4.17. Hence, we obtain the following corollary.

Corollary 4.18. Let W be a doubling weight function on \mathbb{X}^{d+1} such that W(x,t) = W(x,-t) for all $(x,t) \in \mathbb{X}^{d+1}$. Then

$$\lambda_n^E(W;(x,t)) \le c W(\mathbf{c}_{\varrho}((x,t),\frac{1}{n})).$$

Moreover, for $\gamma \geq 0$ and $\mu \geq 0$,

$$\lambda_n^E \left(W_{\gamma,\mu}; (x,t) \right) \ge c \, W_{\gamma,\mu} \left(\mathbf{c}_{\varrho} \left((x,t), \frac{1}{n} \right) \right) = c n^{-d} W_{\gamma,\mu}^{\varrho}(n; x, t).$$

We are now in position to state the positive cubature rule for solid hyperboloid, following which holds for polynomials in $\Pi_n^E(\mathbb{X}^{d+1})$ and under the symmetry assumptions.

Theorem 4.19. Let $d \ge 2$ and $\varrho \ge 0$. Let W be a doubling weight on \mathbb{X}^{d+1} such that W(x,t) = W(x,-t) for all $(x,t) \in \mathbb{X}^{d+1}$. Let Ξ^{ϱ} be a symmetric maximum $\frac{\delta}{n}$ -separated subset of \mathbb{X}^{d+1} . There is a $\delta_0 > 0$ such that for $0 < \delta < \delta_0$ there exist positive numbers $\lambda_{z,r}$, $(z,r) \in \Xi^{\varrho}$, so that

(4.20)
$$\int_{\mathbb{X}^{d+1}} f(x,t)W(x,t)\mathrm{d}x\mathrm{d}t = \sum_{(z,r)\in\Xi^{\varrho}} \lambda_{z,r}f(z,r), \qquad \forall f \in \Pi_n^E(\mathbb{X}^{d+1}).$$

Moreover, $\lambda_{z,r} \sim W(\mathbf{c}_{\varrho}((z,r),\frac{\delta}{n}))$ for all $(z,r) \in \Xi^{\varrho}$.

Again, this follows from [30, Theorem 2.20] when the domain becomes \mathbb{X}_0^{d+1} and it remains valid under symmetry assumptions on the weight and on the polynomials.

4.7. Localized tight frames. The symmetry assumption for the weight and for the ε -separated subset also carries over to the local frame on the solid hyperboloid. Let $\varrho \geq 0$ and let W be a doubling weight on \mathbb{X}^{d+1} that is even in t variable. Let $\mathbf{L}_n^E(W) * f$ denote the operator in $\Pi_{2n}^E(\mathbb{X}^{d+1})$ defined by

$$\mathbf{L}_n^E(W) * f(x) := \int_{\mathbb{X}^{d+1}} f(y,s) \mathbf{L}_n^E(W;(x,t),(y,s)) W(y,s) \mathrm{d}y \mathrm{d}s,$$

where $\mathbf{L}_{n}^{E}(W; \cdot, \cdot)$ is the highly localized kernel defined via a cut-off function \hat{a} that satisfies (3.30). For $j = 0, 1, \ldots$, let Ξ_{j}^{ϱ} be a symmetric maximal $\frac{\delta}{2^{j}}$ -separated subset in \mathbb{X}^{d+1} , so that

$$\int_{\mathbb{X}^{d+1}} f(x,t)W(x,t)\mathrm{d}x\mathrm{d}t = \sum_{(z,r)\in\Xi_j^\varrho} \lambda_{(x,r),j}f(z,r), \quad f\in\Pi_{2^j}^E(\mathbb{X}^{d+1}).$$

For $j = 1, 2, \ldots$, define the operator $F_j^{\varrho}(W)$ by

$$F_j^{\varrho}(W) * f = \mathbf{L}_{2^{j-1}}^E(W) * f$$

and define the frame elements $\psi_{(z,r),j}$ for $(z,r) \in \Xi_j^{\varrho}$ by

$$\psi_{(z,r),j}(x,t) := \sqrt{\lambda_{(z,r),j}} F_j^{\varrho}((x,t),(z,r)), \qquad (x,t) \in \mathbb{X}^{d+1}$$

Then $\Phi = \{\psi_{(z,r),j} : (z,r) \in \Xi_j^{\varrho}, j = 1, 2, 3, ...\}$ is a tight frame by [30, Theorem 2.21].

Theorem 4.20. Let W be a doubling weight on \mathbb{X}^{d+1} even in its t variable. If $f \in L^2(\mathbb{X}^{d+1}, W)$ and f is even in t variable, then

$$f = \sum_{j=0}^{\infty} \sum_{(z,r)\in \Xi_j^\varrho} \langle f, \psi_{(z,r),j} \rangle_{\mathsf{w}} \psi_{(z,r),j} \quad in \ L^2(\mathbb{X}^{d+1}, W)$$

and

$$||f||_{2,W} = \Big(\sum_{j=0}^{\infty} \sum_{(z,r)\in\Xi_j^{\varrho}} |\langle f, \psi_{(z,r),j} \rangle_W|^2 \Big)^{1/2}.$$

Furthermore, for $\gamma \geq 0$ and $\mu \geq 0$, the frame for $W^{\varrho}_{\gamma,\mu}$ is highly localized in the sense that, for every $\kappa > 0$, there exists a constant $c_{\kappa} > 0$ such that

$$(4.21) \quad |\psi_{(z,r),j}(x,t)| \le c_{\kappa} \frac{2^{j(d+1)/2}}{\sqrt{W_{\gamma,\mu}^{\varrho}(2^{j};x,t)}(1+2^{j}\mathsf{d}^{\varrho}_{\mathbb{X}}((x,t),(z,r)))^{\kappa}}, \quad (x,t) \in \mathbb{X}^{d+1}.$$

Like the case on the hyperbolic surface, the frame elements involve only orthogonal polynomials even in t variable. The localization (4.21) follows from Theorem 4.9 and $\lambda_{(z,r),j} \sim 2^{-j(d+1)} W^{\varrho}_{\gamma,\mu}(2^j; x, t)$ which holds for $W^{\varrho}_{\gamma,\mu}$ by Corollary 4.18 and (4.9). We note, however, that the localized tight frame holds for the weight function $W^{\varrho}_{\gamma,\mu}$, which does not include the Lebesgue measure on \mathbb{X}^{d+1} .

4.8. Characterization of best appoximation. For $f \in L^p(\mathbb{X}^{d+1}, W^{\varrho}_{\gamma,\mu})$, we denote by $\mathbf{E}_n(f)_{p,W^{\varrho}_{\gamma,\mu}}$ the best approximation to f from $\Pi^{d+1}_n(\mathbb{X}^{d+1})$ in the norm $\|\cdot\|_{p,W^{\varrho}_{\gamma,\mu}}$; that is,

$$\mathbf{E}_n(f)_{p,W^\varrho_{\gamma,\mu}} := \inf_{g \in \Pi_n(\mathbb{X}^{d+1})} \|f - g\|_{p,W^\varrho_{\gamma,\mu}}, \qquad 1 \leq p \leq \infty.$$

As in the case of hyperbolic surface, if f is even in t variable, then we can choose the polynomial of best approximation from $\Pi_n^E(\mathbb{X}^{d+1})$. We can give a characterization of this quantity for functions that are even in t variable.

For $f \in L^p(\mathbb{X}^{d+1}, W^{\varrho}_{\gamma,\mu})$ and r > 0, the modulus of smoothness is defined by

$$\omega_r(f;\rho)_{p,W^{\varrho}_{\gamma,\mu}} = \sup_{0 \le \theta \le \rho} \left\| \left(I - \mathbf{S}_{\theta,W^{\varrho}_{\gamma,\mu}} \right)^{r/2} f \right\|_{p,W^{\varrho}_{\gamma,\mu}}, \quad 1 \le p \le \infty,$$

where the operator $\mathbf{S}_{\theta, W^{\varrho}_{\gamma,\mu}}$ is defined by, for n = 0, 1, 2, ... and $\lambda = \gamma + \mu + \frac{d}{2}$,

$$\operatorname{proj}_{n}^{E}\left(W_{\gamma,\mu}^{\varrho};\mathbf{S}_{\theta,W_{\gamma,\mu}^{\varrho}}f\right) = R_{n}^{\left(\lambda - \frac{1}{2},\lambda - \frac{1}{2}\right)}(\cos\theta)\operatorname{proj}_{n}^{E}\left(W_{\gamma,\mu}^{\varrho};f\right).$$

In terms of the fractional differential operator $(-\mathfrak{D}_{\gamma,\mu}^{\varrho})^{\frac{r}{2}}$, the *K*-functional for a doubling weight *W*, even in *t* variable on \mathbb{X}^{d+1} , is defined by

$$\mathbf{K}_{r}(f,t)_{p,W} := \inf_{g \in W_{p}^{r}(\mathbb{X}^{d+1},W)} \left\{ \|f - g\|_{p,W} + t^{r} \left\| (-\mathfrak{D}_{\gamma,\mu}^{\varrho})^{\frac{r}{2}} f \right\|_{p,W} \right\}$$

where the Sobolev space $W_p^r(\mathbb{X}^{d+1}, W)$ is the space that consists of $f \in L^p(\mathbb{X}^{d+1}, W)$, even in t variable, so that $\left\| (-\mathfrak{D}_{\gamma,\mu}^{\varrho})^{\frac{r}{2}} f \right\|_{p,W}$ is finite. For $W_{\gamma,\mu}^{\varrho}$, Assertions 1 and 3 hold and we now verify that Assertion 5 holds as well.

For $W^{\varrho}_{\gamma,\mu}$, Assertions 1 and 3 hold and we now verify that Assertion 5 holds as well. By (4.12), the kernel $L_n^{(r)}(\varpi)$ in Assertion 5 becomes

$$\mathbf{L}_{n}^{(r)}\big(W_{\gamma,\mu}^{\varrho};(x,t),(y,s)\big) = \sum_{k=0}^{\infty} \widehat{a}\left(\frac{k}{n}\right) \left(k(k+2\gamma+2\mu+d)\right)^{\frac{r}{2}} \mathbf{P}_{k}^{E}\big(W_{\gamma,\mu}^{\varrho};(x,t),(y,s)\big).$$

Lemma 4.21. Let $\gamma, \mu \geq -\frac{1}{2}$ and $\kappa > 0$. Then, for r > 0 and $(x, t), (y, s) \in \mathbb{X}^{d+1}$,

$$\left|\mathbf{L}_{n}^{(r)}\left(W_{\gamma,\mu}^{\varrho};(x,t),(y,s)\right)\right| \leq c_{\kappa} \frac{n^{d+r+1}}{\sqrt{W_{\gamma,\mu}^{\varrho}(n;t)}\sqrt{W_{\gamma,\mu}^{\varrho}(n;s)}} \left(1 + n\mathsf{d}_{\mathbb{X}}^{\varrho}((x,t),(y,s))\right)^{-\kappa}$$

Proof. This can be proved similarly as in the proof of Lemma 3.22 and using the estimate in Theorem 4.9. We omit the details. \Box

We can now state the characterization of the best approximation by polynomials on the hyperboloid, following [30, Theorem 3.12].

Theorem 4.22. Let $f \in L^p(\mathbb{X}^{d+1}, W)$ if $1 \leq p < \infty$ and $f \in C(\mathbb{X}^{d+1})$ if $p = \infty$. Assume that f satisfies f(x,t) = f(x,-t). Let $\varrho \geq 0$, r > 0 and $n = 1, 2, \ldots$ Then

(i) for $W = W^{\varrho}_{\gamma,\mu}$ with $\gamma, \mu \ge 0$,

$$\mathbf{E}_n(f)_{p,W^{\varrho}_{\gamma,\mu}} \le c \, \mathbf{K}_r(f;n^{-1})_{p,W^{\varrho}_{\gamma,\mu}};$$

(ii) for every doubling weight W that satisfies W(x,t) = W(x,-t) on \mathbb{X}^{d+1} ,

$$\mathbf{K}_{r}(f; n^{-1})_{p,W} \le cn^{-r} \sum_{k=0}^{n} (k+1)^{r-1} \mathbf{E}_{k}(f)_{p,W}.$$

For $W = W^{\varrho}_{\gamma,\mu}$, the K-functional is equivalent to the modulus of smoothness.

Theorem 4.23. Let $\varrho \ge 0$, $\gamma, \mu \ge 0$ and $f \in L_p^r(\mathbb{X}^{d+1}, W_{\gamma,\mu}^{\varrho})$, $1 \le p \le \infty$. Then for $0 < \theta \le \pi/2$ and r > 0

$$c_1 \mathbf{K}_r(f;\theta)_{p,W^{\varrho}_{\gamma,\mu}} \le \omega_r(f;\theta)_{p,W^{\varrho}_{\gamma,\mu}} \le c_2 \mathbf{K}_r(f;\theta)_{p,W^{\varrho}_{\gamma,\mu}}.$$

In particular, the characterization in Theorem 4.22 can be stated in terms of the modulus of smoothness instead.

References

- P. Baldi, G. Kerkyacharian, D. Marinucci, and D. Picard. Asymptotics for spherical needlets, Ann. Statist. 37 (2009), 1150–1171.
- [2] P. Baldi, G. Kerkyacharian, D. Marinucci, and D. Picard. Adaptive density estimation for directional data using needlets, Ann. Statist. 37 (2009), 3362–3395.
- [3] G. Brown, F. Dai, Approximation of smooth functions on compact two-point homogeneous spaces, J. Funct. Anal. 220 (2005), 401–423.
- [4] F. Dai, Multivariate polynomial inequalities with respect to doubling weights and A_∞ weights, J. Funct. Anal. 235 (2006), 137–170.
- [5] F. Dai and A. Prymak, L_p -Bernstein inequalities on C^2 domains, 2020. arXiv:2010.06728
- [6] F. Dai and A. Prymak, On directional Whitney inequality, 2020. arXiv:2010.08374
- [7] F. Dai and H. Wang, Optimal cubature formulas in weighted Besov spaces with A_{∞} weights on multivariate domains, *Const. Approx.* **37** (2013), 167–194.
- [8] F. Dai and Y. Xu, Approximation theory and harmonic analysis on spheres and balls. Springer Monographs in Mathematics, Springer, 2013.
- [9] C. F. Dunkl and Y. Xu, Orthogonal Polynomials of Several Variables. Encyclopedia of Mathematics and its Applications 155, Cambridge University Press, Cambridge, 2014.
- [10] R. A. DeVore and G. G. Lorentz, Constructive approximation, Grundlehren der Mathematischen Wissenschaften, vol. 303, Springer-Verlag, Berlin, 1993.
- [11] G. Kyriazis, P. Petrushev and Y. Xu, Decomposition of weighted Triebel-Lizorkin and Besov spaces on the ball, Proc. London Math. Soc. 97 (2008), 477–513.
- [12] K. Ivanov and P. Petrushev, Fast memory efficient evaluation of spherical polynomials at scattered points, Adv. Comput. Math. 41 (2015), 191–230.
- [13] K. Ivanov and P. Petrushev, Highly effective stable evaluation of bandlimited functions on the sphere, Numer. Algorithms, 71 (2016), 585–611.

- [14] K. Ivanov, P. Petrushev and Y. Xu, Decomposition of spaces of distributions induced by tensor product bases. J. Funct. Anal. 263 (2012), 1147–1197.
- [15] Q. T. Le Gia, I. H. Sloan, Y. G. Wang and R. S. Womersley, Needlet approximation for isotropic random fields on the sphere. J. Approx. Theory 216 (2017), 86–116.
- [16] G. Mastroianni and V. Totik Weighted polynomial inequalities with doubling and A_{∞} weights. Const. Approx. 16 (2000) 37–71.
- [17] F. J. Narcowich, P. Petrushev and J. D. Ward, Localized tight frames on spheres, SIAM J. Math. Anal. 38 (2006), 574–594.
- [18] F. J. Narcowich, P. Petrushev and J. D. Ward, Decomposition of Besov and Triebel-Lizorkin spaces on the sphere, J. Funct. Anal. 238 (2006), 530–564.
- [19] P. Petrushev and Y. Xu, Localized polynomial frames on the ball, Constr. Approx. 27 (2008), 121–148.
- [20] Kh. P. Rustamov, On the approximation of functions on a sphere, (Russian), Izv. Ross. Akad. Nauk Ser. Mat. 57 (1993), 127–148; translation in Russian Acad. Sci. Izv. Math. 43 (1994), no. 2, 311–329.
- [21] E. Stein, Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals Princeton University Press, Princeton, 1993.
- [22] G. Szegő, Orthogonal polynomials. 4th edition, Amer. Math. Soc., Providence, RI. 1975
- [23] V. Totik, Polynomial approximation on polytopes, Mem. Amer. Math. Soc. 232 (2014), no. 1091, vi+112.
- [24] V. Totik, Polynomial approximation in several variables, J. Approx. Theory 252: 105364 (2020).
- [25] Y. G. Wang, Q. T. Le Gia, I. H. Sloan and R. S. Womersley, Fully discrete needlet approximation on the sphere. *Appl. Comput. Harmon. Anal.* 43 (2017), 292–316.
- [26] Y. Xu, Summability of Fourier orthogonal series for Jacobi weight on a ball in R^d, Trans. Amer. Math. Soc., 351 (1999), 2439–2458.
- [27] Y. Xu, Weighted approximation of functions on the unit sphere. Const. Approx. 21 (2005), 1–28.
- [28] Y. Xu, Fourier series in orthogonal polynomials on a cone of revolution, J. Fourier Anal. Appl., 26 (2020), Article number: 36.
- [29] Y. Xu, Orthogonal structure and orthogonal series in and on a double cone or a hyperboloid. Trans. Amer. Math. Soc., 374 (2021), 3603–3657.
- [30] Y. Xu, Approximation and localized polynomial frame on conic domains. arXiv:2011.14180

Department of Mathematics, University of Oregon, Eugene, OR 97403–1222, USA $\mathit{Email}\ address:\ \texttt{yuan@uoregon.edu}$