INDEPENDENT SETS IN SUBGRAPHS OF A SHIFT GRAPH

ANDRII ARMAN, VOJTĚCH RÖDL, AND MARCELO TADEU SALES

ABSTRACT. Erdős, Hajnal and Szemerédi proved that any subset G of vertices of a shift graph Sh_n^k has the property that the independence number of the subgraph induced by Gsatisfies $\alpha(\operatorname{Sh}_n^k[G]) \ge (\frac{1}{2} - \varepsilon) |G|$, where $\varepsilon \to 0$ as $k \to \infty$. In this note we prove that for k = 2 and $n \to \infty$ there are graphs $G \subseteq {\binom{[n]}{2}}$ with $\alpha(\operatorname{Sh}_n^2[G]) \le (\frac{1}{4} + o(1)) |G|$, and $\frac{1}{4}$ is best possible. We also consider a related problem for infinite shift graphs.

1. INTRODUCTION

For $n > k \in \mathbb{N}$ the shift graph Sh_n^k with

$$V(Sh_n^k) = \{ (x_1, \dots, x_k) : 1 \le x_1 < \dots < x_k \le n \}$$

is a graph in which two vertices $\mathbf{x} = (x_1, \ldots, x_k)$ and $\mathbf{y} = (y_1, \ldots, y_k)$ are adjacent if $x_i = y_{i+1}$ for all $i \in \{1, \ldots, k-1\}$ (or $y_i = x_{i+1}$ for all $i \in \{1, \ldots, k-1\}$). Shift graphs were introduced by Erdős and Hajnal [3],[4] and are standard examples of graphs with large chromatic number and large odd girth. More precisely, while the odd girth of Sh_n^k is 2k + 1, they proved* that Sh_n^k has chromatic number $(1 + o(1)) \log^{(k-1)} n$, where $\log^{(k-1)}$ stands for k - 1 times iterated \log_2 .

Shift graphs have another interesting property: For each finite set $G \subseteq V(\operatorname{Sh}_n^k)$ the induced subgraph $\operatorname{Sh}_n^k[G]$ has a relatively large independent set with respect to |G|. In other words, the property "having a large independent subset" is hereditary for Sh_n^k . Namely, for

(1)
$$\alpha_n^k = \min\left\{\frac{\alpha(\operatorname{Sh}_n^k[G])}{|G|} : \emptyset \neq G \subseteq V(\operatorname{Sh}_n^k)\right\},$$

Erdős, Hajnal and Szemerédi [5, Theorem 1] proved the following.

Theorem 1.1 (Erdős, Hajnal, Szemerédi). For positive integers k < n

$$\alpha_n^k \ge \frac{1}{2} - \frac{1}{k}.$$

The second author was supported by NSF grant DMS 1764385.

^{*}In [4] authors considered infinite graphs, however their proof can be adapted for finite case (see [1] and [6] for more detailed description).

As for the upper bound, for $n \ge 2k + 1$ the shift graph Sh_n^k contains an odd cycle and so $\alpha_n^k < 1/2$. Therefore, Theorem 1.1 yields a lower bound which for large values of k is essentially optimal.

Nevertheless, determining the values of α_n^k for fixed k and large n seems to represent an interesting and non-trivial problem. We will concentrate our attention on the case k = 2. In this case the bound from Theorem 1.1 is not optimal, as we observe that $\alpha_n^2 \ge 1/4$ for all n, and prove a matching upper bound.

Theorem 1.2. $\lim_{n \to \infty} \alpha_n^2 = \frac{1}{4}$.

In [2], Czipszer, Erdős and Hajnal proved that the densest independent set of the infinite graph $\operatorname{Sh}^2_{\mathbb{N}}$ has density 1/4 (see Section 3 for precise formulation). We complement their result by showing that the infinite shift graph $\operatorname{Sh}^2_{\mathbb{N}}$ does not have a similar hereditary property, i.e., there exists $G \subseteq V(\operatorname{Sh}^2_{\mathbb{N}})$ such that any independent set in $\operatorname{Sh}^2_{\mathbb{N}}[G]$ has density zero in G (see Theorem 3.1).

2. Proof of Theorem 1.2

Note that $\alpha_n^2 = \min\left\{\frac{\alpha(\operatorname{Sh}_n^2[G])}{|G|} : \emptyset \neq G \subseteq V(\operatorname{Sh}_n^2)\right\}$ is a nonincreasing positive sequence, so the sequence $\{\alpha_n^2\}$ has a limit. Additionally, we will often view $G \subseteq V(\operatorname{Sh}_n^2)$ as a graph with V(G) = [n] and set of edges equal to G. Subsequently |G| will denote both a size of Gas a subset of $V(\operatorname{Sh}_n^2)$, and the number of edges in G when it is viewed as a graph.

2.1. Lower bound. We first show that the value of the limit in Theorem 1.2 is at least 1/4.

Claim 2.1. For every set $G \subseteq V(\operatorname{Sh}_n^2)$ we have $\alpha(\operatorname{Sh}_n^2[G]) \ge \frac{1}{4}|G|$.

Proof. Let $G \subseteq V(\operatorname{Sh}_n^2)$ be given. Consider a random colouring $c : [n] \to \{r, b\}$ such that every $i \in [n]$ is coloured red/blue with probability 1/2 independently of other elements of [n].

Let G_c be a random subset of G defined by

$$G_c = \{(i, j) \in G : i < j, c(i) = b, c(j) = r\}.$$

Then such G_c is always an independent set in Sh_n^2 . Moreover, $\mathbb{P}(e \in G_c) = \frac{1}{4}$ for every $e \in G$, and so $\mathbb{E}(|G_c|) = \frac{1}{4}|G|$. Therefore $\alpha(\operatorname{Sh}_n^2[G]) \ge \frac{1}{4}|G|$. 2.2. Upper bound. We now proceed and prove the upper bound

(2)
$$\lim_{n \to \infty} \alpha_n^2 \leqslant \frac{1}{4}.$$

In what follows for every $\varepsilon > 0$, integer d satisfying $\frac{3+\ln d}{4d} \leq \frac{\varepsilon}{2}$, and for every integer $n \geq n_0(\varepsilon, d)$ that is a multiple of 2^d , we will construct a graph $G_{\varepsilon}(n, d) \subseteq V(\operatorname{Sh}_n^2)$ with

$$\alpha(\operatorname{Sh}_n^2[G_{\varepsilon}(n,d)]) \leqslant \left(\frac{1}{4} + \varepsilon\right) |G_{\varepsilon}(n,d)|.$$

To be more precise, for such ε and d we inductively build $G_{\varepsilon}(n, d)$ satisfying

(3)
$$\frac{\alpha(\operatorname{Sh}_n^2[G_{\varepsilon}(n,d)])}{|G_{\varepsilon}(n,d)|} \leqslant \left(\frac{1}{4} + \frac{3+\ln d}{4d} + \frac{\varepsilon}{2}\right)$$

Since $\{\alpha_n^2\}$ is nonincreasing, (3) implies that $\lim_{n\to\infty} \alpha_n^2 \leq 1/4 + \varepsilon$, which subsequently implies (2) by letting $\varepsilon \to 0$.

While constructing $G_{\varepsilon}(n,d)$ we will use random bipartite graphs. Recall that if G is a graph and $X, Y \subseteq V(G)$ then G[X,Y] is a graph consisting of edges of G with one vertex in X and another in Y. Finally let $e_G(X,Y) = |E(G[X,Y])|$ and we will omit subscript when G is obvious from the context. The following claim can be easily verified by considering a random graph and so the proof of Claim 2.2 is postponed to Appendix.

Claim 2.2. For $\varepsilon > 0$ and $d \in \mathbb{N}$ there is $n_0 = n_0(\varepsilon, d)$ such that for all $n \ge n_0$ that are divisible by 2^d the following holds. Let $[n] = S \cup L$, where $S = \{1, \ldots, \frac{n}{2}\}$ and $L = [n] \setminus S$. There exists a bipartite graph $B_{\varepsilon}(n, d)$ with bipartition $V(B_{\varepsilon}(n, d)) = S \sqcup L$ such that

(i) $|B_{\varepsilon}(n,d)| = \frac{n^2}{2^{d+1}}$. (ii) for all $X \subseteq S$ and $Y \subseteq L$

$$e(X,Y) = \frac{1}{2^{d-1}}|X||Y| \pm \frac{\varepsilon n^2}{2^{d+2}}.$$

Construction of $G_{\varepsilon}(n,d)$.

Definition 2.3. For every even n let $G_{\varepsilon}(n, 1)$ be such that

$$G_{\varepsilon}(n,1) = \{(i,j) : 1 \leq i \leq \frac{n}{2} < j \leq n\},\$$

i.e., $G_{\varepsilon}(n,1)$ is a complete balanced bipartite graph.

For $d \in \mathbb{N}$ define graph $G_{\varepsilon}(n,d)$ recursively for all sufficiently large[†] n such that $2^d | n$. Let $[n] = S \cup L$, where $S = \{1, \ldots, \frac{n}{2}\}$ and $L = [n] \setminus S$. Then define

$$G_{\varepsilon}(n,d) = G_{\varepsilon}(S,d-1) \cup G_{\varepsilon}(L,d-1) \cup B_{\varepsilon}(n,d),$$

 ${}^{\dagger}n \ge 2^i n_0(\varepsilon, d-i)$ for all $i \in \{0, 1, \dots, d-2\}$, where $n_0(\varepsilon, d-i)$ is the number provided by Claim 2.2.

where $G_{\varepsilon}(S, d-1) = G_{\varepsilon}(\frac{n}{2}, d-1), V(G_{\varepsilon}(L, d-1)) = L$ and $G_{\varepsilon}(L, d-1) \cong G_{\varepsilon}(\frac{n}{2}, d-1)$, and $B_{\varepsilon}(n, d)$ is a graph guaranteed by Claim 2.2.

To summarize, every $G_{\varepsilon}(n,d) = G$ satisfies the following properties (for $S_n = \{1, \dots, \frac{n}{2}\}$ and $L_n = [n] \setminus S_n$):

- (i) $e_G(S_n, L_n) = \frac{n^2}{2^{d+1}}$.
- (ii) for all $X \subseteq S_n$ and $Y \subseteq L_n$

$$e(X,Y) = \frac{1}{2^{d-1}}|X||Y| \pm \frac{\varepsilon n^2}{2^{d+2}}$$

(iii) $G[S_n] \cong G[L_n] = G_{\varepsilon}(\frac{n}{2}, d-1)$

Using properties (i) and (iii) and induction on d it is easy to verify that for all $d \in \mathbb{N}$ and n divisible by 2^d

(4)
$$|G_{\varepsilon}(n,d)| = d\frac{n^2}{2^{d+1}}$$

We will now proceed with proving (3). First let $G \subseteq V(\operatorname{Sh}_n^2)$ and let $I \subseteq G$ be an independent set in Sh_n^2 . In other words there is no $1 \leq i < j < k \leq n$ with both (i, j) and (j, k) in I. One can observe that for each such $I \subseteq G$ there exists a 2-colouring $c : [n] \to \{r, b\}$ with c(i) = r and c(j) = b whenever $(i, j) \in I$, and then

(5)
$$I \subseteq G_c = \{(x, y) \in G : x < y, c(x) = b, c(y) = r\}.$$

Therefore, in order to prove (3) we will show that for $G = G_{\varepsilon}(n, d)$ and any $c : [n] \to \{r, b\}$

(6)
$$\frac{|G_c|}{|G|} \leqslant \frac{1}{4} + \frac{3 + \ln d}{4d} + \frac{\varepsilon}{2}$$

For the rest of our calculation let ε be fixed. We will now prove (6) by induction on d. In order to make use of recursive structure of $G_{\varepsilon}(n, d)$ we will prove a version of (6) with an additional assumption that $|\{i : c(i) = b\}| = \alpha n$.

To that end for $d \in \mathbb{N}$, $\alpha \in [0, 1]$ and $n \ge n_0(\varepsilon, d)$ let

(7)
$$f_d^{\alpha}(n) = d \cdot \max_c \left\{ \frac{|G_c|}{|G|} : G = G_{\varepsilon}(n,d), |\{i : c(i) = b\}| = \alpha n \right\}.$$

We will prove the following estimate on $f_d^{\alpha}(n)$.

Claim 2.4. For every $d \in \mathbb{N}$, $\alpha \in [0, 1]$ and $n \ge n_0(\varepsilon, d)$

$$f_d^{\alpha}(n) \leq (d+3)(\alpha-\alpha^2) + \frac{1}{4}\ln d + \frac{d\varepsilon}{2}.$$

From (7) it follows that for $G = G_{\varepsilon}(n, d)$ and any colouring c we have

$$\frac{|G_c|}{|G|} \leqslant \max_{\alpha \in [0,1]} \frac{f_d^{\alpha}(n)}{d}.$$

Then by Claim 2.4 we get

$$\frac{|G_c|}{|G|} \leqslant \frac{1}{4}\frac{d+3}{d} + \frac{\ln d}{4d} + \frac{\varepsilon}{2},$$

establishing (6) and (3). Hence it remains to prove Claim 2.4 in order to finish the proof of the upper bound.

Proof of Claim 2.4. We prove a slightly stronger inequality for all $n \ge n_0(\varepsilon, d)$

(8)
$$f_d^{\alpha}(n) \leq (d+3)(\alpha - \alpha^2) + \frac{1}{4} \sum_{i=3}^{d+1} \frac{1}{i} + \frac{d\varepsilon}{2}.$$

The proof is by induction on d. For d = 1 recall that $G = G_{\varepsilon}(n, 1)$ is a complete bipartite graph between S_n and L_n . Let $c : [n] \to \{r, b\}$ be such that for $B = \{i : c(i) = b\}$ we have $|B| = \alpha n$. Then in view of (5) the maximum value of $|G_c|$ is achieved when $B = [\alpha n]$ and so

$$f_1^{\alpha}(n) = \begin{cases} 2\alpha, & \alpha \in [0, \frac{1}{2}] \\ 2 - 2\alpha, & \alpha \in [\frac{1}{2}, 1] \end{cases}$$

Now it is easy to verify that $f_1^{\alpha} \leq 4(\alpha - \alpha^2)$ for all $\alpha \in [0, 1]$, establishing (8) in the case d = 1.

To prove inductive step let $G = G_{\varepsilon}(n,d)$ and let $c : [n] \to \{r,b\}$ be such that for $B = \{i : c(i) = b\}$ we have $|B| = \alpha n$. As before, let $S = \{1, \ldots, \frac{n}{2}\}$ and $L = [n] \setminus S$. Let B_S, B_L , R_S and R_L denote the set of blue and red vertices in S and L respectively. We will further refine our analysis by assuming that $|B_S| = x\frac{n}{2}$ with some $x \in [0, 2\alpha]$. Since $|B_S| + |R_S| = \frac{n}{2}$, $|B_S| + |B_L| = |B| = \alpha n$, and $|B_L| + |R_L| = \frac{n}{2}$, we have $|R_S| = (1 - x)\frac{n}{2}$, $|B_L| = (2\alpha - x)\frac{n}{2}$, and consequently $|R_L| = (1 - 2\alpha + x)\frac{n}{2}$ (see Figure 1). Then

(9)
$$|G_c| = |G_c[S]| + |G_c[L]| + e_G(B_S, R_L).$$



FIGURE 1. Proportions of red and blue vertices in $G_c[S]$ and $G_c[L]$.

Now, by (iii) $G[S] = G_{\varepsilon}(\frac{n}{2}, d-1)$ and we assumed $|B_S| = x\frac{n}{2}$, so

(10)
$$|G_c[S]| \stackrel{(7)}{\leqslant} \frac{f_{d-1}^x(\frac{n}{2})}{d-1} |G[S]| \stackrel{(4)}{=} \frac{n^2}{2^{d+2}} f_{d-1}^x\left(\frac{n}{2}\right).$$

Similarly, since $|B_L| = (2\alpha - x)\frac{n}{2}$ we have

(11)
$$|G_c[L]| \leq \frac{n^2}{2^{d+2}} f_{d-1}^{2\alpha-x}\left(\frac{n}{2}\right)$$

And finally, since $G = G_{\varepsilon}(n, d)$,

(12)
$$e_G(B_S, R_L) \stackrel{(ii)}{\leq} \frac{1}{2^{d-1}} |B_S|| R_L| + \frac{\varepsilon n^2}{2^{d+2}} = \frac{n^2}{2^{d+1}} \left(x(1 - 2\alpha + x) + \frac{\varepsilon}{2} \right)$$

Combining (9) with (10), (11), and (12) we obtain

$$|G_c| \leq \frac{n^2}{2^{d+1}} \left(\frac{1}{2} \left(f_{d-1}^x \left(\frac{n}{2} \right) + f_{d-1}^{2\alpha - x} \left(\frac{n}{2} \right) \right) + x(1 - 2\alpha + x) + \frac{\varepsilon}{2} \right).$$

Finally, $|G| = |G_{\varepsilon}(n,d)| \stackrel{(4)}{=} d_{\frac{n^2}{2^{d+1}}}$ and so by (7) we deduce that

(13)
$$f_d^{\alpha}(n) \leq \max_{x \in \mathbb{R}} \left\{ \frac{1}{2} \left(f_{d-1}^x \left(\frac{n}{2} \right) + f_{d-1}^{2\alpha - x} \left(\frac{n}{2} \right) \right) + x(1 - 2\alpha + x) + \frac{\varepsilon}{2} \right\}.$$

The last inequality allows us to incorporate induction hypothesis. In particular, by induction hypothesis we have

$$f_{d-1}^{x}(\frac{n}{2}) \leq (d+2)(x-x^{2}) + \frac{1}{4}\sum_{i=3}^{d}\frac{1}{i} + \frac{(d-1)\varepsilon}{2},$$
$$f_{d-1}^{2\alpha-x}(\frac{n}{2}) \leq (d+2)(2\alpha-x)(1-2\alpha+x) + \frac{1}{4}\sum_{i=3}^{d}\frac{1}{i} + \frac{(d-1)\varepsilon}{2},$$

and these two inequalities together with (13), after some simple but tedious algebraic manipulations yield

$$f_d^{\alpha}(n) \le \max_{x \in \mathbb{R}} \left\{ -(d+1)x^2 + (1+2\alpha(d+1))x + (d+2)(\alpha-2\alpha^2) + \frac{1}{4}\sum_{i=3}^d \frac{1}{i} + \frac{d\varepsilon}{2} \right\}$$

In other words $f_d^{\alpha}(n) \leq \max_{x \in \mathbb{R}} \{g(x)\}$, where $g(x) = ax^2 + bx + c$ with a = -(d+1). Since a < 0 we have $\max_{x \in \mathbb{R}} g(x) = g(\frac{-b}{2a}) = c - \frac{b^2}{4a}$. Therefore after another set of algebraic manipulations we obtain

$$f_d^{\alpha}(n) \leq \max_{x \in \mathbb{R}} \{g(x)\} \leq (d+3)(\alpha - \alpha^2) + \frac{1}{4} \sum_{i=3}^{d+1} \frac{1}{i} + \frac{d\varepsilon}{2},$$

finishing the proof of the inductive step and Claim 2.4.

3. Infinite graphs

Recall that Theorem 1.2 states

(14)
$$\lim_{n \to \infty} \min \left\{ \frac{\alpha(\operatorname{Sh}_n^2[G])}{|G|} : \emptyset \neq G \subseteq V(\operatorname{Sh}_n^2) \right\} = \frac{1}{4}.$$

On the other hand, considering $I = \{(i, j) : 1 \leq i \leq \frac{n}{2} < j \leq n\}$ we clearly have $\alpha(\operatorname{Sh}_n^2) \geq \lfloor \frac{n^2}{4} \rfloor$. Moreover $\lfloor \frac{n^2}{4} \rfloor$ is optimal, since any graph $G \subseteq V(\operatorname{Sh}_n^2)$ with $|G| \geq \lfloor \frac{n^2}{4} \rfloor + 1$ contains a triangle and hence such G is not an independent set in Sh_n^2 . Therefore,

(15)
$$\lim_{n \to \infty} \frac{\alpha(\operatorname{Sh}_n^2)}{|\operatorname{Sh}_n^2|} = \frac{1}{2}.$$

It may be interesting to note that infinite version of (15) was considered by Czipszer, Erdős and Hajnal [2] who proved that if I is independent set in countable shift graph $\text{Sh}_{\mathbb{N}}^2$, then the density of I does not exceed 1/4, i.e.

(16)
$$\liminf_{n \to \infty} \frac{\left| I \cap \binom{[n]}{2} \right|}{\binom{n}{2}} \leqslant \frac{1}{4}.$$

(Here $\frac{1}{4}$ is clearly optimal, since $I = \{(i, j) : i < j, i \text{ odd}, j \text{ even}\}$ is independent in $\mathrm{Sh}^2_{\mathbb{N}}$.)

To complete this discussion we provide an infinite variant of (14).

Theorem 3.1. There is $G \subseteq V(Sh_{\mathbb{N}}^2)$ such that if I is an independent set in $Sh_{\mathbb{N}}^2[G]$, then

$$\liminf_{n \to \infty} \frac{\left| I \cap \binom{[n]}{2} \right|}{\left| G \cap \binom{[n]}{2} \right|} = 0.$$

Proof. Consider an infinite ordered tree G with $V(G) = \mathbb{N}$, and with vertices labeled v_i^j , where j denotes the "level" L_j that vertex v_i^j belongs to and i denotes the order in which vertices are listed on the level.

Consider a labeling of vertices of G by integers satisfying $v_i^j < v_i^{j'}$ if j < j' and $v_i^j < v_{i'}^j$ if i < i' such that for all v_i^j the finite set $N^+(v_i^j)$ of all children of v_i^j forms an interval (and these intervals on the level L_{j+1} follow the order of their parents on L_j , see Figure 2). Finally we will assume that for all v_i^j

(17)
$$|N^{+}(v_{i}^{j})| \ge 2^{j} \sum_{v < v_{i}^{j}} |N^{+}(v)|.$$

Now, let $I \subseteq G$ be an infinite independent set in $\operatorname{Sh}_{\mathbb{N}}^2$ and let $(v_k^{j-1}, v_i^j) \in I$, where v_k^{j-1} and v_i^j are parent and child respectively. Let $w = \max\{N^+(v_i^j)\}$ be the largest son of v_i^j and let $W = \{1, \ldots, w\}$ (see Figure 2). Then

(18)
$$G[W] = \bigcup_{v \le v_i^j} \left\{ (v, u) : u \in N^+(v) \right\}.$$



FIGURE 2. Infinite tree G, vertices are ordered top to bottom, left to right. Edges of I are labeled with red.

In particular in view of (17)

(19)
$$|G[W]| \ge |\{(v_i^j, u) : u \in N^+(v)\}| = |N^+(v_i^j)|.$$

On other hand, since $(v_k^{j-1}, v_i^j) \in I$ and I is independent set in $\operatorname{Sh}^2_{\mathbb{N}}$, the set I does not contain any other edge incident to v_i^j . Consequently,

(20)
$$|I[W]| \leq \left| \bigcup_{v < v_i^j} \left\{ (v, u) : u \in N^+(v) \right\} \right| \stackrel{(17)}{\leq} 2^{-j} |N^+(v_i^j)|$$

In view of (19) and (20) we have $|I[W]|/|G[W]| \leq \frac{1}{2^j}$. Now, since I is infinite there are edges $(v_k^{j-1}, v_i^j) \in I$ with sufficiently large j, hence the ratio |I[W]|/|G[W]| can be made arbitrary small, finishing the proof.

4. Concluding remarks

In [5] it was proved[‡] that for any n, k

(21)
$$\alpha_n^k \ge \begin{cases} \frac{1}{2} - \frac{1}{k}, & \text{if } k \text{ is even,} \\ \frac{1}{2} - \frac{1}{2k}, & \text{if } k \text{ is odd.} \end{cases}$$

It remains an open problem to determine for any $k \ge 3$ the exact value of $\lim_{n\to\infty} \alpha_n^k$. For k = 4 we were able to improve the constant in the lower bound (21) from $\frac{1}{4}$ to $\frac{3\$}{8}$ and for k = 3 we believe that estimate in (21) is sharp.

Problem 4.1. Show that $\lim_{n\to\infty} \alpha_n^3 = \frac{1}{3}$.

Finally, all of the results in this paper can be reformulated in terms of subgraphs with no increasing paths of length two. For instance, Theorem 1.2 implies that for any $\varepsilon > 0$ there exists an vertex-ordered graph G such that if $G' \subseteq G$ with $|G'| \ge (\frac{1}{4} + \varepsilon) |G|$, then G' contains an increasing path of length two, i.e. there are i < j < k with $(i, j), (j, k) \in G'$. One can ask similar questions for longer increasing paths.

Problem 4.2. For any $\varepsilon > 0$ does there exist an ordered graph G such that if $G' \subseteq G$ with $|G'| \ge \left(\frac{1}{3} + \varepsilon\right) |G|$, then G' contains an increasing path of length three?

Note that in regards to Problem 4.2, one can consider a random coloring c of V(G) with colors $\{0, 1, 2\}$ and define G' to be the collection of all $(i, j) \in E(G)$ with i < j and c(i) < c(j). Then such G' on average contains $\frac{1}{3}|G|$ edges and has no increasing paths of length three, motivating constant $\frac{1}{3}$ in the problem.

[‡]the result follows from the proof of Theorem 1 in [5]

 $^{{}^{\$}\}alpha(\operatorname{Sh}_{n}^{4}[G]) \geq \frac{3}{8}|G|$ can be proved by considering a random colouring $c:[n] \to \{0,1\}$ and forming an independent set in $\operatorname{Sh}_{n}^{4}$ by taking hyperedges of G of form 1000, 1110, or x01y for some $x, y \in \{0,1\}$.

A. ARMAN, V. RÖDL, AND M. T. SALES

References

- C. Avart, B. Kay, C. Reiher, and V. Rödl, The chromatic number of finite type-graphs, J. Combin. Theory Ser. B 122 (2017), 877–896, DOI 10.1016/j.jctb.2016.10.004. ↑*
- [2] J. Czipszer, P. Erdős, and A. Hajnal, Some extremal problems on infinite graphs, Magyar Tud. Akad. Mat. Kutató Int. Közl. 7 (1962), 441–457 (Russian). [↑]1, 3
- [3] P. Erdős and A. Hajnal, On chromatic number of infinite graphs, *Theory of Graphs* (Proc. Colloq., Tihany, 1966), Academic Press, New York, 1968, pp. 83–98. [↑]1
- [4] P. Erdős and A. Hajnal, Some remarks on set theory. IX. Combinatorial problems in measure theory and set theory, *Michigan Math. J.* 11 (1964), 107–127. [↑]1, *
- [5] P. Erdős, A. Hajnal, and E. Szemerédi, On almost bipartite large chromatic graphs, *Theory and practice of combinatorics*, North-Holland Math. Stud., vol. 60, North-Holland, Amsterdam, 1982, pp. 117–123. [↑]1, 4, [‡]
- [6] C. C. Harner and R. C. Entringer, Arc colorings of digraphs, J. Combinatorial Theory Ser. B 13 (1972), 219–225, DOI 10.1016/0095-8956(72)90057-3. ↑*
- S. Janson, T. Luczak, and A. Rucinski, Random graphs, Wiley-Interscience Series in Discrete Mathematics and Optimization, Wiley-Interscience, New York, 2000. [↑]4

Appendix

Proof of Claim 2.2. Let $B_{\varepsilon}(n,d) = G$, where G is a random graph between S and L obtained by selecting a random subset of size $\frac{n^2}{2^{d+1}}$ without replacement from $K_{S,L}$ (complete bipartite graph between S and L). Then G satisfies (i) and we will show that G satisfies (ii) almost surely.

For every $X \subseteq S$ and $Y \subseteq L$, $e(X,Y) = e_G(X,Y)$ is distributed as a hypergeometric random variable $\operatorname{H}\left(\frac{n^2}{4}, \frac{n^2}{2^{d+1}}, |X||Y|\right)$ with expectation $\frac{1}{2^{d-1}}|X||Y|$. Let $B_{X,Y}$ be the event that

$$\left| e_G(X,Y) - \frac{1}{2^{d-1}} |X| |Y| \right| > \frac{\varepsilon n^2}{2^{d+2}},$$

i.e., $B_{X,Y}$ is the event that (ii) fails for given X and Y.

We will use a concentration inequality for hypergeometric random variables (this version is a corollary of Theorem 2.10 and inequalities (2.5), (2.6) of Janson, Luczak, Rucinski [7]).

Theorem 4.3. Let $Z \sim H(N, m, k)$ be a hypergeometric random variable with the expectation $\mu = \frac{mk}{N}$, then for $t \ge 0$

$$\mathbb{P}(|Z - \mu| > t) \leq 2 \exp\left(\frac{-t^2}{2(\mu + t/3)}\right).$$

For a given $X \subseteq L$ and $Y \subseteq R$, as a consequence of Theorem 4.3 with $Z = e_G(X, Y)$, $t = \frac{\varepsilon n^2}{2^{d+2}}$ and $\mu = \frac{1}{2^{d-1}}|X||Y| \leq \frac{n^2}{2^{d+1}}$ we get

$$\mathbb{P}(B_{X,Y}) = e^{-\Omega(n^2)},$$

where constant in $\Omega()$ term depends on ε and d only. Therefore,

$$\mathbb{P}\left(\bigcup_{X,Y} B_{X,Y}\right) \leqslant \sum_{X,Y} \mathbb{P}\left(B_{X,Y}\right) \leqslant 2^{n} e^{-\Omega(n^{2})} = o(1).$$

In particular, $\mathbb{P}(G \text{ satisfies (ii)}) = \mathbb{P}\left(\bigcap_{X,Y} \overline{B_{X,Y}}\right) = 1 - o(1)$. Hence, G almost surely satisfies (ii).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANITOBA, WINNIPEG, MB R3L0J7, CANADA *Email address*: andrii.arman@umanitoba.ca

DEPARTMENT OF MATHEMATICS, EMORY UNIVERSITY, ATLANTA, GA 30322, USA *Email address:* rodl@mathcs.emory.edu

DEPARTMENT OF MATHEMATICS, EMORY UNIVERSITY, ATLANTA, GA 30322, USA *Email address*: marcelo.tadeu.sales@emory.edu