Invariant Gorenstein Rings On 3-Dimensional Vector Spaces

By: Tamir Buqaie

Supervised by: Prof. Amiram Braun

Thesis submitted as partial fulfillment of the requirements for the masters degree

> University of Haifa Faculty of Natural Sciences Department of Mathematics

> > October, 2018

Approved by: Date: (Supervisor) Approved by: Date:

(Chairperson of M.sc. Committee)

Acknowledgments

I would like to thank my Master thesis advisor, Professor Amiram Braun for his dedication, interest, professionality and human attitude.I appreciate his constructive comments which helped me to learn and improve this paper.

I would like to thank my first mathematics teacher,Osima Shehady for her creative,successful and outstanding education methods. I thank my family and my friends,for their support, understanding and for giving me positive energy.

Contents

Invariant Gorenstein Rings On 3-Dimensional Vector Spaces

Tamir Buqaie

Supervised by: Prof. Amiram Braun

Abstract

For a finite subgroup $G \subset SL(V)$, let $S(V)^G$ denote the subring of G-invariants, where V is a 3– dimensional G – module over F and $S(V)$ the symmetric algebra of V. In this thesis we discuss the modular case of invariant theory,that is,when $charF = p$ divides |G|. The aim of this work is to classify **Gorenstein** ring of the form $S(V)^G$.

A.Braun in [\[2\]](#page-47-1) and Fleischmann-Woodcock in [\[6\]](#page-47-2) proved that if $S(V)^G$ is Cohen-Macaulay ring and $S(V)^{W(G)}$ is a polynomial ring then $S(V)^G$ is Gorenstein if and only if $G/W(G) \subseteq SL(m/m^2)$, where $W(G)$ is the G-subgroup generated by all pseudoreflections and m is the maximal homogenous ideal of $S(V)^{W(G)}$. (See [Theorem [1.2\]](#page-7-1)).

Here, since $G \subset SL(V)$, we have one kind of pseudo-reflection ,called transvection, in other words $W(G) = T(G)$, where $T(G)$ denote the $G-$ subgroup generated by all transvections of G.This work is divide into 7 chapters, the classification of $S(V)^G$, in each chapter, depends on the previous theorem which provides a necessary and suffiecient condition for G such that $S(V)^G$ is Gorenstein. So to apply this theorem we might ask ourselves, whether $S(V)^{T(G)}$ is polynomial ring and what are the $T(G)$ −invariants of this ring?Or at least, what are the degrees of the $T(G)$ −invariants ? Since $dim_F V = 3$, the Cohen-Macaulay property of $S(V)^G$ holds [\[9,](#page-47-3) Proposition 5.6.10].These are some of the classical questions that invariant theorists have studied in the modular case.

We prove the following theorem : Let $G \subset SL(3, \mathbb{F}_p)$ be a finite group and $V = \mathbb{F}_p^3$ is a reducible *G*-module, then $S(V)^G$ is Gorenstein. (See [Theorem [1.3\]](#page-8-0)).

We explain now how this work is organized.

As mentioned above, we divide our work into 7 chapters. In chapters A, B, D, E, F and G we fix a basis for V, and represent each $g \in G$, by matrices, with respect to this basis. Then we compute the ring of $T(G)$ −invariants. We are mainly concered with the case of $S(V)^{T(G)}$ being a polynomial ring. Under the assumptions of each chapter we prove that $S(V)^{T(G)}$ is a polynomial ring . Generally, we can not confirm in chapter D that the polynomial property of $S(V)^{T(G)}$ holds. We provide in chapters A, B, D , and G, a sufficient and necessary condition on G such that $S(V)^G$ is Gorenstein, when $G \subset SL(3, F)$. However, in chapter E and F, we exclusively consider the case $G \subset SL(3, \mathbb{F}_n)$. The technique in this work will be to exhibit the action of the group G on m/m^2 and using [Theorem [1.2\]](#page-7-1).

In order to prove that $S(V)^{T(G)}$ is a polynomial ring , in most cases, we use Kemper's theorem [\[4,](#page-47-4) Theorem 3.7.5].

In chapter A, we consider V, a reducible G – module of the form $V = F v \oplus W$ and Fv , W are G-submodules with $dim_F W = 2$. This is a special chapter, since by **Proposition [2.3](#page-10-0)**, the Gorenstein property of $S(V)^G$ depends on W. So we use the computation of [\[3\]](#page-47-5),and proceed our proof in steps by dividing it into 3 cases, which are:

- 1. W is an irreducible and primitive $T(G)$ module.
- 2. W is a reducible $T(G)$ module.
- 3. W is an irreducible and imprimitive $T(G)$ module.

In fact, the results in chapter A are obtained from the discussion in [\[3\]](#page-47-5). The difference between the situation here and the situation of [\[3\]](#page-47-5), is that in [3], $G \subset SL(2, F)$ and here $G = G|W \subset GL(2, F)$.

In chapter B, we consider V, a reducible, indecomposable G – module which has at least two G-submodules W_1, W_2 where $dim W_i = 2$ for $i = 1, 2$. In addition to the main theorem, we prove, under specific condition, that $S(V)^G$ is Gorenstein in the following cases (see [Theorem [3.5\]](#page-17-0)) :

(i)
$$
p = 3
$$
.

- (ii) $|G|$ is prime to 3.
- (iii) $F = \mathbb{F}_q$, where $q = p^s$ and $g.c.d(3, s) = 1$.

We define a F-vector space X, where $dim_F X \in \{1,2\}$, and separate this chapter into two cases. In the first case, we deal with $dim(span_F X) = 1$. In the second case, we deal with $dim(span_F X) = 2$, where $T(G)$ is a direct product of two transvection groups $T_1(G), T_2(G)$. In Lemma [3.7](#page-18-0), Lemma [3.8](#page-18-1), and Proposition [3.9](#page-19-0) we examine the normality of $T_1(G), T_2(G)$. However this is not needed for **Proposition [3.10](#page-20-0)** ,Proposition [3.13](#page-23-0), and Corollary [3.14](#page-25-0).

Chapter C is different, we provide in this chapter preliminaries to chapters D, E, F and G.

In chapter D, we consider V, a reducible and indecomposable G−module which has a unique G-submodule W with $dim_F W = 2$, and V contains no 1–dimensional G-submodules. In this chapter, we can not conclude that $S(V)^{T(G)}$ is always polynomial ring. So we make specific assumption (see **Assumption [5.2](#page-31-1)**) to prove [Theorem 1.3. This is applicable in case $F = \mathbb{F}_p$.

In chapter E, we consider V, a reducible and indecomposable G – module which has a unique 1–dimensional G–submodule $U = Fv_0$ and V contains no 2–dimensional G–submodules.Hence $W = V/U$ is an irreducible G– module. We exclusively consider here the case $F = \mathbb{F}_p$.

There are two cases of W that Lemma [6.2](#page-36-0) have provided . The first case is when W is an unfaithful irreducible $T(G)$ –module, and the second one is when W is a faithful irreducible $T(G)$ −module.

In chapter F, we consider V, a reducible and indecomposable G−module which has a unique G–submodule W with $dim_F W = 2$, and a unique 1–dimensional G–submodule $F v_0$, and $F v_0 \subset W \subset V$. We exclusively consider here the case $F = \mathbb{F}_p$.

Finally, in chapter G, we consider V, a reducible and indecomposable G – module and contains (at least) two 1–dimensional G –submodules.By using **Proposition [4.3](#page-28-0)**, we provide a sufficient and necessary condition on G for the Gorenstein property of $S(V)^G$, and present the main result in **Corollary [8.1](#page-46-1)**.

Index of notations

- \mathbb{F}_p The prime field of order p.
- $|G|$ The order of group G .
- $F-$ A finite field with $char F = p$.
- \bar{F} − The algebraic closure of F.
- $S(V)$ The symmetric algebra of V.
- $S(V)^G$ The subring of $G-$ invariants.
- $T(G)$ The G–subgroup generated by all transvections.
- $W(G)$ The G–subgroup generated by all pseudoreflections.
- $m-$ The unique homogenous maximal ideal of $S(V)^{T(G)}$.
- V^* The hom-dual $hom_F(V, F)$.
- $GL(V)$ The general linear group of V.
- $GL(n, F)$ The linear automorphisms of $Fⁿ$.
- $SL(V)$ The special linear group of V.
- $SL(n, F)$ The linear automorphisms of $Fⁿ$ of determinant 1.
- $G|W-$ The restriction of G to W.
- $<\sigma_1, ..., \sigma_s>$ The subgroup generated by $\{\sigma_1, ..., \sigma_s\}$.
- $Fix_G(W)$ The elements of G, that acts trivially on subspace W.

1 Introduction

Let $G \subset SL(V)$ be a finite subgroup of SL(V). Let F be a field with $char F = p > 0$, V a 3– dimensional F– vector space and $S(V)$ the symmetric algebra of V. We denote by $S(V)^G$ the subring of G − *invariants*. One of the main purposes of this thesis is to classify Gorenstein rings of the form $S(V)^G$, when p divides |G|.

The following are results of K.Watanabe ,Fleischmann-Woodcock and A.Braun .

Theorem 1.1. $(K. Watanabe [10], [11])$ $(K. Watanabe [10], [11])$ $(K. Watanabe [10], [11])$ $(K. Watanabe [10], [11])$ $(K. Watanabe [10], [11])$

Suppose that $(|G|, p) = 1$ and G contains no pseudo-reflection. Then the following are equivalent:

1. $G \subset SL(V)$.

2. $S(V)^G$ is a Gorenstein ring.

Theorem 1.2. $(A.Braun [2], Fleischmann-Woodcock [6])$ $(A.Braun [2], Fleischmann-Woodcock [6])$ $(A.Braun [2], Fleischmann-Woodcock [6])$ $(A.Braun [2], Fleischmann-Woodcock [6])$ $(A.Braun [2], Fleischmann-Woodcock [6])$

Suppose that $S(V)^G$ is Cohen-Macaulay and $S(V)^{W(G)}$ is a polynomial ring. Then $S(V)^G$ is Gorenstein iff $G/W(G) \subseteq SL(m/m^2)$, where m is the unique homogenous maximal ideal of $S(V)^{W(G)}$ and $W(G)$ is the G – subgroup generated by all pseudo-reflections(of all types).

Here an element $q \in G$ is called pseudoreflection if $rank(q - I) = 1$. If $(|G|, p) = 1$ then every pseudoreflection is diagonalizable with one's on all but a single diagonal entry. In the modular case, that is if p divides $|G|$ we have a new type of pseudoreflection, called transvection, having the additional property $(g - I)^2 = 0$, so it is not diagonalizable.Denote by $T(G)$ the $G - subgroup$ generated by all transvections .

Eagon and Hochster proved in [\[5\]](#page-47-8) that $S(V)^G$ is Cohen-Macaulay for all nonmodular groups G, i.e when $(|G|, p) = 1$. But there are finite $p - groups$, such that $S(V)^G$ is not Cohen-Macaulay.

Example 1.1. Let F be a field of characteristic p and V is a faithful representation of a non-trivial p-group P and mV denote the faithful representation of G formed by taking the direct sum of m copies of V. Then $S(mV)^P$ is not Cohen-Maculay when $m \geq 3$. [\[12,](#page-47-9) Theorem 1.2]

However in this thesis we assume that $dim_F V = 3$ and by [\[9,](#page-47-3) Proposition 5.6.10] $S(V)^G$ is always Cohen-Macaulay ring.

It is shown in [\[3,](#page-47-5) Section 2] that $S(V)^G$ can fail to be Gorenstein even if $G \subset SL(V)$ and $dim V = 2$. Moreover, [\[3\]](#page-47-5) provides necessary and sufficient conditions to ensure the Gorenstein property of $S(V)^G$ in case $dim_F V = 2$ and $G \subset SL(V)$. We mostly examine here the case of $dim_F V = 3, G \subset SL(V)$ and V is a reducible G-module. The analysis is carried out by separation into different cases. A noteworthy consequence is perhaps the following:

Theorem 1.3. Let $G \subset SL(3, \mathbb{F}_p)$ be a finite group and $V = \mathbb{F}_p^3$ is a reducible G-module. Then $S(V)^G$ is Gorenstein.

This extends a similar result in [\[3,](#page-47-5) Corollary (C)].The separation into cases is summed up in the following table.

Convention: Given a basis $\{v_1, ..., v_n\}$ of V, and $g \in G$. Then $g(v_i) = \sum_{j=1}^n a_{ij}v_j, a_{ij} \in F$ and the matrix $A := (a_{ij})$, representing g, acts on coordinates (=row vectors) from the right.

2 Chapter $(A): V = Fv \oplus W$, and Fv, W are G-submodules.

In this chapter we consider V, a reducible $G - module$ of the form $V = F v \oplus W$ and $F v$, W are G – modules with $dim_F W = 2$. **Proposition [2.3](#page-10-0)** provides a sufficient and necessary condition for G as above, to ensure the Gorenstein property of $S(V)^G$. Our proof will proceed in steps by dividing it into 3 cases:

- 1. W is an irreducible and primitive $T(G)$ module.
- 2. W is a reducible $T(G)$ module.
- 3. W is an irreducible and imprimitive $T(G)$ module.

The next two results are usefull in this chapter.

Lemma 2.1. W is a faithful G -module.

Proof. Let $1_G \neq g \in G$, with $g|W = Id$. Then

$$
g = \left(\begin{array}{rr} \lambda(g) & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right).
$$

Where $g(v) = \lambda(g)v$, for all $g \in G$. Since $G \subset SL(V)$, $\lambda(g) = 1$ and therefore

$$
g = \left(\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right).
$$

A contradiction.

The next result does not require the assumption $V = F v \oplus W$.

Lemma 2.2. Suppose that $G \subset SL(V)$ and U a 1-dimensional G -submodule. Then $U \subseteq \text{ker}(\sigma - I)$ for all transvection σ in G. Equivalently, $T(G)$ acts trivially on U.

Proof. Let $\sigma \in T(G)$ be a transvection $\sigma \neq I$, and $M = ker(\sigma - I)$. Suppose that $Fu = U \not\subset M$. Then with respect to a basis $\{u, m_1, m_2\}, m_1, m_2 \in M$, we have: $\sigma =$ $\sqrt{ }$ \mathcal{L} $\lambda(\sigma)$ 0 0 0 1 0 0 0 1 \setminus , where $\sigma(u) = \lambda(\sigma)u$ and $\sigma|M = Id$. But $G \subset SL(V)$, therefore $\lambda(\sigma) = 1$, implying that $\sigma = I$, a contardiction.

 \Box

Proposition 2.3. Let $G \subset SL(V)$ be a finite group and $V = Fv \oplus W$ a decomposition of V into $G-submodules$ with $dim_F W = 2$. Then the following are equivalent:

- 1. $S(V)^G$ is Gorenstein.
- 2. $det(g|W) = det(g|m/m^2)$, for each $g \in G$ where m is the unique homogenous maximal ideal of $S(W)^{T(G)}$.

Proof. By Lemma [2.2](#page-9-1) $S(V)^{T(G)} = S(W)^{T(G)}[v]$. Now by [\[8\]](#page-47-10), $S(W)^{T(G)} = F[a_1, a_2]$ is a polynomial ring, where $m = (a_1, a_2)$. Let P be the unique homogenous maximal ideal of $S(V)^{T(G)}$, then P = (a_1, a_2, v) . By [Theorem [1.2\]](#page-7-1) $S(V)^G$ is Gorenstein if and only if $1 = det(g|P/P^2) = \lambda(g)det(g|m/m^2)$, where $g(v) = \lambda(g)v$, for all $g \in G$. Since $G \subset SL(V)$, $det(g|W) = \lambda(g)^{-1}$, hence $S(V)^G$ is Gorenstein if and only if $det(g|W) = det(g|m/m^2)$, for each $g \in G$.

The transvections group $T(G)|W$ had been classified by Kantor whenever $T(G)|W$ is an irreducible primitive linear group [\[7,](#page-47-11) Theorem 1.5]. So with the above notation one of the following holds:

(i) $T(G)|W = SL(2, \mathbb{F}_q)$, where $p|q$.

(ii)
$$
T(G)|W \cong SL(2, \mathbb{F}_5), T(G)|W \subset SL(2, \mathbb{F}_9)
$$
 and $\mathbb{F}_9 \subseteq F$.

We now consider the above case (i).

Proposition 2.4. Suppose $G \subset SL(V)$ is a finite group with $T(G)|W = SL(2, \mathbb{F}_q)$,where $q = p^s$ and $\mathbb{F}_q \subseteq F$. Let $V = F^3 = Fv \oplus W$ be the natural $SL(3, F)$ – module. Then $S(V)^G$ is Gorenstein if and only if $G|W \subset GL(2, \mathbb{F}_{q^2})$.

Proof. Recall as in [\[3,](#page-47-5) Proposition 3.5] that $S(W)^{T(G)} = S(W)^{SL(2, \mathbb{F}_q)} = F[u, c_{21}], deg(u) = q +$ $1, deg(c_{21}) = q^2 - q$. Let $g \in G$ be arbitrary and set $g|W = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with respect to the basis $\{x,y\}$. So $g^{-1}|W = \frac{1}{\det(g|W)} \left(\begin{array}{cc} d & -b \\ -c & a \end{array}\right)$. Since $gT(G)g^{-1} = T(G)$, we get after restriction to W that $(g|W) \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ $(g^{-1}|W) = \frac{1}{det(g|W)} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} = \frac{1}{det(g|W)} \begin{pmatrix} ad-bc-ac & a^2 \\ -c^2 & ad-bc \end{pmatrix}$ $\frac{-bc-ac}{-c^2}$ $\frac{a^2}{ad-bc+ac}$) is in $SL(2,\mathbb{F}_q)$. Similarly, $(g|W) \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} (g^{-1}|W) = \frac{1}{det(g|W)} \begin{pmatrix} ad-bc+bd & -b^2 \\ d^2 & ad-bc \end{pmatrix}$ $\frac{bc + bd}{d^2} \frac{-b^2}{ad - bc - bd}$ $\in SL(2, \mathbb{F}_q)$. Set $e = det(g|W) =$ $ad - bc$. Then $\frac{ac}{e}, \frac{a^2}{e}$ $\frac{u^2}{e}, \frac{c^2}{e}$ $\frac{b^2}{e}, \frac{b^2}{e}$ $\frac{b^2}{e}, \frac{d^2}{e}$ $\frac{d^2}{e}, \frac{bd}{e} \in \mathbb{F}_q$. Assume firstly that $a \neq 0$. Then $\gamma := \frac{c}{a} = \frac{(\frac{c^2}{c})^2}{(\frac{c^2}{c})^2}$ $\frac{(\frac{ac}{e})}{(\frac{ac}{e})} \in \mathbb{F}_q$. If $b \neq 0$ then $\mu := \frac{d}{b} = \frac{\left(\frac{bd}{e}\right)}{\left(\frac{b^2}{e}\right)}$ $\frac{\sqrt{e}}{\sqrt{b^2}}$ $\in \mathbb{F}_q$. Hence $e = ad - bc = ab\mu - ba\gamma = ab(\mu - \gamma)$ and therefore $\beta := \frac{b}{a} = \frac{ab(\mu - \gamma)}{a^2(\mu - \gamma)} = \frac{e}{a^2(\mu - \gamma)} \in \mathbb{F}_q$, as well as $\delta := \frac{d}{a} = \frac{\mu b}{a} = \mu \beta \in \mathbb{F}_q$. If $b = 0$ then take $\beta = 0$ and $\delta := \frac{d}{a} = \frac{ad}{a^2} = \frac{e}{a^2} \in \mathbb{F}_q$. So in both cases $b = a\beta$, $c = a\gamma$, $d = a\delta$, β , γ , $\delta \in \mathbb{F}_q$. Therefore $g|W = \left(\begin{smallmatrix} a & 0 \\ 0 & a \end{smallmatrix}\right) \left(\begin{smallmatrix} 1 & \beta \\ \gamma & \delta \end{smallmatrix}\right), \text{ with } h := \left(\begin{smallmatrix} 1 & \beta \\ \gamma & \delta \end{smallmatrix}\right) \in GL(2, \mathbb{F}_q).$

If $a = 0$ then $e = det(g|W) = -bc$ implies $b \neq 0 \neq c$. So we get equality $g|W = \begin{pmatrix} c & 0 \\ 0 & c \end{pmatrix} \begin{pmatrix} 0 & \beta \\ 1 & \delta \end{pmatrix}$ with c replacing a and $\beta, \delta \in \mathbb{F}_q$. Consequently if $a \neq 0$ we get: $g(c_{21}) = (aI_W)(h(c_{21})) = a^{q^2-q}c_{21}$, and as in the proof of [\[3,](#page-47-5) Proposition 3.5] we get that $g(u) =$ $(aI_W)(h(u)) = det(h)a^{degu}u = det(h)a^{q+1}u$. But $det(g|W) = a^2det(h)$, hence $g(u) = det(g|W)a^{q-1}u$.

Let $M = Fu + Fc_{21}$ be the 2-dimensional subspace of the polynomial ring $S(W)^{T(G)} = F[u, c_{21}], m =$ (u, c_{21}) , so $m/m^2 \cong M$. Then the matrix representing g|M with respect to the basis $\{u, c_{21}\}$ is $\int_{a}^{a^{q-1} \det(g|W)} \frac{0}{a^2}$ $\frac{e^{ct}(g|W)}{a^{q^2-q}}$. Therefore the condition of **Proposition [2.3](#page-10-0)** : $det(g|W) = det(g|m/m^2) =$ $det(g|M)$ is equivalent to: $det(g|W) = a^{q-1}det(g|W) a^{q^2-q} = a^{q^2-1}det(g|W)$. Hence it is equivalent to $a^{q^2-1}=1$, namely $a \in \mathbb{F}_{q^2}$. Using $h \in GL(2, \mathbb{F}_q)$, this is also equivalent to $g|W = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL(2, \mathbb{F}_{q^2})$. A similar conclusion is obtained if $a = 0$.

We next consider the above case (ii).

Proposition 2.5. Suppose $G \subset SL(3, F)$ is a finite group with $T(G)|W \cong SL(2, \mathbb{F}_5)$, $T(G)|W \subseteq$ $SL(2, \mathbb{F}_9)$ and $\mathbb{F}_9 \subseteq F$ (here $p = 3$). Let $V = F^3 = Fv \oplus W$ be the natural $SL(3, F)$ – module. Then $S(V)^G$ is Gorenstein iff $G|W \subseteq \langle T(G)|W, \eta I_W \rangle$, where η is a 20^{th} -primitive root of unity.

Proof. Let $g \in G$, choose $\xi \in \overline{F}$ such that $\xi^2 det(g|W) = 1$. Hence $\hat{g} := \xi I_W g|W \in SL(2, \overline{F})$ where \overline{F} is the algebraic closure of F.Clearly $\hat{g}(T(G)|W)\hat{g}^{-1} = T(G)|W$. Therefore the computation in [\[3,](#page-47-5) Proposition 3.7], with \hat{g} instead of $g|W$, shows that $\hat{g} \in T(G)|W$. Hence $\binom{\xi}{0} = \hat{g}(g^{-1}|W) \in$ $(T(G)|W)(G|W) = G|W$. Consequently $G|W \subseteq \leq T(G)|W, (\begin{smallmatrix} \lambda & 0 \\ 0 & \lambda \end{smallmatrix})|\lambda \in \overline{F} >$.

Now $(\frac{\xi}{\theta})g|W = \hat{g}|W \in T(G)|W$. Recall from [\[3,](#page-47-5) Proposition 6.6.] that $S(W)^{SL(2,\mathbb{F}_5)}$ = 0 ξ $F[f_{10}, f_{12}]$, where $f_{10} = x_1^9 x_2 - x_1 x_2^9$ and $f_{12} = x_1^{12} + x_1^{10} x_2^2 - x_1^6 x_2^6 + x_1^2 x_2^{10} - x_2^{12}$. Hence $\hat{g}(f_{10}) =$ $f_{10}, \hat{g}(f_{12}) = f_{12}$ implying that $g(f_{10}) = \xi^{-10} f_{10}, g(f_{12}) = \xi^{-12} f_{12}.$

Let $U = F f_{10} + F f_{12}$ be the 2-dimensional subspace of the polynomial ring $S(W)^{SL(2,\mathbb{F}_5)}$ $F[f_{10}, f_{12}], m = (f_{10}, f_{12}),$ so $m/m^2 \cong U$. Then the matrix representing g|U with respect to the basis $\{f_{10}, f_{12}\}\$ is $\left(\begin{array}{cc} \xi^{-10} & 0 \\ 0 & \xi^{-1} \end{array}\right)$ $\frac{0}{\sqrt{2}}$ ($\frac{0}{\sqrt{2}}$). Therefore, by **Proposition [2.3](#page-10-0)** $det(g|W) = det(g|m/m^2) = det(g|U)$ is translated into $\xi^{-2} = \xi^{-22}$ or $\xi^{20} = 1$. Therefore $S(V)^G$ is Gorenstein iff $G|W \subseteq \langle T(G)|W, \eta I_W \rangle$ where $\eta \in \overline{F}$ is a primitive 20 – th root of unity.

We now consider case 2, namely the possibility of W being a reducible $T(G)$ – module.

Proposition 2.6. Let $G \subset SL(3, F)$ be a finite group. Let $V = F^3 = Fv \oplus W$ be the natural $SL(3, F)$ – module and suppose W is a **reducible** $T(G)$ -submodule. Then $S(V)^G$ is Gorenstein if and only if $G|W \subseteq \{(\begin{smallmatrix} a & b \\ 0 & d \end{smallmatrix}) | a^{p^n-1} = 1, a, b, d \in F\}.$

Proof. Set $\hat{g} := \xi I_W g |W$, with $det(\hat{g}) = \xi^2 det(g|W) = 1$. Since W is a reducible $T(G)|W$ module, there exists a basis $\{x_1, x_2\}$ of W with respect to which $T(G)|W \subseteq \left(\begin{smallmatrix} F & F \ 0 & F \end{smallmatrix}\right)$. Every transvection acts trivially on every one dimensional submodule of W by Lemma [2.2](#page-9-1). Hence it must stabilize Fx_2 . Therefore since $T(G)|W \subset SL(W)$, $T(G)|W \subseteq \left(\begin{smallmatrix} 1 & F \\ 0 & 1 \end{smallmatrix}\right)$. Consequently $T(G)|W$ is elementary abelian

 \Box

group.So $T(G)|W=<\sigma_1>\times\cdots\times<\sigma_n$ >, where $\sigma_i=\begin{pmatrix}1 & \alpha_i\\ 0 & 1\end{pmatrix}$, for $i=1,...,n$, and $\{\alpha_1,...,\alpha_n\}$ are linearly independent over \mathbb{F}_p . Let $g|W = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Now the computation in the proof of [\[3,](#page-47-5) Theorem 3.1] with \hat{g} instead of $g|W$ shows that $\xi c = 0$, hence $c = 0$, $\hat{g} = \begin{pmatrix} \lambda & \pi \\ 0 & \lambda^{-1} \end{pmatrix}$.

Hence $\hat{g}|m/m^2 = \begin{pmatrix} \lambda^{p^n} & 0 \\ 0 & \lambda^{-1} \end{pmatrix}$, where $\xi a = \lambda$, $\xi d = \lambda^{-1}$, $\pi \in F$ and $m = (b, x_2)$, where b which is of degree p^n is defined as in the proof of [\[3,](#page-47-5) Theorem 3.1]. Hence $\hat{g}(b) = \lambda^{p^n} b = (\xi a)^{p^n} b = (\xi)^{p^n} g(b)$ and $\hat{g}(x_2) = \lambda^{-1}x_2 = (\xi d)x_2 = \xi g(x_2)$. Therefore $g|m/m^2 = \left(\begin{matrix} a_0^{\xi_0} & 0 \ 0 & d \end{matrix}\right)$. Consequently $det(g|m/m^2) =$ $a^{p^n}d = a^{p^n-1}(ad) = a^{p^n-1}det(g|W)$. Hence, by **Proposition [2.3](#page-10-0)** $\widetilde{S}(V)^G$ is Gorenstein if and only if $a^{p^n-1} = 1$. In other words $S(V)^G$ is Gorenstein if and only if $G|W \subseteq \{ \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} | a^{p^n-1} = 1, b, d \in F \}.$ \Box

Our next objective is to handle case 3, i.e when $T(G)|W$ acts irreducibly and imprimitively on W. It is well known(e.g.[\[13\]](#page-47-12)) that $T(G)|W$ is a monomial subgroup. Recall that $H \subset GL(W)$ is called monomial if W has a basis with respect to which the matrix of each element of H has exactly one non-zero entry in each row and column. If $char F \neq 2$, then by [\[3,](#page-47-5) Lemma 3.9.] $T(G)|W = 1$. This contradicts the assumption that W is an irreducible $T(G)|W$ –module.Hence we only deal with $p = 2$.

Proposition 2.7. Assume that charF = 2. Let $G \subset SL(3, F)$ be a finite group with $T(G)|W$ acting imprimitively and irreducibly on the submodule W and $V = Fv \oplus W$. Then $S(V)^G$ is Gorenstein iff $G|W \subseteq \mathcal{F}(G)|W, \delta I_W >$, where δ is a primitive d – th root of unity and d defined in the proof.

Proof. Let $\{x_1, x_2\}$ be the basis of W with respect to which $T(G)|W$ is irreducible and monomial. Let $\{g_1, ..., g_n\}$ be the set of transvections generating $T(G)|W$. The $T(G)|W-\$ irreducibility of W forces $n \geq 2$. So either $g_i = \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}$ or $\begin{pmatrix} 0 & * \\ * & 0 \end{pmatrix}$. The first possibility leads, as in [\[3,](#page-47-5) Lemma 3.9.] to $g_i|W = I_W$. So $g_i = \begin{pmatrix} 0 & \alpha_i \\ \alpha_i^{-1} & 0 \end{pmatrix}$, for $i = 1, ..., n$, with respect to the basis $\{x_1, x_2\}$ of W. Let n_{ij} be the minimal number such that $(g_i g_j)^{n_{ij}} = 1$, where $i \neq j$, clearly $g_i^{-1} = g_i$ for $i = 1, ..., n$. Also $n_{ij} = n_{ji} > 1$ and since $char F = 2$, n_{ij} is odd. Recall that $S(W)^{T(G)}$ is a polynomial ring [\[8,](#page-47-10) Theorem 2.4]. We next compute the actual generators of $S(W)^{T(G)}$.

Let $[(T(G)|W), (T(G)|W)] = \langle g_i g_j g_i g_j \rangle$ be the commutator subgroup of $T(G)|W$. We have $(g_1g_2)^{n_{12}} = 1$. Therefore $g_2 = (g_1g_2)^{n_{12}}g_2 = (g_1g_2)^{n_{12}-1}(g_1g_2)g_2 = (g_1g_2)^{n_{21}-1}g_1$. But $n_{12}-1$ is even, since n_{12} is odd .Hence $g_2 = (g_1g_2g_1g_2)^{\frac{n_{12}-1}{2}}g_1 \in [T(G)|W,T(G)|W]g_1$. This similarly holds for $g_i, i \geq$ 3, hence $\frac{|(T(G)|W)|}{|([T(G)|W,T(G)|W])|} = 2$. Let $d = l.c.m\{n_{ij}|i < j\}$. Since $g_ig_jg_ig_j = \begin{pmatrix} \alpha_i\alpha_j^{-1}\alpha_i\alpha_j^{-1} & 0 \\ 0 & \alpha_i^{-1}\alpha_i\alpha_j^{-1} \end{pmatrix}$ $\alpha_i \alpha_j \atop 0 \quad \alpha_i^{-1} \alpha_j \alpha_i^{-1} \alpha_j},$ it follows that $[(T(G)|W), (T(G)|W)] = {\binom{\binom{c}{0}}{c^{-1}}}\zeta^d = 1$. Since $(\alpha_i\alpha_j^{-1})^{n_{ij}} = 1$ it follows that $\alpha_i^{n_{ij}} = \alpha_j^{n_{ij}}$. Consequently $\alpha_i^d := \beta$ for $i = \{1, 2, 3, ..., n\}$.

Now we have:

1. $g_i(x_1^d + \alpha_i^d x_2^d) = g_i(x_1)^d + \alpha_i^d g_i(x_2)^d = \alpha_i^d x_2^d + \alpha_i^d (\alpha_i^{-1} x_1)^d = x_1^d + \alpha_i^d x_2^d$. 2. $g_i(x_1x_2) = g_i(x_1)g_i(x_2) = \alpha_i x_2 \cdot \alpha_i^{-1} x_1 = x_1 x_2.$

Hence ${x_1x_2, x_1^d + \beta x_2^d} \in S(W)_{\text{max}}^{T(G)}$ and $deg(x_1x_2)deg(x_1^d + \beta x_2^d) = 2d = |T(G)|W|$. So by [\[4,](#page-47-4) Theorem 3.7.5] we get that $S(W)^{T(G)} = F[x_1x_2, x_1^d + \beta x_2^d]$. Let $g|W = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Set $\hat{g} = \xi I_W g|W$, where $\xi^2 det(g|W) = 1$. Now the computation of [\[3,](#page-47-5) Theorem 3.10], with \hat{g} instead of $g|W$, shows that $\hat{q} \in T(G)|W$.

We have:

(i)
$$
g(x_1x_2) = { \xi^{-1} \ 0 \choose 0 \ \xi^{-1}} \hat{g}(x_1x_2) = \xi^{-2}x_1x_2.
$$

\n(ii) $g(x_1^d + \beta x_2^d) = { \xi^{-1} \ 0 \choose 0 \ \xi^{-1}} \hat{g}(x_1^d + \beta x_2^d) = { \xi^{-1} \ 0 \choose 0 \ \xi^{-1}} (x_1^d + \beta x_2^d) = \xi^{-d}(x_1^d + \beta x_2^d).$

Consequently, $g|m/m^2 = \begin{pmatrix} \xi^{-2} & 0 \\ 0 & \xi^{-2} \end{pmatrix}$ $\int_{0}^{2} \int_{\xi^{-d}}^{0}$, where $m/m^{2} = U = F(x_{1}x_{2}) + F(x_{1}^{d} + \beta x_{2}^{d})$. So $det(g|m/m^{2}) =$ $\xi^{-2}\xi^{-d} = det(g|W)\xi^{-d}$, and $det(g|W) = det(g|m/m^2)$ if and only if $\xi^{-d} = 1$. Consequently, by **Proposition [2.3](#page-10-0)** $S(V)^G$ is Gorenstein if and only if $G|W \subseteq \langle T(G)|W, \delta I_W \rangle$ where δ is a primitive $d - th$ root of unity.

 \Box

Corollary 2.8. Suppose $G \subset SL(3, \mathbb{F}_p)$, and $V = \mathbb{F}_p v \oplus W$ a decomposition of $V = \mathbb{F}_p^3$ into $G-submodules.$ Then $S(V)^G$ is Gorenstein.

Proof. This is a consequence of Proposition [2.4](#page-10-1), Proposition [2.6](#page-11-0) and Proposition [2.7](#page-12-0). \Box

3 Chapter (B) : *V* is indecomposable and contains (at least) two 2− dimensional $G-$ submodules W_1, W_2 .

In this chapter we consider V, a reducible, indecomposable G− module which has at least two Gsubmodules W_1, W_2 where $dim W_i = 2$ for $i = 1, 2$. Fix a basis $\{w_2, w_1, v\}$ such that $W_i = F w_i + F v$. Then $W_1 \cap W_2 = Fv$ is 1-dimensional G-submodule of V. Therefore for every $g \in G$ we have a matrix representation with respect to this basis.

$$
g = \left(\begin{array}{ccc} \lambda_2(g) & 0 & \delta_{13}(g) \\ 0 & \lambda_1(g) & \delta_{23}(g) \\ 0 & 0 & \lambda(g) \end{array}\right).
$$

Let $\sigma \in T(G)$ be a transvection . $\sigma : W_i \to W_i$ for $i = 1, 2$. So $\sigma(w_i) = \alpha_i w_i + \beta_i v$ where $\alpha_i, \beta_i \in F$. Since $| \langle \sigma \rangle | = p$ and σ acts trivially on Fv by Lemma [2.2](#page-9-1), $w_i = \sigma^p(w_i) = \alpha_i^p w_i + \gamma_i v$, hence $\gamma_i = 0$ and $\alpha_i^p = 1$, then $\alpha_i = 1$, for $i = 1, 2$. As a result, σ is represented by the matrix :

$$
\left(\begin{array}{ccc} 1 & 0 & a \\ 0 & 1 & b \\ 0 & 0 & 1 \end{array}\right)
$$

with respect the basis $\{w_2, w_1, v\}.$

Corollary 3.1. $T(G)$ is an elementary abelian group.

Proof. Suppose that τ is another transvection where

$$
\tau = \left(\begin{array}{ccc} 1 & 0 & a_1 \\ 0 & 1 & b_1 \\ 0 & 0 & 1 \end{array}\right).
$$

Then it easy to see that $\tau\sigma = \sigma\tau$. Therefore $T(G)$ is an elementary abelian group. Say it is minimaly generated by $\{\sigma_1, ..., \sigma_n\}$, where

$$
\sigma_i = \left(\begin{array}{ccc} 1 & 0 & a_i \\ 0 & 1 & b_i \\ 0 & 0 & 1 \end{array}\right)
$$

for $i = 1, ..., n$. So $|T(G)| = p^n$.

Let $X := \{ \begin{pmatrix} a_i \\ b_i \end{pmatrix} \in \mathbb{F}_p \mid i = 1, ..., n \},\$ then the elements of X are linearly independent over \mathbb{F}_p . Clearly $dim(span_F X) \in \{1,2\}.$

Case 1: $dim(span_F X) = 1$

Assume that $dim(span_F X) = 1$. So $\binom{a_i}{b_i} = \mu_i \binom{a}{b}$ for $i = 1, ..., n$. Where $\binom{a_1}{b_1} = \binom{a}{b}, \mu_1 = 1$ and $\mu_i \in F$ for $i = 2, ..., n$. Since the elements of X are linearly independent over \mathbb{F}_p , this is equivalent to $\{1, \mu_2, ..., \mu_n\}$ being of linearly independent over \mathbb{F}_n .

Lemma 3.2. $S(V)^{T(G)}$ is a polynomial ring.

Proof. We shall firstly change the basis of V. We have $\sigma_1(w_1) = w_1 + bv$ and $\sigma_1(w_2) = w_2 + av$, where $a = a_1$ and $b = b_1$. Hence $\sigma_1(aw_1 - bw_2) = a\sigma(w_1) - b\sigma(w_2) = a(w_1 + bv) - b(w_2 + av) = aw_1 - bw_2$. So $0 \neq (aw_1 - bw_2) \in S(V)^{< \sigma_1>}.$ We therefore change the basis of V into $\{w_2, aw_1 - bw_2, v\}$, and then σ_1 is represented by the matrix:

$$
\left(\begin{array}{rrr} 1 & 0 & a \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right).
$$

We also have:

 $\sigma_i(aw_1 - bw_2) = a\sigma_i(w_1) - b\sigma_i(w_2) = a(w_1 + b_i v) - b(w_2 + av_i) = aw_1 - bw_2 + (ab_i - ba_i)v$. But $\binom{a_i}{b_i} = \mu_i \binom{a}{b}$, and therefore $(ab_i - ba_i) = 0$. Hence $0 \neq (aw_1 - bw_2) \in S(V)^{< \sigma_i}$, for $i = 1, ..., n$. So each σ_i is represented by the matrix:

$$
\left(\begin{array}{rrr} 1 & 0 & a_i \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right)
$$

where $a_i = \mu_i a$. So we are in the same situation as in the modular binary case (see [\[3\]](#page-47-5)). Consequently, by [\[3,](#page-47-5) Lemma 3.13] $S(V)^{T(G)} = F[x, aw_1 - bw_2, v]$ is a polynomial ring, and x is an homogenous polynomial in $\{v, aw_1 - bw_2\}$ of degree p^n having $w_2^{p^n}$ $_2^p$ as one of its monomial.

We now need to shift to the action of G on $S(V)^{T(G)}$.

Proposition 3.3. $S(V)^G$ is Gorenstein if and only if every element of $G/T(G)$ has order dividing $|T(G)| - 1 = p^{n} - 1.$

Proof. If $a = 0$ or $b = 0$ we argue exactly as in the binary form paper (see[\[3\]](#page-47-5)). So we may assume that $a \neq 0$ and $b \neq 0$. Since $T(G) \lhd G$, $g\sigma_i g^{-1} \in T(G)$, $\forall i = 1, ..., n$. Hence

$$
g\sigma_i g^{-1} = \varphi_i = \left(\begin{array}{ccc} 1 & 0 & \epsilon_i a \\ 0 & 1 & \epsilon_i b \\ 0 & 0 & 1 \end{array}\right), \epsilon_i \in \mathbb{F}_p 1 + \mathbb{F}_p \mu_2 ... + \mathbb{F}_p \mu_n.
$$

So $g\sigma_i = \varphi_i g$ is written explicitly by:

$$
\begin{pmatrix}\n\lambda_2(g) & 0 & \delta_{13}(g) \\
0 & \lambda_1(g) & \delta_{23}(g) \\
0 & 0 & \lambda(g)\n\end{pmatrix}\n\begin{pmatrix}\n1 & 0 & \mu_i a \\
0 & 1 & \mu_i b \\
0 & 0 & 1\n\end{pmatrix}\n=\n\begin{pmatrix}\n1 & 0 & \epsilon_i a \\
0 & 1 & \epsilon_i b \\
0 & 0 & 1\n\end{pmatrix}\n\begin{pmatrix}\n\lambda_2(g) & 0 & \delta_{13}(g) \\
0 & \lambda_1(g) & \delta_{23}(g) \\
0 & 0 & \lambda(g)\n\end{pmatrix}.
$$

By equating the $(1, 3)$ and $(2, 3)$ entries in both sides, we get :

$$
\begin{cases}\n\lambda_2(g)\mu_i a = \epsilon_i a \lambda(g) \\
\lambda_1(g)\mu_i b = \epsilon_i b \lambda(g)\n\end{cases}
$$

Since $a \neq 0 \neq b$ we get in particular that $\lambda_1(g) = \lambda_2(g)$ for each $g \in G$. Recall that $G \subset SL(V)$, hence $\lambda_1(g) \cdot \lambda_2(g) \cdot \lambda(g) = 1$. Let $m = \langle x, aw_1-bw_2, v \rangle$ be the unique graded maximal ideal of $S(V)^{T(G)}$ $F[x, aw_1 - bw_2, v]$. Then,

$$
g|m/m^{2} = \begin{pmatrix} \lambda_2(g)^{p^n} & 0 & 0 \\ 0 & \lambda_1(g) & a\delta_{23}(g) - b\delta_{13}(g) \\ 0 & 0 & \lambda(g) \end{pmatrix}.
$$

So by [Theorem [1.2\]](#page-7-1) $S(V)^G$ is Gorenstein iff $\lambda_2(g)^{p^n} \cdot \lambda_1(g) \cdot \lambda(g) = 1$. Since $\lambda_2(g) = \lambda_1(g)$ and $\lambda_1(g) \cdot \lambda_2(g) \cdot \lambda(g) = 1$, then $S(V)^G$ is Gorenstein if and only if $\lambda_1(g)^{p^n-1} = 1$. Since $\lambda_1(g) =$ $\lambda_2(g), |T(G)| = p^n$, then $|gT(G)| = |g|m/m^2| = |\lambda_1(g)|$, for $g \in G$. So $S(V)^G$ is Gorenstein if and only if every $gT(G) \in G/T(G)$ has order dividing $|T(G)| - 1 = pⁿ - 1$. \Box

Remark 3.4. The difference between the previous result and the results of [\[3\]](#page-47-5), is that in [\[3\]](#page-47-5), $G/T(G)$ was a cyclic group and here it is not necessarily so.

Case $2:dim(span_F X) = 2$

We also assume that $T(G) = \langle \sigma_1, ..., \sigma_{n_1}, \tau_1, ..., \tau_{n_2} \rangle$, where

$$
\sigma_i = \left(\begin{array}{ccc} 1 & 0 & a\mu_i \\ 0 & 1 & b\mu_i \\ 0 & 0 & 1 \end{array}\right)
$$

and

$$
\tau_i = \left(\begin{array}{ccc} 1 & 0 & c\alpha_i \\ 0 & 1 & d\alpha_i \\ 0 & 0 & 1 \end{array} \right),
$$

where $\binom{a}{b}, \binom{c}{d} \in \binom{F}{F}$ are F-linearly independent. Also $\{\mu_1, ..., \mu_{n_1}\} \subset F$ are \mathbb{F}_p -linearly independent and $\{\alpha_1,\ldots,\alpha_{n_2}\}\subset F$ are \mathbb{F}_p -linearly independent. Let $T_1(G) = \langle \sigma_1,\ldots,\sigma_{n_1} \rangle$ and $T_2(G) = \langle$ $\tau_1, ..., \tau_{n_2}$ > .

We shall prove the following result.

Theorem 3.5. Let F be a field with char $F = p > 0$. Let $G \subset SL(3, F)$ be a finite subgroup and $V = F³$, the natural $SL(3, F)$ – module with the above notation, assumption and $\lambda_1(g) = \lambda_2(g) \,\forall g \in G$. Then $S(V)^G$ is Gorenstein in the following cases:

- (*i*) $p = 3$.
- (*ii*) $|G|$ *is prime to* 3.
- (iii) $F = \mathbb{F}_q$, where $q = p^s$ and $g.c.d(3, s) = 1$.

Lemma 3.6. $S(V)^{T(G)}$ is a polynomial ring.

Proof. $\sigma_i(w_1) = w_1 + b\mu_i v$ and $\sigma_i(w_2) = w_2 + a\mu_i v$, implying that : $\sigma_i(aw_1 - bw_2) = aw_1 - bw_2$, for $i = 1, ..., n_1$. Similarly, $\tau_i(cw_1 - dw_2) = cw_1 - dw_2$, for $i = 1, ..., n_2$. Since $\{aw_1 - bw_2, cw_1 - dw_2\}$ are linearly independent (otherwise $\binom{a}{b} = \lambda \binom{c}{d}$, in contradiction to their F-linear independence), $\{aw_1 - bw_2, cw_1 - dw_2, v\}$ is a basis of V and they are F-algebraically independent.Now $\tau_i(aw_1 - bw_2, w_2, w_1)$ $bw_2) = a\tau_i(w_1) - b\tau_i(w_2) = a(w_1 + d\alpha_i v) - b(w_2 + c\alpha_i v) = (aw_1 - bw_2) + (ad - bc)\alpha_i v$. Hence by [\[3,](#page-47-5) Lemma 3.13] $S(F(aw_1 - bw_2) + Fv)^{T(G)} = F[x, v]$ is a polynomial ring, where x is homogenous polynomial (in $(aw_1 - bw_2)$, v) of degree p^{n_2} , having $(aw_1 - bw_2)^{p^{n_2}}$ as one of its monomials. Similarly $\sigma_i(cw_1 - dw_2) = (cw_1 - dw_2) + (cb - da)\mu_i v$, and again $S(F(cw_1 - dw_2) + Fv)^{T(G)} = F[y, v]$, is a polynomial ring, where y is homogenous polynomial (in $(cw_1 - dw_2), v$) of degree p^{n_1} , having $(cw_1 - dw_2)^{p^{n_1}}$ as one of its monomials. Since $\{aw_1 - bw_2, cw_1 - dw_2, v\}$ are algebraic independent, then $\{x, y, v\}$ are algebraically independent. Since $(degx) \cdot (degy) \cdot (degy) = p^{n_2} \cdot p^{n_1} \cdot 1 = |T(G)|$, we have by [\[4,](#page-47-4) Theorem 3.9.4] that $S(V)^{T(G)} = F[x, y, v]$ is a polynomial ring.

In the next three results we examine the normality of $T_1(G), T_2(G)$. This is not needed for **Propo**sition [3.10](#page-20-0) ,Proposition [3.13](#page-23-0), and Corollary [3.14](#page-25-0).

Lemma 3.7. Let $G \subset GL(V)$ be a finite subgroup, $T(G) = T_1(G) \times T_2(G)$ and suppose that $T_k(G) \triangleleft G$ for $k = 1, 2$. Then one of the following holds:

- (a) $\lambda_2(g) = \lambda_1(g) \ \forall g \in G$.
- (b) $\lambda_2(g) = \delta(g)\lambda_1(g)$, where $\delta(g) \in \mathbb{F}_p$ $\forall g \in G$.

Proof. Since g normalizes $T_1(G)$ $\forall g \in G$, we have for each $i = 1, ..., n_1$:

$$
\gamma_i = g \sigma_i g^{-1} = \left(\begin{array}{ccc} 1 & 0 & \sum_{j=1}^{n_1} a \mu_j r_{ji} \\ 0 & 1 & \sum_{j=1}^{n_1} b \mu_j r_{ji} \\ 0 & 0 & 1 \end{array} \right),
$$

where $r_{ji} \in \mathbb{F}_p$. Since $g\sigma_i = \gamma_i g$ we get by equating the $(1, 3)$ and $(2, 3)$ entries in both sides:

(1)
\n
$$
\begin{cases}\n\lambda_2(g)a\mu_i = \lambda(g)(\sum_{j=1}^{n_1} \mu_j r_{ji})a \\
\lambda_1(g)b\mu_i = \lambda(g)(\sum_{j=1}^{n_1} \mu_j r_{ji})b\n\end{cases}
$$

Similarly, since g normalizes $T_2(G)$ $\forall g \in G$, we have for each $i = 1, ..., n_2$:

(2)

$$
\begin{cases}\n\lambda_2(g)c\alpha_i = \lambda(g)(\sum_{j=1}^{n_2} \alpha_j r_{ji})c \\
\lambda_1(g)d\alpha_i = \lambda(g)(\sum_{j=1}^{n_2} \alpha_j r_{ji})d\n\end{cases}
$$

Recall that $\binom{a}{b}, \binom{c}{d} \in \binom{F}{F}$ are F-linearly independent. If $a \neq 0 \neq b$ (or $c \neq 0 \neq d$) then we get by (1)(or (2)) that : $\lambda_1(g) = \lambda_2(g) \,\forall g \in G$. On the other hand, if $\begin{pmatrix} a & c \\ b & d \end{pmatrix} = \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$, then from (1) and (2) we get that $\delta(g) = \frac{\lambda_2(g)}{\lambda_1(g)} = \frac{\alpha_i \sum_{j=1}^{n_1} \mu_j r_{ji}}{\mu_i \sum_{i=1}^{n_2} \alpha_j r_{ji}}$ $\frac{\alpha_i \sum_{j=1}^i \mu_j r_{ji}}{\mu_i \sum_{j=1}^{n_2} \alpha_j r_{ji}}$, and if $\left(\begin{smallmatrix} a & c \\ b & d \end{smallmatrix}\right) = \left(\begin{smallmatrix} 0 & c \\ b & 0 \end{smallmatrix}\right)$, then from (1) and (2) we get that $\delta(g) = \frac{\lambda_2(g)}{\lambda_1(g)} = \frac{\mu_i \sum_{j=1}^{n_2} \alpha_j r_{ji}}{\alpha_i \sum_{i=1}^{n_1} \mu_i r_{ji}}$ $\frac{\mu_i}{\alpha_i \sum_{j=1}^{n_1} \mu_j r_{ji}}$. Since $\mu_i, \alpha_i, r_{ji} \in \mathbb{F}_p$ then $\delta(g) \in \mathbb{F}_p$. \Box

Lemma 3.8. $T_k(G) \triangleleft G$ for $k = 1, 2$ in the two following cases:

- (a) $\lambda_2(q) = \lambda_1(q) \; \forall q \in G$.
- (b) $\begin{pmatrix} a & c \\ b & d \end{pmatrix} \in \{ \begin{pmatrix} 0 & c \\ b & 0 \end{pmatrix}, \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \}.$

Proof. Suppose to the contrary that $T_1(G) \nless G$. Then $gT(G)g^{-1} = T(G)$ implies that $g\sigma_i g^{-1} = T(G)$ $\sigma_1^{e_1} \cdot \ldots \cdot \sigma_{n_1}^{e_{n_1}} \cdot \tau_1^{f_1} \cdot \ldots \cdot \tau_{n_2}^{f_{n_2}}$, where not all of the $f_i's$ are 0. By a simple calculation we have:

$$
g\sigma_i g^{-1} = \begin{pmatrix} 1 & 0 & \lambda(g)^{-1} \mu_i \lambda_2(g) a \\ 0 & 1 & \lambda(g)^{-1} \mu_i \lambda_1(g) b \\ 0 & 0 & 1 \end{pmatrix}.
$$

On the other hand we have :

$$
\sigma_1^{e_1} \cdot \ldots \cdot \sigma_{n_1}^{e_{n_1}} = \left(\begin{array}{ccc} 1 & 0 & \sum_{i=1}^{n_1} a \mu_i e_i \\ 0 & 1 & \sum_{i=1}^{n_1} b \mu_i e_i \\ 0 & 0 & 1 \end{array} \right).
$$

as well as

$$
\tau_1^{f_1} \cdot \ldots \cdot \tau_{n_2}^{f_{n_2}} = \left(\begin{array}{ccc} 1 & 0 & \sum_{i=1}^{n_2} c \alpha_i f_i \\ 0 & 1 & \sum_{i=1}^{n_2} d \alpha_i f_i \\ 0 & 0 & 1 \end{array} \right).
$$

Hence by equating the $(1, 3)$ and $(2, 3)$ entries in both sides :

$$
\begin{cases}\n\lambda(g)^{-1}\mu_i\lambda_2(g)a = \sum_{j=1}^{n_1} a\mu_j e_j + \sum_{j=1}^{n_2} c\alpha_j f_j \\
\lambda(g)^{-1}\mu_i\lambda_1(g)b = \sum_{j=1}^{n_1} b\mu_j e_j + \sum_{j=1}^{n_2} d\alpha_j f_j\n\end{cases}
$$

We firstly assume that $\lambda_2(g) = \lambda_1(g)$. Then $\lambda_2(g)\lambda(g)^{-1}\mu_i\left(\frac{a}{b}\right) = \left(\sum_{j=1}^{n_1} \mu_j e_j\right)\left(\frac{a}{b}\right) + \left(\sum_{j=1}^{n_2} \alpha_j f_j\right)\left(\frac{c}{d}\right)$. So since $\binom{a}{b}$, $\binom{c}{d}$ are *F*-linearly independent we get that $\sum_{j=1}^{n_2} \alpha_j f_j = 0$. Since $\{\alpha_1, ..., \alpha_{n_2}\}$ are \mathbb{F}_p linearly independent , $f_j = 0$ for $j = 1, ..., n_2$. A contradiction, hence $T_1(G) \triangleleft G$.

Assume now that item (b) holds.Say $\begin{pmatrix} a & c \\ b & d \end{pmatrix} = \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$, hence $\begin{pmatrix} \lambda(g)^{-1} \mu_i \lambda_2(g)a \\ 0 \end{pmatrix} = \left(\sum_{j=1}^{n_1} \mu_j e_j \right) \begin{pmatrix} a \\ 0 \end{pmatrix}$ + $(\sum_{j=1}^{n_2} \alpha_j f_j)$ $(\begin{matrix} 0 \\ d \end{matrix})$, so $(\sum_{j=1}^{n_2} \alpha_j f_j) = 0$. Since $\{\alpha_1, ..., \alpha_{n_2}\}\$ are \mathbb{F}_p -linearly independent, $f_j = 0$ for $j =$ 1, ..., n_2 , a contradiction. If $\begin{pmatrix} a & c \\ b & d \end{pmatrix} = \begin{pmatrix} 0 & c \\ b & 0 \end{pmatrix}$, then by the same steps above we get that $\left(\sum_{j=1}^{n_2} \alpha_j f_j\right) = 0$, again $f_j = 0$ for $j = 1, ..., n_2$, again a contradiction. By a similar discussion we get that $T_2(G) \triangleleft G$. \Box

The combination of Lemma [3.7](#page-18-0) and Lemma [3.8](#page-18-1) yields.

Proposition 3.9. The following conditions are equivalent.

- (a) $T_k(G) \triangleleft G$, for $k = 1, 2$.
- (b) one of the following holds:
- 1. $\begin{pmatrix} a & c \\ b & d \end{pmatrix} \in \{ \begin{pmatrix} 0 & c \\ b & 0 \end{pmatrix}, \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \}$
- 2. $\lambda_2(g) = \lambda_1(g)$, for all $g \in G$.

Proof. That (b) implies (a) is a consequence of Lemma [3.8](#page-18-1). That (a) implies (b) follows from the proof of Lemma [3.7](#page-18-0).

We can now give a necessary and sufficient condition for a subgroup G such that $S(V)^G$ is Gorenstein.

Proposition 3.10. Suppose $n_1 < n_2$. Then $S(V)^G$ is Gorenstein if and only if :

- 1. $\lambda_1(g)^{p^{n_1}+p^{n_2}}\lambda(g) = 1$, in case $c \neq 0 \neq d$.
- 2. $\lambda_1(g)^{p^{n_2}} \lambda_2(g)^{p^{n_1}} \lambda(g) = 1$, in case $c = 0 \neq d$.
- 3. $\lambda_2(g)^{p^{n_2}} \lambda_1(g)^{p^{n_1}} \lambda(g) = 1$, in case $c \neq 0 = d$.

The case $n_2 < n_1$ is similar. Note that we do not assume any normality assumption on $T_k(G)$.

Proof. By Lemma [3.6](#page-17-1), we have that $S(V)^{T(G)} = F[x, y, v], deg(y) = p^{n_1}, deg(x) = p^{n_2}$. Since $g^{-1}T(G)g = T(G)$ we have $g(y) \in S(V)^{T(G)}$. Therefore, since g preserve degrees, $g(y) = \alpha y + \beta v^{p^{n_1}}$. Recall that $(cw_1 - dw_2)^{p^{n_1}}$ appears as a monomial in y and all other monomials involve v in addition.

We have :

 $g(cw_1 - dw_2) = cg(w_1) - dg(w_2) = c(\lambda_1(g)w_1 + \delta_{23}(g)v) - d(\lambda_2(g)w_2 + \delta_{13}(g)v) = (c\lambda_1(g)w_1$ $d\lambda_2(g)w_2) + \chi v$, where $\chi = c\delta_{23}(g) - d\delta_{13}(g)$. Hence $g((cw_1 - dw_2)^{p^{n_1}}) = c^{p^{n_1}}\lambda_1(g)^{p^{n_1}}w_1^{p^{n_1}}$ $d^{p^{n_1}}\lambda_2(g)^{p^{n_1}}w_2^{p^{n_1}}+\chi^{p^{n_1}}v^{p^{n_1}}$. So by comparing coefficients of y in $g(y)$ we get:

$$
c^{p^{n_1}} \lambda_1(g)^{p^{n_1}} w_1^{p^{n_1}} - d^{p^{n_1}} \lambda_2(g)^{p^{n_1}} w_2^{p^{n_1}} = \alpha (cw_1 - dw_2)^{p^{n_1}}.
$$
 Consequently
\n(1)
$$
c^{p^{n_1}} \lambda_1(g)^{p^{n_1}} = \alpha c^{p^{n_1}}, d^{p^{n_1}} \lambda_2(g)^{p^{n_1}} = \alpha d^{p^{n_1}}.
$$

Similarly $g(x) = \delta x + \varepsilon y^{p^{n_2 - n_1}} +$ mixed terms (in y, v) + $\xi v^{p^{n_2}}$. Recall that $(aw_1 - bw_2)^{p^{n_2}}$ appears as a monomial in x and all other monomials involve v in addition. So the contribution to the terms with $w_1^{p^{n_2}}$ p^{n_2} in the right hand side of the last equation are $\delta(aw_1 - bw_2)^{p^{n_2}} + \varepsilon((cw_1 - dw_2)^{p^{n_1}})^{p^{n_2-n_1}} =$ $\delta(aw_1 - bw_2)^{p^{n_2}} + \varepsilon (cw_1 - dw_2)^{p^{n_2}}.$

The contribution of the left hand side are from $g((aw_1 - bw_2)^{p^{n_2}}) = a^{p^{n_2}}(\lambda_1(g)w_1 + \delta_{23}(g)v)^{p^{n_2}}$ $b^{p^{n_2}}(\lambda_2(g)w_2+\delta_{13}(g)v)^{p^{n_2}}$. Consequently we get :

$$
(2) \ \ a^{p^{n_2}} \lambda_1(g)^{p^{n_2}} = \delta a^{p^{n_2}} + \varepsilon c^{p^{n_2}}, b^{p^{n_2}} \lambda_2(g)^{p^{n_2}} = \delta b^{p^{n_2}} + \varepsilon d^{p^{n_2}}.
$$

If $c \neq 0 \neq d$, then (1) shows that $\lambda_1(g)^{p^{n_1}} = \alpha = \lambda_2(g)^{p^{n_1}}$, so $\lambda_1(g) = \lambda_2(g)$. Now (2) shows that $a^{p^{n_2}}(\lambda_1(g)^{p^{n_2}} - \delta) = \varepsilon c^{p^{n_2}}, b^{p^{n_2}}(\lambda_2(g)^{p^{n_2}} - \delta) = \varepsilon d^{p^{n_2}}.$

So if $\varepsilon \neq 0$ then $\frac{a^{p^{n_2}}}{\varepsilon c^{p^{n_2}}} = \frac{b^{p^{n_2}}}{\varepsilon d^{p^{n_2}}}$. Hence $\frac{a}{c} = \frac{b}{d}$ or $\frac{a}{b} = \frac{c}{d}$, which is a contradiction,so $\varepsilon = 0$ in case $c \neq 0 \neq d$. Therefore $\lambda_1(g)^{p^{n_2}} = \delta = \lambda_2(g)^{p^{n_2}}$. Let $m = \langle x, y, v \rangle$ be the unique homogenous maximal ideal of $S(V)^{T(G)} = F[x, y, v]$. The matrix representing g on m/m^2 whenever $c \neq 0 \neq d$ is

$$
\left(\begin{array}{ccc} \lambda_1(g)^{p^{n_2}} & 0 & 0 \\ 0 & \lambda_1(g)^{p^{n_1}} & 0 \\ 0 & 0 & \lambda(g) \end{array}\right),
$$

so the condition for $S(V)^G$ to be Gorenstein is $\lambda_1(g)^{p^{n_2}+p^{n_1}}\lambda(g)=1$. Since $G \subset SL(V)$, $\lambda_1(g)=$ $\lambda(g)^{-2}$. Hence $S(V)^G$ is Gorenstein if and only if $\lambda_1(g)^{p^{n_2}+p^{n_1-2}}=1$. Moreover, by **Lemma [3.8](#page-18-1)**, $T_k(G) \lhd G$ for $k = 1, 2$.

Assume now that $c = 0 \neq d$, hence from (1) we get $\lambda_2(g)^{p^{n_1}} = \alpha$ and from (2), since $a \neq 0$ in this case, $\lambda_1(g)^{p^{n_2}} = \delta$. So the matrix representing g on m/m^2 is

$$
\left(\begin{array}{ccc} \lambda_1(g)^{p^{n_2}} & 0 & 0 \\ 0 & \lambda_2(g)^{p^{n_1}} & 0 \\ 0 & 0 & \lambda(g) \end{array}\right),
$$

and $S(V)^G$ is Gorenstein if and only if $\lambda_1(g)^{p^{n_2}}\lambda_2(g)^{p^{n_1}}\lambda(g) = 1$.

Assume now that $c \neq 0 = d$. Then from (1) we get $\lambda_1(g)^{p^{n_1}} = \alpha$ and from (2), since $b \neq 0$ in this case, $\lambda_2(g)^{p^{n_2}} = \delta$. So the matrix representing g on m/m^2 is

$$
\left(\begin{array}{ccc} \lambda_2(g)^{p^{n_2}} & 0 & 0 \\ 0 & \lambda_1(g)^{p^{n_1}} & 0 \\ 0 & 0 & \lambda(g) \end{array}\right),
$$

and $S(V)^G$ is Gorenstein if and only if $\lambda_2(g)^{p^{n_2}}\lambda_1(g)^{p^{n_1}}\lambda(g) = 1$. This settles the case $n_1 < n_2$. If $n_1 > n_2$ then we do the same analysis but with $\{a, b\}$ instead of $\{c, d\}$.

Our next result will be used in the proof of Theorem [3.5](#page-17-0).

Lemma 3.11. Suppose $T_k(G) \triangleleft G$. Then $\left(\frac{\lambda_2(g)}{\lambda(g)}\right)^{p^{n_k}-1} = \left(\frac{\lambda_1(g)}{\lambda(g)}\right)^{p^{n_k}-1} = 1$, for $k = 1, 2$.

Proof. We have by Lemma [3.6](#page-17-1) that $S(F(aw_1 - bw_2) + Fv)^{T_2(G)} = F[x, v]$, where x is homogenous polynomial (in $(aw_1 - bw_2)$, v) of degree p^{n_2} , having $(aw_1 - bw_2)^{p^{n_2}}$ as one of its monomials and all other monomials involve v. In other words $x = (aw_1 - bw_2)^{p^{n_2}} + v^{p^{n_2}-1}(aw_1 - bw_2) +$ monomials involve v . By a simple calculating we get that :

$$
\begin{cases} g(aw_1 - bw_2)^{p^{n_2}} = a^{p^{n_2}} \lambda_1(g)^{p^{n_2}} w_1^{p^{n_2}} - b^{p^{n_2}} \lambda_2(g)^{p^{n_2}} w_2^{p^{n_2}} + (a\delta_{23}(g) - b\delta_{13}(g))^{p^{n_2}} v^{p^{n_2}} \\ g(v^{p^{n_2}-1}(aw_1 - bw_2)) = a\lambda(g)^{p^{n_2}-1} \lambda_1(g) w_1 v^{p^{n_2}-1} - b\lambda(g)^{p^{n_2}-1} \lambda_2(g) w_2 v^{p^{n_2}-1} + \lambda(g)^{p^{n_2}-1} v^{p^{n_2}} (a\delta_{23}(g) - b\delta_{13}(g)). \end{cases}
$$

 \Box

Now $T_2(G) \triangleleft G$ imply that $F[x, v]$ is g-stable. Therefore, since g preserve degrees, we get $g(x) = \rho x + \kappa v^{p^{n_2}}$, $\rho, \kappa \in F$. Assume firstly, using **Lemma [3.7](#page-18-0)**, that $\lambda_1(g) = \lambda_2(g)$. Hence, if $a \neq 0$ then by comparing coefficients of $w_1^{p^{n_2}}$ p^{n_2} and $v^{p^{n_2}-1}w_1$ in $g(x)$ we get $\rho = \lambda_1(g)^{p^{n_2}} = \lambda(g)^{p^{n_2}-1}\lambda_1(g)$. So $\left(\frac{\lambda_1(g)}{\lambda(g)}\right)^{p^{n_2}-1} = 1$, and if $b \neq 0$ then by comparing coefficients of $w_2^{p^{n_2}}$ p^{n_2} and $v^{p^{n_2}-1}w_2$ in $g(x)$ we get $\rho = \lambda_2(g)^{p^{n_2}} = \lambda(g)^{p^{n_2}-1}\lambda_2(g)$. So $\left(\frac{\lambda_2(g)}{\lambda(g)}\right)^{p^{n_2}-1} = 1$.

Assume now, using Lemma [3.7](#page-18-0), that $\lambda_2(g) = \delta(g)\lambda_1(g)$, where $\delta(g) \in \mathbb{F}_p$. Consequently, by **Proposition [3.9](#page-19-0)**, $\begin{pmatrix} a & c \\ b & d \end{pmatrix} \in \{ \begin{pmatrix} 0 & c \\ b & 0 \end{pmatrix}, \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \}.$ Hence if $a \neq 0 = b$, then by comparing coefficient of $w_1^{p^{n_2}}$ p^{n_2} and $v^{p^{n_2}-1}w_1$ in $g(x)$ we get $\rho = \lambda_1(g)^{p^{n_2}} = \lambda(g)^{p^{n_2}-1}\lambda_1(g)$. So $(\frac{\lambda_1(g)}{\lambda(g)})^{p^{n_2}-1} = 1$, hence $\left(\frac{\delta(g)^{-1}\lambda_2(g)}{\lambda(g)}\right)^{p^{n_2}-1}=1.$ By little Fermat's theorem we get that $\delta(g)^{p^{n_2}-1}=1$, then $\left(\frac{\lambda_2(g)}{\lambda(g)}\right)^{p^{n_2}-1}=1.$ If $b \neq 0 = a$, then by comparing coefficient of $w_2^{p^{n_2}}$ p^{n_2} and $v^{p^{n_2}-1}w_2$ in $g(x)$ we get $\rho = \lambda_2(g)^{p^{n_2}} =$ $\lambda(g)^{p^{n_2}-1}\lambda_2(g)$. So $\left(\frac{\lambda_2(g)}{\lambda(g)}\right)^{p^{n_2}-1}=1$, hence $\left(\frac{\delta(g)\lambda_1(g)}{\lambda(g)}\right)^{p^{n_2}-1}=1$. By little Fermat's theorem we get that $\delta(g)^{p^{n_2}-1} = 1$, then $\left(\frac{\lambda_2(g)}{\lambda(g)}\right)^{p^{n_2}-1} = 1$.

Similarly, if we look at $S(F(cw_1 - dw_2) + Fv)^{T_1(G)} = F[y, v]$ from **Lemma [3.6](#page-17-1)**, then the above reasoning shows again that $\left(\frac{\lambda_k(g)}{\lambda(g)}\right)^{p^{n_1}-1} = 1$, for $k = 1, 2$. \Box

Lemma 3.12. $(|G|/|T(G)|, p) = 1$, and if $\lambda_1(g) = \lambda_2(g) \,\forall g \in G$ then $G/T(G)$ is a cyclic group. *Proof.* Since $G \subset SL(V)$, $\lambda(g) = \lambda_1(g)^{-1} \lambda_2(g)^{-1}$. We have:

$$
g^{p} = \begin{pmatrix} \lambda_{2}(g)^{p} & 0 & * \\ 0 & \lambda_{1}(g)^{p} & * \\ 0 & 0 & \lambda_{1}(g)^{-p}\lambda_{2}(g)^{-p} \end{pmatrix}
$$

If $g^p \in T(G)$ then $\lambda_1(g)^p = \lambda_2(g)^p = 1$, and then $\lambda_1(g) = \lambda_2(g) = 1$, so :

$$
g = \left(\begin{array}{ccc} 1 & 0 & * \\ 0 & 1 & * \\ 0 & 0 & 1 \end{array}\right) \in T(G).
$$

This shows that $(|G|/|T(G)|, p) = 1$. Assume now that $\lambda_1(g) = \lambda_2(g) \,\forall g \in G$. Set $U = \mathbb{F}_p \mu_1 + \cdots + \mathbb{F}_p \mu_{n_1}$. By Lemma [3.8](#page-18-1), $T_1(G) \lhd G$ then $g \sigma_i g^{-1} \in T_1(G)$. Now since

$$
g\sigma_i g^{-1} = \begin{pmatrix} 1 & 0 & \lambda(g)^{-1} \mu_i \lambda_2(g) a \\ 0 & 1 & \lambda(g)^{-1} \mu_i \lambda_1(g) b \\ 0 & 0 & 1 \end{pmatrix}
$$

for $i = 1, ..., n_1$, we get that $\lambda(g)^{-1}\lambda_1(g)\mu_i \in U$, hence $\lambda(g)^{-1}\lambda_1(g)U \subseteq U$ Since $G \subset SL(V)$ and $\lambda_1(g) = \lambda_2(g)$ then $\lambda(g)^{-1}\lambda_1(g) = \lambda_1(g)^3$. Consequently by [\[3,](#page-47-5) Lemma 3.12.] $|\lambda_1(g)^3|$ divides $(p^{n_1}-1)$. By a similar discussion on $T_2(G)$ we get that $|\lambda_1(g)^3|$ divides $(p^{n_2}-1)$. Let $H = \{ \epsilon \in F | \epsilon^{3(p^{n_1}-1)} = 1 \}$. Define $\Phi: G/T(G) \longrightarrow H$, by $\Phi(gT(G)) = \lambda_1(g)$, for each $g \in G$. Since

$$
gT(G) = g|m/m^{2} = \begin{pmatrix} \lambda_{1}(g)^{p^{n_{2}}} & 0 & 0 \\ 0 & \lambda_{1}(g)^{p^{n_{1}}} & 0 \\ 0 & 0 & \lambda_{1}(g)^{-2} \end{pmatrix},
$$

it follows that Φ is an injective homomorphism with its image in H since $|g(T(G)| = |\lambda_1(g)|$ divides $3(p^{n_1}-1)$. Since H is cyclic group it follows that $G/T(G)$ is cyclic.

Recall that n_1 and n_2 are the numbers of the generators of $T_1(G)$ and $T_2(G)$ respectively. The next result considers the case $n_1 = n_2$ which was omitted from **Proposition [3.10](#page-20-0).**

Proposition 3.13. Suppose that $n_1 = n_2$. Then $S(V)^G$ is Gorenstein iff $\lambda_2(g)^{p^{n_1}}\lambda_1(g)^{p^{n_1}}\lambda(g) = 1$.

Proof. We have by Lemma [3.6](#page-17-1) that $S(V)^{T(G)} = F[x, y, v], deg(y) = p^{n_1}, deg(x) = p^{n_2}$. Since $g^{-1}T(G)g = T(G)$ we have $g(y), g(x) \in S(V)^{T(G)}$. Hence, since g preserve degrees, we get :

(1) $g(y) = \alpha y + \gamma x + \beta v^{p^{n_1}}, \alpha, \gamma, \beta \in F$.

(2)
$$
g(x) = \delta x + \varepsilon y + \chi v^{p^{n_1}}, \delta, \varepsilon, \chi \in F.
$$

Therefore, $S(V)^G = [S(V)^{T(G)}]^{G/T(G)}$. Let $m = (x, y, v)$ be the unique graded maximal ideal of $S(V)^{T(G)} = F[y, x, v]$, then the matrix of $\bar{g} \in G/T(G)$ on $m/m^2 = F\bar{y} + F\bar{x} + F\bar{v}$ is :

$$
\left(\begin{array}{ccc} \alpha & \gamma & 0 \\ \varepsilon & \delta & 0 \\ 0 & 0 & \lambda(g) \end{array}\right).
$$

So by [Theorem [1.2\]](#page-7-1) $S(V)^G$ is Gorenstein if and only if $(\alpha \delta - \varepsilon \gamma) \lambda(g) = 1 \ \forall g \in G$.

The computation of $\alpha\delta - \varepsilon\gamma$:

We shall assume that $c \neq 0 \neq d$ and $a \neq 0 \neq b$. We know that $(cw_1 - dw_2)^{p^{n_1}}$ appears as a monomial in y, $(aw_1 - bw_2)^{p^{n_1}}$ appears as a monomial in x, and all other monomials involve also v. Therefore the coefficients of $w_1^{p^{n_1}}$ \int_1^{p} in (1) are :

 $c^{p^{n_1}}\lambda_1(g)^{p^{n_1}}$ in the left hand side and $\alpha c^{p^{n_1}} + \gamma a^{p^{n_1}}$ in the right hand side.Hence $c^{p^{n_1}}\lambda_1(g)^{p^{n_1}} =$ $\alpha c^{p^{n_1}} + \gamma a^{p^{n_1}}$. Similarly, for the coefficients of $w_2^{p^{n_1}}$ $q^{p^{n_1}}$, we have $d^{p^{n_1}} \lambda_2(g)^{p^{n_1}} = \alpha d^{p^{n_1}} + \gamma b^{p^{n_1}}$. From (2) we similarly have:

$$
\begin{cases}\na^{p^{n_1}}\lambda_1(g)^{p^{n_1}} = \delta a^{p^{n_1}} + \varepsilon c^{p^{n_1}} \\
b^{p^{n_1}}\lambda_2(g)^{p^{n_1}} = \delta b^{p^{n_1}} + \varepsilon d^{p^{n_1}}\n\end{cases}
$$

Hence from (1) and (2) we have :

$$
\left\{\begin{array}{ll} \lambda_1(g)^{p^{n_1}}=\alpha+\gamma(\frac{a}{c})^{p^{n_1}} \\ \lambda_2(g)^{p^{n_1}}=\alpha+\gamma(\frac{b}{d})^{p^{n_1}} \\ \lambda_1(g)^{p^{n_1}}=\delta+\varepsilon(\frac{c}{d})^{p^{n_1}} \\ \lambda_2(g)^{p^{n_1}}=\delta+\varepsilon(\frac{d}{b})^{p^{n_1}} \end{array}\right.
$$

Hence,

$$
\begin{cases} (\lambda_1(g)-\lambda_2(g))^{p^{n_1}}=\gamma(\frac{a}{c}-\frac{b}{d})^{p^{n_1}}\\ (\lambda_1(g)-\lambda_2(g))^{p^{n_1}}=\varepsilon(\frac{c}{a}-\frac{b}{b})^{p^{n_1}} \end{cases}
$$

It follows that:

(A) :=
$$
(\lambda_1(g) - \lambda_2(g))^{2p^{n_1}} = \gamma \varepsilon ((\frac{a}{c} - \frac{b}{d})(\frac{c}{a} - \frac{d}{b}))^{p^{n_1}}.
$$

Also we have :

$$
\left\{\begin{array}{l}(\frac{c}{a}\lambda_1(g)-\frac{d}{b}\lambda_2(g))^{p^{n_1}}=\alpha(\frac{c}{a}-\frac{d}{b})^{p^{n_1}}\\(\frac{a}{c}\lambda_1(g)-\frac{b}{d}\lambda_2(g))^{p^{n_1}}=\delta(\frac{a}{c}-\frac{b}{d})^{p^{n_1}}\end{array}\right.
$$

And it follows that :

(B) :=
$$
\left(\frac{c}{a}\lambda_1(g) - \frac{d}{b}\lambda_2(g)\right)^{p^{n_1}} \left(\frac{a}{c}\lambda_1(g) - \frac{b}{d}\lambda_2(g)\right)^{p^{n_1}} = \alpha \delta \left[\left(\frac{c}{a} - \frac{d}{b}\right)\left(\frac{a}{c} - \frac{b}{d}\right)\right]^{p^{n_1}}.
$$

So (B)-(A) = $(\alpha \delta - \epsilon \gamma)[(\frac{c}{a} - \frac{d}{b})(\frac{a}{c} - \frac{b}{d})]^{p^{n_1}}$. Observe that $(\frac{c}{a} - \frac{d}{b})(\frac{a}{c} - \frac{b}{d}) = 2 - (\frac{cb}{ad} + \frac{ad}{bc})$. Also $(B)-(A) = (\lambda_1(g)^2 + \lambda_2(g)^2 - (\frac{c\delta}{ad} + \frac{ad}{bc})\lambda_1(g)\lambda_2(g))^{p^{n_1}} - (\lambda_1(g)^2 + \lambda_2(g)^2 - 2\lambda_1(g)\lambda_2(g))^{p^{n_1}} =$ $(\lambda_1(g))^{p^{n_1}} (\lambda_2(g))^{p^{n_1}} [2 - (\frac{cb}{ad} + \frac{ad}{bc})]^{p^{n_1}}.$

Hence $\frac{(B)-(A)}{((\frac{c}{a}-\frac{d}{b})(\frac{a}{c}-\frac{b}{a}))^{p^{n_1}}} = (\alpha\delta - \varepsilon\gamma)$. Therefore $\alpha\delta - \varepsilon\gamma = \lambda_1(g)^{p^{n_1}}\lambda_2(g)^{p^{n_1}}$, hence $S(V)^G$ is Gorenstein if and only if $\lambda_2(g)^{p^{n_1}}\lambda_1(g)^{p^{n_1}}\lambda(g) = 1$.

The cases where one of $\{a, b, c, d\}$ is 0 are easier. For example, if $a = 0$, hence since $\binom{a}{b}$, $\binom{c}{d}$ b) $\setminus d$ are F-linearly independent, $c \neq 0$, Hence $\lambda_1(g)^{p^{n_1}} = \alpha$ and $\varepsilon = 0$, therefore $\delta = \lambda_2(g)^{p^{n_1}}$. Hence $\alpha\delta-\varepsilon\gamma=\alpha\delta=\lambda_1(g)^{p^{n_1}}\lambda_2(g)^{p^{n_1}}$. So again $S(V)^G$ Gorenstein if and only if $\lambda_2(g)^{p^{n_1}}\lambda_1(g)^{p^{n_1}}\lambda(g)=1$. Equivalently $\lambda(g)^{p^{n_1}-1}=1$.

 \Box

We finally arrive at: The proof of Theorem [3.5.](#page-17-0)

Proof. We have by Lemma [3.12](#page-22-0) that $(|G|/|T(G)|, p) = 1$ and $G/T(G) \cong \langle \lambda_1(g) \rangle$ is a cyclic group. Let $\frac{|G|}{|T(G)|} = |\lambda_1(g)| \equiv k$, since $G \subset SL(V)$, $\frac{\lambda_1(g)}{\lambda(g)} = \frac{\lambda_1(g)}{\lambda_1(g)^{-1}\lambda_2(g)^{-1}} = \frac{\lambda_1(g)}{\lambda_1(g)^{-1}\lambda_1(g)^{-1}} = \lambda_1(g)^3$. Hence, by **Lemma [3.11](#page-21-0)**, $(\lambda_1(g)^3)^{p^{n_1}-1} = (\lambda_1(g)^3)^{p^{n_2}-1} = 1$. So $(\lambda_1(g)^3)^{p^{n_1}+p^{n_2}-2} = 1$, which means that k divides $[3 \cdot (p^{n_1} + p^{n_2} - 2)].$

If $p = 3$ then since $(3, k) = 1, k$ divides $(p^{n_1} + p^{n_2} - 2)$, consequently by **Proposition [3.10](#page-20-0)** and **Proposition [3.13](#page-23-0)**, $S(V)^G$ is Gorenstein. This settles item (*i*).

Assume now that $3 \nmid |G|$. The action of g^3 on m/m^2 where m is the unique maximal ideal of $S(V)^{T(G)}$ is :

$$
\left(\begin{array}{ccc} \lambda_1(g)^{3\cdot p^{n_2}} & 0 & 0 \\ 0 & \lambda_1(g)^{3\cdot p^{n_1}} & 0 \\ 0 & 0 & \lambda_1(g)^{-6} \end{array}\right).
$$

Since $|\lambda_1(g)^3|$ divides $p^{n_1}+p^{n_2}-2$, the above reasoning shows that $S(V)^{}=S(m/m^2)^{}$ is Gorenstein. Consequently if $3 \nmid |G|$, then $G = \langle g^3, T(G) \rangle$ and $S(V)^G$ is therefore Gorenstein, so item (ii) is verified.

We have by Lemma [3.8](#page-18-1) that $T_k(G) \lhd G$ for $k = 1, 2$, hence by using the proof of Lemma [3.7](#page-18-0) we get :

(1)

$$
\begin{cases}\n\lambda_2(g)a\mu_i = \lambda(g)(\sum_{j=1}^{n_1} \mu_j r_{ji})a \\
\lambda_1(g)b\mu_i = \lambda(g)(\sum_{j=1}^{n_1} \mu_j r_{ji})b\n\end{cases}
$$

(2)

$$
\begin{cases}\n\lambda_2(g)c\alpha_i = \lambda(g)(\sum_{j=1}^{n_2} \alpha_j r_{ji})c \\
\lambda_1(g)d\alpha_i = \lambda(g)(\sum_{j=1}^{n_2} \alpha_j r_{ji})d\n\end{cases}
$$

Given the assumption of item (iii) . Since $\lambda_1(g) = \lambda_2(g)$, it follows that $(a \neq 0 \neq b)$ or $(c \neq 0 \neq d)$. Say $a, c, d \neq 0$, the equations above imply that $\frac{\lambda(g)}{\lambda_1(g)}(\sum_{j=1}^{n_1} \mu_j r_{ji}) = \mu_i$ and $\frac{\lambda(g)}{\lambda_1(g)}(\sum_{j=1}^{n_2} \alpha_j r_{ji}) = \alpha_i$. Set $M_1 = \mathbb{F}_p \mu_1 + ... + \mathbb{F}_p \mu_{n_1}$ and $M_2 = \mathbb{F}_p \alpha_1 + ... + \mathbb{F}_p \alpha_{n_2}$, so $\dim_{\mathbb{F}_p} M_1 = n_1$ and $\dim_{\mathbb{F}_p} M_2 = n_2$. Thus $\frac{\lambda_1(g)}{\lambda(g)}M_1 = \lambda_1(g)^3M_1 \subseteq M_1$ and $\frac{\lambda_1(g)}{\lambda(g)}M_2 = \lambda_1(g)^3M_2 \subseteq M_2$. Suppose to the contrary that $\lambda_1(g) \notin \mathbb{F}_p(\lambda_1(g)^3)$, then $[\mathbb{F}_p(\lambda_1(g)) : \mathbb{F}_p(\lambda_1(g)^3)] = 3$. But $[F : \mathbb{F}_p] = s$ is relatively prime to 3, a contradiction. Thus $\lambda_1(g) \in \mathbb{F}_p(\lambda_1(g)^3)$, hence $\lambda_1(g)M_1 \subseteq M_1$, similarly, we get that $\lambda_2(g)M_2 \subseteq M_2$. Therefore, by [\[3,](#page-47-5) Lemma 3.12.] $k|(p^{n_1}-1)$ and $k|(p^{n_2}-1)$, hence $k|(p^{n_1}+p^{n_2}-2)$. Consequently, by Proposition [3.10](#page-20-0) item (1) , $S(V)^G$ is Gorenstein and item *(iii)* holds.

We also have here the following.

Corollary 3.14. Suppose $G \subset SL(3, \mathbb{F}_p)$ and $V = \mathbb{F}_p^3$ contains two 2-dimensional G-submodules. Then $S(V)^G$ is Gorenstein.

Proof. Since $F = \mathbb{F}_p$, $dim_{F_p}(span X) \in \{1, 2\}$, and either $T(G) = T_1(G) \times T_2(G)$ or $T(G) = T_1(G)$. If $dim_{\mathbb{F}_p}(span X) = 2$, then $n_1 = n_2 = 1$. So by **Proposition [3.13](#page-23-0)** the condition for being Gorenstein is $\lambda_2(g)^p \lambda_1(g)^p \lambda(g) = 1$, for each $g \in G$. Since $\lambda(g) = \lambda_1(g)^{-1} \lambda_2(g)^{-1}$, it follows that it is equivalent to $\lambda_2(g)^{p-1}\lambda_1(g)^{p-1} = 1$. Now since $\lambda_1(g)$, $\lambda_2(g) \in \mathbb{F}_p \setminus \{0\}$ it follows that $\lambda_1(g)^{p-1} = 1 = \lambda_2(g)^{p-1}$, and the condition is fullfilled. If $dim_{\mathbb{F}_p}(span X) = 1$, then $n = 1$, that is $|T(G)| = p$, so since $|gT(G)| = |\lambda_1(g)|$, it follows from $\lambda_1(g) \in \mathbb{F}_p \setminus \{0\}$ that $|gT(G)|$ divides $|T(G)| - 1$, and again the Gorenstein property holds.

 \Box

4 Chapter(C): Preliminaries to Chapters D,E,F and G.

Let $G \subset GL(V)$ be a group and $W \subset V$ a G -submodule with $dim_F W = dim_F V - 1$. With respesct to a fixed basis, each $g \in G$ is represented by the matrix :

$$
g = \begin{pmatrix} g_{11} & g_{12} & \dots & g_{1n} \\ 0 & g_{22} & \dots & g_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & g_{n2} & \dots & g_{nn} \end{pmatrix}
$$

Lemma 4.1. Suppose $g \in G \cap SL(V)$. Then

$$
g^{-1}hg = \begin{pmatrix} 1 & G_{11}(\alpha_1, ..., \alpha_{n-1})g|W & 0 & \dots & \dots & 0 & 0 \\ 0 & 0 & 1 & 0 & \dots & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & & 0 & \vdots & \vdots & \vdots \\ 0 & \dots & \vdots & 0 & \dots & 0 & 1 & 0 \\ 0 & 0 & 0 & \dots & \dots & \dots & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & G_{11}(\alpha_1, ..., \alpha_{n-1})g|W \\ \vdots & & & \vdots & \vdots \\ 0 & & & & \vdots \\ 0 & & & & & \end{pmatrix}, where
$$

$$
g|W = \begin{pmatrix} g_{22} & g_{23} & \dots & g_{2n} \\ g_{32} & \dots & g_{3n} \\ \vdots & \vdots & \ddots & \vdots \\ g_{n2} & \dots & g_{nn} \end{pmatrix}, G_{11} = det(g|W), and
$$

$$
h = \begin{pmatrix} 1 & \alpha_1 & \dots & \dots & \dots & \alpha_{n-1} \\ 0 & 1 & 0 & \dots & \dots & 0 \\ 0 & 0 & 1 & 0 & \dots & \dots & 0 \\ \vdots & \vdots & 0 & \dots & \vdots & \vdots \\ 0 & \dots & \vdots & 0 & \dots & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & \alpha_1 & \dots & \alpha_{n-1} \\ 0 & 1 & \dots & \alpha_{n-1} \\ \vdots & & & & \vdots \\ 0 & \dots & \vdots & 0 & \dots & \dots & 0 & 1 \end{pmatrix}.
$$

Proof.

$$
g^{-1} = \begin{pmatrix} G_{11} & -G_{21} & \dots & (-1)^{n+1} G_{n1} \\ (-1)^{i+j} G_{ji} & & \\ \vdots & & \vdots & \ddots & \\ 0 & & & \\ 0 & & & \end{pmatrix} = \begin{pmatrix} G_{11} & -G_{21} & \dots & (-1)^{n+1} G_{n1} \\ \vdots & & \\ 0 & & \end{pmatrix}.
$$

 He

ence,
\n
$$
g^{-1}\begin{pmatrix}\n1 & \alpha_1 & \cdots & \cdots & \alpha_{n-1} \\
0 & 1 & 0 & \cdots & \cdots & 0 \\
0 & 0 & 1 & 0 & \cdots & 0 & 0 \\
\vdots & \vdots & 0 & \cdots & 0 & 1 & 0 \\
0 & \cdots & \vdots & 0 & \cdots & 0 & 1\n\end{pmatrix} g = g^{-1}hg =
$$
\n
$$
\begin{pmatrix}\nG_{11} & -G_{21} & \cdots & (-1)^{n+1}G_{n1} \\
0 & g^{-1}|W & 0\n\end{pmatrix}\n\begin{pmatrix}\ng_{11} & g_{12} & \cdots & g_{1n} \\
0 & \vdots & \vdots & \vdots \\
0 & g^{-1}|W & 0\n\end{pmatrix} =
$$
\n
$$
\begin{pmatrix}\nG_{11} & -G_{21} & \cdots & (-1)^{n+1}G_{n1} \\
0 & g^{-1}|W & 0\n\end{pmatrix}\n\begin{pmatrix}\ng_{11} & (g_{12}, ..., g_{1n}) + (\alpha_1, ..., \alpha_{n-1})g|W \\
\vdots & \vdots & \vdots \\
0 & g^{-1}|W & 0\n\end{pmatrix} +
$$
\n
$$
\begin{pmatrix}\nG_{11} & -G_{21} & \cdots & (-1)^{n+1}G_{n1} \\
0 & g^{-1}|W & 0\n\end{pmatrix}\n\begin{pmatrix}\ng_{11} & g_{12} & \cdots & g_{1n} \\
0 & g_{1N} & 0 & \cdots & 0 \\
\vdots & \vdots & \vdots & \ddots & \vdots \\
0 & 0 & \cdots & 0\n\end{pmatrix} =
$$
\n
$$
\begin{pmatrix}\nG_{11} & -G_{21} & \cdots & (-1)^{n+1}G_{n1} \\
0 & g^{-1}|W & 0\n\end{pmatrix}\n\begin{pmatrix}\n0 & (\alpha_1, ..., \alpha_{n-1})g|W & \cdots & 0 \\
0 & 0 & \cdots & 0 \\
\vdots & \vdots & \ddots & \vdots \\
0 & 0 & \cdots & 0\n\end{pmatrix} =
$$
\n
$$
I + \begin{pmatrix}\n0 & G_{11}(\alpha_1, ..., \alpha_{n-1})g|W & \cdots & 0 \\
0 & 0 & \cdots & 0 \\
\
$$

as claimed.

Corollary 4.2. Suppose $g \in G$ is a transvection. Then

$$
g^{-1}hg = \begin{pmatrix} 1 & & & & & (a_1, ..., a_{n-1})g|W & & & & & & & & 0 \\ 0 & & & 1 & 0 & & & & & & 0 & 0 \\ 0 & & & 0 & & 1 & 0 & & & & & 0 \\ \vdots & & & \vdots & & & & & 0 & \vdots & & \vdots & \vdots \\ 0 & & & & & & \vdots & 0 & & & 0 & 1 & 0 \\ 0 & & & & & & & 0 & & 0 & & 0 & & 0 \end{pmatrix} = \begin{pmatrix} 1 & & & & & & & (a_1, ..., a_{n-1})g|W & & & & & & & 0 \\ 0 & & & & & & & \vdots & & & & & 0 \\ \vdots & & & & & & & & \vdots & & & & & 0 \\ 0 & & & & & & & 0 & & 0 & & & 0 & & 0 \end{pmatrix}
$$

Proof. Since g is transvection we have $g \in SL(V)$. Now g|W is either the identity or a transvection on W. Hence in the 1st case there is nothing to prove and in the 2nd case $det(q|W) = 1$ and $q_{11} = 1$. Since $G_{11} = det(g|W)$, the result follows from **Lemma [4.1](#page-26-1)**. \Box

Let V be a G-module and W a G-submodule. Recall that $Fix_G(W) := \{g \in G | g | W = id\}.$

Proposition 4.3. Suppose G is generated by transvections and $dim_F W = dim_F V - 1$. Then $S(V)^{Fix_G(W)} = S(W)[z],$ where $z = v^{p^t} + q_1v^{p^{t-1}} + ... + q_tv, q_i \in S(W)^{G|W}$ is homogenous of degree $p^t - p^{t-i}$ and $|Fix_G(W)| = p^t$.

Proof. We fix a basis $\{v, w_{n-1}, ..., w_1\}$ of V and $\{w_{n-1}, ..., w_1\}$ a basis of W. The action of all elements of G are given by matrices with respect to this basis. So $Fix_G(W) = \langle \sigma_1, ..., \sigma_t \rangle$, where $\{\sigma_1, ..., \sigma_t\}$ is a minimal generating set and

$$
\sigma_i = \begin{pmatrix}\n1 & \alpha_1^{(i)} & \dots & \dots & \dots & \alpha_{n-1}^{(i)} \\
0 & 1 & 0 & \dots & \dots & 0 & 0 \\
\vdots & \vdots & 0 & & \vdots & \vdots & \vdots \\
0 & \dots & \vdots & 0 & \dots & 0 & 1 & 0 \\
0 & 0 & 0 & \dots & \dots & 0 & 1\n\end{pmatrix}\n= \begin{pmatrix}\n1 & \alpha_1^{(i)} & \dots & \alpha_{n-1}^{(i)} \\
0 & & & \vdots & \vdots \\
0 & & & I_{n-1} & \end{pmatrix}, i = 1, \dots, t.
$$

Hence $\{(\alpha_1^{(i)},...,\alpha_{n-1}^{(i)})|i=1,...,t\}$ is a \mathbb{F}_p -linearly independent set. Since, by **Corollary [4.2](#page-28-1)**, $Fix_G(W)$ is a normal subgroup of G, then for a transvection $g \in G$, $\{g^{-1}\sigma_1g, ..., g^{-1}\sigma_tg\}$ is also a minimal generating set of $Fix_G(W)$. Consequently, $span_{\mathbb{F}_p}\{(\alpha_1^{(i)},...,\alpha_{n-1}^{(i)})|i=1,...,t\}=span_{\mathbb{F}_p}\{(\alpha_1^{(i)},...,\alpha_{n-1}^{(i)})(g|W)|i=1,...,t\}.$

Recall that $S(V)^{Fix_G(W)} = S(W)[z]$, where z is obtained via the following procedure:

$$
\begin{cases}\n z_1 = v^p - (\sigma_1 - I)(v)^{p-1}v = v^p - (\alpha_1^{(1)}w_{n-1} + \dots + \alpha_{n-1}^{(1)}w_1)^{p-1}v, \\
 z_i = z_{i-1}^p - (\sigma_i - I)(z_{i-1})^{p-1}z_{i-1}, i = 2, ..., t. \\
 z = z_t.\n\end{cases}
$$

Similarly, set

$$
\sigma_{i,g} = \begin{pmatrix} 1 &&& (\alpha_1^{(i)}, ..., \alpha_{n-1}^{(i)}) g | W \\ 0 && 1 && 0 & \ldots & \ldots & 0 \\ 0 && 0 && 1 & 0 & \ldots & \ldots & 0 \\ \vdots && \vdots && 0 && \vdots && \vdots & \vdots \\ 0 && & \ldots && \vdots & 0 & \ldots & 0 & 1 \\ 0 && 0 && 0 & \ldots & \ldots & \ldots & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 &&& (\alpha_1^{(i)}, ..., \alpha_{n-1}^{(i)}) g | W \\ 0 &&& I_{n-1} && \\ \vdots &&& I_{n-1} && \\ 0 &&& \ldots & \ldots & 0 & 1 \end{pmatrix}
$$

,

for $i = 1, ..., t$. Hence $z'_1 = v^p - (\sigma_{1,g} - I)(v)^{p-1}v = v^p - (\beta_1^{(1)}w_{n-1} + ... + \beta_{n-1}^{(1)}w_1)^{p-1}v$, where $(\beta_1^{(i)},...,\beta_{n-1}^{(i)}) = (\alpha_1^{(i)},...,\alpha_{n-1}^{(i)}) (g|W)$. Hence $g((\sigma_i - I)(v)) = g(\alpha_1^{(i)} w_{n-1} + ... + \alpha_{n-1}^{(i)} w_1) =$ $\beta_1^{(i)} w_{n-1} + \dots + \beta_{n-1}^{(i)} w_1$, for $i = 1, ..., t$. If $z_i' = (z_i')$ $\binom{d}{i-1}^p - [(\sigma_{i,g} - I)(z^{'}_i)]$ $'_{i-1})]^{p-1}z_i^{p}$ z'_{i-1} and $z' = z'_t$ θ_t , then, by induction, starting with $i = 1$, we have :

$$
\label{eq:2} \left\{ \begin{array}{l} z_i=v^{p^i}+\gamma_{1i}v^{p^{i-1}}+\ldots+\gamma_{ii}v,\gamma_{ji}\in S(W), 1\leq j\leq i.\\ z_i^{'}=v^{p^i}+\delta_{1i}v^{p^{i-1}}+\ldots+\delta_{ii}v,\delta_{ji}\in S(W), 1\leq j\leq i.\\ g(\gamma_{ji})=\delta_{ji}, \forall j=1,...,i. \end{array} \right.
$$

Since $S(V)^{Fix_G(W)} = S(W)[z] = S(W)[z]$ and z, z' are monic polynomials in v with 0 as the free coefficient, it follows that $z' = z$ and consequently $\delta_{jt} = \gamma_{jt}$, for $j = 1, ..., t$. Therefore $g(\gamma_{jt}) = \gamma_{jt}$ for $1 \leq j \leq t$, and all transvections $g \in G$. Hence $q_j := \gamma_{jt} \in S(W)^{G|W}$ for $j = 1, ..., t$.

Let V be a F-finite dimensional G-module,then $V^* = Hom_F(V, F)$ is a G-module by the action $(g \cdot f)(v) = f(g^{-1}v)$, for $f \in V^*$, $g \in G$ and $v \in V$. Let $W \subset V$ be a G -submodule. Set $W^{\perp} = \{f \in V^* | f(W) = 0\}$. Clearly W^{\perp} is a G -submodule of V^* and $dim_F W^{\perp} = dim_F V - dim_F W$.

Lemma 4.4. $V^*/W^{\perp} \cong W^*$ as $G-modules$.

Proof. Consider the map $\pi : V^* \longrightarrow W^*$ given by $\pi(f) = f|W$, that is the restriction of f to $W.Ker(\pi) = \{f \in V^* | f | W = 0\} = W^{\perp}$. Hence the induced map $\bar{\pi}: V^* / W^{\perp} \longrightarrow W^*$ is injective. Since $dim_F V^* - dim_F W^{\perp} = dim_F V - dim_F W^{\perp} = dim_F W = dim_F W^*$ we get that $\bar{\pi}$ is an isomorphism. It is also easy to cheak that π and hence $\bar{\pi}$ are G-module maps. □ We also have the following results.

Corollary 4.5. $V/W \cong (W^{\perp})^*$.

Proof. $(W^{\perp})^{\perp} = \{v \in V | g(v) = 0, \forall g \in W^{\perp}\}\.$ Hence $W \subset (W^{\perp})^{\perp}$. Since $\dim_F(W^{\perp})^{\perp} = \dim_F V^*$ $dim_F W^{\perp} = dim_F V^* - (dim_F V - dim_F W) = dim_F W$ we get that $(W^{\perp})^{\perp} = W$. We now apply **Lemma [4.4](#page-29-0)** to the pair $W^{\perp} \subset V^*$. Hence $V/W = (V^*)^*/(W^{\perp})^{\perp} \cong (W^{\perp})^*$.

$$
\Box
$$

Corollary 4.6. ${g \in G | g(V) \subseteq W} = {g \in G | (g|W^{\perp}) = 0}.$

Proof. This is a consequence of **Corollary [4.5](#page-30-0)**, since $\{g \in G | g(V) \subseteq W\} = Ann_GV/W = Ann_G(W^{\perp})^* = Ann_G(W^{\perp})^*$ $Ann_GW^{\perp} = \{g \in G | g | W^{\perp} = 0\}.$ \Box

Corollary 4.7. Let $W \subset V$ be a G -submodule. Then

- 1. W is faithful G-module iff W^* is a faithful G-module iff V^*/W^{\perp} is a faithful G-module.
- 2. W is an irreducible G-module iff W^* is an irreducible G-module iff V^*/W^{\perp} is an irreducible G−module.

Proof. This follows from **Lemma [4.4](#page-29-0)**.

5 Chapter(D): V is indecomposable,contains a unique 2− dimensional G− submodule, but no 1− dimensional G− submodule.

In this chapter we consider V, a reducible and indecomposable G – module which has a unique G-submodule W with $\dim_F W = 2$, and V contains no 1–dimensional G-submodules, hence W is an irreducible $G-model$.

Lemma 5.1. One of the following holds:

- 1. W is a reducible $T(G)$ module and $\sigma | W = id, \forall \sigma \in T(G)$.
- 2. W is an irreducible $T(G)$ − module.

Proof. Assume that W is a reducible $T(G)$ – module. Then there exists $w \in W$ such that Fw is a $T(G)$ − module, and it follows,using Lemma [2.2](#page-9-1), that $\sigma(w) = w \,\forall \sigma \in T(G)$. Let $g \in G$, since $T(G) \lhd G$, $(g^{-1}\sigma g) = \tau$ for some $\tau \in T(G)$, hence $\sigma(g(w)) = g(\tau(w)) = g(w)$. If $g(w) \notin Fw$ then $\{w, g(w)\}\$ is a basis for W and σ acts trivially on $W \forall \sigma \in T(G)$ so (1) is established. On the other hand if $g(w) \in Fw \ \forall g \in G$, then Fw is 1–dimensional G-module. This contradicts the assumption that V contains no 1−dimensional G−modules.

For W an irreducible $T(G)$ − module, the transvection group $T(G)|W$ had been classified by Kantor whenever $T(G)|W$ is an irreducible primitive linear group, see e.g [\[7,](#page-47-11) Theorem 1.5]. So one of the following holds in this case :

- (i) $T(G)|W = SL(2, \mathbb{F}_q)$, where $p|q$.
- (ii) $T(G)|W \cong SL(2, \mathbb{F}_5), T(G)|W \subset SL(2, \mathbb{F}_9)$ and $\mathbb{F}_9 \subset F$.

Let $\{v, w_2, w_1\}$ be a basis of $V = F^3$ where $\{w_1, w_2\}$ is a basis of W. So every $g \in G$ is represented with respect to this basis, by the matrix:

$$
(5.a) \left(\begin{array}{ccc} \lambda(g) & \delta_{12}(g) & \delta_{13}(g) \\ 0 & \delta_{22}(g) & \delta_{23}(g) \\ 0 & \delta_{32}(g) & \delta_{33}(g) \end{array} \right).
$$

We shall make the following assumption:

Assumption 5.2. Let $\alpha : T(G) \longrightarrow (T(G)|W)$ be the map defined by $\alpha(\tau) = (\tau|W)$ and let $N :=$ $ker(\alpha)$. Then $S(V)^{T(G)}$ is a polynomial ring, $S(V)^{T(G)} = S(W)^{T(G)}[n] = F[n, m_1, m_2]$ where $m =$ (m_1, m_2) is the homogenous maximal ideal of $S(W)^{T(G)} = F[m_1, m_2]$, and $S(V)^N = S(W)[n]$.

The next result provides a necessary and sufficient condition on G to make $S(V)^G$ into a Gorenstein ring .

Proposition 5.3. We keep the above notation and **Assumption [5.2](#page-31-1).** Let $G \subset SL(V)$ be a finite group.Then the following are equivalent:

1. $S(V)^G$ is Gorenstein.

2.
$$
det(g|m/m^2) = \lambda(g)^{1-d}det(g|W)
$$
, where $d = deg(n) = \frac{|T(G)|}{deg(m_1) \cdot deg(m_2)}$ and $m = (m_1, m_2)$.

Proof. Suppose firstly that $S(V)^G$ is Gorenstein. Then by [Theorem [1.2\]](#page-7-1) $det(g|p/p^2) = 1$, where $p = (n, m_1, m_2)$ is the homogenous maximal ideal of $S(V)^{T(G)}$. Clearly g acts linearly on p/p^2 and is represented by the matrix:

$$
\left[\begin{array}{c|c}\lambda(g)^d & * & * \\ \hline 0 & g|m/m^2 \\ 0 & & \end{array}\right]
$$

So $det(g|p/p^2) = \lambda(g)^d det(g|m/m^2) = 1$. Since $G \subset SL(V)$, $\lambda(g) det(g|W) = 1$. Consequently, $\lambda(g)^d \det(g|m/m^2) = \lambda(g) \det(g|W)$, hence $\det(g|m/m^2) = \lambda(g)^{1-d} \det(g|W)$. Suppose now that $det(g|m/m^2) = \lambda(g)^{1-d}det(g|W)$. Hence, using $G \subset SL(V)$, $\lambda(g)^d det(g|m/m^2) = \lambda(g)det(g|W) = 1$. So $det(g|p/p^2) = 1$, then by [Theorem [1.2\]](#page-7-1) $S(V)^G$ is Gorenstein.

Theorem 5.4. Suppose $W \subset V$ is an irreducible G-submodule and $\sigma | W = Id$, for all $\sigma \in T(G)$ (as in Lemma [5.1](#page-31-2) (1)). Then $S(V)^G$ is Gorenstein iff $\lambda(g)^{1-d} = 1$. Consequently if $V = \mathbb{F}_p^3$ then $S(V)^G$ is always Gorenstein.

Proof. By assumption $T(G) = \ker(\alpha)$, where $\alpha : T(G) \longrightarrow (T(G)|W)$ is the restriction map. There-fore by Proposition [4.3](#page-28-0) $S(V)^{T(G)} = S(W)[n]$, where $d = deg(n) = p^t$. Hence Assumption [5.2](#page-31-1) holds here with $m_1 = w_1, m_2 = w_2, m = (w_1, w_2), p = (n, w_1, w_2)$. Since $m/m^2 = Fw_1 + Fw_2 = W$ we have that $det(g|m/m^2) = det(g|W)$, implying by **Proposition [5.3](#page-32-0)** that $S(V)^G$ is Gorenstein if and pnly if $\lambda(g)^{1-d} = 1$. Consequently, if $V = \mathbb{F}_p^3$ then , since $\lambda(g) \in \mathbb{F}_p$, $\lambda(g)^{1-d} = 1$.

 \Box

 \Box

The next result applies in case (i), where W is an irreducible primitive $T(G)$ −module.

Proposition 5.5. Suppose $G \subset SL(V)$ is a finite group with $T(G)|W = SL(2, \mathbb{F}_q)$,where $q = p^s$, $\mathbb{F}_q \subseteq F$ and satisfies **Assumption [5.2](#page-31-1)**. Let $V = F^3$ be the natural $SL(3, F)$ – module. Let $g \in G$ be arbitrary and set $g|W = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with respect to the \mathbb{F}_q basis $\{w_1, w_2\}$. Then $S(V)^G$ is Gorenstein if and only if $a^{q^2-1} = \lambda(g)^{1-d}$.

Proof. The computation in **Proposition [2.4](#page-10-1)** shows that the matrix representing $g|m/m^2$ is $\left(\frac{a^{q-1}det(g|W)}{a^2}\right)$ $\frac{e^{t}(g|W)}{0}$ $\frac{0}{a^{q^2-q}}$, hence $det(g|m/m^2) = a^{q^2-1}det(g|W)$. Hence, by **Proposition [5.3](#page-32-0)**, $S(V)^G$ is Gorenstein if and only if $a^{q^2-1}det(g|W) = det(g|W) \lambda(g)^{1-d}$. Thus $S(V)^G$ is Gorenstein if and only if $a^{q^2-1} = \lambda(g)^{1-d}$. \Box

Remark 5.6. For W an irreducible and imprimitive $T(G)$ −module, $T(G)|W$ is a monomial subgroup (e.g.[\[13\]](#page-47-12)).In this case, if char $F \neq 2$, then by [\[3,](#page-47-5) Lemma 3.9.] $T(G)|W = 1$, which contradicts the assumption that W is an irreducible $T(G)-module$. On the other hand, if $p = 2$, then $T(G)|W = \langle$ σ >,where $\sigma = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. This implies that $(w_2 + w_1) \in S(V)^G$. Hence $\mathbb{F}_2(w_2 + w_1)$ is G -module,which contradicts the assumption that W is an irreducible G−module.Hence we deal only with the case when W is an irreducible primitive $T(G)-module$.

Proposition 5.7. Suppose $F = \mathbb{F}_p$ and W is an irreducible $T(G)$ –module. Then **Assumption [5.2](#page-31-1)** holds for G.

Proof. Clearly

$$
N = \ker(\alpha) \subseteq \left(\begin{array}{ccc} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right).
$$

Since $F = \mathbb{F}_p$, we have by **Remark [5.6](#page-33-0)** that W is a primitive $T(G)$ −module, and therefore $T(G)|W =$ $SL(2, \mathbb{F}_p)$. If $N \neq 1$ then there exists

$$
\sigma := \left(\begin{array}{ccc} 1 & \alpha & \beta \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right), (\alpha, \beta) \neq (0, 0), \sigma \in N.
$$

Consequently by Corollary [4.2](#page-28-1) we have :

$$
\left(\begin{array}{ccc} 1 & (\alpha,\beta)SL(2,\mathbb{F}_p) \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right) \subseteq N,
$$

and therefore since $(\alpha, \beta)SL(2, \mathbb{F}_p) = (\mathbb{F}_p, \mathbb{F}_p)$ we conclude that

$$
N = \left(\begin{array}{ccc} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right).
$$

Let

$$
g = \left(\begin{array}{ccc} 1 & \delta_{12}(g) & \delta_{13}(g) \\ 0 & \delta_{22}(g) & \delta_{23}(g) \\ 0 & \delta_{32}(g) & \delta_{33}(g) \end{array}\right) \in T(G),
$$

be an arbitrary element.Then

$$
\begin{pmatrix} 1 & \delta_{12}(g) & \delta_{13}(g) \\ 0 & \delta_{22}(g) & \delta_{23}(g) \\ 0 & \delta_{32}(g) & \delta_{33}(g) \end{pmatrix} \begin{pmatrix} 1 & -\delta_{12}(g) & -\delta_{13}(g) \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \delta_{22}(g) & \delta_{23}(g) \\ 0 & \delta_{32}(g) & \delta_{33}(g) \end{pmatrix} \in T(G)N = T(G).
$$

Consequently, since $T(G)|W = \{g|W = \left(\begin{smallmatrix} \delta_{22}(g) & \delta_{23}(g) \\ \delta_{32}(g) & \delta_{33}(g) \end{smallmatrix}\right)|g \in T(G)\}\$ and $SL(2, \mathbb{F}_p) = T(G)|W$, then

$$
H := \left(\begin{array}{c|c} 1 & 0 & 0 \\ \hline 0 & SL(2, \mathbb{F}_p) \\ 0 & SL(2, \mathbb{F}_p) \end{array}\right) \subseteq T(G).
$$

Moreover $H \cap N = \{1\}$. Hence $|HN| = |H| \cdot |N| = (p^2 - p) \cdot (p+1) \cdot |N| = |T(G)|W| \cdot |N| = \frac{|T(G)|}{|N|}$ $\frac{|G||}{|N|} \cdot |N| =$ $|T(G)|$, implying that $T(G) = HN$. Let $\{v, w_2, w_1\}$ be the basis representing the matrices above, where $W = span_{\mathbb{F}_p}\{w_1, w_2\}$. Now by **Proposition [4.3](#page-28-0)**, there exists $n = v^{p^2} + q_1v^p + q_2v$, $q_i \in S(W)^{T(G)|W}$, with $S(V)^N = S(W)[n]$. Since $h(v) = v$ for each $h \in H$, it follows that $n \in S(V)^H$, and therefore $S(V)^{T(G)} = S(W)[n]^{\dot{T}(G)/N} = S(W)^{H}[n] = S(W)^{SL(2,\mathbb{F}_p)}[n] = F[d_1, d_2, n]$, where $\{d_1, d_2\}$ are the Dickson invariants of degrees $\{p+1, p^2-p\}$ respectively.

Suppose $N = \{1\}$. Hence $T(G) \cong T(G)|W = SL(2, \mathbb{F}_p)$. Now $T(G) = \langle \phi_1, ..., \phi_k \rangle$, where ϕ_i , is a transvection, for $i = 1, ..., k$. Moreover $k \ge 2$ (since $T(G) \cong SL(2, \mathbb{F}_p)$), and we may assume that at least two out of $\{\phi_1|W,...,\phi_k|W\}$ do not commute, say $\{\phi_1|W,\phi_2|W\}$. Therefore $\langle \phi_1|W,\phi_2|W\rangle \cong$ $SL(2, \mathbb{F}_p)$, implying that $\phi_i | W \in < \phi_1 | W, \phi_2 | W >$, for each $i \geq 3$. Therefore, since α is injective, we have: $\phi_i \in \phi_1, \phi_2 >$, for $i \geq 3$, that is $T(G) = \phi_1, \phi_2 >$. Hence $\ker(\phi_1 - I) \cap \ker(\phi_2 - I) := Fv$, is a one dimensional trivial $T(G)$ −module. Moreover, since W is an irrreducible $T(G)$ −module, $v \notin W$, hence $V = F v \oplus W$ as $T(G)$ –modules. Consequently with respect to the basis $\{v, w_1, w_2\}$ (where $\{w_1, w_2\}$ is any basis of W), $T(G)$ is represent by matrices of the form :

$$
H:=\left(\begin{array}{c|c}1 & 0 & 0 \\ \hline 0 & SL(2,\mathbb{F}_p) \\ 0 & \end{array}\right),
$$

and since $|H| = |SL(2, \mathbb{F}_p)| = |T(G)|W| = |T(G)|$ we have $H = T(G)$. Hence $S(V)^{T(G)} = S(W)^{SL(2, \mathbb{F}_p)}[v] =$ $F[d_1, d_2, v]$ and we take $n = v$.

Theorem 5.8. Suppose $G \subset SL(3, \mathbb{F}_p)$ and W is an irreducible $T(G)$ -submodule of $V = \mathbb{F}_p^3$. Then $S(V)^G$ is Gorenstein.

Proof. By Proposition [5.5](#page-32-1) and Proposition [5.7](#page-33-1), we have to check that $a^{p^2-1} = \lambda(g)^{1-d}$, where $d = deg(n) \in \{1, p^2\}$ and $g|W = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. If $N = 1$ then $d = 1$, and if $N \neq 1$, then $d = p^2$. Since $a, \lambda(g) \in \mathbb{F}_p$ we have $a^{p-1} = \lambda(g)^{p-1} = 1$. Hence $a^{p^2-1} = 1 = \lambda(g)^{1-d}$ for $d \in \{1, p^2\}$.

Corollary 5.9. Suppose $G \subset SL(3, \mathbb{F}_p)$, $W \subset V = \mathbb{F}_p^3$ is an irreducible G -submodule and $dim_{\mathbb{F}_p} W =$ 2. Then $S(V)^G$ is Gorenstein.

Proof. This is a combination of Lemma [5.1](#page-31-2), Theorem [5.4](#page-32-2) and Theorem [5.8](#page-34-0)

 \Box

6 Chapter(E): V is indecomposable, contains a unique $1-$ dimensional G− submodule U, but no 2− dimensional G− submodule.

In this chapter we consider V, a 3-dimensional reducible and indecomposable G – module which has a unique 1–dimensional G-submodule $U = Fv_0$ and V contains no 2–dimensional G-submodules.Hence $W = V/U$ is an irreducible G- module. We shall exclusively consider here the case $F = \mathbb{F}_p$.

Let $\{w_1, w_2, v_0\}$ be any basis of $V = F^3$. So every $g \in G$ is represented with respect to this basis, by the matrix:

$$
(6.a) \t g = \begin{pmatrix} \delta_{11}(g) & \delta_{12}(g) & \delta_{13}(g) \\ \delta_{21}(g) & \delta_{22}(g) & \delta_{23}(g) \\ 0 & 0 & \lambda(g) \end{pmatrix}.
$$

Let $\alpha : T(G) \longrightarrow (T(G)|W)$ be the map which is defined by $\alpha(\tau) = \tau|W$, hence $ker(\alpha) = {\sigma \in T(G)}$ $T(G)|\sigma(x) - x \in Fv_0, \forall x \in V$. Consequently,

$$
ker(\alpha) \subseteq \left(\begin{array}{ccc} 1 & 0 & * \\ 0 & 1 & * \\ 0 & 0 & 1 \end{array}\right).
$$

We shall firstly consider the case that $W = V/\mathbb{F}_p v_0$ is an irreducible $T(G)$ – module (on top of being G–irreducible).Since $F = \mathbb{F}_p$, it follows from **Remark [5.6](#page-33-0)**, that $T(G)|W = SL(2, \mathbb{F}_p)$.

Proposition 6.1. Suppose $V = \mathbb{F}_p^3$, and $V/\mathbb{F}_p v_0$ is an unfaithful, irreducible $T(G)$ –module. Then $S(V)^{T(G)}$ is a polynomial ring.

Proof. Clearly $T(G) \subset SL(V)$. Let $N := ker(\alpha)$, then by assumption $N \neq 1$. Set $U = \mathbb{F}_p v_0$. Then by Corollary [4.6](#page-30-1), $N = Ann_{T(G)}V/U = Ann_{T(G)}U^{\perp}.$

 V/U is an irreducible $T(G)$ -module hence by **Corollary [4.7](#page-30-2)**, U^{\perp} is a 2-dimensional irreducible $T(G)$ – submodule of V^* . Therefore by the proof of the early part of **Proposition [5.7](#page-33-1)**, $|N| = p^2$.

Hence,

with generators

$$
N = \left(\begin{array}{ccc} 1 & 0 & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{array}\right),
$$

$$
\left\{ \left(\begin{array}{ccc} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right), \left(\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{array}\right) \right\}.
$$

Let

$$
g = \begin{pmatrix} \delta_{11}(g) & \delta_{12}(g) & \delta_{13}(g) \\ \delta_{21}(g) & \delta_{22}(g) & \delta_{23}(g) \\ 0 & 0 & 1 \end{pmatrix},
$$

be a transvection in $T(G)$. Then

$$
\begin{pmatrix} 1 & 0 & -\delta_{13}(g) \\ 0 & 1 & -\delta_{23}(g) \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \delta_{11}(g) & \delta_{12}(g) & \delta_{13}(g) \\ \delta_{21}(g) & \delta_{22}(g) & \delta_{23}(g) \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} \delta_{11}(g) & \delta_{12}(g) & 0 \\ \delta_{21}(g) & \delta_{22}(g) & 0 \\ 0 & 0 & 1 \end{pmatrix} \in NT(G) = T(G).
$$

Since $\{ \binom{\delta_{11}(g) \ \delta_{12}(g)}{\delta_{21}(g) \ \delta_{22}(g)} | g \in T(G) \} = SL(2, \mathbb{F}_p) = T(G) | W$ then

$$
H := \left(\begin{array}{c|c} SL(2, \mathbb{F}_p) & 0 \\ \hline 0 & 0 & 1 \end{array}\right) \subseteq T(G).
$$

Since $|T(G)| = |SL(2, \mathbb{F}_p)| \cdot |N| = (p+1)(p^2 - p)p^2$, then $T(G) = \langle H, N \rangle$. We have $S(V)^N =$ $\mathbb{F}_p[v_0, a, b]$ where $a = w_1^p - v_0^{p-1}w_1$ and $b = w_2^p - v_0^{p-1}w_2$. Consider $\{\sigma, \tau\}$, generators of H, where

$$
\tau = \left(\begin{array}{ccc} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right), \sigma = \left(\begin{array}{ccc} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right).
$$

Then $\tau(b) = b, \tau(a) = a + b, \sigma(a) = a, \sigma(b) = a + b$. Hence $A = a^p b - b^p a$ and $B = \frac{a^{p^2} b - b^{p^2} a}{A}$, are Dickson invariants of $(S(V)^N)^H = S(V)^{T(G)}$. So $deg(A) \cdot deg(B) \cdot deg(v_0) = p(p+1) \cdot p(p^2-p) = |T(G)|$. Hence by [\[4,](#page-47-4) Theorem 3.7.5] $S(V)^{T(G)} = \mathbb{F}_p[v_0, A, B]$ is a polynomial ring. \Box

Lemma 6.2. Suppose $V = \mathbb{F}_p^3$, and $W = V/\mathbb{F}_p v_0$ is an irreducible $T(G)$ -module. Then one of the following holds.

- 1. W is an unfaithful irreducible $T(G)$ –module and $T(G) = \langle H, N \rangle$.
- 2. W is a faithful irreducible $T(G)$ module and $T(G) = H$, where H is defined in **Proposition** [6.1](#page-35-1).

Proof. If item (1) holds then by **Proposition [6.1](#page-35-1)** the result follows. So we may assume that W is a faithful irreducible $T(G)$ –module. Let $T(G) = \langle \phi_1, ..., \phi_k \rangle$, so $(T(G)|W) = \langle \phi_1 | W, ..., \phi_k | W \rangle$. If $\phi_i|W = Id$ then $\phi_i \in N = 1$. Hence $\phi_i|W$ are non trivial transvections for $i = 1, ..., k$. Since W is an irreducible $T(G)$ –module then there exist at least two transvections that do not commute ,say $(\phi_1|W)$ and $(\phi_2|W)$. But then W is an irreducible $\langle \phi_1|W, \phi_2|W \rangle$ – module. Hence $\langle \phi_1|W, \phi_2|W \rangle \cong$ $SL(2, \mathbb{F}_p)$. Since $|(T(G)|W)| = |SL(2, \mathbb{F}_p)|$ then $\lt \phi_1 |W, \phi_2 |W \gt = (T(G)|W)$. Therefore, $\forall \phi_i \in T(G)$ with $i \geq 3$, $(\phi_i|W)$ is generated by $(\phi_1|W)$ and $(\phi_2|W)$. So $\phi_i = h \cdot Z$ where $h \in \phi_1, \phi_2 >$ and $Z \in N = ker(\alpha)$. Since $N = 1, \phi_i = h$. Therefore $T(G) = \phi_1, \phi_2 > h$.

Consider $U^{\perp} \subset V^*$. Since $W = V/U$ is a faithful and irreducible $T(G)$ -module it follows from Corollary [4.5](#page-30-0), Corollary [4.7](#page-30-2) that U^{\perp} is a faithful and irreducible $T(G)$ -submodule of V^* .

Consider $ker(\phi_1-I) \cap ker(\phi_2-I) \subset V^*$. It is a 1-dimensional subspace (since ϕ_1, ϕ_2 are transvections on V^*). Let $Fg = \ker(\phi_1 - I) \cap \ker(\phi_2 - I) \subset V^*$. Since $T(G) = \phi_1, \phi_2 >$ it follows that Fg is a trivial $T(G)$ -submodule of V^* . Since U^{\perp} is irreducible it follows that $Fg \oplus U^{\perp} = V^*$, a direct sum decomposition of V^* as $T(G)$ -modules. Since U^{\perp} is irreducible and faithful we have, as in **Proposition [5.7](#page-33-1),** a representation of $T(G)$ as matrices of the form :

$$
\left(\begin{array}{cc|c} 1 & 0 & 0 \\ \hline 0 & SL(2, \mathbb{F}_p) \\ 0 & \end{array}\right),
$$

with respect to the basis $\{g, h_1, h_2\}$, where $\{h_1, h_2\}$ is any basis of U^{\perp} . Let $\{x, w_1, w_2\}$ be the dual basis of $\{g, h_1, h_2\}$. So $g(x) = 1, h_1(x) = h_2(x) = 0, h_i(w_j) = \delta_{ij}$ and $g(w_1) = g(w_2) = 0$. Hence $x \in (Fh_1 + Fh_2)^{\perp} = (U^{\perp})^{\perp} = U$ and $\mathbb{F}_p x = \mathbb{F}_p v_0$.

 $(Fg)^{\perp} = Fw_1 + Fw_2$ is a 2-dimensional $T(G)$ -submodule of V and $(Fg)^{\perp} \oplus (U^{\perp})^{\perp} = V$. Hence the matrices representing $T(G)$ with respect to the basis $\{w_1, w_2, v_0\}$ are

$$
H := \left(\begin{array}{c|c} SL(2, \mathbb{F}_p) & 0 \\ \hline 0 & 0 & 1 \end{array}\right).
$$

The following theorem deals with the special case, when $W = V/\mathbb{F}_p v_0$ is an irreducible primitive $T(G)$ –module and $F = \mathbb{F}_p$. Since $F = \mathbb{F}_p$, by [\[7,](#page-47-11) Theorem 1.5] $(T(G)|W) = SL(2, \mathbb{F}_p)$ and $p \neq 2$.

Theorem 6.3. Suppose that $G \subset SL(3, \mathbb{F}_p)$, $W = V/\mathbb{F}_p v_0$ is an irreducible $T(G)$ –module. Then $S(V)^G$ is Gorenstein.

Proof. Assume firstly that W is a faithful irreducible $T(G)$ -module.By Lemma [6.2](#page-36-0) (2), $S(V)^{T(G)}$ = $S(V)^H = \mathbb{F}_p[u, y, v_0]$ is a polynomial ring where $u = w_1 w_2^p - w_2 w_1^p, y = \frac{w_1 w_2^{p^2} - w_2 w_1^{p^2}}{u}$ are the Dickson's invariants. Now for $g \in G$ since $gT(G)g^{-1} = T(G), \mathbb{F}_p[u, y, v_0]$ is g-stable and since g preserve degrees we get $g(u) = \alpha u + \beta v_0^{p+1}$ with $\alpha, \beta \in \mathbb{F}_p$. By Fermat' little theorem and formula (6.a) we get:

 $g(u) = (\delta_{11}(g)w_1 + \delta_{12}(g)w_2 + \delta_{13}(g)v_0)(\delta_{21}(g)w_1^p + \delta_{22}(g)w_2^p + \delta_{23}(g)v_0^p) - (\delta_{21}(g)w_1 + \delta_{22}(g)w_2 +$ $\delta_{23}(g) v_0) (\delta_{11}(g) w_1^p + \delta_{12}(g) w_2^p + \delta_{13}(g) v_0^p) = \delta_{11}(g) \delta_{22}(g) (\overline{w_1 w_2^p} - w_2 w_1^p) + \delta_{11}(g) \delta_{23}(g) (w_1 v_0^p - v_0 w_1^p) + \overline{\delta_{11}(g) \delta_{23}(g)} \overline{w_1 w_0^p} - \overline{\delta_{12}(g)} \overline{w_1 w_0^p} - \overline{\delta_{13}(g)} \overline{w_2 w_0^p})$ $\delta_{12}(g)\delta_{23}(g)(w_2v_0^p-v_0w_2^p)+\delta_{13}(g)\delta_{22}(g)(w_2^pv_0-w_2v_0^p)+\delta_{12}(g)\delta_{21}(g)(w_2w_1^p-w_1w_2^p)+\delta_{13}(g)\delta_{21}(g)(w_1^pv_0-w_2^p)$ $w_1v_0^p = (\delta_{11}(g)\delta_{22}(g) - \delta_{12}(g)\delta_{21}(g))(w_1w_2^p - w_2w_1^p) + (\delta_{11}(g)\delta_{23}(g) - \delta_{13}(g)\delta_{21}(g))(w_1v_0^p - v_0w_1^p) +$ $(\delta_{12}(g)\delta_{23}(g) - \delta_{13}(g)\delta_{22}(g))(w_2v_0^p - v_0^pw_2).$

Now by comparing coefficients of $g(u)$ we get:

(6.b)
$$
\begin{cases} \alpha = \delta_{11}(g)\delta_{22}(g) - \delta_{12}(g)\delta_{21}(g) \\ \beta = 0 \\ \delta_{11}(g)\delta_{23}(g) - \delta_{13}(g)\delta_{21}(g) = 0 \\ \delta_{12}(g)\delta_{23}(g) - \delta_{13}(g)\delta_{22}(g) = 0 \end{cases}
$$

Since $G \subset SL(V)$, $\lambda(g)^{-1} = \delta_{11}(g)\delta_{22}(g) - \delta_{12}(g)\delta_{21}(g)$, then $g(u) = \lambda(g)^{-1}u$. Similarly, $g(y) =$ $\gamma y + q(u, v_0)$, where $\gamma \in \mathbb{F}_p$ and $q(u, v_0)$ is a homogenous polynomial (in u, v_0) with $deg(q)$ $deg(y) = p^2 - p.$

By Fermat' little theorem , formula (6.a) and (6.b) we get:

$$
g(w_1w_2^{p^2} - w_2w_1^{p^2}) = (\delta_{11}(g)\delta_{22}(g) - \delta_{12}(g)\delta_{21}(g))(w_1w_2^{p^2} - w_2w_1^{p^2}) + (\delta_{11}(g)\delta_{23}(g) - \delta_{13}(g)\delta_{21}(g))(w_1w_0^{p^2} - v_0w_1^{p^2}) + (\delta_{12}(g)\delta_{23}(g) - \delta_{13}(g)\delta_{22}(g))(w_2w_0^{p^2} - w_0^{p^2}w_2) = \lambda(g)^{-1}(w_1w_2^{p^2} - w_2w_1^{p^2}) + 0 + 0,
$$

using $\delta_{11}(g)\delta_{22}(g) - \delta_{12}(g)\delta_{21}(g) = \lambda(g)^{-1}$. Since $uy = w_1w_2^{p^2} - w_2w_1^{p^2}$ we get $g(y) = \frac{g(w_1w_2^{p^2} - w_2w_1^{p^2})}{g(y)} = \frac{\lambda(g)^{-1}(w_1w_2^{p^2} - w_2w_1^{p^2})}{\lambda(g)^{-1}w} = \frac{w_1w_2^{p^2} - w_2w_1^{p^2}}{w} = y$. So by comparing coefficients we get that $\gamma = 1$ and $q(u, v_0) = 0$.

Clearly,g acts linearly on $\mathbb{F}_p u + \mathbb{F}_p y + \mathbb{F}_p v_0$ and is represented by the matrix :

Therefore, by [Theorem [1.2\]](#page-7-1) $S(V)^G$ is Gorenstein.

Assume now that W is an unfaithful irreducible $T(G)$ –module.By **Proposition [6.1](#page-35-1)** $S(V)^{T(G)}$ = $\mathbb{F}_p[A, B, v_0]$ is a polynomial ring where $A = a^p b - b^p a$, $B = \frac{a^{p^2} b - b^{p^2} a}{A}$ and $deg(A) = p^2 + p$, $deg(B) =$ $p(p^2 - p)$. We have :

 $A = a^p b - b^p a = (w_1^{p^2} - v_0^{p(p-1}w_1^p)(w_2^p - v_0^{p(p-1)}w_2) - (w_2^{p^2} - v_0^{p(p-1)})w_2^p(w_1^p - v_0^{p-1}w_1) = (w_1^{p^2}w_2^p - w_0^{p(p-1)})w_2^p$ $w_2^{p^2}w_1^p\rangle-v_0^{p-1}(w_1^{p^2}w_2-w_2^{p^2}w_1)+v_0^{p^2-1}(w_1^pw_2-w_2^pw_1)=(-u)^p-v_0^{p-1}(w_1^{p^2}w_2-w_2^{p^2}w_1)+v_0^{p^2-1}(-u),$ where $u = w_1 w_2^p - w_2 w_1^p$. We get in the previous calculation that :

 $\int g(u) = \lambda(g)^{-1}u.$ $g(w_1^{p^2} w_2 - w_2^{p^2} w_1) = \lambda(g)^{-1} (w_1^{p^2} w_2 - w_2^{p^2} w_1).$ Since $g(v_0^{p^2-1}) = (\lambda(g)^{-1})^{p^2-1}v_0^{p^2-1} = v_0^{p^2-1}$, then $g(A) = \lambda(g)^{-1}A$.

We also have :

$$
\begin{array}{c} a^{p^2}b - b^{p^2}a = (w_1^{p^3} - v_0^{p^2(p-1)}w_1^{p^2})(w_2^p - v_0^{p-1}w_2) - (w_2^{p^3}v_0^{p^2(p-1)}w_2^{p^2})(w_1 - v_0^{p-1}w_1) = (w_1^{p^3}w_2^{p^2} - w_2^{p^3}w_1^p) - v_0^{(p-1)}(w_1^{p^3}w_2 - w_2^{p^3}w_1) + v_0^{p^2(p-1)}(w_2^{p^2}w_1^p - w_2^{p^2}w_2^p) + v_0^{(p^2+1)(p-1)}(w_1^{p^2}w_2 - w_2^{p^2}w_1) = \\ (-u)^{p^2} - v_0^{(p-1)}(w_1^{p^3}w_2 - w_2^{p^3}w_1) + v_0^{p^2(p-1)}(u)^p + v_0^{(p^2+1)(p-1)}(-uy).\end{array}
$$

We get in the previous calculation that $g(u) = \lambda(g)u$ and $g(y) = 1$. On the other hand we have :

$$
g(v_0^{(p-1)}(w_1^{p^3}w_2 - w_2^{p^3}w_1)) = (\delta_{11}(g)\delta_{22}(g) - \delta_{12}(g)\delta_{21}(g))(w_1^{p^3}w_2 - w_2^{p^3}w_1) + (\delta_{23}(g)\delta_{11}(g) - \delta_{13}(g)\delta_{21}(g))(v_0^{p^3}w_2 - w_2^{p^3}v_0) + (\delta_{22}(g)\delta_{13}(g) - \delta_{12}(g)\delta_{23}(g))(w_1^{p^3}v_0 - w_2^{p^3}v_0).
$$
 Hence by **(6.b)**, $g(v_0^{(p-1)}(w_1^{p^3}w_2 - w_2^{p^3}w_1)) = \lambda(g)^{-1}(v_0^{(p-1)}(w_1^{p^3}w_2 - w_2^{p^3}w_1).$

Consequently $g(a^{p^2}b - b^{p^2}a) = \lambda(g)^{-1}(a^{p^2}b - b^{p^2}a)$, hence $g(B) = \frac{g(a^{p^2}b - b^{p^2}a)}{g(A)} = \frac{\lambda(g)^{-1}(a^{p^2}b - b^{p^2}a)}{\lambda(g)^{-1}A}$ B. So g acts linearly on $\mathbb{F}_p A + \mathbb{F}_p B + \mathbb{F}_p v_0$ and is represented by the matrix :

$$
\left(\begin{array}{ccc} \lambda(g)^{-1} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \lambda(g) \end{array}\right).
$$

Again, by [Theorem [1.2\]](#page-7-1) $S(V)^G$ is Gorenstein.

We shall now consider the possibility that W is a reducible $T(G)$ –module.

Lemma 6.4. Suppose $W = V/Fv$ is a reducible $T(G)$ −module. Then W is a trivial $T(G)$ −module, that is $ker(\alpha) = T(G)$.

Proof. Let $F\overline{w_1}$ be a $T(G)$ − submodule of W. Hence $\phi(\overline{w_1}) = \overline{w_1}, \forall \phi \in T(G)$. Since W is irreducible G-submodule, $F\bar{w_1}$ is not a G-submodule.Consequently, there exists $g \in G$ such that $g(\bar{w_1}) = \bar{w_2} \notin$ $F\bar{w_1}$. Since $T(G) \lhd G$, $\forall \sigma \in T(G)$, $g^{-1}\sigma g = \tau$, for some $\tau \in T(G)$. Hence $(g^{-1}\sigma g)(\bar{w_1}) = \tau(\bar{w_1}) = \bar{w_1}$, and therefore $\sigma(g(\bar{w_1})) = g(\bar{w_1})$, hence $\sigma(\bar{w_2}) = \bar{w_2}$. But $\{\bar{w_1}, \bar{w_2}\}$ is a basis of W.Consequently W is a trivial $T(G)$ −module.

Corollary 6.5. Suppose $W = V/\mathbb{F}_p v_0$ is a reducible $T(G)$ -module, then $S(V)^{T(G)}$ is a polynomial ring.

Proof.

$$
ker(\alpha) = T(G) \subseteq \left(\begin{array}{ccc} 1 & 0 & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{array} \right).
$$

Hence $|T(G)| \in \{p, p^2\}$. If $|T(G)| = p^2$, then

$$
T(G) = \left(\begin{array}{ccc} 1 & 0 & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{array} \right),
$$

 \Box

with generators :

$$
\left(\begin{array}{rrr}1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1\end{array}\right), \left(\begin{array}{rrr}1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1\end{array}\right).
$$

Hence $S(V)^{T(G)} = \mathbb{F}_p[v_0, x, y]$, where $x = w_1^p - v_0^{p-1}w_1$, $y = w_2^p - v_0^{p-1}w_2$. If $|T(G)| = p$, then $T(G)$ is generated by :

$$
\left(\begin{array}{ccc} 1 & 0 & \alpha \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{array}\right), \alpha, \beta \in \mathbb{F}_p.
$$

Then as in Lemma [3.2](#page-15-0) $S(V)^{T(G)} = \mathbb{F}_p[v_0, \alpha w_2 - \beta w_1, z]$ is a polynomial ring and $z = w_1^p$ - $(\alpha v_0)^{p-1}w_1.$

Corollary 6.6. Suppose $W = V/\mathbb{F}_p v_0$ is a reducible $T(G)$ − module and an irreducible G−module. Then $|T(G)| = p^2$.

Proof. Suppose as in **Corollary [6.5](#page-39-0)**, that $|T(G)| = p$, so

$$
T(G) = < \left(\begin{array}{rrr} 1 & 0 & \alpha \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{array}\right) > .
$$

Since $\mathbb{F}_p v_0, \mathbb{F}_p(\alpha w_1 - \beta w_2)$ are trivial $T(G)$ – modules we have that $A = \mathbb{F}_p v_0 + \mathbb{F}_p(\alpha w_1 - \beta w_2)$ $ker(\sigma - I)$ for each transvection $\sigma \in T(G)$. Let $g \in G$ and σ a transvection in G. Then $g^{-1}\sigma g = \tau$ for some transvection τ . Then for each $a \in A$ we have $(g^{-1}\sigma g)(a) = \tau(a) = a$ and hence $\sigma(g(a)) = g(a)$, implying that $g(a) \in A$ for each $g \in G$. Hence A is a 2-dimensional G−submodule of V. This contradicts the assumption on V.

Theorem 6.7. Suppose $W = V/\mathbb{F}_p v_0$ is a reducible $T(G)$ – module and an irreducible G–module. Then $S(V)^G$ is Gorenstein.

 \Box

Proof. We follow here the argument in **Proposition [3.13](#page-23-0)** with $\begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $\begin{pmatrix} c \\ d \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. Recall that by Corollary [6.5](#page-39-0) and Corollary [6.6](#page-40-0),

$$
T(G) = \left(\begin{array}{ccc} 1 & 0 & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{array} \right),
$$

with generators :

$$
\left(\begin{array}{rrr}1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1\end{array}\right), \left(\begin{array}{rrr}1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1\end{array}\right).
$$

Hence $S(V)^{T(G)} = \mathbb{F}_p[v_0, x, y]$, where $x = w_1^p - v_0^{p-1}w_1$, $y = w_2^p - v_0^{p-1}w_2$. So if $m = (v_0, x, y)$ then $\bar{g} \in G/T(G)$ is represented on $m/m^2 = \mathbb{F}_p \bar{y} + \mathbb{F}_p \bar{x} + \mathbb{F}_p v_0$ by the matrix :

$$
\left(\begin{array}{ccc} \alpha & \gamma & 0 \\ \varepsilon & \delta & 0 \\ 0 & 0 & \lambda(g) \end{array}\right),
$$

and $g \in G$ is represented on $V = \mathbb{F}_p w_2 + \mathbb{F}_p w_1 + \mathbb{F}_p v_0$, by the matrix :

$$
\left(\begin{array}{ccc} \lambda_2(g) & \delta_{12}(g) & \delta_{13}(g) \\ \delta_{21}(g) & \lambda_1(g) & \delta_{23}(g) \\ 0 & 0 & \lambda(g) \end{array}\right).
$$

We shall show that $det(\bar{g}) = 1$, for each $\bar{g} \in G/T(G)$, implying by [Theorem [1.2\]](#page-7-1) that $S(V)^G$ is Gorenstein. Now $g(y) = \alpha y + \gamma x + \beta v_0^p$ and $g(x) = \delta x + \varepsilon y + \chi v_0^p$ (keeping the notation in **Lemma [3.13](#page-23-0)**). Siem. Now $g(y) = \alpha y + \gamma x + \beta v_0$ and $g(x) = \alpha x + \epsilon y + \chi v_0$ (keeping the notation in **Lemma 3.15**).
Since $g(w_2) = \lambda_2(g)w_2 + \delta_{12}(g)w_1 + \delta_{13}(g)v_0$, we have $g(y) = g(w_2)^p - (\lambda(g)v_0)^{p-1}g(w_2) = (\lambda_2(g)w_2 + \lambda_3(g)w_1 + \lambda_4(g)w_2)$ $\delta_{12}(g)w_1 + \delta_{13}(g)v_0)^p - (\lambda(g)v_0)^{p-1}(\lambda_2(g)w_2 + \delta_{12}(g)w_1 + \delta_{13}(g)v_0) = (\lambda_2(g)^pw_2^p - \lambda(g)^{p-1}\lambda_2(g)v_0^{p-1}w_2) +$ $(\delta_{12}(g)^p w_1^p - \lambda(g)^{p-1} \delta_{12}(g) v_0^{p-1} w_1) + (\delta_{13}(g)^p - \lambda(g)^{p-1} \delta_{13}(g)) v_0^p.$

Consequently $\alpha = \lambda_2(g)^p = \lambda(g)^{p-1}\lambda_2(g)$, $\gamma = \delta_{12}(g)^p = \lambda(g)^{p-1}\delta_{12}(g)$. Consequently, since $\lambda_2(g)$, $\delta_{12}(g) \in$ \mathbb{F}_p , it follows that $\alpha = \lambda_2(g)$ and $\gamma = \delta_{12}(g)$.

Similarly using $g(w_1) = \delta_{21}(g)w_2 + \lambda_1(g)w_1 + \delta_{23}(g)v_0$, we have $g(x) = g(w_1)^p - (\lambda(g)v_0)^{p-1}g(w_1) =$ $(\delta_{21}(g)w_2 + \lambda_1(g)w_1 + \delta_{23}(g)v_0)^p - (\lambda(g)v_0)^{p-1}(\delta_{21}(g)w_2 + \lambda_1(g)w_1 + \delta_{23}(g)v_0) = (\delta_{21}(g)^pw_2^p \lambda(g)^{p-1}\delta_{21}(g)v_0^{p-1}w_2)+(\lambda_1(g)^pw_1^p-\lambda(g)^{p-1}\lambda_1(g)v_0^{p-1}w_1)+(\delta_{23}(g)^p-\lambda(g)^{p-1}\delta_{23}(g))v_0^p.$ Therefore $\varepsilon = \delta_{21}(g)^p$, $\delta = \lambda_1(g)^p$, and since $\lambda_1(g)$, $\delta_{21}(g) \in \mathbb{F}_p$, we have $\varepsilon = \delta_{21}(g)$, $\delta = \lambda_1(g)$. Therefore $det(\bar{g}) = (\alpha \delta - \varepsilon \gamma) \lambda(g) = (\lambda_2(g) \lambda_1(g) - \delta_{21}(g) \delta_{12}(g)) \lambda(g) = det(g) = 1$ (since $G \subset SL(3, \mathbb{F}_p)$). \Box

As a consequence of Theorem [6.3](#page-37-0) and Theorem [6.7](#page-40-1) we have :

Corollary 6.8. Suppose $W = V/\mathbb{F}_p v_0$ is an irreducible G-module. Then $S(V)^G$ is Gorenstein.

7 Chapter(F): V is indecomposable and contains two $G-$ submodules $\mathbb{F}_p v_0, W$, where $\mathbb{F}_p v_0 \subset W \subset V$, and $dim_{\mathbb{F}_p} W = 2$.

In this chapter we conside V, a reducible indecomposable G−module,which has a unique 1−dimensional G– submodule Fv_0 , and a unique 2–dimensional G– submodule W, and $Fv_0 \subset W \subset V$. We shall exclusively consider the case $F = \mathbb{F}_p$.

Let $\{w_2, w_1, v_0\}$ be a basis of $V = \mathbb{F}_p^3$, where $W = \mathbb{F}_p w_1 + \mathbb{F}_p v_0$. So every $g \in G$ is represented with respect to this basis , by the matrix :

$$
g = \begin{pmatrix} \lambda_2(g) & \delta_{12}(g) & \delta_{13}(g) \\ 0 & \lambda_1(g) & \delta_{23}(g) \\ 0 & 0 & \lambda(g) \end{pmatrix}.
$$

By Lemma [2.2](#page-9-1), $T(G)$ acts trivially on $\mathbb{F}_p v_0$, and also on $W/\mathbb{F}_p v_0$, hence

$$
T(G) \subseteq \left(\begin{array}{ccc} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{array} \right).
$$

So one of the following three cases holds:

- (i) $|T(G)| = p$,
- (ii) $|T(G)| = p^2$,
- (iii) $|T(G)| = p^3$.

Corollary 7.1. If case (i) holds. Then $S(V)^G$ is Gorenstein.

Proof. Since $|T(G)| = p, T(G) := \langle \sigma \rangle$ or $\langle \tau \rangle$, where

$$
\sigma = \left(\begin{array}{ccc} 1 & 0 & a \\ 0 & 1 & b \\ 0 & 0 & 1 \end{array}\right), \tau = \left(\begin{array}{ccc} 1 & c & d \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right), a, b, c, d \in \mathbb{F}_p.
$$

Assume firstly that $T(G) = \langle \sigma \rangle$. We shall change the basis of V into $\{w_2, aw_1 - bw_2, v_0\}$. We have, $\sigma(aw_1 - bw_2) = a(w_1 + bv_0) - b(w_2 + av_0) = aw_1 - bw_2$. Suppose $a \neq 0$, so $0 \neq w'_1 := (aw_1 - bw_2) \in$ $S(V)^{T(G)}$. Therefore σ is represented with respect to $\{w_2, w'_1, v_0\}$ by the matrix:

$$
\sigma = \left(\begin{array}{rrr} 1 & 0 & a \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right).
$$

Consequently, by [\[3,](#page-47-5) Lemma 3.13] $S(V)^{T(G)} = \mathbb{F}_p[x, w'_1, v_0]$ is a polynomial ring, and $x = w_2^p$ – $(av_0)^{p-1}w_2$. Let $m = (x, w'_1, v_0)$ be the graded maximal homogenous ideal of $S(V)^{T(G)}$.

Since $T(G) \lhd G, \mathbb{F}_p[x, w'_1, v_0]$ is g-stable and since g preserve degrees we get that $g(\mathbb{F}_p w'_1 + \mathbb{F}_p v_0) \in$ $(\mathbb{F}_p w'_1 + \mathbb{F}_p v_0)$ for every $g \in G$. This implies that $(\mathbb{F}_p w'_1 + \mathbb{F}_p v_0)$ is G -module. Therefore G has a triangular presentation with respect to the new basis $\{w_2, w'_1, v_0\}$. Hence we still use the same $\lambda_2(g), \delta_{12}(g), \delta_{13}(g), \lambda_1(g), \delta_{23}(g)$, for the new presentation of g, with respect to the basis $\{w_2, w'_1, v_0\}$. We have:

$$
(7.a) \begin{cases} g(x) = \lambda_2(g)^p w_2^p + \delta_{12}(g)^p (w_1')^p + \delta_{13}(g)^p v_0^p - (a\lambda(g)v_0)^{p-1} (\lambda_2(g)w_2 + \delta_{12}(g)w_1' + \delta_{13}(g)v_0).\\ g(w_2) = \lambda_2(g)w_2 + \delta_{12}(g)w_1' + \delta_{13}(g)v_0.\\ g(w_1') = \lambda_1(g)w_1' + \delta_{23}(g)v_0.\\ g(v_0) = \lambda(g)v_0. \end{cases}
$$

Now since $gT(G)g^{-1} = T(G), \mathbb{F}_p[x, w'_1, v_0]$ is g-stable and since g preserve degrees we get:

$$
(7.b)\left\{\n\begin{array}{l}\ng(x) = \alpha_1(w_2^p - (av_0)^{p-1}w_2) + \alpha_2(w_1')^p + \alpha_3v_0^p + \text{mixed terms in }\{w_1', v_0\}.\ \alpha_1, \alpha_2, \alpha_3 \in \mathbb{F}_p.\n\end{array}\n\right.
$$

Hence by comparing coefficient of w_2^p in $g(x)$, we get that $\alpha_1 = \lambda_2(g)^p$. So g acts linearly on m/m^2 and is represented by the matrix:

$$
\left(\begin{array}{ccc} \lambda_2(g)^p&0&0\\ 0&\lambda_1(g)&\delta_{23}(g)\\ 0&0&\lambda(g) \end{array}\right)=\left(\begin{array}{ccc} \lambda_2(g)&0&0\\ 0&\lambda_1(g)&\delta_{23}(g)\\ 0&0&\lambda(g) \end{array}\right).
$$

Hence by [Theorem [1.2\]](#page-7-1) $S(V)^G$ is Gorenstein. If $a = 0$, then $S(V)^{T(G)} = \mathbb{F}_p[w_2, x, v_0]$ where $x =$ $w_1^p - (bv_0)^{p-1}w_1$. So we have:

$$
(7.c)\begin{cases} g(x) = (\lambda_1(g)w_1 + \delta_{23}(g)v_0)^p - (bv_0)^{p-1}(\lambda_1(g)w_1 + \delta_{23}(g)v_0).\\ g(x) = \gamma_1 x + \gamma_2 w_2^p + \gamma_3 v_0^p + \text{mixed terms in } \{w_2, v_0\}. \qquad \gamma_1, \gamma_2, \gamma_3 \in \mathbb{F}_p. \end{cases}
$$

Hence by comparing coefficient of w_1^p in both expressions, we get that $\gamma_1 = \lambda_1(g)^p = \lambda_1(g)$. Hence by taking $m = (w_2, x, v_0)$, the matrix represinting $g|m/m^2$ with respect to $\overline{w_2}, \overline{x}, \overline{v_0}$ is

$$
\left(\begin{array}{ccc} \lambda_2(g) & \delta_{12}(g) & \delta_{13}(g) \\ 0 & \lambda_1(g)^p & 0 \\ 0 & 0 & \lambda(g) \end{array}\right),
$$

so its determinant is $\lambda_2(g) \cdot \lambda_1(g) \cdot \lambda(g) = 1$, for each $g \in G$, and $S(V)^G$ is Gorenstein.

Assume now that $T(G) = \langle \tau \rangle$. If $c = 0$, then we are exactly in the first case. So we may assume that $c \neq 0$. Consider $w'_1 := cw_1 + dv_0$, then $\{w_2, w'_1, v_0\}$ is a basis of V, with $\{w'_1, v_0\}$ a basis of W. Then the matrix of τ with respect to this new basis is

$$
\tau = \left(\begin{array}{rrr} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right).
$$

Hence $S(V)^{< \tau>} = \mathbb{F}_p[z, w'_1, v_0]$, where $z = w_2^p - (w'_1)^{p-1}w_2$.

By the same discussion as above,we have:

$$
(7.d) \begin{cases} g(z) = \lambda_2(g)^p w_2^p + \delta_{12}(g)^p (w_1')^p + \delta_{13}(g)^p v_0^p + g(-(w_1')^{p-1})(\lambda_2(g)w_2 + \delta_{12}(g)w_1' + \delta_{13}(g)v_0).\\ g(z) = \beta_1 z + \beta_2(w_1')^p + \beta_3 v_0^p + \text{mixed terms in } \{w_1', v_0\}. \quad \beta_1, \beta_2, \beta_3 \in \mathbb{F}_p.\\ g(w_1') = \lambda_1(g)w_1' + \delta_{23}(g)v_0. \end{cases}
$$

Hence by comparing coefficient of w_2^p in $g(z)$, we get $\beta_1 = \lambda_2(g)^p$. Let $m = (z, w'_1, v_0)$ be the graded maximal homogenous ideal of $S(V)^{T(G)}$. So g acts linearly on m/m^2 and is represented by the matrix:

$$
\left(\begin{array}{ccc} \lambda_2(g)^p & 0 & 0 \\ 0 & \lambda_1(g) & \delta_{23}(g) \\ 0 & 0 & \lambda(g) \end{array}\right) = \left(\begin{array}{ccc} \lambda_2(g) & 0 & 0 \\ 0 & \lambda_1(g) & \delta_{23}(g) \\ 0 & 0 & \lambda(g) \end{array}\right).
$$

Hence by [Theorem [1.2\]](#page-7-1) $S(V)^G$ is Gorenstein.

Corollary 7.2. If case (ii) holds. Then $S(V)^G$ is Gorenstein.

Proof. Since $|T(G)| = p^2$, it is elementary abelian, hence

$$
T(G) = \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad or \quad T(G) = \begin{pmatrix} 1 & 0 & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{pmatrix}.
$$

Assume firstly that,

$$
T(G) = \left(\begin{array}{ccc} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right).
$$

Hence by **Proposition [4.3](#page-28-0)**, $S(V)^{T(G)} = S(V)^{Fix_{T(G)}(W)} = S(W)[t]$ is a polynomial ring where $t = w_2^{p^2} + q_1 w_2^p + q_2 w_2$, and $q_1, q_2 \in \mathbb{F}_p[w_1, v_0]$ is homogenous of degrees $p^2 - p$, $p^2 - 1$ respectively. Let $m = (t, w_1, v_0)$ be the graded maximal homogenous ideal of $S(V)^{T(G)}$. By a similar discussion to the one in (7.d) we get that g acts linearly on m/m^2 and is represented by the matrix:

$$
\begin{pmatrix} \lambda_2(g)^{p^2} & 0 & 0 \\ 0 & \lambda_1(g) & \delta_{23}(g) \\ 0 & 0 & \lambda(g) \end{pmatrix} = \begin{pmatrix} \lambda_2(g) & 0 & 0 \\ 0 & \lambda_1(g) & \delta_{23}(g) \\ 0 & 0 & \lambda(g) \end{pmatrix}.
$$

Consequently, by [Theorem [1.2\]](#page-7-1) $S(V)^G$ is Gorenstein.

Assume now that

$$
T(G) = \left(\begin{array}{ccc} 1 & 0 & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{array} \right).
$$

Hence by the **Corollary [6.5](#page-39-0)**, $S(V)^{T(G)} = \mathbb{F}_p[y, x, v_0]$ is a polynomial ring, where $y = w_2^p - v_0^{p-1}w_2$ and $x = w_1^p - v_0^{p-1} w_1$. We have, using $F = \mathbb{F}_p$ (so $\lambda(g)^{p-1} = 1$):

$$
(7.e) \begin{cases} g(y) = (\lambda_2(g)w_2^p + \delta_{12}(g)w_1^p + \delta_{13}(g)v_0^p) - v_0^{p-1}(\lambda_2(g)w_2 + \delta_{12}(g)w_1 + \delta_{13}(g)v_0) = \lambda_2(g)y + \delta_{12}(g)x, \\ g(x) = (\lambda_1(g)w_1^p + \delta_{23}(g)v_0^p) - v_0^{p-1}(\lambda_1(g)w_1 + \delta_{23}(g)v_0) = \lambda_1(g)x. \end{cases}
$$

Let $m = (y, x, v_0)$ be the graded maximal homogenous ideal of $S(V)^{T(G)}$. So g acts linearly on m (resp. m/m^2) and is represented by the matrix:

$$
\left(\begin{array}{ccc} \lambda_2(g) & \delta_{12}(g) & 0 \\ 0 & \lambda_1(g) & 0 \\ 0 & 0 & \lambda(g) \end{array}\right).
$$

Consequently, by [Theorem [1.2\]](#page-7-1) $S(V)^G$ is Gorenstein.

Corollary 7.3. If case (iii) holds. Then $S(V)^G$ is Gorenstein.

Proof. Since $|T(G)| = p^3$,

$$
T(G) = \left(\begin{array}{ccc} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{array}\right).
$$

Hence $T(G) = U_3(\mathbb{F}_p)$, the group of all upper triangular matrices with 1's along the diagonal and entries from the finite field \mathbb{F}_p . This group was shown to have a polynomial ring of invariants by Bertin [\[1\]](#page-47-13).

We shall find explicitly the generators of $S(V)^{T(G)}$. $T(G) = \langle \sigma_1, \sigma_2, \sigma_3 \rangle$, where

$$
\sigma_1 = \left(\begin{array}{ccc} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right), \sigma_2 = \left(\begin{array}{ccc} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right), \sigma_3 = \left(\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{array}\right).
$$

Consequently, by [\[3,](#page-47-5) Lemma 3.13], $S(V)^{< \sigma_1>}= \mathbb{F}_p[a,w_1,v_0]$, is a polynomial ring where $a=w_2^p$ – $v_0^{p-1}w_2$. Define $u = a^p - b^{p-1}a$, where $b := w_1^p - v_0^{p-1}w_1$. Observe that $\sigma_3(b) = b, \sigma_3(a) = a, \sigma_2(b) = b$ $b, \sigma_2(a) = a + b$, hence $\sigma_2(u) = u$. We have, for $i = 1, 2, 3$:

$$
\begin{cases}\n\sigma_i(u) = u.\n\sigma_i(b) = b.\n\sigma_i(v_0) = v_0.\n\end{cases}
$$

The $T(G)$ −invariants $\{u, b, v_0\}$ are algebraically independent over \mathbb{F}_p . Since $deg(u) \cdot deg(v) \cdot deg(v_0) =$ $p^2 \cdot p \cdot 1 = p^3 = |T(G)|$, we have by [\[4,](#page-47-4) Theorem 3.9.4] that $S(V)^{T(G)} = \mathbb{F}_p[u, b, v_0]$ is a polynomial ring.Let $m = (u, b, v_0)$ be the graded maximal homogenous ideal of $S(V)^{T(G)}$. Since $g(b) = \lambda_1(g)^p w_1^p + s = \alpha b + \beta v_0^p$, where $s \in m^2$, and $\alpha, \beta \in \mathbb{F}_p$, we get that $\alpha = \lambda_1(g)^p$. Since $g(u) = \lambda_2(g)^{p^2}u + r = \gamma u + \delta b^p + \pi v_0^{p^2}$ mixed terms in $\{b, v_0\}$, where $r \in m^2$, and $\gamma, \delta, \pi \in \mathbb{F}_p$, we

$$
g|m/m^{2} = \begin{pmatrix} \lambda_{2}(g)^{p^{2}} & 0 & 0 \\ 0 & \lambda_{1}(g)^{p} & 0 \\ 0 & 0 & \lambda(g) \end{pmatrix} = \begin{pmatrix} \lambda_{2}(g) & 0 & 0 \\ 0 & \lambda_{1}(g) & 0 \\ 0 & 0 & \lambda(g) \end{pmatrix}.
$$

Consequently, by [Theorem [1.2\]](#page-7-1) $S(V)^G$ is Gorenstein.

get that $\gamma = \lambda_2(g)^{p^2}$. Hence,

8 Chapter (G) : *V* is indecomposable and contains (at least) two 1−dimensional G− submodules.

Let Fw_1, Fw_2 be the two 1–dimensional G–submodules. Hence $W = Fw_1 + Fw_2$ is G–submodule of V. Let $v \in V \backslash W$, so $\{v, w_1, w_2\}$ is a basis of V. Each $g \in G$ is represented with respect to this basis by the matrix :

$$
g = \left(\begin{array}{ccc} \lambda(g) & \delta_{12}(g) & \delta_{13}(g) \\ 0 & \lambda_2(g) & 0 \\ 0 & 0 & \lambda_1(g) \end{array}\right).
$$

In particular each transvection $\sigma \in T(G)$ is represented by the matrix :

$$
\sigma = \left(\begin{array}{ccc} \lambda(\sigma) & \delta_{12}(\sigma) & \delta_{13}(\sigma) \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right).
$$

Hence $T(G)|W = Id$. Hence

$$
T(G) \subseteq \left(\begin{array}{ccc} 1 & F & F \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right).
$$

Therefore by **Proposition [4.3](#page-28-0)** $S(V)^{T(G)} = S(W)[z]$, where $z = v^{p^t} + q_1 v^{p^{t-1}} + ... + q_t v$, an homogenous polynomial, where $q_i \in S(W)^{T(G)|W} = S(W)$ and $|T(G)| = p^t$. Let $m = (z, w_2, w_1)$. Then $m/m^2 = F\overline{z} + F\overline{w}_2 + F\overline{w}_1$. Now for $g \in G$ we have $g(z) = \alpha z + x$ where x is in m^2 . Therefore $\bar{g}(\bar{z}) = \alpha \bar{z}$, where $\bar{g} = gT(G)$. Also $g(z) = g(v)^{p^t} + g(q_1)g(v)^{p^{t-1}} + ... + g(q_t)g(v) = \lambda(g)^{p^t}v^{p^t} + \text{mixed}$ terms. Hence $\alpha = \lambda(g)^{p^t}$ and therefore \bar{g} is represented on $F\bar{z} + F\bar{w_2} + F\bar{w_1}$ by the matrix :

$$
\left(\begin{array}{ccc} \lambda(g)^{p^t} & 0 & 0 \\ 0 & \lambda_2(g) & 0 \\ 0 & 0 & \lambda_1(g) \end{array}\right).
$$

Therefore by [Theorem [1.2\]](#page-7-1) $S(V)^G$ is Gorenstein iff $\lambda(g)^{p^t} \cdot \lambda_2(g) \cdot \lambda_1(g) = 1$, for each $g \in G$.

Corollary 8.1. Suppose $V = \mathbb{F}_p^3$, contains two 1- dimensional G-submodules and $G \subseteq SL(3, \mathbb{F}_p)$. Then $S(V)^G$ is Gorenstein.

Proof. $\lambda(g) \in \mathbb{F}_p$ hence $\lambda(g)^{p^t} \cdot \lambda_2(g) \cdot \lambda_1(g) = \lambda(g) \cdot \lambda_2(g) \cdot \lambda_1(g) = det(g) = 1$, where the last equality holds since $g \in SL(3, \mathbb{F}_p)$.

9 References

References

- [1] M.J. Bertin. Sous-anneaux d'invariants d'anneaux de polynomes . C.R.Acad.Sci.Paris., 260:5655–5658, 1965.
- [2] A. Braun. The Gorenstein property and Kemper et al.conjecture. David Benson's Front Page.Groups ,Representations and Cohomology Preprint Archive, 2014.
- [3] A. Braun. The Gorenstein property for modular binary forms invariants. J.Algebra, 451:232–247, 2016.
- [4] H. Derksen, G.Kemper. Computational invariant theory,in:Invariant Theory and Algebraic Transformation Groups,I,in:Encyclopaedia Math.Sci. Springer-Verlag,Berlin., 130, 2002.
- [5] J.A. Eagon, M.Hochster. Cohem-Macaulay rings ,invariant theory and the generic perfection of deteminantal loci. ,Amer.J.Math., 93:1020–1058, 1971.
- [6] P. Fleishmann, C.Woodcock. Relative invariants,ideal classes and quasi-canonical modules of modular rings of invariants . J.Algebra, 348:110–134, 2011.
- [7] G. Kemper, G.Malle. The finite irreducible linear groups with polynomial ring of invariants . Transform.Groups., 2 (1):57–89., 1997.
- [8] H. Nakajima. Invariants of Finite Groups Generated by Pseudoreflections in Positive Characteristics . Tsukuba J.Math., 3, (1979) 109-122.
- [9] M.D. Neusel, L.Smith. Invariants Theory of Finite Groups . Math.surveys and Monographs., vol.94, American Mathematical Society, 2002.
- [10] K. Watanabe. Certain invariant subring are Gorenstein 1 . Osaka J.Math., 2:1–8., 1974.
- [11] K. Watanabe. Certain invariant subring are Gorenstein 2 . Osaka J.Math., 2:379–388., 1974.
- [12] CampellH.E.A. WehlauD.L. Non-Cohen-Macaulay Vector Invariants and a Noether Bound for a Gorenstein Ring of Invariats. Canad.Math.Bull, 42 (2):155–161, 1999.
- [13] A.E. Zalesskii, V.N. Serezkin. Linear groups generated by transvections . Math.USSR, Izv., 10:25–46., 1976.