A spectral approach to quenched linear and higher-order response for partially hyperbolic dynamics

Harry Crimmins*and Yushi Nakano[†]

May 25, 2021

Abstract

For smooth random dynamical systems we consider the quenched linear and higherorder response of equivariant physical measures to perturbations of the random dynamics. We show that the spectral perturbation theory of Gouëzel, Keller, and Liverani [36, 32], which has been applied to deterministic systems with great success, may be adapted to study random systems that possess good mixing properties. As a consequence, we obtain general linear and higher-order response results for random dynamical systems that we then apply to random Anosov diffeomorphisms and random U(1) extensions of expanding maps. We emphasise that our results apply to random dynamical systems over a general ergodic base map, and are obtained without resorting to infinite dimensional multiplicative ergodic theory.

1 Introduction

In this paper we study quenched response theory for random dynamical systems (RDSs). The setup is as follows: take M to be a \mathcal{C}^{∞} Riemannian manifold with Riemannian measure m, $(\Omega, \mathcal{F}, \mathbb{P})$ to be a Lebesgue space, and for some $r \geq 1$ and each $\epsilon \in (-1, 1)$ let $\mathcal{T}_{\epsilon} : \Omega \to \mathcal{C}^{r+1}(M, M)$ denote a one-parameter family of random maps with a 'measurable' dependence on ω . After fixing an invertible, \mathbb{P} -ergodic map $\sigma : \Omega \to \Omega$, from each \mathcal{T}_{ϵ} we obtain random dynamical systems $(\mathcal{T}_{\epsilon}, \sigma)$ whose trajectories are random variables of the form

$$x, \mathcal{T}_{\epsilon,\omega}(x), \mathcal{T}_{\epsilon,\omega}^{(2)}(x), \dots, \mathcal{T}_{\epsilon,\omega}^{(n)}(x), \dots,$$

where $\mathcal{T}_{\epsilon,\omega}^{(n)}$ is short for the composition $\mathcal{T}_{\epsilon,\sigma^{n-1}\omega} \circ \cdots \circ \mathcal{T}_{\epsilon,\omega}$. A family of probability measures $\{\mu_{\epsilon,\omega}\}_{\omega\in\Omega}$ on M is said to be *equivariant* for $(\mathcal{T}_{\epsilon},\sigma)$ if $\mu_{\epsilon,\omega} \circ \mathcal{T}_{\epsilon,\omega}^{-1} = \mu_{\epsilon,\sigma\omega}$ for \mathbb{P} -a.e. ω (see Subsection 2.1 for a precise definition). When \mathcal{T}_{ϵ} possesses some (partial) hyperbolicity and good mixing properties one hopes to find a unique *physical* equivariant family of probability measures¹, as such objects describe the *m*-a.e. realized statistical behaviour of the given RDS. Quenched response theory is, broadly speaking, concerned with questions of the regularity of the map $\epsilon \mapsto \{\mu_{\epsilon,\omega}\}_{\omega\in\Omega}$ and, in particular, how this regularity is inherited from that of $\epsilon \mapsto \mathcal{T}_{\epsilon}$. The one-parameter family of random maps $\epsilon \mapsto \mathcal{T}_{\epsilon}$ is said to exhibit quenched linear response if

 $^{^{*}}$ School of Mathematics and Statistics, University of New South Wales, Sydney NSW 2052, Australia. Email: harry.crimmins@unsw.edu.au

[†]Department of Mathematics, Tokai University, 4-1-1 Kitakaname, Hiratuka, Kanagawa, 259-1292, Japan. Email: yushi.nakano@tsc.u-tokai.ac.jp

¹A \mathcal{T} -equivariant family of measure $\{\mu_{\omega}\}_{\omega\in\Omega}$ is physical with respect to m if $n^{-1}\sum_{i=0}^{n-1} \delta_{\mathcal{T}_{\sigma^{-i}\omega^{i}}(x)} \to \mu$ for x in a (possibly ω -dependent) positive m-measure set with \mathbb{P} -probability 1.

the measures $\{\mu_{\epsilon,\omega}\}_{\omega\in\Omega}$ vary differentiability with ϵ in an appropriate topology, with quenched higher-order (e.g. quadratic) response being defined analogously.

Linear and higher-order response theory for deterministic (i.e. non-random) systems is an established area of research, and there are a plethora of methods available for treating various systems (see [6] for a good review). Response theory has been developed for expanding maps in one and many dimensions [6, 7, 46], intermittent systems [10, 2, 37], Anosov diffeomorphisms [44, 45, 32], partially hyperbolic systems [18], and piecewise expanding interval maps [5, 9]. The tools and techniques one may apply to deduce response results are likewise numerous: there are arguments based on structural stability [44], standard pairs [18], the implicit function theorem [46], and on the spectral perturbation theory of Gouëzel, Keller and Liverani [36, 32] (and variants thereof, e.g. [26]).

On the other hand, the literature on quenched response theory for random dynamical systems is relatively small, and has only recently become an active research topic. With a few notable exceptions, most results for random systems have focussed on the continuity of the equivariant random measure [8, 3, 24, 28, 40], although some more generally apply to the continuity of the Oseledets splitting and Lyapunov exponents associated to the RDS's Perron-Frobenius operator cocycle [11, 16]. Quenched linear and higher-order response results are, to the best of our knowledge, limited to [47], where quenched linear and higher-order response is proven for general RDSs of \mathcal{C}^k uniformly expanding maps, and to [20], wherein quenched linear response is proven for RDSs of Anosov maps nearby a fixed Anosov map. The relatively fewer results for response theory in the random case has been largely attributed to the difficulty in finding appropriate generalisations of the tools, techniques and constructions that have succeeded in the deterministic case. While the authors believe this sentiment is generally well-founded, in this paper we find that for quenched linear and higher-order response problems it is possible to directly generalise an approach from the deterministic case to the random case with surprisingly little trouble. In particular, by building on [40] we will show that the application of Gouëzel–Keller–Liverani (GKL) spectral perturbation theory to response problems can be 'lifted' to the random case, allowing one to deduce corresponding quenched response from deterministic response 'for free'.

In the deterministic setting the application of GKL perturbation theory to response problems is part of the more general 'functional analytic' approach to studying dynamical systems, which recasts the investigation of invariant measures and statistical properties of dynamical systems in functional analytic and operator theoretic terms. The hero of this approach is the Perron-Frobenius operator, which for a non-singular² map $T \in C^{r+1}(M, M)$ is denoted by \mathcal{L}_T and defined for $f \in L^1(m)$ by

$$(\mathcal{L}_T f)(x) = \sum_{T(y)=x} \frac{f(y)}{|\det D_y T|}.$$

The key observation is that the statistical properties of T are often encoded in the spectral data of \mathcal{L}_T provided that one consider the operator on an appropriate Banach space [4, 7, 39, 25]. Specifically, one desires a Banach space for which \mathcal{L}_T is bounded and quasi-compact (in addition to some other benign conditions), since in these cases a unique physical invariant measure μ_T for T is often obtained as a fixed point of \mathcal{L}_T . One may then attempt to answer response theory questions by studying the regularity of the map $T \mapsto \mathcal{L}_T$ with a view towards deducing the regularity of $T \mapsto \mu_T$ via some spectral argument. The main obstruction to carrying out such a strategy is that $T \mapsto \mathcal{L}_T$ is usually not continuous in the relevant operator norm, and so standard spectral perturbation theory (e.g. Kato [35]) cannot be applied. Instead, however, one often has that $T \mapsto \mathcal{L}_T$ is continuous (or \mathcal{C}^k) in some weaker topology, and by applying GKL spectral perturbation theory it is then possible to deduce regularity results for $T \mapsto \mu_T$.

²A map $T: M \to M$ is non-singular with respect to m if m(A) = 0 implies that $m(T^{-1}(A)) = 0$.

The main contribution of this paper is to show that the strategy detailed in the previous paragraph may still be applied in the random case to deduce quenched linear and higher-order response results. More precisely, with $\{(\mathcal{T}_{\epsilon}, \sigma)\}_{\epsilon \in (-1,1)}$ denoting the RDSs from earlier, the main (psuedo) Theorem of this paper is the following (see Theorem 3.6 for a precise statement and Section 4 for our application to RDSs):

Theorem A

Suppose that (\mathcal{T}_0, σ) exhibits ω -uniform exponential mixing on M, and that for \mathbb{P} -a.e. ω the hypotheses of GKL perturbation theory are 'uniformly' satisfied for the one-parameter families $\epsilon \mapsto \mathcal{T}_{\epsilon,\omega}$ as in the deterministic case. Then whatever linear and higher-order response results that hold \mathbb{P} -a.e. at $\epsilon = 0$ for the physical invariant probability measures of the one-parameter families $\epsilon \mapsto \mathcal{T}_{\epsilon,\omega}$ also hold in the quenched sense for the equivariant physical probability measures of the one-parameter family $\epsilon \mapsto \{(\mathcal{T}_{\epsilon}, \sigma)\}_{\epsilon \in (-1,1)}$ of RDSs.

We note that despite the mixing requirement placed on (\mathcal{T}_0, σ) in Theorem A we do not require that σ exhibit any mixing behaviour, other than being ergodic. The general strategy behind the proof of Theorem A is to consider for each $\epsilon \in (-1, 1)$ a 'lifted' operator obtained from the Perron-Frobenius operators $\{\mathcal{L}_{\mathcal{T}_{\epsilon,\omega}}\}_{\omega\in\Omega}$ associated to $\{\mathcal{T}_{\epsilon,\omega}\}_{\omega\in\Omega}$. Then, using the fact that the hypotheses of the Gouëzel-Keller-Liverani Theorem (Theorem 2.1) are satisfied 'uniformly' for the Perron-Frobenius operators $\epsilon \mapsto \mathcal{L}_{\mathcal{T}_{\epsilon,\omega}}$ and ω in some \mathbb{P} -full set, we deduce that the Gouëzel-Keller-Liverani Theorem may be applied to the lifted operator. By construction the fixed points of these lifted operators are exactly the equivariant physical probability measures of the corresponding RDS, and so we obtain the claimed linear and higher-order response via the conclusion of the Gouëzel-Keller-Liverani Theorem. Using Theorem A we easily obtain new quenched linear and higher-order response results for random Anosov maps (Theorem 4.8) and for random U(1) extensions of expanding maps (Theorem 4.10). We note that our examples consist of random maps that are uniformly close to a fixed system. However, this is not a strict requirement for the application of our theory and one could also consider 'non-local' examples e.g. it is clear that the arguments in Section 4 are applicable to random systems consisting of arbitrary \mathcal{C}^k expanding maps.

The structure of the paper is as follows. In Section 2 we introduce some conventions that are used throughout the paper and review some preliminary material related to random dynamical systems and the Gouëzel–Keller–Liverani Theorem. In Section 3 we consider random operator cocycles and their 'lifts', and then prove our main abstract result, Theorem 3.6, which is a version of the Gouëzel–Keller–Liverani Theorem for the 'lifts' of certain operator cocycles. In Section 4 we discuss how Theorem 3.6 may be applied to study the quenched linear and higherorder response of general random C^{r+1} dynamical systems, and then consider in detail the cases of random Anosov maps and random U(1) extensions of expanding maps. Lastly, Appendix A contains the proof of a technical lemma from Section 4.

2 Preliminaries

We adopt the following notational conventions:

- 1. The symbol 'C' will, unless otherwise stated, be used to indiscriminately refer to many constants, which are uniform (or almost surely uniform) and whose value may change between usages. If we wish to emphasise that C depends on parameters a_1, \ldots, a_n we may write C_{a_1,\ldots,a_n} instead.
- 2. If X and Y are topological vector spaces such that X is continuously included into Y then we will write $X \hookrightarrow Y$.

- 3. If X and Y are Banach spaces then we denote the set of bounded, linear operators from X to Y by L(X, Y). When X = Y, it is simply written as L(X).
- 4. When X is a metric space we denote the Borel σ -algebra on X by \mathcal{B}_X .
- 5. If $A \in L(X)$ then we denote the spectrum of A by $\sigma(A)$ and the spectral radius by $\rho(A)$. We will frequently consider operators acting on a number of spaces simultaneously, and in such a situation we may denote $\sigma(A)$ and $\rho(A)$ by $\sigma(A|X)$ and $\rho(A|X)$, respectively, for clarity.

2.1 Random dynamical systems

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\sigma : \Omega \to \Omega$ a measurably invertible, measure-preserving map. For a measurable space (Σ, \mathcal{G}) , we say that a measurable map $\Phi : \mathbb{N}_0 \times \Omega \times \Sigma \to \Sigma$ is a random dynamical system (or, for short, an *RDS*) on Σ over the driving system σ if

$$\varphi_{\omega}^{(0)} = \mathrm{id}_{\Sigma} \quad \mathrm{and} \quad \varphi_{\omega}^{(n+m)} = \varphi_{\sigma^m\omega}^{(n)} \circ \varphi_{\omega}^{(m)}$$

for each $n, m \in \mathbb{N}_0$ and $\omega \in \Omega$, with the notation $\varphi_{\omega}^{(n)} = \Phi(n, \omega, \cdot)$ and $\sigma \omega = \sigma(\omega)$, where $\mathbb{N}_0 = \{0\} \cup \mathbb{N}$. A standard reference for random dynamical systems is the monograph by Arnold [1]. It is easy to check that

$$\varphi_{\omega}^{(n)} = \varphi_{\sigma^{n-1}\omega} \circ \varphi_{\sigma^{n-2}\omega} \circ \dots \circ \varphi_{\omega} \tag{1}$$

with the notation $\varphi_{\omega} = \Phi(1, \omega, \cdot)$. Conversely, for each measurable map $\varphi : \Omega \times \Sigma \to \Sigma$: $(\omega, x) \mapsto \varphi_{\omega}(x)$, the measurable map $(n, \omega, x) \mapsto \varphi_{\omega}^{(n)}(x)$ given by (1) is an RDS. We call it an *RDS induced by* φ over σ , and simply denote it by (φ, σ) .

It is easy to see that if we define a skew-product map $\Theta : \Omega \times \Sigma \to \Omega \times \Sigma$ by $\Theta(\omega, x) = (\sigma \omega, \varphi_{\omega}(x))$ for each $(\omega, x) \in \Omega \times \Sigma$, then

$$\Theta^n(\omega, x) = (\sigma^n \omega, \varphi^{(n)}_{\omega}(x)) \quad \text{for all } n \in \mathbb{N}_0.$$

Furthermore, a probability measure μ on $(\Omega \times \Sigma, \mathcal{F} \times \mathcal{G})$ is invariant for Θ (i.e. $\mu \circ \Theta^{-1} = \mu$) and $\mu \circ \pi_{\Omega}^{-1} = \mathbb{P}$, where $\pi_{\Omega}(\omega, x) = \omega$ for each $(\omega, x) \in \Omega \times \Sigma$, if and only if there is a measurable family of probability measure $\{\mu_{\omega}\}_{\omega\in\Omega}$ (i.e. for each $A \in \mathcal{G}$, the map $\omega \mapsto \mu_{\omega}(A)$ is $(\mathcal{F}, \mathcal{B}_{\mathbb{R}})$ -measurable) such that $\mu(A) = \int_{\Omega} \int_{\Sigma} 1_A(\omega, x) \mu_{\omega}(\mathrm{d}x) \mathbb{P}(\mathrm{d}\omega)$ for each $A \in \mathcal{F} \times \mathcal{G}$, and that

$$\mu_{\omega} \circ \varphi_{\omega}^{-1} = \mu_{\sigma\omega} \quad \text{for almost every } \omega \in \Omega \tag{2}$$

(cf. [1, Subsection 1.4]). Hence, we say that a measurable family of probability measure $\{\mu_{\omega}\}_{\omega\in\Omega}$ is equivariant for (φ, σ) if it satisfies (2).

2.2 The Gouëzel–Keller–Liverani Theorem

We recall the statement of the Gouëzel-Keller-Liverani theorem from [7] (although we note that the result first appeared in full generality in [32, 31], and in less generality in [36]). Fix an integer $N \ge 1$ and let E_j , $j \in \{0, ..., N\}$, be Banach spaces with $E_j \hookrightarrow E_{j-1}$ for each $j \in \{1, ..., N\}$. For a family of linear operators $\{A_{\epsilon}\}_{\epsilon \in [-1,1]}$ on these spaces we consider the following conditions:

(GKL1) For all $i \in \{1, \ldots, N\}$ and $|\epsilon| \leq 1$ we have

$$\|A_{\epsilon}\|_{L(E_i)} \le C.$$

- (GKL2) There exists M > 0 such that $||A_{\epsilon}^{n}||_{L(E_{0})} \leq CM^{n}$ for all $|\epsilon| \leq 1$ and $n \in \mathbb{N}$.
- (GKL3) There exists $\alpha < M$ such that for every $|\epsilon| \leq 1, f \in E_1$ and $n \in \mathbb{N}$ we have

$$\|A_{\epsilon}^{n}f\|_{E_{1}} \leq C\alpha^{n} \|f\|_{E_{1}} + CM^{n} \|f\|_{E_{0}}.$$

(GKL4) For every $|\epsilon| \leq 1$ we have

$$\left\|A_{\epsilon} - A_{0}\right\|_{L(E_{N}, E_{N-1})} \le C \left|\epsilon\right|.$$

If $N \ge 2$ we have the following additional requirement:

(GKL5) There exist operators Q_1, \ldots, Q_{N-1} such that for all $j \in \{1, \ldots, N-1\}$ and $i \in \{j, \ldots, N\}$ we have

$$\|Q_j\|_{L(E_i, E_{i-j})} \le C,\tag{3}$$

and that for all $|\epsilon| \leq 1$ and $j \in \{2, \ldots, N\}$ we have

$$\left\| A_{\epsilon} - A_0 - \sum_{k=1}^{j-1} \epsilon^k Q_k \right\|_{L(E_N, E_{N-j})} \le C \left|\epsilon\right|^j.$$

$$\tag{4}$$

Theorem 2.1 (The Gouëzel–Keller–Liverani Theorem, [7, Theorem A.4]). Fix an integer $N \ge 1$ and let E_j , $j \in \{0, \ldots, N\}$, be Banach spaces with $E_j \hookrightarrow E_{j-1}$ for each $j \in \{1, \ldots, N\}$. Suppose that $\{A_{\epsilon}\}_{\epsilon \in [-1,1]}$ satisfies (GKL1)-(GKL4) and if $N \ge 2$ then also (GKL5). For $z \notin \sigma(A_0|E_N)$ set $R_0(z) = (z - A_0)^{-1}$ and define

$$S_{\epsilon}^{(N)}(z) = R_0(z) + \sum_{k=1}^{N-1} \epsilon^k \sum_{\substack{j=1\\l_i \ge 1}}^k \sum_{\substack{l_1 + \dots + l_j = k\\l_i \ge 1}}^k R_0(z) Q_{l_1} R_0(z) \cdots R_0(z) Q_{l_j} R_0(z).$$
(5)

In addition, for any $a > \alpha$ let

$$\eta = \frac{\log(a/\alpha)}{\log(M/\alpha)},$$

and for $\delta > 0$ set

 $\mathcal{V}_{\delta,a}(A_0) = \left\{ z \in \mathbb{C} : |z| \ge a \text{ and } \operatorname{dist}\left(z, \sigma\left(A_0|E_j\right)\right) \ge \delta, \quad \forall j \in \{1, \dots, N\} \right\}.$

There exist $\epsilon_0 > 0$ so that $\mathcal{V}_{\delta,a}(A_0) \cap \sigma(A_{\epsilon}|E_1) = \emptyset$ for every $|\epsilon| \leq \epsilon_0$ and so that, for each $z \in \mathcal{V}_{\delta,a}(A_0)$, we have

$$\|(z - A_{\epsilon})^{-1}\|_{L(E_1)} \le C,$$

and

$$\left\| (z - A_{\epsilon})^{-1} - S_{\epsilon}^{(N)}(z) \right\|_{L(E_N, E_0)} \le C \left| \epsilon \right|^{N-1+\eta}$$

Remark 2.2. While the Gouëzel–Keller–Liverani Theorem as stated in Theorem 2.1 is true, there is an error in the proof of the result in both [30] and [7]. For details of the error we refer the reader to [31], and to the proof of [30, Theorem 3.3] for a corrected argument.

Remark 2.3. We emphasise that the inclusion $E_j \subset E_{j-1}$ need not be compact in Theorem 2.1, which will be important in our application in Section 4.

3 A spectral approach to stability theory for operator cocycles

Let X be a Banach space, and $\mathcal{S}_{L(X)}$ denote the σ -algebra generated by the strong operator topology on L(X). If $A : \Omega \to L(X)$ is $(\mathcal{F}, \mathcal{S}_{L(X)})$ -measurable then we say it is *strongly measurable*. For an overview of the properties of strong measurable maps we refer the reader to [27, Appendix A]. The following lemma records the main properties of strongly measurable maps that we shall use.

Lemma 3.1 ([27, Lemmas A.5 and A.6]). Suppose that X is a separable Banach space and that $(\Omega, \mathcal{F}, \mathbb{P})$ is a Lebesgue space. Then

- 1. The set of strongly measurable maps is closed under (operator) composition i.e. if A_i : $\Omega \to L(X), i \in \{1, 2\}$, are strongly measurable then so to is $A_2A_1 : \Omega \to L(X)$.
- 2. If $A : \Omega \to L(X)$ is strongly measurable and $f : \Omega \to X$ is $(\mathcal{F}, \mathcal{B}_X)$ -measurable then $\omega \mapsto A_{\omega} f_{\omega}$ is $(\mathcal{F}, \mathcal{B}_X)$ -measurable too.
- 3. If $A : \Omega \to L(X)$ is such that $\omega \mapsto A(\omega)f$ is $(\mathcal{F}, \mathcal{B}_X)$ -measurable for every $f \in X$ then A is strongly measurable.

As a slight abuse of notation, for a given strongly measurable map $A : \Omega \to L(X)$, we denote an $(\mathcal{F} \times \mathcal{B}_X, \mathcal{B}_X)$ -measurable map $(\omega, f) \mapsto A(\omega)f$ by A. In light of the previous lemma, we may now formally define the main objects of study for this section.

Definition 3.2. An RDS (A, σ) on X induced by A is called an operator cocycle (or a linear RDS) if $(\Omega, \mathcal{F}, \mathbb{P})$ is a Lebesgue space, $\sigma : \Omega \to \Omega$ is an invertible, ergodic, \mathbb{P} -preserving map, X is a separable Banach space and $A : \Omega \mapsto L(X)$ is strongly measurable. We say that (A, σ) is bounded if $A \in L^{\infty}(\Omega, L(X))$.

Throughout the rest of this paper, we assume that $(\Omega, \mathcal{F}, \mathbb{P})$ is a Lebesgue space and σ : $\Omega \to \Omega$ is an invertible, ergodic, \mathbb{P} -preserving map. An operator cocycle (A, σ) is explicitly written as a measurable map

$$\mathbb{N}_0 \times \Omega \times X \to X : (n, \omega, f) \mapsto A^{(n)}(\omega)f, \quad A^{(n)}(\omega) := A(\sigma^{n-1}\omega) \circ \cdots \circ A(\omega).$$

We denote by X^* the dual space of X.

Definition 3.3. Let $\xi \in X^*$ be non-zero. We say that $A \in L(X)$ is ξ -Markov if $\xi(Af) = \xi(f)$ for every $f \in X$. We say that an operator cocycle (A, σ) is ξ -Markov if A is almost surely ξ -Markov.

Notice that our terminology in Definition 3.3 may be slightly non-standard: in the literature a linear operator $A: X \to X$ is called Markov if $X = L^1(S, \mu)$ for a probability space (S, μ) and A is positive (i.e. $Af \ge 0$ μ -almost everywhere if $f \ge 0$ μ -almost everywhere) and ξ -Markov with $\xi(f) = \int_S f d\mu$ (cf. [38]). See also Definition 4.3 for more general definition of positivity. We do not add the positivity condition to Definition 3.3 to make clear that the result in this section holds without it.

Definition 3.4. Suppose that (A, σ) is a ξ -Markov operator cocycle for some non-zero $\xi \in X^*$. We say that (A, σ) is ξ -mixing with rate $\rho \in [0, 1)$ if for every $n \in \mathbb{N}$ we have

$$\operatorname{ess\,sup}_{\omega\in\Omega} \left\| A^{(n)}(\omega) \right\|_{\ker\xi} \le C\rho^n.$$
(6)

We say that (A, σ) is ξ -mixing if it is ξ -mixing with some rate $\rho \in [0, 1)$.

Fix a non-zero $\xi \in X^*$. We define $\mathcal{X} \equiv \mathcal{X}_{\xi}$ as

$$\mathcal{X} = \{ f \in L^{\infty}(\Omega, X) : \xi(f) \text{ is almost surely constant} \}.$$

Since \mathcal{X} is a closed subspace of $L^{\infty}(\Omega, X)$, it is a Banach space with the usual norm. If (A, σ) is a bounded ξ -Markov operator cocycle then we define $\mathbb{A} : \mathcal{X} \to \mathcal{X}$ by

$$(\mathbb{A}f)(\omega) = A(\sigma^{-1}(\omega))f(\sigma^{-1}(\omega)). \tag{7}$$

We say that \mathbb{A} is the *lift* of (A, σ) . That $\mathbb{A} \in L(\mathcal{X})$ follows from Lemma 3.1 and the boundedness of (A, σ) (see [40] for a possible extension of the lift to the case when σ is not invertible). The following proposition is a natural generalisation of [40, Proposition 2.3].

Proposition 3.5. Fix non-zero $\xi \in X^*$. If (A, σ) is a bounded, ξ -Markov, ξ -mixing operator cocycle with rate $\rho \in [0, 1)$ then 1 is a simple eigenvalue of \mathbb{A} and $\sigma(\mathbb{A}|\mathcal{X}) \setminus \{1\} \subseteq \{z \in \mathbb{C} : |z| \leq \rho\}$.

Proof. For each $c \in \mathbb{C}$, let

$$\mathcal{X}_c = \{ f \in \mathcal{X} : \xi(f) = c \text{ almost surely} \}.$$
(8)

We note that \mathcal{X}_c is non-empty since ξ is assumed to be non-zero. Since (A, σ) is a ξ -Markov operator cocycle, the lift \mathbb{A} preserves \mathcal{X}_c . For any $f, g \in \mathcal{X}_c$ one has $f - g \in \mathcal{X}_0$ (i.e. $f - g \in \ker \xi$ almost surely), and so as (A, σ) is ξ -mixing with rate ρ we have for every $n \in \mathbb{N}$ and almost every $\omega \in \Omega$ that

$$\|(\mathbb{A}^{n}f)(\omega) - (\mathbb{A}^{n}g)(\omega)\| = \|A^{(n)}(\sigma^{-n}(\omega))(f_{\sigma^{-n}(\omega)} - g_{\sigma^{-n}(\omega)})\|$$

$$\leq C\rho^{n} \|f_{\sigma^{-n}(\omega)} - g_{\sigma^{-n}(\omega)}\|.$$
(9)

Upon taking the essential supremum we see that \mathbb{A}^n is a contraction mapping on \mathcal{X}_c for large enough n. Since \mathcal{X}_c is complete, it follows that \mathbb{A} has a unique fixed point v_c in \mathcal{X}_c . Obviously $v_c = cv_1$, and thus 1 is an eigenvalue of \mathbb{A} on \mathcal{X} . Furthermore, $\mathcal{X} = \operatorname{span}\{v_1\} \oplus \mathcal{X}_0$ (indeed, for every $f \in \mathcal{X}$ we can write $f = f_1 + f_0$ where $f_1 = \xi(f)v_1 \in \operatorname{span}\{v_1\}$ and $f_0 = f - f_1 \in \mathcal{X}_0$, and note that $\operatorname{span}\{v_1\}$ and \mathcal{X}_0 are closed subspaces).

Since A preserves both span $\{v_1\}$ and \mathcal{X}_0 , we have

$$\sigma(\mathbb{A}|\mathcal{X}) = \sigma(\mathbb{A}|\operatorname{span}\{v_1\}) \sqcup \sigma(\mathbb{A}|\mathcal{X}_0).$$

It is clear that $\sigma(\mathbb{A}|\operatorname{span}\{v_1\})$ only consists of a simple eigenvalue 1, while $\rho(\mathbb{A}|\mathcal{X}_0) \leq \rho$ since (A, σ) is ξ -mixing with rate ρ . Thus $\sigma(\mathbb{A}|\mathcal{X}) \setminus \{1\} = \sigma(\mathbb{A}|\mathcal{X}_0) \subseteq \{z \in \mathbb{C} : |z| \leq \rho\}$.

3.1 Main result

Given a bounded, ξ -Markov, ξ -mixing operator cocycle (A, σ) we are interested in question of stability (and differentiability) of the ξ -normalised fixed point v of \mathbb{A} . To this end we formulate a number of conditions on operator cocycles that are reminiscent of the conditions of the Gouëzel-Keller-Liverani Theorem.

Fix an integer $N \ge 1$ and let $E_j, j \in \{0, \ldots, N\}$, be Banach spaces. Let $\{(A_{\epsilon}, \sigma)\}_{\epsilon \in [-1,1]}$ be a family of operator cocycles on these spaces.

(QR0) $E_j \hookrightarrow E_{j-1}$ for each $j \in \{1, \ldots, N\}$, E_N is separable and $\|\cdot\|_{E_j}$ -dense in E_j for each $j \in \{0, \ldots, N\}$ (in particular, E_1 is separable). There exists non-zero $\xi \in E_0^*$ such that (A_{ϵ}, σ) is ξ -Markov on E_j for each $|\epsilon| \leq 1, j \in \{0, \ldots, N\}$ and that (A_0, σ) is ξ -mixing on E_j for each $j \in \{1, N\}$.

- (QR1) For all $i \in \{1, \ldots, N\}$ and $|\epsilon| \leq 1$ we have $\operatorname{ess\,sup}_{\omega} ||A_{\epsilon}(\omega)||_{L(E_i)} \leq C$.
- (QR2) There exists M > 0 such that $\operatorname{ess\,sup}_{\omega} \left\| A_{\epsilon}^{(n)}(\omega) \right\|_{L(E_0)} \leq CM^n$ for all $|\epsilon| \leq 1$ and $n \in \mathbb{N}$.
- (QR3) There exists $\alpha < M$ such that for every $f \in E_1$, $|\epsilon| \leq 1$ and $n \in \mathbb{N}$ we have

$$\operatorname{ess\,sup}_{\omega} \left\| A_{\epsilon}^{(n)}(\omega) f \right\|_{E_{1}} \le C \alpha^{n} \left\| f \right\|_{E_{1}} + C M^{n} \left\| f \right\|_{E_{0}}.$$

(QR4) For every $|\epsilon| \leq 1$ we have

$$\operatorname{ess\,sup}_{\omega} \|A_{\epsilon}(\omega) - A_{0}(\omega)\|_{L(E_{N}, E_{N-1})} \leq C |\epsilon|$$

If $N \ge 2$ we have the following additional requirement:

(QR5) There exist operators $Q_1(\omega), \ldots, Q_{N-1}(\omega)$ for each ω such that for all $j \in \{1, \ldots, N-1\}$ and $i \in \{j, \ldots, N\}$ we have

$$\operatorname{ess\,sup} \|Q_j(\omega)\|_{L(E_i, E_{i-j})} \le C,$$

and such that for all $|\epsilon| \leq 1$ and $j \in \{2, \ldots, N\}$ we have

$$\operatorname{ess\,sup}_{\omega} \left\| A_{\epsilon}(\omega) - A_{0}(\omega) - \sum_{k=1}^{j-1} \epsilon^{k} Q_{k}(\omega) \right\|_{L(E_{N}, E_{N-j})} \leq C \left|\epsilon\right|^{j}$$

We need not assume that Q_1, \ldots, Q_{N-1} are measurable³, which will make our application in Section 4 simpler.

Our main theorem for this section is the following.

Theorem 3.6. Fix an integer $N \geq 1$, and let E_j , $j \in \{0, \ldots, N\}$, be Banach spaces and $\{(A_{\epsilon}, \sigma)\}_{\epsilon \in [-1,1]}$ a family of operator cocycles on these spaces. Suppose that $\{(A_{\epsilon}, \sigma)\}_{\epsilon \in [-1,1]}$ satisfies (QR0)-(QR4) and if $N \geq 2$ then also (QR5). Then there exists $\epsilon_0 \in (0, 1]$ such that one can find a unique $v_{\epsilon} \in L^{\infty}(\Omega, E_1)$ such that $A_{\epsilon}(\omega)v_{\epsilon}(\omega) = v_{\epsilon}(\sigma\omega)$ and $\xi(v_{\epsilon}(\omega)) = 1$ for each $\epsilon \in (-\epsilon_0, \epsilon_0)$ and almost every $\omega \in \Omega$, that

$$\sup_{\epsilon|<\epsilon_0} \operatorname{ess\,sup}_{\omega} \|v_{\epsilon}(\omega)\|_{E_1} < \infty,$$

and that (A_{ϵ}, σ) is ξ -mixing whenever $|\epsilon| < \epsilon_0$. Moreover, there exists $\{v_0^{(k)}\}_{k=1}^{N-1} \subset L^{\infty}(\Omega, E_0)$ such that $\xi(v_0^{(k)}) = 0$ almost surely for each k, and that for every $\eta \in (0, \log(1/\alpha)/\log(M/\alpha))$, we have

$$v_{\epsilon} = v_0 + \sum_{k=1}^{N-1} \epsilon^k v_0^{(k)} + O_{\eta}(\epsilon^{N-1+\eta}), \qquad (10)$$

where $O_{\eta}(\epsilon^{N-1+\eta})$ is to be understood as an essentially bounded term in E_0 that possibly depends on η .

Remark 3.7. One is free to take $E_0 = E_1 = \cdots = E_N$ in Theorem 3.6, in which case the conditions (QR0)-(QR3) collapse into a single bound and (QR4)-(QR5) become standard operator norm inequalities. Hence in this simple case one recovers an expected Banach space perturbation result.

³Recall that the essential supremum of a (not necessarily measurable) complex-valued function f on Ω is given as the infimum of $\sup_{\omega \in \Omega_0} |f(\omega)|$ over all \mathbb{P} -full measure sets Ω_0 .

Remark 3.8. We note that Theorem 3.6 has been proven before for the cases where N = 1 and N = 2 in [24] and [20], respectively.

Remark 3.9. The claim that 'there exists $\epsilon_0 \in (0, 1]$ such that (A_{ϵ}, σ) is ξ -mixing whenever $|\epsilon| < \epsilon_0$ ' is exactly the content of [20, Proposition 1] (as well as being an easy corollary of [16, Proposition 3.11]). Upon examining these proofs it is clear that something slightly stronger is true: in the setting of Theorem 3.6, for every $\kappa \in (\rho, 1)$ there exists $\epsilon_{\kappa} > 0$ such that for all $\epsilon \in (-\epsilon_{\kappa}, \epsilon_{\kappa})$ one has

$$\sup_{n \in \mathbb{N}} \kappa^{-n} \operatorname{ess\,sup}_{\omega} \left\| A^{n}_{\epsilon} \right\|_{\ker \xi} \right\|_{L(E_{1})} \leq C.$$
(11)

3.2 The proof of Theorem 3.6

Before detailing the proof of Theorem 3.6 we introduce some basic constructs. For each $j \in \{0, \ldots, N\}$ let

 $\mathcal{E}_j = \{ f \in L^{\infty}(\Omega, E_j) : \xi(f) \text{ is almost surely constant} \}.$

Since $\xi \in E_j^*$ for each $j \in \{0, \ldots, N\}$ we observe that each \mathcal{E}_j is a closed subspace of $L^{\infty}(\Omega, E_j)$, and consequently a Banach space. Moreover, we have $\mathcal{E}_j \hookrightarrow \mathcal{E}_{j-1}$ for $j \in \{1, \ldots, N\}$. For each $j \in \{1, \ldots, N\}$ we may consider the lift $\mathbb{A}_{\epsilon,j}$ of the operator cocycle (A_{ϵ}, σ) on E_j , although we will always omit the subscript j and just write \mathbb{A}_{ϵ} , which will be of no consequence.

The beginning of the proof of Theorem 3.6 is straightforward. First we note that (QR1) implies that (A_0, σ) is bounded on E_j for $j \in \{1, N\}$ and so Proposition 3.5 may be applied to characterise the spectrum of \mathbb{A}_0 on \mathcal{E}_1 and \mathcal{E}_N . Let ρ be the rate of ξ -mixing in (QR0), that is, (A_0, σ) is ξ -mixing on E_j with rate ρ for each $j \in \{1, N\}$. Then it follows from (QR0) and Proposition 3.5 that 1 is a simple eigenvalue of \mathbb{A}_0 when considered on either space, and we have

$$\sigma(\mathbb{A}_0|\mathcal{E}_j) \setminus \{1\} \subseteq \{z \in \mathbb{C} : |z| \le \rho\}$$
(12)

for $j \in \{1, N\}$. For each $j \in \{1, ..., N\}$ one may use basic functional analysis and the fact that $\mathcal{E}_N \hookrightarrow \mathcal{E}_j \hookrightarrow \mathcal{E}_1$ to deduce that 1 is a simple eigenvalue of $\mathbb{A}_0 : \mathcal{E}_j \to \mathcal{E}_j$ and that (12) holds. As a consequence we find a ξ -normalised $v_0 \in \mathcal{E}_N$ that is the unique ξ -normalised fixed point of $\mathbb{A}_0 : \mathcal{E}_j \to \mathcal{E}_j$ for each $j \in \{1, ..., N\}$.

We now turn to constructing the ξ -normalised fixed points of $\mathbb{A}_{\epsilon} : \mathcal{E}_1 \to \mathcal{E}_1$. By Remark 3.9 we may find some $\kappa \in (\rho, 1)$ and $\epsilon_0 > 0$ such that (A_{ϵ}, σ) is ξ -mixing on E_1 with rate κ for every $\epsilon \in (-\epsilon_0, \epsilon_0)$. We note that each (A_{ϵ}, σ) is bounded on E_1 due to (QR1), and so by Proposition 3.5 we find that 1 is a simple eigenvalue of $\mathbb{A}_{\epsilon} : \mathcal{E}_1 \to \mathcal{E}_1$, and that $\sigma(\mathbb{A}_{\epsilon}|\mathcal{E}_1) \setminus \{1\} \subseteq$ $\{z \in \mathbb{C} : |z| \leq \kappa\}$. Thus $\mathbb{A}_{\epsilon} : \mathcal{E}_1 \to \mathcal{E}_1$ has a unique ξ -normalised fixed point $v_{\epsilon} \in \mathcal{E}_1$ for each $\epsilon \in (-\epsilon_0, \epsilon_0)$ by Proposition 3.5. Moreover, by virtue of the uniform bound (11) we may strengthen the conclusion of Proposition 3.5: for all n sufficiently large the family of maps $\{\mathbb{A}^n_{\epsilon}\}_{|\epsilon| < \epsilon_0}$ uniformly contract the set \mathcal{X}_1 from (8). Hence we deduce the bound

$$\sup_{|\epsilon|<\epsilon_0} \operatorname{ess\,sup}_{\omega} \|v_{\epsilon}(\omega)\|_{E_1} < \infty,$$

as required for Theorem 3.6.

Thus to complete the proof of Theorem 3.6 it suffices to prove (10). It may be easily seen from the proof of Proposition 3.5 that the eigenprojection $\Pi_{\epsilon} \in L(\mathcal{E}_1)$ of $\mathbb{A}_{\epsilon} : \mathcal{E}_1 \to \mathcal{E}_1$ onto the eigenspace for 1 is defined for $f \in \mathcal{E}_1$ and $\epsilon \in (-\epsilon_0, \epsilon_0)$ by

$$\Pi_{\epsilon}(f) = \xi(f)v_{\epsilon}.$$

Since each v_{ϵ} is ξ -normalised, we consequently have

$$v_{\epsilon} = v_0 + (\Pi_{\epsilon} - \Pi_0)v_0. \tag{13}$$

If $\delta \in (0, 1 - \kappa)$ then $D_{\delta} = \{z \in \mathbb{C} : |z - 1| = \delta\} \subseteq \mathbb{C} \setminus \sigma(\mathbb{A}_{\epsilon} | \mathcal{E}_1)$ for every $\epsilon \in (-\epsilon_0, \epsilon_0)$. Thus

$$\Pi_{\epsilon} = \int_{D_{\delta}} (z - \mathbb{A}_{\epsilon})^{-1} \,\mathrm{d}z.$$
(14)

Applying (14) to (13) yields

$$v_{\epsilon} = v_0 + \int_{D_{\delta}} \left((z - \mathbb{A}_{\epsilon})^{-1} - (z - \mathbb{A}_0)^{-1} \right) v_0 \, \mathrm{d}z.$$
(15)

The idea is to apply the Gouëzel–Keller–Liverani Theorem to the lifts $\{\mathbb{A}_{\epsilon}\}_{\epsilon \in [-1,1]}$ with Banach spaces $\{\mathcal{E}_j\}_{0 \leq j \leq N}$, and then develop a Taylor expansion in (15). The hypothesis that (QR1)-(QR4) holds for $\{(A_{\epsilon}, \sigma)\}_{\epsilon \in [-1,1]}$ with constants almost surely independent of ω readily implies that the lifts $\{\mathbb{A}_{\epsilon}\}_{\epsilon \in [-1,1]}$ satisfy (GKL1)-(GKL4) for the spaces $\{\mathcal{E}_j\}_{0 \leq j \leq N}$. Hence, in the case where N = 1 we may apply Theorem 2.1 to the lifts $\{\mathbb{A}_{\epsilon}\}_{\epsilon \in [-1,1]}$.

However, the case where $N \geq 2$ is more delicate because the measurability of Q_j is not required in Theorem 3.6. Thus, we introduce the following functional space instead of $L^{\infty}(\Omega, E_j)$, where the objects are defined up to almost everywhere equality but we loosen the measurability requirement. For each $j \in \{0, \ldots, N\}$, let $B(\Omega, E_j)$ denote the set of (not necessarily measurable) bounded E_j -valued functions on Ω equipped with the uniform norm $\|f\|_{B(\Omega, E_j)} = \sup_{\omega \in \Omega} \|f(\omega)\|_{E_j}$ for each $f \in B(\Omega, E_j)$, and let

$$\mathcal{N}_j = \{ f \in B(\Omega, E_j) : f = 0 \text{ almost surely} \}.$$

Then it is straightforward to see that \mathcal{N}_j is a closed subspace of $B(\Omega, E_j)$, and thus we can form a quotient space

$$I^{\infty}(\Omega, E_j) = B(\Omega, E_j) / \mathcal{N}_j.$$

Since $B(\Omega, E_j)$ is a Banach space, $I^{\infty}(\Omega, E_j)$ is also a Banach space with respect to the quotient norm

$$||f||_{I^{\infty}(\Omega,E_j)} = \inf_{h \in \mathcal{N}_j} ||g - h||_{B(\Omega,E_j)}, \quad f \in I^{\infty}(\Omega,E_j),$$

where g is a representative of f. As for $L^{\infty}(\Omega, E_j)$, under the identification of each element of $I^{\infty}(\Omega, E_j)$ with its representative, we have

$$||f||_{I^{\infty}(\Omega, E_j)} = \operatorname{ess\,sup}_{\omega} ||f(\omega)||_{E_j}.$$

Thus under the identification we have $||f||_{L^{\infty}(\Omega, E_j)} = ||f||_{I^{\infty}(\Omega, E_j)}$ for each $f \in L^{\infty}(\Omega, E_j)$, that is, $L^{\infty}(\Omega, E_j)$ isometrically injects into $I^{\infty}(\Omega, E_j)$. Finally let

$$\widetilde{\mathcal{E}}_j = \{ f \in I^\infty(\Omega, E_j) : \xi(f) \text{ is almost surely constant} \}.$$
(16)

We simply write $||f||_{\tilde{\mathcal{E}}_j}$ for $||f||_{I^{\infty}(\Omega, E_j)}$ if $f \in \tilde{\mathcal{E}}_j$. Repeating the previous argument, one can show (GKL1)-(GKL4) for the lifts $\{\mathbb{A}_{\epsilon}\}_{\epsilon \in [-1,1]}$ with respect to the spaces $\{\tilde{\mathcal{E}}_j\}_{0 \le j \le N}$.

We need some auxiliary lemmas. The first lemma is a standard exercise in functional analysis, which will allow us in Lemma 3.11 to characterise the relationship between the spaces $\tilde{\mathcal{E}}_j, j \in \{0, \ldots, N\}$.

Lemma 3.10. Assume the setting of Theorem 3.6. For each $K \in \mathbb{C}$ and $j \in \{0, ..., N\}$ let $U_{K,j} = \{g \in E_j : \xi(g) = K\}$. Then there is a countable subset of $U_{K,j} \cap E_N$ that is $\|\cdot\|_j$ -dense in $U_{K,j}$.

Proof. Fix j. We begin by reducing to the case where K = 0. Since ξ is non-zero on E_0 and E_N is $\|\cdot\|_{E_0}$ -dense in E_0 there exists some $g \in E_N$ such that $\xi(g) \neq 0$. Without loss of generality we may assume that $\xi(g) = 1$, in which case $U_{K,j} = Kg + U_{0,j}$. Thus to obtain a countable, $\|\cdot\|_j$ -dense subset of $U_{K,j}$ inside $U_{K,j} \cap E_N$ it suffices to do so for the case where K = 0 and then translate by Kg. We will now prove the lemma for K = 0. Since E_N is separable and $\xi \in E_N^*$ there is a countable, $\|\cdot\|_{E_N}$ -dense subset G of $U_{0,N} \cap E_N$. We will show that G is $\|\cdot\|_{E_j}$ -dense in $U_{0,j}$ too. Let $u \in U_{0,j}$. As E_N is $\|\cdot\|_{E_j}$ -dense in E_j there exists a sequence $\{u_k\}_{k\in\mathbb{N}} \subseteq E_N$ with $\|\cdot\|_{E_j}$ -limit u. It follows that $\{u_k - \xi(u_k)g\}_{k\in\mathbb{N}}$ is contained in $U_{0,N}$ and satisfies $\lim_{k\to\infty} u_k - \xi(u_k)g = \lim_{k\to\infty} u_k = u$ in E_j since $\xi(u_k) \to 0$. Lastly, G is $\|\cdot\|_{E_j}$ -dense in $U_{0,N}$ and so for each k we may find a $v_k \in G$ so that $\|v_k - (u_k - \xi(u_k)g)\|_{E_N} \to 0$, which implies that $\lim_{k\to\infty} v_k = u$ in E_j . Thus the $\|\cdot\|_{E_j}$ -completion of G is $U_{0,j}$.

Lemma 3.11. Assume the setting of Theorem 3.6. For each $j \in \{1, ..., N\}$ the space $\widetilde{\mathcal{E}}_j$ is equal to the $\|\cdot\|_{\widetilde{\mathcal{E}}_j}$ -completion of $\widetilde{\mathcal{E}}_N$.

Proof. Fix $j \in \{1, \ldots, N\}$ and suppose that $f \in \widetilde{\mathcal{E}}_j$. Recall that $\xi(f)$ is almost surely constant, and let $G \subseteq E_N$ denote a countable, $\|\cdot\|_{E_j}$ -dense subset of $\{g \in E_j : \xi(g) = \xi(f)\}$ as produced by Lemma 3.10. Fix an enumeration $\{g_i\}_{i\in\mathbb{N}}$ of G and for each $i, n \in \mathbb{N}$ set

$$V_{i,n} = \begin{cases} B_{\|\cdot\|_j}(n^{-1}, g_i) & i = 1, \\ B_{\|\cdot\|_j}(n^{-1}, g_i) \setminus \left(\bigcup_{k < i} B_{\|\cdot\|_j}(n^{-1}, g_k)\right) & i > 1. \end{cases}$$

We note that $\{V_{i,n}\}_{i\in\mathbb{N}}$ is a countable partition of E_j for each $n \in \mathbb{N}$, and that each $V_{i,n}$ is measurable in the Borel σ -algebra on E_j . We define a sequence of approximations $\{f_n\}_{n\in\mathbb{N}} \subseteq \widetilde{\mathcal{E}}_N$ by

$$f_n(\omega) = \sum_{i=1}^{\infty} g_i \chi_{f^{-1}(V_{i,n})}(\omega).$$

Notice that $\xi(f_n) = \xi(f)$ almost everywhere and that

$$\operatorname{ess\,sup}_{\omega} \|f_n(\omega) - f(\omega)\|_{E_j} \le n^{-1}.$$

Hence $\lim_{n\to\infty} ||f_n - f||_{\tilde{\mathcal{E}}_j} = 0$, and so f is in the $||\cdot||_{\tilde{\mathcal{E}}_j}$ -completion of $\tilde{\mathcal{E}}_N$. The required claim immediately follows.

With Lemma 3.11 in hand we can deduce (GKL5) for the lifted systems.

Proposition 3.12. Assume the setting of Theorem 3.6 with $N \ge 2$. Then (GKL5) holds with operators \mathbb{Q}_j defined by

$$(\mathbb{Q}_j f)(\omega) = Q_j(\sigma^{-1}\omega)f_{\sigma^{-1}\omega},$$

where $j \in \{1, \dots, N-1\}, i \in \{j, \dots, N\}$ and $f \in \widetilde{\mathcal{E}}_i$.

Proof. It is straightforward to verify the inequalities in (GKL5) from (QR5) if we can show that $\mathbb{Q}_j(\widetilde{\mathcal{E}}_i) \subset \widetilde{\mathcal{E}}_{i-j}$ for each $i \in \{j, \ldots, N\}$ and $j \in \{i, \ldots, N-1\}$ because (A_{ϵ}, σ) is ξ -Markov for every $\epsilon \in [-1, 1]$. Furthermore, for each $f \in \widetilde{\mathcal{E}}_i$ we have

$$\operatorname{ess\,sup}_{\omega} \left\| (\mathbb{Q}_j f)(\omega) \right\|_{E_{i-j}} \le \left(\operatorname{ess\,sup}_{\omega} \left\| Q_j(\omega) \right\|_{L(E_i, E_{i-j})} \right) \operatorname{ess\,sup}_{\omega} \left\| f(\omega) \right\|_{E_i}$$

which is finite due to (QR5). Hence it suffices to show that $\xi(\mathbb{Q}_j f)$ is almost surely constant for each $i \in \{1, \ldots, N\}$, $j \in \{i, \ldots, N-1\}$ and $f \in \widetilde{\mathcal{E}}_i$. By Lemma 3.11 we may find a sequence $\{f_n\}_{n \in \mathbb{N}} \subseteq \widetilde{\mathcal{E}}_N$ such that $||f_n - f||_{\widetilde{\mathcal{E}}_i} \to 0$ as $n \to \infty$. Using (QR5), we have for each $n \in \mathbb{N}$ that

$$\mathbb{Q}_j f_n = \lim_{\epsilon \to 0} \left(\epsilon^{-j} (\mathbb{A}_{\epsilon} - \mathbb{A}_0) f_n - \sum_{k=1}^{j-1} \epsilon^{k-j} \mathbb{Q}_k f_n \right)$$

in E_{N-j-1} almost surely, with the convention $\sum_{k=1}^{0} \epsilon^{k-j} \mathbb{Q}_k f_n = 0$. This implies that

$$\mathbb{Q}_{j}f = \lim_{n \to \infty} \lim_{\epsilon \to 0} \left(\epsilon^{-j} (\mathbb{A}_{\epsilon} - \mathbb{A}_{0}) f_{n} - \sum_{k=1}^{j-1} \epsilon^{k-j} \mathbb{Q}_{k} f_{n} \right)$$
(17)

in E_0 almost surely. Therefore, since (A_{ϵ}, σ) is ξ -Markov for every $\epsilon \in [-1, 1]$ we have almost surely that $\xi(\mathbb{Q}_1 f) = 0$ and

$$\xi(\mathbb{Q}_j f) = -\lim_{n \to \infty} \lim_{\epsilon \to 0} \sum_{k=1}^{j-1} \epsilon^{k-j} \xi(\mathbb{Q}_k f_n).$$

By induction with respect to j we deduce that $\xi(\mathbb{Q}_j f) = 0$ almost surely.

By Proposition 3.12 we have (GKL5) for the lifts $\{\mathbb{A}_{\epsilon}\}_{\epsilon \in [-1,1]}$ with the spaces $\{\widetilde{\mathcal{E}}_j\}_{0 \leq j \leq N}$ whenever $N \geq 2$ in the setting of Theorem 3.6, and so we can apply Theorem 2.1 in this case. As a consequence we may now finish the proof of Theorem 3.6. Let $\eta \in (0, \log(1/\alpha)/\log(M/\alpha))$ and fix $a \in (\alpha, 1)$ so that $\eta = \log(a/\alpha)/\log(M/\alpha)$. Recall δ from (14) and notice that we may take δ to be as small as we like. Henceforth we fix $\delta \in (0, 1 - a)$ and choose some $\delta_0 \in (0, \min\{\delta, 1 - a - \delta\})$. Upon recalling the statement of Theorem 2.1 and our earlier characterisation of $\sigma(\mathbb{A}_0|\widetilde{\mathcal{E}}_j)$ for $j \in \{1, \ldots, N\}$ (recall the remark following (12)), we have

$$D_{\delta} \subseteq \{z \in \mathbb{C} : |z| \ge s \text{ and } |z-1| \ge \delta_0\} \subseteq \mathcal{V}_{\delta_0,a}(\mathbb{A}_0).$$

We now apply Theorem 2.1 to the lifts $\{\mathbb{A}_{\epsilon}\}_{\epsilon \in [-1,1]}$ with Banach spaces $\widetilde{\mathcal{E}}_j$, $j \in \{0, \ldots, N\}$, to deduce the existence of $\epsilon_{\eta} \in (0, \epsilon_0)$ such that for every $\epsilon \in (-\epsilon_{\eta}, \epsilon_{\eta})$ we have $\mathcal{V}_{\delta_0, a}(A_0) \cap \sigma(\mathbb{A}_{\epsilon}|\widetilde{\mathcal{E}}_1) = \emptyset$ and, for each $z \in \mathcal{V}_{\delta_0, a}(\mathbb{A}_0)$, that

$$\left\| (z - \mathbb{A}_{\epsilon})^{-1} - \mathbb{S}_{\epsilon}^{(N)}(z) \right\|_{L(\widetilde{\mathcal{E}}_{N}, \widetilde{\mathcal{E}}_{0})} \leq C \left|\epsilon\right|^{N-1+\eta},\tag{18}$$

where $\mathbb{S}_{\epsilon}^{(N)}(z)$ is defined as in (5). With (18) in hand we may proceed with obtaining (10) via (15). In particular, for $z \in D_{\delta}$ we have

$$\left((z - \mathbb{A}_{\epsilon})^{-1} - (z - \mathbb{A}_{0})^{-1} \right) v_{0} = \sum_{k=1}^{N-1} \epsilon^{k} \sum_{m=1}^{k} \sum_{\substack{l_{1} + \dots + l_{m} = k \\ l_{i} \ge 1}} \left(\prod_{i=1}^{m} (z - \mathbb{A}_{0})^{-1} \mathbb{Q}_{l_{i}} \right) (z - \mathbb{A}_{0})^{-1} v_{0}$$

$$+ \left((z - \mathbb{A}_{\epsilon})^{-1} - \mathbb{S}_{\epsilon}^{(N)} \right) v_{0}.$$

$$(19)$$

For each $k \in \{1, \ldots, N-1\}$ we now define $v_0^{(k)} \in \widetilde{\mathcal{E}}_0$ by

$$v_0^{(k)} = \int_{D_{\delta}} \sum_{m=1}^k \sum_{\substack{l_1 + \dots + l_m = k \\ l_i \ge 1}} \left(\prod_{i=1}^m (z - \mathbb{A}_0)^{-1} \mathbb{Q}_{l_i} \right) (z - \mathbb{A}_0)^{-1} v_0 \, \mathrm{d}z$$

(recall that $v_0 \in \widetilde{\mathcal{E}}_N$). Furthermore, notice that $\xi(v_0^{(k)}) = 0$ almost surely for all k as per the proof of Proposition 3.12. By integrating (19) over D_{δ} and recalling (15) we get

$$v_{\epsilon} = v_0 + \sum_{k=1}^{N-1} \epsilon^k v_0^{(k)} + \int_{D_{\delta}} \left((z - \mathbb{A}_{\epsilon})^{-1} - \mathbb{S}_{\epsilon}^{(N)} \right) v_0 \, \mathrm{d}z \tag{20}$$

in $\widetilde{\mathcal{E}}_0$. Moreover, since $D_{\delta} \subseteq \mathcal{V}_{\delta_0,a}(\mathbb{A}_0)$ it follows from (18) that

$$\left\| \int_{D_{\delta}} \left((z - \mathbb{A}_{\epsilon})^{-1} - \mathbb{S}_{\epsilon}^{(N)} \right) v_0 \, \mathrm{d}z \right\|_{\widetilde{\mathcal{E}}_0} \leq \sup_{z \in \mathcal{V}_{\delta_0, s}(\mathbb{A}_0)} \left\| (z - \mathbb{A}_{\epsilon})^{-1} - \mathbb{S}_{\epsilon}^{(N)}(z) \right\|_{L(\widetilde{\mathcal{E}}_N, \widetilde{\mathcal{E}}_0)} \|v_0\|_{\widetilde{\mathcal{E}}_N}$$

$$\leq C \left\| v_0 \right\|_{\widetilde{\mathcal{E}}_N} |\epsilon|^{N-1+\eta} .$$

$$(21)$$

Finally we show that $v_0^{(k)}$ lies in $\mathcal{E}_0 \subset L^{\infty}(\Omega, E_0)$ for each $k = 1, \ldots, N-1$. Recall that $v_{\epsilon} \in \mathcal{E}_0$ for every $\epsilon \in (-\epsilon_0, \epsilon_0)$. Thus $\epsilon^{-1}(v_{\epsilon} - v_0)$ belongs to \mathcal{E}_0 . Therefore, since \mathcal{E}_0 isometrically injects into $\widetilde{\mathcal{E}}_0$ (recall the argument above (16)), it follows from the Taylor expansion (20), (21) that $\{\epsilon^{-1}(v_{\epsilon} - v_0)\}_{|\epsilon| < \epsilon_0}$ is a Cauchy sequence in \mathcal{E}_0 . Denote its limit by v'_0 , so that $\epsilon^{-1}(v_{\epsilon} - v_0) - v'_0$ lies in \mathcal{E}_0 and $\|\epsilon^{-1}(v_{\epsilon} - v_0) - v'_0\|_{\mathcal{E}_0} \to 0$ as $\epsilon \to 0$. Then by using again the fact that \mathcal{E}_0 isometrically injects into $\widetilde{\mathcal{E}}_0$, we deduce that v'_0 equals the limit of $\{\epsilon^{-1}(v_{\epsilon} - v_0)\}_{|\epsilon| < \epsilon_0}$ in $\widetilde{\mathcal{E}}_0$. Hence, $v'_0 = v_0^{(1)}$ by (20) and (21), which concludes that $v_0^{(1)}$ lies in \mathcal{E}_0 . By induction (by considering $\epsilon^{-k}(v_{\epsilon} - v_0 - \sum_{j=1}^{k-1} \epsilon^j v_0^{(j)})$ instead of $\epsilon^{-1}(v_{\epsilon} - v_0)$), we can show that $v_0^{(k)}$ also lies in \mathcal{E}_0 for each $k = 2, \ldots, N - 1$. This completes the proof of Theorem 3.6 because (20) and (21) hold with \mathcal{E}_j in place of $\widetilde{\mathcal{E}}_j$.

4 Applications to smooth random dynamical systems

In this section we shall apply Theorem 3.6 to smooth random dynamical systems in order to obtain stability and differentiability results for their random equivariant probability measures. In particular, we will treat random Anosov maps and random U(1) extensions of expanding maps. The treatments of these settings have much in common, and so we will discuss some general, abstract details in earlier subsections.

4.1 Equivariant family of measures

Let M be a compact connected \mathcal{C}^{∞} Riemannian manifold and let m denote the associated Riemannian probability measure on M. Fix a Lebesgue space $(\Omega, \mathcal{F}, \mathbb{P})$ and an invertible, ergodic, \mathbb{P} -preserving map $\sigma : \Omega \to \Omega$. For some $r \geq 1$ let $\mathcal{T} : \Omega \to \mathcal{C}^{r+1}(M, M)$ denote a $(\mathcal{F}, \mathcal{B}_{\mathcal{C}^{r+1}(M,M)})$ -measurable map. The RDS (\mathcal{T}, σ) induced by \mathcal{T} over σ is explicitly written as a measurable map

$$\mathbb{N}_0 \times \Omega \times X \ni (n, \omega, x) \mapsto \mathcal{T}_{\omega}^{(n)}(x), \quad \mathcal{T}_{\omega}^{(n)} := \mathcal{T}_{\sigma^{n-1}\omega} \circ \cdots \circ \mathcal{T}_{\omega},$$

and since σ is invertible, the equivariance of a measurable family of probability measures $\{\mu_{\omega}\}_{\omega\in\Omega}$ for (\mathcal{T},σ) is given as

 $\mu_{\omega} \circ \mathcal{T}_{\omega}^{-1} = \mu_{\sigma\omega} \quad \text{for almost every } \omega \in \Omega,$

(recall Subsection 2.1).

We aim to study the regularity of the dependence of $\{\mu_{\omega}\}_{\omega\in\Omega}$ on the map \mathcal{T} as \mathcal{T} is fiberwise varied in a uniformly \mathcal{C}^N way for some $N \leq r$. To do this we shall realise equivariant families of probability measures as fixed points of (the lifts of) certain operator cocycles (linear RDSs) and then apply Theorem 3.6. In particular, we shall consider the Perron-Frobenius operator cocycle associated to the RDS (\mathcal{T}, σ) on an appropriate Banach space. Recall that the *Perron-Frobenius operator* \mathcal{L}_T associated to a non-singular⁴ measurable map $T: M \to M$ is given by

$$\mathcal{L}_T f = \frac{\mathrm{d}[(fm) \circ T]}{\mathrm{d}m} \quad \text{for } f \in L^1(M, m),$$
(22)

where fm is a finite signed measure given by $(fm)(A) = \int_A f \, \mathrm{d}m$ for $A \in \mathcal{A}$ and $\mathrm{d}\mu/\mathrm{d}m$ is the Radon-Nikodym derivative of an absolutely continuous finite signed measure μ . Note that for each M-valued random variable ψ whose distribution is fm with some density function $f \in L^1(M,m), T(\psi)$ has the distribution $(\mathcal{L}_T f)m$ (and thus, \mathcal{L}_T is also called the *transfer* operator associated with T). It is straightforward to see that

$$\int_{M} \mathcal{L}_{T} f g \, \mathrm{d}m = \int_{M} f g \circ T \, \mathrm{d}m \qquad \text{for } f \in L^{1}(M, m) \text{ and } g \in L^{\infty}(M, m),$$
(23)

and that \mathcal{L}_T is an *m*-Markov operator, where in an abuse of notation we are denoting the linear functional $f \in L^1(M, m) \mapsto \int f \, \mathrm{d}m$ by *m*. In addition, \mathcal{L}_T is positive: if $f \in L^1(M, m)$ satisfies $f \geq 0$ almost everywhere then $\mathcal{L}_T f \geq 0$ almost everywhere.

If 5 det $D_{x}T \neq 0$ for any $x \in M$ then since $T \in \mathcal{C}^{r+1}(M, M)$, one has $\mathcal{L}_{T} \in L(\mathcal{C}^{r}(M))$ with

$$(\mathcal{L}_T f)(x) = \sum_{T(y)=x} \frac{f(y)}{|\det D_y T|} \quad \text{for all } f \in \mathcal{C}^r(M).$$

Hence from \mathcal{T} we obtain a map $\mathcal{L}_{\mathcal{T}} : \omega \mapsto \mathcal{L}_{\mathcal{T}_{\omega}} : \Omega \to L(\mathcal{C}^{r}(M))$, which is measurable by virtue of the following proposition. (We postpone its proof until Appendix A because it is technical and standard.) Let $\mathcal{N}^{r+1}(M, M)$ denote the set of $T \in \mathcal{C}^{r+1}(M, M)$ satisfying det $D_x T \neq 0$ for any $x \in M$.

Proposition 4.1. The map $T \mapsto \mathcal{L}_T$ is continuous on $\mathcal{N}^{r+1}(M, M)$ with respect to the strong operator topology on $L(\mathcal{C}^r(M))$.

Thus if we demand that $\mathcal{T} \in \mathcal{N}^{r+1}(M, M)$ almost surely then $(\mathcal{L}_{\mathcal{T}}, \sigma)$ is an *m*-Markov operator cocycle on $\mathcal{C}^{r}(M)$, which we shall call the *Perron-Frobenius operator cocycle* (on $\mathcal{C}^{r}(M)$) associated to \mathcal{T} . In order to apply the theory of Section 3 we require that the Perron-Frobenius operator cocycle is bounded and *m*-mixing. This later condition will entail some mixing hypotheses on our random systems. However, as in the deterministic case, in order to realise the mixing of the RDS in operator theoretic terms we may be forced to consider the Perron-Frobenius operator cocycle on an alternative Banach space. Specifically, we shall seek Banach spaces $(X, \|\cdot\|_X)$ satisfying the following conditions:

- (s1) $\mathcal{C}^r(M)$ is dense in X with $\mathcal{C}^r(M) \hookrightarrow X$;
- (s2) The embedding $\mathcal{C}^r(M) \hookrightarrow (\mathcal{C}^\infty(M))^*$ given by the map $h \in \mathcal{C}^r(M) \mapsto (g \in \mathcal{C}^\infty(M) \mapsto \int gh \, \mathrm{d}m)$ continuously extends to an embedding $X \hookrightarrow (\mathcal{C}^\infty(M))^*$.

It is clear that any X satisfying (s1) must be separable. Moreover, we note that the functional $\varphi \in (\mathcal{C}^{\infty}(M))^* \mapsto \varphi(1_M)$ is continuous on $(\mathcal{C}^{\infty}(M))^*$, and yields *m* when pulled-back via the embedding $\mathcal{C}^r(M) \hookrightarrow (\mathcal{C}^{\infty}(M))^*$ that is described in (s2). Hence, if (s2) holds, and so we have an embedding $X \hookrightarrow (\mathcal{C}^{\infty}(M))^*$ that continuously extends the $\mathcal{C}^r(M) \hookrightarrow (\mathcal{C}^{\infty}(M))^*$, then *m* induces a continuous linear functional on *X*. In particular, we may speak of *m*-Markov operators in L(X). The following proposition gives a sufficient condition for an *m*-Markov operator in $L(\mathcal{C}^r(M))$ to be extended to an *m*-Markov operator in L(X).

⁴Recall that a measurable map $T: M \to M$ is said to be non-singular (with respect to m) if m(A) = 0 implies that $m(T^{-1}(A)) = 0$.

⁵Notice that if det $D_x T \neq 0$ then T is automatically non-singular with respect to m.

Proposition 4.2. Let (A, σ) be a bounded, m-Markov operator cocycle on $C^{r}(M)$ and X a Banach space satisfying (s1) and (s2). Suppose that

$$\underset{\omega}{\operatorname{ess\,sup}} \sup_{\substack{f \in \mathcal{C}^{r}(M) \\ \|f\|_{X} = 1}} \|A(\omega)f\|_{X} < \infty.$$

$$(24)$$

Then A almost surely extends to a unique, bounded operator on X such that $\omega \mapsto A(\omega) : \Omega \to L(X)$ is strongly measurable. Consequently, (A, σ) is a bounded, m-Markov operator cocycle on X such that

$$\operatorname{ess\,sup}_{\omega} \|A(\omega)\|_{L(X)} < \infty. \tag{25}$$

Proof. It is clear that A almost surely extends to a unique, bounded operator on X, and that

$$\underset{\omega}{\operatorname{ess\,sup}} \|A(\omega)\|_{L(X)} = \underset{\omega}{\operatorname{ess\,sup}} \sup_{\substack{f \in \mathcal{C}^r(M) \\ \|f\|_X = 1}} \|A(\omega)f\|_X < \infty.$$

That A is almost surely m-Markov in L(X) follows straightforwardly from the fact that A is almost surely m-Markov in $L(\mathcal{C}^r(M))$, and that m uniquely extends to a continuous linear functional on X. Hence, it only remains to show that $\omega \mapsto A(\omega)$ is strongly measurable in L(X). Suppose that $f \in X$. Then there exists a sequence $\{f_n\}_{n \in \mathbb{N}} \subset \mathcal{C}^r(M)$ with limit f in X. For each n the map $\omega \mapsto A(\omega)f_n$ is $(\mathcal{F}, \mathcal{B}_{\mathcal{C}^r(M)})$ -measurable, and so it must be $(\mathcal{F}, \mathcal{B}_X)$ measurable too due to (s1). Moreover, for almost every ω we have

$$\lim_{n \to \infty} \|A(\omega)f_n - A(\omega)f\|_X = 0,$$

which is to say that $\omega \mapsto A(\omega)f$ is the almost everywhere pointwise limit (in X) of $(\mathcal{F}, \mathcal{B}_X)$ measurable functions. Hence $\omega \mapsto A(\omega)f$ is $(\mathcal{F}, \mathcal{B}_X)$ -measurable since $(\Omega, \mathcal{F}, \mathbb{P})$ is a Lebesgue space (in particular complete). That $\omega \mapsto A(\omega)$ is strongly measurable in L(X) then follows from Lemma 3.1 and the fact that $f \in X$ was arbitrary.

Hence, by Propositions 4.1 and 4.2, if $\mathcal{T} : \Omega \to \mathcal{N}^{r+1}(M, M)$ is measurable and X satisfies (s1) and (s2), then the Perron-Frobenius operator cocycle $(\mathcal{L}_{\mathcal{T}}, \sigma)$ on $\mathcal{C}^r(M)$ can be extended to a bounded, *m*-Markov operator cocycle on X. Compare also (25) with (QR1).

The following proposition will help us to describe the relationship between the equivariant family of probability measures for (\mathcal{T}, σ) and the fixed point of the lift of a bounded, *m*-mixing Perron-Frobenius operator cocycle $(\mathcal{L}_{\mathcal{T}}, \sigma)$.

Definition 4.3. Assume that X satisfies (s1). $A \in L(X)$ is called positive if $A(X_+) \subset X_+$, where X_+ is the completion of $\{f \in C^r(M) : f \ge 0\}$ in $\|\cdot\|_X$. An operator cocycle (A, σ) is called positive if A is almost surely positive. Furthermore, a distribution $f \in (\mathcal{C}^{\infty}(M))^*$ is called positive if $f(g) \ge 0$ for every $g \in \mathcal{C}^{\infty}(M)$ such that $g \ge 0$.

Proposition 4.4. Let X be a Banach space satisfying (s1) and (s2) and (A, σ) a bounded, m-Markov operator cocycle on X. Suppose that (A, σ) is positive and m-mixing and that h is the unique m-normalised fixed point of the lift $\mathbb{A} : \mathcal{X} \to \mathcal{X}$ on $\mathcal{X} \subset L^{\infty}(\Omega, X)$ (recall (7) for its definition). Then there exists a measurable family of Radon probability measures $\{\mu_{\omega}\}_{\omega \in \Omega}$ such that $h(\omega)(g) = \int g d\mu_{\omega}$ for every $g \in \mathcal{C}^{\infty}(M)$ and almost every ω .

Proof. Notice that the set

$$\mathcal{D} = \{ f \in L^{\infty}(\Omega, X) : m(f) = 1 \text{ and } f \in X_+ \text{ almost surely} \}$$

is almost surely invariant under $A(\omega)$ since (A, σ) is bounded, positive, and *m*-Markov. Hence we may carry out the construction of *h* in Proposition 3.5 with \mathcal{D} in place of \mathcal{X}_1 , to conclude that $h \in X_+$ almost surely. Thus, there exists $\{f_k\}_{k \in \mathbb{N}} \subseteq L^{\infty}(\Omega, \mathcal{C}^r(M))$ such that $f_k(\omega) \ge 0$ and $\int f_k(\omega) dm = 1$ for every k and so that $\lim_{k\to\infty} f_k(\omega) = h(\omega)$ in X for almost every ω . As $X \hookrightarrow (\mathcal{C}^{\infty}(M))^*$ it follows that $\lim_{k\to\infty} f_k(\omega) = h(\omega)$ in the sense of distributions as well. Thus for any positive $g \in \mathcal{C}^{\infty}(M)$ we have

$$h(\omega)(g) = \lim_{k \to \infty} f_k(\omega)(g) = \lim_{k \to \infty} \int f_k(\omega) \cdot g \,\mathrm{d}m \tag{26}$$

(recall the embedding of $\mathcal{C}^r(M)$ in (s2)). As $f_k(\omega)$ and g are positive, it follows from (26) that $h(\omega)(g) \geq 0$ for every such g. Hence $h(\omega)$ is a positive distribution for almost every ω . On the other hand, as is well known, for any positive $f \in (\mathcal{C}^\infty(M))^*$, one can find a positive Radon measure μ_f such that $f(g) = \int g d\mu_f$ for every $g \in \mathcal{C}^\infty(M)$. We denote by μ_ω the positive Radon measure corresponding to $h(\omega)$.

To see that μ_{ω} is a probability measure for almost every ω , we note that by (26) and as $\int f_k(\omega) dm = 1$ for every k we have

$$\mu_{\omega}(M) = h(\omega)(1_M) = \lim_{k \to \infty} \int f_k(\omega) \, \mathrm{d}m = 1.$$

Finally, $\{\mu_{\omega}\}_{\omega\in\Omega}$ is a measurable family on the complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$ because for any $A \in \mathcal{B}_M$, by using (26) again we have

$$\mu_{\omega}(A) = h(\omega)(1_A) = \lim_{k \to \infty} \int_A f_k(\omega) \, \mathrm{d}m$$

for almost every ω , while for every $k, \omega \mapsto f_k(\omega) : \Omega \to \mathcal{C}^r(M)$ is measurable and $f \mapsto \int_A f \, \mathrm{d}m : \mathcal{C}^r(M) \mapsto \mathbb{C}$ is continuous, so that $\omega \mapsto \int_A f_k(\omega) \, \mathrm{d}m$ is measurable too.

Hence if X satisfies (s1) and (s2), and the Perron-Frobenius operator cocycle $(\mathcal{L}_{\mathcal{T}}, \sigma)$ on X is *m*-mixing, then we obtain a measurable family of Radon probability measures $\{\mu_{\omega}\}_{\omega\in\Omega}$ such that $h: \omega \mapsto (g \mapsto \int g d\mu_{\omega})$ is in $L^{\infty}(\Omega, X)$. Furthermore, $\{\mu_{\omega}\}_{\omega\in\Omega}$ is equivariant because it follows from (26) that for any $A \in \mathcal{B}_M$ and almost every ω

$$\mu_{\omega}(T_{\omega}^{-1}(A)) = h(\omega) \left(1_{T_{\omega}^{-1}(A)} \right) = \lim_{k \to \infty} \int f_k(\omega) \cdot 1_A \circ T_{\omega} dm$$

Due to (23), the continuity of $\mathcal{L}_{T_{\omega}}: X \to X$ and the fact that h is the fixed point of the lift of $(\mathcal{L}_{\mathcal{T}}, \sigma)$, this coincides with

$$\lim_{k \to \infty} \int \mathcal{L}_{T_{\omega}} f_k(\omega) \cdot 1_A dm = \lim_{k \to \infty} \mathcal{L}_{T_{\omega}} f_k(\omega) (1_A) = \mathcal{L}_{T_{\omega}} h(\omega) (1_A) = h(\sigma \omega) (1_A) = \mu_{\sigma \omega}(A).$$

4.2 The conditions (QR4) and (QR5)

In this subsection we discuss a sufficient condition for a family of Perron-Frobenius operator cocycles $\{(\mathcal{L}_{\mathcal{T}_{\epsilon}}, \sigma)\}_{\epsilon \in [-1,1]}$ to satisfy (QR4) and (QR5). We emphasise that these conditions hold rather independently of how the underlying random dynamics ($\mathcal{T}_{\epsilon}, \sigma$) behaves (see Proposition 4.5 for precise statement), so we treat (QR4) and (QR5) here as a final preparation before specialising to our applications. For simplicity, throughout this subsection we assume that Mis a *d*-dimensional torus \mathbb{T}^d . One may straightforwardly remove this assumption by considering a partition of unity, refer to e.g. [32, 7] (see also Appendix A).

Notice that (QR4) and (QR5) are conditions for a single iteration $\mathcal{L}_{T_{\epsilon,\omega}}$ (not for $\mathcal{L}_{T_{\epsilon,\sigma^{n-1}\omega}} \circ \cdots \circ \mathcal{L}_{T_{\epsilon,\omega}}$, $n \in \mathbb{N}$), and so clear observations may be found in the non-random setting. Fix $r \geq 1$ and $1 \leq s \leq r$, and we will consider $T \in \mathcal{C}^N([-1,1], \mathcal{C}^{r+1}(\mathbb{T}^d, \mathbb{T}^d))$. Let $1 \leq N \leq s$ be an integer, and E_j , $j \in \{0, \ldots, N\}$, Banach spaces with $E_j \hookrightarrow E_{j-1}$ for each $j \in \{1, \ldots, N\}$ satisfying the following conditions:

- (S1) (s1) holds with E_j in place of X for each $j \in \{0, \ldots, N\}$;
- (S2) (s2) holds with E_j in place of X for each $j \in \{0, \ldots, N\}$;
- (S3) There are constants C > 0 and $0 \le \rho \le r N$ such that

$$||uf||_{E_j} \le C ||u||_{\mathcal{C}^{\rho+j}} ||f||_{E_j} \quad \text{for each } u, f \in \mathcal{C}^r(\mathbb{T}^d) \text{ and } j \in \{0, \dots, N\}.$$

(S4) There is a constant C > 0 such that

$$\left\|\frac{\partial}{\partial x_l}f\right\|_{E_{j-1}} \le C\|f\|_{E_j} \quad \text{for each } f \in \mathcal{C}^r(\mathbb{T}^d), \, l \in \{1, \dots, d\} \text{ and } j \in \{1, \dots, N\}.$$

Observe that all conditions (S1)-(S4) are not for the operators $\mathcal{L}_{T_{\epsilon}}$, $\epsilon \in [-1, 1]$, with $T_{\epsilon} := T(\epsilon)$, but for the spaces E_j , $j \in \{0, \ldots, N\}$, so the following proposition is quite useful in our applications. Note that if

$$\|\mathcal{L}_{T_{\epsilon}}f\|_{E_{j}} < C_{\epsilon}\|f\|_{E_{j}} \quad \text{for each } f \in \mathcal{C}^{r}(\mathbb{T}^{d}), \, j \in \{0, \dots, N\} \text{ and } |\epsilon| \le 1,$$

$$(27)$$

then it follows from Proposition 4.2 that $\mathcal{L}_{T_{\epsilon}}$ is a bounded operator on E_j for each $j \in \{0, \ldots, N\}$ and $|\epsilon| \leq 1$.

Proposition 4.5. Let N be a positive integer, $T \in \mathcal{C}^{N}([-1,1], \mathcal{C}^{r+1}(\mathbb{T}^{d}, \mathbb{T}^{d}))$, and $E_{j}, j \in \{0, \ldots, N\}$, Banach spaces with $E_{j} \hookrightarrow E_{j-1}$ for each $j \in \{1, \ldots, N\}$ satisfying (S1)-(S4). Suppose that $T_{\epsilon} \in \mathcal{N}^{r+1}(\mathbb{T}^{d}, \mathbb{T}^{d})$ for each $\epsilon \in [-1, 1]$ and (27) holds. Then $\epsilon \mapsto \mathcal{L}_{T_{\epsilon}}f$ is in $\mathcal{C}^{j}([-1, 1], E_{i-j})$ for each $j \in \{1, \ldots, N\}$, $i \in \{j, \ldots, N\}$ and $f \in E_{i}$.

Before starting the proof of Proposition 4.5, we discuss a consequence of Proposition 4.5 with respect to the conditions (QR4) and (QR5). Let $\{(\mathcal{T}_{\epsilon}, \sigma)\}_{\epsilon \in [-1,1]}$ be a family of RDSs such that for almost every ω , the map $\epsilon \mapsto T_{\epsilon,\omega} := \mathcal{T}_{\epsilon}(\omega)$ is in $\mathcal{C}^{N}([-1,1], \mathcal{C}^{r+1}(\mathbb{T}^{d}, \mathbb{T}^{d}))$. Let E_{j} , $j \in \{0, \ldots, N\}$ be Banach spaces with $E_{j} \hookrightarrow E_{j-1}$ for each $j \in \{1, \ldots, N\}$ satisfying (S1)-(S4). We suppose that

(S5) $T_{\epsilon,\omega} \in \mathcal{N}^{r+1}(\mathbb{T}^d, \mathbb{T}^d)$ for each $\epsilon \in [-1, 1]$ and almost every ω . Furthermore, (24) holds with E_j and $\mathcal{L}_{T_{\epsilon,\omega}}$ in place of X and $A(\omega)$ for every $j \in \{0, \ldots, N\}$ and $|\epsilon| \leq 1$.

Then, it follows from Proposition 4.2, the Perron-Frobenius operator cocycles $(\mathcal{L}_{\mathcal{T}_{\epsilon}}, \sigma), \epsilon \in [-1, 1]$, can be extended to a bounded operator cocycles on each E_j , and (QR1) holds for these operator cocycles by virtue of (25).

For each $j \in \{0, \ldots, N\}$, $i \in \{j, \ldots, N\}$ and almost every ω , it follows from Proposition 4.5 that we can define $Q_j(\omega) : E_i \to E_{i-j}$ by

$$Q_j(\omega)f = \frac{1}{j!} \left(\frac{d^j}{d\epsilon^j} \mathcal{L}_{T_{\epsilon,\omega}} f \right)_{\epsilon=0} \quad \text{for } f \in E_i.$$

By the definition, it is straightforward to see that for all $\epsilon \in [-1, 1]$ and $2 \leq j \leq N$,

$$\operatorname{ess\,sup}_{\omega} \left\| \mathcal{L}_{T_{\epsilon,\omega}} - \mathcal{L}_{T_{0,\omega}} \right\|_{L(E_N, E_{N-1})} \leq C \left| \epsilon \right|,$$

and

$$\operatorname{ess\,sup}_{\omega} \left\| \mathcal{L}_{T_{\epsilon,\omega}} - \mathcal{L}_{T_{0,\omega}} - \sum_{k=1}^{j-1} \epsilon^k Q_k(\omega) \right\|_{L(E_N, E_{N-j})} \leq C \left|\epsilon\right|^j.$$

To summarise the above argument, we conclude the following:

Corollary 4.6. Suppose that (S1)-(S5) hold for the family of Perron-Frobenius operator cocyles $\{(\mathcal{L}_{\mathcal{T}_{\epsilon}}, \sigma)\}_{\epsilon \in [-1,1]}$ on Banach spaces E_j , $j \in \{0, \ldots, N\}$. Then (QR1), (QR4) and (QR5) hold.

We now return to the proof of Proposition 4.5.

Proof of Proposition 4.5. Fix $1 \leq \sigma \leq N$, $1 \leq \rho \leq r$, $f \in \mathcal{C}^{\sigma}([-1,1],\mathcal{C}^{\rho}(\mathbb{T}^d))$, $g \in \mathcal{C}^{\rho}(\mathbb{T}^d)$ and $1 \leq l \leq d$ for the time being (notice that this f is different from f in the statement of Proposition 4.5 in the sense that this f depends on $\epsilon \in [-1,1]$). We simply denote $\frac{d^a}{d\epsilon^a}f \in \mathcal{C}^{\sigma-a}([-1,1],\mathcal{C}^{\rho}(\mathbb{T}^d))$ by $f^{(a)}$ for each integer $a \in [0,\sigma]$. We also simply denote by $\partial_l g$ the partial derivative of g with respect to the l-th coordinate and let $\partial^{\alpha} = \partial_1^{\alpha_1} \cdots \partial_d^{\alpha_d}$ and $|\alpha| = \alpha_1 + \cdots + \alpha_d$ for each multi-index $\alpha = (\alpha_1, \ldots, \alpha_d) \in \mathbb{N}_0^d$. Then for each $\epsilon \in [-1, 1]$ and $x \in \mathbb{T}^d$,

$$\partial_l(f_{\epsilon}^{(1)})(x) = \partial_{\epsilon}\partial_l \hat{f}(\epsilon, x) = \partial_l \partial_{\epsilon} \hat{f}(\epsilon, x)$$

where $\tilde{f}: [-1,1] \times \mathbb{T}^d \to \mathbb{T}^d$ is given by $\tilde{f}(\epsilon, x) = f_{\epsilon}(x)$. (Since $f^{(1)} \in \mathcal{C}^0([-1,1], \mathcal{C}^1(\mathbb{T}^d))$, it is straightforward to see that the first equality holds and $(\epsilon, x) \mapsto \partial_l(f_{\epsilon}^{(1)})(x)$ is continuous. The second equality also immediately follows from these observations together with Schwarz-Clairaut's theorem on equality of mixed partials.) In particular,

$$f^{(1)} = (\epsilon \mapsto \partial_{\epsilon} \tilde{f}(\epsilon, \cdot)) \quad \text{in } \mathcal{C}^{\sigma-1}([-1, 1], \mathcal{C}^{\rho}(\mathbb{T}^d)).$$
(28)

Furthermore, it is also straightforward to see that the map $\epsilon \mapsto \partial_l f_{\epsilon}$ is in $\mathcal{C}^{\sigma}([-1,1], \mathcal{C}^{\rho-1}(\mathbb{T}^d))$, which we denote by $\partial_l f$ as a slight abuse of notation, and that

$$(\partial_l f)^{(1)} = \partial_l (f^{(1)}) \quad \text{in } \mathcal{C}^{\sigma-1}([-1,1], \mathcal{C}^{\rho-1}(\mathbb{T}^d)),$$
 (29)

$$(\det DT)^{(1)} = \det DT^{(1)} \quad \text{in } \mathcal{C}^{N-1}([-1,1],\mathcal{C}^{r-1}(\mathbb{T}^d)).$$
 (30)

Moreover, we denote by $T_{(l)} \in \mathcal{C}^N([-1,1], \mathcal{C}^{r+1}(\mathbb{T}^d))$ the map $\epsilon \mapsto (x \mapsto T_{(l),\epsilon}(x))$, where $T_{(l),\epsilon}(x)$ is the *l*-th coordinate of $T_{\epsilon}(x) \in \mathbb{T}^d$ (under the identification of \mathbb{T}^d with \mathbb{R}^d). Finally we define a map $\mathbf{L}f: [-1,1] \to \mathcal{C}^{\rho}(\mathbb{T}^d)$ by

$$(\mathbf{L}f)_{\epsilon} = \mathcal{L}_{T_{\epsilon}}f_{\epsilon} \text{ for } \epsilon \in [-1, 1],$$

which is well-defined by virtue of (27). The following is the key lemma for the proof of Proposition 4.5.

Lemma 4.7. For each $f \in C^{\sigma}([-1,1], C^{\rho}(\mathbb{T}^d))$ with $\sigma \in [1,N]$ and $\rho \in [1,r]$, $(\mathbf{L}f)^{(1)}$ exists in $C^{\sigma-1}([-1,1], C^{\rho-1}(\mathbb{T}^d))$, and is of the form

$$(\mathbf{L}f)^{(1)} = \mathbf{L}\left(\sum_{|\alpha| \le 1} J_{0,\alpha} \cdot \partial^{\alpha} f + \sum_{|\alpha| \le 1} J_{1,\alpha} \cdot \partial^{\alpha} f^{(1)}\right),\tag{31}$$

where $J_{k,\alpha}$ is in $\mathcal{C}^{N-1}([-1,1],\mathcal{C}^{r-1}(\mathbb{T}^d))$ being a polynomial function of $\partial^{\beta}T_{(l)}$, $\partial^{\beta}T_{(l)}^{(1)}$ $(1 \leq l \leq d, |\beta| \leq 2)$ and $(\det DT)^{-1}$ for each k = 0, 1 and multi-index α with $|\alpha| \leq 1$.

Proof. Observe that det $D_x T_{\epsilon} > 0$ for all $|\epsilon| \leq 1$ and $x \in \mathbb{T}^d$ or det $D_x T_{\epsilon} < 0$ for all $|\epsilon| \leq 1$ and $x \in \mathbb{T}^d$ because $T_{\epsilon} \in \mathcal{N}^{r+1}(\mathbb{T}^d, \mathbb{T}^d)$ for each $|\epsilon| \leq 1$ and $\epsilon \mapsto T_{\epsilon}$ is continuous. We only consider the former case because the other case is similar. Also, we only show (31) around $\epsilon = 0$ to keep our notation simple (the general case can be literally treated). We first note that there is $\epsilon_0 > 0$, $B \in \mathbb{N}$, a finite covering $\{U_\lambda\}_{\lambda \in \Lambda}$ of \mathbb{T}^d and \mathcal{C}^{r+1} maps $(T_{\epsilon})_{\lambda,b}^{-1} : U_{\lambda} \to (T_{\epsilon})_{\lambda,b}^{-1}(U_{\lambda})$ for $|\epsilon| < \epsilon_0, \lambda \in \Lambda$ and $b \in \{1, \ldots, B\}$ such that for each $|\epsilon| < \epsilon_0, \lambda \in \Lambda, b \in \{1, \ldots, B\}$ and $g \in \mathcal{C}^r(\mathbb{T}^d)$,

$$T_{\epsilon} \circ (T_{\epsilon})_{\lambda,b}^{-1}(x) = x \quad \text{on } U_{\lambda}$$
(32)

and

$$\mathcal{M}_1(\epsilon, g)(x) := \sum_{T_\epsilon(y)=x} g(y) = \sum_{b=1}^B g \circ (T_\epsilon)_{\lambda, b}^{-1}(x) \quad \text{on } U_\lambda$$
(33)

(because $T_0 \in \mathcal{N}^{r+1}(\mathbb{T}^d, \mathbb{T}^d)$ and \mathbb{T}^d is compact, see Appendix A for detail). Note also that if we define $\mathcal{M}_2 : [-1, 1] \times \mathcal{C}^r(\mathbb{T}^d) \to \mathcal{C}^r(\mathbb{T}^d)$ by

$$\mathcal{M}_2(\epsilon, g) = \frac{g}{\det DT_{\epsilon}} \quad \text{for } \epsilon \in [-1, 1], \ g \in \mathcal{C}^r(\mathbb{T}^d),$$

then for each $f \in \mathcal{C}^{\sigma}([-1,1],\mathcal{C}^{r}(\mathbb{T}^{d}))$ and $|\epsilon| < \epsilon_{0}$, we have

$$(\mathbf{L}f)_{\epsilon} = \mathcal{M}_1(\epsilon, \mathcal{M}_2(\epsilon, f_{\epsilon})).$$

Notice that both \mathcal{M}_1 and \mathcal{M}_2 are linear with respect to $g \in \mathcal{C}^r(\mathbb{T}^d)$. Hence it follows from the chain rule for (Fréchet) derivatives that

$$(\mathbf{L}f)_{\epsilon}^{(1)} = \partial_{\epsilon}\mathcal{M}_{1}(\epsilon, \mathcal{M}_{2}(\epsilon, f_{\epsilon})) + \mathcal{M}_{1}\left(\epsilon, \partial_{\epsilon}\mathcal{M}_{2}(\epsilon, f_{\epsilon}) + \mathcal{M}_{2}\left(\epsilon, f_{\epsilon}^{(1)}\right)\right)$$
(34)

if the derivatives exists, where $\partial_{\epsilon} = \frac{\partial}{\partial \epsilon}$

Now we calculate $\partial_{\epsilon} \mathcal{M}_1$ and $\partial_{\epsilon} \mathcal{M}_2$. We first show that

$$\partial_{\epsilon} \mathcal{M}_{1}(\epsilon, g) = \mathcal{M}_{1}\left(\epsilon, -\sum_{l=1}^{d} \partial_{l} g \cdot \frac{\sum_{k=1}^{d} (\operatorname{adj}(DT_{\epsilon}))_{l,k} \cdot T_{(k),\epsilon}^{(1)}}{\det DT_{\epsilon}}\right),$$
(35)

where $\operatorname{adj}(A)$ is the adjugate matrix (i.e. the transpose of the cofactor matrix) of a matrix A. By (28) (with $\epsilon \mapsto \mathcal{M}_1(\epsilon, g)$ in place of f), (33) and the chain rule for derivatives, we have

$$\partial_{\epsilon} \mathcal{M}_{1}(\epsilon, g)(x) = \partial_{\epsilon} \widetilde{\mathcal{M}}_{1,g}(\epsilon, x) = \sum_{b=1}^{B} \sum_{l=1}^{d} \partial_{l} g \circ (T_{\epsilon})_{b,\lambda}^{-1}(x) \cdot \partial_{\epsilon} ((T_{\epsilon})_{b,\lambda}^{-1})_{(l)}(x), \quad x \in U_{\lambda},$$
(36)

where $((T_{\epsilon})_{b,\lambda}^{-1})_{(l)}(x)$ is the *l*-th coordinate of $(T_{\epsilon})_{b,\lambda}^{-1}(x)$ and $\widetilde{\mathcal{M}}_{1,g}(\epsilon, x) = \mathcal{M}_1(\epsilon, g)(x)$. On the other hand, by differentiating the *l*-th coordinate of (32) for $1 \leq l \leq d$ we have

$$T_{(\ell),\epsilon}^{(1)}(y) + \sum_{k=1}^{d} \partial_k T_{(\ell),\epsilon}(y) \cdot \partial_{\epsilon}((T_{\epsilon})_{b,\lambda}^{-1})_{(k)}(x) = 0, \quad y = (T_{\epsilon})_{b,\lambda}^{-1}(x), \ x \in U_{\lambda}.$$

In the matrix form (under the identification of \mathbb{T}^d with \mathbb{R}^d), this can be written as

$$T_{\epsilon}^{(1)}(y) + D_y T_{\epsilon} \left[\partial_{\epsilon} (T_{\epsilon})_{b,\lambda}^{-1}(x) \right] = 0, \quad y = (T_{\epsilon})_{b,\lambda}^{-1}(x), \ x \in U_{\lambda},$$

where we see $T_{\epsilon}^{(1)}(y)$ and $\partial_{\epsilon}(T_{\epsilon})_{b,\lambda}^{-1}(x)$ as column vectors. Thus, since $A^{-1} = (\det A)^{-1} \operatorname{adj}(A)$ for any invertible matrix A, we have

$$\partial_{\epsilon}(T_{\epsilon})^{-1}_{b,\lambda}(x) = -(\det D_y T_{\epsilon})^{-1} \operatorname{adj}(D_y T_{\epsilon}) \left[T_{\epsilon}^{(1)}(y)\right], \quad y = (T_{\epsilon})^{-1}_{b,\lambda}(x), \ x \in U_{\lambda}.$$
(37)

(35) immediately follows from (36) and (37). Furthermore, by the quotient rule for derivatives and (30) we have

$$\partial_{\epsilon} \mathcal{M}_2(\epsilon, g) = -\frac{g \cdot \det DT_{\epsilon}^{(1)}}{(\det DT_{\epsilon})^2}$$
(38)

and

$$\partial_l \mathcal{M}_2(\epsilon, g) = \frac{\partial_l g}{\det DT_\epsilon} - \frac{g \cdot \partial_l (\det DT_\epsilon)}{(\det DT_\epsilon)^2}.$$
(39)

The conclusion immediately follows from (34), (35), (38) and (39).

Now we complete the proof of Proposition 4.5. We first consider the case when $f \in \mathcal{C}^r(\mathbb{T}^d)$. We will show by induction that for each $1 \leq k \leq j$, $(\mathbf{L}f)^{(k)}$ exists and is of the form

$$(\mathbf{L}f)^{(k)} = \mathbf{L}\left(\sum_{|\alpha| \le k} \widehat{J}_{k,\alpha} \cdot \partial^{\alpha} f\right),\tag{40}$$

where $\widehat{J}_{k,\alpha}$ is in $\mathcal{C}^{N-k}([-1,1],\mathcal{C}^{r-k}(\mathbb{T}^d))$ being a polynomial function of $\partial^{\beta}T_{(l)}^{(k')}$ $(1 \leq l \leq d, 0 \leq k' \leq k, |\beta| \leq k+1)$ and $(\det DT)^{-1}$ for each multi-index α with $|\alpha| \leq k$. (40) for k = 1 is an immediate consequence of Proposition 4.7 (notice that f in Proposition 4.7 depended on ϵ while f here does not). Suppose that $k \geq 2$ and (40) holds with k-1 instead of k. Then, by Proposition 4.7, we have

$$(\mathbf{L}f)^{(k)} = \mathbf{L}\left(\sum_{|\alpha| \le 1} J_{0,\alpha} \cdot \partial^{\alpha} \left(\sum_{|\beta| \le k-1} \widehat{J}_{k-1,\beta} \cdot \partial^{\beta} f\right) + \sum_{|\alpha| \le 1} J_{1,\alpha} \cdot \partial^{\alpha} \left(\sum_{|\beta| \le k-1} \widehat{J}_{k-1,\beta}^{(1)} \cdot \partial^{\beta} f\right)\right).$$

Therefore, (40) immediately follows from (29) and (30). Furthermore, $\epsilon \mapsto \frac{d^j}{d\epsilon^j} \mathcal{L}_{T_\epsilon} f = (\mathbf{L}f)_{\epsilon}^{(j)}$ exists as an element of $\mathcal{C}^0([-1,1], E_{i-j})$ by (S3) and the fact that $s-j \geq 0$.

We next consider the general case, i.e. the case when $f \in E_i$. By (S1), one can find $\{f_n\}_{n\geq 1} \subset \mathcal{C}^r(\mathbb{T}^d)$ such that $||f - f_n||_{E_i} \to 0$ as $n \to \infty$. By the result in the previous paragraph, $(\mathbf{L}f_n)^{(k)}$ exists as an element of a Banach space $\mathcal{C}^0([-1,1], E_{i-k})$ for all $1 \leq k \leq j$. On the other hand, it follows from (S3), (S4), (27) and (40) that

$$\sup_{\epsilon \in [-1,1]} \left\| (\mathbf{L}f_n)_{\epsilon}^{(k)} - (\mathbf{L}f_m)_{\epsilon}^{(k)} \right\|_{E_{i-k}} \le C \sum_{|\alpha| \le k} \sup_{\epsilon \in [-1,1]} \left\| \widehat{J}_{k,\alpha,\epsilon} \right\|_{\mathcal{C}^{r-k}} \left\| f_n - f_m \right\|_{E_i} \to 0$$

as $n, m \to \infty$. In particular, $\lim_{n\to\infty} (\mathbf{L}f_n)^{(j)}$ exists. In a similar manner, we can show that **L** is a bounded operator from $\mathcal{C}^0([-1,1], E_i)$ to $\mathcal{C}^j([-1,1], E_{i-j})$, so that $\lim_{n\to\infty} (\mathbf{L}f_n)^{(j)} = (\mathbf{L}f)^{(j)}$ in $\mathcal{C}^0([-1,1], E_{i-j})$. In conclusion, $\mathbf{L}f : \epsilon \mapsto \mathcal{L}_{T_\epsilon}f$ is in $\mathcal{C}^j([-1,1], E_{i-j})$.

4.3 Random Anosov maps

Let M be a compact, connected \mathcal{C}^{∞} Riemannian manifold with dimension d. In this section we consider random dynamical systems consisting of Anosov maps lying in a small $\mathcal{C}^{r+1}(M, M)$ neighbourhood of a fixed, topologically transitive Anosov diffeomorphism $T \in \mathcal{C}^{r+1}(M, M)$ for some $r \geq 1$. The setting we consider is very similar to that of [20, Section 4], however we shall obtain more general conclusions than those of [20]. For the remainder of this section we shall fix a topologically transitive Anosov diffeomorphism $T \in \mathcal{C}^{r+1}(M, M)$. Recall that $(\Omega, \mathcal{F}, \mathbb{P})$ is a Lebesgue space and $\sigma : \Omega \to \Omega$ is a measurably invertible, ergodic, measure-preseving map. For every $\eta > 0$ we define

$$\mathcal{O}_{\eta}(T) = \{ S \in \mathcal{C}^{r+1}(M, M) : d_{\mathcal{C}^{r+1}}(S, T) < \eta \}.$$

Recall that if η is sufficiently small then $\mathcal{O}_{\eta}(T) \subset \mathcal{N}^{r+1}(M, M)$ and every $S \in \mathcal{O}_{\eta}(T)$ is an Anosov diffeomorphism. A map $\mathcal{T} : \Omega \to \mathcal{C}^{r+1}(M, M)$ will be said to be measurable if it is $(\mathcal{F}, \mathcal{B}_{\mathcal{C}^{r+1}(M,M)})$ -measurable.

We consider random dynamical systems induced by measurable maps $\mathcal{T} : \Omega \to \mathcal{O}_{\eta}(T)$ for some small, fixed η , over σ . Our main result for this subsection concerns the stability properties of the equivariant family of probability measures associated to such systems. We will formulate our result in the setting developed by Gouëzel and Liverani in [32]. In particular, in [32] it is shown that when a topologically transitive Anosov map is smoothly perturbed the SRB measure varies with similar regularity in certain anisotropic Banach spaces.

A small technical comment is required before proceeding: in [32] the usual metric on M is replaced by an adapted metric for T (which will also be adapted for $S \in \mathcal{O}_{\eta}(T)$ provided that η is sufficiently small); we shall do the same here. We denote by m the Riemannian probability measure induced by the adapted metric on M. For each $q \ge 0$, $p \in \mathbb{N}_0$ with $p \le r$ one obtains a space $B_{p,q}(T)$ by taking the completion of $\mathcal{C}^r(M)$ with respect to anisotropic norms⁶ $\|\cdot\|_{p,q}$ as defined in [32, Section 3]. Since the map T is fixed we will just write $B_{p,q}$ in place of $B_{p,q}(T)$. Our main result for this subsection is the following.

Theorem 4.8. Let $N, p \in \mathbb{N}$ and $q \geq 0$ satisfying that p + q < r - N. Then there exists $\eta_0 > 0$ such that every measurable $\mathcal{T} : \Omega \to \mathcal{O}_{\eta_0}(T)$ has an equivariant measurable family of Radon probability measures $\{\mu_{\omega}^{\mathcal{T}}\}_{\omega\in\Omega}$ and $h_{\mathcal{T}} \in L^{\infty}(\Omega, B_{p+N,q})$ such that $h_{\mathcal{T}}(\omega)(g) = \int g d\mu_{\omega}^{\mathcal{T}}$ for each $g \in \mathcal{C}^{\infty}(M)$ and almost every ω . In addition, if $\{\mathcal{T}_{\epsilon} : \Omega \to \mathcal{O}_{\eta_0}(T)\}_{\epsilon\in[-1,1]}$ is a family of measurable maps such that there is a bounded subset \mathcal{K} of $\mathcal{C}^{N}([-1,1],\mathcal{C}^{r+1}(M,M))$ satisfying that $\epsilon \mapsto \mathcal{T}_{\epsilon}(\omega)$ lies in \mathcal{K} for almost every ω , then the map $\epsilon \mapsto h_{\mathcal{T}_{\epsilon}}(\omega)$ is in $\mathcal{C}^{N-1}([-1,1], B_{p,q+N})$ for almost every ω .

We will use Theorem 3.6 to prove Theorem 4.8, with help of Proposition 4.2, Proposition 4.4 and Corollary 4.6. Therefore, what we should do is to check (S1)-(S5), (QR0), (QR2) and (QR3) for appropriate Banach spaces. We start with the basic properties of the $B_{p,q}$ spaces from [32].

- 1. By definition of $\|\cdot\|_{p,q}$, it is straightforward to see that $\|\partial_l f\|_{p,q} \leq \|f\|_{p+1,q-1}$ for each $f \in \mathcal{C}^r(M)$ and $1 \leq l \leq d$. Furthermore, $B_{p+1,q-1} \hookrightarrow B_{p,q}$.
- 2. [32, Lemma 2.1] If p + q < r then the unit ball in $B_{p+1,q-1}$ is relatively compact in $B_{p,q}$.
- 3. [32, Lemma 3.2] $||uf||_{p,q} \leq C ||u||_{\mathcal{C}^{p+q}} ||f||_{p,q}$ for each $u \in \mathcal{C}^{p+q}(M)$ and $f \in \mathcal{C}^{r}(M)$. In particular, if p+q < r then $\mathcal{C}^{r}(M) \hookrightarrow B_{p,q}$ (see also [32, Remark 4.3]).
- 4. [32, Proposition 4.1] We have $B_{p,q} \hookrightarrow (\mathcal{C}^q(M))^*$. Specifically, for each $h \in \mathcal{C}^r(M)$ one obtains a distribution $\tilde{h} \in (\mathcal{C}^q(M))^*$ defined by $\tilde{h}(g) = \int hg \, \mathrm{d}m$. The map $h \mapsto \tilde{h}$ continuously extends from $\mathcal{C}^r(M)$ to $B_{p,q}$ and yields the required inclusion.

We also remark that there exists injections $B_{p,q} \to B_{p-1,q}$ and $B_{p,q} \to B_{p,q'}$ for q' > q due to [32, Remark 4.2]. By the fourth item of the above list, the functional $h \mapsto \int h \, dm$ on $\mathcal{C}^r(M)$ extends to a continuous functional on $B_{p,q}$, which we shall also denote by m. The following result summarises some facts from [32] and [15] pertaining to the boundedness and mixing of the Perron-Frobenius operator associated to maps in $\mathcal{O}_{\eta}(T)$ for η small. We refer the reader to [32, Lemma 2.2] and the discussion at the beginning of [32, Section 7] for the first and second items, and [15, Proposition 2.10] for the third item (see also [19, Section 3]).

Proposition 4.9. There exists $0 < \eta_0 \leq \eta$ such that for any $p \in \mathbb{N}_0$ and $q \geq 0$ with p + q < r we have:

1. For every sequence $\{T_i\}_{i\in\mathbb{N}}\subseteq \mathcal{O}_{\eta_0}(T)$ and $n\in\mathbb{N}$ we have

$$\left\|\mathcal{L}_{T_n}\circ\cdots\circ\mathcal{L}_{T_1}\right\|_{L(B_{p,q})}\leq C_{p,q}.$$

⁶Actually, the norms in [32, Section 3] are defined on the real Banach space $\mathcal{C}^{r}(M,\mathbb{R})$. Here we consider the complexification, which is of no consequence.

⁷Recall that $\mathcal{C}^{r+1}(M, M)$ is a \mathcal{C}^{r+1} Banach manifold and so for $k \leq r+1$ we may talk of \mathcal{C}^k curves taking values in $\mathcal{C}^{r+1}(M, M)$.

2. There exists $\alpha_{p,q} \in [0,1)$ such that for every $\{T_i\}_{i \in \mathbb{N}} \subseteq \mathcal{O}_{\eta_0}(T)$, $n \in \mathbb{N}$ and $f \in B_{p+1,q}$ we have

$$\|(\mathcal{L}_{T_n} \circ \cdots \circ \mathcal{L}_{T_1})f\|_{p+1,q} \le C_{p,q}\alpha_{p,q}^n \|f\|_{B_{p+1,q}} + C_{p,q} \|f\|_{B_{p,q+1}}.$$

3. There exists a constant $\lambda_{p,q} \in [0,1)$ such that for every sequence $\{T_i\}_{i \in \mathbb{Z}} \subseteq \mathcal{O}_{\eta_2}(T)$ and $n \in \mathbb{N}$ we have

$$\left\| \mathcal{L}_{T_n} \circ \cdots \circ \mathcal{L}_{T_1} \right\|_{V_{p,q}} \leq C_{p,q} \lambda_{p,q}^n,$$

where $V_{p,q} = \ker \left(m \right|_{B_{p,q}} \right) = \{ h \in B_{p,q} \mid m(h) = 0 \}.$

Fix $q \ge 0$ and $p \in \mathbb{N}$ with p + q < r - N. Let $E_j = B_{p+j,q+N-j}$, $j \in \{0, \ldots, N\}$. Then, the conditions (S1)-(S4) on theses Banach spaces immediately follow from the above list (recall that each E_j is the completion of $\mathcal{C}^r(M)$ with respect to $\|\cdot\|_{E_j}$). Furthermore, fix a bounded subset \mathcal{K} of $\mathcal{C}^N([-1,1], \mathcal{C}^{r+1}(M,M))$ and let $\{\mathcal{T}_{\epsilon} : \Omega \to \mathcal{O}_{\eta_0}(T)\}_{\epsilon \in [-1,1]}$ be a family of measurable maps such that $\epsilon \mapsto \mathcal{T}_{\epsilon}(\omega)$ lies in \mathcal{K} for almost every ω . Then, by virtue of Proposition 4.2, the first part of Proposition 4.9 and the above list, Perron-Frobenius operator cocycles $\{(\mathcal{L}_{\mathcal{T}_{\epsilon}},\sigma)\}_{\epsilon \in [-1,1]}$ associated with the random dynamics $\{(\mathcal{T}_{\epsilon},\sigma)\}_{\epsilon \in [-1,1]}$ are precisely defined on these Banach spaces and they satisfy (S5) and (QR0) with $\xi = m$ (see the remark following (s2)) except the *m*-mixing property. In fact, (QR2), (QR3) and the mixing of $\{(\mathcal{L}_{\mathcal{T}_{\epsilon}},\sigma)\}_{\epsilon \in [-1,1]}$ on E_j for $j \in \{1, N\}$ are consequences of each items of Proposition 4.9, respectively. By Corollary 4.6, (QR1), (QR4) and (QR5) also hold for $\{(\mathcal{L}_{\mathcal{T}_{\epsilon}},\sigma)\}_{\epsilon \in [-1,1]}$ on E_j (see also [32, Lemma 7.1] and [32, Section 9]).

By Proposition 4.4 (and the remark following it), for each $\epsilon \in [-1, 1]$, \mathcal{T}_{ϵ} has an equivariant measurable family of Radon probability measures $\{\mu_{\omega}^{\mathcal{T}_{\epsilon}}\}_{\omega\in\Omega}$ and $h_{\mathcal{T}_{\epsilon}} \in L^{\infty}(\Omega, E_N) = L^{\infty}(\Omega, B_{p+N-1,q})$ such that $h_{\mathcal{T}_{\epsilon}}(\omega)(g) = \int g d\mu_{\omega}^{\mathcal{T}_{\epsilon}}$ for each $g \in \mathcal{C}^{\infty}(M)$ and almost every ω . Furthermore, we apply Theorem 3.6 to deduce the claim that $\epsilon \mapsto h_{\mathcal{T}_{\epsilon}}(\omega)$ is in $\mathcal{C}^{N-1}([-1, 1], E_0) = \mathcal{C}^{N-1}([-1, 1], B_{p-1,q+N})$ for almost every ω , which completes the proof of Theorem 3.6.

4.4 Random U(1) extensions of expanding maps

In this section, we will apply Theorem 3.6 to quenched linear response problems for random U(1) extensions of expanding maps. Let \mathcal{U} be the set of \mathcal{C}^{∞} endomorphisms $T : \mathbb{T}^2 \to \mathbb{T}^2$ on the torus $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$ of the form

$$T: \begin{pmatrix} x \\ s \end{pmatrix} \mapsto \begin{pmatrix} E(x) \\ s + \frac{1}{2\pi}\tau(x) \mod 1 \end{pmatrix}, \tag{41}$$

where $E : \mathbb{S}^1 \to \mathbb{S}^1$ is a \mathcal{C}^{∞} orientation-preserving endomorphism on the circle $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$ and $\tau : \mathbb{S}^1 \to \mathbb{R}$ is a \mathcal{C}^{∞} function (*T* is called the U(1) extension of *E* over τ). U(1) extensions of expanding maps can be seen as a toy model of (piecewise) hyperbolic flows such as geodesic flows on negatively curved manifolds or dispersive billiard flows (via suspension flows of hyperbolic maps; see [34, 43]), and has been intensively studied by several authors (see e.g. [17, 22, 42, 41, 12, 23, 13]). When we want to emphasise the dependence of *E* and τ in (41) on *T*, we write them as E_T and τ_T . Fix $T \in \mathcal{U}$ and assume that *E* is an expanding map on \mathbb{S}^1 in the sense that $\min_{x \in \mathbb{S}^1} E'(x) > 1$. Let *r* be a positive integer. For every $\eta > 0$ we define

$$\mathcal{O}_{\eta}(T) = \{ S \in \mathcal{U} \mid d_{\mathcal{C}^{r+1}}(S, T) < \eta \}.$$

Note that $\mathcal{U} \subset \mathcal{N}^{r+1}(\mathbb{T}^2, \mathbb{T}^2)$, and that if η is sufficiently small then E_S is an expanding map for every $S \in \mathcal{O}_{\eta}(T)$.

Recall that $(\Omega, \mathcal{F}, \mathbb{P})$ is a Lebesgue space and $\sigma : \Omega \to \Omega$ is a measurably invertible, ergodic, \mathbb{P} -preserving map on $(\Omega, \mathcal{F}, \mathbb{P})$. When $\tau_T(x) = \alpha$ for any $x \in \mathbb{S}^1$ with some constant α , then obviously T does not admit any mixing physical measure because the rotation $s \mapsto s + \frac{1}{2\pi} \alpha$ mod 1 has no mixing physical measure. However, it is known that if τ satisfies a generic condition, called the *partial captivity* condition, then T admits a unique absolutely continuous invariant probability measure for which correlation functions of T decay exponentially fast (in particular, T is mixing). The partial captivity condition was first introduced by Faure [22] and proven to be generic in [41]. Furthermore, it was shown in [42, Theorem 1.6] that if T satisfies the partial captivity condition, then there is an $\eta_0 > 0$ and an $m_0 \in \mathbb{N}$, only depending on T(see the comment above Proposition 4.11 for more precise choice of η_0 and m_0), such that if $r \geq m_0$ then for any measurable map $\mathcal{T} : \Omega \to \mathcal{O}_{\eta_0}(T)$, the RDS (\mathcal{T}, σ) induced by \mathcal{T} over σ admits a unique equivariant measurable family of absolutely continuous probability measures $\{\mu_{\omega}^{\mathcal{T}}\}_{\omega\in\Omega}$ such that the Radon-Nikodym derivative of $\mu_{\omega}^{\mathcal{T}}$ is in the usual Sobolev space $H^r(\mathbb{T}^2)$ of regularity r for \mathbb{P} -almost every ω and that quenched correlation functions of (\mathcal{T}, σ) for $\{\mu_{\omega}^{\mathcal{T}}\}_{\omega\in\Omega}$ decay exponentially fast. See also [21, 29] for related results.

Assume that T satisfies the transversality condition, and fix such an $\eta_0 > 0$ and an $m_0 \in \mathbb{N}$. Assume also that $r \ge m_0 + 1$. The main result in this section is the following.

Theorem 4.10. Let N be positive integers such that $N \leq r-m_0$. If $\{\mathcal{T}_{\epsilon} : \Omega \to \mathcal{O}_{\eta_0}(T)\}_{\epsilon \in [-1,1]}$ is a family of measurable maps such that there is a bounded subset \mathcal{K} of $\mathcal{C}^N([-1,1],\mathcal{C}^{r+1}(\mathbb{T}^2,\mathbb{T}^2))$ satisfying that $\epsilon \mapsto \mathcal{T}_{\epsilon}(\omega)$ lies in \mathcal{K} for \mathbb{P} -almost every $\omega \in \Omega$, then the map $\epsilon \mapsto \mu_{\omega}^{\mathcal{T}_{\epsilon}}$ is in $\mathcal{C}^{N-1}([-1,1], H^{r-N}(\mathbb{T}^2))$ for \mathbb{P} -almost every $\omega \in \Omega$.

We recall the basic properties of the Sobolev spaces $H^m(\mathbb{T}^2)$ with regularity $m \in \mathbb{N}_0$. Recall that $\|f\|_{H^m}^2 = \sum_{|\alpha| \le m} \|\partial^{\alpha} f\|_{L^2}^2$.

- 1. By definition of $\|\cdot\|_{H^m}$, it is straightforward to see that $\|\partial_l f\|_{H^m} \leq \|f\|_{H^{m+1}}$ for each $f \in \mathcal{C}^{m+1}(\mathbb{T}^2)$ and $1 \leq l \leq d$, and $\|uf\|_{H^m} \leq C \|u\|_{\mathcal{C}^m} \|f\|_{H^m}$ for each $u, f \in \mathcal{C}^m(\mathbb{T}^2)$.
- 2. By Kondrachov embedding theorem, $H^{m+1}(\mathbb{T}^2) \hookrightarrow H^m(\mathbb{T}^2)$ and the unit ball in $H^{m+1}(\mathbb{T}^2)$ is relatively compact in $H^m(\mathbb{T}^2)$.
- 3. $\mathcal{C}^{m'}(\mathbb{T}^2)$ is dense in $H^m(\mathbb{T}^2)$ for each $m' \geq m$ because $\mathcal{C}^{m'}(\mathbb{T}^2) \subset H^m(\mathbb{T}^2) \subset \mathcal{C}^{m-d/2}(\mathbb{T}^2)$ by Sobolev embedding theorem.
- 4. By Cauchy-Schwarz inequality, we have $H^m(\mathbb{T}^2) \hookrightarrow (\mathcal{C}^0(\mathbb{T}^2))^*$ by $h \mapsto \tilde{h}$ given by $\tilde{h}(g) = \int hg \, \mathrm{d}m$ for $g \in \mathcal{C}^0(\mathbb{T}^2)$.

Let $\lambda_0 := (\inf_{S \in \mathcal{O}_{\eta_0}} \min_{x \in \mathbb{S}^1} E'_S(x))^{-1}$, which is less than 1 by taking η_0 small if necessary. Fix $\lambda \in (\lambda_0^{\frac{1}{2}}, 1)$. Let m_0 be a positive integer such that $\lambda^{2m_0+1} > \deg(T)$, where $\deg(T)$ is the degree of T. Let $N \leq r - m_0$ be a positive integer. By taking η_0 small if necessary, we assume that $\inf_{S \in \mathcal{O}_{\eta_0}} \min_{x \in \mathbb{S}^1} E'_S(x) < \lambda$. Fix a family of measurable maps $\{\mathcal{T}_\epsilon : \Omega \to \mathcal{O}_{\eta_0}(T)\}_{\epsilon \in [-1,1]}$ such that there is a bounded subset \mathcal{K} of $\mathcal{C}^N([-1,1], \mathcal{C}^{r+1}(\mathbb{T}^2, \mathbb{T}^2))$ satisfying that $\epsilon \mapsto \mathcal{T}_\epsilon(\omega)$ lies in \mathcal{K} for \mathbb{P} -almost every $\omega \in \Omega$. Let $(\mathcal{L}_{\mathcal{T}_\epsilon}, \sigma)$ be the Perron-Probenius cocycle induced by $(\mathcal{T}_\epsilon, \sigma)$. Then, it follows from [42, §4] that $\mathcal{L}_{\mathcal{T}_\epsilon}$ almost surely extends to a unique, bounded operator on $H^m(\mathbb{T}^2)$ such that $\omega \mapsto \mathcal{L}_{\mathcal{T}_\epsilon}(\omega) : \Omega \to L(H^m(\mathbb{T}^2))$ is strongly measurable for each $\epsilon \in [-1, 1]$. The following estimates were proven in [14, Subsections 2.3 and 2.4]:

Proposition 4.11. There is a constant $\rho \in (0,1)$ (which may depend on T and η_0) such that for all $|\epsilon| \leq 1$ the following holds:

(1) For each $m \ge 0$ and $n \ge 1$,

$$\operatorname{ess\,sup}_{\omega} \|\mathcal{L}_{\mathcal{T}_{\epsilon}}^{(n)}(\omega)\|_{L(H^{m}(\mathbb{T}^{2}))} \leq C.$$

(2) For each $m \ge m_0$, $n \ge 1$ and $f \in H^m(\mathbb{T}^2)$,

$$\underset{\omega}{\mathrm{ess\,sup}} \|\mathcal{L}_{\mathcal{T}_{\epsilon}}^{(n)}(\omega)f\|_{H^{m+1}} \le C\lambda^{n} \|f\|_{H^{m+1}} + C\|f\|_{H^{m}}.$$

(3) For each $m \ge m_0$, $n \ge 1$ and $f \in H^m(\mathbb{T}^2)$ with $\int_{\mathbb{T}^2} f \, \mathrm{d}m = 0$,

$$\operatorname{ess\,sup}_{\omega} \|\mathcal{L}_{\mathcal{T}_{\epsilon}}^{(n)}(\omega)f\|_{H^{m}} \leq C\rho^{n} \|f\|_{H^{m}}.$$

We now prove Theorem 4.10. Let m = r - N. For $j \in \{0, ..., N\}$ set $E_j = H^{m+j}(\mathbb{T}^2)$. Then, in the same manner as one in the proof of Theorem 4.8, we can apply Theorem 3.6, with help of Proposition 4.2, Proposition 4.4 and Corollary 4.6, to deduce the claim that $\epsilon \mapsto \mu_{\omega}^{\mathcal{T}_{\epsilon}}$ is in $\mathcal{C}^{N-1}([-1,1], E_0) = \mathcal{C}^{N-1}([-1,1], H^{r-N}(\mathbb{T}^2))$ for almost every ω , which completes the proof of Theorem 4.10.

A Proof of Proposition 4.1

Let $T \in \mathcal{N}^{r+1}(M, M)$. Then it follows from [33, Corollary 1] that T is a covering map. Hence by a basic property of covering spaces, there is a discrete topological space Γ such that for every $x \in M$ there is a neighborhood U_x of x such that $T^{-1}(\{x\})$ is homeomorphic to Γ and $T^{-1}(U_x)$ is homeomorphic to $U_x \times \Gamma$. In other words, $T^{-1}(U_x)$ is a union of disjoint open sets $\{\widetilde{U}_{b,x}\}_{j=1}^B$ such that $T: \widetilde{U}_{b,x} \to U_x$ is a homeomorphism for each $b = 1, \ldots, B$, where B is the cardinality of Γ . If $B = \infty$, then since $|\det DT|$ is bounded uniformly away from 0 due to the compactness of M, we have

$$m(T^{-1}(U_x)) \ge B \cdot \inf_{y \in M} |\det DT(y)| m(U_x) = \infty,$$

which contradicts to that m is a finite measure. Hence $B < \infty$. Furthermore, there is a small neighborhood \mathcal{U} of T in $\mathcal{N}^{r+1}(M, M)$ such that for each $S \in \mathcal{U}$ and $x \in M$, there are disjoint open sets $\{\widetilde{U}_{b,x}^S\}_{b=1}^B$ such that $S : \widetilde{U}_{b,x}^S \to U_x$ is a \mathcal{C}^{r+1} diffeomorphism for each b and that for each $y \in U_x$,

$$d_M\left(\left(S|_{\widetilde{U}_{b,x}^S}\right)^{-1}(y), \left(T|_{\widetilde{U}_{b,x}}\right)^{-1}(y)\right) \to 0$$

$$\tag{42}$$

as $S \to T$ in $\mathcal{N}^{r+1}(M, M)$.

Since M is compact, there are a finite subfamily $\{U_{\lambda}\}_{\lambda \in \Lambda}$ (with $|\Lambda| < \infty$) of the open covering $\{U_x\}_{x \in M}$ of M. For each $\lambda \in \Lambda$ there are disjoint open sets $\{\widetilde{U}_{b,\lambda}\}_{b=1}^B$ such that $T: \widetilde{U}_{b,\lambda} \to U_{\lambda}$ is a \mathcal{C}^{r+1} diffeomorphism for each $b = 1, \ldots, B$. Notice that for each $\lambda \in \Lambda$, $x \in U_{\lambda}$ and a complex-valued function f on M, it holds that

$$\sum_{T(y)=x} f(y) = \sum_{b=1}^{B} f \circ \left(T|_{\widetilde{U}_{b,\lambda}} \right)^{-1} (x).$$
(43)

Let $\{K_{\lambda}\}_{\lambda \in \Lambda}$ be a closed covering of M such that $K_{\lambda} \subset U_{\lambda}$, and $\{\rho_{\lambda}\}_{\lambda \in \Lambda}$ a partition of unity of M subordinate to the covering $\{K_{\lambda}\}_{\lambda \in \Lambda}$ (that is, ρ_{λ} is a \mathcal{C}^{∞} function on M with values in $[0,1] \subset \mathbb{R}$ such that the support of ρ_{λ} is contained in K_{λ} for each $\lambda \in \Lambda$ and $\sum_{\lambda \in \Lambda} \rho_{\lambda}(x) = 1$ for each $x \in M$). Then, in view of (43) we get that for each $f \in \mathcal{C}^{r}(M)$ and $x \in M$,

$$\mathcal{L}_T f(x) = \sum_{\lambda \in \Lambda} \rho_\lambda(x) \cdot \sum_{T(y)=x} \frac{f(y)}{|\det DT(y)|} = \sum_{\lambda \in \Lambda} \sum_{b=1}^B \rho_\lambda(x) \cdot \frac{f}{|\det DT|} \circ \left(T|_{\widetilde{U}_{b,\lambda}}\right)^{-1}(x)$$

and for each $S \in \mathcal{U}$,

$$\left\|\mathcal{L}_{T}f - \mathcal{L}_{S}f\right\|_{\mathcal{C}^{r}} \leq \sum_{\lambda \in \Lambda} \sum_{b=1}^{B} \left\|\rho_{\lambda}\right\|_{\mathcal{C}^{r}} \left\|\frac{f}{|\det DT|} \circ \left(T|_{\widetilde{U}_{b,\lambda}}\right)^{-1} - \frac{f}{|\det DS|} \circ \left(S|_{\widetilde{U}_{b,\lambda}}\right)^{-1}\right\|_{\mathcal{C}^{r}(K_{\lambda})}.$$

Therefore, since both $|\Lambda|$ and B are finite and $||\det DT|^{-1} - |\det DS|^{-1}||_{\mathcal{C}^r} \to 0$ as $S \to T$ in $\mathcal{N}^{r+1}(M, M)$, it suffices to show that for each $\lambda \in \Lambda$, $b = 1, \ldots, B$ and $f \in \mathcal{C}^r(M)$,

$$\left\| f \circ \left(T|_{\widetilde{U}_{b,\lambda}} \right)^{-1} - f \circ \left(S|_{\widetilde{U}_{b,\lambda}^{S}} \right)^{-1} \right\|_{\mathcal{C}^{r}(K_{\lambda})} \to 0 \quad \text{as } S \to T \text{ in } \mathcal{N}^{r+1}(M,M).$$
(44)

Fix $\lambda \in \Lambda$, b = 1, ..., B and $f \in \mathcal{C}^{r}(M)$. By taking K_{λ} small if necessary, we can assume that K_{λ} is included in a local chart of M, so we assume that K_{λ} is a closed subset of \mathbb{R}^{d} with the dimension d of M. We use the notations $\partial_{i}(\cdot)$, $\partial^{\alpha}(\cdot)$ and $\operatorname{adj}(\cdot)$ given in the proof of Proposition 4.5. Recall that $T|_{\widetilde{U}_{b,\lambda}} : \widetilde{U}_{b,\lambda} \to U_{\lambda}$ is a \mathcal{C}^{r+1} diffeomorphism, so by the inverse function theorem and the fact that $A^{-1} = (\det A)^{-1}\operatorname{adj}(A)$ for any invertible matrix A, we have

$$D\left(\left(T|_{\widetilde{U}_{b,\lambda}}\right)^{-1}\right)(x) = DT(y) = (\det DT(y))^{-1}\operatorname{adj}(DT(y))$$

with $y = (T|_{\widetilde{U}_{b,\lambda}})^{-1}(x)$ for any $x \in U_{\lambda}$. Since each entry of $\operatorname{adj}(DT(y))$ (the transpose of the cofactor matrix of DT(y)) is a polynomial of $\partial_i T$ ($1 \le i \le d$), by the chain rule for derivatives we conclude that for each $i = 1, \ldots, d$,

$$\partial_i \left(f \circ \left(T |_{\widetilde{U}_{b,\lambda}} \right)^{-1} \right) = \sum_{l=1}^d \left(J_l \cdot \partial_l f \right) \circ \left(T |_{\widetilde{U}_{b,\lambda}} \right)^{-1} \quad \text{on } U_\lambda,$$

where J_l is a polynomial function of $\partial_i T_j$ $(1 \le i, j \le d)$ and $(\det DT)^{-1}$. Applying this formula repeatedly, we get that for each multi-index α with $|\alpha| \le r$,

$$\partial^{\alpha} \left(f \circ \left(T |_{\widetilde{U}_{b,\lambda}} \right)^{-1} \right) = \sum_{|\beta| \le |\alpha|} \left(J_{\alpha,\beta} \cdot \partial^{\beta} f \right) \circ \left(T |_{\widetilde{U}_{b,\lambda}} \right)^{-1} \quad \text{on } U_{\lambda};$$

where $J_{\alpha,\beta} = J_{\alpha,\beta}^T$ is a polynomial function of $\partial^{\gamma} T_j$ $(1 \leq j \leq d, |\gamma| \leq |\beta|)$ and $(\det DT)^{-1}$. Now (44) immediately follows from (42), and this completes the proof.

Acknowledgments

Y.N. was partially supported by JSPS KAKENHI Grant Numbers 19K14575 and 19K21834. H.C. is supported by an Australian Government Research Training Program Scholarship, a UNSW Science PhD Writing Scholarship, and by the UNSW School of Mathematics and Statistics.

References

- [1] L. ARNOLD, *Random dynamical systems*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 1998.
- W. BAHSOUN AND B. SAUSSOL, Linear response in the intermittent family: Differentiation in a weighted C⁰-norm, Discrete & Continuous Dynamical Systems-A, 36 (2016), p. 6657.

- [3] V. BALADI, Correlation spectrum of quenched and annealed equilibrium states for random expanding maps, Communications in mathematical physics, 186 (1997), pp. 671–700.
- [4] —, Positive transfer operators and decay of correlations, vol. 16, World scientific, 2000.
- [5] —, On the susceptibility function of piecewise expanding interval maps, Communications in mathematical physics, 275 (2007), pp. 839–859.
- [6] —, *Linear response, or else*, arXiv preprint arXiv:1408.2937, (2014).
- [7] <u>—</u>, Dynamical zeta functions and dynamical determinants for hyperbolic maps, Springer, 2018.
- [8] V. BALADI, A. KONDAH, AND B. SCHMITT, Random correlations for small perturbations of expanding maps, Random and Computational Dynamics, 4 (1996), pp. 179–204.
- [9] V. BALADI AND D. SMANIA, Linear response formula for piecewise expanding unimodal maps, Nonlinearity, 21 (2008), p. 677.
- [10] V. BALADI AND M. TODD, Linear response for intermittent maps, Communications in Mathematical Physics, 347 (2016), pp. 857–874.
- [11] T. BOGENSCHÜTZ, Stochastic stability of invariant subspaces, Ergodic Theory and Dynamical Systems, 20 (2000), pp. 663–680.
- [12] O. BUTTERLEY AND P. ESLAMI, Exponential mixing for skew products with discontinuities, Transactions of the American Mathematical Society, 369 (2017), pp. 783–803.
- [13] J. CHEN AND H. HU, The spectral gap for transfer operators of torus extensions over expanding maps, Nonlinearity, 32 (2018), p. 356.
- [14] Y. M. CHUNG, Y. NAKANO, AND J. WITTSTEN, Quenched limit theorems for random U(1) extensions of expanding maps, arXiv preprint arXiv:2104.01606, (2021).
- [15] J.-P. CONZE AND A. RAUGI, Limit theorems for sequential expanding dynamical systems, in Ergodic Theory and Related Fields: 2004-2006 Chapel Hill Workshops on Probability and Ergodic Theory, University of North Carolina Chapel Hill, North Carolina, vol. 430, American Mathematical Soc., 2007, p. 89.
- [16] H. CRIMMINS, Stability of hyperbolic oseledets splittings for quasi-compact operator cocycles, arXiv preprint arXiv:1912.03008, (2019).
- [17] D. DOLGOPYAT, On mixing properties of compact group extensions of hyperbolic systems, Israel J. Math., 130 (2002), pp. 157–205.
- [18] —, On differentiability of SRB states for partially hyperbolic systems, Inventiones mathematicae, 155 (2004), pp. 389–449.
- [19] D. DRAGIČEVIĆ, G. FROYLAND, C. GONZÁLEZ-TOKMAN, AND S. VAIENTI, A spectral approach for quenched limit theorems for random hyperbolic dynamical systems, Transactions of the American Mathematical Society, 373 (2020), pp. 629–664.
- [20] D. DRAGIČEVIĆ AND J. SEDRO, Statistical stability and linear response for random hyperbolic dynamics, arXiv preprint arXiv:2007.06088, (2020).
- [21] S. DYATLOV AND M. ZWORSKI, Stochastic stability of Pollicott-Ruelle resonances, Nonlinearity, 28 (2015), p. 3511.

- [22] F. FAURE, Semiclassical origin of the spectral gap for transfer operators of a partially expanding map, Nonlinearity, 24 (2011), p. 1473.
- [23] F. FAURE AND T. WEICH, Global normal form and asymptotic spectral gap for open partially expanding maps, Communications in Mathematical Physics, 356 (2017), pp. 755– 822.
- [24] G. FROYLAND, C. GONZÁLEZ-TOKMAN, AND A. QUAS, Stability and approximation of random invariant densities for Lasota-Yorke map cocycles, Nonlinearity, 27 (2014), p. 647.
- [25] S. GALATOLO, Statistical properties of dynamics. Introduction to the functional analytic approach, arXiv preprint arXiv:1510.02615, (2015).
- [26] S. GALATOLO AND J. SEDRO, Quadratic response of random and deterministic dynamical systems, Chaos: An Interdisciplinary Journal of Nonlinear Science, 30 (2020), p. 023113.
- [27] C. GONZÁLEZ-TOKMAN AND A. QUAS, A semi-invertible operator Oseledets theorem, Ergodic Theory and Dynamical Systems, 34 (2014), pp. 1230–1272.
- [28] —, Stability and Collapse of the Lyapunov spectrum for Perron-Frobenius Operator cocycles, arXiv preprint arXiv:1806.08873, (2018).
- [29] L. GOSSART, Flat trace statistics of the transfer operator of a random partially expanding map, Nonlinearity, 33 (2020), p. 3303.
- [30] S. GOUËZEL, Characterization of weak convergence of Birkhoff sums for Gibbs-Markov maps, Israel Journal of Mathematics, 180 (2010), pp. 1–41.
- [31] S. GOUËZEL AND C. LIVERANI, Corrigendum for "Banach spaces adapted to Anosov systems".
- [32] —, Banach spaces adapted to Anosov systems, Ergodic Theory and dynamical systems, 26 (2006), pp. 189–217.
- [33] P. A. GRIFFITHS AND J. A. WOLF, Complete maps and differentiable coverings., Michigan Mathematical Journal, 10 (1963), pp. 253–255.
- [34] B. HASSELBLATT, Ergodic Theory and Negative Curvature, Springer, 2017.
- [35] T. KATO, Perturbation theory for linear operators, Grundlehren der mathematischen Wissenschaften, Springer Berlin Heidelberg, 1966.
- [36] G. KELLER AND C. LIVERANI, Stability of the spectrum for transfer operators, Annali della Scuola Normale Superiore di Pisa-Classe di Scienze, 28 (1999), pp. 141–152.
- [37] A. KOREPANOV, Linear response for intermittent maps with summable and nonsummable decay of correlations, Nonlinearity, 29 (2016), p. 1735.
- [38] A. LASOTA AND M. C. MACKEY, Chaos, fractals, and noise: stochastic aspects of dynamics, vol. 97, Springer Science & Business Media, 2013.
- [39] C. LIVERANI, Invariant measures and their properties. a functional analytic point of view, Dynamical systems. Part II, (2004), pp. 185–237.
- [40] Y. NAKANO, Stochastic stability for fiber expanding maps via a perturbative spectral approach, Stochastics and Dynamics, 16 (2016), p. 1650011.

- [41] Y. NAKANO, M. TSUJII, AND J. WITTSTEN, The partial captivity condition for U(1)
- [42] Y. NAKANO AND J. WITTSTEN, On the spectra of quenched random perturbations of partially expanding maps on the torus, Nonlinearity, 28 (2015), p. 951.

extensions of expanding maps on the circle, Nonlinearity, 29 (2016), p. 1917.

- [43] W. PARRY AND M. POLLICOTT, Zeta functions and the periodic orbit structure of hyperbolic dynamics, Astérisque, 187 (1990), pp. 1–268.
- [44] D. RUELLE, Differentiation of SRB states, Communications in Mathematical Physics, 187 (1997), pp. 227–241.
- [45] <u>—, Nonequilibrium statistical mechanics near equilibrium: computing higher-order terms,</u> Nonlinearity, 11 (1998), p. 5.
- [46] J. SEDRO, A regularity result for fixed points, with applications to linear response, Nonlinearity, 31 (2018), p. 1417.
- [47] J. SEDRO AND H. H. RUGH, Regularity of characteristic exponents and linear response for transfer operator cocycles, arXiv preprint arXiv:2004.10103, (2020).