

ARITHMETIC VOLUMES OF UNITARY SHIMURA VARIETIES

JAN HENDRIK BRUINIER AND BENJAMIN HOWARD

ABSTRACT. The integral model of a $\mathrm{GU}(n-1, 1)$ Shimura variety carries a universal abelian scheme over it, and the dual top exterior power of its Lie algebra carries a natural hermitian metric. We express the arithmetic volume of this metrized line bundle, defined as an iterated self-intersection in the Gillet-Soulé arithmetic Chow ring, in terms of logarithmic derivatives of Dirichlet L -functions. We also determine the arithmetic volumes of Kudla-Rapoport divisors and relate them to coefficients of Eisenstein series.

CONTENTS

1. Introduction	1
2. Eisenstein series and theta functions	10
3. Automorphic Green functions	30
4. Integral models and arithmetic intersection theory	39
5. A Shimura variety and its line bundles	52
6. Kudla-Rapoport divisors at split primes	63
7. The generating series of arithmetic divisors	85
8. The volume calculations	97
Appendix A. Coefficients of modular forms	114
References	117

1. INTRODUCTION

The explicit calculation of arithmetic volumes of Shimura varieties began with the work of Kühn [Kuhn01] and Kramer (independently, in unpublished work). Over the moduli stack $\mathcal{X} \rightarrow \mathrm{Spec}(\mathbb{Z})$ of elliptic curves there is a line bundle of weight one modular forms, characterized as the dual Lie algebra of the universal elliptic curve. This line bundle carries a natural hermitian metric, and so determines a class in the codimension one arithmetic Chow of \mathcal{X} of Gillet-Soulé. Working on a suitable compactification, Kühn and

Date: August 1, 2023.

J.B. was supported in part by the LOEWE research unit USAG and by the DFG Collaborative Research Centre TRR 326 “Geometry and Arithmetic of Uniformized Structures”, project number 444845124. B.H. was supported in part by NSF grants DMS-1801905 and DMS-2101636.

Kramer computed the self-intersection multiplicity of this hermitian line bundle, and gave a simple formula for it in terms of the logarithmic derivative of the Riemann zeta function at $s = -1$.

More generally, if \mathcal{X} is an integral model of a PEL type (or even Hodge type) Shimura variety, then \mathcal{X} carries a *Hodge bundle*: the dual of the top exterior power of the Lie algebra of the universal abelian scheme over \mathcal{X} . The Hodge bundle again carries a natural hermitian metric, and one can ask if its arithmetic volume, defined as the $\dim(\mathcal{X})$ -fold self-intersection in the Gillet-Soulé arithmetic Chow group, is again related to logarithmic derivatives of Dirichlet L -functions.

In some instances this is known. If \mathcal{X} is the integral model of a quaternionic Shimura curve over \mathbb{Q} , the volume was computed by Kudla-Rapoport-Yang [KRY06]. Hilbert modular surfaces and the Siegel threefold were considered by Bruinier-Burgos-Kühn [BBK07] and Jung-von Pippich [JvP], respectively. The arithmetic volumes of $\mathrm{GSpin}(n, 2)$ Shimura varieties, which include all of the above examples as special cases, were computed (up to a \mathbb{Q} -linear combination of logarithms of certain bad primes) by Hörmann [Hor14].

Most recently, Yuan [Yuan22] extended the above mentioned result of Kudla-Rapoport-Yang (using completely different methods) to all quaternionic Shimura curves over totally real fields.

The present paper deals with the case of $\mathrm{GU}(n-1, 1)$ Shimura varieties. The overall strategy is closely related to that of [BBK07] and [Hor14], insofar as it ultimately rests on the use of Borcherds products. However, Borcherds products will make no explicit appearance in this paper; all the arithmetic information we need from the theory has already been extracted in [BHK⁺a], and is recalled here in §7.2.

1.1. The volume formulas. Let $\mathbf{k} \subset \mathbb{C}$ be an imaginary quadratic field of odd discriminant

$$-D = \mathrm{disc}(\mathbf{k}).$$

Given an integer $n \geq 1$, there is a regular Deligne-Mumford stack

$$\mathcal{M}_{(n-1,1)} \rightarrow \mathrm{Spec}(\mathcal{O}_{\mathbf{k}}),$$

flat of relative dimension $n-1$, parametrizing principally polarized abelian schemes A , endowed with an action $\mathcal{O}_{\mathbf{k}} \rightarrow \mathrm{End}(A)$, and extra data encoding a signature $(n-1, 1)$ condition on $\mathrm{Lie}(A)$. See §4.1 for details.

This stack admits a decomposition

$$\mathcal{M}_{(n-1,1)} = \bigsqcup_W \mathcal{M}_W$$

into open and closed substacks indexed by strict similarity classes of relevant \mathbf{k} -hermitian spaces (W, h) of signature $(n-1, 1)$, in such a way that the generic fiber of \mathcal{M}_W is a Shimura variety for the unitary similitude group $\mathrm{GU}(W)$. Here *relevant* means that W contains an $\mathcal{O}_{\mathbf{k}}$ -lattice $\mathfrak{a} \subset W$ that is self-dual, in the sense of (1.5.1). The notion of strict similarity is defined

in §4.1, but the important fact is that there is a dichotomy between the cases of n odd and n even. If n is odd the relevant W form a single strict similarity class, and the disjoint union has a single term. If n is even then strict similarity is equivalent to isometry, and the number of terms in the disjoint union is $2^{o(D)-1}$, where

$$o(D) = \#\{\text{prime divisors of } D\}.$$

Fix a relevant W as above, and denote by $\pi : A \rightarrow \mathcal{M}_W$ the restriction to \mathcal{M}_W of the universal abelian scheme over $\mathcal{M}_{(n-1,1)}$. Its relative dimension is $n = \dim(A)$, and its *metrized Hodge bundle*

$$\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}} \in \widehat{\text{Pic}}(\mathcal{M}_W)$$

is the line bundle $\omega_{A/\mathcal{M}_W}^{\text{Hdg}} = \pi_* \Omega_{A/\mathcal{M}_W}^{\dim(A)}$ endowed with the hermitian metric

$$(1.1.1) \quad \|s_z\|^2 = \left| \frac{1}{(2\pi i)^{\dim(A)}} \int_{A_z(\mathbb{C})} s_z \wedge \bar{s}_z \right|$$

for any $z \in \mathcal{M}_W(\mathbb{C})$ and $s_z \in \omega_{A/\mathcal{M}_W, z}^{\text{Hdg}} \cong H^0(A_z, \Omega_{A_z/\mathbb{C}}^{\dim(A_z)})$.

The stack \mathcal{M}_W has a canonical toroidal compactification $\bar{\mathcal{M}}_W$, with boundary a Cartier divisor smooth over \mathcal{O}_k . Although the Hodge bundle has a distinguished extension to the compactification, the metric on it does not extend smoothly. Instead, the metric is pre-log singular along the boundary, in the sense of Definition 1.20 of [BBK07]. The pre-log singular conditions also allow us to view

$$(1.1.2) \quad \widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}} \in \widehat{\text{CH}}^1(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}})$$

as a class in the codimension one arithmetic Chow group of Burgos-Kramer-Kühn [BKK07].

These arithmetic Chow groups come with intersection pairings, and distinguished linear functionals in codimensions $n-1$ and n . The first of these linear functionals is the *complex degree*, denoted

$$\text{deg}_{\mathbb{C}} : \widehat{\text{CH}}^{n-1}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}) \rightarrow \mathbb{Q},$$

defined by first sending an arithmetic cycle class to its image in the Chow group $\text{CH}^{n-1}(\bar{\mathcal{M}}_W/\mathbb{C})$ of 0-cycles on the complex fiber, and then taking the degree in the usual sense. The second is the more subtle *arithmetic degree*, denoted

$$\widehat{\text{deg}} : \widehat{\text{CH}}^n(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}) \rightarrow \mathbb{R}.$$

We define the *complex volume*

$$\text{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}) = \text{deg}_{\mathbb{C}} \left(\underbrace{\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}} \cdots \widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}}_{n-1 \text{ times}} \right)$$

and the *arithmetic volume*

$$\widehat{\text{vol}}(\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}) = \widehat{\text{deg}}(\underbrace{\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}} \cdots \widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}}_{n \text{ times}})$$

as the complex and arithmetic degrees of the $(n-1)$ -fold and n -fold iterated intersections, respectively.

Our main result is the calculation of these complex and arithmetic volumes in terms of Dirichlet L -functions. To state it, for any place $\ell \leq \infty$ denote by

$$\text{inv}_\ell(W) = (\det W, -D)_\ell \in \{\pm 1\}$$

the local invariant of W , where the right hand side is the usual Hilbert symbol. Our assumption that W contains a self-dual $\mathcal{O}_{\mathbf{k}}$ -lattice implies that $\text{inv}_\ell(W) = 1$ for all finite primes $\ell \nmid D$. As $\text{inv}_\infty(W) = -1$, and the product of all local invariants is 1, we obtain

$$(1.1.3) \quad \prod_{\ell|D} \text{inv}_\ell(W) = -1.$$

The quadratic Dirichlet character determined by \mathbf{k}/\mathbb{Q} is denoted

$$\varepsilon : (\mathbb{Z}/D\mathbb{Z})^\times \rightarrow \{\pm 1\}.$$

For an integer $k \geq 1$ set

$$(1.1.4) \quad \mathbf{a}_k(s) = \frac{D^{k/2} \Gamma(s+k) L(2s+k, \varepsilon^k)}{2^k \pi^{s+k}},$$

where, if k is even, we understand $L(s, \varepsilon^k) = \zeta(s)$. Define

$$(1.1.5) \quad \begin{aligned} \mathbf{A}_W(s) &= \mathbf{a}_1(s) \cdots \mathbf{a}_n(s) \\ &\times \begin{cases} \prod_{\ell|D} \left(1 + \left(\frac{-1}{\ell}\right)^{\frac{n}{2}} \text{inv}_\ell(W) \ell^{-s-\frac{n}{2}}\right) & \text{if } n \text{ is even} \\ 1 & \text{if } n \text{ is odd.} \end{cases} \end{aligned}$$

If n is odd there is no dependence on W beyond its dimension. We note that all $\mathbf{a}_k(0)$ are positive rational numbers, and hence so is $\mathbf{A}_W(0)$.

The following is our main result. It appears in the text as Theorem 8.6.1.

Theorem A. *The metrized Hodge bundle has complex volume*

$$\text{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}) = \mathbf{A}_W(0) \cdot \begin{cases} 2^{n-1} & \text{if } n \text{ is odd} \\ 2^{n-o(D)} & \text{if } n \text{ is even} \end{cases}$$

and arithmetic volume

$$\widehat{\text{vol}}(\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}) = \left(2 \frac{\mathbf{A}'_W(0)}{\mathbf{A}_W(0)} - n C_0(n) + \log(D) \right) \cdot \text{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}),$$

where we have set

$$C_0(n) = 2 \log \left(\frac{4\pi e^\gamma}{\sqrt{D}} \right) + (n-4) \left(\frac{L'(0, \varepsilon)}{L(0, \varepsilon)} + \frac{\log(D)}{2} \right).$$

Here $\gamma = -\Gamma'(1)$ is the Euler-Mascheroni constant.

Remark 1.1.1. Gross [Gro97] associates a motive to a reductive group over a field. The L -function of the motive associated to $U(W)$ is essentially $\mathbf{a}_1(s) \cdots \mathbf{a}_n(s)$, which perhaps provides some explanation for why this particular product appears in the statement of Theorem A.

Remark 1.1.2. There is a slightly different way to express the meromorphic function $\mathbf{A}_W(s)$ as a product. Define $\mathbf{b}_{W,1}(s) = \mathbf{a}_1(s)$. For $k \geq 2$ even, set

$$\mathbf{b}_{W,k}(s) = \mathbf{a}_k(s) \prod_{\ell|D} \left(1 + \left(\frac{-1}{\ell} \right)^{\frac{k}{2}} \text{inv}_\ell(W) \ell^{-s-\frac{k}{2}} \right).$$

For $k \geq 3$ odd, set

$$\mathbf{b}_{W,k}(s) = \mathbf{a}_k(s) \prod_{\ell|D} \left(1 + \left(\frac{-1}{\ell} \right)^{\frac{k-1}{2}} \text{inv}_\ell(W) \ell^{-s+\frac{1-k}{2}} \right)^{-1}.$$

When $k > 1$ we have $\mathbf{b}_{W,k}(s)\mathbf{b}_{W,k+1}(s) = \mathbf{a}_k(s)\mathbf{a}_{k+1}(s)$, which implies that

$$\mathbf{A}_W(s) = \mathbf{b}_{W,1}(s) \cdots \mathbf{b}_{W,n}(s).$$

1.2. The generating series of divisors. Most of the calculations that go into the proof of Theorem A are not carried out on the $\text{GU}(W)$ Shimura variety \mathcal{M}_W , but rather on a slightly different Shimura variety.

Let us again fix a relevant \mathbf{k} -hermitian space V of signature $(n-1, 1)$, and now assume $n \geq 2$. We may present this hermitian space in the form

$$V \cong \text{Hom}_{\mathbf{k}}(W_0, W),$$

where W_0 and W are relevant \mathbf{k} -hermitian space of signature $(1, 0)$ and $(n-1, 1)$, respectively (the right hand side carries a natural \mathbf{k} -hermitian form, and here we require that the isomorphism is an isometry). Following [KR14], we associate to V an open and closed substack

$$\mathcal{S}_V \subset \mathcal{M}_{W_0} \times_{\text{Spec}(\mathcal{O}_{\mathbf{k}})} \mathcal{M}_W.$$

As explained in §2 of [BHK⁺a], the generic fiber of \mathcal{S}_V is a Shimura variety for the subgroup $G \subset \text{GU}(W_0) \times \text{GU}(W)$ consisting of pairs for which the similitude factors are equal, and there is a natural surjection $G \rightarrow U(V)$. This allows one to view the moduli space \mathcal{S}_V as an approximation to the Shimura variety for the unitary group $U(V)$, which is awkward to work with directly because it is of abelian type but not PEL type.

In §5.1 we recall that the Shimura variety \mathcal{S}_V carries a family of Kudla-Rapoport divisors

$$\mathcal{Z}_V(m) \rightarrow \mathcal{S}_V$$

indexed by positive $m \in \mathbb{Z}$. It also carries a distinguished line bundle \mathcal{L}_V , called the *tautological bundle* or the *line bundle of weight one modular forms*, and an *exceptional divisor* $\text{Exc}_V \subset \mathcal{S}_V$ that is proper and supported in characteristics $\ell \mid D$.

The main result of [BHK⁺a] says that the Kudla-Rapoport divisors can be enhanced to arithmetic divisors

$$\widehat{\mathcal{Z}}_V^{\text{tot}}(m) = (\bar{\mathcal{Z}}_V(m) + \mathcal{B}_V(m), \Phi_V(m)) \in \widehat{\text{CH}}^1(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})$$

(for $m > 0$) in such a way that the formal generating series

$$\widehat{\phi}_V(\tau) = \sum_{m \geq 0} \widehat{\mathcal{Z}}_V^{\text{tot}}(m) \cdot q^m \in \widehat{\text{CH}}^1(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})[[q]]$$

is, at least when $n \geq 2$, a modular form. More precisely, it is the q -expansion of an element

$$\widehat{\phi}_V(\tau) \in M_n(\Gamma_0(D), \varepsilon^n) \otimes \widehat{\text{CH}}^1(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}}).$$

Here $\bar{\mathcal{S}}_V$ is the canonical toroidal compactification of \mathcal{S}_V , $\bar{\mathcal{Z}}_V(m) \subset \bar{\mathcal{S}}_V$ is the Zariski closure of $\mathcal{Z}_V(m)$, $\mathcal{B}_V(m)$ is an explicit linear combination of boundary components, and $\Phi_V(m)$ is a Green function for $\bar{\mathcal{Z}}_V(m) + \mathcal{B}_V(m)$ constructed using the theory of regularized theta lifts. The constant term is defined by first endowing \mathcal{L}_V with a particular pre-log singular hermitian metric, and then setting

$$\widehat{\mathcal{Z}}_V^{\text{tot}}(0) = -\widehat{\mathcal{L}}_V + (\text{Exc}_V, -\log(D)).$$

The second term on the right is the exceptional divisor endowed with the constant Green function $-\log(D)$.

Given that the above generating series is modular, the arithmetic divisor

$$(1.2.1) \quad \widehat{\mathcal{K}}_V \stackrel{\text{def}}{=} 2\widehat{\mathcal{L}}_V - (\text{Exc}_V, 0) \in \widehat{\text{CH}}^1(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})$$

determines a modular form

$$(1.2.2) \quad \widehat{\text{deg}}(\widehat{\phi}_V(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) = \sum_{m \geq 0} \widehat{\text{deg}}(\widehat{\mathcal{Z}}_V^{\text{tot}}(m) \cdot \widehat{\mathcal{K}}_V^{n-1}) \cdot q^m.$$

Here $\widehat{\mathcal{K}}_V^{n-1}$ is the $(n-1)$ -fold iterated arithmetic intersection. The peculiar choice of (1.2.1) is motivated by fact (Theorem 5.5.1) that it has trivial arithmetic intersection with every irreducible component of the exceptional divisor Exc_V , which simplifies the calculation of the coefficients of (1.2.2).

Conjecture B. *If $n > 2$, we have the equality of modular forms*

$$(1.2.3) \quad \widehat{\text{deg}}(\widehat{\phi}_V(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) \stackrel{?}{=} -\text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \sum_{r|D} \left(\frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \frac{3}{2} \log(D) - \log(r) \right) \gamma_r E_r(\tau),$$

in which

$$E_r(\tau) \in M_n(\Gamma_0(D), \varepsilon^n)$$

is the Eisenstein series of Proposition 2.7.1, and γ_r is the 4th root of unity (depending on V) defined in Proposition 2.7.3.

Conjecture B is stated in an equivalent form as Conjecture 8.4.1; the equivalence of the two statements follows from Corollary 8.5.2. Because the coefficients of the Eisenstein series $E_r(\tau)$ are given by explicit formulas, Conjecture B provides us with (conjectural) explicit formulas for the arithmetic intersection multiplicities appearing as the coefficients in (1.2.2). These can be found in §8.5, and especially (8.5.3).

The following is our main evidence for Conjecture B.

Theorem C. *If $n > 2$, the modular forms on either side of (1.2.3) have the same m^{th} Fourier coefficient for all positive $m \in \mathbb{Z}$ with $\varepsilon(m) = 1$. Moreover, their difference is a cusp form.*

The final claim of Theorem C allows us to equate the constant terms on both sides of (1.2.3), yielding

$$\widehat{\deg}(\widehat{\mathcal{Z}}_V^{\text{tot}}(0) \cdot \widehat{\mathcal{K}}_V^{n-1}) = -\text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \left(\frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \frac{3}{2} \log(D) \right).$$

This equality is equivalent to the second claim in the following theorem, which appears in the text as Theorems 8.2.3 and 8.4.2.

Theorem D. *The arithmetic divisor (1.2.1) has complex volume*

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) = \frac{|\text{CL}(\mathbf{k})|}{2^{\sigma(D)-n} |\mathcal{O}_{\mathbf{k}}^{\times}|} \cdot \mathbf{A}_V(0),$$

and arithmetic volume

$$\widehat{\text{vol}}(\widehat{\mathcal{K}}_V) = \left(2 \frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \log(D) \right) \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V).$$

In practice we do not deduce the second claim of Theorem D from the cuspidality claim of Theorem C. Instead, the proofs of the two statements are intertwined in a rather complicated way.

In the text, Theorem A is deduced from Theorem D using the relations between (1.1.2) and (1.2.1) found in Theorem 5.5.1. This latter theorem also allows us to deduce formulas for the complex and arithmetic volumes of $\widehat{\mathcal{L}}$ from Theorem D. These are found in Theorem 8.2.3 and Corollary 8.5.1, but the formula for the arithmetic volume is somewhat complicated by correction terms coming from the components of the exceptional divisor $\text{Exc}_V \subset \mathcal{S}_V$.

We end this subsection with some remarks about the restriction $\varepsilon(m) = 1$ in Theorem C. In the proof of that theorem, we first prove the equality for all primes p satisfying $\varepsilon(p) = 1$. The equality for all m with $\varepsilon(m) = 1$ then follows from a result of Bellaïche, found in Appendix A.

When $\varepsilon(p) = 1$, the strategy for the calculation of the coefficients of (1.2.2) is to first relate the Kudla-Rapoport divisor $\mathcal{Z}_V(p)$ to a Shimura variety $\mathcal{S}_{V^{\flat}}$ in one dimension lower, where $V^{\flat} \subset V$ is the orthogonal complement to a vector of hermitian norm p . This is done in §6. The assumption $\varepsilon(p) = 1$ guarantees that the hermitian space V^{\flat} is relevant (i.e. contains

a self-dual lattice), just as V was assumed to be, and so by induction one knows the analogue of Theorem D with V replaced by V^\flat . This formula for the arithmetic volume of $\widehat{\mathcal{K}}_{V^\flat}$ on $\widehat{\mathcal{S}}_{V^\flat}$ then gives a partial formula (Lemma 8.4.3) for the p^{th} coefficient in (1.2.2). Combining this partial formula with the modularity of the generating series $\widehat{\phi}_V(\tau)$ then proves Theorems C and D simultaneously.

In contrast, when $\varepsilon(p) = -1$ the Kudla-Rapoport divisor $\mathcal{Z}_V(p)$ should be related to the Shimura variety defined by a hermitian space V^\flat that is not relevant in the sense above. Thus carrying out the above inductive procedure when $\varepsilon(p) = -1$ would require working with Shimura varieties whose moduli interpretation involves non-principal polarizations, but the results of [BHK⁺a], and in particular the modularity of $\widehat{\phi}_V(\tau)$, are not known in this generality.

In summary, an extension of the modularity results of [BHK⁺a] to Shimura varieties associated to non-relevant hermitian spaces seems to be the missing ingredient needed to remove the restriction $\varepsilon(m) = 1$ in Theorem C.

1.3. Connections with the Kudla-Rapoport conjecture. Theorem D is closely related to a conjecture of Kudla-Rapoport [KR14] on the degrees of arithmetic 0-cycles on integral models of unitary Shimura varieties. Briefly, the Shimura variety \mathcal{S}_V is endowed not just with the family of Kudla-Rapoport divisors of the previous subsection, but with special cycles in all codimensions. In particular, it carries a family of arithmetic 0-cycles $\widehat{\mathcal{Z}}_V(T)$ indexed by $n \times n$ hermitian matrices T . The conjecture of Kudla-Rapoport predicts that the arithmetic degrees of these 0-cycles should agree with the Fourier coefficients of the derivative of an Eisenstein series on the quasi-split unitary group $U(n, n)$.

For those 0-cycles with $\det(T) \neq 0$, the Kudla-Rapoport conjecture is now a theorem of Li-Zhang [LZ19] and He-Li-Shi-Yang [HLSY22], who prove the unramified and ramified cases, respectively. The relevance of Theorem D is to the degenerate cases in which $\det(T) = 0$, and especially to the most degenerate case $T = 0$. In this case the associated arithmetic 0-cycle is, up to some correction factors at the primes $p \mid D$, the n -fold iterated intersection of the hermitian line bundle $\widehat{\mathcal{K}}_V$ (or $\widehat{\mathcal{L}}_V$) on \mathcal{S}_V appearing above. In other words, the Kudla-Rapoport conjecture predicts that the arithmetic volume of $\widehat{\mathcal{K}}_V$ is essentially the constant term of the derivative of an Eisenstein series.

We expect that the proof of Theorem A makes accessible more degenerate cases of the Kudla-Rapoport conjecture, beyond the most degenerate case $T = 0$. For example, consider a matrix of the form

$$T = \begin{pmatrix} T' & \\ & 0_{n-d} \end{pmatrix}$$

with T' a nonsingular $d \times d$ hermitian matrix with $m = \det(T')$ a squarefree product of primes split in \mathbf{k} . One could hope to relate the arithmetic degree

of $\widehat{\mathcal{Z}}_V(T)$ to the arithmetic intersection number

$$\widehat{\deg}(\widehat{\mathcal{Z}}_{V^b}^{\text{tot}}(m) \cdot \widehat{\mathcal{K}}_{V^b}^{n-d})$$

appearing in (1.2.2), where V^b is a hermitian space of signature $(n-d, 1)$. Theorem C might then be used to related these arithmetic degrees to degenerate coefficients of a $U(n, n)$ Eisenstein series.

We hope to return to these questions in future work.

1.4. Outline of the paper. Fix a \mathbf{k} -hermitian space V of signature $(n-1, 1)$, and a self-dual $\mathcal{O}_{\mathbf{k}}$ -lattice $L \subset V$.

We construct in §2 an Eisenstein series $E_L(\tau, s, n)$ of weight n , valued in a finite dimensional representation of $SL_2(\mathbb{Z})$. Its Fourier coefficients can be expressed in terms of local representation densities of the lattice L , which we then compute in order to make the coefficients completely explicit. Evaluating at the non-central point $s_0 = (n-1)/2$ yields a holomorphic Eisenstein series, which we express in terms of the Eisenstein series $E_r(\tau)$ appearing in Conjecture B.

In §3 we introduce a complex Shimura variety $\text{Sh}_K(H, \mathcal{D})$ associated to the unitary group $H = U(V)$. It carries a family of special divisors $Z(m)$ indexed by positive $m \in \mathbb{Z}$, and a metrized line bundle $\widehat{\mathcal{L}}$. To a harmonic Maass form f of weight $2-n$ we attach a divisor $Z(f)$, defined as a linear combination of $Z(m)$'s. We then construct a Green function $\Phi(f)$ for $Z(f)$ using the machinery of regularized theta lifts, and show in Theorem 3.4.2 that the normalized integral

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}})^{-1} \int_{\text{Sh}_K(H, \mathcal{D})} \Phi(f) \text{ch}(\widehat{\mathcal{L}})^{n-1}$$

can be expressed in terms of the Fourier coefficients of the derivative of $E_L(\tau, s, n)$ at $s = s_0$. The proof follows and generalizes ideas of Kudla [Ku03], who computed similar integrals on orthogonal Shimura varieties.

In §4 we work with \mathbf{k} -hermitian spaces W_0 and W of signatures $(1, 0)$ and $(n-1, 1)$. The Shimura varieties associated to their similitude groups have moduli interpretations, which can be used to construct regular integral models \mathcal{M}_{W_0} and \mathcal{M}_W . We recall the definition of these integral models, their toroidal compactifications, and the structure of their reductions at $p \mid D$. Then we recall those aspects of the Gillet-Soulé arithmetic intersection theory (as extended by Burgos-Kramer-Kuhn [BKK07]) that are needed in the sequel.

In §5 we choose W_0 and W so that $V \cong \text{Hom}_{\mathbf{k}}(W_0, W)$ as hermitian spaces. This allows us to define the open and closed substack

$$\mathcal{S}_V \subset \mathcal{M}_{W_0} \times_{\mathcal{O}_{\mathbf{k}}} \mathcal{M}_W$$

appearing in §1.2. It is related to the unitary Shimura variety of §3 by a finite cover

$$\mathcal{S}_V(\mathbb{C}) \rightarrow \text{Sh}_K(H, \mathcal{D}).$$

We define the Kudla-Rapoport divisors on \mathcal{S}_V , and the metrized line bundle $\widehat{\mathcal{L}}_V$. The universal abelian scheme over \mathcal{M}_W pulls back to an abelian scheme $A \rightarrow \mathcal{S}_V$, which determines a metrized Hodge bundle on \mathcal{S}_V . The main result of §5 is Theorem 5.5.1, which explains the precise connection between $\widehat{\mathcal{L}}_V$ and the metrized Hodge bundle. Going back and forth between these line bundles and (1.2.1), which for our purposes is the best behaved of the three, is essential to our methods.

In §6 we prove Theorems 6.1.2 and 6.1.4, which are essential to the induction arguments used to prove Theorems C and D. The idea is that if p is a prime split in \mathbf{k} , the Kudla-Rapoport divisor $\mathcal{Z}_V(p)$ should be closely related to a Shimura variety \mathcal{S}_{V^b} defined in the same way as \mathcal{S}_V , but with $\dim(V^b) = \dim(V) - 1$. This is true, up to some error terms coming from divisors supported in characteristics dividing pD . Using such a relation, we are able to express the height of the divisor $\mathcal{Z}_V(p)$ with respect to $\widehat{\mathcal{K}}_V$ in terms of the arithmetic volume of the metrized line bundle $\widehat{\mathcal{K}}_{V^b}$ on the lower-dimensional Shimura variety \mathcal{S}_{V^b} . Because of the many subtleties in the moduli problem defining the integral models, this is perhaps the most technical part of this work.

In §7 we recall the modularity results of [BHK⁺a]. We also prove Theorem 7.3.1, which can be summarized as saying that the boundary of the toroidal compactification $\bar{\mathcal{S}}_V$ can, for our purposes, be ignored.

In §8, we put everything together to prove the main results.

1.5. Notation and conventions. Throughout the paper, $\mathbf{k} \subset \mathbb{C}$ is a quadratic imaginary field of discriminant $-D$ whose associated quadratic character is denoted

$$\varepsilon : (\mathbb{Z}/D\mathbb{Z})^\times \rightarrow \{\pm 1\}.$$

The symbol V always denotes a \mathbf{k} -hermitian space of signature $(n-1, 1)$ with $n \geq 1$, and the hermitian form is denoted $\langle -, - \rangle$. Hermitian forms are always linear in the first variable and conjugate-linear in the second, and are assumed to be nondegenerate.

Beginning in §4, and continuing for the rest of the paper, we assume that D is odd and the hermitian space admits an $\mathcal{O}_{\mathbf{k}}$ -lattice $L \subset V$ that is *self-dual*, in the sense that

$$(1.5.1) \quad L = \{x \in V : \langle x, L \rangle \subset \mathcal{O}_{\mathbf{k}}\}.$$

These restrictions are not imposed in §2 and §3 unless stated explicitly.

The term *stack* means separated Deligne-Mumford stack.

1.6. Acknowledgements. The authors thank Joel Bellaïche for his help with Appendix A, and Dick Gross for pointing out Remark 1.1.1.

2. EISENSTEIN SERIES AND THETA FUNCTIONS

Fix a \mathbf{k} -hermitian space V of signature $(n-1, 1)$ with $n \geq 1$, and denote the hermitian form by $\langle -, - \rangle$. After reviewing some basics of Eisenstein

series, theta functions, and the Siegel-Weil formula, we attach a particular Eisenstein series to a lattice $L \subset V$, and express its Fourier coefficients in terms of representation densities. We then compute these representation densities in the case of a self-dual lattice, and so obtain explicit formulas for the Fourier coefficients; these formulas have a different shape depending on whether n is even or odd.

2.1. A seesaw dual reductive pair. Let V_0 be the unique symplectic space over \mathbb{Q} of dimension 2, and denote the symplectic form by $\langle x, y \rangle_0$. Set $V_{0\mathbf{k}} = V_0 \otimes_{\mathbb{Q}} \mathbf{k}$, and extend the symplectic form \mathbf{k} -linearly in the first argument and \mathbf{k} -conjugate-linearly in the second argument. This defines a skew-hermitian form on $V_{0\mathbf{k}}$.

The \mathbb{Q} -vector space underlying V carries a \mathbb{Q} -bilinear form

$$(2.1.1) \quad [x, y] = \mathrm{tr}_{\mathbf{k}/\mathbb{Q}} \langle x, y \rangle$$

of signature $(2n - 2, 2)$, with associated quadratic form

$$(2.1.2) \quad Q(x) = \langle x, x \rangle.$$

This data determine a seesaw dual reductive pair

$$\begin{array}{ccc} G = \mathrm{U}(V_{0\mathbf{k}}) & & H' = \mathrm{O}(V) \\ | & \searrow & | \\ G' = \mathrm{Sp}(V_0) & & H = \mathrm{U}(V). \end{array}$$

In particular, there are compatible Weil representations of $G(\mathbb{A}) \times H(\mathbb{A})$ and $G'(\mathbb{A}) \times H'(\mathbb{A})$ on the space $S(V(\mathbb{A}))$ of Schwartz-Bruhat functions on $V(\mathbb{A})$. This will be made explicit in §2.3.

Let e, f be a basis of V_0 with Gram matrix $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, and use this to identify

$$G \cong \left\{ g \in \mathrm{Res}_{\mathbf{k}/\mathbb{Q}} \mathrm{GL}_2 : g \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} {}^t \bar{g} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\}$$

and $G' \cong \mathrm{SL}_2$. Let $P = NM$ be the parabolic subgroup of G with Levi factor

$$M = \left\{ m(a) = \begin{pmatrix} a & 0 \\ 0 & \bar{a}^{-1} \end{pmatrix} : a \in \mathrm{Res}_{\mathbf{k}/\mathbb{Q}} \mathbb{G}_m \right\}$$

and unipotent radical

$$N = \left\{ n(b) = \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} : b \in \mathbb{G}_a \right\}.$$

For a place $p \leq \infty$ of \mathbb{Q} , let $K_p \subset G(\mathbb{Q}_p)$ be the maximal compact subgroup defined by

$$K_p = \begin{cases} G(\mathbb{Q}_p) \cap \mathrm{GL}_2(\mathcal{O}_{\mathbf{k},p}), & \text{if } p < \infty, \\ G(\mathbb{R}) \cap \mathrm{U}(2, \mathbb{R}), & \text{if } p = \infty, \end{cases}$$

and put $K = \prod_{p \leq \infty} K_p$. If we let \mathbf{k}^1 be the torus of norm one elements in $\mathbf{k}^\times = \text{Res}_{\mathbf{k}/\mathbb{Q}} \mathbb{G}_m$, there is a natural homomorphism

$$G' \times \mathbf{k}^1 \xrightarrow{(g,a) \mapsto g \cdot m(a)} G,$$

and the image of \mathbf{k}^1 is the center of G . This homomorphism induces an isomorphism

$$\text{SO}(2, \mathbb{R}) \times \text{U}(1, \mathbb{R}) / \{\pm 1\} \cong K_\infty.$$

Let \mathcal{H} be the complex upper half-plane. The first isomorphism in

$$\mathcal{H} \cong \text{SL}_2(\mathbb{R}) / \text{SO}(2, \mathbb{R}) \cong G(\mathbb{R}) / K_\infty.$$

sends $\tau = u + iv \in \mathcal{H}$ to the element

$$(2.1.3) \quad g_\tau = n(u)m(v^{1/2}) \in \text{SL}_2(\mathbb{R})$$

satisfying $g_\tau \cdot i = \tau$. The second is induced $\text{SL}_2(\mathbb{R}) = G'(\mathbb{R}) \subset G(\mathbb{R})$.

2.2. Eisenstein series. Given an $s \in \mathbb{C}$ and a character $\chi : \mathbb{A}_{\mathbf{k}}^\times / \mathbf{k}^\times \rightarrow \mathbb{C}^\times$, let

$$I(s, \chi) = \text{Ind}_{P(\mathbb{A})}^{G(\mathbb{A})} (\chi | \cdot |_{\mathbb{A}_{\mathbf{k}}}^s)$$

be the induced representation, realized on the space of smooth K -finite functions Φ on $G(\mathbb{A})$ satisfying

$$\Phi(n(b)m(a)g, s) = \chi(a) |a|_{\mathbb{A}_{\mathbf{k}}}^{s+\frac{1}{2}} \Phi(g, s)$$

for $b \in \mathbb{A}$ and $a \in \mathbb{A}_{\mathbf{k}}^\times$. Here $|a|_{\mathbb{A}_{\mathbf{k}}}$ denotes the norm on $\mathbb{A}_{\mathbf{k}}^\times$. In particular, at the archimedean place we have the normalized absolute value $|a|_\infty = a\bar{a}$.

Recall that a section of $I(s, \chi)$ is called *standard* if its restriction to K is independent on s . For a standard section Φ of $I(s, \chi)$, the associated Siegel Eisenstein series

$$E(g, s, \Phi) = \sum_{\gamma \in P(\mathbb{Q}) \backslash G(\mathbb{Q})} \Phi(\gamma g, s)$$

converges absolutely for $\text{Re}(s) > 1/2$, and defines an automorphic form on $G(\mathbb{A})$. The Eisenstein series has meromorphic continuation to all $s \in \mathbb{C}$. We now describe this Eisenstein series in more classical terms, by restricting it to the subgroup $\text{SL}_2 \cong G' \subset G$.

Assume that $\Phi(s) = \Phi_\infty(s) \otimes \Phi_f(s)$ is a factorizable standard section with $\Phi_f = \otimes_{p < \infty} \Phi_p(s)$, and that Φ_∞ is a normalized standard section of weight $\ell \in \mathbb{Z}$. In other words, $\Phi_\infty(1, s) = 1$, and for all

$$k = \begin{pmatrix} a & b \\ -b & a \end{pmatrix} \in \text{SO}(2, \mathbb{R}) \subset K_\infty$$

we have $\Phi_\infty(gk, s) = \Phi_\infty(g, s) \cdot \underline{k}^\ell$, where

$$(2.2.1) \quad \underline{k} = a + ib \in \mathbb{C}^\times.$$

Abbreviate $\Gamma = \text{SL}_2(\mathbb{Z}) \subset G(\mathbb{A})$, and let $\Gamma_\infty = P(\mathbb{Q}) \cap \Gamma$ be the subgroup of upper triangular matrices.

Lemma 2.2.1. *If Φ_∞ is a normalized standard section of weight $\ell \in \mathbb{Z}$ as above, then*

$$E(g_\tau, s, \Phi) = j(g_\tau, i)^{-\ell} \sum_{\gamma \in \Gamma_\infty \setminus \Gamma} \text{Im}(\gamma\tau)^{s+\frac{1}{2}-\frac{\ell}{2}} \cdot j(\gamma, \tau)^{-\ell} \cdot \Phi_f(\gamma).$$

Here $g_\tau \in \text{SL}_2(\mathbb{R})$ is as in (2.1.3), and $j(\gamma, \tau) = c\tau + d$ the usual automorphy factor associated to

$$\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{R}).$$

Proof. For $\gamma \in \Gamma$ we factor $\gamma g_\tau = n(\beta)m(\alpha)k$ with $\beta \in \mathbb{R}$, $\alpha \in \mathbb{R}^+$, and $k \in \text{SO}(2, \mathbb{R})$. Then a computation shows that

$$\alpha = \frac{v^{1/2}}{|c\tau + d|}, \quad \underline{k} = \frac{c\bar{\tau} + d}{|c\tau + d|},$$

and hence

$$\Phi_\infty(\gamma g_\tau, s) = v^{\ell/2} \cdot \frac{v^{s+\frac{1}{2}-\frac{\ell}{2}}}{|c\tau + d|^{2s+1-\ell}} \cdot (c\tau + d)^{-\ell}.$$

The inclusions $\Gamma \subset \text{SL}_2(\mathbb{Q}) \subset G(\mathbb{Q})$ induce bijections

$$\Gamma_\infty \setminus \Gamma \cong (P(\mathbb{Q}) \cap \text{SL}_2(\mathbb{Q})) \setminus \text{SL}_2(\mathbb{Q}) \cong P(\mathbb{Q}) \setminus G(\mathbb{Q}),$$

and hence

$$\begin{aligned} E(g_\tau, s, \Phi) &= \sum_{\gamma \in \Gamma_\infty \setminus \Gamma} \Phi_\infty(\gamma g_\tau, s) \Phi_f(\gamma) \\ &= v^{\ell/2} \sum_{\gamma \in \Gamma_\infty \setminus \Gamma} \text{Im}(\gamma\tau)^{s+\frac{1-\ell}{2}} \cdot (c\tau + d)^{-\ell} \cdot \Phi_f(\gamma) \\ &= j(g_\tau, i)^{-\ell} \sum_{\gamma \in \Gamma_\infty \setminus \Gamma} \text{Im}(\gamma\tau)^{s+\frac{1-\ell}{2}} \cdot j(\gamma, \tau)^{-\ell} \cdot \Phi_f(\gamma), \end{aligned}$$

as desired. \square

2.3. The Siegel-Weil formula. Write ψ for the standard additive character of \mathbb{A}/\mathbb{Q} , satisfying $\psi_\infty(x) = e^{2\pi ix}$. Let

$$\varepsilon_{\mathbb{A}} : \mathbb{A}^\times / \mathbb{Q}^\times \rightarrow \{\pm 1\}$$

be the idele class character determined by the quadratic extension \mathbf{k}/\mathbb{Q} , and fix a character

$$\chi : \mathbb{A}_{\mathbf{k}}^\times / \mathbf{k}^\times \rightarrow \mathbb{C}^\times$$

such that $\chi|_{\mathbb{A}^\times} = \varepsilon_{\mathbb{A}}^n$.

As explained in [HKS96] and [Ich07], the choices of ψ and χ determine a Weil representation $\omega = \omega_{\psi, \chi}$ of the group $G(\mathbb{A}) \times H(\mathbb{A})$ on the space of

Bruhat-Schwartz functions $S(V(\mathbb{A}))$. Recalling the \mathbb{Q} -bilinear form (2.1.1) and associated quadratic form (2.1.2), the action is given by the formulas

$$\begin{aligned}\omega(m(a))\varphi(x) &= \chi(a)|a|_{\mathbb{A}_k}^{n/2}\varphi(xa), \\ \omega(n(b))\varphi(x) &= \psi(bQ(x))\varphi(x), \\ \omega\left(\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}\right)\varphi(x) &= \int_{V(\mathbb{A})} \varphi(y)\psi([x, y]) dy, \\ \omega(h)\varphi(x) &= \varphi(h^{-1}x)\end{aligned}$$

for $\varphi \in S(V(\mathbb{A}))$ and $x \in V(\mathbb{A})$. Here $a \in \mathbb{A}_k^\times$, $b \in \mathbb{A}$, $h \in H(\mathbb{A})$, and the Fourier transform is taken with respect to the self-dual measure dy on $V(\mathbb{A})$. To a Schwartz function φ there is an associated theta function

$$(2.3.1) \quad \theta(g, h, \varphi) = \sum_{x \in V(\mathbb{Q})} \omega(g, h)\varphi(x).$$

Abbreviating $s_0 = (n - 1)/2$, there is a $G(\mathbb{A})$ -intertwining operator

$$\lambda : S(V(\mathbb{A})) \rightarrow I(s_0, \chi)$$

defined by $\lambda(\varphi)(g) = (\omega(g, 1)\varphi)(0)$. We extend $\lambda(\varphi)$ to a standard section of $I(s, \chi)$ by setting

$$\lambda(\varphi)(g, s) = |a(g)|_{\mathbb{A}_k}^{s-s_0} (\omega(g, 1)\varphi)(0).$$

The following theorem is a case of the Siegel-Weil formula; see [Weil65] or Theorem 1.1 of [Ich07].

Theorem 2.3.1. *Assume that $n > 2$, or that V is anisotropic. For $\varphi \in S(V(\mathbb{A}))$ and $g \in G(\mathbb{A})$ we have*

$$E(g, \lambda(\varphi), s_0) = \kappa \int_{H(\mathbb{Q}) \backslash H(\mathbb{A})} \theta(g, h, \varphi) dh,$$

where Haar measure is normalized by $\text{vol}(H(\mathbb{Q}) \backslash H(\mathbb{A})) = 1$, and

$$(2.3.2) \quad \kappa = \begin{cases} 1 & \text{if } n > 1 \\ 2 & \text{if } n = 1. \end{cases}$$

2.4. A special Eisenstein series. We consider Eisenstein series constructed from particular Schwartz functions on $V(\mathbb{A})$. At the archimedean place we take the Gaussian associated to a majorant and at the non-archimedean places we take the characteristic function of a coset of an \mathcal{O}_k -lattice.

Let $L \subset V$ be an \mathcal{O}_k -lattice on which the hermitian form is \mathcal{O}_k -valued. Its dual lattice under the \mathbb{Q} -bilinear form (2.1.1) is denoted

$$(2.4.1) \quad L' \stackrel{\text{def}}{=} \{x \in V : [x, L] \subset \mathbb{Z}\}.$$

For $\mu \in L'/L$ let

$$\varphi_\mu = \text{char}(\mu + \hat{L}) \in S(V(\mathbb{A}_f))$$

be the characteristic function of $\mu + \hat{L} \subset V(\mathbb{A}_f)$, and denote by S_L the subspace of $S(V(\mathbb{A}_f))$ generated by all φ_μ with $\mu \in L'/L$. Hence we may identify

$$S_L \cong \mathbb{C}[L'/L].$$

The restriction to $\Gamma = \mathrm{SL}_2(\mathbb{Z})$ of the Weil representation of $\mathrm{SL}_2(\mathbb{A}_f) \subset G(\mathbb{A}_f)$ takes the subspace S_L to itself, giving rise to a representation

$$(2.4.2) \quad \omega_L : \Gamma \rightarrow \mathrm{Aut}(S_L).$$

The corresponding dual representation

$$\omega_L^\vee : \Gamma \rightarrow \mathrm{Aut}(S_L^\vee)$$

is given by $\omega_L^\vee(\gamma)(f) = f \circ \omega_L^{-1}(\gamma)$ for $f \in S_L^\vee$. On the space S_L we also have the conjugate representation $\bar{\omega}_L$ defined by

$$\bar{\omega}_L(\gamma)(\varphi) = \overline{\omega_L(\gamma)(\bar{\varphi})}$$

for $\varphi \in S_L$. If we use the standard \mathbb{C} -bilinear pairing

$$(2.4.3) \quad \left\langle \sum_{\mu} a_{\mu} \varphi_{\mu}, \sum_{\mu} b_{\mu} \varphi_{\mu} \right\rangle = \sum_{\mu} a_{\mu} b_{\mu}$$

to identify $S_L \cong S_L^\vee$ as \mathbb{C} -vector spaces, then $\bar{\omega}_L = \omega_L^\vee$, and this agrees with the representation denoted ρ_L in [Bo98], [Br02], and [BY09].

For $\ell \in \mathbb{Z}$ and $\tau \in \mathcal{H}$, we define an S_L -valued Eisenstein series

$$(2.4.4) \quad E_L(\tau, s, \ell) = \sum_{\gamma \in \Gamma_\infty \backslash \Gamma} \mathrm{Im}(\gamma\tau)^{s + \frac{1-\ell}{2}} (c\tau + d)^{-\ell} \cdot \bar{\omega}_L(\gamma)^{-1} \varphi_0$$

of weight ℓ and representation $\bar{\omega}_L$. Note that this Eisenstein series has a slightly different normalization than the Eisenstein series in equation (2.17) of [BY09], since it is adapted to the unitary setting.

Recall the Maass lowering and raising operators in weight ℓ defined by

$$L_\ell = -2iv^2 \frac{\partial}{\partial \bar{\tau}}, \quad R_\ell = 2i \frac{\partial}{\partial \tau} + \ell v^{-1}.$$

They lower (respectively raise) the weight of an automorphic form by 2. It is easily seen that

$$(2.4.5) \quad \begin{aligned} L_\ell E_L(\tau, s, \ell) &= \left(s + \frac{1-\ell}{2} \right) E_L(\tau, s, \ell - 2) \\ R_\ell E_L(\tau, s, \ell) &= \left(s + \frac{1+\ell}{2} \right) E_L(\tau, s, \ell + 2). \end{aligned}$$

The Eisenstein series $E_L(\tau, s, \ell)$ is an eigenform of the hyperbolic Laplacian Δ_ℓ of weight ℓ (normalized as in (3.1) of [BF04]) with eigenvalue $(\frac{\ell-1}{2})^2 - s^2$. The following is a consequence of (2.4.5).

Lemma 2.4.1. *Assume that $n > 2$, or that V is anisotropic. The Eisenstein series $E_L(\tau, s_0, n)$ of weight $n = \dim(V)$ at $s_0 = (n - 1)/2$ is holomorphic in τ . Its derivative*

$$E'_L(\tau, s_0, n) \stackrel{\text{def}}{=} \frac{\partial}{\partial s} E'_L(\tau, s, n) \Big|_{s=s_0}$$

satisfies

$$L_n(E'_L(\tau, s_0, n)) = E_L(\tau, s_0, n - 2).$$

Remark 2.4.2. The Eisenstein series $E_L(\tau, s, n - 2)$ is coherent as it arises via the Siegel-Weil formula from the global hermitian space V . On the other hand, the Eisenstein series $E_L(\tau, s, n)$ is incoherent as it is associated to the incoherent collection of local hermitian spaces obtained by replacing V_∞ by the positive definite hermitian space of dimension n and keeping the non-archimedean part. In particular, at $s = 0$, the center of symmetry of the functional equation, the Eisenstein series $E_L(\tau, s, n)$ vanishes identically.

2.5. Coefficients of Eisenstein series. When the weight of the Eisenstein series (2.4.4) is $n = \dim(V)$, we denote its Fourier expansion by

$$(2.5.1) \quad E_L(\tau, s, n) = \sum_{\mu \in L'/L} \sum_{m \in \mathbb{Z} + \langle \mu, \mu \rangle} B(m, \mu; s, v) q^m \varphi_\mu,$$

where $v = \text{Im}(\tau)$ is the imaginary part of τ . We summarize some well-known facts about the coefficients; details can be found in [BrKu03] and [KY10].

If $m \neq 0$, there is a factorization

$$(2.5.2) \quad B(m, \mu; s, v) = B(m, \mu, s) \cdot \mathcal{W}_m(s, v)$$

in which the non-archimedean contribution $B(m, \mu, s)$ is independent of v . The second factor is the archimedean Whittaker function

$$(2.5.3) \quad \mathcal{W}_m(s, v) = (4\pi|m|v)^{-n/2} e^{2\pi m v} \cdot W_{\text{sgn}(m)\frac{n}{2}, s}(4\pi|m|v),$$

where

$$W_{\kappa, \mu}(z) = e^{-z/2} z^{1/2 + \mu} U\left(\frac{1}{2} + \mu - \kappa, 1 + 2\mu, z\right)$$

denotes the classical confluent hypergeometric function as in [AbSt84, Chapter 13], and

$$(2.5.4) \quad U(a, b, z) = \frac{1}{\Gamma(a)} \int_0^\infty e^{-zt} t^{a-1} (1+t)^{b-a-1} dt$$

for $\text{Re}(a) > 0$ and $|\arg(z)| < \pi/2$.

If $m = 0$, the Fourier coefficient has the form

$$B(0, \mu; s, v) = \delta_{\mu, 0} \cdot v^{s + \frac{1}{2} - \frac{n}{2}} + B(0, \mu, s) \cdot v^{-s + \frac{1}{2} - \frac{n}{2}},$$

where $B(0, \mu, s)$ is independent of v , and $\delta_{\mu, 0}$ is the Kronecker delta.

Remark 2.5.1. Note that our normalization of the archimedean Whittaker function in (2.5.3) differs from the normalization in [KY10, Proposition 2.3]. The normalizing factor $\Gamma(s + \frac{n}{2} + \frac{1}{2})^{-1}$ for $m > 0$, and $\Gamma(s - \frac{n}{2} + \frac{1}{2})^{-1}$ for $m < 0$, appears in loc. cit. In our case this factor is included in the normalization of the $B(m, \mu, s)$, since this leads to slightly cleaner formulas in Theorem 3.4.2.

The integral representation (2.5.4) can be used to describe the asymptotic behavior of the Whittaker function as $v \rightarrow \infty$, which is given by

$$\mathcal{W}_m(s, v) = \begin{cases} 1 + O(v^{-1}) & \text{if } m > 0 \\ O(e^{-4\pi|m|v}) & \text{if } m < 0, \end{cases}$$

locally uniformly in s and m .

We will also require the following lemma on the value at $s_0 = (n-1)/2$ of the Whittaker function and its derivative $\mathcal{W}'_m(s_0, v) = \frac{\partial}{\partial s} \mathcal{W}_m(s, v) |_{s=s_0}$.

Lemma 2.5.2. *The special value at s_0 of the Whittaker function is*

$$\mathcal{W}_m(s_0, v) = \begin{cases} 1 & \text{if } m > 0 \\ \Gamma(1-n, 4\pi|m|v) & \text{if } m < 0. \end{cases}$$

For $m > 0$ we have

$$\mathcal{W}'_m(s_0, v) = \sum_{j=1}^{n-1} \binom{n-1}{j} \frac{\Gamma(j)}{(4\pi mv)^j}.$$

In particular,

$$\mathcal{W}'_m(s_0, v) = O(v^{-1}), \quad \text{for } v \rightarrow \infty.$$

Proof. We only prove this for $m > 0$ and leave the similar case when $m < 0$ to the reader. According to (2.5.3) and (2.5.4), we have for $m > 0$ that

$$\begin{aligned} \mathcal{W}_m(s, v) &= \frac{(4\pi mv)^{s-s_0}}{\Gamma(s-s_0)} \cdot \int_0^\infty e^{-4\pi mvt} t^{s-s_0-1} (1+t)^{s+s_0} dt \\ &= 1 + \frac{(4\pi mv)^{s-s_0}}{\Gamma(s-s_0)} \cdot \int_0^\infty e^{-4\pi mvt} t^{s-s_0-1} ((1+t)^{s+s_0} - 1) dt, \end{aligned}$$

where the latter integral converges absolutely for $\text{Re}(s) > s_0 - 1$. We immediately see that $\mathcal{W}_m(s_0, v) = 1$, and in addition that

$$\begin{aligned} \mathcal{W}'_m(s_0, v) &= \int_0^\infty e^{-4\pi mvt} ((1+t)^{n-1} - 1) \frac{dt}{t} \\ &= \sum_{j=1}^{n-1} \binom{n-1}{j} \int_0^\infty e^{-4\pi mvt} t^j \frac{dt}{t} \\ &= \sum_{j=1}^{n-1} \binom{n-1}{j} \frac{\Gamma(j)}{(4\pi mv)^j}. \end{aligned}$$

This gives the claimed formula and the bound for $v \rightarrow \infty$. \square

We now turn to the explicit calculation of the factor $B(m, \mu, s)$ in (2.5.2), using formulas from [BrKu03] and [KY10]. Fix $\mu \in L'/L$, and let

$$d_\mu = \min\{b \in \mathbb{Z}^+ : b\mu = 0\}$$

be the order of μ . For $m \in \mathbb{Z} + Q(\mu)$ and $a \in \mathbb{Z}$, we denote by $N_{m,\mu}(a)$ the modulo a representation number

$$N_{m,\mu}(a) = \#\{r \in L/aL : Q(r + \mu) \equiv m \pmod{a\mathbb{Z}}\}.$$

For a prime p , set

$$(2.5.5) \quad L_{m,\mu}^{(p)}(p^{-s}) = (1 - p^{-s+2n-1}) \sum_{\nu=0}^{\infty} N_{m,\mu}(p^\nu) p^{-\nu s}$$

so that

$$\sum_{a=1}^{\infty} N_{m,\mu}(a) a^{-s} = \zeta(s - 2n + 1) \prod_p L_{m,\mu}^{(p)}(p^{-s}).$$

Proposition 2.5.3. *For nonzero $m \in \mathbb{Z} + Q(\mu)$ we have*

$$B(m, \mu, s) = -\frac{2^n \pi^{s + \frac{n}{2} + \frac{1}{2}}}{\sqrt{|L'/L|}} \cdot \begin{cases} \frac{|m|^{s + \frac{n}{2} - \frac{1}{2}}}{\Gamma(s + \frac{n}{2} + \frac{1}{2})} \prod_p L_{m,\mu}^{(p)}(p^{-2s-n}) & \text{if } m > 0 \\ \frac{|m|^{s + \frac{n}{2} - \frac{1}{2}}}{\Gamma(s - \frac{n}{2} + \frac{1}{2})} \prod_p L_{m,\mu}^{(p)}(p^{-2s-n}) & \text{if } m < 0. \end{cases}$$

Proof. This is obtained from Proposition 3.2 of [BrKu03] by noticing that our Eisenstein series $E_L(\tau, s, n)$ is $1/2$ the Eisenstein series $E_0(\tau, s - \frac{n}{2} + \frac{1}{2})$ of [BrKu03] with $\kappa = n$ and $r = 2n$. \square

According to equation (3.20) of [BrKu03], the Euler factors $L_{m,\mu}^{(p)}(p^{-s})$ are polynomials in p^{-s} . Moreover, by (3.23) of [BrKu03], if p does not divide $d_\mu^2 m |L'/L|$ then

$$L_{m,\mu}^{(p)}(p^{-s}) = 1 - \chi_F(p) p^{n-1-s},$$

where χ_F is the quadratic Dirichlet character associated to the quadratic extension $\mathbb{Q}(\sqrt{(-1)^n |L'/L|})$. This implies that the Euler product in Proposition 2.5.3 converges absolutely for $\text{Re}(s) > 0$ and that $B(m, \mu, s_0)$ is rational if $n > 1$ (so that $s_0 = \frac{n-1}{2}$ lies in the region of convergence). In this case it is nonpositive for positive m and vanishes for negative m .

Later we will require the following lemma on the behavior of the coefficients of the Eisenstein series in the limit $v \rightarrow \infty$.

Lemma 2.5.4. *Assume that $n > 2$, or that V is anisotropic. The coefficients $B'(m, \mu; s_0, v)$ of $E'_L(\tau, s_0, n)$ satisfy*

$$\lim_{v \rightarrow \infty} (B'(m, \mu; s_0, v) - \kappa \delta_{m,0} \delta_{\mu,0} \log v) = \begin{cases} B'(m, \mu, s_0) & \text{if } m > 0 \\ B'(0, \mu, s_0) & \text{if } m = 0 \text{ and } n = 1 \\ 0 & \text{if } m = 0 \text{ and } n > 1 \\ 0 & \text{if } m < 0. \end{cases}$$

Proof. For $m \neq 0$ the assertion follows from Proposition 2.5.3 together with Lemma 2.5.2 on the asymptotic behavior of the Whittaker function $\mathcal{W}_m(s, v)$ and its derivative at $s = s_0$.

For $m = 0$, we consider the Laurent expansion of $B(0, \mu; s, v)$ at $s = s_0$, which is given by

$$\begin{aligned} B(0, \mu; s, v) &= \delta_{\mu,0} + B(0, \mu, s_0)v^{-2s_0} \\ &\quad + (\delta_{\mu,0} \log(v) - B(0, \mu, s_0) \log(v)v^{-2s_0} + B'(0, \mu, s_0)v^{-2s_0})(s - s_0) \\ &\quad + O((s - s_0)^2). \end{aligned}$$

If $n > 1$, then the holomorphicity of $E_L(\tau, s_0, n)$ implies that $B(0, \mu, s_0) = 0$. Since $2s_0 > 0$, the term involving v^{-2s_0} vanishes in the limit $v \rightarrow \infty$, and we obtain the assertion.

If $n = 1$ then $s_0 = 0$ is the center of symmetry of the functional equation of the incoherent Eisenstein series, and hence $E_L(\tau, 0, n)$ vanishes. The vanishing of the constant Fourier coefficient implies

$$B(0, \mu, 0) = -\delta_{\mu,0},$$

and consequently

$$B'(0, \mu; 0, v) = 2\delta_{\mu,0} \log(v) + B'(0, \mu, s_0).$$

This concludes the proof of the lemma. \square

2.6. Self-dual lattices. In this subsection we assume the $\mathcal{O}_{\mathbf{k}}$ -lattice $L \subset V$ is self-dual (1.5.1) under the hermitian form, and that $D = -\text{disc}(\mathbf{k})$ is odd. The first of these assumptions implies that (2.4.1) satisfies $L' = \mathfrak{d}_{\mathbf{k}}^{-1}L$, where $\mathfrak{d}_{\mathbf{k}} \subset \mathcal{O}_{\mathbf{k}}$ is the different. We will determine the coefficient $B(m, \mu, s)$ from (2.5.2) more explicitly.

To compute the Euler factors $L_{m,\mu}^{(p)}(p^{-s})$ in Proposition 2.5.3, we need to determine the representation numbers $N_{m,\mu}(p^\nu)$. We now derive explicit formulas for these quantities using finite Fourier transforms and formulas for certain lattice Gauss sums. As we will only compute the coefficients for $\mu = 0$, we abbreviate $N_m(p^\nu) = N_{m,0}(p^\nu)$.

Fix an $a \in \mathbb{Z}$. For $c \in \mathbb{Z}^+$, consider the Gauss sum

$$G(a, c) = \sum_{x \in \mathcal{O}_{\mathbf{k}}/c\mathcal{O}_{\mathbf{k}}} e\left(\frac{aN(x)}{c}\right).$$

We are mainly interested in the case when $c = p^\nu$ is a prime power.

Lemma 2.6.1. *Let p^ν be a prime power. If we write $\gcd(a, p^\nu) = p^\alpha$ with $0 \leq \alpha \leq \nu$ and $a = p^\alpha a'$ with $a' \in \mathbb{Z}$, then*

$$G(a, p^\nu) = \begin{cases} p^{2\nu} & \text{if } \alpha = \nu \\ p^{\alpha+\nu} \left(\frac{-D}{p}\right)^{\nu-\alpha} & \text{if } \alpha < \nu \text{ and } p \nmid D \\ \epsilon_p p^{\alpha+\nu+\frac{1}{2}} \left(\frac{a'}{p}\right) \left(\frac{D'}{p}\right)^{\nu-\alpha-1} & \text{if } \alpha < \nu \text{ and } p \mid D. \end{cases}$$

Here, in the final case, we have set $D' = D/p$ and

$$\epsilon_r = \begin{cases} 1 & \text{if } r \equiv 1 \pmod{4} \\ i & \text{if } r \equiv 3 \pmod{4}. \end{cases}$$

for any odd positive integer r .

Proof. It is easily seen that

$$G(a, p^\nu) = p^{2\alpha} G(a', p^{\nu-\alpha}).$$

Hence we may assume that a is coprime to p . Now, if p is odd, the assertion can be easily deduced from the classical Gauss sum

$$\sum_{x \in \mathbb{Z}/c\mathbb{Z}} e\left(\frac{ax^2}{c}\right) = \epsilon_c \sqrt{c} \left(\frac{a}{c}\right)$$

for odd positive c with $\gcd(c, a) = 1$.

For $p = 2$ and a odd we have

$$(2.6.1) \quad G(a, 2^\nu) = \sum_{x, y \in \mathbb{Z}/2^\nu\mathbb{Z}} e\left(a \frac{x^2 + xy + \frac{1+D}{4}y^2}{2^\nu}\right),$$

where we have used that D is odd. If $\nu = 1$ the asserted formula is easily checked. Hence, assume that $\nu \geq 2$. Then we use the classical Gauss sum

$$\sum_{x \in \mathbb{Z}/2^\nu\mathbb{Z}} e\left(a \frac{x^2 + xy}{2^\nu}\right) = \begin{cases} 0, & 2 \nmid y, \\ e\left(-a \frac{y^2}{2^{\nu+2}}\right) \sum_{x \in \mathbb{Z}/2^\nu\mathbb{Z}} e\left(a \frac{x^2}{2^\nu}\right), & 2 \mid y, \end{cases}$$

to rewrite (2.6.1) as follows:

$$G(a, 2^\nu) = \frac{1}{2} \sum_{x, y \in \mathbb{Z}/2^\nu\mathbb{Z}} e\left(a \frac{x^2 + Dy^2}{2^\nu}\right).$$

Inserting the evaluation of the Gauss sum

$$\sum_{x \in \mathbb{Z}/2^\nu\mathbb{Z}} e\left(a \frac{x^2}{2^\nu}\right) = (1+i)\epsilon_a^{-1} 2^{\nu/2} \left(\frac{2^\nu}{a}\right),$$

we finally find that

$$G(a, 2^\nu) = 2^\nu \left(\frac{2^\nu}{D}\right) = 2^\nu \left(\frac{-D}{2}\right)^\nu,$$

concluding the proof of the lemma. \square

Lemma 2.6.2. *Let p^ν be a prime power. If we write $\gcd(a, p^\nu) = p^\alpha$ with $0 \leq \alpha \leq \nu$ and $a = p^\alpha a'$ with $a' \in \mathbb{Z}$, then the Gauss sum*

$$G_L(a, p^\nu) = \sum_{x \in L/p^\nu L} e\left(\frac{a\langle x, x \rangle}{p^\nu}\right)$$

is given by

$$G_L(a, p^\nu) = \begin{cases} p^{2n\nu} & \text{if } \alpha = \nu \\ p^{n\alpha+n\nu} \left(\frac{-D}{p}\right)^{n(\nu-\alpha)} & \text{if } \alpha < \nu \text{ and } p \nmid D \\ \epsilon_p^n \operatorname{inv}_p(V) p^{n\alpha+n\nu+\frac{n}{2}} \left(\frac{a'}{p}\right)^n \left(\frac{D'}{p}\right)^{n(\nu-\alpha-1)} & \text{if } \alpha < \nu \text{ and } p \mid D, \end{cases}$$

where ϵ_p and D' have the same meaning as in Lemma 2.6.1.

Proof. The Gauss sum $G_L(a, p^\nu)$ only depends on $L_p = L \otimes_{\mathbb{Z}} \mathbb{Z}_p$. If we let b_1, \dots, b_n be an orthogonal $\mathcal{O}_{\mathbf{k}, p}$ -module basis of L_p (which exists because of our hypothesis that D is odd), then

$$G_L(a, p^\nu) = G(a\langle b_1, b_1 \rangle, p^\nu) \cdots G(a\langle b_n, b_n \rangle, p^\nu),$$

and the claim follows from Lemma 2.6.1. Note that in the case $p \mid D$, standard formulas for the Hilbert symbol imply

$$\operatorname{inv}_p(V) = \left(\frac{\langle b_1, b_1 \rangle}{p}\right) \cdots \left(\frac{\langle b_n, b_n \rangle}{p}\right). \quad \square$$

Proposition 2.6.3. *Assume that n is even, and $m \in \mathbb{Z}$ is nonzero. Fix a prime p , and factor $m = p^\beta m'$ with $\gcd(m', p) = 1$. If $p \nmid D$ then*

$$L_m^{(p)}(p^{-s}) = (1 - p^{n-s-1}) \sum_{\gamma=0}^{\beta} p^{(n-s)\gamma}.$$

If $p \mid D$ then

$$\begin{aligned} L_m^{(p)}(p^{-s}) &= 1 - \left(\frac{-1}{p}\right)^{\frac{n}{2}} \operatorname{inv}_p(V) p^{\frac{n}{2}} p^{n-1-s} \\ &\quad + \left(\frac{-1}{p}\right)^{\frac{n}{2}} \operatorname{inv}_p(V) p^{\frac{n}{2}} (1 - p^{n-1-s}) \sum_{\gamma=1}^{\beta} p^{(n-s)\gamma}. \end{aligned}$$

Proof. We compute the representation number $N_m(p^\nu)$ using the identity

$$\begin{aligned} N_m(p^\nu) &= \frac{1}{p^\nu} \sum_{a \in \mathbb{Z}/p^\nu \mathbb{Z}} G_L(a, p^\nu) e\left(-\frac{am}{p^\nu}\right) \\ &= \frac{1}{p^\nu} \sum_{\alpha=0}^{\nu} \sum_{a' \in (\mathbb{Z}/p^{\nu-\alpha} \mathbb{Z})^\times} G_L(p^\alpha a', p^\nu) e\left(-\frac{a'm}{p^{\nu-\alpha}}\right). \end{aligned}$$

As n is even, Lemma 2.6.2 implies

$$G_L(a, p^\nu) = \begin{cases} p^{2n\nu} & \text{if } \alpha = \nu \\ p^{n\alpha+n\nu} & \text{if } \alpha < \nu \text{ and } p \nmid D \\ \left(\frac{-1}{p}\right)^{n/2} \operatorname{inv}_p(V) p^{n\alpha+n\nu+\frac{n}{2}} & \text{if } \alpha < \nu \text{ and } p \mid D. \end{cases}$$

If $p \nmid D$ we obtain

$$\begin{aligned}
N_m(p^\nu) &= p^{n\nu-\nu} \sum_{\alpha=0}^{\nu} p^{n\alpha} \sum_{a' \in (\mathbb{Z}/p^{\nu-\alpha}\mathbb{Z})^\times} e\left(-\frac{a'm}{p^{\nu-\alpha}}\right) \\
&= p^{n\nu-\nu} \sum_{\alpha=0}^{\nu} p^{n\alpha} \sum_{d|\gcd(p^{\nu-\alpha}, m)} \mu\left(\frac{p^{\nu-\alpha}}{d}\right) d \\
&= p^{n\nu-\nu} \sum_{\alpha=0}^{\nu} p^{n\alpha} \sum_{\gamma=0}^{\min(\nu-\alpha, \beta)} \mu(p^{\nu-\alpha-\gamma}) p^\gamma.
\end{aligned}$$

Here we have used the evaluation of the Ramanujan sum. Consequently, the Euler factor (2.5.5) is

$$\begin{aligned}
L_m^{(p)}(p^{-s}) &= (1 - p^{2n-s-1}) \sum_{\nu=0}^{\infty} N_m(p^\nu) p^{-\nu s} \\
&= (1 - p^{2n-s-1}) \sum_{\gamma=0}^{\beta} \sum_{\alpha=0}^{\infty} \sum_{\nu=\alpha+\gamma}^{\infty} p^{n\alpha+\gamma} \mu(p^{\nu-\alpha-\gamma}) p^{(n-1-s)\nu} \\
&= (1 - p^{n-s-1}) \sum_{\gamma=0}^{\beta} p^{(n-s)\gamma}.
\end{aligned}$$

In the case $p \mid D$ we find

$$\begin{aligned}
N_m(p^\nu) &= p^{2n\nu-\nu} + \left(\frac{-1}{p}\right)^{n/2} \operatorname{inv}_p(V) p^{n\nu-\nu+\frac{n}{2}} \sum_{\alpha=0}^{\nu-1} p^{n\alpha} \sum_{a' \in (\mathbb{Z}/p^{\nu-\alpha}\mathbb{Z})^\times} e\left(-\frac{a'm}{p^{\nu-\alpha}}\right) \\
&= p^{2n\nu-\nu} + \left(\frac{-1}{p}\right)^{n/2} \operatorname{inv}_p(V) p^{n\nu-\nu+\frac{n}{2}} \sum_{\alpha=0}^{\nu-1} p^{n\alpha} \sum_{d|\gcd(p^{\nu-\alpha}, m)} \mu\left(\frac{p^{\nu-\alpha}}{d}\right) d \\
&= p^{2n\nu-\nu} + \left(\frac{-1}{p}\right)^{n/2} \operatorname{inv}_p(V) p^{n\nu-\nu+\frac{n}{2}} \sum_{\alpha=0}^{\nu-1} p^{n\alpha} \sum_{\gamma=0}^{\min(\nu-\alpha, \beta)} \mu(p^{\nu-\alpha-\gamma}) p^\gamma.
\end{aligned}$$

Consequently, the Euler factor $L_m^{(p)}(p^{-s})$ of (2.5.5) is

$$\begin{aligned}
&(1 - p^{2n-s-1}) \sum_{\nu=0}^{\infty} N_m(p^\nu) p^{-\nu s} \\
&= 1 + \left(\frac{-1}{p}\right)^{\frac{n}{2}} \operatorname{inv}_p(V) p^{\frac{n}{2}} (1 - p^{2n-s-1}) \sum_{\nu=1}^{\infty} p^{(n-1-s)\nu} \sum_{\alpha=0}^{\nu-1} \sum_{\gamma=0}^{\min(\nu-\alpha, \beta)} p^{n\alpha+\gamma} \mu(p^{\nu-\alpha-\gamma}) \\
&= 1 + \left(\frac{-1}{p}\right)^{\frac{n}{2}} \operatorname{inv}_p(V) p^{\frac{n}{2}} (1 - p^{2n-s-1}) \sum_{\gamma=0}^{\beta} \sum_{\alpha=0}^{\infty} \sum_{\substack{\nu \geq \alpha+\gamma \\ \nu \geq \alpha+1}} p^{(n-1-s)\nu} p^{n\alpha+\gamma} \mu(p^{\nu-\alpha-\gamma}).
\end{aligned}$$

The contribution of $\gamma = 0$ to the latter sum is given by

$$(1 - p^{2n-s-1}) \sum_{\alpha=0}^{\infty} \sum_{\nu \geq \alpha+1} p^{(n-1-s)\nu} p^{n\alpha} \mu(p^{\nu-\alpha}) = -p^{n-1-s}.$$

The contribution of $\gamma \geq 1$ is equal to

$$\begin{aligned} & (1 - p^{2n-s-1}) \sum_{\gamma=1}^{\beta} \sum_{\alpha=0}^{\infty} \sum_{\nu \geq \alpha+\gamma} p^{n\alpha+\gamma} p^{(n-1-s)\nu} \mu(p^{\nu-\alpha-\gamma}) \\ &= (1 - p^{2n-s-1}) \sum_{\gamma=1}^{\beta} \sum_{\alpha=0}^{\infty} p^{n\alpha+\gamma} \left(p^{(n-1-s)(\alpha+\gamma)} - p^{(n-1-s)(\alpha+\gamma+1)} \right) \\ &= (1 - p^{n-1-s}) \sum_{\gamma=1}^{\beta} p^{(n-s)\gamma}. \end{aligned}$$

Putting the terms together, we find

$$\begin{aligned} L_m^{(p)}(p^{-s}) &= 1 - \left(\frac{-1}{p} \right)^{\frac{n}{2}} \operatorname{inv}_p(V) p^{\frac{n}{2}} p^{n-1-s} \\ &\quad + \left(\frac{-1}{p} \right)^{\frac{n}{2}} \operatorname{inv}_p(V) p^{\frac{n}{2}} (1 - p^{n-1-s}) \sum_{\gamma=1}^{\beta} p^{(n-s)\gamma}. \end{aligned}$$

This concludes the proof of the proposition. \square

Proposition 2.6.4. *Assume n is odd, and $m \in \mathbb{Z}$ is nonzero. Fix a prime p , and factor $m = p^\beta m'$ with $\gcd(m', p) = 1$. If $p \nmid D$ then*

$$L_m^{(p)}(p^{-s}) = \left(1 - \left(\frac{-D}{p} \right) p^{n-s-1} \right) \sum_{\gamma=0}^{\beta} \left(\frac{-D}{p} \right)^{\gamma} p^{(n-s)\gamma}.$$

If $p \mid D$ then

$$L_m^{(p)}(p^{-s}) = 1 + \operatorname{inv}_p(V) \left(\frac{-1}{p} \right)^{\frac{n-1}{2}} \left(\frac{D'}{p} \right)^{\beta} \left(\frac{m'}{p} \right) p^{(\beta+3/2)n-1/2-(\beta+1)s},$$

where we have set $D' = D/p$.

Proof. We argue in the same way as in the proof of Proposition 2.6.3, using the identity

$$N_m(p^\nu) = \frac{1}{p^\nu} \sum_{a \in \mathbb{Z}/p^\nu \mathbb{Z}} G_L(a, p^\nu) e\left(-\frac{am}{p^\nu}\right)$$

and the value of the Gauss sum $G_L(a, p^\nu)$ computed in Lemma 2.6.2. If we write $a = p^\alpha a'$ with a' coprime to p , we have

$$G_L(a, p^\nu) = \begin{cases} p^{2n\nu} & \text{if } \alpha = \nu \\ p^{n\alpha+n\nu} \left(\frac{-D}{p}\right)^{\nu-\alpha} & \text{if } \alpha < \nu \text{ and } p \nmid D \\ \epsilon_p^n \operatorname{inv}_p(V) p^{n\alpha+n\nu+\frac{n}{2}} \left(\frac{a'}{p}\right) \left(\frac{D'}{p}\right)^{\nu-\alpha-1} & \text{if } \alpha < \nu \text{ and } p \mid D. \end{cases}$$

If $p \nmid D$ we obtain

$$\begin{aligned} N_m(p^\nu) &= p^{n\nu-\nu} \sum_{\alpha=0}^{\nu} p^{n\alpha} \left(\frac{-D}{p}\right)^{\nu-\alpha} \sum_{a' \in (\mathbb{Z}/p^{\nu-\alpha}\mathbb{Z})^\times} e\left(-\frac{a'm}{p^{\nu-\alpha}}\right) \\ &= p^{n\nu-\nu} \sum_{\alpha=0}^{\nu} p^{n\alpha} \left(\frac{-D}{p}\right)^{\nu-\alpha} \sum_{d \mid \gcd(p^{\nu-\alpha}, m)} \mu\left(\frac{p^{\nu-\alpha}}{d}\right) d \\ &= p^{n\nu-\nu} \sum_{\alpha=0}^{\nu} p^{n\alpha} \left(\frac{-D}{p}\right)^{\nu-\alpha} \sum_{\gamma=0}^{\min(\nu-\alpha, \beta)} \mu(p^{\nu-\alpha-\gamma}) p^\gamma. \end{aligned}$$

Here we have used the evaluation of the Ramanujan sum. Hence, the Euler factor (2.5.5) is

$$\begin{aligned} L_m^{(p)}(p^{-s}) &= (1 - p^{2n-s-1}) \sum_{\nu=0}^{\infty} N_m(p^\nu) p^{-\nu s} \\ &= (1 - p^{2n-s-1}) \sum_{\gamma=0}^{\beta} \sum_{\alpha=0}^{\infty} \sum_{\nu=\alpha+\gamma}^{\infty} p^{n\alpha+\gamma} \left(\frac{-D}{p}\right)^{\nu-\alpha} \mu(p^{\nu-\alpha-\gamma}) p^{(n-1-s)\nu} \\ &= \left(1 - \left(\frac{-D}{p}\right) p^{n-s-1}\right) \sum_{\gamma=0}^{\beta} \left(\frac{-D}{p}\right)^{\gamma} p^{(n-s)\gamma}. \end{aligned}$$

When $p \mid D$ we find

$$\begin{aligned} N_m(p^\nu) &= p^{2n\nu-\nu} + \epsilon_p^n \operatorname{inv}_p(V) p^{n\nu-\nu+\frac{n}{2}} \\ &\quad \times \sum_{\alpha=0}^{\nu-1} \left(\frac{D'}{p}\right)^{\nu-\alpha-1} p^{n\alpha} \sum_{a' \in (\mathbb{Z}/p^{\nu-\alpha}\mathbb{Z})^\times} \left(\frac{a'}{p}\right) e\left(-\frac{a'm}{p^{\nu-\alpha}}\right). \end{aligned}$$

The latter Gauss sum is equal to $p^{\nu-\alpha-\frac{1}{2}} \epsilon_p \left(\frac{-m'}{p}\right)$ if $\beta = \nu - \alpha - 1$ and is zero otherwise. Inserting this we find that $N_m(p^\nu)$ is equal to

$$p^{2n\nu-\nu} \cdot \begin{cases} 1 + \operatorname{inv}_p(V) \left(\frac{-1}{p}\right)^{\frac{n+1}{2}} \left(\frac{D'}{p}\right)^\beta \left(\frac{-m'}{p}\right) p^{(1-n)(\beta+1/2)} & \text{if } \nu > \beta \\ 1 & \text{if } \nu \leq \beta, \end{cases}$$

and hence $L_m^{(p)}(p^{-s})$ is equal to

$$\begin{aligned} & (1 - p^{2n-s-1}) \sum_{\nu=0}^{\infty} N_m(p^\nu) p^{-\nu s} \\ &= 1 + \text{inv}_p(V) \left(\frac{-1}{p}\right)^{\frac{n+1}{2}} \left(\frac{D'}{p}\right)^\beta \left(\frac{-m'}{p}\right) p^{(1-n)(\beta+1/2)} p^{(2n-1-s)(\beta+1)} \\ &= 1 + \text{inv}_p(V) \left(\frac{-1}{p}\right)^{\frac{n-1}{2}} \left(\frac{D'}{p}\right)^\beta \left(\frac{m'}{p}\right) p^{(\beta+3/2)n-1/2-(\beta+1)s}. \end{aligned}$$

This concludes the proof of the proposition. \square

Recall the function $\mathbf{a}_n(s)$ of (1.1.4), and abbreviate $s_0 = (n-1)/2$. We now express the coefficients $B(m, 0, s)$ in terms of

$$\mathbf{a}_n(s - s_0) = \frac{D^{n/2} \Gamma\left(s + \frac{n}{2} + \frac{1}{2}\right) L(2s + 1, \varepsilon^n)}{2^n \pi^{s + \frac{n}{2} + \frac{1}{2}}}.$$

Corollary 2.6.5. *Assume that $n \geq 2$ is even. If $m > 0$ then*

$$B(m, 0, s) = -\frac{m^{s+s_0}}{\mathbf{a}_n(s - s_0)} \cdot \prod_{p \nmid D} \sum_{\gamma=0}^{v_p(m)} p^{-2s\gamma} \cdot \prod_{p|D} \frac{L_m^{(p)}(p^{-2s-n})}{1 - p^{-2s-1}}.$$

If $m > 0$ is prime to D then

$$B(m, 0, s) = -\frac{m^{s+s_0} \sigma_{-2s}(m)}{\mathbf{a}_n(s - s_0)} \cdot \prod_{p|D} \frac{1 - \left(\frac{-1}{p}\right)^{\frac{n}{2}} \text{inv}_p(V) p^{\frac{n}{2}-1-2s}}{1 - p^{-2s-1}},$$

where $\sigma_s(m) = \sum_{d|m} d^s$ is the usual divisor sum.

Proof. Combine Propositions 2.5.3 and 2.6.3. \square

Corollary 2.6.6. *Assume that $n \geq 1$ is odd. If $m > 0$ then*

$$B(m, 0, s) = -\frac{m^{s+s_0}}{\mathbf{a}_n(s - s_0)} \cdot \prod_{p \nmid D} \sum_{\gamma=0}^{v_p(m)} \varepsilon(p)^\gamma p^{-2s\gamma} \cdot \prod_{p|D} L_m^{(p)}(p^{-2s-n}).$$

If $m > 0$ is prime to D , then

$$B(m, 0, s) = -\frac{m^{s+s_0} \sigma_{-2s, \varepsilon}(m)}{\mathbf{a}_n(s - s_0)} \cdot \prod_{p|D} \left(1 + \left(\frac{-1}{p}\right)^{\frac{n-1}{2}} \left(\frac{m}{p}\right) \text{inv}_p(V) p^{\frac{n-1}{2}-2s}\right),$$

where $\sigma_{s, \varepsilon}(m) = \sum_{d|m} \varepsilon(d) d^s$ is the usual divisor sum.

Proof. Combine Propositions 2.5.3 and 2.6.4. \square

Remark 2.6.7. If $n = 1$, then $s_0 = 0$ is the center of symmetry of the functional equation of the incoherent Eisenstein series $E_L(\tau, s, 1)$. Hence $E_L(\tau, s, 1)$ vanishes identically (to odd order) at $s = 0$. For all positive m this implies that the coefficients $B(m, \mu, s_0)$ also vanish. Consequently,

in view of Proposition 2.5.3, there must be at least one prime $p < \infty$ for which the local factor $L_m^{(p)}(p^{-1})$ is zero. According to Proposition 2.6.4, this happens exactly when

$$\text{inv}_p(V) = - \begin{cases} \left(\frac{-D}{p}\right)^\beta, & \text{if } p \nmid D, \\ \left(\frac{D'}{p}\right)^\beta \left(\frac{m'}{p}\right), & \text{if } p \mid D, \end{cases}$$

where we have used the notation of the cited proposition. In terms of the local Hilbert symbol this is equivalent to the condition $\text{inv}_p(V) = -(-D, m)_p$, which means that V_p does not represent m .

The derivatives $B'(m, 0, s_0)$ of the coefficients can be computed as follows. Let \mathbb{V} be the incoherent hermitian space over \mathbb{A}_k whose non-archimedean contribution is given by $V \otimes_{\mathbb{Q}} \mathbb{A}_f$ and whose archimedean contribution is $V \otimes_{\mathbb{Q}} \mathbb{R}$ with the positive definite hermitian form $-\langle \cdot, \cdot \rangle$. Following [Ku97] we define the ‘Diff set’ associated with \mathbb{V} and $m \in \mathbb{Q}^\times$ by

$$\text{Diff}(\mathbb{V}, m) = \{p \leq \infty \mid \mathbb{V}_p \text{ does not represent } m\}.$$

This is a finite set of places of \mathbb{Q} of odd cardinality. When $m > 0$ it only consists of finite primes. The above argument shows that $L_m^{(p)}(p^{-1}) = 0$ if and only if $p \in \text{Diff}(\mathbb{V}, m)$. Hence $B'(m, 0, s_0) \neq 0$ if and only if $\text{Diff}(\mathbb{V}, m)$ consists of a single prime q . By Proposition 2.5.3 we find for $m \in \mathbb{Z}^+$ in this case that

$$B'(m, 0, 0) = -\frac{2\pi}{\sqrt{D}} \cdot \prod_{\substack{p \text{ prime} \\ p \neq q}} L_m^{(p)}(p^{-1}) \cdot \frac{d}{ds} L_m^{(q)}(q^{-2s-1}) \Big|_{s=0}.$$

This expression can be evaluated explicitly by means of Proposition 2.6.4. For instance, when $(m, D) = 1$ and $m = q^\beta m'$ with m' coprime to q , we see that $\text{inv}_q(V) = 1$, the prime q must be inert in k , and β must be odd. Under these conditions we find that

$$B'(m, 0, 0) = -\frac{2^{o(D)+1} \pi \sigma_{0,\varepsilon}(m')}{\sqrt{DL}(\varepsilon, 1)} (\beta + 1) \cdot \log(q),$$

where $o(D)$ denotes the number of prime factors of D .

Regardless of whether n is even or odd, we have the following lower bound for the coefficients of the Eisenstein series $E_L(\tau, s_0, n)$.

Corollary 2.6.8. *Assume that $n \geq 2$. There is a constant $C > 0$, depending only on L , such that*

$$-B(m, 0, s_0) > C \cdot m^{n-1}$$

for all $m \in \mathbb{Z}^+$.

Proof. This follows easily from Corollary 2.6.5 and Corollary 2.6.6. \square

2.7. Decomposing the Eisenstein series. As in the previous subsection, we assume the $\mathcal{O}_{\mathbf{k}}$ -lattice $L \subset V$ is self-dual under the hermitian form, and that $D = -\text{disc}(\mathbf{k})$ is odd. We also assume that V has dimension $n \geq 2$.

Consider again the holomorphic Eisenstein series

$$E_L(\tau, s_0, n) = \sum_{m \geq 0} \sum_{\mu \in L'/L} B(m, \mu, s_0) q^m \varphi_\mu$$

valued in $S_L = \mathbb{C}[L'/L]$ obtained by evaluating (2.5.1) at $s_0 = (n-1)/2$. Applying the linear functional $S_L \rightarrow \mathbb{C}$ sending $\varphi_0 \mapsto 1$ and $\varphi_\mu \mapsto 0$ for $\mu \neq 0$ yields an Eisenstein series

$$\sum_{m \geq 0} B(m, 0, s_0) \cdot q^m \in M_n(\Gamma_0(D), \varepsilon^n),$$

which can be decomposed as a linear combination of classical Eisenstein series indexed by the cusps

$$\infty_r = \frac{r}{D} \in \Gamma_0(D) \backslash \mathbb{P}^1(\mathbb{Q})$$

as $r \mid D$ varies. In this subsection we make this decomposition explicit.

For any divisor $r \mid D$ set $r' = D/r$. Our assumption that D is odd implies that the quadratic character $\varepsilon : (\mathbb{Z}/D\mathbb{Z})^\times \rightarrow \{\pm 1\}$ determined by \mathbf{k} is

$$\varepsilon(a) = \left(\frac{a}{D} \right).$$

Hence we may factor $\varepsilon = \varepsilon_r \cdot \varepsilon_{r'}$ with

$$\varepsilon_r(a) = \left(\frac{a}{r} \right) \quad \text{and} \quad \varepsilon_{r'}(a) = \left(\frac{a}{r'} \right).$$

Define the quadratic Gauss sum

$$\tau(\varepsilon_r) = \sum_{a \in (\mathbb{Z}/r\mathbb{Z})^\times} \varepsilon_r(a) e^{2\pi i a/r} = \begin{cases} \sqrt{r} & \text{if } r \equiv 1 \pmod{4} \\ i\sqrt{r} & \text{if } r \equiv 3 \pmod{4}, \end{cases}$$

and similarly with r replaced by r' .

Proposition 2.7.1. *For every divisor $r \mid D$ there is an Eisenstein series*

$$E_r(\tau) = \sum_{m \geq 0} e_r(m) \cdot q^m \in M_n(\Gamma_0(D), \varepsilon^n)$$

whose Fourier coefficients are as follows. The constant term is

$$e_r(0) = \begin{cases} 1 & \text{if } r = 1 \\ 0 & \text{otherwise.} \end{cases}$$

If n is even the coefficients indexed by $m > 0$ are

$$e_r(m) = \frac{r^{n/2} (-2\pi i)^n}{D^n \Gamma(n) L_D(n, \varepsilon^n)} \sum_{\substack{c \mid m \\ c > 0 \\ \gcd(m/c, r) = 1}} c^{n-1} \sum_{d \mid \gcd(c, r')} d \mu(r'/d).$$

If n is odd the coefficients indexed by $m > 0$ are

$$e_r(m) = \varepsilon_r(r') \frac{r^{n/2} (-2\pi i)^n \tau(\varepsilon_{r'})}{D^n \Gamma(n) L_D(n, \varepsilon^n)} \sum_{\substack{c|m \\ c>0 \\ \gcd(m/c, r)=1}} \varepsilon_r(m/c) \varepsilon_{r'}(c) \cdot c^{n-1}.$$

In both formulas $L_D(s, \varepsilon^n)$ is the Dirichlet L -function with Euler factors at all $\ell \mid D$ removed.

Proof. For each $s, t \in \mathbb{Z}/D\mathbb{Z}$ define an Eisenstein series

$$G_{(s,t)}(z) = \sum_{\substack{c,d \in \mathbb{Z} \\ (c,d) \neq (0,0) \\ (c,d) \equiv (s,t) \pmod{D}}} (cz + d)^{-n}.$$

Theorem 7.1.3 of [Miy89] implies that

$$E_r(z) = \frac{\varepsilon_r^n(r')}{2r^{n/2} L_D(n, \varepsilon^n)} \sum_{s,t \in \mathbb{Z}/D\mathbb{Z}} \varepsilon_r^n(s) \varepsilon_{r'}^n(t) G_{(s,t)}(r'z)$$

has the desired Fourier expansion. \square

Remark 2.7.2. The Eisenstein series of Proposition 2.7.1 agree with the Eisenstein series denoted the same way in §4.4 of [BHK⁺a]. In particular, E_r is nonvanishing at the cusp $\infty_r = r/D$, and vanishes at all other cusps.

Proposition 2.7.3. *We have the equality*

$$\sum_{r|D} \gamma_r E_r(\tau) = \sum_{m \geq 0} B(m, 0, s_0) \cdot q^m$$

of forms in $M_n(\Gamma_0(D), \varepsilon^n)$. Here for a prime $\ell \mid D$ we have set

$$\gamma_\ell = (D, \ell)_\ell^n \cdot \text{inv}_\ell(V) \cdot \begin{cases} 1 & \text{if } \ell \equiv 1 \pmod{4} \\ i^{-n} & \text{if } \ell \equiv 3 \pmod{4}, \end{cases}$$

where $(-, -)_\ell$ is the usual Hilbert symbol at ℓ , and for a divisor $r \mid D$ we have set

$$\gamma_r = \prod_{\ell|r} \gamma_\ell.$$

Proof. We use the \mathbb{C} -bilinear pairing $\langle \cdot, \cdot \rangle$ on S_L to write

$$\sum_{m \geq 0} B(m, 0, s_0) \cdot q^m = \langle E_L, \varphi_0 \rangle.$$

On the other hand, using (2.4.4) and the scalar Eisenstein series

$$E = \sum_{\gamma \in \Gamma_\infty \setminus \Gamma_0(D)} \varepsilon^n(d) \cdot (1|_n \gamma)$$

of weight n for the cusp ∞ in $M_n(\Gamma_0(D), \varepsilon^n)$, we obtain

$$\begin{aligned} E_L = E_L(\tau, s_0, n) &= \sum_{\gamma \in \Gamma_\infty \backslash \Gamma} (c\tau + d)^{-n} \cdot \bar{\omega}_L(\gamma)^{-1} \varphi_0 \\ &= \sum_{\gamma \in \Gamma_0(D) \backslash \Gamma} (E \mid_n \gamma) \cdot \bar{\omega}_L(\gamma)^{-1} \varphi_0. \end{aligned}$$

Hence

$$\langle E_L, \varphi_0 \rangle = \sum_{\gamma \in \Gamma_0(D) \backslash \Gamma} (E \mid_n \gamma) \cdot \langle \bar{\omega}_L(\gamma)^{-1} \varphi_0, \varphi_0 \rangle.$$

We compute this finite sum by means of the left coset decomposition

$$\Gamma = \bigsqcup_{r|D} \bigsqcup_{c \in \mathbb{Z}/r\mathbb{Z}} \Gamma_0(D) R_r \begin{pmatrix} 1 & c \\ 0 & 1 \end{pmatrix},$$

see e.g. equation (8.2.1) in [BHK⁺a]. Using the notation of [BHK⁺a] we find

$$\begin{aligned} \langle E_L, \varphi_0 \rangle &= \sum_{\gamma \in \Gamma_0(D) \backslash \Gamma} (E \mid_n \gamma) \cdot \langle \varphi_0, \omega_L(\gamma) \varphi_0 \rangle \\ &= \sum_{r|D} \sum_{c \in \mathbb{Z}/r\mathbb{Z}} (E \mid_n R_r \begin{pmatrix} 1 & c \\ 0 & 1 \end{pmatrix}) \cdot \langle \varphi_0, \omega_L(R_r) \omega(\begin{pmatrix} 1 & c \\ 0 & 1 \end{pmatrix}) \varphi_0 \rangle \\ &= \sum_{r|D} \sum_{c \in \mathbb{Z}/r\mathbb{Z}} (E \mid_n W_r \begin{pmatrix} 1 & c \\ 0 & r \end{pmatrix}) \cdot \langle \varphi_0, \omega_L(R_r) \varphi_0 \rangle. \end{aligned}$$

In terms of the U_r operators on $M_n(\Gamma_0(D), \varepsilon^n)$, given by $\sum_n a(n)q^n \mid_n U_r = \sum_n a(rn)q^n$, we may rewrite this as

$$\langle E_L, \varphi_0 \rangle = \sum_{r|D} \langle \varphi_0, \omega_L(R_r) \varphi_0 \rangle \cdot r^{1-n/2} (E \mid_n W_r \mid_n U_r).$$

By means of equation (8.2.3) of [BHK⁺a], we obtain for the matrix coefficients

$$\langle \varphi_0, \omega_L(R_r) \varphi_0 \rangle = \varepsilon_r^n(\alpha) \varepsilon_r^n(-\beta) \gamma_r r^{-n/2},$$

where we have written $rr' = D$. Inserting the definition

$$E_r = \varepsilon_r^n(-\beta) \varepsilon_r^n(\alpha r) (E \mid_n W_r)$$

of the Eisenstein series for the cusp $\infty_r = r/D$, we find

$$\langle E_L, \varphi_0 \rangle = \sum_{r|D} \varepsilon_r^n(r) \gamma_r r^{1-n} (E_r \mid_n U_r).$$

It follows from the Fourier expansion of E_r given in Proposition 2.7.1 that

$$E_r \mid_n U_r = \varepsilon_r^n(r) r^{n-1} E_r.$$

Inserting this, we finally obtain

$$\langle E_L, \varphi_0 \rangle = \sum_{r|D} \gamma_r E_r,$$

concluding the proof of the proposition. \square

The following fact about the coefficients of the Eisenstein series of Proposition 2.7.1 will be needed much later, in the proof of Lemma 8.4.4.

Proposition 2.7.4. *For any positive $m \in \mathbb{Z}$ with $\gcd(m, D) = 1$, we have*

$$\sum_{r|D} \gamma_r e_r(m) \log(D/r) = \left(\sum_{\ell|D} \frac{\log(\ell)}{1 + \beta_\ell \cdot \left(\frac{m}{\ell}\right)^n} \right) \sum_{r|D} \gamma_r e_r(m).$$

Here both sums over $r \mid D$ are over all positive divisors of D , the sum over $\ell \mid D$ is over only the prime divisors, γ_r is as in Proposition 2.7.3, and we abbreviate

$$\beta_\ell = (-1)^{n+1} \operatorname{inv}_\ell(V) \cdot \begin{cases} \left(\frac{-1}{\ell}\right)^{\frac{n}{2}} \ell^{\frac{n}{2}} & \text{if } n \text{ is even} \\ \left(\frac{-1}{\ell}\right)^{\frac{n-1}{2}} \ell^{\frac{n-1}{2}} & \text{if } n \text{ is odd.} \end{cases}.$$

Proof. Directly from the formulas of Proposition 2.7.1, one can see that there is a factorization

$$\gamma_r e_r(m) = c(m) f_r(m)$$

in which $c(m)$ is some function independent of r , and

$$f_r(m) = \left(\frac{m}{r}\right)^n \cdot \prod_{\ell|r} \beta_\ell.$$

The relation $f_r(m) = \prod_{\ell|r} f_\ell(m)$ implies the combinatorial identity

$$\sum_{r|D} f_r(m) \log(D/r) = \left(\sum_{\ell|D} \frac{\log(\ell)}{1 + f_\ell(m)} \right) \sum_{r|D} f_r(m),$$

and multiplying both sides by $c(m)$ proves the claim. \square

3. AUTOMORPHIC GREEN FUNCTIONS

As in §2, we fix a \mathbf{k} -hermitian space V of signature $(n-1, 1)$ with $n \geq 1$. We recall the family of special divisors on the complex Shimura variety associated to the unitary group $U(V)$, and derive a geometric variant of the Siegel-Weil formula. Using this, we compute the integrals of automorphic Green functions for the special divisors.

3.1. A complex Shimura variety. The fixed embedding $\mathbf{k} \subset \mathbb{C}$ identifies $\mathbf{k} \otimes_{\mathbb{Q}} \mathbb{R} \cong \mathbb{C}$, and allows us to distinguish between the two orthogonal idempotents $\epsilon, \bar{\epsilon} \in \mathbf{k} \otimes_{\mathbb{Q}} \mathbb{C}$, which we label in such a way that

$$(\alpha \otimes 1)\epsilon = (1 \otimes \alpha)\epsilon, \quad (\alpha \otimes 1)\bar{\epsilon} = (1 \otimes \bar{\alpha})\bar{\epsilon}$$

for all $\alpha \in \mathbf{k}$.

Viewing $V(\mathbb{R}) = V \otimes_{\mathbb{Q}} \mathbb{R}$ as \mathbb{C} -hermitian space of signature $(n-1, 1)$, define a hermitian symmetric domain¹

$$(3.1.1) \quad \mathcal{D} = \{\text{negative definite complex lines } z \subset V(\mathbb{R})\}.$$

The natural map

$$V(\mathbb{R}) \hookrightarrow V(\mathbb{C}) = \epsilon V(\mathbb{C}) \oplus \bar{\epsilon} V(\mathbb{C}) \xrightarrow{\text{proj}} \epsilon V(\mathbb{C})$$

is a \mathbb{C} -linear isomorphism, and identifies

$$(3.1.2) \quad \mathcal{D} = \{z \in \mathbb{P}(\epsilon V(\mathbb{C})) : [z, \bar{z}] < 0\}.$$

Here we have endowed $V(\mathbb{C}) = V \otimes_{\mathbb{Q}} \mathbb{C}$ with the \mathbb{C} -bilinear extension of the \mathbb{Q} -bilinear form $[-, -]$ of (2.1.1), and $z \mapsto \bar{z}$ is the complex conjugation on the second factor in the tensor product. Under either interpretation, there is an evident action of the real points of

$$H = \text{U}(V)$$

on \mathcal{D} by holomorphic automorphisms.

Fix an $\mathcal{O}_{\mathbf{k}}$ -lattice $L \subset V$ on which the hermitian form is $\mathcal{O}_{\mathbf{k}}$ -valued. Choose a compact open subgroup $K \subset H(\mathbb{A}_f)$ that stabilizes L , and hence acts on the finite dimensional vector space $S_L = \mathbb{C}[L'/L]$ of §2.4. This choice determines a complex Shimura variety

$$(3.1.3) \quad \text{Sh}_K(H, \mathcal{D}) = H(\mathbb{Q}) \backslash \mathcal{D} \times H(\mathbb{A}_f) / K,$$

which we view as a complex orbifold of dimension $n-1$. If we write $H(\mathbb{A}_f) = \bigsqcup_j H(\mathbb{Q})h_jK$ as a finite disjoint union, then

$$(3.1.4) \quad \text{Sh}_K(H, \mathcal{D}) = \bigsqcup_j (H(\mathbb{Q}) \cap K_j) \backslash \mathcal{D},$$

where $K_j = h_j K h_j^{-1}$ stabilizes the lattice $L_j = h_j L$.

The Shimura variety carries a family of special divisors

$$Z(m) \in \text{Div}(\text{Sh}_K(H, \mathcal{D}))$$

indexed by positive integers m , which we describe in terms of the decomposition (3.1.4). For each $x \in V$ with $\langle x, x \rangle > 0$, define an analytic divisor

$$\mathcal{D}(x) = \{z \in \mathcal{D} : x \perp z\}$$

¹We take this opportunity to point out a misstatement in (2.1.2) of [BHK⁺a]: the word “planes” should be replaced by “lines” in the definition of the hermitian symmetric domain.

where \perp means orthogonal with respect to the hermitian or bilinear form, depending on whether the reader prefers the model (3.1.1) or (3.1.2). The pullback of $Z(m)$ via the uniformization

$$\mathcal{D} \rightarrow (H(\mathbb{Q}) \cap K_j) \backslash \mathcal{D} \subset \text{Sh}_K(H, \mathcal{D})$$

is the locally finite analytic divisor

$$\sum_{\substack{x \in L_j \\ \langle x, x \rangle = m}} \mathcal{D}(x) \in \text{Div}(\mathcal{D}).$$

More generally, given a positive $m \in \mathbb{Q}$ and a K -fixed function $\varphi \in S_L$, there is a special divisor

$$Z(m, \varphi) \in \text{Div}_{\mathbb{C}}(\text{Sh}_K(H, \mathcal{D}))$$

whose pullback via the above uniformization is

$$\sum_{\mu \in L'/L} \varphi(\mu) \sum_{\substack{x \in \mu_j + L_j \\ \langle x, x \rangle = m}} \mathcal{D}(x) \in \text{Div}_{\mathbb{C}}(\mathcal{D}),$$

where μ_j is the image of μ under $h_j : L'/L \rightarrow L'_j/L_j$. Taking $\varphi = \varphi_0$ to be the characteristic function of $0 \in L'/L$ recovers $Z(m)$.

Let \mathcal{L} be the tautological line bundle over \mathcal{D} whose fiber over a point $z \in \mathcal{D}$ is the complex line $\mathcal{L}_z = z \subset V(\mathbb{R})$. Denote by $\widehat{\mathcal{L}}$ the tautological line bundle endowed with the hermitian metric $\|\cdot\|$ defined as follows: if we use the interpretation (3.1.1) to view $z \in \mathcal{D}$ as a line in $V(\mathbb{R})$, then

$$\|s\|^2 = -\frac{\langle s, s \rangle}{4\pi e^\gamma}$$

for every $s \in \mathcal{L}_z$. If we instead use (3.1.2) to view z as a line in $\epsilon V(\mathbb{C})$, the metric is the same, but with $\langle -, - \rangle$ replaced by $[-, -]$. The Chern form $\text{ch}(\widehat{\mathcal{L}})$ is an $H(\mathbb{R})$ -invariant Kähler form on \mathcal{D} . The metrized tautological bundle descends to $\text{Sh}_K(H, \mathcal{D})$, and its holomorphic sections are hermitian modular forms of weight 1. As in the introduction, we abbreviate

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}) = \int_{\text{Sh}_K(H, \mathcal{D})} \text{ch}(\widehat{\mathcal{L}})^{n-1}.$$

In the subsections below we will recall the construction of Green functions for certain linear combinations of special divisors, and compute their integrals with respect to the volume form $\text{ch}(\widehat{\mathcal{L}})^{n-1}$.

Remark 3.1.1. Exactly as in the case of orthogonal Shimura varieties [Ku03], one can show that

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}})^{-1} \int_{Z(m, \varphi)} \text{ch}(\widehat{\mathcal{L}})^{n-2} = - \sum_{\mu \in L'/L} \varphi(\mu) B(m, \mu, s_0).$$

On the right hand side the notation is that of §2.5.

3.2. A geometric variant of the Siegel-Weil formula. For $z \in \mathcal{D}$ and $x \in V(\mathbb{R})$, let

$$\varphi_\infty(z, x) = \exp(-2\pi\langle x_{z^\perp}, x_{z^\perp} \rangle + 2\pi\langle x_z, x_z \rangle)$$

be the Gaussian. Here we are using the model (3.1.1), so that z determines an orthogonal decomposition $V(\mathbb{R}) = z^\perp \oplus z$, and the vectors x_{z^\perp} and x_z are the projections of x to the two summands.

As in §2.1, let $G \cong \mathrm{U}(1, 1)$ be the quasi-split unitary group over \mathbb{Q} , so that $\mathrm{SL}_2 \cong G' \subset G$. Fix a base point $z_0 \in \mathcal{D}$ and consider the Schwartz function

$$\varphi_\infty(x) \stackrel{\text{def}}{=} \varphi_\infty(z_0, x) \in S(V(\mathbb{R})).$$

It has weight $n - 2$ under the action of the Weil representation (§2.3) of $\mathrm{SL}_2(\mathbb{R})$. In other words, using the notation of (2.2.1),

$$\omega(k)(\varphi_\infty) = \underline{k}^{n-2}\varphi_\infty$$

for all $k \in \mathrm{SO}(2, \mathbb{R})$.

A choice of $\varphi_f \in S(V(\mathbb{A}_f))$ determines a Schwartz function $\varphi_\infty \otimes \varphi_f \in S(V(\mathbb{A}))$. Applying the construction (2.3.1) yields a corresponding theta function $\theta(g, h, \varphi_\infty \otimes \varphi_f)$ of the variables $(g, h) \in G(\mathbb{A}) \times H(\mathbb{A})$. Restricting the first variable to $g \in \mathrm{SL}_2(\mathbb{R}) \subset G(\mathbb{A})$, we obtain the Siegel theta function

$$\theta(g, h, \varphi_\infty \otimes \varphi_f) = \sum_{x \in V(\mathbb{Q})} \omega(g, 1)\varphi_\infty(z_0, h_\infty^{-1}x) \cdot \varphi_f(h_f^{-1}x),$$

where $h = h_\infty h_f \in H(\mathbb{A})$.

We may view this theta function as a function of the variables

$$(\tau, z, h_f) \in \mathcal{H} \times \mathcal{D} \times H(\mathbb{A}_f)$$

as follows. Let $g_\tau \in \mathrm{SL}_2(\mathbb{R})$ be as in (2.1.3), and choose an $h_\infty \in H(\mathbb{R})$ with $h_\infty z_0 = z$, so that $\varphi_\infty(z_0, h_\infty^{-1}x) = \varphi_\infty(z, x)$. Now set

$$(3.2.1) \quad \begin{aligned} \theta(\tau, z, h_f, \varphi_f) &= v^{1-\frac{n}{2}}\theta(g_\tau, h_\infty h_f, \varphi_\infty \otimes \varphi_f) \\ &= v \sum_{x \in V(\mathbb{Q})} e(\langle x_{z^\perp}, x_{z^\perp} \rangle \tau - \langle x_z, x_z \rangle \bar{\tau}) \cdot \varphi_f(h_f^{-1}x). \end{aligned}$$

In the variable τ , we have the transformation law

$$(3.2.2) \quad \theta(\gamma\tau, z, h_f, \varphi_f) = (c\tau + d)^{n-2}\theta(\tau, z, h_f, \omega(\gamma)^{-1}\varphi_f)$$

for $\gamma \in \Gamma$. In the variable z , we have the transformation law

$$(3.2.3) \quad \theta(\tau, \delta z, \delta h_f, \varphi_f) = \theta(\tau, z, h_f, \varphi_f).$$

for $\delta \in H(\mathbb{Q})$,

Now suppose that $\varphi_f \in S_L = \mathbb{C}[L'/L]$ is fixed by the action of K . Using (3.2.1) and (3.2.3), we see that the Siegel theta function (3.2.1) descends

to a function on $\mathcal{H} \times \text{Sh}_K(H, \mathcal{D})$. We are interested in the geometric theta integral

$$(3.2.4) \quad \mathcal{I}(\tau, \varphi_f) = \text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}})^{-1} \int_{\text{Sh}_K(H, \mathcal{D})} \theta(\tau, z, h_f, \varphi_f) \text{ch}(\widehat{\mathcal{L}})^{n-1}$$

as a function of $\tau \in \mathcal{H}$.

Proposition 3.2.1. *Assume that $n > 2$ or that V is anisotropic. Recalling the notation of (2.3.2) and (2.4.4), we have*

$$\kappa \cdot \mathcal{I}(\tau, \varphi_f) = \langle E_L(\tau, s_0, n-2), \varphi_f \rangle,$$

where the pairing on the right is (2.4.3).

Proof. We rewrite the integral in (3.2.4) as an integral over the quotient $H(\mathbb{Q}) \backslash H(\mathbb{A})$. To this end, we normalize the Haar measure on the compact subgroup $H(\mathbb{R})_{z_0} \subset H(\mathbb{R})$ so that the volume is 1. Then we normalize the Haar measure dh_{∞} on $H(\mathbb{R})$ so that the induced measure on the quotient $H(\mathbb{R})/H(\mathbb{R})_{z_0} \cong \mathcal{D}$ agrees with the measure defined by $\text{ch}(\widehat{\mathcal{L}})^{n-1}$. Finally, we normalize the Haar measure dh_f on $H(\mathbb{A}_f)$ so that $\text{vol}(H(\mathbb{Q}) \backslash H(\mathbb{A})) = 1$. Setting

$$\text{vol}(K) = \int_K dh_f,$$

we then have the identities

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}) = \int_{H(\mathbb{Q}) \backslash H(\mathbb{A}) / H(\mathbb{R})_{z_0} K} dh_{\infty} dh_f = \text{vol}(K)^{-1}.$$

Using (3.2.1) we obtain

$$\begin{aligned} \mathcal{I}(\tau, \varphi) &= \text{vol}(K) \int_{\text{Sh}_K(H, \mathcal{D})} \theta(\tau, z, h_f, \varphi_f) \text{ch}(\widehat{\mathcal{L}})^{n-1} \\ &= v^{1-\frac{n}{2}} \int_{H(\mathbb{Q}) \backslash H(\mathbb{A})} \theta(g_{\tau}, h_{\infty} h_f, \varphi_{\infty} \otimes \varphi_f) dh_{\infty} dh_f, \end{aligned}$$

and applying the Siegel-Weil formula of Theorem 2.3.1 shows

$$\kappa \cdot \mathcal{I}(\tau, \varphi) = v^{1-\frac{n}{2}} E(g_{\tau}, \lambda(\varphi_{\infty} \otimes \varphi_f), s_0).$$

As φ_{∞} has weight $n-2$, Lemma 2.2.1 implies

$$\begin{aligned} &E(g_{\tau}, \lambda(\varphi_{\infty} \otimes \varphi_f), s_0) \\ &= v^{\frac{n}{2}-1} \sum_{\gamma \in \Gamma_{\infty} \backslash \Gamma} \text{Im}(\gamma\tau)^{s_0 + \frac{1}{2} - \frac{n-2}{2}} (c\tau + d)^{2-n} \cdot (\omega_L(\gamma)\varphi_f)(0) \\ &= v^{\frac{n}{2}-1} \sum_{\gamma \in \Gamma_{\infty} \backslash \Gamma} \text{Im}(\gamma\tau)^{s_0 + \frac{1}{2} - \frac{n-2}{2}} (c\tau + d)^{2-n} \cdot (\omega_L^{\vee}(\gamma)^{-1}\delta_0)(\varphi_f). \end{aligned}$$

Here $\delta_0 \in S_L^{\vee}$ denotes the functional $\varphi_f \mapsto \varphi_f(0)$. As the Weil representation is unitary, we have

$$(\omega_L^{\vee}(\gamma)^{-1}\delta_0)(\varphi_f) = \langle \bar{\omega}_L(\gamma)^{-1}\varphi_0, \varphi_f \rangle.$$

To complete the proof, substitute this equality into the final expression above, and recall the definition of $E_L(\tau, s, n-2)$ from (2.4.4). \square

3.3. Automorphic Green functions. Let σ be a finite dimensional representation of $\Gamma = \mathrm{SL}_2(\mathbb{Z})$ on a complex vector space V_σ , and assume that σ factors through a finite quotient. In our applications, σ will be the Weil representation

$$\omega_L : \Gamma \rightarrow \mathrm{Aut}(S_L)$$

of (2.4.2), or its complex conjugate, or its dual.

For $k \in \mathbb{Z}$, denote by $H_k(\sigma)$ the vector space of (weak) harmonic Maass forms of weight k for the group Γ with representation σ as in [BY09]. We write

$$S_k(\sigma) \subset M_k(\sigma) \subset M_k^!(\sigma)$$

for the spaces of cusp forms, holomorphic modular forms, and weakly holomorphic modular forms, respectively.

A harmonic Maass form $f \in H_k(\sigma)$ has a Fourier expansion

$$(3.3.1) \quad f(\tau) = \sum_{\substack{m \in \mathbb{Q} \\ m \gg -\infty}} c^+(m)q^m + \sum_{\substack{m \in \mathbb{Q} \\ m < 0}} c^-(m)\Gamma(1-k, 4\pi|m|v)q^m$$

with coefficients $c^\pm(m) \in V_\sigma$. Here $\tau = i + iv \in \mathcal{H}$, $q = e^{2\pi i\tau}$, and

$$\Gamma(s, x) = \int_x^\infty e^{-t}t^{s-1}dt$$

is the incomplete gamma function. The coefficients are supported on rational numbers with uniformly bounded denominators. The first summand on the right hand side of (3.3.1) is denoted by f^+ and is called the holomorphic part of f , the second summand is denoted by f^- and is called the non-holomorphic part.

As in [BF04], there is a surjective conjugate-linear differential operator

$$(3.3.2) \quad \xi_k : H_k(\omega_L) \rightarrow S_{2-k}(\bar{\omega}_L)$$

defined by

$$\xi_k(f)(\tau) = 2iv^k \frac{\overline{\partial f}}{\partial \bar{\tau}}.$$

Its kernel is $M_k^!(\omega_L)$.

Suppose $f \in H_{2-n}(\omega_L)$ is K -fixed with Fourier coefficients $c^\pm(m) \in S_L$. Define the special divisor associated to f by

$$(3.3.3) \quad Z(f) = \sum_{m>0} \sum_{\mu \in L'/L} c^+(-m, \mu)Z(m, \varphi_\mu) \in \mathrm{Div}_{\mathbb{C}}(\mathrm{Sh}_K(H, \mathcal{D})).$$

Here $c^\pm(m, \mu) \in \mathbb{C}$ is the value of $c^\pm(m)$ at $\mu \in L'/L$, and φ_μ is the characteristic function of the coset $\mu + \hat{L}$.

We recall from §3 of [BHY15] the construction of an automorphic Green function for $Z(f)$. Using (3.2.1), define an S_L -valued Siegel theta function

$$(3.3.4) \quad \theta_L(\tau, z, h) = \sum_{\mu \in L'/L} \theta(\tau, z, h, \varphi_\mu) \varphi_\mu$$

for $\tau \in \mathcal{H}$, $z \in \mathcal{D}$, and $h \in H(\mathbb{A}_f)$. The transformation law (3.2.2) implies that it defines a non-holomorphic modular form for Γ of weight $n-2$ with representation $\bar{\omega}_L$, and so, using the notation (2.4.3), the pairing $\langle f, \theta_L(\tau, z, h) \rangle$ is Γ -invariant as a function of $\tau \in \mathcal{H}$.

Following [Bo98] and [BF04], we consider the regularized theta lift

$$(3.3.5) \quad \Phi(z, h, f) = \int_{\Gamma \backslash \mathcal{H}}^{\text{reg}} \langle f, \theta_L(\tau, z, h) \rangle d\mu(\tau)$$

of f , where $d\mu(\tau) = \frac{du dv}{v^2}$ is the invariant measure. Here the integral is regularized as in [Bo98] and [BHY15]. The main properties of

$$\Phi(f) = \Phi(z, h, f),$$

viewed as a function on $\text{Sh}_K(H, \mathcal{D})$, are summarized in the following theorem, see Theorem 3.3.1 and Proposition 3.3.4 of [BHY15].

Theorem 3.3.1. *The Green function $\Phi(z, h, f)$ is smooth on $\text{Sh}_K(H, \mathcal{D}) \setminus Z(f)$. It has logarithmic singularities along $Z(f)$, and it is integrable over $\text{Sh}_K(H, \mathcal{D})$ if $n > 2$ or if V is anisotropic. The corresponding current on smooth compactly supported $(n-1, n-1)$ -forms on $\text{Sh}_K(H, \mathcal{D})$ satisfies the Green equation*

$$dd^c[\Phi(f)] + \delta_{Z(f)} = [dd^c\Phi(f)].$$

3.4. Integrals of Green functions. In this subsection we assume that either $n > 2$, or that $n \geq 1$ and V is anisotropic. Choose an S_L -valued harmonic Maass form

$$f \in H_{2-n}(\omega_L)$$

fixed by the action of the compact open subgroup $K \subset H(\mathbb{A}_f)$ on S_L .

The main result of this subsection expresses the integral

$$(3.4.1) \quad \mathcal{I}(f) = \text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}})^{-1} \int_{\text{Sh}_K(H, \mathcal{D})} \Phi(f) \text{ch}(\widehat{\mathcal{L}})^{n-1}$$

in terms of the derivatives $B'(m, \mu, s_0)$ at $s_0 = (n-1)/2$ of the coefficients (2.5.2) of $E_L(\tau, s, n)$. The proof follows the argument of [Ku03] in the orthogonal case, which is based on the Siegel-Weil formula. In fact, we slightly generalize this argument by allowing harmonic Maass forms for the inputs of the regularized theta lift. Let $\kappa \in \{1, 2\}$ be as in (2.3.2).

Lemma 3.4.1. *We have*

$$\mathcal{I}(f) = \lim_{T \rightarrow \infty} \left(\frac{1}{\kappa} \int_{\mathcal{F}_T} \langle f, E_L(\tau, s_0, n-2) \rangle d\mu(\tau) - c^+(0, 0) \log(T) \right).$$

Here $\mathcal{F}_T = \{\tau \in \mathcal{F} : \text{Im}(\tau) \leq T\}$ is the truncation at height T of the standard fundamental domain \mathcal{F} for the action of Γ on the complex upper half plane.

Proof. Exactly as in Proposition 2.5 of [Ku03], on $\text{Sh}_K(H, \mathcal{D}) \setminus Z(f)$ we have the equality

$$\Phi(z, h, f) = \lim_{T \rightarrow \infty} \left(\int_{\mathcal{F}_T} \langle f, \theta_L(\tau, z, h) \rangle d\mu(\tau) - c^+(0, 0) \log(T) \right).$$

The integral $\mathcal{I}(f)$ is therefore given by the limit as $T \rightarrow \infty$ of

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}})^{-1} \int_{\text{Sh}_K(H, \mathcal{D})} \int_{\mathcal{F}_T} \langle f, \theta_L(\tau, z, h) \rangle d\mu(\tau) \text{ch}(\widehat{\mathcal{L}})^{n-1} - c^+(0, 0) \log(T).$$

As \mathcal{F}_T is compact, we may interchange the order of integration. Inserting (3.3.4) and (3.2.4), we find that $\mathcal{I}(f)$ is the limit as $T \rightarrow \infty$ of

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}})^{-1} \int_{\mathcal{F}_T} \left\langle f, \int_{\text{Sh}_K(H, \mathcal{D})} \theta_L(\tau, z, h) \text{ch}(\widehat{\mathcal{L}})^{n-1} \right\rangle d\mu(\tau) - c^+(0, 0) \log(T),$$

which we rewrite as

$$\int_{\mathcal{F}_T} \sum_{\mu \in L'/L} f(\tau)(\mu) \cdot \mathcal{I}(\tau, \varphi_{\mu}) d\mu(\tau) - c^+(0, 0) \log(T).$$

Applying Proposition 3.2.1 completes the proof. \square

Theorem 3.4.2. *If $n > 2$, or if $n = 2$ and V is anisotropic, the integral (3.4.1) satisfies*

$$\mathcal{I}(f) = \sum_{\mu \in L'/L} \sum_{\substack{m \in \mathbb{Q} \\ m > 0}} c^+(-m, \mu) B'(m, \mu, s_0).$$

If $n = 1$, the integral (3.4.1) satisfies

$$\mathcal{I}(f) = \frac{1}{2} \sum_{\mu \in L'/L} \sum_{\substack{m \in \mathbb{Q} \\ m \geq 0}} c^+(-m, \mu) B'(m, \mu, s_0).$$

Proof. Abbreviate

$$A_T(f) = \int_{\mathcal{F}_T} \langle f, E_L(\tau, s_0, n-2) \rangle d\mu(\tau),$$

so that Lemma 3.4.1 becomes

$$(3.4.2) \quad \mathcal{I}(f) = \lim_{T \rightarrow \infty} \left(\frac{A_T(f)}{\kappa} - c^+(0, 0) \log(T) \right).$$

Using Lemma 2.4.1 we see that

$$E_L(\tau, s_0, n-2) d\mu(\tau) = (L_n E'_L(\tau, s_0, n)) d\mu(\tau) = -\bar{\partial} E'_L(\tau, s_0, n) d\tau,$$

and so

$$\begin{aligned} A_T(f) &= - \int_{\mathcal{F}_T} \langle f, \bar{\partial} E'_L(\tau, s_0, n) \rangle d\tau \\ &= - \int_{\mathcal{F}_T} d \langle \langle f, E'_L(\tau, s_0, n) \rangle \rangle d\tau + \int_{\mathcal{F}_T} \langle (\bar{\partial} f), E'_L(\tau, s_0, n) \rangle d\tau. \end{aligned}$$

Applying Stokes' theorem to the first term, and

$$(\bar{\partial}f)d\tau = -(L_{2-n}f) d\mu(\tau)$$

to the second, we obtain

$$\begin{aligned} A_T(f) &= - \int_{\partial\mathcal{F}_T} \langle f, E'_L(\tau, s_0, n) \rangle d\tau - \int_{\mathcal{F}_T} \langle L_{2-n}f, E'_L(\tau, s_0, n) \rangle d\mu(\tau) \\ &= \int_{u=0}^1 \langle f(u+iT), E'_L(u+iT, s_0, n) \rangle du \\ &\quad - \int_{\mathcal{F}_T} \left\langle \overline{\xi_{2-n}(f)}, E'_L(\tau, s_0, n) \right\rangle v^n d\mu(\tau), \end{aligned}$$

where ξ_{2-n} is the differential operator (3.3.2). Inserting the Fourier expansions (3.3.1) and (2.5.1) of f and the Eisenstein series yields

$$\begin{aligned} A_T(f) &= \sum_{\substack{\mu \in L'/L \\ m \in \mathbb{Q}}} c^+(-m, \mu) B'(m, \mu; s_0, T) \\ &\quad + \sum_{\substack{\mu \in L'/L \\ m \in \mathbb{Q}^+}} c^-(-m, \mu) \Gamma(n-1, 4\pi|m|T) B'(m, \mu; s_0, T) \\ &\quad - \int_{\mathcal{F}_T} \left\langle \overline{\xi_{2-n}(f)}, E'_L(\tau, s_0, n) \right\rangle v^n d\mu(\tau). \end{aligned}$$

The exponential decay of the incomplete gamma function and the polynomial growth of the coefficients $c^-(m, \mu)$ imply that the second term goes to zero in the limit $T \rightarrow \infty$. The third term converges to the Petersson scalar product of $\xi_{2-n}(f)$ and $E'_L(\tau, s_0, n)$. But since $\xi_{2-n}(f)$ is cuspidal and hence orthogonal to Eisenstein series, this Petersson scalar product vanishes. Inserting this into (3.4.2), we find

$$\mathcal{I}(f) = \lim_{T \rightarrow \infty} \left(\frac{1}{\kappa} \sum_{\substack{\mu \in L'/L \\ m \in \mathbb{Q}}} c^+(-m, \mu) B'(m, \mu; s_0, T) - c^+(0, 0) \log(T) \right).$$

The exponential decay of the coefficients $B'(m, \mu; s_0, T)$ in (2.5.2) for $m < 0$ and the subexponential growth of the coefficients $c^+(-m, \mu)$ imply that the contribution of all $m < 0$ in the above term vanishes. If $n > 1$ we obtain by virtue of Lemma 2.5.4 that

$$\mathcal{I}(f) = \frac{1}{\kappa} \sum_{\substack{\mu \in L'/L \\ m > 0}} c^+(-m, \mu) B'(m, \mu, s_0),$$

as desired. When $n = 1$, again by Lemma 2.5.4, there is an additional contribution to the sum from $m = 0$. \square

4. INTEGRAL MODELS AND ARITHMETIC INTERSECTION THEORY

We recall the integral models of $\mathrm{GU}(n-1, 1)$ Shimura varieties constructed by Pappas and Krämer, and the arithmetic intersection theory of Gillet-Soulé and Burgos-Kramer-Kühn on their toroidal compactifications.

From here until the end of the paper, we assume $D = -\mathrm{disc}(\mathbf{k})$ is odd.

4.1. Moduli problems. For any $n \geq 1$, work of Pappas [Pap00] and Krämer [Kra03], as summarized in §2.3 of [BHK⁺a], provides us with a regular and flat $\mathcal{O}_{\mathbf{k}}$ -stack

$$\mathcal{M}_{(n-1,1)} \rightarrow \mathrm{Spec}(\mathcal{O}_{\mathbf{k}})$$

with reduced fibers. For an $\mathcal{O}_{\mathbf{k}}$ -scheme S , the objects of the groupoid $\mathcal{M}_{(n-1,1)}(S)$ are quadruples $(A, \iota, \psi, \mathcal{F})$ in which

- $A \rightarrow S$ is an abelian scheme of dimension n ,
- $\iota : \mathcal{O}_{\mathbf{k}} \rightarrow \mathrm{End}(A)$ is an $\mathcal{O}_{\mathbf{k}}$ -action,
- $\psi : A \rightarrow A^{\vee}$ is a principal polarization whose induced Rosati involution restricts to complex conjugation on the image of ι ,
- $\mathcal{F} \subset \mathrm{Lie}(A)$ is an $\mathcal{O}_{\mathbf{k}}$ -stable hyperplane² satisfying the signature $(n-1, 1)$ condition of Krämer [Kra03]: the actions of $\mathcal{O}_{\mathbf{k}}$ on \mathcal{F} and on $\mathrm{Lie}(A)/\mathcal{F}$ are through the structure morphism $i_S : \mathcal{O}_{\mathbf{k}} \rightarrow \mathcal{O}_S$ and its complex conjugate, respectively.

Remark 4.1.1. The stack $\mathcal{M}_{(n-1,1)}$ is denoted $\mathcal{M}_{(n-1,1)}^{\mathrm{Kra}}$ in [BHK⁺a].

Remark 4.1.2. One does not know a good theory of integral models $\mathcal{M}_{(n-1,1)}$ if 2 is ramified in \mathbf{k} , and this lack of knowledge is the main reason for restricting to quadratic imaginary fields of odd discriminant.

Definition 4.1.3. A finite dimensional \mathbf{k} -hermitian space W is *relevant* if it admits a self-dual $\mathcal{O}_{\mathbf{k}}$ -lattice $\mathfrak{a} \subset W$. Two relevant hermitian spaces W and W' are *strictly similar* if there is a \mathbf{k} -linear isomorphism $W \cong W'$ identifying the hermitian forms up to scaling by a positive rational number.

Proposition 4.1.4. *Let $o(D)$ denote the number of prime divisors of D .*

- (1) *Any two self-dual $\mathcal{O}_{\mathbf{k}}$ -lattices in a relevant \mathbf{k} -hermitian space are isometric everywhere locally.*
- (2) *There are $2^{o(D)-1}$ isomorphism classes of relevant hermitian spaces of any fixed signature (r, s) . If $r + s$ is odd all lie in the same strict similarity class, but if $r + s$ is even all lie in different strict similarity classes.*

Proof. The first claim is a theorem of Jacobowitz [Jac62], which uses our assumption that D is odd. The second claim is Lemma 2.11 of [KR14]. \square

²That is to say, an \mathcal{O}_S -module local direct summand of rank $n-1$

As in Proposition 2.12(i) of [KR14], there is a decomposition

$$(4.1.1) \quad \mathcal{M}_{(n-1,1)} = \bigsqcup_W \mathcal{M}_W,$$

where the disjoint union is over the strict similarity classes of relevant hermitian spaces W of signature $(n-1, 1)$, and the generic fiber of each \mathcal{M}_W is a Shimura variety of type $\mathrm{GU}(W)$. When n is odd, Proposition 4.1.4 implies that the disjoint union has a single term. When n is even, the disjoint union is over the $2^{\sigma(D)-1}$ isomorphism classes of relevant \mathbf{k} -hermitian spaces.

Denote by

$$\mathcal{M}_{(1,0)} \rightarrow \mathrm{Spec}(\mathcal{O}_{\mathbf{k}})$$

the moduli stack of elliptic curves with CM by $\mathcal{O}_{\mathbf{k}}$. More precisely, the functor of points assigns to an $\mathcal{O}_{\mathbf{k}}$ -scheme S the groupoid $\mathcal{M}_{(1,0)}(S)$ whose objects are pairs (A_0, ι_0) in which

- $A_0 \rightarrow S$ is an elliptic curve,
- $\iota_0 : \mathcal{O}_{\mathbf{k}} \rightarrow \mathrm{End}(A_0)$ is an $\mathcal{O}_{\mathbf{k}}$ -action satisfying the signature $(1, 0)$ condition: the induced action of $\mathcal{O}_{\mathbf{k}}$ on $\mathrm{Lie}(A_0)$ is through the structure morphism $i_S : \mathcal{O}_{\mathbf{k}} \rightarrow \mathcal{O}_S$.

As A_0 is an elliptic curve, there is a unique principal polarization

- $\psi_0 : A_0 \rightarrow A_0^\vee$,

and the induced Rosati involution on $\mathrm{End}(A_0)$ restricts to complex conjugation on the image of ι_0 . By Proposition 5.1 of [KRY99] or Proposition 2.1.2 of [How15], $\mathcal{M}_{(1,0)}$ is finite étale over $\mathcal{O}_{\mathbf{k}}$.

Remark 4.1.5. For ease of notation, we usually write objects of $\mathcal{M}_{(1,0)}$ simply as A_0 , suppressing the remaining data from the notation. Similarly, objects of $\mathcal{M}_{(n-1,1)}$ will usually be written simply as A .

When $n = 1$, the data of ψ and \mathcal{F} in the moduli problem $\mathcal{M}_{(n-1,1)}$ are uniquely determined, and $\mathcal{M}_{(0,1)}$ classifies pairs (A, ι) over $\mathcal{O}_{\mathbf{k}}$ -schemes S in which $A \rightarrow S$ is an elliptic curve and $\iota : \mathcal{O}_{\mathbf{k}} \rightarrow \mathrm{End}(A)$ is an action such that the induced action of $\mathcal{O}_{\mathbf{k}}$ on the \mathcal{O}_S -module $\mathrm{Lie}(A)$ is through the conjugate of the structure morphism $\mathcal{O}_{\mathbf{k}} \rightarrow \mathcal{O}_S$. Replacing ι by its conjugate therefore defines a canonical isomorphism

$$\mathcal{M}_{(0,1)} \cong \mathcal{M}_{(1,0)}.$$

In particular, there is a decomposition of $\mathcal{M}_{(1,0)}$ analogous to (4.1.1), but this statement is vacuous, as the decomposition has a single term by Proposition 4.1.4. If W_0 denotes the unique, up to strict similarity, relevant \mathbf{k} -hermitian space of signature $(1, 0)$, it will be convenient to set

$$\mathcal{M}_{W_0} = \mathcal{M}_{(1,0)}.$$

Fix relevant \mathbf{k} -hermitian spaces (W_0, h_0) and (W, h) of signatures $(1, 0)$ and $(n-1, 1)$, and self-dual $\mathcal{O}_{\mathbf{k}}$ -lattices $\mathfrak{a}_0 \subset W_0$ and $\mathfrak{a} \subset W$. The hermitian

forms induce alternating \mathbb{Q} -bilinear forms

$$(4.1.2) \quad e_0(a, b) = \mathrm{Tr}_{\mathbf{k}/\mathbb{Q}} \left(\frac{h_0(a, b)}{\sqrt{-D}} \right), \quad e(a, b) = \mathrm{Tr}_{\mathbf{k}/\mathbb{Q}} \left(\frac{h(a, b)}{\sqrt{-D}} \right)$$

on W_0 and W , and \mathfrak{a}_0 and \mathfrak{a} are self-dual with respect to these forms.

Definition 4.1.6. Suppose we are given an integer $N \geq 1$ and an $\mathcal{O}_{\mathbf{k}}$ -scheme S with $N \in \mathcal{O}_S^\times$. A *level N -structure* on $A_0 \in \mathcal{M}_{W_0}(S)$ is a pair (η_0, ζ_0) consisting of isomorphisms

$$\eta_0 : A_0[N] \cong \underline{\mathfrak{a}_0/N\mathfrak{a}_0}, \quad \zeta_0 : \mu_N \cong \underline{\mathbb{Z}/N\mathbb{Z}}$$

of étale sheaves on S that identify the Weil pairing on $A_0[N]$ induced by the principal polarization with the pairing on $\mathfrak{a}_0/N\mathfrak{a}_0$ induced by e_0 . Similarly, a *level N -structure* on $A \in \mathcal{M}_W(S)$ is a pair (η, ζ) consisting of isomorphisms

$$\eta : A[N] \cong \underline{\mathfrak{a}/N\mathfrak{a}}, \quad \zeta : \mu_N \cong \underline{\mathbb{Z}/N\mathbb{Z}}$$

that identify the Weil pairing on $A[N]$ with the alternating pairing on $\mathfrak{a}/N\mathfrak{a}$.

Over $\mathcal{O}_{\mathbf{k}}[1/N]$, we obtain finite étale covers

$$(4.1.3) \quad \mathcal{M}_{W_0}(N) \rightarrow \mathcal{M}_{W_0/\mathcal{O}_{\mathbf{k}}[1/N]}, \quad \mathcal{M}_W(N) \rightarrow \mathcal{M}_{W/\mathcal{O}_{\mathbf{k}}[1/N]}$$

by adding level structures to the moduli problems. By the first claim of Proposition 4.1.4, these are independent of the choices of \mathfrak{a}_0 and \mathfrak{a} .

4.2. The exceptional divisor. Now assume $n \geq 2$. We recall some more constructions from §2.3 of [BHK⁺a], following work of Pappas [Pap00] and Krämer [Kra03].

There is a normal and flat $\mathcal{O}_{\mathbf{k}}$ -stack

$$\mathcal{M}_{(n-1,1)}^{\mathrm{Pap}} \rightarrow \mathrm{Spec}(\mathcal{O}_{\mathbf{k}})$$

defined as the moduli space of triples (A, ι, ψ) whose components are as in the definition of $\mathcal{M}_{(n-1,1)}$. However, instead of including a hyperplane $\mathcal{F} \subset \mathrm{Lie}(A)$ as part of the moduli problem, we demand that the action of $\mathcal{O}_{\mathbf{k}}$ on $\mathrm{Lie}(A)$ satisfy a Kottwitz-style signature $(n-1, 1)$ condition, and the wedge conditions of Pappas. We refer the reader to §2.3 of [BHK⁺a] for the precise definitions.

Forgetting the hyperplane $\mathcal{F} \subset \mathrm{Lie}(A)$ defines a morphism

$$(4.2.1) \quad \mathcal{M}_{(n-1,1)} \rightarrow \mathcal{M}_{(n-1,1)}^{\mathrm{Pap}},$$

which can be realized as a blow-up. To interpret it as such, define the *singular locus*

$$(4.2.2) \quad \mathrm{Sing}_{(n-1,1)} \subset \mathcal{M}_{(n-1,1)}^{\mathrm{Pap}},$$

as the reduced locus of nonsmooth points. It is a proper $\mathcal{O}_{\mathbf{k}}$ -stack of dimension 0, supported in characteristics dividing D .

The morphism (4.2.1) can be identified with the blow-up of the singular locus, and so there is a canonical cartesian diagram

$$(4.2.3) \quad \begin{array}{ccc} \mathrm{Exc}_{(n-1,1)} & \longrightarrow & \mathcal{M}_{(n-1,1)} \\ \downarrow & & \downarrow \\ \mathrm{Sing}_{(n-1,1)} & \longrightarrow & \mathcal{M}_{(n-1,1)}^{\mathrm{Pap}} \end{array}$$

in which the upper left corner is the *exceptional divisor*. In particular, (4.2.1) is relatively representable and projective, and restricts to an isomorphism

$$\mathcal{M}_{(n-1,1)} \setminus \mathrm{Exc}_{(n-1,1)} \cong \mathcal{M}_{(n-1,1)}^{\mathrm{Pap}} \setminus \mathrm{Sing}_{(n-1,1)}.$$

Remark 4.2.1. If $\mathfrak{p} \subset \mathcal{O}_{\mathbf{k}}$ is a prime above $p \mid D$ with residue field $\mathbb{F}_{\mathfrak{p}}$, the $\mathbb{F}_{\mathfrak{p}}^{\mathrm{alg}}$ -points of the singular locus are those

$$A \in \mathcal{M}_{(n-1,1)}^{\mathrm{Pap}}(\mathbb{F}_{\mathfrak{p}}^{\mathrm{alg}})$$

for which the action of $\mathcal{O}_{\mathbf{k}}$ on the $\mathbb{F}_{\mathfrak{p}}^{\mathrm{alg}}$ -vector space $\mathrm{Lie}(A)$ is through the reduction map $\mathcal{O}_{\mathbf{k}} \rightarrow \mathbb{F}_{\mathfrak{p}}^{\mathrm{alg}}$. At such a point, *any* hyperplane $\mathcal{F} \subset \mathrm{Lie}(A)$ is $\mathcal{O}_{\mathbf{k}}$ -stable and satisfies Kramer's signature condition, and the fiber over A of the left vertical morphism in (4.2.3) is the projective space parametrizing all such \mathcal{F} .

4.3. Toroidal compactification. We turn to the study of compactifications of $\mathcal{M}_{(n-1,1)}$. When $n = 1$ this stack is finite (hence proper) over $\mathcal{O}_{\mathbf{k}}$, and so we set

$$\bar{\mathcal{M}}_{(0,1)} = \mathcal{M}_{(0,1)}.$$

For the rest of this subsection, assume $n \geq 2$.

One finds in §2 of [How15] the construction of a toroidal compactification

$$\mathcal{M}_{(n-1,1)} \subset \bar{\mathcal{M}}_{(n-1,1)},$$

obtained by imitating the constructions of Faltings and Chai [FC90]. If one works over $\mathcal{O}_{\mathbf{k}}[1/D]$, these are special cases of the constructions of Lan [Lan13]. The compactification is regular, and is smooth outside the exceptional divisor of (4.2.3).

The universal abelian scheme $A \rightarrow \mathcal{M}_{(n-1,1)}$ extends uniquely to a semi-abelian scheme

$$\bar{A} \rightarrow \bar{\mathcal{M}}_{(n-1,1)}$$

with $\mathcal{O}_{\mathbf{k}}$ -action. At a geometric point of the boundary, this semi-abelian scheme is an extension of an abelian variety by a torus of rank two. The universal hyperplane $\mathcal{F} \subset \mathrm{Lie}(A)$ over $\mathcal{M}_{(n-1,1)}$ extends uniquely to an $\mathcal{O}_{\mathbf{k}}$ -stable hyperplane

$$(4.3.1) \quad \bar{\mathcal{F}} \subset \mathrm{Lie}(\bar{A}),$$

which again satisfies Kramer's signature $(n - 1, 1)$ condition.

There is a similar toroidal compactification of $\mathcal{M}_{(n-1,1)}^{\text{Pap}}$ (now only normal instead of regular), and a Cartesian diagram

$$(4.3.2) \quad \begin{array}{ccc} \text{Exc}_{(n-1,1)} & \longrightarrow & \bar{\mathcal{M}}_{(n-1,1)} \\ \downarrow & & \downarrow \\ \text{Sing}_{(n-1,1)} & \longrightarrow & \bar{\mathcal{M}}_{(n-1,1)}^{\text{Pap}} \end{array}$$

extending (4.2.3). The vertical arrow on the right is the blow-up along the closed immersion in the bottom row.

Remark 4.3.1. For general Shimura varieties, a toroidal compactification depends on a choice of rational polyhedral cone decomposition of a convex cone sitting inside the real points of a \mathbb{Q} -vector space. For our $\text{GU}(n-1,1)$ Shimura varieties, the vector space in question is one-dimensional. Hence the rational polyhedral cone decomposition is uniquely determined, and the toroidal compactifications are canonical.

Let \mathcal{A}_n be the moduli stack of principally polarized abelian schemes of dimension n . There are canonical morphisms

$$(4.3.3) \quad \mathcal{M}_{(n-1,1)} \xrightarrow{(4.2.1)} \mathcal{M}_{(n-1,1)}^{\text{Pap}} \rightarrow \mathcal{A}_n/\mathcal{O}_{\mathbf{k}},$$

in which the first arrow is projective, and the second is finite.

Lemma 4.3.2. *If $\bar{\mathcal{A}}_n$ is any choice of smooth Faltings-Chai toroidal compactification, $\bar{\mathcal{M}}_{(n-1,1)}^{\text{Pap}}$ is canonically identified with the normalization of*

$$(4.3.4) \quad \mathcal{M}_{(n-1,1)}^{\text{Pap}} \rightarrow \bar{\mathcal{A}}_n/\mathcal{O}_{\mathbf{k}}.$$

See [Sta18, Tag 0BAK] for normalization.

Proof. By [How15], the universal abelian scheme $A \rightarrow \mathcal{M}_{(n-1,1)}^{\text{Pap}}$ extends (necessarily uniquely) to a semi-abelian scheme \bar{A} over the compactification. This extension satisfies a universal property: Suppose S is an irreducible normal scheme with generic point $\eta \rightarrow S$, and we are given a morphism

$$\eta \rightarrow \mathcal{M}_{(n-1,1)}^{\text{Pap}}.$$

If the pullback A_η extends to a semi-abelian scheme over S , then this extension is the pullback of \bar{A} via a (necessarily unique) morphism

$$S \rightarrow \bar{\mathcal{M}}_{(n-1,1)}^{\text{Pap}}$$

restricting to the given morphism at the generic point.

This extension property is analogous to Theorem IV.5.7(5) of [FC90], but the statement is simplified by the uniqueness of the rational polyhedral cone decomposition of Remark 4.3.1. If one works over $\mathcal{O}_{\mathbf{k}}[1/D]$, the extension property is a special case of Theorem 6.4.1.1(6) of [Lan13]. The proof over $\mathcal{O}_{\mathbf{k}}$ is exactly the same.

On the other hand, Theorem 11.4 of [Lan16] shows that the compactification of $\mathcal{M}_{(n-1,1)}^{\text{Pap}}$ defined by normalization also carries a semi-abelian scheme satisfying the same universal property, and the lemma follows. \square

Fix an \mathcal{M}_W as in (4.1.1). We need compactifications of the étale covers

$$\mathcal{M}_W(N) \rightarrow \mathcal{M}_{W/\mathcal{O}_k[1/N]}$$

of (4.1.3). Rather than repeat the constructions of [How15] with level structure, we will appeal to the results of [Lan16] and [MP19], in which integral models and compactifications of general Hodge-type Shimura varieties are constructed as normalizations of Faltings-Chai compactifications of Siegel moduli spaces. These results cannot be applied directly to the covers above, as the failure of the first arrow in (4.3.3) to be finite (it contracts the entire exceptional divisor to a 0-dimensional substack) implies that Lemma 4.3.2 is false if $\mathcal{M}_{(n-1,1)}^{\text{Pap}}$ is replaced by $\mathcal{M}_{(n-1,1)}$. Thus the proof of Proposition 4.3.3 below requires the roundabout step of first compactifying covers of $\mathcal{M}_{(n-1,1)}^{\text{Pap}}$.

For any $N \geq 1$, define $\bar{\mathcal{M}}_W(N)$ as the normalization of

$$\mathcal{M}_W(N) \rightarrow \bar{\mathcal{M}}_{(n-1,1)/\mathcal{O}_k[1/N]}.$$

Proposition 4.3.3. *The stack $\bar{\mathcal{M}}_W(N)$ is regular, flat, and proper over $\mathcal{O}_k[1/N]$, and is a projective scheme if $N \geq 3$. It is smooth away from its exceptional divisor*

$$\text{Exc}_{(n-1,1)} \times_{\bar{\mathcal{M}}_{(n-1,1)}} \bar{\mathcal{M}}_W(N).$$

In particular, it is smooth in a neighborhood of its boundary, which is a Cartier divisor smooth over $\mathcal{O}_k[1/N]$.

Proof. There is a decomposition

$$\mathcal{M}_{(n-1,1)}^{\text{Pap}} = \bigsqcup_W \mathcal{M}_W^{\text{Pap}},$$

exactly as in (4.1.1). We may add $N \geq 1$ level structure, exactly as in Definition 4.1.6, to obtain a finite étale cover

$$\mathcal{M}_W^{\text{Pap}}(N) \rightarrow \mathcal{M}_{W/\mathcal{O}_k[1/N]}^{\text{Pap}}.$$

This cover has a compactification $\bar{\mathcal{M}}_W^{\text{Pap}}(N)$ defined as the normalization of

$$\mathcal{M}_W^{\text{Pap}}(N) \rightarrow \bar{\mathcal{M}}_{(n-1,1)/\mathcal{O}_k[1/N]}^{\text{Pap}},$$

which, by Lemma 4.3.2, is the same as the normalization of

$$(4.3.5) \quad \mathcal{M}_W^{\text{Pap}}(N) \rightarrow \bar{\mathcal{A}}_{n/\mathcal{O}_k[1/N]}.$$

The properness, flatness, and normality of $\bar{\mathcal{M}}_W^{\text{Pap}}(N)$ follow from its construction as the normalization of a proper, flat, and smooth $\mathcal{O}_k[1/N]$ -stack.

It is not regular (as this is not true even of its interior). However, its singular locus

$$\mathrm{Sing}_W(N) = \mathrm{Sing}_{(n-1,1)} \times_{\mathcal{M}_{(n-1,1)}^{\mathrm{Pap}}} \mathcal{M}_W^{\mathrm{Pap}}(N)$$

is 0-dimensional and proper over $\mathcal{O}_{\mathbf{k}}[1/N]$, and the smoothness of

$$\mathcal{M}_W^{\mathrm{Pap}}(N) \setminus \mathrm{Sing}_W(N)$$

implies the smoothness of

$$\bar{\mathcal{M}}_W^{\mathrm{Pap}}(N) \setminus \mathrm{Sing}_W(N).$$

This is a consequence of the following principle, found in §14 of [Lan16] and Theorem 1 of [MP19]: if $U \subset \bar{\mathcal{M}}_W^{\mathrm{Pap}}(N)$ is an open neighborhood of the boundary $\partial \bar{\mathcal{M}}_W^{\mathrm{Pap}}(N)$, then the singularities of U are no worse than the singularities of $U \setminus \partial \bar{\mathcal{M}}_W^{\mathrm{Pap}}(N)$.

By the same principle (and its proof), the smoothness of $\bar{\mathcal{M}}_W^{\mathrm{Pap}}(N)$ near the boundary implies the smoothness of the boundary divisor itself. This is actually a particular feature of our $\mathrm{GU}(W)$ Shimura varieties. For more general Shimura varieties the boundary has a canonical stratification by locally closed substacks (see Theorem 4.1.5 of [MP19] or §9 of [Lan16]), and the above cited principle tells us that each *stratum* is smooth. In our special case each boundary stratum is a divisor, and is simply a union of connected components of $\partial \bar{\mathcal{M}}_W^{\mathrm{Pap}}(N)$. Hence the entire boundary is a Cartier divisor smooth over $\mathcal{O}_{\mathbf{k}}[1/N]$.

Using the characterization of (4.3.2) as a blow-up, one may identify $\bar{\mathcal{M}}_W(N)$ with the blow-up of $\bar{\mathcal{M}}_W^{\mathrm{Pap}}(N)$ along its singular locus. As the singular locus does not meet the boundary, it follows that $\bar{\mathcal{M}}_W(N)$ is itself proper and flat with boundary a smooth Cartier divisor. Moreover, this shows that $\bar{\mathcal{M}}_W(N)$ is smooth away from the exceptional divisor of the blow-up, and in particular in an open neighborhood of its boundary. As the interior is regular, being étale over the regular stack \mathcal{M}_W , we find that $\bar{\mathcal{M}}_W(N)$ is regular.

Finally, when $N \geq 3$ we claim that both the source and target of

$$\bar{\mathcal{M}}_W(N) \rightarrow \bar{\mathcal{M}}_W^{\mathrm{Pap}}(N)$$

are projective schemes. As this morphism is a blow-up, it suffices to prove this for the target. For this, we use the results of Chapter V.5 of [FC90]. Adding principal level structure yields a finite étale cover

$$\mathcal{A}_n(N) \rightarrow \mathcal{A}_{n/\mathbb{Z}[1/N]},$$

and the compactification $\bar{\mathcal{A}}_n$ may be chosen so that (4.3.5) factors as

$$\mathcal{M}_W^{\mathrm{Pap}}(N) \rightarrow \bar{\mathcal{A}}_n(N) \rightarrow \bar{\mathcal{A}}_{n/\mathcal{O}_{\mathbf{k}}[1/N]},$$

where $\bar{\mathcal{A}}_n(N)$ is the smooth projective scheme obtained by normalizing

$$\mathcal{A}_n(N) \rightarrow \bar{\mathcal{A}}_{n/\mathcal{O}_{\mathbf{k}}[1/N]}.$$

This realizes $\bar{\mathcal{M}}_W^{\text{Pap}}(N)$ as the normalization of the first arrow in the composition, and hence provides us with a finite and relatively representable morphism

$$\bar{\mathcal{M}}_W^{\text{Pap}}(N) \rightarrow \bar{\mathcal{A}}_n(N).$$

As the target is a projective scheme, so is the source. \square

Remark 4.3.4. One could avoid the constructions of [How15] and [Lan13] altogether, and define $\bar{\mathcal{M}}_{(n-1,1)}$ from the start as a blow-up of the normalization of (4.3.4). All properties that we need to know about this compactification could then be deduced from [Lan16] and [MP19], except for the extension (4.3.1) of the universal hyperplane across the boundary (which is needed in the proof of Proposition 5.2.1). For this one needs detailed information about the local charts near the boundary, as found in [How15].

4.4. Arithmetic Chow groups. We summarize the arithmetic intersection theory of Gillet-Soulé [GS90, SABK] and Burgos-Kramer-Kühn [BBK07, BKK07] for the Shimura varieties \mathcal{M}_W of (4.1.1), mostly for the purposes of fixing normalizations of arithmetic degrees, heights, and volumes.

As usual, denote by $\widehat{\text{Pic}}(\mathcal{M}_W)$ the group of isomorphism classes of hermitian line bundles on \mathcal{M}_W , and similarly for the toroidal compactification $\bar{\mathcal{M}}_W$. In order to account for metrics that do not extend smoothly across the boundary, when we work on the compactification $\bar{\mathcal{M}}_W$ we must relax the notion of a hermitian metric.

Accordingly, denote by $\widehat{\text{Pic}}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}})$ the group of isomorphism classes of pre-log singular hermitian line bundles on $\bar{\mathcal{M}}_W$, in the sense of Definition 1.20 of [BBK07]. Exactly as in Proposition 5.2.1 of [How20], the natural maps

$$(4.4.1) \quad \widehat{\text{Pic}}(\bar{\mathcal{M}}_W) \rightarrow \widehat{\text{Pic}}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}) \rightarrow \widehat{\text{Pic}}(\mathcal{M}_W)$$

are injective.

Fix an integer $N \geq 3$, so that the toroidal compactification $\bar{\mathcal{M}}_W(N)$ of Proposition 4.3.3 is a regular scheme of dimension n , projective and flat over $\mathcal{O}_k[1/N]$. For $0 \leq d \leq n$, we have a homomorphism of arithmetic Chow groups

$$(4.4.2) \quad \widehat{\text{CH}}_{\text{GS}}^d(\bar{\mathcal{M}}_W(N)) \rightarrow \widehat{\text{CH}}^d(\bar{\mathcal{M}}_W(N), \mathcal{D}_{\text{pre}})$$

from Theorem 6.23 and (7.49) of [BKK07].

The domain of (4.4.2) is the arithmetic Chow group of Gillet-Soulé [GS90, SABK], defined as the rational equivalence classes of arithmetic cycles (\mathcal{Z}, g) in which \mathcal{Z} is a codimension d cycle on $\bar{\mathcal{M}}_W(N)$, and g is a $(d-1, d-1)$ current on $\bar{\mathcal{M}}_W(N)(\mathbb{C})$ satisfying the Green equation

$$dd^c g + \delta_{\mathcal{Z}} = [\omega]$$

of currents for some smooth (d, d) form ω .

The codomain of (4.4.2) is the pre-log-log arithmetic Chow group of Burgos-Kramer-Kühn [BBK07, BKK07]. It is defined as rational equivalence classes of arithmetic cycles $(\mathcal{Z}, \mathfrak{g})$ where \mathcal{Z} is as above, and \mathfrak{g} is as in Definition 1.15 of [BBK07]. For our purposes it is enough to know that $\mathfrak{g} = (\omega, g)$ is a pair consisting of a $(d-1, d-1)$ current on $\bar{\mathcal{M}}_W(N)(\mathbb{C})$ and a smooth (d, d) form ω on the interior $\mathcal{M}_W(N)(\mathbb{C})$, again related by the above Green equation (by Proposition 7.44 of [BKK07]). There are further constraints: The form ω is required to have pre-log-log growth along the boundary of the toroidal compactification (which implies local integrability on the compactification, so that the Green equation makes sense). The current g is required to be represented by a smooth form defined on the complement of some codimension d cycle on $\mathcal{M}_W(N)(\mathbb{C})$, again satisfying certain growth conditions. Moreover, the relation imposed between g and ω is stronger than the Green equation. Nevertheless, the object \mathfrak{g} is completely determined by the current g , so we usually just write (\mathcal{Z}, g) for the arithmetic cycle.

Whenever $N \mid N'$ the map $\bar{\mathcal{M}}_W(N') \rightarrow \bar{\mathcal{M}}_W(N)$ induces a pullback

$$\widehat{\text{CH}}^d(\bar{\mathcal{M}}_W(N), \mathcal{D}_{\text{pre}}) \rightarrow \widehat{\text{CH}}^d(\bar{\mathcal{M}}_W(N'), \mathcal{D}_{\text{pre}}),$$

allowing us to define, exactly as in §6.3 of [BKK07],

$$\widehat{\text{CH}}^d(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}) = \varprojlim_{N \geq 3} \widehat{\text{CH}}^d(\bar{\mathcal{M}}_W(N), \mathcal{D}_{\text{pre}}).$$

There is an *arithmetic intersection pairing*

$$\widehat{\text{CH}}^{d_1}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}) \otimes \widehat{\text{CH}}^{d_2}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}) \rightarrow \widehat{\text{CH}}^{d_1+d_2}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}})_{\mathbb{Q}},$$

an *arithmetic Chern class map*

$$\widehat{\text{Pic}}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}) \rightarrow \widehat{\text{CH}}^1(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}),$$

and an *arithmetic degree*

$$\widehat{\text{deg}} : \widehat{\text{CH}}^n(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}) \rightarrow \mathbb{R}$$

induced by the analogous structures on the arithmetic Chow group of each scheme $\bar{\mathcal{M}}_W(N)$ with $N \geq 3$.

Remark 4.4.1. The conventions for delta currents, currents associated to smooth forms, and Green forms used in [BKK07] and [BBK07] differ from those of Gillet-Soulé [GS90, SABK] by powers of 2 and $2\pi i$. We will always use the conventions of Gillet-Soulé. For example, the arithmetic Chern class map sends a pre-log singular hermitian line bundle $\widehat{\mathcal{L}}$ to the arithmetic divisor

$$\widehat{\text{div}}(s) = (\text{div}(s), -\log \|s\|^2)$$

for any nonzero rational section s .

To make explicit the normalization of the arithmetic degree, first note that for each $N \geq 3$, the finite étale cover $\mathcal{M}_W(N) \rightarrow \mathcal{M}_{W/\mathcal{O}_k[1/N]}$ has constant fiber degree, in the sense that there is a $d_N \in \mathbb{Z}$ satisfying

$$(4.4.3) \quad \frac{d_N}{\#\text{Aut}(x)} = \#\{\text{geometric points } y \rightarrow \mathcal{M}_W(N) \text{ above } x\}$$

for every geometric point $x \rightarrow \mathcal{M}_{W/\mathcal{O}_k[1/N]}$. Now suppose we have a class

$$\widehat{\mathcal{Z}} = (\mathcal{Z}_N, g_N)_{N \geq 3} \in \widehat{\text{CH}}^n(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}).$$

If we write each $\mathcal{Z}_N = \sum_i m_{N,i} \mathcal{Z}_{N,i}$ as a \mathbb{Z} -linear combination of 0-dimensional irreducible closed subschemes on $\bar{\mathcal{M}}_W(N)$, then the real number

$$\widehat{\text{deg}}_N(\mathcal{Z}_N, g_N) = \sum_i m_{N,i} \cdot \#\mathcal{Z}_{N,i}(\mathbb{F}_i^{\text{alg}}) \cdot \log(\#\mathbb{F}_i) + \int_{\mathcal{M}_W(N)(\mathbb{C})} g_N,$$

where \mathbb{F}_i is the residue field of the unique prime of $\mathcal{O}_k[1/N]$ below $\mathcal{Z}_{N,i}$, is well-defined up to adding a \mathbb{Q} -linear combination of $\{\log(p) : p \mid N\}$, and

$$(4.4.4) \quad \widehat{\text{deg}}(\widehat{\mathcal{Z}}) = \frac{1}{d_N} \cdot \widehat{\text{deg}}_N(\mathcal{Z}_N, g_N)$$

up to the same ambiguity.

Remark 4.4.2. There is no $1/2$ in front of the integral, in apparent disagreement with §3.4.3 of [GS90]. In fact there is no disagreement. For us $\mathcal{M}_W(N)(\mathbb{C})$ means the set of all morphisms $\text{Spec}(\mathbb{C}) \rightarrow \mathcal{M}_W(N)$ as k -schemes, whereas Gillet-Soulé would take morphisms as \mathbb{Q} -schemes. Thus for Gillet-Soulé the domain of integration would be our $\mathcal{M}_W(N)(\mathbb{C})$ *together with its complex conjugate*, yielding an integral twice as big as ours.

We will also need the *complex degree*

$$\text{deg}_{\mathbb{C}} : \widehat{\text{CH}}^{n-1}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}) \rightarrow \mathbb{Q}$$

sending a class

$$\widehat{\mathcal{Z}} = (\mathcal{Z}_N, g_N)_{N \geq 3} \in \widehat{\text{CH}}^{n-1}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}})$$

to

$$\text{deg}_{\mathbb{C}}(\widehat{\mathcal{Z}}) = \frac{1}{d_N} \cdot \text{deg}(\mathcal{Z}_N/\mathbb{C}),$$

where the degree on the right hand side is the usual degree of a 0-cycle on the complex fiber $\bar{\mathcal{M}}_W(N)_{/\mathbb{C}}$. Equivalently,

$$(4.4.5) \quad \text{deg}_{\mathbb{C}}(\widehat{\mathcal{Z}}) = \frac{1}{d_N} \cdot \int_{\mathcal{M}_W(N)(\mathbb{C})} \omega_N$$

is the integral of the $(n-1, n-1)$ form determined by the Green equation.

$$dd^c g_N + \delta_{\mathcal{Z}_N} = [\omega_N].$$

4.5. **Volumes and heights.** For a pre-log singular hermitian line bundle

$$\widehat{\mathcal{L}} \in \widehat{\text{Pic}}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}),$$

define the *complex volume*

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}) = \text{deg}_{\mathbb{C}}(\widehat{\mathcal{L}}^{n-1})$$

by applying the complex degree of §4.4 to the $(n-1)$ -fold iterated arithmetic intersection of the arithmetic Chern class

$$\widehat{\mathcal{L}} \in \widehat{\text{CH}}^1(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}).$$

Of course this only depends on the restriction of $\widehat{\mathcal{L}}$ to a line bundle on the complex fiber $\bar{\mathcal{M}}_{W/\mathbb{C}}$. Similarly, define the *arithmetic volume*

$$\widehat{\text{vol}}(\widehat{\mathcal{L}}) = \widehat{\text{deg}}(\widehat{\mathcal{L}}^n)$$

using the iterated arithmetic intersection and arithmetic degree of §4.4.

Remark 4.5.1. The pre-log singularity of the metric on $\widehat{\mathcal{L}}$ implies the integrability of its Chern form: the smooth $(1, 1)$ form on $\mathcal{M}_W(\mathbb{C})$ defined locally by

$$\text{ch}(\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}) = \frac{1}{2\pi i} \partial \bar{\partial} \log \|s\|^2 = -dd^c \log \|s\|^2$$

for any nonzero holomorphic section s . We will frequently use the equivalent characterization of the complex volume

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}) = \int_{\mathcal{M}_W(\mathbb{C})} \text{ch}(\widehat{\mathcal{L}})^{n-1}$$

as the (orbifold) integral of its top exterior power, which follows from (4.4.5).

Let \mathcal{Z} be a divisor on $\bar{\mathcal{M}}_W$, and assume that \mathcal{Z} intersects the boundary $\partial \bar{\mathcal{M}}_W$ properly in the generic fiber. As in the discussion following Theorem 7.58 of [BKK07], one can then define the *arithmetic height* (or *Faltings height*) of \mathcal{Z} with respect to $\widehat{\mathcal{L}}$, denoted

$$\text{ht}_{\widehat{\mathcal{L}}}(\mathcal{Z}) \in \mathbb{R}.$$

To make it explicit, fix $N \geq 3$. Choose a pair $(\mathcal{Y}, g_{\mathcal{Y}})$ representing the iterated intersection

$$\widehat{\mathcal{L}}^{n-1} \in \widehat{\text{CH}}^{n-1}(\bar{\mathcal{M}}_W(N), \mathcal{D}_{\text{pre}}),$$

and do this in such a way that the supports of \mathcal{Z}_N and \mathcal{Y} do not intersect in the generic fiber, where \mathcal{Z}_N is the pullback of \mathcal{Z} via

$$\bar{\mathcal{M}}_W(N) \rightarrow \bar{\mathcal{M}}_{W/\mathcal{O}_k[1/N]}.$$

Now form the intersection

$$\mathcal{Z}_N \cdot \mathcal{Y} \in \text{CH}_{\mathcal{Z}_N \cap \mathcal{Y}}^n(\bar{\mathcal{M}}_W(N))_{\mathbb{Q}}$$

in the Chow group with support along the closed subset $\text{Sppt}(\mathcal{Z}_N) \cap \text{Sppt}(\mathcal{Y})$. As this closed subset is supported in finitely many nonzero characteristics, there is a natural change-of-support map

$$\text{CH}_{\mathcal{Z} \cap \mathcal{Y}}^n(\bar{\mathcal{M}}_W(N)) \rightarrow \bigoplus_{\substack{\mathfrak{p} \subset \mathcal{O}_k \\ \mathfrak{p} \nmid N\mathcal{O}_k}} \text{CH}_{\mathfrak{p}}^n(\bar{\mathcal{M}}_W(N))$$

where $\text{CH}_{\mathfrak{p}}^n(\bar{\mathcal{M}}_W(N))$ is the Chow group with support in the mod \mathfrak{p} fiber of $\bar{\mathcal{M}}_W(N)$. For each \mathfrak{p} there is a degree map

$$(4.5.1) \quad \text{deg}_{\mathfrak{p}} : \text{CH}_{\mathfrak{p}}^n(\bar{\mathcal{M}}_W(N)) \rightarrow \mathbb{Z}$$

sending a reduced and irreducible codimension n subscheme (a.k.a., a closed point) $\mathcal{C} \subset \bar{\mathcal{M}}_W(N)$ supported at \mathfrak{p} to

$$\text{deg}_{\mathfrak{p}}(\mathcal{C}) = \#\mathcal{C}(\mathbb{F}_{\mathfrak{p}}^{\text{alg}})$$

where $\mathbb{F}_{\mathfrak{p}} = \mathcal{O}_k/\mathfrak{p}$. The arithmetic height satisfies

$$(4.5.2) \quad d_N \cdot \text{ht}_{\hat{\mathcal{L}}}(\mathcal{Z}) = \sum_{\substack{\mathfrak{p} \subset \mathcal{O}_k \\ \mathfrak{p} \nmid N\mathcal{O}_k}} \text{deg}_{\mathfrak{p}}(\mathcal{Z}_N \cdot \mathcal{Y}) \cdot \log(\#\mathbb{F}_{\mathfrak{p}}) + \int_{\mathcal{Z}_N(\mathbb{C})} g_{\mathcal{Y}}$$

up to a \mathbb{Q} -linear combination of $\{\log(p) : p \mid N\}$, where d_N is as in (4.4.3).

A choice of pre-log singular hermitian metric on the line bundle $\mathcal{O}(\mathcal{Z})$ determines a class

$$\hat{\mathcal{Z}} \in \widehat{\text{Pic}}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}).$$

If we view the constant function 1 as a global section of $\mathcal{O}(\mathcal{Z})$ and set $g_{\mathcal{Z}} = -\log \|1\|^2$, the arithmetic Chern class of this hermitian line bundle is

$$(\mathcal{Z}, g_{\mathcal{Z}}) \in \widehat{\text{CH}}^1(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}),$$

and Proposition 7.56 of [BKK07] gives the relation

$$(4.5.3) \quad \widehat{\text{deg}}(\hat{\mathcal{Z}} \cdot \hat{\mathcal{L}}^{n-1}) = \text{ht}_{\hat{\mathcal{L}}}(\mathcal{Z}) + \int_{\mathcal{M}_W(\mathbb{C})} g_{\mathcal{Z}} \cdot \text{ch}(\hat{\mathcal{L}})^{n-1}.$$

From the point of view of arithmetic volumes and heights, some hermitian line bundles are essentially indistinguishable.

Definition 4.5.2. We say that

$$\hat{\mathcal{L}}_1, \hat{\mathcal{L}}_2 \in \widehat{\text{Pic}}(\mathcal{M}_W)$$

are *numerically equivalent* if the difference $\hat{\mathcal{E}} = \hat{\mathcal{L}}_1 - \hat{\mathcal{L}}_2$ lies in the subgroup

$$\widehat{\text{Pic}}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}) \subset \widehat{\text{Pic}}(\mathcal{M}_W)$$

of (4.4.1) and satisfies $\widehat{\text{deg}}(\hat{\mathcal{Z}} \cdot \hat{\mathcal{E}}) = 0$ for all $\hat{\mathcal{Z}} \in \widehat{\text{CH}}^{n-1}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}})$.

Lemma 4.5.3. *If $\widehat{\mathcal{L}}_1$ and $\widehat{\mathcal{L}}_2$ are numerically equivalent, then their Chern forms are equal. Moreover,*

$$(4.5.4) \quad \widehat{\text{vol}}(\widehat{\mathcal{L}}_1) = \widehat{\text{vol}}(\widehat{\mathcal{L}}_2) \quad \text{and} \quad \text{ht}_{\widehat{\mathcal{L}}_1}(\mathcal{Z}) = \text{ht}_{\widehat{\mathcal{L}}_2}(\mathcal{Z})$$

for any divisor \mathcal{Z} as above.

Proof. Set $\widehat{\mathcal{E}} = \widehat{\mathcal{L}}_1 - \widehat{\mathcal{L}}_2$. If g is any smooth $(n-2, n-2)$ form on $\bar{\mathcal{M}}_W(\mathbb{C})$, the arithmetic cycle class

$$\widehat{\mathcal{Z}} = (0, g) \in \widehat{\text{CH}}^{n-1}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}})$$

satisfies

$$0 = \widehat{\text{deg}}(\widehat{\mathcal{Z}} \cdot \widehat{\mathcal{E}}) = \int_{\mathcal{M}_W(\mathbb{C})} \text{ch}(\widehat{\mathcal{E}}) \wedge g.$$

As this holds for all choices of g , we must have $\text{ch}(\widehat{\mathcal{E}}) = 0$. This proves the desired equality of Chern forms. The first equality of (4.5.4) is clear from

$$\widehat{\text{vol}}(\widehat{\mathcal{L}}_1) = \sum_{i=0}^n \widehat{\text{deg}}(\widehat{\mathcal{L}}_2^{n-i} \cdot \widehat{\mathcal{E}}^i),$$

as every term with $i > 0$ vanishes. The second equality follows from (4.5.3) and the equality of Chern forms already proved. \square

If \mathcal{Z} is a Cartier divisor on $\bar{\mathcal{M}}_W$ supported in nonzero characteristics, the line bundle $\mathcal{O}(\mathcal{Z})$ is canonically trivialized in the generic fiber by the constant function 1. For any real number c , denote by

$$(4.5.5) \quad (\mathcal{Z}, c) \in \widehat{\text{Pic}}(\bar{\mathcal{M}}_W)$$

the line bundle $\mathcal{O}(\mathcal{Z})$ endowed with the constant metric characterized by $-\log \|1\|^2 = c$. In particular, adding $(0, c)$ to a hermitian line bundle just rescales $\|\cdot\|^2$ by e^{-c} . The following lemma describes the effect of this on arithmetic volumes.

Lemma 4.5.4. *For any $\widehat{\mathcal{L}} \in \widehat{\text{Pic}}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}})$ and any constant $c \in \mathbb{R}$, we have*

$$\widehat{\text{vol}}(\widehat{\mathcal{L}} + (0, c)) = \widehat{\text{vol}}(\widehat{\mathcal{L}}) + nc \int_{\mathcal{M}_W(\mathbb{C})} \text{ch}(\widehat{\mathcal{L}})^{n-1},$$

where $\text{ch}(\widehat{\mathcal{L}})$ is the Chern form of Remark 4.5.1.

Proof. A trivial induction argument shows that

$$\widehat{\mathcal{L}}^{k-1} \cdot (0, c) = (0, c \cdot \text{ch}(\widehat{\mathcal{L}})^{k-1}) \in \widehat{\text{CH}}^k(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}})$$

for all $k > 0$. As $(0, c) \cdot (0, c) = 0$ we deduce

$$\begin{aligned} (\widehat{\mathcal{L}} + (0, c))^k &= \widehat{\mathcal{L}}^k + k\widehat{\mathcal{L}}^{k-1} \cdot (0, c) \\ &= \widehat{\mathcal{L}}^k + k(0, c \cdot \text{ch}(\widehat{\mathcal{L}})^{k-1}). \end{aligned}$$

Setting $k = n$ and taking the arithmetic degree completes the proof. \square

5. A SHIMURA VARIETY AND ITS LINE BUNDLES

For the rest of this paper we fix a \mathbf{k} -hermitian space V of signature $(n-1, 1)$, with $n \geq 1$, and assume that V admits a self-dual $\mathcal{O}_{\mathbf{k}}$ -lattice (1.5.1). The hermitian form on V is denoted $\langle -, - \rangle$. We will associate to V an n -dimensional open and closed substack

$$\mathcal{S}_V \subset \mathcal{M}_{(1,0)} \times_{\mathcal{O}_{\mathbf{k}}} \mathcal{M}_{(n-1,1)},$$

construct some interesting hermitian line bundles on it, and explain the relations between them.

5.1. A special Shimura variety. To attach a Shimura variety to our fixed V , choose relevant (Definition 4.1.3) hermitian spaces (W_0, h_0) and (W, h) of signatures $(1, 0)$ and $(n-1, 1)$, respectively, in such a way that

$$(5.1.1) \quad V \cong \mathrm{Hom}_{\mathbf{k}}(W_0, W)$$

as \mathbf{k} -hermitian spaces, where the hermitian form $\langle -, - \rangle$ on the right is defined by the relation

$$(5.1.2) \quad \langle x, y \rangle \cdot h_0(w_0, w'_0) = h(x(w_0), y(w'_0))$$

for all $w_0, w'_0 \in W_0$ and $x, y \in V$.

Remark 5.1.1. Such W_0 and W always exist, and in fact there is a distinguished choice: take $W_0 = \mathbf{k}$ with its norm form, and $W = V$. Any other choice of W_0 will be strictly similar to this distinguished choice (this is a trivial case of Proposition 4.1.4), and the relation (5.1.1) then forces W to be strictly similar to V . In particular, the Shimura varieties

$$\mathcal{M}_{W_0} \cong \mathcal{M}_{(1,0)} \quad \text{and} \quad \mathcal{M}_W \cong \mathcal{M}_V$$

depend only on V , and not on the particular W_0 and W chosen in (5.1.1).

As in §2.2 of [KR14], if S is a connected $\mathcal{O}_{\mathbf{k}}$ -scheme and

$$(A_0, A) \in \mathcal{M}_{W_0}(S) \times \mathcal{M}_W(S),$$

then $\mathrm{Hom}_{\mathcal{O}_{\mathbf{k}}}(A_0, A)$ carries a positive definite hermitian form

$$(5.1.3) \quad \langle x, y \rangle = \psi_0^{-1} \circ y^\vee \circ \psi \circ x \in \mathrm{End}_{\mathcal{O}_{\mathbf{k}}}(A_0) \cong \mathcal{O}_{\mathbf{k}},$$

where $\psi_0 : A_0 \rightarrow A_0^\vee$ and $\psi : A \rightarrow A^\vee$ are the principal polarizations.

As in §2.3 of [BHK⁺a], there is an open and closed substack

$$(5.1.4) \quad \mathcal{S}_V \subset \mathcal{M}_{W_0} \times_{\mathcal{O}_{\mathbf{k}}} \mathcal{M}_W$$

characterized by its points valued in algebraically closed fields, which are those pairs

$$(A_0, A) \in \mathcal{M}_{W_0}(F) \times \mathcal{M}_W(F)$$

for which there exists an isometry

$$V \otimes \mathbb{Q}_\ell \cong \mathrm{Hom}_{\mathcal{O}_{\mathbf{k}}}(T_\ell(A_0), T_\ell(A)) \otimes \mathbb{Q}_\ell$$

of \mathbf{k}_ℓ -hermitian spaces for every $\ell \neq \mathrm{char}(F)$. Here the hermitian form on the right is defined as in (5.1.3). When $F = \mathbb{C}$ this is equivalent (by

the Hasse principle for hermitian spaces) to the existence of an isometry of \mathbf{k} -hermitian spaces

$$V \cong \mathrm{Hom}_{\mathbf{k}}(H_1(A_0, \mathbb{Q}), H_1(A, \mathbb{Q})),$$

where the hermitian form on the right hand side is again defined by imitating the construction (5.1.3) in Betti homology.

Remark 5.1.2. The projection $\mathcal{S}_V \rightarrow \mathcal{M}_W$ is a finite étale surjection, and the fiber over a geometric point $x \in \mathcal{M}_W(F)$ satisfies

$$\sum_{y \in \mathcal{S}_{V,x}(F)} \frac{1}{|\mathrm{Aut}(y)|} = \frac{|\mathrm{CL}(\mathbf{k})|}{|\mathcal{O}_{\mathbf{k}}^\times|} \cdot \begin{cases} 1 & \text{if } n \text{ is even} \\ 2^{1-o(D)} & \text{if } n \text{ is odd.} \end{cases}$$

Remark 5.1.3. The stack \mathcal{S}_V is denoted $\mathcal{S}_{\mathrm{Kra}}$ in [BHK⁺a]. Later we will want to vary V , and so we have included it in the notation to avoid confusion. As explained in [loc. cit.], the generic fiber of \mathcal{S}_V is a Shimura variety for the subgroup $G \subset \mathrm{GU}(W_0) \times \mathrm{GU}(W)$ of pairs (g_0, g) for which the similitude factors of the two components are equal.

Definition 5.1.4. If $n \geq 2$, define the *exceptional divisor*

$$\mathrm{Exc}_V \subset \mathcal{S}_V$$

as the pullback of the exceptional divisor (4.2.3) via $\mathcal{S}_V \rightarrow \mathcal{M}_{(n-1,1)}$. Equivalently, it is defined by the cartesian diagram

$$(5.1.5) \quad \begin{array}{ccc} \mathrm{Exc}_V & \longrightarrow & \mathcal{S}_V \\ \downarrow & & \downarrow \\ \mathrm{Sing}_{(n-1,1)} & \longrightarrow & \mathcal{M}_{(n-1,1)}^{\mathrm{Pap}}. \end{array}$$

Definition 5.1.5 ([KR14]). For any positive $m \in \mathbb{Z}$, the *Kudla-Rapoport divisor* $\mathcal{Z}_V(m)$ is the $\mathcal{O}_{\mathbf{k}}$ -stack classifying triples (A_0, A, x) consisting of a pair

$$(A_0, A) \in \mathcal{S}_V(S)$$

and an $x \in \mathrm{Hom}_{\mathcal{O}_{\mathbf{k}}}(A_0, A)$ satisfying $\langle x, x \rangle = m$.

The natural forgetful morphism

$$\mathcal{Z}_V(m) \rightarrow \mathcal{S}_V$$

is finite and unramified, with image of codimension 1. Denote by

$$\bar{\mathcal{Z}}_V(m) \rightarrow \bar{\mathcal{S}}_V$$

the normalization of $\mathcal{Z}_V(m) \rightarrow \mathcal{S}_V$.

Remark 5.1.6. The morphism $\mathcal{Z}_V(m) \rightarrow \mathcal{S}_V$ determines a Weil divisor on \mathcal{S}_V in the usual way: each irreducible component of $\mathcal{Z}_V(m)$ contributes an irreducible component to the divisor, counted with multiplicity equal to the length of the local ring of its generic point. By abuse of notation, we denote

this Weil divisor again by $\mathcal{Z}_V(m)$. Similarly there is a divisor determined by $\bar{\mathcal{Z}}_V(m)$, denoted the same way; it agrees with the Zariski closure in $\bar{\mathcal{S}}_V$ of the divisor $\mathcal{Z}_V(m)$.

Loosely speaking, each Kudla-Rapoport divisor is a union of unitary Shimura varieties associated to \mathbf{k} -hermitian spaces of signature $(n-2, 1)$. In §6 we will make this more precise, at least when m is a prime split in \mathbf{k} .

Definition 5.1.7. Let N be a positive integer, and let S be an $\mathcal{O}_{\mathbf{k}}$ -scheme with $N \in \mathcal{O}_S^\times$. A *level N -structure* on a pair $(A_0, A) \in \mathcal{S}_V(S)$ consists of level N -structures (η_0, ξ_0) and (η, ξ) on A_0 and A , in the sense of Definition 4.1.6, such that $\xi_0 = \xi$.

Adding level structure to pairs (A_0, A) defines a finite étale cover

$$\mathcal{S}_V(N) \rightarrow \mathcal{S}_{V/\mathcal{O}_{\mathbf{k}}[1/N]},$$

and (5.1.4) lifts to a canonical open and closed immersion

$$\mathcal{S}_V(N) \subset \mathcal{M}_{W_0}(N) \times_{\mathcal{O}_{\mathbf{k}}[1/N]} \mathcal{M}_W(N).$$

Define a toroidal compactification

$$(5.1.6) \quad \bar{\mathcal{S}}_V(N) \subset \mathcal{M}_{W_0}(N) \times_{\mathcal{O}_{\mathbf{k}}[1/N]} \bar{\mathcal{M}}_W(N)$$

as the Zariski closure of $\mathcal{S}_V(N)$. When $N = 1$ we abbreviate

$$\bar{\mathcal{S}}_V = \bar{\mathcal{S}}_V(1).$$

Note that $\bar{\mathcal{S}}_V(N)$ is also characterized as the normalization of

$$\mathcal{S}_V(N) \rightarrow \bar{\mathcal{M}}_{W/\mathcal{O}_{\mathbf{k}}[1/N]}.$$

Remark 5.1.8. As (5.1.6) is an open and closed immersion, and $\mathcal{M}_{W_0}(N)$ is finite étale over $\mathcal{O}_{\mathbf{k}}[1/N]$, the compactification $\bar{\mathcal{S}}_V(N)$ inherits all the nice properties of $\bar{\mathcal{M}}_W(N)$. In particular, Proposition 4.3.3 holds word-for-word with \mathcal{M}_W replaced by \mathcal{S}_V , and the same is true of the entire discussion of arithmetic intersection theory in §4.4 and §4.5.

5.2. Construction of hermitian line bundles. The construction (1.1.1) associates to the universal CM elliptic curve $A_0 \rightarrow \mathcal{M}_{W_0} = \mathcal{M}_{(1,0)}$ its metrized Hodge bundle

$$(5.2.1) \quad \widehat{\omega}_{A_0/\mathcal{M}_{W_0}}^{\text{Hdg}} \in \widehat{\text{Pic}}(\mathcal{M}_{W_0}).$$

Similarly, the universal $A \rightarrow \mathcal{M}_W$ determines a metrized Hodge bundle

$$(5.2.2) \quad \widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}} \in \widehat{\text{Pic}}(\mathcal{M}_W).$$

Pulling back the universal objects via projection to the two factors in (5.1.4) yields a universal pair (A_0, A) of polarized abelian schemes (of relative dimensions 1 and n) over \mathcal{S}_V . As in §2.4 of [BHK⁺a] there is a *metrized line bundle of modular forms*

$$(5.2.3) \quad \widehat{\mathcal{L}}_V \in \widehat{\text{Pic}}(\mathcal{S}_V).$$

The line bundle underlying (5.2.3) has inverse

$$(5.2.4) \quad \mathcal{L}_V^{-1} = \text{Lie}(A_0) \otimes \text{Lie}(A)/\mathcal{F},$$

where $\mathcal{F} \subset \text{Lie}(A)$ is the universal hyperplane satisfying Krämer's signature condition (§4.1). The hermitian metric is defined as in §7.2 of [BHK⁺a]: if we use Proposition 2.4.2 of [BHK⁺a] to identify

$$(5.2.5) \quad \mathcal{L}_{V,z} \subset \text{Hom}_{\mathbf{k}}(H_1(A_{0,z}, \mathbb{Q}), H_1(A_z, \mathbb{Q})) \otimes_{\mathbb{Q}} \mathbb{C} \cong V \otimes_{\mathbb{Q}} \mathbb{C}$$

at a complex point $z \in \mathcal{S}_V(\mathbb{C})$, the line $\mathcal{L}_{V,z}$ is isotropic with respect to the \mathbb{C} -bilinear extension of the \mathbb{Q} -bilinear form $[x, y] = \text{Tr}_{\mathbf{k}/\mathbb{Q}}\langle x, y \rangle$ on V , and

$$(5.2.6) \quad \|s\|^2 = -\frac{[s, \bar{s}]}{4\pi e^\gamma}$$

for any $s \in \mathcal{L}_{V,z}$. Note that our $\widehat{\mathcal{L}}_V$ is denoted $\widehat{\omega}$ in [BHK⁺a, BHK⁺b].

Proposition 5.2.1. *The hermitian line bundles (5.2.2) and (5.2.3) lie in the subgroups*

$$\widehat{\text{Pic}}(\bar{\mathcal{M}}_W, \mathcal{D}_{\text{pre}}) \subset \widehat{\text{Pic}}(\mathcal{M}_W) \quad \text{and} \quad \widehat{\text{Pic}}(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}}) \subset \widehat{\text{Pic}}(\mathcal{S}_V),$$

respectively, of (4.4.1).

Proof. The extension to $\bar{\mathcal{M}}_W$ of the line bundle

$$\omega_{A/\mathcal{M}_W}^{\text{Hdg}} \cong \det(\text{Lie}(A))^{-1}$$

underlying (5.2.2) is part of (4.3.1). The fact that the hermitian metric has a pre-log singularity along the boundary is a special case of Theorem 6.16 of [BKK05]. Note that this also uses Proposition 3.2 of [Fr09], which shows that any rank one log-singular hermitian vector bundle in the sense of [BKK05] is also a pre-log-singular hermitian line bundle in the sense of Definition 1.20 of [BBK07]. This proves the claim for (5.2.2), and the proof for (5.2.3) is the same. \square

Pulling back the hermitian line bundles (5.2.1) and (5.2.2) via projection to the two factors in (5.1.6), we obtain three hermitian line bundles

$$(5.2.7) \quad \widehat{\omega}_{A_0/\mathcal{S}_V}^{\text{Hdg}}, \widehat{\omega}_{A/\mathcal{S}_V}^{\text{Hdg}}, \widehat{\mathcal{L}}_V \in \widehat{\text{Pic}}(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}}).$$

In the remaining subsections we will make explicit the relations between them. The reader may wish to skip directly to Theorem 5.5.1 for the main results.

5.3. An application of the Chowla-Selberg formula. We will prove that, up to numerical equivalence (Definition 4.5.2), the first line bundle in (5.2.7) is just the trivial line bundle with a constant metric. The constant defining the metric is an interesting quantity in its own right. Recall the Chowla-Selberg formula: the Faltings height, normalized as in §5.2 of [BHK⁺b], of any elliptic curve with CM by $\mathcal{O}_{\mathbf{k}}$ is

$$(5.3.1) \quad h_{\mathbf{k}}^{\text{Falt}} = -\frac{1}{2} \frac{L'(0, \varepsilon)}{L(0, \varepsilon)} - \frac{1}{4} \log(4\pi^2 D).$$

Proposition 5.3.1. *The metrized line bundles*

$$\widehat{\omega}_{A_0/\mathcal{S}_V}^{\text{Hdg}} \in \widehat{\text{Pic}}(\mathcal{S}_V) \quad \text{and} \quad (0, C_1) \in \widehat{\text{Pic}}(\mathcal{S}_V)$$

are numerically equivalent, where $C_1 = \log(2\pi) + 2h_{\mathbf{k}}^{\text{Falt}}$, and we are using the notation of (4.5.5).

Proof. Fix an $N \geq 3$. As $\mathcal{M}_{W_0}(N)$ is finite étale over $\mathcal{O}_{\mathbf{k}}[1/N]$, there is an isomorphism

$$\mathcal{M}_{W_0}(N) \cong \bigsqcup_i \mathcal{X}_i$$

with each $\mathcal{X}_i \cong \text{Spec}(\mathcal{O}_{\mathbf{k}_i}[1/N])$ for a finite field extension \mathbf{k}_i/\mathbf{k} unramified outside N . For each \mathcal{X}_i , consider the metrized Hodge bundle

$$\widehat{\omega}_{A_0/\mathcal{X}_i}^{\text{Hdg}} \in \widehat{\text{Pic}}(\mathcal{X}_i)$$

of the universal $A_0 \rightarrow \mathcal{X}_i$. The underlying line bundle can be identified with an element in the ideal class group of $\mathcal{O}_{\mathbf{k}_i}[1/N]$, and hence some power of it admits a trivializing section

$$s_i \in H^0(\mathcal{X}_i, (\omega_{A_0/\mathcal{X}_i}^{\text{Hdg}})^{\otimes d_i}) \cong H^0(A_0, \Omega_{A_0/\mathcal{X}_i}^{\otimes d_i}).$$

Comparing (1.1.1) with the definition of the Faltings height (as in §5.2 of [BHK⁺b]) shows that

$$\frac{-1}{\#\mathcal{X}_i(\mathbb{C})} \sum_{x \in \mathcal{X}_i(\mathbb{C})} \log \|s_{i,x}\|^2 = d_i \cdot C_1,$$

up to a \mathbb{Q} -linear combination of $\{\log(p) : p \mid N\}$.

Letting d be the least common multiple of all d_i 's, we see that

$$(\widehat{\omega}_{A_0/\mathcal{S}_V(N)}^{\text{Hdg}})^{\otimes d} \in \widehat{\text{Pic}}(\bar{\mathcal{S}}_V(N))$$

admits a trivializing section s with $-\log \|s\|^2$ constant on every connected component of $\bar{\mathcal{S}}_V(N)(\mathbb{C})$, and such that the average value of $-\log \|s\|^2$ over any $\text{Aut}(\mathbb{C}/\mathbf{k})$ -orbit of components is $d \cdot C_1$, up to a \mathbb{Q} -linear combination of $\{\log(p) : p \mid N\}$.

If we represent

$$d \cdot \widehat{\omega}_{A_0/\mathcal{S}_V(N)}^{\text{Hdg}} \in \widehat{\text{CH}}^1(\bar{\mathcal{S}}_V(N))$$

by the arithmetic divisor $\widehat{\text{div}}(s) = (0, -\log \|s\|^2)$, then for any

$$(\mathcal{Z}_N, g_N) \in \widehat{\text{CH}}^{n-1}(\bar{\mathcal{S}}_V(N))$$

the vanishing of the Chern form of $-\log \|s\|^2$ implies the $*$ -product formula

$$[-\log \|s\|^2] * g_N = -\log \|s\|^2 \wedge \delta_{\mathcal{Z}_N},$$

which implies the intersection formula

$$d \cdot \widehat{\text{deg}}_N((\mathcal{Z}_N, g_N) \cdot \widehat{\omega}_{A_0/\mathcal{S}_V(N)}^{\text{Hdg}}) = - \sum_{x \in \mathcal{Z}_N(\mathbb{C})} \log \|s_x\|^2$$

up to a \mathbb{Q} -linear combination of $\{\log(p) : p \mid N\}$.

Let $L \subset \mathbb{C}$ be a finite Galois extension of \mathbf{k} large enough that all complex points of \mathcal{Z}_N are defined over L , and rewrite the equality above as

$$d \cdot \widehat{\deg}_N((\mathcal{Z}_N, g_N) \cdot \widehat{\omega}_{A_0/S_V(N)}^{\text{Hdg}}) = \frac{-1}{[L : \mathbf{k}]} \sum_{\substack{x \in \mathcal{Z}_N(L) \\ \sigma \in \text{Gal}(L/\mathbf{k})}} \log \|s_{x\sigma}\|^2.$$

The right hand side is $dC_1 \cdot \#\mathcal{Z}_N(\mathbb{C})$, and hence

$$\widehat{\deg}_N((\mathcal{Z}_N, g_N) \cdot \widehat{\omega}_{A_0/S_V(N)}^{\text{Hdg}}) = C_1 \cdot \#\mathcal{Z}_N(\mathbb{C}) = \widehat{\deg}_N((\mathcal{Z}_N, g_N) \cdot (0, C_1))$$

up to a \mathbb{Q} -linear combination of $\{\log(p) : p \mid N\}$. Varying N shows that

$$\widehat{\deg}(\widehat{\mathcal{Z}} \cdot \widehat{\omega}_{A_0/S_V}^{\text{Hdg}}) = \widehat{\deg}(\widehat{\mathcal{Z}} \cdot (0, C_1))$$

for every $\widehat{\mathcal{Z}} \in \widehat{\text{CH}}^{n-1}(\bar{\mathcal{S}}_V)$, and the claim follows. \square

5.4. Another hermitian line bundle. In order to relate the hermitian line bundles of (5.2.7), we recall from §5.1 of [BHK⁺b] a fourth hermitian line bundle. Denote by

$$H_{\text{dR}}^1(A) = \mathbb{R}^1 \pi_* \Omega_{A/S_V}^\bullet$$

the first relative algebraic deRham cohomology of $\pi : A \rightarrow \mathcal{S}_V$, a rank $2n$ vector bundle on \mathcal{S}_V . The action of $\mathcal{O}_{\mathbf{k}}$ on A induces an action on $H_{\text{dR}}^1(A) = H_{\text{dR}}^1(A)^\vee$, which is locally free of rank n over $\mathcal{O}_{\mathbf{k}} \otimes_{\mathbb{Z}} \mathcal{O}_S$. If

$$H_1^{\text{dR}}(A) \rightarrow \mathcal{V}$$

denotes the largest quotient on which the action of $\mathcal{O}_{\mathbf{k}}$ is through the structure morphism $\mathcal{O}_{\mathbf{k}} \rightarrow \mathcal{O}_{\mathcal{S}_V}$, then \mathcal{V} is a rank n vector bundle on \mathcal{S}_V , equipped with a morphism

$$\det(\mathcal{V})^{-1} \rightarrow \bigwedge^n H_{\text{dR}}^1(A) \rightarrow H_{\text{dR}}^n(A).$$

Given a complex point $z \in \mathcal{S}_V(\mathbb{C})$ and a vector $s_z \in \det(\mathcal{V}_z)^{-1}$, we view s_z as an element of $H^n(A_z, \mathbb{C})$, and define

$$\|s_z\|^2 = \left| \int_{A_z(\mathbb{C})} s_z \wedge \bar{s}_z \right|.$$

This defines a hermitian metric on $\det(\mathcal{V})^{-1}$, and hence we obtain a hermitian line bundle

$$(5.4.1) \quad \det(\mathcal{V}) \in \widehat{\text{Pic}}(\mathcal{S}_V).$$

Proposition 5.4.1. *Assume $n \geq 2$. Recalling the exceptional divisor of Definition 5.1.4 and the notation (4.5.5), we have the equality*

$$2\widehat{\mathcal{L}}_V = \widehat{\omega}_{A/S_V}^{\text{Hdg}} + 2\widehat{\omega}_{A_0/S_V}^{\text{Hdg}} + \det(\mathcal{V}) + (\text{Exc}_V, C_2)$$

in $\widehat{\text{Pic}}(\mathcal{S}_V)$, where

$$C_2 = 2 \log \left(\frac{2e^\gamma}{D} \right) + (2 - n) \log(2\pi).$$

In particular, $\det(\mathcal{V}) \in \widehat{\text{Pic}}(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})$ by (5.2.7).

Proof. This is a restatement of Proposition 5.1.2 of [BHK⁺b], keeping in mind that the metric on

$$\det(\text{Lie}(A))^{-1} = \omega_{A/\bar{\mathcal{S}}_V}^{\text{Hdg}}$$

used in [loc. cit.] differs from (1.1.1) by a power of 2π . \square

It is an observation of Gross [Gro78] that the hermitian line bundle $\det(\mathcal{V})$ behaves in the generic fiber, for all arithmetic purposes, like the trivial bundle with a constant metric. This observation was extended to integral models in §5.3 of [BHK⁺b], whose results are the basis of the following proposition.

Proposition 5.4.2. *The Chern form of $\det(\mathcal{V})$ is identically 0. If $n > 2$ then, up to numerical equivalence,*

$$\det(\mathcal{V}) = (0, C_3),$$

where

$$C_3 = (4 - 2n)h_{\mathbf{k}}^{\text{Falt}} + \log(4\pi^2 D).$$

Proof. Fix an integer $N \geq 3$. As in Theorem 1 of [Gro78], the \mathbb{C} -vector space $H^0(\mathcal{S}_V(N)_{/\mathbb{C}}, \det(\mathcal{V}))$ has dimension 1, and the norm of any global section is a locally constant function on $\mathcal{S}_V(N)(\mathbb{C})$. The vanishing of the Chern form follows. Moreover, one can choose a nonzero global section t defined over a finite Galois extension \mathbf{k}'/\mathbf{k} , and then for every \mathbf{k} -algebra embedding $\sigma : \mathbf{k}' \rightarrow \mathbb{C}$ the norm $\|t^\sigma\|$ must again be locally constant.

If $n > 2$ then³ Theorem 5.3.1 of [BHK⁺b] allows us to choose t so that it extends to a nowhere vanishing section

$$t \in H^0(\mathcal{S}_V(N)_{/\mathcal{O}_{\mathbf{k}'[1/N]}}, \det(\mathcal{V})).$$

Setting $d = [\mathbf{k}' : \mathbf{k}]$ and taking the tensor product of all $\text{Gal}(\mathbf{k}'/\mathbf{k})$ -conjugates of t , we obtain a section

$$s \in H^0(\mathcal{S}_V(N), \det(\mathcal{V})^{\otimes d})$$

such that $\text{div}(s) = 0$, and such that $-\log \|s\|$ is locally constant. Let

$$c_X = -\log \|s\|^2$$

denote its value on the connected component $X \subset \mathcal{S}_V(N)_{/\mathbb{C}}$.

Fix a finite Galois extension L/\mathbf{k} contained \mathbb{C} large enough that every component $X \subset \mathcal{S}_V(N)_{/\mathbb{C}}$ is defined over L . We may further enlarge L to assume that each X admits an L -point

$$x \in X(L) \subset \mathcal{S}_V(N)(L)$$

³The assumption $n > 2$ is mistakenly omitted in Theorem 5.3.1 of [BHK⁺b], whose proof requires that $\mathcal{M}_{(n-1,1)}^{\text{Pap}}$ has geometrically normal fibers. The normality is a theorem of Pappas when $n > 2$, but is false when $n = 2$.

that extends to

$$\underline{x} : \text{Spec}(\mathcal{O}_L[1/N]) \rightarrow \mathcal{S}_V(N).$$

For example, start by fixing a complex point $x \in X(\mathbb{C})$ corresponding to a pair (A_0, A) for which A has complex multiplication. Then enlarge L so that both A_0 and A (along with their level N -structures) are defined over L and have everywhere good reduction.

Applying Corollary 5.3.2 and Proposition 5.3.3 of [BHK⁺b] to \underline{x} , we find

$$\frac{-1}{[L : \mathbf{k}]} \sum_{\sigma \in \text{Gal}(L/\mathbf{k})} \log \|s_{x^\sigma}\|^2 = dC_3$$

up to a \mathbb{Q} -linear combination of $\{\log(p) : p \mid N\}$, and hence

$$\frac{1}{[L : \mathbf{k}]} \sum_{\sigma \in \text{Gal}(L/\mathbf{k})} c_{X^\sigma} = dC_3$$

up to the same ambiguity.

We have now shown that the average value of $-\log \|s\|^2$ over any $\text{Aut}(\mathbb{C}/\mathbf{k})$ orbit of connected components in $\mathcal{S}_V(N)(\mathbb{C})$ is dC_3 , up to a \mathbb{Q} -linear combination of $\{\log(p) : p \mid N\}$. The proposition follows from this, by the same argument used in the proof of Proposition 5.3.1. \square

5.5. Comparison of hermitian line bundles. We now come to the main results of §5.

The exceptional divisor $\text{Exc}_V \subset \bar{\mathcal{S}}_V$ of Definition 5.1.4 is a recurring nuisance, in part because it has nontrivial arithmetic intersection with $\hat{\mathcal{L}}_V$. This will be explored more fully in §5.6 below. The following proposition allows us to avoid this nuisance by slightly modifying $\hat{\mathcal{L}}_V$, and also clarifies the relation between the second and third line bundles in (5.2.7).

Theorem 5.5.1. *Assume $n \geq 2$. Recalling the notation (4.5.5), the hermitian line bundle*

$$(5.5.1) \quad \hat{\mathcal{K}}_V \stackrel{\text{def}}{=} 2\hat{\mathcal{L}}_V - (\text{Exc}_V, 0) \in \widehat{\text{Pic}}(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})$$

enjoys the following properties.

- (1) *Every irreducible component $E \subset \text{Exc}_V$ satisfies*

$$(E, 0) \cdot \hat{\mathcal{K}}_V = 0$$

in $\widehat{\text{CH}}^2(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})_{\mathbb{Q}}$, as well as the height relation

$$\text{ht}_{\hat{\mathcal{K}}_V}(E) = 0.$$

The same equalities hold if we replace $\hat{\mathcal{K}}_V$ with $\hat{\omega}_{A/\mathcal{S}_V}^{\text{Hdg}}$ or $\hat{\omega}_{A_0/\mathcal{S}_V}^{\text{Hdg}}$.

- (2) *We have the equality of Chern forms*

$$\text{ch}(\hat{\omega}_{A/\mathcal{S}_V}^{\text{Hdg}}) = 2 \text{ch}(\hat{\mathcal{L}}_V) = \text{ch}(\hat{\mathcal{K}}_V).$$

(3) If $n > 2$ then, up to numerical equivalence,

$$\widehat{\mathcal{K}}_V = \widehat{\omega}_{A/\mathcal{S}_V}^{\text{Hdg}} + (0, C_0(n)) \in \widehat{\text{Pic}}(\mathcal{S}_V),$$

where $C_0(n)$ is the constant of Theorem A. In particular, up to numerical equivalence,

$$2\widehat{\mathcal{L}}_V = \widehat{\omega}_{A/\mathcal{S}_V}^{\text{Hdg}} + (\text{Exc}_V, C_0(n)).$$

Proof. For (1), the key observation is that the abelian scheme $A \rightarrow \mathcal{S}_V$ is a pullback via the vertical arrow on the right in (5.1.5). In particular, $\omega_{A/\mathcal{S}_V}^{\text{Hdg}}$ is isomorphic to the pullback of

$$\omega_{A/\mathcal{M}_{(n-1,1)}^{\text{Pap}}}^{\text{Hdg}} \in \text{Pic}(\mathcal{M}_{(n-1,1)}^{\text{Pap}}).$$

If $\mathcal{M}_{(n-1,1)}^{\text{Pap}}$ were a scheme we could trivialize this latter line bundle over a Zariski open neighborhood of the (0-dimensional) singular locus. This would pull-back to a trivialization of $\omega_{A/\mathcal{S}_V}^{\text{Hdg}}$ over an open neighborhood of Exc_V , and in particular over an open neighborhood of any irreducible component $E \subset \text{Exc}_V$.

To account for the stackiness, simply fix an integer $N \geq 3$ and apply the same reasoning with level N -structure to see that $\omega_{A/\mathcal{S}_V(N)}^{\text{Hdg}}$ is trivial in some Zariski open neighborhood of

$$E(N) = E \times_{\mathcal{S}_V} \mathcal{S}_V(N).$$

This implies the arithmetic intersection formula $(E(N), 0) \cdot \widehat{\omega}_{A/\mathcal{S}_V(N)}^{\text{Hdg}} = 0$ and varying N proves

$$(E, 0) \cdot \widehat{\omega}_{A/\mathcal{S}_V}^{\text{Hdg}} = 0.$$

The line bundle (5.4.1) is also a pullback via the vertical arrow on the right in (5.1.5), hence the same argument shows

$$(E, 0) \cdot \det(\mathcal{V}) = 0.$$

The proof of Proposition 5.3.1 shows that, for any $N \geq 3$, there is a positive multiple of

$$\widehat{\omega}_{A_0/\mathcal{S}_V(N)}^{\text{Hdg}} \in \widehat{\text{CH}}^1(\bar{\mathcal{S}}_V(N), \mathcal{D}_{\text{pre}})$$

that can be represented by a purely archimedean arithmetic divisor $(0, g)$. Any such arithmetic divisor satisfies $(E(N), 0) \cdot (0, g) = 0$, and varying N shows that

$$(E, 0) \cdot \widehat{\omega}_{A_0/\mathcal{S}_V}^{\text{Hdg}} = 0.$$

Rewriting the relation of Proposition 5.4.1 as

$$(5.5.2) \quad \widehat{\mathcal{K}}_V = \widehat{\omega}_{A/\mathcal{S}_V}^{\text{Hdg}} + 2\widehat{\omega}_{A_0/\mathcal{S}_V}^{\text{Hdg}} + \det(\mathcal{V}) + (0, C_2),$$

we have shown that the right hand side has trivial arithmetic intersection with $(E, 0)$, and hence so does the left hand side. This proves the first equality of (1). The second is a formal consequence of this and (4.5.3).

The first equality of (2) follows from (5.5.2), as the Chern forms of the final three terms on the right vanish by Lemma 4.5.3, Proposition 5.3.1, and Proposition 5.4.2. The second equality of (2) is clear from the definitions.

Claim (3) also follows from (5.5.2), using Proposition 5.3.1, Proposition 5.4.2, and the equality $C_0(n) = 2C_1 + C_2 + C_3$. \square

Remark 5.5.2. If $n > 2$ then part (3) of Theorem 5.5.1 implies

$$\begin{aligned} 2^n \cdot \widehat{\text{vol}}(\widehat{\mathcal{K}}_V) &= \widehat{\text{vol}}(\widehat{\omega}_{A/S_V}^{\text{Hdg}} + (0, C_0(n))) \\ &= \widehat{\text{vol}}(\widehat{\omega}_{A/S_V}^{\text{Hdg}}) + nC_0(n) \int_{\mathcal{S}_V(\mathbb{C})} \text{ch}(\widehat{\omega}_{A/S_V}^{\text{Hdg}})^{n-1} \end{aligned}$$

where we have used Lemma 4.5.3 for the first equality, and Lemma 4.5.4 for the second. In Proposition 6.7.1 we will show that this equality also holds when $n = 2$.

The final claim of Theorem 5.5.1 has the following variant.

Proposition 5.5.3. *Fix an $N \geq 3$. There is a positive integer r such that the r^{th} power of $\mathcal{L}_V^{\otimes 2}$ becomes isomorphic to the r^{th} power of $\omega_{A/S_V}^{\text{Hdg}} \otimes \mathcal{O}(\text{Exc}_V)$ after pulling both line bundles back via the morphism*

$$\bar{\mathcal{S}}_V(N) \rightarrow \bar{\mathcal{S}}_{V/\mathcal{O}_k[1/N]}.$$

Proof. We begin with the equality

$$2\widehat{\mathcal{L}}_V = \widehat{\omega}_{A/S_V}^{\text{Hdg}} + 2\widehat{\omega}_{A_0/S_V}^{\text{Hdg}} + \det(\mathcal{V}) + (\text{Exc}_V, C_2)$$

of metrized line bundles in $\widehat{\text{Pic}}(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})$ from Proposition 5.4.1. In particular, after forgetting the metrics we obtain the same equality in $\text{Pic}(\bar{\mathcal{S}}_V)$.

After pullback to $\bar{\mathcal{S}}_V(N)$, the proof of Proposition 5.3.1 shows that some power of $\omega_{A_0/S_V}^{\text{Hdg}}$ is trivial. Similarly, the proof of Proposition 5.4.2 shows that some power of $\det(\mathcal{V})$ admits a trivializing section s over the interior $\mathcal{S}_V(N)$ such that $\|s\|$ is a locally constant function on the complex fiber. This constancy of the norm implies that s , when viewed as a rational section on the toroidal compactification $\bar{\mathcal{S}}_V(N)$, still has no zeros or poles; the argument for this is exactly as in the proof of Proposition 5.2.1 of [How20], and relies on the pre-log singularity of the metric on $\det(\mathcal{V})$ proved in Proposition 5.4.1. Thus some power of $\det(\mathcal{V})$ is trivial, and the claim follows. \square

5.6. Volume of the exceptional divisor. In this subsection we assume $n \geq 2$, and fix an irreducible (= connected) component E of the exceptional divisor $\text{Exc}_V \subset \bar{\mathcal{S}}_V$ of Definition 5.1.4. Recalling the notation (4.5.5), our goal is to compute the arithmetic volume of

$$(E, 0) \in \widehat{\text{Pic}}(\bar{\mathcal{S}}_V).$$

By definition of the exceptional divisor, there is a commutative diagram with cartesian squares

$$(5.6.1) \quad \begin{array}{ccc} E & \longrightarrow & e \\ \downarrow & & \downarrow \\ \mathrm{Exc}_V & \longrightarrow & \mathcal{M}_{(1,0)} \times_{\mathcal{O}_{\mathbf{k}}} \mathrm{Sing}_{(n-1,1)} \\ \downarrow & & \downarrow \\ \mathcal{S}_V & \longrightarrow & \mathcal{M}_{(1,0)} \times_{\mathcal{O}_{\mathbf{k}}} \mathcal{M}_{(n-1,1)}^{\mathrm{Pap}} \end{array}$$

in which e is a connected component of the 0-dimensional reduced and irreducible $\mathcal{O}_{\mathbf{k}}$ -stack $\mathcal{M}_{(1,0)} \times_{\mathcal{O}_{\mathbf{k}}} \mathrm{Sing}_{(n-1,1)}$. In particular, e is supported in a single characteristic $p \mid D$, and admits a presentation as a stack quotient

$$e \cong \Delta \backslash \mathrm{Spec}(\mathbb{F}'_{\mathfrak{p}}),$$

in which $\mathfrak{p} \subset \mathcal{O}_{\mathbf{k}}$ is the prime above p , $\mathbb{F}'_{\mathfrak{p}}$ is a finite extension of $\mathbb{F}_{\mathfrak{p}} = \mathcal{O}_{\mathbf{k}}/\mathfrak{p}$, and Δ is a finite group acting on $\mathbb{F}'_{\mathfrak{p}}$. Define a rational number

$$m_E \stackrel{\mathrm{def}}{=} \sum_{z \in e(\mathbb{F}_{\mathfrak{p}}^{\mathrm{alg}})} \frac{1}{|\mathrm{Aut}(z)|} = \frac{[\mathbb{F}'_{\mathfrak{p}} : \mathbb{F}_{\mathfrak{p}}]}{|\Delta|}.$$

Proposition 5.6.1. *The iterated intersection*

$$\widehat{\mathcal{L}}_V^{-1} \cdots \widehat{\mathcal{L}}_V^{-1} \cdot (E, 0) \in \widehat{\mathrm{CH}}^n(\overline{\mathcal{S}}_V, \mathcal{D}_{\mathrm{pre}})$$

has arithmetic degree $\widehat{m}_E = m_E \log(p)$

Proof. After Remark 4.2.1, one might expect that E is isomorphic to the projective space of hyperplanes in an n -dimensional vector space. In particular, there should be a universal hyperplane $\mathcal{F} \subset \mathcal{O}_E^n$, with the property that the restriction of \mathcal{L}_V^{-1} to E is isomorphic to $\mathcal{O}_E^n/\mathcal{F}$.

The obstruction to this being true is due to stacky issues, which can be removed by fixing an integer $N \geq 3$ prime to p and adding level N -structure to the universal pair (A_0, A) over e . Let $e(N) \rightarrow e$ be the scheme classifying such level structures, and consider the cartesian diagram (this is the definition of the upper left corner)

$$\begin{array}{ccc} E(N) & \longrightarrow & e(N) \\ \downarrow & & \downarrow \\ E & \longrightarrow & e. \end{array}$$

The scheme $e(N)$, being a reduced scheme finite over $\mathbb{F}_{\mathfrak{p}}$, is a disjoint union of finitely many spectra of finite extensions of $\mathbb{F}_{\mathfrak{p}}$.

Fix a connected component $e' \subset e(N)$, and let

$$\mathrm{Spec}(\mathbb{F}'_{\mathfrak{p}}) = e' \subset e(N)$$

be the connected component below it. As in Remark 4.2.1, E' is precisely the projective space over e' classifying hyperplanes $\mathcal{F} \subset \text{Lie}(A|_{e'})$. Moreover, after fixing a trivialization $\text{Lie}(A_0|_{e'}) \cong \mathbb{F}'_{\mathfrak{p}}$, the isomorphism (5.2.4) identifies

$$\mathcal{L}_V^{-1}|_{E'} = \text{Lie}(A|_{E'})/\mathcal{F} \in \text{Pic}(E') \cong \text{CH}^1(E')$$

where $\mathcal{F} \subset \text{Lie}(A|_{E'})$ is the universal hyperplane. A routine exercise then shows that the iterated intersection

$$(\mathcal{L}_V^{-1}|_{E'}) \cdots (\mathcal{L}_V^{-1}|_{E'}) \in \text{CH}^{n-1}(E') \cong \mathbb{Z}$$

is represented by the cycle class of any $\mathbb{F}'_{\mathfrak{p}}$ -valued point of E' . In other words, the cycle class of any section to $E' \rightarrow e'$.

Now allow the connected component $E' \subset E(N)$ to vary. If we fix any section to $E(N) \rightarrow e(N)$, and use it to view $e(N)$ as a 0-cycle on $E(N)$, then

$$e(N) = (\mathcal{L}_V^{-1}|_{E(N)}) \cdots (\mathcal{L}_V^{-1}|_{E(N)}) \in \text{CH}^{n-1}(E(N)).$$

This implies the arithmetic intersection formula

$$(e(N), 0) = \widehat{\mathcal{L}}_V^{-1} \cdots \widehat{\mathcal{L}}_V^{-1} \cdot (E(N), 0) \in \widehat{\text{CH}}^n(\bar{\mathcal{S}}_V(N), \mathcal{D}_{\text{pre}})_{\mathbb{Q}}.$$

Finally, recalling (4.4.4), we deduce

$$\widehat{\text{deg}}(\widehat{\mathcal{L}}_V^{-1} \cdots \widehat{\mathcal{L}}_V^{-1} \cdot (E, 0)) = \frac{\#e(N)(\mathbb{F}_{\mathfrak{p}}^{\text{alg}})}{d_N} \log(p) = \sum_{z \in e(\mathbb{F}_{\mathfrak{p}}^{\text{alg}})} \frac{\log(p)}{|\text{Aut}(z)|}$$

up to a \mathbb{Q} -linear combination of $\{\log(p) : p \mid N\}$. Varying N completes the proof. \square

Corollary 5.6.2. *The hermitian line bundle $(E, 0) \in \widehat{\text{Pic}}(\bar{\mathcal{S}}_V)$ has arithmetic volume*

$$\widehat{\text{vol}}(E, 0) = (-2)^{n-1} \cdot \widehat{m}_E.$$

Proof. Part (1) of Theorem 5.5.1 implies the second equality in

$$(E, 0) \cdot (E, 0) = (\text{Exc}_V, 0) \cdot (E, 0) = 2\widehat{\mathcal{L}}_V \cdot (E, 0) \in \widehat{\text{CH}}^2(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})_{\mathbb{Q}}.$$

This implies the iterated intersection formula

$$(E, 0) \cdots (E, 0) = 2^{n-1} \widehat{\mathcal{L}}_V \cdots \widehat{\mathcal{L}}_V \cdot (E, 0) \in \widehat{\text{CH}}^n(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})_{\mathbb{Q}},$$

and claim follows using Proposition 5.6.1. \square

6. KUDLA-RAPOPORT DIVISORS AT SPLIT PRIMES

Let \mathcal{S}_V be the Shimura variety (5.1.4) associated to a \mathbf{k} -hermitian space V of signature $(n-1, 1)$ containing a self-dual lattice. Throughout this section we assume $n > 2$.

Fix a prime p split in \mathbf{k} , and factor

$$p\mathcal{O}_{\mathbf{k}} = \mathfrak{p}\bar{\mathfrak{p}}.$$

We explain how the Kudla-Rapoport divisor $\mathcal{Z}_V(p) \rightarrow \mathcal{S}_V$ of Definition 5.1.5 is related to a Shimura variety \mathcal{S}_{V^b} with V^b of signature $(n-2, 1)$.

6.1. Statement of the results. Let V^b denote the \mathbf{k} -hermitian space of signature $(n-2, 1)$ whose local invariants satisfy

$$(6.1.1) \quad \text{inv}_\ell(V^b) = (p, -D)_\ell \cdot \text{inv}_\ell(V)$$

for all places $\ell \leq \infty$. Using standard formulas for the Hilbert symbol, this is equivalent to

$$\text{inv}_\ell(V^b) = \left(\frac{p}{\ell}\right) \cdot \text{inv}_\ell(V).$$

Equivalently, V^b is the orthogonal complement to the \mathbf{k} -span of any $x \in V$ of hermitian norm $\langle x, x \rangle = p$. Obviously V^b depends on the prime p fixed above, but this prime will not change and we suppress the dependence from the notation.

Lemma 6.1.1. *The hermitian space V^b admits a self-dual $\mathcal{O}_{\mathbf{k}}$ -lattice (so is relevant in the sense of Definition 4.1.3).*

Proof. Suppose ℓ is a rational prime unramified in \mathbf{k} . If $\ell \neq p$ then $p \in \mathbb{Z}_\ell^\times$, and hence is a norm from the unramified extension \mathbf{k}_ℓ . If $\ell = p$ then p is again a norm from $\mathbf{k}_\ell \cong \mathbb{Q}_p \times \mathbb{Q}_p$. In either case, $(p, -D)_\ell = 1$, and so the local invariants of V and V^b agree at all unramified primes.

In general, a \mathbf{k} -hermitian space admits a self-dual $\mathcal{O}_{\mathbf{k}}$ -lattice if and only if it has local invariant 1 at every prime unramified in \mathbf{k} . As V satisfies this condition by hypothesis, so does V^b . \square

Lemma 6.1.1 allows us to form the $\mathcal{O}_{\mathbf{k}}$ -stacks

$$\mathcal{S}_{V^b} \subset \bar{\mathcal{S}}_{V^b} \subset \mathcal{M}_{(1,0)} \times_{\mathcal{O}_{\mathbf{k}}} \bar{\mathcal{M}}_{(n-2,1)}$$

analogous to

$$\mathcal{S}_V \subset \bar{\mathcal{S}}_V \subset \mathcal{M}_{(1,0)} \times_{\mathcal{O}_{\mathbf{k}}} \bar{\mathcal{M}}_{(n-1,1)},$$

but in one dimension lower. The stack \mathcal{S}_{V^b} is endowed with its own hermitian line bundle $\widehat{\mathcal{L}}_{V^b}$ as in (5.2.3), its own exceptional divisor Exc_{V^b} as in Definition 5.1.4, and its own universal pair (A_0^b, A^b) of polarized abelian schemes with $\mathcal{O}_{\mathbf{k}}$ -actions.

The following theorems, whose proofs will occupy the remainder of §6, lie at the core of the inductive arguments of §8. In both theorems, $\bar{\mathcal{Z}}_V(p)$ is the divisor on $\bar{\mathcal{S}}_V$ of Remark 5.1.6, and the arithmetic heights and volumes are those of §4.5.

Theorem 6.1.2. *If we set*

$$\widehat{\mathcal{K}}_V = 2\widehat{\mathcal{L}}_V - (\text{Exc}_V, 0) \in \widehat{\text{Pic}}(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})$$

as in Theorem 5.5.1, and similarly with V replaced by V^b , then

$$\int_{\bar{\mathcal{Z}}_V(p)(\mathbb{C})} \text{ch}(\widehat{\mathcal{K}}_V)^{n-2} = (p^{n-1} + 1) \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_{V^b}).$$

Moreover, there is a rational number $a(p)$ such that

$$\frac{\mathrm{ht}_{\widehat{\mathcal{K}}_V}(\bar{\mathcal{Z}}_V(p))}{p^{n-1} + 1} = \widehat{\mathrm{vol}}(\widehat{\mathcal{K}}_{V^b}) + a(p) \log(p).$$

Remark 6.1.3. The proof of Theorem 6.1.2 gives no hint as to the value of the rational number $a(p)$. It will be determined later in Theorem 8.4.2.

Theorem 6.1.4. *There is a rational number $b(p)$ such that*

$$\begin{aligned} \frac{\mathrm{ht}_{\widehat{\omega}_{A/S_V}^{\mathrm{Hdg}}}(\bar{\mathcal{Z}}_V(p))}{p^{n-1} + 1} &= \widehat{\mathrm{vol}}(\widehat{\omega}_{A^b/S_{V^b}}^{\mathrm{Hdg}}) + b(p) \log(p) \\ &+ (1 - n) \left(\frac{L'(0, \varepsilon)}{L(0, \varepsilon)} + \frac{\log(D)}{2} \right) \mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A^b/S_{V^b}}^{\mathrm{Hdg}}). \end{aligned}$$

The proofs are rather long, so we summarize now the key steps. The central idea is to make precise the impressionistic relation

$$\mathcal{Z}_V(p) \text{ “} = \text{” } (p^{n-1} + 1) \cdot \mathcal{S}_{V^b},$$

by decomposing

$$(6.1.2) \quad \mathcal{Z}_V(p)_{/\mathcal{O}_{\mathbf{k}}[1/p]} = \mathcal{U}_0 \sqcup \mathcal{U}_{\mathfrak{p}} \sqcup \mathcal{U}_{\bar{\mathfrak{p}}}$$

as a disjoint union of open and closed substacks (Proposition 6.2.2 below). The stacks on the right hand side are related to \mathcal{S}_{V^b} by closed immersions

$$i_{\mathfrak{p}} : \mathcal{S}_{V^b}/\mathcal{O}_{\mathbf{k}}[1/p] \rightarrow \mathcal{U}_{\mathfrak{p}}, \quad i_{\bar{\mathfrak{p}}} : \mathcal{S}_{V^b}/\mathcal{O}_{\mathbf{k}}[1/p] \rightarrow \mathcal{U}_{\bar{\mathfrak{p}}},$$

and by a diagram

$$\mathcal{S}_{V^b}/\mathcal{O}_{\mathbf{k}}[1/p] \longleftarrow \mathcal{T}_{V^b} \xrightarrow{i_0} \mathcal{U}_0$$

in which the leftward arrow is a finite étale surjection of degree $p^{n-1} - 1$, and the rightward arrow is a closed immersion. See (6.5.11) for the definition of the stack \mathcal{T}_{V^b} .

Recalling the exceptional locus of Definition 5.1.4, denote by

$$\mathcal{S}_V^{\mathrm{nexc}} = \mathcal{S}_V \setminus \mathrm{Exc}_V$$

the (open) nonexceptional locus of \mathcal{S}_V , set

$$\mathcal{Z}_V^{\mathrm{nexc}}(p) = \mathcal{Z}_V(p) \times_{\mathcal{S}_V} \mathcal{S}_V^{\mathrm{nexc}},$$

and make the same definitions with V replaced by V^b . Set

$$\mathcal{U}_{\square}^{\mathrm{nexc}} = \mathcal{U}_{\square} \cap \mathcal{Z}_V^{\mathrm{nexc}}(p)$$

for $\square \in \{0, \mathfrak{p}, \bar{\mathfrak{p}}\}$, and

$$\mathcal{T}_{V^b}^{\mathrm{nexc}} = \mathcal{T}_{V^b} \times_{\mathcal{S}_{V^b}} \mathcal{S}_{V^b}^{\mathrm{nexc}}.$$

We will show that the closed immersions i_0 , $i_{\mathfrak{p}}$, and $i_{\bar{\mathfrak{p}}}$ are close to being isomorphisms. More precisely, i_0 restricts to an isomorphism

$$i_0 : \mathcal{T}_{V^b}^{\mathrm{nexc}} \cong \mathcal{U}_0^{\mathrm{nexc}},$$

while $i_{\mathfrak{p}}$ and $i_{\bar{\mathfrak{p}}}$ restrict to isomorphisms

$$i_{\mathfrak{p}} : \mathcal{S}_{V^{\flat}/\mathcal{O}_k[1/p]}^{\text{nexc}} \cong \mathcal{U}_{\mathfrak{p}}^{\text{nexc}} \quad \text{and} \quad i_{\bar{\mathfrak{p}}} : \mathcal{S}_{V^{\flat}/\mathcal{O}_k[1/p]}^{\text{nexc}} \cong \mathcal{U}_{\bar{\mathfrak{p}}}^{\text{nexc}}.$$

After taking compactifications into account, both theorems above will follow easily from these isomorphisms. Note that it suffices to work only over the nonexceptional locus, as part (1) of Theorem 5.5.1 guarantees that the hermitian line bundles in questions have trivial arithmetic intersection with all components of the exceptional divisor. This why we work with $\widehat{\mathcal{K}}_V$ in Theorem 6.1.2 instead of $\widehat{\mathcal{L}}_V$.

6.2. Decomposing the Kudla-Rapoport divisor. The decomposition (6.1.2) is a geometric reflection of a result of linear algebra.

As in (5.1.1), choose an isomorphism

$$V \cong \text{Hom}_{\mathbf{k}}(W_0, W)$$

in which (W_0, h_0) and (W, h) are relevant hermitian spaces of signatures $(1, 0)$ and $(n-1, 1)$. Fix self-dual $\mathcal{O}_{\mathbf{k}}$ -lattices $\mathfrak{a}_0 \subset W_0$ and $\mathfrak{a} \subset W$. Any vector

$$x \in \text{Hom}_{\mathcal{O}_{\mathbf{k}}}(\mathfrak{a}_0, \mathfrak{a}) \subset V$$

with $\langle x, x \rangle = p$ determines an orthogonal decomposition

$$W = \tilde{W}_0 \oplus W^{\flat}$$

with $\tilde{W}_0 = x(W_0)$, and a corresponding decomposition

$$V = \mathbf{k}x \oplus V^{\flat}$$

with $V^{\flat} = \text{Hom}_{\mathbf{k}}(W_0, W^{\flat})$.

Lemma 6.2.1. *The $\mathcal{O}_{\mathbf{k}}$ -lattice $\tilde{\mathfrak{a}}_0 = \mathfrak{a} \cap \tilde{W}_0$ satisfies exactly one of*

$$\tilde{\mathfrak{a}}_0 = x(\mathfrak{a}_0), \quad \tilde{\mathfrak{a}}_0 = \mathfrak{p}^{-1}x(\mathfrak{a}_0), \quad \tilde{\mathfrak{a}}_0 = \bar{\mathfrak{p}}^{-1}x(\mathfrak{a}_0).$$

Proof. Use x to identify $W_0 = \tilde{W}_0$, and hence $\mathfrak{a}_0 \subset \tilde{\mathfrak{a}}_0 \subset \mathfrak{a}$. Recalling the symplectic forms

$$e_0 : W_0 \times W_0 \rightarrow \mathbb{Q}, \quad e : W \times W \rightarrow \mathbb{Q}$$

of (4.1.2), the relation (5.1.2) implies that $e|_{W_0} = p \cdot e_0$. The inclusion

$$e_0(p\tilde{\mathfrak{a}}_0, \mathfrak{a}_0) = e(\tilde{\mathfrak{a}}_0, \mathfrak{a}_0) \subset e(\mathfrak{a}, \mathfrak{a}) = \mathbb{Z},$$

together with the self-duality of \mathfrak{a}_0 under e_0 , shows that $p\tilde{\mathfrak{a}}_0 \subset \mathfrak{a}_0$. If equality held we would have

$$e_0(\mathfrak{a}_0, \mathfrak{a}_0) = p^{-1}e(\mathfrak{a}_0, \mathfrak{a}_0) \subset pe(\tilde{\mathfrak{a}}_0, \tilde{\mathfrak{a}}_0) \subset pe(\mathfrak{a}, \mathfrak{a}) \subset p\mathbb{Z},$$

contradicting \mathfrak{a}_0 being self-dual under e_0 .

As $\tilde{\mathfrak{a}}_0$ is $\mathcal{O}_{\mathbf{k}}$ -stable with $\mathfrak{a}_0 \subset \tilde{\mathfrak{a}}_0 \subsetneq \mathfrak{p}^{-1}\mathfrak{a}_0$, it is \mathfrak{a}_0 , $\mathfrak{p}^{-1}\mathfrak{a}_0$, or $\bar{\mathfrak{p}}^{-1}\mathfrak{a}_0$. \square

Proposition 6.2.2. *Let (A_0, A, x) be the universal triple over $\mathcal{Z}_V(p)/\mathcal{O}_k[1/p]$, so that $x \in \text{Hom}_{\mathcal{O}_{\mathbf{k}}}(A_0, A)$ satisfies $\langle x, x \rangle = p$. There is a decomposition (6.1.2) into open and closed substacks in which*

- \mathcal{U}_0 is the locus of points where $\ker(x) = 0$,

- $\mathcal{U}_{\mathfrak{p}}$ is the locus of points where $\ker(x) = A_0[\mathfrak{p}]$,
- $\mathcal{U}_{\bar{\mathfrak{p}}}$ is the locus of points where $\ker(x) = A_0[\bar{\mathfrak{p}}]$.

Proof. Recalling (5.1.3), the relation $\langle x, x \rangle = p$ implies that multiplication-by- p factors as

$$A_0 \xrightarrow{x} A \cong A^\vee \xrightarrow{x^\vee} A_0^\vee \cong A_0,$$

and so $\ker(x) \subset A_0[p]$. Both of these group schemes are finite étale over $\mathcal{Z}_V(p)/\mathcal{O}_{\mathbf{k}[1/p]}$, which implies that each of \mathcal{U}_0 , $\mathcal{U}_{\mathfrak{p}}$, and $\mathcal{U}_{\bar{\mathfrak{p}}}$ is open and closed. It only remains to prove that every geometric point $s \rightarrow \mathcal{Z}_V(p)/\mathcal{O}_{\mathbf{k}[1/p]}$ is contained in one of them.

Abbreviate $T_0 = T_p(A_{0s})$ and $T = T_p(A_s)$. Applying the snake lemma to the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & T_0 & \longrightarrow & T_0 \otimes \mathbb{Q}_p & \longrightarrow & A_{0s}[p^\infty] \longrightarrow 0 \\ & & \downarrow x_s & & \downarrow x_s & & \downarrow x_s \\ 0 & \longrightarrow & T & \longrightarrow & T \otimes \mathbb{Q}_p & \longrightarrow & A_s[p^\infty] \longrightarrow 0, \end{array}$$

and using the vertical arrow on the left to identify $T_0 \subset T$, we find that

$$\ker(x_s) \cong \tilde{T}_0/T_0,$$

where $\tilde{T}_0 = T \cap T_0\mathbb{Q}_p$.

After fixing an isomorphism $\mathbb{Z}_p \cong \mathbb{Z}_p(1)$ of étale sheaves on s , there are unique $\mathcal{O}_{\mathbf{k},p}$ -valued hermitian forms h_0 and h on T_0 and T , respectively, related to the Weil pairings e_0 and e by (4.1.2). Thus we may apply Lemma 6.2.1 with \mathfrak{a}_0 and \mathfrak{a} replaced by T_0 and T , to see that \tilde{T}_0 must be one of T_0 , $\mathfrak{p}^{-1}T_0$, or $\bar{\mathfrak{p}}^{-1}T_0$. These three cases correspond to $\ker(x_s)$ being trivial, $A_{0s}[\mathfrak{p}]$, or $A_{0s}[\bar{\mathfrak{p}}]$. \square

6.3. Analysis of $\mathcal{U}_{\mathfrak{p}}$. In this subsection we study the structure of the substack $\mathcal{U}_{\mathfrak{p}}$ of (6.1.2), and make explicit its relation to $\mathcal{S}_{V^{\flat}}$. The analogous analysis of $\mathcal{U}_{\bar{\mathfrak{p}}}$ is obtained by replacing \mathfrak{p} by $\bar{\mathfrak{p}}$ everywhere.

Return to the situation of Lemma 6.2.1, so that $x \in \mathrm{Hom}_{\mathcal{O}_{\mathbf{k}}}(\mathfrak{a}_0, \mathfrak{a})$ with $\langle x, x \rangle = p$ determines an orthogonal decomposition

$$W = \tilde{W}_0 \oplus W^{\flat}.$$

Set $\tilde{\mathfrak{a}}_0 = \mathfrak{a} \cap \tilde{W}_0$.

Lemma 6.3.1. *If $\tilde{\mathfrak{a}}_0 = \mathfrak{p}^{-1}x(\mathfrak{a}_0)$, there is an orthogonal decomposition*

$$\mathfrak{a} = \tilde{\mathfrak{a}}_0 \oplus \mathfrak{a}^{\flat} \subset W$$

in which $\mathfrak{a}^{\flat} = \mathfrak{a} \cap W^{\flat}$.

Proof. If we use x to identify $W_0 = \tilde{W}_0$, the assumption $\tilde{\mathfrak{a}}_0 = \mathfrak{p}^{-1}\mathfrak{a}_0$ implies that $\tilde{\mathfrak{a}}_0$ is self-dual with respect to the hermitian form $h|_{\tilde{W}_0} = ph_0$. The desired decomposition then follows by elementary linear algebra. \square

The lemma suggests that if $(A_0, A, x) \in \mathcal{U}_{\mathfrak{p}}(S)$ for an $\mathcal{O}_{\mathfrak{k}}[1/p]$ -scheme S , then $x : A_0 \rightarrow A$ should determine an $\mathcal{O}_{\mathfrak{k}}$ -linear splitting

$$(6.3.1) \quad A = \tilde{A}_0 \times A^{\flat}$$

of principally polarized abelian schemes. Indeed, this is the case. If we set

$$\tilde{A}_0 = A_0/A_0[\mathfrak{p}]$$

and recall that $\ker(x) = A_0[\mathfrak{p}]$, the morphism $x : A_0 \rightarrow A$ factors as

$$A_0 \rightarrow \tilde{A}_0 \xrightarrow{y} A$$

for some $y \in \mathrm{Hom}_{\mathcal{O}_{\mathfrak{k}}}(\tilde{A}_0, A)$ satisfying $\langle y, y \rangle = 1$. In other words, the composition

$$\tilde{A}_0 \xrightarrow{y} A \cong A^{\vee} \xrightarrow{y^{\vee}} \tilde{A}_0^{\vee} \cong \tilde{A}_0$$

is the identity. This implies that the composition

$$A \cong A^{\vee} \xrightarrow{y^{\vee}} \tilde{A}_0^{\vee} \cong \tilde{A}_0 \xrightarrow{y} A$$

is a Rosati-fixed idempotent in $\mathrm{End}_{\mathcal{O}_{\mathfrak{k}}}(A)$, and A admits a unique splitting (6.3.1) of principally polarized abelian schemes over S such that this idempotent is the projection to the first factor.

Apply the above construction to the universal triple (A_0, A, x) over $\mathcal{U}_{\mathfrak{p}}$, and recall that A comes equipped with an $\mathcal{O}_{\mathfrak{k}}$ -stable hyperplane $\mathcal{F} \subset \mathrm{Lie}(A)$ satisfying Kramer's signature condition. Using the decomposition

$$\mathrm{Lie}(A) = \mathrm{Lie}(\tilde{A}_0) \oplus \mathrm{Lie}(A^{\flat})$$

of vector bundles on S , denote by $\mathcal{U}_{\mathfrak{p}}^{\dagger} \subset \mathcal{U}_{\mathfrak{p}}$ the largest closed substack over which $\mathrm{Lie}(\tilde{A}_0) \subset \mathcal{F}$.

Proposition 6.3.2. *There is a canonical isomorphism*

$$i_{\mathfrak{p}} : \mathcal{S}_{V^{\flat}/\mathcal{O}_{\mathfrak{k}}[1/p]} \cong \mathcal{U}_{\mathfrak{p}}^{\dagger}.$$

Proof. As above, let S be an $\mathcal{O}_{\mathfrak{k}}[1/p]$ -scheme. Given a point $(A_0, A, x) \in \mathcal{U}_{\mathfrak{p}}^{\dagger}(S)$, the splitting (6.3.1) determines

$$A^{\flat} \in \mathcal{M}_{(n-2,1)}(S),$$

where we have endowed A^{\flat} with the $\mathcal{O}_{\mathfrak{k}}$ -stable hyperplane

$$\mathcal{F}^{\flat} = \mathcal{F}/\mathrm{Lie}(\tilde{A}_0) \subset \mathrm{Lie}(A)/\mathrm{Lie}(\tilde{A}_0) \cong \mathrm{Lie}(A^{\flat}).$$

satisfying Kramer's condition. The pair (A_0, A^{\flat}) defines an S -point of

$$\mathcal{S}_{V^{\flat}} \subset \mathcal{M}_{(1,0)} \times_{\mathcal{O}_{\mathfrak{k}}} \mathcal{M}_{(n-2,1)},$$

and we have now constructed a morphism

$$(6.3.2) \quad \mathcal{U}_{\mathfrak{p}}^{\dagger} \xrightarrow{(A_0, A, x) \rightarrow (A_0, A^{\flat})} \mathcal{S}_{V^{\flat}/\mathcal{O}_{\mathfrak{k}}[1/p]}.$$

Conversely, start with an S -point

$$(A_0^{\flat}, A^{\flat}) \in \mathcal{S}_{V^{\flat}}(S).$$

First define elliptic curves $A_0 = A_0^b$ and $\tilde{A}_0 = A_0/A_0[\mathfrak{p}]$. Then define an abelian scheme A by (6.3.1), and endow A with its product principal polarization and product \mathcal{O}_k -action. Recalling that A^b comes equipped with a hyperplane $\mathcal{F}^b \subset \text{Lie}(A^b)$ satisfying Kramer's condition, we endow A with the hyperplane

$$\mathcal{F} = \text{Lie}(\tilde{A}_0) \oplus \mathcal{F}^b \subset \text{Lie}(A).$$

It is easy to check that A , with its extra data, defines an S -point of $\mathcal{M}_{(n-1,1)}$, and that (A_0, A) defines an S -point of the open and closed substack

$$\mathcal{S}_V \subset \mathcal{M}_{(1,0)} \times_{\mathcal{O}_k} \mathcal{M}_{(n-2,1)}.$$

If we define $x \in \text{Hom}_{\mathcal{O}_k}(A_0, A)$ as the composition

$$A_0 \rightarrow \tilde{A}_0 \hookrightarrow \tilde{A}_0 \times A^b = A,$$

where the first arrow is the quotient map, then $\langle x, x \rangle = p$ and $\ker(x) = A_0[\mathfrak{p}]$. The triple (A_0, A, x) defines an S -point of \mathcal{U}_p^\dagger , and the morphism

$$\mathcal{S}_{V^b}/\mathcal{O}_k[1/p] \xrightarrow{(A_0^b, A^b) \rightarrow (A_0, A, x)} \mathcal{U}_p^\dagger$$

is inverse to (6.3.2). \square

Proposition 6.3.2 gives us a commutative diagram (6.3.3)

$$\begin{array}{ccc} \mathcal{S}_{V^b}/\mathcal{O}_k[1/p] & \xrightarrow{\quad\quad\quad} & \mathcal{S}_V/\mathcal{O}_k[1/p] \\ \downarrow & & \downarrow \\ (\mathcal{M}_{(1,0)} \times_{\mathcal{O}_k} \mathcal{M}_{(n-2,1)}^{\text{Pap}})/\mathcal{O}_k[1/p] & \longrightarrow & (\mathcal{M}_{(1,0)} \times_{\mathcal{O}_k} \mathcal{M}_{(n-1,1)}^{\text{Pap}})/\mathcal{O}_k[1/p] \end{array}$$

in which the top horizontal arrow is the composition

$$\mathcal{S}_{V^b}/\mathcal{O}_k[1/p] \xrightarrow{i_p} \mathcal{U}_p^\dagger \hookrightarrow \mathcal{Z}_V(p)/\mathcal{O}_k[1/p] \rightarrow \mathcal{S}_V/\mathcal{O}_k[1/p],$$

and the bottom horizontal arrow sends $(A_0^b, A^b) \mapsto (A_0, A)$, where

$$(6.3.4) \quad A_0 = A_0^b, \quad \tilde{A}_0 = A_0/A_0[\mathfrak{p}], \quad A = \tilde{A}_0 \times A^b.$$

Both horizontal arrows are finite and unramified.

Proposition 6.3.3. *The homomorphism*

$$\widehat{\text{Pic}}(\mathcal{S}_V/\mathcal{O}_k[1/p])_{\mathbb{Q}} \rightarrow \widehat{\text{Pic}}(\mathcal{S}_{V^b}/\mathcal{O}_k[1/p])_{\mathbb{Q}}$$

induced by (6.3.3) sends

$$\widehat{\omega}_{A/\mathcal{S}_V}^{\text{Hdg}} \mapsto \widehat{\omega}_{A_0^b/\mathcal{S}_{V^b}}^{\text{Hdg}} + \widehat{\omega}_{A^b/\mathcal{S}_{V^b}}^{\text{Hdg}},$$

where (A_0^b, A^b) is the universal pair over \mathcal{S}_{V^b} . The same map also sends

$$\widehat{\mathcal{L}}_V \mapsto \widehat{\mathcal{L}}_{V^b} \quad \text{and} \quad (\text{Exc}_V, 0) \mapsto (\text{Exc}_{V^b}, 0).$$

Proof. It follows from (6.3.4) that the top horizontal arrow in (6.3.3) sends

$$\widehat{\omega}_{A/\mathcal{S}_V}^{\text{Hdg}} \mapsto \widehat{\omega}_{\tilde{A}_0/\mathcal{S}_{V^b}}^{\text{Hdg}} + \widehat{\omega}_{A^b/\mathcal{S}_{V^b}}^{\text{Hdg}}.$$

As we have inverted p on the base, the degree p quotient map $A_0^b \rightarrow \tilde{A}_0$ induces an isomorphism on Lie algebras, and hence an isomorphism

$$\omega_{\tilde{A}_0/\mathcal{S}_{V^b}}^{\text{Hdg}} \cong \omega_{A_0^b/\mathcal{S}_{V^b}}^{\text{Hdg}}.$$

This isomorphism does not respect the metrics (1.1.1), but one can easily check that

$$\widehat{\omega}_{\tilde{A}_0/\mathcal{S}_{V^b}}^{\text{Hdg}} = \widehat{\omega}_{A_0^b/\mathcal{S}_{V^b}}^{\text{Hdg}} + (0, -\log(p))$$

as elements of $\widehat{\text{Pic}}(\mathcal{S}_{V^b}/\mathcal{O}_{\mathbf{k}}[1/p])$. The correction term $(0, -\log(p))$ is torsion in the arithmetic Picard group, as

$$2(0, -\log(p)) = (\text{div}(p), -\log(p^2)) = 0,$$

and the first claim follows.

For the second claim, let S be an $\mathcal{O}_{\mathbf{k}}[1/p]$ -scheme, fix an S -point $(A_0^b, A^b) \in \mathcal{S}_{V^b}(S)$, and let $(A_0, A) \in \mathcal{S}_V(S)$ be its image under the top horizontal arrow in (6.3.3). Tracing through the construction of $i_{\mathfrak{p}}$ in Proposition 6.3.2, we see that

$$\text{Lie}(A)/\mathcal{F} \cong \text{Lie}(A^b)/\mathcal{F}^b.$$

It follows immediately from this and (5.2.4) that $\mathcal{L}_V|_S \cong \mathcal{L}_{V^b}|_S$. At a complex point $s \in S(\mathbb{C})$ we have a decomposition

$$H_1(A_s, \mathbb{Q}) = H_1(\tilde{A}_{0s}, \mathbb{Q}) \oplus H_1(A_s^b, \mathbb{Q}),$$

which induces an isometric embedding

$$\text{Hom}_{\mathbf{k}}(H_1(A_{0s}, \mathbb{Q}), H_1(A_s^b, \mathbb{Q})) \subset \text{Hom}_{\mathbf{k}}(H_1(A_{0s}, \mathbb{Q}), H_1(A_s, \mathbb{Q}))$$

of \mathbf{k} -hermitian spaces. Recalling (5.2.5), this isometric embedding restricts to the isomorphism $\mathcal{L}_{V^b,s} \cong \mathcal{L}_{V,s}$ just constructed, which therefore preserves the metrics defined by (5.2.6).

Finally, we show that under the top horizontal arrow in (6.3.3), the exceptional divisor $\text{Exc}_V \subset \mathcal{S}_V$ pulls back to the exceptional divisor $\text{Exc}_{V^b} \subset \mathcal{S}_{V^b}$. These divisors are, by Definition 5.1.4, the pullbacks of the singular loci

$$(6.3.5) \quad \mathcal{M}_{(1,0)} \times_{\mathcal{O}_{\mathbf{k}}} \text{Sing}_{(n-1,1)} \subset \mathcal{M}_{(1,0)} \times_{\mathcal{O}_{\mathbf{k}}} \mathcal{M}_{(n-1,1)}^{\text{Pap}}$$

$$(6.3.6) \quad \mathcal{M}_{(1,0)} \times_{\mathcal{O}_{\mathbf{k}}} \text{Sing}_{(n-2,1)} \subset \mathcal{M}_{(1,0)} \times_{\mathcal{O}_{\mathbf{k}}} \mathcal{M}_{(n-2,1)}^{\text{Pap}}$$

of (4.2.2) under the vertical arrows in (6.3.3), and so it suffices to prove that the first singular locus (6.3.5) pulls back to the second singular locus (6.3.6) under the bottom horizontal arrow in (6.3.3). Moreover, as each of (6.3.5) and (6.3.6) is a reduced $\mathcal{O}_{\mathbf{k}}$ -stack of dimension 0, and as the bottom horizontal arrow in (6.3.3) is finite and unramified, it suffices to check this on the level of geometric points. This is an easy exercise in linear algebra, using the characterization of the singular locus found in Remark 4.2.1. \square

Proposition 6.3.4. *The nonexceptional locus $\mathcal{U}_{\mathfrak{p}}^{\text{nexc}} \subset \mathcal{U}_{\mathfrak{p}}$ satisfies*

$$\mathcal{U}_{\mathfrak{p}}^{\text{nexc}} \subset \mathcal{U}_{\mathfrak{p}}^{\dagger},$$

and the isomorphism of Proposition 6.3.2 restricts to an isomorphism

$$\mathcal{S}_{V^{\flat}/\mathcal{O}_{\mathbf{k}}[1/p]}^{\text{nexc}} \cong \mathcal{U}_{\mathfrak{p}}^{\text{nexc}}.$$

Proof. Suppose S is an $\mathcal{O}_{\mathbf{k}}[1/p]$ -scheme,

$$(A_0, A, x) \in \mathcal{U}_{\mathfrak{p}}^{\text{nexc}}(S),$$

and let A^{\flat} be the abelian scheme of (6.3.1). In particular,

$$\text{Lie}(A) = \text{Lie}(\tilde{A}_0) \oplus \text{Lie}(A^{\flat}).$$

The natural map

$$\mathcal{U}_{\mathfrak{p}}^{\text{nexc}} \rightarrow \mathcal{M}_{(n-1,1)} \setminus \text{Exc}_{(n-1,1)}$$

endows A with a hyperplane $\mathcal{F} \subset \text{Lie}(A)$, and results of Krämer, as summarized in Theorem 2.3.3 of [BHK⁺a], shows that this hyperplane is actually determined by the $\mathcal{O}_{\mathbf{k}}$ -action on A using the following recipe: If we fix a $\pi \in \mathcal{O}_{\mathbf{k}}$ such that $\mathcal{O}_{\mathbf{k}} = \mathbb{Z}[\pi]$, and let

$$\bar{\epsilon}_S = \bar{\pi} \otimes 1 - 1 \otimes i_S(\bar{\pi}) \in \mathcal{O}_{\mathbf{k}} \otimes_{\mathbb{Z}} \mathcal{O}_S,$$

where $i_S : \mathcal{O}_{\mathbf{k}} \rightarrow \mathcal{O}_S$ is the structure map, then

$$\mathcal{F} = \ker(\bar{\epsilon}_S : \text{Lie}(A) \rightarrow \text{Lie}(A)).$$

On the other hand, the action of $\mathcal{O}_{\mathbf{k}}$ on the Lie algebra of A_0 is through the structure morphism i_S , and so the same is true of $\tilde{A}_0 = A_0/A_0[\mathfrak{p}]$. This implies that $\bar{\epsilon}_S$ annihilates $\text{Lie}(\tilde{A}_0)$, and hence $\text{Lie}(\tilde{A}_0) \subset \mathcal{F}$. Having shown

$$(A_0, A, x) \in \mathcal{U}_{\mathfrak{p}}^{\dagger}(S),$$

the first claim of the proposition is proved. The second claim follows from the first and Proposition 6.3.3. \square

Proposition 6.3.5. *The closed immersion $i_{\mathfrak{p}} : \mathcal{S}_{V^{\flat}/\mathcal{O}_{\mathbf{k}}[1/p]} \rightarrow \mathcal{U}_{\mathfrak{p}}$ of Proposition 6.3.2 extends uniquely to a proper morphism*

$$\bar{\mathcal{S}}_{V^{\flat}/\mathcal{O}_{\mathbf{k}}[1/p]} \rightarrow \bar{\mathcal{U}}_{\mathfrak{p}},$$

where the codomain is defined as the Zariski closure of

$$\mathcal{U}_{\mathfrak{p}} \subset \bar{\mathcal{Z}}_V(p)/\mathcal{O}_{\mathbf{k}}[1/p].$$

or, equivalently, as the normalization of $\mathcal{U}_{\mathfrak{p}} \rightarrow \bar{\mathcal{S}}_{V/\mathcal{O}_{\mathbf{k}}[1/p]}$.

Proof. For ease of notation, we omit all subscripts $\mathcal{O}_{\mathbf{k}}[1/p]$ in the proof.

As the exceptional locus of $\bar{\mathcal{S}}_{V^{\flat}}$ does not meet the boundary, it suffices to construct the extension over its complement

$$\bar{\mathcal{S}}_{V^{\flat}}^{\text{nexc}} = \bar{\mathcal{S}}_{V^{\flat}} \setminus \text{Exc}_{V^{\flat}}.$$

Consider the universal pair (A_0, A^b) over $\mathcal{S}_{V^b}^{\text{nexc}}$, and let

$$x : A_0 \rightarrow A = \tilde{A}_0 \times A^b$$

be as in the proof of Proposition 6.3.2. In other words, the triple (A_0, A, x) over $\mathcal{S}_{V^b}^{\text{nexc}}$ determines the isomorphism $i_{\mathfrak{p}} : \mathcal{S}_{V^b}^{\text{nexc}} \cong \mathcal{U}_{\mathfrak{p}}^{\text{nexc}}$ of Proposition 6.3.4.

The discussion of §4.3 shows that A^b extends to a semi-abelian scheme over $\bar{\mathcal{S}}_{V^b}^{\text{nexc}}$. The elliptic curve A_0 extends to an elliptic curve $\bar{\mathcal{S}}_{V^b}^{\text{nexc}}$, and so the same is true of its quotient \tilde{A}_0 . It follows that A extends to a semi-abelian scheme over $\bar{\mathcal{S}}_{V^b}^{\text{nexc}}$, and the extension property used in the proof of Lemma 4.3.2 provides us with a morphism

$$i_{\mathfrak{p}} : \bar{\mathcal{S}}_{V^b}^{\text{nexc}} \rightarrow \mathcal{M}_{(1,0)} \times_{\mathcal{O}_k} \bar{\mathcal{M}}_{(n-1,1)}^{\text{Pap}}$$

taking values in the open subscheme

$$\mathcal{M}_{(1,0)} \times_{\mathcal{O}_k} (\bar{\mathcal{M}}_{(n-1,1)}^{\text{Pap}} \setminus \text{Sing}_{(n-1,1)}) \cong \mathcal{M}_{(1,0)} \times_{\mathcal{O}_k} (\bar{\mathcal{M}}_{(n-1,1)} \setminus \text{Exc}_{(n-1,1)}).$$

This provides us with a morphism

$$i_{\mathfrak{p}} : \bar{\mathcal{S}}_{V^b}^{\text{nexc}} \rightarrow \mathcal{M}_{(1,0)} \times_{\mathcal{O}_k} \bar{\mathcal{M}}_{(n-1,1)},$$

taking values in the open and closed substack $\bar{\mathcal{S}}_V$.

We now have a commutative diagram of solid arrows

$$\begin{array}{ccc} \mathcal{S}_{V^b}^{\text{nexc}} & \xrightarrow{i_{\mathfrak{p}}} & \bar{\mathcal{U}}_{\mathfrak{p}} \\ \downarrow & \nearrow \text{dotted} & \downarrow \\ \bar{\mathcal{S}}_{V^b}^{\text{nexc}} & \longrightarrow & \bar{\mathcal{S}}_V \end{array}$$

in which the vertical arrow on the right is integral, and hence affine, by its construction as a normalization; see [Sta18, Tag 035I]. To complete the proof of the proposition, it suffices to show that there is a unique dotted arrow making the diagram commute. This immediately reduces to the corresponding claim for affine schemes, in which we are given homomorphisms of rings

$$\begin{array}{ccc} R^b & \longleftarrow & B \\ \uparrow & \nearrow \text{dotted} & \uparrow \\ \bar{R}^b & \longleftarrow & A \end{array}$$

with B integral over A , and $\bar{R}^b \subset R^b$ is an inclusion of normal domains with the same field of fractions. The image of any $b \in B$ under $B \rightarrow R^b$ is integral over \bar{R}^b , hence contained in \bar{R}^b . Thus there is a unique dotted arrow making the diagram commute. \square

6.4. Two lemmas on abelian schemes. For use in the next subsection, we prove two lemmas on abelian schemes. The first is a criterion for determining when a polarization descends to a quotient. The second is a criterion for the existence of an extension as a semi-abelian scheme.

Lemma 6.4.1. *Suppose $f : X \rightarrow Y$ is an isogeny of abelian schemes over a Noetherian scheme S , and $\psi_X : X \rightarrow X^\vee$ is a polarization whose degree d is invertible in \mathcal{O}_S . The following are equivalent.*

- (1) *The kernel of f is contained in $\ker(\psi_X)$, and is totally isotropic with respect to the Weil pairing*

$$e : \ker(\psi_X) \times \ker(\psi_X) \rightarrow \mu_d.$$

- (2) *There exists a (necessarily unique) polarization $\psi_Y : Y \rightarrow Y^\vee$ making the diagram*

$$\begin{array}{ccc} X & \xrightarrow{\psi_X} & X^\vee \\ f \downarrow & & \uparrow f^\vee \\ Y & \xrightarrow{\psi_Y} & Y^\vee \end{array}$$

commute.

Proof. This is routine. See, for example, Proposition 10.4 of [Pol03]. \square

Lemma 6.4.2. *Suppose S is a normal Noetherian scheme, $U \subset S$ is a dense open subscheme, and $f : X \rightarrow Y$ is an isogeny of abelian schemes over U whose degree d is invertible in \mathcal{O}_S . If X extends to a semi-abelian scheme over S then so does Y , and both extensions are unique.*

Proof. The uniqueness claim follows from Proposition I.2.7 of [FC90], so we only need to prove the existence of a semi-abelian extension of Y .

Consider the morphisms $\ker(f) \rightarrow X[d] \rightarrow U$. The second arrow and the composition are both étale, as $d \in \mathcal{O}_U^\times$, and so the first arrow is also étale by [Sta18, Tag 02GW]. As the first arrow is also a closed immersion, we deduce that $\ker(f) \subset X[d]$ is a union of connected components.

If $X^* \rightarrow S$ denotes the semi-abelian extension of X , the multiplication map $[d] : X^* \rightarrow X^*$ is quasi-finite and flat. Indeed, by [Sta18, Tag 039E] flatness can be checked fiberwise on S , where it follows by combining Proposition 3.11 of Exposé VI.B of [SGA70] with the surjectivity of $[d]$ on any extension of an abelian variety by a torus. The group scheme $X^*[d] \rightarrow S$ is therefore quasi-finite and flat. Using our assumption that $d \in \mathcal{O}_S^\times$, and [Sta18, Tag 02GM], we deduce that it is étale.

As we assume that S is normal, so is $X^*[d]$. As the connected components of a normal scheme are the same as its irreducible components, it follows that the Zariski closure of $\ker(f)$ in $X^*[d]$ is an open and closed subscheme $\ker(f)^* \subset X^*[d]$. In particular $\ker(f)^* \rightarrow S$ is quasi-finite étale.

By Lemma IV.7.1.2 of [MB85], the fppf quotient $Y^* = X^*/\ker(f)^*$ provides a semi-abelian extension of Y to S . \square

6.5. Analysis of \mathcal{U}_0 . In this subsection we study the substack \mathcal{U}_0 of (6.1.2), and make explicit its relation with \mathcal{S}_{V^b} .

Return to the situation of Lemma 6.2.1, so that $x \in \text{Hom}_{\mathcal{O}_k}(\mathfrak{a}_0, \mathfrak{a})$ with $\langle x, x \rangle = p$ determines an orthogonal decomposition

$$(6.5.1) \quad W = \tilde{W}_0 \oplus W^b.$$

Set $\tilde{\mathfrak{a}}_0 = \mathfrak{a} \cap \tilde{W}_0$.

Lemma 6.5.1. *Assume that $\tilde{\mathfrak{a}}_0 = x(\mathfrak{a}_0)$. If we set $\mathfrak{b} = \mathfrak{a} \cap W^b$ and let $\mathfrak{b}^\vee \subset W^b$ be its dual lattice with respect to $h|_{W^b}$, the \mathcal{O}_k -lattice*

$$\mathfrak{a}^b = \mathfrak{p} \cdot \mathfrak{b}^\vee + \mathfrak{b},$$

is self-dual with respect to $h|_{W^b}$. Moreover, there are inclusions of \mathcal{O}_k -lattices

$$(6.5.2) \quad \tilde{\mathfrak{a}}_0 \oplus \mathfrak{a}^b \supseteq^p \tilde{\mathfrak{a}}_0 \oplus \mathfrak{b} \subset^{p^2} \mathfrak{a}$$

in W of the indicated indices, and a canonical \mathcal{O}_k -linear injection

$$(6.5.3) \quad \mathfrak{p}^{-1}\tilde{\mathfrak{a}}_0/\tilde{\mathfrak{a}}_0 \cong \mathfrak{p}^{-1}\mathfrak{b}/\mathfrak{b} \xrightarrow{t} \mathfrak{p}^{-1}\mathfrak{a}^b/\mathfrak{a}^b.$$

Proof. Throughout the proof, we use x to identify $W_0 = \tilde{W}_0$, so that $h|_{W_0} = ph_0$. The projections to the two factors in (6.5.1) induce isomorphisms

$$\mathfrak{a}/(\mathfrak{a}_0 \oplus \mathfrak{b}) \rightarrow \mathfrak{a}_0^\vee/\mathfrak{a}_0, \quad \mathfrak{a}/(\mathfrak{a}_0 \oplus \mathfrak{b}) \rightarrow \mathfrak{b}^\vee/\mathfrak{b},$$

where $\mathfrak{a}_0^\vee = p^{-1}\mathfrak{a}_0$ is the dual lattice of \mathfrak{a}_0 with respect to $h|_{W_0}$. Composing these isomorphisms yields an \mathcal{O}_k -linear isomorphism

$$(6.5.4) \quad p^{-1}\mathfrak{a}_0/\mathfrak{a}_0 \cong \mathfrak{b}^\vee/\mathfrak{b}$$

respecting the $p^{-1}\mathcal{O}_k/\mathcal{O}_k$ -valued hermitian forms induced by $h|_{W_0} = ph_0$ and $h|_{W^b}$. It follows that $\mathfrak{p} \cdot (\mathfrak{b}^\vee/\mathfrak{b}) \subset \mathfrak{b}^\vee/\mathfrak{b}$ is maximal isotropic with respect to $h|_{W^b}$, and so \mathfrak{a}^b is self-dual under $h|_{W^b}$.

Consider the inclusions $\mathfrak{b} \subset \mathfrak{a}^b \subset \mathfrak{b}^\vee$ of \mathcal{O}_k -modules. The first is an isomorphism everywhere locally except at $\bar{\mathfrak{p}}$, while the second is an isomorphism everywhere locally except at \mathfrak{p} . In particular, the first induces an isomorphism

$$\mathfrak{p}^{-1}\mathfrak{b}/\mathfrak{b} \cong \mathfrak{p}^{-1}\mathfrak{a}^b/\mathfrak{a}^b.$$

Restricting (6.5.4) to an isomorphism of \mathfrak{p} -torsion submodules defines (6.5.3). The inclusions of (6.5.2), with the indicated indices, are clear from what we have said. \square

Just as Lemma 6.5.1 is more complicated than Lemma 6.3.1, the analysis of \mathcal{U}_0 is more complicated than that of \mathcal{U}_p .

Let S be an $\mathcal{O}_k[1/p]$ -scheme. Given Lemma 6.5.1, one expects that any S -point $(A_0, A, x) \in \mathcal{U}_0(S)$ should determine a diagram of abelian schemes with \mathcal{O}_k -actions

$$(6.5.5) \quad \tilde{A}_0 \times A^b \xleftarrow{\text{deg}=p} \tilde{A}_0 \times B \xrightarrow{\text{deg}=p^2} A$$

in which the arrows are \mathcal{O}_k -linear isogenies of the indicated degrees. There should also be a canonical \mathcal{O}_k -linear closed immersion

$$(6.5.6) \quad \tilde{A}_0[\mathfrak{p}] \cong B[\mathfrak{p}] \xrightarrow{t} A^b[\mathfrak{p}],$$

and the morphism $x : A_0 \rightarrow A$ should factor as

$$A_0 \cong \tilde{A}_0 \hookrightarrow \tilde{A}_0 \times B \rightarrow A.$$

Moreover, these abelian schemes should come with polarizations with the following properties. The elliptic curve \tilde{A}_0 is equipped with its unique polarization of degree p^2 , B is equipped with a polarization of degree p^2 , and A^b is equipped with a principal polarization. The pullback of the product polarization on $\tilde{A}_0 \times A^b$ via the leftwards arrow in (6.5.5) is the product polarization on $\tilde{A}_0 \times B$, and similarly for the the pullback of the principal polarization on A via the rightward arrow.

Here is how to construct the data above from (A_0, A, x) . As in the proof of Proposition 6.2.2, at any geometric point $s \rightarrow S$ we may apply Lemma 6.5.1 with \mathfrak{a}_0 and \mathfrak{a} replaced by the p -adic Tate modules of A_{0s} and A_s to obtain \mathcal{O}_k -lattices

$$(6.5.7) \quad T_p(\tilde{A}_{0s}) \oplus T_p(A_s^b) \supset T_p(\tilde{A}_{0s}) \oplus T_p(B_s) \subset T_p(A_s).$$

analogous to (6.5.2), along with an \mathcal{O}_k -linear injection

$$(6.5.8) \quad \mathfrak{p}^{-1}T_p(\tilde{A}_{0s})/T_p(\tilde{A}_{0s}) \cong \mathfrak{p}^{-1}T_p(B_s)/T_p(B_s) \xrightarrow{t} \mathfrak{p}^{-1}T_p(A_s^b)/T_p(A_s^b).$$

To be clear, we have not yet constructed the abelian schemes \tilde{A}_0 , A^b , and B , only lattices in $T_p(A_s)[1/p]$ that we choose to call $T_p(\tilde{A}_s)$, $T_p(A_s^b)$, and $T_p(B_s)$. However, as $p \in \mathcal{O}_S^\times$ and both inclusions in (6.5.7) have finite index, there are abelian schemes C and D over S with \mathcal{O}_k -actions and \mathcal{O}_k -linear isogenies

$$D \xleftarrow{p} C \xrightarrow{p^2} A$$

of the indicated degrees that identify

$$(6.5.9) \quad T_p(C_s) = T_p(\tilde{A}_{0s}) \oplus T_p(B_s), \quad T_p(D_s) = T_p(\tilde{A}_{0s}) \oplus T_p(A_s^b).$$

The condition that $\langle x, x \rangle = p$ implies that

$$A_0 \xrightarrow{x} A \cong A^\vee \xrightarrow{x^\vee} A_0^\vee \cong A_0$$

is multiplication by p , which implies that the composition

$$A \cong A^\vee \xrightarrow{x^\vee} A_0^\vee \cong A_0 \xrightarrow{x} A$$

is a Rosati-fixed element $\alpha \in \text{End}_{\mathcal{O}_k}(A)[1/p]$ such that $\alpha \circ \alpha = p\alpha$. In particular, we obtain an idempotent

$$p^{-1}\alpha \in \text{End}_{\mathcal{O}_k}(A)[1/p] \cong \text{End}_{\mathcal{O}_k}(C)[1/p] \cong \text{End}_{\mathcal{O}_k}(D)[1/p],$$

whose induced actions on $T_p(C_s)[1/p]$ and $T_p(D_s)[1/p]$ are just the projections to the first summands in (6.5.9). It follows that we in fact have

$$p^{-1}\alpha \in \text{End}_{\mathcal{O}_k}(C) \cong \text{End}_{\mathcal{O}_k}(D).$$

These idempotents are the projections to the first factors for unique decompositions

$$C = \tilde{A}_0 \times B, \quad D = \tilde{A}_0 \times A^b$$

of abelian schemes with \mathcal{O}_k -actions, and the image of x under

$$\mathrm{Hom}_{\mathcal{O}_k}(A_0, A)[1/p] \cong \mathrm{Hom}_{\mathcal{O}_k}(A_0, C)[1/p] \rightarrow \mathrm{Hom}_{\mathcal{O}_k}(A_0, \tilde{A}_0)[1/p]$$

is an isomorphism $A_0 \cong \tilde{A}_0$. In particular, we now have \mathcal{O}_k -linear isogenies (6.5.5) such that taking p -adic Tate modules recovers (6.5.7), and (6.5.8) induces the closed immersion (6.5.6).

If we pullback the principal polarization via $\tilde{A}_0 \times B \rightarrow A$, we obtain a polarization on $\tilde{A}_0 \times B$ of degree p^4 . As the idempotent $p^{-1}\alpha$ constructed above is Rosati-fixed, this polarization of $\tilde{A}_0 \times B$ splits as the product of polarizations $\tilde{A}_0 \rightarrow \tilde{A}_0^\vee$ and $B \rightarrow B^\vee$. By examining the induced Weil pairing on p -adic Tate modules, one can show that each polarization has kernel isomorphic, étale locally on S , to $\mathcal{O}_k/(p)$, and that

$$\ker(B \rightarrow A^b) = \ker(B \rightarrow B^\vee)[\mathfrak{p}]$$

is totally isotropic under the Weil pairing on $\ker(B \rightarrow B^\vee)$. Hence, by Lemma 6.4.1, $B \rightarrow B^\vee$ descends to a principal polarization on A^b . This completes the construction of the abelian schemes (6.5.5) with all of their expected extra structure.

Now apply this construction to the universal triple (A_0, A, x) over \mathcal{U}_0 . As $p \in \mathcal{O}_{\mathcal{U}_0}^\times$, the p -power isogenies (6.5.5) induce an isomorphism

$$(6.5.10) \quad \mathrm{Lie}(A) \cong \mathrm{Lie}(\tilde{A}_0) \times \mathrm{Lie}(A^b)$$

of rank n vector bundles on \mathcal{U}_0 . Recalling that A comes equipped with an \mathcal{O}_k -stable hyperplane $\mathcal{F} \subset \mathrm{Lie}(A)$, denote by

$$\mathcal{U}_0^\dagger \subset \mathcal{U}_0$$

the largest closed substack over which $\mathrm{Lie}(\tilde{A}_0) \subset \mathcal{F}$

Define a finite étale cover

$$(6.5.11) \quad \mathcal{T}_{V^b} \rightarrow \mathcal{S}_{V^b/\mathcal{O}_k[1/p]}$$

of degree $p^{n-1} - 1$ as follows: for any $\mathcal{O}_k[1/p]$ -scheme S , let $\mathcal{T}_{V^b}(S)$ be the groupoid of triples (A_0^b, A^b, t) in which

$$(A_0^b, A^b) \in \mathcal{S}_{V^b}(S) \quad \text{and} \quad t : A_0^b[\mathfrak{p}] \rightarrow A^b[\mathfrak{p}]$$

is an \mathcal{O}_k -linear closed immersion of group schemes over S .

Proposition 6.5.2. *There is a canonical isomorphism $i_0 : \mathcal{T}_{V^b} \cong \mathcal{U}_0^\dagger$.*

Proof. The morphism $\mathcal{U}_0^\dagger \rightarrow \mathcal{T}_{V^b}$ is essentially described above. Suppose S is an $\mathcal{O}_k[1/p]$ -scheme and

$$(A_0, A, x) \in \mathcal{U}_0^\dagger(S).$$

We can use the isomorphism (6.5.10) to endow the abelian scheme A^b of (6.5.5) with the rank one local direct summand

$$\mathcal{F}^b = \mathcal{F} / \text{Lie}(\tilde{A}_0) \subset \text{Lie}(A) / \text{Lie}(\tilde{A}_0) = \text{Lie}(A^b)$$

to obtain a point $A^b \in \mathcal{M}_{(n-2,1)}(S)$. Setting $A_0^b = \tilde{A}_0$, this defines a morphism

$$\mathcal{U}_0^\dagger \xrightarrow{(A_0, A, x) \mapsto (A_0^b, A^b)} \mathcal{S}_{V^b / \mathcal{O}_k[1/p]},$$

and the closed immersion (6.5.6) defines the desired lift to $\mathcal{U}_0^\dagger \rightarrow \mathcal{T}_{V^b}$.

Now we construct the inverse. Start with a point $(A_0^b, A^b, t) \in \mathcal{T}_{V^b}(S)$. Denote by B the abelian scheme dual to

$$B^\vee = A^b / \text{Im}(t).$$

Using the principal polarization to identify A^b with its dual, the quotient map $A^b \rightarrow B^\vee$ dualizes to a morphism $B \rightarrow A^b$, and the composition

$$B \rightarrow A^b \rightarrow B^\vee$$

is a degree p^2 polarization (it is the pullback via $B \rightarrow A^b$ of the principal polarization on A^b). It has the property that each factor on the right hand side of

$$\ker(B \rightarrow B^\vee) = \ker(B \rightarrow B^\vee)[\mathfrak{p}] \times \ker(B \rightarrow B^\vee)[\bar{\mathfrak{p}}]$$

has order p , and the second factor is the kernel of $B \rightarrow A^b$. In particular, the induced map $B[\mathfrak{p}] \rightarrow A^b[\mathfrak{p}]$ is an isomorphism, and the closed immersion

$$A_0^b[\mathfrak{p}] \xrightarrow{t} A^b[\mathfrak{p}] \cong B[\mathfrak{p}]$$

has image $\ker(B \rightarrow B^\vee)[\mathfrak{p}]$. Setting $\tilde{A}_0 = A_0^b$, the resulting isomorphism

$$\tilde{A}_0[\mathfrak{p}] \cong \ker(B \rightarrow B^\vee)[\mathfrak{p}]$$

admits a unique \mathcal{O}_k -linear extension to an isomorphism

$$t : \tilde{A}_0[p] \cong \ker(B \rightarrow B^\vee)$$

identifying the Weil pairings on source and target.

The antidiagonal subgroup

$$\Delta \stackrel{\text{def}}{=} \tilde{A}_0[p] \xrightarrow{a_0 \mapsto (-a_0, t(a_0))} \tilde{A}_0 \times B.$$

is totally isotropic under the Weil pairing induced by the product polarization (where \tilde{A}_0 is given the polarization of degree p^2), which therefore descends, by Lemma 6.4.1, to a principal polarization on the quotient

$$A = (\tilde{A}_0 \times B) / \Delta.$$

As $p \in \mathcal{O}_S^\times$, there is an induced isomorphism of Lie algebras (6.5.10), which allows us to define a corank one local direct summand $\mathcal{F} \subset \text{Lie}(A)$ as the product

$$\mathcal{F} = \text{Lie}(\tilde{A}_0) \times \mathcal{F}^b.$$

This defines a point $A \in \mathcal{M}_{(n-1,1)}(S)$. Setting $A_0 = \tilde{A}_0$, the composition

$$A_0 = \tilde{A}_0 \hookrightarrow \tilde{A}_0 \times B \rightarrow A$$

defines $x \in \mathrm{Hom}_{\mathcal{O}_k}(A_0, A)$ with $\langle x, x \rangle = p$

The above construction determines a morphism

$$\mathcal{T}_{V^b} \xrightarrow{(A_0^b, A^b, t) \rightarrow (A_0, A, x)} \mathcal{Z}_V(p)$$

taking values in the open substack \mathcal{U}_0^\dagger , which is inverse to the map $\mathcal{U}_0^\dagger \rightarrow \mathcal{T}_{V^b}$ constructed above. \square

Proposition 6.5.2 gives us morphisms

$$(6.5.12) \quad \mathcal{S}_{V^b}/\mathcal{O}_k[1/p] \xleftarrow{(6.5.11)} \mathcal{T}_{V^b} \longrightarrow \mathcal{S}_V/\mathcal{O}_k[1/p]$$

in which the arrow on the right is the (finite and unramified) composition

$$\mathcal{T}_{V^b} \xrightarrow{i_0} \mathcal{U}_0^\dagger \hookrightarrow \mathcal{Z}_V(p)/\mathcal{O}_k[1/p] \rightarrow \mathcal{S}_V/\mathcal{O}_k[1/p],$$

sending $(A_0^b, A^b, t) \mapsto (A_0, A)$, where $A_0 = A_0^b$, and A is related to A^b by a diagram (6.5.5).

Proposition 6.5.3. *The hermitian line bundles*

$$\widehat{\omega}_{A_0^b/\mathcal{S}_{V^b}}^{\mathrm{Hdg}} + \widehat{\omega}_{A^b/\mathcal{S}_{V^b}}^{\mathrm{Hdg}} \in \widehat{\mathrm{Pic}}(\mathcal{S}_{V^b})$$

and $\widehat{\omega}_{A/\mathcal{S}_V}^{\mathrm{Hdg}} \in \widehat{\mathrm{Pic}}(\mathcal{S}_V)$ have the same images under the homomorphisms

$$\widehat{\mathrm{Pic}}(\mathcal{S}_{V^b}) \longrightarrow \widehat{\mathrm{Pic}}(\mathcal{T}_{V^b})_{\mathbb{Q}} \longleftarrow \widehat{\mathrm{Pic}}(\mathcal{S}_V)$$

induced by (6.5.12). The same is true of $(\mathrm{Exc}_{V^b}, 0)$ and $(\mathrm{Exc}_V, 0)$, and of $\widehat{\mathcal{L}}_{V^b}$ and $\widehat{\mathcal{L}}_V$.

Proof. If (A_0^b, A^b) denotes the pullback of the universal object via the left arrow in (6.5.12), and (A_0, A) denotes the pullback of the universal object via the right arrow in (6.5.12), examination of the proof of Proposition 6.5.2 shows that there is a quasi-isogeny

$$f \in \mathrm{Hom}(A_0^b \times A^b, A)[1/p]$$

of degree $\deg(f) = p$. As in the proof of Proposition 6.3.3, this implies that

$$\widehat{\omega}_{A/\mathcal{T}_{V^b}}^{\mathrm{Hdg}} = \widehat{\omega}_{A_0^b/\mathcal{T}_{V^b}}^{\mathrm{Hdg}} + \widehat{\omega}_{A^b/\mathcal{T}_{V^b}}^{\mathrm{Hdg}} + (0, -\log(p))$$

as elements of $\widehat{\mathrm{Pic}}(\mathcal{T}_{V^b})$, and the term $(0, -\log(p))$ is torsion. The first claim follows from this. The remaining claims are essentially the same as in Proposition 6.3.3, and we leave the details to the reader. \square

Proposition 6.5.4. *The nonexceptional locus $\mathcal{U}_0^{\text{nexc}} \subset \mathcal{U}_0$ satisfies*

$$\mathcal{U}_0^{\text{nexc}} \subset \mathcal{U}_0^\dagger.$$

Moreover, the morphism of Proposition 6.5.2 restricts to an isomorphism

$$\mathcal{S}_{V^b/\mathcal{O}_k[1/p]}^{\text{nexc}} \cong \mathcal{U}_0^{\text{nexc}}.$$

Proof. The proof is essentially the same as Proposition 6.3.4, and the details are left to the reader. \square

Proposition 6.5.5. *Denote by $\bar{\mathcal{T}}_{V^b}$ the normalization of $\mathcal{T}_{V^b} \rightarrow \bar{\mathcal{S}}_{V^b/\mathcal{O}_k[1/p]}$. The closed immersion $i_0 : \mathcal{T}_{V^b} \rightarrow \mathcal{U}_0$ of Proposition 6.5.2 extends to a proper morphism*

$$\bar{\mathcal{T}}_{V^b} \rightarrow \bar{\mathcal{U}}_0,$$

where the codomain is defined as the Zariski closure of

$$\mathcal{U}_0 \subset \bar{\mathcal{Z}}_V(p)/\mathcal{O}_k[1/p],$$

or, equivalently, as the normalization of $\mathcal{U}_0 \rightarrow \bar{\mathcal{S}}_{V/\mathcal{O}_k[1/p]}$.

Proof. Consider the universal triple (A_0, A^b, t) over \mathcal{T}_{V^b} , and let (A_0, A) be the pullback of the universal pair over \mathcal{S}_V via the composition

$$\mathcal{T}_{V^b} \xrightarrow{i_0} \mathcal{U}_0 \rightarrow \mathcal{S}_V.$$

We know that A^b extends to a semi-abelian scheme over $\bar{\mathcal{T}}_{V^b}$, obtained as a pullback via

$$\bar{\mathcal{T}}_{V^b} \rightarrow \bar{\mathcal{S}}_{V^b} \rightarrow \bar{\mathcal{M}}_{(n-2,1)},$$

and that A_0 extends to an elliptic curve over $\bar{\mathcal{T}}_{V^b}$, obtained as a pullback via

$$\bar{\mathcal{T}}_{V^b} \rightarrow \bar{\mathcal{S}}_{V^b} \rightarrow \mathcal{M}_{(1,0)},$$

Using Lemma 6.4.2 and the isogenies (6.5.5), we deduce that A also extends to a semi-abelian scheme over $\bar{\mathcal{T}}_{V^b}$. With this extension in hand, the proof is essentially identical to that of Proposition 6.3.5. \square

6.6. Proof of Theorems 6.1.2 and 6.1.4. Define an $\mathcal{O}_k[1/p]$ -stack

$$\mathcal{Z}_V^\dagger(p) = \mathcal{T}_{V^b} \sqcup \mathcal{S}_{V^b/\mathcal{O}_k[1/p]} \sqcup \mathcal{S}_{V^b/\mathcal{O}_k[1/p]}.$$

Propositions 6.3.2 and 6.5.2 provide us with a canonical closed immersion

$$\mathcal{Z}_V^\dagger(p) \xrightarrow{i=i_0 \sqcup i_p \sqcup i_{\bar{p}}} \mathcal{U}_0 \sqcup \mathcal{U}_p \sqcup \mathcal{U}_{\bar{p}} = \mathcal{Z}_V(p)/\mathcal{O}_k[1/p]$$

with image the closed substack $\mathcal{U}_0^\dagger \sqcup \mathcal{U}_p^\dagger \sqcup \mathcal{U}_{\bar{p}}^\dagger$. Hence we have a diagram

$$(6.6.1) \quad \begin{array}{ccc} \mathcal{Z}_V^\dagger(p) & \xrightarrow{i} & \mathcal{Z}_V(p)/\mathcal{O}_k[1/p] \\ \alpha \downarrow & & \downarrow \beta \\ \mathcal{S}_{V^b/\mathcal{O}_k[1/p]} & & \mathcal{S}_{V/\mathcal{O}_k[1/p]} \end{array}$$

in which α is a finite étale surjection of degree $p^{n-1} + 1$, and β is finite.

Lemma 6.6.1. *As divisors on $\mathcal{S}_{V/\mathcal{O}_k[1/p]}$, we have*

$$\mathcal{Z}_V(p)_{/\mathcal{O}_k[1/p]} = \mathcal{Z}_V^\dagger(p) + E,$$

where $\mathcal{Z}_V^\dagger(p)$ has no irreducible components contained in the exceptional divisor $\text{Exc}_{V/\mathcal{O}_k[1/p]} \subset \mathcal{S}_{V/\mathcal{O}_k[1/p]}$ of Definition 5.1.4, and E is supported entirely on the exceptional divisor.

Proof. The *exceptional locus* of $\mathcal{Z}_V^\dagger(p)$ is the preimage of the exceptional divisor under α , and an irreducible component of it is *exceptional* if it is contained in the exceptional locus. Similarly, the *exceptional locus* of $\mathcal{Z}_V(p)_{/\mathcal{O}_k[1/p]}$ is the preimage of the exceptional divisor under β , and an irreducible component of it is *exceptional* if it is contained in the exceptional locus.

The final claims of Propositions 6.3.4 and 6.5.4 show that i restricts to an isomorphism between the non-exceptional loci of the source and target. In particular, it establishes a bijection between the generic points of non-exceptional irreducible components, and corresponding non-exceptional generic have local rings of the same (finite) length.

However, the morphism α is finite étale, so no irreducible component of the source can map into the exceptional divisor of the target. Thus the closed immersion i actually establishes a bijection between irreducible components of the source and non-exceptional irreducible components of the target, in a way that matches up their multiplicities. The claim follows immediately. \square

Propositions 6.3.5 and 6.5.5 imply that the diagram above extends to

$$(6.6.2) \quad \begin{array}{ccc} \bar{\mathcal{Z}}_V^\dagger(p) & \xrightarrow{i} & \bar{\mathcal{Z}}_V(p)_{/\mathcal{O}_k[1/p]} \\ \alpha \downarrow & & \downarrow \beta \\ \bar{\mathcal{S}}_{V^b/\mathcal{O}_k[1/p]} & & \bar{\mathcal{S}}_{V/\mathcal{O}_k[1/p]}, \end{array}$$

where we have defined

$$(6.6.3) \quad \bar{\mathcal{Z}}_V^\dagger(p) = \bar{\mathcal{T}}_{V^b} \sqcup \bar{\mathcal{S}}_{V^b/\mathcal{O}_k[1/p]} \sqcup \bar{\mathcal{S}}_{V/\mathcal{O}_k[1/p]}.$$

Equivalently, this is the normalization of

$$\alpha : \mathcal{Z}_V^\dagger(p) \rightarrow \bar{\mathcal{S}}_{V^b/\mathcal{O}_k[1/p]}.$$

For any $N \geq 1$ prime to p , we can add level structure to $\bar{\mathcal{T}}_{V^b}$ by defining

$$\mathcal{T}_{V^b}(N) = \mathcal{T}_{V^b} \times_{\mathcal{S}_{V^b}} \mathcal{S}_{V^b}(N),$$

and letting $\bar{\mathcal{T}}_{V^b}(N)$ be the normalization of

$$\mathcal{T}_{V^b}(N) \rightarrow \bar{\mathcal{T}}_{V^b/\mathcal{O}_k[1/N]}.$$

Equivalently, this is the normalization of

$$\mathcal{T}_{V^b}(N) \rightarrow \bar{\mathcal{S}}_{V^b}(N)_{/\mathcal{O}_k[\frac{1}{Np}]}.$$

Lemma 6.6.2. *The stack $\bar{\mathcal{T}}_{V^b}(N)$ is regular, flat, and proper over $\mathcal{O}_k[\frac{1}{Np}]$, and is a projective scheme if $N \geq 3$. It is smooth in a neighborhood of its boundary*

$$\partial\bar{\mathcal{T}}_{V^b}(N) = \bar{\mathcal{T}}_{V^b}(N) \setminus \mathcal{T}_{V^b}(N),$$

which is a Cartier divisor smooth over $\mathcal{O}_k[\frac{1}{Np}]$.

Proof. The proof is similar to that of Proposition 4.3.3, and is again based on the results of [Lan16] and [MP19].

Recall from Remark 5.1.3 that the generic fiber of \mathcal{S}_{V^b} is the Shimura variety associated to a reductive group G^b and a certain compact open subgroup of $G^b(\mathbb{A}_f)$, and one can easily check that the generic fiber of the finite étale cover

$$(6.6.4) \quad \mathcal{T}_{V^b}(N) \rightarrow \mathcal{S}_{V^b}(N)_{/\mathcal{O}_k[\frac{1}{Np}]}$$

is obtained by shrinking compact open subgroup.

Recall from §5.1 that $\mathcal{S}_{V^b}(N)$ is constructed as an open and closed sub-stack

$$\mathcal{S}_{V^b}(N) \subset \mathcal{M}_{W_0}(N) \times_{\mathcal{O}_k[1/N]} \mathcal{M}_{W^b}(N).$$

In this construction, we may replace $\mathcal{M}_{W^b}(N)$ with the stack $\mathcal{M}_{W^b}^{\text{Pap}}(N)$ appearing in the proof of Proposition 4.3.3 to obtain a blow-down

$$\mathcal{S}_{V^b}^{\text{Pap}}(N) \subset \mathcal{M}_{W_0}(N) \times_{\mathcal{O}_k[1/N]} \mathcal{M}_{W^b}^{\text{Pap}}(N).$$

Repeating the construction of (6.6.4) with $\mathcal{S}_{V^b}(N)$ replaced by $\mathcal{S}_{V^b}^{\text{Pap}}(N)$ yields a finite étale cover

$$\mathcal{T}_{V^b}^{\text{Pap}}(N) \rightarrow \mathcal{S}_{V^b}^{\text{Pap}}(N)_{/\mathcal{O}_k[\frac{1}{Np}]},$$

and $\mathcal{T}_{V^b}(N)$ is the blow-up of the normal stack $\mathcal{T}_{V^b}^{\text{Pap}}(N)$ along its proper and 0-dimensional locus of nonsmooth points.

As in the proof of Proposition 4.3.3, we now have morphisms

$$\mathcal{T}_{V^b}^{\text{Pap}}(N) \rightarrow \mathcal{S}_{V^b}^{\text{Pap}}(N)_{/\mathcal{O}_k[\frac{1}{Np}]} \rightarrow \mathcal{M}_{W^b}^{\text{Pap}}(N)_{/\mathcal{O}_k[\frac{1}{Np}]} \rightarrow \bar{\mathcal{A}}_{/\mathcal{O}_k[\frac{1}{Np}]},$$

and we may define $\bar{\mathcal{T}}_{V^b}^{\text{Pap}}(N)$ as the normalization of the composition. We may now apply the results of [Lan16] and [MP19] to deduce properties of $\bar{\mathcal{T}}_{V^b}^{\text{Pap}}(N)$ from properties of its interior. Arguing exactly as in the proof of Proposition 4.3.3, this compactification is normal and flat, and is a projective scheme if $N \geq 3$. Its boundary is a smooth Cartier divisor. The nonsmooth locus has dimension 0, and does not meet the boundary. As $\bar{\mathcal{T}}_{V^b}(N)$ can be identified with the blow-up of $\bar{\mathcal{T}}_{V^b}^{\text{Pap}}(N)$ along its nonsmooth locus, it has all the same properties, and is also regular (being smooth near the boundary and regular in the interior). \square

Lemma 6.6.2 and Remark 5.1.8 provide us with an $\mathcal{O}_k[\frac{1}{Np}]$ -stack

$$\bar{\mathcal{Z}}_V^\dagger(p, N) \stackrel{\text{def}}{=} \bar{\mathcal{T}}_{V^b}(N) \sqcup \bar{\mathcal{S}}_{V^b}(N)_{/\mathcal{O}_k[\frac{1}{Np}]} \sqcup \bar{\mathcal{S}}_{V^b}(N)_{/\mathcal{O}_k[\frac{1}{Np}]},$$

with all the nice properties enumerated in Proposition 4.3.3. Thus (6.6.3) has its own theory of pre-log singular hermitian line bundles and Burgos-Kramer-Kühn arithmetic Chow groups

$$\widehat{\text{CH}}^d(\bar{\mathcal{Z}}_V^\dagger(p), \mathcal{D}_{\text{pre}}) = \varprojlim_{\substack{N \geq 3 \\ p \nmid N}} \widehat{\text{CH}}^d(\bar{\mathcal{Z}}_V^\dagger(p, N), \mathcal{D}_{\text{pre}}),$$

exactly as in §4.4. These Chow groups include notions of arithmetic heights and volumes as in §4.5, taking values in the abelian group $\mathbb{R}/\mathbb{Q} \log(p)$.

One can repeat the construction of the diagram (6.6.2) with level structures, and so obtain pullbacks

$$\widehat{\text{CH}}^d(\bar{\mathcal{S}}_{V^b}, \mathcal{D}_{\text{pre}}) \xrightarrow{\alpha^*} \widehat{\text{CH}}^d(\bar{\mathcal{Z}}_V^\dagger(p), \mathcal{D}_{\text{pre}}) \xleftarrow{(\beta \circ i)^*} \widehat{\text{CH}}^d(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})$$

and

$$\widehat{\text{Pic}}(\bar{\mathcal{S}}_{V^b}, \mathcal{D}_{\text{pre}})_{\mathbb{Q}} \xrightarrow{\alpha^*} \widehat{\text{Pic}}(\bar{\mathcal{Z}}_V^\dagger(p), \mathcal{D}_{\text{pre}})_{\mathbb{Q}} \xleftarrow{(\beta \circ i)^*} \widehat{\text{Pic}}(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})_{\mathbb{Q}}$$

Lemma 6.6.3. *If two pre-log singular hermitian line bundles*

$$\widehat{\mathcal{P}}^b \in \widehat{\text{Pic}}(\bar{\mathcal{S}}_{V^b}, \mathcal{D}_{\text{pre}}) \quad \text{and} \quad \widehat{\mathcal{P}} \in \widehat{\text{Pic}}(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})$$

have the same pullback to $\widehat{\text{Pic}}(\bar{\mathcal{Z}}_V^\dagger(p), \mathcal{D}_{\text{pre}})_{\mathbb{Q}}$ then

$$\int_{\mathcal{Z}_V(p)(\mathbb{C})} \text{ch}(\widehat{\mathcal{P}})^{n-2} = (p^{n-1} + 1) \int_{\mathcal{S}_{V^b}(\mathbb{C})} \text{ch}(\widehat{\mathcal{P}}^b)^{n-2}.$$

Moreover, there is an $a(p) \in \mathbb{Q}$ such that

$$\text{ht}_{\widehat{\mathcal{P}}}(\bar{\mathcal{Z}}_V(p)) = (p^{n-1} + 1) \cdot \widehat{\text{vol}}(\widehat{\mathcal{P}}^b) + \text{ht}_{\widehat{\mathcal{P}}}(E) + a(p) \log(p),$$

where E is the divisor of Lemma 6.6.1.

Proof. As in the proof of Lemma 6.6.1, the closed immersion i in (6.6.1) restricts to an isomorphism of non-exceptional loci, and hence induces an isomorphism in the generic fiber. Thus

$$\begin{aligned} \int_{\mathcal{Z}_V(p)(\mathbb{C})} \beta^* \text{ch}(\widehat{\mathcal{P}})^{n-2} &= \int_{\mathcal{Z}_V^\dagger(p)(\mathbb{C})} (\beta \circ i)^* \text{ch}(\widehat{\mathcal{P}})^{n-2} \\ &= \int_{\mathcal{Z}_V^\dagger(p)(\mathbb{C})} \alpha^* \text{ch}(\widehat{\mathcal{P}}^b)^{n-2} \\ &= (p^{n-1} + 1) \int_{\mathcal{S}_{V^b}(\mathbb{C})} \text{ch}(\widehat{\mathcal{P}}^b)^{n-2}. \end{aligned}$$

The second claim follows from Lemma 6.6.1 and the equalities

$$\text{ht}_{\widehat{\mathcal{P}}}(\bar{\mathcal{Z}}_V^\dagger(p)) = \widehat{\text{vol}}((\beta \circ i)^* \widehat{\mathcal{P}}) = \widehat{\text{vol}}(\alpha^* \widehat{\mathcal{P}}^b) = (p^{n-1} + 1) \cdot \widehat{\text{vol}}(\widehat{\mathcal{P}}^b)$$

in $\mathbb{R}/\mathbb{Q} \log(p)$, where the first and last equalities are obtained directly by unpacking the definitions of pullbacks, heights, and arithmetic intersections in [BKK07], and using the fact that α is (away from the boundary) a finite étale surjection of degree $p^{n-1} + 1$. \square

Proof of Theorems 6.1.2 and 6.1.4. Propositions 6.3.3 and 6.5.3 imply that the hypotheses of Lemma 6.6.3 are satisfied by

$$\widehat{\mathcal{P}}^b = \widehat{\mathcal{K}}_{V^b} \quad \text{and} \quad \widehat{\mathcal{P}} = \widehat{\mathcal{K}}_V$$

which therefore gives the equality

$$\int_{\mathcal{Z}_V(p)(\mathbb{C})} \text{ch}(\widehat{\mathcal{K}}_V)^{n-2} = (p^{n-1} + 1) \int_{\mathcal{S}_{V^b}(\mathbb{C})} \text{ch}(\widehat{\mathcal{K}}_{V^b})^{n-2},$$

and the equality

$$\text{ht}_{\widehat{\mathcal{K}}_V}(\bar{\mathcal{Z}}_V(p)) = (p^{n-1} + 1) \widehat{\text{vol}}(\widehat{\mathcal{K}}_{V^b}) + \text{ht}_{\widehat{\mathcal{K}}_V}(E)$$

up to a rational multiple of $\log(p)$. The second term on the right vanishes by claim (1) of Theorem 5.5.1, completing the proof of Theorem 6.1.2.

Similarly, Propositions 6.3.3 and 6.5.3 show that the hypotheses of Lemma 6.6.3 are satisfied by

$$\widehat{\mathcal{P}}^b = \widehat{\omega}_{A_0^b/S_{V^b}}^{\text{Hdg}} + \widehat{\omega}_{A^b/S_{V^b}}^{\text{Hdg}} \quad \text{and} \quad \widehat{\mathcal{P}} = \widehat{\omega}_{A/S_V}^{\text{Hdg}},$$

and so

$$\text{ht}_{\widehat{\omega}_{A/S_V}^{\text{Hdg}}}(\bar{\mathcal{Z}}_V(p)) = (p^{n-1} + 1) \cdot \widehat{\text{vol}}(\widehat{\omega}_{A_0^b/S_{V^b}}^{\text{Hdg}} + \widehat{\omega}_{A^b/S_{V^b}}^{\text{Hdg}}) + \text{ht}_{\widehat{\omega}_{A/S_V}^{\text{Hdg}}}(E)$$

up to a rational multiple of $\log(p)$. One again, the second term on the right vanishes by claim (1) of Theorem 5.5.1. For the first term on the right, Proposition 5.3.1 and Lemmas 4.5.3 and 4.5.4 imply

$$\begin{aligned} \widehat{\text{vol}}(\widehat{\omega}_{A_0^b/S_{V^b}}^{\text{Hdg}} + \widehat{\omega}_{A^b/S_{V^b}}^{\text{Hdg}}) &= \widehat{\text{vol}}((0, C_1) + \widehat{\omega}_{A^b/S_{V^b}}^{\text{Hdg}}) \\ &= \widehat{\text{vol}}(\widehat{\omega}_{A^b/S_{V^b}}^{\text{Hdg}}) + (n-1)C_1 \int_{\mathcal{S}_{V^b}(\mathbb{C})} \text{ch}(\widehat{\omega}_{A^b/S_{V^b}}^{\text{Hdg}})^{n-2}, \end{aligned}$$

where

$$C_1 = \log(2\pi) + 2h_{\mathbf{k}}^{\text{Falt}} = -\frac{L'(0, \varepsilon)}{L(0, \varepsilon)} - \frac{\log(D)}{2}.$$

This proves Theorem 6.1.4. \square

6.7. Comparing volumes. Recall from Remark 5.5.2 that when V has signature $(n-1, 1)$ with $n > 2$ (an assumption in force throughout §6) there is an explicit relation between the arithmetic volumes of $\widehat{\omega}_{A/S_V}^{\text{Hdg}}$ and $\widehat{\mathcal{K}}_V$.

The restriction to $n > 2$, which originated in Proposition 5.4.2, can now be removed using Theorems 6.1.2 and 6.1.4.

Proposition 6.7.1. *The arithmetic volume relation*

$$\widehat{\text{vol}}(\widehat{\mathcal{K}}_V) = \widehat{\text{vol}}(\widehat{\omega}_{A/S_V}^{\text{Hdg}}) + nC_0(n) \text{vol}_{\mathbb{C}}(\widehat{\omega}_{A/S_V}^{\text{Hdg}}),$$

known by Remark 5.5.2 when $n > 2$, also holds for $n = 2$.

Proof. Assume that V has signature $(1, 1)$, and fix a prime p split in \mathbf{k} .

Let V^\sharp be the \mathbf{k} -hermitian space of signature $(2, 1)$ whose local invariants satisfy

$$\mathrm{inv}_\ell(V) = (p, -D)_\ell \cdot \mathrm{inv}_\ell(V^\sharp)$$

for all places $\ell \leq \infty$, so that the discussion of §6.1 applies with the pair (V^\sharp, V) replacing (V, V^b) . Define $H_p \in \mathbb{R}$ by the relation

$$(p+1)H_p = \mathrm{ht}_{\widehat{\mathcal{K}}_{V^\sharp}}(\bar{\mathcal{Z}}_{V^\sharp}(p)) - \mathrm{ht}_{\widehat{\omega}_{A^\sharp/S_{V^\sharp}}^{\mathrm{Hdg}}}(\bar{\mathcal{Z}}_{V^\sharp}(p)).$$

On the one hand, part (3) of Theorem 5.5.1 and Lemma 4.5.3 imply

$$(6.7.1) \quad \mathrm{ht}_{\widehat{\mathcal{K}}_{V^\sharp}}(\bar{\mathcal{Z}}_{V^\sharp}(p)) = \mathrm{ht}_{\widehat{\omega}_{A^\sharp/S_{V^\sharp}}^{\mathrm{Hdg}} + (0, C_0(3))}(\bar{\mathcal{Z}}_{V^\sharp}(p)).$$

In the codimension 2 arithmetic Chow group of $\bar{\mathcal{S}}_{V^\sharp}$ we have the relation

$$\left(\widehat{\omega}_{A^\sharp/S_{V^\sharp}}^{\mathrm{Hdg}} + (0, C_0(3))\right)^2 = \left(\widehat{\omega}_{A^\sharp/S_{V^\sharp}}^{\mathrm{Hdg}}\right)^2 + \left(0, 2C_0(3) \mathrm{ch}(\widehat{\omega}_{A^\sharp/S_{V^\sharp}}^{\mathrm{Hdg}})\right),$$

as in the proof of Lemma 4.5.4. Directly from the definition of arithmetic height in §4.5, it follows that the right hand side of (6.7.1) is equal to

$$\mathrm{ht}_{\widehat{\omega}_{A^\sharp/S_{V^\sharp}}^{\mathrm{Hdg}}}(\bar{\mathcal{Z}}_{V^\sharp}(p)) + 2C_0(3) \int_{\bar{\mathcal{Z}}_{V^\sharp}(p)(\mathbb{C})} \mathrm{ch}(\widehat{\omega}_{A^\sharp/S_{V^\sharp}}^{\mathrm{Hdg}}).$$

Combining the first claim of Theorem 6.1.2 with the equality of Chern forms of Theorem 5.5.1, we find

$$\int_{\bar{\mathcal{Z}}_{V^\sharp}(p)(\mathbb{C})} \mathrm{ch}(\widehat{\omega}_{A^\sharp/S_{V^\sharp}}^{\mathrm{Hdg}}) = (p+1) \mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A/S_V}^{\mathrm{Hdg}}).$$

Putting all of this together allows us to rewrite (6.7.1) as

$$\mathrm{ht}_{\widehat{\mathcal{K}}_{V^\sharp}}(\bar{\mathcal{Z}}_{V^\sharp}(p)) = \mathrm{ht}_{\widehat{\omega}_{A^\sharp/S_{V^\sharp}}^{\mathrm{Hdg}}}(\bar{\mathcal{Z}}_{V^\sharp}(p)) + 2C_0(3)(p+1) \mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A/S_V}^{\mathrm{Hdg}}),$$

which is equivalent to

$$(6.7.2) \quad H_p = 2C_0(3) \mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A/S_V}^{\mathrm{Hdg}}).$$

On the other hand, the height formulas of Theorems 6.1.2 and 6.1.4 imply

$$\begin{aligned} H_p &= \widehat{\mathrm{vol}}(\widehat{\mathcal{K}}_V) - \widehat{\mathrm{vol}}(\widehat{\omega}_{A/S_V}^{\mathrm{Hdg}}) + 2 \left(\frac{L'(0, \varepsilon)}{L(0, \varepsilon)} + \frac{\log(D)}{2} \right) \mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A/S_V}^{\mathrm{Hdg}}) \\ &= \widehat{\mathrm{vol}}(\widehat{\mathcal{K}}_V) - \widehat{\mathrm{vol}}(\widehat{\omega}_{A/S_V}^{\mathrm{Hdg}}) + 2(C_0(3) - C_0(2)) \mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A/S_V}^{\mathrm{Hdg}}) \end{aligned}$$

up to a rational multiple of $\log(p)$, and combining this with (6.7.2) proves

$$\widehat{\mathrm{vol}}(\widehat{\mathcal{K}}_V) - \widehat{\mathrm{vol}}(\widehat{\omega}_{A/S_V}^{\mathrm{Hdg}}) - 2C_0(2) \mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A/S_V}^{\mathrm{Hdg}}) \in \mathbb{Q} \log(p).$$

The only way this can hold for all primes p split in \mathbf{k} is if this real number is 0, completing the proof when $n = 2$. \square

7. THE GENERATING SERIES OF ARITHMETIC DIVISORS

We continue to work with the Shimura variety \mathcal{S}_V of (5.1.4) associated to a \mathbf{k} -hermitian space V of signature $(n-1, 1)$ containing a self-dual $\mathcal{O}_{\mathbf{k}}$ -lattice. Throughout this section we assume $n > 2$.

After explaining the connection between the complex orbifold $\mathcal{S}_V(\mathbb{C})$ and the Shimura variety (3.1.3) associated to the unitary group $U(V)$, we will use the results of §3.3 to construct Green functions for the Kudla-Rapoport divisors $\mathcal{Z}_V(m) \rightarrow \mathcal{S}_V$.

7.1. Arithmetic Kudla-Rapoport divisors. We want to construct Green functions for the special divisors $\mathcal{Z}_V(m) \rightarrow \mathcal{S}_V$ using the machinery of regularized theta lifts from §3.3. In order to do this, we must make explicit the relation between the complex orbifold $\mathcal{S}_V(\mathbb{C})$ and the Shimura variety (3.1.3).

Recall from §5.1 that in the construction of \mathcal{S}_V we fixed an isomorphism

$$V \cong \mathrm{Hom}_{\mathbf{k}}(W_0, W),$$

where W_0 and W are \mathbf{k} -hermitian spaces of signatures $(1, 0)$ and $(n-1, 1)$, respectively. The subgroup $G \subset \mathrm{GU}(W_0) \times \mathrm{GU}(W)$ of Remark 5.1.3 acts on both W_0 and W via unitary similitudes. The group G also acts on V , defining a surjective homomorphism

$$G \rightarrow H = U(V),$$

whose kernel is the central $\mathrm{Res}_{\mathbf{k}/\mathbb{Q}}\mathbb{G}_m$ in G .

Fix self-dual $\mathcal{O}_{\mathbf{k}}$ -lattices $\mathfrak{a}_0 \subset W_0$ and $\mathfrak{a} \subset W$ as in §4.1, and define a self-dual $\mathcal{O}_{\mathbf{k}}$ -lattice

$$L = \mathrm{Hom}_{\mathcal{O}_{\mathbf{k}}}(\mathfrak{a}_0, \mathfrak{a}) \subset V.$$

Denote by $K \subset G(\mathbb{A}_f)$ the largest compact open subgroup fixing the lattices \mathfrak{a}_0 and \mathfrak{a} . By abuse of notation, denote in the same way the image of K under $G(\mathbb{A}_f) \rightarrow H(\mathbb{A}_f)$. We now have a commutative diagram

$$(7.1.1) \quad \begin{array}{ccc} \mathcal{S}_V(\mathbb{C}) & \longrightarrow & \mathrm{Sh}_K(H, \mathcal{D}) \\ \parallel & & \parallel \text{(3.1.3)} \\ G(\mathbb{Q}) \backslash \mathcal{D} \times G(\mathbb{A}_f) / K & \longrightarrow & H(\mathbb{Q}) \backslash \mathcal{D} \times H(\mathbb{A}_f) / K, \end{array}$$

in which the identification on the left is that of §2 of [BHK⁺a].

Let $H_{2-n}^{\infty}(D, \varepsilon^n)$ denote the space of \mathbb{C} -valued harmonic Maass forms f of weight $2-n$, level $\Gamma_0(D)$, and character ε^n such that

- f is bounded at all cusps of $\Gamma_0(D)$ different from the cusp ∞ ,
- f has polynomial growth at ∞ , in sense that there is a

$$P_f = \sum_{m \leq 0} c^+(m) q^m \in \mathbb{C}[q^{-1}]$$

such that $f - P_f = o(1)$ as q goes to 0.

Such a harmonic Maass form has a Fourier expansion analogous to (3.3.1) with Fourier coefficients $c^\pm(m) \in \mathbb{C}$.

Fix an $f \in H_{2-n}^\infty(D, \varepsilon^n)$ with Fourier coefficients $c^\pm(m)$. This form can be lifted to a vector valued harmonic Maass form, in the sense of §3.3, by setting

$$(7.1.2) \quad \tilde{f} = \sum_{\gamma \in \Gamma_0(D) \backslash \mathrm{SL}_2(\mathbb{Z})} (f|_{2-n}\gamma)(\omega_L(\gamma)^{-1}\varphi_0) \in H_{2-n}(\omega_L),$$

where $\varphi_0 \in S_L = \mathbb{C}[L'/L]$ is the characteristic function of $0 \in L'/L$. We denote the Fourier coefficients of \tilde{f} by $\tilde{c}^\pm(m, \mu)$ for $\mu \in L'/L$ and $m \in \mathbb{Q}$. The coefficients of \tilde{f} can be computed in terms of the coefficients of f , and for $m < 0$ we have

$$(7.1.3) \quad \tilde{c}^+(m, \mu) = \begin{cases} c^+(m) & \text{if } \mu = 0 \\ 0 & \text{if } \mu \neq 0 \end{cases}$$

as in Proposition 6.1.2 of [BHK⁺a] or §5 of [Sch09].

Under the covering map (7.1.1), the divisors $Z(m)$ and the hermitian line bundle $\hat{\mathcal{L}}$ defined in §3.1 pull back to the divisors $\mathcal{Z}_V(m)$ and $\hat{\mathcal{L}}_V$ defined in §5.1. This allows us to apply the construction (3.3.5) to the vector valued form (7.1.2) to obtain a Green function $\Phi(z, h, f)$ for the analytic divisor

$$\sum_{m>0} c^+(-m)Z(m) \in \mathrm{Div}_{\mathbb{C}}(\mathrm{Sh}_K(H, \mathcal{D}))$$

of (3.3.3), and then pull it back to a Green function $\Phi_V(f)$ for the divisor

$$\sum_{m>0} c^+(-m)\mathcal{Z}_V(m) \in \mathrm{Div}_{\mathbb{C}}(\mathcal{S}_V).$$

We will apply this construction for particular choices of f . As in §7.3 of [BHK⁺a], for every $m > 0$ there is a unique $f_m(\tau) \in H_{2-n}^\infty(D, \varepsilon^n)$ with q -expansion

$$f_m(\tau) = q^{-m} + O(1).$$

For any $f \in H_{2-n}^\infty(D, \varepsilon^n)$ with Fourier coefficients $c^\pm(m)$, we then have

$$f(\tau) = \sum_{m>0} c^+(-m)f_m(\tau).$$

Applying the above constructions to these particular harmonic forms yield Green functions

$$\Phi_V(m) \stackrel{\mathrm{def}}{=} \Phi_V(f_m)$$

for the divisors $\mathcal{Z}_V(m)$ on \mathcal{S}_V .

Recall from Remark 5.1.6 that $\bar{\mathcal{Z}}_V(m)$ is the Zariski closure of $\mathcal{Z}_V(m)$ in the toroidal compactification $\bar{\mathcal{S}}_V$ of §5.1. The following proposition shows that one can add boundary components to $\bar{\mathcal{Z}}_V(m)$ in order to obtain a class in the arithmetic Chow group (§4.4) of $\bar{\mathcal{S}}_V$. We will see in Theorem 7.3.1 below that these boundary components are (for our purposes) negligible.

Proposition 7.1.1. *Fix an integer $m > 0$, and recall the divisor $\bar{Z}_V(m)$ on \bar{S}_V from Remark 5.1.6. There is a unique divisor $\mathcal{B}_V(m)$ on \bar{S}_V that is supported on the boundary and satisfies*

$$\widehat{Z}_V^{\text{tot}}(m) \stackrel{\text{def}}{=} (\bar{Z}_V(m) + \mathcal{B}_V(m), \Phi_V(m)) \in \widehat{\text{CH}}^1(\bar{S}_V, \mathcal{D}_{\text{pre}}).$$

Proof. This follows from the analysis of $\Phi_V(m)$ near the boundary of \bar{S}_V carried out in [BHY15]; see Theorem 7.2.3 of [BHK⁺a]. A precise description of the boundary divisor $\mathcal{B}_V(m)$ is given by (5.3.3) of [BHK⁺a]. \square

Remark 7.1.2. As we assume $n > 2$, Theorem 3.4.2 and (7.1.3) imply the integral formula

$$\int_{\mathcal{S}_V(\mathbb{C})} \Phi_V(m) \text{ch}(\widehat{\mathcal{L}}_V)^{n-1} = \text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_V) B'(m, 0, s_0),$$

where $B'(m, 0, s_0)$ is the derivative at $s_0 = (n-1)/2$ of the Eisenstein series coefficient of (2.5.2), at $\mu = 0$, determined by the self-dual lattice $L \subset V$.

7.2. The modularity theorem. Proposition 7.1.1 defines arithmetic divisors

$$\widehat{Z}_V^{\text{tot}}(m) \in \widehat{\text{CH}}^1(\bar{S}_V, \mathcal{D}_{\text{pre}})$$

for all $m > 0$. We extend the definition to $m = 0$ by setting

$$\widehat{Z}_V^{\text{tot}}(0) = \widehat{\mathcal{L}}_V^{-1} + (\text{Exc}_V, -\log(D)),$$

where the second term is the exceptional divisor $\text{Exc}_V \subset \bar{S}_V$ of Definition 5.1.4 endowed with the constant Green function $-\log(D)$. The following is the main result of [BHK⁺a].

Theorem 7.2.1 (Theorem 7.3.1 of [BHK⁺a]). *The generating series*

$$(7.2.1) \quad \widehat{\phi}_V(\tau) = \sum_{m \geq 0} \widehat{Z}_V^{\text{tot}}(m) \cdot q^m \in \widehat{\text{CH}}^1(\bar{S}_V, \mathcal{D}_{\text{pre}})[[q]]$$

is the q -expansion of a modular form in $M_n(\Gamma_0(D), \varepsilon^n) \otimes \widehat{\text{CH}}^1(\bar{S}_V, \mathcal{D}_{\text{pre}})$.

A close examination of the proof of Theorem 7.2.1 reveals more information about the modular form (7.2.1). It admits a natural decomposition

$$(7.2.2) \quad \widehat{\phi}_V = \widehat{\phi}_V^{\text{eis}} + \widehat{\phi}_V^{\text{exc}} + \widehat{\phi}_V^{\text{cusp}}$$

as a sum of three forms. This decomposition will play an important role in §8.4, and so the remainder of this subsection is devoted to describing the three terms on the right.

Recall from Proposition 2.7.1 the Eisenstein series

$$E_r(\tau) = \sum_{m \geq 0} e_r(m) q^m \in M_n(\Gamma_0(D), \varepsilon^n)$$

indexed by a positive divisor $r \mid D$. In Proposition 2.7.3 we defined, for every divisor $r \mid D$, a constant $\gamma_r \in \{\pm 1, \pm i\}$. Using these, define the *Eisenstein part* of (7.2.1) by

$$\widehat{\phi}_V^{\text{eis}}(\tau) = \sum_{r \mid D} \gamma_r E_r(\tau) \cdot \left(\widehat{\mathcal{Z}}_V^{\text{tot}}(0) - \frac{1}{2}(\text{Exc}_V, 0) + \sum_{\ell \mid r} (\bar{\mathcal{S}}_{V/\mathbb{F}_\ell}, 0) \right),$$

where $\bar{\mathcal{S}}_{V/\mathbb{F}_\ell}$ is the special fiber of $\bar{\mathcal{S}}_V$ at the unique prime $\mathfrak{l} \subset \mathcal{O}_K$ above $\ell \mid r$. We may view $\ell \in \mathbf{k}^\times$ as a rational section of the trivial hermitian line bundle on $\bar{\mathcal{S}}_V$, and doing so shows that the class

$$(\text{div}(\ell), -\log |\ell|^2) \in \widehat{\text{Pic}}(\bar{\mathcal{S}}_V)_\mathbb{Q}$$

is trivial; here we are using the notation (4.5.5). This implies $(\bar{\mathcal{S}}_{V/\mathbb{F}_\ell}, 0) = (0, \log(\ell))$ in the arithmetic Chow group, and hence

$$(7.2.3) \quad \widehat{\phi}_V^{\text{eis}}(\tau) = \sum_{r \mid D} \gamma_r E_r(\tau) \left(\widehat{\mathcal{L}}_V^{-1} + \frac{1}{2}(\text{Exc}_V, -2 \log(D/r)) \right).$$

The *exceptional part* of (7.2.1) is

$$(7.2.4) \quad \widehat{\phi}_V^{\text{exc}}(\tau) = \frac{1}{2} \sum_{E \subset \text{Exc}_V} \vartheta_E(\tau) \cdot (E, 0),$$

where the sum extends over the irreducible components $E \subset \text{Exc}_V$ of the exceptional divisor on \mathcal{S}_V , and the scalar valued theta series $\vartheta_E(\tau)$ counts vectors in a positive definite rank n hermitian lattice determined by the component E . More precisely, recall that the top horizontal arrow in (5.6.1) collapses E to a connected 0-dimensional closed substack

$$e \subset \mathcal{M}_{(1,0)} \times_{\mathcal{O}_k} \mathcal{M}_{(n-1,1)}^{\text{Pap}}.$$

The hermitian lattice associated to E is then defined by

$$L_E = \text{Hom}_{\mathcal{O}_k}(A_{0,s}, A_s)$$

for any geometric point $s \rightarrow e$, as in §2.6 of [BHK⁺a]. For our purposes, all we need to know is that the coefficients of $\vartheta_E(\tau)$ are scalar multiples of arithmetic divisors of the form $(E, 0)$.

Having defined all but one of the four modular forms appearing in (7.2.2), we complete the definitions by setting

$$(7.2.5) \quad \widehat{\phi}_V^{\text{cusp}} = \widehat{\phi}_V - \widehat{\phi}_V^{\text{eis}} - \widehat{\phi}_V^{\text{exc}}.$$

The following result justifies calling this the *cuspidal part* of (7.2.1).

Proposition 7.2.2. *The generating series (7.2.5) is cuspidal, in the sense that it lies in the subspace*

$$S_n(\Gamma_0(D), \varepsilon^n) \otimes \widehat{\text{CH}}^1(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}}) \subset M_n(\Gamma_0(D), \varepsilon^n) \otimes \widehat{\text{CH}}^1(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}}).$$

Proof. The proposition is implicit in the proof of Theorem 7.3.1 of [BHK⁺a]. The essential point is that Theorem 4.2.3 of [BHK⁺a], a variant of a result of Borchers, gives a criterion for determining not just when a formal q -expansion is modular, but when it lies in the subspace

$$M_n^\infty(\Gamma_0(D), \varepsilon^n) \subset M_n(\Gamma_0(D), \varepsilon^n)$$

of forms that vanish at every cusp except possibly ∞ .

In the proof of Theorem 7.3.1 of [BHK⁺a], this criterion is used to show that the formal generating series

$$\widehat{\phi}_V - \widehat{\phi}_V^{\text{exc}} - \sum_{\substack{r|D \\ r>1}} \gamma_r E_r(\tau) \cdot \left(\widehat{\mathcal{Z}}_V^{\text{tot}}(0) - \frac{1}{2}(\text{Exc}_V, 0) + \sum_{\ell|r} (\bar{\mathcal{S}}_V/\mathbb{F}_\ell, 0) \right)$$

lies in

$$M_n^\infty(\Gamma_0(D), \varepsilon^n) \otimes \widehat{\text{CH}}^1(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})$$

(which immediately implies that $\widehat{\phi}_V$ is modular). This is spelled out more directly in the proof of Theorem 7.1.5 of [BHK⁺a], but for the generating series, valued in the usual Chow group of $\bar{\mathcal{S}}_V$, obtained from $\widehat{\phi}_V$ by forgetting the Green functions.

In other words,

$$\widehat{\phi}_V^{\text{cusp}}(\tau) + E_1(\tau) \cdot \left(\widehat{\mathcal{Z}}_V^{\text{tot}}(0) - \frac{1}{2}(\text{Exc}_V, 0) \right)$$

vanishes at every cusp except possibly ∞ . The Eisenstein series $E_1(\tau)$ has this same property (Remark 2.7.2), and therefore so also does $\widehat{\phi}_V^{\text{cusp}}(\tau)$. On the other hand, one can check directly (using Proposition 2.7.1 for the Eisenstein part) that the constant term in the q -expansion of (7.2.5) also vanishes, proving its cuspidality. \square

7.3. Heights of arithmetic divisors. Recall from (4.5.3) that if the divisor \mathcal{Z} underlying

$$\widehat{\mathcal{Z}} = (\mathcal{Z}, g_{\mathcal{Z}}) \in \widehat{\text{CH}}^1(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})$$

meets the boundary of $\bar{\mathcal{S}}_V$ properly in the generic fiber, then

$$\widehat{\text{deg}}(\widehat{\mathcal{Z}} \cdot \widehat{\mathcal{L}}_V^{n-1}) = \text{ht}_{\widehat{\mathcal{L}}_V}(\mathcal{Z}) + \int_{\mathcal{S}_V(\mathbb{C})} g_{\mathcal{Z}} \cdot \text{ch}(\widehat{\mathcal{L}}_V)^{n-1}.$$

We cannot apply this directly to the arithmetic divisor of Proposition 7.1.1, because the underlying divisor $\bar{\mathcal{Z}}_V(m) + \mathcal{B}_V(m)$ typically contains boundary components with nonzero multiplicities. The following result, as in §3.2 of [BBK07] or §3.4 of [Hor14], shows that even when $\mathcal{B}_V(m) \neq 0$ these boundary components are negligible for our purposes.

Theorem 7.3.1. *The arithmetic divisor of Proposition 7.1.1 satisfies*

$$\widehat{\text{deg}}(\widehat{\mathcal{Z}}_V^{\text{tot}}(m) \cdot \widehat{\mathcal{L}}_V^{n-1}) = \text{ht}_{\widehat{\mathcal{L}}_V}(\bar{\mathcal{Z}}_V(m)) + \int_{\mathcal{S}_V(\mathbb{C})} \Phi_V(m) \text{ch}(\widehat{\mathcal{L}}_V)^{n-1}.$$

The same relation holds if $\widehat{\mathcal{L}}_V$ is replaced by $\widehat{\omega}_{A/S_V}^{\text{Hdg}}$ or the hermitian line bundle \widehat{K}_V of Theorem 5.5.1.

The proof of Theorem 7.3.1 requires quite a bit of preparation, and will occupy the remainder of this subsection.

Fix an $N \geq 3$, and recall from §5.1 the finite morphism of $\mathcal{O}_k[1/N]$ -schemes

$$\bar{\mathcal{S}}_V(N) \rightarrow \bar{\mathcal{S}}_{V/\mathcal{O}_k[1/N]}$$

obtained by adding level structure and compactifying. To ease notation, the pullback of the special divisor $\bar{\mathcal{Z}}_V(m)$ via this map is denoted the same way, and similarly for \mathcal{L}_V and $\omega_{A/S_V}^{\text{Hdg}}$.

Proposition 7.3.2. *There is a positive integer k and sections*

$$f_1, \dots, f_{n-1} \in H^0(\bar{\mathcal{S}}_V(N), \omega_{A/S_V}^{\text{Hdg}, \otimes k})$$

with the following properties:

- (1) All possible intersections of

$$\bar{\mathcal{Z}}_V(m), \text{div}(f_1), \dots, \text{div}(f_{n-1})$$

are proper.

- (2) In the generic fiber, the divisor $\text{div}(f_1)$ is disjoint from the boundary.
 (3) The codimension 2 cycle $\text{div}(f_1) \cap \text{div}(f_2)$ is disjoint from the boundary.

Proof. This uses the theory of Baily-Borel (or minimal) compactifications of integral models developed in §5 of [MP19]. For the reasons explained after Lemma 4.3.2, this theory cannot be applied directly to the integral models $\mathcal{M}_W(N)$ or $\mathcal{S}_V(N)$, but it does apply to the integral models

$$\mathcal{M}_W^{\text{Pap}}(N) \subset \bar{\mathcal{M}}_W^{\text{Pap}}(N)$$

used in the proof of Proposition 4.3.3.

As in §5.2.1 of [MP19], the universal abelian scheme $A \rightarrow \mathcal{M}_W^{\text{Pap}}(N)$ extends uniquely to a semi-abelian scheme over its toroidal compactification, and the inverse of the determinant of its Lie algebra defines an extension of the Hodge bundle

$$\omega \stackrel{\text{def}}{=} \omega_{A/\mathcal{M}_W^{\text{Pap}}(N)}^{\text{Hdg}}$$

to an ample line bundle on the toroidal compactification. This allows us to define the Baily-Borel compactification

$$\mathcal{M}_W^{\text{Pap}}(N)^{\min} = \text{Proj} \left(\bigoplus_{r \geq 0} H^0(\bar{\mathcal{M}}_W^{\text{Pap}}(N), \omega^{\otimes r}) \right).$$

Theorem 5.2.11 of [MP19] provides us with a morphism

$$\bar{\mathcal{M}}_W^{\text{Pap}}(N) \rightarrow \mathcal{M}_W^{\text{Pap}}(N)^{\min}$$

of projective $\mathcal{O}_k[1/N]$ -schemes such that the composition

$$\mathcal{M}_W^{\text{Pap}}(N) \rightarrow \bar{\mathcal{M}}_W^{\text{Pap}}(N) \rightarrow \mathcal{M}_W^{\text{Pap}}(N)^{\min}$$

is an open immersion whose complement (the Baily-Borel boundary) is flat over $\mathcal{O}_k[1/N]$. In particular, as the classical Baily-Borel compactification of the complex fiber of $\mathcal{M}_W^{\text{Pap}}(N)$ has dimension 0, the boundary of the integral Baily-Borel compactification has dimension 1.

Moreover, the Hodge bundle ω on $\bar{\mathcal{M}}_W^{\text{Pap}}(N)$ is the pullback of a uniquely determined ample line bundle on the Baily-Borel compactification (denoted the same way), whose pullback via the composition

$$\bar{\mathcal{S}}_V(N) \xrightarrow{(5.1.6)} \bar{\mathcal{M}}_W(N) \rightarrow \bar{\mathcal{M}}_W^{\text{Pap}}(N) \rightarrow \mathcal{M}_W^{\text{Pap}}(N)^{\min}$$

agrees with $\omega_{A/S_V}^{\text{Hdg}}$.

Fix a positive integer k large enough that ω^k defines a closed immersion of $\mathcal{S}_W^{\text{Pap}}(N)^{\min}$ into projective space, and choose a hyperplane section

$$s_1 \in H^0(\mathcal{M}_W^{\text{Pap}}(N)^{\min}, \omega^{\otimes k})$$

whose divisor does not contain any of the (finitely many) boundary points of the generic fiber of $\mathcal{S}_W^{\text{Pap}}(N)^{\min}$, and does not contain the image of any generic point of the divisor $\bar{\mathcal{Z}}_V(m) \subset \bar{\mathcal{S}}_V(N)$. By construction, the divisor of the pullback

$$f_1 \in H^0(\bar{\mathcal{S}}_V(N), \omega_{A/S_V}^{\text{Hdg}, \otimes k})$$

of s_1 does not meet the toroidal boundary in the generic fiber, and intersects $\bar{\mathcal{Z}}_V(m)$ properly.

Now choose a hyperplane section

$$s_2 \in H^0(\mathcal{M}_W^{\text{Pap}}(N)^{\min}, \omega^{\otimes k})$$

whose divisor does not contain any point of the intersection $\text{div}(f_1)$ with the Baily-Borel boundary (this intersection is 0-dimensional, because the boundary is 1-dimensional), and does not contain the image of any generic point of $\text{div}(f_1)$, $\bar{\mathcal{Z}}_V(m)$, or their intersection. Now define

$$f_2 \in H^0(\bar{\mathcal{S}}_V(N), \omega_{A/S_V}^{\text{Hdg}, \otimes k})$$

as the pullback of s_2 , and continue in this fashion. \square

Corollary 7.3.3. *There is a positive integer k and sections*

$$F_1, \dots, F_{n-1} \in H^0(\bar{\mathcal{S}}_V(N), \mathcal{L}_V^{\otimes k})$$

with the following properties:

- (1) *On the complement of the exceptional divisor*

$$\text{Exc}_V(N) \stackrel{\text{def}}{=} \bar{\mathcal{S}}_V(N) \times_{\bar{\mathcal{S}}_V} \text{Exc}_V \subset \bar{\mathcal{S}}_V(N),$$

all possible intersections of

$$\bar{\mathcal{Z}}_V(m), \text{div}(F_1), \dots, \text{div}(F_{n-1})$$

- are proper. In particular, this holds on the generic fiber of $\bar{\mathcal{S}}_V(N)$.
- (2) In the generic fiber, the divisor $\text{div}(F_1)$ is disjoint from the boundary.
 - (3) The codimension 2 cycle $\text{div}(F_1) \cap \text{div}(F_2)$ is disjoint from the boundary.

Proof. This is clear from Propositions 5.5.3 and 7.3.2, and the fact that the exceptional divisor $\text{Exc}_V \subset \bar{\mathcal{S}}_V$ has empty generic fiber and does not intersect the boundary. \square

Proof of Theorem 7.3.1. Fix sections F_1, \dots, F_{n-1} as in Corollary 7.3.3. Each determines an arithmetic divisor

$$(\mathcal{Z}_i, g_i) \stackrel{\text{def}}{=} (\text{div}(F_i), -\log \|F_i\|^2)$$

representing $k \cdot \widehat{\mathcal{L}}_V \in \widehat{\text{CH}}^1(\bar{\mathcal{S}}_V(N), \mathcal{D}_{\text{pre}})$, and satisfying the Green equation

$$dd^c[g_i] + \delta_{\mathcal{Z}_i} = k[\text{ch}(\widehat{\mathcal{L}}_V)].$$

As the divisors $\mathcal{Z}_1, \dots, \mathcal{Z}_{n-1}$ intersect properly in the generic fiber, a direct examination of the construction of the arithmetic intersection pairing in §4.3 of [BKK07] shows that the iterated intersection

$$k^{n-1} \widehat{\mathcal{L}}_V^{n-1} = (\mathcal{Z}_1, g_1) \cdots (\mathcal{Z}_{n-1}, g_{n-1}) \in \widehat{\text{CH}}^{n-1}(\bar{\mathcal{S}}_V(N), \mathcal{D}_{\text{pre}})$$

is represented by a codimension $n - 1$ arithmetic cycle

$$\widehat{\mathcal{Y}} = (\mathcal{Y}, g_{\mathcal{Y}}),$$

in which the support of \mathcal{Y} is contained in the naive scheme-theoretic intersection $\mathcal{Z}_1 \cap \cdots \cap \mathcal{Z}_{n-1}$, and

$$g_{\mathcal{Y}} = g_1 * \cdots * g_{n-1}$$

is the iterated star product. (Really, this means the iterated star product of the pre-log-log Green objects

$$\mathfrak{g}_i = (k \text{ch}(\widehat{\mathcal{L}}_V), g_i) \in \widehat{H}_{\mathcal{D}_{\text{pre}}, \mathcal{Z}_i}^2(\bar{\mathcal{S}}_V(N)(\mathbb{C}), 1),$$

in the sense of [BBK07, BKK07], but we continue to suppress this extra notational complexity.)

Our choice of sections F_i guarantees that \mathcal{Y} does not intersect the boundary of $\bar{\mathcal{S}}_V(N)$, and does not intersect $\bar{\mathcal{Z}}_V(m)$ in the generic fiber. Thus

$$\begin{aligned} & k^{n-1} \widehat{\text{deg}}_N(\widehat{\mathcal{Z}}_V^{\text{tot}}(m) \cdot \widehat{\mathcal{L}}_V^{n-1}) \\ &= \widehat{\text{deg}}_N(\widehat{\mathcal{Z}}_V^{\text{tot}}(m) \cdot \widehat{\mathcal{Y}}) \\ (7.3.1) \quad &= \sum_{\substack{\mathfrak{p} \subset \mathcal{O}_{\mathbf{k}} \\ \mathfrak{p} \nmid N \mathcal{O}_{\mathbf{k}}}} \text{deg}_{\mathfrak{p}}(\bar{\mathcal{Z}}_V(m) \cdot \mathcal{Y}) \cdot \log(\#\mathcal{O}_{\mathbf{k}}/\mathfrak{p}) + \int_{\bar{\mathcal{S}}_V(N)(\mathbb{C})} \Phi_V(m) * g_{\mathcal{Y}}. \end{aligned}$$

Here we are working in the arithmetic Chow group of the $\mathcal{O}_{\mathbf{k}}[1/N]$ -scheme $\bar{\mathcal{S}}_V(N)$, as in §4.4, so these equalities are understood up to a \mathbb{Q} -linear combination of $\{\log(p) : p \mid N\}$. The degree $\text{deg}_{\mathfrak{p}}$ is that of (4.5.1), applied to

the 0-cycle

$$\bar{\mathcal{Z}}_V(m) \cdot \mathcal{Y} \in \mathrm{CH}_{\bar{\mathcal{Z}}_V(m) \cap \mathcal{Y}}^n(\bar{\mathcal{S}}_V(N))$$

as in the discussion of §4.5.

The integral on the right hand side of (7.3.1) is examined in Proposition 7.3.4 below. Substituting the expression found there into (7.3.1), and comparing with (4.5.2), we find the equality stated in Theorem 7.3.1 holds up to a \mathbb{Q} -linear combination of $\{\log(p) : p \mid N\}$. The theorem follows by allowing $N \geq 3$ to vary. \square

It remains to prove the formula for the integral in (7.3.1) used above.

Proposition 7.3.4. *Continuing with the notation in the above proof, we have*

$$\int_{\bar{\mathcal{S}}_V(N)(\mathbb{C})} \Phi_V(m) * g_{\mathcal{Y}} = k^{n-1} \int_{\mathcal{S}_V(N)(\mathbb{C})} \Phi_V(m) \mathrm{ch}(\widehat{\mathcal{L}}_V)^{n-1} + \int_{\mathcal{Z}_V(m)(\mathbb{C})} g_{\mathcal{Y}}.$$

Here, continuing with our abuse of notation, the final integral is really over the pullback of $\mathcal{Z}_V(m)(\mathbb{C})$ via $\mathcal{S}_V(N)(\mathbb{C}) \rightarrow \mathcal{S}_V(\mathbb{C})$.

Proof. We argue as in the proof of Theorem 3.3 of [BBK07]. See also Theorem 3.4.3 of [Hor14]. Abbreviate

$$\Omega = \mathrm{ch}(\widehat{\mathcal{L}}_V),$$

and denote by $D = \partial\bar{\mathcal{S}}_V(N)(\mathbb{C})$ the toroidal boundary divisor. For $\varepsilon > 0$ denote by $B_\varepsilon(D)$ an ε -neighborhood of D .

As in the proof above, the iterated arithmetic intersection

$$\widehat{\mathcal{Z}} \stackrel{\mathrm{def}}{=} (\mathcal{Z}_2, g_2) \cdots (\mathcal{Z}_{n-1}, g_{n-1}) \in \widehat{\mathrm{CH}}^{n-2}(\bar{\mathcal{S}}_V(N), \mathcal{D}_{\mathrm{pre}})$$

is represented by a codimension $n - 2$ arithmetic cycle

$$\widehat{\mathcal{Z}} = (\mathcal{Z}, g_{\mathcal{Z}}),$$

in which the support of \mathcal{Z} is contained in the naive scheme-theoretic intersection $\mathcal{Z}_2 \cap \cdots \cap \mathcal{Z}_{n-1}$, and the iterated star product

$$g_{\mathcal{Z}} = g_2 * \cdots * g_{n-1}$$

is a $(n - 3, n - 3)$ -form with a logarithmic singularity along $\mathcal{Z}(\mathbb{C})$ and a pre-log-log singularity along D , satisfying the Green equation

$$dd^c[g_{\mathcal{Z}}] + \delta_{\mathcal{Z}} = k^{n-2}[\Omega^{n-2}].$$

Recall from [BBK07, Section 1.3] that the star product

$$g_{\mathcal{Y}} = g_1 * g_{\mathcal{Z}}$$

can be represented as follows: Arguing as in [Bu97, p. 362] there exists an embedded resolution of singularities of $\mathcal{Z}_1(\mathbb{C}) \cup \mathcal{Z}(\mathbb{C})$, denoted

$$\pi : \tilde{X} \rightarrow \bar{\mathcal{S}}_V(N)(\mathbb{C}),$$

which factors through embedded resolutions of $\mathcal{Z}_1(\mathbb{C})$, $\mathcal{Z}(\mathbb{C})$, and $\mathcal{Z}_1(\mathbb{C}) \cup \mathcal{Z}(\mathbb{C})$, and such that the strict transforms of $\mathcal{Z}_1(\mathbb{C})$ and $\mathcal{Z}(\mathbb{C})$ are disjoint.

By the above assumptions on the F_i , the strict transform of $\mathcal{Z}_1(\mathbb{C})$ is also disjoint from $\pi^{-1}(D)$. Let $\sigma_{\mathcal{Z}}, 1 - \sigma_{\mathcal{Z}}$ be a partition of the unity on \tilde{X} for which $\sigma_{\mathcal{Z}} = 1$ in a neighborhood of the strict transform of $\mathcal{Z}(\mathbb{C}) \cup D$ and $\sigma_{\mathcal{Z}} = 0$ in a neighborhood of the strict transform of $\mathcal{Z}_1(\mathbb{C})$. Then, according to identity (1.5) of [BBK07],

$$g_{\mathcal{Y}} = k^{n-2} \sigma_{\mathcal{Z}} g_1 \wedge \Omega^{n-2} + (dd^c(1 - \sigma_{\mathcal{Z}}) g_1) \wedge g_{\mathcal{Z}}.$$

In particular, if ε is sufficiently small then on $B_\varepsilon(D)$ we have

$$(7.3.2) \quad g_{\mathcal{Y}} = k^{n-2} g_1 \wedge \Omega^{n-2}.$$

We may now apply [BBK07, Theorem 1.14] with $y = \mathcal{Z}_V^{\text{tot}}(m)(\mathbb{C})$, $g_y = \Phi_V(m)$, $z = \mathcal{Y}$, and $g_z = g_{\mathcal{Y}}$ to see that

$$\begin{aligned} \int_{\tilde{\mathcal{S}}_V(N)(\mathbb{C})} \Phi_V(m) * g_{\mathcal{Y}} &= \int_{\mathcal{Z}_V(m)(\mathbb{C})} g_{\mathcal{Y}} + k^{n-1} \int_{\tilde{\mathcal{S}}_V(N)(\mathbb{C})} \Phi_V(m) \Omega^{n-1} \\ &\quad - \lim_{\varepsilon \rightarrow 0} \int_{\partial B_\varepsilon(D)} (g_{\mathcal{Y}} \wedge d^c \Phi_V(m) - \Phi_V(m) \wedge d^c g_{\mathcal{Y}}). \end{aligned}$$

By (7.3.2) the limit on the right hand side is equal to

$$k^{n-2} \lim_{\varepsilon \rightarrow 0} \left(\int_{\partial B_\varepsilon(D)} g_1 \wedge \Omega^{n-2} \wedge d^c \Phi_V(m) - \Phi_V(m) \wedge d^c g_1 \wedge \Omega^{n-2} \right),$$

which vanishes by Lemma 7.3.5 below. \square

Lemma 7.3.5. *If F_1 is any meromorphic section of the line bundle $\mathcal{L}_V^{\otimes k}$ on $\tilde{\mathcal{S}}_V(N)(\mathbb{C})$, whose divisor is disjoint from the boundary, then*

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_{\partial B_\varepsilon(D)} \log \|F_1\| \wedge \Omega^{n-2} \wedge d^c \Phi_V(m) &= 0, \\ \lim_{\varepsilon \rightarrow 0} \int_{\partial B_\varepsilon(D)} \Phi_V(m) \wedge d^c \log \|F_1\| \wedge \Omega^{n-2} &= 0. \end{aligned}$$

Before we prove this lemma, we need to recall from Section 4.3 of [BHY15] the complex analytic description of the boundary of $\tilde{\mathcal{S}}_V(\mathbb{C})$. Using (7.1.1), it suffices to do this for the boundary of $\text{Sh}_K(H, \mathcal{D})$. By (3.1.4) the variety $\text{Sh}_K(H, \mathcal{D})$ is a finite disjoint union of connected Shimura varieties of the form $X_\Gamma = \Gamma \backslash \mathcal{D}$, where $\Gamma \subset H = \text{U}(V)$ is an arithmetic subgroup stabilizing a self-dual hermitian lattice in V . Hence it suffices to consider the boundary of the canonical toroidal compactification \tilde{X}_Γ of X_Γ .

The boundary of \tilde{X}_Γ is a smooth divisor. Its connected components correspond to Γ -orbits of isotropic lines in V . Fix an isotropic line $I \subset V$. Let $\ell \in I$ be a generator, and let $\tilde{\ell} \in V$ be isotropic such that $\langle \ell, \tilde{\ell} \rangle = 1$. The orthogonal complement $W = \ell^\perp \cap \tilde{\ell}^\perp$ is a positive definite hermitian space over \mathbf{k} of dimension $n - 2$, and we have

$$V = W \oplus \mathbf{k}\ell \oplus \mathbf{k}\tilde{\ell}.$$

Every $z \in \mathcal{D}$, viewed as a negative definite \mathbb{C} -line in $V(\mathbb{R})$ as in (3.1.1), has a unique basis vector of the form

$$\mathfrak{z} + \tau\sqrt{-D}\ell + \tilde{\ell}$$

with $\mathfrak{z} \in W(\mathbb{R})$ and $\tau \in \mathbb{C}$. The condition that z is negative definite is equivalent to the real number

$$N(\mathfrak{z}, \tau) \stackrel{\text{def}}{=} -\langle (\mathfrak{z}, \tau), (\mathfrak{z}, \tau) \rangle = 2\sqrt{D} \cdot \text{Im}(\tau) - \langle \mathfrak{z}, \mathfrak{z} \rangle$$

being positive. Consequently, \mathcal{D} is isomorphic to

$$\mathcal{H}_{\ell, \tilde{\ell}} \stackrel{\text{def}}{=} \{(\mathfrak{z}, \tau) \in W(\mathbb{R}) \times \mathcal{H} : 2\sqrt{D} \cdot \text{Im}(\tau) > \langle \mathfrak{z}, \mathfrak{z} \rangle\}.$$

The Chern form Ω is given in these coordinates by

$$(7.3.3) \quad \Omega = -dd^c \log N(\mathfrak{z}, \tau).$$

For $\varepsilon > 0$ we define

$$\begin{aligned} U_\varepsilon(\ell) &= \left\{ z \in \mathcal{D} : -\frac{\langle z, z \rangle}{|\langle z, \ell \rangle|^2} > \frac{1}{\varepsilon} \right\} \\ &\cong \{(\mathfrak{z}, \tau) \in \mathcal{H}_{\ell, \tilde{\ell}} : N(\mathfrak{z}, \tau) > 1/\varepsilon\}. \end{aligned}$$

The stabilizer $U(V)_\ell$ of $I = \mathbf{k}\ell$ acts on this subset. Let $\Gamma_\ell = \Gamma \cap U(V)_\ell$. If ε is sufficiently small, then

$$\Gamma_\ell \backslash U_\varepsilon(\ell) \longrightarrow X_\Gamma$$

is an open immersion. The center of $U(V)_\ell$ is given by the subgroup of translations T_a for $a \in \mathbb{Q}$, where

$$T_a(x) = x + a\langle x, \ell \rangle \sqrt{-D}\ell$$

for $x \in V$. The action of these translations on $\mathcal{H}_{\ell, \tilde{\ell}}$ is given by $T_a(\mathfrak{z}, \tau) = (\mathfrak{z}, \tau + a)$. The center of Γ_ℓ is of the form

$$\Gamma_{\ell, T} = \{T_a : a \in r\mathbb{Z}\}$$

for a unique $r \in \mathbb{Q}_{>0}$. If we put $q_r = e^{2\pi i \tau / r}$, then $(\mathfrak{z}, \tau) \mapsto (\mathfrak{z}, q_r)$ defines an isomorphism from $\Gamma_{\ell, T} \backslash U_\varepsilon(\ell)$ to

$$V_\varepsilon(\ell) = \left\{ (\mathfrak{z}, q_r) \in \mathbb{C}^{n-2} \times \mathbb{C} : 0 < |q_r| < \exp\left(-\frac{\pi}{r\sqrt{D}}(\langle \mathfrak{z}, \mathfrak{z} \rangle + 1/\varepsilon)\right) \right\}.$$

Adding the origin to this punctured disc bundle yields the disc bundle

$$\tilde{V}_\varepsilon(\ell) = \left\{ (\mathfrak{z}, q_r) \in \mathbb{C}^{n-2} \times \mathbb{C} : |q_r| < \exp\left(-\frac{\pi}{r\sqrt{D}}(\langle \mathfrak{z}, \mathfrak{z} \rangle + 1/\varepsilon)\right) \right\}.$$

The action of $\Gamma_\ell / \Gamma_{\ell, T}$ on $V_\varepsilon(\ell)$ extends to a properly discontinuous action on $\tilde{V}_\varepsilon(\ell)$, which leaves the boundary divisor $q_r = 0$ invariant. We obtain an open immersion of orbifolds

$$\Gamma_\ell \backslash U_\varepsilon(\ell) \longrightarrow (\Gamma_\ell / \Gamma_{\ell, T}) \backslash \tilde{V}_\varepsilon(\ell).$$

It can be used to glue the right hand side to X_Γ to obtain a partial compactification near the cusp I . The family of these open immersions for small

$\varepsilon > 0$ defines a base of open neighborhoods of the boundary divisor D_I associated with I .

Any boundary point $z \in D_I$ has a representative of the form $(\mathfrak{z}_0, 0) \in \tilde{V}_\varepsilon(\ell)$. The images of the sets

$$(7.3.4) \quad B_\delta(\mathfrak{z}_0, 0) = \left\{ (\mathfrak{z}, q_r) \in \tilde{V}_\varepsilon(\ell) : \langle \mathfrak{z} - \mathfrak{z}_0, \mathfrak{z} - \mathfrak{z}_0 \rangle < \delta, |q_r| < \delta \right\}.$$

in \tilde{X}_Γ for small $\delta > 0$ define a base of open neighborhoods of the boundary point.

Proof of Lemma 7.3.5. In view of the above local description of the boundary, it suffices to show that for every isotropic line $I \subset V$ and for a small neighborhood

$$Y = B_\delta(\mathfrak{z}_0, 0)$$

of any boundary point $(\mathfrak{z}_0, 0)$ of D_I we have

$$(7.3.5) \quad \lim_{\varepsilon \rightarrow 0} \int_{\partial B_\varepsilon(D) \cap Y} \log \|F_1\| \wedge \Omega^{n-2} \wedge d^c \Phi_V(m) = 0,$$

$$(7.3.6) \quad \lim_{\varepsilon \rightarrow 0} \int_{\partial B_\varepsilon(D) \cap Y} \Phi_V(m) \wedge d^c \log \|F_1\| \wedge \Omega^{n-2} = 0.$$

Using the coordinates of (7.3.4), we have

$$N(\mathfrak{z}, \tau) = -\frac{\sqrt{Dr}}{\pi} \log |q_r| - \langle \mathfrak{z}, \mathfrak{z} \rangle.$$

By (7.3.3) we find

$$\Omega = \frac{1}{N(\mathfrak{z}, \tau)^2} (dN(\mathfrak{z}, \tau)) \wedge (d^c N(\mathfrak{z}, \tau)) + \frac{1}{N(\mathfrak{z}, \tau)} dd^c \langle \mathfrak{z}, \mathfrak{z} \rangle.$$

If we write in addition $q_r = e^{2\pi i \tau / r} = t \cdot e^{i\rho}$ in polar coordinates with $t \geq 0$ and $\rho \in [0, 2\pi)$, and employ the identities $d \log |q_r| = \frac{dt}{t}$ and $d^c \log |q_r| = \frac{1}{4\pi} d\rho$, we obtain

$$\begin{aligned} \Omega &= \frac{1}{N(\mathfrak{z}, \tau)^2} \left(-\frac{\sqrt{Dr}}{\pi} \frac{dt}{t} - d\langle \mathfrak{z}, \mathfrak{z} \rangle \right) \wedge \left(-\frac{\sqrt{Dr}}{4\pi^2} d\rho - d^c \langle \mathfrak{z}, \mathfrak{z} \rangle \right) \\ &\quad + \frac{1}{N(\mathfrak{z}, \tau)} dd^c \langle \mathfrak{z}, \mathfrak{z} \rangle. \end{aligned}$$

Hence the restriction of Ω to $\partial B_\varepsilon(D) \cap Y$ is given by

$$\Omega|_{\partial B_\varepsilon(D) \cap Y} = \frac{1}{N(\mathfrak{z}, \tau)^2} d\langle \mathfrak{z}, \mathfrak{z} \rangle \wedge \left(\frac{\sqrt{Dr}}{4\pi^2} d\rho + d^c \langle \mathfrak{z}, \mathfrak{z} \rangle \right) + \frac{1}{N(\mathfrak{z}, \tau)} dd^c \langle \mathfrak{z}, \mathfrak{z} \rangle.$$

The differential form $d^c \log \|F_1\|$ is the sum of a smooth 1-form on Y and a constant multiple of $\frac{1}{N(\mathfrak{z}, \tau)} d^c N(\mathfrak{z}, \tau)$. This implies that

$$\Omega^{n-2} \wedge d^c \log \|F_1\| |_{\partial B_\varepsilon(D) \cap Y} = \frac{h(t, \rho, \mathfrak{z})}{|\log t|^{n-1}} d\rho \wedge d\mathfrak{z}_1 \dots d\mathfrak{z}_{n-2} \wedge d\bar{\mathfrak{z}}_1 \dots d\bar{\mathfrak{z}}_{n-2},$$

where h is a continuous function in $t \in [0, 1)$, $\rho \in [0, 2\pi]$ and \mathfrak{z} , which is smooth for $t > 0$.

On the other hand, according to [BHY15, Theorem 4.10], the restriction of the Green function $\Phi_V(m)$ to Y is the sum of a continuous function and constant multiples of the functions

$$\log t, \quad \log |\log t|, \quad \sum_{x \in S_m} \log |\langle x, \mathfrak{z} + \tilde{\ell} \rangle|,$$

where S_m denotes the finite set

$$S_m = \{x \in L \cap \ell^\perp : \langle x, \mathfrak{z}_0 + \tilde{\ell} \rangle = 0, \langle x, x \rangle = m\}.$$

This implies that

$$\int_{\partial B_\varepsilon(D) \cap Y} \Phi_V(m) \wedge d^c \log \|F_1\| \wedge \Omega^{n-2} \leq C \frac{1}{|\log \varepsilon|^{n-2}}$$

for some positive constant C . Since $n \geq 3$, we find that the limit in (7.3.6) vanishes.

For the limit in (7.3.5) we may argue similarly. Here [BHY15, Theorem 4.10] implies that the restriction of $d^c \Phi_V(m)$ to Y is the sum of a 1-form with log-log growth along $D \cap Y$ and constant multiples of the differentials

$$d\rho, \quad \frac{d\rho}{|\log t|}, \quad \sum_{x \in S_m} d^c \log |\langle x, \mathfrak{z} + \tilde{\ell} \rangle|.$$

On the other hand, $\log \|F_1\|$ is the sum of a continuous function and $\log |\log t|$. Putting the terms together, we obtain

$$\int_{\partial B_\varepsilon(D) \cap Y} \log \|F_1\| \wedge \Omega^{n-2} \wedge d^c \Phi_V(m) \leq C' \frac{\log |\log \varepsilon|}{|\log \varepsilon|^{n-2}}$$

for some positive constant C' . Again the limit as $\varepsilon \rightarrow 0$ vanishes. \square

Lemma 7.3.5 completes the proof of Proposition 7.3.4, which completes the proof of Theorem 7.3.1.

8. THE VOLUME CALCULATIONS

This section contains our main results. We continue to work with the Shimura variety \mathcal{S}_V of §5.1.4 defined by a \mathbf{k} -hermitian space V of signature $(n-1, 1)$ with $n \geq 1$, containing a self-dual $\mathcal{O}_{\mathbf{k}}$ -lattice $L \subset V$. We maintain the assumption that D is odd imposed since §4.

We will use induction on n to compute the complex and arithmetic volumes (as defined in §4.5) of the hermitian line bundle

$$\widehat{\mathcal{K}}_V = 2\widehat{\mathcal{L}}_V - (\text{Exc}_V, 0) \in \widehat{\text{Pic}}(\overline{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})$$

of Theorem 5.5.1. All results stated in the introduction will follow as consequences of this calculation.

8.1. Volumes of Shimura curves. First we consider the case $n = 2$, so that V has signature $(1, 1)$ and $\bar{\mathcal{S}}_V(\mathbb{C})$ has dimension one. The following proposition is a consequence of the main result of [How20], which is itself a consequence of the calculation of arithmetic volumes of modular curves and quaternion Shimura curves due to Kühn [Kuhn01], Bost (unpublished), and Kudla-Rapoport-Yang [KRY06].

Proposition 8.1.1. *Suppose $n = 2$. The metrized Hodge bundle of $A \rightarrow \mathcal{S}_V$ has complex volume*

$$\mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{S}_V}^{\mathrm{Hdg}}) = \frac{|\mathrm{CL}(\mathbf{k})|^2}{2^{o(D)-1} \cdot 12 \cdot |\mathcal{O}_{\mathbf{k}}^{\times}|^2} \prod_{\ell|D} (1 + \ell^*)$$

and arithmetic volume

$$\widehat{\mathrm{vol}}(\widehat{\omega}_{A/\mathcal{S}_V}^{\mathrm{Hdg}}) = \left(-2 - \frac{4\zeta'(-1)}{\zeta(-1)} - \sum_{\ell|D} \frac{1 - \ell^*}{1 + \ell^*} \cdot \log(\ell) \right) \mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{S}_V}^{\mathrm{Hdg}}),$$

where $\mathrm{CL}(\mathbf{k})$ is the class group of \mathbf{k} , $o(D)$ is the number of prime divisors of D , and we abbreviate

$$\ell^* = \left(\frac{-1}{\ell} \right) \mathrm{inv}_{\ell}(V)\ell.$$

Proof. As in (5.1.4), we realize

$$\mathcal{S}_V \subset \mathcal{M}_{W_0} \times_{\mathcal{O}_{\mathbf{k}}} \mathcal{M}_W$$

as a union of connected components, where W_0 and W have signatures $(1, 0)$ and $(1, 1)$, respectively.

The $\mathrm{GU}(W)$ Shimura datum defining the generic fibers of $\mathcal{M}_W^{\mathrm{Pap}}$ and \mathcal{M}_W actually has reflex field \mathbb{Q} rather than \mathbf{k} , and in §4 of [How20] one finds the definition of a regular and flat Deligne-Mumford stack

$$\mathcal{X}_W \rightarrow \mathrm{Spec}(\mathbb{Z})$$

such that $\mathcal{M}_W^{\mathrm{Pap}} \cong \mathcal{X}_{W/\mathcal{O}_{\mathbf{k}}}$, and such that the universal abelian surface over $\mathcal{M}_W^{\mathrm{Pap}}$ descends to $A \rightarrow \mathcal{X}_W$. The stack \mathcal{X}_W has its own canonical toroidal compactification and its own theory of arithmetic Chow groups, and Theorem A of [How20] implies

$$(8.1.1) \quad \mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{X}_W}^{\mathrm{Hdg}}) = \frac{|\mathrm{CL}(\mathbf{k})|}{2^{o(D)-1} \cdot 12 \cdot |\mathcal{O}_{\mathbf{k}}^{\times}|} \prod_{\ell|D} (1 + \ell^*)$$

and

$$(8.1.2) \quad \widehat{\mathrm{vol}}(\widehat{\omega}_{A/\mathcal{X}_W}^{\mathrm{Hdg}}) = \left(-1 - \frac{2\zeta'(-1)}{\zeta(-1)} - \frac{1}{2} \sum_{\ell|D} \frac{1 - \ell^*}{1 + \ell^*} \cdot \log(\ell) \right) \mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{X}_W}^{\mathrm{Hdg}}).$$

The composition

$$\mathcal{S}_V \rightarrow \mathcal{M}_W \rightarrow \mathcal{M}_W^{\text{Pap}} \cong \mathcal{X}_{W/\mathcal{O}_k}$$

extends to a proper morphism

$$\varphi : \bar{\mathcal{S}}_V \rightarrow \bar{\mathcal{X}}_{V/\mathcal{O}_k},$$

which, using Remark 5.1.2, restricts to a finite étale surjection of degree

$$\deg(\varphi) = |\text{CL}(\mathbf{k})|/|\mathcal{O}_k^\times|$$

from $\bar{\mathcal{S}}_V \setminus \text{Exc}_V$ to smooth locus of $\bar{\mathcal{X}}_{V/\mathcal{O}_k}$.

If $\widehat{\mathcal{L}}$ is any pre-log singular hermitian line bundle on $\bar{\mathcal{X}}_V$, and $\widehat{\mathcal{L}}_{/\mathcal{O}_k}$ is its base change to $\bar{\mathcal{X}}_{V/\mathcal{O}_k}$, then unpacking the definitions shows that⁴

$$\text{vol}_{\mathbb{C}}(\varphi^* \widehat{\mathcal{L}}_{/\mathcal{O}_k}) = \deg(\varphi) \cdot \text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}})$$

and

$$\widehat{\text{vol}}(\varphi^* \widehat{\mathcal{L}}_{/\mathcal{O}_k}) = 2 \cdot \deg(\varphi) \cdot \widehat{\text{vol}}(\widehat{\mathcal{L}}).$$

The proposition follows by applying these to the hermitian line bundles

$$\widehat{\mathcal{L}} = \widehat{\omega}_{A/\mathcal{X}_V}^{\text{Hdg}}, \quad \varphi^* \widehat{\mathcal{L}}_{/\mathcal{O}_k} = \widehat{\omega}_{A/\mathcal{S}_V}^{\text{Hdg}}$$

and using (8.1.1) and (8.1.2). \square

We now express the complex and arithmetic volumes of $\widehat{\mathcal{K}}_V$ in terms of the meromorphic function $\mathbf{A}_V(s)$ defined by (1.1.5). This will form the base case of the inductive proofs of Theorems 8.2.3 and 8.4.2 below.

Corollary 8.1.2. *Suppose $n = 2$. The hermitian line bundle (5.5.1) has complex volume*

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) = \frac{|\text{CL}(\mathbf{k})|}{2^{\sigma(D)-2} |\mathcal{O}_k^\times|} \cdot \mathbf{A}_V(0),$$

and arithmetic volume

$$\widehat{\text{vol}}(\widehat{\mathcal{K}}_V) = \left(2 \frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \log(D) \right) \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V).$$

Proof. Replacing W by V in (1.1.3), and using $D \equiv 3 \pmod{4}$, we find

$$\prod_{\ell|D} \left(\frac{-1}{\ell} \right) \text{inv}_{\ell}(V) = 1.$$

⁴The factor of 2 appears in the second equality because it is the degree of the finite flat morphism $\mathcal{X}_{W/\mathcal{O}_k} \rightarrow \mathcal{X}_W$. It does not appear in the first equality because, under the conventions of Remark 4.4.2, this morphism induces an isomorphism on complex points.

Recalling (1.1.5), this proves the second equality in

$$\begin{aligned} \mathbf{A}_V(0) &= \frac{D^{\frac{3}{2}}L(1, \varepsilon)\zeta(2)}{8\pi^3} \prod_{\ell|D} \left(1 + \left(\frac{-1}{\ell}\right) \text{inv}_\ell(V)\ell^{-1}\right) \\ &= \frac{D^{\frac{1}{2}}L(1, \varepsilon)\zeta(2)}{8\pi^3} \prod_{\ell|D} (1 + \ell^*) \\ &= \frac{|\text{CL}(\mathbf{k})|}{24 \cdot |\mathcal{O}_{\mathbf{k}}^\times|} \prod_{\ell|D} (1 + \ell^*). \end{aligned}$$

The final equality follows from $\zeta(2) = \pi^2/6$ and Dirichlet's class number formula

$$D^{\frac{1}{2}}L(1, \varepsilon) = 2\pi|\text{CL}(\mathbf{k})|/|\mathcal{O}_{\mathbf{k}}^\times|.$$

Comparing this with Proposition 8.1.1, and using the equality of Chern forms of Theorem (5.5.1), shows

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) = \text{vol}_{\mathbb{C}}(\widehat{\omega}_{A/S_V}^{\text{Hdg}}) = \frac{|\text{CL}(\mathbf{k})|}{2^{o(D)-2}|\mathcal{O}_{\mathbf{k}}^\times|} \cdot \mathbf{A}_V(0).$$

For the second claim, taking the logarithmic derivative of (1.1.5) yields

$$\begin{aligned} \frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} &= \frac{\mathbf{a}'_1(0)}{\mathbf{a}_1(0)} + \frac{\mathbf{a}'_2(0)}{\mathbf{a}_2(0)} - \sum_{\ell|D} \frac{\log(\ell)}{1 + \ell^*} \\ &= \frac{\mathbf{a}'_1(0)}{\mathbf{a}_1(0)} + \frac{\mathbf{a}'_2(0)}{\mathbf{a}_2(0)} - \frac{1}{2} \log(D) - \frac{1}{2} \sum_{\ell|D} \left(\frac{1 - \ell^*}{1 + \ell^*}\right) \log(\ell). \end{aligned}$$

One can use the functional equation of $L(s, \varepsilon^k)$ to see that

$$(8.1.3) \quad \frac{\mathbf{a}'_k(0)}{\mathbf{a}_k(0)} = -2 \frac{L'(1-k, \varepsilon^k)}{L(1-k, \varepsilon^k)} - \frac{\Gamma'(k)}{\Gamma(k)} + \log\left(\frac{4\pi}{D}\right) + (-1)^k \log(D)$$

for all $k \geq 1$. Recalling the constant $C_0(2)$ of Theorem A, and using the well-known formulas $\Gamma'(1)/\Gamma(1) = -\gamma$ and $\Gamma'(2)/\Gamma(2) = 1 - \gamma$, we find

$$\frac{\mathbf{a}'_1(0)}{\mathbf{a}_1(0)} + \frac{\mathbf{a}'_2(0)}{\mathbf{a}_2(0)} = -1 + C_0(2) - 2 \frac{\zeta'(-1)}{\zeta(-1)}.$$

Putting this all together gives

$$2 \frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} - 2C_0(2) + \log(D) = -2 - 4 \frac{\zeta'(-1)}{\zeta(-1)} - \sum_{\ell|D} \left(\frac{1 - \ell^*}{1 + \ell^*}\right) \log(\ell),$$

and comparing with Proposition 8.1.1 shows that

$$\widehat{\text{vol}}(\widehat{\omega}_{A/S_V}^{\text{Hdg}}) = \left(2 \frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} - 2C_0(2) + \log(D)\right) \text{vol}_{\mathbb{C}}(\widehat{\omega}_{A/S_V}^{\text{Hdg}}).$$

The second claim follows from this using Proposition 6.7.1 and the equality of Chern forms of Theorem 5.5.1. \square

8.2. The complex volume formula. We now use induction on $n \geq 2$ to compute the complex volumes of $\widehat{\mathcal{L}}_V$ and $\widehat{\mathcal{K}}_V$ on the Shimura variety $\bar{\mathcal{S}}_V$ associated to a relevant hermitian space V of signature $(n-1, 1)$.

Taking arithmetic intersection against the fixed arithmetic cycle class

$$\widehat{\mathcal{L}}_V^{n-2} \in \widehat{\text{CH}}^{n-2}(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})_{\mathbb{Q}}$$

and then applying the complex degree of §4.5 defines a linear functional

$$\widehat{\text{CH}}^1(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}}) \rightarrow \mathbb{Q},$$

denoted

$$(8.2.1) \quad \widehat{\mathcal{Z}} \mapsto \deg_{\mathbb{C}}(\widehat{\mathcal{Z}} \cdot \widehat{\mathcal{L}}_V^{n-2})$$

and called *complex intersection against $\widehat{\mathcal{L}}_V^{n-2}$* . Of course this only depends on the underlying divisor \mathcal{Z} and line bundle \mathcal{L}_V on $\bar{\mathcal{S}}_V$, and in fact only on their restrictions to the complex fiber.

Assuming that $n > 2$ (an assumption imposed throughout §7), we may apply (8.2.1) coefficient-by-coefficient to the generating series of special divisors (7.2.1). This yields a modular form

$$\deg_{\mathbb{C}}(\widehat{\phi}_V(\tau) \cdot \widehat{\mathcal{L}}_V^{n-2}) = \sum_{m \geq 0} \deg_{\mathbb{C}}(\widehat{\mathcal{Z}}_V^{\text{tot}}(m) \cdot \widehat{\mathcal{L}}_V^{n-2}) \cdot q^m \in M_n(\Gamma_0(D), \varepsilon^n)$$

with rational coefficients, and similarly for the other forms appearing in the decomposition

$$\widehat{\phi}_V(\tau) = \widehat{\phi}_V^{\text{eis}}(\tau) + \widehat{\phi}_V^{\text{exc}}(\tau) + \widehat{\phi}_V^{\text{cusp}}(\tau)$$

of (7.2.2). Note that we have the obvious equality

$$(8.2.2) \quad \deg_{\mathbb{C}}(\widehat{\phi}_V^{\text{exc}}(\tau) \cdot \widehat{\mathcal{L}}_V^{n-2}) = 0,$$

as the coefficients of the generating series (7.2.4) lie in the kernel of

$$\widehat{\text{CH}}^1(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}}) \rightarrow \text{CH}^1(\bar{\mathcal{S}}_{V/\mathbb{C}}).$$

Proposition 8.2.1. *For $n > 2$, we have the equalities of modular forms*

$$\begin{aligned} \deg_{\mathbb{C}}(\widehat{\phi}_V(\tau) \cdot \widehat{\mathcal{L}}_V^{n-2}) &= -\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_V) + \sum_{m > 0} \left(\int_{\mathcal{Z}_V(m)(\mathbb{C})} \text{ch}(\widehat{\mathcal{L}}_V)^{n-2} \right) \cdot q^m \\ \deg_{\mathbb{C}}(\widehat{\phi}_V^{\text{eis}}(\tau) \cdot \widehat{\mathcal{L}}_V^{n-2}) &= -\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_V) \sum_{r|D} \gamma_r E_r(\tau). \end{aligned}$$

Proof. The second claim follows easily from (7.2.3) and the definitions.

For the first claim, we compare the two modular forms coefficient-by-coefficient. As the equality of constant terms is obvious, it suffices to prove that

$$(8.2.3) \quad \deg_{\mathbb{C}}(\widehat{\mathcal{Z}}_V^{\text{tot}}(m) \cdot \widehat{\mathcal{L}}_V^{n-2}) = \int_{\mathcal{Z}_V(m)(\mathbb{C})} \text{ch}(\widehat{\mathcal{L}}_V)^{n-2}$$

for all $m > 0$. The left hand side is equal to the image of

$$(\bar{\mathcal{Z}}_V(m) + \mathcal{B}_V(m)) \cdot \mathcal{L}_V^{n-2} \in \text{CH}^{n-1}(\bar{\mathcal{S}}_V)$$

under the composition

$$(8.2.4) \quad \text{CH}^{n-1}(\bar{\mathcal{S}}_V) \rightarrow \text{CH}^{n-1}(\bar{\mathcal{S}}_{V/\mathbb{C}}) \xrightarrow{\text{deg}} \mathbb{Q}$$

of the restriction to the complex fiber and the usual degree of a 0-cycle. As in the proof of Theorem 7.3.1, one may represent the cycle class $\mathcal{L}_V^{n-2} \in \text{CH}^{n-2}(\bar{\mathcal{S}}_V)$ by a 1-cycle that does not meet the boundary in the generic fiber, and hence the left hand side of (8.2.3) is equal to the image of

$$\bar{\mathcal{Z}}_V(m) \cdot \mathcal{L}_V^{n-2} \in \text{CH}^{n-1}(\bar{\mathcal{S}}_V)$$

under the composition (8.2.4).

Fix a smooth (on all of $\bar{\mathcal{S}}_V(\mathbb{C})$, not just the interior) metric on $\widehat{\mathcal{L}}_V$, and let φ be the Chern form of this metric. The exterior power

$$\varphi^{n-2} \in H^{2n-4}(\bar{\mathcal{S}}_V(\mathbb{C}), \mathbb{C})$$

agrees with the cycle class of \mathcal{L}_V^{n-2} . As the second arrow in (8.2.4) factors through the cycle class map to cohomology, it follows that

$$\text{deg}_{\mathbb{C}}(\widehat{\mathcal{Z}}_V^{\text{tot}}(m) \cdot \widehat{\mathcal{L}}_V^{n-2}) = \int_{\mathcal{Z}_V(m)(\mathbb{C})} \varphi^{n-2}.$$

We cannot apply this directly to $\varphi = \text{ch}(\widehat{\mathcal{L}}_V)$ because the metric on $\widehat{\mathcal{L}}_V$ does not extend smoothly across the boundary, but it is easy to see that

$$\varphi^{n-2} = \text{ch}(\widehat{\mathcal{L}}_V)^{n-2} + d\alpha$$

for a pre-log-log form α on $\mathcal{S}_V(\mathbb{C})$. Propositions 7.6 and 7.12 of [BKK07] imply that the pullback of α to $\mathcal{Z}_V(m)(\mathbb{C})$ is locally integrable on the closure $\bar{\mathcal{Z}}_V(m)(\mathbb{C})$, and satisfies the equality of currents $[d\alpha] = d[\alpha]$. Evaluating both sides of this equation at the constant function 1 yields

$$\int_{\mathcal{Z}_V(m)(\mathbb{C})} d\alpha = 0,$$

completing the proof. \square

Recall from §2.6 and §2.7 the holomorphic Eisenstein series

$$\sum_{m \geq 0} B(m, 0, s_0) \cdot q^m \in M_n(\Gamma_0(D), \varepsilon^n)$$

associated to the self-dual hermitian lattice $L \subset V$.

Proposition 8.2.2. *Assume $n > 2$. We have the equality of modular forms*

$$\text{deg}_{\mathbb{C}}(\widehat{\phi}_V(\tau) \cdot \widehat{\mathcal{L}}_V^{n-2}) = \text{deg}_{\mathbb{C}}(\widehat{\phi}_V^{\text{eis}}(\tau) \cdot \widehat{\mathcal{L}}_V^{n-2}).$$

Equivalently, given (8.2.2),

$$\text{deg}_{\mathbb{C}}(\widehat{\phi}_V^{\text{cusp}}(\tau) \cdot \widehat{\mathcal{L}}_V^{n-2}) = 0.$$

Proof. Given Proposition 8.2.1 and the equality

$$\sum_{r|D} \gamma_r E_r(\tau) = \sum_{m \geq 0} B(m, 0, s_0) \cdot q^m$$

of Proposition 2.7.3, the first claim is just a restatement of the equality

$$\int_{\mathcal{Z}_V(m)(\mathbb{C})} \text{ch}(\widehat{\mathcal{L}}_V)^{n-2} = -\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_V) B(m, 0, s_0)$$

of Remark 3.1.1. Here we are using the discussion of §7.1, and especially the diagram (7.1.1), to relate the Shimura variety for the unitary group $H = \text{U}(V)$ used throughout §3.1 to the complex orbifold $\mathcal{S}_V(\mathbb{C})$ currently under consideration. \square

Theorem 8.2.3. *If $n \geq 2$, the complex volumes of (5.2.3) and (5.5.1) are*

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_V) = \frac{|\text{CL}(\mathbf{k})|}{2^{o(D)-1} |\mathcal{O}_{\mathbf{k}}^{\times}|} \cdot \mathbf{A}_V(0)$$

and

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) = \frac{|\text{CL}(\mathbf{k})|}{2^{o(D)-n} |\mathcal{O}_{\mathbf{k}}^{\times}|} \cdot \mathbf{A}_V(0).$$

Proof. The two claims are equivalent, by the equality of Chern forms of Theorem 5.5.1, and both hold when $n = 2$ by Corollary 8.1.2. It therefore suffices to assume that $n > 2$, and that the first equality holds for all relevant (Definition 4.1.3) hermitian spaces V^{\flat} of signature $(n-2, 1)$.

Let p be a prime split in \mathbf{k} , and let V^{\flat} be the hermitian space of signature $(n-2, 1)$ defined by (6.1.1). Using the first claim of Theorem 6.1.2 and the equality of Chern forms of Theorem 5.5.1, we find

$$\int_{\mathcal{Z}_V(p)(\mathbb{C})} \text{ch}(\widehat{\mathcal{L}}_V)^{n-2} = (p^{n-1} + 1) \cdot \text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_{V^{\flat}}).$$

On the other hand, we have the equality

$$\int_{\mathcal{Z}_V(p)(\mathbb{C})} \text{ch}(\widehat{\mathcal{L}}_V)^{n-2} = -B(p, 0, s_0) \cdot \text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_V)$$

from the proof of Proposition 8.2.2. Directly from the definition (1.1.5) and the explicit formulas of Corollaries 2.6.5 and 2.6.6, we find that

$$\frac{B(p, 0, s_0)}{p^{n-1} + 1} = -\frac{\mathbf{A}_{V^{\flat}}(0)}{\mathbf{A}_V(0)}.$$

Combining the three formulas in the previous paragraph, one obtains

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_{V^{\flat}}) = \frac{\mathbf{A}_{V^{\flat}}(0)}{\mathbf{A}_V(0)} \cdot \text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_V).$$

The theorem follows from this and our induction hypothesis

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_{V^{\flat}}) = \frac{|\text{CL}(\mathbf{k})|}{2^{o(D)-1} |\mathcal{O}_{\mathbf{k}}^{\times}|} \cdot \mathbf{A}_{V^{\flat}}(0). \quad \square$$

Remark 8.2.4. Theorem 8.2.3 is also true when $n = 1$. In this case

$$\mathcal{S}_V(\mathbb{C}) \subset \mathcal{M}_{1,0}(\mathbb{C}) \times \mathcal{M}_{0,1}(\mathbb{C})$$

is a complex orbifold of dimension 0 parametrizing certain pairs of elliptic curves with complex multiplication by $\mathcal{O}_{\mathbf{k}}$, the Chern form of $\widehat{\mathcal{L}}_V$ is (obviously) trivial, and the theory of complex multiplication implies

$$\mathrm{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_V) = \int_{\mathcal{S}_V(\mathbb{C})} 1 = \sum_{s \in \mathcal{S}_V(\mathbb{C})} \frac{1}{|\mathrm{Aut}(s)|} = \frac{|\mathrm{CL}(\mathbf{k})|^2}{2^{o(D)-1} \cdot |\mathcal{O}_{\mathbf{k}}^\times|^2}.$$

This agrees with

$$\frac{|\mathrm{CL}(\mathbf{k})|}{2^{o(D)-1} |\mathcal{O}_{\mathbf{k}}^\times|} \cdot \mathbf{A}_V(0) = \frac{|\mathrm{CL}(\mathbf{k})|}{2^{o(D)-1} |\mathcal{O}_{\mathbf{k}}^\times|} \cdot \mathbf{a}_1(0) = \frac{|\mathrm{CL}(\mathbf{k})|^2}{2^{o(D)-1} \cdot |\mathcal{O}_{\mathbf{k}}^\times|^2},$$

where the final equality is by Dirichlet's class number formula.

8.3. An alternate proof of the complex volume formula. We have invoked Remark 3.1.1 in the proofs of Proposition 8.2.2 and Theorem 8.2.3, but we have enough information at this point to give an independent proof of Theorem 8.2.3, and deduce much of Remark 3.1.1 and Proposition 8.2.2 as a consequence. This is not logically necessary in what follows, but the same method will be used in the proof of Theorem 8.4.2 below, and so we explain the argument here.

As before, the proof of Theorem 8.2.3 is by induction on $n \geq 2$, with the case $n = 2$ already known by Corollary 8.1.2. Thus we assume $n > 2$.

Suppose p is a prime split in \mathbf{k} , and let V^b denote the \mathbf{k} -hermitian space of signature $(n-2, 1)$ characterized by (6.1.1). It follows from Theorem 6.1.2 and the induction hypothesis that

$$\begin{aligned} \int_{\mathcal{Z}_V(p)(\mathbb{C})} \mathrm{ch}(\widehat{\mathcal{L}}_V)^{n-2} &= (p^{n-1} + 1) \mathrm{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_{V^b}) \\ &= \frac{|\mathrm{CL}(\mathbf{k})|}{2^{o(D)-1} |\mathcal{O}_{\mathbf{k}}^\times|} \cdot (p^{n-1} + 1) \cdot \mathbf{A}_{V^b}(0). \end{aligned}$$

Exactly as in the proof of Theorem 8.2.3, Corollaries 2.6.5 and 2.6.6 imply

$$-\frac{B(p, 0, s_0)}{p^{n-1} + 1} = \frac{\mathbf{A}_{V^b}(0)}{\mathbf{A}_V(0)},$$

and therefore

$$\int_{\mathcal{Z}_V(p)(\mathbb{C})} \mathrm{ch}(\widehat{\mathcal{L}}_V)^{n-2} = -\frac{|\mathrm{CL}(\mathbf{k})|}{2^{o(D)-1} |\mathcal{O}_{\mathbf{k}}^\times|} \cdot \mathbf{A}_V(0) \cdot B(p, 0, s_0).$$

Combining this with Proposition 8.2.1 shows that the p^{th} coefficient of the *cuspidal* modular form

$$(8.3.1) \quad \deg_{\mathbb{C}}(\widehat{\phi}_V^{\mathrm{cusp}}(\tau) \cdot \widehat{\mathcal{L}}_V^{n-2}) = \deg_{\mathbb{C}}(\widehat{\phi}_V(\tau) \cdot \widehat{\mathcal{L}}_V^{n-2}) - \deg_{\mathbb{C}}(\widehat{\phi}_V^{\mathrm{eis}}(\tau) \cdot \widehat{\mathcal{L}}_V^{n-2})$$

is

$$(8.3.2) \quad \left(\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_V) - \frac{|\text{CL}(\mathbf{k})|}{2^{o(D)-1} |\mathcal{O}_{\mathbf{k}}^{\times}|} \cdot \mathbf{A}_V(0) \right) \cdot B(p, 0, s_0).$$

The essential thing is that the quantity in parentheses in (8.3.2) is independent of the choice of split prime p , which we now allow to vary. The Hecke bound for coefficients of cusp forms implies that the absolute value of (8.3.2) is $O(p^{n/2})$ as $p \rightarrow \infty$. On the other hand, Corollary 2.6.8 shows that the absolute value of $B(p, 0, s_0)$ is greater than a constant times p^{n-1} . As $n > 2$, we conclude from this that the quantity in parentheses in (8.3.2) vanishes, completing our alternate proof of Theorem 8.2.3.

Moreover, having given a new proof that the p^{th} Fourier coefficient of (8.3.1) vanishes for all primes p split in \mathbf{k} , we deduce from Corollary A.2 that the m^{th} Fourier coefficient vanishes for all m with $\varepsilon(m) = 1$. Again invoking Proposition 8.2.1, we deduce that

$$\int_{\mathcal{Z}_V(m)(\mathbb{C})} \text{ch}(\widehat{\mathcal{L}}_V)^{n-2} = -\text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_V) B(m, 0, s_0)$$

for all such m . In this way, our alternate proof recovers much of Remark 3.1.1 and Proposition 8.2.2 in a new way.

8.4. The arithmetic volume formula. The arithmetic intersection and arithmetic degree of §4.4 provide us with a bilinear pairing

$$\widehat{\text{CH}}^1(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}}) \otimes \widehat{\text{CH}}^{n-1}(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}}) \rightarrow \mathbb{R},$$

sending $\widehat{\mathcal{Z}}_1 \otimes \widehat{\mathcal{Z}}_2 \mapsto \widehat{\text{deg}}(\widehat{\mathcal{Z}}_1 \cdot \widehat{\mathcal{Z}}_2)$. The hermitian line bundle

$$\widehat{\mathcal{K}}_V = 2\widehat{\mathcal{L}}_V - (\text{Exc}_V, 0) \in \widehat{\text{Pic}}(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}})$$

of Theorem 5.5.1 therefore determines a linear functional

$$\widehat{\text{CH}}^1(\bar{\mathcal{S}}_V, \mathcal{D}_{\text{pre}}) \rightarrow \mathbb{R},$$

called *arithmetic intersection against $\widehat{\mathcal{K}}_V^{n-1}$* , given by $\widehat{\mathcal{Z}} \mapsto \widehat{\text{deg}}(\widehat{\mathcal{Z}} \cdot \widehat{\mathcal{K}}_V^{n-1})$.

Assuming (as throughout §7) that $n > 2$, we have the modular form

$$(8.4.1) \quad \widehat{\text{deg}}(\widehat{\phi}_V(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) = \sum_{m \geq 0} \widehat{\text{deg}}(\widehat{\mathcal{Z}}^{\text{tot}}(m) \cdot \widehat{\mathcal{K}}_V^{n-1}) \cdot q^m$$

in $M_n(\Gamma_0(D), \varepsilon^n)$ obtained by applying this linear functional coefficient-by-coefficient to (7.2.1), and similarly for the other forms appearing in the decomposition

$$\widehat{\phi}_V(\tau) = \widehat{\phi}_V^{\text{eis}}(\tau) + \widehat{\phi}_V^{\text{exc}}(\tau) + \widehat{\phi}_V^{\text{cusp}}(\tau)$$

of (7.2.2). As $\widehat{\mathcal{K}}_V$ has trivial arithmetic intersection with every irreducible component of the exceptional divisor $\text{Exc}_V \subset \bar{\mathcal{S}}_V$ by Theorem 5.5.1, it has trivial arithmetic intersection with every coefficient of 7.2.4, and

$$(8.4.2) \quad \widehat{\text{deg}}(\widehat{\phi}_V^{\text{exc}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) = 0.$$

Motivated by Proposition 8.2.2, we make the following conjecture.

Conjecture 8.4.1. *Assuming $n > 2$, we have the equality of modular forms*

$$\widehat{\deg}(\widehat{\phi}_V(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) = \widehat{\deg}(\widehat{\phi}_V^{\text{eis}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}).$$

Equivalently, given (8.4.2),

$$\widehat{\deg}(\widehat{\phi}_V^{\text{cusp}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) = 0.$$

At long last we come to the central result of this work: the calculation of the arithmetic volume of $\widehat{\mathcal{K}}_V$ via induction. The calculation will simultaneously yield evidence for Conjecture 8.4.1, and also provide an explicit value for the mysterious rational numbers $a(p)$ of Theorem 6.1.2.

Theorem 8.4.2. *The hermitian line bundle $\widehat{\mathcal{K}}_V$ satisfies the following properties.*

- (1) *If $n \geq 2$, its arithmetic volume (in the sense of §4.5) is*

$$\widehat{\text{vol}}(\widehat{\mathcal{K}}_V) = \left(2 \frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \log(D) \right) \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V).$$

- (2) *If $n > 2$, then for any prime p split in \mathbf{k} the rational number $a(p)$ of Theorem 6.1.2 is given by*

$$a(p) = 2 \cdot \frac{p^{n-1} - 1}{p^{n-1} + 1} \cdot \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_{V^{\flat}}),$$

where V^{\flat} the hermitian space of signature $(n-2, 1)$ with local invariants determined by (6.1.1).

- (3) *If $n > 2$ then for any positive $m \in \mathbb{Z}$ with $\varepsilon(m) = 1$, the m^{th} Fourier coefficient of*

$$\widehat{\deg}(\widehat{\phi}_V^{\text{cusp}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) \in S_n(\Gamma_0(D), \varepsilon^n)$$

vanishes.

Proof. The proof is by induction on n . As we already know the first claim for $n = 2$ by Corollary 8.1.2, it suffices to assume that $n > 2$, and that the first claim holds all relevant (Definition 4.1.3) hermitian spaces V^{\flat} of signature $(n-2, 1)$.

We define $C(V)$ by the relation

$$\widehat{\text{vol}}(\widehat{\mathcal{K}}_V) = C(V) \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V),$$

and similarly for any V^{\flat} as above, so that our induction hypothesis is

$$(8.4.3) \quad C(V^{\flat}) = 2 \frac{\mathbf{A}'_{V^{\flat}}(0)}{\mathbf{A}_{V^{\flat}}(0)} + \log(D).$$

Recall from Proposition 2.7.3 the Eisenstein series

$$\sum_{r|D} \gamma_r E_r(\tau) = \sum_{m \geq 0} B(m, 0, s_0) q^m \in M_n(\Gamma_0(D), \varepsilon^n).$$

Recall also from Proposition 2.7.4 the rational number

$$\beta_\ell = (-1)^{n+1} \operatorname{inv}_\ell(V) \cdot \begin{cases} \left(\frac{-1}{\ell}\right)^{\frac{n}{2}} \ell^{\frac{n}{2}} & \text{if } n \text{ is even} \\ \left(\frac{-1}{\ell}\right)^{\frac{n-1}{2}} \ell^{\frac{n-1}{2}} & \text{if } n \text{ is odd} \end{cases}$$

for every prime $\ell \mid D$.

Lemma 8.4.3. *Let p be a prime split in \mathbf{k} , let V^\flat the hermitian space of signature $(n-2, 1)$ with local invariants determined by (6.1.1). If we abbreviate*

$$\begin{aligned} \kappa(p) &= 2 \frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \log(D) + \sum_{\ell \mid D} \frac{2 \log(\ell)}{1 + \beta_\ell \cdot \left(\frac{p}{\ell}\right)^n}, \\ \alpha(p) &= (p^{n-1} + 1)a(p) - 2(p^{n-1} - 1) \operatorname{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_{V^\flat}), \end{aligned}$$

then the p^{th} coefficient of the modular form

$$\widehat{\operatorname{deg}}(\widehat{\phi}_V(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) \in M_n(\Gamma_0(D), \varepsilon^n)$$

is equal to

$$-\frac{\kappa(p)}{2} \cdot B(p, 0, s_0) \cdot \operatorname{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) + \alpha(p) \cdot \log(p).$$

Proof. By Theorem 7.3.1, the p^{th} coefficient of (8.4.1) is

$$(8.4.4) \quad \widehat{\operatorname{deg}}(\widehat{\mathcal{Z}}_V^{\text{tot}}(p) \cdot \widehat{\mathcal{K}}_V^{n-1}) = \operatorname{ht}_{\widehat{\mathcal{K}}_V}(\widehat{\mathcal{Z}}_V(p)) + \int_{S_V(\mathbb{C})} \Phi_V(p) \operatorname{ch}(\widehat{\mathcal{K}}_V)^{n-1}.$$

Using Theorem 6.1.2, we rewrite the first term on the right hand side as

$$(8.4.5) \quad \begin{aligned} \operatorname{ht}_{\widehat{\mathcal{K}}_V}(\widehat{\mathcal{Z}}_V(p)) &= (p^{n-1} + 1) \left(\widehat{\operatorname{vol}}(\widehat{\mathcal{K}}_{V^\flat}) + a(p) \log(p) \right) \\ &= (p^{n-1} + 1) \left(C(V^\flat) \operatorname{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_{V^\flat}) + a(p) \log(p) \right). \end{aligned}$$

Now consider the second term on the right hand side of (8.4.4). By Remark 7.1.2 and the equality $2 \operatorname{ch}(\widehat{\mathcal{L}}_V) = \operatorname{ch}(\widehat{\mathcal{K}}_V)$ of Theorem 5.5.1, it is equal to

$$\int_{S_V(\mathbb{C})} \Phi_V(p) \operatorname{ch}(\widehat{\mathcal{K}}_V)^{n-1} = \operatorname{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) B'(p, 0, s_0).$$

Theorem 8.2.3 implies the equality

$$(8.4.6) \quad \operatorname{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) = 2 \cdot \frac{\mathbf{A}_V(0)}{\mathbf{A}_{V^\flat}(0)} \cdot \operatorname{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_{V^\flat}),$$

which we use to rewrite the integral above as

$$(8.4.7) \quad \int_{S_V(\mathbb{C})} \Phi_V(p) \operatorname{ch}(\widehat{\mathcal{K}}_V)^{n-1} = C_1 \cdot (p^{n-1} + 1) \cdot \operatorname{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_{V^\flat}),$$

in which we have set

$$C_1 = 2 \cdot \frac{B'(p, 0, s_0)}{p^{n-1} + 1} \cdot \frac{\mathbf{A}_V(0)}{\mathbf{A}_{V^\flat}(0)}.$$

Plugging (8.4.5) and (8.4.7) into (8.4.4), we obtain

$$\widehat{\deg}(\widehat{\mathcal{Z}}_V^{\text{tot}}(p) \cdot \widehat{\mathcal{K}}_V^{n-1}) = (p^{n-1} + 1)(C_1 + C(V^b)) \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_{V^b}) \\ + (p^{n-1} + 1)a(p) \log(p).$$

By direct calculation, Corollaries 2.6.5 and 2.6.6 (along with the definition (1.1.5) of $\mathbf{A}_V(s)$ and $\mathbf{A}_{V^b}(s)$) give the explicit formulas

$$(8.4.8) \quad \frac{B(p, 0, s_0)}{p^{n-1} + 1} = -\frac{\mathbf{A}_{V^b}(0)}{\mathbf{A}_V(0)}$$

and

$$\frac{B'(p, 0, s_0)}{B(p, 0, s_0)} = \frac{\mathbf{A}'_{V^b}(0)}{\mathbf{A}_{V^b}(0)} - \frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \frac{p^{n-1} - 1}{p^{n-1} + 1} \log(p) - \sum_{\ell|D} \frac{\log(\ell)}{1 + \beta_{\ell} \cdot \left(\frac{p}{\ell}\right)^n}.$$

Using these equalities we see that the constant C_1 can be rewritten as

$$C_1 = -2 \frac{\mathbf{A}'_{V^b}(0)}{\mathbf{A}_{V^b}(0)} + 2 \frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} - 2 \frac{p^{n-1} - 1}{p^{n-1} + 1} \log(p) + \sum_{\ell|D} \frac{2 \log(\ell)}{1 + \beta_{\ell} \cdot \left(\frac{p}{\ell}\right)^n}.$$

Combining this with the induction hypothesis (8.4.3) shows that

$$C_1 + C(V^b) = \kappa(p) - 2 \frac{p^{n-1} - 1}{p^{n-1} + 1} \log(p),$$

and we obtain

$$\widehat{\deg}(\widehat{\mathcal{Z}}_V^{\text{tot}}(p) \cdot \widehat{\mathcal{K}}_V^{n-1}) = (p^{n-1} + 1)\kappa(p) \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_{V^b}) + \alpha(p) \log(p).$$

To complete the proof of the lemma, substitute the equality

$$\text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_{V^b}) \stackrel{(8.4.6)}{=} \frac{\mathbf{A}_{V^b}(0)}{2\mathbf{A}_V(0)} \cdot \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \stackrel{(8.4.8)}{=} -\frac{B(p, 0, s_0)}{2(p^{n-1} + 1)} \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V)$$

into the first term on the right hand side. \square

Lemma 8.4.4. *We have the equality of modular forms*

$$(8.4.9) \quad \widehat{\deg}(\widehat{\phi}_V^{\text{eis}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) = -\frac{1}{2} \widehat{\text{vol}}(\widehat{\mathcal{K}}_V) \sum_{r|D} \gamma_r E_r(\tau) \\ - \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \sum_{r|D} \log(D/r) \gamma_r E_r(\tau).$$

Moreover, for positive $m \in \mathbb{Z}$ with $\gcd(m, D) = 1$ the m^{th} coefficient of this modular form is

$$-\frac{1}{2} \left(\widehat{\text{vol}}(\widehat{\mathcal{K}}_V) + \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \sum_{\ell|D} \frac{2 \log(\ell)}{1 + \beta_{\ell} \cdot \left(\frac{m}{\ell}\right)^n} \right) B(m, 0, s_0).$$

Proof. Directly from the definition (7.2.3) of $\widehat{\phi}_V^{\text{eis}}(\tau)$ we find

$$\begin{aligned} \widehat{\deg}(\widehat{\phi}_V^{\text{eis}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) &= -\widehat{\deg}(\widehat{\mathcal{L}}_V \cdot \widehat{\mathcal{K}}_V^{n-1}) \sum_{r|D} \gamma_r E_r(\tau) \\ &\quad - \sum_{r|D} \widehat{\deg}((\text{Exc}_V, \log(D/r)) \cdot \widehat{\mathcal{K}}_V^{n-1}) \gamma_r E_r(\tau). \end{aligned}$$

For the first term on the right hand side, we use Theorem 5.5.1 to see that

$$\widehat{\deg}(\widehat{\mathcal{L}}_V \cdot \widehat{\mathcal{K}}_V^{n-1}) = 2^{-1} \cdot \widehat{\deg}(\widehat{\mathcal{K}}_V \cdot \widehat{\mathcal{K}}_V^{n-1}) = 2^{-1} \cdot \widehat{\text{vol}}(\widehat{\mathcal{K}}_V).$$

For the second term on the right hand side, it follows from Theorem 5.5.1 that $(\text{Exc}_V, 0) \cdot \widehat{\mathcal{K}}_V^{n-1} = 0$. As in the proof of Lemma 4.5.4, we have the easy arithmetic intersection formula

$$(0, \log(D/r)) \cdot \widehat{\mathcal{K}}_V^{n-1} = (0, \log(D/r) \text{ch}(\widehat{\mathcal{K}}_V^{n-1})) \in \widehat{\text{CH}}^n(\widehat{\mathcal{S}}_V, \mathcal{D}_{\text{pre}}),$$

and hence, directly from the definition (4.4.4) of arithmetic degree,

$$\begin{aligned} \widehat{\deg}((0, \log(D/r)) \cdot \widehat{\mathcal{K}}_V^{n-1}) &= \log(D/r) \int_{\mathcal{S}_V(\mathbb{C})} \text{ch}(\widehat{\mathcal{K}}_V)^{n-1} \\ &= \log(D/r) \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V). \end{aligned}$$

Thus

$$\widehat{\deg}((\text{Exc}_V, \log(D/r)) \cdot \widehat{\mathcal{K}}_V^{n-1}) = \log(D/r) \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V).$$

The equality (8.4.9) follows immediately from these formulas.

For the second claim, using Proposition 2.7.3 and Proposition 2.7.4, the m^{th} coefficient of the modular form on the right hand side of (8.4.9) is

$$\begin{aligned} &-\frac{1}{2} \widehat{\text{vol}}(\widehat{\mathcal{K}}_V) \sum_{r|D} \gamma_r e_r(m) - \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \sum_{r|D} \gamma_r e_r(m) \log(D/r) \\ &= -\frac{1}{2} \widehat{\text{vol}}(\widehat{\mathcal{K}}_V) B(m, 0, s_0) - \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \left(\sum_{\ell|D} \frac{\log(\ell)}{1 + \beta_{\ell} \cdot \left(\frac{m}{\ell}\right)^n} \right) B(m, 0, s_0), \end{aligned}$$

as desired. \square

The proof of Theorem 8.4.2 now proceeds by examining the Fourier coefficients of the modular form

$$(8.4.10) \quad \widehat{\deg}(\widehat{\phi}_V^{\text{cusp}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) = \widehat{\deg}(\widehat{\phi}_V(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) - \widehat{\deg}(\widehat{\phi}_V^{\text{eis}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}).$$

If we abbreviate

$$c = \frac{\widehat{\text{vol}}(\widehat{\mathcal{K}}_V)}{2} - \left(\frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \frac{\log(D)}{2} \right) \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V),$$

then comparing Lemma 8.4.3 and Lemma 8.4.4 shows that the p^{th} coefficient of (8.4.10) is

$$cB(p, 0, s_0) + \alpha(p) \log(p)$$

for all primes p split in \mathbf{k} . In particular, the modular form

$$\widehat{\deg}(\widehat{\phi}_V^{\text{cusp}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) - c \sum_{m \geq 0} B(m, 0, s_0)$$

has the property that its p^{th} coefficient is $\alpha(p) \log(p) \in \mathbb{Q} \log(p)$ for all such primes. It follows from Corollary A.4 that $\alpha(p) = 0$, proving the second claim of Theorem 8.4.2.

We have now seen the p^{th} coefficient of the *cuspidal* modular form (8.4.10) is $cB(p, 0, s_0)$ for all primes p split in \mathbf{k} , and may argue as in §8.3. The Hecke bound for coefficients of cusp forms implies that the absolute value of (8.3.2) is $O(p^{n/2})$ as $p \rightarrow \infty$, while Corollary 2.6.8 shows that the absolute value of $B(p, 0, s_0)$ is greater than a constant times p^{n-1} . We conclude that $c = 0$, proving the first claim of Theorem 8.4.2.

Finally, having now shown that the p^{th} coefficient of (8.4.10) vanishes for all primes p split in \mathbf{k} , Corollary A.2 shows that the m^{th} coefficient vanishes whenever $\varepsilon(m) = 1$. This proves the third claim of Theorem 8.4.2. \square

8.5. Consequences of Theorem 8.4.2. We now collect many corollaries of Theorem 8.4.2. The first of these is explicit formulas for the arithmetic volumes of the other (perhaps more natural) hermitian line bundles on \mathcal{S}_V defined in §5.2.

Corollary 8.5.1. *Assuming $n \geq 2$, we have the arithmetic volume formulas*

$$\widehat{\text{vol}}(\widehat{\omega}_{A/S_V}^{\text{Hdg}}) = \left(2 \frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} - nC_0(n) + \log(D) \right) \text{vol}_{\mathbb{C}}(\widehat{\omega}_{A/S_V}^{\text{Hdg}})$$

and

$$\widehat{\text{vol}}(\widehat{\mathcal{L}}_V) = \left(\frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \frac{\log(D)}{2} \right) \text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_V) + \frac{(-1)^{n-1}}{2} \sum_{E \subset \text{Exc}_V} \widehat{m}_E.$$

In the second equality the sum is over all irreducible components $E \subset \text{Exc}_V$ of the exceptional divisor of Definition 5.1.4 and, if $p \mid D$ denotes the prime at which E is supported, $\widehat{m}_E = m_E \log(p)$ is the constant of §5.6.

Proof. The first equality follows from Theorem 8.4.2, using Proposition 6.7.1 and the equality of Chern forms of Theorem 5.5.1.

For the second equality, combining 5.5.1 with the intersection formula

$$\widehat{\mathcal{K}}_V \cdot (\text{Exc}_V, 0) = 0$$

of Theorem 5.5.1 shows that

$$\begin{aligned} 2^n \cdot \widehat{\text{vol}}(\widehat{\mathcal{L}}_V) &= \widehat{\text{vol}}(\widehat{\mathcal{K}}_V + (\text{Exc}_V, 0)) \\ &= \widehat{\text{vol}}(\widehat{\mathcal{K}}_V) + \widehat{\text{vol}}(\text{Exc}_V, 0). \end{aligned}$$

Combining this with Theorem 8.4.2 and the equality of Chern forms of Theorem 5.5.1 yields

$$\widehat{\text{vol}}(\widehat{\mathcal{L}}_V) = \left(\frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \frac{\log(D)}{2} \right) \text{vol}_{\mathbb{C}}(\widehat{\mathcal{L}}_V) + \frac{\widehat{\text{vol}}(\text{Exc}_V, 0)}{2^n}.$$

Now use Corollary 5.6.2. \square

The most interesting aspect of the expected equality

$$\sum_{m \geq 0} \widehat{\deg}(\widehat{\mathcal{Z}}^{\text{tot}}(m) \cdot \widehat{\mathcal{K}}_V^{n-1}) \cdot q^m \stackrel{?}{=} \widehat{\deg}(\widehat{\phi}_V^{\text{eis}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1})$$

of Conjecture 8.4.1 is that the coefficients of the modular form on the right hand side can be made completely explicit, by combining the formulas of §2.7 and Theorem 8.2.3 with the following corollary of Theorem 8.4.2.

Corollary 8.5.2. *Assuming $n > 2$, we have the equality of modular forms*

$$\begin{aligned} & \widehat{\deg}(\widehat{\phi}_V^{\text{eis}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) \\ &= -\text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \sum_{r|D} \left(\frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \frac{3}{2} \log(D) - \log(r) \right) \gamma_r E_r(\tau). \end{aligned}$$

Proof. Start with the equality

$$\begin{aligned} \widehat{\deg}(\widehat{\phi}_V^{\text{eis}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) &= -\frac{1}{2} \widehat{\text{vol}}(\widehat{\mathcal{K}}_V) \sum_{r|D} \gamma_r E_r(\tau) \\ &\quad - \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \sum_{r|D} \log(D/r) \gamma_r E_r(\tau) \end{aligned}$$

of Lemma 8.4.4, and substitute the arithmetic volume formula of Theorem 8.4.2 into the first term on the right hand side. \square

Assume now that $n > 2$. In light of Corollary 8.5.2, Conjecture 8.4.1 is equivalent to

$$(8.5.1) \quad \widehat{\deg}(\widehat{\mathcal{Z}}^{\text{tot}}(m) \cdot \widehat{\mathcal{K}}_V^{n-1}) \stackrel{?}{=} -\text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \sum_{r|D} \left(\frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \frac{3}{2} \log(D) - \log(r) \right) \gamma_r e_r(m),$$

for all positive $m \in \mathbb{Z}$. Of course the right hand side can be made explicit using the formulas of §2.7.

For example, let us now suppose $\gcd(m, D) = 1$, and abbreviate

$$\kappa(m) = 2 \frac{\mathbf{A}'_V(0)}{\mathbf{A}_V(0)} + \log(D) + \sum_{\ell|D} \frac{2 \log(\ell)}{1 + \beta_{\ell} \cdot \left(\frac{m}{\ell}\right)^n},$$

where

$$\beta_{\ell} = (-1)^{n+1} \text{inv}_{\ell}(V) \cdot \begin{cases} \left(\frac{-1}{\ell}\right)^{\frac{n}{2}} \ell^{\frac{n}{2}} & \text{if } n \text{ is even} \\ \left(\frac{-1}{\ell}\right)^{\frac{n-1}{2}} \ell^{\frac{n-1}{2}} & \text{if } n \text{ is odd.} \end{cases}$$

We can use Propositions 2.7.3 and 2.7.4 to rewrite (8.5.1) as

$$\widehat{\deg}(\widehat{\mathcal{Z}}^{\text{tot}}(m) \cdot \widehat{\mathcal{K}}_V^{n-1}) \stackrel{?}{=} -\frac{\kappa(m)}{2} \cdot \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \cdot B(m, 0, s_0).$$

As in the proof of Proposition 8.2.2, Remark 3.1.1 implies

$$(8.5.2) \quad \int_{\mathcal{Z}_V(m)(\mathbb{C})} \text{ch}(\widehat{\mathcal{K}}_V)^{n-2} = -\frac{1}{2} \cdot \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \cdot B(m, 0, s_0)$$

(the factor of $1/2$ appears because of the relation $2 \text{ch}(\widehat{\mathcal{L}}_V) = \text{ch}(\widehat{\mathcal{K}}_V)$ of Theorem 5.5.1), and we find that (8.5.1) is equivalent to

$$(8.5.3) \quad \widehat{\text{deg}}(\widehat{\mathcal{Z}}^{\text{tot}}(m) \cdot \widehat{\mathcal{K}}_V^{n-1}) \stackrel{?}{=} \kappa(m) \int_{\mathcal{Z}_V(m)(\mathbb{C})} \text{ch}(\widehat{\mathcal{K}}_V)^{n-2}.$$

The following corollary of Theorem 8.4.2 shows that this equality holds for many values of m .

Corollary 8.5.3. *Assume $n > 2$. The equality (8.5.3) holds for all positive $m \in \mathbb{Z}$ with $\varepsilon(m) = 1$.*

Proof. The final claim of Theorem 8.4.2 shows that the m^{th} Fourier coefficients of

$$\widehat{\text{deg}}(\widehat{\phi}_V(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1}) \quad \text{and} \quad \widehat{\text{deg}}(\widehat{\phi}_V^{\text{eis}}(\tau) \cdot \widehat{\mathcal{K}}_V^{n-1})$$

agree for all m with $\varepsilon(m) = 1$, and hence (by Corollary 8.5.2) the equality (8.5.1) holds for all such m . The discussion above shows that this is equivalent to (8.5.3). \square

Again suppose that $n > 2$ and $\gcd(m, D) = 1$, and recall the equality

$$\widehat{\text{deg}}(\widehat{\mathcal{Z}}_V^{\text{tot}}(m) \cdot \widehat{\mathcal{K}}_V^{n-1}) = \text{ht}_{\widehat{\mathcal{K}}_V}(\bar{\mathcal{Z}}_V(m)) + \int_{S_V(\mathbb{C})} \Phi_V(m) \text{ch}(\widehat{\mathcal{K}}_V)^{n-1}$$

of Theorem 7.3.1. As in the proof of Lemma 8.4.3, the integral on the right hand side is

$$\begin{aligned} \int_{S_V(\mathbb{C})} \Phi_V(m) \text{ch}(\widehat{\mathcal{K}}_V)^{n-1} &= B'(m, 0, s_0) \cdot \text{vol}_{\mathbb{C}}(\widehat{\mathcal{K}}_V) \\ &\stackrel{(8.5.2)}{=} -2 \frac{B'(m, 0, s_0)}{B(m, 0, s_0)} \int_{\mathcal{Z}_V(m)(\mathbb{C})} \text{ch}(\widehat{\mathcal{K}}_V)^{n-2}. \end{aligned}$$

Solving for the height term, and find that the conjectural equality of (8.5.3) is equivalent to

$$\text{ht}_{\widehat{\mathcal{K}}_V}(\bar{\mathcal{Z}}_V(m)) \stackrel{?}{=} \left(\kappa(m) + 2 \frac{B'(m, 0, s_0)}{B(m, 0, s_0)} \right) \int_{\mathcal{Z}_V(m)(\mathbb{C})} \text{ch}(\widehat{\mathcal{K}}_V)^{n-2}.$$

By the explicit calculation of $B(m, 0, s)$ found in Corollaries 2.6.5 and 2.6.6, this last conjectural equality is equivalent to

$$(8.5.4) \quad \text{ht}_{\widehat{\mathcal{K}}_V}(\bar{\mathcal{Z}}_V(m)) \stackrel{?}{=} \left(2 \frac{\mathbf{A}'_{V,m}(0)}{\mathbf{A}_{V,m}(0)} + \log(D) \right) \int_{\mathcal{Z}_V(m)(\mathbb{C})} \text{ch}(\widehat{\mathcal{K}}_V)^{n-2}.$$

Here if n is even we have set

$$\mathbf{A}_{V,m}(s) = \mathbf{a}_1(s) \cdots \mathbf{a}_{n-1}(s) \cdot m^{s+n-1} \sigma_{-2s-n+1}(m),$$

while if n is odd we have set

$$\begin{aligned} \mathbf{A}_{V,m}(s) &= \mathbf{a}_1(s) \cdots \mathbf{a}_{n-1}(s) \cdot m^{s+n-1} \sigma_{-2s-n+1,\varepsilon}(m) \\ &\quad \times \prod_{\ell|D} \left(1 + \left(\frac{-1}{\ell} \right)^{\frac{n-1}{2}} \left(\frac{m}{\ell} \right) \text{inv}_\ell(V) \ell^{-s-\frac{n-1}{2}} \right). \end{aligned}$$

Corollary 8.5.4. *Assume $n > 2$. The equality (8.5.4) holds for all positive $m \in \mathbb{Z}$ with $\varepsilon(m) = 1$.*

Proof. The discussion above shows that (8.5.4) is equivalent to (8.5.3), which is known when $\varepsilon(m) = 1$ by Corollary 8.5.3. \square

8.6. Volume of the Hodge bundle. We now derive Theorem A of the introduction as a consequence of Theorem 8.4.2. Let W be a \mathbf{k} -hermitian space of signature $(n-1, 1)$ with $n \geq 1$, containing a self-dual $\mathcal{O}_{\mathbf{k}}$ -lattice. Let \mathcal{M}_W be the moduli space of (4.1.1), and recall that in §5.2 we associated a metrized Hodge bundle

$$\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}} \in \widehat{\text{Pic}}(\overline{\mathcal{M}}_W, \mathcal{D}_{\text{pre}})$$

to the universal abelian scheme $A \rightarrow \mathcal{M}_W$.

Theorem 8.6.1. *The metrized Hodge bundle has complex volume*

$$\text{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}) = \mathbf{A}_W(0) \times \begin{cases} 2^{n-1} & \text{if } n \text{ is odd} \\ 2^{n-o(D)} & \text{if } n \text{ is even} \end{cases}$$

and arithmetic volume

$$\widehat{\text{vol}}(\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}) = \left(2 \frac{\mathbf{A}'_W(0)}{\mathbf{A}_W(0)} - nC_0(n) + \log(D) \right) \text{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}),$$

where $C_0(n)$ is the constant defined in Theorem A.

Proof. First assume $n \geq 2$. Let $W_0 = \mathbf{k}$ endowed with its hermitian form $h(x, y) = x\overline{y}$, and define a hermitian form on $V = \text{Hom}_{\mathbf{k}}(W_0, W)$ as in (5.1.2). In particular $W \cong V$ as hermitian spaces.

As in Remark 5.1.2, there is a finite étale surjection $\mathcal{S}_V \rightarrow \mathcal{M}_W$ of degree

$$t_n = \frac{|\text{CL}(\mathbf{k})|}{|\mathcal{O}_{\mathbf{k}}^\times|} \times \begin{cases} 1 & \text{if } n \text{ is even} \\ 2^{1-o(D)} & \text{if } n \text{ is odd.} \end{cases}$$

Denote again by $A \rightarrow \mathcal{S}_V$ the pullback of $A \rightarrow \mathcal{M}_W$ via this morphism, so that

$$\text{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}) = \frac{1}{t_n} \text{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{S}_V}^{\text{Hdg}})$$

and

$$\widehat{\text{vol}}(\widehat{\omega}_{A/\mathcal{M}_W}^{\text{Hdg}}) = \frac{1}{t_n} \widehat{\text{vol}}(\widehat{\omega}_{A/\mathcal{S}_V}^{\text{Hdg}}).$$

Using these formulas, the theorem for $n > 2$ follows from Theorem 8.2.3 and Corollary 8.5.1, along with the equality of Chern forms of Theorem 5.5.1.

It remains to treat the case $n = 1$, so that

$$\mathcal{M}_W \cong \mathcal{M}_{(0,1)} \cong \mathcal{M}_{(1,0)}.$$

In this case $\mathbf{A}_W(s) = \mathbf{a}_1(s)$. Dirichlet's class number formula implies

$$\mathbf{a}_1(0) = \frac{|\mathrm{CL}(\mathbf{k})|}{|\mathcal{O}_{\mathbf{k}}^\times|},$$

while (8.1.3) and (5.3.1) imply

$$2 \frac{\mathbf{a}'_1(0)}{\mathbf{a}_1(0)} - C_0(1) + \log(D) = -\frac{L'(0, \varepsilon)}{L(0, \varepsilon)} - \frac{\log(D)}{2} = \log(2\pi) + 2h_{\mathbf{k}}^{\mathrm{Falt}}.$$

The theory of complex multiplication implies

$$\mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{M}_{(1,0)}}^{\mathrm{Hdg}}) = \int_{\mathcal{M}_{(1,0)}(\mathbb{C})} 1 = \sum_{x \in \mathcal{M}_{(1,0)}(\mathbb{C})} \frac{1}{|\mathrm{Aut}(x)|} = \frac{|\mathrm{CL}(\mathbf{k})|}{|\mathcal{O}_{\mathbf{k}}^\times|} = \mathbf{A}_W(0),$$

while the argument of Proposition 5.3.1, which is essentially the Chowla-Selberg formula, implies the first equality in

$$\begin{aligned} \widehat{\mathrm{deg}}(\widehat{\omega}_{A/\mathcal{M}_{(1,0)}}^{\mathrm{Hdg}}) &= \widehat{\mathrm{deg}}(0, \log(2\pi) + 2h_{\mathbf{k}}^{\mathrm{Falt}}) \\ &= (\log(2\pi) + 2h_{\mathbf{k}}^{\mathrm{Falt}}) \int_{\mathcal{M}_{(1,0)}(\mathbb{C})} 1 \\ &= \left(2 \frac{\mathbf{A}'_W(0)}{\mathbf{A}_W(0)} - C_0(1) + \log(D) \right) \mathrm{vol}_{\mathbb{C}}(\widehat{\omega}_{A/\mathcal{M}_{(1,0)}}^{\mathrm{Hdg}}). \end{aligned}$$

This completes the proof. \square

APPENDIX A. COEFFICIENTS OF MODULAR FORMS

In this appendix we quote a theorem of Bellaïche, and derive some consequences that play an essential role in the proof of Theorem 8.4.2. We thank Joel Bellaïche for his help. Let N be a positive integer and let χ be a Dirichlet character modulo N . Fix a modular form

$$f(\tau) = \sum_{m \geq 0} c(m) \cdot q^m \in M_k(\Gamma_0(N), \chi)$$

of weight $k \in \mathbb{Z}_{>0}$ with nebentypus character χ (we do not assume that f is a Hecke eigenform).

Theorem A.1 (Bellaïche). *Suppose the coefficients of $f(\tau)$ satisfy*

- (i) $c(m) = 0$ for all integers m with $\mathrm{gcd}(m, N) > 1$, and
- (ii) $c(p) = 0$ for a set of primes p of density 1.

Then $f(\tau)$ vanishes identically.

Proof. If the nebentypus character χ is trivial, this is a direct consequence of [Be20, Theorem I], which actually gives an analogous result in positive characteristic. An inspection of the proof shows that the result equally holds for general nebentypus.

For the convenience of the reader, we also sketch a more direct proof that was provided to us by Bellaïche. Let L be the field generated by the coefficients of f . Without loss of generality, possibly replacing L by an extension, we may view L as an extension of \mathbb{Q}_p for any prime p . We may also assume that L contains all N -th roots of unity. We first consider the case when $N > 1$. Choose a prime p that divides N .

Let M_f be the subspace of $M_k(\Gamma_0(N), \chi, L)$ generated by f under the action of the Hecke operators T_ℓ for ℓ prime not dividing N . Let A_f be the L -subalgebra of $\text{End}_L(M_f)$ generated by the T_ℓ for ℓ prime not dividing N . Notice that A_f contains all the T_n for n prime to N . This is because any such T_n is a polynomial in the T_ℓ (ℓ prime not dividing N) and the diamond operators, which are just scalar in L , and hence belong to A_f . (Here we use the hypothesis that f has a nebentypus).

Let $G = G_{\mathbb{Q}, N}$ be the Galois group of the maximal algebraic extension of \mathbb{Q} unramified outside N . There exists a continuous pseudo-representation $(\tau, \delta) : G \rightarrow A_f$ of dimension 2 which sends Frob_ℓ (for ℓ prime not dividing N) to T_ℓ , see e.g. [Be20, Section 10] or [BS15, Section 3.10]. Here δ is $w^{k-1}\chi$, where w denotes the p -adic cyclotomic character, and hence takes values in L^\times .

We claim that the linear subspace generated by $\tau(G)$ in A_f has to be A_f . This follows from the formula $\tau(gh) + \delta(h)\tau(gh^{-1}) = \tau(g)\tau(h)$, a formula true for all pseudo-representations of dimension 2, since it is true for $\tau =$ the trace, and $\delta =$ the determinant in the algebra $\text{Mat}_2(\mathbb{C})$. Thus the linear subspace generated by $\tau(G)$ is a sub-algebra containing all the T_ℓ (ℓ not dividing N), hence is A_f .

Let V be the kernel of the linear map $A_f \rightarrow L$ given by $T \mapsto c_T f(1)$. By the hypothesis (ii) we know that $c_f(\ell) = c_{T_\ell f}(1) = 0$ for a dense set of primes ℓ . Hence $\tau(\text{Frob}_\ell)$ is in V for a dense set of ℓ . Since V is closed, this means by the Chebotarev density theorem that $\tau(G)$ is contained in V . By the above claim, we find that $V = A_f$. This implies that $T_n \in V$ for all n prime to N , and hence that $c_f(n) = 0$ for all n prime to N . On the other hand, by hypothesis (i) we know that $c_f(n) = 0$ for all n not prime to N . Hence we have $f = 0$, concluding the proof in the case $N > 1$.

If $N = 1$, we chose an auxiliary prime q and replace f by

$$f' = \sum_{\substack{n \geq 0 \\ (n, q) = 1}} c(n)q^n,$$

which belongs to $M_k(\Gamma_0(q^2), L)$ by [Miy89, Lemma 4.3.10]. Applying the above argument to f' we find $c(n) = 0$ for all n coprime to q . But since f has level 1, this means that $f = 0$. \square

Corollary A.2. *Let ψ be a quadratic Dirichlet character modulo N . Fix $\delta \in \{\pm 1\}$, and suppose the coefficients of $f(\tau)$ satisfy $c(p) = 0$ for all but finitely many primes p with $\psi(p) = \delta$. Then $c(m) = 0$ for all integers m with $\psi(m) = \delta$.*

Proof. Recall from [Miy89, Lemma 4.3.10] that if ψ is any quadratic Dirichlet character modulo N , then the twist

$$f_\psi(\tau) = \sum_{m \geq 0} \psi(m)c(m) \cdot q^m$$

of $f(\tau)$ defines an element of $M_k(\Gamma_0(N^2), \chi)$. As the Fourier coefficients of

$$\sum_{\substack{m \geq 0 \\ \psi(m) = \delta}} c(m) \cdot q^m = \frac{1}{2} \cdot (f_{\psi^2}(\tau) + \delta f_\psi(\tau)) \in M_k(\Gamma_0(N^2), \chi)$$

satisfy the hypotheses of Theorem A.1, this modular form vanishes identically. The claim follows. \square

We now apply these results to the space of modular forms $M_n(\Gamma_0(D), \varepsilon^n)$ which is relevant for the present paper. Let us write

$$M \subset M_n(\Gamma_0(D), \varepsilon^n)$$

for the \mathbb{Q} -subspace of forms with rational Fourier coefficients. The \mathbb{Q} -linear dual $M^\vee = \text{Hom}_{\mathbb{Q}}(M, \mathbb{Q})$ is generated by the coefficient extraction functionals

$$a_m : M \rightarrow \mathbb{Q},$$

sending a modular form $f(\tau)$ to its m^{th} Fourier coefficient.

Corollary A.3. *For $\delta \in \{\pm 1\}$ and any positive $t \in \mathbb{Z}$, the subspaces*

$$\begin{aligned} A_\delta &= \text{Span}_{\mathbb{Q}}\{a_m : m \in \mathbb{Z}_{>0} \text{ with } \varepsilon(m) = \delta\} \\ P_{\delta,t} &= \text{Span}_{\mathbb{Q}}\{a_p : p \text{ prime with } \varepsilon(p) = \delta \text{ and } p > t\} \end{aligned}$$

of M^\vee are equal.

Proof. For any subspace $B \subset M^\vee$ we write $B^\perp \subset M$ for its orthogonal complement with respect to the natural non-degenerate pairing $M^\vee \times M \rightarrow \mathbb{Q}$. It suffices to show that the inclusion $A_\delta^\perp \subset P_{\delta,t}^\perp$ is an equality, so fix an

$$f(\tau) = \sum_{m \geq 0} c(m) \cdot q^m \in P_{\delta,t}^\perp \subset M.$$

By definition of $P_{\delta,t}^\perp$, we have $c(p) = 0$ for all primes $p > t$ with $\varepsilon(p) = \delta$, and so Corollary A.2 implies that $c(m) = 0$ for all $m \in \mathbb{Z}_+$ with $\varepsilon(m) = \delta$. In other words, $f(\tau) \in A_\delta^\perp$. \square

Corollary A.4. *Fix $\delta \in \{\pm 1\}$. If*

$$f(\tau) = \sum_{m \geq 0} c(m) \cdot q^m \in M_n(\Gamma_0(D), \varepsilon^n)$$

is a modular form with the property that

$$c(p) \in \mathbb{Q} \log(p)$$

for all primes p with $\varepsilon(p) = \delta$, then $c(p) = 0$ for all such primes.

Proof. Fix a prime p with $\varepsilon(p) = \delta$, and consider the coefficient extraction functional $a_p : M \rightarrow \mathbb{Q}$. Taking $t = p$ in Corollary A.3, we find that there are rational numbers $\alpha_1, \dots, \alpha_r$ and primes $p < p_1 < \dots < p_r$ with all $\varepsilon(p_i) = \delta$, such that

$$a_p = \sum_{i=1}^r \alpha_i \cdot a_{p_i}$$

holds as an equality of \mathbb{Q} -linear functionals on M . We now view this as an equality of \mathbb{C} -linear functionals on $M_n(\Gamma_0(D), \varepsilon^n) = M \otimes_{\mathbb{Q}} \mathbb{C}$, and apply this linear functional to $f(\tau)$. The result is

$$c(p) = \sum_{i=1}^r \alpha_i \cdot c(p_i) \in \sum_{i=1}^r \mathbb{Q} \log(p_i),$$

and the claim follows from the \mathbb{Q} -linear independence of the set

$$\{\log(p), \log(p_1), \dots, \log(p_r)\}. \quad \square$$

Remark A.5. Of course Corollary A.2 allows us to strengthen the conclusion of Corollary A.4 to $c(m) = 0$ whenever $\varepsilon(m) = \delta$.

REFERENCES

- [AbSt84] M. Abramowitz and I. Stegun. *Pocketbook of Mathematical Functions*. Verlag Harri Deutsch, Thun (1984).
- [Be20] Joël Bellaïche. Image of pseudo-representations and coefficients of modular forms modulo p . *Advances in Math.* **353** (2020), 647–721.
- [BS15] Joël Bellaïche and Kannan Soundararajan. The number of nonzero coefficients of modular forms (mod p). *Algebra Number Theory* **9** (2015), 1825–1856.
- [Bo98] Richard Borcherds. Automorphic forms with singularities on Grassmannians. *Invent. Math.* **132** (1998), 491–562.
- [Br02] Jan H. Bruinier. Borcherds products on $O(2, l)$ and Chern classes of Heegner divisors. Springer Lecture Notes in Mathematics **1780**, Springer-Verlag (2002).
- [BBK07] Jan H. Bruinier, José I. Burgos Gil, and Ulf Kühn. Borcherds products and arithmetic intersection theory on Hilbert modular surfaces. *Duke Math. J.*, **139** No. 1, (2007), 1–88.
- [BF04] Jan H. Bruinier and Jens Funke. On two geometric theta lifts. *Duke Math. J.* **125** (2004), 45–90.
- [BHY15] Jan H. Bruinier, Benjamin Howard, and Tonghai Yang. Heights of Kudla-Rapoport divisors and derivatives of L-functions. *Invent. Math.* **201** (2015), 1–95.
- [BrKu03] Jan H. Bruinier and Ulf Kühn. Integrals of automorphic Green’s functions associated to Heegner divisors. *Int. Math. Res. Not.* **31** (2003), 1687–1729.
- [BY09] Jan H. Bruinier and Tonghai Yang. Faltings heights of CM cycles and derivatives of L-functions. *Invent. Math.* **177** (2009), 631–681.
- [Bu97] José I. Burgos Gil. Arithmetic Chow rings and Deligne-Beilinson cohomology. *J. Algebr. Geom.*, **6** (1997), 335–377.
- [BKK07] José I. Burgos Gil, J. Kramer, and Ulf Kühn. Cohomological arithmetic Chow rings. *J. Inst. Math. Jussieu*, **6**(1):1–172, 2007.
- [BKK05] José I. Burgos Gil, J. Kramer, and Ulf Kühn. Arithmetic characteristic classes of automorphic vector bundles *Doc. Math.*, **10** (2005), 619–716.

- [BHK⁺a] Jan H. Bruinier, Benjamin Howard, Steven S. Kudla, Michael Rapoport, and Tonghai Yang. Modularity of generating series of divisors on unitary Shimura varieties. *Asterisque*, **421** (2020), 8–125.
- [BHK⁺b] Jan H. Bruinier, Benjamin Howard, Steven S. Kudla, Michael Rapoport, and Tonghai Yang. Modularity of generating series of divisors on unitary Shimura varieties II: arithmetic applications. *Asterisque*, **421** (2020), 127–186.
- [FC90] Gerd Faltings and Ching-Li Chai. *Degeneration of abelian varieties*, volume 22 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, (1990). With an appendix by David Mumford.
- [Fr09] Gerard Freixas i Montplet. Heights and metrics with logarithmic singularities. *Journal für die Reine und Angewandte Mathematik*, **627** (2009), 97–153.
- [GS90] Henri Gillet and Christophe Soulé. Arithmetic intersection theory. *Inst. Hautes Études Sci. Publ. Math.*, **72** (1991) 93–174. 1990.
- [Gro78] Benedict Gross. On the periods of abelian integrals and a formula of Chowla and Selberg. *Invent. Math.* **45** (1978), 193–211.
- [Gro97] Benedict Gross. On the motive of a reductive group. *Invent. Math.* **130** (1997), 287–313.
- [HKS96] Michael Harris, Steven S. Kudla and William J. Sweet. Theta dichotomy for unitary groups. *J. Amer. Math. Soc.* **9** (1996), 941–1004.
- [HLSY22] Qiao He, Chao Li, Yousheng Shi, and Tonghai Yang A proof of the Kudla-Rapoport conjecture for Krämer models *Preprint* arXiv:2208.07988 (2022).
- [Hor14] Fritz Hörmann, *The geometric and arithmetic volume of Shimura varieties of orthogonal type*, CRM Monograph Series, vol. 35, American Mathematical Society, Providence, RI, (2014).
- [How15] Benjamin Howard. Complex multiplication cycles and Kudla-Rapoport divisors II. *Amer. J. Math.*, 137(3):639–698, 2015.
- [How20] Benjamin Howard. Arithmetic volumes of unitary Shimura curves. To appear in *Theta Series: Representation Theory, Geometry, and Arithmetic*, Fields Institute Communications Series. *Preprint* arXiv:2010.07362.
- [Ich07] Atsushi Ichino. On the Siegel-Weil formula for unitary groups, *Math. Z.* **255** (2007), 721–729.
- [Jac62] Ronald Jacobowitz. Hermitian forms over local fields. *Amer. J. Math.*, **84** (1962), 441–465.
- [JvP] Barbara Jung and Anna-Maria Von Pippich. The arithmetic volume of the moduli space of abelian surfaces. arXiv:2205.11864.
- [Ku97] Stephen S. Kudla. Central derivatives of Eisenstein series and height pairings, *Ann. of Math* **146** (1997), 545–646.
- [Ku03] Stephen S. Kudla. Integrals of Borchers forms. *Compos. Math.* **137** (2003), 293–349.
- [KR14] Stephen S. Kudla and Michael Rapoport. Special cycles on unitary Shimura varieties II: Global theory. *J. Reine Angew. Math.*, **697** (2014), 91–157.
- [KRY99] Stephen S. Kudla, Michael Rapoport, and Tonghai Yang. On the derivative of an Eisenstein series of weight one. *Internat. Math. Res. Notices*, (7):347–385, 1999.
- [KRY06] Stephen S. Kudla, Michael Rapoport, and Tonghai Yang. *Modular forms and special cycles on Shimura curves*, volume 161 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ (2006).
- [KY10] Steven S. Kudla and Tonghai Yang. Eisenstein series for SL_2 , *Science China Mathematics*, a special volume in honor of Yuan Wang’s 80th birthday. **53** (2010), 2275–2316.
- [Kuhn01] Ulf Kühn. Generalized arithmetic intersection numbers. *J. Reine Angew. Math.*, **534** (2001), 209–236.

- [Kra03] Nicole Krämer. Local models for ramified unitary groups. *Abh. Math. Sem. Univ. Hamburg*, 73:67–80, 2003.
- [Lan13] Kai-Wen Lan. *Arithmetic compactifications of PEL-type Shimura varieties*, volume 36 of *London Mathematical Society Monographs Series*. Princeton University Press, Princeton, NJ, (2013).
- [Lan16] Kai-Wen Lan. Compactifications of PEL-type Shimura varieties in ramified characteristics. *Forum Math. Sigma*, 4 (2016), Paper No. e1, 98.
- [LZ19] Chao Li and Wei Zhang. Kudla-Rapoport cycles and derivatives of local densities. *newblock J. Amer. Math. Soc.*, 35 (2022), no. 3, 705–797.
- [MP19] Keerthi Madapusi Pera. Toroidal compactifications of integral models of Shimura varieties of Hodge type. *Ann. Sci. Éc. Norm. Supér.*, 52 No. 2 (2019), 93–514.
- [Miy89] Toshitune Miyake. *Modular forms*. Translated from the Japanese by Yoshitaka Maeda. Springer-Verlag, Berlin, (1989).
- [MB85] Laurent Moret-Bailly. Pinceaux de variétés abéliennes. *Astérisque*, 129 (1985).
- [Pap00] Georgios Pappas. On the arithmetic moduli schemes of PEL Shimura varieties. *J. Algebraic Geom.*, 9 No. 3 (2000), 577–605.
- [Pol03] Alexander Polishchuk. *Abelian varieties, theta functions and the Fourier transform* volume 153 of *Cambridge tracts in mathematics*. Cambridge University Press, Cambridge (2003).
- [Sch09] Nils R. Scheithauer. The Weil representation of $SL_2(\mathbb{Z})$ and some applications *IMRN* 8 (2009), 1488–1545.
- [SABK] C. Soulé, D. Abramovich, J.-F. Burnol, and J. Kramer. *Lectures on Arakelov Geometry*. Cambridge Studies in Advanced Mathematics 33, Cambridge University Press, Cambridge (1992).
- [SGA70] *Schémas en groupes. I: Propriétés générales des schémas en groupes*. Séminaire de Géométrie Algébrique du Bois Marie 1962/64 (SGA 3). Dirigé par M. Demazure et A. Grothendieck. Lecture Notes in Mathematics, Vol. 151. Springer-Verlag, Berlin-New York, 1970.
- [Sta18] The Stacks Project Authors. *Stacks Project*. <https://stacks.math.columbia.edu>, 2018.
- [Weil65] A. Weil. Sur la formule de Siegel dans la théorie des groupes classiques. *Acta Math.* 113 (1965), 1–87.
- [Yuan22] X. Yuan. Modular Heights of Quaternionic Shimura Curves. arXiv:2205.13995.

FACHBEREICH MATHEMATIK, TECHNISCHE UNIVERSITÄT DARMSTADT, SCHLOSSGARTEN-STRASSE 7, D-64289 DARMSTADT, GERMANY

Email address: `bruinier@mathematik.tu-darmstadt.de`

DEPARTMENT OF MATHEMATICS, BOSTON COLLEGE, 140 COMMONWEALTH AVE, CHESTNUT HILL, MA 02467, USA

Email address: `howardbe@bc.edu`