To the memories of Jacob Feldman and Gian-Carlo Rota A spectral theory for combinatorial dynamics

James Propp

ABSTRACT. This article proposes a framework for the study of periodic maps T from a (typically finite) set X to itself when the set X is equipped with one or more real- or complex-valued functions. The main idea, inspired by the time-evolution operator construction from ergodic theory, is the introduction of a vector space that contains the given functions and is closed under composition with T, along with a time-evolution operator on that vector space. I show that the invariant functions and 0-mesic functions span complementary subspaces associated respectively with the eigenvalue 1 and the other eigenvalues. Alongside other examples, I give an explicit description of the spectrum of the evolution operator when X is the set of k-element multisets with elements in $\{0, 1, \ldots, n-1\}$, T increments each element of a multiset by 1 mod n, and $g_i : X \to \mathbb{R}$ (with $1 \leq i \leq k$) maps a multiset to its *i*th smallest element.

1. INTRODUCTION

Recent work in dynamical algebraic combinatorics has paid a great deal of attention to *homomesies:* numerical functions on a dynamical system with the property that the average of the function over an orbit doesn't depend on which orbit one takes. Missing from this work is attention to *invariants:* quantities that are constant on orbits. It would be conceptually helpful to bring homomesies and invariants into a uniform framework. The article [PR] that introduced the concept of homomesy provided this framework in the special case of linear actions (see section 2.4 of that article), but most actions of interest to combinatorialists are nonlinear (piecewise linear, birational, or purely combinatorial). Here I introduce a vector space that may be the proper setting for a unified treatment of homomesies and invariants; a periodic action induces a diagonalizable linear map on the vector space, and the spectrum of the map carries dynamical information about homomesies and invariants for the action and its powers.

2. LINEARIZATION

Given a set X, an invertible map T from X to itself satisfying $T^n = \text{Id}_X$ (that is, $T^n(x) = x$ for all $x \in X$) with $n \ge 1$, and a collection of functions g_1, \ldots, g_k from X to \mathbb{C} , let V be the linear span of all functions of the form $g_i \circ T^j$ with $1 \le i \le k$, $0 \le j < n$. I will sometimes refer to the g_i 's as *statistics* (or sometimes as the "original" statistics as opposed to the time-shifted statistics $g_i \circ T^j$). V is a finite-dimensional space of dimension at most kn.

This paper has been prepared using ALCO author class on May 26, 2021.

KEYWORDS. combinatorics, dynamics, invariant, homomesy, spectrum, ergodic.

The role played by the g_i 's is crucial, much as the choice of an algebra of measurable sets is crucial in ergodic theory. When we enlarge our set of initial statistics we potentially enlarge V, and when we restrict our set of initial statistics we potentially reduce V. As an extreme case, if our collection of statistics is empty, V is 0-dimensional regardless of the dynamics of the map T.

For all $f \in V$ define $Uf \in V$ by (Uf)(x) = f(T(x)). We will call the action of Uon V the *linearization* of the original action of T on X. It is easy to check that Usends V to itself (e.g., U sends $g_i \circ T^j$ to $g_i \circ T^{j+1}$, which in the case j = n - 1 is g_i); that U is linear; that U^n is the identity map on V; and that U has inverse U^{n-1} . V is the *dynamical span* of g_1, \ldots, g_k in the sense that it is the smallest vector space that contains g_1, \ldots, g_k and is closed under the action of U. We will also call V the *dynamical closure* of the linear span of g_1, \ldots, g_k . The function $g_i \circ T^j$ can also be written as $U^j g_i$.

Let us say that a function $f \in V$ is homomesic if $\frac{1}{n} \sum_{j=0}^{n-1} U^j f$ is a constant function, and more specifically *c*-mesic if $\frac{1}{n} \sum_{j=0}^{n-1} U^j f = c$ for all $x \in X$, that is, if $\frac{1}{n} \sum_{j=0}^{n-1} f(T^j x) = c$ for all $x \in X$. Let us say that a function $f \in V$ is invariant if Uf = f, that is, if f(Tx) = f(x) for all $x \in X$.

When X is finite (as it will be except in sections 5 and 6) we create an |X|-by-kn matrix M whose rows correspond to elements $x \in X$ (the order in which the elements of X are listed is unimportant) and whose columns from left to right correspond to the respective functions $g_1, \ldots, g_k, Ug_1, \ldots, Ug_k, \ldots, U^{n-1}g_1, \ldots, U^{n-1}g_k$, where the entry in the x row and the U^jg_i column is $(U^jg_i)(x) = g_i(T^jx)$; call M the presenting matrix of U, and note that V can be identified with the span of the columns of M. In particular, dim V is the rank of M.

Since $U^n = I$ (the identity map from V to itself), U is a diagonalizable operator on V whose eigenvalues are *n*th roots of unity; there exists a basis for V whose elements are eigenvectors for U. (Typically our statistics are real-valued, but if we want to look at the eigenspaces associated with eigenvalues other than 1 and -1 we need to treat V as a vector space over \mathbb{C} .) Let V_1 be the span of the 1-eigenvectors (i.e., the nonzero elements $f \in V$ satisfying Uf = 1f = f) and let V_1^{\perp} be the span of the other (non-unital) eigenvectors, so that $V = V_1 \oplus V_1^{\perp}$ and dim $V = \dim V_1 + \dim V_1^{\perp}$. (Note that despite the notation no inner product is involved; that is, V_1^{\perp} is a complement but not an orthocomplement.)

PROPOSITION 2.1. V_1 is the space of invariants and V_1^{\perp} is the space of 0-mesies.

Proof. If $f \in V_1$ then Uf = 1f = f and vice versa; hence V_1 is the space of invariants. If $f \in V_1^{\perp}$ so that f is in the span of the non-unital eigenvectors then $f + Uf + U^2f + \cdots + U^{n-1}f = 0$ (since in the case where f is a ζ -eigenvector with $\zeta \neq 1$ we have $f + Uf + U^2f + \cdots + U^{n-1}f = (1 + \zeta + \zeta^2 + \cdots + \zeta^{n-1})f = 0f$). Conversely, given f satisfying $f + Uf + U^2f + \cdots + U^{n-1}f = 0$, if we write $f = f_1 + f_2$ with $f_1 \in V_1$ and $f_2 \in V_1^{\perp}$ and we apply $I + U + U^2 + \cdots + U^{n-1}$ to $f_1 + f_2$, we get nf_1 , which can only vanish if f_1 does, implying $f \in V_1^{\perp}$.

It follows that the multiplicity of 1 as an eigenvalue of U gives the number of linearly independent invariants in V while the sum of the multiplicities of the nonunital eigenvalues of U gives the number of linearly independent 0-mesies in V. It is easy to show that every homomesic function f can be written uniquely as the sum of a constant function and a 0-mesic function, specifically the constant function $g = (1/n)(f + Uf + U^2f + ... + U^{n-1}f)$ and the 0-mesic function h = f - g.

In view of the complementary nature of V_1 and V_1^{\perp} , it might be appropriate to refer to the elements of V_1^{\perp} as "coinvariants" rather than 0-mesies, but this word

is already in use with a different meaning. Another term for elements of V_1^{\perp} that seems apt is "survariants", which has no existing meaning and in some ways seems preferable to "0-mesies"; however, the terms "homomesy" and "homomesic" seem to have been adopted to the point where a change in nomenclature might be confusing.

It is possible for V_1 to be trivial; for instance, if $X = \{1, -1\}$ with T(x) = -x of order n = 2 equipped with the identity statistic $g_1(x) = x$, then V is 1-dimensional with dim $V_1 = 0$ and dim $V_1^{\perp} = 1$. However, for all examples considered in this paper V_1 will contain the constant functions and hence have dimension ≥ 1 . As long as there is at least one homomesic function f that is not a 0-mesic, the nonzero constant function $\sum_{j=0}^{n-1} U^j f$ is in V, so that the constant functions form a 1-dimensional subspace of V_1 .

I proceed to give alternative characterizations of V_1 and V_1^{\perp} .

PROPOSITION 2.2. $f \in V_1$ if and only if there exists $g \in V$ with $f = g + Ug + U^2g + \cdots + U^{n-1}g$.

Proof. If $f = g + Ug + U^2g + \dots + U^{n-1}g$ then $Uf = Ug + U^2g + U^3g + \dots + U^ng = f$ (since $U^ng = g$), so $f \in V_1$. Conversely, suppose f is in V_1 . Then putting g = (1/n)fwe have $f = g + Ug + U^2g + \dots + U^{n-1}g$.

Consequently, the space of invariants is spanned by the sums $f_i := g_i + Ug_i + \cdots + U^{n-1}g_i$; call these the *spanning invariants*. Since these k functions span V_1 , dim $V_1 \leq k$. If we divide the presenting matrix into n blocks of width k and sum those blocks, we obtain a matrix M_1 whose column span is V_1 .

I next generalize Proposition 2.2 to eigenfunctions, exploiting the discrete Fourier transform. Given $\zeta \in \mathbb{C}$ with $\zeta^n = 1$, let V_{ζ} be the space of $f \in V$ with $Uf = \zeta f$ (or equivalently $\overline{\zeta}Uf = f$), so that V_1^{\perp} is the direct sum of the spaces V_{ζ} with $\zeta \neq 1$.

PROPOSITION 2.3. For $\zeta^n = 1$, $f \in V_{\zeta}$ if and only if there exists $g \in V$ with $f = g + \overline{\zeta} Ug + \cdots + \overline{\zeta}^{n-1} U^{n-1}g$.

Proof. The proof of Proposition 2.2 applies to any linear operator U' on the \mathbb{C} -vector space V that satisfies $(U')^n = I$, and in particular applies to $U' = \overline{\zeta}U$.

Consequently, the ζ -eigenspace is spanned by the functions $g_i + \bar{\zeta} U g_i + \cdots + \bar{\zeta}^{n-1} U^{n-1} g_i$; let us call these the *spanning* ζ -eigenfunctions. We have dim $V_{\zeta} \leq k$. If we take the blocks discussed following the proof of Proposition 2.2 and sum them with respective coefficients $1, \bar{\zeta}, \bar{\zeta}^2, \ldots$, we obtain a matrix M_{ζ} whose column span is V_{ζ} .

I provide an alternate characterization of the elements of $V_1^{\perp} = \bigoplus_{\zeta^n=1, \zeta \neq 1} V_{\zeta}$:

PROPOSITION 2.4. $f \in V_1^{\perp}$ if and only if there exists $g \in V$ with f = g - Ug.

(Note that we could just as well have used Ug - g as g. A function of the form Ug - g is called a coboundary in dynamical system theory; it measures the increase in g from one moment to the next.)

Proof. If f = g - Ug, then $f + Uf + \dots + U^{n-1}f = (g - Ug) + (Ug - U^2g) + \dots + (U^{n-1}g - U^ng) = g - U^ng = 0$ (since $U^ng = g$), so f is 0-mesic and belongs to V_1^{\perp} .

Algebraic Combinatorics, draft (May 26, 2021)

Conversely, suppose f is in V_1^{\perp} . Let $h = 1f + 2Uf + 3U^2f + \dots + nU^{n-1}f$. Then

$$\begin{split} h - Uh &= (1U^0 f + 2U^1 f + 3U^2 f + \dots + (n-1)U^{n-2} f + nU^{n-1} f) \\ &- (1U^1 f + 2U^2 f + 3U^3 f + \dots + (n-1)U^{n-1} f + nU^n f) \\ &= (U^0 f + U^1 f + U^2 f + \dots + U^{n-1} f) - nU^n f \\ &= 0 - nf \quad (\text{because } f \text{ is } 0\text{-mesic and } U^n f = f) \\ &= -nf, \end{split}$$

so putting g = -(1/n)h we have f = g - Ug.

If we are interested in counting all nonunital eigenvalues (with multiplicity), we can just take the rank of the matrix M - M' where M is the presenting matrix and M' is obtained from M by cyclically shifting the columns k positions to the right.

It should be stressed that in the framework being proposed here, 0-mesies can be seen either as coboundaries of arbitrary functions or as combinations of eigenfunctions of the time-evolution operator U. In particular, real 0-mesies can occur as combinations of complex eigenfunctions.

It may be helpful to note that the 0-mesies form the kernel of $I+U+U^2+\cdots+U^{n-1}$, which Proposition 2.4 identifies as the image of I-U; likewise, the invariants form the kernel of I-U, which Proposition 2.2 identifies as the image of $I+U+U^2+\cdots+U^{n-1}$; and more generally the ζ -eigenfunctions form the kernel of $I-\bar{\zeta}U$, which Proposition 2.3 identifies as the image of $I + \bar{\zeta}U + \bar{\zeta}^2U^2 + \cdots + \bar{\zeta}^{n-1}U^{n-1}$. In each case we have a short exact sequence of vector spaces. Two function g_1, g_2 give rise to the same element of V_1 via Proposition 2.2 iff they differ by an element of V_1^{\perp} , and two function g_1, g_2 give rise to the same element of V_1^{\perp} via Proposition 2.4 iff they differ by an element of V_1 .

Let us define $V^{(0)}$ as the intersection of V_1^{\perp} with the span of g_1, \ldots, g_k . There is a sense in which $V^{(0)}$ determines V_1^{\perp} , namely, the linear combination $\sum_{i,j} a_{i,j} U^j g_i$ is in V_1^{\perp} if and only if the linear combination $\sum_{i,j} a_{i,j} g_i$ (in which $U^j g_i$ is replaced by g_i) is in V_1^{\perp} . That is because every function of the form $U^j g - g$ is a coboundary (note that $U^j g - g = U(g + Ug + \cdots + U^{j-1}g) - (g + Ug + \cdots + U^{j-1}g)$).

The sum of the multiplicities of the non-unital eigenvalues (that is, the eigenvalues unequal to 1) gives us the dimension of the space of 0-mesies. The dynamical significance of the specific multiplicities of individual non-unital eigenvalues is subtler. For every d dividing n, the sum of the multiplicities of the eigenvalues ζ satisfying $\zeta^d = 1$ is the dimension of the space of invariants of T^d , while the sum of the multiplicities of those ζ with $\zeta^d \neq 1$ is the dimension of the space of 0-mesies of T^d . Additional meaning of the multiplicities appears if, in the spirit of classical invariant theory, one extends V to a ring R, introducing statistics that are products of the g_i 's. Invariant functions in the ring R can arise from noninvariant eigenfunctions in V associated with complex eigenvalues whose product is 1; see the paragraph following the proof of Proposition 4.1. However, it should be noted that the ring R need not be graded by polynomial degree; e.g., in the example treated in Proposition 4.1, each g_i takes values in $\{0, 1\}$ and hence satisfies $g_i^2 = g_i$. Also note that when the set X is finite, the ring R (being a set of functions with domain X), viewed as a vector space, must be finite-dimensional.

3. Example: Rowmotion in a Chain

Fix $n, k \ge 2$ and let $X_{n,k}$ be the set of k-element multisets with elements belonging to $\{0, 1, \ldots, n-1\}$. Let us denote a generic element of $X_{n,k}$ by $x = (x_1, x_2, \ldots, x_k)$ with $x_1 \le x_2 \le \cdots \le x_k$. We can represent an element of

 $X_{n,k}$ by writing down its elements with appropriate multiplicities, in weakly increasing order, with parentheses and intervening commas omitted. For instance, $X_{3,3} = \{000, 001, 002, 011, 012, 022, 111, 112, 122, 222\}$. I will use superscripts to indicate repetition, e.g., I write $X_{n,k} = \{0^k, 0^{k-1}1, \ldots, (n-1)^k\}$. It is well-known that $|X_{n,k}| = \binom{n+k-1}{k}$.

Given $x = (x_1, x_2, \ldots, x_k)$, put $x_0 = 0$ and $x_{k+1} = n-1$, and for $1 \leq i \leq k$ define the reflections $\rho_i : X_{n,k} \to X_{n,k}$ by $\rho_i(x) = x'$ where $x'_i = x_{i-1} + x_{i+1} - x_i$ and $x'_j = x_j$ for all $j \neq i$. The ρ_i 's satisfy the Coxeter relations $\rho_i^2 = 1$, $(\rho_i \rho_{i+1})^3 = 1$, and $(\rho_i \rho_j)^2 = 1$ for |i-j| > 1, and thus form a representation of the Coxeter group A_k .

Any product of all the ρ_i 's, each taken one at a time, is a Coxeter element γ satisfying $\gamma^{k+1} = 1$. For convenience, we take $\gamma_{n,k} = \rho_k \circ \cdots \circ \rho_1$ which updates entries from left to right. For instance, $\gamma_{3,3}$ sends 000 to 002 to 022 to 222 to 000 and sends 001 to 012 to 122 to 111 to 001 and sends 011 to 112 to 011. We have $\gamma_{n,k}(x) = x'$ where $x'_i = x'_{i-1} + x_{i+1} - x_i$; that is, the new value in location *i* equals the new value in location i - 1 plus the old value in location i + 1 minus the old value in location *i*. The operation γ can be seen as a special case of piecwise-linear rowmotion on the order polytope of a product of two chains in the case where one of the chains is of length 1. For more on piecewise-linear rowmotion, see [EP2] and section 5 of this article.

For $x = (x_1, x_2, \ldots, x_k)$ and $0 \leq i \leq k+1$ let $g_i(x) = x_i$, so that for instance $g_1(001) = 0, g_2(001) = 0$, and $g_3(001) = 1$. The maps $g_i : X \to \mathbb{R}$ $(1 \leq i \leq k)$ and $T : X \to X$ are all linear with respect to x_0, \ldots, x_{k+1} so the dynamical span V of g_1, \ldots, g_k is easy to determine; it is spanned by g_1, \ldots, g_k along with the constant functions. Define U as above. The relation $x'_i = x'_{i-1} + x_{i+1} - x_i$ gives $Ug_i = Ug_{i-1} + g_{i+1} - g_i$. If we apply U^j to the preceding relation and sum as j goes from 0 to k, we find that the orbit-averages of g_{i-1}, g_i , and g_{i+1} are in arithmetic progression. Since the orbit-averages of g_0 and g_{k+1} are 0 and n-1 respectively, we see that g_i is c_i -mesic with $c_i = i\frac{n-1}{k+1}$. The k functions $g_i - c_i$ $(1 \leq i \leq k)$ are the 0-mesies of this action while the constant functions are the only invariants; together they span the full (k+1)-dimensional vector space dynamically spanned by the g_i 's. The action of U on this space is a simple rotation whose spectrum assigns multiplicity 1 to each k + 1st root of 1. Concretely, U acts as a cyclic rotation on the functions $g_1 - g_0, g_2 - g_1, \ldots, g_{k+1} - g_k$.

4. Example: Multiset Rotation

Fix $n, k \ge 2$, and define $X_{n,k}$ as in section 3.

Let us define the rotation operator $T_{n,k}: X_{n,k} \to X_{n,k}$ that increments each element of x by 1 mod n, sending i to i + 1 for i < n - 1 and sending n - 1 to 0; for instance, $T_{3,3}(001) = 112$ and $T_{3,3}(112) = 022$ (note that the 2 in 112 has become a 0 and has moved to the left). It is obvious that $T_{n,k}^n$ is the identity on $X_{n,k}$. In contexts where it is safe to do so without confusion, I will omit subscripts on X and T. Let $g_i(x)$ denote the *i*th smallest element of x as before, and let $U_{n,k}$ denote the time-evolution operator for the action $f \mapsto f \circ T_{n,k}$ on the dynamical closure of the linear span of g_1, \ldots, g_k .

It has been known for several years, as part of the unwritten lore of dynamical algebraic combinatorics, that $g_i + g_j$ takes average value n - 1 on each orbit of T whenever i + j = k + 1 (including the case i = j = (k + 1)/2 when k is odd). When we pass to the dynamical span we find the invariants that were "missing" from earlier treatments of this example.

The case n = 2 behaves differently than the general case, so we give it separate consideration. Here the possible eigenvalues are 1 and -1.

PROPOSITION 4.1. The multiplicities of the eigenvalues 1 and -1 in the spectrum of $U_{2,k}$ are $\lceil (k+1)/2 \rceil$ and $\lfloor (k+1)/2 \rfloor$ respectively.

Proof. X has k + 1 elements; if k is odd, X consists of (k + 1)/2 orbits of size 2, and if k is even, X consists of k/2 orbits of size 2 and the fixed point $0^{k/2}1^{k/2}$. Here I show the presenting matrix for k = 3 whose columns correspond to the six functions $g_1, g_2, g_3, Ug_1, Ug_2$, and Ug_3 , and the presenting matrix for k = 4 whose columns correspond to the eight functions $g_1, g_2, g_3, g_4, Ug_1, Ug_2, Ug_3$, and Ug_4 :

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 \end{pmatrix}$$
$$\begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}$$

It is easy to see that for general k the k+1 rows of the presenting matrix are linearly independent, so that the matrix has rank k+1, which implies that the dynamical span of the original vector space (the column-span of the matrix) is (k+1)-dimensional. If we add the left half of the (k+1)-by-2k presenting matrix to the right half, we get a matrix whose columns correspond to the spanning invariants $g_1 + Ug_1, g_2 + Ug_2, \ldots$ and therefore span the space of invariants. In the cases k = 3 and k = 4, these matrices are $\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$

and

$$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix} + \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 \\ 0 & 1 & 2 \\ 1 & 1 & 1 \end{pmatrix}$$
$$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix} + \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 2 \\ 0 & 0 & 2 & 2 \\ 0 & 1 & 1 & 2 \\ 1 & 1 & 1 & 1 \end{pmatrix}$$

For general k, the first $\lceil (k+1)/2 \rceil$ rows of the matrix are linearly independent and the remaining rows repeat earlier rows, so the rank is $\lceil (k+1)/2 \rceil$, so dim $V_1 = \lceil (k+1)/2 \rceil$ as claimed. To find dim $V_1^{\perp} = \dim V_{-1}$, we subtract the two half-matrices instead of adding, obtaining

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} - \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} -1 & -1 & -1 \\ 0 & -1 & 0 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}$$
$$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix} - \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} -1 & -1 & -1 & -1 \\ 0 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{pmatrix}$$

and

in the cases k = 3 and k = 4. The first $\lfloor (k+1)/2 \rfloor$ rows of the matrix are linearly independent and each of the remaining rows is either the zero vector or the negative of an earlier row, so the rank is $\lfloor (k+1)/2 \rfloor$, so dim $V_{-1} = \lfloor (k+1)/2 \rfloor$ as claimed. \Box

Following up on my earlier remark about invariant theory at the end of section 2, let us pause to consider the quadratic invariant $(g_1 + ... + g_k)(Ug_1 + ... + Ug_k)$, equal to the number of 1's in x times the number of 1's in Tx (or, equivalently, the number of 0's in x times the number of 1's in x). If for all i we put $s_i = (g_i + Ug_i)/2$ (an eigenfunction for the eigenvalue 1) and $a_i = (g_i - Ug_i)/2$ (an eigenfunction for the eigenvalue -1) so that $g_i = s_i + a_i$ and $Ug_i = s_i - a_i$, then putting $S = s_1 + ... + s_k$ and $A = a_1 + ... + a_k$ we see that the quadratic invariant equals $(S + A)(S - A) = S^2 - A^2$. This representation shows us explicitly how the quadratic invariant arises from all the possible pairs of eigenfunction s_i, s_j with the eigenvalue 1 and all the possible pairs of eigenfunction a_i, a_j with the eigenvalue -1.

We now consider n > 2.

PROPOSITION 4.2. For n > 2, the multiplicity of the eigenvalue ζ in the spectrum of $U_{n,k}$ is $\lfloor k/2 \rfloor + 1$ when $\zeta = 1$ and k when $\zeta \neq 1$.

Proof. I prove the second claim first. Given $\zeta \neq 1$ with $\zeta^n = 1$, define $h_i = g_i + \bar{\zeta} U g_i + \bar{\zeta}^2 U^2 g_i + \cdots + \bar{\zeta}^{n-1} U^{n-1} g_i$ for $1 \leq i \leq k$ (the spanning ζ -eigenfunctions; see Proposition 2.3). Since these functions span V_{ζ} , dim $V_{\zeta} \leq k$. To prove the reverse inequality and thereby prove equality it suffices to find a k-element subset of X, call it $Y = \{y^{(1)}, \ldots, y^{(k)}\}$, so that the vectors $(g_i(y^{(1)}), \ldots, g_i(y^{(k)}))^t$ (with $1 \leq i \leq k$) are linearly independent; that is, we need the k-by-k matrix M whose i, jth entry is $g_i(y^{(j)})$ to be nonsingular. Let $y^{(j)} = 0^{k-j+1}1^{j-1}$. Then M has only four distinct entries:

$$W = 0 + 1\zeta + 2\zeta^{2} + \dots + (n-2)\zeta^{n-2} + 0\zeta^{n-1}$$

$$N = 0 + 1\zeta + 2\zeta^{2} + \dots + (n-2)\zeta^{n-2} + (n-1)\zeta^{n-1}$$

$$S = 1 + 2\zeta + 3\zeta^{2} + \dots + (n-1)\zeta^{n-2} + 0\zeta^{n-1}$$

$$E = 1 + 2\zeta + 3\zeta^{2} + \dots + (n-1)\zeta^{n-2} + (n-1)\zeta^{n-1}$$

(For later use, note that

$$(1-\zeta)(S-N) = (1-\zeta)(1+\zeta+\zeta^2+\dots+\zeta^{n-2}-(n-1)\zeta^{n-1})$$

= 1-\zeta^{n-1}-(n-1)\zeta^{n-1}+(n-1)\zeta^n
= n(1-\zeta^{n-1})

which is nonvanishing for all n > 1 and that

$$(1-\zeta)(E-W) = (1-\zeta)(1+\zeta+\zeta^2+\dots+\zeta^{n-2}+(n-1)\zeta^{n-1})$$

= 1-\zeta^{n-1}+(n-1)\zeta^{n-1}-(n-1)\zeta^n
= (n-2)(\zeta^{n-1}-1)

which vanishes for n = 2 and is nonvanishing for n > 2.)

When k is odd, M has the form illustrated below for k = 3 and k = 5.

$$\begin{pmatrix} N & N & N \\ W & N & E \\ W & S & E \end{pmatrix} \qquad \begin{pmatrix} N & N & N & N & N \\ W & N & N & N & E \\ W & W & N & E & E \\ W & W & S & E & E \\ W & S & S & S & E \end{pmatrix}$$

Algebraic Combinatorics, draft (May 26, 2021)

Note that each matrix is split by diagonal and antidiagonal lines into four zones, each of which has all entries equal. Similarly, when k is even, M has the form illustrated below for k = 4 and k = 6.

$$\begin{pmatrix} N & N & N & N \\ W & N & N & E \\ W & W & E & E \\ W & S & S & E \end{pmatrix} \qquad \begin{pmatrix} N & N & N & N & N \\ W & N & N & N & E \\ W & W & N & N & E \\ W & W & N & N & E \\ W & W & W & E & E \\ W & W & S & S & E \\ W & S & S & S & S & E \end{pmatrix}$$

Let $D_k(N, E, S, W)$ denote the determinant of the k-by-k matrix of the above form, where for now N, E, S, W are to be treated as formal indeterminates. There are many ways to evaluate these determinants, but my favorite is the following argument, communicated to me by Joe Buhler. If we subtract W/E times the last column from the first column, so that the only nonvanishing entry in the first column is N - (E/W)N = N(E - W)/E, we obtain the recurrence

$$D_k(N, E, S, W) = (N(E - W)/E) D_{k-1}(E, S, W, N)$$

(where the cyclic rotation of the arguments corresponds to 90 degree rotation of the submatrix). With this recurrence and the initial condition $D_2(N, E, S, W) = N(E - W)$ it is easy to prove the general formula

$$D_{2i+r}(N, E, S, W) = \begin{cases} (-1)^i N(N-S)^{i-1} (W-E)^i & \text{if } r = 0, \\ (-1)^i N(N-S)^i (W-E)^i & \text{if } r = 1 \end{cases}$$

by induction. When $\zeta \neq 1$ we have $N - S \neq 0$ and $W - E \neq 0$ as noted earlier so $D_k(N, E, S, W) \neq 0$, proving that the matrix is nonsingular as claimed. (This is the part of the proof that assumes n > 2; we have W = E when n = 2.)

It remains to consider $\zeta = 1$. In this case we have N = S so the matrix is singular. Indeed, the last $\lfloor (k-1)/2 \rfloor$ rows of M coincide with earlier rows, showing that the corank is at least $\lfloor (k-1)/2 \rfloor$, or equivalently that the rank is at most $\lceil (k+1)/2 \rceil = \lfloor k/2 \rfloor + 1$. But it is easy to see that the first $\lfloor k/2 \rfloor + 1$ rows of M are linearly independent, so equality holds, as claimed.

This proof was greatly facilitated by the fact that we did not need to work in a basis, but were able to find a manageable spanning set of vectors.

It is worth noting that all the non-unital eigenvalues in our example have equal multiplicity. Of course, the rationality of the entries of the matrix implies that two Galois-conjugate roots of 1 (i.e., roots that are primitive *m*th roots for the same *m*) must have the same multiplicity, but a priori roots that are not Galois-conjugate could have different multiplicities. Indeed, consider the rotation action $T_{4,2}$ restricted to multisets in which all elements are distinct; this set consists of the six elements 01, 02, 03, 12, 13, and 23, and rotation sends sends 01 to 12 to 23 to 03 to 01 and sends 02 to 13 to 02. The interested reader can verify that 1 has multiplicity 2, -1 has multiplicity 2, and *i* and -i each have multiplicity 1, so in this case the non-unital eigenvalues do not all have the same multiplicity.

5. Example: Rowmotion on a product of two chains

Recall from [S1] the definitions of a partially ordered set P, order ideals (or downsets), filters (or upsets), and antichains, and from [S2] and [EP1] the definitions of the order polytope $\mathcal{O}(P)$, reverse order polytope $\tilde{\mathcal{O}}(P)$, and chain polytope $\mathcal{A}(P)$ of a poset. For notation, I remind the reader that the order polytope of a finite poset P is the set of order-preserving maps $\lambda: P \to [0, 1]$, viewed as a subset of $\mathbb{R}^{|P|}$ in the natural way. Put $X = \mathcal{O}(P)$, $Y = \mathcal{A}(P)$, $Z = \tilde{\mathcal{O}}(P)$, and $T = \rho_{\mathcal{P}}$. Piecewise-linear rowmotion $\rho_{\mathcal{P}}$ (defined in [EP2]) is a volume-preserving invertible map from the order polytope to itself related to Striker and Williams' original definition of rowmotion for order ideals [SW]. When P is $[a] \times [b]$, $\rho_{\mathcal{P}}$ is of order n := a + b. For each $p \in P$, define the evaluation statistic $1_p(\lambda) = \lambda(p) \in [0, 1]$, so that we have statistics of the form $1_p \circ T^j$ $(0 \leq j \leq n - 1, p \in P)$ spanning a vector space V of dimension at most *nab*. Let U be the time-evolution operator on the space V sending $f \in V$ to $f \circ T \in V$.

Einstein and Propp, in unpublished work that was later generalized by Joseph and Roby (see sections 4 and 6 of the September 1, 2018 version of [EP1] and Theorem 5.12 in [JR]), showed that piecewise-linear rowmotion, although originally defined as a composition of piecewise-linear involutions (toggle operators), can be obtained as a composition of three transfer operators $\nabla : X \to Y$, $\Delta^{-1} : Y \to Z$, and $\Theta : Z \to X$, mirroring the original definition of rowmotion in the work of Brouwer and Schrijver [BS]. My goal in this section is to assert that linearization is "functorial" in the sense that the linearization of piecewise-linear rowmotion can be expressed as a composition of three linearized transfer operators. It is not my goal here to prove this assertion nor even to make the assertion plausible; I merely aim to make the *content* of the assertion clear by demonstrating it in the case a = b = 2 (the simplest case in which the piecewise-linear map T is not merely linear or affine).

In the transfer operator formulation, ρ_P for an arbitrary finite poset P is expressed as the composition $\Theta \circ \Delta^{-1} \circ \nabla$ where ∇ (down-transfer) is a piecewise-linear map from X to Y, Δ^{-1} (inverse up-transfer) is a piecewise-linear map from Y to Z, and Θ (complementation) is an affine map from Z back to X. In the case a = b = 2, for $x \in X$ put $y = \nabla(x)$, $z = \Delta^{-1}(y)$, and $x' = \Theta(z)$ so that Tx = x'. Explicitly,

$$\nabla(x_1, x_2, x_3, x_4) = (x_1 - 0, x_2 - x_1, x_3 - x_1, x_4 - \max(x_2, x_3)),$$

$$\Delta^{-1}(y_1, y_2, y_3, y_4) = (y_1 + \max(y_2, y_3) + y_4, y_2 + y_4, y_3 + y_4, y_4), \text{ and}$$

$$\Theta(z_1, z_2, z_3, z_4) = (1 - z_1, 1 - z_2, 1 - z_3, 1 - z_4).$$

To reduce the complexity of the notation let us use x_1, x_2, x_3, x_4 to denote the coordinate functions mapping X to \mathbb{R} , and likewise for the y and z variables. The dynamical closure of x_1, x_2, x_3, x_4 in X under the action of $\Theta \circ \Delta^{-1} \circ \nabla$ is 6-dimensional, with additional basis functions $x_5 = \max(x_2, x_3)$ and the constant function $x_6 = 1$. The dynamical closure of y_1, y_2, y_3, y_4 in Y under the action of $\nabla \circ \Theta \circ \Delta^{-1}$ is 6-dimensional, with additional basis functions $y_5 = \max(y_2, y_3)$ and $y_6 = 1$. The dynamical closure of z_1, z_2, z_3, z_4 in Z under the action of $\Delta^{-1} \circ \nabla \circ \Theta$ is 6-dimensional, with additional basis functions $z_5 = \max(z_2, z_3)$ and $z_6 = 1$. With these extra variables we have

$$U_{\nabla}(x_1, x_2, x_3, x_4, x_5, x_6) = (x_1, x_2 - x_1, x_3 - x_1, x_4 - x_5, x_5 - x_1, x_6),$$

$$U_{\Delta^{-1}}(y_1, y_2, y_3, y_4, y_5, y_6) = (y_1 + y_4 + y_5, y_2 + y_4, y_3 + y_4, y_4, y_4 + y_5, y_6), \text{ and}$$

$$U_{\Theta}(z_1, z_2, z_3, z_4, z_5, z_6) = (z_6 - z_1, z_6 - z_2, z_6 - z_3, z_6 - z_4, z_5 + z_6 - z_2 - z_3, z_6)$$

which demonstrates linear relations among all the quantities. (Those who wish to check the formulas should be mindful that at one point one needs to use the identity $\max(s,t) + \min(s,t) = s + t$.) Note that $U_{\Theta \circ \Delta^{-1} \circ \nabla} = U_{\nabla} U_{\Delta^{-1}} U_{\Theta}$.

A similar picture prevails for birational rowmotion (see [EP2]), where now the monoid is multiplicative rather than additive, along the lines of the example presented in the next section.

6. EXAMPLE: THE LYNESS 5-CYCLE

Here I revisit an example from section 2.6 of [PR]. The Lyness 5-cycle (the smallest nontrivial cluster algebra) exhibits a non-obvious homomesy: if X is the set of all

(x, y) in \mathbb{R}^2 with x, y, x + 1, y + 1, and x + y + 1 all nonzero, and $T : X \to X$ is the period-5 map sending (x, y) to (y, (y + 1)/x), and $g : X \to \mathbb{R}$ is the map sending (x, y) to $\log |x^{-1} + x^{-2}|$, then g is 0-mesic.

To fit this into the proposed framework, let us use the multiplicative monoid generated by the functions x, y, x + 1, y + 1, and x + y + 1; technically this is not a vector space unless we allow fractional exponents, but for present purposes this turns out not to matter. The map T sends $x^a y^b (x + 1)^c (y + 1)^d (x + y + 1)^e$ to $x^{a'} y^{b'} (x + 1)^{c'} (y + 1)^{d'} (x + y + 1)^{e'}$ where

$$(a' b' c' d' e') = (a b c d e) \begin{pmatrix} 0 & -1 & 0 & -1 & -1 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}.$$

This 5-by-5 matrix M satisfies $M^5 = I$, and each of the 5th roots of unity occurs in the spectrum with multiplicity 1.

The Lyness 5-cycle is associated with four-row frieze patterns. The above analysis can also be applied to frieze patterns with more than four rows; the details differ slightly according to whether the number of rows is odd or even. The eigenvalues still have dynamical meaning, inasmuch as they determine the dimensionalities of spaces of invariants for powers of the shift-map; however, the eigenvalues do not appear to be associated with anything like eigenvectors, since it is unclear how to to make sense of expressions like $x^{\alpha}y^{\beta}(x+1)^{\gamma}(y+1)^{\delta}(x+y+1)^{\epsilon}$ when $\alpha, \beta, \gamma, \delta, \epsilon$ are algebraic numbers.

7. FLATNESS

I have already remarked that, for a map T of period n, the rationality of the entries of the presenting matrix implies that two Galois-conjugate nth roots of 1 must have the same multiplicity. When n is prime, this implies that all non-unital eigenvalues have the same multiplicity, but when n is composite, the function that maps eigenvalues to multiplicities (hereafter the *spectral multiplicity function*) could a priori be far from constant. It is therefore somewhat surprising that for many examples, the spectral multiplicity function is "flat" in a logarithmic sense, meaning that its values are bounded by constant multiples of one another. For instance, in Proposition 4.1 the multiplicity takes on two values whose ratio approaches 1 as k gets large, while in Proposition 4.2 the multiplicity takes on two values whose ratio approaches 2 as kgets large.

I do not have a quantitative conjecture, but I suggest that this phenomenon – the flatness of the spectral multiplicity function – might apply in many situations. To the extent that this flatness is prevalent, it would provide a loose explanation for the relative paucity of invariants in comparison with homomesies. For, if the multiplicities of distinct eigenvalues always have ratio between 1/c and c (with c > 1), then the multiplicity of 1 divided by the sum of all the multiplicities must also be between 1/c and c, implying that dim $V/\dim V_1^{\perp}$ is between n/c and nc. That is, homomesies outnumber invariants roughly by a factor of n-1.

ACKNOWLEDGMENTS: The author thanks Joe Buhler, David Einstein, Darij Grinberg, Michael Joseph, and Tom Roby for helpful suggestions.

References

Algebraic Combinatorics, draft (May 26, 2021)

[BS] Andries Brouwer and Alexander Schrijver, On the period of an operator, defined on antichains. Math. Centrum report ZW 24/74 (1974).

[EP1] David Einstein and James Propp, Combinatorial, piecewise-linear, and birational homomesy for products of two chains. Preprint; https://arxiv.org/abs/1310.5294.

[EP2] David Einstein and James Propp, Combinatorial, piecewise-linear, and birational homomesy for products of two chains. To appear in *Algebraic Combinatorics*.

[JR] Michael Joseph and Tom Roby, Birational and noncommutative lifts of antichain toggling and rowmotion. To appear in *Algebraic Combinatorics*.

[PR] James Propp and Tom Roby, Homomesy in Products of Two Chains. *Electronic Journal of Combinatorics*, Volume 22, Issue 3 (2015), article P3.4.

[S1] Richard Stanley, Enumerative Combinatorics, vol 1.

[S2] Richard Stanley, Two Poset Polytopes. Discrete Comput. Geom. 1 (1986), no. 1, 9–23.

[SW] Jessica Striker and Nathan Williams, Promotion and rowmotion. *European Journal of Combinatorics* **33** (2012), 1919–1942.

University of Massachusetts Lowell, Department of Mathematical Sciences $\mathit{URL}: \texttt{https://jamespropp.org}$