On the weak norm of \mathscr{U}_p -residuals of all subgroups of a finite group^{*}

Yubo Lv[†] Yangming Li^{‡§}

Abstract

Let \mathscr{F} be a formation and G a finite group. The weak norm of a subgroup Hin G with respect to \mathscr{F} is defined by $N_{\mathscr{F}}(G, H) = \bigcap_{T \leq H} N_G(T^{\mathscr{F}})$. In particular, $N_{\mathscr{F}}(G) = N_{\mathscr{F}}(G,G)$. Let $N^i_{\mathscr{F}}(G), i \geq 1$, be a upper series of G by setting $N^{\mathfrak{F}}_{\mathscr{F}}(G) = 1, N^{i+1}_{\mathscr{F}}(G)/N^i_{\mathscr{F}}(G) = N_{\mathscr{F}}(G/N^i_{\mathscr{F}}(G))$ and denoted by $N^{\infty}_{\mathscr{F}}(G)$ the terminal term of the series. In this paper, for the case $\mathscr{F} \in \{\mathscr{U}_p, \mathscr{U}\}$, where $\mathscr{U}_p(\mathscr{U}, \text{respectively})$ is the class of all finite p-supersolvable groups(supersolvable groups, respectively), we characterize the structure of some given finite groups by the properties of weak norm of some subgroups in G with respect to \mathscr{F} . Some of our main results may regard as a continuation of many nice previous work.

Keywords: Weak norm; Normalizer; supersolvable residual; *p*-supersolvable residual; *p*-length; *p*-Fitting length.

MSC 2020: 20D10, 20D25.

1 Introduction

Throughout this paper, all groups are finite. We use the standard terminology and notations as in [1]. For the specific, we denote

• G: a finite group.

^{*}Supported by the major project of Basic and Applied Research (Natural Science) in Guangdong Province, China (Grant Number: 2017KZDXM058) and the Science and Technology Program of Guangzhou Municipality, China (Grant number: 201804010088).

[†]School of Mathematical Sciences, Guizhou Normal University, Guiyang, 550001, China, lvyubo341281@163.com

[‡]Corresponding author

[§]College of Mathematics, Guizhou Normal University, Guiyang, 550001, China; Dept. of Math., Guangdong University of Education, Guangzhou, 510310, China, liyangming@gdei.edu.cn

- |G|: the order of G.
- $\pi(G)$: the set of prime divisors of |G|.
- p: a prime.
- G_p : the Sylow *p*-subgroup of *G*.
- $F_p(G)$: the *p*-Fitting subgroup of *G*.
- \mathfrak{G} : the class of all finite groups.
- \mathscr{F} : a formation of groups, that is a class of finite groups satisfying the following: (1) if $G \in \mathscr{F}$ and N is a normal subgroup of G, then $G/N \in \mathscr{F}$, and (2) if N_1, N_2 are normal subgroups of G such that $G/N_i \in \mathscr{F}(i = 1, 2)$, then $G/(N_1 \cap N_2) \in \mathscr{F}$.
- $G^{\mathscr{F}}$: the \mathscr{F} -residual of G, that is the intersection of all those normal subgroups N of G such that $G/N \in \mathscr{F}$.
- $\mathscr{F}_1\mathscr{F}_2$: the formation product or Gaschëtz product of \mathscr{F}_1 and \mathscr{F}_2 , that is the class $\{G \in \mathfrak{G} \mid G^{\mathscr{F}_2} \in \mathscr{F}_1\}$. In particular, denote $\mathscr{F}^2 = \mathscr{F}\mathscr{F}$.
- \mathscr{A} : the class of all Abelian groups.
- \mathcal{N} : the class of all nilpotent groups.
- \mathcal{N}_p : the class of all *p*-nilpotent groups.
- \mathscr{U} : the class of all supersolvable groups.
- \mathscr{U}_p : the class of all *p*-supersolvable groups.
- $l_p(G)$: the *p*-length of a *p*-solvable group *G*, that is the number of the *p*-factor groups in the upper *p*-series of *G*:

$$1 = P_0(G) \trianglelefteq M_0(G) \trianglelefteq P_1(G) \trianglelefteq M_1(G) \trianglelefteq \cdots \trianglelefteq P_n(G) \trianglelefteq M_n(G) = G$$

such that $M_i(G)/P_i(G) = O_{p'}(G/P_i(G))$ and $P_i(G)/M_{i-1}(G) = O_p(G/M_{i-1}(G))$.

• $h_p(G)$: the *p*-Fitting length of a *p*-solvable group *G*, that is the smallest positive integer *n* such that

$$1 = F_p^0(G) \le F_p^1(G) \le \dots \le F_p^{n-1}(G) \le F_p^n(G) = G,$$

where $F_p^1(G) = F_p(G)$ and $F_p^{i+1}(G)/F_p^i(G) = F_p(G/F_p^i(G))$ for $i = 1, 2, \dots, n-1$.

- h(G): the Fitting length of a solvable group G, that is the positive integer n such that $1 = F^0(G) \leq F^1(G) \leq \cdots \leq F^{n-1}(G) \leq F^n(G) = G$, where $F^1(G) = F(G)$ and $F^{i+1}(G)/F^i(G) = F(G/F^i(G))$ for $i = 1, 2, \cdots, n-1$.
- N(G): the intersection of the normalizers of all subgroups of G.
- $S(G)(\text{or } N_{\mathscr{N}}(G))$: the intersection of the normalizers of the nilpotent residuals of all subgroups of G.
- $N^{\mathcal{N}_p}(G)$: the intersection of the normalizers of the *p*-nilpotent residuals of all subgroups of G.

It is interesting to characterize the structure of a given group by using of some special subgroups. For example, Gashëtz and N. Itô [1, Satz 5.7,p.436] proved that G is solvable with Fitting length at most 3 if all minimal subgroups of G are normal. Also, it is well know that G is nilpotent if G' normalizers each subgroup of G(see Baer's theorem in [2]). Further more, R. Baer in [3] defined the subgroup N(G), the norm of a group G. Obviously, a group G is a Dedeking group if and only if G = N(G). The norm of a group has many other good properties and has studied further by many scholars. In recent years, some weaker versions of the concepts of norm of groups have been introduced. Let \mathscr{F} be a non-empty formation. Recently, Su and Wang in [4, 5] introduced the subgroup $N_{\mathscr{F}}(G)$, the norm of \mathscr{F} -residual of a group G as follows

$$N_{\mathscr{F}}(G) = \bigcap_{H \le G} N_G(H^{\mathscr{F}}). \tag{(\star)}$$

As in [4], we set $N^0_{\mathscr{F}}(G) = 1$ and if $N^i_{\mathscr{F}}(G)$ is defined, set $N^{i+1}_{\mathscr{F}}(G)/N^i_{\mathscr{F}}(G) = N_{\mathscr{F}}(G/N^i_{\mathscr{F}}(G))$. The subgroup $N^\infty_{\mathscr{F}}(G)$ is the terminal term of the ascending series. In fact, $N^\infty_{\mathscr{F}}(G) = N^k_{\mathscr{F}}(G)$ for some integer k such that $N^k_{\mathscr{F}}(G) = N^{k+1}_{\mathscr{F}}(G) = N^{k+2}_{\mathscr{F}}(G) = \cdots$.

Many scholars also call $N_{\mathscr{F}}(G)(\mathscr{F} = \mathscr{A}, \mathscr{N}, \mathscr{N}_p$, respectively) the generalized norm of group G. Obviously, $N_{\mathscr{F}}(G)$ is a characteristic subgroup and every element of $N_{\mathscr{F}}(G)$ normalize the \mathscr{F} -residual of each subgroup of G. The so called norm of \mathscr{F} -residual has many other nice properties and also closely related to the global properties of a given group. For some given formation \mathscr{F} , there are many papers devoted to study the *p*-length, Fitting length, solvability, (*p*-)nilpotency and so on. For example, for the case $\mathscr{F} = \mathscr{A}$, Li and Shen in [6] denoted $N_{\mathscr{A}}(G)$ by D(G). They fingered out that G is solvable with Fitting length at most 3 if all elements of G of prime order are in D(G)(see [6, Theorem 4.1]). It is a dual problem of Gashëtz and N. Itô[1, Satz 5.7, p.436]. Li and Shen also defined the D-group, i.e., G = D(G), they characterized the relationship between D(G) and G. Shen, Shi and Qian in [7] considered the case $\mathscr{F} = \mathscr{N}$, they denoted $N_{\mathscr{N}}(G)$ by S(G). Shen et al., deeply studied the dual problem of Gashëtz and N. Itô and characterized the \mathscr{F}_{nn} -groups by means of the subgroup $S^{\infty}(G)$, where $S^{\infty}(G) = N^{\infty}_{\mathscr{N}}(G)$ and \mathscr{F}_{nn} -groups are class of groups belong to $\mathscr{N}\mathscr{N}$. They also introduced the S-group, i.e., G = S(G) and given some sufficient and necessary conditions involved S-groups. Meanwhile, Gong and Guo in [8] also consider the case $\mathscr{F} = \mathscr{N}$ and given some meaningful conclusions. In the case $\mathscr{F} = \mathscr{N}_p$, Guo and Li in [9] introduced the norm of \mathscr{N}_p -residual of a group G, they denoted $N_{\mathscr{N}_p}(G)$ by $N^{\mathscr{N}_p}(G)$. As a local version of Gong's results, Li and Guo characterized the relationship between $C_G(G^{\mathscr{N}_p})$ and $N^{\mathscr{N}_p}(G)$. In particular, Li and Guo in [9] also investigated the relationship between $N_{\mathscr{N}}(G)$ and $N^{\mathscr{N}_p}(G)$. For more detail and other relevant conclusions about the norm of \mathscr{F} -residual, please see [4-13].

We wonder whether the above conclusions hold for general formations. A natural idea is to replace $\mathscr{F}(\mathscr{F} \in {\mathscr{N}_p, \mathscr{N}, \mathscr{A}})$ by \mathscr{U}_p or \mathscr{U} in (\star) .

Remark 1.1. (1) In general, for a group G and some $p \in \pi(G)$, $N_{\mathscr{U}_p}(G) \neq N_{\mathscr{F}}(G)$ is possible, where $\mathscr{F} \in \{\mathscr{N}_p, \mathscr{N}, \mathscr{A}\}$. For example, let $G = S_4$, the symmetric group of degree 4. Obviously, G is a 3-supersolvable non-3-nilpotent group, so $G = N_{\mathscr{U}_3}(G)$. Pick a subgroup $H \cong S_3$, the symmetric group of degree 3, but $N_G(H^{\mathscr{N}_3}) = N_G(S_3^{\mathscr{N}_3}) =$ $N_G(C_3) = S_3 < S_4$, so $N_{\mathscr{N}_3}(G) < G$.

(2) The condition in definition (\star) that the intersection of the \mathscr{F} -residuals of all subgroups of G may be too strong and some of the subgroups may be redundant whenever $\mathscr{F} = \mathscr{U}_p$ (see Example 1.3 below).

(3) The case that $N_{\mathscr{U}}(G) = N_{\mathscr{U}_p}(G) = 1$ is possible for a solvable group G and $p \in \pi(G)$. Let $H = \langle a, b \mid a^3 = b^3 = 1, [a, b] = 1 \rangle \cong C_3 \times C_3$ and $Q_8 = \langle c, d \mid c^4 = 1, c^2 = d^2 = e, c^d = c^{-1} \rangle$. Considering the irreducible action of Q_8 on H by $a^c = a^{-1}b, b^c = ab, a^d = b^{-1}, b^d = a$, denote $T = H \rtimes Q_8 \cong (C_3 \times C_3) \rtimes Q_8$. Let $C = \langle f \rangle$ be a cyclic group of order 3 and let C act on T by $a^f = b^{-1}, b^f = ab^{-1}, c^f = d^3, d^f = cd$. Then $G = \langle a, b, c, d, e, f \rangle \cong ((C_3 \times C_3) \rtimes Q_8) \rtimes C_3$ is solvable(IdGroup=[216,153]). It is easy to see that $G^{\mathscr{U}_2} = \langle a, b, c, d, e \rangle \cong (C_3 \times C_3) \rtimes Q_8$ and $C_G(G^{\mathscr{U}_2}) = 1$. By Theorem 3.4, $N_{\mathscr{U}_2}(G) = 1$. Further more, by Theorem 3.14, $N_{\mathscr{U}}(G) = 1$. So we have $N_{\mathscr{U}}(G) = N_{\mathscr{U}_2}(G) = 1$.

Recall that the weak centralizer of H in G, $C_G^*(H)$, introduced in [14, P.33, Definitions], is defined by

$$C_G^*(H) = \bigcap \{ N_G(K) : K \le H \}.$$

In the above investigation, we introduce the following more general and interesting definition.

Definition 1.2. Let G be a group and \mathscr{F} a formation. We define $N_{\mathscr{F}}(G, H)$, the weak norm of H in G with respect to \mathscr{F} as follows:

$$N_{\mathscr{F}}(G,H) = \bigcap_{T \le H} N_G(T^{\mathscr{F}}).$$

In particular, $N_{\mathscr{U}}(G, H) = \bigcap_{T \leq H} N_G(T^{\mathscr{U}})$ and $N_{\mathscr{U}_p}(G, H) = \bigcap_{T \leq H} N_G(T^{\mathscr{U}_p}).$

Example 1.3. Let $G = \langle a, b, c, d \rangle \rtimes \langle e, f \rangle \cong C_2^4 \rtimes C_6$, where $e^2 = f^3 = 1$ and $a^e = ac, b^e = bd, c^e = c, d^e = d, a^f = b, b^f = ab, c^f = d, d^f = cd$ (IdGroup=[96,70] in GAP [15]). Let $H = \langle a, b, f \rangle$, then $N_{\mathscr{U}_2}(G) = N_{\mathscr{U}_2}(G, H) = \langle a, b, c, d, f \rangle \cong C_2^4 \rtimes C_3 < G$.

Obviously, $N_{\mathscr{F}}(G) = N_{\mathscr{F}}(G,G) \leq N_{\mathscr{F}}(G,H) \leq G$. Without causing confusion, we call $N_{\mathscr{F}}(G) = N_{\mathscr{F}}(G,G)$ the norm of G with respect to \mathscr{F} . Moreover, for a subgroup H of G, we call G is an \mathscr{N}_1 -group with respect to H and \mathscr{F} if G = $N_{\mathscr{F}}(G,H)$. In this paper, we mainly investigate properties of $N_{\mathscr{F}}(G,H)$ and the influence of $N_{\mathscr{F}}(G,H)$ on the structure of group G. Actually, we mainly consider the case $\mathscr{F} \in \{\mathscr{U}, \mathscr{U}_p\}$. Our main work may be regard as the continuation of some conclusions in [6–9].

2 Preliminaries

In this section, we always assume that \mathscr{F} is a non-empty formation and G is a group. We first give some important lemmas.

Lemma 2.1. Let H, K, N be subgroups of G and $N \leq G$. Then

(1) If $H \leq K$, then $N_{\mathscr{F}}(G, K) \leq N_{\mathscr{F}}(G, H)$; (2) $K \cap N_{\mathscr{F}}(G, H) \leq N_{\mathscr{F}}(K, K \cap H)$, in particular, if $H \leq K$, then $K \cap N_{\mathscr{F}}(G, H) \leq N_{\mathscr{F}}(K, H)$. (3) If $N \leq H$, then $N_{\mathscr{F}}(G, H)N/N \leq N_{\mathscr{F}}(G/N, H/N)$.

Proof. (1) By definition, $N_{\mathscr{F}}(G, K) = \bigcap_{T \leq K} N_G(T^{\mathscr{F}}) \leq \bigcap_{T \leq H} N_G(T^{\mathscr{F}}) = N_{\mathscr{F}}(G, H).$ (2) Obviously, $K \cap N_{\mathscr{F}}(G, H) = K \cap (\bigcap_{T \leq H} N_G(T^{\mathscr{F}})) \leq \bigcap_{T \leq H \cap K} N_K(T^{\mathscr{F}}) = N_{\mathscr{F}}(K, H \cap K).$ In particular, if $H \leq K$, then $K \cap N_{\mathscr{F}}(G, H) \leq N_{\mathscr{F}}(K, H).$

(3) Let $x \in N_{\mathscr{F}}(G, H)$, then x normalize $T^{\mathscr{F}}$ for every $T \leq H$, so xN normalize $T^{\mathscr{F}}N/N = (TN/N)^{\mathscr{F}}$. Thus every element of $N_{\mathscr{F}}(G, H)N/N$ normalize $(T/N)^{\mathscr{F}}$ for all subgroups T/N of H/N, so $N_{\mathscr{F}}(G, H)N/N \leq N_{\mathscr{F}}(G/N, H/N)$.

As a corollary, we have

Lemma 2.2. Let G be a group, let K be a subgroup of G and N a normal subgroup of G, then

(1) $K \cap N_{\mathscr{F}}(G) \le N_{\mathscr{F}}(K).$ (2) $N_{\mathscr{F}}(G)N/N \le N_{\mathscr{F}}(G/N).$

Lemma 2.3. [4, Lemmas 2.2,2.3] Let G be a group, let K be a subgroup of G and N a normal subgroup of G, then

- (1) $K \cap N^{\infty}_{\mathscr{F}}(G) \leq N^{\infty}_{\mathscr{F}}(K).$ (2) $N^{\infty}_{\mathscr{F}}(G)N/N \leq N^{\infty}_{\mathscr{F}}(G/N).$
- $(2) N_{\mathcal{F}}(G)N/N \leq N_{\mathcal{F}}(G/N).$

(3) If $N \leq N^{\infty}_{\mathscr{F}}(G)$, then $N^{\infty}_{\mathscr{F}}(G)/N = N^{\infty}_{\mathscr{F}}(G/N)$.

Lemma 2.4. Let G be a p-solvable group.

(1) If $N \leq G$, then $l_p(G/N) \leq l_p(G)$.

(2) If $U \leq G$, then $l_p(U) \leq l_p(G)$.

(3) Let N_1 and N_2 be two normal subgroups of G, then $l_p(G/(N_1 \cap N_2)) \leq max\{l_p(G/N_1), l_p(G/N_2)\}.$

- (4) $l_p(G/\Phi(G)) = l_p(G).$
- (5) If N is a normal p'-group of G, then $l_p(G/N) = l_p(G)$.

Proof. The proof of (1)-(4) follows from [1, VI, 6.4] and (5) is obviously.

Lemma 2.5. Let G be a group and N a normal subgroup of G. Then

(1) if $N/N \cap \Phi(G)$ is p-nilpotent, then N is p-nilpotent;

(2) Let H be a subgroup of G and N be a p'-group, if HN/N is p-nilpotent, then H is p-nilpotent.

Proof. The statement (1) is directly form [16, Lemma 2.5], and (2) is easy.

Lemma 2.6. Let G be a p-solvable group and H a subgroup of G, let H, N, A, B be subgroups of G and N, A, B are normal in G, then

(1) $h_p(H) \le h_p(G)$. (2) $h_p(G/N) \le h_p(G)$. (3) If $G = A \times B$, then $h_p(G) = \max\{h_p(A), h_p(B)\}$. (4) If $h_p(G/A) \le k$ and $h_p(G/N) \le k$, then $h_p(G/(A \cap B)) \le k$. (5) $h_p(G/\Phi(G))) = h_p(G)$.

(6) If N is a p'-group, then $h_p(G/N) = h_p(G)$.

Proof. Let $1 = N_0 \leq N_1 \leq N_2 \leq \cdots \leq N_r = G$ be the shortest normal chain of G with p-nilpotent factors N_i/N_{i-1} for all $i = 1, 2, \cdots, r$.

(1) If $H \leq G$, then $1 = N_0 \cap H \leq N_1 \cap H \leq N_2 \cap H \leq \cdots \leq N_r \cap H = G \cap H = H$ is a normal chain of H with $N_i \cap H/N_{i-1} \cap H \cong (N_i \cap H)N_{i-1}/N_{i-1}$ a p-nilpotent factors for all $i = 1, 2, \cdots, r$, so $h_p(H) \leq h_p(G)$. (2) If $N \leq G$, obviously, $\overline{1} = N_0 N/N \leq N_1 N/N \leq N_2 N/N \leq \cdots \leq N_r/N = G/N$ is a normal chain of G/N. Note that $N_i N/N/N_{i-1} N/N = N_i N/N_{i-1}N \cong N_i/N_{i-1}(N_i \cap N) \leq N_i/N_{i-1}$ is *p*-nilpotent, so $h_p(G/N) \leq h_p(G)$.

(3) Let $1 = A_0 \leq A_1 \leq A_2 \leq \cdots \leq A_r = A$ and $1 = B_0 \leq B_1 \leq B_2 \leq \cdots \leq B_t = B$ be the shortest normal chain of A and B with p-nilpotent factors respectively. Without loss of generality, assume $r \leq t$, then $1 = A_0B_0 \leq A_1B_1 \leq A_2B_2 \leq \cdots \leq A_rB_r \leq A_rB_{r+1} \leq A_rB_{r+2} \leq A_rB_t = AB$ is a normal chain. Since $A_iB_i/A_{i-1}B_{i-1} \cong A_i/A_{i-1}(A_i \cap B_{i-1}) \cdot B_i/B_{i-1}(B_i \cap A_{i-1})$ whenever $i \leq r$ and $A_rB_j/A_rB_{j-1} \cong B_j/B_{j-1}(B_j \cap A_i)$ whenever $r < j \leq t$ are p-nilpotent. So $h_p(G) = \max\{h_p(A), h_p(B)\}$.

(4) Since $G/(A \cap B) \cong G/A \times G/B$, the result follows from (1) and (3).

(5) Assume that $\overline{1} = \Phi(G)/\Phi(G) = T_0/\Phi(G) \leq T_1/\Phi(G) \leq T_2/\Phi(G) \leq \cdots \leq T_s/\Phi(G) = G/\Phi(G)$ is the normal chain of G with p-nilpotent factors, so $s \leq r$ and $T_i/T_{i-1} \cong T_i/\Phi(G)/T_{i-1}/\Phi(G)$ is p-nilpotent. Since $T_1/\Phi(G)$ is p-nilpotent, then T_1 is p-nilpotent by Lemma 2.5, so $1 \leq T_1 \leq T_2 \leq \cdots \leq T_s = G$ is a normal chain with p-nilpotent factors. Now we have $r \leq s$, hence s = r.

(6) Let $\overline{1} = N_0/N \leq N_1/N \leq N_2/N \leq \cdots \leq N_r/N = G/N$ be a normal chain of G/N with p-nilpotent factors $N_i/N/N_{i-1}/N \cong N_i/N_{i-1}$ for all $i = 1, 2, \cdots, r$. Now by Lemma 2.5, N_1 is p-nilpotent since N is a p'-group, so $1 \leq N_1 \leq N_2 \leq \cdots \leq N_r = G$ is a normal chain of G with p-nilpotent factors N_i/N_{i-1} for all $i = 1, 2, \cdots, r$, which implies that $h_p(G) \leq h_p(G/N)$, now by (1), we have $h_p(G/N) = h_p(G)$.

As a local vision of [17, Lemma 3.2], we have

Lemma 2.7. Let G be a solvable group, then $l_p(G) \leq 1$ if $h(G) \leq 2$ for every $p \in \pi(G)$.

Lemma 2.8. [18, Theorem 1 and Proposition 1] Let \mathscr{F} be a saturated formation.

(1) Assume that G is a group such that G does not belong to \mathscr{F} , but all its proper subgroups belong to \mathscr{F} . Then $F'(G)/\Phi(G)$ is the unique minimal normal subgroup of $G/\Phi(G)$, where $F'(G) = Soc(G \mod \Phi(G))$, and $F'(G) = G^{\mathscr{F}}\Phi(G)$. In addition, if the derived subgroup of $G^{\mathscr{F}}$ is a proper subgroup of $G^{\mathscr{F}}$, then $G^{\mathscr{F}}$ is a soluble group. Furthermore, if $G^{\mathscr{F}}$ is soluble, then F'(G) = F(G), the Fitting subgroup of G. Moreover $(G^{\mathscr{F}})' = T \cap G^{\mathscr{F}}$ for every maximal subgroup T of G such that $G/T_G \notin \mathscr{F}$ and F'(G)T = G.

(2) Assume that G is a group such that G does not belong to \mathscr{F} and there exists a maximal subgroup M of G such that $M \in \mathscr{F}$ and G = MF(G). Then $G^{\mathscr{F}}/(G^{\mathscr{F}})'$ is a chief factor of G, $G^{\mathscr{F}}$ is a p-group for some prime p, $G^{\mathscr{F}}$ has exponent p if p > 2 and exponent at most 4 if p = 2. Moreover, either $G^{\mathscr{F}}$ is elementary abelian or $(G^{\mathscr{F}})' = Z(G^{\mathscr{F}}) = \Phi(G^{\mathscr{F}})$ is an elementary abelian group.

Lemma 2.9. Let G be a p-solvable minimal non-p-supersolvable group, then $G^{\mathscr{U}_p}$ is a p-group.

Proof. By Lemma 2.8(1), $G/\Phi(G)$ has the unique minimal normal subgroup $F'(G)/\Phi(G)$, where $F'(G) = G^{\mathscr{U}_p}\Phi(G)$, so $F'(G)/\Phi(G)$ is an elementary abelian *p*-group since *G* is a *p*-solvable group. Otherwise, $F'(G)/\Phi(G)$ is a *p'*-group and the *p*-supersolvability of $G/\Phi(G)/F'(G)/\Phi(G)$ implies that $G/\Phi(G)$ is *p*-supersolvable, thus *G* is *p*-supersolvable, a contradiction. Furthermore, $\Phi(G) = 1$. Now we have $G^{\mathscr{U}_p}$ is solvable and F'(G) =F(G), G = F(G)T for some maximal subgroup *T* of *G*. Now by Lemma 2.8(2), $G^{\mathscr{U}_p}$ is a *p*-group.

Lemma 2.10. Let G be a group and $\pi(G) = \{p_1, p_2, \cdots, p_n\}$, then $G^{\mathscr{U}} = G^{\mathscr{U}_{p_1}} G^{\mathscr{U}_{p_2}} \cdots G^{\mathscr{U}_{p_n}}$.

Proof. It follows from the supersolvability of $G/G^{\mathscr{U}}$ that $G^{\mathscr{U}_{p_i}} \leq G^{\mathscr{U}}$ for all $p_i \in \pi(G)$, so $G^{\mathscr{U}_{p_1}}G^{\mathscr{U}_{p_2}}\cdots G^{\mathscr{U}_{p_n}} \leq G^{\mathscr{U}}$. Conversely, obviously, $G/(G^{\mathscr{U}_{p_1}}G^{\mathscr{U}_{p_2}}\cdots G^{\mathscr{U}_{p_n}}) \in \mathscr{U}_{p_i}$ for any $p_i \in \pi(G)$, so $G^{\mathscr{U}} \leq G^{\mathscr{U}_{p_1}}G^{\mathscr{U}_{p_2}}\cdots G^{\mathscr{U}_{p_n}}$, as desired. \Box

3 Basic properties

Now we give some elementary properties. Firstly, we have

Proposition 3.1. Let $G = A \times B$ with (|A|, |B|) = 1 and $H \leq G$, then $N_{\mathscr{U}_p}(G, H) = N_{\mathscr{U}_p}(A, A \cap H) \times B$ whenever $p \in \pi(A)$ or $A \times N_{\mathscr{U}_p}(B, B \cap H)$ whenever $p \in \pi(B)$.

Proof. By hypothesis, we may assume that $p \in \pi(A) \setminus \pi(B)$. Note that $H = (H \cap A) \times (H \cap B)$ and $T = (T \cap A) \times (T \cap B)$, so $T^{\mathscr{U}_p} = (T \cap A)^{\mathscr{U}_p} \times (T \cap B)^{\mathscr{U}_p} = (T \cap A)^{\mathscr{U}_p}$, where $T \leq H$. Now $N_G(T^{\mathscr{U}_p}) = N_{A \times B}(T^{\mathscr{U}_p}) = N_A((T \cap A)^{\mathscr{U}_p}) \times B$. We have

$$N_{\mathscr{U}_p}(G, H) = \bigcap_{T \le H} N_G(T^{\mathscr{U}_p})$$
$$= \left(\bigcap_{T \le H} N_A((T \cap A)^{\mathscr{U}_p})\right) \times B$$
$$= \left(\bigcap_{T \le H \cap A} N_A(T^{\mathscr{U}_p})\right) \times B$$
$$= N_{\mathscr{U}_p}(A, A \cap H) \times B.$$

Similarly, if $p \in \pi(B) \setminus \pi(A)$, we have $N_{\mathscr{U}_p}(G, H) = A \times N_{\mathscr{U}_p}(B, B \cap H)$.

Proposition 3.2. Let G be a p-solvable group, then the \mathscr{U}_p -residual of $N_{\mathscr{U}_p}(G)$ is a p-group.

Proof. Denote $X = N_{\mathscr{U}_p}(G)$. We first prove this result in the case that X = G. Assume $G \in \mathscr{U}_p$, obviously $G^{\mathscr{U}_p} = 1$ is a *p*-group. Assume now that $G \notin \mathscr{U}_p$, if $O_p(G) > 1$, then by Lemma 2.2(2), $G/O_p(G) = N_{\mathscr{U}_p}(G/O_p(G))$. By induction on $|G|, (G/O_p(G))^{\mathscr{U}_p} = N_{\mathscr{U}_p}^{\mathscr{U}_p}(G/O_p(G))$ is a *p*-group. Obviously $G^{\mathscr{U}_p}$ is a *p*-group. If $O_p(G) = 1$, then for any K < G, by Lemma 2.2(1), $K = N_{\mathscr{U}_p}(K)$, so by induction, $K^{\mathscr{U}_p}$ is a *p*-group. Note that $K^{\mathscr{U}_p} \leq G$, so $K^{\mathscr{U}_p} \leq O_p(G) = 1$, this implies that G is a *p*-solvable minimal non-*p*-supersolvable group, now by Lemma 2.9, $G^{\mathscr{U}_p}$ is a *p*-group.

Assume now that X < G, by Lemma 2.2(1), $X = X \cap N_{\mathscr{U}_p}(G) \leq N_{\mathscr{U}_p}(X) \leq X$, so $X = N_{\mathscr{U}_p}(X)$. As a similar argument above, we also have the conclusion.

Denote $Z_{\infty}(G)$ be the terminal term of the ascending central series of G. As we know, $Z_{\infty}(G) = \bigcap \{N \leq G \mid Z(G/N) = 1\}$, we have the following similar result.

Theorem 3.3. Let G be a group, then $N^{\infty}_{\mathscr{U}_p}(G) = \bigcap \{N \leq G \mid N_{\mathscr{U}_p}(G/N) = 1\}.$

Proof. Let $N \leq G$ such that $N_{\mathscr{U}_p}(G/N) = 1$, then $N^{\infty}_{\mathscr{U}_p}(G/N) = 1$. By Lemma 2.3(2), $N^{\infty}_{\mathscr{U}_p}(G)N/N \leq N^{\infty}_{\mathscr{U}_p}(G/N) = 1$, so $N^{\infty}_{\mathscr{U}_p}(G) \leq N$, thus $N^{\infty}_{\mathscr{U}_p}(G) \leq \bigcap \{N \leq G \mid N_{\mathscr{U}_p}(G/N) = 1\}$.

Conversely, choice the least positive integer n such that $N^{\infty}_{\mathscr{U}_p}(G) = N^n_{\mathscr{U}_p}(G)$. By definition, $N_{\mathscr{U}_p}(G/N^n_{\mathscr{U}_p}(G)) = N^{n+1}_{\mathscr{U}_p}(G)/N^n_{\mathscr{U}_p}(G) = N^n_{\mathscr{U}_p}(G)/N^n_{\mathscr{U}_p}(G) = 1$. Obviously, $\bigcap \{N \leq G \mid N_{\mathscr{U}_p}(G/N) = 1\} \leq N^{\infty}_{\mathscr{U}_p}(G)$.

Now we give some characterizations between $N_{\mathscr{U}_p}(G, H)$ and $C_G(H^{\mathscr{U}_p})$.

Theorem 3.4. Let G be a p-solvable group and H a normal subgroup of G, then $N_{\mathscr{U}_p}(G, H) = 1$ if and only if $C_G(H^{\mathscr{U}_p}) = 1$.

Proof. As we know, $C_G(H^{\mathscr{U}_p}) \leq N_{\mathscr{U}_p}(G, H)$, the necessary is obviously. Assume now $C_G(H^{\mathscr{U}_p}) = 1$, we prove that $N_{\mathscr{U}_p}(G, H) = 1$. If not, let N be a minimal normal subgroup of G contained in $N_{\mathscr{U}_p}(G, H)$, if $N \not\leq H$, then N normalize $H^{\mathscr{U}_p}$ by definition, we have $[N, H^{\mathscr{U}_p}] \leq N \cap H^{\mathscr{U}_p} = 1$ by the minimal normality of N, so $N \leq C_G(H^{\mathscr{U}_p}) = 1$, a contradiction. If H < G, then (H, H) satisfies our hypothesis. By induction on |G||H| and Lemma 2.1(2), $N \leq H \cap N_{\mathscr{U}_p}(G, H) \leq N_{\mathscr{U}_p}(H, H) = 1$, a contradiction, thus H = G. Now by the p-solvability of G, N is a p'-group or an elementary abelian p-group. If N is a p'-group, then $G/C_G(N) \in \mathscr{U}_p$, otherwise, there is a minimal non-p-supersolvable subgroup $T/C_G(N)$ of $G/C_G(N)$. Let $T = C_G(N)L$ such that $C_G(N) \cap L \leq \Phi(L)$. Since $L/\Phi(L) \leq T/C_G(N) \cong L/L \cap C_G(N)$ is a minimal non-p-supersolvable group, we have $(L/\Phi(L))^{\mathscr{U}_p} = L^{\mathscr{U}_p}\Phi(L)/\Phi(L)$ is a pgroup by Lemma 2.9. Now let $P \in Syl_p(L^{\mathscr{U}_p})$, then $L^{\mathscr{U}_p} = P(L^{\mathscr{U}_p} \cap \Phi(L))$. By Frattini argument, $L = N_L(P)L^{\mathscr{U}_p} = N_L(P)P(L^{\mathscr{U}_p} \cap \Phi(L)) = N_L(P)$, we have $P \leq L^{\mathscr{U}_p}$ and N normalize P, so $P \leq C_G(N) \cap L \leq \Phi(L)$, thus $(L/\Phi(L))^{\mathscr{U}_p} = 1, L \in \mathscr{U}_p$, contrary to our choice of $T/C_G(N)$. Consequently, $G^{\mathscr{U}_p} \leq C_G(N)$, hence $N \leq C_G(G^{\mathscr{U}_p}) = 1$, a contradiction. So N is an elementary abelian p-group and $N \leq C_G(N)$. Note that $\Phi(G) \leq F(G) \leq C_G(N)$, we consider the following two cases:

Case 1. $\Phi(G) = C_G(N)$.

In this case, $F(G) = C_G(N) = \Phi(G) \leq F_p(G)$, so $N \leq \Phi(G)$. By the *p*-solvability of G, $F_p(G/\Phi(G)) = F_p(G)/\Phi(G) \neq 1$, so $\Phi(G) = F(G) < F_p(G)$. Now follows from $F_p(G)/O_{p'}(G) = O_{p'p}(G)/O_{p'}(G) = O_p(G/O_{p'}(G))$ that $O_{p'}(G) > 1$, so $O_{p'}(G) \leq C_G(N) = \Phi(G)$. On the other hand, Let $P \in Syl_p(F_p(G))$, then $F_p(G) = [O_{p'}(G)]P$. By the Frattini argument, we have $G = F_p(G)N_G(P) = O_{p'}(G)N_G(P) = N_G(P)$, so $F_p(G) = P \times O_{p'}(G) = F(G)$, a contradiction.

Case 2. $\Phi(G) < C_G(N)$.

In this case, if $N \not\leq \Phi(G)$, then there exists some maximal subgroup M of G such that G = NM and $N \cap M = 1$. Obviously, $(G/N)^{\mathscr{U}_p} = G^{\mathscr{U}_p} N/N \cong G^{\mathscr{U}_p}/G^{\mathscr{U}_p} \cap N \cong$ $M^{\mathscr{U}_p}$ and N normalize $M^{\mathscr{U}_p}$. By the minimal normality of N, we have $G^{\mathscr{U}_p} \cap N = 1$ or $G^{\mathscr{U}_p} \cap N = N$. If the former holds, then $N \leq C_G(G^{\mathscr{U}_p}) = 1$, a contradiction. If the later holds, then $G^{\mathscr{U}_p}/N \cong M^{\mathscr{U}_p}$. On the other hand, $[N, M^{\mathscr{U}_p}] = 1$, so $N \leq C_G(M^{\mathscr{U}_p}) \cap C_G(N) = C_G(G^{\mathscr{U}_p}) = 1$ since $G^{\mathscr{U}_p} = NM^{\mathscr{U}_p} = N \times M^{\mathscr{U}_p}$, a contradiction. Now if $G = C_G(N)$, then $N \leq C_G(G^{\mathscr{U}_p}) = 1$, a contradiction. So $C_G(N) < G$ and we may chose a non-p-supersolvable subgroup L of G such that $G = C_G(N)L$ and $C_G(N) \cap L \leq \Phi(L)$. Obviously, N normalize $L^{\mathscr{U}_p}$ and $N \cap L^{\mathscr{U}_p} \leq G$. Since N is a minimal normal subgroup of G, we have $N \cap L^{\mathscr{U}_p} = 1$ or $N \cap L^{\mathscr{U}_p} = N$. If $N \cap L^{\mathscr{U}_p} = 1$, then $L^{\mathscr{U}_p} \leq C_G(N) \cap L$, so $G/C_G(N) \cong L/L \cap C_G(N) \in \mathscr{U}_p$, thus $G^{\mathscr{U}_p} \leq C_G(N)$ and hence $N \leq C_G(G^{\mathscr{U}_p}) = 1$, a contradiction. Which implies that $N \leq L^{\mathscr{U}_p}$. Now by the fact $G = C_G(N)L$ that N is a minimal normal subgroup of L. So $F(L) = C_L(N) = C_G(N) \cap L = \Phi(L)$ and $F(L) < F_p(L)$, as a similar argument of **Case 1**, we also have a contradiction that $F_p(L) = F(L)$.

As corollaries of Theorem 3.4, we have

Corollary 3.5. Let G be a solvable group and H a normal subgroup of G, then $N_{\mathscr{U}}(G, H) = 1$ if and only if $C_G(H^{\mathscr{U}}) = 1$.

Corollary 3.6. Let G be a p-solvable group. Then $N_{\mathscr{U}_p}(G) = 1$ if and only if $C_G(G^{\mathscr{U}_p}) = 1$.

Corollary 3.7. Let G be a p-solvable group, if $Z(G^{\mathscr{U}_p}) = 1$, then $C_G(G^{\mathscr{U}_p}) = N_{\mathscr{U}_p}(G)$.

Proof. By Corollary 3.6, we may assume that $C_G(G^{\mathscr{U}_p}) > 1$. Moreover, obviously,

 $C_G(G^{\mathscr{U}_p}) \leq N_{\mathscr{U}_p}(G)$. Now consider $G/C_G(G^{\mathscr{U}_p})$, let

$$gC_G(G^{\mathscr{U}_p}) \in C_{G/C_G(G^{\mathscr{U}_p})}((G/C_G(G^{\mathscr{U}_p})^{\mathscr{U}_p}) = C_{G/C_G(G^{\mathscr{U}_p})}(G^{\mathscr{U}_p}C_G(G^{\mathscr{U}_p})/C_G(G^{\mathscr{U}_p})),$$

then $[g, G^{\mathscr{U}_p}] \leq C_G(G^{\mathscr{U}_p}) \cap G^{\mathscr{U}_p} = Z(G^{\mathscr{U}_p}) = 1$, thus $g \in C_G(G^{\mathscr{U}_p})$ and $C_{G/C_G(G^{\mathscr{U}_p})}(G/C_G(G^{\mathscr{U}_p})^{\mathscr{U}_p}) = \overline{1}$. By Corollary 3.6, we have $N_{\mathscr{U}_p}(G/C_G(G^{\mathscr{U}_p})) = \overline{1}$, so by Lemma 2.2(2), $N_{\mathscr{U}_p}(G) \leq C_G(G^{\mathscr{U}_p})$, as desired.

Corollary 3.8. Let G be a p-solvable group, then $Z(G^{\mathscr{U}_p}) = 1$ if and only if $G^{\mathscr{U}_p} \cap N_{\mathscr{U}_p}(G) = 1$.

Proof. Since $Z(G^{\mathscr{U}_p}) \leq G^{\mathscr{U}_p} \cap N_{\mathscr{U}_p}(G)$, so the "only if" part is obviously. Now assume that $Z(G^{\mathscr{U}_p}) = 1$, it follows form Corollary 3.7 that $C_G(G^{\mathscr{U}_p}) = N_{\mathscr{U}_p}(G)$, so $G^{\mathscr{U}_p} \cap N_{\mathscr{U}_p}(G) = G^{\mathscr{U}_p} \cap C_G(G^{\mathscr{U}_p}) = Z(G^{\mathscr{U}_p}) = 1$, as desired.

Theorem 3.9. Let G be a group, then $Z_{\infty}(G^{\mathscr{U}_p}) \leq N_{\mathscr{U}_p}^{\infty}(G)$.

Proof. If $Z(G^{\mathscr{U}_p}) = 1$, obviously the result holds. So we assume $Z(G^{\mathscr{U}_p}) > 1$. Now, $Z(G^{\mathscr{U}_p}) \leq N_{\mathscr{U}_p}(G)$ by definition. By induction on |G|, $Z_{\infty}(G^{\mathscr{U}_p}/Z(G^{\mathscr{U}_p})) = Z_{\infty}((G/Z(G^{\mathscr{U}_p}))^{\mathscr{U}_p}) \leq N_{\mathscr{U}_p}^{\infty}(G/Z(G^{\mathscr{U}_p}))$, so $Z_{\infty}(G^{\mathscr{U}_p}) \leq N_{\mathscr{U}_p}^{\infty}(G)$ by Lemma 2.3(3). \Box

As a similar argument above, we have

Corollary 3.10. Let G be a group, then $Z_{\infty}(G^{\mathscr{U}}) \leq N^{\infty}_{\mathscr{U}_p}(G)$ for every $p \in \pi(G)$.

At the end of this section, we investigate the relationship between $N_{\mathscr{U}_p}(G, H)$ and $N_{\mathscr{U}}(G, H)$.

Lemma 3.11. Let G be a group and H a subgroup of G, then $\bigcap_{p \in \pi(H)} N_{\mathscr{U}_p}(G, H) \leq N_{\mathscr{U}}(G, H).$

Proof. Denote $T = \bigcap_{p \in \pi(G)} N_{\mathscr{U}_p}(G, H)$, then for any subgroup K of H, by definition, T normalize $K^{\mathscr{U}_p}$ for every $p \in \pi(K)$. Now by Lemma 2.10, T normalize $K^{\mathscr{U}}$ for any $K \leq H$, so $T \leq N_{\mathscr{U}}(G, H)$, as desired.

Lemma 3.12. Let G be a group and H a subgroup of G. If $H^{\mathscr{U}}$ is a p-group for some $p \in \pi(H)$, where $\pi(H) = \{p_1, p_2, \cdots, p_n\}$, then $\bigcap_{p_i \in \pi(H)} N_{\mathscr{U}_{p_i}}(G, H) = N_{\mathscr{U}}(G, H)$.

Proof. By Lemma 3.11, we only need to prove $N_{\mathscr{U}}(G, H) \leq \bigcap_{p_i \in \pi(H)} N_{\mathscr{U}_{p_i}}(G, H)$. Without loss of generality, we assume $p = p_1$. By Lemma 2.10, $H^{\mathscr{U}_{p_i}}$ is a *p*-group for every $p_i \in \pi(H)$. Now if $j \neq 1$, the p_j -supersolvability of $H/H^{\mathscr{U}_{p_j}}$ implies that $H \in \mathscr{U}_{p_j}$. Thus by Lemma 2.10, $K^{\mathscr{U}} = K^{\mathscr{U}_p}$ for every subgroup K of H and $G = N_{\mathscr{U}_{p_i}}(G, H)$ for every $p \neq p_j \in \pi(H)$, so $\bigcap_{p_i \in \pi(H)} N_{\mathscr{U}_{p_i}}(G, H) = N_{\mathscr{U}_p}(G, H) = N_{\mathscr{U}}(G, H)$, as desired.

Theorem 3.13. Let G be a solvable group and H a normal subgroup of G, then $\bigcap_{p \in \pi(H)} N_{\mathscr{U}_p}(G, H) = G \text{ if and only if } N_{\mathscr{U}}(G, H) = G.$

Proof. Denote $\pi(H) = \{p_1, p_2, \cdots, p_n\}$, the " \Rightarrow " part of this result follows from Lemma 3.11. Now we prove the " \Leftarrow " part, this is equivalent to prove that $N_{\mathscr{U}_{p_i}}(G, H) =$ G for arbitrary $p_i \in \pi(H)$. Assume false, then there is at least one prime, say p in $\pi(H)$ such that $N_{\mathscr{U}_p}(G,H) < G$. If $p \notin \pi(H^{\mathscr{U}_p})$, then the *p*-supersolvability of $H/H^{\mathscr{U}_p}$ implies that $H \in \mathscr{U}_p$, then $H^{\mathscr{U}_p} = 1$, so $G = N_{\mathscr{U}_p}(G, H)$, a contradiction. By Lemma 2.2(1), $H = N_{\mathscr{U}}(H)$. So $H^{\mathscr{U}}$ is nilpotent by [5, Theorem B]. Note that $p \in \pi(H^{\mathscr{U}})$, so there is a minimal normal subgroup N of G contained in $O_p(H^{\mathscr{U}})$. By Lemma 2.1, $G/N = N_{\mathscr{U}}(G/N, H/N)$. By induction on |G||H|, we have $G/N = N_{\mathscr{U}_{p_i}}(G/N, H/N)$ for any $p_i \in \pi(H)$. Now by Proposition 3.2, $(H/N)^{\mathscr{U}_{p_i}}$ is a p_i -group for any $p_i \in \pi(H)$. In particular, $H^{\mathscr{U}_p}$ is a p-group. If there exists some $p \neq p_i \in \pi(H)$ such that $H^{\mathscr{U}_{p_i}} \neq 1$, then as a similar proof above, there is a minimal normal subgroup T of G contained in $O_{p_i}(H^{\mathscr{U}_{p_i}})$ such that $G/T = N_{\mathscr{U}_p}(G/T, H/T)$. For any subgroup K of H, we have $(KT/T)^{\mathscr{U}_p} \trianglelefteq G/T$, so $K^{\mathscr{U}_p}T \trianglelefteq G$. Note that $[K^{\mathscr{U}_p},T] \leq [H^{\mathscr{U}_p},T] \leq H^{\mathscr{U}_p} \cap T = 1$ since $p \neq p_i$, hence $K^{\mathscr{U}_p}$ char $K^{\mathscr{U}_p}T \leq G$, which implies that $K^{\mathscr{U}_p} \trianglelefteq G$. This is also contrary to that $G = N_{\mathscr{U}_p}(G, H)$. So $H^{\mathscr{U}_{p_i}} = 1$ for any $p \neq p_i \in \pi(H)$, thus by Lemma 2.10, $H^{\mathscr{U}} = H^{\mathscr{U}_{p_1}} H^{\mathscr{U}_{p_2}} \cdots H^{\mathscr{U}_{p_n}} = H^{\mathscr{U}_p}$ is a *p*-group, now by Lemma 3.12, we have $\bigcap_{p_i \in \pi(H)} N_{\mathscr{U}_{p_i}}(G, H) = N_{\mathscr{U}}(G, H)$, a contradiction.

Theorem 3.14. Let G be a solvable group and H a normal subgroup of G, then $\bigcap_{p \in \pi(H)} N_{\mathscr{U}_p}(G, H) = 1 \text{ if and only if } N_{\mathscr{U}}(G, H) = 1.$

Proof. Assume that $N_{\mathscr{U}}(G, H) = 1$, then by Lemma 3.11, $\bigcap_{p \in \pi(H)} N_{\mathscr{U}_p}(G, H) = 1$. Now assume that $\bigcap_{p \in \pi(H)} N_{\mathscr{U}_p}(G, H) = 1$, since $C_G(H^{\mathscr{U}}) \leq C_G(H^{\mathscr{U}_p}) \leq N_{\mathscr{U}_p}(G, H)$, so $1 = C_G(H^{\mathscr{U}}) \leq \bigcap_{p \in \pi(H)} N_{\mathscr{U}_p}(G, H)$. By Corollary 3.5, $N_{\mathscr{U}}(G, H) = 1$.

4 \mathcal{N}_1 -group

Let G be a group and H a subgroup of G. We now consider the case that $G = N_{\mathscr{U}_p}(G, H)$, i.e., G is an \mathscr{N}_1 -group with respect to H and \mathscr{U}_p . Obviously, G is an

 \mathcal{N}_1 -group with respect to H and \mathcal{U}_p if and only if $K^{\mathcal{U}_p} \trianglelefteq G$ for every $K \le H$. There are many groups which are \mathcal{N}_1 -groups, here we list some of them.

Proposition 4.1. The following groups are \mathcal{N}_1 -groups with respect to some subgroup H and \mathcal{U}_p :

(1) All (p-)supersolvable groups.

(2) Groups with H a (p-)supersolvable subgroup.

(3) Groups all of whose non-(p-)supersolvable subgroups are normal.

(4) Groups with $G^{\mathscr{U}_p}$ a cyclic subgroup.

(5) Groups with a normal minimal non-p-supersolvable subgroup H.

Proposition 4.2. Let G be an \mathcal{N}_1 -group with respect to H and \mathcal{U}_p , then

(1) For any $K \leq G$, K is an \mathcal{N}_1 -group with respect to $H \cap K$ and \mathcal{U}_p ;

(2) Let $N \leq G$ and $N \leq H$, then G/N is an \mathcal{N}_1 -group with respect to H/N and \mathcal{U}_p ;

Proof. The proof of (1) and (2) follows from Lemmas 2.1(2)(3) respectively.

In general, if both of a normal subgroup N and corresponding quotient group G/N are \mathscr{N}_1 -groups with respect to themselves and \mathscr{U}_p , then G may be not an \mathscr{N}_1 -group.

Example 4.3. Let $G = \langle a, b, c, d \rangle \rtimes \langle e, f \rangle \cong C_2^4 \rtimes C_6$, where $e^2 = f^3 = 1$ and $a^e = ac, b^e = bd, c^e = c, d^e = d, a^f = b, b^f = ab, c^f = d, d^f = cd$ (IdGroup=[96,70] in GAP [15]). Let $N = \langle a, b, c, d, e \rangle$, then $N = N_{\mathscr{U}_2}(N)$ and $G/N = N_{\mathscr{U}_2}(G/N)$. Now let $H = \langle a, b, f \rangle \cong A_4$, the alternating group of degree 4, then $H^{\mathscr{U}_2} = \langle a, b \rangle$ is not normal in G, so G is not an \mathscr{N}_1 -group with respect to itself and \mathscr{U}_2 .

In the case $\mathscr{F} \in {\mathscr{A}, \mathscr{N}}$, the solvability of $N_{\mathscr{F}}(G)$ has been investigated by many scholars. For example, see [6, 7]. For the case $\mathscr{F} = \mathscr{U}$, we have

Theorem 4.4. Let G be an \mathcal{N}_1 -group with respect to itself and \mathscr{U} . Then G is solvable. In particular, $N_{\mathscr{U}}(G)$ and $N_{\mathscr{U}}^{\infty}$ are solvable.

Proof. Obviously, every subgroup and every homomorphic image of G satisfies our hypothesis. If G is not solvable, then G is a non-abelian simple group. Let H be a proper subgroup of G, then $H^{\mathscr{U}} \leq G$, so $H^{\mathscr{U}} = 1$ and H is supersolvable. This implies that G is a minimal non-supersolvable group. Now by Doerk's result [19], G is solvable, a contradiction.

Let H be a subgroup of G, for the case of $\mathscr{F} = \mathscr{U}_p$, the solvability of $N_{\mathscr{U}_p}(G, H)$ does not always holds. Here, we have

Theorem 4.5. Let G be a group and H a subgroup of G, if G is an \mathcal{N}_1 -group with respect to H and \mathcal{U}_p , where p is the smallest prime in $\pi(H)$, then H is p-solvable.

Proof. Assume that the result is false and let the pair (G, H) be a counterexample with |G||H| minimal. By Feit-Thompson's theorem, we may assume that p = 2. If H < G, then by Lemma 2.1(2), $H = N_{\mathscr{U}_2}(H, H) = N_{\mathscr{U}_2}(H)$, so H is 2-solvable by the choice of (G, H), a contradiction. Now we assume that H = G, by hypothesis, $G = N_{\mathscr{U}_2}(G)$. If G is a non-abelian simple group, we prove the contradiction that G is a solvable simple group. Let K be any proper subgroup of G, then $K^{\mathscr{U}_2} \leq G$, so $K^{\mathscr{U}_2} = 1$ and K is 2-supersolvable. Which implies that G is a non-2-supersolvable group all of whose proper subgroups are 2-supersolvable. Now by [20, 1.1], G is solvable, a contradiction. Let N be a minimal normal subgroup of G, then Lemma 2.2 implies that $G/N = N_{\mathscr{U}_2}(G/N)$ and $N = N_{\mathscr{U}_2}(N)$, thus G/N and N are 2-solvable by our choice of (G, H), hence G is 2-solvable, the finial contradiction complement our proof. □

Corollary 4.6. Let G be a group and p is the smallest prime in $\pi(G)$. If G is an \mathcal{N}_1 -group with respect to itself and \mathcal{U}_p , then G is p-solvable.

In Theorem 4.5, the restriction that p is smallest prime in $\pi(H)$ is necessary. Furthermore, in general, we can't obtain the *p*-solvability of G even though p = 2.

Example 4.7. Let G = SL(2,5)(IdGroup=[120,5]), see [15]), then $\pi(G) = \{2,3,5\}$ and G is not 2-solvable and not 3-solvable. But $G = N_{\mathscr{U}_3}(G)$.

Furthermore, G has a normal subgroup, say $H \cong C_2$, obviously, $G = N_{\mathscr{U}_2}(G, H)$. But we have the following obvious conclusions:

Theorem 4.8. Let G be a group and H a subgroup of G. Assume G is an \mathcal{N}_1 -group with respect to H and \mathcal{U}_p , where p is the smallest prime divisor of |H|. Then G is p-solvable if one of the statement holds:

(1) $H \trianglelefteq G$ and G/H is p-solvable; (2) $G^{\mathscr{U}_p} \le H$;

Theorem 4.9. Let G be a group and P a Sylow p-subgroup of G. Assume that $P \trianglelefteq G$ and G is an \mathcal{N}_1 -group with itself and \mathcal{U}_p , then G is solvable.

Theorem 4.10. Let G be a p-solvable \mathcal{N}_1 -group with respect to itself and \mathcal{U}_p , then

(1) G^{U_p} is a p-group.
(2) N_{U_p}(G/N) > 1 for any proper normal subgroup N of G.
(3) h_p(G) ≤ 3;

Proof. (1) It follows from Proposition 3.2.

(2) Let N be a proper normal subgroup of G, by Lemma 2.2(2), $G/N = N_{\mathscr{U}_p}(G/N)$, so $N_{\mathscr{U}_p}(G/N) > 1$ by our choice of N.

(3) By (1), $G^{\mathscr{U}_p}$ is a *p*-group, then $G^{\mathscr{U}_p} \leq F_p(G)$. Consider $\overline{G} = G/F_p(G)$, then \overline{G} is *p*-supersolvable, thus $\overline{G}' = G'F_p(G)/F_p(G)$ is *p*-nilpotent. So $h_p(G) \leq h_p(\overline{G}) + 1 \leq h_p(\overline{G}/\overline{G}') + h_p(\overline{G}') + 1 = 3$.

As a corollary of Theorem 4.10, we have

Corollary 4.11. Let G be a \mathcal{N}_1 -group with respect to itself and \mathcal{U}_p for every $p \in \pi(G)$, then

(1) G is solvable.
(2) G = N_𝔅(G).
(3) G^𝔅 is nilpotent.
(4) G/N ∈ 𝔊𝔅 for every normal subgroup N of G.
(5) h(G) ≤ 3.
(6) l_r(G) ≤ 2 for every r ∈ π(G).

Proof. (1) Let $p \in \pi(G)$ be the smallest prime, if p > 2, obviously G is solvable by the well know Feit-Thompson Theorem. If p = 2, then by Theorem 4.5, G is 2-solvable, so G is solvable, as desired.

(2) By hypothesis, $G = N_{\mathscr{U}_p}(G)$ for every $p \in \pi(G)$. Now by Theorem 3.13, $G = N_{\mathscr{U}}(G)$.

(3) Let $p \in \pi(G)$, if $G \in \mathscr{U}_p$, then $G^{\mathscr{U}_p} = 1$, if $G \notin \mathscr{U}_p$, by Theorem 4.10(1), $G^{\mathscr{U}_p}$ is a p-group. Now assume $\pi(G) = \{p_1, p_2, \cdots, p_n\}$, then by Lemma 2.10, $G^{\mathscr{U}} = G^{\mathscr{U}_{p_1}}G^{\mathscr{U}_{p_2}}\cdots G^{\mathscr{U}_{p_n}}$, so $G^{\mathscr{U}}$ is nilpotent.

(4) By (2), $G^{\mathscr{U}}$ is nilpotent. Let N be a proper normal subgroup of G, then $(G/N)^{\mathscr{U}} = G^{\mathscr{U}}N/N \cong G^{\mathscr{U}}/G^{\mathscr{U}} \cap N$ is nilpotent, as desired.

(5) It follows from Theorem 4.10(3).

(6) If $O_{r'}(G) > 1$, then $N_{\mathscr{U}_p}(G/O_{r'}(G)) = G/O_{r'}(G)$ for every $p \in \pi(G)$ by Lemma 2.2(2), so $G/O_{r'}(G)$ satisfies our hypothesis. By induction, $l_r(G/O_{r'}(G)) \leq 2$, which implies that $l_r(G) \leq 2$. As a similarly argument above and by Lemma 2.4, we have $l_r(G) = l_r(G/\Phi(G)) \leq 2$, hence we may assume that $O_{r'}(G) = \Phi(G) = 1$, so $F(G) = F_r(G) = O_r(G)$. Now consider G/F(G), by (5), $h(G/F(G)) \leq 2$. Now by Lemma 2.7, $l_r(G/F(G)) = l_r(G/O_r(G)) \leq 1$, so $l_r(G) \leq l_r(G/O_r(G)) + 1 \leq 2$, as desired.

Recall that the well-know results of P.Hall and D.J.S. Robinson as follows:

Theorem 4.12. Let G be a group and N a nilpotent normal subgroup of G. Then (1) (P. Hall) If G/N' is nilpotent, then G is nilpotent.

(2) (D.J.S. Robinson). If G/N' is supersolvable, then G is supersolvable.

Here we have

Theorem 4.13. Let G be a p-solvable \mathcal{N}_1 -group with respect to itself and \mathcal{U}_p . Then $G \in \mathcal{N}$ if $G^{\mathcal{N}} \leq G^{\mathcal{U}_p^2}$.

Theorem 4.14. Let G be an \mathcal{N}_1 -group with respect to itself and \mathscr{U} . Then $G \in \mathscr{U}$ if $G^{\mathscr{U}} \leq G^{\mathscr{U}^2}$.

5 Applications

Theorem 5.1. Let G be a p-solvable group, then the following statements are equivalent:

(1) $G \in \mathcal{N}_p \mathcal{U}_p.$ (2) $G/N_{\mathcal{U}_p}(G) \in \mathcal{N}_p \mathcal{U}_p.$

Proof. $(1) \Rightarrow (2)$. It is obviously.

 $(2) \Rightarrow (1)$. Assume the result is false and let G be a counterexample of minimal order. If $N_{\mathscr{U}_p}(G) = 1$, there is nothing to prove. Now we assume that $N_{\mathscr{U}_p}(G) > 1$. Let N be a minimal normal subgroup of G contained in $N_{\mathscr{U}_p}(G)$. Then either N is an elementary abelian p-group or a p'-group. Note that $G/N/N_{\mathscr{U}_p}(G/N) \leq G/N/N_{\mathscr{U}_p}(G)/N \cong G/N_{\mathscr{U}_p}(G)$ by Lemma 2.2(2), then $G/N \in \mathscr{N}_p \mathscr{U}_p$ by the choice of G. So we may assume that N is an elementary abelian p-group, otherwise, $G^{\mathscr{U}_p}N/N \cong G^{\mathscr{U}_p}/(G^{\mathscr{U}_p} \cap N) \in \mathscr{N}_p$, so $G \in \mathscr{N}_p \mathscr{U}_p$ by Lemma 2.5(2), a contradiction. Furthermore, if $N \leq \Phi(G)$, then $G/\Phi(G) \in \mathscr{N}_p \mathscr{U}_p$, so $G^{\mathscr{U}_p}\Phi(G)/\Phi(G) \in \mathscr{N}_p$, hence $G \in \mathscr{N}_p \mathscr{U}_p$ by Lemma 2.5(1), a contradiction again. Now there exist some maximal subgroup Mof G such that G = MN and $M \cap N = 1$. Also we have $M^{\mathscr{U}_p} = (G/N)^{\mathscr{U}_p} \in \mathscr{N}_p$. Note that $N \leq N_{\mathscr{U}_p}(G)$, then N normalize $M^{\mathscr{U}_p}$, thus $[N, M^{\mathscr{U}_p}] = 1$ and $NM^{\mathscr{U}_p} = N \times M^{\mathscr{U}_p} \leq G$. Since $G/NM^{\mathscr{U}_p} = MN/NM^{\mathscr{U}_p} = MNM^{\mathscr{U}_p}/NM^{\mathscr{U}_p} \cong M/M \cap NM^{\mathscr{U}_p} = M/M^{\mathscr{U}_p} \in \mathscr{U}_p$, then $G^{\mathscr{U}_p} \leq NM^{\mathscr{U}_p} \in \mathscr{N}_p$, a contradiction.

Corollary 5.2. Let G be a p-solvable group, if $G = N^{\infty}_{\mathscr{U}_{p}}(G)$, then (1) $G \in \mathscr{N}_{p}\mathscr{U}_{p}$. (2) $N^{\infty}_{\mathscr{U}_{p}}(G/N) > 1$ for any proper normal subgroup N of G. (3) $h_{p}(G) \leq 3$;

Proof. (1) Set $G = N^n_{\mathscr{U}_p}(G)$ for some positive integer n. If n = 1, the result follows from Theorem 4.10(1). If $n \geq 2$, then $N_{\mathscr{U}_p}(G) < G$. By Lemma 2.3, $G/N_{\mathscr{U}_p}(G) = N^{\infty}_{\mathscr{U}_p}(G/N_{\mathscr{U}_p}(G))$, so $G/N_{\mathscr{U}_p}(G)$ satisfies our hypothesis, thus $G/N_{\mathscr{U}_p}(G) \in \mathscr{N}_p \mathscr{U}_p$ by induction on |G|. Now by Theorem 5.1, $G \in \mathscr{N}_p \mathscr{U}_p$, as desired. (2) If there exists some proper normal subgroup N of G such that $N^{\infty}_{\mathscr{U}_p}(G/N) = 1$, then $G = N^{\infty}_{\mathscr{U}_p}(G) \leq N$ by Lemma 2.3(3), that is impossible.

The proof of (3) is similar to Theorem 4.10(3).

We have proved in Theorem 4.4 that G is solvable if $G = N^{\infty}_{\mathscr{U}}(G)$. Now as similar argument above, we have

Corollary 5.3. Let G be a group, if $G = N^{\infty}_{\mathscr{Y}}(G)$, then

- (1) $G \in \mathcal{N} \mathcal{U}$.
- (2) $h(G) \le 3$.

(3) $l_p(G) \leq 2$ for every $p \in \pi(G)$.

Theorem 5.4. Let G be a group and $p \in \pi(G)$.

(1) Assume G is p-solvable, then $h_p(G) \leq k$ if and only if $h_p(G/N^{\infty}_{\mathscr{U}_p}(G)) \leq k$, where $k \geq 3$.

(2) Assume G is solvable, then $l_p(G) \leq k$ if and only if $l_p(G/N^{\infty}_{\mathscr{U}}(G)) \leq k$, where $k \geq 2$.

Proof. We only prove (1) since the proof of (2) is similar and based on Corollary 5.3(4). The " \Rightarrow " part of this theorem is obviously, so we prove the " \Leftarrow " part. Setting $N^{\infty}_{\mathscr{U}_p}(G) = N^n_{\mathscr{U}_p}(G)$ for some positive integer n. If $G = N^n_{\mathscr{U}_p}(G)$, then the result follows form Corollary 5.2(3). So we assume that $N^n_{\mathscr{U}_p}(G) < G$.

Assume that the result is false and let the pair (G, n) be a counterexample with n|G| minimal. For the case n = 1, we have $h_p(G/N_{\mathscr{U}_p}(G)) \leq k$ by hypothesis. Let N be a minimal normal subgroup of G, then by Lemma 2.2(2), $G/N/N_{\mathscr{U}_p}(G/N) \leq$ $G/N/N_{\mathscr{U}_p}(G)N/N \cong G/N_{\mathscr{U}_p}(G)N \cong G/N_{\mathscr{U}_p}(G)/N_{\mathscr{U}_p}(G)N/N_{\mathscr{U}_p}(G), \text{ so } h_p(G/N/N_{\mathscr{U}_p}(G/N)) \leq 1$ k by Lemma 2.6(2), thus $h_p(G/N) \leq k$ by the choice of (G, 1). Suppose that N is not unique, that is G has at least two minimal normal subgroups, say N_1 , N_2 and $N_1 \neq N_2$, then as above, $h_p(G/N_1), h_p(G/N_2) \leq k$, so $h_p(G) \leq k$ by Lemma 2.6(4), a contradiction. With similar argument and by using of Lemma 2.6(5), we have $\Phi(G) = 1$. Furthermore, if $O_{p'}(G) > 1$, then it follows from Lemma 2.6(6) that $h_p(G/O_{p'}(G)) = h_p(G) \leq k$, a contradiction. So $F_p(G) = F(G) = C_G(N) = N \leq C_G(N)$ $N_{\mathscr{U}_p}(G)$. Therefore, $G = MN, M \cap N = 1$ for some maximal subgroup M of G. If $M^{\mathscr{U}_p} = 1$, that is G/N is p-supersolvable, so $G/N = N_{\mathscr{U}_p}(G/N) = N_{\mathscr{U}_p}^{\infty}(G/N) =$ $N^{\infty}_{\mathscr{U}_p}(G)/N$, thus $G = N^{\infty}_{\mathscr{U}_p}(G)$, a contradiction. If $M^{\mathscr{U}_p} > 1$, then N normalize $M^{\mathscr{U}_p}$, so $[N, M^{\mathscr{U}_p}] = 1$ and hence $M^{\mathscr{U}_p} \leq C_G(N) = N, M^{\mathscr{U}_p} \leq N \cap M = 1$, a contradiction again. Now assume that n > 1, since $G/N^n_{\mathscr{U}_p}(G) \cong G/N^{n-1}_{\mathscr{U}_p}(G)/N^n_{\mathscr{U}_p}(G)/N^{n-1}_{\mathscr{U}_p}(G) =$ $G/N^{n-1}_{\mathscr{U}_p}(G)/N_{\mathscr{U}_p}(G/N^{n-1}_{\mathscr{U}_p}(G))$, so by induction and combining the proof of the case n = 1, we have that $h_p(G/N^{n-1}_{\mathscr{U}_p}(G)) \leq k$, hence $h_p(G) \leq k$ by our choice of (G, n), a contradiction.

Theorem 5.5. Let G be a p-solvable group. If all elements of order p and order 4(if p = 2) of G are in $N^{\infty}_{\mathscr{U}_{p}}(G)$, then $h_{p}(G) \leq 3$.

Proof. Assume that the result is false and let G be a counterexample with minimal order, we prove this result by several steps:

Step 1. Let H < G, then $h_p(H) \leq 3$.

Let H be a proper subgroup of G, then all elements of H of order p and order 4(if p = 2) are contained in $H \cap N^{\infty}_{\mathscr{U}_p}(G) \leq N^{\infty}_{\mathscr{U}_p}(H)$ by Lemma 2.3(1), so H satisfies our hypothesis, $h_p(H) \leq 3$ by the choice of G.

Step 2. $O_{p'}(G) = 1$, $C_G(O_p(G)) \le O_p(G) = F(G)$.

Let $T = O_{p'}(G)$, if $T \neq 1$, for any $xT \in N^{\infty}_{\mathscr{U}_p}(G)T/T$ with |x| = p or |x| = 4 (if p = 2), $xT \leq N^{\infty}_{\mathscr{U}_p}(G/T)$ by Lemma 2.3(3), so G/T satisfies our hypothesis, thus $h_p(G/T) \leq 3$ by the choice of G. It is easy to see that $h_p(G) \leq 3$, a contradiction. Further more, we have $C_G(O_p(G)) \leq O_p(G) = F(G)$ since G is p-solvable and $O_{p'}(G) = 1$.

Step 3. $\Phi(G) < O_p(G)$.

By (2), $\Phi(G) \leq O_p(G)$, if $O_p(G) = \Phi(G)$, then $O_{pp'}(G)/\Phi(G) = O_{p'}(G/\Phi(G)) > 1$ since G is p-solvable and $O_p(G/\Phi(G)) = 1$. Let $O_{pp'}(G) = \Phi(G)T$, where T > 1 is a p'-group, then $G = N_G(T)O_{pp'}(G) = N_G(T)$ by Frattini argument, it follows that $T \leq G$ and then $T \leq O_{p'}(G) = 1$ by **Step** 2, a contradiction.

Step 4. Finial contradiction.

Denote by \mathscr{F}_p the class of all *p*-solvable groups whose *p*-Fitting length are less than 3, then by Lemma 2.6, \mathscr{F}_p is a saturated formation. Now by **Step** 1, *G* is an \mathscr{F}_p -critical group, i.e, *G* is not belongs to \mathscr{F}_p , but all proper subgroups of *G* belong to \mathscr{F}_p . Now by **Steps** 1,3, there exists some maximal subgroup *M* of *G* such that $G = O_p(G)M = F(G)M$ and $M \in \mathscr{F}_p$, so by Lemma 2.8(2), $G^{\mathscr{F}_p}$ is a *p*-group and $G^{\mathscr{F}_p}$ has exponent *p* if p > 2 and exponent at most 4 if p = 2, hence $G^{\mathscr{F}_p} \leq O_p(G)$ and $G^{\mathscr{F}_p} \leq N^{\infty}_{\mathscr{U}_p}(G)$, it follows that $h_p(G/N^{\infty}_{\mathscr{U}_p}(G)) \leq 3$. Now by Theorem 5.4(2), $h_p(G) \leq 3$, a contradiction.

Corollary 5.6. Let G be a p-solvable group, if every cyclic subgroup of order prime p and order 4(if p = 2) of G is contained in $N_{\mathscr{H}_p}(G)$, then $h_p(G) \leq 3$.

Combining Theorem 5.4, Lemma 2.4 and Lemma 2.8, we have the following Theorem 5.7. It's proof is similar to Theorem 5.5.

Theorem 5.7. Let G be a solvable group, if every cyclic subgroup of order prime p and order 4(if p = 2) of G is contained in $N^{\infty}_{\mathscr{U}}(G)$, then $l_p(G) \leq 2$.

As a local vision of Theorem 5.7, we have the following question.

Question 5.8. Let G be a p-solvable group, if every cyclic subgroup of order prime p and order 4(if p = 2) of G is contained in $N^{\infty}_{\mathscr{U}_p}(G)$, does $l_p(G) \leq 2$?

Remark 5.9. (1) In Theorem 5.5, Corollary 5.6 and Question 5.8, the Theorem 4.5 implies the hypothesises that G is p-solvable are necessary. For example, consider $G = A_5$, the alternating group of degree 5, then $G = N_{\mathscr{U}_3}(G) = N_{\mathscr{U}_5}(G)$, so G satisfies our condition, but G is simple.

(2) In Theorem 5.7, the solvability of G is necessary. For example, choice the non-solvable group $G = C_7 \times A_5(IdGroup=[420,13])$ and let p = 7. Then $N_{\mathscr{U}}(G) = N_{\mathscr{U}}^{\infty}(G) = C_7$ and every cyclic subgroup of order 7 is in C_7 .

(3) In Question 5.8, the integer 2 is the minimum upper bound. For example, let $G = C_5 \times S_4(IdGroup=[120,37])$ in GAP). Obviously, G is 2-solvable and the proper non-2-supersolvable groups of G isomorphism to one of $A_4, S_4, C_5 \times A_4$. As $A_4, S_4, C_5 \times A_4$ are normal in G, so $G = N_{\mathscr{U}_2}(G)$. Note that $1 < C_5 < C_{10} \times C_2 < C_5 \times A_4 < C_5 \times S_4 = G$ is the upper 2-series of G, so $l_2(G) = 2$.

References

- [1] B. Huppert. (1967). Endliche Gruppen, Springer, Berlin-Heidelberg-New York.
- [2] R. Baer. (1956). Norm and hypernorm, Publ. Math. Debrecen 4:347-356.
- [3] R. Baer, Der Kern. (1935). eine charkteristishe Untergruppe, Compos. Math. 1:254-283.
- [4] Su Ning, Wang Yanming.(2013). On the normalizers of ℱ-residuals of all subgroups of a finite group, Journal of Algebra, 185-198.
- [5] Su, Ning, Wang, Yanming.(2014). On the Intersection of the Normalizers of the *F*-Residuals of Subgroups of a Finite Group. Algebras and Representation Theory. 17:507-518.
- [6] Li, Shirong, Shen, Zhencai. (2010). On the intersection of the normalizers of derived subgroups of all subgroups of a finite group. Journal of Algebra. 323:1349-1357.
- [7] Shen Zhencai, Shi Wujie, Qian Guohua. (2012). On the norm of the nilpotent residuals of all subgroups of a finite group, Journal of Algebra. 352:290-298.
- [8] Gong, Lü, Guo, Xiuyun. (2013). On the Intersection of the Normalizers of the Nilpotent Residuals of All Subgroups of a Finite Group. Algebra Colloquium. 20(2):349-360.
- [9] Li, Xuan, Guo, Xiuyun.(2015). On the normalizers of *p*-nilpotency-residuals of all subgroups in a finite group. Journal of Algebra and Its Applications. 14(10).1550146.
- [10] Shen, Zhencai, Li, Shirong, Shi, Wujie. (2014). On the norm of the derived subgroups of all subgroups of a finite group. Bulletin of the Iranian Mathematical Society. 40:281-291.

- [11] Li, Xuan, Guo, Xiuyun.(2016).On Generalized Norms of Finite Groups. Communications in Algebra. 44:1088-1095.
- [12] Wang, Junxin, Guo, Xiuyun.(2007). On the Norm of Finite Groups. Algebra Colloquium. 14(4):605-612.
- [13] Hu, Bin, Huang, Jianhong, Skiba, Alexander. (2020). On the σ -nilpotent norm and the σ -nilpotnet length of a finite group. Glasgow Mathematical Journal. 63:1-12.
- [14] Weinstein, M., Between Nilpotent and Solvable, Passaic/New York/Jersey:Polygonal Publishing House, 1982.
- [15] The GAP Group: GAPgroups, algorithms, and programming.(2020). Version 4.11.0. http://www.gap-system.org.
- [16] A. N. Skiba.(2015). On σ -subnormal and σ -permutable subgroups of finite groups, Journal of Algebra. 436:1-16.
- [17] E. Jabara.(2018). The fitting length of a product of mutually permutable finite groups. Acta Math. Hungar.
- [18] A. Ballester-Bolinches, M.C.Pedraza-Aguilera.(1996). On minimal subgroups of finite groups, Acta Math. Hungar.73(4):335-342.
- [19] Doerk, K.(1966). Minimal nicht überauflösbare, endliche Gruppen. Math. Z. 91, 198C205.
- [20] Tuccillo F.(1992). On finite minimal non-*p*-supersoluble groups. Colloquium Mathematicum,63:119-131.