

**A NUMERICAL CHARACTERIZATION OF THE  
EXTREMAL BETTI NUMBERS OF  $t$ -SPREAD STRONGLY  
STABLE IDEALS**

Luca Amata, Antonino Ficarra, Marilena Crupi

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCES, PHYSICS AND EARTH  
SCIENCES, UNIVERSITY OF MESSINA, VIALE FERDINANDO STAGNO D'ALCONTRES 31,  
98166 MESSINA, ITALY

*E-mail addresses:* lamata@unime.it; antficarra@unime.it; mcrupi@unime.it

**Abstract**

Let  $K$  be a field and let  $S = K[x_1, \dots, x_n]$  be a standard polynomial ring over a field  $K$ . We characterize the extremal Betti numbers, values as well positions, of a  $t$ -spread strongly stable ideal of  $S$ . Our approach is constructive. Indeed, given some positive integers  $a_1, \dots, a_r$  and some pairs of positive integers  $(k_1, \ell_1), \dots, (k_r, \ell_r)$ , we are able to determine under which conditions there exist a  $t$ -spread strongly stable ideal  $I$  of  $S$  with  $\beta_{k_i, k_i \ell_i}(I) = a_i$ ,  $i = 1, \dots, r$ , as extremal Betti numbers, and then to construct it.

1. INTRODUCTION

Let  $K$  be a field and let  $S = K[x_1, \dots, x_n]$  be a polynomial ring. A square-free monomial ideal of  $S$ , known also as Stanley–Reisner ideal, is a monomial ideal generated by squarefree monomials. The intimate relationship of this algebraic object with simplicial topology determines its combinatorial nature. The importance of such a class of monomial ideals is due to Richard Stanley. In fact, in 1975, Stanley used the theory of Cohen–Macaulay rings to solve the upper bound conjecture for spheres [20]. More in details, he used tools from commutative algebra to study simplicial complexes by considering the Hilbert function of Stanley–Reisner rings, whose defining ideals are squarefree ideals.

Recently, a generalization of the notion of squarefree ideal has been given in [16] by the definition of  $t$ -spread monomial ideal. Let  $t \geq 0$  an integer and let  $u = x_{i_1} x_{i_2} \cdots x_{i_d}$  be a monomial of  $S$ , with  $1 \leq i_1 \leq i_2 \leq \cdots \leq i_d \leq n$ .  $u$

---

*Keywords:* monomial ideals, minimal graded resolution, extremal Betti numbers,  $t$ -spread ideals.

*2020 Mathematics Subject Classification:* 05E40, 13B25, 13D02, 16W50, 68W30.

is called  $t$ -spread if  $i_{j+1} - i_j \geq t$ , for all  $j = 1, \dots, d-1$ . A monomial ideal  $I$  is a  $t$ -spread monomial ideal if it is generated by  $t$ -spread monomials. Clearly, when  $t = 1$ , a 1-spread monomial ideal is a squarefree monomial ideal. Many results which are true for squarefree ideals can be suitably rewritten for the class of  $t$ -spread ideals. For example, the classical Kruskal–Katona theorem that characterizes the  $f$ -vectors of simplicial complexes has been generalized by C. Andrei–Ciobanu for the class of  $t$ -spread strongly stable ideals [7].

Among the algebraic invariants of a graded ideal  $I$  of  $S$ , the graded Betti numbers  $\beta_{i,j}(I)$  are probably the most important: the integer  $\beta_{i,j}(I)$  gives the dimension, as a  $K$ -vector space, of the  $j$ -th component of the  $i$ -th free module in the minimal graded free resolution of  $I$ . They are usually displayed in a table called the *Betti table* of  $I$ :

Table 1: Betti table

The columns are indexed from left to right by homological degree starting with homological degree zero. The rows are indexed increasingly from top to bottom starting with the minimal degree of an element in a minimal system of homogeneous generators of  $I$ . The graded Betti numbers which appear in the outside corners of the dashed line are called extremal Betti numbers. This definition has been introduced by Bayer, Charalambous and Popescu in [9].

In this paper, our attention is devoted to  $t$ -spread strongly stable ideals (Definition 2.2). There is a combinatorial formula [16, Corollary 1.12] that allows us to easily compute the graded Betti numbers of such a class of ideals. Moreover, thanks to this formula, one can give a characterization of these invariants (Theorem 2.7) which is useful for the studying of their behavior. Such a result will be a pivotal tool throughout the paper.

Our goal is to solve the following question:

(\*) *Given  $n, r, t$  positive integers,  $r$  positive integers  $a_1, \dots, a_r$  and  $r$  positive pairs of integers  $(k_1, \ell_1), \dots, (k_r, \ell_r)$ , under which conditions there exist a  $t$ -spread strongly stable ideal  $I$  of  $S = K[x_1, \dots, x_n]$  such that*

$$\beta_{k_1, k_1 + \ell_1}(I) = a_1, \dots, \beta_{k_r, k_r + \ell_r}(I) = a_r$$

are its extremal Betti numbers?

For  $t = 1$ , a positive answer can be found in [3]; whereas in [5] the maximal number  $r$  of extremal Betti numbers allowed for a  $t$ -spread strongly stable ideals, for  $t \geq 2$ , has been computed. In this paper, we are able to give a positive answer to question (\*) for  $t \geq 2$ .

Our approach is mainly constructive. We give a numerical characterization of the extremal Betti numbers of a  $t$ -spread strongly stable ideal and if the ideal that solves the problem does exist, then we provide the combinatorial tools to determine it.

The outline of the paper is the following. Section 2, contains preliminary notions and results. In Section 3, we introduce the notion of  $j$ -gap $_t$  of a  $t$ -spread monomial (Definition 3.7),  $j$  is a positive integer, and the notion of Borel  $t$ -shadow of a set of  $t$ -spread monomials (Definition 3.14). These definitions will have a key role throughout the paper. Section 4 contains a numerical characterization (positions and values) of the extremal Betti numbers of a  $t$ -strongly stable ideal (Theorem 4.4). Finally, Section 5 contains an example which illustrates our methods.

All the examples have been verified using specific packages of *Macaulay2* some of which developed by the authors of the paper.

## 2. PRELIMINARIES

Let  $S = K[x_1, \dots, x_n]$  be the standard polynomial ring in  $n$  variables with coefficients in a field  $K$ .  $S$  is an  $\mathbb{N}$ -graded ring where  $\deg x_i = 1$ , for all  $i = 1, \dots, n$ . A monomial ideal  $I$  of  $S$  is an ideal generated by monomials. If  $I$  is a monomial ideal, we denote by  $G(I)$  the unique minimal set of monomial generators of  $I$  and we set

$$G(I)_\ell = \{u \in G(I) : \deg u = \ell\}.$$

For a monomial  $u \in S$ ,  $u \neq 1$ , we denote by  $\text{supp}(u)$  the set of all index  $i$  for which  $x_i$  divides  $u$  and by  $\max(u)$  ( $\min(u)$ ) the maximal (minimal) index  $i$  for which  $x_i$  divides  $u$ .

If  $u = 1$ , for convenience, we set  $\max(1) = \min(1) = 0$ .

The next definitions have been introduced in [16].

**Definition 2.1** If  $t \geq 0$  is an integer, a monomial  $x_{i_1}x_{i_2} \cdots x_{i_\ell}$  of  $S$  with  $1 \leq i_1 \leq i_2 \leq \dots \leq i_\ell \leq n$  is called  $t$ -spread, if  $i_{j+1} - i_j \geq t$ , for all  $j = 1, \dots, \ell - 1$ . A  $t$ -spread monomial ideal is a monomial ideal generated by  $t$ -spread monomials.

For example, every monomial ideal of the polynomial ring  $S$  is 0-spread and every squarefree monomial ideal is a 1-spread monomial ideal. Furthermore, one can observe that if  $t \geq 1$ , then every  $t$ -spread monomial is a

squarefree monomial.

A special class of  $t$ -spread monomial ideals consists of the  $t$ -spread strongly stable ideals.

**Definition 2.2** A  $t$ -spread strongly stable ideal  $I$  of  $S$  is a  $t$ -spread monomial ideal such that for all  $t$ -spread monomials  $u \in I$  and for all  $i, j$  such that  $1 \leq i < j \leq n$ ,  $x_j$  divides  $u$  and  $x_i(u/x_j)$  is  $t$ -spread, it follows that  $x_i(u/x_j) \in I$ .

**Remark 2.3** One can observe that the defining property of a  $t$ -strongly stable ideal needs to be checked only for the set of monomial generators of a  $t$ -spread monomial ideal [16, Lemma 1.2]. In fact, let  $I$  be a  $t$ -spread monomial ideal and suppose that for all  $u \in G(I)$ , for all integers  $i < j$  with  $j \in \text{supp}(u)$  and such that  $x_i(u/x_j)$  is a  $t$ -spread monomial, one has  $x_i(u/x_j) \in I$ . Then  $I$  is a  $t$ -spread strongly stable ideal.

It is clear that the notion of  $t$ -spread strongly stable ideal generalizes the concept of strongly stable and squarefree strongly stable ideal.

There is an equivalent definition for a  $t$ -spread strongly stable ideal [7, Definition 1.1]. Following the same notations as in [7], we will denote by  $[I_j]_t$  the set of all  $t$ -spread monomials of degree  $j$  of an arbitrary monomial ideal  $I$ .

Let us denote by  $M_{n,d,t}$  the set of all  $t$ -spread monomials of degree  $d$  of the ring  $S$ .

**Definition 2.4** A subset  $L$  of  $M_{n,d,t}$  is called a  $t$ -spread strongly stable set, if for all  $t$ -spread monomials  $u \in L$ , all  $j \in \text{supp}(u)$  and all  $1 \leq i < j$  such that  $x_i(u/x_j)$  is a  $t$ -spread monomial, it follows that  $x_i(u/x_j) \in L$ .

A  $t$ -spread monomial ideal  $I$  is a  $t$ -spread strongly stable ideal if  $[I_j]_t$  is a  $t$ -spread strongly stable set for all  $j$ .

Let  $u_1, \dots, u_r$  be  $t$ -spread monomials of  $S$ . The unique  $t$ -spread strongly stable ideal containing  $u_1, \dots, u_r$  will be denoted by  $B_t(u_1, \dots, u_r)$ , [16]. The monomials  $u_1, \dots, u_r$  are called  $t$ -spread Borel generators.

Furthermore, if  $u_1, \dots, u_r$  are  $t$ -spread monomials of degree  $d$ , the smallest  $t$ -spread strongly stable set of  $M_{n,d,t}$  containing the monomials  $u_1, \dots, u_r$ , will be denoted by  $B_t\{u_1, \dots, u_r\}$ .

Now, for a nonempty subset  $L$  of  $M_{n,d,t}$ , we define the  $t$ -shadow of  $L$

$$\text{Shad}_t(L) = \{x_i w : w \in L \text{ and } x_i w \text{ is } t\text{-spread monomial, } i = 1, \dots, n\}.$$

Throughout the paper, we assume that  $t \geq 1$  and that  $M_{n,d,t}$  is endowed with the *squarefree lexicographic order*,  $>_{\text{slex}}$  [8], i.e., let  $u = x_{i_1} x_{i_2} \cdots x_{i_d}$ ,

$v = x_{j_1}x_{j_2}\cdots x_{j_d}$ , be  $t$ -spread monomials of degree  $d$ , with  $1 \leq i_1 < i_2 < \cdots < i_d \leq n$ , and  $1 \leq j_1 < j_2 < \cdots < j_d \leq n$ , then  $u >_{\text{slex}} v$  if  $i_1 = j_1, \dots, i_{s-1} = j_{s-1}$ , and  $i_s < j_s$ , for some  $1 \leq s \leq d$ .

For a nonempty subset  $T$  of  $M_{n,d,t}$ , we denote by  $\max T$  ( $\min T$ ) the maximal (minimum) monomial  $w \in T$ , with respect to  $>_{\text{slex}}$ .

As for the class of (squarefree) strongly stable ideals of  $S$ , there exists an explicit formula due to Herzog, Ene and Qureshi [16, Corollary 1.12] for the graded Betti numbers of a  $t$ -spread strongly stable ideal given by

$$\beta_{k,k+\ell}(I) = \sum_{u \in G(I)_\ell} \binom{\max(u) - t(\ell - 1) - 1}{k}. \quad (1)$$

In order to apply such a formula, we just need to know the minimal set of monomial generators of  $I$ .

**Remark 2.5** If we put  $t = 0$  in (1), then we obtain the well-known formula of Eliahou and Kervaire [14] for the (strongly) stable ideals; whereas, if we put  $t = 1$  in (1), then we obtain the Aramova, Herzog and Hibi formula for squarefree (strongly) stable ideals [8].

**Definition 2.6** ([9]) A graded Betti number  $\beta_{k,k+\ell}(I) \neq 0$  is called *extremal* if  $\beta_{i,i+j}(I) = 0$  for all  $i \geq k, j \geq \ell, (i, j) \neq (k, \ell)$ .

If  $\beta_{k,k+\ell}(I)$  is extremal, the pair  $(k, \ell)$  is called a *corner* of  $I$ . If  $(k_1, \ell_1), \dots, (k_r, \ell_r)$ , with  $n-1 \geq k_1 > k_2 > \cdots > k_r \geq 1$  and  $1 \leq \ell_1 < \ell_2 < \cdots < \ell_r$ , are all the corners of a graded ideal  $I$  of  $S$ , the set

$$\text{Corn}(I) = \left\{ (k_1, \ell_1), (k_2, \ell_2), \dots, (k_r, \ell_r) \right\}$$

is called the *corner sequence* of  $I$  [10]. The  $r$ -tuple

$$a(I) = (\beta_{k_1, k_1 + \ell_1}(I), \beta_{k_2, k_2 + \ell_2}(I), \dots, \beta_{k_r, k_r + \ell_r}(I))$$

is called the *corner values sequence* of  $I$  [10].

As a consequence of formula (1), in [4] the following two results were stated.

**Theorem 2.7** ([4, Theorem 1]) *Let  $I$  be a  $t$ -spread strongly stable ideal of  $S$ .*

*The following conditions are equivalent:*

- (a)  $\beta_{k,k+\ell}(I)$  is extremal;
- (b)  $k + t(\ell - 1) + 1 = \max \{ \max(u) : u \in G(I)_\ell \}$  and  $\max(u) < k + t(j - 1) + 1$ , for all  $j > \ell$  and for all  $u \in G(I)_j$ .

**Corollary 2.8** ([4, Corollary 2]) *Let  $I$  be a  $t$ -spread strongly stable ideal of  $S$  and let  $\beta_{k,k+\ell}(I)$  be an extremal Betti number of  $I$ . Then*

$$\beta_{k,k+\ell}(I) = \left| \left\{ u \in G(I)_\ell : \max(u) = k + t(\ell - 1) + 1 \right\} \right|.$$

We will see that Corollary 2.8 will be fundamental in order to obtain the numerical characterization of the extremal Betti numbers of a  $t$ -spread strongly stable ideal.

### 3. COMBINATORICS OF $t$ -SPREAD MONOMIALS

In this Section, we generalize some notions and tools from [3]. Our purpose is to suitably manipulate the  $t$ -spread monomials for  $t \geq 1$ .

Let  $t \geq 1$  and  $n$  be a positive integer. Let  $(k, \ell)$  be a pair of positive integers such that  $k + t(\ell - 1) + 1 \leq n$ , we define the set

$$A^t(k, \ell) := \{u \in M_{n,\ell,t} : \max(u) = k + t(\ell - 1) + 1\}.$$

**Remark 3.1** In [3], if  $t = 1$ , the set  $A^1(k, \ell)$  is denoted by  $A^s(k, \ell)$  and consists of all squarefree monomials  $u \in S$  of degree  $\ell$  such that  $\max(u) = k + \ell$ .

**Example 3.2** Let  $t = 3$ ,  $(k, \ell) = (5, 2)$  and  $n \geq k + t(\ell - 1) + 1 = 9$ .

If  $n = 9$ , then

$$A^3(5, 2) = \{u \in M_{9,2,3} : \max(u) = 9\} = \{x_1x_9, x_2x_9, x_3x_9, x_4x_9, x_5x_9, x_6x_9\}$$

From Corollary 2.8, it is clear that in order to obtain a numerical characterization of the extremal Betti numbers of a  $t$ -spread strongly stable ideal  $I$  of  $S$  one has to consider the sets  $A^t(k, \ell)$ .

Indeed, if  $\beta_{k,k+\ell}(I) = a$  is an extremal Betti number of  $I$ , then

$$a = \beta_{k,k+\ell}(I) = \left| \left\{ u \in G(I)_\ell : \max(u) = k + t(\ell - 1) + 1 \right\} \right|.$$

Furthermore, since

$$\{u \in G(I)_\ell : \max(u) = k + t(\ell - 1) + 1\} \subseteq A^t(k, \ell),$$

then

$$a = \beta_{k,k+\ell}(I) \leq |A^t(k, \ell)|.$$

Hence, our first aim is to determine the cardinality of the sets  $A^t(k, \ell)$ .

From [16, Theorem 2.3] (see also [7]), the cardinality of  $M_{n,d,t}$  is given by

$$|M_{n,d,t}| = \binom{n - (d-1)(t-1)}{d}. \quad (2)$$

**Lemma 3.3** *Let  $k, \ell, t$  be three positive integers. Then  $|A^t(k, \ell)| = \binom{k+\ell-1}{\ell-1}$ .*

*Proof.* Let  $u \in A^t(k, \ell)$ , then  $u = x_{i_1}x_{i_2} \cdots x_{i_\ell}$  is a  $t$ -spread monomial of degree  $\ell$ , with  $\max(u) = k + (\ell - 1)t + 1$ . Set  $\tilde{u} = x_{i_1}x_{i_2} \cdots x_{i_{\ell-1}}$ , then  $\tilde{u}$  is a  $t$ -spread monomial of degree  $\ell - 1$  with  $\max(\tilde{u}) \leq \max(u) - t = k + (\ell - 2)t + 1$ . Hence,

$$|A^t(k, \ell)| = |M_{k+(\ell-2)t+1, \ell-1, t}|.$$

Therefore, by (2), we have

$$\begin{aligned} |A^t(k, \ell)| = |M_{k+(\ell-2)t+1, \ell-1, t}| &= \binom{k + (\ell - 2)t + 1 - (\ell - 1 - 1)(t - 1)}{\ell - 1} \\ &= \binom{k + \ell - 1}{\ell - 1}. \end{aligned}$$

□

For instance, if we consider Example 3.2, then  $A^3(5, 2)$  consists of six monomials. On the other hand, by Lemma 3.3,  $|A^3(5, 2)| = \binom{5+2-1}{2-1} = 6$ .

Setting  $A^t(k, \ell) = \{u_1, \dots, u_q\}$ ,  $q = |A^t(k, \ell)| = \binom{k+\ell-1}{\ell-1}$ , we can suppose, after a permutation of the indices, that

$$u_1 >_{\text{slex}} u_2 >_{\text{slex}} \cdots >_{\text{slex}} u_q. \quad (3)$$

Thus, for the  $i$ -th ( $1 \leq i \leq q$ ) monomial  $u$  of degree  $\ell$  with  $\max(u) = k + (\ell - 1)t + 1$ , we mean the monomial of  $A^t(k, \ell)$  that appears in the  $i$ -th position of (3).

Now, let  $(k, \ell)$  be a corner of a  $t$ -spread strongly stable ideal  $I$  of  $S$ , and let  $\beta_{k, k+\ell}(I) = a$ . The bound  $a \leq |A^t(k, \ell)|$  can be improved. More precisely, let  $u$  be the smallest  $t$ -spread monomial belonging to  $G(I)_\ell$  with  $\max(u) = k + t(\ell - 1) + 1$ , with respect to  $>_{\text{slex}}$ . Thus,  $u$  is equal to some  $u_j \in A^t(k, \ell)$ ,  $1 \leq j \leq |A^t(k, \ell)|$ . Therefore, by Corollary 2.8,

$$\begin{aligned} a = \beta_{k, k+\ell}(I) &= \left| \{u \in G(I)_\ell : \max(u) = k + t(\ell - 1) + 1\} \right| \\ &\leq \left| \{v \in A^t(k, \ell) : v \geq_{\text{slex}} u_j\} \right|. \end{aligned} \quad (4)$$

**Remark 3.4** A subset  $L$  of  $M_{n, d, t}$  is called  $t$ -spread lex set, if for all  $u \in L$  and for all  $v \in M_{n, d, t}$  with  $v \geq_{\text{slex}} u$ , it follows that  $v \in L$ . We observe that in (4), the equality holds if and only if  $G(I)_\ell$  is a  $t$ -spread lex set.

Now, we want to compute the cardinality of the set  $\{v \in A^t(k, \ell) : v \geq_{\text{slex}} u_j\}$ .

To do this, we introduce the following subsets of  $A^t(k, \ell)$ . Let  $u_i, u_j$ ,  $i < j$ , be two monomials of (3), the sets

$$\begin{aligned} [u_i, u_j] &:= \{w \in A^t(k, \ell) : u_i \geq_{\text{slex}} w \geq_{\text{slex}} u_j\}, \\ [u_i, u_j) &:= \{w \in A^t(k, \ell) : u_i \geq_{\text{slex}} w >_{\text{slex}} u_j\} \end{aligned}$$

are called *segment of  $A^t(k, \ell)$  of initial element  $u_i$  and final element  $u_j$*  and *left segment of  $A^t(k, \ell)$  of initial element  $u_i$  and final element  $u_j$* , respectively.

Hence, the set  $\{v \in A^t(k, \ell) : v \geq_{\text{slex}} u_j\}$  can be written as follow:

$$[x_1 x_{1+t} \cdots x_{1+t(\ell-2)} x_{k+t(\ell-1)+1}, u_j].$$

Indeed,  $x_1 x_{1+t} \cdots x_{1+t(\ell-2)} x_{k+t(\ell-1)+1}$  is the greatest monomial of  $A^t(k, \ell)$  with respect to  $>_{\text{slex}}$ .

In order to compute the cardinality of  $[x_1 x_{1+t} \cdots x_{1+t(\ell-2)} x_{k+t(\ell-1)+1}, u_j]$ , we illustrate how one may decompose  $|A^t(k, \ell)|$  via  $\min(u)$ , for all  $u \in A^t(k, \ell)$ .

**Proposition 3.5** [3, Lemma 4.3] *Let  $k, \ell$  be positive integers. Then*

$$\begin{aligned} \binom{k+\ell-1}{\ell-1} &= \sum_{i=1}^{k+1} \binom{k+\ell-1-i}{\ell-2} \\ &= \binom{k+\ell-2}{\ell-2} + \binom{k+\ell-3}{\ell-2} + \cdots + \binom{\ell-2}{\ell-2}. \end{aligned} \tag{5}$$

**Remark 3.6** It is important to discuss the binomial coefficients described in Proposition 3.5.

By Lemma 3.3,  $\binom{k+\ell-1}{\ell-1} = |A^t(k, \ell)|$ . Therefore  $\binom{k+\ell-1}{\ell-1}$  gives the number of  $t$ -spread monomials  $u$  of degree  $\ell$  with  $\max(u) = k + t(\ell - 1) + 1$ .

Note that for a monomial  $u \in A^t(k, \ell)$  one has  $\min(u) \leq k + 1$ . Indeed, the smallest monomial of  $A^t(k, \ell)$  with respect to  $>_{\text{slex}}$  is

$$x_{k+1} x_{k+t+1} x_{k+2t+1} \cdots x_{k+(\ell-1)t+1}.$$

Hence

$$|A^t(k, \ell)| = \binom{k+\ell-1}{\ell-1} = \sum_{i=1}^{k+1} b_i,$$

where  $b_i = |\{u \in A^t(k, \ell) : \min(u) = i\}|$ , for all  $i = 1, \dots, k + 1$ .

Now, fix  $i \in \{1, \dots, k + 1\}$ . The integer  $b_i$  is the number of all monomials  $u$  of  $A^t(k, \ell)$  with  $\min(u) = i$ . If  $u$  is a monomial of this type, then  $u$  can be written as  $u = x_{i_1} x_{i_2} \cdots x_{i_\ell}$ , with  $i_1 = i$ ,  $i_\ell = k + t(\ell - 1) + 1$  and  $i_2 \geq i_1 + t = i + t$ . Setting  $\tilde{u} = x_{i_2-(i+t)+1} x_{i_3-(i+t)+1} \cdots x_{i_\ell-(i+t)+1}$ , then  $i_2 - (i + t) + 1 \geq 1$ . Therefore, computing the number of all monomials



$u$  of  $A^t(k, \ell)$  with  $\min(u) = i$  is equivalent to determining the number of all  $t$ -spread monomials  $\tilde{u}$  of degree  $\ell - 1$  with  $\min(u) \geq 1$  and  $\max(u) = i_\ell - (i + t) + 1 = (k - i + 1) + (\ell - 2)t + 1$ ; such a number is the cardinality of the set  $A^t(k - i + 1, \ell - 1)$ . Therefore

$$b_i = |A^t(k - i + 1, \ell - 1)| = \binom{k - i + 1 + \ell - 1 - 1}{\ell - 1 - 1} = \binom{k + \ell - 1 - i}{\ell - 2}.$$

For a positive integer  $q$ , we set  $[q] = \{1, \dots, q\}$ . Following [3], we introduce the following definition that will have a key role throughout the paper.

**Definition 3.7** Let  $u = x_{i_1}x_{i_2}\cdots x_{i_d}$  be a  $t$ -spread monomial of degree  $d$  of  $S$ ,  $1 \leq i_1 < i_2 < \dots < i_d \leq n$ . Let  $j \in [d - 1]$ . We say that  $u$  has a  $j$ -gap $_t$  if

$$i_{j+1} - i_j - t > 0.$$

If  $u$  has a  $j$ -gap $_t$ , then we set

$$\text{wd}(j\text{-gap}_t)(u) := i_{j+1} - i_j - t$$

and we call it the *width of the  $j$ -gap $_t$* .

We define  $\text{Gap}_t(u)$  by

$$\text{Gap}_t(u) := \{j \in [d - 1] : \text{there exists a } j\text{-gap}_t(u)\}.$$

The following example illustrates the *combinatorial meaning* of the width of a  $j$ -gap $_t$ .

**Example 3.8** Let  $t = 2$  and  $u = x_{i_1}x_{i_2}x_{i_3}x_{i_4}x_{i_5} = x_2x_4x_6x_{13}x_{15} \in K[x_1, \dots, x_{15}]$  be a 2-spread monomial of degree 5. We have  $\text{Gap}_2(u) = \{3\}$  and  $\text{wd}(3\text{-gap}_2)(u) = 13 - 6 - 2 = 5$ . The width has the following meaning: *there exist exactly five variables (each of them has index smaller than 13) that may take the place of  $x_{i_4} = x_{13}$  in  $u$  in order to still obtain a 2-spread monomial*. These variables are  $x_{12}, x_{11}, x_{10}, x_9, x_8$ . Indeed,

$$\begin{aligned} x_{12}(u/x_{13}) &= x_2x_4x_6x_{12}x_{15}, \\ x_{11}(u/x_{13}) &= x_2x_4x_6x_{11}x_{15}, & x_9(u/x_{13}) &= x_2x_4x_6x_9x_{15}, \\ x_{10}(u/x_{13}) &= x_2x_4x_6x_{10}x_{15}, & x_8(u/x_{13}) &= x_2x_4x_6x_8x_{15}, \end{aligned}$$

are all 2-spread monomials.

There is another reason to consider the  $j$ -gap $_t$  of a  $t$ -spread monomial  $u \in A^t(k, \ell)$ , the set  $\text{Gap}_t(u)$  allow us to find the largest  $t$ -spread monomial of  $A^t(k, \ell)$  that follows  $u$  in the order  $>_{\text{slex}}$ , as the next result shows.

**Proposition 3.9** *Let  $n, t$  be positive integers. Let  $(k, \ell)$  be a pair of positive integers such that  $k + t(\ell - 1) + 1 \leq n$ . Let  $u = x_{i_1}x_{i_2} \cdots x_{i_\ell}$ ,  $1 \leq i_1 < i_2 < \cdots < i_\ell \leq n$  be a  $t$ -spread monomial of  $A^t(k, \ell)$ , that is  $i_\ell = k + t(\ell - 1) + 1$ . If  $\text{Gap}_t(u) = \emptyset$ , then  $u$  is the smallest  $t$ -spread monomial of  $A^t(k, \ell)$  with respect to  $>_{\text{slex}}$ . Otherwise, if  $p = \max \text{Gap}_t(u)$ , then the greatest monomial of  $A^t(k, \ell)$  following  $u$  in the squarefree lexicographic order is*

$$x_{i_1} \cdots x_{i_{p-1}} x_{i_p+1} x_{i_p+1+t} \cdots x_{i_p+1+t(\ell-p-1)} x_{k+t(\ell-1)+1}.$$

*Proof.* The proof is similar to the proof of [5, Lemma 3.1]. We include it for the reader's convenience.

If  $\text{Gap}_t(u) = \emptyset$ , then  $u = x_k x_{k+t+1} \cdots x_{k+t(\ell-2)+1} x_{k+t(\ell-1)+1}$ . Hence,  $u$  is the smallest  $t$ -spread monomial of degree  $d$ .

Now, let  $\text{Gap}_t(u) \neq \emptyset$  and let  $p = \max \text{Gap}_t(u)$ . If  $w = x_{s_1} x_{s_2} \cdots x_{s_\ell}$  is a monomial of  $A^t(k, \ell)$  with  $u >_{\text{slex}} w$ , then  $i_1 = s_1, \dots, i_{j-1} = s_{j-1}$  and  $i_j < s_j$  for some index  $j$ . We have  $j \leq p$ . Indeed, if  $j > p$ , then  $i_{p+2} - i_{p+1} = \cdots = i_\ell - i_{\ell-1} = t$ . Hence,  $i_\ell - i_j = t(\ell - j)$ ,  $i_\ell = s_\ell = k + t(\ell - 1) + 1$ ,  $s_\ell - s_j \geq t(\ell - j)$  and  $i_j < s_j$ . Therefore, we have

$$t(\ell - j) \leq s_\ell - s_j = i_\ell - s_j < i_\ell - i_j = t(\ell - j),$$

and consequently  $t(\ell - j) < t(\ell - j)$ . This is absurd. So  $j \leq p$ .

Let

$$v = x_{j_1} x_{j_2} \cdots x_{j_d} = x_{i_1} \cdots x_{i_{p-1}} x_{i_p+1} x_{i_p+1+t} \cdots x_{i_p+1+t(\ell-p-1)} x_{k+t(\ell-1)+1}.$$

Since  $i_1 = j_1, \dots, i_{p-1} = j_{p-1}$  and  $i_p < j_p = i_p + 1$ , then  $u >_{\text{slex}} v$ . Clearly,  $v$  is the greatest monomial of  $A^t(k, \ell)$  following  $u$  in the squarefree lexicographic order.  $\square$

For instance, if  $u = x_{i_1} x_{i_2} x_{i_3} x_{i_4} x_{i_5} = x_2 x_6 x_{11} x_{14} x_{17} \in A^3(4, 5) = A^t(k, \ell)$ , then  $p = \max \text{Gap}_3(u) = 2$ , and the greatest monomial  $v$  of  $A^3(4, 5)$  that follows  $u$ , with respect to  $>_{\text{slex}}$ , is

$$\begin{aligned} v &= x_{i_1} \cdots x_{i_{p-1}} x_{i_p+1} x_{i_p+1+t} \cdots x_{i_p+1+t(\ell-p-1)} x_{k+t(\ell-1)+1} \\ &= x_{i_1} x_{i_2+1} x_{i_2+1+t} x_{i_2+1+2t} x_{k+t(\ell-1)+1} \\ &= x_2 x_7 x_{10} x_{13} x_{17}. \end{aligned}$$

The following result shows that the cardinality of the segment

$$[x_1 x_{1+t} \cdots x_{1+t(\ell-2)} x_{k+t(\ell-1)+1}, u]$$

depends on the structure of the monomial  $u$ . More precisely, in order to compute  $|[x_1 x_{1+t} \cdots x_{1+t(\ell-2)} x_{k+t(\ell-1)+1}, u]|$ , one needs to analyze the set  $\text{Gap}_t(u/x_{\max(u)})$ .

**Theorem 3.10** Let  $(k, \ell)$  be a pair of positive integers with  $\ell \geq 2$  and let  $u = x_{i_1}x_{i_2} \cdots x_{i_{\ell-1}}x_{i_\ell}$  be a monomial of  $A^t(k, \ell)$ ,  $t \geq 1$ .

Setting  $\tilde{u} = x_{i_1}x_{i_2} \cdots x_{i_{\ell-1}}$ , then  $|[x_1x_{1+t} \cdots x_{1+t(\ell-2)}x_{k+t(\ell-1)+1}, u]|$  is a sum of  $r$  suitable binomial coefficients, where

$$r = \begin{cases} i_1, & \text{if } \text{Gap}_t(\tilde{u}) = \emptyset, \\ i_1 + \sum_{j=1}^p \text{wd}(g_j\text{-gap}_t)(\tilde{u}), & \text{if } \text{Gap}_t(\tilde{u}) = \{g_1, \dots, g_p\} \neq \emptyset. \end{cases}$$

*Proof.* Let  $m = |[x_1x_{1+t} \cdots x_{1+t(\ell-2)}x_{k+t(\ell-1)+1}, u]|$ , i.e.,  $m$  is the number of all monomials  $w \in A^t(k, \ell)$  such that  $w \geq_{\text{slex}} u$ . By Proposition 3.5, the binomial coefficient  $\binom{k+\ell-1}{\ell-1}$  can be decomposed as a sum of  $k+1$  binomial coefficients:

$$\begin{aligned} \binom{k+\ell-1}{\ell-1} &= \sum_{j=1}^{k+1} \binom{k+\ell-1-j}{\ell-2} \\ &= \binom{k+\ell-2}{\ell-2} + \binom{k+\ell-3}{\ell-2} + \cdots + \binom{\ell-2}{\ell-2}. \end{aligned} \tag{6}$$

From Remark 3.6, for all  $j = 1, \dots, k+1$ , the  $j$ -th binomial coefficient of (6),  $\binom{k+\ell-1-j}{\ell-2}$ , counts the number of  $t$ -spread monomials  $w$  of degree  $\ell$  with  $\min(w) = j$  and  $\max(w) = k + t(\ell - 1) + 1$ . Now, every monomial  $w \in A^t(k, \ell)$  with  $\min(w) < i_1 = \min(\tilde{u}) = \min(u)$  is greater than  $u$  with respect to  $>_{\text{slex}}$ . Therefore the first  $i_1 - 1$  binomial coefficients in (6) give a contribution to the computation of  $m$ .

We need to distinguish two cases:  $\text{Gap}_t(\tilde{u}) = \emptyset$ ,  $\text{Gap}_t(\tilde{u}) \neq \emptyset$ .

**Case 1.** Let  $\text{Gap}_t(\tilde{u}) = \emptyset$ . In such a case,  $u$  is the greatest monomial of  $A^t(k, \ell)$  with  $\min(u) = i_1$ . Therefore, the following sum of binomial coefficients

$$\sum_{j=1}^{i_1-1} \binom{k+\ell-1-j}{\ell-2} \tag{7}$$

gives the number of all monomials  $w \in A^t(k, \ell)$  greater than  $u$ . In fact, since  $i_{j+1} - i_j = t$ , for all  $j = 1, \dots, \ell - 1$ , then the other monomials greater than  $u$  which are different from the  $w$ 's counted by (7) do not exist. Hence,

$$\begin{aligned} m = |[x_1x_{1+t} \cdots x_{1+t(\ell-2)}x_{k+t(\ell-1)+1}, u]| &= \sum_{j=1}^{i_1-1} \binom{k+\ell-1-j}{\ell-2} + 1 \\ &= \sum_{j=1}^{i_1-1} \binom{k+\ell-1-j}{\ell-2} + \binom{0}{0}. \end{aligned}$$

Therefore,  $m$  is the sum of  $r = i_1 - 1 + 1 = i_1 = \min(\tilde{u}) = \min(u)$  binomial coefficients.

**Case 2.** Assume  $\text{Gap}_t(\tilde{u}) = \{g_1, \dots, g_p\} \neq \emptyset, p \geq 1$ . Let  $w \in A^t(k, \ell)$  such that  $w = x_{j_1}x_{j_2} \cdots x_{j_\ell}, w >_{\text{slex}} u$ . Since  $i_\ell = j_\ell$ , then there exists an integer  $s \in \{1, \dots, \ell - 1\}$  such that

$$j_1 = i_1, \quad j_2 = i_2, \dots, j_{s-1} = i_{s-1}, \quad j_s < i_s.$$

If  $s = 1$ , then  $w$  belongs to the set of monomials counted by (7). Let  $s > 1$ , then  $s - 1 \in \text{Gap}_t(\tilde{u})$ . Indeed,

$$i_s - i_{s-1} - t > j_s - j_{s-1} - t = \text{wd}((s-1)\text{-gap}_t)(w) \geq 0,$$

and so  $\text{wd}((s-1)\text{-gap}_t)(\tilde{u}) > 0$ . Therefore  $s-1 = g_h$  for some  $h \in \{1, \dots, p\}$ . Since,  $j_s < i_s$ , then  $j_s \leq i_s - 1 = i_{(g_h+1)} - 1$  and  $w = x_{i_1}x_{i_2} \cdots x_{i_{g_h}}z$ , with  $z$  a  $t$ -spread monomial of degree  $\ell - g_h$ , with

$$\begin{aligned} \max(z) &= k + t(\ell - 1) + 1, \\ \min(z) &= j_s \leq i_s - 1 = i_{(g_h+1)} - 1, \end{aligned}$$

and, furthermore,  $\min(z) \geq i_{g_h} + t$ . Thus,  $\min(z)$  belongs to the set

$$\begin{aligned} \{i_{g_h} + t, \dots, i_{(g_h+1)} - 1\} &= \{i_{g_h} + t, \dots, i_{(g_h+1)} - i_{g_h} - t + i_{g_h} + t - 1\} \\ &= \{i_{g_h} + t, \dots, i_{g_h} + t + \text{wd}(h\text{-gap}_t)(\tilde{u}) - 1\}. \end{aligned} \quad (8)$$

It follows that  $\min(z)$  may assume the  $\text{wd}(h\text{-gap}_t)(\tilde{u})$  possible values described in (8).

Now, let us consider the  $i_1$ -th binomial in (6):

$$\binom{k + \ell - 1 - i_1}{\ell - 2} = \sum_{j=1}^{k-i_1+2} \binom{k + \ell - 1 - i_1 - j}{\ell - 3}. \quad (9)$$

The  $j$ -th binomial coefficient of this sum counts the number of all monomials  $w = x_{j_1}x_{j_2} \cdots x_{j_\ell} \in A^t(k, \ell)$  such that  $\min(w) = j_1 = i_1$  and  $j_2 = i_2 + t + j - 1$ . Hence, if  $1 \in \text{Gap}_t(\tilde{u})$ , by the same arguments as before, the sum of the first  $\text{wd}(1\text{-gap}_t)(\tilde{u})$  in (9) gives the number of monomials  $w = x_{i_1}z \in A^t(k, \ell)$  greater than  $u$ . Thus, we consider the  $(\text{wd}(1\text{-gap}_t)(\tilde{u}) + 1)$ -th binomial coefficient and we decompose it; if  $2 \in \text{Gap}_t(\tilde{u})$ , then the sum of the first  $\text{wd}(2\text{-gap}_t)(\tilde{u})$  binomial coefficients of this new binomial decomposition gives the number of monomials  $x_{i_1}x_{i_2}z \in A^t(k, \ell)$  greater than  $u$ .

Otherwise, if  $1 \notin \text{Gap}_t(\tilde{u})$ , let us consider the first binomial in (9):

$$\binom{k + \ell - 2 - i_1}{\ell - 3} = \sum_{j=1}^{k-i_1+2} \binom{k + \ell - 2 - i_1 - j}{\ell - 4}.$$

The  $j$ -th binomial coefficient of this sum counts the number of all monomials  $w = x_{j_1}x_{j_2}\cdots x_{j_\ell} \in A^t(k, \ell)$  with  $\min(w) = j_1 = i_1$ ,  $j_2 = i_2$  and  $j_3 = i_3 + t + j - 1$ . If  $2 \notin \text{Gap}_t(\tilde{u})$ , then we consider the first binomial of this sum and we decompose it. We continue to evaluate successive binomial decomposition until we reach a gap  $g_j \in \text{Gap}_t(\tilde{u})$ , then the first  $\text{wd}(g_j\text{-gap}_t)(\tilde{u})$  binomial coefficients of the last evaluated binomial decomposition will give a contribute to the computation of  $m$ . This procedure can be iterated for all  $g_j \in \text{Gap}_t(\tilde{u})$ ,  $j \geq 3$ .

So,  $|[x_1x_{1+t}\cdots x_{1+t(\ell-2)}x_{k+t(\ell-1)+1}, u]|$  is the sum of

$$i_1 - 1 + \sum_{j=1}^p \text{wd}(g_j\text{-gap}_t)(\tilde{u})$$

suitable binomial coefficients.

Finally, in order to get  $m = |[x_1x_{1+t}\cdots x_{1+t(\ell-2)}x_{k+t(\ell-1)+1}, u]|$ , we must take into account the binomial  $\binom{0}{0} = 1$  which counts the monomial  $u$ :

$$r = i_1 - 1 + \sum_{j=1}^p \text{wd}(g_j\text{-gap}_t)(\tilde{u}) + 1 = i_1 + \sum_{j=1}^p \text{wd}(g_j\text{-gap}_t)(\tilde{u}).$$

The assertion follows.  $\square$

The next example illustrates Theorem 3.10.

**Example 3.11** Let  $S = K[x_1, \dots, x_{16}]$ ,  $t = 3$ ,  $(k, \ell) = (6, 4)$ , then  $k + t(\ell - 1) + 1 = 16$ . Let  $u = x_4x_9x_{13}x_{16} \in A^3(6, 4)$ . The greatest monomial of  $A^3(6, 4)$ , with respect to  $>_{\text{slex}}$ , is  $v = x_1x_4x_7x_{16}$ . We want to evaluate  $m = |[v, u]|$ .

From Lemma 3.3,  $|A^3(6, 4)| = \binom{9}{3} = 84$ .

Using the notation as in Theorem 3.10,  $\tilde{u} = x_4x_9x_{13}$ ,  $\text{Gap}_3(\tilde{u}) = \{1, 2\}$  and

$$\begin{aligned} i_1 &= 4, \\ \text{wd}(1\text{-gap}_3)(\tilde{u}) &= 9 - 4 - 3 = 2, \\ \text{wd}(2\text{-gap}_3)(\tilde{u}) &= 13 - 9 - 3 = 1. \end{aligned}$$

In order to compute  $m = |[x_1x_4x_7x_{16}, u]|$ , we consider the following binomial decomposition (Proposition 3.5):

$$\binom{9}{3} = \binom{8}{2} + \binom{7}{2} + \binom{6}{2} + \binom{5}{2} + \binom{4}{2} + \binom{3}{2} + \binom{2}{2}. \quad (10)$$

Since  $\min(u) = \min(\tilde{u}) = i_1 = 4$ , then all monomials  $w \in A^3(6, 4)$  with  $\min(w) \leq i_1 - 1 = 3$  are greater than  $u$ . Hence, for the computation of  $m$  we must take into account the sum of the first three binomial coefficients in (10), i.e.,  $\boxed{\binom{8}{2} + \binom{7}{2} + \binom{6}{2} = 64}$ .

Now, let us consider the following binomial decomposition:

$$\binom{5}{2} = \binom{4}{1} + \binom{3}{1} + \binom{2}{1} + \binom{1}{1}.$$

Since  $\text{wd}(1\text{-gap}_3)(\tilde{u}) = 2$ , the sum  $\boxed{\binom{4}{1} + \binom{3}{1} = 7}$  gives the number of all monomials of type  $x_{i_1}z = x_4z \in A^3(6, 4)$ , with  $z$  a  $t$ -spread monomial of degree 3 such that  $\max(z) = 16$  and  $\min(z) \in \{7, 8\}$ .

Up to this point we have got  $\boxed{64 + 7 = 71}$  monomials.

The next decomposition we need to consider is the following one

$$\binom{2}{1} = \binom{1}{0} + \binom{0}{0}.$$

Since  $2 \in \text{Gap}_3(\tilde{u})$  and  $\text{wd}(2\text{-gap}_3)(\tilde{u}) = 1$ , we must take into account  $\boxed{\binom{1}{0} = 1}$  monomial of the type  $x_{i_1}x_{i_2}z = x_4x_9z \in A^3(6, 4)$  with  $\max(z) = 16$  and  $\min(z) = 12$ .

Finally, we have obtained  $\boxed{71 + 1 = 72}$  monomials of  $A^3(6, 4)$  greater than  $u$ , and so  $m = |[x_1x_4x_7x_{16}, u]| = 72 + 1 = 73$ .

The following scheme summarizes the previous calculation.

$$\begin{aligned} \binom{9}{3} &= \boxed{\binom{8}{2} + \binom{7}{2} + \binom{6}{2}} + \binom{5}{2} + \binom{4}{2} + \binom{3}{2} + \binom{2}{2} \\ \binom{5}{2} &= \boxed{\binom{4}{1} + \binom{3}{1}} + \binom{2}{1} + \binom{1}{1} \\ \binom{2}{1} &= \boxed{\binom{1}{0}} + \binom{0}{0}. \end{aligned}$$

Figure 1 shows the list of all monomials which come into play to determine  $[v, u]$ :

$$\begin{array}{l}
\left. \begin{array}{l}
x_1x_4x_7x_{16}, x_1x_4x_8x_{16}, x_1x_4x_9x_{16}, x_1x_4x_{10}x_{16}, x_1x_4x_{11}x_{16}, x_1x_4x_{12}x_{16}, x_1x_4x_{13}x_{16} \\
x_1x_5x_8x_{16}, x_1x_5x_9x_{16}, x_1x_5x_{10}x_{16}, x_1x_5x_{11}x_{16}, x_1x_5x_{12}x_{16}, x_1x_5x_{13}x_{16} \\
x_1x_6x_9x_{16}, x_1x_6x_{10}x_{16}, x_1x_6x_{11}x_{16}, x_1x_6x_{12}x_{16}, x_1x_6x_{13}x_{16} \\
x_1x_7x_{10}x_{16}, x_1x_7x_{11}x_{16}, x_1x_7x_{12}x_{16}, x_1x_7x_{13}x_{16} \\
x_1x_8x_{11}x_{16}, x_1x_8x_{12}x_{16}, x_1x_8x_{13}x_{16} \\
x_1x_9x_{12}x_{16}, x_1x_9x_{13}x_{16} \\
x_1x_{10}x_{13}x_{16}
\end{array} \right\} \begin{array}{l} (8) \\ (2) \end{array} \\
\left. \begin{array}{l}
x_2x_5x_8x_{16}, x_2x_5x_9x_{16}, x_2x_5x_{10}x_{16}, x_2x_5x_{11}x_{16}, x_2x_5x_{12}x_{16}, x_2x_5x_{13}x_{16} \\
x_2x_6x_9x_{16}, x_2x_6x_{10}x_{16}, x_2x_6x_{11}x_{16}, x_2x_6x_{12}x_{16}, x_2x_6x_{13}x_{16} \\
x_2x_7x_{10}x_{16}, x_2x_7x_{11}x_{16}, x_2x_7x_{12}x_{16}, x_2x_7x_{13}x_{16} \\
x_2x_8x_{11}x_{16}, x_2x_8x_{12}x_{16}, x_2x_8x_{13}x_{16} \\
x_2x_9x_{12}x_{16}, x_2x_9x_{13}x_{16} \\
x_2x_{10}x_{13}x_{16}
\end{array} \right\} \begin{array}{l} (7) \\ (2) \end{array} \\
\left. \begin{array}{l}
x_3x_6x_9x_{16}, x_3x_6x_{10}x_{16}, x_3x_6x_{11}x_{16}, x_3x_6x_{12}x_{16}, x_3x_6x_{13}x_{16} \\
x_3x_7x_{10}x_{16}, x_3x_7x_{11}x_{16}, x_3x_7x_{12}x_{16}, x_3x_7x_{13}x_{16} \\
x_3x_8x_{11}x_{16}, x_3x_8x_{12}x_{16}, x_3x_8x_{13}x_{16} \\
x_3x_9x_{12}x_{16}, x_3x_9x_{13}x_{16} \\
x_3x_{10}x_{13}x_{16}
\end{array} \right\} \begin{array}{l} (6) \\ (2) \end{array} \\
\left. \begin{array}{l}
x_4x_7x_{10}x_{16}, x_4x_7x_{11}x_{16}, x_4x_7x_{12}x_{16}, x_4x_7x_{13}x_{16} \\
x_4x_8x_{11}x_{16}, x_4x_8x_{12}x_{16}, x_4x_8x_{13}x_{16}
\end{array} \right\} \begin{array}{l} (4) + (3) \\ (1) + (1) \end{array} \\
\left. \begin{array}{l}
x_4x_9x_{12}x_{16} \\
x_4x_9x_{13}x_{16}
\end{array} \right\} \begin{array}{l} (1) \\ (0) \end{array} \\
\left. \begin{array}{l}
x_4x_9x_{13}x_{16}
\end{array} \right\} \begin{array}{l} (0) \\ (0) \end{array}
\end{array}$$

Figure 1: Monomials of  $[v, u]$ .

**Remark 3.12** Let  $I$  be a  $t$ -spread strongly stable ideal of  $S = K[x_1, \dots, x_n]$  generated in degree  $\ell$ , and let  $k$  and  $a$  two positive integers. Then,  $I$  has extremal Betti number  $\beta_{k, k+\ell}(I) = a$  if and only if  $k + t(\ell - 1) + 1 \leq n$  and  $a \leq |A^t(k, \ell)|$ .

If  $A^t(k, \ell) = \{u_1, \dots, u_q\}$ , with  $u_1, \dots, u_q$  as in (3), and  $u_j$  is the smallest monomial of  $A^t(k, \ell)$  belonging to  $G(I)_\ell$ , then

$$\begin{aligned}
a &\leq |\{v \in A^t(k, \ell) : v \geq_{\text{slex}} u_j\}| \\
&= |[x_1x_{1+t} \cdots x_{1+t(\ell-2)+1}x_{k+t(\ell-1)+1}, u_j]| = m,
\end{aligned}$$

and Theorem 3.10 allows to determine  $m$ . Therefore, if the ideal  $I$  is generated in one degree  $\ell$ , then the proposed question (\*) (see, the section Introduction) is solved.

If  $I$  is a  $t$ -spread strongly stable ideal generated in several degrees and  $\beta_{k, k+\ell}(I) = a$  is an extremal Betti number, the bound  $a \leq |A^t(k, \ell)|$  is not sufficient to solve the question (\*), as next example shows.

**Example 3.13** Let  $n = 13$ ,  $t = 2$ . We will show that a  $t$ -spread strongly stable ideal  $I$  of  $S = K[x_1, \dots, x_{13}]$  with corners  $(k_1, \ell_1) = (5, 2)$ ,  $(k_2, \ell_2) = (3, 4)$ ,  $k_1 > k_2$ ,  $2 \leq \ell_1 < \ell_2$ ,  $k_i + t(\ell_i - 1) + 1 \leq 13$  ( $i = 1, 2$ ) and extremal

Betti numbers  $\beta_{k_1, k_1 + \ell_1}(I) = a_1 = 3$  and  $\beta_{k_2, k_2 + \ell_2}(I) = a_2 = 10$  does not exist.

First of all, we write down the monomials of the sets  $A^t(k_i, \ell_i)$ ,  $i = 1, 2$ :

$$\begin{aligned} A^2(5, 2) &= \{x_1x_8, x_2x_8, x_3x_8, x_4x_8, x_5x_8, x_6x_8\}, \\ A^2(3, 4) &= \{x_1x_3x_5x_{10}, x_1x_3x_6x_{10}, x_1x_3x_7x_{10}, x_1x_3x_8x_{10}, x_1x_4x_6x_{10}, \\ &\quad x_1x_4x_7x_{10}, x_1x_4x_8x_{10}, x_1x_5x_7x_{10}, x_1x_5x_8x_{10}, x_1x_6x_8x_{10}, \\ &\quad x_2x_4x_6x_{10}, x_2x_4x_7x_{10}, x_2x_4x_8x_{10}, x_2x_5x_7x_{10}, x_2x_5x_8x_{10}, \\ &\quad x_2x_6x_8x_{10}, x_3x_5x_7x_{10}, x_3x_5x_8x_{10}, x_3x_6x_8x_{10}, x_4x_6x_8x_{10}\}. \end{aligned}$$

Assume that the sets  $A^t(k_i, \ell_i)$  are ordered by  $>_{\text{slex}}$ .

We observe that for the  $a_i$ 's ( $i = 1, 2$ ) the following upper bounds hold

$$\begin{aligned} a_1 &\leq |A^t(k_1, \ell_1)| = \binom{5+2-1}{2-1} = 6, \\ a_2 &\leq |A^t(k_2, \ell_2)| = \binom{3+4-1}{4-1} = 20. \end{aligned}$$

From Corollary 2.8,

$$\beta_{k_1, k_1 + \ell_1}(I) = \beta_{5,7}(I) = \left| \left\{ u \in G(I)_2 : \max(u) = 8 \right\} \right|.$$

$G(I)_{\ell_1} = G(I)_2$  has to be a 2-spread strongly stable set such that  $|G(I)_2 \cap A^2(5, 2)| = a_1 = 3$ . Hence,  $G(I)_2 \cap A^2(5, 2) = \{x_1x_8, x_2x_8, x_3x_8\}$ . Therefore we set,

$$G(I)_2 = B_2\{x_1x_8, x_2x_8, x_3x_8\}.$$

Now, we have to determine  $G(I)_{\ell_2} = G(I)_4$  such that  $\beta_{k_2, k_2 + \ell_2}(I) = a_2 = 10$ . So, we have to find 10 monomials of  $A^2(3, 4)$  not belonging to  $\text{Shad}_2^{\ell_2 - \ell_1}(G(I)_{\ell_1})$ . Since  $\min \text{Shad}_2^{\ell_2 - \ell_1}(G(I)_{\ell_1}) = x_3x_6x_8x_{10}$ , then the only monomial of  $A^t(k_2, \ell_2) = A^2(3, 4)$  that may belong to  $G(I)_{\ell_2} = G(I)_4$  is  $x_4x_6x_8x_{10}$ . Hence, the value of  $\beta_{k_2, k_2 + \ell_2}(I) = \beta_{3,7}(I)$  can be at most 1. Thus, the value  $a_2 = 10$  cannot be reached.

Finally, a 2-spread strongly stable ideal  $I$  of  $S$  with extremal Betti numbers  $\beta_{k_1, k_1 + \ell_1}(I) = a_1 = 3$  and  $\beta_{k_2, k_2 + \ell_2}(I) = a_2 = 10$  does not exist.

We close the Section by the next definition that will be useful for solving our problem.

**Definition 3.14** Let  $(k_1, \ell_1), (k_2, \ell_2)$  be two pairs of positive integers such that  $k_1 > k_2, \ell_1 < \ell_2$ , and  $k_i + t(\ell_i - 1) + 1 \leq n$ ,  $i = 1, 2$ . If  $T = \{u_1, \dots, u_r\}$  is a subset of monomials of  $A^t(k_1, \ell_1)$ , we define the *Borel  $t$ -shadow* of  $T = \{u_1, \dots, u_r\}$  with respect to the corner  $(k_2, \ell_2)$  as follows

$$\text{BShad}_t(T)_{(k_2, \ell_2)} = \left\{ v \in \text{Shad}_t^{\ell_2 - \ell_1}(B_t\{T\}) : \max(v) \leq k_2 + t(\ell_2 - 1) + 1 \right\}.$$



**Remark 3.15** It is clear that  $\text{BShad}_t(u_1, \dots, u_r)_{(k_2, \ell_2)}$  is a  $t$ -spread strongly stable set. Moreover, one can easily verify that, if  $u_1 >_{\text{slex}} u_2 >_{\text{slex}} \dots >_{\text{slex}} u_r$ , then

$$\min \text{BShad}_t(u_1, \dots, u_r)_{(k_2, \ell_2)} = \min \text{BShad}_t(u_r)_{(k_2, \ell_2)}.$$

#### 4. A NUMERICAL CHARACTERIZATION

In this Section we give a numerical characterization of the extremal Betti numbers of a  $t$ -spread strongly stable ideal by generalizing the methods in [3].

In order to find optimal bounds for the values of the extremal Betti numbers of a  $t$ -spread strongly stable ideal, we need some comments.

**Discussion 4.1** Let  $t \geq 1$  and  $S = K[x_1, \dots, x_n]$ . Let  $(k_1, \ell_1), (k_2, \ell_2)$  be two pairs of positive integers with  $k_1 > k_2$ ,  $2 \leq \ell_1 < \ell_2$  and  $k_i + t(\ell_i - 1) + 1 \leq n$  ( $i = 1, 2$ ), and let  $a_1, a_2$  be two positive integers.

Now, let  $J_2 = [u_{2,1}, u_{2,a_2}]$  be a segment of  $A^t(k_2, \ell_2)$  of cardinality  $a_2$ . We want to find the admissible values of  $a_1$  such that there is a segment  $J_1 = [u_{1,1}, u_{1,a_1}]$  of  $A^t(k_1, \ell_1)$  of cardinality  $a_1$  with

$$J_2 \cap \text{BShad}_t([u_{1,1}, u_{1,a_1}])_{(k_2, \ell_2)} = \emptyset.$$

First of all,  $a_1 \leq |A^t(k_1, \ell_1)| = \binom{k_1 + \ell_1 - 1}{\ell_1 - 1}$ . The inequality must be strict, otherwise  $\text{BShad}_t([u_{1,1}, u_{1,a_1}])_{(k_2, \ell_2)} = \text{BShad}_t(A^t(k_1, \ell_1))_{(k_2, \ell_2)} = A^t(k_2, \ell_2)$ , and thus  $J_2 \cap \text{BShad}_t([u_{1,1}, u_{1,a_1}])_{(k_2, \ell_2)} = J_2 \neq \emptyset$ .

Suppose  $a_1$  is such that  $J_2 \cap \text{BShad}_t([u_{1,1}, u_{1,a_1}])_{(k_2, \ell_2)} = \emptyset$ . Let

$$v_1 := \min \{u \in A^t(k_1, \ell_1) : \max J_2 = u_{2,1} \notin \text{BShad}_t(u)\}.$$

Clearly,  $u_{1,a_1} \geq_{\text{slex}} v_1$ . By Theorem 3.10, we can calculate the following cardinalities:

$$\begin{aligned} n_1 &:= |\{u \in A^t(k_1, \ell_1) : u \geq_{\text{slex}} v_1\}| = |[x_1 x_{1+t} \cdots x_{1+t(\ell_1-2)} x_{k_1+t(\ell_1-1)+1}, v_1]|, \\ p_1 &:= |\{v \in A^t(k_1, \ell_1) : v >_{\text{slex}} u_{1,1}\}| = |[x_1 x_{1+t} \cdots x_{1+t(\ell_1-2)} x_{k_1+t(\ell_1-1)+1}, u_{1,1}]|. \end{aligned}$$

Since  $[u_{1,1}, u_{1,a_1}] \subseteq [x_1 x_{1+t} \cdots x_{1+t(\ell_1-2)} x_{k_1+t(\ell_1-1)+1}, v_1]$ , then

$$a_1 \leq n_1.$$

We can improve such a bound for  $a_1$ . Indeed,  $[u_{1,1}, u_{1,a_1}] \subseteq [u_{1,1}, v_1]$  and since

$$[u_{1,1}, v_1] = [x_1 \cdots x_{1+t(\ell_1-2)} x_{k_1+t(\ell_1-1)+1}, v_1] \setminus [x_1 \cdots x_{1+t(\ell_1-2)} x_{k_1+t(\ell_1-1)+1}, u_{1,1}),$$

then we have  $|[u_{1,1}, v_1]| = n_1 - p_1$ . Finally,

$$a_1 \leq n_1 - p_1.$$

It is clear that if  $u_{1,1} = x_1 x_{1+t} \cdots x_{1+t(\ell_1-2)} x_{k_1+t(\ell_1-1)+1} = \max A^t(k_1, \ell_1)$ , then  $p_1 = 0$ , and the bound  $a_1 \leq n_1$  is an optimal bound.

We want to review Example 3.13 *via* Discussion 4.1.

**Example 4.2** Let  $n = 13$ ,  $t = 2$ ,  $(k_1, \ell_1) = (5, 2)$ ,  $(k_2, \ell_2) = (3, 4)$ ,  $k_1 > k_2$ ,  $2 \leq \ell_1 < \ell_2$ ,  $k_i + t(\ell_i - 1) + 1 \leq 13$ ,  $i = 1, 2$ , and  $a_1 \geq 1$ ,  $a_2 = 10$ . Let  $J_2 = [x_2 x_4 x_6 x_{10}, x_4 x_6 x_8 x_{10}]$  be a segment of  $A^t(k_2, \ell_2)$  of cardinality  $a_2 = 10$ . We consider this segment since its monomials are the smallest of  $A^t(k_2, \ell_2)$  with respect to  $>_{\text{slex}}$ . Such a choice gives us the best chance to construct a  $t$ -spread strongly stable ideal of  $S = K[x_1, \dots, x_{13}]$  with extremal Betti numbers  $\beta_{k_1, k_1+\ell_1}(I) = a_1$  and  $\beta_{k_2, k_2+\ell_2}(I) = a_2 = 10$ .

Let us determine the admissible values of  $a_1$ . Let

$$\begin{aligned} v_1 &= \min \{u \in A^t(k_1, \ell_1) : \max J_2 = u_{2,1} \notin \text{BShad}_t(u)\} \\ &= \min \{u \in A^2(5, 2) : x_2 x_4 x_6 x_{10} \notin \text{BShad}_t(u)\} \\ &= x_1 x_8. \end{aligned}$$

Hence,  $n_1 = |[x_1 x_8, x_1 x_8]| = 1$ . Let  $J_1 = [u_{1,1}, u_{1,a_1}]$  be a segment of  $A^t(k_1, \ell_1)$  of cardinality  $a_1$ . If  $u_{1,1} = x_1 x_8$ , then  $p_1 = 0$ , and  $a_1 \leq n_1 - p_1 = 1$ . In such a case  $a_1 = 1$ , and we can construct the ideal  $I$  as follows:

$$\begin{aligned} G(I)_{\ell_1} &= G(I)_2 = B_2\{x_1 x_8\}, \\ G(I)_{\ell_2} &= G(I)_4 = B_2\{[x_2 x_4 x_6 x_{10}, x_4 x_6 x_8 x_{10}]\} \setminus \text{Shad}_2^2([I]_2). \end{aligned}$$

The ideal  $I$  has two extremal Betti numbers,  $\beta_{k_1, k_1+\ell_1}(I) = \beta_{5,7}(I) = 1 = a_1$  and  $\beta_{k_2, k_2+\ell_2}(I) = \beta_{3,7}(I) = 10 = a_2$ .

If  $u_{1,1} <_{\text{slex}} x_1 x_8$ , then  $p_1 \geq 1$ , and  $a_1 \leq n_1 - p_1 \leq 0$  and in such a case the ideal  $I$  does not exist. One can immediately observe that a  $t$ -spread strongly stable ideal  $I$  of  $S$  with extremal Betti numbers  $\beta_{k_1, k_1+\ell_1}(I) = a_1 = 3$  and  $\beta_{k_2, k_2+\ell_2}(I) = a_2 = 10$  does not exist, since  $a_1$  can be at most 1.

Discussion 4.1 will be useful for the aim of this article.

Recall that the floor function of a real number  $x$  is defined as follows:

$$\lfloor x \rfloor = \max\{n \in \mathbb{Z} : n \leq x\}.$$

In particular, if  $-1 \leq x < 0$  then  $\lfloor x \rfloor = -1$  and if  $0 \leq x < 1$  then  $\lfloor x \rfloor = 0$ .

We quote the next result from [5]. Such a result gives the maximal number of corners allowed for a  $t$ -spread strongly stable ideal.

**Theorem 4.3** ([5, Theorem 4.1, Theorem 5.1]) *Let  $n, t, k$  be three positive integers such that  $n, t \geq 2$  and  $k \geq 3$ . Assume*

$$n = d + kt, \quad 1 \leq d \leq t.$$

*Then, for  $k \geq 3$ , every  $t$ -spread strongly stable ideal  $I$  of  $S = K[x_1, \dots, x_n]$  of initial degree  $\ell_1$ ,  $2 \leq \ell_1 \leq k + \lfloor \frac{d-2}{t} \rfloor + 1$ , and with a corner in degree  $\ell_1$  can have at most*

$$\begin{aligned} k + \lfloor \frac{d-3}{t} \rfloor \text{ corners,} & \quad \text{if } \ell_1 = 2, \\ k + \lfloor \frac{d-2}{t} \rfloor - (\ell_1 - 2) \text{ corners,} & \quad \text{if } \ell_1 \geq 3. \end{aligned}$$

Now we are in position to prove the main result in the paper.

**Theorem 4.4** *Let  $n, \ell_1, r$  three positive integer, such that  $n = d + kt$ ,  $t \geq 1$ ,  $1 \leq d \leq t$ ,  $k \geq 3$ ,  $2 \leq \ell_1 \leq k + \lfloor \frac{d-2}{t} \rfloor + 1$ , and  $r$  an integer such that*

$$\begin{aligned} 1 \leq r \leq k + \lfloor \frac{d-3}{t} \rfloor, & \quad \text{if } \ell_1 = 2, \\ 1 \leq r \leq k + \lfloor \frac{d-2}{t} \rfloor - (\ell_1 - 2), & \quad \text{if } \ell_1 \geq 3. \end{aligned}$$

*Let  $(k_1, \ell_1), \dots, (k_r, \ell_r)$  be  $r$  pairs of positive integers such that  $n - t - 1 \geq k_1 > k_2 > \dots > k_r \geq 0$ ,  $k_i + t(\ell_i - 1) + 1 \leq n$ , for all  $i = 1, \dots, r$ ,*

$$\begin{aligned} 2 = \ell_1 < \ell_2 < \dots < \ell_r \leq k + \lfloor \frac{d-3}{t} \rfloor + 1, & \quad \text{if } \ell_1 = 2, \\ 3 \leq \ell_1 < \ell_2 < \dots < \ell_r \leq k + \lfloor \frac{d-2}{t} \rfloor + 1, & \quad \text{if } \ell_1 \geq 3, \end{aligned}$$

*and let  $a_1, \dots, a_r$  be  $r$  positive integers. The following conditions are equivalent:*

- (1) *There exists a  $t$ -spread strongly stable ideal  $I$  of  $S = K[x_1, \dots, x_n]$  with*

$$\beta_{k_1, k_1 + \ell_1}(I) = a_1, \dots, \beta_{k_r, k_r + \ell_r}(I) = a_r$$

*as extremal Betti numbers.*

- (2) *Setting*

- (i)  $v_r = x_{k_r+1}x_{k_r+t+1} \cdots x_{k_r+t(\ell_r-1)+1}$  and  $A_r = [w_r, v_r]$ , with  $w_r \in A^t(k_r, \ell_r)$  such that  $|A_r| = a_r$ ;
- (ii)  $v_{r-i} = \min \{ u \in A^t(k_{r-i}, \ell_{r-i}) : \max A_{r-i+1} \notin \text{BShad}_t(u)_{(k_{r-i+1}, \ell_{r-i+1})} \}$ ,  $A_{r-i} = [w_{r-i}, v_{r-i}]$ , with  $w_{r-i} \in A^t(k_{r-i}, \ell_{r-i})$  such that  $|A_{r-i}| = a_{r-i}$ ;

(iii) for  $i = 1, \dots, r$ ,  $n_i = |\{u \in A^t(k_i, \ell_i) : u \geq v_i\}|$ , then the integers  $a_i$  satisfy the following bound:

$$a_i \leq n_i.$$

Moreover, if  $a_i = |[u_{i,1}, u_{i,a_i}]|$ , with  $u_{i,j} \in A^t(k_i, \ell_i)$  for all  $i = 1, \dots, r$ , and all  $j = 1, \dots, a_i$ , setting  $p_i = |\{v \in A^t(k_i, \ell_i) : v > u_{i,1}\}|$ , then  $a_i$  satisfy the bound  $a_i \leq n_i - p_i$ , for all  $i = 1, \dots, r$ .

*Proof.* (1)  $\implies$  (2). It follows by applying iteratively Discussion 4.1.  
(2)  $\implies$  (1). We construct a  $t$ -spread strongly stable ideal  $I$  of  $S = K[x_1, \dots, x_n]$  generated in degrees  $\ell_1, \dots, \ell_r$ , with

$$\beta_{k_1, k_1 + \ell_1}(I) = a_1, \dots, \beta_{k_r, k_r + \ell_r}(I) = a_r$$

as extremal Betti numbers.

First of all, we set

$$G(I)_{\ell_1} := B_t\{u_{1,1}, \dots, u_{1,a_1}\},$$

where  $u_{1,1} >_{\text{slex}} \dots >_{\text{slex}} u_{1,a_1}$  are the first  $a_1$  monomials of  $A^t(k_1, \ell_1)$  with respect to the squarefree lexicographic order. The existence of such monomials is guaranteed by the hypothesis  $a_1 \leq n_1 \leq |A^t(k_1, \ell_1)|$ . We observe that  $[I_{\ell_1}]_t = G(I)_{\ell_1}$  is a  $t$ -spread strongly stable set, and by Corollary 2.8

$$\beta_{k_1, k_1 + \ell_1}(I) = \left| \left\{ u \in G(I)_{\ell_1} : \max(u) = k_1 + t(\ell_1 - 1) + 1 \right\} \right| = |[u_{1,1}, u_{1,a_1}]| = a_1.$$

Now, we construct  $G(I)_{\ell_2}$ . Setting  $p_1 = |\{v \in A^t(k_1, \ell_1) : v > u_{1,1}\}|$ , then  $p_1 = 0$  because  $u_{1,1} = x_1 x_{1+t} \dots x_{1+t(\ell_2-2)} x_{k_1+t(\ell_2-1)+1} = \max A^t(k_1, \ell_1)$ . By hypothesis (iii),  $a_1 \leq n_1 - p_1 = n_1$ , i.e.

$$\begin{aligned} |[u_{1,1}, u_{1,a_1}]| &\leq |[x_1 x_{1+t} \dots x_{1+t(\ell_2-2)} x_{k_1+t(\ell_2-1)+1}, v_1]| \\ &= |[u_{1,1}, v_1]|. \end{aligned}$$

It follows that  $u_{1,a_1} \geq_{\text{slex}} v_1 = \min \{v \in A^t(k_1, \ell_1) : \max A_2 \notin \text{BShad}_t(v)_{(k_2, \ell_2)}\}$ . Thus,  $\max A_2 \notin \text{BShad}_t(u_{1,a_1})_{(k_2, \ell_2)}$ , and  $\min \text{BShad}_t(u_{1,a_1})_{(k_2, \ell_2)} >_{\text{slex}} \max A_2$ . By Remark 3.15,

$$\min \text{BShad}_t(B_t\{[u_{1,1}, u_{1,a_1}]\})_{(k_2, \ell_2)} = \min \text{BShad}_t(B_t\{u_{1,a_1}\})_{(k_2, \ell_2)}.$$

Moreover, since  $[I_{\ell_1}]_t = B_t\{[u_{1,1}, u_{1,a_1}]\}$ , we have

$$\min \text{BShad}_t(B_t\{[u_{1,1}, u_{1,a_1}]\})_{(k_2, \ell_2)} = \min \text{BShad}_t([I_{\ell_1}]_t)_{(k_2, \ell_2)} >_{\text{slex}} \max A_2.$$

Then,  $A_2 \cap \text{BShad}_t([I_{\ell_1}]_t)_{(k_2, \ell_2)} = \emptyset$  and so we can find at least  $a_2$  monomials of  $A^t(k_2, \ell_2)$  not belonging to  $\text{BShad}_t([I_{\ell_1}]_t)_{(k_2, \ell_2)}$ . Let  $u_{2,1}, \dots, u_{2,a_2}$  be the

first  $t$ -spread monomials of  $A^t(k_2, \ell_2)$  not belonging to  $\text{BShad}_t([I_{\ell_1}]_t)_{(k_2, \ell_2)}$ , with respect to  $>_{\text{slex}}$ . We set

$$G(I)_{\ell_2} := B_t\{u_{2,1}, \dots, u_{2,a_2}\} \setminus \text{Shad}_t^{\ell_2 - \ell_1}([I_{\ell_1}]_t).$$

By our assumptions, from Corollary 2.8, we have

$$\beta_{k_2, k_2 + \ell_2}(I) = \left| \left\{ u \in G(I)_{\ell_2} : \max(u) = k_2 + t(\ell_2 - 1) + 1 \right\} \right| = |[u_{2,1}, u_{2,a_2}]| = a_2.$$

One can note that  $[I_{\ell_2}]_t$  is a  $t$ -spread strongly stable set. In fact,

$$[I_{\ell_2}]_t = B_t\{u_{2,1}, \dots, u_{2,a_2}\} \cup \text{Shad}_t^{\ell_2 - \ell_1}([I_{\ell_1}]_t),$$

and the union of two  $t$ -spread strongly stable sets is still a  $t$ -spread strongly stable set.

Now, we construct  $G(I)_{\ell_3}$ . Let  $a_2 = |[u_{2,1}, u_{2,a_2}]|$  and let  $p_2 = |\{p \in A^t(k_2, \ell_2) : v >_{\text{slex}} u_{2,1}\}|$ . By hypothesis (iii),  $a_2 \leq n_2 - p_2$ . Hence,

$$\begin{aligned} |[u_{2,1}, u_{2,a_2}]| &\leq |[x_1 x_{1+t} \cdots x_{1+t(\ell_2-2)+1} x_{k_2+t(\ell_2-1)+1}, v_2]| \\ &\quad - |[x_1 x_{1+t} \cdots x_{1+t(\ell_2-2)+1} x_{k_2+t(\ell_2-1)+1}, u_{2,1}]| \\ &= |[u_{2,1}, v_2]|. \end{aligned}$$

It follows that  $u_{2,a_2} \geq_{\text{slex}} v_2 = \min \{v \in A^t(k_2, \ell_2) : \max A_3 \notin \text{BShad}_t(v)_{(k_3, \ell_3)}\}$ , and thus  $\max A_3 \notin \text{BShad}_t(u_{2,a_2})_{(k_3, \ell_3)}$  and  $\min \text{BShad}_t(u_{2,a_2})_{(k_3, \ell_3)} >_{\text{slex}} \max A_3$ . By Remark 3.15,

$$\min \text{BShad}_t(B_t([u_{2,1}, u_{2,a_2}]))_{(k_3, \ell_3)} = \min \text{BShad}_t(B_t(u_{2,a_2}))_{(k_3, \ell_3)}$$

and, by construction, we have

$$\min \text{BShad}_t([I_{\ell_2}]_t)_{(k_3, \ell_3)} \geq_{\text{slex}} \min \text{BShad}_t(B_t([u_{2,1}, u_{2,a_2}]))_{(k_3, \ell_3)} >_{\text{slex}} \max A_3.$$

Therefore,  $A_3 \cap \text{BShad}_t([I_{\ell_2}]_t)_{(k_3, \ell_3)} = \emptyset$  and we can find at least  $a_3$  monomials of  $A^t(k_3, \ell_3)$  not belonging to  $\text{BShad}_t([I_{\ell_2}]_t)_{(k_3, \ell_3)}$ . Let  $u_{3,1}, \dots, u_{3,a_3}$  be the first  $t$ -spread monomials of  $A^t(k_3, \ell_3)$  not belonging to  $\text{BShad}_t([I_{\ell_2}]_t)_{(k_3, \ell_3)}$ , with respect to  $>_{\text{slex}}$ .

Setting

$$G(I)_{\ell_3} := B_t\{u_{3,1}, \dots, u_{3,a_3}\} \setminus \text{Shad}_t^{\ell_3 - \ell_2}([I_{\ell_2}]_t),$$

we have

$$\beta_{k_3, k_3 + \ell_3}(I) = \left| \left\{ u \in G(I)_{\ell_3} : \max(u) = k_3 + t(\ell_3 - 1) + 1 \right\} \right| = |[u_{3,1}, u_{3,a_3}]| = a_3.$$

Furthermore,  $[I_{\ell_3}]_t$  is  $t$ -spread strongly stable set. Indeed,

$$[I_{\ell_3}]_t = B_t\{u_{3,1}, \dots, u_{3,a_3}\} \cup \text{Shad}_t^{\ell_3 - \ell_2}([I_{\ell_2}]_t).$$

Clearly, we can iterate this construction, setting

$$G(I)_{\ell_i} := B_t\{u_{i,1}, \dots, u_{i,a_i}\} \setminus \text{Shad}_t^{\ell_i - \ell_{i-1}}([I_{\ell_{i-1}}]_t),$$

for  $i = 4, \dots, r$ . At the end, we will obtain a  $t$ -spread strongly stable ideal  $I$  of  $S = K[x_1, \dots, x_n]$  generated in degrees  $\ell_1, \dots, \ell_r$  with  $\beta_{k_1, k_1 + \ell_1}(I) = a_1, \dots, \beta_{k_r, k_r + \ell_r}(I) = a_r$  as extremal Betti numbers.  $\square$

We close this Section with a method for computing the minimum of a Borel  $t$ -shadow. The determination of such a monomial has a key role in the proof of Theorem 4.4.

Firstly, we consider a simple case.

**Lemma 4.5** *Let  $(k_1, \ell_1), (k_2, \ell_2)$  be a pair of positive integers, with  $k_1 > k_2$ ,  $\ell_1 \geq 2$ ,  $\ell_2 = \ell_1 + 1$  and  $k_i + t(\ell_i - 1) + 1 \leq n$ ,  $i = 1, 2$ . If  $u = x_{i_1} x_{i_2} \dots x_{i_{\ell_1}}$  is a monomial of  $A^t(k_1, \ell_1)$ , then*

$$\min \text{BShad}_t(u)_{(k_2, \ell_1 + 1)} = x_{i_1} \dots x_{i_{(\ell_1 - 2 - m)}} \left( \prod_{\nu = \ell_1 - m - 1}^{\ell_1 + 1} x_{k_2 + t(\nu - 1) + 1} \right),$$

with

$$m = \min \left\{ j \in \mathbb{Z}_{\geq -1} : (k_2 + t(\ell_2 - 1) + 1 - i_{(\ell_1 - 1)}) + \sum_{s=0}^j \text{wd}((\ell_1 - 2 - s)\text{-gap}_t)(u) \geq t \right\}.$$

*Proof.* Firstly, we have that  $i_{\ell_1} = k_1 + t(\ell_1 - 1) + 1$ . Moreover, since  $k_1 > k_2$ , we may consider the monomial

$$v = x_{i_1} x_{i_2} \dots x_{i_{(\ell_1 - 1)}} x_{k_2 + t(\ell_1 - 1) + 1} x_{k_2 + t((\ell_1 + 1) - 1) + 1}. \quad (11)$$

If  $k_2 + t(\ell_1 - 1) + 1 - i_{(\ell_1 - 1)} \geq t$ , then  $v$  is the minimum of  $\text{BShad}_t(u)_{(k_2, \ell_1 + 1)}$ .

Otherwise, if  $k_2 + t(\ell_1 - 1) + 1 - i_{(\ell_1 - 1)} < t$ , we proceed as follows.

Let us define the following nonnegative integer

$$\overline{m} = \min \left\{ j \in \mathbb{Z}_{\geq 0} : (k_2 + t((\ell_1 + 1) - 1) + 1 - i_{(\ell_1 - 1)}) + \sum_{s=0}^j \text{wd}((\ell_1 - 2 - s)\text{-gap}_t)(u) \geq t \right\}.$$

Clearly,  $\overline{m} \in \{0, \dots, \ell_1 - 3\}$ . Hence, the minimum of  $\text{BShad}_t(u)_{(k_2, \ell_1 + 1)}$  is the  $t$ -spread monomial,

$$x_{i_1} \dots x_{i_{(\ell_1 - 2 - \overline{m})}} \left( \prod_{\nu = \ell_1 - \overline{m} - 1}^{\ell_1 + 1} x_{k_2 + t(\nu - 1) + 1} \right). \quad (12)$$

More in details, in order to obtain  $\min(\text{BShad}_t(u)_{(k_2, \ell_1 + 1)})$ , we decrease the minimum admissible number of variables to the right of  $x_{k_2 + t(\ell_1 - 1) + 1}$  until we obtain a  $t$ -spread monomial.

Finally, setting

$$m = \min \left\{ j \in \mathbb{Z}_{\geq -1} : (k_2 + t(\ell_2 - 1) + 1 - i_{(\ell_1 - 1)}) + \sum_{s=0}^j \text{wd}((\ell_1 - 2 - s)\text{-gap}_t)(u) \geq t \right\},$$

the assertion follows. Note that if we set  $\bar{m} = -1$  in (12), then we obtain the monomial  $v$  in (11). This is the case when  $k_2 + t(\ell_1 - 1) + 1 - i_{(\ell_1 - 1)} \geq t$ .  $\square$

**Remark 4.6** Observe that the integer  $m$  in Lemma 4.5 can be written in a more suitable way. In fact, setting

$$\tilde{m} = (k_2 + t((\ell_1 + 1) - 1) + 1 - i_{(\ell_1 - 1)}) + \sum_{s=0}^j \text{wd}((\ell_1 - 2 - s)\text{-gap}_t)(u),$$

one has

$$\begin{aligned} \tilde{m} &= k_2 + t\ell_1 + 1 - i_{(\ell_1 - 1)} + \sum_{s=0}^j [i_{(\ell_1 - 1 - s)} - i_{(\ell_1 - 2 - s)} - t] \\ &= k_2 + t\ell_1 + 1 - i_{(\ell_1 - 2 - j)} - (j + 1)t. \end{aligned}$$

Thus,

$$m = \min \left\{ j \in \mathbb{Z}_{\geq -1} : k_2 + t\ell_1 + 1 - i_{(\ell_1 - 2 - j)} \geq (j + 2)t \right\}.$$

Lemma 4.5 assures the correctness of the following algorithm.

**Construction 4.7** Let  $n, t$  be two positive integers. Let  $(k_1, \ell_1), (k_2, \ell_2)$  be two pairs of positive integers such that  $k_1 > k_2$ ,  $2 \leq \ell_1 < \ell_2$  and  $k_i + t(\ell_i - 1) + 1 \leq n$ , for  $i = 1, 2$ .

Let  $u = x_{i_1} x_{i_2} \cdots x_{i_{\ell_1}}$ ,  $1 \leq i_1 < i_2 < \cdots < i_{\ell_1} \leq n$  be a monomial of  $A^t(k_1, \ell_1)$ . Hence,  $i_{\ell_1} = k_1 + t(\ell_1 - 1) + 1$ . We want to determine

$$v = \min \text{BShad}_t(u)_{(k_2, \ell_2)}.$$

By Lemma 4.5, we can get the monomial

$$\tilde{v} = \min \text{BShad}_t(u)_{(k_2, \ell_1 + 1)} = x_{i_1} \cdots x_{i_{(\ell_1 - 2 - m)}} \left( \prod_{\nu=\ell_1 - m - 1}^{\ell_1 + 1} x_{k_2 + t(\nu - 1) + 1} \right),$$

with  $m = \min \left\{ j \in \mathbb{Z}_{\geq -1} : k_2 + t\ell_1 + 1 - i_{(\ell_1 - 2 - m)} \geq (m + 2)t \right\}$ . Hence, if  $\ell_2 = \ell_1 + 1$ , then  $v = \tilde{v}$ .

Otherwise,

$$\begin{aligned} v = \min \text{BShad}_t(u)_{(k_2, \ell_2)} &= \tilde{v} \cdot x_{k_2+t[(\ell_1+2)-1]+1} \cdots x_{k_2+t(\ell_2-1)+1} \\ &= x_{i_1} \cdots x_{i_{\ell_1-2-m}} \left( \prod_{\nu=\ell_1-m-1}^{\ell_1+1} x_{k_2+t(\nu-1)+1} \right) x_{k_2+t[(\ell_1+2)-1]+1} \cdots x_{k_2+t(\ell_2-1)+1}. \end{aligned}$$

Finally,

$$\min \text{BShad}_t(u)_{(k_2, \ell_2)} = x_{i_1} \cdots x_{i_{\ell_1-2-m}} \left( \prod_{\nu=\ell_1-m-1}^{\ell_2} x_{k_2+t(\nu-1)+1} \right).$$

**Example 4.8** Let  $S = K[x_1, \dots, x_{19}]$ ,  $t = 4$ ,  $(k_1, \ell_1) = (7, 3)$ ,  $(k_2, \ell_2) = (6, 4)$ . Let  $u = x_{i_1}x_{i_2}x_{i_3} = x_6x_{10}x_{16}$  be a monomial of  $A^t(k_1, \ell_1) = A^4(7, 3)$ . We want to determine

$$\min \text{BShad}_t(u)_{(k_2, \ell_2)} = \min \text{BShad}_4(x_6x_{10}x_{16})_{(6,4)}.$$

We will use Construction 4.7. One can observe that

$$k_2 + t(\ell_2 - 1) + 1 - i_{\ell_1-1} = 19 - 10 = 9 > 4 = t.$$

Hence, we have  $m = -1$  and so we do not need to consider the widths of any  $j$ -gap $_t$  of  $u$ . Therefore,

$$\begin{aligned} \min \text{BShad}_4(x_6x_{10}x_{16})_{(6,4)} &= x_{i_1} \cdots x_{i_{\ell_1-2-m}} \left( \prod_{\nu=\ell_1-m-1}^{\ell_2} x_{k_2+t(\nu-1)+1} \right) \\ &= x_{i_1}x_{i_2} \left( \prod_{\nu=3}^4 x_{6+4(\nu-1)+1} \right) \\ &= x_6x_{10}x_{15}x_{19}. \end{aligned}$$

## 5. AN EXAMPLE

In this Section, we provide an example that illustrates Theorem 4.4.

**Example 5.1** Let  $n = 25$ ,  $t = 3$ ,

$$\mathcal{C} = \{(k_1, \ell_1), (k_2, \ell_2), (k_3, \ell_3), (k_4, \ell_4)\} = \{(6, 2), (5, 4), (4, 5), (3, 7)\}$$

and

$$a = (a_1, a_2, a_3, a_4) = (2, 1, 3, 2).$$

Firstly, we *decompose*  $n$  as the sum  $d + kt$  with  $d = 1$  and  $k = 8$ , *i.e.*,  $n = 25 = 1 + 8 \cdot 3$ .



On the other hand,  $n - t - 1 = 21$  and, since  $\ell_1 = 2$ , we have to consider the integer  $k + \lfloor \frac{d-3}{t} \rfloor + 1 = 8$ . It follows that the required bounds

$$n - t - 1 \geq k_1 > k_2 > k_3 > k_4 \geq 1, \quad 2 \leq \ell_1 < \ell_2 < \ell_3 < \ell_4 \leq k + \left\lfloor \frac{d-3}{t} \right\rfloor + 1$$

are satisfied. According to Theorem 4.4, a 3-spread strongly stable ideal  $I$  of  $S = K[x_1, \dots, x_{25}]$  with corner sequence  $\mathcal{C}$  and corner values sequence  $a$  does exist if and only if condition (2) in the theorem is satisfied. Hence, we need to verify the bounds  $a_i \leq n_i$  for all  $i = 1, \dots, 4$ .

First of all, let  $v_4 = x_{k_4+1} \cdots x_{k_4+t(\ell_4-1)+1} = x_4 x_7 x_{10} x_{13} x_{16} x_{19} x_{22}$  be the smallest monomial of  $A^t(k_4, \ell_4) = A^3(3, 7)$ . Let  $w_4 \in A^3(3, 7)$  such that  $A_4 = [w_4, v_4]$  has cardinality  $a_4 = 2$ . We have  $w_4 = x_3 x_7 x_{10} x_{13} x_{16} x_{19} x_{22}$ . On the other hand,

$$\begin{aligned} n_4 &= |\{u \in A^3(3, 7) : u \geq v_4\}| = |[x_1 x_4 x_7 x_9 x_{11} x_{14} x_{22}, x_3 x_7 x_{10} x_{13} x_{16} x_{19} x_{22}]| \\ &= |A^3(3, 7)| = \binom{3+7-1}{7-1} = \binom{9}{6} = 84, \end{aligned}$$

and  $2 = a_4 \leq n_4 = 84$ .

Let

$$\begin{aligned} v_3 &= \min \{u \in A^t(k_3, \ell_3) : \max A_4 \notin \text{BShad}_t(u)_{(k_4, \ell_4)}\} \\ &= \min \{u \in A^3(4, 5) : x_3 x_7 x_{10} x_{13} x_{16} x_{19} x_{22} \notin \text{BShad}_3(u)_{(3, 7)}\} \\ &= x_3 x_6 x_{11} x_{14} x_{17}. \end{aligned}$$

Let  $w_3 \in A^3(4, 5)$  be the monomial such that  $|A_3| = |[w_3, v_3]| = a_3 = 3$ . It is  $w_3 = x_3 x_6 x_{10} x_{13} x_{17}$ . Hence,

$$A_3 = \{x_3 x_6 x_{10} x_{13} x_{17}, x_3 x_6 x_{10} x_{14} x_{17}, x_3 x_6 x_{11} x_{14} x_{17}\}.$$

In order to calculate

$$n_3 = |\{u \in A^3(4, 5) : u \geq v_3\}| = |[x_1 x_4 x_7 x_{10} x_{17}, x_3 x_6 x_{11} x_{14} x_{17}]|,$$

we consider the highlighted binomial coefficients of the following binomial decompositions:

$$\begin{aligned} \binom{k_3+\ell_3-1}{\ell_3-1} &= \binom{8}{4} = \boxed{\binom{7}{3}} + \boxed{\binom{6}{3}} + \binom{5}{3} + \binom{4}{3} + \binom{3}{3} \\ &\qquad \qquad \qquad \binom{5}{3} = \binom{4}{2} + \binom{3}{2} + \binom{2}{2} \\ &\qquad \qquad \qquad \binom{4}{2} = \boxed{\binom{3}{1}} + \boxed{\binom{2}{1}} + \binom{1}{1}. \end{aligned}$$

Hence,  $n_3 = \binom{7}{3} + \binom{6}{3} + \binom{3}{1} + \binom{2}{1} + \binom{0}{0} = 61$  and  $3 = a_3 \leq n_3 = 61$ .

Let

$$\begin{aligned}
v_2 &= \min \{u \in A^t(k_2, \ell_2) : \max A_3 \notin \text{BShad}_t(u)_{(k_3, \ell_3)}\} \\
&= \min \{u \in A^3(5, 4) : x_3x_6x_{10}x_{13}x_{17} \notin \text{BShad}_3(u)_{(4,5)}\} \\
&= x_3x_6x_9x_{15}.
\end{aligned}$$

Since  $a_2 = 1$ , we have  $A_2 = [w_2, v_2] = \{v_2\}$ , *i.e.*,  $w_2 = v_2$ . Now, we evaluate  $n_2 = |\{u \in A^3(5, 4) : u \geq v_2\}| = |[x_1x_4x_7x_{15}, x_3x_6x_9x_{15}]|$ . We consider the following binomial decompositions:

$$\begin{aligned}
\binom{k_2+\ell_2-1}{\ell_2-1} &= \binom{8}{3} = \boxed{\binom{7}{2} + \binom{6}{2}} + \binom{5}{2} + \binom{4}{2} + \binom{3}{2} + \binom{2}{2} \\
\binom{5}{2} &= \binom{4}{1} + \binom{3}{1} + \binom{2}{1} + \binom{1}{1} \\
\binom{4}{1} &= \binom{3}{0} + \binom{2}{0} + \binom{1}{0} + \binom{0}{0}.
\end{aligned} \tag{13}$$

Hence,  $n_2 = \binom{7}{2} + \binom{6}{2} + \binom{0}{0} = 37$  and  $1 = a_2 \leq n_2 = 37$ .

Finally,  $v_1 = \min \{u \in A^t(k_1, \ell_1) : \max A_2 \notin \text{BShad}_t(u)_{(k_2, \ell_2)}\} = x_2x_{10}$ . Since  $a_1 = 2$ , we have  $A_2 = [x_1x_{10}, x_2x_{10}] = \{x_1x_{10}, x_2x_{10}\}$ , and  $a_1 = 2 = n_1$ .

Now, we construct a 3-spread strongly stable ideal  $I$  with corner sequence  $\text{Corn}(I) = \mathcal{C}$  and  $a(I) = a$  by refining the bounds for the  $a_i$ 's (see Theorem 4.4, (iii)).

-  $a_1 = 2$ , since  $p_1 = 0$ ,  $a_1 = n_1 - p_1 = n_1 = 2$ , we set

$$G(I)_{\ell_1} = G(I)_2 = B_3\{x_1x_{10}, x_2x_{10}\}.$$

Note that  $x_1x_{10}, x_2x_{10}$  are the greatest monomials of  $A^t(k_1, \ell_1) = A^3(6, 2)$ , with respect to  $>_{\text{slex}}$ .

- Let us consider the corner  $(k_2, \ell_2) = (5, 4)$ . Since  $\min \text{BShad}_3([I_2]_3)_{(5,4)} = x_2x_9x_{12}x_{15}$ , then  $\max(A^t(k_2, \ell_2) \setminus \text{BShad}_t([I_1]_2)_{(5,4)})$  is the largest monomial of  $A^t(k_2, \ell_2)$  following  $x_2x_9x_{12}x_{15}$  respect to  $>_{\text{slex}}$ , *i.e.*,  $u_{2,1} = x_3x_6x_9x_{15}$ . In order to calculate  $p_2 = |\{v \in A^3(5, 4) : v > u_{2,1}\}|$ , we need to consider the same binomial decompositions of (13). Thus,  $p_2 = \binom{7}{2} + \binom{6}{2} = 36$ ,  $a_2 = 1 \leq n_2 - p_2 = 37 - 36 = 1$ . Hence, we set

$$G(I)_{\ell_2} = G(I)_4 = B_3\{x_3x_6x_9x_{15}\} \setminus \text{Shad}_3^2([I_2]_3).$$

- Let us consider the corner  $(k_3, \ell_3) = (4, 5)$ . Since  $\min \text{BShad}_3([I_4]_3)_{(4,5)} = x_3x_6x_9x_{14}x_{17}$ , then  $\max(A^t(k_3, \ell_3) \setminus \text{BShad}_3([I_4]_3)_{(4,5)})$  is the largest monomial of  $A^t(k_3, \ell_3)$  that follows  $x_3x_6x_9x_{14}x_{17}$  respect to  $>_{\text{slex}}$ . It is  $u_{3,1} = x_3x_6x_{10}x_{13}x_{17}$ . In order to evaluate  $p_3 = |\{v \in A^3(4, 5) : v >$

$u_{3,1}\}$ , we consider the following suitable binomial decompositions:

$$\begin{aligned} \binom{k_3+\ell_3-1}{\ell_3-1} &= \binom{8}{4} = \boxed{\binom{7}{3} + \binom{6}{3}} + \binom{5}{3} + \binom{4}{3} + \binom{3}{3} \\ \binom{5}{3} &= \binom{4}{2} + \binom{3}{2} + \binom{2}{2} \\ \binom{4}{2} &= \boxed{\binom{3}{1}} + \binom{2}{1} + \binom{1}{1}. \end{aligned}$$

Therefore,  $p_3 = \binom{7}{3} + \binom{6}{3} + \binom{3}{1} = 58$  and  $3 = a_3 \leq n_3 - p_3 = 61 - 58 = 3$ , and we can construct  $G(I)_{\ell_3} = G(I)_5$ . We set

$$G(I)_5 = B_3\{x_3x_6x_{10}x_{13}x_{17}, x_3x_6x_{10}x_{14}x_{17}, x_3x_6x_{11}x_{14}x_{17}\} \setminus \text{Shad}_3^1([I_4]_3).$$

- Finally, we consider the corner  $(k_4, \ell_4) = (3, 7)$ . The smallest monomial of  $\text{BShad}_3([I_5]_3)_{(3,7)}$  is  $x_3x_6x_{10}x_{13}x_{16}x_{19}x_{22}$ , then the largest monomial of  $A^t(k_4, \ell_4) \setminus \text{BShad}_3([I_5]_3)_{(3,7)}$  is  $u_{4,1} = x_3x_7x_{10}x_{13}x_{16}x_{19}x_{22}$ . In order to compute  $p_4 = |\{v \in A^3(3, 7) : v > u_{4,1}\}|$ , we consider the following binomial decompositions:

$$\begin{aligned} \binom{9}{6} &= \boxed{\binom{8}{5} + \binom{7}{5}} + \binom{6}{5} + \binom{5}{5} \\ \binom{6}{5} &= \boxed{\binom{5}{4}} + \binom{4}{4}. \end{aligned}$$

Hence  $p_4 = \binom{8}{5} + \binom{7}{5} + \binom{5}{4} = 82$ , and  $2 = a_4 \leq n_4 - p_4 = 84 - 82 = 2$ . Thus, we can construct  $G(I)_{\ell_4} = G(I)_7$ . We set

$$G(I)_7 = B_3\{x_3x_7x_{10}x_{13}x_{16}x_{19}x_{22}, x_4x_7x_{10}x_{13}x_{16}x_{19}x_{22}\} \setminus \text{Shad}_3^2([I_5]_3).$$

Finally, we have constructed a 3-spread strongly stable ideal

$$\begin{aligned} I = & B_3(x_1x_{10}, \mathbf{x_2x_{10}}, \mathbf{x_3x_6x_9x_{15}}, x_3x_6x_{10}x_{13}x_{17}, x_3x_6x_{10}x_{14}x_{17}, \mathbf{x_3x_6x_{11}x_{14}x_{17}}, \\ & x_3x_7x_{10}x_{13}x_{16}x_{19}x_{22}, \mathbf{x_4x_7x_{10}x_{13}x_{16}x_{19}x_{22}}) \\ = & (x_1x_4, x_1x_5, x_1x_6, x_1x_7, x_1x_8, x_1x_9, x_1x_{10}, x_2x_5, x_2x_6, x_2x_7, x_2x_8, x_2x_9, \mathbf{x_2x_{10}}, \\ & x_3x_6x_9x_{12}, x_3x_6x_9x_{13}, x_3x_6x_9x_{14}, \mathbf{x_3x_6x_9x_{15}}, \\ & x_3x_6x_{10}x_{13}x_{16}, x_3x_6x_{10}x_{13}x_{17}, x_3x_6x_{10}x_{14}x_{17}, \mathbf{x_3x_6x_{11}x_{14}x_{17}}, \\ & x_3x_7x_{10}x_{13}x_{16}x_{19}x_{22}, \mathbf{x_4x_7x_{10}x_{13}x_{16}x_{19}x_{22}}). \end{aligned}$$

with corner sequence  $\mathcal{C}$  and corner values sequence  $a$ . The highlighted monomial are the 3-spread Borel generators of  $I$ . The Betti table of  $I$  is

	0	1	2	3	4	5	6
Tot	: 23	77	117	100	51	15	2
2	: 13	42	70	70	42	14	2
3	: -	-	-	-	-	-	-
4	: 4	14	20	15	6	1	-
5	: 4	15	21	13	3	-	-
6	: -	-	-	-	-	-	-
7	: 2	6	6	2	-	-	-

**Remark 5.2** All the algorithmic and constructive methods analyzed in this paper have been implemented into a *Macaulay2* packages “TSpreadIdeals.m2”, by the authors themselves, and tested with *Macaulay2* 1.17.2.1. To the best of our knowledge, specific packages for manage  $t$ -spread ideals are not yet available in the most well-known symbolic algebra systems (*e.g.*, *Macaulay2*, CoCoA or Singular). Hence, we believe that such a package could be useful since the topics covered have been introduced recently and represent a new line of research ([4, 6, 7, 16, 13]).

All the examples in this paper have been checked or constructed by this package.

#### REFERENCES

- [1] J. Abbott, A. M. Bigatti, L. Robbiano, *CoCoA: a system for doing computations in Commutative Algebra*. Available at <http://cocoa.dima.unige.it>.
- [2] L. Amata, M. Crupi. Computation of graded ideals with given extremal Betti numbers in a polynomial ring. *J. Symbolic Computation* **93** (2019), 120–132.
- [3] L. Amata, M. Crupi. On the extremal Betti numbers of squarefree monomial ideals. *Int. Electron. J. Algebra* **30** (2021), 168-202.
- [4] L. Amata, M. Crupi. Extremal Betti Numbers of  $t$ -Spread Strongly Stable Ideals. *Mathematics* **7** (2019), 695.
- [5] L. Amata, A. Ficarra, M. Crupi. Upper bounds for Extremal Betti Numbers of  $t$ -Spread Strongly Stable Ideals. *Bull. Math. Soc. Sci. Math. Roumanie (N.S.)*, to appear 2021.
- [6] C. Andrei, V. Ene, B. Lajmiri. Powers of  $t$ -spread principal Borel ideals. *Archiv der Mathematik* **112** (2019), 587–597.
- [7] C. Andrei-Ciobanu, Kruskal–Katona Theorem for  $t$ -spread strongly stable ideals. *Bull. Math. Soc. Sci. Math. Roumanie (N.S.)* **62(110)(2)** (2019), 107–122.
- [8] A. Aramova, J. Herzog, T. Hibi. Squarefree lexsegment ideals. *Math.Z.* **228** (1998), 353–378.
- [9] D. Bayer, H. Charalambous, S. Popescu. Extremal Betti numbers and Applications to Monomial Ideals. *J. Algebra* **221** (1999), 497–512.
- [10] M. Crupi. Extremal Betti numbers of graded modules, *J. Pure Appl. Algebra* **220**, (2016), 2277–2288.

- [11] M. Crupi. Computing general strongly stable modules with given extremal Betti numbers. *J. Com. Alg.* **12(1)** (2020), 53–70
- [12] M. Crupi, C. Ferrò. Squarefree monomial modules and extremal Betti numbers, *Algebra Colloq.* **23(3)** (2016), 519–530.
- [13] R. Dinu. Gorenstein  $T$ -spread Veronese algebras. *Osaka J. Math.* **57(4)** (2020), 935–947.
- [14] S. Eliahou, M. Kervaire. Minimal resolutions of some monomial ideals, *J. Algebra* **129** (1990), 1–25.
- [15] D. Eisenbud, *Commutative Algebra with a view toward Algebraic Geometry*, Springer-Verlag, 1995.
- [16] V. Ene, J. Herzog, A. A. Qureshi.  $t$ -spread strongly stable monomial ideals\*, *Communications in Algebra* **47(12)** (2019), 5303–5316.
- [17] D. R. Grayson, M. E. Stillman, *Macaulay2, a software system for research in algebraic geometry*. Available at <http://www.math.uiuc.edu/Macaulay2>.
- [18] J. Herzog, T. Hibi, *Monomial ideals*, Graduate texts in Mathematics **260**, Springer-Verlag, 2011.
- [19] J. Herzog, L. Sharifan, M. Varbaro. The possible extremal Betti numbers of a homogeneous ideal. *Proceedings of the AMS* **142** (2014), 1875–1891.
- [20] R. P. Stanley. The upper bound conjecture and Cohen–Macaulay rings, *Studies in Appl. Math.* **54(2)** (1975), 135–142.
- [21] E. Miller, B. Sturmfels. *Combinatorial Commutative Algebra*, Graduate Texts in Mathematics, Springer-Verlag **227**, 2005.