

Riemann-Roch inequality for smooth tropical toric surfaces

Ken Sumi

Abstract

For a divisor D on a tropical variety X , we define two amounts in order to estimate the value of $h^0(X, D)$, which are described by terms of global sections and computed more easily than $h^0(X, D)$. As an application of its estimation, we show that a Riemann-Roch inequality holds for smooth tropical toric surfaces.

1 Introduction

In this paper, we prove a Riemann-Roch inequality for smooth tropical toric surfaces by calculating $h^0(X, D)$, which is a higher dimensional generalization of the rank of divisor.

For a divisor D , the rank of divisor $r(D)$ on a finite graph is defined by Baker-Norine[3]. Gathmann-Kerber[8] and Mikhakin-Zharkov[16] generalized the definition of the rank of divisor for tropical curves and showed a Riemann-Roch theorem for compact tropical curves.

The rank of divisor is defined as the maximum number k such that $|D - E| \neq \emptyset$ for arbitrary effective divisor E of degree k . It is an alternative notion of the dimension of the projectivization of $H^0(X, \mathcal{O}(D))$, however, the rank of divisor $r(D)$ cannot be given only by the correspondence tropical module $\Gamma(X, \mathcal{O}(D))$. For example, [20][Example 6.5] says that there are two pairs (C, D) and (C', D') of tropical curves and divisors such that spaces of global sections $H^0(C, \mathcal{O}(D))$, $H^0(C', \mathcal{O}(D'))$ are isomorphic, while $r(D) \neq r(D')$.

A higher dimensional generalization of $r(D)$ (precisely, $r(D) + 1$) is introduced in Cartwright[6], denoted by $h^0(X, D)$, and formulate a Riemann-Roch inequality for tropical surfaces. The author computed the value of $h^0(X, D)$ for a divisor D on a tropical torus X and showed that the Riemann-Roch inequality formulated by Cartwright holds for tropical Abelian surfaces in [21].

In this paper, we generalize the method of the computation of the value of $h^0(X, D)$ used in the author's previous work [21] for general tropical varieties and divisors. Concretely, we define two amounts h_a^0 and h_b^0 . We define h_a^0 as the maximal number k such that there is a tropical submodule of $\Gamma(X, \mathcal{O}(D))$ satisfying that it is generated by k elements and its generators satisfies the identity theorem, that is, two generators are distinct if and only if these are

distinct on any open subset of X . In general, the identity theorem does not hold in tropical geometry since tropical regular function means a piecewise-linear convex function with integral slopes. We define h_b^0 as the maximal number k such that for all $x \in X$ there exist k sections satisfying that each two sections of these k sections are distinct ‘around’ x .

Then we get the following inequality.

Proposition 1 (Proposition 22).

$$h_a^0(X, D) \leq h^0(X, D) \leq h_b^0(X, D)$$

When we have a concrete minimal generating set of $\Gamma(X, \mathcal{O}(D))$, we can compute these amounts h_a^0 and h_b^0 more easily than computing $h^0(X, D)$ since $h^0(X, D)$ is defined by terms of divisors.

Tropical toric varieties are studied by many mathematicians. For example, Mayer[13] and Shaw [18] studied intersection theory on tropical toric varieties, especially smooth tropical toric surfaces.

The author shows the following basic statement.

Proposition 2 (Proposition 36). Let X be a smooth tropical toric variety and D be a divisor on X . Then

$$\Gamma(X, \mathcal{O}(D)) \cong \left\langle \sum_{\text{div}(x^m)+D \geq 0} \mathbb{T} \cdot x^m \right\rangle.$$

By using the above two propositions and classical Riemann-Roch inequality for smooth tropical toric surfaces, we get the value of $h^0(X, D)$ and show the Riemann-Roch inequality for smooth tropical toric surfaces.

2 Preliminaries

2.1 Tropical modules

The tropical semifield \mathbb{T} is the set $\mathbb{R} \cup \{-\infty\}$ equipped with the tropical sum “ $x + y$ ” = $\max\{x, y\}$ and the tropical product “ xy ” = $x + y$ for $x, y \in \mathbb{T}$. A commutative semigroup V equipped with the unit $\{-\infty\}$ and a scalar product $\mathbb{T} \times V \rightarrow V$ is called a *tropical module* or a \mathbb{T} -*module* if the following conditions are satisfied:

- “ $(x + y)v$ ” = “ $xv + yv$ ” for any $x, y \in \mathbb{T}$, any $v \in V$;
- “ $x(v + w)$ ” = “ $xv + xw$ ” for any $x \in \mathbb{T}$, any $v, w \in V$;
- “ $x(yv)$ ” = “ $(xy)v$ ” for any $x, y \in \mathbb{T}$, any $v \in V$;
- “ $0 \cdot v$ ” = v for the multiplicative unit $0 \in \mathbb{T}$ and any $v \in V$;
- if “ xv ” = “ yv ” for some $x, y \in \mathbb{T}$ and $v \in \mathbb{T}$, then $x = y$ or $v = -\infty$.

Let V, W be tropical modules. A map $f: V \rightarrow W$ is called a *tropical linear morphism* if $f("v_1 + v_2") = "f(v_1) + f(v_2)"$ and $f("tv") = "tf(v)"$ for any $v, v_1, v_2 \in V, t \in \mathbb{T}$. If there is the set theoretical inverse map f^{-1} and it is also a tropical linear morphism, then we call f an isomorphism.

A function $f: \mathbb{R}^n \rightarrow \mathbb{T}$ is a *tropical polynomial* (resp. *tropical Laurent polynomial*) if f is a constant map to $\{-\infty\}$ or is of the form $f(x) = \max_{j \in S} \{a_j + j \cdot x\}$, where S is a finite subset of $(\mathbb{Z}_{\geq 0})^n$ (resp. \mathbb{Z}^n) and $a_j \in \mathbb{R}$. A tropical polynomial naturally extends to a function from \mathbb{T}^n to \mathbb{T} . The set of tropical polynomials on \mathbb{R}^n is naturally a tropical module.

A \mathbb{Z} -*affine linear function* is a tropical Laurent monomial, that is, an affine linear function on \mathbb{R}^n whose slope is in $(\mathbb{Z}^n)^*$.

For a matrix $(t_{ij})_{1 \leq i, j \leq l}$ with $t_{ij} \in \mathbb{T}$, we define the tropical determinant $\det_{\text{trop}}(t_{i,j})$ to be

$$\max_{\sigma \in \mathfrak{S}_l} \left\{ \sum_{i=1}^l t_{\sigma(i)i} \right\},$$

where \mathfrak{S}_l is the symmetric group of $\{1, \dots, l\}$.

Definition 3. Let M be a tropical module. An element $m \in M$ is an *extremal* if there exist no elements $a, b \in M$ such that " $a + b$ " = m and $a, b \neq m$

2.2 Tropical varieties

A tropical variety is defined as a balanced polyhedral complex with weight 1 whose each cells are locally a convex rational polyhedron in \mathbb{T}^n .

Definition 4. The *sedentarity* of a point $t = (t_1, \dots, t_n) \in \mathbb{T}^n$ is the set $I := \{i \in \{1, \dots, n\} \mid t_i = -\infty\}$. A *face* of \mathbb{T}^n of sedentarity I is a set $\{t \in \mathbb{T}^n \mid t_i = \infty \text{ for all } i \in I\}$. The face of sedentarity \emptyset is \mathbb{T}^n itself.

Definition 5. An *rational polyhedron* P in \mathbb{T}^n is a finite intersection of a face of \mathbb{T}^n and half-spaces defined by finite inequalities $a_i \cdot x + b_i \geq 0, i \in I$ where $x \in \mathbb{R}^n, a \in \mathbb{Z}^n$, and $b \in \mathbb{R}$. A *face of* P is an intersection of P and a face of \mathbb{T}^n or a hyperplane $a \cdot x + b = 0, a \in \mathbb{Z}^n, b \in \mathbb{R}$ appearing the definition of P .

A face or a point is *inner* if it is contained in \mathbb{R}^n .

Definition 6. An *rational polyhedral complex* in \mathbb{T}^n is a union of a collection of rational polyhedra in \mathbb{T}^n such that the intersection of any subcollection is a common face of all rational polyhedra in the subcollection.

A *cell of rational polyhedral complex* \mathcal{P} is a face of a rational polyhedron in the collection. A *facet of* \mathcal{P} is face which is not contained in any other face.

The *dimension* of an rational polyhedral complex is the maximal dimension of rational polyhedra in the collection. If all facets of an rational polyhedral complex is n -dimensional, then the complex is said to be pure n -dimensional or pure of dimension n .

An *n -dimensional weighted rational polyhedral complex* is a pure n -dimensional rational polyhedral complex with a weight function which is a \mathbb{Z} -valued function

on the set of n -dimensional cells. Then we can consider an rational polyhedral complex as a weighted complex whose weight function is the constant 1.

Definition 7. Let P be a d -dimensional weighted rational polyhedral complex in \mathbb{T}^n with a weight function w and E be a $(d-1)$ -dimensional cell of P with $E \cap \mathbb{R}^n \neq \emptyset$. Let F_1, \dots, F_k be the d -dimensional cells adjacent to E and $\pi: \mathbb{R}^n \rightarrow \mathbb{R}^{n-d+1}$ be the projection whose kernel is parallel to E . Then $\pi(F_1), \dots, \pi(F_k)$ are rays adjacent to the point $\pi(E)$. Let $v_1, \dots, v_k \in \mathbb{Z}^{n-d+1}$ be the primitive vectors parallel to $\pi(F_1), \dots, \pi(F_k)$ respectively. Then P is *balanced at E* if $\sum_{i=1, \dots, k} w(F_i)v_i = 1$ and P is *balanced* if P is balanced at E for all $(d-1)$ -dimensional faces E of P .

Definition 8. A *tropical variety* is a topological space X with an atlas $\{(U_\alpha, \Phi_\alpha)\}_\alpha$ such that

- U_α is homeomorphic to an open subset of an n -dimensional balanced rational polyhedral complex with weight 1 in \mathbb{T}^{n_α} by $\Phi_\alpha: U_\alpha \rightarrow \mathbb{T}^{n_\alpha}$, which is an isomorphism to its image $\Phi(U_\alpha)$;
- $\Phi_\alpha(U_\alpha) \cap \mathbb{R}^{n_\alpha} \neq \emptyset$;
- the transition function on the “finite part” $\iota_{n_\alpha}^{-1} \circ \Phi_\alpha \circ \Phi_\beta^{-1} \circ \iota_{n_\beta}$ is a restriction of a \mathbb{Z} -affine linear map from \mathbb{R}^{n_β} to \mathbb{R}^{n_α} . Here $\iota_n: \mathbb{R}^n \rightarrow \mathbb{T}^n$ is the natural embedding.

If every U_α is homeomorphic to an open subset of an n -dimensional balanced rational polyhedral complex, a tropical variety X is called *n -dimensional*. Especially, 1-dimensional tropical varieties are called *tropical curves* and 2-dimensional tropical varieties are called *tropical surfaces*.

Definition 9. Let U be an open subset of a tropical variety. A continuous function $f: U \rightarrow \mathbb{T} \cup \{-\infty\}$ is said to be *rational* if there exist charts $\{(U_{\alpha_i}, \Phi_{\alpha_i})\}_i$ and tropical polynomials $g_{\alpha_i}, h_{\alpha_i}$ on \mathbb{T}^{α_i} satisfying that $U = \bigcup U_{\alpha_i}$ and $f \circ \Phi_{\alpha_i}^{-1} = g_{\alpha_i} - h_{\alpha_i}$ on $\Phi_{\alpha_i}(U_{\alpha_i})$.

It is said to be *regular* if we can take a \mathbb{Z} -affine linear function as h_{α_i} in the above notation.

Definition 10. The *sedentarity number* of a point p of a tropical variety X is the minimal cardinal of the sedentarity of $\Phi_\alpha(p) \in U_\alpha$ when α runs..

A point p of a tropical variety X is said to be *inner* if the sedentarity number of p is zero. We denote the set of inner points of X by X° .

2.3 Cartier divisors and line bundles

Definition 11. A (Cartier) divisor on X is a collection of pairs $D = \{(U_\alpha, f_\alpha)\}_\alpha$ of an open subset U_α and a rational function f_α on U_α satisfying that $\bigcup_\alpha U_\alpha = X$ for any two charts U_α and U_β with $U_\alpha \cap U_\beta \neq \emptyset$ there is an invertible tropical function $d_{\alpha, \beta}$ on $U_\alpha \cap U_\beta$ which coincides with $f_\alpha - f_\beta$ at any inner point of $U_\alpha \cap U_\beta$. We identify two divisors $\{(U_\alpha, f_\alpha)\}_\alpha, \{(V_\beta, g_\beta)\}_\beta$ if $\{(U_\alpha, f_\alpha)\}_\alpha \cup \{(V_\beta, g_\beta)\}_\beta$ is also a divisor.

We can define the sum of two divisors $D = \{(U_\alpha, f_\alpha)\}_\alpha$ and $D' = \{(V_\beta, g_\beta)\}_\beta$ to be a divisor $D + D' := \{(U_\alpha \cap V_\beta, f_\alpha + g_\beta)\}_{\alpha, \beta}$.

For a divisor $D = \{(U_\alpha, f_\alpha)\}_\alpha$, $\text{supp } D$ is a set of non-smooth points of f_α for some α . We call that a divisor $D = \{(U_\alpha, f_\alpha)\}_\alpha$ *passes* p if $p \in \text{supp } D$.

Remark 12. For a divisor $\{(U_\alpha, f_\alpha)\}_\alpha$, each (U_α, f_α) defines a weighted balanced polyhedral complex in U_α of codimension one and we can glue these complexes. That is, a Cartier divisor defines a weighted balanced polyhedral complex in X of codimension one. Conversely a weighted balanced polyhedral complex in X of codimension one defines a Cartier divisor. Thus we can identify an equivalent class of a Cartier divisor and a weighted balanced polyhedral complex in X of codimension one.

Definition 13. A divisor D is *effective* if there exists a representation $\{(U_\alpha, f_\alpha)\}_\alpha$ of D such that all f_α are regular functions.

Two divisors $\{(U_\alpha, f_\alpha)\}_\alpha$ and $\{(V_\beta, g_\beta)\}_\beta$ are *linearly equivalent* if there exists a rational function h on X such that $f_\alpha = g_\beta + h$ on each $U_\alpha \cap V_\beta \neq \emptyset$.

For a divisor D , we denote a set of effective divisors linearly equivalent to D by $|D|$.

Definition 14 ([2], Definitions 1.5). Let X be an n -dimensional tropical variety. A *tropical line bundle* on X is a tuple $(L, \pi, \{U_i, \Psi_i\}_i)$ of a topological space L , a continuous surjection $\pi : L \rightarrow M$, an open covering $\{U_i\}$ called the trivializing covering and homeomorphisms $\Psi_i : \pi^{-1}(U_i) \cong U_i \times \mathbb{T}$ called trivializations which satisfy:

- The following diagram is commute:

$$\begin{array}{ccc} \pi^{-1}(U_i) & \xrightarrow{\Psi_i} & U_i \times \mathbb{T} \\ & \searrow \pi & \downarrow \\ & & U_i \end{array}$$

Here the vertical map is the first projection;

- For every i, j with $U_i \cap U_j \neq \emptyset$, there exists an invertible regular function $\varphi_{ij} : U_i \cap U_j \rightarrow \mathbb{R}$ such that $\Psi_j \circ \Psi_i^{-1} : (U_i \cap U_j) \times \mathbb{T} \rightarrow (U_i \cap U_j) \times \mathbb{T}$ is given by $(x, t) \mapsto (x, \varphi_{ij}(x)t)$. These φ_{ij} are called a *transition functions* of L .

It is clear that transition functions φ_{ij} satisfies the cocycle condition “ $\varphi_{ij} \cdot \varphi_{jk} = \varphi_{ik}$ ”. We identify two tropical line bundles $(L, \pi, \{U_i, \Psi_i\}_{i \in I})$ and $(L, \pi, \{U_j, \Psi_j\}_{j \in J})$ if $(L, \pi, \{U_i, \Psi_i\}_{i \in I \sqcup J})$ is a tropical line bundle on M , that is, satisfies the second condition above. We often write L for $(L, \pi, \{U_i, \Psi_i\}_i)$ to avoid heavy notation.

Definition 15. We take two line bundles $(L_1, \pi_1, \{U_{1i}, \Psi_{1i}\}), (L_2, \pi_2, \{U_{2j}, \Psi_{2j}\})$. Let $\varphi_{ij}^1, \varphi_{ij}^2$ be the correspondence transition functions. The tensor product

$L_1 \otimes L_2$ is defined to be the tropical line bundle on X whose transition functions are “ $\varphi_{ij}^1 \varphi_{ij}^2$ ” = $\varphi_{ij}^1 + \varphi_{ij}^2$.

The inverse L_1^{-1} of the line bundle L_1 is defined to be the tropical line bundle on X whose transition functions are “ $(\varphi_{ij}^1)^{-1}$ ” = $-\varphi_{ij}^1$.

Definition 16. Let $\pi: L \rightarrow X$ be a line bundle on a tropical variety X and U be an open subset of X . A function $s: U \rightarrow \pi^{-1}(U)$ is a *regular section* (resp. *rational section*) on U of L if $\pi \circ s$ is the identity map on U and $p_i \circ \Psi_i \circ s$ is a regular (resp. rational) function, where $p_i: U_i \times \mathbb{T} \rightarrow \mathbb{T}$ is the second projection. If $U = M$ then we call s a global regular (resp. rational) section or simply a regular (resp. rational) section. The set $\Gamma(U, L)$ of regular sections on U of L naturally has the structure of a tropical module.

A rational section s defines a divisor $\text{div } s = \{(U_i, p_i \circ \Psi_i \circ s)\}_i$. Conversely, a divisor $D = \{(U_\alpha, f_\alpha)\}_\alpha$ defines a line bundle $\mathcal{O}(D)$. This correspondence is compatible for a sum of divisors and a tensor product of line bundles.

Assume $\Gamma(X, \mathcal{O}(D)) \neq \emptyset$ and fix a regular section $s \in \Gamma(X, \mathcal{O}(D))$. For any regular section s' , we can define a rational function $f = \frac{s'}{s}$. Since s and s' are regular, $\text{div}(f) + D \geq 0$. Then we get a map $\Gamma(X, \mathcal{O}(D)) \rightarrow \{f \in \mathcal{M}(X) \mid \text{div}(f) + D \geq 0\}$. It is clear that this map is an isomorphism of \mathbb{T} -modules. Thus we get the following proposition.

Proposition 17.

$$\Gamma(X, \mathcal{O}(D)) \cong \{f \in \mathcal{M}(X) \mid \text{div}(f) + D \geq 0\}.$$

At the end of this section, we note the following proposition.

Proposition 18. [[11], Corollary 8] If a submodule M of the tropical module $\Gamma(X, \mathcal{O}(D))$ is finitely generated, this is generated by extremals. This generating set is minimal and unique up to tropical scalar multiplication.

Precisely this proposition is a generalization of [11], Corollary 8, and proved as the same way.

3 Rank of divisor

We consider a tropical variety X with an atlas $\mathcal{V} = \{V_\beta, \Phi_\beta\}$ and a Cartier divisor D on X .

Definition 19 (Cartwright [6], Definition 3.1). Let X be a tropical variety and D be a divisor on X . We define $h^0(X, D)$ as

$$h^0(X, D) := \min \left\{ k \in \mathbb{Z}_{\geq 0} \mid \begin{array}{l} \text{there exist } k \text{ points } p_1, \dots, p_k, \text{ such that} \\ \text{there is no divisor } E \in |D| \text{ which passes all } p_i \end{array} \right\}.$$

If there is no such k , then we define $h^0(X, D)$ as ∞ .

The value of $h^0(X, D)$ is difficult to compute in general. In order to estimate this, we define two amounts $h_a^0(X, D)$ and $h_b^0(X, D)$.

Let M be a finitely generated submodule of $\Gamma(X, \mathcal{O}(D))$ and s_1, \dots, s_k be a minimal generating set of M . By Proposition 18, this minimal generating set is unique up to tropical multiplications. Thus we can define $U(M)$ to be the complement of $\cup_{i=1}^k \text{supp}(\text{div } s_i)$ and let $\{U(M)_\alpha\}$ to be the set of connected components of $U(M)$.

Then restriction of generators $s_i|_{U(M)_\alpha \cap V_\beta}$ are affine linear functions on each open subset $U(M)_\alpha \cap V_\beta$. Thus we can assume that $M|_{U(M)_\alpha \cap V_\beta}$ is a tropical submodule of $\mathbb{T}[x_1, \dots, x_n]$. Here x_1, \dots, x_n are the coordinates of the ambient space of $\text{Im } \varphi_\beta$.

For $y \in U(M)_\alpha \cap V_\beta$, we define $l_M(y)$ to be the number of monomials in $M|_{U(M)_\alpha \cap V_\beta}$ up to tropical multiplications. It is clear that this number is independent for the choice of α, β . For $x \notin U(M)$, we define $U(M)_x$ to be the union of U_α such that its closure contains x , and define

$$l_M(x) := \max_{y \in U_x} l_M(y)$$

Definition 20. We define two numbers $h_a^0(X, D)$ and $h_b^0(X, D)$ as the following;

$$h_a^0(X, D) := \max \left\{ k \in \mathbb{Z}_{\geq 0} \mid \begin{array}{l} \exists k \text{ sections } s_1, \dots, s_k, \text{ s.t.} \\ \forall x \in X, l_{\langle s_1, \dots, s_k \rangle}(x) = k \end{array} \right\}$$

$$\begin{aligned} h_b^0(X, D) &:= \max \left\{ k \in \mathbb{Z}_{\geq 0} \mid \begin{array}{l} \forall x \in X, \exists k \text{ sections } s_1, \dots, s_k, \\ \text{s.t. } l_{\langle s_1, \dots, s_k \rangle}(x) = k \end{array} \right\} \\ &= \min_{x \in X} \{l_{\Gamma(X, \mathcal{O}(D))}(x)\} \end{aligned}$$

Remark 21. These numbers are more easy to compute than $h^0(X, D)$ when we have a concrete expression of a minimal generating set of $\Gamma(X, \mathcal{O}(D))$. Moreover, these coincide the cardinal of a minimal generating set of $\Gamma(X, \mathcal{O}(D))$ when the generators of $\Gamma(X, \mathcal{O}(D))$ have different slopes each other around any point $x \in X$.

Proposition 22 (Main result 1).

$$h_a^0(X, D) \leq h^0(X, D) \leq h_b^0(X, D)$$

Proof. When $\Gamma(X, \mathcal{O}(D)) = \{-\infty\}$ or this is generated only one element, it is clearly that $h^0(X, D) = h_a^0(X, D) = h_b^0(X, D)$. Thus we assume that $\Gamma(X, \mathcal{O}(D)) \neq \{-\infty\}$ and $\Gamma(X, \mathcal{O}(D))$ is generated by 2 or more elements.

Let $l := h_a^0(X, D)$. By definition of $h_a^0(X, D)$, we can take l sections s_1, \dots, s_l satisfying that $l_{\langle s_1, \dots, s_l \rangle}(x) = l$ for all $x \in X$. In order to show $h^0(X, D) \geq l$, we take arbitrary $l - 1$ -points $p_1, \dots, p_{l-1} \in X$.

Now we define a regular section of $\mathcal{O}(D)$ by the Vandermonde determinant

$$\begin{aligned} V_D(x; p_1, \dots, p_{l-1}) &:= \det_{\text{trop}} \begin{pmatrix} s_1(x) & \cdots & s_l(x) \\ s_1(p_1) & \cdots & s_l(p_1) \\ \vdots & \ddots & \vdots \\ s_1(p_{l-1}) & \cdots & s_l(p_{l-1}) \end{pmatrix} \\ &= \max_{\sigma \in \mathfrak{S}_l} \left\{ s_{\sigma(1)}(x) + \sum_{i=2}^l s_{\sigma(i)}(p_{i-1}) \right\} \\ &= \max_{i \in \{1, \dots, l\}} \left\{ s_i(x) + \max_{\sigma \in \mathfrak{S}_l, \sigma(1)=i} \sum_{j=2}^l s_{\sigma(j)}(p_{j-1}) \right\}. \end{aligned}$$

Lemma 23. $\text{div } V_D(x; p_1, \dots, p_{l-1})$ passes through the $l-1$ points p_1, \dots, p_{l-1} .

Proof. Let us show that $\text{div } V_D(x; p_1, \dots, p_{l-1})$ passes through p_1 . It is clear when the value $\text{div } V_D(p_1; p_1, \dots, p_{l-1})$ is infinite, thus we assume that $\text{div } V_D(p_1; p_1, \dots, p_{l-1})$ is finite.

Suppose that the maximum in $\text{div } V_D(p_1; p_1, \dots, p_{l-1})$ is realized by just one element i_{max} . We can assume this i_{max} is 1. Let σ_1 be an element satisfying that $\max_{\sigma \in \mathfrak{S}_l, \sigma(1)=1} \sum_{i=2}^l s_{\sigma(i)}(p_{i-1}) = \sum_{i=2}^l s_{\sigma_1(i)}(p_{i-1})$, that is, $\text{div } V_D(p_1; p_1, \dots, p_{l-1}) = s_1(p_1) + s_{\sigma_1(1)}(p_1) + s_{\sigma_1(2)}(p_2) + \cdots + s_{\sigma_1(l-1)}(p_{l-1})$. Then the following equality holds;

$$\begin{aligned} V_D(p_1; p_1, \dots, p_{l-1}) &= s_1(p_1) + \max_{\sigma \in \mathfrak{S}_l, \sigma(1)=1} \sum_{j=2}^l s_{\sigma(j)}(p_{j-1}) \\ &= s_{\sigma_1(2)}(p_1) + \max_{\sigma \in \mathfrak{S}_l, \sigma(1)=\sigma_1(2)} \sum_{j=2}^l s_{\sigma(j)}(p_{j-1}). \end{aligned}$$

That is, the maximum in $\text{div } V_D(p_1; p_1, \dots, p_{l-1})$ is realized by two elements $i = 1$ and $i = \sigma_1(2)$. This is contradiction.

Thus the maximum in $\text{div } V_D(p_1; p_1, \dots, p_{l-1})$ is realized by two elements i_1, i_2 . Then $\text{div } V_D(x; p_1, \dots, p_{l-1})$ passes p_1 since the slopes of s_{i_1} and s_{i_2} are distinct by definition of s_i 's. \square

By the above lemma, we get $h^0(X, D) \geq l$.

Next, we show that $h^0(X, D) \leq h_b^0(X, D)$. Let $L := h_b^0(X, D)$. By definition of $h_b^0(X, D)$ and $l_{\Gamma(X, \mathcal{O}(D))}(x)$, there is a point $x' \in U(\Gamma(X, \mathcal{O}(D)))$ satisfying that $l_{\Gamma(X, \mathcal{O}(D))}(x') = L$.

Suppose $h^0(X, D) \geq L + 1$ and take general L points p_1, \dots, p_L near x' . That is, if $x' \in U(\Gamma(X, \mathcal{O}(D)))_\alpha$, we take L points in the same component $U(\Gamma(X, \mathcal{O}(D)))_\alpha$. Then we can take L sections s_1, \dots, s_L and a global section $s := \max_{1 \leq i \leq L} \{s_i(x) + t_i\}$ passing through all p_i .

Now we define a multigraph Γ whose vertex set is $\{1, \dots, L\}$. For each $k = 1, \dots, L$, we can choose two distinct elements $i_k \neq j_k \in \{1, \dots, L\}$ such

that $s_{i_k}(p_k) + t_{i_k} = s_{j_k}(p_k) + t_{j_k}$. We connect i_k and j_k by an edge for each $k = 1, \dots, L$. Then this multigraph Γ has L vertices and L edges, thus it has a cycle.

Assume $1, \dots, k$ make a cycle. Then we get

$$\begin{aligned} s_1(p_{i_1}) + t_1 &= s_2(p_{i_1}) + t_2 \\ s_2(p_{i_2}) + t_2 &= s_3(p_{i_2}) + t_3 \\ &\vdots \\ s_{k-1}(p_{i_{k-1}}) + t_{k-1} &= s_k(p_{i_{k-1}}) + t_k \\ s_k(p_{i_k}) + t_k &= s_1(p_{i_k}) + t_1 \end{aligned}$$

for some p_{i_1}, \dots, p_{i_k} . By combining these equations, we get

$$(s_1(p_{i_1}) - s_2(p_{i_1})) + \dots + (s_k(p_{i_k}) - s_1(p_{i_k})) = 0.$$

More generally, we define the functions l_{i_1, \dots, i_k} for $k > 1$ and a subset $\{i_1, \dots, i_k\} \subset \{1, \dots, L\}$ as follows:

$$l_{i_1, \dots, i_k}(x_1, \dots, x_k) := \sum_{j=1}^k (s_{i_j}(x_j) - s_{i_{j+1}}(x_j)),$$

where $i_{k+1} := i_1$. These functions are rational functions on X^k . To summarize the above argument, we obtain the following lemma.

Lemma 24. Suppose $h^0(X, D) \geq L + 1$.

Then for general L points p_1, \dots, p_L near x' , there exist ordered sequences $\{i_1, \dots, i_k\}, \{j_1, \dots, j_k\} \subset \{1, \dots, L\}$ such that $l_{i_1, \dots, i_k}(p_{j_1}, \dots, p_{j_k}) = 0$.

The equations $l_{i_1, \dots, i_k}(x_{j_1}, \dots, x_{j_k}) = 0$ define a codimension 1 topological subspace of X^L . Thus when we take sufficiently general L points p_1, \dots, p_L near x' , for any ordered sequences $\{i_1, \dots, i_k\}$ and $\{j_1, \dots, j_k\}$ of $\{1, \dots, L\}$, $l_{i_1, \dots, i_k}(p_{j_1}, \dots, p_{j_k}) \neq 0$.

This contradicts to the above Lemma. Thus we get $h^0(X, D) < L + 1$.

We define subsets $H_1, H_k \subset (X)^L$, $k = 2, \dots, L$ as below.

$$H_1 := \{(x_1, \dots, x_L) \in (X)^L \mid x_i \notin U(\Gamma(X, \mathcal{O}(D))) \text{ for some } i\}.$$

$$H_k := \left\{ (x_1, \dots, x_L) \in (X)^L \mid \begin{array}{l} \exists \text{an ordered sequence } \{i_1, \dots, i_k\} \text{ of elements in } \{1, \dots, L\} \\ \text{and a subset } \{x_{j_1}, \dots, x_{j_k}\} \subset \{x_1, \dots, x_L\} \\ \text{satisfying } l_{i_1, \dots, i_k}(x_{j_1}, \dots, x_{j_k}) = 0 \end{array} \right\}.$$

It is clear that the dimension of H_1 is $Ln - 1$. The dimension of H_k , for $k = 2, \dots, l$ is also $Ln - 1$ since functions l_{i_1, \dots, i_k} are rational functions on X^L . Thus $X^L \setminus \cup_{k=1}^L H_k$ is nonempty. That is, for general L points p_1, \dots, p_L in $X^L \setminus \cup_{k=1}^L H_k$, $l_{i_1, \dots, i_k}(p_{j_1}, \dots, p_{j_k}) \neq 0$ for any subset $\{p_{j_1}, \dots, p_{j_k}\}$ of these L points and any ordered sequence $\{i_1, \dots, i_k\}$ of $\{1, \dots, L\}$. This contradicts Lemma 24. Thus we get $h^0(X, D) < L + 1$. \square

Remark 25. In tropical geometry, the Identity theorem does not hold. this facts seems to be related to whether or not $h^0(X, D)$ is equal to $\dim \Gamma(X, \mathcal{O}(D))$.

4 Tropical toric varieties

Let $N \cong \mathbb{Z}^n$ be a lattice, $M := N^\vee = \text{hom}(N, \mathbb{Z})$ be the dual lattice of N , and $N_{\mathbb{R}} := N \otimes \mathbb{R}$, $M_{\mathbb{R}} := M \otimes \mathbb{R}$ be the correspondence real vector spaces.

Definition 26. Let $V = \mathbb{Z}^n \otimes \mathbb{R}$ be a real vector space with a lattice \mathbb{Z}^n . A *rational polyhedral cone* in V is a cone in V defined by an intersection of finite half planes $\{x \in V \mid a_i(x) \geq 0\}$ for some $a_i \in (\mathbb{Z}^n)^\vee$.

A *face* of σ is a subset $\tau \subset \sigma$ such that $\psi_i(\tau \cap U_i)$ is an intersection of $\text{Im } \psi_i$ and a face of ρ_i in \mathbb{R}^n for each i . It is denoted by $\tau \prec \sigma$.

A *fan* Σ in V is a finite collection of rational polyhedral cones in V satisfying the following:

- Any face of $\sigma \in \Sigma$ belongs to Σ ;
- For $\sigma, \tau \in \Sigma$ with $\sigma \cap \tau \neq \emptyset$, the intersection is a face of both σ and τ .

We denote $\Sigma^{[k]}$ the set of cones in Σ of dimension k .

A cone σ is said to be *smooth* if it is generated by a subset of a basis of \mathbb{Z}^n .

A fan Σ is said to be *smooth* if every cone of Σ is smooth.

A *dual cone* σ^\vee of a cone σ is a cone defined by $\{a \in V^\vee \mid a(x) \geq 0 \forall x \in \sigma\}$.

If σ is a rational polyhedral cone, then σ^\vee is the same. We define S_σ to be $\sigma^\vee \cap M$.

Definition 27 ([13], Definition 1.12). Let $S \subset \mathbb{Z}^n$ be a semigroup generating by finite sets $\{g_1, \dots, g_k\}$ and $R \subset \mathbb{Z}^k$ be a generating set of $\{r \in \mathbb{Z}^k \mid \sum r_i g_i = 0\}$. We define

$$K(S) := \{x \in \mathbb{T}^k \mid x \cdot r^+ = x \cdot r^- \forall r \in R\}$$

where $r^+ := (\max(r_1, 0), \dots, \max(r_k, 0))$ and $r^- := (\max(-r_1, 0), \dots, \max(-r_k, 0))$ for $r = (r_1, \dots, r_k)$.

By Lemma 1.13 of [13], a set $U_\sigma := \text{Hom}(S_\sigma, \mathbb{T})$ is identified with $K(S_\sigma) \subset \mathbb{T}^k$ naturally and this identification gives $\text{Hom}(S_\sigma, \mathbb{T})$ a structure of tropical variety. We call this tropical variety U_σ a *tropical affine toric variety*.

Let τ be a face of a cone σ . Then the inclusion $\tau \hookrightarrow \sigma$ induces an inclusion $\iota_{\tau, \sigma}: U_\tau \hookrightarrow U_\sigma$.

Definition 28. Let Σ be a fan on $N_{\mathbb{R}}$. A *tropical toric variety* X_Σ is a tropical variety $(\coprod U_\sigma) / \sim$, where $x \sim y$ means that there exist three cones σ_1, σ_2, τ and $z \in U_\tau$ such that $\tau \subset \sigma_1 \cap \sigma_2$, $x \in U_{\sigma_1}$, $y \in U_{\sigma_2}$, $\iota_{\tau, \sigma_1}(z) = x$ and $\iota_{\tau, \sigma_2}(z) = y$. A tropical toric variety X_Σ is *smooth* if each cone of Σ is smooth.

In the following, we consider only smooth tropical toric varieties.

Definition 29. Let σ be a smooth cone in $N_{\mathbb{R}}$ and $\{\rho_1, \dots, \rho_k\}$ be the set of rays of σ . Let $e_{\rho_i} \in N$ be a primitive generator of ρ_i . We fix a dual frame $\{m_{\rho_1}, \dots, m_{\rho_k}\} \subset M$ of $\{e_{\rho_1}, \dots, e_{\rho_k}\}$, that is, $m_{\rho_i} \in M$ be an element $m \in M$ satisfying that $m \cdot e_{\rho_i} = 1$ and $m \cdot e_{\rho_j} = 0$ for $j \neq i$. Then we define a regular function $\varphi_{\rho_i, \sigma}$ on U_{σ} by $\varphi_{\rho_i, \sigma}(x) := x(m_{\rho_i})$ for $x \in U_{\sigma} = \text{Hom}(\sigma^{\vee} \cap M, \mathbb{T})$.

Let Σ be a smooth fan, ρ be a ray of Σ , $\sigma_1, \dots, \sigma_l$ be cones of Σ with $\rho \prec \sigma_i$. Then on $U_{\sigma_i} \cap U_{\sigma_j}$ for $i \neq j$, the difference of regular functions φ_{ρ, σ_i} and φ_{ρ, σ_j} are an invertible regular function on $U_{\sigma_i} \cap U_{\sigma_j}$. Thus we define a *ray divisor* D_{ρ_i} on X_{Σ} to be a divisor represented by $(U_{\sigma}, \varphi_{\rho_i, \sigma})$ and a *toric divisor* to be a divisor of the form $\sum_{\rho \in \Sigma^1} a_{\rho} D_{\rho}$, $a_{\rho} \in \mathbb{Z}$.

Definition 30. Let Σ be a smooth fan with rays $\{\rho_1, \dots, \rho_d\}$. The *canonical divisor* $K_{X_{\Sigma}}$ on X_{Σ} is a divisor $-\sum_{i=1}^d D_{\rho_i}$.

Remark 31. Mikhalkin[15] defines the canonical class for general tropical varieties. The above definition is the restriction of Mikhalkin's definition for tropical toric varieties.

The following lemma allows us to consider only toric divisors.

Lemma 32 ([13], Lemma 2.42). Any Cartier divisor on a smooth tropical toric variety is linearly equivalent to a toric divisor.

Let $D = \sum_{\rho} a_{\rho} D_{\rho}$ be a toric divisor of a tropical toric variety X_{Σ} . Then Proposition 17 said $\Gamma(X, \mathcal{O}(D)) \cong \{f \in \mathcal{M}(X) \mid \text{div}(f) + D \geq 0\}$. Let f be a rational function on X with $\text{div}(f) + D \geq 0$. Then $f|_{U_0}$ is a regular function on $U_0 = N_{\mathbb{R}}$, thus we can consider that f is written as " $\sum_{m \in A} c_m x^m$ " for some finite subset $A \subset M$.

In order to describe $\Gamma(X, \mathcal{O}(D))$ by its generators, we show the following lemma.

Lemma 33.

$$\text{div}(x^m) = \sum_{\rho} \langle m, e_{\rho} \rangle D_{\rho}$$

Proof. It is enough to show the above equation on U_{σ} for a maximal cone σ . Let ρ_1, \dots, ρ_k be rays of σ , e_{ρ_i} be the primitive generator of ρ_i , and $\{m_{\rho_1}, \dots, m_{\rho_k}\}$ be the dual basis of $\{e_{\rho_1}, \dots, e_{\rho_k}\}$. Then $m \in M$ can be written as $\sum \langle m_{\rho_i}, e_{\rho_i} \rangle m_{\rho_i}$. Thus $\text{div}(x^m) = \sum \langle m_{\rho_i}, e_{\rho_i} \rangle D_{\rho_i}$ by the definition of $\varphi_{\rho, \sigma}$. \square

Definition 34. Let $f = \sum_{m \in A} c_m x^m$ be a rational function on X_{Σ} . Take a generic point p of $\text{supp } D_{\rho}$ and a neighborhood V of p such that $f|_{V \cap N_{\mathbb{R}}}$ is an affine linear function " $c_m \cdot x^m = c_m + \langle m, x \rangle$ ". Then the *degree of f along a ray divisor D_{ρ}* , denoted by $\text{deg}(f)_{\rho}$, is defined as $\langle m, e_{\rho} \rangle$. This is not depend on the choice of a generic point p since the degree of f along D_{ρ} coincides with $\min_{m \in A} \{\langle m, e_{\rho} \rangle \mid c_m \neq -\infty\}$.

Then we get the following lemma.

Lemma 35. $\operatorname{div}(f) = \operatorname{div}(f|_{\mathbb{R}^n}) + \sum_{\rho \in \Sigma^{[1]}} \operatorname{deg}(f)_\rho D_\rho$.

Proposition 36. Let D be a toric divisor on a smooth tropical toric variety X . Then

$$\Gamma(X, \mathcal{O}(D)) \cong \left\langle \sum_{\operatorname{div}(x^m) + D \geq 0} \mathbb{T} \cdot x^m \right\rangle.$$

Proof. It is clear that

$$\Gamma(X, \mathcal{O}(D)) \cong \{f \in \mathcal{M}(X) \mid \operatorname{div}(f) + D \geq 0\} \supseteq \left\langle \sum_{\operatorname{div}(x^m) + D \geq 0} \mathbb{T} \cdot x^m \right\rangle$$

We show the inverse inclusion. Let f be a rational function on X with $\operatorname{div}(f) + D \geq 0$. This f can be written as $\left\langle \sum_{m \in A} c_m x^m \right\rangle = \max_{m \in A} \{m \cdot x + c_m\}$, where A is a finite subset of M satisfying that for all proper subset $A' \subset A$, $\left\langle \sum_{m \in A'} c_m x^m \right\rangle \neq f$.

Suppose that there exists $m' \in A$ satisfying $\operatorname{div}(x^{m'}) + D \not\geq 0$. Then there exists a ray ρ such that $\langle m', e_\rho \rangle + a_\rho < 0$, although $\operatorname{deg}(f)_\rho + a_\rho = \min_{m \in A} \langle m, e_\rho \rangle + a_\rho \leq \langle m', e_\rho \rangle + a_\rho < 0$. This contradicts to $\operatorname{div}(f) + D \geq 0$. \square

For a toric divisor $D = \sum a_\rho D_\rho$, we define a polyhedron $P(D)$ by $P(D) := \{m \in M_{\mathbb{R}} \mid \langle m, e_\rho \rangle + a_\rho \geq 0\}$. Then by Proposition 36, we get $h_a^0(X, D) = h_b^0(X, D) = |P_D \cap M|$. Thus we get the following result by Proposition 22.

Proposition 37. For a toric divisor D on a smooth tropical toric variety X ,

$$h^0(X, D) = |P_D \cap M|.$$

This result is compatible for the classical result.

5 Intersection number of divisors on tropical toric surfaces

By the intersection theory on tropical toric surface described in [18], we get the following proposition.

Proposition 38. Let Σ be a two dimensional smooth fan in $N_{\mathbb{R}}$. Let ρ_1 and ρ_2 be two distinct rays of Σ . Then the intersection number $D_{\rho_1} D_{\rho_2}$ of these is one if there exists a cone σ such that $\rho_1 \prec \sigma$ and $\rho_2 \prec \sigma$ and zero if else.

Let ρ be a ray of Σ with the primitive generating vector $u \in N$ and ρ_1, ρ_2 be distinct two rays of Σ adjacent to ρ with the primitive generating vectors $u_1, u_2 \in N$. Then the self-intersection number of D_ρ is the number b satisfying that $u_1 + u_2 + bu = 0$.

Remark 39. The latter of the above proposition is different to Corollary 3.2.6 of Shaw [18] at a glance, but these are equal.

Theorem 40 (Main result 2). Let Σ be a smooth fan in $N_{\mathbb{R}}$, D be a divisor on X_{Σ} , $K_{X_{\Sigma}}$ be the canonical divisor of X_{Σ} and $\chi(X_{\Sigma})$ be the topological Euler characteristic of X_{Σ} . Then

$$h^0(X_{\Sigma}, D) + h^0(X_{\Sigma}, K_{X_{\Sigma}} - D) \geq \chi(X_{\Sigma}) + \frac{1}{2}D(D - K_{X_{\Sigma}})$$

Proof. By Lemma ??, we can assume that D is a toric divisor $\sum_{\rho} a_{\rho} D_{\rho}$.

Let X_{Σ}^{an} be a complex toric variety associated to the fan Σ , D_{ρ}^{an} be the classical ray divisor associated to a ray ρ , D^{an} be the classical divisor $\sum_{\rho} a_{\rho} D_{\rho}^{an}$ corresponding to D and $K_{X_{\Sigma}^{an}}$ be a divisor $-\sum_{\rho} D_{\rho}^{an}$ corresponding to the torus-invariant canonical divisor $K_{\Sigma} = -\sum_{\rho} D_{\rho}$ of X_{Σ}^{an} . Then classical Riemann-Roch inequality for compact smooth toric surfaces says that

$$h^0(X_{\Sigma}^{an}, D^{an}) + h^0(X_{\Sigma}^{an}, K_{X_{\Sigma}^{an}} - D^{an}) \geq \chi(\mathcal{O}_{X_{\Sigma}^{an}}) + \frac{1}{2}D^{an}(D^{an} - K_{X_{\Sigma}^{an}})$$

Proposition 37 tells us the fact that the value of the right-hand side of the above inequality coincides with the values of corresponding classical one. That is,

$$h^0(X_{\Sigma}, D) + h^0(X_{\Sigma}, K_{X_{\Sigma}} - D) = h^0(X_{\Sigma}^{an}, D^{an}) + h^0(X_{\Sigma}^{an}, K_{X_{\Sigma}^{an}} - D^{an}).$$

On the other hand, by Proposition 38 we get

$$\chi(X_{\Sigma}) + \frac{1}{2}D(D - K_{X_{\Sigma}}) = \chi(\mathcal{O}_{X_{\Sigma}^{an}}) + \frac{1}{2}D^{an}(D^{an} - K_{X_{\Sigma}^{an}}).$$

Here we use $\chi(X_{\Sigma}) = \chi(\mathcal{O}_{X_{\Sigma}^{an}}) = 1$. Thus we get the inequality by the classical inequality. \square

References

- [1] Lars Allermann and Johannes Rau, First steps in tropical intersection theory, *Mathematische Zeitschrift*, 264.3, 633-670 (2010)
- [2] Lars Allermann, Chern classes of tropical vector bundles, *Arkiv for Matematik*, 50.2, 237-258 (2012)
- [3] Matthew Baker and Serguei Norine, Riemann-Roch and Abel-Jacobi theory on a finite graph, *Advances in Mathematics*, 215.2, 766-788 (2007)
- [4] Christina Birkenhake and Herbert Lange, *Complex Abelian Varieties*, 2nd ed., Springer Grundlehren, 302 (2004)
- [5] Dustin Cartwright, Combinatorial tropical surfaces, arXiv preprint, arXiv1506.02023 (2015)
- [6] Dustin Cartwright, A specialization inequality for tropical complexes. arXiv preprint, arXiv 1511.00650 (2015)

- [7] Melody Chan, Combinatorics of the tropical Torelli map, *Algebra and Number Theory*, 6.6, 1133-1169 (2012)
- [8] Andreas Gathmann and Michael Kerber, A Riemann-Roch theorem in tropical geometry, *Mathematische Zeitschrift*, 259.1, 217-230 (2008)
- [9] Andreas Gathmann, Michael Kerber and Hannah Markwig, Tropical fans and the moduli spaces of tropical curves, *Compositio Mathematica*, 145.1, 173-195 (2009)
- [10] Mark Gross, Tropical geometry and mirror symmetry, *Journal of the American Mathematical Society*, 114 (2011)
- [11] Christian Haase, Gregg Musiker and Josephine Yu, Linear systems on tropical curves, *Mathematische Zeitschrift*, 270.3, 1111-1140 (2012)
- [12] Philipp Jell, Johannes Rau and Kristin Shaw, Lefschetz $(1, 1)$ -theorem in tropical geometry, arXiv preprint, arXiv:1711.07900 (2017).
- [13] Henning Meyer, *Intersection Theory on Tropical Toric Varieties and Compactifications of Tropical Parameter Spaces* (2011)
- [14] Grigory Mikhalkin, Enumerative tropical algebraic geometry in \mathbb{R}^2 , *Journal of the American Mathematical Society*, 18.2, 313-377 (2005)
- [15] Grigory Mikhalkin, Tropical geometry and its applications, *Proceedings of the International. Congress of Mathematicians*, 2, 827-852 (2006)
- [16] Grigory Mikhalkin and Ilia Zharkov, Tropical curves, their Jacobians and theta functions, *Curves and abelian varieties*, 465, 203-230 (2008)
- [17] Grigory Mikhalkin, Ilia Zharkov, Tropical eigenwave and intermediate Jacobians, *Homological mirror symmetry and tropical geometry*, Springer International Publishing, 309-349 (2014)
- [18] Kristin Shaw, *Tropical Intersection Theory and Surfaces*, Ph.D. Thesis, Université de Genève, 2011
- [19] Kristin Shaw, Tropical surfaces, arXiv preprint, arXiv:1506.07407 (2015)
- [20] Yoshitomi Shuhei, Generators of modules in tropical geometry, arXiv preprint arXiv:1001.0448 (2010).
- [21] Ken Sumi, Tropical theta functions and Riemann-Roch inequality for tropical Abelian surfaces, *Mathematische Zeitschrift* 297.3 (2021): 1329-1351.