A new Menon-type identity derived from group actions

Marius Tărnăuceanu

May 26, 2021

Abstract

In this short note, we give a new Menon-type identity involving the sum of element orders and the sum of cyclic subgroup orders of a finite group. It is based on applying the weighted form of Burnside's lemma to a natural group action.

MSC 2020: Primary 11A25; Secondary 20D60.

Key words : Menon's identity, weighted Burnside's lemma, group action, sum of element orders, sum of cyclic subgroup orders.

1 Introduction

One of the most interesting arithmetical identities is due to P.K. Menon [5].

Menon's identity. For every positive integer n we have

$$\sum_{n \in \mathbb{Z}_n^*} \gcd(a-1, n) = \varphi(n) \,\tau(n),$$

where \mathbb{Z}_n^* is the group of units of the ring $\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$, gcd(,) represents the greatest common divisor, φ is the Euler's totient function and $\tau(n)$ is the number of divisors of n.

There are several approaches to Menon's identity and many generalisations. One of the methods used to prove Menon-type identities is based on the Burnside's Lemma concerning group actions (see e.g. [5–10]). In what follows, we will use a generalization of this result, called the Weighted Form of Burnside's Lemma (see e.g. [2]). Weighted Form of Burnside's Lemma. Given a finite group G acting on a finite set X, we denote

$$Fix(g) = \{x \in X : g \circ x = x\}, \, \forall g \in G.$$

Let R be a commutative ring containing the rationals and $w : X \longrightarrow R$ be a weight function that is constant on the distinct orbits $O_{x_1}, ..., O_{x_k}$ of X. For every i = 1, ..., k, let $w(O_{x_i}) = w(x)$, where $x \in O_{x_i}$. Then

$$\sum_{i=1}^{k} w(O_{x_i}) = \frac{1}{|G|} \sum_{g \in G} \sum_{x \in Fix(g)} w(x).$$
(1)

Note that the Burnside's Lemma is obtained from (1) by taking the weight function $w(x) = 1, \forall x \in X$.

Next we will consider a finite group G of order n and the functions

$$\psi(G) = \sum_{g \in G} o(g)$$
 and $\sigma(G) = \sum_{H \in C(G)} |H|,$

where o(g) is the order of $g \in G$ and C(G) is the set of cyclic subgroups of G.¹ Also, for every divisor m of n, we will denote $G_m = \{g \in G : g^m = 1\}$.

Our main result is stated as follows.

Theorem 1. Under the above notations, we have

$$\sum_{a \in \mathbb{Z}_n^*} \psi(G_{\gcd(a-1,n)}) = \varphi(n) \,\sigma(G).$$
⁽²⁾

Clearly, (2) gives a new connection between the above functions $\psi(G)$ and $\sigma(G)$. We remark that an alternative way of writing (2) is

$$\sum_{a \in \mathbb{Z}_n^*} \sum_{d | \gcd(a-1,n)} d\varphi(d) n_d(G) = \varphi(n) \,\sigma(G),\tag{3}$$

where $n_d(G)$ denotes the number of cyclic subgroups of order d in G, for all d dividing n.

 $^{^1\,{\}rm For}$ more details concerning these functions, we refer the reader to [1,3] and [4], respectively.

For $G = \mathbb{Z}_n$, Theorem 1 leads to the following corollary.

Corollary 2. We have

$$\sum_{a \in \mathbb{Z}_n^*} \psi(\mathbb{Z}_{\gcd(a-1,n)}) = \varphi(n) \,\sigma(n), \tag{4}$$

where $\sigma(n)$ is the sum of divisors of n.

Finally, since $\psi(\mathbb{Z}_n) \geq \frac{q}{p+1}n^2$, where q and p are the smallest and the largest prime divisor of $n \geq 2$ (see the proof of Lemma 2.9(2) in [3]), from (4) we infer the following inequalities.

Corollary 3. We have

$$\frac{q}{p+1}\frac{1}{\varphi(n)}\sum_{a\in\mathbb{Z}_n^*}\gcd(a-1,n)^2\leq\sigma(n)\leq\frac{1}{\varphi(n)}\sum_{a\in\mathbb{Z}_n^*}\gcd(a-1,n)^2.$$
 (5)

2 Proof of Theorem 1

Let $\mathbb{Z}_n^* = \{a \in \mathbb{N} : 1 \leq a \leq n, \gcd(a, n) = 1\}$ be the group of units (mod n). The natural action of \mathbb{Z}_n^* on G is defined by

$$a \circ g = g^a, \,\forall \, (a,g) \in \mathbb{Z}_n^* \times G.$$

Then two elements of G belong to the same orbit if and only if they generate the same cyclic subgroup. This shows that the weight function $w: G \longrightarrow \mathbb{R}$, $w(g) = o(g), \forall g \in G$, is constant on the distinct orbits O_{g_1}, \ldots, O_{g_k} of G. Thus we can apply the Weighted Form of Burnside's Lemma.

First of all, we observe that $w(O_{g_i}) = o(g_i) = |\langle g_i \rangle|, \forall i = 1, ..., k$, and therefore the left side of (1) is $\sigma(G)$.

Next we will prove that $Fix(a) = G_{gcd(a-1,n)}$, for any $a \in \mathbb{Z}_n^*$. Indeed, if $g \in Fix(a)$ then $g^a = g$, that is $g^{a-1} = 1$. Since |G| = n, we also have $g^n = 1$. Consequently, $g^{gcd(a-1,n)} = 1$, i.e. $g \in G_{gcd(a-1,n)}$. The converse inclusion is obvious.

Now (1) becomes

$$\sigma(G) = \frac{1}{\varphi(n)} \sum_{a \in \mathbb{Z}_n^*} \sum_{g \in G_{\gcd(a-1,n)}} o(g) = \frac{1}{\varphi(n)} \sum_{a \in \mathbb{Z}_n^*} \psi(G_{\gcd(a-1,n)}),$$

as desired.

References

- H. Amiri, S.M. Jafarian Amiri and I.M. Isaacs, Sums of element orders in finite groups, Comm. Algebra 37 (2009), 2978–2980.
- [2] F. Harary and E.M. Palmer, *Graphical enumeration*, Academic Press, New York, 1973.
- [3] M. Herzog, P. Longobardi and M. Maj, An exact upper bound for sums of element orders in non-cyclic finite groups, J. Pure Appl. Algebra 222 (2018), 1628–1642.
- [4] T. De Medts and Tărnăuceanu, Finite groups determined by an inequality of the orders of their subgroups, Bull. Belg. Math. Soc. Simon Stevin 15 (2008), 699–704.
- [5] P.K. Menon, On the sum $\sum (a-1,n)[(a,n)=1]$, J. Indian Math. Soc. **29** (1965) 155-163.
- [6] I. M. Richards, A remark on the number of cyclic subgroups of a finite group, Amer. Math. Monthly 91 (1984), 571–572.
- B. Sury, Some number-theoretic identities from group actions, Rend. Circ. Mat. Palermo 58 (2009), 99-108.
- [8] M. Tărnăuceanu, A generalization of Menon's identity, J. Number Theory 132 (2012), 2568-2573.
- [9] M. Tărnăuceanu, The number of cyclic subgroups of finite abelian groups and Menon's identity, Bull. Aust. Math. Soc. 101 (2020), 201-206.
- [10] L. Tóth, Menon's identity and arithmetical sums representing functions of several variables, Rend. Semin. Mat. Univ. Politec. Torino 69 (2011), 97-110.

Marius Tărnăuceanu Faculty of Mathematics "Al.I. Cuza" University Iași, Romania e-mail: tarnauc@uaic.ro