

SELF CONSISTENT TRANSFER OPERATORS. INVARIANT MEASURES, CONVERGENCE TO EQUILIBRIUM, LINEAR RESPONSE AND CONTROL OF THE STATISTICAL PROPERTIES.

STEFANO GALATOLO¹

ABSTRACT. We describe a general approach to the theory of self consistent transfer operators. These operators have been introduced as tools for the study of the statistical properties of a large number of all to all interacting dynamical systems subjected to a mean field coupling.

We consider a large class of self consistent transfer operators and prove general statements about existence of invariant measures, speed of convergence to equilibrium, statistical stability and linear response.

While most of the results presented in the paper are valid in a weak coupling regime, the existence results for the invariant measures we show also hold outside the weak coupling regime.

We apply the general statements to examples of different nature: coupled continuous maps, coupled expanding maps, coupled systems with additive noise, systems made of *different maps* coupled by a mean field interaction and other examples of self consistent transfer operators not coming from coupled maps.

We also consider the problem of finding the optimal coupling between maps in order to change the statistical properties of the system in a prescribed way.

CONTENTS

1. Introduction	2
2. Self consistent transfer operators for coupled circle maps, heuristics and formalization	6
3. Self consistent operators, the existence of the invariant measure.	8
4. Self consistent operators, exponential convergence to equilibrium	14
5. Statistical stability and linear response for nonlinear perturbations	22
6. Coupled expanding circle maps	26
7. Mean field coupled continuous maps	33
8. Maps with additive noise on \mathbb{S}^1	34
9. Self consistent operators not coming from coupled map networks	42
10. Coupling different maps	46
11. The optimal coupling	48
11.1. Some reminders on optimization of a linear function on a convex set	49
11.2. Optimizing the response of the expectation of an observable	50
References	52

2010 *Mathematics Subject Classification.* 37A10; 37A60; 37C30; 37C60; 37H05.

Key words and phrases. Self consistent transfer operators, invariant measures, convergence to equilibrium, linear response.

1. INTRODUCTION

Suppose we have a normed real or complex vector space B_w and a collection of linear operators $L_{\delta,f} : B_w \rightarrow B_w$ depending on some parameter $\delta \in \mathbb{R}$ and $f \in B_w$. One can consider the nonlinear function $\mathcal{L}_\delta : B_w \rightarrow B_w$ defined by

$$(1) \quad \mathcal{L}_\delta(f) := L_{\delta,f}f.$$

These kind of nonlinear functions have been also called *self consistent transfer operators* (as the operator itself depends on the point at which it is calculated). In many examples where these operators are used the family $L_{\delta,f}$ depends in a Lipschitz way on f and the Lipschitz constant is proportional to δ . In this context the parameter δ represents in some sense a measure of the strength of the nonlinearity of the function \mathcal{L} . These concepts have been introduced and studied as models to describe the collective behavior of a network of interacting systems coupled by a mean field interaction. In the case of operators modeling coupled extended systems the parameter δ often represents the strength of the coupling or of the interaction between the systems.

These operators, their invariant measures and their properties have been studied by different techniques in classes of examples. In this paper we attempt a general approach to the study of this kind of operators, and the statistical properties of their dynamics. The main goal is to investigate under which assumptions we can establish some basic important properties of the self consistent system as the existence of the invariant measure in a certain regularity class, the convergence to equilibrium, the statistical stability and response to perturbation of the system.

We show the flexibility and the effectiveness of the approach applying it to several kinds of self consistent operators coming from coupled expanding maps, coupled random systems and other examples.

We study the behavior of the invariant measures of these operators, their convergence to equilibrium and their statistical stability mostly in the "weak coupling" regime, in the sense that most results will hold for intervals of values of the type $\delta \in [0, \bar{\delta}]$ for a relatively small $\bar{\delta}$ (with an estimate for the size of $\bar{\delta}$). Some of the results presented however will hold even for large values of δ . In particular, under suitable assumptions, we prove the existence of some invariant probability measure for the self consistent operator \mathcal{L}_δ , also providing estimates on its regularity imposing no restrictions on the size of δ (Theorem 2), thus these results also hold in a *strong coupling* regime, for which very few results are known. Sufficient assumptions for the uniqueness of the invariant measure are then shown in the case of weak coupling regime (Theorem 3). Still in the weak coupling regime we study the attractiveness of the invariant measure as a fixed point of \mathcal{L}_δ , providing exponential convergence to equilibrium results (Theorem 6) and study the response of the invariant measure of the system to changes in the function defining the mean field coupling interaction in the zero coupling limit (Theorems 12 and 14). We also investigate these questions from an optimal control point of view. Suppose we have an initial uncoupled system and we want to introduce a coupling which maximizes certain aspects of the statistical properties of the coupled system, as for example the average of a given observable. What is the best coupling to be introduced in order to do so? This is a problem related to the control of the statistical properties of chaotic and complex systems.

In this paper we have to deal with several concepts: networks of coupled systems; self consistent transfer operators; linear response; random and deterministic systems; optimal response and control of the statistical properties. To help the reader, each main section dealing with these concepts will have an introductory part trying to explain the concept, the main ideas behind and giving some additional references for its deeper understanding.

Transfer operators. An efficient method for the study of transport and the statistical properties of a dynamical system is to associate to the system a certain transfer operator describing how the dynamics act on suitable spaces of measures or probability distributions on the phase space. Important properties of the original system are related to fixed points and other properties of these transfer operators. The transfer operator which is convenient to associate to a dynamical system is usually a linear operator. Self consistent transfer operators are nonlinear operators.

As mentioned before, these operators arise as natural models of extended systems in which there is a large set of interacting dynamical systems and we consider the dynamics of each element of the large set (the local dynamics) as being influenced or perturbed by the state of the other elements in a mean field coupling. This means that the perturbation we apply to the dynamics of each local system depends on the distribution of the states of all the other elements of the large system. This global state will be represented by a probability measure, representing the probability of finding a local system in a given state of the phase space. If now we consider the transfer operator associated to the dynamics of each local system we have that this linear operator depends on the current global state of the system. One can furthermore suppose all the local systems to be homogeneous and consider again the measure representing the global state of the system as a representative for the probability of finding a local system in a given state¹. Applying the transfer operator to see how this probability measure evolves, we have then a transfer operator depending on a certain measure and acting on the measure itself. This naturally brings us to the formalization presented in (1). From a formal point of view this give rise to a nonlinear function to be applied to a certain functional space of measures. In the weak coupling regime however this nonlinear function is a small nonlinear perturbation of a linear one, simplifying the situation and the understanding of the properties of this function.

The use of self consistent operators for the study of networks of coupled maps was developed from a mathematical point of view in [29] and [5]. In Section 2 we explain some of the heuristics behind the use of these operators for the study of coupled maps. We refer to [7] and [35] for a further discussion on the scientific context in which these concepts appear and for an accurate bibliography on the subject (see also [12] and [27] for other approaches to maps in a global coupling). For introductory material we also recommend the reading of the paper [34].

Overview of the main results. In Section 3 we show a set of general assumptions on the family of operators $L_{\delta,f}$, ensuring that the nonlinear operator \mathcal{L}_δ has a fixed point of a certain type and hence the associated system has some invariant probability measure (see Theorem 2). This result is obtained by topological

¹We could also consider in a similar way interacting systems of different types, where instead of a single measure representing the distribution of the states of the systems in a certain phase space we will have a vector of measures representing the states of systems of different type (see Section 10 for more details).

methods, applying the Brouwer fixed point theorem to a suitable sequence of finite dimensional nonlinear operators approximating \mathcal{L}_δ . The assumptions required for this result are related to the regularity of the family of linear operators $L_{\delta,f}$ when f varies in a strong-weak topology, the regularity of its invariant measures (see the assumptions *(Exi1)*, *(Exi1.b)*, *(Exi2)* in Theorem 2) and the existence of a suitable finite dimensional projection, allowing to apply a kind of finite element reduction of the problem. The result also holds outside the weak coupling regime. We also discuss the uniqueness of the invariant probability measure. This will be proved in the weak coupling regime (see Theorem 3). The set of assumptions for the uniqueness, essentially require that the operators $L_{\delta,f}$ and their fixed points depend on f in a Lipschitz way (see assumptions *(Exi3)* in Theorem 3). The assumptions required to apply these results are not difficult to be verified, and in the following sections we show how to apply this general framework to interacting random and deterministic systems, together with examples of different kind.

In section 4 we take the same point of view with the goal of investigating the convergence to equilibrium: the attractiveness of the invariant measure as a fixed point of \mathcal{L}_δ and in the weak coupling regime we show assumptions under which we can prove exponential speed of convergence to equilibrium for a general class of self consistent transfer operators (see Theorem 6). The assumptions we require are related to convergence to equilibrium and a common "one step" Lasota Yorke inequality satisfied by each transfer operator in the family $L_{\delta,f}$ (see assumptions *(Con1)*, ..., *(Con3)*). The assumptions made are in a certain sense natural when considering suitable coupled dynamical systems like expanding maps or random systems with additive noise, and in the next sections we apply these general results to several classes of examples.

In Section 5, after an introduction to the concept of Linear Response and some related bibliography, we prove a general statistical stability result (see Theorem 12) and a linear response result for nonlinear perturbations of linear transfer operators (see Theorem 14), describing the first order change in the invariant measure of the system when an infinitesimal perturbation leading to a nonlinear operator is applied. We remark that this result is similar in the statement and in the proof to many other general linear response results proved for linear transfer operators (see e.g. [14]).

The methods used to establish the general statements in sections 3, 4, 5 are related to the classical transfer operator approach, letting the transfer operator associated to the system to act on stronger and weaker spaces (in a way similar to the classical reference [30]), exploiting the fact that the perturbations we are interested to apply to our systems are small when considered in a kind of mixed norm, from the stronger to the weaker space.

We then show the flexibility of this general approach applying it to systems of different kind. In particular we will consider coupled deterministic expanding maps and random maps with additive noise, coupling identical maps or different ones in a mean field regime. For these examples we will use simple spaces of functions as L^1, C^k , the Sobolev spaces $W^{k,1}, W^{k,2}$ or the space of Borel signed measures equipped with the total variation or the Wasserstein distance.

In Section 6, after recalling several useful classical results on expanding maps we show that the self consistent transfer operator associated to a network of coupled expanding maps has an invariant measure in a suitable Sobolev space $W^{k,1}$ and

we show an estimate for its Sobolev norm (see Theorem 20). In the small coupling regime we also show exponential convergence to equilibrium for this kind of systems. This will allow to apply our general linear response statement and get a linear response statement for the zero coupling limit of such systems. Similar results for this kind of systems in the weak coupling regime appear in [34], the spaces used and the methods of proof however are quite different.

In Section 7 we consider coupled continuous maps on the circle without asking for expansiveness or small coupling and we prove the existence of an invariant probability measure for the associated self consistent transfer operators, providing a sort of Krylov Bogoliubov theorem for this kinds of systems.

In Section 8 we consider coupled random maps and we apply our general framework to this case. More precisely, we consider maps with additive noise in which at every iterate of the dynamics a certain deterministic map is applied and then a random i.i.d. perturbation is added. Due to the regularizing effect of the noise at the level of the associated transfer operators we do not need to put particular restrictions on the maps considered. These examples are then particularly interesting for the applications. After recalling the basic properties of these systems and the associated transfer operators we define a self consistent transfer operator representing the global behavior of a network of coupled random maps. We prove the existence of invariant measures for this self consistent operator and show an estimate for its C^k norm which is uniform when varying the coupling strength. In the case of weak coupling, we also prove exponential speed of convergence to equilibrium for this globally coupled system. We then apply the general linear response results to this system, obtaining again a linear response result for the system in the zero coupling limit.

In Section 9 we consider a class of self consistent transfer operators where the deterministic part of the dynamics is driven by a certain map whose slope depends on the average of a given observable, in some sense similar to the examples studied in [36]. For these systems we study the existence, uniqueness of the invariant measures and linear response, similarly to what is proved for the systems coming from coupled maps.

In Section 10 we consider suitable self consistent transfer operators to model a *mean field interaction of different maps*. For simplicity we consider two types of maps. We show that the general framework we are considering also applies to this case, showing the existence and uniqueness of the invariant measure in a weak coupling regime.

In Section 11 we consider the linear response results we proved from an optimal control point of view. Suppose we want to introduce in the system a coupling which changes the statistical properties of the dynamics in some desired way. What is the optimal coupling to be considered? Given some observable whose average is meant to be optimized and a convex set P of allowed infinitesimal couplings to be applied, we show conditions under which the problem has a solution in P and this solution is unique. We remark that in [42] the research in this direction of research was motivated, with the goal of the management of the statistical properties of complex systems and in this direction several results for probabilistic cellular automata were shown.

2. SELF CONSISTENT TRANSFER OPERATORS FOR COUPLED CIRCLE MAPS, HEURISTICS AND FORMALIZATION

Since the study of self consistent transfer operators is strongly motivated by the applications to systems of globally interacting maps, in this section we briefly introduce a model representing the dynamics of a large number of coupled maps in a global mean field interaction and the associated *self consistent transfer operators*. We will see how the formalization leads to the study of a self consistent transfer operator of the kind defined at beginning of the introduction.

We remark that in this paper we only consider discrete time dynamical systems. In the continuous time case, the models one is lead to consider are related to the topic of Vlasov-type differential equations, we suggest the recent surveys [11], [22] and the references therein for an introduction to the subject.

We are now going to define more precisely the self consistent transfer operators associated to a set of dynamical systems coupled by a mean field interaction. One can think the set of interacting systems as a continuum, endowed with a measure, as for instance a swarm of interacting particles distributed by a certain density in different parts of the space. We take this point of view and we consider case in which the set of systems we consider is a measurable space M with a probability p . The set M can be finite or infinite and in each case we can define the self consistent transfer operator associated to the system.

We remark that one could see the case where M is infinite as a suitable limit of finite sets and define the self consistent transfer operator associated to the global coupling of infinitely many systems by a suitable limit of finitely many couplings (see [5], [29], [34] and Note 2 for further details on this approach).

Let us fix some notation and terminology: let us consider two metric spaces X, Y , the spaces of Borel probability measures $PM(X)$, $PM(Y)$ on X and Y , and a Borel measurable $F : X \rightarrow Y$. We denote the pushforward of F as $L_F : PM(X) \rightarrow PM(Y)$, defined by the relation

$$[L_F(\mu)](A) = \mu(F^{-1}(A))$$

for each $\mu \in PM(X)$ and measurable set $A \subseteq Y$. The pushforward can be extended as a linear function $L_F : SM(X) \rightarrow SM(Y)$ from the vector space of Borel signed measures on X to the same space on Y . In this case L_F is linear and it will be also called as the transfer operator associated to the function F .

We now define a model for the dynamics of a (possibly infinite) family of dynamical systems interacting in the mean field. For simplicity we will suppose as a phase space for each interacting system the unit circle \mathbb{S}^1 and we will equip \mathbb{S}^1 with the Borel σ -algebra. Let us consider an additional metric spaces M equipped with the Borel σ -algebra and a probability measure $p \in PM(M)$. Let us consider a collection of *identical dynamical systems* $(\mathbb{S}^1, T)_i$, with $i \in M$ and $T : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ being a Borel measurable function. The set M can be finite or infinite. In the case $M = M_n := \{1, \dots, n\}$ is finite we consider a finite set of interacting systems. In this case a natural choice is to set p as the uniform distribution p_n giving to each system the same weight $\frac{1}{n}$.

The initial state of this collection of interacting systems can be identified by a point $x(0) = (x_i(0))_{i \in M} \in (\mathbb{S}^1)^M$ (we suppose $i \rightarrow x_i(0)$ being measurable). We now define the dynamics of the interacting systems by defining a global map

$\mathcal{T} : (\mathbb{S}^1)^M \rightarrow (\mathbb{S}^1)^M$ and global trajectory of the system by

$$x(t+1) := \mathcal{T}(x(t))$$

where $x(t+1)$ is defined on every coordinate by applying at each step the local dynamics T , plus a perturbation given by the mean field interaction with the other systems, by

$$(2) \quad x_i(t+1) = \Phi_{\delta, x(t)} \circ T(x_i(t))$$

for each $i \in M$, where $\Phi_{\delta, x(t)} : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ represents the perturbation provided by the global mean field coupling with strength $\delta \geq 0$, defined in the following way: let $\pi_{\mathbb{S}^1} : \mathbb{R} \rightarrow \mathbb{S}^1$ be the universal covering projection, let us consider some continuous function $h : \mathbb{S}^1 \times \mathbb{S}^1 \rightarrow \mathbb{R}$, representing the way in which the presence of some system in the state y perturbs the systems in the state x ; we define $\Phi_{\delta, x(t)} : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ as

$$\Phi_{\delta, x(t)}(x) = x + \pi_{\mathbb{S}^1} \left(\delta \int_M h(x, x_j(t)) dp(j) \right)$$

($j \rightarrow h(x, x_j(t))$ can be viewed as a function $: M \rightarrow \mathbb{R}$). We remark that with this definition, for each $t \in \mathbb{N}$, $i \rightarrow x_i(t)$ is also measurable². We say that the global state $x(t)$ of the system is represented by a probability measure $\mu_{x(t)} \in PM(\mathbb{S}^1)$ if

$$\int_M g(x_j(t)) dp(j) = \int_{\mathbb{S}^1} g(y) d\mu_{x(t)}(y)$$

for each continuous $g : \mathbb{S}^1 \rightarrow \mathbb{R}$ (i.e. μ is the pushforward of p by the measurable function $i \rightarrow x_i(t)$). Now we see how the measures representing given initial conditions evolve with the dynamics.

Lemma 1. *Let us consider the system $((\mathbb{S}^1)^M, \mathcal{T})$ defined above. Let $\mu \in PM(\mathbb{S}^1)$, let us consider*

$$\Phi_{\delta, \mu}(x) := x + \pi_{\mathbb{S}^1} \left(\delta \int_{\mathbb{S}^1} h(x, y) d\mu(y) \right).$$

Suppose the initial condition of the system $x(0)$ is represented by a measure μ , then $x(1) = \mathcal{T}(x(0))$ is represented by $L_{\Phi_{\delta, \mu} \circ T}(\mu)$.

Proof. Indeed, considering a continuous function $g : \mathbb{S}^1 \rightarrow \mathbb{R}$, we have

$$\begin{aligned} \int_M g(x_j(1)) dp(j) &= \int_M g(\Phi_{\delta, x(0)} \circ T(x_j(0))) dp(j) \\ &= \int_{\mathbb{S}^1} g \circ \Phi_{\delta, x(0)} \circ T(y) d\mu(y) \\ &= \int_{\mathbb{S}^1} g dL_{\Phi_{\delta, x(0)} \circ T}(\mu). \end{aligned}$$

²We remark that in the case where $M = M_n$ is finite and $p = p_n$ the above formula becomes

$$(3) \quad \Phi_{\delta, x(t)}(x) = x + \pi_{\mathbb{S}^1} \left(\frac{\delta}{n} \sum_{j=1}^n h(x, x_j(t)) \right).$$

One approach to the definition of the dynamics of system made of infinitely many globally interacting maps is to start by the case of n interacting maps and then considering the limit for $n \rightarrow \infty$. The perturbation $\Phi_{\delta, x(t)}$ induced by the interaction between the systems is defined as a suitable limit of 3. This might raise some technical problem in selecting states and the assumptions for which the limit converge. With our approach we might also consider an infinite space of interacting systems as a limit of a finite interacting family. In this case it is sufficient to see (M, p) as a suitable limit of (M_n, p_n) .

But since $x(0)$ is represented by μ

$$\begin{aligned}\Phi_{\delta,x(0)}(x) &= x + \pi_{\mathbb{S}^1}(\delta \int_M h(x, x_j(0)) dp(j)) \\ &= \Phi_{\delta,\mu}(x).\end{aligned}$$

leading to the statement. \square

Hence the measure representing the current state of the system fully determines the measure which represents the next state of the system, defining a function between measures

$$\mu \rightarrow L_{\Phi_{\delta,\mu} \circ T}(\mu).$$

This function is an example of what in the following section we will consider as a self consistent transfer operator. In the case of coupled systems $((\mathbb{S}^1)^M, \mathcal{T})$ described above, to describe the evolution of a certain probability measure representing the global state of the system we hence apply at each time a transfer operator from a family of the kind

$$L_{\delta,\mu} := L_{\Phi_{\delta,\mu} \circ T} = L_{\Phi_{\delta,\mu}} L_T.$$

Each operator $L_{\delta,\mu}$ can be seen as the transfer operator associated to the dynamics of a given node of the network of coupled systems, given that the distribution of the states of the other nodes in the network is represented by the measure μ .

We remark that the extended system $((\mathbb{S}^1)^M, \mathcal{T})$ above described can be identified by the choice of the phase space \mathbb{S}^1 , the local dynamics T , the strength of coupling δ and the coupling function h . Hence it can be identified as the quadruple $(\mathbb{S}^1, T, \delta, h)$.

3. SELF CONSISTENT OPERATORS, THE EXISTENCE OF THE INVARIANT MEASURE.

Motivated by the class of examples described in the previous section, given a compact metric space X we consider a family of Markov operators $L_{\delta,\mu} : SM(X) \rightarrow SM(X)$ depending on a probability measure $\mu \in PM(X)$ and $\delta \geq 0$. In our statements, we will apply the operators $L_{\delta,\mu}$ to different strong and weak spaces of measures which are subspaces of $PM(X)$. We now introduce the notations and the basic assumptions to formalize this. Let $(B_w, \|\cdot\|_w) \subseteq SM(X)$ be a normed vector subspace of $SM(X)$. In the paper we will suppose that the weak norm $\|\cdot\|_w$ is strong enough so that the function $\mu \rightarrow \mu(X)$ is continuous as a function $: B_w \rightarrow \mathbb{R}$ and that $\|\mu_n - \mu\|_w \rightarrow 0$ for a sequence of positive measures μ_n implies that μ is positive.

A *self consistent transfer operator* in our context will be the given of a family of Markov linear operators such that $L_{\delta,\mu} : B_w \rightarrow B_w$ for each $\mu \in P_w := B_w \cap PM(X)$, some $\delta \geq 0$ and the dynamical system (A, \mathcal{L}_δ) where $A \subseteq P_w$ and $\mathcal{L}_\delta : A \rightarrow A$ is defined by

$$(4) \quad \mathcal{L}_\delta(\mu) := L_{\delta,\mu}(\mu).$$

In the notation \mathcal{L}_δ we emphasize the dependence on δ as in the following we will be interested in the behavior of these operators for certain sets of values of δ or in the limit $\delta \rightarrow 0$. We also point out that here and in the following we will use the calligraphic notation \mathcal{L} to denote some operator which is not necessarily linear and the notation L to denote linear operators.

In the case where $L_{\delta,\mu}$ is the transfer operator associated to a map $T_{\delta,\mu} : X \rightarrow X$ to this dynamical system one can also associate the skew product dynamical system $(A \times X, F)$ on $A \times X$ where $F : A \times X \rightarrow A \times X$ is defined by

$$F(\mu, x) = (\mathcal{L}_\delta(\mu), T_{\delta,\mu}(x))$$

(see also [4]). One can remark that in the case μ is a fixed point for \mathcal{L}_δ the associated dynamics will be nontrivial only on the second coordinate, where $T_{\delta,\mu}$ represents a map for which μ is an invariant measure. Hence by the classical ergodic theory results, finding the fixed points of \mathcal{L}_δ gives important information on the statistical behavior of the second coordinate of the system F .

We will hence be interested in the dynamics \mathcal{L}_δ considered on a space of measures, and on the properties of its fixed points. In particular we will be interested in the attractiveness of these fixed points (which will determine the convergence to equilibrium of the global system) and to the stability or response of these fixed points with respect to perturbations of the global system.

Notation. If A, B are two normed vector spaces and $L : A \rightarrow B$ is a linear operator we denote the mixed norm $\|L\|_{A \rightarrow B}$ as

$$\|L\|_{A \rightarrow B} := \sup_{f \in A, \|f\|_A \leq 1} \|Lf\|_B.$$

Standing assumptions. In this section we will use the following standing assumptions and notations.

Let B_s, B_w be normed vector spaces such that $(B_s, \|\cdot\|_s) \subseteq (B_w, \|\cdot\|_w) \subseteq SM(X)$. Suppose $\|\cdot\|_s \geq \|\cdot\|_w$. Let P_w the set of probability measures in B_w . We will suppose that P_w is a complete metric space. We also denote by $P_s := P_w \cap B_s$ the set of probability measures in B_s . We suppose $P_s \neq \emptyset$. We will also suppose that there is $M \geq 0$ such that as μ varies in P_w the family $L_{\delta,\mu}$ is such that $\|L_{\delta,\mu}\|_{B_w \rightarrow B_w} \leq M$ and $\|L_{\delta,\mu}\|_{B_s \rightarrow B_s} \leq M$.

We now prove general statements regarding the existence and uniqueness of regular (and then physically meaningful) invariant measures for self consistent transfer operators. We remark that since our transfer operators are not linear, the normalization of the measure to a probability one is important in this context. In the case in which we put no restrictions on the size of the parameter δ representing the nonlinearity strength, by a topological reasoning we prove a general result (see Theorem 2) on the existence of invariant probability measures. We then suppose that the parameter δ is below a certain threshold, and in this weak coupling regime we also prove some unique existence result (see Theorem 3). We remark that in the weak coupling regime similar results have been proved in several cases of extended systems (see e.g. [5], [44], [46]), also showing the uniqueness of the invariant measure in a certain class. It is known on the other hand that as the coupling strength grows, phase transitions phenomena can occur, leading to the presence of multiple invariant measures (see [45], and [36] for a case not arising from coupled maps in which the uniqueness of absolutely continuous invariant measures is lost for each $\delta > 0$).

We will hence consider a family of linear Markov operators $L_{\delta,\mu}$ and look for a solution of the nonlinear equation $L_{\delta,\mu}(\mu) = \mu$. About this we prove the following:

Theorem 2. *Suppose that there exists $\pi_n : B_w \rightarrow B_s$, a linear projection of finite rank n which is a Markov operator with the following properties: there is $M \geq 0$*

and a decreasing sequence $a(n) \rightarrow 0$ such that for each $n \geq 0$

$$(5) \quad \begin{aligned} \|\pi_n\|_{B_w \rightarrow B_w} &< M, \\ \|\pi_n\|_{B_s \rightarrow B_s} &< M \end{aligned}$$

and

$$(6) \quad \|\pi_n f - f\|_w \leq a(n) \|f\|_s.$$

Let us suppose that $\pi_n(P_w) \subseteq P_s$ and $\pi_n(P_w)$ is bounded in B_s . Let us fix $\delta \geq 0$ and suppose furthermore that:

Exi1: there is $M_1 \geq 0$ such that $\forall \mu_1 \in P_w$ and $f \in P_w$ which is a fixed point of L_{δ, μ_1} it holds

$$\|f\|_s \leq M_1;$$

Exi1.b: $\forall \mu_1 \in P_w$, $n \in \mathbb{N}$ and for each $f \in P_w$ which is a fixed point for the finite rank approximation $\pi_n L_{\delta, \pi_n \mu_1} \pi_n$ of L_{δ, μ_1} it holds

$$\|f\|_s \leq M_1;$$

Exi2: there is $K_1 \geq 0$ such that $\forall \mu_1, \mu_2 \in P_w$

$$\|L_{\delta, \mu_1} - L_{\delta, \mu_2}\|_{B_s \rightarrow B_w} \leq \delta K_1 \|\mu_1 - \mu_2\|_w.$$

Then there is $\mu \in P_s$ such that

$$\mathcal{L}_\delta \mu = \mu.$$

and

$$(7) \quad \|\mu\|_s \leq M_1.$$

To understand the assumptions made we suggest to think of B_w as a weak space, for example L^1 and of B_s as a stronger space in which regular fixed points of the linear transfer operators $L_{\delta, \mu}$ are contained, for example, in the case of transfer operators associated to expanding maps, one can think of B_s as some Sobolev space. The projection π_n allows to reduce the problem to a finite dimensional one and find fixed points of the finite dimensional reduced operators by the Brouwer fixed point theorem. In concrete examples π_n could be a finite dimensional Markov discretization, as the Ulam discretization or similar. The assumption (Exi2) in some sense says that the family of operators $L_{\delta, \mu}$ depends on μ in a Lipschitz way, considering a (weak) mixed norm topology. We remark that the assumptions (Exi1) and (Exi2) are quite natural for a family of transfer operators depending on a parameter.

Proof of Theorem 2. Without loss of generality we can suppose that each operator $L_{\delta, \mu}$ is such that $\|L_{\delta, \mu}\|_{B_s \rightarrow B_s} \leq M$, $\|L_{\delta, \mu}\|_{B_w \rightarrow B_w} \leq M$. First we prove that under the assumptions of the theorem, given a sequence μ_i of probability measures which is bounded in B_s there is a converging subsequence $\mu_{i_k} \rightarrow \mu \in P_w$, converging in the weak topology. Indeed let μ_i such a sequence, with $\|\mu_i\|_s \leq M_2$, let us consider $\nu_{n,i} := \pi_n \mu_i$. Since π_n is Markov this is a sequence of probability measures. By (5) this sequence is bounded in B_w and belongs to the finite dimensional space $\pi_n B_w$, then it has a converging subsequence $\nu_{n, i_{n,k}} \rightarrow \nu_n$ where we denote by $i_{n,k}$ a sequence of indices for which we have this convergence. We remark that this indices can depend on n . By the standing assumptions on the norm $\|\cdot\|_w$ stated at the beginning of the section, we also have that $\nu_n \in P_w$. By (6), for each n and k we have $\|\nu_{n, i_{n,k}} - \mu_{i_{n,k}}\|_w \leq a(n) M_2$. Without loss of generality we can

suppose that $i_{n,k}$ is a subsequence of $i_{n-1,k}$ (suppose we selected the sequence of indices $i_{n-1,k}$, then we can select the subsequence $\nu_{n,i_{n,k}} \rightarrow \nu_n$ only from the indices belonging to $i_{n-1,k}$ since $\nu_{n-1,i_{n-1,k}}$ is also a bounded sequence, and so on for each n by induction). In this case, for $m \geq n$ we have

$$\begin{aligned} \|\nu_{m,i_{m,k}} - \nu_{n,i_{m,k}}\|_w &\leq \|\nu_{m,i_{m,k}} - \mu_{i_{m,k}}\|_w + \|\nu_{n,i_{m,k}} - \mu_{i_{m,k}}\|_w \\ &\leq 2a(n)M_2. \end{aligned}$$

Since this is true for each k , by taking the limits it holds that $\|\nu_n - \nu_m\|_w \leq 2a(n)M_2$ and hence ν_n is a Cauchy sequence of probability measures in P_w . By the completeness of P_w , the sequence ν_n will then converge to some $\nu \in P_w$. We also have that $\nu_{k,i_{k,k}} \rightarrow \nu$ in B_w and since $\|\nu_{k,i_{k,k}} - \mu_{i_{k,k}}\|_w \leq a(k)M_2$ we also have $\mu_{i_{k,k}} \rightarrow \nu$ in B_w , finding a converging subsequence as claimed.

Let us consider a finite rank approximation of \mathcal{L}_δ , defined by

$$\mathcal{L}_{\delta,n}(\mu) := \pi_n L_{\delta,\pi_n \mu} \pi_n(\mu).$$

We now prove that $\mathcal{L}_{\delta,n}$ is a continuous function $P_w \rightarrow P_w$, indeed let $\mu \in P_w, \nu \in B_w$ such that $\mu + \nu \in P_w$, we have

$$\begin{aligned} \|\mathcal{L}_{\delta,n}(\mu + \nu) - \mathcal{L}_{\delta,n}(\mu)\|_w &\leq \|\pi_n L_{\delta,\pi_n(\mu+\nu)} \pi_n(\mu + \nu) - \pi_n L_{\delta,\pi_n \mu} \pi_n(\mu)\|_w \\ &\leq \|\pi_n\|_{B_w \rightarrow B_w} [\|L_{\delta,\pi_n(\mu+\nu)} \pi_n(\mu + \nu) - L_{\delta,\pi_n \mu} \pi_n(\mu + \nu)\|_w \\ &\quad + \|L_{\delta,\pi_n \mu} \pi_n(\mu + \nu) - L_{\delta,\pi_n \mu} \pi_n(\mu)\|_w] \\ &\leq \|\pi_n\|_{B_w \rightarrow B_w} [\|L_{\delta,\pi_n(\mu+\nu)} \pi_n(\mu) + L_{\delta,\pi_n(\mu+\nu)} \pi_n(\nu) \\ &\quad - L_{\delta,\pi_n \mu} \pi_n(\mu) - L_{\delta,\pi_n \mu} \pi_n(\nu)\|_w \\ &\quad + \|L_{\delta,\pi_n \mu} \pi_n(\mu + \nu) - L_{\delta,\pi_n \mu} \pi_n(\mu)\|_w] \end{aligned}$$

and

$$\|L_{\delta,\pi_n \mu} \pi_n(\mu + \nu) - L_{\delta,\pi_n \mu} \pi_n(\mu)\|_w \leq M^2 \|\nu\|_w$$

while using (Exi2)

$$\begin{aligned} \|L_{\delta,\pi_n(\mu+\nu)} \pi_n(\mu) - L_{\delta,\pi_n \mu} \pi_n(\mu)\|_w &\leq \delta K_1 \|\pi_n \nu\|_w \|\pi_n \mu\|_s \\ \|L_{\delta,\pi_n(\mu+\nu)} \pi_n(\nu) - L_{\delta,\pi_n \mu} \pi_n(\nu)\|_w &\leq 2M^2 \|\pi_n \nu\|_w \end{aligned}$$

hence

$$\|\mathcal{L}_{\delta,n}(\mu + \nu) - \mathcal{L}_{\delta,n}(\mu)\|_w \leq M[M + \delta K_1 \|\pi_n \mu\|_s + 2M^2] \|\nu\|_w.$$

By assumption $\pi_n P_w$ is bounded in B_s , this shows that $\|\pi_n \mu\|_s$ is uniformly bounded as μ ranges in P_w and then $\mathcal{L}_{\delta,n}(\mu)$ is Lipschitz continuous $P_w \rightarrow P_w$. Note that since P_w is a convex set, $\pi_n P_w$ is a finite dimensional convex and bounded set. Now let us see that this is also a closed set in B_w . We will deduce that it is compact. Suppose $p_i \in \pi_n P_w \subseteq P_w$ is a Cauchy sequence for the B_w norm. Since P_w is complete this will converge to a point w of P_w . But $\pi_n B_w$, being a finite dimensional vector space, is a closed space, then $w \in \pi_n B_w$. Suppose $w = \pi_n(u)$ with $u \in B_w$, since $w = \pi_n(\pi_n u) = \pi_n(w)$ and $w \in P_w$ then $w \in \pi_n P_w$. By this $\pi_n P_w$ is a closed subspace of B_w . Since $\pi_n P_w$ is a bounded, convex and closed subset of a finite dimensional space, then it is homeomorphic to a closed disc (see e.g. [13], Corollary 1.1.1). We have that $\mathcal{L}_{\delta,n}$ is continuous on $\pi_n P_w$ and $\mathcal{L}_{\delta,n}(\pi_n P_w) \subseteq \pi_n P_w$. Then by the Brouwer fixed point theorem for each n there is $\mu_n \in \pi_n P_w \subseteq P_w$ such that

$$\mathcal{L}_{\delta,n}(\mu_n) = \mu_n.$$

This means that $\pi_n L_{\delta, \pi_n \mu_n} \pi_n(\mu_n) = \mu_n$ and then by *Exi1.b* we have that for each $n \in \mathbb{N}$, $\|\mu_n\|_s \leq M_1$. As we proved above μ_n has then a converging subsequence $\mu_{n_k} \rightarrow \hat{\mu}$ in the weak norm to some element $\hat{\mu} \in P_w$.

Now let us prove that

$$\mathcal{L}_\delta(\hat{\mu}) = \hat{\mu}.$$

In fact we have for each $k \geq 0$

$$(8) \quad \mathcal{L}_\delta \hat{\mu} = L_{\delta, \hat{\mu}}(\hat{\mu} - \mu_{n_k}) + L_{\delta, \hat{\mu}}(\mu_{n_k})$$

Since $\|\hat{\mu} - \mu_{n_k}\|_w \rightarrow 0$, and the operator $L_{\delta, \hat{\mu}}$ is bounded then

$$L_{\delta, \hat{\mu}}(\mu_{n_k}) \rightarrow \mathcal{L}_\delta \hat{\mu}$$

in the weak norm. By *Exi2*

$$\|L_{\delta, \hat{\mu}}(\mu_{n_k}) - L_{\delta, \mu_{n_k}}(\mu_{n_k})\|_w \leq \delta K_1 \|\hat{\mu} - \mu_{n_k}\|_w \|\mu_{n_k}\|_s$$

which by *Exi1.b* becomes

$$\|L_{\delta, \hat{\mu}}(\mu_{n_k}) - L_{\delta, \mu_{n_k}}(\mu_{n_k})\|_w \leq \delta K_1 M_1 \|\hat{\mu} - \mu_{n_k}\|_w$$

and then

$$L_{\delta, \mu_{n_k}}(\mu_{n_k}) \rightarrow \mathcal{L}_\delta \hat{\mu}$$

in the weak norm. Since $\mu_{n_k} = \pi_{n_k} \mu_{n_k}$ we also have that

$$L_{\delta, \mu_{n_k}}(\mu_{n_k}) - \mathcal{L}_{\delta, n_k} \mu_{n_k} = L_{\delta, \mu_{n_k}}(\mu_{n_k}) - \pi_{n_k} L_{\delta, \mu_{n_k}} \mu_{n_k}$$

and then by (6)

$$\begin{aligned} \|L_{\delta, \mu_{n_k}}(\mu_{n_k}) - \mathcal{L}_{\delta, n_k} \mu_{n_k}\|_w &\leq a(n_k) \|L_{\delta, \mu_{n_k}} \mu_{n_k}\|_s \\ &\leq a(n_k) M M_1 \rightarrow 0. \end{aligned}$$

We then proved that

$$\mathcal{L}_{\delta, n_k} \mu_{n_k} \rightarrow \mathcal{L}_\delta \hat{\mu}.$$

Since

$$\mathcal{L}_{\delta, n_k} \mu_{n_k} = \mu_{n_k}$$

then we get

$$\hat{\mu} = \lim_{k \rightarrow \infty} \mu_{n_k} = \mathcal{L}_\delta(\hat{\mu})$$

proving the invariance of $\hat{\mu}$.

Now we are only left to prove that $\|\hat{\mu}\|_s \leq M_1$. Since $L_{\delta, \hat{\mu}}(\hat{\mu}) = \hat{\mu}$ this directly follows from (*Exi1*). \square

Theorem 2 give general, sufficient conditions for the existence of the invariant probability measure of a self consistent operator, but it is hard to apply it constructively to approximate the invariant measure. Furthermore it does not give information about the uniqueness. Now we prove a kind of constructive existence and uniqueness result in the case of weak coupling. We remark that the setting is slightly different from the previous result. In particular we ask a strong stability property (see (*Exi3*)) for the invariant measures of the operators $L_{\delta, \mu}$ as μ vary.

Theorem 3. *Suppose that the family $L_{\delta, \mu}$ satisfies (*Exi1*) and (*Exi2*). Suppose that P_w contains some probability measure μ with $\|\mu\|_w \leq M_1$ (where M_1 is the constant coming from (*Exi1*)) and there is $\bar{\delta} \geq 0$ such that for each $0 \leq \delta < \bar{\delta}$ and $\mu \in P_w$ with $\|\mu\|_w \leq M_1$, $L_{\delta, \mu}$ has a unique fixed probability measure in P_w which we denote by f_μ and that the family $L_{\delta, \mu}$ satisfies the following:*

Exi3: there is $K_2 \geq 1$ such that $\forall \mu_1, \mu_2 \in P_w$ with $\max(\|\mu_1\|_w, \|\mu_2\|_w) \leq M_1$

$$\|f_{\mu_1} - f_{\mu_2}\|_w \leq \delta K_2 \|\mu_1 - \mu_2\|_w.$$

Then for each $0 \leq \delta \leq \min(\bar{\delta}, \frac{1}{K_2})$, there is a unique $\mu \in P_w$ such that

$$\mathcal{L}_\delta(\mu) = \mu.$$

Furthermore $\mu = \lim_{k \rightarrow \infty} \mu_k$ where μ_k is any sequence defined inductively in the following way: let μ_0 be some probability measure in P_w with $\|\mu_0\|_w \leq M_1$, then μ_1 is the fixed probability measure of L_{δ, μ_0} , μ_i in P_w ; μ_i is the fixed probability measure of $L_{\delta, \mu_{i-1}}$ in P_w and so on.

The assumption (*Exi3*) correspond to a Lipschitz quantitative stability for the fixed points of the operators in the family $L_{\delta, \mu}$ when the operators are perturbed by changing μ . This is a strong assumption which is however satisfied for many interesting systems, as expanding and uniformly hyperbolic or many random ones, but it is not satisfied for other systems like piecewise expanding maps for perturbations changing their turning points. We remark that indeed self consistent transfer operators arising from piecewise expanding maps show a complicated behavior from the point of view of the uniqueness of the invariant measure ([36]), confirming the importance of some strong assumption as (*Exi3*).

Proof of Theorem 3. Without loss of generality we can consider $\bar{\delta} > 0$ such that

$$(9) \quad K_2 \bar{\delta} < 1.$$

Now let us consider some $0 < \delta \leq \bar{\delta}$. Let $f_0 \in P_w$ with $\|f_0\|_w \leq M_1$. Let $f_1 \in P_s$ be the fixed probability measure of L_{δ, f_0} in B_w , again $\|f_1\|_w \leq M_1$. Now, L_{δ, f_1} has a fixed probability measure which we will denote by f_2 . We also have $\|f_2\|_w \leq M_1$. By (*Exi3*)

$$\|f_1 - f_2\|_w \leq \delta K_2 \|f_0 - f_1\|_w.$$

Now let us consider the linear operator L_{δ, f_2} , this operator has a fixed probability measure $f_3 \in B_s$ with $\|f_3\|_w \leq M_1$. We get

$$\|f_3 - f_2\|_w \leq K_2 \delta \|f_2 - f_1\|_w \leq (K_2 \delta)^2 \|f_0 - f_1\|_w.$$

Continuing as before, this will lead to a new fixed probability measure f_4 with $\|f_4 - f_3\|_w \leq (K_2 \delta)^3 \|f_0 - f_1\|_w$ and so on, defining a sequence f_k with $\|f_k\|_w \leq M_1$ and $\|f_k - f_{k-1}\|_w \leq (K_2 \delta)^{k-1} \|f_0 - f_1\|_w$. Since $(K_2 \delta)^k$ is summable, f_k is a Cauchy sequence in P_w .

Since P_w is complete this sequence has a limit. Let $f := \lim_{k \rightarrow \infty} f_k \in P_w$. By (*Exi1*), f_k is also uniformly bounded in B_s . Now we can prove that $\mathcal{L}_\delta(f) = L_{\delta, f}(f) = f$. Indeed

$$\begin{aligned} L_{\delta, f}(f) &= L_{\delta, f}\left(\lim_{k \rightarrow \infty} f_k\right) \\ &= \lim_{k \rightarrow \infty} L_{\delta, f}(f_k) \end{aligned}$$

because of the continuity of $L_{\delta, f}$ in the weak norm. Furthermore

$$\begin{aligned} \lim_{k \rightarrow \infty} L_{\delta, f}(f_k) &= \lim_{k \rightarrow \infty} L_{\delta, f}(f_k) - L_{\delta, f_{k-1}}(f_k) + L_{\delta, f_{k-1}}(f_k) \\ &= \lim_{k \rightarrow \infty} L_{\delta, f}(f_k) - L_{\delta, f_{k-1}}(f_k) + f_k \end{aligned}$$

because $L_{\delta, f_{k-1}}(f_k) = f_k$. However, by (Exi2) there is $K_1 \geq 0$ such that $\|L_{\delta, \mu_1} - L_{\delta, \mu_2}\|_{B_s \rightarrow B_w} \leq K_1 \delta \|\mu_1 - \mu_2\|_w$ and using this together with (Exi1) we get

$$\begin{aligned} \|L_{\delta, f}(f_k) - L_{\delta, f_{k-1}}(f_k)\|_w &\leq K_1 \delta \|f - f_k\|_w \|f_k\|_s \\ &\leq K_1 \delta M_1 \|f - f_k\|_w \xrightarrow{k \rightarrow \infty} 0. \end{aligned}$$

Then in the B_w topology

$$L_{\delta, f}(f) = \lim_{k \rightarrow \infty} L_{\delta, f}(f_k) = \lim_{k \rightarrow \infty} f_k = f.$$

Regarding the uniqueness, suppose $\mu_1, \mu_2 \in P_w$ are invariant for \mathcal{L}_δ . Then $L_{\delta, \mu_1}(\mu_1) = \mu_1$ and $L_{\delta, \mu_2}(\mu_2) = \mu_2$. By (Exi1) we have $\max(\|\mu_1\|_w, \|\mu_2\|_w) \leq M_1$ and then by (Exi3) we have $\|\mu_1 - \mu_2\|_w \leq \delta K_2 \|\mu_1 - \mu_2\|_w$, implying $\|\mu_1 - \mu_2\|_w = 0$ because $K_2 \delta < 1$. \square

Remark 4. We remark that while the method used in the proof of Theorem 2 to find an invariant measure for the self consistent transfer operator relies on a compactness argument and the Brouwer fixed point theorem, the way the fixed point f is found in the previous proof is constructive, once we have a mean of finding the invariant measures of the various operators L_{δ, f_k} . In this case we only have to construct the sequence $f_k \rightarrow f$ and we also have an explicit way to estimate its speed of convergence to the invariant probability measure.

Remark 5. As a general remark on the uniqueness, we show that when δ is small and the operators $L_{\delta, \mu}$ are statistically stable in some sense, different invariant probability measures in B_w of \mathcal{L}_δ must be near each other. Indeed, suppose that each operator of the family $L_{\delta, \mu}$ with $\delta \geq 0$ and $\mu \in P_w$ has a unique fixed probability measure in B_w which we denote by $f_\mu \in P_w$ and there is $F : \mathbb{R} \rightarrow \mathbb{R}$ such that $\forall \mu_1, \mu_2 \in P_w$

$$\|f_{\mu_1} - f_{\mu_2}\|_w \leq F(\delta).$$

Suppose that $\mu, \nu \in P_w$ are invariant measures, as in the previous proof this implies that $\mu = f_\mu$ and $\nu = f_\nu$. Then we have

$$\|\mu - \nu\|_w = \|f_\nu - f_\mu\|_w \leq F(\delta).$$

In the case $\lim_{\delta \rightarrow 0} F(\delta) = 0$ we see that in the weak coupling case, different invariant measures must be near each other.

4. SELF CONSISTENT OPERATORS, EXPONENTIAL CONVERGENCE TO EQUILIBRIUM

Theorems 2 and 3 give information about the existence of fixed probability measures for the self consistent operators but gives no information on whether they are attractive fixed points. In this section we address this question, giving general sufficient conditions for this to hold. In the case were the invariant probability measure is attractive, we have that the associated system has convergence to equilibrium in some sense, since iterates of some initial probability measure will converge to the invariant one. It is important to estimate the speed of this convergence. In the case of weak coupling, we will show a set of general conditions implying exponential speed of convergence to equilibrium for self consistent transfer operators.

Standing assumptions. In this section we will consider a setup similar as the one in the previous section, with strong and weak spaces B_s and B_w and we will consider a stronger space $(B_{ss}, \|\cdot\|_{ss})$ with norm satisfying $\|\cdot\|_{ss} \geq \|\cdot\|_s$. We denote

by P_{ss} the set of probability measures in B_{ss} . We will suppose that P_w is bounded for the B_w norm. Again let us consider a family of Markov bounded operators $L_{\delta,\mu} : B_i \rightarrow B_i$, $i \in \{ss, s, w\}$ depending on $\mu \in P_w$ and parameter $\delta \geq 0$. We will consider furthermore the following standing assumptions:

Con1 The operators $L_{\delta,\mu}$ satisfy a common "one step" Lasota Yorke inequality. There exist constants $\hat{\delta}, B, \lambda_1 \geq 0$ with $\lambda_1 < 1$ such that $\forall f \in B_s, \mu \in P_w$, $0 \leq \delta \leq \hat{\delta}$

$$(10) \quad \|L_{\delta,\mu}f\|_w \leq \|f\|_w$$

$$(11) \quad \|L_{\delta,\mu}f\|_s \leq \lambda_1 \|f\|_s + B \|f\|_w.$$

Con2 The family of operators satisfy an extended (*Exi2*) property: there is $K \geq 1$ such that $\forall f \in B_s, \mu, \nu \in P_w, 0 \leq \delta \leq \hat{\delta}$

$$(12) \quad \|(L_{\delta,\mu} - L_{\delta,\nu})(f)\|_w \leq \delta K \|\mu - \nu\|_w \|f\|_s$$

and $\forall f \in B_{ss}, \mu, \nu \in P_w$

$$\|(L_{\delta,\mu} - L_{\delta,\nu})(f)\|_s \leq \delta K \|\mu - \nu\|_w \|f\|_{ss}.$$

We remark that by (12), when $\delta = 0$ $L_{\delta,\mu}, L_{\delta,\nu} : B_s \rightarrow B_s$ are identical operators for each $\mu, \nu \in P_w$. We hence denote this operator as L_0 . We also suppose that $\forall f \in B_s, \nu \in P_w, 0 \leq \delta \leq \hat{\delta}$

$$(13) \quad \|(L_0 - L_{\delta,\nu})(f)\|_w \leq \delta K \|\nu\|_w \|f\|_s.$$

Con3 The operator $L_0 : B_s \rightarrow B_s$ has convergence to equilibrium: there exists $a_n \geq 0$ with $a_n \rightarrow 0$ such that for each $n \in \mathbb{N}$ and $v \in V_s$

$$(14) \quad \|L_0^n(v)\|_w \leq a_n \|v\|_s$$

where

$$V_s = \{\mu \in B_s | \mu(X) = 0\}^3.$$

We remark that the assumption (*Con1*) implies that the family $L_{\delta,\mu}$ of operators is uniformly bounded as operators $L_{\delta,\mu} : B_s \rightarrow B_s$ and $L_{\delta,\mu} : B_w \rightarrow B_w$ as μ varies in P_w .

We also remark that the convergence to equilibrium assumption is sometimes not trivial to be proved in a given system, but it is somehow expected in systems having some sort of indecomposability and chaotic behavior (for instance some kind of topological mixing, expansion, hyperbolicity or presence of noise, see also Remark 38).

The following statement estimates the speed of convergence to equilibrium for self consistent transfer operators \mathcal{L}_δ when δ is small.

Theorem 6. *Let $L_{\delta,\mu}$ be a family of Markov operators $L_{\delta,\mu} : B_i \rightarrow B_i$, $i \in \{ss, s, w\}$ with $\mu \in P_w$ and Let us suppose that the operators satisfy assumptions (*Con1*), ..., (*Con3*) for some $\hat{\delta} > 0$ and that*

$$(15) \quad \sup_{\mu \in P_w, \delta \leq \hat{\delta}} \|L_{\delta,\mu}\|_{B_{ss} \rightarrow B_{ss}} < +\infty.$$

³We recall that since $\mu \rightarrow \mu(X)$ is continuous, V_s is closed and $\forall \mu \in A$ $L_{\delta,\mu}(V_s) \subseteq V_s$.

Let us consider for each $\delta \leq \hat{\delta}$ the self consistent operator $\mathcal{L}_\delta : B_s \rightarrow B_s$ defined as in (4), suppose that for each such δ there is an invariant probability measure $\mu_\delta \in P_{ss}$ for \mathcal{L}_δ and suppose that

$$(16) \quad \sup_{\delta \leq \hat{\delta}} \|\mu_\delta\|_{ss} < +\infty.$$

Then there exists $\bar{\delta}$ such that $0 < \bar{\delta} < \hat{\delta}$ and there are $C, \gamma \geq 0$ such that for each $n \in \mathbb{N}$, $0 < \delta < \bar{\delta}$, $\nu \in P_{ss}$ we have

$$(17) \quad \|\mathcal{L}_\delta^n(\nu) - \mu_\delta\|_s \leq Ce^{-\gamma n} \|\nu - \mu_\delta\|_s.$$

We remark that the convergence speed estimates provided in (17) are in the strong norm. These estimates are uniform for δ small enough and uniform in ν . We also remark that since there is the strong norm on both sides of the inequality, (17) is similar to a spectral gap estimate, rather than a convergence to equilibrium estimate (where the regularity of the measure is estimated in the strong norm and the convergence is in the weak one, resulting in a weaker estimate).

Before the proof of Theorem 6 we prove several results on the convergence to equilibrium of sequential composition of operators, in particular it will be useful to prove a Lasota Yorke inequality for such kind of composition of operators in the family $L_{\delta, \mu}$.

Lemma 7. *Let $L_{\delta, \mu}$ be a family of Markov operators satisfying (Con1). Let $\mu_1, \dots, \mu_n \in P_w$ and*

$$(18) \quad L(n) := L_{\delta, \mu_n} \circ L_{\delta, \mu_{n-1}} \circ \dots \circ L_{\delta, \mu_1}$$

be a sequential composition of operators in such family, then

$$(19) \quad \|L(n)f\|_w \leq \|f\|_w$$

and

$$(20) \quad \|L(n)f\|_s \leq \lambda_1^n \|f\|_s + \frac{B}{1 - \lambda_1} \|f\|_w.$$

Proof. The first equation is straightforward from (Con1). Let us now prove (20). We have

$$\|L_{\delta, \mu_1} f\|_s \leq \lambda_1 \|f\|_s + B \|f\|_w$$

thus

$$\begin{aligned} \|L_{\delta, \mu_2} \circ L_{\delta, \mu_1}(f)\|_s &\leq \lambda_1 \|L_{\delta, \mu_2} f\|_s + B \|L_{\delta, \mu_2} f\|_w \\ &\leq \lambda_1^2 \|f\|_s + \lambda_1 B \|f\|_w + B \|f\|_w \\ &\leq \lambda_1^2 \|f\|_s + (1 + \lambda_1) B \|f\|_w \end{aligned}$$

Continuing the composition we get (20). \square

Lemma 8. *Let $\delta \geq 0$ and let $L(n)$ be a sequential composition of operators L_{δ, μ_i} as in 18 with $i \in \{1, \dots, n\}$ and $\mu_i \in P_w$ satisfying the above assumptions (Con1) and (Con2). Let L_0 be the operator in the family for $\delta = 0$ as in (Con3). Since P_w is bounded, let us denote by $Q := \sup_{\mu \in P_w} \|\mu\|_w$. Then there is $C \geq 0$ such that $\forall g \in B_s, \forall n \geq 0$*

$$(21) \quad \|L(n)g - L_0^n g\|_w \leq \delta Q K(C \|g\|_s + n \frac{B}{1 - \lambda} \|g\|_w).$$

where B is the second coefficient of the Lasota Yorke inequality (10).

Proof. To shorten notation let us denote for $i \in \{1, \dots, n\}$, $L_i := L_{\delta, \mu_i}$. By (Con2), equation (13) we get

$$\|L_0 g - L_j g\|_w \leq \delta K \|\mu_j\|_w \|g\|_s \leq \delta Q K \|g\|_s.$$

The case $n = 1$ of (21) directly follows from (13). Let us now suppose inductively

$$\|L(n-1)g - L_0^{n-1}g\|_w \leq \delta Q K (C_{n-1} \|g\|_s + (n-1) \frac{B}{1-\lambda} \|g\|_w)$$

then

$$\begin{aligned} \|L_n L(n-1)g - L_0^n g\|_w &\leq \|L_n L(n-1)g - L_n L_0^{n-1}g + L_n L_0^{n-1}g - L_0^n g\|_w \\ &\leq \|L_n L(n-1)g - L_n L_0^{n-1}g\|_w + \|L_n L_0^{n-1}g - L_0^n g\|_w \\ &\leq \delta Q K (C_{n-1} \|g\|_s + (n-1) \frac{B}{1-\lambda} \|g\|_w) + \|[L_n - L_0](L_0^{n-1}g)\|_w \\ &\leq \delta Q K (C_{n-1} \|g\|_s + (n-1) \frac{B}{1-\lambda} \|g\|_w) + \delta Q K \|L_0^{n-1}g\|_s \\ &\leq \delta Q K (C_{n-1} \|g\|_s + (n-1) \frac{B}{1-\lambda} \|g\|_w) + \delta Q K (\lambda^{n-1} \|g\|_s + \frac{B}{1-\lambda} \|g\|_w) \\ &\leq \delta Q K [(C_{n-1} + \lambda^{n-1}) \|g\|_s] + n \frac{B}{1-\lambda} K \|g\|_w. \end{aligned}$$

The statement follows from the observation that continuing the composition, C_n remains being bounded by the sum of a geometric series. \square

Next statement is inspired to the methods developed in [17] and allows to estimate the speed of convergence to equilibrium of a sequential composition of linear operators satisfying (Con1), ..., (Con3). The statement is in some sense homologous to Proposition 2.7 in [10].

Theorem 9. *Let us consider $\delta \geq 0$, a family of operators $L_{\delta, \mu}$ with $\mu \in P_w$ satisfying (Con1), (Con2), let us consider a sequential composition $L(n)$ as above. Suppose that L_0 has convergence to equilibrium (see (Con3)). Let us fix $n_1 > 0$ and consider the 2×2 matrix M defined by*

$$M := \begin{pmatrix} \lambda_1^{n_1} & \frac{B}{1-\lambda_1} \\ \delta Q K C + a_{n_1} & \delta Q K n_1 \frac{B}{1-\lambda_1} \end{pmatrix}.$$

Under the previous assumptions for any $g \in V_s$ the following holds:

(i) for each integer $i \geq 0$ the norms of the iterates $L(in_1)g$ are bounded by

$$\begin{pmatrix} \|L(in_1)g\|_s \\ \|L(in_1)g\|_w \end{pmatrix} \leq M^i \begin{pmatrix} \|g\|_s \\ \|g\|_w \end{pmatrix}.$$

Here \leq indicates the componentwise \leq relation (both coordinates are less or equal).

(ii) If ρ is the maximum eigenvalue of M^T , with eigenvector $\begin{pmatrix} a \\ b \end{pmatrix}$ suppose $a, b \geq 0$ and $a + b = 1$, let us define the (a, b) balanced-norm as

$$\|g\|_{(a,b)} := a \|g\|_s + b \|g\|_w.$$

In this case we have

$$(22) \quad \|L(in_1)g\|_{(a,b)} \leq \rho^i \|g\|_{(a,b)}.$$

Furthermore, the situation in which $\rho < 1$, $a, b \geq 0$ can be achieved if n_1 is big enough and δ small enough. More precisely, fixing n_1 large enough we have that $\rho = \rho(\delta)$ can be seen as a function of δ . There is some $\delta_1 < 1$ such that

$$(23) \quad \rho_1 = \sup_{\delta \leq \delta_1} \rho(\delta) < 1$$

and there is a positive eigenvector of $\rho(\delta)$ for $\delta \leq \delta_1$.

As a consequence we also have

$$\|L(in_1)g\|_s \leq (1/a)\rho^i \|g\|_s,$$

and

$$\|L(in_1)g\|_w \leq (1/b)\rho^i \|g\|_s.$$

For the proof of Theorem 9 the following lemma will be useful

Lemma 10. *Let us consider real sequences a_n, b_n such that $a_n \geq 0$, $b_n \geq 0$ for each $n \in \mathbb{N}$ and $a_n, b_n \rightarrow 0$, real numbers $\delta, A, B, C \geq 0$ and a real matrix of the form*

$$\begin{pmatrix} b_n & \delta B + a_n \\ A & \delta n C \end{pmatrix}.$$

Then there is $n_1 \geq 0$, $\bar{\delta} \geq 0$ and $0 \leq \bar{\rho} < 1$ such that for each $0 \leq \delta \leq \bar{\delta}$ the matrix

$$\begin{pmatrix} b_{n_1} & \delta B + a_{n_1} \\ A & \delta n_1 C \end{pmatrix}$$

has largest eigenvalue ρ such that $0 \leq \rho \leq \bar{\rho}$ and an associated eigenvector (a, b) , such that $a, b \geq 0$.

Proof. Fixing n and letting $\delta \rightarrow 0$, the matrix $\begin{pmatrix} b_n & a_n \\ A & 0 \end{pmatrix}$, has maximum right eigenvalue $\frac{1}{2}b_n + \frac{1}{2}\sqrt{b_n^2 + 4Aa_n}$ with eigenvector $\begin{pmatrix} \frac{1}{2A}(b_n + \sqrt{b_n^2 + 4Aa_n}) \\ 1 \end{pmatrix}$.

Now if we take n_1 big enough we can let $0 \leq \frac{1}{2}b_{n_1} + \frac{1}{2}\sqrt{b_{n_1}^2 + 4Aa_{n_1}} < 1$ and then for sufficiently small δ the statement holds. \square

Now we are ready to prove Theorem 9.

Proof of Theorem 9. For the proof of (i): let us consider $n_1 \geq 0$ and $g_0 \in V_s$ and let us denote $g_i = L(in_1)g_0$. By Lemma 7 we have

$$(24) \quad \|g_{i+1}\|_s \leq \lambda_1^{n_1} \|g_i\|_s + \frac{B}{1 - \lambda_1} \|g_i\|_w.$$

By Lemma 8, assumption (Con3) and (14) we get

$$(25) \quad \begin{aligned} \|g_{i+1}\|_w &\leq \|L_0^{n_1} g_i\|_w + \delta QK(C\|g_i\|_s + n_1 \frac{B}{1 - \lambda_1} \|g_i\|_w) \\ &\leq a_{n_1} \|g_i\|_s + \delta QK(C\|g_i\|_s + n_1 \frac{B}{1 - \lambda_1} \|g_i\|_w). \end{aligned}$$

Compacting (24) and (25) into a vector notation, setting $v_i = \begin{pmatrix} \|g_i\|_s \\ \|g_i\|_w \end{pmatrix}$ we get

$$(26) \quad v_{i+1} \preceq \begin{pmatrix} \lambda_1^{n_1} & \frac{B}{1 - \lambda_1} \\ \delta QK C + a_{n_1} & \delta QK n_1 \frac{B}{1 - \lambda_1} \end{pmatrix} v_i = M v_i.$$

We remark that the matrix M does not depend on g_0 and depend on the operators in the family $L_{\delta,\mu}$, composing the sequential composition $L(n)$ only by their common coefficients λ_1, a_{n_1}, K, B coming from the assumptions (Con1), ..., (Con3). Furthermore, since M is positive, $v_1 \preceq v_2$ implies $Mv_1 \preceq Mv_2$. Hence the inequality can be iterated and we have

$$v_1 \preceq Mv_0, v_2 \preceq Mv_1 \preceq M^2v_0 \dots$$

proving (i). To prove (ii) let us consider the (a, b) balanced-norm: $\|g\|_{(a,b)} = a\|g\|_s + b\|g\|_w$. The statement (i) implies

$$\begin{aligned} \|L(in_1)g_0\|_{(a,b)} &= (a, b) \cdot \begin{pmatrix} \|g_i\|_s \\ \|g_i\|_w \end{pmatrix} \\ &\leq (a, b) \cdot M^i \cdot \begin{pmatrix} \|g_0\|_s \\ \|g_0\|_w \end{pmatrix}, \\ &\leq [((a, b) \cdot M^i)^T]^T \cdot \begin{pmatrix} \|g_0\|_s \\ \|g_0\|_w \end{pmatrix}, \\ &\leq [(M^{iT} \cdot (a, b)^T]^T \cdot \begin{pmatrix} \|g_0\|_s \\ \|g_0\|_w \end{pmatrix}, \\ &\leq [\rho^i \cdot (a, b)^T]^T \cdot \begin{pmatrix} \|g_0\|_s \\ \|g_0\|_w \end{pmatrix}, \end{aligned}$$

hence

$$\|L(in_1)g_0\|_{(a,b)} \leq \rho^i \|g_0\|_{(a,b)}$$

proving (ii). The remaining part of the statement is a direct consequence of Lemma 10. \square

We are ready to prove the main statement of this section.

Proof of Theorem 6. We need to estimate $\|\mathcal{L}_\delta^n(\nu) - \mu_\delta\|_s$. Let us denote by ν_n the sequence of probability measures where $\nu_1 = \nu$ and $\nu_n = L_{\delta,\nu_{n-1}}\nu_{n-1}$. The sequence $\mathcal{L}_\delta^n(\nu)$ can be seen as a sequential composition

$$\mathcal{L}_\delta^n(\nu) = L(n)(\nu)$$

where using the same notations as in 18

$$L(n) = L_{\delta,\nu_n} \circ L_{\delta,\nu_{n-1}} \circ \dots \circ L_{\delta,\nu_1}.$$

We remark that by the assumptions, $\|L_{\delta,\nu_i}\|_{B_s \rightarrow B_s}$ are uniformly bounded. Let us estimate this by

$$(27) \quad \|L(n)(\nu) - \mathcal{L}_\delta^n(\mu_\delta)\|_s \leq \|L(n)(\nu) - L(n)(\mu_\delta)\|_s + \|L(n)(\mu_\delta) - \mathcal{L}_\delta^n(\mu_\delta)\|_s.$$

Since our operators satisfy (Con1), ..., (Con3) and $\nu - \mu_\delta \in V$ we can estimate

$$(28) \quad \|L(n)(\nu) - L(n)(\mu_\delta)\|_s = \|L(n)(\nu - \mu_\delta)\|_s$$

using Theorem 9.⁴

⁴The proof is quite technical. We are going to explain its idea informally to help the reader to understand the motivation of various estimates: by Theorem 9 we get that $\|L(n)(\nu) - L(n)(\mu_\delta)\|_s$ decreases exponentially in n . The remaining term $\|L(n)(\mu_\delta) - \mathcal{L}_\delta^n(\mu_\delta)\|_s$ is small when δ is small and ν_1, \dots, ν_n are close to μ_δ because the operators involved in the composition $L(n)$ are all near to L_{δ,μ_δ} .

The idea is to use the balanced norm $\|\cdot\|_{(a,b)}$ to estimate $\|\cdot\|_s$, and exploit the fact that after n_1 iterates $\|L(n_1)(\nu) - L(n_1)(\mu_\delta)\|_{(a,b)}$ is contracted by a certain factor $\rho_1 < 1$.

Let n_1, δ_1, ρ_1 and $\|\cdot\|_{(a,b)}$ the parameters and the norm found applying Theorem 9 (see in particular (23)) to (28). Let us consider $\delta \leq \delta_1$. We remark that the norm $\|\cdot\|_{(a,b)}$ also depends on δ .

To simplify notations let us define a general constant that will be used in the estimates. Let

$$M_\delta := \max(1 + B, \|\mu_\delta\|_{ss}, \sup_{\mu \in P_w} \|L_{\delta,\mu}\|_{B_{ss} \rightarrow B_{ss}}, \sup_{\mu \in P_w} \|L_{\delta,\mu}\|_{B_s \rightarrow B_s})$$

and

$$M_1 = \sup_{\delta \leq \delta_1} (M_\delta).$$

By the assumptions (15), (16) we have that $M_1 < \infty$. To find $\bar{\delta} \leq \delta_1$ satisfying our statement we are going to impose a further condition to the parameter δ which is again satisfied for δ small enough. Let us state this condition: let us define for each $n \geq 0$, by induction the following sequence

$$(29) \quad C_0 = 1, \quad C_n = M_1^n C_{n-1}.$$

Let

$$(30) \quad M_2 := KC_{n_1} n_1 (KM_1 + 1)^{n_1}.$$

Now let us fix $\bar{\delta} \geq 0$ such that

$$(31) \quad \rho_2 = (\rho_1 + \bar{\delta} M_2) < 1.$$

We now see why this condition is sufficient for our statement to hold. We have indeed

$$\begin{aligned} \|L(n)(\mu_\delta) - \mathcal{L}_\delta^n(\mu_\delta)\|_{(a,b)} &= \|L_{\delta,\nu_n} \dots L_{\delta,\nu_1} \mu_\delta - L_{\delta,\mu_\delta}^n(\mu_\delta)\|_{(a,b)} \\ &\leq \|L_{\delta,\nu_n} \dots L_{\delta,\nu_1} \mu_\delta - L_{\delta,\mu_\delta} L_{\delta,\nu_{n-1}} \dots L_{\delta,\nu_1} \mu_\delta\|_{(a,b)} \\ &\quad + \|L_{\delta,\mu_\delta} L_{\delta,\nu_{n-1}} \dots L_{\delta,\nu_1} \mu_\delta - L_{\delta,\mu_\delta}^n(\mu_\delta)\|_{(a,b)}. \end{aligned}$$

We recall that by (Con2)

$$\begin{aligned} \|(L_{\delta,\nu_i} - L_{\delta,\nu_j})(\omega)\|_{(a,b)} &= a\|(L_{\delta,\nu_i} - L_{\delta,\nu_j})(\omega)\|_w + b\|(L_{\delta,\nu_i} - L_{\delta,\nu_j})(\omega)\|_s \\ &\leq a\delta K \|\nu_i - \nu_j\|_w \|\omega\|_s + b\delta K \|\nu_i - \nu_j\|_w \|\omega\|_{ss} \\ &\leq \delta K \|\nu_i - \nu_j\|_w \|\omega\|_{ss}. \end{aligned}$$

Suppose inductively that

$$(32) \quad \|L_{\delta,\nu_{n-1}} \dots L_{\delta,\nu_1} \mu_\delta - L_{\delta,\mu_\delta}^{n-1}(\mu_\delta)\|_{(a,b)} \leq \delta K C_{n-1} (\|\nu_{n-1} - \mu_\delta\|_w + \dots + \|\nu_1 - \mu_\delta\|_w)$$

(where $C_n \geq 1$ as defined in (29)) then

$$\begin{aligned} \|L_{\delta,\nu_n} L_{\delta,\nu_{n-1}} \dots L_{\delta,\nu_1} \mu_\delta - L_{\delta,\mu_\delta} L_{\delta,\nu_{n-1}} \dots L_{\delta,\nu_1} \mu_\delta\|_{(a,b)} &\leq \delta K \|\nu_n - \mu_\delta\|_w \|L_{\delta,\nu_{n-1}} \dots L_{\delta,\nu_1} \mu_\delta\|_{ss} \\ &\leq \delta M_1^n K \|\nu_n - \mu_\delta\|_w \end{aligned}$$

and

$$\|L_{\delta,\mu} L_{\delta,\nu_{n-1}} \dots L_{\delta,\nu_1} \mu_\delta - L_{\delta,\mu_\delta}^n(\mu_\delta)\|_{(a,b)} \leq \|L_{\delta,\mu_\delta}\|_{(a,b)} \|L_{\delta,\nu_{n-1}} \dots L_{\delta,\nu_1} \mu_\delta - L_{\delta,\mu_\delta}^{n-1}(\mu_\delta)\|_{(a,b)}$$

and by 32

$$\|L_{\delta,\mu} \dots L_{\delta,\nu_1} \mu_\delta - L_{\delta,\mu_\delta}^n(\mu_\delta)\|_{(a,b)} \leq \|L_{\delta,\mu_\delta}\|_{(a,b)} \delta K C_{n-1} (\|\nu_{n-1} - \mu_\delta\|_w + \dots + \|\nu_1 - \mu_\delta\|_w)$$

If we prove that δ can be made small enough so that $\|L(n_1)(\mu_\delta) - L_{\delta,\mu}^{n_1}(\mu_\delta)\|_{(a,b)}$ is not relevant, then we have that also $\|L(n_1)(\nu) - \mathcal{L}_\delta^{n_1}(\mu_\delta)\|_{(a,b)}$ is contracted. Hence continuing the iteration we have an exponential decrease of this norm, which implies exponential decrease of the $\|\cdot\|_s$ norm.

putting the two estimates together

$$\begin{aligned}
\|L_{\delta,\nu_n}\dots L_{\delta,\nu_1}\mu_\delta - L_{\delta,\mu_\delta}^n(\mu_\delta)\|_{(a,b)} &\leq \delta M_1^n K \|\nu_n - \mu_\delta\|_w \\
&\quad + \delta M_1 K C_{n-1} (\|\nu_{n-1} - \mu_\delta\|_w + \dots + \|\nu_1 - \mu_\delta\|_w) \\
&\leq \delta M_1^n K C_{n-1} (\|\nu_n - \mu_\delta\|_w + \|\nu_{n-1} - \mu_\delta\|_w + \dots \\
&\quad \dots + \|\nu_1 - \mu_\delta\|_w) \\
&\leq \delta K C_n (\|\nu_n - \mu_\delta\|_w + \|\nu_{n-1} - \mu_\delta\|_w + \dots + \|\nu_1 - \mu_\delta\|_w).
\end{aligned}$$

Now we find a coarse estimate for $\|\nu_{n_1} - \mu_\delta\|_w, \|\nu_{n_1-1} - \mu_\delta\|_w, \dots, \|\nu_1 - \mu_\delta\|_w$ which will be sufficient for our purposes. Recalling that $\nu_n = L_{\delta,\nu_{n-1}}\nu_{n-1}$ we have

$$\begin{aligned}
\|\nu_n - \mu_\delta\|_w &\leq \|L_{\delta,\nu_{n-1}}\nu_{n-1} - L_{\delta,\mu_\delta}\mu_\delta\|_w \\
&\leq \|L_{\delta,\nu_{n-1}}\nu_{n-1} - L_{\delta,\nu_{n-1}}\mu_\delta\|_w + \|L_{\delta,\nu_{n-1}}\mu_\delta - L_{\delta,\mu_\delta}\mu_\delta\|_w
\end{aligned}$$

then

$$\begin{aligned}
\|L_{\delta,\nu_{n-1}}\mu_\delta - L_{\delta,\mu_\delta}\mu_\delta\|_w &\leq \delta K \|\nu_{n-1} - \mu_\delta\|_w \|\mu_\delta\|_s \\
\|L_{\delta,\nu_{n-1}}\nu_{n-1} - L_{\delta,\nu_{n-1}}\mu_\delta\|_w &\leq \|\nu_{n-1} - \mu_\delta\|_w
\end{aligned}$$

and

$$\begin{aligned}
\|\nu_n - \mu_\delta\|_w &\leq \|\nu_{n-1} - \mu_\delta\|_w (\delta K \|\mu_\delta\|_s + 1) \\
&\leq \|\nu_{n-1} - \mu_\delta\|_w (\delta K M_1 + 1)
\end{aligned}$$

and then

$$\max(\|\nu_n - \mu_\delta\|_w, \|\nu_{n-1} - \mu_\delta\|_w, \dots, \|\nu_1 - \mu_\delta\|_w) \leq \|\nu - \mu_\delta\|_w (\delta K M_1 + 1)^n.$$

Finally we have an estimate for $\|L_{\delta,\nu_n}\dots L_{\delta,\nu_1}\mu_\delta - L_{\delta,\mu}^n(\mu_\delta)\|_{(a,b)}$:

$$\|L_{\delta,\nu_n}\dots L_{\delta,\nu_1}\mu_\delta - L_{\delta,\mu}^n(\mu_\delta)\|_{(a,b)} \leq \delta K C_n n \|\nu - \mu_\delta\|_w (\delta K M_1 + 1)^n.$$

Now the main estimates are ready. Let us apply Theorem 9 to (28). We get

$$\begin{aligned}
\|L(n_1)(\nu) - L(n_1)(\mu_\delta)\|_{(a,b)} &\leq \|L(n_1)(\mu_\delta - \nu)\|_{(a,b)} \\
&\leq \rho_1 \|\mu_\delta - \nu\|_{(a,b)}
\end{aligned}$$

with $\rho_1 < 1$ and then

$$(33) \quad \|L(n_1)(\nu) - L_{\delta,\mu}^{n_1}(\mu_\delta)\|_{(a,b)} \leq \|L(n_1)(\nu) - L(n_1)(\mu_\delta)\|_{(a,b)}$$

$$(34) \quad \quad \quad + \|L(n_1)(\mu_\delta) - L_{\delta,\mu_\delta}^{n_1}(\mu_\delta)\|_{(a,b)}$$

$$(35) \quad \quad \quad \leq \rho_1 \|\mu_\delta - \nu\|_{(a,b)}$$

$$(36) \quad \quad \quad + \delta K C_{n_1} n_1 \|\nu - \mu_\delta\|_w (\delta K M_1 + 1)^{n_1}$$

$$(37) \quad \quad \quad \leq \|(\mu_\delta - \nu)\|_{(a,b)} (\rho_1 + \delta K C_{n_1} n_1 (K M_1 + 1)^{n_1})$$

$$(37) \quad \quad \quad \leq \|(\mu_\delta - \nu)\|_{(a,b)} (\rho_1 + \delta M_2)$$

where M_2 is defined as in (30). But by (31)

$$(38) \quad \quad \quad \rho_2 = (\rho_1 + \bar{\delta} M_2) < 1.$$

Taking $\delta \leq \bar{\delta}$ we hence get that for each $i \geq 1$

$$\|L(i n_1)(\nu) - L_{\delta,\mu}^{i n_1}(\mu_\delta)\|_{(a,b)} \leq \rho_2^i \|(\mu_\delta - \nu)\|_{(a,b)}$$

proving the statement. \square

Remark 11. *We remark that if in the previous proof instead of considering 27 we considered the estimate*

$$\begin{aligned} \|L(n)(\nu) - \mathcal{L}_\delta^n(\mu_\delta)\|_s &= \|L(n)(\nu) - L_{\delta, \mu_\delta}^n(\mu_\delta)\|_s \\ &\leq \|L(n)(\nu) - L_{\delta, \mu_\delta}^n(\nu)\|_s + \|L_{\delta, \mu_\delta}^n(\nu) - L_{\delta, \mu_\delta}^n(\mu_\delta)\|_s \end{aligned}$$

we would have a much easier estimate for the summand

$$\|L_{\delta, \mu_\delta}^n(\nu) - L_{\delta, \mu_\delta}^n(\mu_\delta)\|_s = \|L_{\delta, \mu_\delta}^n(\nu - \mu_\delta)\|_s,$$

but estimating $\|L(n)(\nu) - L_{\delta, \mu_\delta}^n(\nu)\|_s$ by our assumptions (Con1), ..., (Con3) would involve a term of the kind $\|\nu\|_{ss}$, which would result in a weaker final statement.

5. STATISTICAL STABILITY AND LINEAR RESPONSE FOR NONLINEAR PERTURBATIONS

The concept of *Linear Response* intends to quantify the response of the statistical properties of the system when the system is submitted to a certain infinitesimal perturbation. This will be measured in some sense by the derivative of the invariant measure of the system with respect to the perturbation. Let $(\mathcal{L}_\delta)_{\delta \geq 0}$ be a one parameter family of transfer operators associated to a family of perturbations of an initial operator \mathcal{L}_0 , with strength δ , and let us suppose that μ_δ is the unique invariant probability measure of the operator \mathcal{L}_δ in a certain space B_{ss} . The linear response of the invariant measure of \mathcal{L}_0 under the given perturbation is defined by the limit

$$(39) \quad \dot{\mu} := \lim_{\delta \rightarrow 0} \frac{\mu_\delta - \mu_0}{\delta}$$

where the topology where this convergence takes place may depend on the system and on the kind of perturbation applied. The linear response to the perturbation hence represents the first order term of the response of a system to a perturbation and when it holds, a linear response formula can be written: $\mu_\delta = \mu_0 + \dot{\mu}\delta + o(\delta)$, holding in some weaker or stronger sense.

We remark that given an observable function $c : X \rightarrow \mathbb{R}$ if the convergence in (39) is strong enough with respect to the regularity of c we get

$$(40) \quad \lim_{t \rightarrow 0} \frac{\int c d\mu_t - \int c d\mu_0}{t} = \int c d\dot{\mu}$$

showing how the linear response of the invariant measure controls the behavior of observable averages. For instance the convergence in (40) hold when $c \in L^\infty$ and the convergence of the linear response is in L^1 .

Linear response results in the context of deterministic dynamics have been obtained first in the case of uniformly hyperbolic systems in [43]. Nowadays linear response results are known for many other kinds of systems outside the uniformly hyperbolic case and also in the random case (see [3] for a survey mostly related to deterministic systems and the introduction of [14] for an overview of the mathematical results in the random case).

In the case of coupled hyperbolic map lattices, results on the smooth dependence of the SRB measure were obtained in [38], [39]. Linear response results in the case of all-to-all coupled maps with mean field interaction were shown in [34]. Still in the context of all-to-all coupled maps, the works [40] and [41] show numerical

evidence of the fact that it is possible for a network of coupled maps to exhibit linear response, even if its units do not.

The interest of the study of the self consistent transfer operators in a weak coupling regime motivates the study of the response to *nonlinear perturbations* of linear operators. In this section we prove some stability and linear response results for the invariant measures of a family \mathcal{L}_δ of such operators in the limit $\delta \rightarrow 0$ in the case where the limit operator \mathcal{L}_0 is linear. We remark that in [33] is proved an abstract result which can be also applied to the linear response of fixed points of nonlinear operators under suitable perturbations.

Standing assumptions. In this section we consider the following general setting similar to the one considered in [14] (see also [15] and [23]) for families of linear operators. Let X be a compact metric space. In the following we consider three normed vector subspaces of $SM(X)$, the spaces $(B_{ss}, \|\cdot\|_{ss}) \subseteq (B_s, \|\cdot\|_s) \subseteq (B_w, \|\cdot\|_w) \subseteq SM(X)$ with norms satisfying

$$\|\cdot\|_w \leq \|\cdot\|_s \leq \|\cdot\|_{ss}.$$

We remark that, a priori, some of these spaces can be taken equal. Their actual choice depends on the type of system and perturbation under study. Again, we will assume that the linear form $\mu \rightarrow \mu(X)$ is continuous on B_i , for $i \in \{ss, s, w\}$. Since we will mainly consider positive, integral preserving operators acting on these spaces, the following closed invariant spaces $V_{ss} \subseteq V_s \subseteq V_w$ of zero average measures defined as:

$$V_i := \{\mu \in B_i \mid \mu(X) = 0\}$$

where $i \in \{ss, s, w\}$, will play an important role (we recall that we already found V_s in (Con3)). Suppose hence we have a one parameter family of functions $\mathcal{L}_\delta : B_i \rightarrow B_i$, $\delta \in [0, \bar{\delta})$ such that:

- each \mathcal{L}_δ preserves positive measures,
- for each $\mu \in SM(X)$ it holds $[\mathcal{L}_\delta(\mu)](X) = \mu(X)$.

We call such a family, a family of "*nonlinear*" Markov operators. The following is a "statistical stability" statement for a suitable family of such operators, showing sufficient conditions under which the invariant probability measures of these operators are stable under small perturbations of the operators.

Theorem 12. *Let $\mathcal{L}_\delta : B_i \rightarrow B_i$ with $\delta \in [0, \bar{\delta})$ be a family of "nonlinear" Markov operators. Suppose that \mathcal{L}_0 is linear and $\mathcal{L}_0 : B_s \rightarrow B_s$ is bounded. Suppose that for each $\delta \in [0, \bar{\delta})$ there is a probability measure $h_\delta \in B_{ss}$ such that $\mathcal{L}_\delta h_\delta = h_\delta$. Suppose furthermore that:*

(SS1) (regularity bounds) there is $M \geq 0$ such that for each $\delta \in [0, \bar{\delta})$

$$\|h_\delta\|_{ss} \leq M.$$

(SS2) (convergence to equilibrium for the unperturbed operator) There is a sequence $a_n \geq 0$ with $a_n \rightarrow 0$ such that for each $g \in V_{ss}$

$$\|\mathcal{L}_0^n g\|_s \leq a_n \|g\|_{ss};$$

(SS3) (small perturbation) Let $B_{2M} = \{x \in B_{ss}, \|x\|_{ss} \leq 2M\}$. There is $K \geq 0$ such that and $\mathcal{L}_0 - \mathcal{L}_\delta : B_{2M} \rightarrow B_s$ is $K\delta$ -Lipschitz.

Then

$$\lim_{\delta \rightarrow 0} \|h_\delta - h_0\|_s = 0.$$

Remark 13. *The convergence to equilibrium assumption at Item (SS2) is required only for the unperturbed operator \mathcal{L}_0 . We remark that this is a linear operator. We remark that under this assumption h_0 is the unique invariant probability measure of \mathcal{L}_0 in B_{ss} .*

Proof. Let us estimate $\|h_\delta - h_0\|_s$ exploiting $\mathcal{L}_\delta h_\delta = h_\delta$ in the following way:

$$\begin{aligned} \|h_\delta - h_0\|_s &\leq \|\mathcal{L}_\delta^N h_\delta - \mathcal{L}_0^N h_0\|_s \\ &\leq \|\mathcal{L}_\delta^N h_\delta - \mathcal{L}_0^N h_\delta\|_s + \|\mathcal{L}_0^N h_\delta - \mathcal{L}_0^N h_0\|_s. \end{aligned}$$

Since h_δ, h_0 are probability measures, $h_\delta - h_0 \in V_{ss}$ and by (SS1), $\|h_\delta - h_0\|_{ss} \leq 2M$, then because of the assumption (SS2) we have

$$\|\mathcal{L}_0^N h_\delta - \mathcal{L}_0^N h_0\|_s \leq Q(N)$$

with $Q(N) = 2a_N M \rightarrow 0$ (not depending on δ). This implies

$$\|h_\delta - h_0\|_s \leq \|\mathcal{L}_\delta^N h_\delta - \mathcal{L}_0^N h_\delta\|_s + Q(N).$$

To estimate $\|\mathcal{L}_\delta^N h_\delta - \mathcal{L}_0^N h_\delta\|_s$ we rewrite the sum $\mathcal{L}_0^N - \mathcal{L}_\delta^N$ telescopically so that

$$\begin{aligned} (\mathcal{L}_\delta^N - \mathcal{L}_0^N)h_\delta &= \sum_{k=1}^N \mathcal{L}_0^{N-k} (\mathcal{L}_\delta - \mathcal{L}_0) \mathcal{L}_\delta^{k-1} h_\delta \\ &= \sum_{k=1}^N \mathcal{L}_0^{N-k} (\mathcal{L}_\delta - \mathcal{L}_0) h_\delta \end{aligned}$$

(note that only the linearity of \mathcal{L}_0 is used here). The assumption that $\|h_\delta\|_{ss} \leq M$, together with the small perturbation assumption (SS3) implies that $\|(\mathcal{L}_\delta - \mathcal{L}_0)h_\delta\|_s \leq \delta KM$ as $\delta \rightarrow 0$. Thus

$$(41) \quad \|h_\delta - h_0\|_s \leq Q(N) + NM_2(N)[\delta KM]$$

where $M_2(N) = \max_{i \leq N} (1, \|\mathcal{L}_0\|_{B_s \rightarrow B_s}^i)$. Choosing first N big enough to let $Q(N)$ be close to 0 and then δ small enough we can make $NM_2(N)[\delta KM]$ as small as wanted, proving the statistical stability in B_s as wanted. \square

We now show a general result about the linear response of fixed points of Markov operators under suitable nonlinear perturbations, the result will be applied to self consistent transfer operators in the following sections.

Theorem 14 (Linear Response). *Let $\mathcal{L}_\delta : B_s \rightarrow B_s$, $\mathcal{L}_\delta : B_{ss} \rightarrow B_{ss}$ with $\delta \in [0, \bar{\delta})$ be a family of nonlinear Markov operators. Suppose that \mathcal{L}_0 is linear and bounded : $B_i \rightarrow B_i$ for $i \in \{w, s, ss\}$. Suppose that the family satisfy (SS1), (SS2), (SS3). Suppose furthermore that the family \mathcal{L}_δ satisfy*

(LR1) *(resolvent of the unperturbed operator) $(Id - \mathcal{L}_0)^{-1} := \sum_{i=0}^{\infty} \mathcal{L}_0^i$ is a bounded operator $V_w \rightarrow V_w$.*

(LR2) *(small perturbation and derivative operator) Let $\bar{B}_{2M} = \{x \in B_s, \|x\|_s \leq 2M\}$. There is $K \geq 0$ such that $\mathcal{L}_0 - \mathcal{L}_\delta : \bar{B}_{2M} \rightarrow B_w$ is $K\delta$ -Lipschitz. Furthermore, there is $\dot{\mathcal{L}}h_0 \in V_w$ such that*

$$(42) \quad \lim_{\delta \rightarrow 0} \left\| \frac{(\mathcal{L}_\delta - \mathcal{L}_0)h_0 - \dot{\mathcal{L}}h_0}{\delta} \right\|_w = 0.$$

Then we have the following Linear Response formula

$$(43) \quad \lim_{\delta \rightarrow 0} \left\| \frac{h_\delta - h_0}{\delta} - (Id - \mathcal{L}_0)^{-1} \dot{\mathcal{L}} h_0 \right\|_w = 0.$$

Remark 15. The assumption (LR3) on the existence of the resolvent is asked only for the unperturbed transfer operator, which is linear. This allows a large class of perturbations. In many systems it will result from the presence of a spectral gap (compactness or quasi-compactness of \mathcal{L}_0 acting on B_w).

Proof of Theorem 14. By Theorem 12 the system has strong statistical stability in B_s , that is

$$(44) \quad \lim_{\delta \rightarrow 0} \|h_\delta - h_0\|_s = 0.$$

Let us now consider $(Id - \mathcal{L}_0)^{-1}$ as a continuous linear operator $V_w \rightarrow V_w$. Remark that since $\dot{\mathcal{L}} h_0 \in V_w$, the resolvent can be computed at $\dot{\mathcal{L}} h_0$. By using that h_0 and h_δ are fixed points of their respective operators we obtain that

$$(Id - \mathcal{L}_0) \frac{h_\delta - h_0}{\delta} = \frac{1}{\delta} (\mathcal{L}_\delta - \mathcal{L}_0) h_\delta.$$

Since the operators preserve probability measures, $(\mathcal{L}_\delta - \mathcal{L}_0) h_\delta \in V_w$. By applying the resolvent to both sides

$$\begin{aligned} (Id - \mathcal{L}_0)^{-1} (Id - \mathcal{L}_0) \frac{h_\delta - h_0}{\delta} &= (Id - \mathcal{L}_0)^{-1} \frac{\mathcal{L}_\delta - \mathcal{L}_0}{\delta} h_\delta \\ &= (Id - \mathcal{L}_0)^{-1} \frac{\mathcal{L}_\delta - \mathcal{L}_0}{\delta} h_0 \\ &\quad + (Id - \mathcal{L}_0)^{-1} \left[\frac{\mathcal{L}_\delta - \mathcal{L}_0}{\delta} h_\delta - \frac{\mathcal{L}_\delta - \mathcal{L}_0}{\delta} h_0 \right] \end{aligned}$$

we obtain that the left hand side is equal to $\frac{1}{\delta}(h_\delta - h_0)$. Moreover, with respect to the right hand side we observe that, applying assumption (LR2) eventually, as $\delta \rightarrow 0$

$$\left\| (Id - \mathcal{L}_0)^{-1} \left[\frac{\mathcal{L}_\delta - \mathcal{L}_0}{\delta} h_\delta - \frac{\mathcal{L}_\delta - \mathcal{L}_0}{\delta} h_0 \right] \right\|_w \leq \| (Id - \mathcal{L}_0)^{-1} \|_{V_w \rightarrow V_w} K \| h_\delta - h_0 \|_s$$

which goes to zero thanks to Theorem 12. Thus considering the limit $\delta \rightarrow 0$ we are left with

$$\lim_{\delta \rightarrow 0} \frac{h_\delta - h_0}{\delta} = (Id - \mathcal{L}_0)^{-1} \dot{\mathcal{L}} h_0.$$

converging in the $\|\cdot\|_w$ norm, which proves our claim. \square

In Sections 3 and 4 we considered nonlinear self consistent transfer operators of the type

$$\mathcal{L}_\delta(\mu) = L_{\delta, \mu}(\mu)$$

for $\mu \in A \subseteq P_w$. These functions are positive and integral preserving. In many cases these functions can be extended to nonlinear Markov operators $B_i \rightarrow B_i$ for $i \in \{w, s, ss\}$ and the above statistical stability theorems can be applied, as it will be shown in the next sections.

6. COUPLED EXPANDING CIRCLE MAPS

In this section we consider self consistent operators modeling a network of all to all coupled expanding maps, we will prove existence of the invariant measure and exponential convergence to equilibrium for this kind of systems. Similar results appeared in [29] where the rigorous study of maps coupled by mean field interaction was started and in [6], [34] in a more general setting. We will also consider the zero-coupling limit and the related response. We show that the transfer operators in this limit satisfy the assumptions of our general theorems considering as a strong and weak spaces suitable Sobolev spaces $W^{k,1}(\mathbb{S}^1)$ of measures having a density whose k -th derivative is in $L^1(\mathbb{S}^1)$.

Let $k > 1$ and $T_0 \in C^k(\mathbb{S}^1, \mathbb{S}^1)$ be a nonsingular map⁵ of the circle. Let us denote the transfer operator associated to T_0 by L_{T_0} . We recall that the transfer operator associated to a map can be defined on signed measures by the pushforward of the map, however when the map is nonsingular, this operator preserves measures having a density with respect to the Lebesgue measure, $L^1(\mathbb{S}^1, \mathbb{R})$ and then with a small abuse of notation, identifying a measure μ with its density $h_\mu = \frac{d\mu}{dm}$ with respect to the Lebesgue measure m , the same operator can be also considered as $L_{T_0} : L^1(\mathbb{S}^1, \mathbb{R}) \rightarrow L^1(\mathbb{S}^1, \mathbb{R})$. In this case, given any density $\phi \in L^1(\mathbb{S}^1, \mathbb{R})$ the action of the operator on the density can then be described by the explicit formula

$$[L_0(\phi)](x) := \sum_{y \in T_0^{-1}(x)} \frac{\phi(y)}{|T_0'(y)|}.$$

Given $h \in C^k(\mathbb{S}^1 \times \mathbb{S}^1, \mathbb{R})$, $\delta \geq 0$ and (a probability density) $\psi \in L^1(\mathbb{S}^1, \mathbb{R})$, let $\pi : \mathbb{S}^1 \rightarrow \mathbb{R}$ be the natural, universal covering projection, coherently with Section 2, we define $\Phi_{\delta, \psi} : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ as

$$\Phi_{\delta, \psi}(x) = x + \pi(\delta \int_{\mathbb{S}^1} h(x, y) \psi(y) dy).$$

We will always consider δ small enough such that $\Phi_{\delta, \psi}$ is a diffeomorphism. Denote by $Q_{\delta, \psi}$ the transfer operator associated to $\Phi_{\delta, \psi}$, defined as

$$(45) \quad [Q_{\delta, \psi}(\phi)](x) = \frac{\phi(\Phi_{\delta, \psi}^{-1}(x))}{|\Phi_{\delta, \psi}'(\Phi_{\delta, \psi}^{-1}(x))|}$$

for any $\phi \in L^1(\mathbb{S}^1, \mathbb{R})$ (we remark that $Q_{\delta, \psi}$ depends on the product $\delta\psi$).

We will consider expanding maps $T_0 : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ satisfying the following assumptions: there is $\alpha < 1$ such that

- (1) $T_0 \in C^6$,
- (2) $\forall x \in \mathbb{S}^1, |T_0'(x)| \geq \alpha^{-1} > 1$.

Definition 16. A set $A_{M,L}$ of expanding maps is called a uniform family with parameters $M \geq 0$ and $L > 1$ if it satisfies uniformly the expansiveness and regularity condition: $\forall T \in A_{M,L}$

$$\|T\|_{C^6} \leq M, \quad \inf_{x \in \mathbb{S}^1} |T'(x)| \geq L.$$

⁵A nonsingular map T is a map such that for any Lebesgue measurable set A we have $m(A) = 0 \iff m(T^{-1}(A)) = 0$, where m is the Lebesgue measure. If T is nonsingular its associated pushforward map induces a function $L^1(\mathbb{S}^1, \mathbb{R}) \rightarrow L^1(\mathbb{S}^1, \mathbb{R})$.

It is well known that the transfer operator associated to a smooth expanding map has spectral gap and it is quasicompact when acting on suitable Sobolev spaces (see e.g.[23]). In the following we recall some particularly important related estimates we will use in this paper. We start by recalling the fact that such transfer operators satisfy some one step Lasota Yorke inequalities over these Sobolev spaces (see [14], Lemma 29 and its proof). This will be useful when applying the results of Section 4.

Lemma 17. *Let $A_{M,L}$ be a uniform family of expanding maps, the transfer operators L_T associated to a $T \in A_{M,L}$ satisfy a uniform Lasota-Yorke inequality on $W^{k,1}(\mathbb{S}^1)$: for each $1 \leq k \leq 5$ there are $\alpha < 1$, $A_k, B_k \geq 0$ such that for each $n \geq 0$, $T \in A_{M,L}$*

$$(46) \quad \|L_T^n f\|_{W^{k-1,1}} \leq A_k \|f\|_{W^{k-1,1}}$$

$$(47) \quad \|L_T^n f\|_{W^{k,1}} \leq \alpha^{kn} \|f\|_{W^{k,1}} + B_k \|f\|_{W^{k-1,1}}.$$

From this result, it is classically deduced that the transfer operator L_T of a C^6 expanding map T is *quasi-compact* on each $W^{k,1}(\mathbb{S}^1)$, with $1 \leq k \leq 5$. Furthermore, by the topological transitivity of expanding maps, 1 is the only eigenvalue on the unit circle and this implies the following result. (see [14] Proposition 30).

Proposition 18. *For each $T \in A_{M,L}$, there is $C \geq 0$ such that for each*

$$g \in V_k := \{g \in W^{k,1}(\mathbb{S}^1) \text{ s.t. } \int_{\mathbb{S}^1} g \, dm = 0\}$$

with $1 \leq k \leq 5$ and $n \geq 0$ it holds

$$\|L_T^n g\|_{W^{k,1}} \leq C \rho^n \|g\|_{W^{k,1}}.$$

In particular, the resolvent $R(1, L) := (Id - L_T)^{-1} = \sum_{i=0}^{\infty} L_T^i$ is a well-defined and bounded operator on V_k .

Now we recall some estimates relative to small perturbations of expanding maps and their associated transfer operators. These will be useful to apply our general framework to self consistent transfer operators representing a family of coupled expanding maps. The estimates will be useful to check that the assumptions of our general theorems are satisfied. We will again identify absolutely continuous measures with their densities and consider the spaces $W^{3,1}(\mathbb{S}^1)$, $W^{2,1}(\mathbb{S}^1)$, $W^{1,1}(\mathbb{S}^1)$ or $W^{2,1}(\mathbb{S}^1)$, $W^{1,1}(\mathbb{S}^1)$, $L^1(\mathbb{S}^1)$ as strongest, strong and weak space.

Proposition 19. *If L_0 and L_1 are transfer operators associated to expanding maps T_0 and T_1 , then there is a $C \in \mathbb{R}$ such that $\forall k \in \{1, 2, 3\}$, $\forall f \in W^{k,1}$:*

$$(48) \quad \|(L_1 - L_0)f\|_{W^{k-1,1}} \leq C \|T_1 - T_0\|_{C^{k+2}} \|f\|_{W^{k,1}}$$

Proof. In the case $k = 1$ the proof of this statement can be found for example in [15], Proposition 26. When $k = 2$ we have to prove that

$$(49) \quad \|((L_1 - L_0)f)'\|_{L^1} \leq C \|T_1 - T_0\|_{C^4} \|f\|_{W^{2,1}}$$

we have the well known formula (see [15] Equation 3) valid for $i \in \{0, 1\}$

$$(50) \quad (L_i f)' = L_i \left(\frac{1}{T'} f' \right) - L_i \left(\frac{T''}{(T')^2} f \right).$$

By (50) we have

$$\begin{aligned} \|((L_1 - L_0)f)'\|_1 &\leq \left\| L_1 \left(\frac{1}{T_1'} f' \right) - L_0 \left(\frac{1}{T_0'} f' \right) \right\|_1 \\ &\quad + \left\| L_1 \left(\frac{T_1''}{(T_1')^2} f \right) - L_0 \left(\frac{T_0''}{(T_0')^2} f \right) \right\|_1. \end{aligned}$$

Considering each summand and applying the statement for the case $k = 1$ we get

$$\begin{aligned} \left\| L_1 \left(\frac{1}{T_1'} f' \right) - L_0 \left(\frac{1}{T_0'} f' \right) \right\|_1 &\leq \left\| L_1 \left(\frac{1}{T_1'} f' \right) - L_1 \left(\frac{1}{T_0'} f' \right) \right\|_1 \\ &\quad + \left\| L_1 \left(\frac{1}{T_0'} f' \right) - L_0 \left(\frac{1}{T_0'} f' \right) \right\|_1 \\ &\leq \|T_1 - T_0\|_{C^4} K_1 \|f'\|_1 + \|T_1 - T_0\|_{C^4} C \left\| \frac{1}{T_0'} f' \right\|_{W^{1,1}} \\ &\leq \|T_1 - T_0\|_{C^4} [K_1 \|f'\|_1 + C \left\| \frac{1}{T_0'} \right\|_{C^1} \|f'\|_{W^{1,1}}] \end{aligned}$$

for some constant $K_1 \geq 0$. Similarly

$$\begin{aligned} \left\| L_1 \left(\frac{T_1''}{(T_1')^2} f \right) - L_0 \left(\frac{T_0''}{(T_0')^2} f \right) \right\|_1 &\leq \left\| L_1 \left(\frac{T_1''}{(T_1')^2} f \right) - L_1 \left(\frac{T_0''}{(T_0')^2} f \right) \right\|_1 \\ &\quad + \left\| L_1 \left(\frac{T_0''}{(T_0')^2} f \right) - L_0 \left(\frac{T_0''}{(T_0')^2} f \right) \right\|_1 \\ &\leq \left\| \frac{T_1''}{(T_1')^2} - \frac{T_0''}{(T_0')^2} \right\|_\infty \|f\|_1 \\ &\quad + \|T_1 - T_0\|_{C^4} C \left\| \frac{T_0''}{(T_0')^2} f \right\|_{W^{1,1}} \\ &\leq \|T_1 - T_0\|_{C^4} [K_2 \|f\|_1 + \delta C \left\| \frac{T_0''}{(T_0')^2} \right\|_{C^1} \|f\|_{W^{1,1}}] \end{aligned}$$

for some constant $K_2 \geq 0$. Proving the statement. We remark that $\left\| \frac{T_0''}{(T_0')^2} \right\|_{C^1}$ involves the third derivative of T_0 .

When $k = 3$ we have to prove that

$$(51) \quad \|((L_1 - L_0)f)''\|_1 \leq C \|T_1 - T_0\|_{C^5} \|f\|_{W^{3,1}}$$

taking a further derivative in (50) for a transfer operator L_1 we get

$$((L_1 f)')' = \left(L_1 \left(\frac{1}{T_1'} f' \right) \right)' - \left(L_1 \left(\frac{T_1''}{(T_1')^2} f \right) \right)'$$

where

$$\begin{aligned} \left(L_1 \left(\frac{1}{T_1'} f' \right) \right)' &= L_1 \left(\frac{1}{T_1'} \left(\frac{1}{T_1'} f' \right)' \right) - L_1 \left(\frac{T_1''}{(T_1')^2} \left(\frac{1}{T_1'} f' \right) \right) \\ &= L_1 \left(\frac{1}{T_1'} \left(\frac{-T_1''}{(T_1')^2} f' + \frac{1}{T_1'} f'' \right) \right) - L_1 \left(\frac{T_1''}{(T_1')^2} \left(\frac{1}{T_1'} f' \right) \right) \end{aligned}$$

and

$$\begin{aligned} \left(L_1 \left(\frac{T''}{(T')^2} f \right) \right)' &= L_1 \left(\frac{1}{T'} \left(\frac{T''}{(T')^2} f \right)' \right) - L_1 \left(\frac{T''}{(T')^2} \left(\frac{T''}{(T')^2} f \right) \right) \\ &= L_1 \left(\frac{1}{T'} \left(\left(\frac{T''}{(T')^2} \right)' f + \frac{T''}{(T')^2} f' \right) \right) - L_1 \left(\frac{T''}{(T')^2} \left(\frac{T''}{(T')^2} f \right) \right) \end{aligned}$$

and the proof can be concluded as before. \square

Now we can prove that the self consistent transfer operator associated to a family of coupled expanding maps has a regular invariant measure whose regularity we can estimate.

Let us consider our expanding map T_0 and denote with $L_0 = L_{T_0}$ its transfer operator, consider $\delta \geq 0$ and a coupling function $\delta h \in C^6(\mathbb{S}^1 \times \mathbb{S}^1 \rightarrow \mathbb{R})$, consider the extended system $(\mathbb{S}^1, T_0, \delta h)$ in which these maps are coupled by δh as explained in Section 2 and the associated self consistent transfer operator $\mathcal{L}_\delta : L^1(\mathbb{S}^1, \mathbb{R}) \rightarrow L^1(\mathbb{S}^1, \mathbb{R})$

$$(52) \quad \mathcal{L}_\delta(\phi) = Q_{\delta, \phi}(L_0(\phi))$$

as defined at (4). We show that this transfer operator has under suitable assumptions a fixed probability density in $W^{3,1}$ which is unique when δ is small enough.

Proposition 20 (Existence and uniqueness of the invariant measure). *Let $T_0 \in C^6$ be an expanding circle map, let $h \in C^6, \delta \geq 0$ and the associated coupled system modeled by the transfer operator \mathcal{L}_δ as described above. There is $\bar{\delta}$ such that for each $\delta \leq \bar{\delta}$ there is a unique $f_\delta \in L^1$ such that*

$$\mathcal{L}_\delta(f_\delta) = f_\delta$$

furthermore $f_\delta \in W^{3,1}$.

Now let us consider a stronger coupling case and suppose δ is such that the set $\bigcup_{\phi \in P_w} \{\Phi_{\delta, \phi} \circ T_0\} \subseteq A_{M,L}$ is contained in a uniform family of expanding maps with parameters M, L (see Definition 16) and $L > 2$. Then there is at least one probability density $f_\delta \in W^{3,1}$ such that

$$\mathcal{L}_\delta(f_\delta) = f_\delta.$$

Furthermore, for each such invariant measure, $\|f_\delta\|_{W^{3,1}} \leq C(M, L)$, the $W^{3,1}$ norm is bounded by constant only depending on M and L .

Before the proof of this proposition we need to collect some further preliminary result. The following Lemma is about the nowadays well known statistical stability of expanding maps (see e.g. [15] Section 4 and Section 7.4. for more details).

Lemma 21. *There is $K \geq 1$ such that for each $T_1, T_2 \in A_{M,L}$ having $f_1, f_2 \in W^{3,1}$ as invariant densities it holds*

$$\|f_1 - f_2\|_{L^1} \leq K \|T_1 - T_2\|_{C^6}.$$

Now we estimate how the transfer operator $Q_{\delta, \psi}$ changes when changing ψ . This will allow to apply Lemma 21 in the proof of Proposition 3.

Lemma 22. *If $T_0, h \in C^k$ there are K_1 and $K_2 \geq 1$ such that for each $\psi, \phi \in L^1$*

$$\begin{aligned} \|\Phi_{\delta, \psi} - \Phi_{\delta, \phi}\|_{C^k} &\leq \delta K_1 \|\psi - \phi\|_{L^1} \\ \|\Phi_{\delta, \psi} \circ T_0 - \Phi_{\delta, \phi} \circ T_0\|_{C^k} &\leq \delta K_2 \|\psi - \phi\|_{L^1} \end{aligned}$$

Proof. We have that $\Phi_{\delta,\psi}(x) = x + \pi(\delta \int_{\mathbb{S}^1} h(x,y)\psi(y)dy)$ and $\Phi_{\delta,\phi}(x) = x + \pi(\delta \int_{\mathbb{S}^1} h(x,y)\phi(y)dy)$, hence

$$\begin{aligned} |\Phi_{\delta,\psi}(x) - \Phi_{\delta,\phi}(x)| &= |x + \pi(\delta \int_{\mathbb{S}^1} h(x,y)\psi(y)dy) - x + \pi(\delta \int_{\mathbb{S}^1} h(x,y)\phi(y)dy)| \\ &= |\delta \int_{\mathbb{S}^1} h(x,y)[\psi(y) - \phi(y)]dy| \leq \delta \|h\|_{L^\infty} \|\phi - \psi\|_{L^1} \end{aligned}$$

Considering the derivative with respect to x we get $\Phi'_{\delta,\psi}(x) = 1 + \pi(\delta \int_{\mathbb{S}^1} \frac{\partial h(x,y)}{\partial x} \psi(y)dy)$ and similarly for $\Phi_{\delta,\phi}(x)$. We have then

$$|\Phi'_{\delta,\psi}(x) - \Phi'_{\delta,\phi}(x)| = |\delta \int_{\mathbb{S}^1} \frac{\partial h(x,y)}{\partial x} [\psi(y) - \phi(y)]dy|$$

and

$$|\Phi'_{\delta,\psi}(x) - \Phi'_{\delta,\phi}(x)| \leq \delta \left\| \frac{\partial h}{\partial x} \right\|_{L^\infty} \|\phi - \psi\|_{L^1}.$$

similarly, we get the same estimate for the further derivatives and so on, proving the statement. \square

Now we are ready for the proof of Proposition 20.

Proof of Proposition 20. For the first part of the statement (the weak coupling case) we will apply Theorem 3 with $B_s = W^{1,1}[\mathbb{S}^1]$ and $B_w = L^1[\mathbb{S}^1]$.

By Lemma 22 Lemma 17 when δ is small enough all the operators in the family $L_{\delta,\mu}$ with $\mu \in P_w$ are the transfer operators associated to a uniform family of expanding maps and satisfy a uniform Lasota Yorke inequality on $W^{1,1}$ and L^1 , by this each one of these operators has a unique invariant probability measure in $W^{1,1}$ with uniformly bounded norm and (Exi1) is verified.

By Lemma 22 and Proposition 19 we get

$$(\text{spr}) \quad \|(L_{\delta,\mu_1} - L_{\delta,\mu_2})f\|_{L^1} \leq \text{Const} \|\mu_1 - \mu_2\|_{L^1} \|f\|_{W^{1,1}}$$

verifying (Exi2) in this case.

By Lemma 21, Lemma 22 also (Exi3) is verified. Then we can apply Theorem 3 to get the existence and uniqueness for small δ .

Now we consider the second part of the statement and the existence of an invariant measure in the stronger coupling case. The existence in L^1 of a fixed probability measure for \mathcal{L}_δ in this case follows from Theorem 2, applying it with $B_s = BV[\mathbb{S}^1]$ and $B_w = L^1[\mathbb{S}^1]$ to the family of operators $L_{\delta,\mu} = Q_{\delta,\mu} \circ L_{T_0}$. We now verify that the required assumptions hold.

Since the maps $\Phi_{\delta,\mu} \circ T_0$ involved in the system are a uniform family of expanding maps and $L > 2$, the operators $L_{\delta,\mu}$ satisfy a common Lasota Yorke inequality on $BV[\mathbb{S}^1]$ and $L^1[\mathbb{S}^1]$ (see e.g. [15], Section 9.1) and this gives a family of invariant measures for the operators $L_{\delta,\mu}$ which is uniformly bounded in $BV[\mathbb{S}^1]$ hence (Exi1) is verified also in this case.

We have to verify (Exi2) for the BV norm. Let $f \in BV$, consider $f_\epsilon \in W^{1,1}$ with $\|f_\epsilon\|_{W^{1,1}} \leq \|f\|_{BV} + \epsilon$ and $\|f_\epsilon - f\|_{L^1} \leq \epsilon$.

$$\begin{aligned} \|(L_{\delta,\mu_1} - L_{\delta,\mu_2})f\|_{L^1} &= \|(L_{\delta,\mu_1} - L_{\delta,\mu_2})f - f_\epsilon + f_\epsilon\|_{L^1} \\ &\leq \|(L_{\delta,\mu_1} - L_{\delta,\mu_2})f - f_\epsilon\|_{L^1} + \|(L_{\delta,\mu_1} - L_{\delta,\mu_2})f_\epsilon\|_{L^1} \\ &\leq 2M\epsilon + C\|\mu_1 - \mu_2\|_{L^1} \|f_\epsilon\|_{W^{1,1}} \\ &= 2M\epsilon + C\|\mu_1 - \mu_2\|_{L^1} (\|f_\epsilon\|_{BV} + \epsilon) \end{aligned}$$

and since ϵ is arbitrary, also (Exi2) is verified in this case.

Let \mathcal{P}_n be the partition subdividing the circle into n equal intervals. We can consider $\pi_n : L^1(\mathbb{S}^1) \rightarrow L^1(\mathbb{S}^1)$ to be the Ulam discretization defined as $\pi_n(f) = \mathbf{E}(f|\mathcal{P}_n)$, where the conditional expectation is made using the Lebesgue measure, projecting to piecewise constant functions.

For the Ulam projection it is well known that $\|\pi_n f - f\|_{L^1} \leq \frac{K}{n} \|f\|_{BV}$, $\|\pi_n\|_{L^1 \rightarrow L^1} \leq 1$, $\|\pi_n\|_{BV \rightarrow BV} \leq 1$ (see [18] or the proof of [24, Lemma 4.1] e.g.) and then the discretized operators $\pi_n L_{\delta, \mu} \pi_n$ satisfy a uniform Lasota Yorke inequality on BV and L^1 (see e.g. [15], Section 9.3). By this the assumption Exi1.b is satisfied and we can apply Theorem 2, and get the existence of an invariant probability density f in BV . Since $T, h \in C^6$ and $L_{\delta, f} f = f$ we get that $f \in W^{3,1}$ and its norm can be uniformly estimated by the uniform Lasota Yorke inequality on $W^{3,1}$ and $W^{2,1}$ and then on $W^{2,1}$ and $W^{1,1}$ all the transfer operators related to the uniform family of maps $A_{M,L}$ satisfy. \square

The following statement is an estimate for the speed of convergence to equilibrium of mean field coupled expanding maps (see [29], Theorem 4 or [34] Theorem 1.1 for similar statements).

Proposition 23 (Exponential convergence to equilibrium). *Let \mathcal{L}_δ be the family of self consistent transfer operators arising from $T_0 \in C^6$ and $h \in C^6$ as above. Let $f_\delta \in W^{1,1}$ be an invariant probability density of \mathcal{L}_δ . Then there exists $\bar{\delta} > 0$ and $C, \gamma \geq 0$ such that for each $0 < \delta < \bar{\delta}$, and each probability density $\nu \in W^{1,1}$ we have*

$$\|\mathcal{L}_\delta^n(\nu) - f_\delta\|_{W^{1,1}} \leq C e^{-\gamma n} \|\nu - f_\delta\|_{W^{1,1}}.$$

Proof. The proof is an application of Theorem 6, considering $B_{ss} = W^{2,1}$, $B_s = W^{1,1}$, $B_w = L^1$. Let $L_{\delta f}$ be the family of transfer operators associated to this system.

By the Lasota Yorke inequalities we have that $L_{\delta, \mu} : W^{2,1} \rightarrow W^{2,1}$, $L_{\delta, \mu} : W^{1,1} \rightarrow W^{1,1}$, $L_{\delta, \mu} : L^1 \rightarrow L^1$ with $\mu \in B_w(0,1) \subseteq B_w$ are bounded uniformly on δ . By Lemma 17 they satisfy (Con1). Furthermore by Lemma 22 and Proposition 19 they satisfy (Con2).

By Proposition 20 the invariant measures f_δ satisfy $\lim_{\delta \rightarrow 0} \|f_\delta\|_{W^{2,1}} < +\infty$. Since the circle expanding map T_0 has convergence to equilibrium then (Con3) is satisfied. We can then apply Theorem 6 directly implying the statement. \square

To get some useful formula for the linear response for expanding maps coupled in a mean field regime, let us now consider small perturbations of expanding maps $T : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ by left composition with a family of diffeomorphisms $(D_\delta)_{\delta \in [-\epsilon, \epsilon]}$. More precisely, let $D_\delta : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ be a diffeomorphism, with

$$(53) \quad D_\delta = \pi_{\mathbb{S}^1} \circ (Id + \delta S)$$

and $S \in C^6(\mathbb{S}^1, \mathbb{R})$. In this setting let us define the perturbed transfer operators as

$$L_\delta = L_{D_\delta} \circ L_T$$

(remark that here $L_0 = L_T$). These kinds of perturbations are of the type induced by a mean field coupling, they satisfy the "small perturbation" and "existence of a derivative operator" assumptions of our general theorems like (Con2) or (LR2) of Theorem 14 we have indeed (see [14, Proposition 35, 36]):

Theorem 24. *Let $(D_\delta)_{\delta \in [0, \bar{\delta}]}$ be as in (53), and $T : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ be a C^6 uniformly expanding map. Let us define $\dot{L} : W^{4,1}(\mathbb{S}^1) \rightarrow W^{3,1}(\mathbb{S}^1)$ by*

$$(54) \quad \dot{L}(f) := -(S \cdot L_T(f))'.$$

Then one has that for each $1 \leq k \leq 4$ and $f \in W^{k,1}$

$$(55) \quad \left\| \frac{L_\delta - L_0}{\delta}(f) - \dot{L}(f) \right\|_{W^{k-1,1}} \xrightarrow{\delta \rightarrow 0} 0.$$

We have now all the ingredients to prove a result regarding the Linear Response of the coupled system in the small coupling regime.

Theorem 25 (Linear Response for coupled expanding maps (zero coupling limit)). *Consider the family of self consistent transfer operators \mathcal{L}_δ associated to a C^6 expanding map T and a coupling driven by the function h , with $h \in C^6$. Let h_0 be the unique invariant probability measure in L^1 for \mathcal{L}_0 and h_δ some invariant probability measure for \mathcal{L}_δ . Then for $\delta \rightarrow 0$ we have the following Linear Response formula*

$$(56) \quad \lim_{\delta \rightarrow 0} \left\| \frac{h_\delta - h_0}{\delta} + (Id - L_0)^{-1} (h_0 \int_{\mathbb{S}^1} h(x, y) h_0(y) dy)' \right\|_{W^{1,1}} = 0.$$

Proof. The proof is a direct application of Theorem 14 to our case with $B_{s_s} = W^{3,1}(\mathbb{S}^1) \subset B_s = W^{2,1}(\mathbb{S}^1) \subset B_w = W^{1,1}(\mathbb{S}^1)$. Let us see why the assumptions needed to apply the theorem are satisfied. We recall that the transfer operators $\mathcal{L}_\delta : W^{3,1}(\mathbb{S}^1) \rightarrow W^{3,1}(\mathbb{S}^1)$ involved are defined by

$$\mathcal{L}_\delta(\phi) = Q_{\delta, \phi}(L_{T_0}(\phi)).$$

The assumption (SS1) (regularity bounds), is implied by Proposition 20. The assumption (SS2) (convergence to equilibrium for the unperturbed operator), is well known to be verified, as it stands for the unperturbed transfer operator \mathcal{L}_0 which is the transfer operator associated to a smooth expanding map. The assumption (LR1) regarding the existence of the resolvent of the unperturbed operator on the weak space $W^{1,1}$ follows from Proposition 18.

As \mathcal{L}_δ is a small perturbation of \mathcal{L}_0 given by the composition of the transfer operator $Q_{\delta, \phi}$ associated to a diffeomorphism near to the identity, the assumption (SS3) and the first part of (LR2) (small perturbation) follows from Theorem 24, Proposition 19 and Lemma 22 as before. Let us prove indeed that there is $K \geq 0$ such that and $\mathcal{L}_0 - \mathcal{L}_\delta$ is $K\delta$ -Lipschitz when considered as a function $\overline{B}_{2M} \rightarrow B_w$ and $B_{2M} \rightarrow B_s$. In the first case we have to prove that for $\phi_1, \phi_2 \in W^{2,1}$

$$(57) \quad \|[Q_{\delta, \phi_1}(L_{T_0}(\phi_1)) - L_{T_0}(\phi_1)] - [Q_{\delta, \phi_2}(L_{T_0}(\phi_2)) - L_{T_0}(\phi_2)]\|_{W^{1,1}} \leq K\delta \|\phi_1 - \phi_2\|_{W^{2,1}}.$$

Developing the formula we get

$$\begin{aligned} & \|Q_{\delta, \phi_1}(L_{T_0}(\phi_1)) - L_{T_0}(\phi_1) - Q_{\delta, \phi_2}(L_{T_0}(\phi_2)) + L_{T_0}(\phi_2)\|_{W^{1,1}} \\ & \leq \|Q_{\delta, \phi_1}(L_{T_0}(\phi_1)) - L_{T_0}(\phi_1) - Q_{\delta, \phi_1}(L_{T_0}(\phi_2)) \\ & \quad + Q_{\delta, \phi_1}(L_{T_0}(\phi_2)) - Q_{\delta, \phi_2}(L_{T_0}(\phi_2)) + L_{T_0}(\phi_2)\|_{W^{1,1}} \end{aligned}$$

And

$$\begin{aligned} & \|Q_{\delta, \phi_1}(L_{T_0}(\phi_1)) - L_{T_0}(\phi_1) - Q_{\delta, \phi_1}(L_{T_0}(\phi_2)) + L_{T_0}(\phi_2)\|_{W^{1,1}} \\ & \leq \|Q_{\delta, \phi_1}(L_{T_0}(\phi_1 - \phi_2)) - L_{T_0}(\phi_1 - \phi_2)\|_{W^{1,1}} \\ & \leq CK_2 2M\delta \|\phi_1 - \phi_2\|_{W^{2,1}} \end{aligned}$$

by applying Lemma 22 with $\psi = 0$ and $\phi = \phi_1$ and Proposition 19. The other term can be estimated as

$$\begin{aligned} \|Q_{\delta, \phi_1}(L_{T_0}(\phi_2)) - Q_{\delta, \phi_2}(L_{T_0}(\phi_2))\|_{W^{1,1}} &\leq \|Q_{\delta, \phi_1} \circ L_{T_0} - Q_{\delta, \phi_2} \circ L_{T_0}\|_{W^{2,1} \rightarrow W^{1,1}} \|\phi_2\|_{W^{2,1}} \\ &\leq \delta CK_2 2M \|\phi_1 - \phi_2\|_{W^{2,1}} \end{aligned}$$

using Proposition 19 and Lemma 22 and proving the Lipschitz assumption in the $\overline{B}_{2M} \rightarrow B_w$ case. The case $B_{2M} \rightarrow B_s$ is similar.

The assumption (LR2) on the derivative operator follows from Theorem 24. Let us apply it and find an expression for $\dot{\mathcal{L}}h_0$ in our case. In this case the perturbing operator to be considered is Q_{δ, h_0} associated with the diffeomorphism Φ_{δ, h_0} . With the notation (53) we have $D_\delta = \Phi_{\delta, h_0} = Id + \delta S$ with

$$S(x) = \int_{S^1} h(x, y) h_0(y) dy$$

and then

$$(58) \quad \dot{\mathcal{L}}(h_0) = -(S L_{T_0} h_0)' = -(h_0 \int_{S^1} h(x, y) h_0(y) dy)'.$$

Applying Theorem 14, we then get

$$(59) \quad \lim_{\delta \rightarrow 0} \left\| \frac{h_\delta - h_0}{\delta} - (Id - L_{T_0})^{-1} \dot{\mathcal{L}}(h_0) \right\|_{W^{1,1}} = 0$$

as in our case $W^{1,1}$ is the weak space. Substituting (58) in (59) we get (56). \square

Remark 26. From (40) we see that this response result with convergence in the quite strong topology $W^{1,1}$ gives information on the behavior of a large class of observables, for example we can consider observables in L^∞ or L^1 or even distributions in the dual of $W^{1,1}$.

7. MEAN FIELD COUPLED CONTINUOUS MAPS

We show the flexibility of Theorem 2 proving the existence of invariant measure in the general case of continuous maps interacting by a Lipschitz coupling function h .

Proposition 27. *Let us consider a coupled maps system as described in Section 2 with a map $T_0 \in C^0(\mathbb{S}^1 \rightarrow \mathbb{S}^1)$, $h \in Lip(\mathbb{S}^1 \times \mathbb{S}^1 \rightarrow \mathbb{R})$ and $\delta \geq 0$, then there is $\mu \in PM(\mathbb{S}^1)$ such that*

$$\mathcal{L}_\delta(\mu) = \mu.$$

Proof. Let us consider the space of signed Borel measures $SM(\mathbb{S}^1)$, we put two different norms on this space. $\|\cdot\|_w, \|\cdot\|_s$ defined by $\|\mu\|_w = \sup_{g \in Lip(\mathbb{S}^1 \rightarrow \mathbb{R}), \|g\|_{Lip} \leq 1} \int g d\mu$ and $\|\mu\|_s = \mu^+(\mathbb{S}^1) + \mu^-(\mathbb{S}^1)$ (the total variation norm). We apply Theorem 2 with $(SM(\mathbb{S}^1), \|\cdot\|_w)$, $(SM(\mathbb{S}^1), \|\cdot\|_s)$ as a weak and strong space. We remark that by the Prokhorov's theorem, P_w is complete when considered with the $\|\cdot\|_w$ norm.

Now let us define a projection π_n as requested by Theorem 2. Let us consider $n \in \mathbb{N}$ and divide \mathbb{S}^1 into n equal intervals I_1, \dots, I_n , with $I_i = [x_i, x_{i+1})$. Let us consider a partition of unity $\{\phi_1, \dots, \phi_n\}$ made of hat functions ϕ_i which are

linear on each interval of the partition and supported on $I_{i \bmod(n)} \cup I_{i+1 \bmod(n)}$ let us consider the projection $\pi_n : SM(\mathbb{S}^1) \rightarrow SM(\mathbb{S}^1)$ defined by

$$\pi_n(\mu) = \sum_{i \leq n} \delta_{x_{i+1}} \int \phi_i d\mu$$

we have that this projection is linear, preserves probability measures, and $\|\pi_n(\mu)\|_s \leq \|\mu\|_s$, $\|\pi_n(\mu)\|_w \leq \|\mu\|_w$ (the first inequality is straightforward, for the second see [20, Proposition 9.4]) furthermore since each element of measure composing μ is moved by π_n by a distance $\leq \frac{2}{n}$ in the weak norm, we get

$$(60) \quad \|\pi_n(\mu) - \mu\|_w \leq \frac{2}{n} \|\mu\|_s.$$

Each invariant probability measure μ for each $L_{\delta, \mu}$ is such that $\|\mu\|_s \leq 1$. The same can be said for the finite dimensional reduced operator $\pi_n L_{\delta, \pi_n \mu} \pi_n$, hence *Exi1*, *Exi1.b* are satisfied.

To verify *Exi2* we have to verify that

$$(61) \quad \|[L_{\delta, \mu_1} - L_{\delta, \mu_2}]\mu\|_w \leq \delta K \|\mu\|_s \|\mu_1 - \mu_2\|_w$$

we remark that if h is K Lipschitz

$$\begin{aligned} |\Phi_{\delta, \mu_1}(x) - \Phi_{\delta, \mu_2}(x)| &= \delta \int h(x, y) d[\mu_1 - \mu_2](y) \\ &\leq \delta K \|\mu_1 - \mu_2\|_w. \end{aligned}$$

Hence

$$\begin{aligned} \|[L_{\delta, \mu_1} - L_{\delta, \mu_2}]\mu\|_w &= \|[L_{\Phi_{\delta, \mu_1}} - L_{\Phi_{\delta, \mu_2}}]L_T \mu\|_w \\ &\leq \delta K \|\mu_1 - \mu_2\|_w \|L_T \mu\|_s \\ &\leq \delta K \|\mu_1 - \mu_2\|_w \|\mu\|_s \end{aligned}$$

leading to 61.

We can then apply Theorem 2 leading to the existence of an invariant probability measure for \mathcal{L}_δ . \square

Remark 28. For simplicity the statement is proved for maps on \mathbb{S}^1 . It seems that the statement can be generalized with the same idea to maps on compact metric spaces for which there is a sequence of Lipschitz partitions of unity, which can be used to define suitable projections π_n on combinations of delta measures placed on some sequences of ϵ -nets covering the space.

8. MAPS WITH ADDITIVE NOISE ON \mathbb{S}^1

We illustrate the flexibility of our approach with an application to a system of coupled random maps. For simplicity we will consider a class of random dynamical systems on the unit circle \mathbb{S}^1 . Informally speaking, a random dynamics on \mathbb{S}^1 is defined by the iteration of maps chosen randomly in the family $T_\omega : \mathbb{S}^1 \rightarrow \mathbb{S}^1$, $\omega \in \Omega$ according to a certain probability distribution p defined on Ω . In our case we will model this random choice as independent and identically distributed at each time.

Let $T_0: \mathbb{S}^1 \rightarrow \mathbb{S}^1$ a continuous and piecewise C^1 , nonsingular map⁶. We consider a random dynamical system, corresponding to the stochastic process $(X_n)_{n \in \mathbb{N}}$ defined by

$$(62) \quad X_{n+1} = T_0(X_n) + \Omega_n \pmod{1}$$

where $(\Omega_n)_{n \in \mathbb{N}}$ are i.i.d random variables distributed according to some smooth kernel ρ . We will call T_0 the deterministic part of the system and ρ the noise kernel of the system.

Remark 29. *We remark that the maps considered here are quite general. We do not require expansiveness or hyperbolicity, allowing many examples of random maps coming as models of natural phenomena (see e.g. [19],[8],[31]).*

We will consider the annealed transfer operators associated to these systems (see [37], Section 5 for more details about transfer operators associated to this kind of systems or [14]). Let $SM(\mathbb{S}^1)$ be the space of signed Borel measures in \mathbb{S}^1 . The (annealed) transfer operator $L : SM(\mathbb{S}^1) \rightarrow SM(\mathbb{S}^1)$ associated to the system will be defined by

$$(63) \quad L(\mu) = \int_{\Omega} L_{T_\omega}(\mu) dp$$

where $L_{T_\omega}(\mu) : SM(\mathbb{S}^1) \rightarrow SM(\mathbb{S}^1)$ is the transfer operator associated to T_ω , hence taking the average of the pushforward maps with respect to p . For some class of random dynamical systems L is defined as an operator $: L^1(\mathbb{S}^1) \rightarrow L^1(\mathbb{S}^1)$ and sometime this operator is a *kernel* operator: let $k \in L^\infty(\mathbb{S}^1 \times \mathbb{S}^1)$ (the kernel), consider the operator L defined in the following way: for $f \in L^1(\mathbb{S}^1)$

$$(64) \quad Lf(x) = \int k(x, y)f(y)dy.$$

This kind of operators naturally appear when the random dynamics is defined by the action of a deterministic map and some additive noise. Since the effect of the additive noise is to perturb the deterministic map by a translation, the annealed transfer operator will be an average of translations, i.e. a convolution. The well known regularization properties of convolutions then imply that the annealed transfer operator associated to a system with additive noise is a regularizing one.

Let us consider a probability density $\rho : \mathbb{R} \rightarrow \mathbb{R}$ representing how the noise is distributed. For simplicity we will suppose ρ being such that $\rho(x) = \rho(-x)$ for each $x \in \mathbb{R}$ and being a Schwartz function, hence and $\rho \in C^k$ for each $k \geq 1$. The periodization $\tilde{\rho} : \mathbb{S}^1 \rightarrow \mathbb{R}$ of such a function is defined as

$$\tilde{\rho}(x) = \sum_{k \in \mathbb{Z}} \rho(x + k)$$

which clearly converge for a rapidly decreasing function as ρ .

Definition 30. *Let $f \in L^1(\mathbb{S}^1)$ and ρ as before. We define the convolution $\rho * f$ by*

$$(65) \quad \rho * f(x) := \int_{\mathbb{S}^1} \tilde{\rho}(x - y)f(y)dy.$$

⁶We mean that \mathbb{S}^1 can be partitioned in a finite set of intervals where T is C^1 and that the associated pushforward map T_* sends a measure which is absolutely continuous with respect to the Lebesgue measure to another measure which is absolutely continuous with respect to the Lebesgue measure.

To a system with additive noise as defined in (62) we then associate the annealed transfer operator $L : L^1 \rightarrow L^1$ defined by

$$(66) \quad L(\phi) := \rho * L_{T_0}(\phi)$$

where L_{T_0} is the transfer operator associated to the deterministic map T_0 and $\phi \in L^1$.

We now define the self consistent transfer operator associated to an infinite collection of interacting random maps in a mean field coupling. Like in the deterministic case let us consider $h \in C^k(\mathbb{S}^1 \times \mathbb{S}^1, \mathbb{R})$, $\delta \geq 0$, for some $k \geq 1$, a probability density $\psi \in L^1$, let $\pi : \mathbb{S}^1 \rightarrow \mathbb{R}$ be the natural projection, and define $\Phi_{\delta, \psi} : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ again as

$$\Phi_{\delta, \psi}(x) = x + \pi(\delta \int_{\mathbb{S}^1} h(x, y) \psi(y) dy).$$

Denote by $Q_{\delta, \psi}$ the transfer operator associated to $\Phi_{\delta, \psi}$, as in (45). We consider the following family of operators depending on $\phi \in L^1$ and $\delta \geq 0$ defined as

$$(67) \quad L_{\delta, \phi}(\psi) = \rho * [Q_{\delta, \phi}(L_{T_0}(\psi))]$$

for each $\psi \in L^1$. Finally we define the nonlinear self consistent transfer operator associated to this system of coupled random maps as $\mathcal{L}_\delta : L^1(\mathbb{S}^1, \mathbb{R}) \rightarrow L^1(\mathbb{S}^1, \mathbb{R})$ defined by

$$(68) \quad \mathcal{L}_\delta(\phi) = \rho * [Q_{\delta, \phi}(L_{T_0}(\phi))].$$

This represents the idea that a certain initial condition is first moved by the deterministic part of the dynamics represented by the map T_0 and by the mean field perturbation $\Phi_{\delta, \phi}$, then a further (external and independent of the dynamics) random perturbation is applied by the noise to every coupled map. In the remaining part of the section we will apply our general theory to this kind of operators.

The following proposition contains some of the regularization properties for the convolution we need (see [14] Proposition 15 for the proof and details).

Proposition 31. *Let $f \in L^1$ and ρ be as before. The convolution $\rho * f$ has the following properties:*

- (1) *For each $k \geq 1$, $\rho * f : \mathbb{S}^1 \rightarrow \mathbb{R}$ is C^k , and $(\rho * f)^{(i)} = \rho^{(i)} * f$ for any $i \leq k$.*
- (2) *One has the following regularization inequality:*

$$(69) \quad \|\rho * f\|_{C^k} \leq \|\rho\|_{C^k} \|f\|_{L^1}.$$

These regularization properties together with the Ascoli Arzela theory imply that a linear operator $L_0 : L^1 \rightarrow L^1$ of the kind

$$(70) \quad L_0(\phi) = \rho * L_{T_0}(\phi)$$

is a compact operator. If we suppose that the system satisfy a convergence to equilibrium assumption, this will allow to obtain the spectral gap and the existence of the resolvent operators, required to apply Theorem 14. More precisely, let us suppose that L_0 satisfy a convergence to equilibrium condition: for each $g \in C^k$ such that $\int g dm = 0$

$$(71) \quad \lim_{n \rightarrow \infty} \|L_0^n g\|_{L^1} = 0.$$

This easily implies that $\lim_{n \rightarrow \infty} \|L_0^n g\|_{C^k} = 0$ for each $g \in L^1$ such that $\int g dm = 0$ and then some iterate of the transfer operator is a uniform contraction on the space

of C^k densities with zero average. By this the operator has spectral gap, implying the existence of the resolvent operator (see for the details [14], section IV).

Corollary 32. *Let L_0 be the annealed transfer operator associated to a map T_0 with additive noise distributed with a kernel ρ as in (70) and suppose it has convergence to equilibrium (71) then for each $k \geq 1$, $(Id - L_0)^{-1} := \sum_{i=0}^{\infty} L_0^i$ is a bounded operator $C^k \rightarrow C^k$.*

By [14, proposition 18 and 19] and their simple proof it also directly follows that

Lemma 33. *Let us consider transfer operators L_0, L_1 associated to dynamical systems with additive noise having noise kernel ρ , deterministic part given by continuous maps T_0 and T_1 and $k \geq 0$. Then there is $C \geq 0$ such that for all such T_0, T_1 and $f \in L^1$*

$$\|L_1 f - L_0 f\|_{C^{k-1}} \leq C \|\rho\|_{C^k} \|T_0 - T_1\|_{C^0} \|f\|_1.$$

Now let us consider the self consistent transfer operator \mathcal{L}_δ as defined in (68). We state a result analogous to Proposition 20 in the case of systems with additive noise. The application to this case is simpler due to the regularizing effect of the noise.

Proposition 34. *Suppose T_0 is $\mathbb{S}^1 \rightarrow \mathbb{S}^1$ continuous, nonsingular and piecewise C^1 . Suppose $h \in C^k$ with $k \geq 1$, let \mathcal{L}_δ be the self consistent family of operators associated to this coupled system as defined in (68), then for each $\delta \geq 0$ there is $f_\delta \in C^\infty$ such that for each $k \geq 1$*

$$(72) \quad \|f_\delta\|_{C^k} \leq \|\rho\|_{C^k}$$

and

$$\mathcal{L}_\delta(f_\delta) = f_\delta.$$

Let us suppose that the (linear) operator \mathcal{L}_0 has convergence to equilibrium when considered as acting on the spaces C^1 and L^1 (see (71)) then there is $\bar{\delta} > 0$ such that for each $\delta \in [0, \bar{\delta}]$ f_δ is unique.

Proof. We sketch the proof, whose arguments are similar to the ones of Proposition 20. We will obtain the statement applying Theorems 2 and 3 to the family of operators

$$L_{\delta, \phi} = \rho * [Q_{\delta, \phi} \circ L_{T_0}]$$

as defined in (67). First we will apply theorem 2 with $B_s = BV$ and $B_w = L^1$. We remark that given $k \geq 1$, (69) implies a Lasota Yorke inequality which is uniformly satisfied by these operators, indeed

$$(73) \quad \begin{aligned} \|L_{\delta, \phi}(\psi)\|_{C^k} &= \|\rho * [Q_{\delta, \phi} \circ L_{T_0}(\psi)]\|_{C^k} \\ &\leq 0 \|\psi\|_{C^k} + \|\rho\|_{C^k} \|[Q_{\delta, \phi} \circ L_{T_0}(\psi)]\|_1 \\ &\leq 0 \|\psi\|_{C^k} + \|\rho\|_{C^k} \|\psi\|_1. \end{aligned}$$

This implies that the transfer operators in the family are uniformly bounded as operators $L^1 \rightarrow C^k$ and hence each invariant probability measure $f_{\delta, \phi}$ of $L_{\delta, \phi}$ is such that $\|f_{\delta, \phi}\|_{BV} \leq \|f_{\delta, \phi}\|_{C^k} \leq \|\rho\|_{C^k}$ and then the family of operators satisfy (Exi1).

The assumption (Exi2) (or (12)) is provided similarly as a consequence of the Lemma 22 and Lemma 33.

In order to apply Theorem 2 we consider a suitable projection π_n . As in the proof of Proposition 20, let \mathcal{P}_n be the partition subdividing the circle into n equal intervals. Consider π_n defined as in the proof of Proposition 20 by $\pi_n(f) = \mathbf{E}(f|\mathcal{P}_n)$, where the conditional expectation is made using the Lebesgue measure, projecting to piecewise constant functions. Again, the discretized operators satisfy a uniform Lasota Yorke inequality on BV and L^1 , indeed

$$\begin{aligned} \|\pi_n L_{\delta,\phi} \pi_n(\psi)\|_{BV} &\leq \|L_{\delta,\phi} \pi_n(\psi)\|_{BV} \\ &\leq \|L_{\delta,\phi} \pi_n(\psi)\|_{C^k} \\ &\leq \|\rho * [Q_{\delta,\phi} \circ L_{T_0}(\pi_n(\psi))]\|_{C^k} \\ &\leq \|\rho\|_{C^k} \|\pi_n(\psi)\|_1 \\ &\leq \|\rho\|_{C^k} \|\psi\|_1 \end{aligned}$$

and *Exi1.b* is satisfied. We can then apply Theorem 2, and get the existence of an invariant probability density f in BV . Since $\rho \in C^k$ and $L_{\delta,f} f = f$ we get that $f \in C^k$ for each $k \geq 1$, also proving (72).

Now we apply Theorem 3 to get the uniqueness. In this case we consider $B_s = C^1$ and $B_w = L^1$. We first have to prove that for δ small enough each operator $L_{\delta,\phi}$ with $\phi \in P_w$ has a unique invariant probability measure in P_w . Since L_0 has convergence to equilibrium, is regularizing and C^1 is compactly immersed in L^1 it is standard to find that this operator has a unique invariant probability measure. From the convergence to equilibrium, the small perturbation assumption (*Exi2*) we verified above and the regularization property (69) we get that there is $\gamma \geq 0$ such that $L_{\delta,\phi}$ has convergence to equilibrium for each $\delta \leq \gamma$ and $\phi \in P_w$. Indeed let us consider $f \in V_s$ and suppose that by convergence to equilibrium n is such that $\|L_0^n f\|_{L^1} \leq \frac{1}{2\|\rho\|_{C^1}} \|f\|_{C^1}$ then

$$\|L_0^{n+1} f\|_{C^1} \leq \|\rho\|_{C^1} \|L_{T_0}(L_0^n f)\|_{L^1} \leq \frac{1}{2} \|f\|_{C^1}.$$

This implies that L_0^{n+1} is a contraction of V_s . Now let us consider $\phi \in P_w$, $\gamma \leq \frac{1}{4\|\rho\|_{C^1}K(C+nB)}$ and $\delta \leq \gamma$. By Lemma 8 (remark that $Q = 1$ in the case $B_w = L^1$ and by (73), $\lambda = 0$)

$$(74) \quad \|L_{\delta,\phi}^n g - L_0^n g\|_{L^1} \leq \delta K(C\|g\|_{C^1} + nB\|g\|_{L^1})$$

$$(75) \quad \leq \delta K(C + nB)\|g\|_{C^1}$$

$$(76) \quad \leq \frac{1}{4\|\rho\|_{C^1}} \|g\|_{C^1}$$

and then $\|L_{\delta,\phi}^n g\|_{L^1} \leq \frac{3}{4\|\rho\|_{C^1}} \|g\|_{C^1}$, thus repeating the same reasoning as before $L_{\delta,\phi}^{n+1}$ also is a contraction of V_s . Hence we have that each $L_{\delta,\phi}$ has convergence to equilibrium. It follows that $L_{\delta,\phi}$ also has spectral gap on C^1 and on L^1 . Indeed for each $f \in V_{L^1} := \{f \in L^1, \int f = 0\}$ we get $\|L_{\delta,\phi} f\|_{C^1} \leq \|\rho\|_{C^1} \|f\|_{L^1}$ and if we have if n is such that $\|L_{\delta,\phi}^n f\|_{L^1} \leq \frac{3}{4\|\rho\|_{C^1}} \|f\|_{C^1}$ then again

$$\|L_{\delta,\phi}^{n+1} f\|_{L^1} \leq \frac{3}{4\|\rho\|_{C^1}} \|L_{\delta,\phi} f\|_{C^1} \leq \frac{3}{4} \|f\|_{L^1}.$$

Thus each $L_{\delta,\phi}$ has a unique invariant probability measure $f_\phi \in P_w$. Furthermore, the resolvent of $L_{\delta,\phi}$ is defined on V_{L^1} .

Now we can prove that (Exi3) holds. Let us consider probability measures ϕ_1 and $\phi_2 \in P_w$ and the operators L_{δ, ϕ_1} L_{δ, ϕ_2} we have seen that when $\delta \leq \gamma$ these operators have unique fixed probability densities f_{ϕ_1}, f_{ϕ_2} . We want to prove that

$$\|f_{\phi_1} - f_{\phi_2}\|_{L^1} \leq K_2 \|\phi_1 - \phi_2\|_{L^1}.$$

We hence apply a construction similar to the proof of Theorem 14 to the family of operators $\hat{L}_\epsilon : L^1 \rightarrow L^1$ defined by

$$\hat{L}_\epsilon = L_{\delta, \phi_1} + \epsilon[L_{\delta, \phi_2} - L_{\delta, \phi_1}].$$

Consider

$$\begin{aligned} (Id - L_{\delta, \phi_2})(f_{\phi_2} - f_{\phi_1}) &= f_{\phi_2} - L_{\delta, \phi_2}f_{\phi_2} - f_{\phi_1} + L_{\delta, \phi_2}f_{\phi_1} \\ &= (L_{\delta, \phi_2} - L_{\delta, \phi_1})f_{\phi_1}. \end{aligned}$$

We have that

$$(f_{\phi_2} - f_{\phi_1}) = (Id - L_{\delta, \phi_2})^{-1}(L_{\delta, \phi_2} - L_{\delta, \phi_1})f_{\phi_1}.$$

By the fact that $(Id - L_{\delta, \phi_2})^{-1}$ is well defined and continuous on V_{L^1} remarked before and by the fact that $\|f_{\phi_1}\|_{C^1} \leq \|\rho\|_{C^1}$ and (Exi2) we get

$$\|f_{\phi_2} - f_{\phi_1}\|_{L^1} \leq \delta K \|\rho\|_{C^1} \|\phi_2 - \phi_1\|_{L^1}$$

and then (Exi3) is verified. Applying Theorem 3 we then get the uniqueness for δ small enough. \square

Proposition 35 (Exponential convergence to equilibrium). *Let \mathcal{L}_δ be the family of self consistent transfer operators arising from a map T_0 , a kernel ρ and $h \in C^k$ as above. Suppose the uncoupled system \mathcal{L}_0 has convergence to equilibrium. Let f_δ be an invariant probability measure of \mathcal{L}_δ . Let $k \geq 1$, then there exists $\bar{\delta} > 0$ and $C, \gamma \geq 0$ such that for each $0 < \delta < \bar{\delta}$ and each probability density $\nu \in C^k$ we have*

$$\|\mathcal{L}_\delta^n(\nu) - f_\delta\|_{C^k} \leq C e^{-\gamma n} \|\nu - f_\delta\|_{C^k}.$$

Proof. Again the proof is a direct application of Theorem 6 to \mathcal{L}_δ considering the spaces $B_{ss} = C^{k+1}$, $B_s = C^k$ and $B_w = L^1$. The assumption (Con1) for this kind of systems is already verified in (73). The assumption (Con2) is as a direct consequence of Lemma 22 and Lemma 33. The assumption (Con3) is required as an assumption in this statement. \square

Let us now consider a linear response result for the invariant measure of \mathcal{L}_δ when $\delta \rightarrow 0$ in the case of coupled maps with additive noise.

Theorem 36 (Linear Response for coupled maps with additive noise). *Let \mathcal{L}_δ be the family of self consistent transfer operators arising from a map T_0 , a kernel ρ and $h \in C^{k+1}$ as above. Suppose that $k \geq 2$ and the uncoupled initial transfer operator \mathcal{L}_0 has convergence to equilibrium in the sense of (71). Let $f_0 \in C^{k+1}$ be the unique invariant probability density of L_0 and $f_\delta \in C^{k+1}$ be the invariant probability density of L_δ (unique when δ is small enough). Then we have the following Linear Response formula*

$$(77) \quad \lim_{\delta \rightarrow 0} \left\| \frac{f_\delta - f_0}{\delta} + (Id - L_0)^{-1} \rho_\xi * (L_{T_0}(h_0) \int_{S^1} h(x, y) h_0(y) dy)' \right\|_{C^k} = 0.$$

Before the proof, we recall some preliminary result on the response of systems with additive noise. We now consider small perturbations of our random maps with additive noise by composition with a map D_δ , which is when δ is small, a diffeomorphism near to the identity. Consider a map $S \in C^2(\mathbb{S}^1, \mathbb{R})$ and the map $D_\delta : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ defined by $D_\delta = Id + \delta S$. Let us consider then the perturbation of T_0 by composition with D_δ defined by $T_\delta = T_0 \circ D_\delta$. Starting from this family of maps and a kernel ρ we can consider a family of dynamical systems with additive noise as in (62). Since $L_{T_\delta} = L_{D_\delta} \circ L_{T_0}$, to this system we associate the annealed transfer operator defined by

$$(78) \quad L_\delta := \rho * L_{D_\delta \circ T_0} = \rho * (L_{D_\delta} \circ L_{T_0}).$$

Now, in order to apply our general theorems we recall some estimates for this kinds of perturbations which will allow to check that the "small perturbation" assumptions like (Con2) or (LR2) applies to this case. It will be also useful to consider the derivative operator \dot{L} for the family of operators $(L_{D_\delta})_{\delta \in [0, \bar{\delta}]}$. In this direction, the following result ([14, Theorem 24]) gives some useful estimates.

Theorem 37. *Let $(L_\delta)_{\delta \in [0, \bar{\delta}]}$ be the family of transfer operators as described in (78). For each $k \geq 2$*

$$(79) \quad \lim_{\delta \rightarrow 0} \left\| \frac{L_\delta - L_0}{\delta} - \dot{L} \right\|_{C^k \rightarrow C^{k-1}} = 0$$

where $\dot{L} : C^k(\mathbb{S}^1) \rightarrow C^{k-1}(\mathbb{S}^1)$ is defined by:

$$\dot{L}(f) = -\rho * (S \cdot L_{T_0} f)'$$

Proof of Theorem 36. Similar to the proof of Theorem 25, the proof is a direct application of Theorem 14. We apply Theorem 14 to the family of operators $\mathcal{L}_\delta(\phi) = \rho * [Q_{\delta, \phi}(L_{T_0}(\phi))]$ considering the spaces $B_{ss} = C^{k+2}(\mathbb{S}^1) \subseteq B_s = C^{k+1}(\mathbb{S}^1) \subset B_w = C^k(\mathbb{S}^1)$. Let us now verify that the assumptions needed to apply the theorem are satisfied. The assumption (SS1) (regularity bounds), is implied by Proposition 34.

The assumption (SS2)(convergence to equilibrium for the unperturbed operator), is supposed to hold in the statement of Theorem 36. The assumption (SS3) and the first part of the assumption (LR2) (small perturbation) follows from Lemma 22 and 33. Indeed we have to prove that there is $K \geq 0$ such that $\mathcal{L}_0 - \mathcal{L}_\delta$ is $K\delta$ -Lipschitz when considered as a function $\overline{B}_{2M} \rightarrow B_w$ and $B_{2M} \rightarrow B_s$. In the first case we have to prove that for $\phi_1, \phi_2 \in \{\phi \in C^{k+1}, \|\phi\|_{C^{k+1}} \leq 2M\}$

$$\|\rho * [Q_{\delta, \phi_1}(L_{T_0}(\phi_1)) - L_{T_0}(\phi_1)] - \rho * [Q_{\delta, \phi_2}(L_{T_0}(\phi_2)) - L_{T_0}(\phi_2)]\|_{C^k} \leq K\delta \|\phi_1 - \phi_2\|_{C^{k+1}}$$

We have

$$\begin{aligned} & \|\rho * [Q_{\delta, \phi_1}(L_{T_0}(\phi_1)) - L_{T_0}(\phi_1)] - \rho * [Q_{\delta, \phi_2}(L_{T_0}(\phi_2)) + L_{T_0}(\phi_2)]\|_{C^k} \\ & \leq \|\rho * Q_{\delta, \phi_1}(L_{T_0}(\phi_1)) - \rho * L_{T_0}(\phi_1) - \rho * Q_{\delta, \phi_1}(L_{T_0}(\phi_2)) \\ & \quad + \rho * Q_{\delta, \phi_1}(L_{T_0}(\phi_2)) - \rho * Q_{\delta, \phi_2}(L_{T_0}(\phi_2)) + \rho * L_{T_0}(\phi_2)\|_{C^k} \end{aligned}$$

and

$$\begin{aligned} & \|\rho * Q_{\delta, \phi_1}(L_{T_0}(\phi_1)) - \rho * L_{T_0}(\phi_1) - \rho * Q_{\delta, \phi_1}(L_{T_0}(\phi_2)) + \rho * L_{T_0}(\phi_2)\|_{C^k} \\ & \leq \|\rho * Q_{\delta, \phi_1}(L_{T_0}(\phi_1 - \phi_2)) - \rho * L_{T_0}(\phi_1 - \phi_2)\|_{C^k} \\ & \leq \delta CK_1 2M \|\phi_1 - \phi_2\|_{C^{k+1}} \end{aligned}$$

applying Lemma 22 with $\psi = 0$ and $\phi = \phi_1$ and Lemma 33.

Furthermore

$$\|\rho * Q_{\delta, \phi_1}(L_{T_0}(\phi_2)) - \rho * Q_{\delta, \phi_2}(L_{T_0}(\phi_2))\|_{C^k} \leq \delta CK_1 2M \|\phi_1 - \phi_2\|_{C^{k+1}}$$

again by Lemma 22 and Lemma 33.

In the second case have to prove that for $\phi_1, \phi_2 \in \{\phi \in C^{k+2}, \|\phi\|_{C^{k+2}} \leq 2M\}$

$$\|\rho * [Q_{\delta, \phi_1}(L_0(\phi_1)) - L_0(\phi_1)] - \rho * [Q_{\delta, \phi_2}(L_0(\phi_2)) - L_0(\phi_2)]\|_{C^{k+1}} \leq k\delta \|\phi_1 - \phi_2\|_{C^{k+2}}$$

which can be proved similarly as before.

The assumption (LR1) (resolvent of the unperturbed operator) follows from Corollary 32. The second part of assumption (LR2) (derivative operator) follows from Theorem 37 in a way similar to what is done in the proof of Theorem 25. Applying Theorem 14, we then get

$$\lim_{\delta \rightarrow 0} \left\| \frac{h_\delta - h_0}{\delta} - (Id - \mathcal{L}_0)^{-1} \dot{\mathcal{L}} h_0 \right\|_{C^k} = 0.$$

We can now let the formula be more explicit by finding an expression for $\dot{\mathcal{L}}$. In our case

$$S(x) = \int_{S^1} h(x, y) h_0(y) dy$$

and then

$$\dot{\mathcal{L}}(h_0) = \rho * (-h_0 S)' = \rho * (-L_{T_0}(h_0) \int_{S^1} h(x, y) h_0(y) dy)'.$$

□

Remark 38. *The convergence to equilibrium assumption (71) required in Proposition 35 and Theorem 36 for the uncoupled transfer operator \mathcal{L}_0 is easy to be verified in many examples of systems whose deterministic part has some kind of topological mixing and the noise is distributed by a smooth kernel or it has large support in some sense, see [25, Corollary 5.7.1], [21, Lemma 41] or [1, Remarks 2.3 and 6.4]. In more complicated situations it can be also verified by computer aided estimates ([19]).*

Remark 39. *We remark that another meaningful definition for the transfer operator associated to family of random maps coupled in the mean field could be the following*

$$(80) \quad \mathcal{L}_\delta(\phi) = [Q_{\delta, \phi}(\rho * L_{T_0}(\phi))].$$

*In this case one applies the coupling directly to the annealed transfer operator of the random maps. Here for small δ the application of our theory seems to be possible by estimates similar to the ones shown in this section and in Section 6. Indeed the transfer operator realizing the coupling $Q_{\delta, \phi}$ is applied after the convolution. Considering $\phi \in L^1$, by (69) we get that $(\rho * L_{T_0}(\phi))$ is regularized to the regularity of the kernel ρ . If h is smooth enough and δ small enough, this regularity is preserved by the application of $Q_{\delta, \phi}$ leading to the verification of regularity properties like Exi1, Exi1.b, Con1 and SS1.*

*The verification of small perturbation properties like Exi2, Con2, SS3 and LR2 for the family of transfer operators associated to $L_{\delta, \phi} = Q_{\delta, \phi}(\rho * L_{T_0})$ relies on the estimation of the distance of Q_{δ, ϕ_1} from Q_{δ, ϕ_2} on a mixed norm which can be done in a way similar to the use of Lemma 19 and Lemma 22 as done in Section*

6. *The form of the derivative operator \dot{L} is probably similar to the one given at Theorem 24.*

9. SELF CONSISTENT OPERATORS NOT COMING FROM COUPLED MAP NETWORKS

In this section we consider examples of self consistent transfer operators not coming from networks of coupled maps, giving other examples of application of our general theory. The examples are inspired to some examples studied in [36] and [4], where we have a map whose slope depends on the average of a certain observable during the iterates. We add noise to simplify the functional analytic properties of the system. Let us consider again a family of random maps on $[0, 1]$ depending on a probability measure μ and on a parameter $\delta \geq 0$.

Let us consider the classical tent map $T : [0, 1] \rightarrow [0, 1]$, defined by $T(x) = \min(2x, 2 - 2x)$, the family of maps $T_{\delta, \mu} : [0, 1] \rightarrow [0, 1]$ we consider as a self consistent perturbation of the tent map are defined by

$$T_{\delta, \mu}(x) = \frac{T(x)}{1 + \delta \int x d\mu}.$$

Then adding a noise-like perturbation to the map $T_{\delta, \mu}$ we consider the process $(X_n)_{n \in \mathbb{N}}$ defined on $[0, 1]$ by

$$(81) \quad X_{n+1} = T_{\delta, \mu}(X_n) \hat{+} \Omega_n \pmod{1}$$

where $(\Omega_n)_{n \in \mathbb{N}}$ are i.i.d random variables distributed according to a kernel $\rho \in Lip(\mathbb{R})$, supported on $[-1, 1]$ with Lipschitz constant L and where $\hat{+}$ is the ‘‘boundary reflecting’’ sum, defined by $a \hat{+} b := \pi(a + b)$, and $\pi : \mathbb{R} \rightarrow [0, 1]$ is the piecewise linear map $\pi(x) = \min_{i \in \mathbb{Z}} |x - 2i|$. This is a model of a system on $[0, 1]$ with reflecting boundary conditions. When the noise sends a point outside the space $[0, 1]$ the projection π is applied to let the image of the map again in $[0, 1]$. Let us denote as L_π the transfer operator $L_\pi : L^1(\mathbb{R}) \rightarrow L^1([0, 1])$ associated to the map π . Let $b \in \mathbb{R}$ and $g \in Lip(\mathbb{R})$ consider the translation operator τ_b defined by $(\tau_b g)(y) := g(y + b)$.

The annealed transfer operator associated to the random dynamical system (81) is a Markov operator and is given by the following kernel operator (for details see [1], Section 6):

$$(82) \quad L_{\delta, \mu} f(x) = \int k_{\delta, \mu}(x, y) f(y) dy,$$

where

$$(83) \quad k_{\delta, \mu}(x, y) = (L_\pi \tau_{-T_{\delta, \mu}(y)} \rho)(x)$$

and $x, y \in [0, 1]$. Since the perturbation induced on the system with additive noise by increasing the parameter δ is not coming from the composition with a diffeomorphism we cannot use the estimates from the previous sections directly. We hence take a slightly different point of view on systems with additive noise, and related basic estimates which were developed in [1].

In this case we will consider $B_w = L^2[0, 1]$. Let P_w be the set of probability densities in L^2 . The nonlinear self consistent operator we consider in this case hence is given by $\mathcal{L}_\delta : P_w \rightarrow P_w$ defined as

$$(84) \quad \mathcal{L}_\delta \mu = L_{\delta, \mu} \mu$$

for each $\mu \in P_w$. We remark that since $\rho \in Lip(\mathbb{R})$ and it is supported on $[-1, 1]$ the kernel of this operator is bounded: $k_{\delta, \mu} \in L^\infty([0, 1]^2)$. Let us recall some classical and useful facts about kernel operators.

- If $k_{\delta, \mu} \in L^\infty([0, 1]^2)$, then

$$(85) \quad \|L_{\delta, \mu} f\|_\infty \leq \|k_{\delta, \mu}\|_{L^\infty([0, 1]^2)} \|f\|_1$$

and the operator $L_{\delta, \mu} : L^1 \rightarrow L^\infty$ is bounded. Furthermore, $\|L_{\delta, \mu}\|_{L^p \rightarrow L^\infty} \leq \|k_{\delta, \mu}\|_{L^\infty([0, 1]^2)}$ for $1 \leq p \leq \infty$.

- The operator $L_{\delta, \mu} : L^2 \rightarrow L^2$ is compact and

$$(86) \quad \|L_{\delta, \mu} f\|_2 \leq \|k_{\delta, \mu}\|_{L^2([0, 1]^2)} \|f\|_2$$

(see Proposition 4.7 in II.§4 [9] or [28]).

It is also well known that these Markov operators have invariant measures in L^2 (see e.g. [1] Theorem 2.2). Since $k_{\delta, \mu} \in L^\infty([0, 1]^2)$, by (85) we also have that any invariant probability measure $f_{\delta, \mu}$ for this operator satisfies

$$(87) \quad \|f_{\delta, \mu}\|_\infty \leq \|k_{\delta, \mu}\|_{L^\infty([0, 1]^2)} \leq \|\rho\|_{L^\infty[0, 1]}.$$

In [1, Section 6] the following estimates are proved for such kernel operators coming from maps with additive noise and reflecting boundaries conditions (see Propositions 6.2, and 6.3):

Proposition 40. *Assume that $k_{\delta, \mu}$ arising from the system with additive noise $(T_{\delta, \mu}, \rho)$ is given by (81). Let us fix δ , suppose that the family of interval maps $\{T_\epsilon\}_{\epsilon \in [0, \bar{\epsilon}]}$ satisfies*

$$(88) \quad \begin{aligned} T_0 &= T_{\delta, \mu} \\ T_\epsilon &= T_0 + \epsilon \cdot \dot{T} + T_\epsilon, \end{aligned}$$

where $\dot{T}, t_\epsilon \in L^2$ and $\|T_\epsilon\|_2 = o(\epsilon)$. Consider the transfer operator L_0 associated to the unperturbed system $(T_{\delta, \mu}, \rho)$ and L_ϵ be the transfer operator associated to (T_ϵ, ρ) then there is $K \geq 0$ such that for $\epsilon \in [0, \bar{\epsilon}]$

$$(89) \quad \|L_0 - L_\epsilon\|_{L^2 \rightarrow L^2} \leq \epsilon K.$$

and

$$(90) \quad \lim_{\epsilon \rightarrow 0} \frac{L_\epsilon - L_0}{\epsilon} f_0 = - \int_0^1 \left(L_\pi \left(\tau_{-T_0(y)} \frac{d\rho}{dx} \right) \right) (x) \dot{T}(y) f_0(y) dy,$$

with convergence in L^2 .

The equation (89) shows that the perturbations we are interested to apply to the transfer operators associated to this kind of systems are small perturbations in the $L^2 \rightarrow L^2$ topology. We will then consider the transfer operators associated to this kind of systems as operators acting on $L^2[0, 1]$ and in this subsection we will apply our general statements with the choice $B_{ss} = B_s = B_w = L^2[0, 1]$. We now can apply Theorem 3 and prove

Proposition 41. *Let $(T_{\delta, \mu}, \rho)$ be the self consistent system described above, let \mathcal{L}_δ be its associated transfer operator as defined in (84). There is $\bar{\delta} \geq 0$ such that for each $\delta \in [0, \bar{\delta}]$ there is a unique $f_\delta \in P_w$ such that*

$$\mathcal{L}_\delta(f_\delta) = f_\delta.$$

Furthermore there is $M \geq 0$ such that for each $\delta \in [0, \bar{\delta}]$

$$\|f_\delta\|_{L^2} \leq M.$$

Before the proof of Proposition 41 we need a couple of preliminary results

Proposition 42. *There is $C \geq 0$ such that for each $\mu_1, \mu_2 \in P_w \subseteq L^2$,*

$$\begin{aligned} \|L_{\delta, \mu_2} - L_{\delta, \mu_1}\|_{L^2 \rightarrow L^2} &\leq \delta C \|\mu_1 - \mu_2\|_{L^2} \\ \|L_{0, \mu_1} - L_{\delta, \mu_1}\|_{L^2 \rightarrow L^2} &\leq \delta C \|\mu_1\|_{L^2} \end{aligned}$$

Proof. The proof follows by (86), estimating the difference of the associated kernels. Let us first consider $\|L_{\delta, \mu_2} - L_{\delta, \mu_1}\|_{L^2 \rightarrow L^2}$. We have

$$\begin{aligned} \|L_{\delta, \mu_2} - L_{\delta, \mu_1}\|_{L^2 \rightarrow L^2} &\leq \|k_{\delta, \mu_1} - k_{\delta, \mu_2}\|_{L^2([0,1]^2)} \\ &= \left(\int_{[0,1]^2} (k_{\delta, \mu_1}(x, y) - k_{\delta, \mu_2}(x, y))^2 dx dy \right)^{\frac{1}{2}}. \end{aligned}$$

To estimate this we first estimate the distance between the two deterministic parts of the dynamics. For each $y \in [0, 1]$ we get

$$\begin{aligned} |T_{\delta, \mu_2}(y) - T_{\delta, \mu_1}(y)| &\leq \left| \frac{T(y)}{1 + \delta \int x d\mu_1(x)} - \frac{T(y)}{1 + \delta \int x d\mu_2(x)} \right| \\ &\leq \left| \frac{T(y)(1 + \delta \int x d\mu_2(x)) - T(y)(1 + \delta \int x d\mu_1(x))}{(1 + \delta \int x d\mu_1(x))(1 + \delta \int x d\mu_2(x))} \right| \\ &\leq \delta \left| \int x d\mu_2(x) - \int x d\mu_1(x) \right| \\ &\leq \delta \|\mu_1 - \mu_2\|_{L^2}. \end{aligned}$$

Now let us suppose that ρ is L -Lipschitz. Since ρ is supported in $[-1, 1]$ we get that $\tau_{-T_{\delta, \mu}(y)}\rho(x)$ is supported in $[-1, 2]$ for each μ . By this $L_\pi(\tau_{-T_{\delta, \mu}(y)}\rho(x))$ is the sum of at most three non zero contributions for each $x \in [0, 1]$, hence

$$\begin{aligned} |[k_{\delta, \mu_2} - k_{\delta, \mu_1}](x, y)| &\leq |L_\pi[\tau_{-T_{\delta, \mu_2}(y)}\rho(x) - \tau_{-T_{\delta, \mu_1}(y)}\rho(x)]| \\ &\leq 3 \sup_{x \in [-1, 2], y \in [0, 1]} |[\tau_{-T_{\delta, \mu_2}(y)}\rho(x) - \tau_{-T_{\delta, \mu_1}(y)}\rho(x)]| \\ &\leq 3\delta L \|\mu_1 - \mu_2\|_{L^2} \end{aligned}$$

proving the statement. The estimate for $\|L_{0, \mu_1} - L_{\delta, \mu_1}\|_{L^2 \rightarrow L^2}$ is similar. We have

$$\begin{aligned} |T_{\delta, \mu_1}(y) - T_{0, \mu_1}(y)| &\leq \left| \frac{T(y)}{1 + \delta \int x d\mu_1(x)} - T(y) \right| \\ &\leq \left| \frac{T(y)(1 + \delta \int x d\mu_2(x)) - T(y)}{(1 + \delta \int x d\mu_1(x))} \right| \\ &\leq \delta \left| \int x d\mu_1(x) \right| \\ &\leq \delta \|\mu_1\|_{L^2} \end{aligned}$$

and the statement is obtained as before. \square

Proposition 43. *Consider a self consistent operator as defined in (84), consider $V_{L^2} := \{v \in L^2, \int v dm = 0\}$. Suppose \mathcal{L}_0 has convergence to equilibrium in the sense that there is n such that for each $v \in V_{L^2}$ it holds $\|\mathcal{L}_0^n(v)\|_{L^2} \leq \frac{1}{2}\|v\|_{L^2}$. Then there are $K, \bar{\delta} \geq 0$ such that for each $\delta \in [0, \bar{\delta})$, and probability measures μ_1 ,*

$\mu_2 \in L^2$ such that $\max(\mu_1, \mu_2) \leq \sup(\rho)$, L_{δ, μ_i} has convergence to equilibrium and hence a unique invariant probability measure, furthermore the following holds: let f_{μ_1} the unique invariant probability measure of L_{δ, μ_1} , let f_{μ_2} be the unique invariant probability measure of L_{δ, μ_2} , then

$$\|f_{\mu_1} - f_{\mu_2}\|_{L^2} \leq \delta K \|\mu_1 - \mu_2\|_{L^2}.$$

Proof. The proof is similar to part of the proof of Proposition 34. Since $\|L_0^n(v)\|_{L^2} \leq \frac{1}{2}\|v\|_{L^2}$, by (89) for δ small enough we have $\|L_{\delta, \mu_1}^n(v)\|_{L^2} \leq \frac{3}{4}\|v\|_{L^2}$ for each μ_1 with $\|\mu_1\|_w \leq M_1$ and $v \in V_{L^2}$. Then we can define the resolvent for each such operator L_{δ, μ_1} with a uniform bound on its L^2 norm.

By this, since

$$\begin{aligned} (Id - L_{\delta, \mu_2})(f_{\mu_2} - f_{\mu_1}) &= f_{\mu_2} - L_{\delta, \mu_2}f_{\mu_2} - f_{\mu_1} + L_{\delta, \mu_2}f_{\mu_1} \\ &= (L_{\delta, \mu_2} - L_{\delta, \mu_1})f_{\mu_1}. \end{aligned}$$

We have that

$$(f_{\mu_2} - f_{\mu_1}) = (Id - L_{\delta, \mu_2})^{-1}(L_{\delta, \mu_2} - L_{\delta, \mu_1})f_{\mu_1}.$$

Since $\|(Id - L_{\delta, \mu_2})^{-1}\|_{L^2 \rightarrow L^2}$ is uniformly bounded and $\|f_{\mu_1}\|_{L^2} \leq \|f_{\mu_1}\|_{L^\infty} \leq \|\rho\|_\infty$, applying Proposition 42 we get the statement. \square

Proof of Proposition 41. We apply Theorem 3 with $B_s = B_w = L^2[0, 1]$. The assumption (Exi1) is provided by (87), (Exi2) is provided by, Proposition 42, (89) and (Exi3) is provided by Proposition 43. The unique fixed point we find is in L^2 and since the kernels we consider in the construction are uniformly bounded in L^∞ the L^2 norm of the fixed point is uniformly bounded as δ varies. \square

Now we can prove the linear response formula for these self consistent operators in the small nonlinear perturbation regime.

Theorem 44 (Linear Response). *Consider the family of self consistent transfer operators $\mathcal{L}_\delta : L^2 \rightarrow L^2$ as described before. We have the following Linear Response formula*

$$(91) \quad \lim_{\delta \rightarrow 0} \frac{f_\delta - f_0}{\delta} = (Id - L_0)^{-1} \int_0^1 \left(L_\pi \left(\tau_{-T_0(y)} \frac{d\rho}{dx} \right) \right) (x) aT(y) f_0(y) dy,$$

where $a = \int t df_0(t)$ and the limit is converging in L^2 .

Proof. The proof is an application of Theorem 14 where the spaces considered in this case are $B_{ss} = B_s = B_w = L^2$. The assumption (SS1) (regularity bounds), is implied by Proposition 41. Let us remark that the unperturbed system is a noisy tent map, hence it has convergence to equilibrium (by [1, Remarks 6.4]) and the assumption (SS2) is satisfied. To verify assumption (SS3) (small perturbation) we need to verify that, considering $B_{2M} = \{x \in L^2, \|x\| \leq 2M\}$. There is $K \geq 0$ such that and $\mathcal{L}_0 - \mathcal{L}_\delta : B_{2M} \rightarrow L^2$ is $K\delta$ -Lipschitz. We have to verify that for each $\mu_1, \mu_2 \in B_{2M}$

$$(92) \quad \|(\mathcal{L}_\delta - \mathcal{L}_0)\mu_1 - (\mathcal{L}_\delta - \mathcal{L}_0)\mu_2\|_{L^2} \leq K\delta \|\mu_1 - \mu_2\|_{L^2}.$$

Recalling that by Proposition 42 we have $L_{0, \mu_1} = L_{0, \mu_2} := L_0$, we have

$$\begin{aligned} (\mathcal{L}_\delta - \mathcal{L}_0)\mu_1 - (\mathcal{L}_\delta - \mathcal{L}_0)\mu_2 &= L_{\delta, \mu_1}\mu_1 - L_{0, \mu_1}\mu_1 - L_{\delta, \mu_2}\mu_2 + L_{0, \mu_2}\mu_2 \\ &= L_{\delta, \mu_1}\mu_1 - L_{\delta, \mu_1}\mu_2 + L_{\delta, \mu_1}\mu_2 - L_{0, \mu_1}\mu_1 - L_{\delta, \mu_2}\mu_2 + L_{0, \mu_2}\mu_2 \\ &= [L_{\delta, \mu_1} - L_0](\mu_1 - \mu_2) + [L_{\delta, \mu_1} - L_{\delta, \mu_2}]\mu_2 \end{aligned}$$

Now by Proposition 42

$$\| [L_{\delta, \mu_1} - L_0](\mu_1 - \mu_2) \|_{L^2} \leq \delta C \|\mu_1\|_{L^2} \|\mu_1 - \mu_2\|_{L^2}$$

and

$$\| [L_{\delta, \mu_1} - L_{\delta, \mu_2}]\mu_2 \|_{L^2} \leq \delta C \|\mu_2\|_{L^2} \|\mu_1 - \mu_2\|_{L^2}$$

proving the statement. Now we can apply Theorem 12 and deduce that $f_\delta \rightarrow f_0$ in L^2 . The assumption (LR1) on the existence of the resolvent is equivalent to (SS2) since we consider only one space L^2 and for the same reason the first part of (LR2) is equivalent to (SS3). We now only need to compute the derivative operator. When the self consistent operator is considered, as δ increases, the effect of the perturbation on the system is only on the map defining the deterministic part of the dynamics. We then use (90), from Proposition 40. We remark that this perturbation on the deterministic part of the dynamics depends on the invariant measure f_δ of the system as δ changes, however we will see that since $f_\delta \rightarrow f_0$ in L^2 this also give rise to a family of maps with additive noise of the type

$$T_\delta = T_0 + \delta \cdot \dot{T} + t_\delta$$

as in (88). Indeed let us compute \dot{T} in this case. Considering that $T_{\delta, f_\delta}(x) = \frac{T(x)}{1 + \delta \int x df_\delta}$ we get

$$\begin{aligned} \frac{T_\delta - T_0}{\delta} &= \frac{1}{\delta} \left[\frac{T(x)}{1 + \delta \int x df_\delta} - T(x) \right] \\ &= T(x) \frac{-\delta \int x df_\delta}{\delta + \delta^2 \int x df_\delta} \\ &= -T(x) \frac{\int x df_\delta}{1 + \delta \int x df_\delta}. \end{aligned}$$

Since $f_\delta \rightarrow f_0$ in L^2 hence

$$\dot{T} = \lim_{\delta \rightarrow 0} \frac{T_\delta - T_0}{\delta} = -T(x) \int x df_0$$

and we have the expression for $\dot{L}f_0$ from (90). Applying Theorem 14 then, we then get

$$\lim_{\delta \rightarrow 0} \frac{f_\delta - f_0}{\delta} = (Id - L_0)^{-1} \int_0^1 \left(L_\pi \left(\tau_{-T_0(y)} \frac{d\rho}{dx} \right) \right) (x) aT(y) f_0(y) dy$$

where $a = \int t df_0(t)$. □

10. COUPLING DIFFERENT MAPS

In this section we show how one can use a self consistent transfer operator approach as a model for the behavior of networks of coupled maps of different types. We will see that our general theoretical framework naturally includes this case. For simplicity we will consider only two types of maps, also for simplicity we will consider coupled expanding maps on the circle. Let us consider two different C^6 expanding maps of the circle $(T_1, S^1), (T_2, S^1)$. Given two probability densities $\psi_1, \psi_2 \in L^1(\mathbb{S}^1, \mathbb{R})$ representing the distribution of probability of the states in the two systems, two coupling functions $h_1, h_2 \in C^6(\mathbb{S}^1 \times \mathbb{S}^1, \mathbb{R})$ and $\delta \in [0, \epsilon_0]$ representing the way in which these distributions perturb the dynamics (which can be

different for the two different systems), let $\pi : \mathbb{S}^1 \rightarrow \mathbb{R}$ be the natural, universal covering projection, let us define $\Phi_{\delta, \psi_1, \psi_2} : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ with $i \in \{1, 2\}$ as

$$\Phi_{\delta, \psi_1, \psi_2}(x) = x + \pi\left(\delta \int_{\mathbb{S}^1} h_1(x, y)\psi_1(y)dy + \delta \int_{\mathbb{S}^1} h_2(x, y)\psi_2(y)dy\right)$$

(here for simplicity we suppose that the diffeomorphism perturbing the two different maps is the same though with different contributions for the two different maps, but one can consider different ways to define Φ_{δ, ψ_i} for each map) the maps will hence be perturbed by the combined action of the two densities ψ_1 and ψ_2 .

Again we assume ϵ_0 is so small that $\Phi_{\delta, \psi_1, \psi_2}$ is a diffeomorphism for each $\delta \in [0, \epsilon_0]$ and $\Phi'_{\delta, \psi_1, \psi_2} > 0$. Denote by $Q_{\delta, \psi_1, \psi_2}$ the transfer operator associated to $\Phi_{\delta, \psi_1, \psi_2}$, defined as

$$[Q_{\delta, \psi_1, \psi_2}(\phi)](x) = \frac{\phi(\Phi_{\delta, \psi_1, \psi_2}^{-1}(x))}{|\Phi'_{\delta, \psi_1, \psi_2}(\Phi_{\delta, \psi_1, \psi_2}^{-1}(x))|}$$

for any $\phi \in L^1(\mathbb{S}^1, \mathbb{R})$. Now we consider the action of the two maps by considering a global system $(S^1 \times S^1, F_{\delta, \psi_1, \psi_2})$ with

$$F_{\delta, \psi_1, \psi_2}(x_1, x_2) = (\Phi_{\delta, \psi_1, \psi_2} \circ T_1(x_1), \Phi_{\delta, \psi_1, \psi_2} \circ T_2(x_2)).$$

Finally let us consider the space of functions $B_1 := \{(f_1, f_2) \in L^1(\mathbb{S}^1) \times L^1(\mathbb{S}^1)\}$ with the norm $\|(f_1, f_2)\|_{B_1} = \|f_1\|_{L^1} + \|f_2\|_{L^1}$ (this is also called the direct sum $L^1(\mathbb{S}^1) \oplus L^1(\mathbb{S}^1)$), the space P_1 of probability elements in B_1 , $P_1 = \{(f_1, f_2) \in B_1 \text{ s.t. } \forall i \in \{1, 2\}, f_i \geq 0, \int f_i dm = 1\}$ and the stronger spaces $B_2 := \{(f_1, f_2) \in W^{1,1}(\mathbb{S}^1) \times W^{1,1}(\mathbb{S}^1)\}$ with the norm $\|(f_1, f_2)\|_{B_2} = \|f_1\|_{W^{1,1}} + \|f_2\|_{W^{1,1}}$ and $B_3 := \{(f_1, f_2) \in W^{2,1}(\mathbb{S}^1) \times W^{2,1}(\mathbb{S}^1)\}$ with the norm $\|(f_1, f_2)\|_{B_3} = \|f_1\|_{W^{2,1}} + \|f_2\|_{W^{2,1}}$ (again direct sums of Sobolev spaces). These sets can be trivially endowed with a structure of normed vector space. Coherently with the previous sections we define a family of transfer operators $L_{\delta, \phi_1, \phi_2} : B_w \rightarrow B_w$ depending on elements of the weaker space $(\phi_1, \phi_2) \in P_1$ as

$$(93) \quad L_{\delta, \phi_1, \phi_2}((f_1, f_2)) = (Q_{\delta, \phi_1, \phi_2}(L_{T_1}(f_1)), Q_{\delta, \phi_1, \phi_2}(L_{T_2}(f_2))).$$

By this we can define the self consistent transfer operator $\mathcal{L}_\delta : B_w \rightarrow B_w$ associated to this system as

$$(94) \quad \mathcal{L}_\delta((f_1, f_2)) = L_{\delta, f_1, f_2}((f_1, f_2)).$$

We remark that B_1 can be identified with a closed subset of $L^1(\mathbb{S}^1 \times \mathbb{S}^1)$ by $(f_1, f_2) \rightarrow f$ where f is defined by $f(x, y) = f_1(x)f_2(y)$ and \mathcal{L}_δ preserves this subspace.

We now prove the existence and uniqueness of the invariant measure for this kind of self consistent operators for small δ , applying our general statement, Theorem 3.

Theorem 45. *Let T_1, T_2 be two C^6 expanding maps and let $h_1, h_2 \in C^6(\mathbb{S}^1 \times \mathbb{S}^1, \mathbb{R})$. Let us consider a globally coupled system as defined above. There is some $\bar{\delta}$ such that for each $\delta \in [0, \bar{\delta}]$ there is a unique $(f_{1,\delta}, f_{2,\delta}) \in B_2$ such that*

$$\mathcal{L}_\delta((f_{1,\delta}, f_{2,\delta})) = (f_{1,\delta}, f_{2,\delta}).$$

Furthermore there is $M \geq 0$ such that for each $\delta \in [0, \bar{\delta}]$

$$\|(f_{1,\delta}, f_{2,\delta})\|_{B_2} \leq M.$$

Proof. The proof follows by the application of Theorem 3 with $B_w = B_1$ and $B_s = B_2$. We verify the needed assumptions; the assumption (*Exi1*) is trivial, indeed given $\mu = (\phi_1, \phi_2) \in B_1$ for any δ small enough the invariant measure of $L_{\delta, \phi_1, \phi_2}$, which is a system which is the product of two expanding maps, is trivially in B_2 and if we let (ϕ_1, ϕ_2) range in P_1 , the B_2 norm of the associated invariant measure is uniformly bounded.

The assumption (*Exi2*) can be easily deduced by Proposition 19 as done in the case of system made by coupling identical maps obtaining that there is some $K_1 \geq 0$ such that

$$\|L_{\delta, \phi_1, \phi_2} - L_{\delta, \phi_3, \phi_4}\|_{B_2 \rightarrow B_1} \leq \delta K_1 \|(\phi_1, \phi_2) - (\phi_3, \phi_4)\|_{B_1}$$

for δ ranging in some neighborhood of the origin and $(\phi_1, \phi_2), (\phi_3, \phi_4) \in P_1$. Again, applying Lemma 21, like done in Proposition 20 we verify (*Exi3*) for this product system. Then Theorem 3 can be applied, giving the statement. \square

It seems that it is possible to extend all the results we proved for coupled expanding maps to this kind of systems, with the same ideas and estimates (but longer formulas and computations, as we have two coordinates). This work however would be quite long and outside of the scope of this paper.

11. THE OPTIMAL COUPLING

In this section we study the problem of finding an optimal small coupling functions $\delta \hat{h}$ in order to maximize the average of a given observable. This is an optimal control problem in which the goal is to change the statistical properties of the system in an certain direction, in some optimal way. In this case we consider an initial uncoupled system and introduce a small perturbation by a coupling function $\delta \hat{h}$ and we look for the response of the system to this small perturbation like in Theorem 14. We suppose the direction of perturbation \hat{h} can vary in some (infinite dimensional) set P , and in this set we look for an optimal one. In the context of extended systems this kind of problems were also defined as "management of the statistical properties of the complex system" ([42]). In some sense this is an inverse problem related to the linear response, in which the goal is to find the optimal perturbation giving a certain kind of response. Related problems in which the focus is more on the realization of a given fixed response have also been called "linear request" problems (see [16] and [26]).

The problem of finding an optimal infinitesimal perturbation, in order to maximize the average of a given observable and other statistical properties of dynamics was investigated in the case of finite Markov chains in [2] and for a class of random dynamical systems whose transfer operators are Hilbert Schmidt operators in [1].

In this section we start the investigation of these kind of problems in the case of self consistent transfer operators. We obtain existence and uniqueness of the optimal solution under assumptions similar to the ones used in [1]. We will focus on the question of finding the best coupling in order to optimize the behavior of a given observable. Let us explain more precisely but still a bit informally the kind of problem we are going to consider: given a certain system, we consider a set P of allowed infinitesimal perturbations we can put in the system. It is natural to think of the set of allowed perturbations P as a convex set because if two different perturbations of the system are possible, then their convex combination (applying the two perturbations with different intensities) should also be possible.

We will also consider P as a subset of some Hilbert space \mathcal{H} (as it is useful for optimization purposes). Let $\mu_{\dot{h},\delta}$ be the invariant measure of the system after applying a perturbation in the direction $\dot{h} \in P$ with intensity δ (we will formalize later what we mean by direction and intensity in our case). Let the response to this perturbation be denoted as

$$R(\dot{h}) = \lim_{\delta \rightarrow 0} \frac{\mu_{\dot{h},\delta} - \mu_0}{\delta}.$$

Let us consider $c : [0, 1] \rightarrow \mathbb{R}$. We are interested in the rate of increasing of the expectation of c

$$\left. \frac{d(\int c d\mu_{\dot{h},\delta})}{d\delta} \right|_{\delta=0}$$

and the element $\dot{h} \in P$ for which this is maximized, thus we are interested in finding \dot{h}_{opt} such that

$$(95) \quad \left. \frac{d(\int c d\mu_{\dot{h}_{opt},\delta})}{d\delta} \right|_{\delta=0} = \max_{\dot{h} \in P} \left. \frac{d(\int c d\mu_{\dot{h},\delta})}{d\delta} \right|_{\delta=0}.$$

By (39) and (40), under the suitable assumptions, this turns out to be equivalent to finding \dot{h}_{opt} such that

$$(96) \quad \int c dR(\dot{h}_{opt}) = \max_{\dot{h} \in P} \int c dR(\dot{h}).$$

This is hence the maximization of a certain linear function on the set P .

11.1. Some reminders on optimization of a linear function on a convex set . The optimal perturbation problem we mean to consider is related to the maximization of a continuous linear function on the set of allowed infinitesimal perturbations P . The existence and uniqueness of an optimal perturbation hence depends on the properties of the convex bounded set P . We now recall some general results, adapted for our purposes, on optimizing a linear continuous function on convex sets. Let $\mathcal{J} : \mathcal{H} \rightarrow \mathbb{R}$ be a continuous linear function, where \mathcal{H} is a separable Hilbert space and $P \subset \mathcal{H}$.

The abstract problem we consider then is to find $\dot{h}_{opt} \in P$ such that

$$(97) \quad \mathcal{J}(\dot{h}_{opt}) = \max_{\dot{h} \in P} \mathcal{J}(\dot{h}).$$

The following propositions summarizes some efficient criteria for the existence and uniqueness of the solution of such problem (see [1], Section 4 for more details and the proofs).

Proposition 46 (Existence of the optimal solution). *Let P be bounded, convex, and closed in \mathcal{H} . Then, problem (97) has at least one solution.*

Uniqueness of the optimal solution will be provided by strict convexity of P .

Definition 47. *We say that a convex closed set $A \subseteq \mathcal{H}$ is strictly convex if for each pair $x, y \in A$ and for all $0 < \gamma < 1$, the points $\gamma x + (1 - \gamma)y \in \text{int}(A)$, where the relative interior⁷ is meant.*

⁷The relative interior of a closed convex set C is the interior of C relative to the closed affine hull of C .

Proposition 48 (Uniqueness of the optimal solution). *Suppose P is closed, bounded, and strictly convex subset of \mathcal{H} , and that P contains the zero vector in its relative interior. If \mathcal{J} is not uniformly vanishing on P then the optimal solution to (97) is unique.*

We remark that in the case \mathcal{J} is uniformly vanishing, all the elements of P are solutions of the problem (96).

11.2. Optimizing the response of the expectation of an observable. Let $c \in L^1$ be a given observable. We consider the problem of finding an infinitesimal perturbation that maximizes the expectation of c . As motivated before, we want to solve the problem stated in (96). Suppose that P is a closed, bounded, convex subset of \mathcal{H} containing the zero perturbation, and that \mathcal{J} is not uniformly vanishing on P . Let us consider the function $\mathcal{J}(\dot{h}) = \int c dR(\dot{h})$. When this function is continuous as a map from $(P, \|\cdot\|_{\mathcal{H}})$ to \mathbb{R} , we may immediately apply Proposition 46 to obtain that there exists a solution to the problem considered in (96). If, in addition, P is strictly convex and \mathcal{J} is nonvanishing, then by Proposition 48 the solution to (96) is unique.

In the following subsections we hence apply these remarks to find the existence and uniqueness of the optimal coupling in the case of coupled expanding map and maps with additive noise.

11.2.1. The optimal coupling for expanding maps. We consider self consistent transfer operators coming from a system of coupled maps as in Section 6, where \mathcal{L}_0 is the uncoupled operator and \mathcal{L}_δ is the self consistent operator with coupling driven by a function $\dot{h} : \mathbb{S}^1 \times \mathbb{S}^1 \rightarrow \mathbb{R}$ and with strength δ . We proved in 25 (see (56)) that the response of the invariant measure of the system as δ increases is given by

$$R(\dot{h}) = (Id - L_0)^{-1} \left(h_0 \int_{S^1} \dot{h}(x, y) h_0(y) dy \right)'.$$

Given some observable $c \in L^1$ and some convex set of allowed perturbations P we now apply the previous results to the problem of finding the optimal coupling $\dot{h}_{opt} \in P$ solving the problem (96) for this response function $R(\dot{h})$. From Remark 26 (see also (40)) we know that the rate of change of the average of c can be estimated by the linear response when the convergence of the linear response is in $W^{1,1}$, as provided in Theorem 14.

We remark that to apply the general results of Section 11.1 we need P being a subset of a Hilbert space. Since to apply Theorem 25 we need $\dot{h} \in C^6$ we consider a Hilbert space of perturbations which is included in C^6 . A simple choice is $W^{7,2}$. We hence consider a system with coupled expanding maps, the Hilbert space $W^{7,2}$ and a convex set $P \subseteq W^{7,2}(S^1 \times S^1)$.

Proposition 49. *Under the above assumptions, supposing that P is a closed bounded convex set in $W^{7,2}$, Problem (96) has a solution in P . If furthermore P is strictly convex either the optimal solution is unique or every $\dot{h} \in P$ is the optimal solution.*

Proof. The result directly follows applying Propositions 46 and 48. In order to apply the propositions we have to check that $\dot{h} \rightarrow \int c dR(\dot{h})$ is continuous on P . Since

$$R(\dot{h}) = (Id - L_0)^{-1} \left(h_0 \int_{S^1} \dot{h}(x, y) h_0(y) dy \right)'$$

we have

$$\begin{aligned}
\int c \, dR(\dot{h}) &\leq \|c\|_1 \|(Id - L_0)^{-1} (h_0 \int_{S^1} \dot{h}(x, y) h_0(y) dy)'\|_\infty \\
&\leq \|c\|_1 \|(Id - L_0)^{-1}\|_{W^{1,1} \rightarrow W^{1,1}} \|(h_0 \int_{S^1} \dot{h}(x, y) h_0(y) dy)'\|_{W^{1,1}} \\
&\leq \|c\|_1 \|(Id - L_0)^{-1}\|_{W^{1,1} \rightarrow W^{1,1}} \|h_0\|_{C^3}^2 \|\dot{h}\|_{W^{2,1}}.
\end{aligned}$$

Now the result follow by a direct application of Propositions 46 and 48. \square

11.2.2. *The optimal coupling for systems with additive noise.* Now we consider the optimal coupling in order to maximize the average of one observable c in the case of the coupled maps with additive noise as described in Section 8. Since Theorem 36 gives a convergence of the linear response in the strong space C^k , by Remark 26 we know that we can consider very general observables. For simplicity we will consider $c \in L^1$ but in fact we could consider even weaker spaces as distribution spaces (the dual of C^k). For simplicity we also take $P \subseteq W^{1,2}$ to let (77) make sense. The response formula in this case is

$$R(\dot{h}) = (Id - L_0)^{-1} \rho_\xi * (L_{T_0}(h_0) \int_{S^1} \dot{h}(x, y) h_0(y) dy)'$$

We will hence consider the problem (96) with this response function. Similarly to the expanding maps case we get the following statement.

Proposition 50. *Under the above assumptions, supposing that P is a closed bounded convex set in $W^{1,2}$, Problem (96) has a solution in P . If furthermore P is strictly convex either the solution is unique or every $\dot{h} \in P$ is the optimal solution.*

Proof. The result again directly follows applying Propositions 46 and 48. In order to apply the propositions we check that $\dot{h} \rightarrow \int c \, dR(\dot{h})$ is continuous on P . Since in this case

$$R(\dot{h}) = (Id - L_0)^{-1} \rho_\xi * (h_0 \int_{S^1} \dot{h}(x, y) h_0(y) dy)'$$

we have

$$\begin{aligned}
\int c \, dR(\dot{h}) &\leq \|c\|_1 \|(Id - L_0)^{-1} \rho_\xi * (h_0 \int_{S^1} \dot{h}(x, y) h_0(y) dy)'\|_\infty \\
&\leq \|c\|_1 \|h_0\|_\infty \|\dot{h}\|_{W^{2,1}}
\end{aligned}$$

establishing the continuity of $\dot{h} \rightarrow \int c \, dR(\dot{h})$. \square

Acknowledgments. S.G. is partially supported by the research project PRIN 2017S35EHN_004 "Regular and stochastic behavior in dynamical systems" of the Italian Ministry of Education and Research. The author wish to thank F. Sélley, S. Vaienti, M. Tanzi and C. Liverani for fruitful discussions during the preparation of the work. The author wish also to thank the anonymous reviewers for their patience and the help in revising this long text.

REFERENCES

- [1] F. Antown, G. Froyland, S. Galatolo *Optimal linear response for Markov Hilbert-Schmidt integral operators and stochastic dynamical systems* arXiv:2101.09411
- [2] F. Antown, D. Dragicevic, and G. Froyland. *Optimal linear responses for Markov chains and stochastically perturbed dynamical systems*. J. Stat. Phys., 170(6): 1051–1087, (2018).
- [3] V. Baladi, Linear response, or else, *ICM proceedings Seoul 2014*, vol III, pp525-545
- [4] M. Blank. Ergodic averaging with and without invariant measures. Nonlinearity, 30(12):4649, (2017).
- [5] M. L. Blank. *Self-consistent mappings and systems of interacting particles*. Doklady Math., 83, pp 49–52. Springer, 2011.
- [6] Balint, P., Keller, G., Selley, F. M., & Toth, I. P. *Synchronization versus stability of the invariant distribution for a class of globally coupled maps*. Nonlinearity, 31(8), 3770. (2018)
- [7] Jean-René Chazottes and Bastien Fernandez. *Dynamics of coupled map lattices and of related spatially extended systems*, volume 671. Springer Science & Business Media, (2005).
- [8] T. Chihara, Y. Sato, I Nisoli, S Galatolo *Existence of multiple noise-induced transitions in a Lasota-Mackey map* arXiv:2102.11715 (2021)
- [9] J. Conway. *A course in functional analysis*, volume 96. Springer Science & Business Media, 2013
- [10] J.P. Conze, A Raugi, *Limit theorems for sequential expanding dynamical systems* on , Ergodic theory and related fields, Contemp. Math., vol. 430, Amer. Math. Soc., Providence, RI, 2007, pp. 89–121.
- [11] Dietert, H., & Fernandez, B. (2018). *The mathematics of asymptotic stability in the Kuramoto model*. Proceedings of the Royal Society A, 474(2220), 20180467.
- [12] Fernandez, B. *Breaking of ergodicity in expanding systems of globally coupled piecewise a ne circle maps*. Journal of Statistical Physics 154, 4 , 999-1029 (2014).
- [13] M. Florenzano *General Equilibrium Analysis: Existence and Optimality Properties of Equilibria* Springer, ISBN 9781402075124, (2003)
- [14] S. Galatolo, J. Sedro *Quadratic response of random and deterministic dynamical systems* Chaos 30, 023113 (2020); <https://doi.org/10.1063/1.5122658>
- [15] S. Galatolo *Statistical properties of dynamics. Introduction to the functional analytic approach* arXiv:1510.02615
- [16] S. Galatolo, M. Pollicott *Controlling the statistical properties of expanding maps*. Nonlinearity 30, 7 (2017), 2737.
- [17] S. Galatolo, I. Nisoli, B. Saussol. *An elementary way to rigorously estimate convergence to equilibrium and escape rates*. J. Comput. Dyn., 2015, 2 (1) : 51-64. doi: 10.3934/jcd.2015.2.51
- [18] S. Galatolo, I. Nisoli *An Elementary Approach to Rigorous Approximation of Invariant Measures* SIAM J. Appl. Dyn. Syst. 13-2 (2014), pp. 958-985
- [19] S Galatolo, M Monge, I Nisoli *Existence of Noise Induced Order, a Computer Aided Proof* Nonlinearity, 33(9):4237–4276, (2020).
- [20] S. Galatolo, M. Monge, I. Nisoli *Rigorous approximation of stationary measures and convergence to equilibrium for iterated function systems* J. Phys. A: Math. Theor. 49 (2016) 274001
- [21] S Galatolo and P Giulietti *A linear response for dynamical systems with additive noise* Nonlinearity 32 2269 (2019)
- [22] Golse F. *On the dynamics of large particle systems in the mean-field limit*. In Macroscopic and large scale phenomena: coarse graining, mean field limits and ergodicity (eds A Muntean, J Rademacher, A Zagari). Lecture Notes Application Mathematics and Mechanics, vol. 3, pp. 1–144. Springer (2016).
- [23] C. Liverani, *Invariant measures and their properties. A functional analytic point of view*, Dynamical Systems. Part II: Topological Geometrical and Ergodic Properties of Dynamics. Centro di Ricerca Matematica “Ennio De Giorgi”: Proceedings. Published by the Scuola Normale Superiore in Pisa (2004).
- [24] C. Liverani *Rigorous numerical investigation of the statistical properties of piecewise expanding maps. A feasibility study*. Nonlinearity 14 463–490 (2001)
- [25] A. Lasota and M. Mackey. *Probabilistic properties of deterministic systems*. Cambridge university press, 1985

- [26] Kloeckner, B. *The linear request problem*. Proceedings of the American Mathematical Society 146, 7 (2018), 2953-2962.
- [27] K. Kaneko. *Globally coupled chaos violates the law of large numbers but not the central limit theorem*. Physical review letters, 65(12):1391 (1990).
- [28] A. Kolmogorov and S. Fomin. *Elements of the Theory of Functions and Functional Analysis*. Volume 2: Measure. The Lebesgue Integral. Hilbert Space. Graylock, 1961.
- [29] G. Keller. *An ergodic theoretic approach to mean field coupled maps*. In Fractal Geometry and Stochastics II, pages 183–208. Springer, 2000.
- [30] G. Keller ; C. Liverani *Stability of the spectrum for transfer operators* Annali della Scuola Normale Superiore di Pisa, 4, 28 (1999) no. 1, pp. 141-152.
- [31] Marangio, L.; Sedro, J.; Galatolo, S.; Di Garbo, A.; Ghil, M. *Arnold maps with noise: differentiability and non-monotonicity of the rotation number*. J. Stat. Phys. 179, 1594–1624, (2020)
- [32] W. Ott, M. Stenlund, L-S Young. *Memory loss for time-dependent dynamical systems*. Mathematical research letters, 16(2):463–475, 2009.
- [33] J. Sedro *A regularity result for fixed points, with applications to linear response*. Nonlinearity 31 1417 (2018).
- [34] F. M. Sélley, M. Tanzi *Linear Response for a Family of Self-Consistent Transfer Operators* Comm. Math. Phys. 382(3), 1601-1624 (2021).
- [35] F. M. Sélley. *Asymptotic properties of mean field coupled maps*. PhD thesis, (2019).
- [36] F. M. Sélley *A self-consistent dynamical system with multiple absolutely continuous invariant measures* Journal of Computational Dynamics, 8(1), 932 (2021).
- [37] Viana, M.: *Lectures on Lyapunov Exponents*. Cambridge Studies in Advanced Mathematics 145, Cambridge University Press (2014)
- [38] Jiang, M., and de la Llave, R. *Smooth dependence of thermodynamic limits of SRB-measures*. Communications in Mathematical Physics 211, 2 (2000), 303-333.
- [39] Jiang, M., and de la Llave, R. *Linear response function for coupled hyperbolic attractors*. Communications in Mathematical Physics 261, 2 (2006), 379-404.
- [40] Wormell, C. L., and Gottwald, G. A. *On the validity of linear response theory in high-dimensional deterministic dynamical systems*. Journal of Statistical Physics 172, 6 (2018), 1479-1498.
- [41] Wormell, C. L., and Gottwald, G. A. *Linear response for macroscopic observables in high-dimensional systems*. Chaos: An Interdisciplinary Journal of Nonlinear Science 29, 11 (2019), 113127.
- [42] MacKay, R. *Management of complex dynamical systems*. Nonlinearity 31, 2 (2018), R52.
- [43] Ruelle, D., Differentiation of SRB states, *Comm. Math. Phys.* 187 (1997) 227–241.
- [44] G. Keller, C. Liverani. *A spectral gap for a one-dimensional lattice of coupled piecewise expanding interval maps*. In Dynamics of coupled map lattices and of related spatially extended systems, pp 115–151. Springer, 2005.
- [45] J.B. Bardet, G. Keller, R. Zweimüller. *Stochastically stable globally coupled maps with bistable thermodynamic limit*. Comm. Math. Phys. 292 (1), :237–270, (2009).
- [46] M. Jiang, Y. B. Pesin. *Equilibrium measures for coupled map lattices: Existence, uniqueness and finite-dimensional approximations*. Comm. Math. Phys. 193(3):675–711 (1998).

Email address: ¹ stefano.galatolo@unipi.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITA DI PISA, VIA BUONARROTI 1, PISA - ITALY