

# A note on induced Turán numbers

Freddie Illingworth\*

## Abstract

Loh, Tait, Timmons and Zhou introduced the notion of induced Turán numbers, defining  $\text{ex}(n, \{H, F\text{-ind}\})$  to be the greatest number of edges in an  $n$ -vertex graph with no copy of  $H$  and no induced copy of  $F$ . Their and subsequent work has focussed on  $F$  being a complete bipartite graph. In this short note, we complement this focus by asymptotically determining the induced Turán number whenever  $H$  is not bipartite and  $F$  is not an independent set nor a complete bipartite graph.

## 1 Introduction

Consider the following induced version of the classical Turán problem. What is the most edges in an  $n$ -vertex graph which does not contain  $F$  as an induced subgraph? This question has a simple answer: if  $F$  is a clique, then it is the original subgraph problem (and so answered by Turán's theorem [8]), and if  $F$  is not complete, then the maximum is plainly  $\binom{n}{2}$ , as witnessed by the  $n$ -vertex complete graph  $K_n$ .

In an insightful paper, Loh, Tait, Timmons and Zhou [5] recovered an interesting and natural induced Turán problem by forbidding both an induced  $F$  and a (not necessarily induced) copy of  $H$ . This removes the possibility of  $K_n$  being extremal and so rules out the humdrum answer of  $\binom{n}{2}$ . To be precise, for a positive integer  $n$  and graphs  $H$  and  $F$ , they defined *the induced Turán number*

$$\text{ex}(n, \{H, F\text{-ind}\}),$$

to be the maximum number of edges in an  $n$ -vertex graph with no copy of  $H$  and no induced copy of  $F$ .

In [5], Loh, Tait, Timmons and Zhou focussed on complete bipartite  $F$  proving general bounds for  $\text{ex}(n, \{H, K_{s,t}\text{-ind}\})$  with some sharper results when  $s = 2$ . Nikiforov, Tait and Timmons [6] proved spectral improvements of their results. Ergemlidze, Györi and Methuku [3] asymptotically determined  $\text{ex}(n, \{H, F\text{-ind}\})$  in the important case where  $H$  is an odd cycle and  $F$  is  $K_{2,t}$  or  $K_{3,3}$  (except when  $H = C_5$  and  $F = K_{2,2}$  where they strengthened the upper bound). In [4], the author gave further improvements to the upper bounds for  $\text{ex}(n, \{H, K_{2,t}\text{-ind}\})$ .

All the work to date has focussed on complete bipartite  $F$ . The following theorem complements this focus.

**Theorem 1.** *Let  $H$  and  $F$  be graphs with chromatic numbers  $r + 1$  and  $s + 1$  respectively. Then*

$$\text{ex}(n, \{H, F\text{-ind}\}) = \begin{cases} (1 - \frac{1}{r} + o(1)) \binom{n}{2} & \text{if } s \geq r \text{ or } F \text{ is not complete multipartite,} \\ (1 - \frac{1}{s} + o(1)) \binom{n}{2} & \text{if } s < r \text{ and } F \text{ is complete multipartite.} \end{cases}$$

This asymptotically determines the induced Turán number except if  $H$  is bipartite or  $F$  is an independent set or a complete bipartite graph.

---

\*DPMMS, University of Cambridge, UK. *E-mail:* fci21@cam.ac.uk. Research supported by an EPSRC grant. 2020 MSC: 05C35.

## 2 Proof of Theorem 1

We will use  $T_r(n)$  to denote the  $r$ -partite Turán graph on  $n$  vertices. This is complete  $r$ -partite with parts as equal in size as possible. We remind the reader that  $T_r(n)$  has  $(1 - 1/r + o(1))\binom{n}{2}$  edges.

*Proof of Theorem 1.* If  $s \geq r$  or  $F$  is not complete multipartite, then  $F$  is not an induced subgraph of the Turán graph  $T_r(n)$ . This also does not contain  $H$ , so

$$\text{ex}(n, \{H, F\text{-ind}\}) \geq e(T_r(n)) = \left(1 - \frac{1}{r} + o(1)\right)\binom{n}{2}.$$

On the other hand, the Erdős-Simonovits theorem [1] says that any  $n$ -vertex graph not containing  $H$  has at most  $(1 - 1/r + o(1))\binom{n}{2}$  edges. This completes the first part of the theorem.

Now suppose that  $s < r$  and  $F$  is complete  $(s + 1)$ -partite. Firstly the Turán graph  $T_s(n)$  contains neither  $H$  nor  $F$  as subgraphs (let alone induced ones) so,

$$\text{ex}(n, \{H, F\text{-ind}\}) \geq e(T_s(n)) = \left(1 - \frac{1}{s} + o(1)\right)\binom{n}{2}.$$

As  $F$  is complete  $(s + 1)$ -partite, there is a positive integer  $t$  such that  $F$  is an induced subgraph of  $K_{s+1}(t) = T_{s+1}(t(s + 1))$ . By Ramsey's theorem [7], there is a positive integer  $m$  such that any  $m$ -vertex graph contains either an independent  $t$ -set or a copy of  $H$ .

Fix  $\varepsilon > 0$  and let  $G$  be an  $n$ -vertex graph with at least  $(1 - 1/s + \varepsilon)\binom{n}{2}$  edges where  $n$  is sufficiently large. By the Erdős-Stone theorem [2],  $G$  contains a copy of  $K_{s+1}(m)$ . Let the parts of the  $K_{s+1}(m)$  be  $V_1, \dots, V_{s+1}$  so each one has  $m$  vertices. If any  $V_i$  contains  $H$ , then  $G$  does. Otherwise, by the definition of  $m$ , each  $V_i$  contains an independent set of size  $t$ . Thus  $G$  contains an induced copy of  $K_{s+1}(t)$  and so an induced copy of  $F$ . Thus, for all large  $n$ ,

$$\text{ex}(n, \{H, F\text{-ind}\}) \leq \left(1 - \frac{1}{s} + \varepsilon\right)\binom{n}{2}.$$

This holds for arbitrary  $\varepsilon > 0$ , as required. □

## References

- [1] P. Erdős and M. Simonovits. A limit theorem in graph theory. *Studia Scientiarum Mathematicarum Hungarica*, 1:51–57, 1966.
- [2] P. Erdős and A. H. Stone. On the structure of linear graphs. *Bulletin of American Mathematical Society*, 52:1087–1091, 1946.
- [3] B. Ergemlidze, E. Győri, and A. Methuku. Turán number of an induced complete bipartite graph plus an odd cycle. *Combinatorics, Probability and Computing*, 28(2):241–252, March 2019.
- [4] F. Illingworth. Graphs with no induced  $K_{2,t}$ . *Electronic Journal of Combinatorics*, 28(1.19), January 2021.
- [5] P.-S. Loh, M. Tait, C. Timmons, and R. M. Zhou. Induced Turán numbers. *Combinatorics, Probability and Computing*, 27(2):274–288, March 2018.
- [6] V. Nikiforov, M. Tait, and C. Timmons. Degenerate Turán problems for hereditary properties. *Electronic Journal of Combinatorics*, 25(4.39), November 2018.
- [7] F. P. Ramsey. On a problem of formal logic. *Proceedings of the London Mathematical Society*, s2-30(1):264–286, 1930.
- [8] P. Turán. Eine Extremalaufgabe aus der Graphentheorie. *Matematikai és Fizikai Lapok*, 48:436–452, 1941.