

Nonsingular Morse-Smale flows of n -manifolds with attractor-repeller dynamics

O. V. Pochinka, D. D. Shubin

May 28, 2021

Abstract

In the present paper the exhaustive topological classification of nonsingular Morse-Smale flows of n -manifolds with two limit cycles. Hyperbolicity of periodic orbits implies that among them one is attracting and another is repelling. Due to Poincaré–Hopf theorem Euler characteristic of closed manifold M^n which admits the considered flows is equal to zero. Only torus and Klein bottle can be ambient manifolds for such flows in case of $n = 2$. Authors established that there exist exactly two classes of topological equivalence of such flows of torus and three of the Klein bottle. There are no constraints for odd-dimensional manifolds which follow from the fact that Euler characteristic is zero. However, it is known that orientable 3-manifold admits a flow of considered class if and only if it is a lens space. In this paper, it is proved that up to topological equivalence each of \mathbb{S}^3 and $\mathbb{R}P^3$ admit one such flow and other lens spaces two flows each. Also, it is shown that the only non-orientable n -manifold (for $n > 2$), which admits considered flows is the twisted I -bundle over $(n-1)$ -sphere. Moreover, there are exactly two classes of topological equivalence of such flows. Among orientable n -manifolds only the product of $(n-1)$ -sphere and the circle can be ambient manifold of a considered flow and the flows are split into two classes of topological equivalence.

1 Introduction and statement of results

This article will focus on so-called *nonsingular Morse-Smale flows* (abbreviated as *NMS-flows*) which are Morse-Smale flows without fixed points. There is close connection between such flows and topology of ambient manifold. Authors obtained the exhaustive topological classification of NMS-flows with two periodic orbit.

The general theory of hyperbolic dynamical systems (see e.g. [1]) implies that ambient manifold M^n is the union of stable and unstable manifolds of this orbits. It immediately follows that on of this orbits if attracting (denote it with A) and one is repelling R .

Let $G_2(M^n)$ be the class on NMS-flows with exactly two periodic orbits. In cases where the results are fundamentally different for orientable and non-

orientable manifolds we will use notation M_+^n, M_-^n for orientable and non-orientable manifolds respectively.

Recall that periodic orbit is called *twisted* if at least one of its invariant manifolds is non-orientable. Otherwise, we call the orbit *non-twisted*. Since dimension of stable (unstable) manifold of attracting (repelling) periodic orbit is equal to ambient manifold dimension the orbits of the flow $f^t \in G_2(M_-^n)$ are both twisted, whereas orbits of the flow $f \in G_2(M_+^n)$ are both untwisted. So we specify this types further.

Poincaré–Hopf theorem implies that Euler characteristic of a NMS-flow ambient manifold equals to zero. Considering two-dimensional surfaces we immediately get that this constraint leaves us only torus and Klein bottle (actually, both admit NMS-flow). Classification of such flows follows from classification of Morse–Smale flows of surfaces (see e.g. [2], [3], [4]). We provide an independent classification of class $G_2(M^2)$ in section 4.

Theorem 1. *There are five classes of topological equivalence of NMS-flows of two-dimensional manifolds with to periodic orbits (see e.g. 1 and 2). Moreover, two of them are the flows of the torus and three of the Klein bottle (two with twisted orbits and one with non-twisted).*

The fact that the Euler characteristic of an ambient manifold equals to zero does not imply any constraints on odd-dimensional manifold, because Euler characteristic of all such manifolds is zero. Necessary and sufficient condition of topological equivalence of three-dimensional NMS-flows follows from the paper of Umansky [5], where wider class of Morse–Smale flows was considered. However, the realization was not provided in that paper, so one cannot judge whether certain manifolds admits one or the other NMS-flow. In case of a small number of periodic orbits topology of the ambient manifold and exhaustive topological classification can be established.

Recall, that a *lens space* is defined as the topological space obtained by gluing to solid tori by a homeomorphism of their boundaries and is denoted as $L_{p,q}$, $p, q \in \mathbb{Z}$, where $\langle p, q \rangle$ is homotopy type of the meridian image under the gluing homeomorphism. Some common 3-manifolds are really lens spaces, for instance three-dimensional sphere $S^3 = L_{1,0}$, manifold $S^2 \times S^1 = L_{0,1}$ and projective space $\mathbb{R}P^3 = L_{1,2}$.

It turns out that only lens spaces can be ambient manifolds of NMS-flows in case of small number of periodic orbits.

Proposition 1 ([6]). *Let M be an orientable, simple¹, compact 3-manifold without boundary. If M admits an NMS-flow with 0 or 1 saddle periodic orbit, then M is a lens space.*

Existence and uniqueness up to topological equivalence of flows of the class $G_2(S^3)$ follows from the next proposition.

¹ n -manifold is called *simple*, if it is impossible to represent it as connected sum of two n -manifolds each of which is not S^n .




Figure 1: Phase portraits of not topologically equivalent flows on the torus




Figure 2: Phase portraits of not topologically equivalent flows on the Klein bottle: 1. with untwisted orbits; 2-3. with twisted ones

Proposition 2 ([7]). *Up to topological equivalence, there exists exactly one NMS-flow f^t whose periodic orbits are composed of an attractor A and a repeller R . Moreover, the periodic orbits $A \sqcup R$ form a Hopf link in \mathbb{S}^3 .*

In the present paper exhaustive topological classification of class $G_2(M^3)$ is developed.

Theorem 2.

1. *Manifold M_+^3 admits a flow $f^t \in G_2(M_+^3)$ if and only if M_+^3 is a lens space. Moreover, there are exactly one flow of \mathbb{S}^3 (see e.g. 3) and \mathbb{RP}^3 and exactly two flows on each of the rest lens spaces (see e.g. 4).*




Figure 3: Phase portraits of equivalent flows on 3-sphere

2. The only non-orientable manifolds which admits flow of the class $G_2(M_-^3)$ is twisted I -bundle of two-sphere $\mathbb{S}^2 \tilde{\times} \mathbb{S}^1$. Both periodic orbits of such flow are twisted. Moreover, there are exactly two classes of topological equivalence in $G_2(\mathbb{S}^2 \tilde{\times} \mathbb{S}^1)$.




Figure 4: Phase portraits of inequivalent flows on $\mathbb{S}^2 \times \mathbb{S}^1$

Exhaustive classification of $G_2(M^n)$, $n > 3$ follows from the next theorem, proved in this paper.

Theorem 3.

1. A manifold M_+^n admits a flow from $G_2(M_+^n)$ if and only if $M_+^n = \mathbb{S}^{n-1} \times \mathbb{S}^1$. Moreover, there are exactly two classes of topological equivalence in $G_2(\mathbb{S}^{n-1} \times \mathbb{S}^1)$.
2. A manifold M_-^n admits a flow from $G_2(M_-^n)$ if and only if $M_-^n = \mathbb{S}^{n-1} \tilde{\times} \mathbb{S}^1$. Both periodic orbits of such flow are twisted. Moreover, there are exactly two classes of topological equivalence in $G_2(\mathbb{S}^{n-1} \tilde{\times} \mathbb{S}^1)$.

Acknowledgment. This work was supported by the Russian Science Foundation (project 21-11-00010) except for the section 3 which is partially supported by Laboratory of Dynamical Systems and Applications NRU HSE, by Ministry of Science and Higher Education of the Russian Federation (ag. 075-15-2019-1931) and section 4 which was prepared within the framework of the Academic Fund Program at the HSE University in 2021-2022 (grant № 21-04-004).

2 General properties of NMS-flows

In this section we provide NMS-flows properties which are necessary for the further proofs.

Flows f^t and f'^t of the manifold M^n are said to be *topologically equivalent* if there is a homeomorphism $h: M^n \rightarrow M^n$ which sends orbits of f^t into orbits of f'^t and preserves the orientation on the orbits.

To describe behaviour of a flow $f^t: M^n \rightarrow M^n$ in a neighbourhood of attracting or repelling hyperbolic orbit we use the following construction of *suspension*.

Define diffeomorphism $a_{\pm}: \mathbb{R}^{n-1} \rightarrow \mathbb{R}^{n-1}$ by the formula

$$a_{\pm}(x_1, x_2, \dots, x_{n-1}) = (\pm 2x_1, 2x_2, \dots, 2x_{n-1}).$$

Let $g_{\pm}: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a diffeomorphism defined by the formula

$$g_{\pm}(x, r) = (a_{\pm}(x), r - 1).$$

Set $\Pi_{\pm} = \mathbb{R}^n / \langle g_{\pm} \rangle$ and denote the natural projection by $v_{\pm}: \mathbb{R}^n \rightarrow \Pi_{\pm}$. Define the flow \bar{b}^t on \mathbb{R}^n by the system of differential equations:

$$\begin{cases} \dot{x}_1 = 0, \\ \dots, \\ \dot{x}_{n-1} = 0, \\ \dot{x}_n = 1. \end{cases}$$

The natural projection v_{\pm} induces the flow $b_{\pm}^t = v_{\pm} \bar{b}^t v_{\pm}^{-1}: \Pi_{\pm} \rightarrow \Pi_{\pm}$.

Proposition 3 ([8]). *Hyperbolic repelling orbit R of the flow $f^t: M^n \rightarrow M^n$ possesses unstable manifold $W_R^u = \{x \in S \mid f^t(x) \rightarrow R \text{ as } t \rightarrow -\infty\}$ with the following properties:*


1. *there is a map $h_R: W_R^u \rightarrow \Pi_{\pm}$ which is topologically equivalent homeomorphism of the flows $f^t|_{W_R^u}$ and b_{\pm}^t are equivalence. The orbit R is twisted, if $f^t|_{W_R^u}$ is equivalent to b_{\pm}^t and is untwisted otherwise.*
2. *W_R^u is diffeomorphic to $\mathbb{R}^{n-1} \times \mathbb{S}^1$ if R is untwisted and is diffeomorphic to $\mathbb{R}^{n-1} \tilde{\times} \mathbb{S}^1$ if R is twisted.*

Similar theorem holds for the stable manifold of an hyperbolic attracting orbit A which states that a flow f^t is equivalent to b_{\pm}^{-t} in $W_A^s = \{x \in S \mid f^t(x) \rightarrow A \text{ as } t \rightarrow +\infty\}$ by means of the homeomorphism $h: W_A^s \rightarrow \Pi_{\pm}$.


For $r > 0$ let (see Fig. 5)

$$\bar{V}_r^n = \left\{ (x_1, \dots, x_n) \in \mathbb{R}^n \mid x_1^2 + \dots + x_{n-1}^2 \leq r^2 2^{-x_n} \right\}, \quad \mathbb{V}_{\pm r}^n = v_{\pm}(\bar{V}_r^n).$$

By the construction, the quotient space \mathbb{V}_{+r}^n is homeomorphic to *generalized solid torus* $\mathbb{D}^{n-1} \times \mathbb{S}^1$ and quotient \mathbb{V}_{-r}^n is homeomorphic to *generalized solid*



Untwisted orbit of the flow of
two-dimensional manifold



Twisted orbit of the flow of
two-dimensional manifold




Figure 5: The set \bar{V}_r^n

Klein bottle $\mathbb{D}^{n-1} \widetilde{\times} \mathbb{S}^1$. Note, that there are flows $b_{\pm}^t, t \geq 0$ and $b_{\pm}^{-t}, t \leq 0$ correctly defined on the quotients.

Let $\bar{V}^n = \bar{V}_1^n, \mathbb{V}_{\pm}^n = v_{\pm}(\bar{V}^n), \mathbb{L}_{\pm}^n = v_{\pm}(Ox_n)$. Consider homeomorphism $j: \partial \mathbb{V}_{\pm}^n \rightarrow \partial \mathbb{V}_{\pm}^n$, two copies $\mathbb{V}_{\pm}^n \times \mathbb{Z}_2, \mathbb{Z}_2 = \{0, 1\}$ of the manifold \mathbb{V}_{\pm}^n and the homeomorphism $J: \partial \mathbb{V}_{\pm}^n \times \{0\} \rightarrow \partial \mathbb{V}_{\pm}^n \times \{1\}$, defined by the formula $J(s, 0) = (j(s), 1)$. Set

$$M_j^n = \mathbb{V}_{\pm}^n \times \{0\} \cup_J \mathbb{V}_{\pm}^n \times \{1\}.$$

Denote natural projection by $p_j: \mathbb{V}_{\pm}^n \times \mathbb{Z}_2 \rightarrow M_j^n$ and let $f_j^t: M_j^n \rightarrow M_j^n$ be topological flow, defined by the formula:

$$f_j^t(x) = \begin{cases} p_j b_{\pm}^t (p_j|_{\mathbb{V}_{\pm}^n \times \{0\}})^{-1}(x), & x \in p_j|_{\mathbb{V}_{\pm}^n \times \{0\}}, t \leq 0 \\ p_j b_{\pm}^{-t} (p_j|_{\mathbb{V}_{\pm}^n \times \{1\}})^{-1}(x), & x \in p_j|_{\mathbb{V}_{\pm}^n \times \{1\}}, t \geq 0. \end{cases}$$

We will call the constructed flows *n-dimensional model flows*. For any model flow f_j^t defined on the manifold M_j^n let (see Fig. 6)

$$R_j = p_j(\mathbb{L}_{\pm}^n \times \{0\}), A_j = p_j(\mathbb{L}_{\pm}^n \times \{1\}),$$




Figure 6: Phase portrait of model flow on the torus

$$V_{R_j} = p_j(\mathbb{V}_{\pm}^n \times \{0\}), V_{A_j} = p_j(\mathbb{V}_{\pm}^n \times \{1\}),$$

$$\Sigma_j = p_j(\partial\mathbb{V}_{\pm}^n \times \{0\}) = p_j(\partial\mathbb{V}_{\pm}^n \times \{1\}).$$

Lemma 1. *Every flow $f^t \in G_2(M^n)$ is topologically equivalent to some model flow $f^t: M_j^n \rightarrow M_j^n$.*

Proof. Let $f^t \in G_2(M^n)$ and A, R be attracting and repelling hyperbolic orbits of the flow f^t respectively. Due to proposition 3 there is a homeomorphism $h_R: W_R^u \rightarrow \Pi_{\pm}$ which is equivalence homeomorphism for the flows $f^t|_{W_R^u}$ and b_{\pm}^t . Also, there is a homeomorphism $h_A: W_A^s \rightarrow \Pi_{\pm}$ which is equivalence homeomorphism for the flows $f^t|_{W_A^s}$ and b_{\pm}^{-t} . Let $V_A = h_A^{-1}(\mathbb{V}_{\pm}^n)$ and $H_A = h_A|_{\mathbb{V}_{\pm}^n}$. We can choose $r > 0$ such that neighbourhood $V'_R = h_R^{-1}(\mathbb{V}_{\pm r}^n)$ of R is disjoint with V_A . Since non-wandering set consists of A and R then (see e.g. [1])

$$M^n = W_R^u \cup A = W_A^s \cup R$$

and consequently set $M^n \setminus \text{int}(V_A \cup V'_R)$ consists of segments of wandering trajectories of the flow f^t , which has their boundary points on different connected components of the boundary $\partial M^n \setminus \text{int}(V_A \cup V'_R)$ (see e.g. 7). Let $V_R = M^n \setminus \text{int} V_A$. Then the homeomorphism $h_R|_{\mathbb{V}_{\pm r}^n}$ can be extended to the homeomorphism $H_R: \mathbb{V}_{\pm}^n \rightarrow V_R$ which is equivalence homeomorphism for b_{\pm}^t and f^t . Define the homeomorphism $j: \partial\mathbb{V}_{\pm}^n \rightarrow \partial\mathbb{V}_{\pm}^n$ by the formula

$$j = H_A^{-1} H_R|_{\partial\mathbb{V}_{\pm}^n}.$$

Then let homeomorphism $H: M_j^n \rightarrow M^n$, be defined by the formula

$$H(x) = \begin{cases} H_A p_j^{-1}(x), & x \in V_{A_j} \\ H_R p_j^{-1}(x), & x \in V_{R_j} \end{cases}.$$

It is easy to see, that H is an equivalence homeomorphism for f_j^t and f^t . \square




Figure 7: The neighbourhood V_A and V'_R on M^n

Thereby, classification of the flows with the attractor-repeller dynamics can be reduced to the classification of model flows. We give the classification in the further sections. It is easy to show, that it is sufficient to consider some special class of equivalence homeomorphisms by the methods similar to previous proof.

Lemma 2. *If the model flows $f_j^t: M_j^n \rightarrow M_j^n$ and $f_{j'}^t: M_{j'}^n \rightarrow M_{j'}^n$ are topologically equivalent, then there is equivalence homeomorphism $H: M_j^n \rightarrow M_{j'}^n$ for this flows such that $H(\Sigma_j) = \Sigma_{j'}$.*

3 Criteria of topological equivalence of the model flows

In this section we give a criteria for topological equivalence of model flows, which will imply full description of equivalence classes of $G_2(M^n)$.

Denote the connected component of the set $\partial V^n \cap O x_2 x_n$ containing the point $(0, 1, \dots, 0)$ by $\bar{\alpha}$. We consider $\bar{\alpha}$ as curve oriented in the direction of increasing x_n coordinate. Let $\bar{\beta} = \partial V^n \cap O x_1 \dots x_{n-1}$ (see Fig. 8) and

$$\alpha_{\pm} = v_{\pm}(\bar{\alpha}), \beta_{\pm} = v_{\pm}(\bar{\beta}).$$

Denote inclusion map by $i: \partial \mathbb{V}_{\pm}^n \rightarrow \mathbb{V}_{\pm}^n$ and isomorphism induced by it by $i_*: \pi_1(\partial \mathbb{V}_{\pm}^n) \rightarrow \pi_1(\mathbb{V}_{\pm}^n)$. Since the group $\langle g_{\pm} \rangle$ is isomorphic to \mathbb{Z} and acts freely and discontinuously on simple-connected space \bar{V}^n fundamental group $\pi_1(\mathbb{V}_{\pm}^n)$ is also isomorphic to the group \mathbb{Z} (see e.g. [11]) and α_{\pm} is its generator².

²The space $\partial \mathbb{V}_{\pm}^2$ consist of two connected components. In this case i_* is a map, composed of this components fundamental group automorphisms.




Figure 8: $\bar{\alpha}$ and $\bar{\beta}$ on \bar{V}^n

Theorem 4 (Criteria of topological equivalence). *The flows $f_j^t: M_j^n \rightarrow M_j^n$, $f_{j'}^t: M_{j'}^n \rightarrow M_{j'}^n$ are topologically equivalent if and only if there is a homeomorphism $h_0: \partial\mathbb{V}_\pm^n \rightarrow \partial\mathbb{V}_\pm^n$ such that*

$$i_* h_{0*} = i_* \quad (1)$$

and homeomorphism $h_1 = j' h_0 j^{-1}$ is such that

$$i_* h_{1*} = i_* \quad (2)$$

Proof. Necessity. Let the flows f_j^t and $f_{j'}^t$ be topologically equivalent by means of the homeomorphism $H: M_j^n \rightarrow M_{j'}^n$. By the lemma 2 without limitation of generality we assume that $H(\Sigma_j) = \Sigma_{j'}$. Let homeomorphism $H_k: \mathbb{V}_\pm^n \rightarrow \mathbb{V}_\pm^n$, $k \in \{0, 1\}$ be defined such that $(H_k(s), k) = p_{j'}^{-1} H p_j|_{\mathbb{V}_\pm^n \times \{k\}}(s, k): \mathbb{V}_\pm^n \times \{k\} \rightarrow \mathbb{V}_\pm^n \times \{k\}$. Set $h_k = H_k|_{\partial\mathbb{V}_\pm^n}$.

It is easy to see that curves \mathbb{L}_\pm^n and α_\pm are homotopic in \mathbb{V}_\pm^n due to the fact that they bound two-dimensional annulus $v_\pm(\bar{V}^n \cap O x_2 x_n)$ in \mathbb{V}_\pm^n . Since H is an equivalence homeomorphism for flows f_j^t and $f_{j'}^t$, then $H_0(R_j) = R_{j'}$. It implies $H_{0*}([R_j]) = [R_{j'}]$. Considering the fact that $\pi_1(\mathbb{V}_\pm^n) \cong \langle \alpha_\pm \rangle \cong \mathbb{Z}$ we get that $H_{0*} = id$ which implies (1).

$h_1 = j' h_0 j^{-1}$ follows from the definition of the model flow. The condition (2) $i_* h_{1*} = i_*$ for the map h_1 is proved with the same method.

Sufficiency. Assume there is a homeomorphism $h_0: \partial\mathbb{V}_\pm^n \rightarrow \partial\mathbb{V}_\pm^n$ such that $i_* h_{0*} = i_*$ and the homeomorphism $h_1 = j' h_0 j^{-1}$ satisfies $i_* h_{1*} = i_*$. Since $i_* h_{0*} = i_*$ and due to [9][proposition 10.2.26], homeomorphism h_0 admits a lift




Figure 9: Construction of the disk B'

$\bar{h}_0: \partial V^n \rightarrow \partial \bar{V}^n$, which commutes with g_{\pm} . Let $\bar{\beta}' = \bar{h}_0(\bar{\beta})$. Choose positive integer n_0 such that $\bar{\beta}' \subset \{(x_1, \dots, x_{n-1}, x_n) \in \mathbb{R}^n : 0 < x_n < n_0\}$ (see Fig. 9). Extend the homeomorphism \bar{h}_0 to a homeomorphism $\bar{H}_0: \bar{V}^n \rightarrow V^n$ commuting with g_{\pm} which implements equivalence of the flow \bar{b}^t with itself.

Let $y = (x_1, \dots, x_{n-1})$ and define a map $p_0: Oy x_n \rightarrow Oy$ by the formula

$$p_0(y, x_n) = y.$$

By the construction $p_0|_{\partial \bar{V}_r^n}: \partial \bar{V}_r^n \rightarrow Oy \setminus O$ is a diffeomorphism for any $r > 0$. Define the homeomorphism $\bar{w}: Oy \setminus O \rightarrow Oy \setminus O$ by the formula

$$\bar{w} = p_0 \bar{h}_0 (p_0|_{\partial \bar{V}_1^n})^{-1}.$$

Since $i_* h_{0*} = i_*$ the homeomorphism \bar{w} can be continuously extended to the point O by the formula $\bar{w}(O) = O$. Assume that $y' = \bar{w}(y)$, $y \in Oy$, $\bar{\beta}'_0 = p_0(\bar{\beta}')$ and $\bar{\beta}_0 = p_0(g_{\pm}^{n_0}(\bar{\beta}))$. Let \bar{B} , \bar{B}_0 , \bar{B}'_0 denote the disks in Oy which are bounded by spheres $\bar{\beta}$, $\bar{\beta}_0$, $\bar{\beta}'_0$ respectively. Thus, $\bar{w}(\bar{B}) = \bar{B}'_0$.

By virtue of the annulus conjecture (see e.g. [9]) and the fact that spheres $\bar{\beta}_0$ and $\bar{\beta}'_0$ are disjoint and are cylindrically embedded they bound an annulus $K_0 \subset Ox_1 \dots x_{n-1}$. Let $\tau: \mathbb{S}^{n-2} \times [0, 1] \rightarrow K_0$ be a homeomorphism such that $\tau(\mathbb{S}^{n-2} \times \{0\}) = \bar{\beta}_0$ and $\tau(\mathbb{S}^{n-2} \times \{1\}) = \bar{\beta}'_0$. For $t \in [0, 1]$ put $c_t = \tau(\mathbb{S}^{n-2} \times \{t\})$, $r_t = t + 2^{-n_0}(1-t)$ and $C_t = (p_0|_{\partial \bar{V}_{r_t}^3})^{-1}(c_t)$. Define the disk \bar{B}' by the formula

$$\bar{B}' = \bigcup_{t \in [0, 1]} C_t \cup \bar{B}_0.$$

Since $i_* h_{0*} = i_*$, homeomorphism \bar{h}_0 sends the part $\bar{\Gamma}$ of the cylinder $\partial \bar{V}^n$ lying between spheres $\bar{\beta}$, $g_{\pm}^{-1}(\bar{\beta})$ to the part $\bar{\Gamma}'$ of cylinder $\partial \bar{V}^n$ lying between the spheres $\bar{\beta}'$, $g_{\pm}^{-1}(\bar{\beta}')$. For every $y_0 \in Oy$ let $L_{y_0} = \{(y, x_n) \in \mathbb{R}^n : y = y_0\}$. For $(y_0, 0) \in \bar{B}$ let $I_{y_0} = L_{y_0} \cap \bar{W}$ and $I'_{y'_0} = L_{y'_0} \cap \bar{W}'$. Denote boundary points of segments I_{y_0} and $I'_{y'_0}$ by A_{y_0}, B_{y_0} and $A'_{y'_0}, B'_{y'_0}$ where $A_{y_0} = (y_0, 0)$, $B_{y_0} = (y_0, b_{y_0})$ and $A'_{y'_0} = (y'_0, a'_{y'_0})$, $B'_{y'_0} = (y'_0, b'_{y'_0})$, $a'_{y'_0} \leq b'_{y'_0}$. Define a homeomorphism $\bar{h}_{y_0} : I_{y_0} \rightarrow I'_{y'_0}$ by the formula

$$\bar{h}_{y_0}(y_0, x_n) = \left(y'_0, x_n \frac{b'_{y'_0} - a'_{y'_0}}{b_{y_0}} + a'_{y'_0} \right).$$

By virtue of the fact that $\bar{W} = \bigcup_{y_0 \in \bar{B}} I_{y_0}$, \bar{h}_{y_0} is equivalence homeomorphism of the flow \bar{b}^t with itself on \bar{W} . Extend it to \bar{V}^n by the formula

$$\bar{H}_0(x_1, \dots, x_{n-1}, x_n) = g^{-[x_n]}(\bar{h}_{y_0}|_{\bar{W}}(g^{[x_n]}(x_1, \dots, x_{n-1}, x_n))),$$

where $[x]$ denotes the integer part of number x . By the construction $\bar{H}_0 g_{\pm} = g_{\pm} \bar{H}_0$. By virtue of [9] this fact implies, that $H_0 = v_{\pm}^{-1} \bar{H}_0 v_{\pm}$ is a homeomorphism and the following equality holds $H_0 b^t = b^t H_0$.

By the similar methods the homeomorphism h_1 extends to the homeomorphism $H_1 : \mathbb{V}_{\pm}^n \rightarrow \mathbb{V}_{\pm}^n$, which implements equivalence of the flow b_{\pm}^{-t} to itself. Thus, sought homeomorphism $H : M_j^n \rightarrow M_j^n$ is defined by the formula

$$H(x) = \begin{cases} p_j' H_0 p_j^{-1}(x), & x \in p_j(\mathbb{V}_{\pm}^n \times \{0\}) \\ p_j' H_1 p_j^{-1}(x), & x \in p_j(\mathbb{V}_{\pm}^n \times \{1\}) \end{cases}.$$

□

4 Classification of model flows of surfaces

In this section we prove the theorem 1.

Let $f_j^t : M_j^2 \rightarrow M_j^2$ be two-dimensional model flow. Then ambient surface M_j^2 has the form $M_j^2 = \mathbb{V}_{\pm}^2 \times \{0\} \cup_J \mathbb{V}_{\pm}^2 \times \{1\}$, where $J : \partial \mathbb{V}_{\pm}^2 \times \{0\} \rightarrow \partial \mathbb{V}_{\pm}^2 \times \{1\}$ is a homeomorphism, defined as $J(s, 0) = (j(s), 1)$ for some homeomorphism $j : \partial \mathbb{V}_{\pm}^2 \rightarrow \partial \mathbb{V}_{\pm}^2$.

If the periodic orbits are untwisted then tabular neighbourhood is an annulus and its boundary has two connected components each of which is homeomorphic to the circle. If the periodic orbits are twisted then tabular neighbourhood of each is Möbius band and its boundary is homeomorphic to the circle. Let $\mathbb{S}^1 = \{e^{i\varphi}, \varphi \in \mathbb{R}\}$, $\mathbb{S}^0 = \{-1, +1\}$ and define diffeomorphisms on the manifold $\partial \mathbb{V}_{\pm}^2 \cong \mathbb{S}^1 \times \mathbb{S}^0$:

1. $j_1(\varphi, \pm 1) = (\varphi, \pm 1)$;
2. $j_2(\varphi, \pm 1) = (-\varphi, \pm 1)$;
3. $j_3(\varphi, \pm 1) = (\pm \varphi, \pm 1)$.

Define diffeomorphisms on the manifold $\partial\mathbb{V}_-^2 \cong \mathbb{S}^1$:

4. $j_4(\varphi) = \varphi$;
5. $j_5(\varphi) = -\varphi$.

Pictures 10, 11 provide phase portraits of model flows corresponding to given maps. Sign "+" means the gluing with the map φ and the sign "-" with the map $-\varphi$.




Figure 10: Phase portraits of not topologically equivalent flows on the torus




Figure 11: Phase portraits of not topologically equivalent flows on the Klein bottle: 1. with untwisted orbits; 2-3. with twisted ones

Lemma 3.

1. Every model flow f_j^t of torus is topologically equivalent to one of the two: $f_{j_1}^t, f_{j_2}^t$ which are inequivalent to each other.
2. Every model flow f_j^t of Klein bottle is topologically equivalent to one of the three $f_{j_3}^t, f_{j_4}^t, f_{j_5}^t$ which are pairwise inequivalent.

Proof. Since fundamental group of the circle is isomorphic to the group \mathbb{Z} then any orientation-preserving homeomorphism induces identical action in

the fundamental group and orientation-reversing induces action which changes class of the curve to the opposite. Then, by virtue of the theorem 4 flows $f_j^t: M_j^2 \rightarrow M_j^2$, $f_{j'}^t: M_{j'}^2 \rightarrow M_{j'}^2$ are topologically equivalent iff there is orientation-preserving homeomorphism $h_0: \partial\mathbb{V}_\pm^2 \rightarrow \partial\mathbb{V}_\pm^2$ such that the homeomorphism $h_1 = j'h_0j^{-1}$ preserves orientation. That is equivalent to the fact that $j'j^{-1}$ preserves orientation.

Exhaustive search of all the possible combinations of orientability of the homeomorphism j on the connected components gives that homeomorphism $j_i j^{-1}$ preserves orientation exactly for only one value of $i \in \{1, \dots, 5\}$. Moreover, if $i = 1, 2$ the ambient manifold is the torus, but for $i = 3, 4, 5$ is the Klein bottle. Finally, if $i = 4, 5$ both orbits are twisted. \square

5 Classification of 3-dimensional model flows

Let $f_j^t: M_j^3 \rightarrow M_j^3$ be a three-dimensional model flow. Then ambient manifold M_j^3 has the form $M_j^3 = \mathbb{V}_\pm^3 \times \{0\} \cup_J \mathbb{V}_\pm^3 \times \{1\}$ where $J: \partial\mathbb{V}_\pm^3 \times \{0\} \rightarrow \partial\mathbb{V}_\pm^3 \times \{1\}$ is the homeomorphism defined by the formula $J(s, 0) = (j(s), 1)$ for some homeomorphism $j: \partial\mathbb{V}_\pm^3 \rightarrow \partial\mathbb{V}_\pm^3$. It is easy to note that the manifold \mathbb{V}_+^3 is solid torus whereas \mathbb{V}_-^3 is the solid Klein bottle. In the first case ambient manifold M_j^3 is a lens space, which is orientable and (see proposition 4 below) there are countably many pairwise non-homeomorphic lens spaces. All the manifolds obtained by gluing solid Klein bottles, on the contrary, are homeomorphic to the $S^2 \widetilde{\times} S^1$ (see e.g. [10, section 3.1(c)]).

Let us prove the theorem 2 separately for orientable and non-orientable manifolds.

5.1 Orientable case

On the torus $\partial\mathbb{V}_+^3$ the curves α_+ and β_+ are generators of the torus fundamental group. Oriented curves α_+, β_+ on the torus $\partial\mathbb{V}_+^3$ are said to be *longitude and meridian* respectively. Then the action of the homeomorphism j in the fundamental group of the torus is uniquely defined by unimodular integer matrix

$$j_* = \begin{pmatrix} r & p \\ s & q \end{pmatrix}.$$

The classification of the lens spaces is well known (see e.g. [12]) via relations on parameters p, q .

Proposition 4 (3-dimensional lens space classification). *Two lens spaces M_j^3 and $M_{j'}^3$ are homeomorphic iff the induces isomorphisms $j_* = \begin{pmatrix} r & p \\ s & q \end{pmatrix}$, $j'_* = \begin{pmatrix} r' & p' \\ s' & q' \end{pmatrix}$ satisfy the conditions $|p'| = |p|$ and $q' \equiv \pm q \pmod{|p|}$.*




Figure 12: Longitude and meridian pre-images on \bar{V}^3

Lemma 4. Homeomorphism $h_0: \partial\mathbb{V}_+^3 \rightarrow \partial\mathbb{V}_+^3$ such that for h_0 and $h_1 = j'h_0j^{-1}$ the equality $i_*h_{k*} = i_*$, $k = 0, 1$ holds there is iff induced isomorphisms $j_* = \begin{pmatrix} r & p \\ s & q \end{pmatrix}$, $j'_* = \begin{pmatrix} r' & p' \\ s' & q' \end{pmatrix}$ satisfy the conditions $|p'| = |p|, r' \equiv r \pmod{|p|}$.

Proof.

Necessity. Assume there is homeomorphism $h_0: \partial\mathbb{V}_+^3 \rightarrow \partial\mathbb{V}_+^3$ such that for h_0 and $h_1 = j'h_0j^{-1}$ the condition $i_*h_{k*} = i_*$, $k = 0, 1$ holds. Then the homeomorphism h_k acts in the fundamental group of torus with the matrix

$$h_{k*} = \begin{pmatrix} 1 & 0 \\ m_k & \pm 1 \end{pmatrix}$$

and

$$h_{1*}j_* = j'_*h_{0*}.$$

The last relation can be written in matrix form:

$$\begin{pmatrix} 1 & 0 \\ m_1 & \pm 1 \end{pmatrix} \begin{pmatrix} r & p \\ s & q \end{pmatrix} = \begin{pmatrix} r' & p' \\ s' & q' \end{pmatrix} \begin{pmatrix} 1 & 0 \\ m_0 & \pm 1 \end{pmatrix}. \quad (3)$$

So, we get the equalities $p = \pm p'$, $r = r' + m_0 p'$, which are equivalent to $|p'| = |p|$ and $r' \equiv r \pmod{|p|}$.

Sufficiency. Let elements of matrices $j_* = \begin{pmatrix} r & p \\ s & q \end{pmatrix}$, $j'_* = \begin{pmatrix} r' & p' \\ s' & q' \end{pmatrix}$ satisfy the relations $|p'| = |p|$ and $r' \equiv r \pmod{|p|}$. Thus

$$p = \delta_0 p', \quad r = r' + m_0 p' \quad (4)$$

for some $\delta_0 \in \{-1, 1\}$, $m_0 \in \mathbb{Z}$. Let $h_0: \partial\mathbb{V}_+^3 \rightarrow \partial\mathbb{V}_+^3$ be algebraic torus automorphism defined by the matrix $h_{0*} = \begin{pmatrix} 1 & 0 \\ m_0 & \delta_0 \end{pmatrix}$. Then $i_* h_{0*} = i_*$.

The formula (3) and the fact that all the matrix are unimodular we get, that homeomorphism $h_1 = j' h_0 j^{-1}: \partial\mathbb{V}_+^3 \rightarrow \partial\mathbb{V}_+^3$ induces isomorphism, defined by the matrix $h_{1*} = \begin{pmatrix} 1 & 0 \\ m_1 & \delta_1 \end{pmatrix}$ for some $\delta_1 \in \{-1, 1\}$, $m_1 \in \mathbb{Z}$. Thus $i_* h_{1*} = i_*$. \square

Lemma 5. *Up to topological equivalence there is only one flow on both \mathbb{S}^3 and $\mathbb{R}P^3$ and two on the rest of the lens spaces.*

Proof. Considering the fact, that matrices j_* , j'_* are unimodular and the proposition 4 it is easy to establish one more criteria for the lens spaces to be homeomorphic. Namely, two lens spaces M_j^3 and $M_{j'}^3$ are homeomorphic iff induced isomorphisms $j_* = \begin{pmatrix} r & p \\ s & q \end{pmatrix}$, $j'_* = \begin{pmatrix} r' & p' \\ s' & q' \end{pmatrix}$ satisfy the relations $|p'| = |p|$ and $r' \equiv \pm r \pmod{|p|}$.

By virtue of the lemma 4 the previous fact implies that on the same lens space there are either one or two classes of topological equivalence of flows from $G_2(M_+^3)$ and the cases are distinguished by the following condition: whether for two coprime numbers $p \geq 0$, $r \in \mathbb{Z}_p$ there are two numbers n_1, n_2 such that

$$r + n_1 p = -r + n_2 p \quad (5)$$

Check whether the condition (5) holds for all the possible values of p .

1. if $p = 0$ than condition (5) does not hold, since the equality is true only if $r = 0$ but in this case r, p are not coprime.
2. if $p = 1$ the condition holds for $r = 0$ which means that up to topological equivalence there is unique flow in considered class on the sphere \mathbb{S}^3
3. if $p = 2k$ then $r = k(n_2 - n_1)$. Considering the fact that $(r, p) = 1$ get that $k = 1, r = 1$, which means that up to topological equivalence there is unique flow from the considered class on the $\mathbb{R}P^3$
4. for $p = 2k + 1, k > 0$ condition (5) never holds since $n_2 - n_1$ is even and $(r, p) \neq 1$.

\square

6 Non-orientable case

By virtue of the results [13] every homeomorphism $j: \partial\mathbb{V}_-^3 \rightarrow \partial\mathbb{V}_-^3$ satisfy either $i_*(j_*([c])) = i_*([c])$ or $i_*(j_*([c])) = -i_*([c])$. Then the theorem 4 implies that there are two classes of topological conjugacy of the considered flows on the manifold $\mathbb{S}^2 \tilde{\times} S^1$.

7 Classification of model flows on n -manifolds for $n > 3$

According to [14] the only manifold obtained by gluing two copies of \mathbb{V}_+^n is $\mathbb{S}^{n-1} \times \mathbb{S}^1$.

Similarly, the results of the paper [15] imply that the only manifold obtained by gluing two copies of \mathbb{V}_-^n along the boundary is $\mathbb{S}^{n-1} \tilde{\times} \mathbb{S}^1$.

Since fundamental group $\pi_1(\partial\mathbb{V}_\pm^n)$ is isomorphic to the group \mathbb{Z} then any homeomorphism $j: \partial\mathbb{V}_\pm^n \rightarrow \partial\mathbb{V}_\pm^n$ satisfy either condition $i_*(j_*([c])) = i_*([c])$ or $i_*(j_*([c])) = -i_*([c])$. Thus, the theorem 4 implies that each of the manifolds $\mathbb{S}^{n-1} \times \mathbb{S}^1$ and $\mathbb{S}^{n-1} \tilde{\times} \mathbb{S}^1$ admits two classes of topological equivalence of the considered flows.

References

- [1] S. Smale, “Differentiable dynamical systems”, Bull. Amer. Math. Soc., **73**, 1967), 747–817.
- [2] M.M. Peixoto, “On a classification of flows on 2-manifolds”, Proc. Symp. Dyn. Syst. Salvador, 1973, 389–492.
- [3] Oshemkov A. A., Sharko V. V. Classification of Morse-Smale flows on two-dimensional manifolds //Sbornik: Mathematics. - 1998. – T. 189. - №. 8. - C. 1205.
- [4] Kruglov, Vladislav and Malyshev, Dmitriy and Pochinka, O., 2017, 06, *Topological classification of Ω -stable flows on surfaces by means of effectively distinguishable multigraphs*, 38, Discrete & Continuous Dynamical Systems - A, doi: 10.3934/dcds.2018188
- [5] Umanskii Y. L. Necessary and sufficient conditions for topological equivalence of three-dimensional Morse-Smale dynamical systems with a finite number of singular trajectories //Matematicheskii Sbornik. - 1990. - T. 181. - №. 2. - C. 212-239.
- [6] Campos, Beatriz & Cordero, Alicia & Martinez-Alfaro, Jose & Vindel, P.. (2004). NMS Flows on Three-Dimensional Manifolds with One Saddle Periodic Orbit. Acta Mathematica Sinica-english Series - ACTA MATH SIN-ENGLISH SERIES. 20. 47-56. 10.1007/s10114-003-0305-z.
- [7] Yu. Bin, “Behavior 0 nonsingular Morse Smale flows on S^3 ”, Discrete Contin. Dyn. Syst. **36:1**, 2016), 509–540.
- [8] Irwin, M.: A Classification of Elementary Cycles. Topology 9, 35–48 (1970).
- [9] Grines V. Z., Medvedev T. V., Pochinka O. V. Dynamical systems on 2-and 3-manifolds. - Cham: Springer, 2016. – T. 46.

- [10] Gómez-Larrañaga, J. & Wolfgang, Heil & González-Acuña, F.. (2004). 3-Manifolds that are covered by two open Bundles. Boletín de la Sociedad Matemática Mexicana. Third Series. 10.
- [11] Kosniowski C. A first course in algebraic topology. – CUP Archive, 1980.
- [12] Allen Hatcher. Notes on basic 3-manifold topology. Notes available on the author's webpage, 2007.
- [13] Lickorish W. B. R. Homeomorphisms of non-orientable two-manifolds //Mathematical Proceedings of the Cambridge Philosophical Society. – Cambridge University Press, 1963. – T. 59. – №. 2. – C. 307-317.
- [14] Max, Nelson L. Homeomorphisms of $S^n \times S^1$. Bull. Amer. Math. Soc. 73 (1967), no. 6, 939–942.
- [15] Jähren B., Kwasik S. Free involutions on $S^n \times S^1$ //Mathematische Annalen. – 2011. – T. 351. – №. 2. – C. 281-303.