

Infrared spectra of both isomers of CO₂-CO in the CO₂ ν_3 region

A.J. Barclay,^a A.R.W. McKellar,^b and N. Moazzen-Ahmadi^a

^a *Department of Physics and Astronomy, University of Calgary, 2500 University Drive North West, Calgary, Alberta T2N 1N4, Canada*

^b *National Research Council of Canada, Ottawa, Ontario K1A 0R6, Canada*

Abstract

Extensive infrared spectra of the weakly-bound CO₂-CO dimer are observed in the carbon dioxide ν_3 asymmetric stretch region (≈ 2350 cm⁻¹) using a tunable infrared OPO laser source to probe a pulsed slit jet supersonic expansion. Both C-bonded and O-bonded isomers are analyzed for the normal isotopologue as well as for ¹³CO₂-CO and ¹⁶O¹³C¹⁸O-CO, the latter being the first observation of an asymmetrically substituted form for which all values of K_a are allowed.

Combination bands involving the lowest in-plane intermolecular bending modes are also studied for both isomers. Weak bands near 2337 cm⁻¹ are assigned to CO₂ hot band transitions (ν_1, ν_2^{l2}, ν_3) = (01¹1) \leftarrow (01¹0), yielding the splitting of the degenerate CO₂ ν_2 bend into in-plane and out-of-plane components due to the presence of the CO. This splitting has rather different values for the C- and O- bonded isomers, 4.56 and 1.59 cm⁻¹, respectively, with the out-of-plane mode higher in energy than the in-plane for both cases.

1. Introduction

The weakly-bound van der Waals dimer CO₂-CO was first observed by high-resolution microwave [1] and infrared [1-4] spectroscopy in the form of a planar T-shaped C-bonded structure ('isomer 1'), with CO₂ forming the 'top' of the T and CO forming the 'stem' and an intermolecular center of mass separation of about 3.9 Å. More recently, spectra of a second O-bonded isomer of CO₂-CO were observed [5]. This 'isomer 2' has a similar planar T-shaped structure, but with the CO flipped by 180° and a shorter intermolecular distance of about 3.6 Å. In addition to these gas-phase (supersonic jet) studies, infrared spectra of matrix-isolated CO₂-CO have been studied [6-8], but their relevance to the isolated dimer may be limited since interactions with the matrix itself are comparable in strength to the interaction between CO₂ and CO [8]. There have also been a number of theoretical studies of the CO₂ – CO interaction [9,10], including two very recent high level *ab initio* determinations of the four-dimensional potential surface [11,12].

Previously observed infrared spectra of CO₂-CO include those of isomer 1 in the CO₂ ν_3 region (CO₂-CO and CO₂-¹³CO) [1] and the CO stretch region (CO₂-CO and C¹⁸O₂-CO) [2-34]. Isomer 2 was observed only in the CO stretch region (CO₂-CO and C¹⁸O₂-CO) [4,5]. (Isotope numbers for C and O are 12 and 16 unless otherwise given.) In addition, combination bands involving low frequency intermolecular modes had been detected for both isomers in the CO stretch region [4,5]. In the present paper, we report extensive new data for CO₂-CO as observed in the CO₂ ν_3 region. These include the first observations in this region of isomer 2 and of intermolecular combination bands, and the first observation of an asymmetrically substituted dimer (¹⁶O¹³C¹⁸O-CO). Most importantly, we also detect spectra of both isomers corresponding to the weak $(\nu_1, \nu_2^{l2}, \nu_3) = (01^11) \leftarrow (01^10)$ hot band of CO₂, and thus observe the splitting of the

degenerate CO₂ ν_2 bending mode into its in-plane and out-of-plane components due to the presence of the nearby CO molecule. This splitting, turns out to be rather different for isomers 1 and 2.

As mentioned, both CO₂-CO isomers have planar T-shaped equilibrium structures with C_{2v} point group symmetry. The a -inertial axis coincides with the CO axis and is perpendicular to the CO₂ axis. Since rotation around this axis interchanges the carbon dioxide O-atoms, only levels with $K_a = \text{even}$ are allowed by nuclear spin statistics for ground vibrational state dimers containing C¹⁶O₂ or C¹⁸O₂. The b -axis is parallel to the CO₂ axis, so fundamental bands in the CO₂ ν_3 (asymmetric stretch) region have b -type rotational selection rules. The c axis is perpendicular to the plane of the dimer. Each isomer has four intermolecular vibrational modes: van der Waals stretch (A_1 symmetry), out-of-plane rock (B_1), and two in-plane bends (B_2). The in-plane bends can be described as a CO rock and CO₂ rock, or else as a geared and anti-geared bend.

2. Observed spectra

Spectra were recorded at the University of Calgary as described previously [13-15]. A pulsed supersonic slit jet expansion is probed using a rapid-scan optical parametric oscillator source. A typical expansion gas mixture contains about 0.08% carbon dioxide plus 0.3% carbon monoxide in helium carrier gas with a backing pressure of about 11 atmospheres. Wavenumber calibration is carried out by simultaneously recording signals from a fixed etalon and a reference gas cell containing room temperature CO₂. Spectral simulation and fitting are made using the PGOPHER software [16].

2.1. Fundamental bands

Legon and Suckley [1] previously studied the fundamental band of CO₂-CO (and CO₂-¹³CO) in the CO₂ ν_3 region. Our current results for isomer 1 are similar, except that we observe the $K_a = 3 \leftarrow 4$ and $5 \leftarrow 4$ subbands in addition to the previously observed $1 \leftarrow 2$, $1 \leftarrow 0$, and $3 \leftarrow 2$ subbands. We also observe isomer 2, the O-bonded form, for the first time in this region. The spectrum is shown in Fig. 1, and results from its analysis are in Table 1. Note that the band origins of the two isomers are separated by only 0.230 cm⁻¹, so their spectra overlap almost completely. In contrast, the two isomer bands are separated by 7.950 cm⁻¹ in the CO stretch region [4,5], making it easier to detect and analyze the weaker isomer 2 spectrum. The *S*-reduced form of the asymmetric rotor Hamiltonian [16] is used here for consistency with Refs. 2-5. For isomer 1, the ground state parameters are fixed at previously determined [3,5] values which incorporate precise microwave data [1]. We assign 92 observed isomer 1 lines in terms of 117 transitions with *J*-values up to 12 (some lines are unresolved asymmetry doublets or other blends). There were 81 assigned transitions in Ref. 1. For the weaker isomer 2 fundamental, ground and excited state parameters are varied, and we assign 47 lines (59 transitions) belonging to the $K_a = 3 \leftarrow 4$, $1 \leftarrow 2$, $1 \leftarrow 0$, $3 \leftarrow 2$, and $5 \leftarrow 4$ subbands with *J*-values up to 9. The fundamental and combination (see below) bands were fitted simultaneously, with root mean square (rms) residuals of 0.00040 and 0.00036 cm⁻¹ for isomers 1 and 2, respectively.

We also study the ¹³CO₂-CO isotopologue, using isotopically enriched ¹³CO₂ in the expansion gas mix. The results are very similar to those of the normal species (see Fig. 2 and Table 2). For isomer 1, 77 lines (93 transitions) are observed in the $K_a = 3 \leftarrow 4$, $1 \leftarrow 2$, $1 \leftarrow 0$, and $3 \leftarrow 2$ subbands and fitted with an rms residual of 0.00079 cm⁻¹. For isomer 2, 38 lines (47 transitions) are observed in the same subbands and fitted with an rms residual of 0.00021 cm⁻¹.

For this isotopologue there are no previous data, and the ground state parameters are thus determined for the first time. They are very similar to those of the normal isotopologue, reflecting the fact that the ^{13}C atom is located close to the dimer center of mass.

While using the $^{13}\text{CO}_2$ gas mix, a weak spectrum was detected in the 2266 cm^{-1} region which we eventually realized must be the fundamental band of $^{16}\text{O}^{13}\text{C}^{18}\text{O-CO}$. For this species, there is no nuclear spin statistical effect, and all values of K_a are possible. Thus the spectrum, shown in Fig. 3, looks different than any previously studied $\text{CO}_2\text{-CO}$ spectrum. The fitted parameters are given in Table 3. For isomer 1, 79 lines (86 transitions) are assigned in the $K_a = 1 \leftarrow 2$, $0 \leftarrow 1$, $1 \leftarrow 0$, $2 \leftarrow 1$, and $3 \leftarrow 2$ subbands and fitted with an rms residual of 0.00037 cm^{-1} . For isomer 2, 42 lines (47 transitions) are assigned in the same subbands and fitted with a residual of 0.00058 cm^{-1} . The natural abundance of ^{18}O is 0.2%, and since there are two equivalent O atoms, the natural abundance of $^{16}\text{O}^{13}\text{C}^{18}\text{O}$ is 0.4%. But the strength of the $^{16}\text{O}^{13}\text{C}^{18}\text{O-CO}$ spectrum is spread among all K_a values, not just even ones, so 0.2% remains its expected intensity ratio relative to $^{13}\text{CO}_2\text{-CO}$ for the strongest lines in the band. We estimate that the strongest $^{16}\text{O}^{13}\text{C}^{18}\text{O-CO}$ transitions (Fig. 3) actually had about 0.5% the intensity of the strongest $^{13}\text{CO}_2\text{-CO}$ transitions (Fig. 2), though this is only approximate because laser power and jet expansion conditions may change between regions. Thus the observed strength of $^{16}\text{O}^{13}\text{C}^{18}\text{O-CO}$ is roughly twice that which might be expected. A possible explanation for this strength may simply be that our enriched $^{13}\text{CO}_2$ gas sample is also somewhat enriched in ^{18}O .

2.2. Combination bands

In the CO stretch region, combination bands involving intermolecular modes have been studied previously for both isomers of $\text{CO}_2\text{-CO}$ [4,5]. The two modes observed were the lower frequency in-plane bend (B_2 symmetry), with values of 24.34 (isomer 1) and 14.19 (isomer 2)

cm^{-1} , and the out-of-plane rock (B_1 symmetry) with values of about 43.96 (isomer 1) and 22.68 cm^{-1} (isomer 2).

In the CO_2 ν_3 region, we observe the in-plane bend for both isomers. This intermolecular mode and the CO_2 ν_3 intramolecular mode both have B_1 symmetry, so their combination has A_1 symmetry and the combination bands originating in the ground state have a -type selection rules. Results for isomer 1 are illustrated in Fig. 4 for both CO_2 -CO and $^{13}\text{CO}_2$ -CO, with fitted parameters given in Tables 1 and 2. The resulting value for the in-plane bending mode frequency in the excited CO_2 ν_3 state is 24.510 cm^{-1} for CO_2 -CO (and 24.501 cm^{-1} for $^{13}\text{CO}_2$ -CO), close to the value of 24.34 cm^{-1} observed in the CO stretch region. The experimental value is also close to theoretical predictions of 24.44 and 24.45 cm^{-1} (in the ground intramolecular state) from recent *ab initio* studies [11, 12]. (There are some inconsistencies in the intermolecular frequencies reported in Ref. 11; we use here values obtained from their Table 2). The very weak combination band of isomer 2 is not shown here, but its fitted parameters are included in Table 1 (this band was not observed for $^{13}\text{CO}_2$ -CO). The resulting intermolecular mode frequency is 14.372 cm^{-1} . Again, we find that this value is quite close to that observed in the CO stretch region (14.174 cm^{-1}) [5] and to those from recent theoretical predictions [11, 12] (14.94 and 14.68 cm^{-1}).

Small perturbations are evident in the isomer 1 combination band spectra (Fig. 4) which clearly involve the upper state. For example, in the normal CO_2 -CO spectrum, the $J_{KaKc} = 4_{04} \leftarrow 5_{05}$ and $4_{04} \leftarrow 3_{03}$ transitions both have a ‘satellite’ line which is 0.0106 cm^{-1} higher and about 0.13 times the strength of the main line. Similarly, $6_{06} \leftarrow 7_{07}$ and $6_{06} \leftarrow 5_{05}$ show a satellite 0.0075 cm^{-1} higher with about 0.33 times the strength. As well, the upper state levels 5_{24} and 7_{26} are split into two components separated by about 0.007 and 0.004 cm^{-1} , respectively. In the

$^{13}\text{CO}_2\text{-CO}$ spectrum, perturbations only seem to affect upper state levels with $K_a = 2$ and $J < 5$. For example, the 3_{22} level is split into two components separated by 0.0136 cm^{-1} . Looking at the possibilities, it seems likely that the source of these perturbations is the excited intramolecular CO stretch mode of $\text{CO}_2\text{-CO}$ combined with a highly excited ($\approx 230\text{ cm}^{-1}$) intermolecular mode.

2.3. The $(01^11) \leftarrow (01^10)$ hot band

We observed a relatively weak but very interesting spectrum around 2337 cm^{-1} , shown in Fig. 5, whose explanation was not apparent at first. Eventually, we realized that it was due to $\text{CO}_2\text{-CO}$ dimers with CO_2 undergoing the hot band transition $(\nu_1, \nu_2^{l2}, \nu_3) = (01^11) \leftarrow (01^10)$. One source of confusion here was that isomer 2 was stronger than isomer 1, opposite to all our other observed spectra. This point is discussed below in Sec. 3. The other source of confusion was that the spectrum is strongly influenced by Coriolis interactions affecting its upper and lower vibrational states. The degenerate ν_2 bending mode of CO_2 splits into two components in the $\text{CO}_2\text{-CO}$ dimer, and these in-plane and out-of-plane components are mixed by the Coriolis interaction. Observations of this effect in weakly-bound van der Waals complexes are surprisingly rare, probably because the bending fundamental of small linear molecules like CO_2 tend to lie in the more difficult far infrared region. But notably one such investigation was made by Ohshima et al. [17], who studied the $\text{C}_2\text{H}_2\text{-Ar}$ dimer in the region of the C_2H_2 ν_5 bending vibration ($\approx 730\text{ cm}^{-1}$). We have now also observed the $(01^11) \leftarrow (01^10)$ hot band in $\text{CO}_2\text{-Ar}$ [18] and $\text{CO}_2\text{-N}_2$ [19], which are simpler than $\text{CO}_2\text{-CO}$ since there is only one isomer to deal with.

The in-plane (i-p) and out-of-plane (o-p) $\text{CO}_2\text{-CO}$ vibrations corresponding to our lower state (CO_2 (01^10)) have A_1 and B_1 symmetries, respectively, for both isomers. In the upper state, (01^11) , we multiply by B_2 (for the CO_2 ν_3 mode) and obtain B_2 and A_2 symmetries for the i-p and o-p modes, respectively. We expect b -type selection rules, just like the fundamental (Fig. 1). The

i-p component transition is $B_2 \leftarrow A_1$ with $K_a = \text{odd} \leftarrow \text{even}$ subbands, and the o-p component is $A_2 \leftarrow B_1$ with $K_a = \text{even} \leftarrow \text{odd}$ subbands. We also expect strong Coriolis mixing between the i-p and o-p modes characterized by a matrix element

$$\langle \text{i-p}, J, k | H | \text{o-p}, J, k \pm 1 \rangle = \frac{1}{2} \xi_b \times [J(J+1) - k(k \pm 1)]^{1/2},$$

where k is signed K_a , and ξ_b is the b -type Coriolis interaction parameter, related to the dimensionless Coriolis parameter ζ by the relation $\xi_b = 2B\zeta$, where B is the B rotational constant and ζ can take values between zero (no coupling) and unity (complete coupling).

The details of our analysis are given in Table 4, and illustrated by the simulated spectra in Fig. 5. The stronger isomer 2 spectrum is also the one most affected by Coriolis mixing. One result is that the prominent Q -branch of the i-p $K_a = 1 \leftarrow 0$ subband ($\approx 2337.4 \text{ cm}^{-1}$) is compact and runs backwards, with higher J transitions at lower wavenumbers. In contrast, the Q -branch of the o-p $K_a = 0 \leftarrow 1$ subband ($\approx 2336.5 - 2336.8 \text{ cm}^{-1}$) runs in the normal direction and is quite spread out. It turns out that the i-p and o-p modes of isomer 2 are split by about 1.58 cm^{-1} , with i-p lower in energy, and that the Coriolis mixing is virtually complete ($\zeta \approx 1$). Once this is established, the spectrum can be very well fitted with rotational constants that are similar to those already known for isomer 2. We assigned 101 lines (109 transitions) in all subbands from $K_a = 2 \leftarrow 3$, to $3 \leftarrow 2$ (recall that odd \leftarrow even subbands are in-plane and even \leftarrow odd are out-of-plane). The rms residual was 0.00036 cm^{-1} . In addition to the expected transitions within the i-p or o-p stacks, we observed many ‘forbidden’ transitions between stacks which become allowed due to the Coriolis mixing. They follow selection rules $\Delta K_a = 0, \pm 2$, $\Delta K_c = 0, \pm 2$, and help to accurately determine the i-p to o-p splitting.

The isomer 1 spectrum is weaker, but more ‘normal’ in structure, as shown for example by the Q -branch of the i-p $K_a = 1 \leftarrow 0$ subband at $2337.2 - 2337.3 \text{ cm}^{-1}$. The o-p component was difficult to assign because of its weakness. This weakness and the ‘normal’ band structure are explained to the fact that the splitting between i-p and o-p components, about 4.6 cm^{-1} , is much larger here than for isomer 2. This means that there is less population in the o-p mode (hence the weakness) and less Coriolis interaction between modes (hence the normality). We assigned 77 lines (85 transitions) in all subbands from $K_a = 2 \leftarrow 3$ to $3 \leftarrow 2$, and the rms residual of the fit was 0.00034 cm^{-1} . But only 18 of these transitions were for the weak o-p mode, and hence it was necessary to fix the B and C parameters for this mode at ground state values. Alternatively, a similar fit could be obtained by varying B and C and fixing the Coriolis parameters, ξ_b , but we think the result given in Table 4 is more realistic. ‘Forbidden’ transitions between i-p and o-p levels were not observed for isomer 1 since the Coriolis mixing is less strong than for isomer 2, and as a result the splitting of the two modes is less precisely determined in this case.

The lower state of the hot band transitions corresponds to the first excited CO_2 bending state, $(v_1, v_2^{l2}, v_3) = (01^10)$. This is located at 667.380 cm^{-1} for the free CO_2 molecule, but the present spectra do not tell us how much it is shifted in $\text{CO}_2\text{-CO}$. In Table 4, we call these unknown shifts X and Y for isomers 1 and 2, respectively. They are likely to be smaller than a few cm^{-1} , and direct observation of the v_2 fundamental in the 667 cm^{-1} region will be required to determine them. However, the present spectra *do* tell us how much the $(01^11) \leftarrow (01^10)$ hot band origins of the dimer shift relative to free CO_2 . These shifts are equal to $+0.239$ and $+0.469 \text{ cm}^{-1}$ for the i-p modes of isomers 1 and 2 respectively. The o-p mode shifts are very similar, since the i-p to o-p splittings are similar in the upper and lower states, (01^11) and (01^10) . These hot band shifts are similar to, but slightly larger than, those for the fundamental, which are $+0.211$ and

+0.441 cm⁻¹. Table 5 gives a summary of band origin shifts determined here; note that the current shifts in the CO₂ ν_3 region are considerably smaller in magnitude than those in the CO stretch region [4,5].

3. Discussion and conclusions

The observed splitting of the CO₂ ν_2 mode into i-p and o-p components is very different for the two CO₂-CO isomers: 4.56 cm⁻¹ for isomer 1 (C-bonded) and 1.59 cm⁻¹ for isomer 2 (O-bonded). This difference could perhaps be rationalized by noting that isomer 1 is more strongly bound than isomer 2 (calculated zero point binding energies are 287 and 222 cm⁻¹, respectively [12]). But this simple explanation is somewhat undermined by the fact that the CO₂ ν_3 vibrational shift is actually larger for isomer 2 than for isomer 1 (Table 5). For comparison, in the case of CO₂-Ar the ν_2 splitting is 0.877 cm⁻¹ [18], less than those of either CO₂-CO isomer. In the case of CO₂-N₂, the ν_2 splitting is 2.307 cm⁻¹ [19], intermediate between the two CO₂-CO isomers, as might be expected.

The ξ_b parameters (Table 4) give values around 1.00 to 1.02 for the dimensionless Coriolis parameter ζ , similar to those determined for CO₂-Ar and CO₂-N₂. The maximum value of ζ should normally be unity within the harmonic approximation, but in the analogous case of C₂H₂-Ar, a ζ -value of 1.52 was determined and this anomaly was ascribed to large amplitude intermolecular motions [17]. This difference between CO₂-CO (or CO₂-Ar) and C₂H₂-Ar is consistent with the fact that the latter is indeed very nonrigid, as shown by its very low energy intermolecular bending mode (6.2 cm⁻¹) [20] as compared to CO₂-CO (24.8 or 14.2 cm⁻¹).

CO₂-CO isomer 2 is about 65 cm⁻¹ less strongly bound than isomer 1, and was not detected in Refs. 1-3. Based on similar cases [14], we think that observation of isomer 2 requires

using a supersonic expansion with helium carrier gas and a very dilute CO/CO₂ mixture. As the expansion begins, some dimers are ‘trapped’ in the higher energy state and subsequent collisions with helium are inefficient at promoting relaxation into the lower energy state. Thus the exact relative strength of the isomer spectra probably depends on details of the gas mixture and supersonic expansion conditions. (There could also be modest differences in infrared transition probabilities.) For the fundamental bands observed here (Fig. 1), isomer 1 is approximately 3.5 times stronger than isomer 2, and in our previous study in the CO stretch region, isomer 1 was about twice as strong.

However, as we have seen, isomer 2 is actually stronger than isomer 1 in the (01¹1) ← (01¹0) spectrum of Fig. 5 (the exact ratio is a bit uncertain), even though the gas mixture and expansion conditions were similar to those of Fig. 1. How can we understand the prominence of isomer 2 in this hot band spectrum? We speculate that ‘hot’ dimers containing CO₂ in the (01¹0) state, formed early in the supersonic expansion and surviving until the end, are relatively more likely also to contain the ‘hot’ isomer 2. The reason could be that dimers which happen to undergo significant energy changing collisions which relax the CO₂ vibration from (01¹0) to (000) are also more likely to be relaxed from isomer 2 to isomer 1.

In conclusion, spectra of both C- and O-bonded isomers of the CO₂-CO van der Waals complex have been studied in the CO₂ ν_3 fundamental region (≈ 2350 cm⁻¹) using a tunable infrared OPO laser source to probe a pulsed slit jet supersonic expansion. Spectra were assigned for the normal isotopologue as well as for ¹³C¹⁶O₂-CO and ¹⁶O¹³C¹⁸O-CO, the latter being the first observation of an asymmetrically substituted form for which all values of K_a are allowed. In addition to the fundamental bands, combination bands involving the lowest in-plane intermolecular mode were also observed for isomer 1 (C-bonded) of CO₂-CO and ¹³CO₂-CO and

for isomer 2 of CO₂-CO. A weak CO₂-CO spectrum near 2337 cm⁻¹ was assigned to the CO₂ hot band transition $(\nu_1, \nu_2^{f2}, \nu_3) = (01^11) \leftarrow (01^10)$. Its analysis yielded the splitting of the degenerate CO₂ ν_2 bending vibration into in-plane and out-of-plane modes due to the presence of the nearby CO molecule, which turns out to have values of 4.56 and 1.59 cm⁻¹ for isomers 1 and 2, respectively.

Acknowledgements

The financial support of the Natural Sciences and Engineering Research Council of Canada is gratefully acknowledged.

References

- [1] A.C. Legon, A.P. Suckley, *J. Chem. Phys.* 91 (1989) 4440.
- [2] R.W. Randall, J.P.L. Summersgill, B.J. Howard, *J. Chem. Soc. Faraday Trans.* 86 (1990) 1943.
- [3] Y. Xu, A.R.W. McKellar, B.J. Howard, *J. Mol. Spectrosc.* 179 (1996) 345.
- [4] A.J. Barclay, S. Sheybani-Deloui, K.H. Michaelian, A.R.W. McKellar, N. Moazzen-Ahmadi, *Chem. Phys. Lett.* 651 (2016) 62-65.
- [5] S. Sheybani-Deloui, A.J. Barclay, K.H. Michaelian, A.R.W. McKellar, N. Moazzen-Ahmadi, *J. Chem. Phys.* 143 (2015) 121101.
- [6] V. Raducu, D. Jasmin, R. Dahoo, P. Brosset, B. Gauthier-Roy, L. Abouaf-Marguin, *J. Chem. Phys.* 102 (1995) 9235.
- [7] V. Raducu, B. Gauthier-Roy, R. Dahoo, L. Abouaf-Marguin, J. Langlet, J. Caillet, M. Allavena, *J. Chem. Phys.* 105 (1996) 10092.
- [8] J. Langlet, J. Caillet, M. Allavena, V. Raducu, B. Gauthier-Roy, R. Dahoo, L. Abouaf-Marguin, *J. Mol. Struct.* 484, (1999) 145.
- [9] Z.-H. Cao, Y.-X. Bu, *Chem. J. Chinese Universities* 19 (2003) 94.
- [10] M. Venayagamoorthy, T.A. Ford, *J. Mol. Struct: THEOCHEM* 717 (2005) 111.
- [11] A. Badri, L. Shirkov, N.-E. Jaidanea, M. Hochlaf, *Phys. Chem. Chem. Phys.* 21 (2019) 15871.
- [12] E. Castro-Juarez, X.-G. Wang, T. Carrington, Jr., E. Quintas-Sanchez, R. Dawes, *J. Chem. Phys.* 151 (2019) 084307.
- [13] Dehghany, M., Afshari, M., Abusara, Z., Van Eck, C., Moazzen-Ahmadi, N., *J. Mol. Spectrosc.* **2008**, 247, 123-127.

- [14] Moazzen-Ahmadi, N., McKellar, A.R.W., *Int. Rev. Phys. Chem.* **2013**, *32*, 611-650.
- [15] Rezaei, M., Sheybani-Deloui, S., Moazzen-Ahmadi, N., Michaelian, K.H., McKellar, A.R.W., *J. Phys. Chem. A* **2013**, *117*, 9612-9620.
- [16] Western, C.M. PGOPHER, a program for simulating rotational structure version 8.0, 2014, University of Bristol Research Data Repository, [doi:10.5523/bris.huflggvpcuc1zvliqed497r2](https://doi.org/10.5523/bris.huflggvpcuc1zvliqed497r2)
- [17] Y. Ohshima, Y. Matsumoto, M. Takami, and K. Kuchitsu, *J. Chem. Phys.* **99**, 8385 (1993).
- [18] T.A. Gartner, A.J. Barclay, A.R.W. McKellar, and N. Moazzen-Ahmadi, *Phys. Chem. Chem. Phys.* **22**, 21488-21493 (2020).
- [19] A.J. Barclay, A.R.W. McKellar, and N. Moazzen-Ahmadi, *J. Chem. Phys.* **153**, 014303 (2020).
- [20] R.J. Bemish, P.A. Block, L.G. Pedersen, W.T. Yang, and R.E. Miller, *J. Chem. Phys.* **99**, 8585 (1993).

Table 1. Molecular parameters for CO₂ - CO (in cm⁻¹).^a

	Isomer 1			Isomer 2		
	Ground state [3]	Excited state fundamental	Excited state <i>a</i> -type combination	Ground state [5]	Excited state fundamental	Excited state <i>a</i> -type combination
σ_0	0.0	2349.3539(1)	2373.8637(1)	0.0 ^b	2349.5845(1)	2363.9569(2)
<i>A</i>	[0.3956764]	0.392602(21)	0.392334(52)	0.397038(29)	0.393928(33)	0.356143(48)
<i>B</i>	[0.06281439]	0.0627934(28)	0.0635355(97)	0.074488(16)	0.074430(14)	0.076105(20)
<i>C</i>	[0.05383985]	0.0537620(19)	0.0538731(81)	0.062128(17)	0.061992(14)	0.062431(15)
$10^5 D_K$	[-1.167]	-0.921(87)	+1.89(31)	-2.42(14)	-2.42(14) ^c	
$10^5 D_{JK}$	[1.16239]	[1.16239]	[1.16239]	2.838(68)	2.838(68) ^c	
$10^7 D_J$	[2.2148]	[2.2148]	[2.2148]	4.5(15)	4.5(15) ^c	
$10^8 d_I$	[-4.280]	[-4.280]	[-4.280]			
$10^8 d_2$	[-2.512]	[-2.512]	[-2.512]			

^a Uncertainties (1σ) in parentheses are in units of the last quoted digit, and parameters in square brackets were fixed at the indicated values. Ground state parameters were fixed at previously determined values as shown.

^b The energy of isomer 2 relative to isomer 1 is calculated to be about 65 cm⁻¹ [11,12].

^c These excited state fundamental distortion parameters were constrained to equal those of the ground state.

Table 2. Molecular parameters for $^{13}\text{CO}_2 - \text{CO}$ (in cm^{-1}).^a

	Isomer 1			Isomer 2	
	Ground state	Excited state fundamental	Excited state α -type combination	Ground state	Excited state fundamental
σ_0	0.0	2283.6863(1)	2308.1876(2)	0.0	2283.8997(3)
A	0.395634(30)	0.391640(31)	0.392031(45)	0.397024(88)	0.394025(84)
B	0.062297(10)	0.0621619(80)	0.063114(14)	0.073806(46)	0.073809(42)
C	0.053450(10)	0.0533350(85)	0.053438(16)	0.061795(40)	0.061413(45)
$10^5 D_K$	[-1.167]	[-1.167]	[-1.167]		
$10^5 D_{JK}$	1.088(85)	0.886(68)	1.21(13)	1.27(31)	1.83(33)
$10^7 D_J$	2.02(55)	1.49(62)	3.02(92)	13.7(39)	-11.5(64)
$10^8 d_I$	[-4.280]	[-4.280]	[-4.280]		
$10^8 d_2$	[-2.512]	[-2.512]	[-2.512]		

^a Uncertainties (1σ) in parentheses are in units of the last quoted digit, and parameters in square brackets were fixed at the indicated values. The isomer 1 parameters D_K , d_I , and d_2 were fixed at their $^{12}\text{CO}_2 - \text{CO}$ values (Table 1).

Table 3. Molecular parameters for $^{16}\text{O}^{13}\text{C}^{18}\text{O}$ - CO (in cm^{-1}).^a

	Isomer 1		Isomer 2	
	Ground state	Excited state fundamental	Ground state	Excited state fundamental
σ_0	0.0	2266.1751(2)	0.0	2266.3984(2)
A	0.37312(21)	0.370321(82)	0.37465(13)	0.371852(65)
B	0.061271(24)	0.061256(19)	0.072760(26)	0.072681(34)
C	0.052307(22)	0.052217(21)	0.060341(25)	0.060129(29)
$10^5 D_K$	-3.8(44)	-1.95(89)		
$10^5 D_{JK}$	7.2(28)	0.94(15)	2.58(63)	2.74(26)
$10^7 D_J$	2.0(26)	9.(22)	9.5(27)	-3.4(42)

^a Uncertainties (1σ) in parentheses are in units of the last quoted digit.

Table 4. Molecular parameters for the (01¹1)-(01¹0) hot band of CO₂ - CO (in cm⁻¹).^a

	(01 ¹ 0) i-p	(01 ¹ 0) o-p	(01 ¹ 1) i-p	(01 ¹ 1) o-p
Isomer 1				
σ_0	X	4.557(40)+X	2336.8720(1)+X	2341.442(40)+X
<i>A</i>	0.396583(42)	0.395515(32)	0.393387(30)	0.392398(68)
<i>B</i>	0.062855(32)	[0.06281] ^b	0.062689(40)	[0.06281] ^b
<i>C</i>	0.053910(17)	[0.05384] ^b	0.0538513(65)	[0.05384] ^b
ξ_b	0.12770(94)		0.12512(100)	
Isomer 2				
σ_0	Y	1.5856(4)+Y	2337.1023(2)+Y	2338.6802(4)+Y
<i>A</i>	0.397497(48)	0.402802(39)	0.394458(44)	0.393474(75)
<i>B</i>	0.074213(70)	0.074553(45)	0.073822(12)	0.074864(31)
<i>C</i>	0.062074(34)	0.062046(17)	0.0620012(67)	0.062095(16)
$10^5 D_{JK}$	1.54(25)		2.13(24)	4.78(44)
ξ_b	0.15044(19)		0.148846(60)	

^a Uncertainties (1 σ) in parentheses are in units of the last quoted digit. Centrifugal distortion parameters were fixed to their ground state values from Table 1 except as shown. σ_0 is the term value (vibrational energy) of the state. X and Y are equal to the free CO₂ ν_2 frequency (667.380 cm⁻¹) plus or minus unknown vibrational shifts which are unlikely to be more than a few cm⁻¹. i-p = in-plane; o-p = out-of-plane.

^b *B* and *C* parameters for the weak isomer 1 o-p band were fixed at these ground state values.

Table 5. Vibrational shifts of CO₂-CO relative to the free CO₂ molecule (in cm⁻¹).

	Isomer 1	Isomer 2
CO ₂ -CO, CO stretch region [4,5]	+4.970	-2.982
CO ₂ -CO, CO ₂ ν_3 fundamental	+0.211	+0.411
CO ₂ -CO, CO ₂ ν_3 hot band ^a	+0.239	+0.469
¹³ CO ₂ -CO, CO ₂ ν_3 fundamental	+0.199	+0.412
¹⁶ O ¹³ C ¹⁸ O-CO, CO ₂ ν_3 fundamental	+0.204	+0.427

^a These are for the i-p modes; the o-p values are similar.

Figure Captions

- Fig. 1. Observed and simulated spectra of CO₂-CO in the region of the CO₂ ν_3 fundamental band. Simulated spectra are shown for the two isomers of CO₂-CO and for CO₂ dimer, with an effective rotational temperature of 2.2 K. The strongest lines belong to the $K_a = 1 \leftarrow 0$ Q -branch of isomer 1 at 2349.7 cm⁻¹. Here and in the other spectra shown, blank regions correspond to known lines of CO₂ monomer and CO₂-He dimer. Observed peaks at 2349.17 and 2349.47 cm⁻¹ are believed to be due to larger clusters.
- Fig. 2. Observed and simulated spectra of ¹³CO₂-CO in the region of the CO₂ ν_3 fundamental band, recorded using an enriched ¹³CO₂ sample.
- Fig. 3. Observed and simulated spectra of ¹⁶O¹³C¹⁸O-CO in the region of the CO₂ ν_3 fundamental band, recorded using an enriched ¹³CO₂ sample. The spectrum has a different appearance than those in Figs. 1 and 2 because all values of K_a are allowed in the ground and excited states of this asymmetrically substituted isotopologue.
- Fig. 4. Observed and simulated spectra of isomer 1 of CO₂-CO and ¹³CO₂-CO in the region of the combination band involving the lowest in-plane intermolecular bending mode. Note their almost identical appearances. The central peak (2373.85 cm⁻¹) in the top panel is the $K_a = 2 \leftarrow 2$ Q -branch, and the small peak to its left (2373.80 cm⁻¹) is the $K_a = 4 \leftarrow 4$ Q -branch. The ¹³CO₂-CO $K_a = 2 \leftarrow 2$ Q -branch in the lower panel is obscured by a CO₂ monomer line.
- Fig. 5. Observed and simulated spectra of CO₂-CO corresponding to the $(\nu_1, \nu_2^{l2}, \nu_3) = (01^11) \leftarrow (01^10)$ hot band of CO₂. These bands are strongly affected by Coriolis interaction between the in-plane and out-of-plane components of the CO₂ ν_2 bend, especially for isomer 2. The

out-of-plane component of isomer 1 is very weak because it lies 4.5 cm^{-1} above the in-plane component and thus has little population.




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5