4-Adic Complexity of Interleaved Quaternary Sequences

Shiyuan Qiang, Xiaoyan Jing, Minghui Yang, and Keqin Feng

Abstract

Tang and Ding [11] present a series of quaternary sequences w(a,b) interleaved by two binary sequences a and b with ideal autocorrelation and show that such interleaved quaternary sequences have optimal autocorrelation. In this paper we consider the 4-adic complexity $FC_w(4)$ of such quaternary sequence w = w(a,b). We present a general formula on $FC_w(4)$, w = w(a,b). As a direct consequence, we obtain a general lower bound $FC_w(4) \ge \log_4(4^n - 1)$ where 2n is the period of the sequence w. By taking a and b to be several types of known binary sequences with ideal autocorrelation (*m*-sequences, twin-prime, Legendre, Hall sequences and their complement, shift or sample sequences), we compute the exact values of $FC_w(4)$, w = w(a,b) and show that in most cases $FC_w(4)$ reaches or nearly reaches the maximum value $\log_4(4^{2n} - 1)$. Our results show that the 4-adic complexity of the quaternary sequences defined in [11] are large enough to resist the attack of the rational approximation algorithm.

Index Terms

quaternary sequences, 4-adic complexity, interleaved sequences, ideal autocorrelation

I. INTRODUCTION

Binary and quaternary sequences with good autocorrelation have been widely used in digital communication and cryptography ([1], [2]). For a binary sequence $s = \{s_{\lambda}\}_{\lambda=0}^{n-1}$ with period $n, s_{\lambda} \in \{0, 1\}$, the

Keqin Feng was supported by the National Natural Science Foundation of China (NSFC) under Grant 12031011.

Shiyuan Qiang is with the Department of Applied Mathematics, China Agricultural University, Beijing 100083, China (e-mail: qsycau_18@163.com).

Xiaoyan Jing is with School of Mathematics, Northwest University, Xi'an 710127, China (e-mail: jxymg@126.com).

Minghui Yang is with State Key Laboratory of Information Security, Institute of Information Engineering, Chinese Academy of Sciences, Beijing 100093, China (e-mail: yangminghui6688@163.com).

Keqin Feng is with the Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China (e-mail: fengkq@tsinghua.edu.cn).

autocorrelation function of s is defined by

$$A_s(\tau) = \sum_{\lambda=0}^{n-1} (-1)^{s_{\lambda}+s_{\lambda+\tau}} \in \mathbb{Z} \ (0 \le \tau \le n-1),$$

 $A_s(0) = n$ and in many applications we need that the absolute values $|A_s(\tau)|$ $(1 \le \tau \le n-1)$ are small. It is easy to see that $A_s(\tau) \equiv n \pmod{4}$ for all $0 \le \tau \le n-1$. We call a binary sequence *s* with ideal autocorrelation if $n \equiv 3 \pmod{4}$ and $A_s(\tau) = -1$ for all $1 \le \tau \le n-1$. Many series of binary sequences with ideal autocorrelation have been found (see [11], Section (II, *C*)). For all known binary sequences with ideal autocorrelation, their period *n* is $2^m - 1 \pmod{2}$, prime number *p* or product of twin primes *p* and *p*+2.

For a quaternary sequence $s = \{s_{\lambda}\}_{\lambda=0}^{n-1}$ with period $n, s_{\lambda} \in \{0, 1, 2, 3\}$, the autocorrelation function of s is defined by

$$A_s(\tau) = \sum_{\lambda=0}^{n-1} \omega^{s_{\lambda+\tau}-s_{\lambda}} \in \mathbb{Z}[\sqrt{-1}],$$

where $\omega = \sqrt{-1}$ is a complex primitive 4-th root of unity. In many applications we need the absolute values $|A_s(\tau)|$ for all out-of-phase autocorrelation $A_s(\tau)$ $(1 \le \tau \le n-1)$ are small.

The feedback with carry shift register (FCSR) is a kind of nonlinear sequences generator. The *d*-adic complexity $FC_s(d)$ measures the smallest length of FCSR which generates the sequence *s* over $\mathbb{Z}/(d)$. In cryptographic applications, sequences should not only have low autocorrelation values, but also have larger 4-adic complexity. Particularly, a quaternary sequence *s* with period *N* can be generated by a feedback shift register with 4-adic carry and length at least $[FC_s(4)]$, and the 4-adic complexity of such sequences should exceed $\frac{N-16}{4}$ in order to resist the attack of the rational approximation algorithm, see [6], [7].

Definition 1. For a quaternary sequence $s = \{s_{\lambda}\}_{\lambda=0}^{n-1}$ with period $n, s_{\lambda} \in \{0, 1, 2, 3\}$, the 4-adic complexity of s is defined by

$$FC_s(4) = \log_4 \frac{4^n - 1}{d},$$

where $d = \operatorname{gcd}(S(4), 4^n - 1)$, $S(4) = \sum_{\lambda=0}^{n-1} s_{\lambda} 4^{\lambda} \in \mathbb{Z}$.

Several quaternary sequences with good autocorrelation have been constructed ([4], [5], [8], [10]). Tang and Ding [11] construct a series of quaternary sequences with period 2n and optimal autocorrelation by interleaving a pair of binary sequences with ideal autocorrelation and period $n \equiv 3 \pmod{4}$. In this paper we compute the 4-adic complexity of such quaternary sequences.

In Section II we introduce the interleaving construction of quaternary sequence with period 2n by a pair of binary sequences with the same odd period *n*, compute the value of $S(4) \pmod{4^{2n}-1}$ for

such quaternary sequences, and present a general result on the 4-adic complexity of quaternary sequences interleaved by a pair of binary sequences with ideal autocorrelation. In Section III we determine the exact value of the 4-adic complexity for quaternary sequences interleaved by some particular types of binary sequences with ideal autocorrelation (*m*-sequences, Legendre sequences, twin-prime sequences and Hall sequences). The last section is a conclusion.

II. INTERLEAVED QUATERNARY SEQUENCES

In this section we introduce the interleaving construction of quaternary sequences with period 2n by a pair of binary sequences with the same odd period n. For general theory and applications of interleaved sequences we refer to Gong [3]. In this paper we use the following simple form.

Let *n* be an odd integer, $n \ge 3$, $a = \{a_{\lambda}\}_{\lambda=0}^{n-1}$ and $b = \{b_{\lambda}\}_{\lambda=0}^{n-1}$ be a pair of binary sequences, $a_{\lambda}, b_{\lambda} \in \{0,1\}$, with the same period *n*. Consider the following two $n \times 2$ matrices over $\mathbb{F}_2 = \{0,1\}$ (1+1=0)

$$M_{a} = \begin{pmatrix} a_{0} & a_{\frac{n+1}{2}} \\ a_{1} & a_{\frac{n+1}{2}+1} \\ \vdots & \vdots \\ a_{n-1} & a_{\frac{n+1}{2}+n-1} \end{pmatrix}, \quad M_{b}' = \begin{pmatrix} b_{0} & 1-b_{\frac{n+1}{2}} \\ b_{1} & 1-b_{\frac{n+1}{2}+1} \\ \vdots & \vdots \\ b_{n-1} & 1-b_{\frac{n+1}{2}+n-1} \end{pmatrix}$$

Then we get an interleaved binary sequence with period 2n from the matrix M_a

$$c = \{c_i\}_{i=0}^{2n-1}$$

= $(c_0, c_1, c_2, c_3, \cdots, c_{2n-2}, c_{2n-1})$
= $(a_0, a_{\frac{n+1}{2}}, a_1, a_{\frac{n+1}{2}+1}, \cdots, a_{n-1}, a_{\frac{n+1}{2}+n-1}(=a_{\frac{n-1}{2}})).$ (1)

By the Chinese Remainder Theorem we have the isomorphism of rings

$$f: \mathbb{Z}_{2n} \cong \mathbb{Z}_2 \oplus \mathbb{Z}_n, \quad i(\mod 2n) \mapsto (i(\mod 2), i(\mod n)).$$

For $(\mu, \lambda) \in \mathbb{Z}_2 \times \mathbb{Z}_n$, we have $f^{-1}((\mu, \lambda)) = n\mu + (1 - n)\lambda$ $(0 \le \mu \le 1, 0 \le \lambda \le n - 1)$. It is easy to see from the definition (1) of the sequence *c* that for $i = n\mu + (1 - n)\lambda$,

$$c_{i} = \begin{cases} a_{\frac{i}{2}} = a_{\lambda(\frac{1-n}{2})}, & \text{if } 2 \mid i \pmod{\mu} = 0; \\ a_{\frac{n+1}{2} + \frac{i-1}{2}} = a_{\lambda(\frac{1-n}{2})}, & \text{if } 2 \nmid i \pmod{\mu} = 1. \end{cases}$$

(Since for $\mu = 1$, $\frac{n+1}{2} + \frac{i-1}{2} = \frac{n+1}{2} + \frac{n+(1-n)\lambda-1}{2} = n + \lambda(\frac{1-n}{2}) \equiv \lambda(\frac{1-n}{2}) \pmod{n}$).

Similarly we have another interleaved binary sequence with period 2n from the matrix M'_b

$$d = \{d_i\}_{i=0}^{2n-1}$$

= $(d_0, d_1, d_2, d_3, \cdots, d_{2n-2}, d_{2n-1})$

$$=(b_0,1-b_{\frac{n+1}{2}},b_1,1-b_{\frac{n+1}{2}+1},\cdots,b_{n-1},1-b_{\frac{n+1}{2}+n-1}(=1+b_{\frac{n-1}{2}})).$$

Namely, for $i = n\mu + (1 - n)\lambda \pmod{2n}$, $0 \le \mu \le 1$, $0 \le \lambda \le n - 1$, we have

$$d_{i} = \begin{cases} b_{\lambda(\frac{1-n}{2})}, & \text{if } \mu = 0; \\ 1 - b_{\lambda(\frac{1-n}{2})}, & \text{if } \mu = 1. \end{cases}$$

Finally, we use the Gray mapping

$$\phi:\mathbb{Z}_2\oplus\mathbb{Z}_2\cong\mathbb{Z}_4,$$

$$\phi(0,0) = 0, \ \phi(0,1) = 1, \ \phi(1,1) = 2, \ \phi(1,0) = 3$$

to construct a quaternary sequence with period 2n

$$w = w(a,b) = \{w_i\}_{i=0}^{2n-1}, \quad w_i = \phi(c_i, d_i).$$
⁽²⁾

Tang and Ding [11] proved that for any binary sequences a and b with ideal autocorrelation and the same odd period n, the interleaved quaternary sequence w = w(a,b) has optimal autocorrelation $A_w(\tau) \in \{0,-2\}$ for all $1 \le \tau \le 2n-1$. Let $W(4) = \sum_{i=0}^{2n-1} w_i 4^i$. In the remaining part of this section we present a general formula on the value $W(4) \pmod{4^{2n}-1}$ for any binary sequences a and b. Then we restrict to binary sequences a and b having ideal autocorrelation and give a general result on the 4-adic complexity of the quaternary sequences w = w(a,b).

Lemma 1. Let $n \ge 3$ be an odd integer, $a = \{a_{\lambda}\}_{\lambda=0}^{n-1}$ and $b = \{b_{\lambda}\}_{\lambda=0}^{n-1}$ be binary sequences with period $n, a_{\lambda}, b_{\lambda} \in \mathbb{F}_2 = \{0, 1\}, w = w(a, b) = \{w_i\}_{i=0}^{2n-1}, w_i \in \{0, 1, 2, 3\}$ be the interleaved quaternary sequence given by (2), $W(4) = \sum_{i=0}^{2n-1} w_i 4^i \in \mathbb{Z}$. Then

$$\begin{split} W(4) &\equiv 2(4^{n}-1)\sum_{\lambda=0}^{n-1}a_{\lambda}b_{\lambda}4^{2\lambda} + (4^{n}+3)\sum_{\lambda=0}^{n-1}a_{\lambda}4^{2\lambda} + (1-4^{n})\sum_{\lambda=0}^{n-1}b_{\lambda}4^{2\lambda} + 4\cdot\frac{4^{2n}-1}{15} \pmod{4^{2n}-1} \\ &\equiv \begin{cases} -4\sum_{\lambda=0}^{n-1}a_{\lambda}b_{\lambda}4^{2\lambda} + 2\sum_{\lambda=0}^{n-1}(a_{\lambda}+b_{\lambda})4^{2\lambda} - \frac{4^{n}+1}{5} \pmod{4^{n}+1} \\ 4\sum_{\lambda=0}^{n-1}a_{\lambda}4^{2\lambda} + \frac{4^{n}-1}{3} \pmod{4^{n}-1}. \end{cases} \end{split}$$

Proof. From the definition (2) of w = w(a,b),

$$W(4) = \sum_{i=0}^{2n-1} w_i 4^i \equiv \sum_{\lambda=0}^{n-1} \sum_{\mu=0}^{1} \phi(\mu, \lambda) 4^{n\mu+(1-n)\lambda} \pmod{4^{2n}-1}$$
$$\equiv \sum_{\lambda=0}^{n-1} \sum_{\mu=0}^{1} 4^{n\mu+(1-n)\lambda} \left(\sum_{\substack{c_i=0\\d_i=1}} 1 + 2\sum_{\substack{c_i=1\\d_i=1}} 1 + 3\sum_{\substack{c_i=1\\d_i=0}} 1 \right) \pmod{4^{2n}-1}$$
$$(\text{where } i = n\mu + (1-n)\lambda)$$
$$\equiv \sum_{\lambda=0}^{n-1} \sum_{\mu=0}^{1} 4^{n\mu+(1-n)\lambda} \left(2\sum_{\substack{c_i=1\\c_i=1}} 1 + \sum_{\substack{c_i=1\\d_i=1}} 1 \right) \pmod{4^{2n}-1}$$

From *n* is odd and $4^n \equiv 4 \pmod{15}$, we have

$$4^{n} \sum_{\lambda=0}^{n-1} 4^{2\lambda} \equiv 4^{n} \cdot \frac{4^{2n} - 1}{15} \equiv 4 \cdot \frac{4^{2n} - 1}{15} \pmod{4^{2n} - 1}$$

and

$$\begin{split} W(4) &\equiv 2(4^{n}-1)\sum_{\lambda=0}^{n-1}a_{\lambda}b_{\lambda}4^{2\lambda} + (4^{n}+3)\sum_{\lambda=0}^{n-1}a_{\lambda}4^{2\lambda} + (1-4^{n})\sum_{\lambda=0}^{n-1}b_{\lambda}4^{2\lambda} + 4\cdot\frac{4^{2n}-1}{15} \pmod{4^{2n}-1} \\ &\equiv \begin{cases} -4\sum_{\lambda=0}^{n-1}a_{\lambda}b_{\lambda}4^{2\lambda} + 2\sum_{\lambda=0}^{n-1}(a_{\lambda}+b_{\lambda})4^{2\lambda} + 4\cdot\frac{4^{2n}-1}{15} \pmod{4^{n}+1} \\ 4\sum_{\lambda=0}^{n-1}a_{\lambda}4^{2\lambda} + 4\cdot\frac{4^{2n}-1}{15} \pmod{4^{n}-1}. \end{cases}$$

From $4^n \equiv 4 \pmod{5}$ we have $4 \cdot \frac{4^n - 1}{3} \equiv -1 \pmod{5}$. Then we obtain

$$4 \cdot \frac{4^{2n} - 1}{15} = \frac{4(4^n - 1)}{3} \cdot \frac{4^n + 1}{5} \equiv -\frac{4^n + 1}{5} \pmod{4^n + 1}.$$

By $4^n \equiv 1 \pmod{3}$ we know $4 \cdot \frac{4^n+1}{5} \equiv 1 \pmod{3}$. Then we have

$$4 \cdot \frac{4^{2n} - 1}{15} = \frac{4(4^n + 1)}{5} \cdot \frac{4^n - 1}{3} \equiv \frac{4^n - 1}{3} \pmod{4^n - 1}.$$

This completes the proof of Lemma 1.

Remark 1. From Lemma 1 we know that $W(4) \pmod{4^n-1}$ is independent of the sequence b.

Let $s_{\lambda} = (-1)^{a_{\lambda}}$, $t_{\lambda} = (-1)^{b_{\lambda}}$. The value $W(4) = \sum_{\lambda=0}^{2n-1} w_{\lambda} 4^{\lambda} \pmod{4^{2n}-1}$ can be expressed in terms of the $\{\pm 1\}$ -sequences $\{s_{\lambda}\}_{\lambda=0}^{n-1}$ and $\{t_{\lambda}\}_{\lambda=0}^{n-1}$.

Corollary 2. Let $a = \{a_{\lambda}\}_{\lambda=0}^{n-1}$ and $b = \{b_{\lambda}\}_{\lambda=0}^{n-1}$ be binary sequences with odd period $n, a_{\lambda}, b_{\lambda} \in \{0, 1\}, s_{\lambda} = (-1)^{a_{\lambda}}, t_{\lambda} = (-1)^{b_{\lambda}}$. Then for quaternary sequence $w = w(a,b) = \{w_i\}_{i=0}^{2n-1}, w_i \in \{0,1,2,3\}, W(4) = \sum_{i=0}^{2n-1} w_i 4^i \in \mathbb{Z}$, we have

$$W(4) \equiv \begin{cases} -\sum_{\lambda=0}^{n-1} s_{\lambda} t_{\lambda} 4^{2\lambda} \pmod{4^n + 1} \\ -2\sum_{\lambda=0}^{n-1} s_{\lambda} 4^{2\lambda} \pmod{4^n - 1}. \end{cases}$$

Proof. We have $a_{\lambda} = \frac{1}{2}(1 - s_{\lambda})$, $b_{\lambda} = \frac{1}{2}(1 - t_{\lambda})$. Then by Lemma 1,

$$W(4) \equiv -4\sum_{\lambda=0}^{n-1} a_{\lambda} b_{\lambda} 4^{2\lambda} + 2\sum_{\lambda=0}^{n-1} (a_{\lambda} + b_{\lambda}) 4^{2\lambda} - \frac{4^{n} + 1}{5} \pmod{4^{n} + 1}$$
$$\equiv -\sum_{\lambda=0}^{n-1} (1 - s_{\lambda}) (1 - t_{\lambda}) 4^{2\lambda} + \sum_{\lambda=0}^{n-1} (2 - s_{\lambda} - t_{\lambda}) 4^{2\lambda} - \frac{4^{n} + 1}{5} \pmod{4^{n} + 1}$$
$$\equiv \sum_{\lambda=0}^{n-1} 4^{2\lambda} - \sum_{\lambda=0}^{n-1} s_{\lambda} t_{\lambda} 4^{2\lambda} - \frac{4^{n} + 1}{5} \pmod{4^{n} + 1}$$
$$\equiv \frac{4^{2n} - 1}{15} - \sum_{\lambda=0}^{n-1} s_{\lambda} t_{\lambda} 4^{2\lambda} - \frac{4^{n} + 1}{5} \equiv -\sum_{\lambda=0}^{n-1} s_{\lambda} t_{\lambda} 4^{2\lambda} \pmod{4^{n} + 1}$$

and

$$W(4) \equiv 4\sum_{\lambda=0}^{n-1} a_{\lambda} 4^{2\lambda} + \frac{4^{n} - 1}{3} \equiv 2\sum_{\lambda=0}^{n-1} (1 - s_{\lambda}) 4^{2\lambda} + \frac{4^{n} - 1}{3} \pmod{4^{n} - 1}$$
$$\equiv 2 \cdot \frac{4^{2n} - 1}{15} - 2\sum_{\lambda=0}^{n-1} s_{\lambda} 4^{2\lambda} + \frac{4^{n} - 1}{3} \equiv -2\sum_{\lambda=0}^{n-1} s_{\lambda} 4^{2\lambda} \pmod{4^{n} - 1}.$$

Now we present a general formula on the 4-adic complexity of the quaternary interleaved sequence $w = w(a,b) = \{w_i\}_{i=0}^{2n-1}$, where *a*, *b* are binary sequences with period $n (\equiv 3 \pmod{4})$ and ideal autocorrelation.

From now on we denote by $\Sigma(n)$ the set of binary sequences with period $n \equiv 3 \pmod{4}$ and ideal autocorrelation. By the definition of ideal autocorrelation, for $a = \{a_{\lambda}\}_{\lambda=0}^{n-1} \in \Sigma(n), a_{\lambda} \in \{0,1\}, s_{\lambda} = (-1)^{a_{\lambda}} \in \{\pm 1\}$, we have

$$A_{a}(\tau) = \sum_{\lambda=0}^{n-1} (-1)^{a_{\lambda}+a_{\lambda+\tau}} = \sum_{\lambda=0}^{n-1} s_{\lambda} s_{\lambda+\tau} = -1 \ (1 \le \tau \le n-1).$$

Theorem 3. Let $n \equiv 3 \pmod{4}$, $a = \{a_{\lambda}\}_{\lambda=0}^{n-1}$ and $b = \{b_{\lambda}\}_{\lambda=0}^{n-1}$ be known binary sequences in $\sum(n)$, $s_{\lambda} = (-1)^{a_{\lambda}}$, $t_{\lambda} = (-1)^{b_{\lambda}}$. Then the 4-adic complexity of the quaternary interleaved sequence w = w(a,b) is

$$FC_w(4) = \log_4\left((4^n - 1) \cdot \frac{4^n + 1}{d_+}\right)$$

where $d_+ = \operatorname{gcd}\left(\sum_{\lambda=0}^{n-1} s_{\lambda} t_{\lambda} 4^{2\lambda}, 4^n + 1\right)$.

Before proving Theorem 3, we need the following result.

Lemma 4. Let $n \equiv 3 \pmod{4}$ be the period of known binary sequences in $\sum(n)$, n > 3. Namely, $n = 2^m - 1$ $(m \ge 2)$, n is a prime, and n = pq where p and q = p + 2 are prime numbers. Then n + 1 and $4^n - 1$ have no common prime divisor $\pi > 3$.

Proof. (A) For $n = 2^m - 1$ $(m \ge 2)$, we have $gcd(n + 1, 4^n - 1) = gcd(2^m, 4^n - 1) = 1$.

(B) Let n = p be a prime, $p \equiv 3 \pmod{4}$, p > 3. Suppose that p + 1 and $4^p - 1$ have a common (odd) prime divisor $\pi > 3$. Then $\pi \ge 7$, $\pi \mid p+1$ so that $\pi \le \frac{p+1}{4}$. From $4^p \equiv 1 \pmod{\pi}$ and $\pi \ne 3$ we know that the order of 4 (mod π) is p. Therefore $p \mid \pi - 1$ by Fermat's Theorem which implies $p \le (\pi - 1)$. Then we get a contradiction $4\pi \le p+1 \le 1 + (\pi - 1) = \pi$.

(C) Let n = p(p+2), a product of twin-primes. Suppose that $n+1 = (p+1)^2$ and $4^n - 1$ have a common prime divisor $\pi > 3$. Then $4^{p(p+2)} \equiv 1 \pmod{\pi}$. From $\pi \neq 3$ we know that the order r of 4 (mod π) is at least p. We obtain $p \le r \mid \frac{\pi-1}{2}$. From $\pi \mid n+1 = (p+1)^2$ we have $\pi \mid p+1$ and $\pi \le p+1$. Then we obtain a contradiction $p \le \frac{\pi-1}{2} \le \frac{p+1-1}{2} = \frac{p}{2}$. This completes the proof of Lemma 4.

Proof of Theorem 3 By Definition 1, the 4-adic complexity of w = w(a,b) is

$$FC_w(4) = \log_4\left(\frac{4^{2n}-1}{d}\right), \ d = \gcd(W(4), 4^{2n}-1), \ W(4) = \sum_{i=0}^{2n-1} w_i 4^i \in \mathbb{Z}$$

Let

$$d_{+} = \gcd(W(4), 4^{n} + 1), \ d_{-} = \gcd(W(4), 4^{n} - 1).$$

From $gcd(4^n+1,4^n-1)=1$ we know that $gcd(d_+,d_-)=1$ and $d=d_+d_-$.

Firstly we show $d_{-} = 1$. Suppose that $d_{-} = \gcd(W(4), 4^{n} - 1) > 1$. Let π be a common (odd) prime divisor of W(4) and $4^{n} - 1$. If $\pi = 3$, by Corollary 2, $0 \equiv W(4) \equiv -2\sum_{\lambda=0}^{n-1} s_{\lambda} 4^{2\lambda} \equiv \sum_{\lambda=0}^{n-1} s_{\lambda} \pmod{3}$ and then we obtain a contradiction

$$0 \equiv \left(\sum_{\lambda=0}^{n-1} s_{\lambda}\right)^{2} = \sum_{\lambda,\mu=0}^{n-1} s_{\lambda} s_{\mu} = \sum_{\tau=0}^{n-1} \sum_{\lambda=0}^{n-1} s_{\lambda} s_{\lambda+\tau} = n + \sum_{\tau=1}^{n-1} (-1) = 1 \pmod{3}.$$

Therefore $\pi \neq 3$. From $4^n - 1 = (-1)^n - 1 \equiv 3 \pmod{5}$ we know that $\pi \neq 5$. Therefore $\pi \geq 7$. By $0 \equiv W(4) \equiv -2\sum_{\lambda=0}^{n-1} s_{\lambda} 4^{2\lambda} \pmod{\pi}$ we have

$$0 \equiv \left(\sum_{\lambda=0}^{n-1} s_{\lambda} 4^{2\lambda}\right) \left(\sum_{\mu=0}^{n-1} s_{\mu} 4^{-2\mu}\right) \equiv \sum_{\lambda,\mu=0}^{n-1} s_{\lambda} s_{\mu} 4^{2(\lambda-\mu)} \pmod{\pi}$$

$$\equiv \sum_{\tau=0}^{n-1} 4^{2\tau} \sum_{\mu=0}^{n-1} s_{\mu+\tau} s_{\mu} = n + \sum_{\tau=1}^{n-1} (-4^{2\tau}) \equiv n - \left(\frac{4^{2n}-1}{15} - 1\right) \equiv n+1 \pmod{\pi} \quad (\text{since } \pi \ge 7)$$

This implies that $\pi | \gcd(n+1,4^n-1)$. But this contradicts to Lemma 4. Therefore $d_- = 1$ and $d = d_+ = \gcd(W(4),4^n+1)$. By Corollary 2, $W(4) \equiv -\sum_{\lambda=0}^{n-1} s_{\lambda} t_{\lambda} 4^{2\lambda} \pmod{4^n+1}$. We have $d_+ = \gcd(\sum_{\lambda=0}^{n-1} s_{\lambda} t_{\lambda} 4^{2\lambda}, 4^n+1)$. This completes the proof of Theorem 3.

Theorem 3 has the following consequences. Firstly, from $d_+ = \gcd(\sum_{\lambda=0}^{n-1} s_{\lambda} t_{\lambda} 4^{2\lambda}, 4^n + 1)$, $FC_w(4) = \log_4\left(\frac{4^{2n}-1}{d_+}\right)$ and $\sum_{\lambda=0}^{n-1} s_{\lambda} t_{\lambda} 4^{2\lambda} = \sum_{\lambda=0}^{n-1} t_{\lambda} s_{\lambda} 4^{2\lambda}$ we know that

Corollary 5. Let $a, b \in \Sigma(n)$. Then w(a, b) and w(b, a) have the same 4-adic complexity.

Next, we present a transformation group on the set $\Sigma(n)$.

Definition 2. For a binary sequence $a = \{a_i\}_{i=0}^{N-1}$ with period N, $a_i \in \{0,1\}$.

(1). The complement sequence of a is $C(a) = \overline{a} = \{\overline{a}_i\}_{i=0}^{N-1}$ where $\overline{a}_i = 1 - a_i \ (0 \le i \le N-1)$.

(2). For $e \in \mathbb{Z}_N$, the e-shift sequence of a is $L^e(a) = \{b_i\}_{i=0}^{N-1}$ where $b_i = L^e(a_i) = a_{i+e}$ $(0 \le i \le N-1)$.

(3). For $r \in \mathbb{Z}_N^*$, the r-sample sequence of a is $M_r(a) = \{c_i\}_{i=0}^{N-1}$ where $c_i = M_r(a_i) = a_{ri} \ (0 \le i \le N-1)$.

It is easy to see that the sequences C(a), $L^e(a)$ and $M_r(a)$ have the same period N, $C^2 = L^0 = M_1 = Id$, $L^{e_1}L^{e_2} = L^{e_1+e_2}$, $M_{r_1}M_{r_2} = M_{r_1r_2}$. Since

$$M_r L^e(a_i) = M_r(a_{i+e}) = a_{ri+re} = L^{re}(a_{ri}) = L^{re} M_r(a_i),$$

we have $M_r L^e = L^{re} M_r$. Therefore C, L^e $(e \in \mathbb{Z}_N)$ and M_r $(r \in \mathbb{Z}_N^*)$ generate a group G. Next consequence shows that G is a transformation group on the set $\sum(n)$, and the action of some transforms on related quaternary interleaved sequences keeps the 4-adic complexity.

Corollary 6. Let $a, b \in \Sigma(n)$. Then

(1). For any $\sigma \in G$, $\sigma(a) \in \Sigma(n)$.

(2). For $\sigma = L^e$, the quaternary sequences w(a,b) and $\sigma(w) = (\sigma(a), \sigma(b))$ have the same 4-adic complexity.

(3). w(a,b) and $w(\overline{a},b)$ have the same 4-adic complexity.

Proof. Since the group G is generated by C, L^e $(e \in \mathbb{Z}_n)$ and M_r $(r \in \mathbb{Z}_n^*)$. We need to prove (1) for $\sigma = C$, L^e and M_r . Let $a = \{a_i\}_{i=0}^{n-1}$, $b = \{b_i\}_{i=0}^{n-1}$. (1). For each $\tau, 0 \le \tau \le n-1$,

$$A_{\overline{a}}(\tau) = \sum_{i=0}^{n-1} (-1)^{\overline{a}_i + \overline{a}_{i+\tau}} = \sum_{i=0}^{n-1} (-1)^{a_i + 1 + a_{i+\tau} + 1} = \sum_{i=0}^{n-1} (-1)^{a_i + a_{i+\tau}} = A_a(\tau)$$

$$A_{L^{e}(a)}(\tau) = \sum_{i=0}^{n-1} (-1)^{a_{i+e}+a_{i+e+\tau}} = \sum_{j=0}^{n-1} (-1)^{a_{j}+a_{j+\tau}} = A_{a}(\tau).$$
$$A_{M_{r}(a)}(\tau) = \sum_{i=0}^{n-1} (-1)^{a_{ir}+a_{(i+\tau)r}} = \sum_{i=0}^{n-1} (-1)^{a_{ir}+a_{ir+\tau r}} = A_{a}(\tau r).$$

From these formulas and gcd(r,n) = 1 we know that if $a \in \sum(n)$, then $C(a) = \overline{a}$, $L^e(a)$ and $M_r(a)$ belong to $\sum(n)$.

(2) and (3). Let
$$s_i = (-1)^{a_i}, t_i = (-1)^{b_i}$$
, and for $\sigma \in G$, $\sigma(s_i) = (-1)^{\sigma(a_i)}$. Then we have

$$\sum_{i=0}^{n-1} L^{e}(s_{i})L^{e}(t_{i})4^{2i} = \sum_{i=0}^{n-1} s_{i+e}t_{i+e}4^{2i} \equiv 4^{-2e}\sum_{j=0}^{n-1} s_{j}t_{j}4^{2j} \pmod{4^{2n}-1}$$

and

$$\sum_{i=0}^{n-1} (-1)^{\overline{a}_i} t_i 4^{2i} = \sum_{i=0}^{n-1} (-s_i) t_i 4^{2i} = -\sum_{i=0}^{n-1} s_i t_i 4^{2i}$$

which imply (2) and (3) by Theorem 3.

Corollary 7. For $a \in \sum(n)$, w = w(a, a), we have $FC_w(4) = \log_4(5(4^n - 1))$.

Proof. Let $s_i = (-1)^{a_i}$. Then

$$\sum_{i=0}^{n-1} s_i s_i 4^{2i} = \sum_{i=0}^{n-1} 4^{2i} = \frac{4^{2n} - 1}{15} = \frac{4^n + 1}{5} \cdot \frac{4^n - 1}{3}.$$

From $\frac{4^n-1}{3} \equiv \frac{(-1)^n-1}{3} \equiv -2 \cdot 3^{-1} \equiv -4 \equiv 1 \pmod{5}$, we obtain

$$\sum_{i=0}^{n-1} s_i s_i 4^{2i} \equiv \frac{4^n + 1}{5} \pmod{4^n + 1}.$$

Therefore for w = w(a, a), $d_+ = \gcd(\sum_{i=0}^{n-1} s_i^2 4^{2i}, 4^n + 1) = \frac{4^n + 1}{5}$ and $FC_w(4) = \log_4(\frac{4^{2n} - 1}{d_+}) = \log_4(5(4^n - 1))$.

For arbitrary $a, b \in \sum(n)$, in order to determine the exact value of the 4-adic complexity of w(a,b), we need to compute $\sum_{i=0}^{n-1} s_i t_i 4^{2i} \pmod{4^n + 1}$ which is not easy in general case. In Section III, we will consider several particular series of binary sequences in $\sum(n)$ (*m*-sequences, Legendre, twin-prime and Hall sequences). For each *a* in such particular series, we determine the 4-adic complexity of w(b,c)where $b = \sigma(a), c = \tau(a)$ for $\sigma, \tau \in G$.

III. INTERLEAVING BY SEVERAL TYPES OF KNOWN SEQUENCES

In the last section we present a general formula on the 4-adic complexity $FC_w(4)$ of the quaternary sequences w = w(a,b) with period 2*n* interleaved by two binary sequences *a* and *b* in $\Sigma(n)$. Particularly we get a general lower bound $FC_w(4) \ge \log_4(4^n - 1)$. In this section we compute the exact value of $FC_w(4)$, w = w(a,b) by taking *a* and *b* to be some particular types of known binary sequences in $\Sigma(n)$. Namely,

we consider a and b as m-sequences, Legendre, twin-prime and Hall sequences and their complement, shift or sample sequences.

A. m-sequences

Let $q = 2^k$ $(k \ge 2)$, n = q - 1, θ be a primitive element of \mathbb{F}_q , $\mathbb{F}_q^* = \langle \theta \rangle$, and $T : \mathbb{F}_q \to \mathbb{F}_2$ be the trace mapping. The *m*-sequence with period *n* is the binary sequence $m = \{m_i\}_{i=0}^{n-1}$ defined by

$$m_i = T(\theta^i) \in \mathbb{F}_2 = \{0, 1\} \quad (0 \le i \le n - 1)$$

It is well-known that $m \in \Sigma(n)$.

Theorem 8. Let $m = \{m_i\}_{i=0}^{n-1}$ be the m-sequence with period $n = 2^k - 1$ $(k \ge 2), 1 \le l \le n-1, w = w(m, L^l(m))$. Then $FC_w(4)$ reaches the maximum value $\log_4(4^{2n} - 1)$.

Proof. From Theorem 3 we know that

$$FC_w(4) = \log_4\left(\frac{4^{2n}-1}{d_+}\right), \ d_+ = \gcd(M, 4^n+1),$$

where

$$M = \sum_{\lambda=0}^{n-1} (-1)^{T(\theta^{\lambda}) + T(\theta^{\lambda+l})} 4^{2\lambda} = \sum_{\lambda=0}^{n-1} (-1)^{T(\theta^{\lambda}(1+\theta^{l}))} 4^{2\lambda}$$

Since the order of θ is *n* and $1 \le l \le n-1$, we know that $\theta^l + 1 \ne 0$ and $\theta^l + 1 = \theta^r$ for some $r \in \mathbb{Z}$. Then

$$M = \sum_{\lambda=0}^{n-1} (-1)^{T(\theta^{\lambda+r})} 4^{2\lambda} \equiv 4^{-2r} \sum_{\lambda=0}^{n-1} (-1)^{T(\theta^{\lambda})} 4^{2\lambda} \pmod{4^n + 1}.$$
(3)

The following result shows that $R = \sum_{\lambda=0}^{n-1} (-1)^{T(\theta^{\lambda})} 4^{2\lambda} \in \mathbb{Z}$ has a property like usual Gauss sum.

Lemma 9. Let $\overline{R} = \sum_{\lambda=0}^{n-1} (-1)^{T(\theta^{\lambda})} 4^{-2\lambda}$. Then

$$\overline{R}R \equiv n+1 - \frac{4^n+1}{5} \pmod{4^n+1}.$$

Proof.

$$\begin{split} \overline{R}R &= \sum_{\lambda,\mu=0}^{n-1} (-1)^{T(\theta^{\lambda}+\theta^{\mu})} 4^{2(\lambda-\mu)} \\ &\equiv \sum_{\tau=0}^{n-1} 4^{2\tau} \sum_{\mu=0}^{n-1} (-1)^{T(\theta^{\mu+\tau}+\theta^{\mu})} \pmod{4^{2n}-1} \\ &\equiv n + \sum_{\tau=1}^{n-1} 4^{2\tau} \sum_{\mu=0}^{n-1} (-1)^{T(\theta^{\mu}(1+\theta^{\tau}))} \pmod{4^{2n}-1}. \end{split}$$

For $1 \le \tau \le n-1, 1+\theta^{\tau} \in \mathbb{F}_q^*$ $(q=n+1=2^k)$. Therefore

$$\sum_{\mu=0}^{n-1} (-1)^{T(\theta^{\mu}(1+\theta^{\tau}))} = \sum_{x \in \mathbb{F}_q^*} (-1)^{T(x(1+\theta^{\tau}))} = \sum_{x \in \mathbb{F}_q^*} (-1)^{T(x)} = -1.$$

Therefore

$$\overline{R}R \equiv n - \sum_{\tau=1}^{n-1} 4^{2\tau} \equiv n - \left(\frac{4^{2n} - 1}{15} - 1\right) \equiv n + 1 - \frac{4^{2n} - 1}{15} \pmod{4^{2n} - 1}.$$

Then from $\frac{4^n-1}{3} \equiv -2 \cdot 3^{-1} \equiv 1 \pmod{5}$, we obtain

$$\overline{R}R \equiv n+1 - \frac{4^n+1}{5} \pmod{4^n+1}.$$

We continue to prove Theorem 8. From (3) we know that $d_+ = \gcd(4^{-2r}R, 4^n + 1)$. Firstly we claim that d_+ has no prime divisor $\pi \neq 5$. Assume that π is a prime, $\pi \neq 5$ and $\pi | d_+$. Then $\pi | 4^n + 1$ and $0 \equiv d_+ \equiv 4^{-2r}R \pmod{\pi}$. (mod π). Therefore $R \equiv 0 \pmod{\pi}$ and $0 \equiv \overline{RR} \equiv n+1 \equiv 2^k \pmod{\pi}$ which is a contradiction. Therefore $d_+ = 5^u \pmod{2}$. From

$$M \equiv 4^{-2r} \sum_{i=0}^{n-1} (-1)^{T(\theta^i)} 4^{2i} \equiv \sum_{i=0}^{n-1} (-1)^{T(\theta^i)} = -1 + \sum_{x \in \mathbb{F}_q} (-1)^{T(x)} = -1 \pmod{5}$$

we have $5 \nmid d_{+} = 5^{u}$. Therefore $d_{+} = 1$ and $FC_{w}(4) = \log_{4}(4^{2n} - 1)$.

For $s \in \mathbb{Z}_n^*$, $M_s(m) = \{c_i\}_{i=0}^{n-1}$ where $c_i = m_{si} = T(\Theta^{si})$. The sequence $M_s(m)$ is also an *m*-sequence with respect to the primitive element Θ^s . If $s \not\equiv 2^t \pmod{n}$ $(0 \le t \le k-1)$, the sequences *m* and $M_s(m)$ are not shift-equivalent, and the 4-adic complexity of $w = w(m, M_s(m))$ is $FC_w(4) = \log_4\left(\frac{4^{2n}-1}{d_+}\right)$, $d_+ = \gcd(M, 4^n + 1)$, and $M = \sum_{i=0}^{n-1} (-1)^{T(\Theta^i + \Theta^{is})} 4^{2i} = \sum_{x \in \mathbb{F}_q^*} (-1)^{T(x+x^s)} 4^{2i} \in \mathbb{Z}$. It seems that to compute the exponential sum *M* is not easy in general case.

B. Twin-Prime sequences

Let p and q = p+2 be prime numbers, n = pq. The ring $\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$ has a partition $\mathbb{Z}_{pq} = \{0\} \bigcup P \bigcup Q \bigcup \mathbb{Z}_{pq}^*$ where $P = \{p, 2p, \dots, (q-1)p\}, Q = \{q, 2q, \dots, (p-1)q\},$

$$\mathbb{Z}_n^* = \{a(\bmod n) | \gcd(a, n) = 1\}$$
$$= \{ip + jq | 1 \le i \le q - 1, 1 \le j \le p - 1\}.$$

The twin-prime sequence $t = {t_{\lambda}}_{\lambda=0}^{n-1}$ with period n = pq is defined by, for $0 \le \lambda \le n-1$,

$$t_{\lambda} = \begin{cases} 0, & \text{if } \lambda = 0; \\ 0, & \text{if } \lambda \in Q; \\ 1, & \text{if } \lambda \in P; \\ \frac{1}{2}(1 - (\frac{\lambda}{p})(\frac{\lambda}{q})), & \text{if } \lambda \in \mathbb{Z}_n^* , \end{cases}$$

where $(\frac{\lambda}{p})$ is the Legendre symbol. Let $T_{\lambda} = (-1)^{t_{\lambda}}$, then

$$T_{\lambda} = \begin{cases} 1, & \text{if } \lambda = 0; \\ 1, & \text{if } \lambda \in Q; \\ -1, & \text{if } \lambda \in P; \\ (\frac{\lambda}{p})(\frac{\lambda}{q}), & \text{if } \lambda \in \mathbb{Z}_{n}^{*} \end{cases}$$

It is known that t and its complement, shift and sample sequences belong to $\sum(n)$. For $l \in \mathbb{Z}_n^*$, we have sample sequence $M_l(t) = \{M_l(t_\lambda)\}_{\lambda=0}^{n-1}$ where $M_l(t_\lambda) = t_{l\lambda}$. In fact, there are only two sample sequences of t :

$$M_l(t) = \begin{cases} t, & \text{if } (\frac{l}{p})(\frac{l}{q}) = 1; \\ \tau(t), & \text{if } (\frac{l}{p})(\frac{l}{q}) = -1, \end{cases}$$

where $\tau(t) = {\tau(t_{\lambda})}_{\lambda=0}^{n-1}$ and

$$\tau(t_{\lambda}) = \begin{cases} \frac{1}{2} (1 - (\frac{l\lambda}{p})(\frac{l\lambda}{q})) = \frac{1}{2} (1 + (\frac{\lambda}{p})(\frac{\lambda}{q})), & \text{if } \lambda \in \mathbb{Z}_{n}^{*}.\\ t_{\lambda}, & \text{otherwise} \end{cases}$$

Since $\left(\frac{-1}{p}\right)\left(\frac{-1}{q}\right) = (-1)^{\frac{p-1}{2} + \frac{q-1}{2}} = (-1)^{\frac{p-1}{2} + \frac{p+1}{2}} = (-1)^p = -1$, we know that $\tau(t) = M_{-1}(t)$. Let $\tau(T_{\lambda}) = (-1)^{\tau(t_{\lambda})}$, then

$$\mathfrak{r}(T_{\lambda}) = \left\{ egin{array}{ll} -T_{\lambda}, & ext{if } \lambda \in \mathbb{Z}_n^*; \ T_{\lambda}, & ext{otherwise.} \end{array}
ight.$$

Now we compute $FC_w(4)$ for w(a,b), $a,b \in \{t, \tau(t)\}$ and $a \neq b$.

Since w(a,b) and w(b,a) have the same 4-adic complexity, we need to compute $FC_w(4)$ where w = w(a,b) for $(a,b) = (t,\tau(t))$.

Theorem 10. Let t be the twin-prime sequence with period n = pq (q = p + 2). Then

$$FC_w(4) = \log_4(4^{2n} - 1), \text{ for } w = w(t, \tau(t)).$$

Proof. Let $t = \{t_{\lambda}\}_{\lambda=0}^{n-1}$, $\tau(t) = \{\tau(t_{\lambda})\}_{\lambda=0}^{n-1}$, $T_{\lambda} = (-1)^{t_{\lambda}}$ and $\tau(T_{\lambda}) = (-1)^{\tau(t_{\lambda})}$. Then by Theorem 3 for $w = w(t, \tau(t))$, we have $FC_w(4) = \log_4(\frac{4^{2n}-1}{d})$ where $d = \gcd(S_w(4), 4^n + 1)$, $S_w(4) = \sum_{\lambda=0}^{n-1} T_{\lambda}\tau(T_{\lambda}) 4^{2\lambda}$.

It is easy to see that $5|4^{n}+1$, $5|4^{p}+1$ and $5|4^{q}+1$. By

$$gcd(4^{p}+1,4^{q}+1) = gcd(4^{p}+1,4^{2}\cdot 4^{p}+1) = gcd(4^{p}+1,15) = 5,$$

we know that $N = \frac{(4^n+1)\cdot 5}{(4^p+1)(4^q+1)} \in \mathbb{Z}$ and $4^n + 1 = N \cdot \frac{4^p+1}{5} \cdot \frac{4^q+1}{5} \cdot 5$. Therefore

$$d = \gcd(S_w(4), 4^n + 1) | \gcd(S_w(4), 5) \cdot \gcd(S_w(4), N) \cdot \gcd(S_w(4), \frac{4^p + 1}{5}) \cdot \gcd(S_w(4), \frac{4^q + 1}{5}).$$
(4)

For $w = w(t, \tau(t))$,

$$\begin{split} S_w(4) \\ &= \sum_{\lambda=0}^{n-1} T_{\lambda} \tau(T_{\lambda}) 4^{2\lambda} \\ &= \sum_{\lambda \in \mathbb{Z}_n \setminus \mathbb{Z}_n^*} 4^{2\lambda} - \sum_{\lambda \in \mathbb{Z}_n^*} 4^{2\lambda} \\ &= 2 \sum_{\lambda \in \{0\} \cup \mathcal{P} \cup \mathcal{Q}} 4^{2\lambda} - \sum_{\lambda \in \mathbb{Z}_n} 4^{2\lambda} \\ &= 2 (\sum_{i=0}^{p-1} 4^{2iq} + \sum_{j=0}^{q-1} 4^{2jp} - 1) - \frac{4^{2n} - 1}{15} \\ &= 2 \left(\frac{4^{2n} - 1}{4^{2p} - 1} + \frac{4^{2n} - 1}{4^{2q} - 1} - 1 \right) - \frac{4^{2n} - 1}{15} \\ &\equiv 2 \left(\frac{4^n + 1}{4^p + 1} \cdot \frac{4^n - 1}{4^p - 1} + \frac{4^n + 1}{4^q + 1} \cdot \frac{4^n - 1}{4^q - 1} - 1 \right) - \frac{4^n + 1}{5} \pmod{N} \\ &\equiv \left\{ \begin{array}{c} 2 \left(\frac{N(4^q + 1)}{5} \cdot \frac{4^n - 1}{4^p - 1} + \frac{N(4^p + 1)}{5} \cdot \frac{4^n - 1}{4^q - 1} - 1 \right) \pmod{N} \\ 2 \left(\sum_{i=0}^{q-1} (-4^p)^i \cdot \sum_{i=0}^{q-1} 4^{pi} + N \cdot \frac{4^p + 1}{5} \cdot \frac{4^n - 1}{4^q - 1} - 1 \right) \pmod{N} \\ 2 \left(\sum_{i=0}^{q-1} (-4^p)^i \cdot \sum_{i=0}^{p-1} 4^{pi} + N \cdot \frac{4^p + 1}{5} \cdot \frac{4^n - 1}{4^q - 1} - 1 \right) \pmod{M} \\ 2 \left(\sum_{i=0}^{q-1} (-4^q)^i \cdot \sum_{i=0}^{p-1} 4^{qi} - 1 \right) \pmod{M} \\ = \left\{ \begin{array}{c} -2 \neq 0 \pmod{N} \qquad N = \frac{(4^n + 1) \cdot 5}{(4^n + 1) (4^n + 1)} = \frac{4^n + 1}{5} / (\frac{4^p + 1}{5} \cdot \frac{4^q + 1}{5}) \\ 2(0 + p - 1) = 2(p - 1) \pmod{\frac{4^p + 1}{5}} \end{array} \right\} \end{aligned} \right\}$$

and

$$S_w(4) \equiv |\mathbb{Z}_n \setminus \mathbb{Z}_n^*| - |\mathbb{Z}_n^*| = n - 2|\mathbb{Z}_n^*| = pq - 2(p-1)(p+1) \equiv -(p^2 - 2p - 2) \pmod{5}$$

If $5|S_w(4)$ then $(p-1)^2 = p^2 - 2p + 1 \equiv 3 \pmod{5}$ which contradicts to $(\frac{3}{5}) = -1$. Therefore $gcd(S_w(4), 5) = 1$. Assume that $\pi|gcd(S_w(4), \frac{4^p+1}{5}) = gcd(2(p+1), \frac{4^p+1}{5})$. By Fermat's Theorem, we have $4^{\pi-1} \equiv 1 \pmod{\pi}$. From $4^p \equiv -1 \pmod{\pi}$ and $\pi \neq 3,5$ we know the order of 4 $(\mod \pi)$ is 2p which implies $2p|\pi-1$. This contradicts to $\pi|p+1$. Therefore $gcd(S_w(4), \frac{4^p+1}{5}) = gcd(2(p+1), \frac{4^p+1}{5}) = 1$. By similar argument we can obtain $gcd(S_w(4), \frac{4^q+1}{5}) = gcd((p-1), \frac{4^q+1}{5}) = 1$. By (4) and Theorem 3 we have $FC_w(4) = \log_4(4^{2n}-1)$ for $w = (t, \tau(t))$.

C. Legendre sequences

Let $p \equiv 3 \pmod{4}$ be a prime number. The Legendre sequences with period p are $\ell = (\ell_{\lambda})_{\lambda=0}^{p-1}$ and $\ell' = (\ell'_{\lambda})_{\lambda=0}^{p-1}$ defined by

$$\ell_0 = 0, \ell'_0 = 1, \ell_\lambda = \ell'_\lambda = \frac{1}{2} \left(1 - \left(\frac{\lambda}{p}\right) \right) \in \{0, 1\} \quad for \ 1 \le \lambda \le p - 1$$

Let $L_{\lambda} = (-1)^{\ell_{\lambda}}, L'_{\lambda} = (-1)^{\ell'_{\lambda}}$, then

$$L_0 = 1, L'_0 = -1, L_\lambda = L'_\lambda = \left(\frac{\lambda}{p}\right) \quad for \ 1 \le \lambda \le p - 1.$$

It is known that ℓ, ℓ' and their complement, shift and sample sequences belong to $\Sigma(p)$. ℓ has only two sample sequences ℓ and $\tau(\ell) = M_{-1}(\ell)$, ℓ' has only two sample sequences ℓ' and $\tau(\ell') = M_{-1}(\ell')$. Let $\tau(L_i) = (-1)^{\tau(\ell_i)}$ and $\tau(L'_i) = (-1)^{\tau(\ell'_i)}$. Then

$$\tau(L_0) = 1, \tau(L'_0) = -1, \tau(L_\lambda) = -L_\lambda, \tau(L'_\lambda) = -L'_\lambda \quad (1 \le \lambda \le p-1)$$

Like in the twin-prime sequences case, for $a, b \in \{\ell, \ell', \tau(\ell), \tau(\ell')\}$, w(a, b), w(b, a), and $w(\tau(a), \tau(b))$ have the same 4-adic complexity.

Theorem 11. Let $p \equiv 3 \pmod{4}$ be a prime number, ℓ and ℓ' be the Legendre sequences with period p. *Then*

$$FC_{w}(4) = \begin{cases} \log_{4}(4^{2p}-1), & \text{if } 5 \nmid p-2 \text{ and } w \in \{w(\ell,\ell'), w(\ell,\tau(\ell)), w(\ell',\tau(\ell'))\} \\ \log_{4}(\frac{4^{2p}-1}{5}), & \text{if } 5 \mid p-2 \text{ and } w \in \{w(\ell,\ell'), w(\ell,\tau(\ell)), w(\ell',\tau(\ell'))\} \\ \log_{4}(5(4^{p}-1)), & \text{if } w = w(\ell,\tau(\ell')). \end{cases}$$

Proof. (1). For $w = w(\ell, \ell')$, since

$$4^{p} + 1 = (5-1)^{p} + 1 \equiv -1 + 5p + 1 \equiv 5p \pmod{25}$$

and $p \equiv 3 \pmod{4}$ we have $5 ||4^p + 1 (5|4^p + 1, 5^2 \nmid 4^p + 1)$ and

$$S_w(4) = \sum_{\lambda=0}^{p-1} L_{\lambda} L_{\lambda}' 4^{2\lambda} = -1 + \sum_{\lambda=1}^{p-1} 4^{2\lambda} = \frac{4^{2p} - 1}{15} - 2 \equiv \begin{cases} -2 \pmod{\frac{4^{p} + 1}{5}} \\ p - 2 \pmod{5} \end{cases}$$

If $5 \nmid p-2$, then $gcd(S_w(4), 4^p+1) = gcd(S_w(4), \frac{4^p+1}{5}) \cdot gcd(S_w(4), 5) = 1$. Therefore $FC_w(4) = \log_4(4^{2p}-1)$. 1). For $5 \mid p-2$, $gcd(S_w(4), 4^p+1) = 1 \cdot 5 = 5$, we have $FC_w(4) = \log_4(\frac{4^{2p}-1}{5})$.

(2). For $w = w(\ell, \tau(\ell))$ and $w(\ell', \tau(\ell'))$,

$$S_w(4) = \sum_{\lambda=0}^{p-1} L_\lambda \tau(L_\lambda) 4^{2\lambda} = \sum_{\lambda=0}^{p-1} L'_\lambda \tau(L'_\lambda) 4^{2\lambda} = 1 - \sum_{\lambda=1}^{p-1} 4^{2\lambda} = 2 - \frac{4^{2p} - 1}{15}$$

Then we obtain the same result as (1).

(3). For $w = w(\ell, \tau(\ell'),$

$$S_w(4) = \sum_{\lambda=0}^{p-1} L_{\lambda} \tau(L'_{\lambda}) 4^{2\lambda} = -\sum_{\lambda=0}^{p-1} 4^{2\lambda} = -\frac{4^{2p}-1}{15}.$$

From $\frac{4^p-1}{3} \equiv 1 \pmod{5}$, we know $S_w(4) \equiv -\frac{4^p+1}{5} \pmod{4^p+1}$. Then we have $d = \gcd(S_w(4), 4^p+1) = \frac{4^p+1}{5}$ and $FC_w(4) = \log_4 \frac{4^{2p}-1}{d} = \log_4(5(4^p-1))$ by Theorem 3.

D. Interleaving by Hall sequences

Let $p = 4x^2 + 27$ be a prime number, $x \equiv 1 \pmod{3}$. Then $6 \mid p - 1$. Let $\mathbb{F}_p^* = \langle g \rangle$, $D = \langle g^6 \rangle$ and $D_{\lambda} = g^{\lambda}D \ (0 \leq \lambda \leq 5)$ be the cyclotomic classes of order 6 in \mathbb{F}_p^* . The Hall sequence with period p is the binary sequence $h = \{h_{\lambda}\}_{\lambda=0}^{p-1}$ defined by

$$h_{\lambda} = \begin{cases} 1, & \text{if } \lambda \in D_0 \cup D_1 \cup D_3; \\ 0, & \text{otherwise.} \end{cases}$$

It is known that *h* and complement, shift, sample of *h* belong to $\Sigma(p)$ (see [9], p.73-74, Theorem 18, where it is proved as an equivalent form: $D_0 \cup D_1 \cup D_3$ is a difference set of the group $(\mathbb{F}_p, +)$). For $1 \le r \le p-1$, there exists six sample sequences of *h*:

$$M_r(h) = h^{(l)} = \{h_{\lambda}^{(l)}\}_{\lambda=0}^{p-1} \text{ if } r \in D_l,$$

where for $0 \le \lambda \le p - 1$,

$$h_{\lambda}^{(l)} = \begin{cases} 1, & \text{if } \lambda \in D_l \cup D_{1+l} \cup D_{3+l}; \\ 0, & \text{otherwise.} \end{cases}$$

Particularly, we take r = -1. From $-1 = g^{\frac{p-1}{2}}$ and $\frac{p-1}{2} = 2x^2 + 13 \equiv 3 \pmod{6}$, we have $-1 \in D_3$. Thus $M_{-1}(h) = h^{(3)} = \{h_{\lambda}^{(3)}\}_{\lambda=0}^{p-1}$ where for $0 \le \lambda \le p-1$,

$$h_{\lambda}^{(3)} = \begin{cases} 1, & \text{if } \lambda \in D_3 \cup D_4 \cup D_0; \\ 0, & \text{otherwise.} \end{cases}$$

In order to determine the 4-adic complexity of $w = w(h, M_r(h))$, we need some facts on cyclotomic numbers and "Gauss periods".

Definition 3. Let $p \equiv 1 \pmod{6}$, p-1 = 6f, $\mathbb{F}_p^* = \langle g \rangle$, $D = \langle g^6 \rangle$ and $D_{\lambda} = g^{\lambda}D \ (0 \le \lambda \le 5)$ be the cyclotomic classes of order 6.

(1). The "Gauss periods" of order 6 in \mathbb{F}_p^* and valued in \mathbb{Z}_N $(N = 4^{2p} - 1)$ are defined by

$$\xi_{\lambda} = \sum_{i \in D_{\lambda}} 4^{2i} \pmod{4^{2p} - 1} \quad (0 \le \lambda \le 5).$$

Similarly we have the cyclotomic classes of order 3 in \mathbb{F}_p^*

$$C_{\lambda} = g^{\lambda}C, \quad C = \langle g^3 \rangle \ (0 \le \lambda \le 2), \quad |C_{\lambda}| = \frac{p-1}{3}$$

and "Gauss periods" of order 3 $\eta_{\lambda} = \sum_{i \in C_{\lambda}} 4^{2i} \pmod{4^{2p} - 1}$. We have

$$C_{\lambda} = D_{\lambda} \cup D_{\lambda+3}, \quad \eta_{\lambda} = \xi_{\lambda} + \xi_{\lambda+3},$$

(2). The cyclotomic numbers of order 3 over \mathbb{F}_p is defined by

$$(i,j) = |(C_i+1) \cap C_j| = \{a \in C_i \mid a+1 \in C_j\} \ (0 \le i, j \le 2).$$

Now we show that the 4-adic complexity of $w = w(h, M_{-1}(h))$ can be determined by using cyclotomic numbers and "Gauss periods" of order 3. (For other sample sequence $M_r(h)$ of h $(r \in \mathbb{Z}_p^*, r \neq -1)$, in order to determine the 4-adic complexity of $w(h, M_r(h))$ we need more information on cyclotomic numbers and "Gauss periods" of order 6.) We use the following results on cyclotomic numbers (i, j) $(0 \le i, j \le 2)$ and "Gauss periods" η_{λ} $(0 \le \lambda \le 2)$. It is known that any prime number $p \equiv 1 \pmod{3}$ can be expressed as $4p = c^2 + 27d^2$ where $c, d \in \mathbb{Z}$, c is uniquely determined by $c \equiv 1 \pmod{3}$ and d is determined up to sign.

Lemma 12. Let $p \equiv 1 \pmod{3}$ be a prime number, $4p = c^2 + 27d^2$, $c \equiv 1 \pmod{3}$.

(1). ([9], Lemma 7, p.32-35) The cyclotomic numbers (i, j) $(0 \le i, j \le 2)$ of order 3 over \mathbb{F}_p are listed in the following table

$$(i,j)$$
 $j=0$ 12 $9A=p-8+c$ $i=0$ ABC $18B=2p-4-c-9d$ 1BCD $18C=2p-4-c+9d$ 2CDB $9D=p+1+c$

(2). The "Gauss periods" $\eta_{\lambda}~(0 \leq \lambda \leq 2)$ have the following relation

$$\eta_m \eta_{m+k} = \sum_{h=0}^{2} (k,h) \eta_{h+m} + \frac{p-1}{3} \delta_k \pmod{\frac{4^{2p}-1}{15}} \ (m,k \in \{0,1,2\})$$

where $\delta_0 = 1$, $\delta_1 = \delta_2 = 0$.

Proof. (2). By using similar proof given in [9] (Lemma 8, p.38-39) for usual "Gauss periods".

Theorem 13. Let $p = 4x^2 + 27$ be a prime number, $h = \{h_{\lambda}\}_{\lambda=0}^{p-1}$ be the Hall sequence with period p. Then for the quaternary interleaved sequence $w = w(h, M_{-1}(h))$,

$$FC_{w}(4) = \begin{cases} \log_{4}(\frac{4^{2p}-1}{5}), & \text{if } p \equiv 3 \pmod{5}; \\ \log_{4}(4^{2p}-1), & \text{otherwise.} \end{cases}$$

Proof. By Theorem 3 we know that $FC_w(4) = \log_4(\frac{4^{2p}-1}{d_+})$, where $d_+ = \gcd(S_w(4), 4^p + 1)$ and

$$S_w(4) \equiv \sum_{\lambda=0}^{p-1} H_{\lambda} M_{-1}(H_{\lambda}) 4^{2\lambda} \pmod{4^{2p}-1}$$

By the definition as before, for $0 \le \lambda \le p - 1$,

$$H_{\lambda} = (-1)^{h_{\lambda}} = \begin{cases} -1, & \text{if } \lambda \in D_0 \cup D_1 \cup D_3; \\ 1, & \text{otherwise.} \end{cases} \qquad M_{-1}(H_{\lambda}) = \begin{cases} -1, & \text{if } \lambda \in D_3 \cup D_4 \cup D_0; \\ 1, & \text{otherwise.} \end{cases}$$

We have

$$S_{w}(4) \equiv \sum_{\lambda \in \mathbb{F}_{p} \setminus D_{1} \cup D_{4}} 4^{2\lambda} - \sum_{\lambda \in D_{1} \cup D_{4}} 4^{2\lambda} = \sum_{\lambda=0}^{p-1} 4^{2\lambda} - 2\sum_{\lambda \in D_{1} \cup D_{4}} 4^{2\lambda} \pmod{4^{2p} - 1}$$
$$\equiv \frac{4^{2p} - 1}{15} - 2\eta_{1} \equiv \frac{4^{p} + 1}{5} - 2\eta_{1} \pmod{4^{p} + 1}.$$

Therefore $d_+ = \operatorname{gcd}\left(\frac{4^p+1}{5} - 2\eta_1, 4^p+1\right)$. Let $d_+ = 5^l \cdot d'$, $5 \nmid d'$. Firstly we show that d' = 1. Assume that d' > 1. Let π be a prime divisor of d'. Then $\pi > 5$, $\pi \mid 4^p + 1$ and $0 \equiv \frac{4^p+1}{5} - 2\eta_1 \equiv -2\eta_1 \pmod{\pi}$, namely, $\pi \mid \eta_1$. By Lemma 12 we obtain the following linear equations over the finite field \mathbb{F}_{π}

$$C\eta_0 + A\eta_1 + B\eta_2 \equiv \eta_1^2 - \frac{p-1}{3} \equiv -\frac{p-1}{3} \pmod{\pi},$$

$$B\eta_0 + C\eta_1 + D\eta_2 \equiv \eta_0\eta_1 \equiv 0 \pmod{\pi},$$

$$D\eta_0 + B\eta_1 + C\eta_2 \equiv \eta_2\eta_1 \equiv 0 \pmod{\pi}.$$

To solve this system of linear equations in variables (η_0, η_1, η_2) , we obtain $M\eta_1 \equiv M' \pmod{\pi}$ where M is the determinant of coefficient matrix and

$$\begin{split} M' &= \begin{vmatrix} C & -\frac{p-1}{3} & B \\ B & 0 & D \\ D & 0 & C \end{vmatrix} = \frac{p-1}{3} \begin{vmatrix} B & D \\ D & C \end{vmatrix} = \frac{p-1}{3} (BC - D^2) \\ &= \frac{p-1}{3} \left[\frac{1}{18^2} (2p - 4 - c - 9d)(2p - 4 - c + 9d) - \frac{1}{9^2} (p + 1 + c)^2 \right] \\ &= \frac{p-1}{3^4} (1 - 3p - pc). \end{split}$$

From $\pi \mid \eta_1$ we get $\frac{p-1}{3^4}(1-3p-pc) \equiv M\eta_1 \equiv 0 \pmod{\pi}$. From $4^p \equiv -1 \pmod{\pi}$ and $\pi \neq 3,5$ we know that order of 4 (mod π) is 2*p*. By Fermat's Theorem, $2p \mid \pi - 1, \pi - 1 = 2pt$ where *t* is a positive number and then $\pi > p - 1$. Therefore $\pi \nmid p - 1$ and $1 - 3p - pc \equiv 0 \pmod{\pi}$. Multiple by 2*t*, we obtain

$$0 \equiv 2t(pc+3p-1) \equiv (\pi-1)(c+3) - 2t \equiv -(c+3+2t) \pmod{\pi}.$$

From $p = 4x^2 + 27 \ge 31$, $\pi = 1 + 2tp > 2p = 62$, $4p = c^2 + 27d^2$ we have

$$|c+3+2t| \le \sqrt{4p} + 3 + \frac{\pi - 1}{p} \le \sqrt{2\pi} + 3 + \frac{\pi}{31} < \pi$$

By $c+3+2t \equiv 0 \pmod{\pi}$ we obtain c+3+2t = 0. Then by $p \equiv c \equiv 1 \pmod{3}$ we have

$$\pi = 2tp + 1 = -p(c+3) + 1 \equiv -pc + 1 \equiv 0 \pmod{3}$$

which is a contradiction since π is a prime and $\pi \neq 3$. Therefore d' = 1 and $d_+ = 5^l$. At last we determine d_+ . Firstly, by the proof of Theorem 11 we have

$$5 \parallel 4^p + 1$$
, if $p \neq 5$; and $25 \mid 4^5 + 1$.

Then from

$$S_w(4) = \sum_{\lambda=0}^{p-1} 4^{2\lambda} - 2\sum_{\lambda \in D_1 \cup D_4} 4^{2\lambda} \equiv p - 2|D_1 \cup D_4| = p - \frac{p-1}{3} = \frac{2p+1}{3} \pmod{5}$$

we obtain

 $5 \mid S_w(4)$ if and only if $p \equiv 2 \pmod{5}$.

Therefore, if $p \equiv 2 \pmod{5}$ then $d_+ = 5$ we have $FC_w(4) = \log_4(\frac{4^{2p}-1}{5})$. Otherwise, $5 \nmid S_w(4)$, then $d_+ = 1$ and $FC_w(4) = \log_4(4^{2p}-1)$.

Remark 2. To resist the attack of the rational algorithm, the 4-adic complexity of a quaternary sequence w(a,b) with period N should exceed $\frac{N-16}{6}$. From Theorem 3 we can see that the quaternary interleaved sequences defined in [11] are safe enough to resist the attack of the rational approximation algorithm. By Theorem 8, 10, 11, 13, we can see that by considering a and b to be several types of known binary sequences with ideal autocorrelation (m-sequences, twin-prime sequences, Legendre and Hall sequences and their complement, shift or sample sequences), the 4-adic complexity of a quaternary sequence w(a,b) reaches or nearly reaches the maximum in most cases.

IV. CONCLUSION

Tang and Ding [11] present a series of quaternary sequences w(a, b) interleaved by two binary sequences a and b with ideal autocorrelation and show that such interleaved quaternary sequences have optimal autocorrelation. In this paper we consider the 4-adic complexity $FC_w(4)$ of such interleaved quaternary sequences w = w(a, b). Firstly we present a general formula on $FC_w(4)$, w = w(a, b). As a direct consequence, we obtain lower bound $FC_w(4) \ge \log_4(4^n - 1)$ where 2n is the period of w = w(a, b). Then by considering a and b to be several types of known binary sequences (*m*-sequences, twin-prime sequences, Legendre and Hall sequences and their complement, shift or sample sequences), we obtain the exact value of $FC_w(4)$. The results show that in most cases, $FC_w(4)$ reaches or nearly reaches the maximum value $\log_4(4^{2n} - 1)$. Our results show that the quaternary sequences defined in [11] are safe enough from the viewpoint of the 4-adic complexity. Many constructions on binary sequences with period $n \equiv 3 \pmod{4}$ and ideal autocorrelation have been found. It seems to use more skill in order to determine the exact values of the 4-adic complexity of the quaternary sequences interleaved by other types of binary sequences with ideal autocorrelation.

REFERENCES

- [1] T. W. Cusick, C. Ding, and A. Renvall, Stream Ciphers and Number Theory, (Revised Edition), Elsevier, 2004.
- [2] S. W. Golomb and G. Gong, Signal Design for Good Correlation: for Wireless Communication, Cryptography and Radar, Cambridge University Press, 2005.
- [3] G. Gong, "Theory and applications of *q*-ary interleaved sequences," *IEEE Trans. Inf. Theory*, vol. 41, no. 2, pp. 400-411, 1995.
- [4] J. Jang, Y. Kim, S. Kim, and J. No, "New quaternary sequences with ideal autocorrelation constructed from binary sequences with ideal autocorrelation," in *Proc. IEEE Int. Symp. Inf. Theory*, Seoul, Korea, June 28-July 3, pp. 278-281, 2009.
- [5] Y. Kim, J. Jang, S. Kim, and J. No, "New construction of quaternary sequences with ideal autocorrelation from Legendre sequences," in *Proc. IEEE Int. Symp. Inf. Theory*, Seoul, Korea, June 28-July 3, pp. 282-285, 2009.
- [6] A. Klapper, "A survey of feedback with carry shift registers," in *Proc. Sequences and Their Applications*, Seoul, Korea, October 24-October 28, pp. 56-71, 2004.
- [7] A. Klapper and J. Xu, "Register synthesis for algebraic feedback shift registers based on non-primes," *Designs, Codes, and Crypt.*, vol. 31, no. 3, pp. 227-250, 2004.
- [8] N. Li, X. Tang, and T. Helleseth, "New *M*-ary sequences with low autocorrelation from interleaved technique," *Designs*, *Codes and Cryptogr.*, vol. 73, pp. 237-249, 2014.
- [9] T. Storer, Cyclotomy and Difference Sets, Lecture in Advanced Mathematics, Chicago, Markham, 1967.
- [10] W. Su, Y. Yang, Z. Zhou, and X. Tang, "New quaternary sequences of even length with optimal autocorrelation," *Sci. China Inf. Sci.*, vol. 61, no. 2: 022308:1-022308:13, 2018.
- [11] X. Tang and C. Ding, "New classes of belanced quartenary and almost balanced binary sequences with optimal autocorrelation value," *IEEE Trans. Inf. Theory*, vol. 56, no. 2, pp. 6398-6405, 2010.