

# ON PLANE ALGEBRAIC CURVES PASSING THROUGH $n$ -INDEPENDENT NODES

H. A. Hakopian, H. M. Kloyan, D. S. Voskanyan

*Department of Informatics and Applied Mathematics,  
Yerevan State University*

## Abstract

Let a set of nodes  $\mathcal{X}$  in the plane be  $n$ -independent, i.e., each node has a fundamental polynomial of degree  $n$ . Assume that  $\#\mathcal{X} = d(n, k) + 3 = (n + 1) + n + \dots + (n - k + 5) + 3$  and  $4 \leq k \leq n - 1$ . In this paper we prove that there are at most seven linearly independent curves of degree less than or equal to  $k$  that pass through all the nodes of  $\mathcal{X}$ . We provide a characterization of the case when there are exactly seven such curves. Namely, we prove that then the set  $\mathcal{X}$  has a very special construction: all its nodes but three belong to a (maximal) curve of degree  $k - 3$ . Let us mention that in a series of such results this is the third one. At the end an important application to the bivariate polynomial interpolation is provided, which is essential also for the study of the Gasca-Maeztu conjecture.

**MSC2010:** 14H50, 41A05, 41A63.

**Keywords:** algebraic curves, maximal curves, bivariate polynomial interpolation, fundamental polynomial,  $n$ -independent nodes.

## 1 Introduction

Denote the space of all bivariate polynomials of total degree not exceeding  $n$  by

$$\Pi_n = \left\{ \sum_{i+j \leq n} a_{ij} x^i y^j \right\}.$$

We have that

$$N := N_n := \dim \Pi_n = (1/2)(n + 1)(n + 2).$$

Denote by  $\Pi$  the space of all bivariate polynomials.

Consider a set of  $s$  distinct nodes  $\mathcal{X} = \mathcal{X}_s = \{(x_1, y_1), (x_2, y_2), \dots, (x_s, y_s)\}$ . The problem of finding a polynomial  $p \in \Pi_n$ , which satisfies the conditions

$$p(x_i, y_i) = c_i, \quad i = 1, \dots, s, \quad (1)$$

is called interpolation problem.

A polynomial  $p \in \Pi_n$  is called a fundamental polynomial for a node  $A \in \mathcal{X}$  if  $p(A) = 1$  and  $p|_{\mathcal{X} \setminus \{A\}} = 0$ , where  $p|_{\mathcal{X}}$  means the restriction of  $p$  on  $\mathcal{X}$ . We denote this  $n$ -fundamental polynomial by  $p_A^* := p_{A, \mathcal{X}}^*$ . Sometimes we call fundamental also a polynomial that vanishes at all nodes of  $\mathcal{X}$  but one, since it is a nonzero constant times a fundamental polynomial.

**Definition 1.1.** *The interpolation problem with a set of nodes  $\mathcal{X}_s$  is called  $n$ -poised if for any data  $(c_1, \dots, c_s)$  there is a unique polynomial  $p \in \Pi_n$  satisfying the interpolation conditions (1).*

A necessary condition of poisedness is  $\#\mathcal{X}_s = s = N$ .

**Proposition 1.2.** *A set of nodes  $\mathcal{X}_N$  is  $n$ -poised if and only if*

$$p \in \Pi_n \quad \text{and} \quad p|_{\mathcal{X}_N} = 0 \quad \implies \quad p = 0.$$

Next, let us consider the concept of  $n$ -independence (see [1, 2]).

**Definition 1.3.** *A set of nodes  $\mathcal{X}_s$  is called  $n$ -independent, if all its nodes have  $n$ -fundamental polynomials. Otherwise, it is called  $n$ -dependent.*

Fundamental polynomials are linearly independent. Therefore a necessary condition of  $n$ -independence for  $\mathcal{X}_s$  is  $s \leq N$ .

## 1.1 Some Properties of $n$ -Independent Nodes

Let us start with the following

**Lemma 1.4** (Lemma 2.2, [3]). *Suppose that a set of nodes  $\mathcal{X}$  is  $n$ -independent and the nodes of another set  $\mathcal{Y}$  have  $n$ -fundamental polynomials with respect to the set  $\mathcal{Z} = \mathcal{X} \cup \mathcal{Y}$ . Then the set  $\mathcal{Z}$  is  $n$ -independent too.*

Denote the distance between the points  $A$  and  $B$  by  $\rho(A, B)$ . Let us recall the following (see [4, 5])

**Lemma 1.5.** *Suppose that  $\mathcal{X}_s = \{A_i\}_{i=1}^s$  is an  $n$ -independent set. Then there is a number  $\epsilon > 0$  such that any set  $\mathcal{X}'_s = \{A'_i\}_{i=1}^s$ , with the property that  $\rho(A_i, A'_i) < \epsilon$ ,  $i = 1, \dots, s$ , is  $n$ -independent too.*

Next result concerns the extension of  $n$ -independent sets.

**Lemma 1.6** (Lemma 2.1, [2]). *Any  $n$ -independent set  $\mathcal{X}$  with  $\#\mathcal{X} < N$  can be enlarged to an  $n$ -poised set.*

Denote the linear space of polynomials of total degree at most  $n$  vanishing on  $\mathcal{X}$  by

$$\mathcal{P}_{n, \mathcal{X}} = \{p \in \Pi_n : p|_{\mathcal{X}} = 0\}.$$

The following two propositions are well-known (see, e.g., [2]).

**Proposition 1.7.** *For any node set  $\mathcal{X}$  we have that*

$$\dim \mathcal{P}_{n,\mathcal{X}} = N - \#\mathcal{Y},$$

where  $\mathcal{Y}$  is a maximal  $n$ -independent subset of  $\mathcal{X}$ .

The following result is well-known.

**Proposition 1.8.** *If a polynomial  $p \in \Pi_n$  vanishes at  $n+1$  points of a line  $\ell$ , then we have that  $p = \ell r$ , where  $r \in \Pi_{n-1}$ .*

A plane algebraic curve is the zero set of some bivariate polynomial of degree  $\geq 1$ . To simplify notation, we shall use the same letter  $p$ , say, to denote the polynomial  $p$  of degree  $\geq 1$  and the curve given by the equation  $p(x, y) = 0$ .

In the sequel we will need the following

**Proposition 1.9** (Prop. 1.10, [3]). *Let  $\mathcal{X}$  be a set of nodes. Then the following two conditions are equivalent:*

*i) There is no algebraic curve of degree  $k$  passing through all the points of a set  $\mathcal{X}$ ;*

*ii) There is a  $k$ -poised subset of  $\mathcal{X}$ .*

Set  $d(n, k) := N_n - N_{n-k} = (1/2)k(2n + 3 - k)$ . The following is a generalization of Proposition 1.8.

**Proposition 1.10** (Prop. 3.1, [6]). *Let  $q$  be an algebraic curve of degree  $k \leq n$  without multiple components. Then the following hold:*

*i) any subset of  $q$  containing more than  $d(n, k)$  nodes is  $n$ -dependent;*

*ii) any subset  $\mathcal{X}_d$  of  $q$  containing exactly  $d = d(n, k)$  nodes is  $n$ -independent if and only if the following condition holds:*

$$p \in \Pi_n \quad \text{and} \quad p|_{\mathcal{X}_d} = 0 \implies p = qr, \quad \text{where } r \in \Pi_{n-k}. \quad (2)$$

Thus, according to Proposition 1.10, *i*), at most  $d(n, k)$  nodes of  $\mathcal{X}$  can lie in a curve  $q$  of degree  $k \leq n$ . This motivates the following

**Definition 1.11** (Def. 3.1, [6]). *Given an  $n$ -independent set of nodes  $\mathcal{X}_s$  with  $s \geq d(n, k)$ . A curve of degree  $k \leq n$  passing through  $d(n, k)$  points of  $\mathcal{X}_s$  is called maximal.*

We say that a node  $A$  of an  $n$ -poised set  $\mathcal{X}$  uses a curve  $q \in \Pi_s$ , if the latter divides the  $n$ -fundamental polynomial of  $A$ , i.e.,  $p_A^* = qr$  for some  $r \in \Pi_{n-s}$ .

Let us bring a characterization of maximal curves:

**Proposition 1.12** (Prop. 3.3, [6]). *Let a node set  $\mathcal{X}_N$  be  $n$ -poised. Then a curve  $\mu$  of degree  $k$ ,  $k \leq n$ , is a maximal curve if and only if it is used by all nodes in  $\mathcal{X}_N \setminus \mu$ .*

Next result concerns maximal independent sets in curves.

**Proposition 1.13** (Prop. 3.5, [5]). *Assume that  $\sigma$  is an algebraic curve of degree  $k$  without multiple components and  $\mathcal{X}_s \subset \sigma$  is any  $n$ -independent node set of cardinality  $s$ ,  $s < d(n, k)$ . Then the set  $\mathcal{X}_s$  can be extended to a maximal  $n$ -independent set  $\mathcal{X}_d \subset \sigma$  of cardinality  $d = d(n, k)$ .*

Below a replacement of a node in an  $n$ -independent set is described such that the set remains  $n$ -independent.

**Lemma 1.14** (Lemma 6, [7]). *Assume that  $\mathcal{X}$  is an  $n$ -independent node set and a node  $A \in \mathcal{X}$  has an  $n$ -fundamental polynomial  $p_A^*$  such that  $p_A^*(A') \neq 0$ . Then we can replace the node  $A$  with  $A'$  such that the resulted set  $\mathcal{X}' := \mathcal{X} \cup \{A'\} \setminus \{A\}$  is again  $n$ -independent. In particular, such replacement can be done in the following two cases:*

*i) if a node  $A \in \mathcal{X}$  belongs to several components of  $\sigma$ , then we can replace it with a node  $A'$ , which belongs to only one (desired) component of  $\sigma$ ;*

*ii) if a curve  $q$  is not a component of an  $n$ -fundamental polynomial  $p_A^*$  then we can replace the node  $A$  with a node  $A'$  lying in  $q$ .*

Finally, let us bring a well-known

**Lemma 1.15.** *Suppose that  $m$  linearly independent curves pass through all the nodes of  $\mathcal{X}$ . Then for any node  $A \notin \mathcal{X}$  there are  $m - 1$  linearly independent curves in the linear span of given curves, passing through  $A$  and all the nodes of  $\mathcal{X}$ .*

## 2 A series of results and main results

Let us start with the first result:

**Theorem 2.1** (Th. 1, [8]). *Assume that  $\mathcal{X}$  is an  $n$ -independent set of  $d(n, k - 1) + 2$  nodes lying in a curve of degree  $k$  with  $k \leq n$ . Then the curve is determined uniquely by these nodes.*

The second result in this series is the following

**Theorem 2.2** (Th. 4.2, [5]). *Assume that  $\mathcal{X}$  is an  $n$ -independent set of  $d(n, k - 1) + 1$  nodes with  $2 \leq k \leq n - 1$ . Then at most two different curves of degree  $\leq k$  may pass through all the nodes of  $\mathcal{X}$ . Moreover, there are such two curves for the set  $\mathcal{X}$  if and only if all the nodes of  $\mathcal{X}$  but one lie in a maximal curve of degree  $k - 1$ .*

Next result is the following

**Theorem 2.3** (Th. 3, [7]). *Assume that  $\mathcal{X}$  is an  $n$ -independent set of  $d(n, k - 2) + 2$  nodes with  $3 \leq k \leq n - 1$ . Then at most four linearly independent curves of degree  $\leq k$  may pass through all the nodes of  $\mathcal{X}$ . Moreover, there are such four curves for the set  $\mathcal{X}$  if and only if all the nodes of  $\mathcal{X}$  but one lie in a maximal curve of degree  $k - 2$ .*

Now let us present the main result of this paper:

**Theorem 2.4.** *Assume that  $\mathcal{X}$  is an  $n$ -independent set of  $d(n, k - 3) + 3$  nodes with  $4 \leq k \leq n - 1$ . Then at most seven linearly independent curves of degree  $\leq k$  may pass through all the nodes of  $\mathcal{X}$ . Moreover, there are such seven curves for the set  $\mathcal{X}$  if and only if all the nodes of  $\mathcal{X}$  but three lie in a maximal curve of degree  $k - 3$ .*

Let us mention that the inverse implication in the ‘‘Moreover’’ part is straightforward. Indeed, assume that  $d(n, k - 3)$  nodes of  $\mathcal{X}$  are located in a curve  $\mu$  of degree  $k - 3$ . Therefore, the curve  $\mu$  is maximal and the remaining three nodes of  $\mathcal{X}$ , denoted by  $A, B$  and  $C$ , are outside of it:  $A, B, C \notin \mu$ . Hence we have that

$$\mathcal{P}_{k, \mathcal{X}} = \{p : p \in \Pi_k, p_{\mathcal{X}} = 0\} = \{q\mu : q \in \Pi_3, q(A) = q(B) = q(C) = 0\}.$$

Thus we get readily that  $\dim \mathcal{P}_{k, \mathcal{X}} = \dim \{q \in \Pi_3 : q(A) = q(B) = q(C) = 0\} = \dim \mathcal{P}_{3, \{A, B, C\}} = 10 - 3 = 7$ . Note that in the last equality we use Proposition 1.7 and the fact that any three nodes are 3-independent.

We get also that it is enough to prove only the ‘‘Moreover’’ part. Indeed assume that the ‘‘Moreover’’ part is proved. Then assume that there are  $\geq 7$  linearly independent curves satisfying the conditions mentioned in Theorem 2.4. Then as we proved above we have that  $\dim \mathcal{P}_{k, \mathcal{X}} = 7$ , i.e., there are no more such curves, Q.E.D.

Let us mention that in the proof of Theorem 2.4 we find some interesting generalization of Theorem 2.3, where we increase by one the number of nodes and decrease by one the number of linearly independent curves:

**Theorem 2.5.** *Assume that  $\mathcal{X}$  is an  $n$ -independent set of  $d(n, k - 2) + 3$  nodes with  $3 \leq k \leq n - 2$ . Then at most three linearly independent curves of degree  $\leq k$  may pass through all the nodes of  $\mathcal{X}$ . Moreover, there are such three curves for the set  $\mathcal{X}$  if and only if all the nodes of  $\mathcal{X}$  lie in a curve of degree  $k - 1$ , or all the nodes of  $\mathcal{X}$  but three lie in a (maximal) curve of degree  $k - 2$ .*

### 3 Some preliminaries

Before we start the proof of Theorem 2.4, i.e., the proof of the ‘‘Moreover’’ part, we need to do considerable preliminary work.

**Lemma 3.1.** *Assume that the hypotheses of Theorem 2.4 hold and assume additionally that there is a curve  $\sigma \in \Pi_{k-2}$  passing through all the nodes of  $\mathcal{X}$ . Then all the nodes of  $\mathcal{X}$  but three (collinear) lie in a maximal curve  $\mu$  of degree  $k - 3$ .*

*Proof.* First note that the curve  $\sigma$  is of exact degree  $k - 2$ , since it passes through more than  $d(n, k - 3)$   $n$ -independent nodes. This implies also that  $\sigma$  has no multiple component. Therefore, in view of Proposition 1.13, we can extend the set  $\mathcal{X}$  till a maximal  $n$ -independent set  $\mathcal{Z} \subset \sigma$ , by adding  $d(n, k - 2) - d(n, k - 3) - 3 = n - k + 1$  nodes, i.e.,

$$\mathcal{Z} = \mathcal{X} \cup \mathcal{A}, \text{ where } \mathcal{A} = \{A_0, \dots, A_{n-k}\}.$$

In view of Lemma 1.14, *i*), we may suppose that the nodes from  $\mathcal{A}$  are not intersection points of the components of the curve  $\sigma$ .

Next, we are going to prove that these  $n - k + 1$  nodes are collinear together with  $m \geq 3$  nodes from  $\mathcal{X}$ . To this end denote the line through the nodes  $A_0$  and  $A_1$  by  $\ell_{01}$ . Then for each  $i = 2, \dots, n - k$ , choose a line  $\ell_i$  passing through the node  $A_i$ , which is not a component of  $\sigma$ . We require also that each line passes through only one of the mentioned nodes and therefore the lines are distinct.

Now suppose that  $p \in \Pi_k$  vanishes on  $\mathcal{X}$ . Consider the polynomial  $r = p\ell_{01}\ell_2 \cdots \ell_{n-k}$ . We have that  $r \in \Pi_n$  and  $r$  vanishes on the node set  $\mathcal{Z}$ , which is a maximal  $n$ -independent set in the curve  $\sigma$ . Therefore, we obtain that  $r = \sigma s$ , where  $s \in \Pi_{n-k+2}$ . Thus we have that

$$p\ell_{01}\ell_2 \cdots \ell_{n-k} = \sigma s, \text{ where } s \in \Pi_{n-k+2}.$$

The lines  $\ell_i$ ,  $i = 2, \dots, n - k$ , are not components of  $\sigma$ . Therefore, they are components of the polynomial  $s$ . Hence we obtain that

$$p\ell_{01} = \sigma\gamma, \text{ where } \gamma \in \Pi_3.$$

Now let us verify that  $\ell_{01}$  is a component of  $q_{k-1}$ . Indeed, otherwise it is a component of the cubic  $\gamma$  and we get that

$$p \in \Pi_k, p|_{\mathcal{X}} = 0 \implies p = \sigma\beta, \text{ where } \beta \in \Pi_2.$$

Therefore, we get  $\dim \mathcal{P}_{k,\mathcal{X}} \leq 6$ , which contradicts the hypothesis.

Thus we conclude that

$$\sigma = \ell_{01}q_{k-3}, \text{ where } q_{k-3} \in \Pi_{k-3}. \quad (3)$$

Now let us show that all the nodes of  $\mathcal{A}$  belong to  $\ell_{01}$ . Suppose conversely that a node from  $\mathcal{A}$ , say  $A_2$ , does not belong to the line  $\ell_{01}$ . Then in the same way as in the case of the line  $\ell_{01}$  we get that  $\ell_{02}$  is a component

of  $\sigma$ . Therefore the node  $A_0$  is an intersection point of two components of  $\sigma$ , i.e.,  $\ell_{01}$  and  $\ell_{02}$ , which contradicts our assumption.

Thus we get that  $\mathcal{A} \subset \ell_{01}$ . Next, let us verify that when extending the set  $\mathcal{X} \subset \sigma$  till  $n$ -maximal set one has to locate the added nodes outside the component  $q_{k-3}$ . Indeed, what was proved already implies that the only possible location of such a node in  $q_{k-3}$  is an intersection point with  $\ell_{01}$ . But in the latter case, by using Lemma 1.14, we can replace the node with one belonging only to the component  $q_{k-3}$ , which is a contradiction.

Hence, in view of Proposition 1.13 we get that  $\mu = q_{k-3}$  is a maximal curve for  $\mathcal{X}$ . Therefore, it vanishes at exactly  $d(n, k-3)$  nodes of  $\mathcal{X}$ . The remaining three nodes, according to (3), belong to the line  $\ell_{01}$ .  $\square$

**Lemma 3.2.** *Assume that  $p_1, p_2 \in \Pi$ , and  $p_1$  has no multiple factors. Then, for sufficiently small  $\epsilon$ , the polynomial  $p_1 + \epsilon p_2$  has no multiple factors either.*

*Proof.* Assume by way of contradiction that there is a strictly decreasing sequence  $\epsilon_n$  such that

$$p_1 + \epsilon_n p_2 = q_n r_n^2, \text{ where } q_n, r_n \in \Pi \text{ and } \epsilon_n \rightarrow 0. \quad (4)$$

We have that  $\deg(p_1 + \epsilon_n p_2) \leq \max(\deg p_1, \deg p_2)$ , and hence

$$\deg q_n + 2 \deg r_n \leq \max(\deg p_1, \deg p_2).$$

We deduce from here that there is a subsequence  $n_k$  such that

$$\deg q_{n_k} = m_1 = \text{const. and } \deg r_{n_k} = m_2 = \text{const.}$$

Without loss of generality assume that this condition holds for all the sequence, i.e.,

$$\{\epsilon_n\} \equiv \{\epsilon_{n_k}\}. \quad (5)$$

Thus we have that

$$q_n = \sum_{i+j \leq m_1} a_{ij}^{(n)} x^i y^j, \quad r_n = \sum_{i+j \leq m_2} b_{ij}^{(n)} x^i y^j.$$

In view of (4), by a normalization of  $r_n$ , i.e., by multiplying it by a constant  $c$  and dividing  $q_n$  by  $c^2$ , we may assume that

$$\max |b_{ij}^{(n)}| = 1 \quad \forall n. \quad (6)$$

Now, let us denote  $M_n := \max |a_{ij}(n)|$ .

Case 1. Assume that (a subsequence of)  $M_n$  is bounded:  $M_n \leq M$ . Note that in the case of the subsequence we may use again a replacement (5) and have that the whole sequence  $M_n$  is bounded. In this case, by using the Bolzano–Weierstrass theorem, we have for a subsequence  $\{n_k\}$  that

$$a_{ij}^{(n_k)} \rightarrow a_{ij}^0 \text{ and } b_{ij}^{(n_k)} \rightarrow b_{ij}^0, \quad \forall i, j.$$

Here, we use the fact that number of the coefficients  $a_{ij}$  and  $b_{ij}$  is finite.

By setting  $n = n_k$  in (4) and tending  $k \rightarrow \infty$  we obtain that

$$p_1 = q_0 r_0^2,$$

where

$$q_0 = \sum_{i+j \leq m_1} a_{ij}^0 x^i y^j, \quad r_0 = \sum_{i+j \leq m_2} b_{ij}^0 x^i y^j.$$

This contradicts the hypothesis for  $p_1$ .

Case 2. By taking into account a replacement (5) it remains to consider the case  $M_n \rightarrow +\infty$ .

In this case we have that for some subsequence  $n = \{n_k\}$ , we have that

$$|a_{i_0 j_0}^{(n_k)}| = \max_{i,j} |a_{ij}^{(n_k)}| \text{ and } |b_{i_1 j_1}^{(n_k)}| = \max_{i,j} |b_{ij}^{(n_k)}| = 1 \quad \forall k. \quad (7)$$

Here, we use again the fact that number of the coefficients  $a_{ij}$  and  $b_{ij}$  is finite. Also, in the last equality we take into account (6).

Now, let us set  $n = n_k$  in (4) and divide both sides by  $M_{n_k}$  to get

$$\frac{1}{M_{n_k}} p_1 + \frac{\epsilon_{n_k}}{M_{n_k}} p_2 = \left( \frac{1}{M_{n_k}} q_{n_k} \right) r_{n_k}^2. \quad (8)$$

Evidently, the left hand side here tends to zero. For the right hand side we have that the coefficients of the polynomials  $\frac{1}{M_{n_k}} q_{n_k}$  and  $r_{n_k}$  are bounded by 1. As above by using the Bolzano–Weierstrass theorem and passing to a new subsequence  $\{n'_k\} \subset \{n_k\}$  we have that

$$\frac{1}{M_{n'_k}} a_{ij}^{(n'_k)} \rightarrow a_{ij}^* \text{ and } b_{ij}^{(n'_k)} \rightarrow b_{ij}^*, \quad \forall i, j.$$

In view of (7) we have that

$$|a_{i_0 j_0}^*| = 1 \text{ and } |b_{i_1 j_1}^*| = 1. \quad (9)$$

Now, by setting  $n = n'_k$  in (4) and tending  $k \rightarrow \infty$  we get that

$$0 = q_* r_*^2,$$

where

$$q_* = \sum_{i+j \leq m_1} a_{ij}^* x^i y^j, \quad r_* = \sum_{i+j \leq m_2} b_{ij}^* x^i y^j.$$

In view of (9) this is a contradiction.  $\square$

**Remark 3.3.** *In the same way one can prove the following statement:*

*Assume that  $p_1, p_2 \in \Pi$ , and  $p_1$  is not reducible. Then, for sufficiently small  $\epsilon$ , the polynomial  $p_1 + \epsilon p_2$  is not reducible either.*



**Proposition 3.4.** *Suppose that  $\sigma_i$ ,  $i = 0, \dots, 6$  are seven linearly independent polynomials from  $\Pi_k$  passing through all the nodes of the set  $\mathcal{X}$  given in Theorem 2.4. Then, without loss of generality, one may assume that each of these polynomials is of exact degree  $k$  and has no multiple factors, or, alternatively there are three linearly independent polynomials from  $\Pi_{k-1}$  passing through all the nodes of  $\mathcal{X}$ .*

*Proof.* Clearly we may assume that a polynomial, say  $\sigma_0$ , is of exact degree  $k$ . Indeed, assume conversely that the degree of each of these seven polynomials is less than or equal to  $k-1$ . Then the conclusion of Proposition takes place (or we have also that this clearly contradicts Theorem 2.3).

Then we may assume that all the polynomials  $\sigma_i$ ,  $0 \leq i \leq 6$  are of exact degree  $k$ . Indeed, it suffices to replace these polynomials with the seven polynomials  $\sigma_0$  and  $\sigma_i + \epsilon\sigma_i$ ,  $1 \leq i \leq 6$ , for a sufficiently small  $\epsilon$ .

Next, we may assume that a polynomial, say  $\sigma_0$ , has no multiple factors, or, alternatively there are three linearly independent polynomials from  $\Pi_{k-1}$  passing through all the nodes of  $\mathcal{X}$ . Indeed assume conversely that each of these seven polynomials has a multiple factor. In view of Lemma 3.1 each of these multiple factors is a line with multiplicity two. Thus, we have that

$$\sigma_i = \ell_i^2 q_i, \quad 0 \leq i \leq 6, \quad \text{where } \ell_i \in \Pi_1, \quad q_i \in \Pi_{k-2}. \quad (10)$$

Then we replace these polynomials with the seven polynomials  $\check{\sigma}_i = \ell_i q_i \in \Pi_{k-1}$ ,  $0 \leq i \leq 6$ , which clearly vanish at the node set  $\mathcal{X}$ . Let us verify that among these latter seven polynomials there are at least three linearly independent ones. Conversely assume that the seven polynomials are linear combinations of two of them, say  $\check{\sigma}_i = \ell_i q_i$ ,  $i = 0, 1$ . Then we get readily that the seven linearly independent polynomials in (10) are linear combinations of the following six polynomials:

$$\ell_i q_i, x\ell_i q_i, y\ell_i q_i, \quad i = 0, 1,$$

which is a contradiction. Indeed, assume that  $\ell_i = A_i x + B_i y + C_i$ . Then for  $i = 1, 2$  we have that

$$\ell_i^2 q_i = (A_i x + B_i y + C_i) \ell_i q_i = A_i x \ell_i q_i + B_i y \ell_i q_i + C_i \ell_i q_i.$$

While, for  $i = 2, 6$  by assuming  $\ell_i q_i = a_i \ell_0 q_0 + b_i \ell_1 q_1$ , we have that

$$\begin{aligned} \ell_i^2 q_i &= (A_i x + B_i y + C_i) \ell_i q_i = (A_i x + B_i y + C_i)(a_i \ell_1 q_1 + b_i \ell_2 q_2) \\ &= a_i A_i x \ell_1 q_1 + a_i B_i y \ell_1 q_1 + a_i C_i \ell_1 q_1 + b_i A_i x \ell_2 q_2 + b_i B_i y \ell_2 q_2 + b_i C_i \ell_2 q_2. \end{aligned}$$

Finally, by assuming that  $\sigma_0$ , has no multiple factors, let us replace the seven polynomials  $\sigma_i$ ,  $0 \leq i \leq 6$ , with the seven polynomials  $\sigma_0$  and  $\sigma_i + \epsilon\sigma_i$ ,  $1 \leq i \leq 6$ , for a sufficiently small  $\epsilon$ . This, in view of Lemma 3.2, completes the proof.  $\square$

**Proposition 3.5.** *Suppose that  $\sigma_i$ ,  $i = 0, \dots, 6$ , are seven linearly independent polynomials from  $\Pi_k$  passing through all the nodes of the set  $\mathcal{X}$  given in Theorem 2.4. In addition suppose that each of these polynomials is of exact degree  $k$  and has no multiple factors. Then there is a polynomial in the linear span of  $\sigma_i$ ,  $i = 1, \dots, 6$ , which has no multiple factors and differs from  $\sigma_0$  with a factor of degree at least three.*

Let us prove first

**Lemma 3.6.** *Suppose that  $\sigma_0, s_1, s_2$  are linearly independent polynomials of degree  $k$ . In addition suppose that each of these polynomials has no multiple factors. Suppose that any polynomial in the linear span of  $s_i$ ,  $i = 1, 2$ , differs from  $\sigma_0$  with a factor of degree at most two. Then we have that*

$$\sigma_0 = \tilde{\sigma}_0 \beta_0, \quad s_1 = \tilde{\sigma}_0 \beta_1, \quad s_2 = \tilde{\sigma}_0 \beta_2, \quad \text{where } \beta_i \in \Pi_2. \quad (11)$$

Moreover, if  $\beta_0$  has a common factor with  $\beta_1$  as well as with  $\beta_2$  then following alternative takes place: either,

- (i)  $\beta_i = \ell \ell_i$ ,  $i = 0, 1, 2$ , i.e., they have a common linear factor,
- (ii)  $\beta_0$  and  $\beta_1 + \epsilon \beta_2$  are relatively prime  $\forall \epsilon > 0$ .

Furthermore,  $\tilde{\sigma}_0$  is uniquely determined from the first two relations in (11), if  $\beta_0$  and  $\beta_1$  are of degree two and relatively prime.

*Proof.* Consider the polynomials  $\sigma_0, s_1$  and  $s_2$ . In view of the hypotheses and Lemma 3.2 for sufficiently small  $\epsilon$  we have that

$$(s_1 + \epsilon s_2) \beta(c) = \sigma_0 \beta'(c), \quad (12)$$

where  $\beta_c, \beta'_c \in \Pi_2$  are relatively prime.

Note that  $\beta(c)$  is a linear or conic component of  $\sigma_0$ . Suppose that  $\sigma_0$  has  $k$  such components. By considering  $k + 1$  sufficiently small values of  $c$  we derive that there are constants  $c_1$  and  $c_2$  such that  $\beta(c_1) = \beta(c_2) =: \beta_0$ .

Then we readily get from (12) that

$$s_1 \beta_0 = \sigma_0 \beta_1 \quad \text{and} \quad s_2 \beta_0 = \sigma_0 \beta_2, \quad (13)$$

where  $\beta_1, \beta_2 \in \Pi_2$ .

In the case when  $\beta_0$  is relatively prime with  $\beta_1$  or  $\beta_2$  then it clearly divides  $\sigma_0$ . Note that this evidently takes place if  $\beta_0$  is linear. Then by denoting  $\sigma_0 = \tilde{\sigma}_0 \beta_0$  we get that

$$\sigma_0 = \tilde{\sigma}_0 \beta_0, \quad s_1 = \tilde{\sigma}_0 \beta_1, \quad s_2 = \tilde{\sigma}_0 \beta_2. \quad (14)$$

It remains to consider the case when  $\beta_0$  is a reducible conic and has a common linear component with  $\beta_1$  as well as with  $\beta_2$ . Thus suppose that  $\beta_0 = \ell_0 \ell'_0$ . After a cancellation with a linear polynomial in (13) two cases are possible:

Case 1.  $s_1\ell_0 = \sigma_0\ell_1$  and  $s_2\ell'_0 = \sigma_0\ell_2$

Case 2.  $s_1\ell_0 = \sigma_0\ell_1$  and  $s_2\ell_0 = \sigma_0\ell_2$

In Case 1  $\beta_0 = \ell_0\ell'_0$  again divides  $\sigma_0$  and we get (14).

In Case 2  $\beta_0 = \ell_0$  divides  $\sigma_0$  and we get (14), where  $\beta_0$  therefore  $\beta_1$  and  $\beta_2$  are linear. Thus (11) is proved.

Now, consider the ‘‘Moreover’’ case, i.e., each pair in  $\beta_i$ ,  $i = 1, 2, 3$ , has a common factor. Assume that  $\beta_0 = \ell\ell_0$  and  $\beta_1 = \ell\ell_1$ . Then we have that either  $\beta_2 = \ell\ell_2$ , or  $\beta_2 = \ell_0\ell_1$ . The first case reduces to the item (i). Let us consider the second case. In is easily seen that the polynomials  $\beta_0 = \ell\ell_0$  and  $\beta_0 + \epsilon\beta_2 = \ell_1(\ell + \epsilon\ell_0)$  have no common factor.

Finally note that if  $\beta_0$  and  $\beta_1$  have no common factor then  $\tilde{\sigma}_0$  is the greatest common divisor of  $\sigma_0$  and  $s_1$ . Hence it is uniquely determined from the first two relations in (11).  $\square$

Now we are in a position to present

*Proof of Proposition 3.5.* Assume by way of contradiction that any polynomial from

$$S := \text{Linear span}\{\sigma_1, \dots, \sigma_6\},$$

differs from  $\sigma_0$  with a factor of degree at most two.

According to Lemma 3.6 for the polynomial  $\sigma_0$  and any two polynomials from  $S$  the relation (11) takes place.

Case 1. Assume that there is a polynomial  $s_1 \in S$ , say it is  $s_1 = \sigma_1$ , for which the relation (11) holds with  $\beta_1$  having degree two and being relatively prime with  $\beta_0$ .

Then, according to Lemma 3.6,  $\tilde{\sigma}_0$  is determined uniquely.

Now, we apply Lemma 3.6 successively with the triples of polynomials  $\sigma_0, \sigma_1, \sigma_i$ ,  $i = 2, \dots, 6$ , and get that

$$\sigma_i = \tilde{\sigma}_0\beta_i, \quad i = 0, \dots, 6, \quad \text{where } \beta_i \in \Pi_2.$$

Clearly the seven polynomials  $\beta_i$  here, and consequently the seven polynomials  $\sigma_i$  are linearly dependent, which contradicts our assumption.

Case 2. Now, assume that for any triple of polynomials  $\sigma_0, s_1 := \sigma_i, s_2 := \sigma_j$  the relation (11) holds with  $\beta_0$  has a common factor with  $\beta_i$  as well as with  $\beta_j$  and all three have degree two.

If for some triple the alternative (ii) holds then we easily arrive to Case 1 with  $s_1 := \sigma_i + \epsilon\sigma_j$ .

Thus we conclude that the case of alternative (i) holds:

$$\beta_0 = \ell\ell_0, \beta_i = \ell\ell_i, \beta_j = \ell\ell_j.$$

In this case (11) holds also with  $\beta$ 's being linear polynomials:

$$\sigma_0 = \hat{\sigma}_0\ell_0, \sigma_i = \hat{\sigma}_0\ell_i, \sigma_j = \hat{\sigma}_0\ell_j, \quad \text{where } \hat{\sigma} = \tilde{\sigma}\ell.$$

Thus we conclude that for all triples  $\sigma_0, \sigma_i, \sigma_j$  (11) holds with linear  $\beta$ s. In this case  $\tilde{\sigma}_0$  is uniquely determined from the first two relations in (11).

Now, we apply Lemma 3.6, successively with the triples of polynomials  $\sigma_0, s_1 := \sigma_1, s_2 := \sigma_i, i = 2, 3$ , and get that

$$\sigma_i = \tilde{\sigma}_0 \ell_i, \quad i = 0, \dots, 3, \quad \text{where } \ell_i \in \Pi_1.$$

Clearly the four linear polynomials  $\ell_i$  here, and consequently the four polynomials  $\sigma_i$  are linearly dependent, which contradicts our assumption.  $\square$

## 4 The existence of three curves of degree $k - 1$

**Proposition 4.1.** *Assume that the hypotheses of Theorem 2.4 hold. Then, there are three linearly independent curves of degree  $k - 1$  passing through all the nodes of the set  $\mathcal{X}$ .*

*Proof.* Let  $\sigma_0, \dots, \sigma_6$  be the seven curves of degree  $\leq k$  that pass through all the nodes of the  $n$ -independent set  $\mathcal{X}$  with  $\#\mathcal{X} = d(n, k - 3) + 3$ .

In view of Proposition 3.4, without loss of generality, assume that each of these polynomials is of exact degree  $k$  and has no multiple factors.

**Step 1.** Here we will prove that there is at least one curve of degree  $k - 1$  passing through all the nodes of the set  $\mathcal{X}$ .

We start by choosing two nodes  $B_1, B_2 \notin \mathcal{X}$  such that the following two conditions are satisfied:

- i)* the set  $\mathcal{X} \cup \{B_1, B_2\}$  is  $n$ -independent;
- ii)* the line through  $B_1$  and  $B_2$  does not pass through any node from  $\mathcal{X}$ ;

Let us verify that one can find such nodes. Indeed, in view of Lemma 1.6, we can start by choosing some nodes  $B_i = B'_i, i = 1, 2$ , satisfying the condition *i)*. Then, according to Lemma 1.5, for some positive  $\epsilon$  all the nodes in  $\epsilon$  neighborhoods of  $B'_i, i = 1, 2$ , satisfy the condition *i)*.

Finally, from these neighborhoods we can choose the nodes  $B_i, i = 1, 2$ , satisfying the condition *ii)* too.

Denote by  $\ell_0$  the line passing through  $B_1$  and  $B_2$ .

Next we find one more node  $B_3 \in \ell_0$  such that the set  $\mathcal{X} \cup \{B_1, B_2, B_3\}$  is  $n$ -independent. Indeed, if there is no such node then we obtain that

$$p \in \Pi_k, \quad p|_{\mathcal{X} \cup \{B_1, B_2\}} = 0 \Rightarrow p|_{\ell_0} = 0.$$

Therefore  $p = \ell_0 q$ , where  $q \in \Pi_{k-1}$  and  $q|_{\mathcal{X}} = 0$ .

Hence, if there is no  $B_3$  then, according to Lemma 1.15, there are five linearly independent polynomials  $p \in \Pi_k$  satisfying the condition  $p|_{\mathcal{X} \cup \{B_1, B_2\}} = 0$ . Therefore, there are five linearly independent  $q \in \Pi_{k-1}$  satisfying the condition  $q|_{\mathcal{X}} = 0$ .

Next, we find successively two nodes  $B_4, B_5 \in \ell_0$  such that the set  $\mathcal{X} \cup \mathcal{B}_5$  is  $n$ -independent, where  $\mathcal{B}_5 := \{B_1, B_2, B_3, B_4, B_5\}$ .

Indeed, if one cannot find the nodes  $B_4$  or  $B_5$ , then, in the same way as above, there are four or three linearly independent polynomials  $q \in \Pi_{k-1}$  satisfying the condition  $q|_{\mathcal{X}} = 0$ , respectively.

Next, in view of Lemma 1.15, there are two curves of degree at most  $k$ , which pass through all the nodes of  $\mathcal{X} \cup \mathcal{B}_5$ .

Denote these curves by  $\sigma_0, \sigma'_0$ . Consider the curve  $\sigma_0$ . Notice that  $\sigma_0$  passes through more than  $d(n, k-3)$  nodes and, therefore, its degree equals either to  $k-2, k-1$  or  $k$ .

The case of degree  $k-2$  can be excluded in view of Lemma 3.1.

If the curve  $\sigma_0$  or  $\sigma'_0$  is of degree  $k-1$  then we are done.

Then, if a curve  $\sigma_0, \sigma'_0$  has a multiple factor then by throwing away the excessed factor we get again a polynomial from  $\Pi_{k-1}$  vanishing on  $\mathcal{X}$ , Q.E.D.

Thus we may assume that both curves  $\sigma_0$  and  $\sigma'_0$  are of degree  $k$  and have no multiple factors.

Consider again the curve  $\sigma_0$ . In view of Proposition 1.13, we can extend the set  $\mathcal{X} \cup \mathcal{B}_5$  till a maximal  $n$ -independent set  $\mathcal{Z} \subset \sigma_0$ , by adding  $d(n, k) - (d(n, k-3) + 3) - 5 = 3(n-k) + 1$  nodes, i.e.,

$$\mathcal{Z} = \mathcal{X} \cup \mathcal{B}_5 \cup \mathcal{A}, \text{ where } \#\mathcal{A} = 3(n-k) + 1 = [3(n-k-1) - 1] + 5.$$

Let us start with the description of the choice of  $3(n-k-1) - 1$  nodes from  $\mathcal{A}$ .

By using Proposition 3.5 we find a curve  $\sigma$  in the linear span of  $\sigma_i, i = 1, \dots, 6$ , which has no multiple factors and differs from  $\sigma_0$  with a factor of degree at least three:

$\sigma = \gamma q, \sigma_0 = \gamma_0 q, q \in \Pi_{k-d}$  with  $d = \deg \gamma = \deg \gamma_0 \geq 3$ . We have that  $\gamma$  and  $\sigma_0$  are relatively prime.

Consider a line  $\ell_1$  intersecting  $\gamma_0$  at three different points, say  $A_1, A_2$  and  $A_3$ , which do not belong to  $\sigma \cup \ell_0$ . By using a continuity argument we may assume that the line  $\ell_1$  is not parallel to any line component of  $\sigma_0$ . Let  $\ell_i, i = 1, \dots, n-k-1$ , be lines parallel to  $\ell_1$  and enough close to it so that each of them intersects  $\gamma_0$  at three different points, which do not belong to  $\sigma \cup \ell_0$ . We assume also that these intersection points are different from the nodes of  $\mathcal{X} \cup \mathcal{B}_5$ . Let us dismiss an intersection point, say  $A_1$  and denote the desired set of the remaining  $3(n-k-1) - 1$  intersection nodes by  $\mathcal{A}(-1)$ .

Let us prove that the set  $\mathcal{Y} := \mathcal{X} \cup \mathcal{B}_5 \cup \mathcal{A}(-1)$  is  $n$ -independent.

We have that the set  $\mathcal{A}(-1)$  is a subset of Berzolari-Radon construction of degree  $n-k-1$ . Hence it is  $(n-k-1)$ -independent. Now suppose that  $p_{A, \mathcal{A}(-1)}^*$  is a fundamental polynomial of a node  $A \in \mathcal{A}(-1)$  of degree  $n-k-1$ . Then the polynomial  $\sigma \ell_0 p_{A, \mathcal{A}(-1)}^*$  is an  $n$ -fundamental polynomial of the node  $A$  for the set  $\mathcal{Z}$ . Thus, according to Lemma 1.4, the set  $\mathcal{Z}$  is  $n$ -independent.

Finally, according to Proposition 1.13, we extend the set  $\mathcal{Y} \subset \sigma_0$  with the last 5 nodes till a maximal  $n$ -independent set  $\mathcal{Z} \subset \sigma_0$ . The set of these 5 points we denote by  $\mathcal{A}_5$ .

Thus the set  $\mathcal{Z} := \mathcal{Y} \cup \mathcal{A}_5$  is a maximal  $n$ -independent set in  $\sigma_0$ . We have also that  $\mathcal{A} = \mathcal{A}(-1) \cup \mathcal{A}_5$ .

Now suppose that  $\sigma^* \in \Pi_k$  vanishes on  $\mathcal{X}$  and  $\mathcal{A}_5$ . According to Lemma 1.15 there are  $2 = 7 - 5$  such polynomials. Consider the polynomial

$$p = \sigma^* \ell_0 \ell_1 \cdots \ell_{n-k-1}.$$

We have that  $p \in \Pi_n$  and it vanishes on the constructed maximal  $n$ -independent set  $\mathcal{Z} \subset \sigma_0$ . Therefore, we have that

$$\sigma^* \ell_0 \ell_1 \cdots \ell_{n-k-1} = \sigma_0 s, \text{ where } s \in \Pi_{n-k}.$$

The lines  $\ell_i$ ,  $i = 1, \dots, n - k - 1$ , are not components of  $\sigma_0$ . Therefore, they are components of the polynomial  $s$ . Thus we obtain that

$$\sigma^* \ell_0 = \sigma_0 \ell, \text{ where } \ell \in \Pi_1.$$

Since there are two linearly independent polynomials  $\sigma^*$  therefore  $\ell_0 \neq \ell$  and we have that  $\ell_0$  is a component of  $\sigma_0$  :

$$\sigma_0 = \ell_0 q, \text{ where } q \in \Pi_{k-1}.$$

We conclude from here that  $q$  vanishes on  $\mathcal{X}$ .

**Step 2.** Here we will prove that there are three linearly independent curves of degree  $k - 1$  passing through all the nodes of the set  $\mathcal{X}$ .

We find a line  $\ell_0$  and collinear nodes  $B_1, \dots, B_4 \in \ell_0$ , as in the Step 1, such that  $\ell_0 \cap \mathcal{X} = \emptyset$  and the set  $\mathcal{X} \cup \mathcal{B}_4$  is  $n$ -independent, where  $\mathcal{B}_4 := \{B_1, B_2, B_3, B_4\}$ .

Recall that if one cannot find the nodes then there are four linearly independent curves of degree  $\leq k - 1$  passing through all the nodes of  $\mathcal{X}$ .

Next, in view of Proposition 1.15, there are three linearly independent curves of degree at most  $k$ , which pass through all the nodes of the set  $\mathcal{X} \cup \mathcal{B}_4$ .

Denote these curves by  $\sigma_0, \sigma'_0, \sigma''_0$ .

If a curve here, say  $\sigma_0$  is of degree  $\leq k - 1$  and has no multiple components then instead of given triple of curves we consider the curves  $\ell_0 \sigma_0, \ell_1 \sigma_0, \ell_2 \sigma_0$ , where the lines  $\ell_i$  are chosen such that these three curves are linearly independent and have no multiple factors.

Next, if a curve  $\sigma_0, \sigma'_0, \sigma''_0$  has a multiple factor then by throwing away the excessed factor we get again a polynomial from  $\Pi_{k-1}$  vanishing on  $\mathcal{X} \cup \mathcal{B}_4$ , i.e., we are in the situation considered in the previous paragraph.

Hence, we may consider only the case when each of the given three polynomials is of exact degree  $k$  and has no multiple components.

Now consider the curve  $\sigma_0$ . In view of Proposition 1.13, we can extend the set  $\mathcal{X} \cup \mathcal{B}_4$  till a maximal  $n$ -independent set  $\mathcal{Z} \subset \sigma_0$ , by adding  $d(n, k) - (d(n, k - 3) + 3) - 4 = 3(n - k) + 2$  nodes, i.e.,

$$\mathcal{Z} = \mathcal{X} \cup \mathcal{B}_4 \cup \mathcal{A}, \text{ where } \#\mathcal{A} = 3(n - k) + 2 = [3(n - k - 1) - 1] + 1 + 5.$$

We find the set of  $3(n - k - 1) - 1$  points from  $\mathcal{A}$  in the same way as in Step 1 and denote it again by  $\mathcal{A}(-1)$ .

Then, in the same way as in Step 1, we prove the independence of the set  $\mathcal{Y} := \mathcal{X} \cup \mathcal{B}_4 \cup \mathcal{A}(-1)$ .

Next, we choose a node  $\tilde{A}_1 \in \ell_1$  such that  $\tilde{A}_1 \in \sigma_0 \setminus q$ , where  $q$  is one of the two polynomials of degree  $k - 1$  vanishing on  $\mathcal{X}$ , found in Step 1. We assume also that  $\tilde{A}_1$  does not belong to the line  $\ell_0$ .

Now let us prove the independence of the set  $\mathcal{Y}' := \mathcal{Y} \cup \{\tilde{A}_1\}$ . For this end, in view of Lemma 1.4, it suffices to find a fundamental polynomial of the node  $\tilde{A}_1$  with respect to the set  $\mathcal{Y}'$ .

We readily obtain that

$$p_{\tilde{A}_1, \mathcal{Y}'}^* = q\ell_0\ell_2 \cdots \ell_{n-k-1}\ell'\ell'',$$

where  $\ell'$  and  $\ell''$  are lines different from  $\ell_1$  and pass through the nodes  $A_2$  and  $A_3$ , respectively. Recall from Step 1 that these latter two nodes belong to  $\mathcal{A}(-1) \cap \ell_1$ .

Finally, as in Step 1, we extend the set  $\mathcal{Y}' \subset \sigma_0$ , according to Proposition 1.13, with the set of last 5 nodes, denoted by  $\mathcal{A}_5$ , till a maximal  $n$ -independent set in  $\sigma_0$ .

Thus the set  $\mathcal{Z} := \mathcal{Y}' \cup \mathcal{A}_5$  is a maximal  $n$ -independent set in  $\sigma_0$ .

Now suppose that  $\sigma^* \in \Pi_k$  vanishes on  $\mathcal{X}$  and the 5 nodes of  $\mathcal{A}_5$ . According to Lemma 1.15 there are at least two such polynomials. Consider the polynomial  $r = \sigma^*\ell_0\ell_1 \cdots \ell_{n-k-1}$ . We have that  $r \in \Pi_n$  and  $r$  vanishes on the node set  $\mathcal{Z}$ , which is a maximal  $n$ -independent set in the curve  $\sigma_0$ . Therefore, we have that

$$\sigma^*\ell_0\ell_1 \cdots \ell_{n-k-1} = \sigma_0 s, \text{ where } s \in \Pi_{n-k}.$$

The lines  $\ell_i$ ,  $i = 1, \dots, n - k - 1$ , are not components of  $\sigma_0$ . Therefore, they are components of the polynomial  $s$ . Thus we obtain that

$$\sigma^*\ell_0 = \sigma_0 \ell, \text{ where } \ell \in \Pi_1.$$

Since there are two linearly independent polynomials  $p$  therefore  $\ell_0 \neq \ell$  and we have that  $\ell_0$  is a component of  $\sigma_0$ .

Thus we conclude that

$$\sigma_0 = \ell_0 q_{k-1}, \text{ where } q_{k-1} \in \Pi_{k-1}.$$

In the same way by considering the curves  $\sigma'_0$  and  $\sigma''_0$  we get

$$\sigma'_0 = \ell_0 q'_{k-1}, \text{ where } q'_{k-1} \in \Pi_{k-1}.$$

$$\sigma''_0 = \ell_0 q''_{k-1}, \text{ where } q''_{k-1} \in \Pi_{k-1}.$$

Thus we obtain three linearly independent curves  $q_{k-1}, q'_{k-1}, q''_{k-1} \in \Pi_{k-1}$  vanishing at all the nodes of  $\mathcal{X}$ .  $\square$

## 5 Proofs of Theorems 2.4 and 2.5

Let us start with

*Proof.* Proof of Theorem 2.5.

Assume by way of contradiction that there are four curves passing through all the nodes of the set  $\mathcal{X}$ . Then, according to Theorem 2.3, all the nodes of  $\mathcal{X}$  but three belong to a maximal curve  $\mu$  of degree  $k-2$ . The curve  $\mu$  is maximal and the remaining three nodes of  $\mathcal{X}$ , denoted by  $A, B$  and  $C$ , are outside of it:  $A, B, C \notin \mu$ . Hence we have that

$$\mathcal{P}_{k,\mathcal{X}} = \{p : p \in \Pi_k, p_{\mathcal{X}} = 0\} = \{q\mu : q \in \Pi_2, q(A) = q(B) = q(C) = 0\}.$$

Thus we get readily that  $\dim \mathcal{P}_{k,\mathcal{X}} = \dim \{q \in \Pi_2 : q(A) = q(B) = q(C) = 0\} = \dim \mathcal{P}_{2,\{A,B,C\}} = 6 - 3 = 3$ , which contradicts our assumption. Note that in the last equality we use Proposition 1.7 and the fact that any three nodes are 2-independent.

Consider the three curves  $\sigma_i \in \Pi_k$   $i = 0, 1, 2$ , passing through all the nodes of  $\mathcal{X}$ . If one of them is of degree  $k-1$  then the conclusion of Theorem is satisfied and we are done. Thus, we may assume that each curve is of degree  $k$  and has no multiple component.

Now consider the curve  $\sigma_0$ .

By using Proposition 1.13 let us extend the set  $\mathcal{X}$  till a maximal  $n$ -independent set  $\mathcal{Z} \subset \sigma_0$ . Since  $\#\mathcal{Z} = d(n, k)$ , we need to add a set of  $d(n, k) - (d(n, k-2) + 3) = 2(n-k) + 2$  nodes, denoted by

$$\mathcal{A} := \{A_1, \dots, A_{2(n-k)+2}\}.$$

Thus we have that  $\mathcal{Z} := \mathcal{X} \cup \mathcal{A}$ .

In view of Lemma 1.14, *i*), we require that each node of  $\mathcal{A}$  may belong only to one component of the curve  $\sigma_0$ .

**Case 1**,  $n = k + 2$ ,  $\mathcal{A} := \{A_1, \dots, A_6\}$ .

Consider 5 nodes from  $\mathcal{A}$  and a conic  $\beta^*$  passing through them. Denote the sixth node by  $A^*$ .

We have three polynomials from  $\Pi_k$  vanishing on  $\mathcal{X}$ . By using Lemma 1.15 we get two linearly independent curves denoted by  $\sigma^*$ , of degree at most  $k$ , that pass through all the nodes of  $\mathcal{X}$  and the node  $A^* \in \mathcal{A}$ .



Thus we may consider a curve  $\sigma^*$  by assuming that  $\sigma^* \neq \sigma_0$ . Now, notice that the polynomial  $\sigma^* \beta^*$  of degree  $n$  vanishes at all the nodes of  $\mathcal{Z} \subset \sigma_0$ . Consequently, according to Proposition 1.10,  $\sigma_0$  divides this polynomial:

$$\sigma^* \beta^* = \sigma_0 \beta, \quad \beta \in \Pi_2. \quad (15)$$

We have that  $\beta^* \neq \beta$  since  $\sigma^* \neq \sigma_0$ .

Hence if  $\beta^*$  is irreducible then it divides  $\sigma_0$ .

Now suppose that  $\beta^*$  is reducible:  $\beta^* = \ell_1 \ell_2$ , where  $\ell_i \in \Pi_1$ .

Then we have that both lines  $\ell_1, \ell_2$  cannot divide  $\beta$ , hence we obtain that either both lines are component of  $\sigma_0$  or only one of the lines  $\ell_1, \ell_2$  is a component of  $\sigma_0$ .

Let us consider the latter case. Suppose that only the line  $\ell_1$  is a component of  $\sigma_0$  and  $\ell_2$  is a component of  $\beta$ . Then we get from (15) that

$$\sigma^* \ell_1 = \sigma_0 \ell, \quad \text{where } \ell \in \Pi_1. \quad (16)$$

Now, we have that  $\sigma_0 = \ell_1 \bar{\sigma}_0$ , where  $\bar{\sigma}_0 \in \Pi_{k-1}$ .

Therefore we get from (16) that  $\sigma^* = \ell \bar{\sigma}_0$ .

From the last two equalities we conclude that  $\mathcal{X} \subset \bar{\sigma}_0 \cup \{E\}$ , where  $E = \ell_1 \cap \ell$ .

Therefore all the nodes of  $\mathcal{X}$ , except possibly  $E$ , belong to the curve  $\bar{\sigma}_0 \in \Pi_{k-1}$ . Here  $\bar{\sigma}_0 \in \Pi_{k-1}$  is a component of  $\sigma_0$  and  $E$  belongs to a line component of  $\sigma_0$ .

We briefly express the above conditions by saying that the line component  $\ell_1$  of  $\sigma_0$  satisfies  $(-1)$  node condition for  $\mathcal{X}$ .

At the end we will see that if this property holds for all three given curves then we can readily complete the proof of Theorem.

Therefore, from now on we may assume that the equality (15) implies that  $\deg \beta^* = 2$  and  $\beta^*$  is a component of  $\sigma_0$ . We obtain also that  $\beta^*$  is determined uniquely by the 5 nodes from  $\mathcal{A}$ .

Next, we are going to prove that there is a conic passing through all the six nodes of  $\mathcal{A}$ . Assume by way of contradiction that there is no such conic. Denote by  $\beta_i$  the conic passing through the five nodes of  $\mathcal{A}_6 \setminus \{A_i\}$ ,  $i = 1, 2$ .

We readily get that these two conics are different. Then, by the above argument, we have that  $\beta_1$  and  $\beta_2$  are components of  $\sigma_0$ . First assume that one of these two conics, say,  $\beta_1$ , is irreducible. Then consider a common node of  $\beta_1$  and  $\beta_2$ , say,  $A_3$ . It is easily seen that  $A_3$  belongs to two different components of  $\sigma_0$ . One is  $\beta_1$  and another is  $\beta_2$  if it is irreducible or a line component of  $\beta_2$  if it is reducible.

Now, assume that both  $\beta_1$  and  $\beta_2$  are reducible:

$$\beta_1 = \ell_1 \ell'_1, \quad \beta_2 = \ell_2 \ell'_2.$$

Without loss of generality assume that

$$\ell_1 \neq \ell_2, \quad \ell_1 \neq \ell'_2. \quad (17)$$

Now we have that  $\ell_1$  passes through at least one of the common nodes  $A_3, \dots, A_6$ , say  $A_3$ . Then  $A_3$  belongs either to  $\ell_2$  or  $\ell'_2$ . In both cases, in view of (17), we have that  $A_3$  belongs to two different line components of  $\sigma_0$ , which is a contradiction.

Thus we proved that  $\mathcal{A} \subset \beta_0$ , where  $\beta_0 \in \Pi_2$ . Next let us show that  $\beta_0$  divides  $\sigma_0$  or the conclusion of Theorem holds.

Consider a polynomial  $\sigma \in \Pi_k$  that vanishes on  $\mathcal{X}$ . Notice that the following polynomial  $\sigma \beta_0$  of degree  $k + 2 \leq n$  vanishes at all the  $d(n, k)$  nodes of  $\mathcal{Z} \subset \sigma_0$ . Consequently, according to Proposition 1.10,  $\sigma_0$  divides this polynomial:

$$\sigma \beta_0 = \sigma_0 \beta, \quad \beta \in \Pi_2. \quad (18)$$

This is a type (15) equality which, as we mentioned above, implies that  $\deg \beta_0 = 2$  and  $\beta_0$  is a component of  $\sigma_0$  :

$$\sigma = \beta_0 q, \quad q \in \Pi_{k-2}.$$

We conclude also that  $\beta_0$  is uniquely determined by any 5 nodes from  $\mathcal{A}$ .

Thus we get that when completing the set  $\mathcal{X} \subset \sigma_0$  till a maximal  $n$ -independent set  $\mathcal{Z} = \mathcal{X} \cup \mathcal{A}$  we have to add all the six nodes of  $\mathcal{A}$  to the conic  $\beta_0$ . Let us verify that the added nodes cannot belong to another component  $q$ . Indeed, suppose conversely that a node belongs to  $\beta_0 \cap q$ . Then, in view of Lemma 1.14, we can move the node to  $q \setminus \beta_0$  such that the resulted set is also  $n$ -independent. This is a contradiction, since now clearly the six nodes do not belong to a conic. Indeed, the five nodes determine a unique conic and the sixth node is outside of it. This shows that the factor  $r \in \Pi_{k-2}$  to which one can not add a new independent node is merely maximal with respect to  $\mathcal{X}$ . This means that  $r$  passes through exactly  $d(n, k)$  nodes of  $\mathcal{X}$ .

**Case 2,  $n \geq k + 3$ .**

Consider a subset of  $\mathcal{A}$  of cardinality 4 and denote it by  $\mathcal{A}_4$ . Denote also by  $\bar{\mathcal{A}} := \mathcal{A} \setminus \mathcal{A}_4$ . We have that  $\#\bar{\mathcal{A}} = 2(n - k) - 2$ .

We have three polynomials from  $\Pi_k$  vanishing on  $\mathcal{X}$ . By using Lemma 1.15 we get two linearly independent curves denoted by  $\sigma^*$ , of degree at most  $k$ , that pass through all the nodes of  $\mathcal{X}$  and an arbitrary node  $A^* \in \bar{\mathcal{A}}$ , which will be specified below. We call this latter node associated with  $\sigma^*$ . Thus we may assume that  $\sigma^* \neq \sigma_0$ .

We associate another node  $A' \in \bar{\mathcal{A}}$  with the set  $\mathcal{A}_4$  and denote by  $\beta'$  a conic that passes through the four nodes of  $\mathcal{A}_4$  and  $A'$ .

For any line component  $\ell$  of  $\sigma_0$  denote by  $r_\ell$  the polynomial from  $\Pi_{k-1}$  for which

$$\sigma_0 = \ell r_\ell \quad (19)$$

holds.

Assume that a line component  $\ell$  of the curve  $\sigma_0$ , passes through exactly  $m$  nodes from  $\mathcal{X}$ , at which  $r_\ell$  does not vanish. Then we obtain from (19)

that  $r_\ell \in \Pi_{k-1}$  vanishes at the all nodes of the set  $\mathcal{X}$  except  $m$  nodes, which belong to  $\ell$ .

Thus, either for a line component we have that  $m \leq 1$ , or for all line components of  $\sigma_0$  we have that  $m \geq 2$ , meaning that the following condition takes place:

(C) Any line component of the curve  $\sigma_0$ , passes through at least two nodes from  $\mathcal{X}$ , at which  $r_\ell$  does not vanish.

It is easily seen that the condition  $m \leq 1$ , for  $\ell$  means that the line component  $\ell$  of  $\sigma_0$  satisfies the  $(-1)$  node condition for  $\mathcal{X}$ .

Later, by using the condition (C), we intend to divide the set of nodes  $\bar{\mathcal{A}}$  into  $n - k - 2$  pairs such that the lines  $\ell_1, \dots, \ell_{n-k-2}$ , through them, respectively, are not components of  $\sigma_0$ . The remaining two nodes denoted by  $A^*$  and  $A'$  we associate with the curve  $\sigma^*$  and  $\mathcal{A}_4$ , respectively.

Now, let us continue the proof by assuming that the above-described division of  $\bar{\mathcal{A}}$  is established.

Notice that the following polynomial  $\sigma^* \beta' \ell_1 \dots \ell_{n-k-2}$  of degree  $n$  vanishes at all the  $d(n, k)$  nodes of  $\mathcal{Z} \subset \sigma_0$ . Consequently, according to Proposition 1.10,  $\sigma_0$  divides this polynomial:

$$\sigma^* \beta' \ell_1 \ell_2 \dots \ell_{n-k-2} = \sigma_0 r, \quad r \in \Pi_{n-k}. \quad (20)$$

The distinct lines  $\ell_1, \ell_2, \dots, \ell_{n-k-2}$  do not divide the polynomial  $\sigma_0 \in \Pi_k$ , therefore, all they have to divide  $r$ . Hence, we get from (20):

$$\sigma^* \beta' = \sigma_0 \beta, \quad \text{where } \beta \in \Pi_2. \quad (21)$$

Now, we have that  $\beta' \neq \beta$  since  $\sigma^* \neq \sigma_0$ . Then, in the same way as in the Case 1 we obtain that  $\beta'$  divides  $\sigma_0$ :

$$\sigma_0 = \beta' q \quad \text{where } q \in \Pi_{k-2}. \quad (22)$$

Next, we are going to prove that there is a conic passing through all the nodes of  $\mathcal{A}$ . Assume by way of contradiction that there is no such conic. Then, in view of Proposition 1.9, we have that there is a set of six nodes, say  $\mathcal{A}_6 := \{A_1, \dots, A_6\} \subset \mathcal{A}$ , that does not lie in a conic, i.e., it is 2-posed.

Now, let us choose three noncollinear nodes in  $\mathcal{A}_6$ , say  $A_1, A_2, A_3$ , and consider the following sets of four nodes:

$$A_1, A_2, A_3, A_4; \quad A_1, A_2, A_3, A_5; \quad A_1, A_2, A_3, A_6.$$

Now, consider these three sets with the respective associated nodes:

$$A_1, A_2, A_3, A_4, A'; \quad A_1, A_2, A_3, A_5, A''; \quad A_1, A_2, A_3, A_6, A'''. \quad (23)$$

We have that the three conics through these sets are components of  $\sigma_0$ . Since  $\mathcal{A}_6$  is 2-posed we obtain that these three conics cannot coincide. Hence there

are two different conics, say the conics  $\beta', \beta''$ , passing through the first two sets in (23), respectively.

First assume that one of these two conics, say,  $\beta'$ , is irreducible. Then consider a common node, say,  $A_1$ . It is easily seen that  $A_1$  belongs to two different components of  $\sigma_0$ . One is  $\beta'$  and another is  $\beta''$ , if it is irreducible too, or a line component of  $\beta''$ , if it is reducible.

Next, assume that both  $\beta'$  and  $\beta''$  are reducible:

$$\beta' = \ell_1 \ell'_1, \quad \beta'' = \ell_2 \ell'_2.$$

Without loss of generality assume that

$$\ell_1 \neq \ell_2, \quad \ell_1 \neq \ell'_2. \quad (24)$$

Note that  $\ell_1$  passes through at least one of the common nodes  $A_1, A_2, A_3$ , say  $A_1$ . Indeed, if  $\ell_1$  passes through only  $A'$  and  $A_4$  then we obtain that  $\ell'_1$  passes through the three nodes  $A_1, A_2, A_3$ . This is a contradiction since these three nodes are not collinear. Now, we have that  $A_1$  belongs either to  $\ell_2$  or  $\ell'_2$ . In both cases, in view of (24), we have that  $A_1$  belongs to two different line components of  $\sigma_0$ , which is a contradiction.

Thus we proved that  $\mathcal{A} \subset \beta_0$ , where  $\beta_0 \in \Pi_2$ . Next, in the same way as in Case 1, we show that  $\beta_0$  divides  $\sigma_0$ :

$$\sigma_0 = \beta_0 q, \quad q \in \Pi_{k-2}.$$

Also we have that  $\beta_0$  is uniquely determined by the nodes of  $\mathcal{A} \setminus \{A\}$ ,  $\forall A \in \mathcal{A}$ .

Indeed, assume that  $\beta_0$  is not uniquely determined by the nodes from  $\mathcal{A} \setminus \{A_0\}$ , where  $A_0 \in \mathcal{A}$ . Therefore there are infinitely many conics  $\beta_0$  passing through the nodes of  $\mathcal{A} \setminus \{A_0\}$ . Recall that for (any)  $A_0$  one can find a curve, denoted by  $\sigma^*$ , of degree at most  $k$ , that passes through all the nodes of  $\mathcal{X}$  and is different from  $\sigma_0$ .

Then, as in Case 1, we readily get  $\sigma^* \beta_0 = \sigma_0 \beta$ , where  $\beta \in \Pi_2$ . This implies that  $\beta_0$  is a component of  $\sigma_0$ . Hence we get that  $\sigma_0$  has infinitely many components, which is a contradiction.

Thus we get that when completing the set  $\mathcal{X} \subset \sigma_0$  till a maximal  $n$ -independent set  $\mathcal{Z} = \mathcal{X} \cup \mathcal{A}$  we have to add all the nodes of  $\mathcal{A}$  to the conic  $\beta_0$ . Let us verify that added nodes cannot belong to another component  $q$ . Indeed, suppose conversely that a node  $A_0 \in \mathcal{A}$  belongs to  $\beta_0 \cap q$ . Then, in view of Lemma 1.14, we can move  $A_0$  to  $q \setminus \beta_0$  such that the resulted set  $\mathcal{A}$  remains  $n$ -independent. This is a contradiction, since now clearly the nodes of  $\mathcal{A}$  do not belong to a conic. Indeed, the nodes  $\mathcal{A} \setminus \{A_0\}$  determine a unique conic and the moved node is outside of it. This shows that the factor  $r \in \Pi_{k-2}$  to which one cannot add a new independent node is merely maximal with respect to  $\mathcal{X}$ . Therefore  $r$  passes through exactly  $d(n, k)$  nodes of  $\mathcal{X}$ .

At the end, before establishing the division of the set  $\bar{\mathcal{A}}$ , it remains to consider the case when the division may be not possible for all three curves  $\sigma_0$ , i.e., the case when the condition (C) does not hold. Then, we obtain three curves  $q_i$ ,  $i = 0, 1, 2$ , which are components of degree  $k - 1$  of the curves  $\sigma_i$ ,  $i = 0, 1, 2$ , respectively, passing through all the nodes of  $\mathcal{X}$  except possibly one. Since  $\#\mathcal{X} = d(n, k-2) + 3$  we get that  $\deg q_i = k - 1$ ,  $i = 0, 1, 2$ ,

Assume that  $q_i$  passes through all the nodes of  $\mathcal{X}$  except  $E_i$ ,  $E_i \in \mathcal{X}$ ,  $i = 0, 1, 2$ .

First assume that two of these three nodes are different, say  $E_0 \neq E_1$ .

We have that  $q_0$  and  $q_1$  pass through all the nodes of the set  $\mathcal{Y} := \mathcal{X} \setminus \{E_0, E_1\}$ ,  $\#\mathcal{Y} = d(n, k - 3) + 1$ .

If  $q_0 \neq q_1$  then, according to Theorem 2.2, all the nodes of  $\mathcal{Y}$  except one belong to a (maximal) curve  $\mu$  of degree  $k - 2$ . Thus all the nodes of  $\mathcal{X}$  except three belong to  $\mu$ .

Now assume that  $q_0 = q_1 =: q$ . Then we have that  $E_0 = E_1$ , contradicting our assumption.

It remains to consider the case  $E_0 = E_1 = E_2 =: E$ .

Then we have that  $q_0, q_1$  and  $q_2$  pass through all the nodes of the set  $\mathcal{Y} := \mathcal{X} \setminus \{E\}$ ,  $\#\mathcal{Y} = d(n, k - 2) + 2$ . We get from Theorem 2.1 that  $q_0 = q_1 = q_2 =: q$ .

Next, in view of the condition (C), we get that  $\sigma_i = \ell_i q$ ,  $i = 0, 1, 2$ , where  $\ell_i \in \Pi_1$ . This contradicts the linear independence of  $\sigma_i$ ,  $i = 0, 1, 2$ , since we have that  $E \in \ell_i$ ,  $i = 0, 1, 2$ .  $\square$

## 5.1 The division of the set $\bar{\mathcal{A}}$

Next let us establish the above mentioned division of the node set  $\mathcal{A}$  in the case  $n \geq k + 3$ . Note that this is the case when we need the diision.

Recall that each node of  $\mathcal{A}$  belongs only to one component of the curve  $\sigma_0$ . Particularly, the line components of  $\sigma_0$  do not intersect at the nodes of  $\mathcal{A}$ .

By using induction on  $m$  one can prove easily the following

**Lemma 5.1.** *Suppose that a finite set of lines  $\mathcal{L}$  and  $2m$  nodes lying in these lines are given. Suppose also that no node is an intersection point of two lines. Then one can divide the node set into  $m$  pairs such that no pair belongs to the same line from  $\mathcal{L}$  if and only if each line from  $\mathcal{L}$  contains no more than  $m$  nodes.*

Thus the above mentioned division of the node set  $\bar{\mathcal{A}}$  into  $n - k - 2$  pairs is possible if and only if no  $n - k - 1$  nodes of  $\bar{\mathcal{A}}_0 := \mathcal{A} \setminus \{A^*, A_0\}$  are located in a line component of  $\sigma_0$ , where the nodes  $A^*$  and  $A_0$  are the nodes associated with the curve  $\sigma^*$  and  $\mathcal{A}_4$ , respectively. Observe also that we may associate any two nodes  $A^*$  and  $A_0$  of  $\mathcal{A}$  with  $\sigma^*$  and  $\mathcal{A}_4$ ,

Now notice that, in view of  $\#\bar{\mathcal{A}} = 2(n - k - 1)$ , there can be at most two undesirable line components for the set  $\bar{\mathcal{A}}$ , i.e., lines containing at least  $n - k - 1$  nodes from it. In this case a node from each line we assign as associated and leave in the two lines  $\leq n - k - 2$  nodes.

Then assume that we have one undesirable line component for the set  $\bar{\mathcal{A}}$ , containing  $\leq n - k$  nodes from it. In this case two nodes from this line we nominate as associated and leave in the line  $\leq n - k - 2$  nodes.

Finally consider the case of one undesirable line component  $\ell$  of  $\sigma_0$  with  $m \geq n - k + 1$  nodes. We have that

$$\sigma_0 = \ell r_\ell, \text{ where } r_\ell \in \Pi_{k-1}.$$

Now we are going to move  $m - n + k$  nodes, one by one, from  $\ell$  to the other component  $r_\ell$  such that the set  $\mathcal{Z} := \mathcal{X} \cup \mathcal{A}$  remains  $n$ -independent. Again, in view of Lemma 1.14, i), we require that each moved node belongs only to one component of the curve  $\sigma_0$ .

To establish each described movement, in view of Lemma 1.14, ii), it suffices to prove that during this process each node  $A \in \ell \cap \mathcal{A}$ , has no  $n$ -fundamental polynomial for which the curve  $r_\ell$  is a component. Suppose conversely that  $p_A^* = r_\ell s$ ,  $s \in \Pi_{n-k+1}$ . Now, we have that  $s$  vanishes at  $\geq n - k$  nodes in  $\ell \cap \mathcal{A} \setminus \{A\}$ . Indeed, the nodes of the set  $\mathcal{A}$  in the line  $\ell$  do not belong to another component. Therefore,  $r_\ell$  does not vanish at these nodes and hence  $s$  vanishes. According to the condition (C)  $r_\ell$  does not vanish also at least at two nodes from  $\ell \cap \mathcal{X}$ , and hence  $s$  vanishes there too. Thus the number of zeroes of  $s$  in the line  $\ell$  is greater or equal to  $n - k + 2$  and  $s$  together with  $p_A^*$  vanishes at the whole line  $\ell$ , including the node  $A$ , which is a contradiction.

It remains to note that there will be no more undesirable lines, except  $\ell$ , in the resulted set  $\mathcal{A}$ , after the described movement of the nodes, since we finish by keeping exactly  $n - k$  nodes in  $\ell \cap \mathcal{A}$  and outside of it there are only  $n - k - 2$  nodes.

**Proof of Theorem 2.4.** It is easily seen that Theorem 2.4 follows from Proposition 4.1 and Theorem 2.5.

## 6 An application to bivariate interpolation

Recall that a node  $A \in \mathcal{X}$  uses a line  $\ell$  means that  $\ell$  is a factor of the fundamental polynomial  $p = p_A^*$ , i.e.,  $p = \ell r$  for some  $r \in \Pi_{n-1}$ .

A  $GC_n$ -set in the plane is an  $n$ -poised set of nodes, where the fundamental polynomial of each node is a product of  $n$  linear factors. The Gasca-Maeztu conjecture states that any  $GC_n$ -set possesses a subset of  $n+1$  collinear nodes.

It was proved in [9], that any line passing through exactly 2 nodes of a  $GC_n$ -set  $\mathcal{X}$  can be used at most by one node from  $\mathcal{X}$ .

Next, it was proved in [8] that any used line passing through exactly 3 nodes of a  $GC_n$ -set  $\mathcal{X}$  can be used either by exactly one or three nodes from  $\mathcal{X}$ .

In [7] was proved the following result concerning the lines passing through exactly 4 nodes.

**Corollary 6.1.** *Let  $\mathcal{X}$  be an  $n$ -poised set of nodes and  $\ell$  be a line, which passes through exactly 4 nodes. Suppose  $\ell$  is used by at least four nodes from  $\mathcal{X}$ . Then it is used by exactly six nodes from  $\mathcal{X}$ . Moreover, if it is used by six nodes, then they form a 2-poised set. Furthermore, in the latter case, if  $\mathcal{X}$  is a  $GC_n$  set, then the six nodes form a  $GC_2$  set.*

Below we consider the case of lines passing through exactly 5 nodes.

**Corollary 6.2.** *Let  $\mathcal{X}$  be an  $n$ -poised set of nodes and  $\ell$  be a line which passes through exactly 5 nodes. Suppose  $\ell$  is used by at least seven nodes from  $\mathcal{X}$ . Then it is used by exactly ten nodes from  $\mathcal{X}$ . Moreover, if it is used by ten nodes, then they form a 3-poised set. Furthermore, in the latter case, if  $\mathcal{X}$  is a  $GC_n$  set then the ten nodes form a  $GC_3$  set.*

*Proof.* Assume that  $\ell \cap \mathcal{X} = \{A_1, \dots, A_5\} =: \mathcal{A}$ . Assume also that the seven nodes in  $\mathcal{B} := \{B_1, \dots, B_7\} \in \mathcal{X}$  use the line  $\ell : p_{B_i}^* = \ell q_i$ ,  $i = 1, \dots, 7$ , where  $q_i \in \Pi_{n-1}$ .

The polynomials  $q_1, \dots, q_7$ , vanish at  $N - 12$  nodes of the set  $\mathcal{X}' := \mathcal{X} \setminus (\mathcal{A} \cup \mathcal{B})$ . Hence through these  $N - 12 = d(n, n - 4) + 3$  nodes pass seven linearly independent curves of degree  $n - 1$ . By Theorem 2.4 there exists a maximal curve  $\mu$  of degree  $n - 4$  passing through  $N - 15$  nodes of  $\mathcal{X}'$  and the remaining three nodes denoted by  $C_1, C_2, C_3$ , are outside of it. Now, according to Proposition 1.12, the nodes  $C_1, C_2, C_3$ , use  $\mu$  :

$$p_{C_i}^* = \mu r_i, \quad r_i \in \Pi_4, \quad i = 1, 2, 3.$$

These polynomials  $r_i$  have to vanish at the five nodes of  $\mathcal{A} \subset \ell$ . Hence  $q_i = \ell \gamma_i$ ,  $i = 1, 2, 3$ , with  $\gamma_i \in \Pi_3$ . Therefore, the nodes  $C_1, C_2, C_3$ , use the line  $\ell$  :

$$p_{C_i}^* = \mu \ell \gamma_i, \quad i = 1, 2, 3.$$

Hence if seven nodes in  $\mathcal{B} \subset \mathcal{X}$  use the line  $\ell$  then there exist three more nodes  $C_1, C_2, C_3 \in \mathcal{X}$  using it and all the nodes of  $\mathcal{Y} := \mathcal{X} \setminus (\mathcal{A} \cup \mathcal{B} \cup \{C_1, C_2, C_3\})$  lie in a maximal curve  $\mu$  of degree  $n - 4$  :

$$\mathcal{Y} \subset \mu. \tag{25}$$

Next, let us show that there is no eleventh node using  $\ell$ . Assume by way of contradiction that except of the ten nodes in

$$\mathcal{S} := \{B_1, \dots, B_7, C_1, C_2, C_3\},$$

there is an eleventh node  $D$  using  $\ell$ . Of course we have that  $D \in \mathcal{Y}$ .

Then we have that the seven nodes  $B_1, \dots, B_6$  and  $D$  are using  $\ell$  therefore, as was proved above, there exist three more nodes  $E_1, E_2, E_3 \in \mathcal{X}$  (which may coincide or not with  $B_7$  or  $C_1, C_2, C_3$ ) using it and all the nodes of  $\mathcal{Y}' := \mathcal{X} \setminus (\mathcal{A} \cup \{B_1, \dots, B_6, D, E_1, E_2, E_3\})$  lie in a maximal curve  $\mu'$  of degree  $n - 4$ .

We have also that

$$p_D^* = \mu' q', \quad q' \in \Pi_4. \quad (26)$$

Now, notice that both the curves  $\mu$  and  $\mu'$  pass through all the nodes of the set  $\mathcal{Z} := \mathcal{X} \setminus (\mathcal{A} \cup \mathcal{B} \cup \{C_1, C_2, C_3, D, E_1, E_2, E_3\})$  with  $|\mathcal{Z}| \geq N - 19$ .

Then, we get from Theorem 2.1, with  $k = n - 5$ , that  $N - 19 = d(n, n - 5) + 2$  nodes determine the curve of degree  $n - 4$  passing through them uniquely. Thus  $\mu$  and  $\mu'$  coincide.

Therefore, in view of (25) and (26),  $p_D^*$  vanishes at all the nodes of  $\mathcal{Y}$ , which is a contradiction since  $D \in \mathcal{Y}$ .

Now, let us verify the last “moreover” statement. Suppose ten nodes in  $\mathcal{S} \subset \mathcal{X}$  use the line  $\ell$ . Then, as we obtained earlier, the nodes  $\mathcal{Y} := \mathcal{X} \setminus (\mathcal{A} \cup \mathcal{S})$  are located in a maximal curve  $\mu$  of degree  $n - 4$ . Therefore the fundamental polynomial of each  $A \in \mathcal{S}$  uses  $\mu$  and hence  $\ell$ :

$$p_A^* = \mu \ell q_A, \quad \text{where } q_A \in \Pi_3.$$

It is easily seen that  $q_A$  is a 3-fundamental polynomial of  $A \in \mathcal{S}$ . □

## REFERENCES

- [1] Eisenbud D., Green M., Harris J. Cayley-Bacharach Theorems and Conjectures. *Bull. Amer. Math. Soc. (N.S.)*, **33**:3 (1996), 295–324.
- [2] Hakopian H., Jetter K., Zimmermann G. Vandermonde Matrices for Intersection Points of Curves. *Jaen J. Approx.*, **1** (2009), 67–81.
- [3] Hakopian H., Malinyan A. Characterization of  $n$ -Independent Sets with No More Than  $3n$  Points. *Jaen J. Approx.*, **4**:2 (2012), 121–136.
- [4] Hakopian H. Multivariate Divided Differences and Multivariate Interpolation of Lagrange and Hermite Type. *J. Approx. Theory*, **34** (1982), 286–305.
- [5] Hakopian H., Toroyan S. On the Uniqueness of Algebraic Curves Passing Through  $n$ -Independent Nodes. *New York J. Math.*, **22** (2016), 441–452.
- [6] Rafayelyan L. Poised Nodes Set Constructions on Algebraic Curves. *East J. Approx.*, **17**:3 (2011), 285–298.
- [7] Hakopian H., Kloyan H. On the dimension of spaces of algebraic curves passing through  $N$ -independent nodes. *Proceedings of YSU. Physical and Mathematical Sciences*, no. 2 (2019), 3–13.
- [8] Hakopian H., Toroyan S. On the Minimal Number of Nodes Determining Uniquely Algebraic Curves. *Proceedings of YSU. Physical and Mathematical Sciences*, no. 3 (2015), 17–22.
- [9] Carnicer J.M., Gasca M. On Chung and Yao’s Geometric Characterization for Bivariate Polynomial Interpolation. In: *Curve and Surface Design* (2002), 21–30.