

ON THE MULTIPLICITIES OF THE CHARACTER CODEGREES OF FINITE GROUPS

ZEINAB AKHLAGHI, MEHDI EBRAHIMI, AND MARYAM KHATAMI

ABSTRACT. Let G be a finite group and χ be an irreducible character of G , the number $\text{cod}(\chi) = |G : \ker(\chi)|/\chi(1)$ is called the codegree of χ . Also, $\text{cod}(G) = \{\text{cod}(\chi) \mid \chi \in \text{Irr}(G)\}$. For $d \in \text{cod}(G)$, the multiplicity of d in G , denoted by $m'_G(d)$, is the number of irreducible characters of G having codegree d . A finite group G is called a T'_k -group for some integer $k \geq 1$, if there exists $d_0 \in \text{cod}(G)$ such that $m'_G(d_0) = k$ and for every $d \in \text{cod}(G) - \{d_0\}$, we have $m'_G(d) = 1$. In this note we characterize finite T'_k -groups completely, where $k \geq 1$ is an integer.

1. INTRODUCTION

Let G be a finite group. Also let $\text{cd}(G)$ be the set of all irreducible character degrees of G , that is, $\text{cd}(G) = \{\chi(1) \mid \chi \in \text{Irr}(G)\}$, where $\text{Irr}(G)$ is the set of all complex irreducible characters of G . For a character χ of G , the codegree of χ is defined as $\text{cod}(\chi) := |G : \ker(\chi)|/\chi(1)$ (see [14]). Set $\text{cod}(G) := \{\text{cod}(\chi) \mid \chi \in \text{Irr}(G)\}$. It is well known that $\text{cod}(G)$ may be used to provide information on the structure of the group G (see for instance [7], [11] and [14]).

For $d \in \text{cd}(G)$, the multiplicity of degree d , denoted by $m_G(d)$, is the number of irreducible characters of G having the same degree d , i.e., $m_G(d) = |\{\chi \in \text{Irr}(G) \mid \chi(1) = d\}|$. Many people considered the relations between the multiplicities of irreducible character degrees of finite groups and the structure of the groups. Let $\text{Irr}_1(G)$ be the set of nonlinear irreducible characters in $\text{Irr}(G)$ and $\text{cd}_1(G)$ be the set of degrees of the characters in $\text{Irr}_1(G)$. If for a finite group G , there exists a nonnegative integer n such that $|\text{cd}_1(G)| = |\text{Irr}_1(G)| - n$, then G is called a D_n -group. In [2], Berkovich et al. gave the classification of D_0 -groups. Then, Berkovich and Kazarin in [1] and [4] classified all D_1 -groups. Also D_2 -groups have been classified by Liu and Lu (see [12] and [13]).

Inspired by the definition of D_n -groups, we define new families of finite groups called D'_n -groups, that are the groups G satisfying $|\text{cod}(G)| = |\text{Irr}(G)| - n$, for some nonnegative integer n . The second author classified D'_0 -groups, and showed that they are C_2 and S_3 (see [8]).

A finite group G is said to be a T_k -group for some integer $k \geq 1$, if there exists a nontrivial degree $d_0 \in \text{cd}(G)$ such that $m_G(d_0) = k$ and that for every $d \in \text{cd}(G) - \{1, d_0\}$, the multiplicity of d in G is trivial, that is, $m_G(d) = 1$. Tong-Viet in [16] showed that if G is a non-solvable T_k -group for some integer $k \geq 1$, then $k = 2$ and G is isomorphic to $\text{PSL}_2(5)$ or $\text{PSL}_2(7)$. Analogously, one can define a T'_k -group for some integer $k \geq 1$ based on the set $\text{cod}(G)$, that is a finite group G in which there exists $d_0 \in \text{cod}(G)$ such that $m'_G(d_0) = k$ and for every $d \in \text{cod}(G) - \{d_0\}$, we have $m'_G(d) = 1$. In this paper we wish to study T'_k -groups, for some integer $k \geq 1$. In fact, we characterize these groups completely and we prove the following theorem:

2000 *Mathematics Subject Classification.* 20C15.

Key words and phrases. finite group, codegree, character degree.

The first author is supported by a grant from IPM (No. 1400200028). The second author is supported by a grant from the School of Mathematics, Institute for Research in Fundamental Sciences (IPM)..

Main Theorem. *Let G be a finite T'_k -group, for some integer $k \geq 1$. Then one of the following occurs:*

- (a) G is non-solvable if and only if $k = 2$ and $G \cong \text{PSL}_2(5)$ or $G \cong \text{PSL}_2(7)$.
- (b) G is solvable if and only if one of the following holds:
 - (1) $G \cong C_4; S_4; \text{Suz}(2) \cong C_5 \rtimes C_4$ or $\text{Dic}_{12} \cong \text{SmallGroup}(12,1)$. Moreover $k = 2$.
 - (2) $G \cong C_2 \times S_3; D_{18}$ or Frobenius group $C_3^2 \rtimes Q_8$. Moreover $k = 3$.
 - (3) G is an elementary abelian p -group, for some prime p , or G is an extra-special 2-group. In particular $k = |G/G'| - 1$.
 - (4) G is isomorphic to the Frobenius group $C_2^\beta \rtimes C_{2^\beta - 1}$, for some integer β , where $p = 2^\beta - 1$ is prime. Moreover $k = p - 1$.
 - (5) G is isomorphic to the Frobenius group $C_p^\beta \rtimes C_2$, for some prime p and integer β . Moreover $k = (p^\beta - 1)/2$.

If $N \trianglelefteq G$ and $\theta \in \text{Irr}(N)$, then $I_G(\theta)$ denotes the inertia group of θ in G , and $\text{Irr}(G|\theta)$ denotes the set of all irreducible constituents of θ^G . Moreover, $\text{Irr}(G|N) = \text{Irr}(G) - \text{Irr}(G/N)$. By a rational group, we mean a group whose all irreducible characters are rational-valued. By n_p we mean, the p -part of n , where n is a natural number while, by $n_q(G)$ we mean the number of Sylow q -subgroups of G . For the rest of the notations, we follow [9].

2. PRELIMINARIES

In this section, we recall some known results of irreducible characters which will be needed throughout the proofs.

Remark 2.1. *Let K be a normal abelian subgroup of G . Then $|G : I_G(\lambda)| \leq |K| - 1$, for each $\lambda \in \text{Irr}(K)$.*

The above remark follows by the fact that there is a correspondence between G -orbits of $\text{Irr}(K)$ and G -conjugacy classes of K . We use this remark, several times through the proofs.

Lemma 2.2. *(see [15, Lemma 2.2]) Let G be a finite group acting on a module M over a finite field and $1 \in S \subseteq M$ be a union of some G -orbits of M . Also, assume q is a prime divisor of the order of G such that for every $v \in M \setminus S$, $C_G(v)$ contains a Sylow q -subgroup of G as a normal subgroup. Assume for each $1 \neq x \in S$, q divides $|\text{orb}_G(x)|$. Then, $|M| - |S| = n_q(G)(|C_M(Q)| - 1)$ for some $Q \in \text{Syl}_q(G)$.*

Lemma 2.3. *(see [16, Lemma 2.3]) Let S be a finite non-abelian simple group.*

(1) *If S is a sporadic simple group, the Tits group or an alternating group of degree at least 7, then S has two nontrivial irreducible characters, with distinct degrees and both extendible to $\text{Aut}(S)$.*

(2) *If S is a simple group of Lie type in characteristic p and $S \not\cong {}^2F_4(2)'$, then the Steinberg character of S , denoted by St_S , of degree $|S|_p$, is extendible to $\text{Aut}(S)$. Furthermore, if $S \not\cong \text{PSL}_2(3^f)$ for some integer $f \geq 2$, then S possesses an irreducible character θ such that $\theta(1) \neq |S|_p$ and θ also extends to $\text{Aut}(S)$.*

Lemma 2.4. *(see [5, Lemma 5]) Let N be a minimal normal subgroup of G such that $N = S_1 \times \cdots \times S_t$, where $S_i \cong S$ is a non-abelian simple group. If $\sigma \in \text{Irr}(S)$ extends to $\text{Aut}(S)$, then $\sigma \times \cdots \times \sigma \in \text{Irr}(N)$ extends to G .*

Lemma 2.5. *(see [3, Theorem E]) Let $N \trianglelefteq G$, where N is a p -group and G/N is solvable. Let $\theta \in \text{Irr}(N)$ be invariant in G , and assume that the members of $\text{Irr}(G|\theta)$ have distinct degrees. Then, G is a p -group and $\text{Irr}(G|\theta) = \{\chi\}$. In particular, $\chi(1) = \sqrt{|G|/|N|}$.*

Lemma 2.6. *Let G be a finite T'_k -group, where $k \geq 1$ is an integer. Then G is abelian if and only if G is an elementary abelian p -group, for some prime p ; or $G \cong C_4$.*

Proof. Since the 'if' part of the statement is clear, we only prove the 'only if' part. Note that for every $\lambda \in \text{Irr}(G)$, $\text{cod}(\lambda) = o(\lambda)$, where $o(\lambda)$ is the order of λ in $\text{Irr}(G) \cong G$. So, G does not have a homomorphic image isomorphic to $C_p \times C_q$, for distinct primes p and q . Thus, G is a p -group, for some prime p . First assume that $p \neq 2$. If G has a factor group isomorphic to C_{p^2} , then $m'_G(p)$ and $m'_G(p^2)$ are greater than one, a contradiction. So G is an elementary abelian p -group. Now assume that $p = 2$. If G contains an element of order greater than 2, then by easy calculation we can see that $G \cong C_4$. So in this case either G is an elementary abelian 2-group or $G \cong C_4$, and the result follows. \blacksquare

Lemma 2.7. *Let G be a finite group and K be a nontrivial normal subgroup of G . In addition, let $\chi \in \text{Irr}(G|K)$, $\delta \in \text{Irr}(K)$ and $\nu \in \text{Irr}(I_G(\delta)|\delta)$.*

- (a) *There exist $\lambda \in \text{Irr}(K)$, $\theta \in \text{Irr}(I_G(\lambda)|\lambda)$, such that $\text{cod}(\chi) = |I_G(\lambda)|/(|\ker(\chi)|\theta(1))$ and also $|I_G(\delta)|/(|\ker(\nu^G)|\nu(1)) \in \text{cod}(G)$.*
- (b) *If $\ker(\chi) \cap K = 1$, then $|K|$ divides $\text{cod}(\chi)$.*
- (c) *If χ is faithful, then $\text{cod}(\chi) \neq p$, where p is a prime divisor of $|G|/|K|$.*

Proof. (a) It is an immediate consequence of Clifford's theorem, [9, Theorem 6.11].

(b) First, note that we can consider χ as an irreducible character of $\overline{G} = G/\ker(\chi)$, and adopt the bar convention for this quotient group. Since $K \cap \ker(\chi) = 1$, we have $\overline{K} = K\ker(\chi)/\ker(\chi) \cong K$. Now, using part (a), we have

$$\text{cod}(\chi) = \frac{|I_{\overline{G}}(\lambda)|}{\theta(1)} = |\overline{K}| \frac{|I_{\overline{G}}(\lambda)|}{|\overline{K}|\theta(1)} = |K| \frac{|I_{\overline{G}}(\lambda)|}{|K|\theta(1)},$$

for some $\lambda \in \text{Irr}(\overline{K})$ and $\theta \in \text{Irr}(I_{\overline{G}}(\lambda)|\lambda)$. By, [9, Theorem 6.11], $\theta(1)$ is a divisor of $|I_{\overline{G}}(\lambda)|/|K|$, so $|K|$ divides $\text{cod}(\chi)$.

(c) Now, assume $\ker(\chi) = 1$, and $\text{cod}(\chi) = p$, for some prime p dividing $|G/K|$. Hence, using (a) and (b), we have $|K| = p$ and there exist $\lambda \in \text{Irr}(K)$, and $\theta \in \text{Irr}(I_G(\lambda)|\lambda)$, such that $|I_G(\lambda)/K| = \theta(1)$. It follows that $K \leq \mathbf{Z}(P)$, for every $P \in \text{Syl}_p(G)$. On the other hand, the argument in [9, page 84] implies that $|I_G(\lambda)/K| = \theta(1) = 1$. Then, $I_G(\lambda) = K$, which is not possible, as $P \leq I_G(\lambda)$. \blacksquare

Lemma 2.8. *Let G be a finite solvable T'_k -group and K be an abelian normal subgroup of G of order r^a , for some prime r and positive integers a and k . Assume, also $m'_G(n) > 1$, for some integer n .*

- (a) *Let $\lambda \in \text{Irr}(K)$. Assume $\text{cod}(\chi) \neq n$ and χ is faithful, for all $\chi \in \text{Irr}(G|\lambda)$. Then there exists integer b , such that $I_G(\lambda)/K$ is a group of order r^{2b} .*
- (b) *Assume $\text{cod}(\chi) \neq n$ and χ is faithful for all $\chi \in \text{Irr}(G|K)$. Then, Sylow r -subgroups of G/K have order r^{2c} for some integer c , and Hall r' -subgroups of G/K act on K , Frobeniusly.*

Proof. (a) As G is a T'_k -group, and $\text{cod}(\chi) \neq n$, for all $\chi \in \text{Irr}(G|\lambda)$, using Lemma 2.7(a), it follows that the members of $\text{Irr}(I_G(\lambda)|\lambda)$ must have distinct degrees. Now using Lemma 2.5, we have $|I_G(\lambda)|/|K| = r^{2b}$, for some integer b .

(b) As K is a normal subgroup of G , then for each $R \in \text{Syl}_r(G)$, we have $K \cap \mathbf{Z}(R) \neq 1$. Therefore, as $K \cong \text{Irr}(K)$, there exists $1 \neq \lambda \in \text{Irr}(K)$ such that $I_G(\lambda)/K$, contains R/K . Now, using part (a), $|R/K|$ must be an even power of r . Also $I_G(\lambda)/K$ does not contain any r' -element for each nontrivial $\lambda \in \text{Irr}(K)$, so it follows that Hall r' -subgroups of G/K act on K , Frobeniusly. \blacksquare

3. NON-SOLVABLE T'_k -GROUPS

In this section, we aim to characterize non-solvable T'_k -groups, where $k \geq 1$ is an integer, via the following lemmas.

Lemma 3.1. *Let S be a finite non-abelian simple T'_k -group, for some integer $k \geq 1$. Then $S \cong \text{PSL}_2(5)$ or $\text{PSL}_2(7)$.*

Proof. Since S is simple, for every nontrivial $\chi \in \text{Irr}(S)$, $\ker(\chi) = 1$ and so $\text{cod}(\chi) = |G|/\chi(1)$. So S is a T_k -group and using [16, Theorem 3.2] the result follows. ■

Lemma 3.2. *Let G be a finite group and M be a non-abelian minimal normal subgroup of G such that $G/M \cong \text{PSL}_2(5)$ or $\text{PSL}_2(7)$. Then G is not a T'_k -group, for some integer $k \geq 1$.*

Proof. We argue by the contradiction and suppose that G is a T'_k -group, for some integer $k \geq 1$. By Lemmas 2.3 and 2.4 we get that, M contains an irreducible character λ which is extendible to G . Since G/M contains two distinct irreducible characters whose degrees are 3, then according to Gallagher's theorem ([9, Corollary 6.17]), there are $\chi_1, \chi_2 \in \text{Irr}(G|\lambda)$ whose degrees are $3\lambda(1)$. We show that $\ker(\chi_i) = 1$, for each $i \in \{1, 2\}$. If $\ker(\chi_i) \neq 1$ for some $i \in \{1, 2\}$, then $G = \ker(\chi_i) \times M$, implying that M is a non-abelian simple group. Since $G/\ker(\chi_i)$ is a T'_l -group, for some integer l , then by Lemma 3.1, it should be isomorphic to $\text{PSL}_2(5)$ or $\text{PSL}_2(7)$. Hence $G \cong \text{PSL}_2(q)^2$ for some $q \in \{5, 7\}$ or $G \cong \text{PSL}_2(7) \times \text{PSL}_2(5)$. In all cases G has 4 distinct faithful characters of degree 9, and 2 distinct irreducible characters with degree 3 and kernel M , implying that $m'_G(|G|/9) = 4$ and $m'_G(|G|/(3|M|)) = 2$, which is a contradiction. So $\ker(\chi_1) = \ker(\chi_2) = 1$ and $\text{cod}(\chi_1) = \text{cod}(\chi_2) = |G|/(3\lambda(1))$. On the other hand, G/M has two distinct irreducible characters with the same codegree $|G|/(3|M|)$ which is different from $|G|/(3\lambda(1))$, causing a contradiction. ■

Lemma 3.3. *Let G be a finite group and M be a minimal normal subgroup of G such that $G/M \cong \text{PSL}_2(5)$ or $\text{PSL}_2(7)$. Then G is not a T'_k -group, for some integer $k \geq 1$.*

Proof. Note that by Lemma 3.2, M is abelian. Let G be a counterexample. First we prove that G is perfect. If $G' \cap M = 1$, then $G = G' \times M$. Let $\lambda \in \text{Irr}(M)$ be a nontrivial irreducible character and $\chi_i \in \text{Irr}(G')$, for $i \in \{1, 2\}$, be characters with degree 3. Then, $\ker(\chi_i \times \lambda) = 1$, for $i = 1, 2$, and so we get a contradiction by our assumption as $m'_G(|G|/(3|M|)) > 1$ and $m'_G(|G|/3) > 1$. Therefore, $M \leq G'$ and G is perfect. Let $1_M \neq \lambda \in \text{Irr}(M)$ and $\chi \in \text{Irr}(G|\lambda)$. If $\ker(\chi) \neq 1$, then $G = \ker(\chi) \times M$, which is a contradiction, as G is perfect. Hence all characters in $\text{Irr}(G|M)$ are faithful. Then, by Lemma 2.7(b), $|M|$ is a divisor of $\text{cod}(\chi)$.

We remark that $C_G(M) = M$, since otherwise $M \leq \mathbb{M}(G/M)$, where $\mathbb{M}(G/M)$ is the Schur multiplier of G/M , and $G \cong \text{SL}_2(q)$, for $q \in \{5, 7\}$. It is a contradiction, as G is a T'_k -group.

Note that $|G|/(3|M|) \in \text{cod}(G/M)$ has nontrivial multiplicity. We claim that $\text{cod}(\chi) \neq |G|/(3|M|)$, for every $\chi \in \text{Irr}(G|\lambda)$. Suppose $\text{cod}(\chi) = |G|/(3|M|)$, for some $\chi \in \text{Irr}(G|\lambda)$ and $\lambda \in \text{Irr}(M)$. Since M is an elementary abelian group, it follows that $|M| = 2, 4, 5, 7$ or 8 . As $C_G(M) = M$, by the normalizer-centralizer theorem, the only possibility is $G/M \cong \text{PSL}_2(7)$ and $|M| = 8$. Therefore, by Lemma 2.7(a), $\text{cod}(\chi) = |G|/(3|M|) = 8 \times 7 = |M| \times |I_G(\lambda)|/(|M|\theta(1))$, for some $\theta \in \text{Irr}(I_G(\lambda)|\lambda)$. Note that, $A = I_G(\lambda)/M$ is a subgroup of $\text{PSL}_2(7)$. Hence, looking at the subgroups of $\text{PSL}_2(7)$ either $|A| = 7$ or $|A| = 21$, implying that $8 = |M| \leq |G : I_G(\lambda)| \in \{8, 8 \times 3\}$, which is impossible, by Remark 2.1. Therefore, all characters in $\text{Irr}(G|M)$ have distinct codegrees and so different degrees. As, $C_G(M) = M$, we deduce that $I_G(\lambda) \neq G$, for every $1 \neq \lambda \in \text{Irr}(M)$. So $I_G(\lambda)/M$ is a proper subgroup of $\text{PSL}_2(5)$

or $\mathrm{PSL}_2(7)$ and so it is solvable, for every $1 \neq \lambda \in \mathrm{Irr}(M)$. Now using Lemma 2.8(a), we get that $I_G(\lambda)/M = 1$ or C_2^2 , for all $1 \neq \lambda \in \mathrm{Irr}(M)$. Therefore, applying Lemma 2.5, $\mathrm{cd}(G|M) \subseteq \{30, 60\}$ when $G/M \cong \mathrm{PSL}_2(5)$, and $\mathrm{cd}(G|M) \subseteq \{84, 168\}$ when $G/M \cong \mathrm{PSL}_2(7)$. Hence $\mathrm{cd}(G/M) \cap \mathrm{cd}(G|M) = \emptyset$. Thus, G is a T_2 -group, and using [16] we get a contradiction. ■

Proof of the Main Theorem case (a): Let $G \cong \mathrm{PSL}_2(5)$ or $\mathrm{PSL}_2(7)$. Then for every nontrivial $\chi \in \mathrm{Irr}(G)$, $\ker(\chi) = 1$ and so $\mathrm{cod}(\chi) = |G|/\chi(1)$. By [16, Theorem 3.2], G is a T_2 -group and so G is a T'_2 -group. Now, we concentrate on the 'only if' part. Let G be a counterexample with minimal order. Let M be the last term of the derived series of G , and N be a normal subgroup of G contained in M such that M/N is a chief factor of G .

First, we claim that G is perfect. Suppose false. Hence M is a proper subgroup of G . Note that, G/N is a finite non-solvable T'_l -group, for some integer $l \geq 1$. If $N \neq 1$, then by the minimality of $|G|$, $G/N \cong \mathrm{PSL}_2(5)$ or $\mathrm{PSL}_2(7)$, and consequently $G = M$, a contradiction. Hence, $N = 1$. Let $C = C_G(M)$. If $C \neq 1$, then by the minimality of $|G|$ we deduce that $M \cong G/C \cong \mathrm{PSL}_2(5)$ or $\mathrm{PSL}_2(7)$ and so $G \cong M \times C$. Since C is solvable, it has a nontrivial linear character, say ζ . Let also χ_1 and χ_2 be two distinct irreducible characters of M of degree 3. Then $\chi_1 \times \zeta$ and $\chi_2 \times \zeta$ are two irreducible characters with the same degree 3 whose kernels are isomorphic to $\ker(\zeta)$. Therefore, they have the same codegree $|G|/(3|\ker(\zeta)|)$ and it is different from $|G|/(3|C|) = \mathrm{cod}(\chi_i \times 1)$ for $i \in \{1, 2\}$, a contradiction. Hence $C = 1$ and G is a subgroup of $\mathrm{Aut}(M)$.

Using Lemmas 2.3 and 2.4, there exists $\lambda \in \mathrm{Irr}(M)$ which is extendible to G . We claim that G/M is a D_0 -group or an abelian group. Otherwise, G/M possesses at least two distinct nonlinear irreducible characters with the same degree $d > 1$, and also since G/M is solvable, it has at least two distinct linear characters. So by Gallagher's theorem, there exists four distinct characters $\lambda_1, \lambda_2, \psi_1$ and ψ_2 in $\mathrm{Irr}(G|\lambda)$, such that $\lambda_1(1) = \lambda_2(1) = \lambda(1)$ and $\psi_1(1) = \psi_2(1) = \lambda(1)d$. Note that, as M is the only minimal normal subgroup of G , $\ker(\chi) = 1$, for all $\chi \in \{\lambda_1, \lambda_2, \psi_1, \psi_2\}$. Hence, $m'_G(\mathrm{cod}(\lambda_1)) > 1$ and $m'_G(\mathrm{cod}(\psi_1)) > 1$, implying that G is not a T'_k -group, which is a contradiction. Therefore, G/M is a D_0 -group or an abelian group, as desired. Now we claim that $|G/G'| = 2$. Otherwise, there exist at least two distinct nontrivial irreducible characters in $\mathrm{Irr}(G/G')$ with the same codegree. On the other hand, by Gallagher's theorem, there exist at least two distinct extensions of λ to G with the same codegree, which causes a contradiction. So G/G' is a cyclic group of order 2. Therefore, looking at the classification of the D_0 -groups (see [2]), we get that the only possibilities for G/M is being isomorphic to either S_3 or C_2 . Recall that, by Gallagher's theorem, the multiplicity of $\mathrm{cod}(\chi)$, for some $\chi \in \mathrm{Irr}(G|\lambda)$ is greater than one, where χ is an extension of λ . Noting that $\ker(\psi) = 1$, for all $\psi \in \mathrm{Irr}(G|M)$, we deduce the multiplicity of every character degree in $\mathrm{cd}(G|M) \setminus \{\lambda(1)\}$ must be trivial, as G is a T'_k -group, for some integer $k \geq 1$. As $\mathrm{cd}(G/M) \subseteq \{2\}$, we deduce that $\mathrm{cd}(G|M) \cap \mathrm{cd}(G/M) = \emptyset$. Then, we conclude that G should be a T_k -group, for some integer $k \geq 1$. Now, looking at the structure of non-solvable T_k -groups (see [16]), we get a contradiction. □

Hence $G = M$ is perfect. Let K be a maximal normal subgroup of G . Then G/K is a simple T'_r -group, for some integer $r \geq 1$. Lemma 3.1 implies that $G/K \in \{\mathrm{PSL}_2(5), \mathrm{PSL}_2(7)\}$. If $K = 1$, then we are done, so we may assume K is not trivial. Let L be a normal subgroup of G contained in K such that K/L is a chief factor of G . Then we work on $\overline{G} = G/L$ and we use the bar convention for this quotient group. As, $\overline{G}/\overline{K} \in \{\mathrm{PSL}_2(5), \mathrm{PSL}_2(7)\}$, by Lemma 3.3, we get a contradiction.

4. SOLVABEL T'_k -GROUP

In this section we aim to study solvable T'_k -groups for some integer $k \geq 1$, and prove Case (b) of the main theorem, through Lemmas 4.1-4.9.

Lemma 4.1. *Let G be a finite non-abelian p -group, for some prime p . Then G is a T'_k -group for some integer $k \geq 1$, if and only if $p = 2$ and G is an extra-special group.*

Proof. First, we prove that G is a rational group and $p = 2$. Let $\chi \in \text{Irr}(G|G')$ and ϵ be a $|G|$ th root of unity and $\delta \in \text{Gal}(\mathbb{Q}(\epsilon)/\mathbb{Q})$. Then $\chi^\delta \in \text{Irr}(G)$ and $\chi^\delta(1) = \chi(1)$. As $|\ker(\chi^\delta)| = |\ker(\chi)|$, we have $\text{cod}(\chi^\delta) = \text{cod}(\chi)$. We show that $|G|/|\ker(\chi)G'|$ is not trivial. Suppose false, and take K to be a normal subgroup of G , contained in G' , such that G'/K is a chief factor of G and $\ker(\chi) \cap G' \leq K$. Hence,

$$\frac{G}{K} \cong \frac{G'}{K} \times \frac{\ker(\chi)K}{K}$$

is an abelian group, implying that $G' \leq K$, a contradiction. Therefore by Lemma 2.7(c), $\text{cod}(\chi) \neq p$. So, as G is a T'_k -group, for some integer k , and $m'_G(p) > 1$, we get that $\chi^\delta = \chi$. Hence $\chi(x)$ is rational for all $x \in G$. So, all nonlinear irreducible characters of G are rational-valued, implying that $|G|$ is even by [9, (3.16), p. 46], and it follows that $p = 2$. Note that by Lemma 2.6, either $G/G' \cong C_4$ or G/G' is an elementary abelian 2-group. Since G is a 2-group, the first case is not possible and so linear characters of G are also rational-valued. Hence G is a rational group, as wanted.

Assume K is a normal subgroup of G , contained in G' , such that G'/K is a chief factor of G . Write $\overline{G} = G/K$ and adopt the bar convention for this quotient group. Remark that \overline{G}' is a central subgroup of \overline{G} of order 2. Then, $\mathbf{Z}(\overline{G}) \cong \overline{N} \times \overline{G}'$, where \overline{N} is an elementary abelian group, by [6, Theorem 3.6].

We prove that $\mathbf{Z}(\overline{G}/\overline{N}) = \overline{G}'\overline{N}/\overline{N}$. Suppose false, then there exists an element x out of $\overline{G}'\overline{N}$, such that for all elements $y \in \overline{G}$, $[x, y] \in \overline{N}$. On the other hand $[x, y] \in \overline{G}'$, implying that $[x, y] = 1$ and so $x \in \mathbf{Z}(\overline{G}) = \overline{G}'\overline{N}$, a contradiction, as wanted. Therefore, $\overline{G}/\overline{N}$ is an extra-special group. Now, we aim to prove that $\overline{N} = 1$. Suppose false. Let $\overline{N}_1, \overline{N}_2$ be two different subgroups of $\mathbf{Z}(\overline{G})$ of order $|\overline{N}|$, not containing \overline{G}' . Then, replacing \overline{N} with each one of \overline{N}_i 's, for $i = 1, 2$, similar to the above discussion, we have $\overline{G}/\overline{N}_i$, for $i = 1, 2$, are extra-special groups with the same order. Therefore, \overline{G} has two distinct nonlinear irreducible characters with the same codegree different from 2, a contradiction. Thus, $\overline{N} = 1$. Hence \overline{G} is an extra-special group. If K is trivial, we are done. Hence, we may assume K is a minimal normal subgroup of G . Then $K \leq \mathbf{Z}(G)$.

Note that, $K = \mathbf{Z}(G)$, as otherwise $\mathbf{Z}(G) = K \times T$, where $TK/K = \mathbf{Z}(\overline{G})$ (as $\mathbf{Z}(G)/K \leq \mathbf{Z}(G/K)$ and $|\mathbf{Z}(G/K)| = |G'/K| = 2$), hence replacing K by T in the above discussion, we have G/T and G/K are isomorphic and so G has two distinct nonlinear irreducible characters with the same codegree, a contradiction. Then, $\mathbf{Z}(G) = K$ and χ is faithful for all $\chi \in \text{Irr}(G|K)$, yielding that $|G/K|$ is a square, by Lemmas 2.7(c) and 2.8(b), a contradiction. So $K = 1$, as wanted. ■

Lemma 4.2. *Let G be a finite solvable T'_k -group, for some integer $k \geq 1$, and $G/G' \cong C_p^\alpha$, for some prime p and integer α . Assume also G' is a minimal normal subgroup of G whose order is co-prime to $|G/G'|$. Then one of the following occurs:*

- (1) $G \cong C_2 \times S_3$;
- (2) $G \cong D_{2q}$ for some prime $q \geq 3$; or
- (3) G is a Frobenius group $C_2^\beta \rtimes C_{2^\beta-1}$, where $p = 2^\beta - 1$. In particular β is odd.

Proof. Let $|G'| = q^\beta$, for some prime q and integer β and $C_G(G') = G' \times C$, where C is an elementary abelian p -group. Then $C \leq \mathbf{Z}(G)$ and $G = C \times T$, where T is a subgroup of G containing G' . Remark that T is not isomorphic to G' . Note that, χ is faithful, for all $\chi \in \text{Irr}(T|T')$, as $\ker(\chi) \leq C_T(T') = T' = G'$ and $\ker(\chi) \cap T' = 1$.

Firstly, assume that $|T|_p > 2$, then $m'_T(p) > 1$. Hence, by Lemma 2.7(c), we get that all characters in $\text{Irr}(T|T')$ have distinct degrees and so T is a D_0 -group. Looking at the structure of D_0 -groups we

get that T is a Frobenius group, with cyclic Frobenius complement of order p and $p = q^\beta - 1$. This is not possible, unless $q = 2$ and β is odd. Hence $p \neq 2$. Now we show that in this case C is trivial. Assume, $C = C_p^{\alpha_1}$ and $\alpha_1 \geq 1$. Since $p \neq 2$, we deduce that $|\text{Irr}(C) \setminus \{1_C\}| > 1$. Let $\lambda_1, \lambda_2 \in \text{Irr}(C)$ be two distinct nontrivial characters and $\chi \in \text{Irr}(T)$ be the only nonlinear irreducible character of T . Then $|\ker(\lambda_1 \times \chi)| = |\ker(\lambda_2 \times \chi)| = |C|/p$ and $\lambda_1 \times \chi(1) = \lambda_2 \times \chi(1) = \chi(1)$, so $\text{cod}(\lambda_1 \times \chi) = \text{cod}(\lambda_2 \times \chi)$. Note that $\text{cod}(\lambda_1 \times \chi)$ is not equal to p , by Lemma 2.7(c), a contradiction, as both multiplicities of $\text{cod}(\lambda_1 \times \chi)$ and p are not trivial. Hence $C = 1$. So G is the described group in case (3) of the Lemma.

Now we assume $|T|_p = p = 2$. Recall that, $C_T(T') = T'$ and T' is a minimal normal subgroup of $T \cong G/C$, where T is a Frobenius group whose Frobenius complements are cyclic groups of order 2. Therefore, $T \cong G/C \cong D_{2q}$. First assume $q > 3$. We prove that C is trivial. Suppose false. In this case $\text{Irr}(G/C)$ contains two distinct nonlinear irreducible characters χ_1 and χ_2 . Let $\lambda \in \text{Irr}(C)$ be a nontrivial irreducible character. Then, $|\ker(\chi_1 \times \lambda)| = |\ker(\chi_2 \times \lambda)| = |\ker(\lambda)| = |C|/2$ and so $\text{cod}(\chi_1 \times \lambda) = \text{cod}(\chi_2 \times \lambda) = 2\text{cod}(\chi_1)$, hence $m'_G(2\text{cod}(\chi_1)) > 1$. Noting that $m'_T(\text{cod}(\chi_1)) = (q-1)/2 > 1$, we get a contradiction. Therefore $C = 1$. So $G \cong D_{2q}$, a described group in case (2) of the lemma.

The last possibility is that $q = 3$ and $G \cong C_2^{\alpha-1} \times D_6$. Hence, in this case $m'_G(6) = 2^{\alpha-1} - 1$ and $m'_G(2) = 2^\alpha - 1$. Therefore, either $\alpha = 1$ or $\alpha = 2$, which means that either $G \cong C_2 \times S_3$ or $G \cong D_6$, which are the groups in case (1) and (2) of the lemma. ■

Lemma 4.3. *Let G be a finite solvable T'_k -group for some integer $k \geq 1$ and $K < G' < G$ be a normal series of G . Assume also G/K is isomorphic to the Frobenius group $C_{2^\beta} \rtimes C_p$, for some integer β and a prime number $p = 2^\beta - 1$. Then, $K = 1$.*

Proof. Let G be a counterexample with minimal order. Therefore, we may assume K is a minimal normal subgroup of G . First of all, we show that $\ker(\chi) = 1$, for all $\chi \in \text{Irr}(G|K)$. Suppose false. Then, there exists $\chi \in \text{Irr}(G|K)$ such that $\ker(\chi)K/K$ is a nontrivial normal subgroup of G/K , implying that $G' \cong K \times T$, where $T \cong \ker(\chi)$ (Note that $\ker(\chi) \not\cong G/K$, as $|G/G'|$ is a prime). Now, we work on group $\overline{G} = G/T$, and we adopt bar convention for this quotient group. Note that $\overline{K} = \overline{G}'$. As $|G/G'|$ is prime, we deduce that $(|\overline{G}'|, |\overline{G}/\overline{G}'|) = 1$. So the hypothesis of the Lemma 4.2, is provided for \overline{G} . Applying Lemma 4.2, $|K| = |\overline{G}'| = 2^\beta$. Hence both groups G/K and G/T have faithful irreducible characters with the same codegree 2^β and so $m'_G(2^\beta) > 1$, which is a contradiction, as $m'_G(p) > 1$. So $T = 1$, as wanted.

Hence, by Lemmas 2.7(c) and 2.8(b), we get that G/K acts on K Frobeniusly, which is a contradiction, as G/K is a Frobenius group itself and it is well-known that a complement of a Frobenius group is not Frobenius. ■

Lemma 4.4. *Let G be a finite solvable T'_k -group for some integer $k \geq 1$, and $K < G' < G$ be a normal series of G , such that $G/K \cong C_2 \times S_3$. Then $K = 1$.*

Proof. Let G be a counterexample with minimal order. Then, we may assume K is a minimal normal subgroup of G of order p^α , for some prime p and integer α . Let $T = \ker(\chi)$ for some $\chi \in \text{Irr}(G|K)$. Then, T is isomorphic to one of the groups in $\{S_3, C_6, C_2, C_3, 1\}$.

Let $p \neq 2$. Assume $T \in \{S_3, C_6, C_3\}$. Then $G' = T_0 \times K$, where $T_0 = T \cap G' \cong C_3$. Then, G/T_0 satisfies the hypothesis of Lemma 4.2. Therefore, $G/T_0 \cong C_2 \times S_3$ and $K \cong C_3$. Note that, $m'_G(2) > 1$ and both G/T_0 and G/K contain faithful irreducible characters whose codegrees are 3. So $m'_G(3) > 1$, a contradiction. Therefore, $T \cong C_2$ or 1. Set $S = \prod_{\chi_0 \in \text{Irr}(G|K)} \ker(\chi_0)$. Now, we consider the group $\overline{G} = G/S$ and we adopt the bar convention for this quotient group. Then $\ker(\chi_1) = 1$, for all $\chi_1 \in \text{Irr}(\overline{G}|\overline{K})$. Note that $\overline{G}/\overline{K} \cong S_3$ or $C_2 \times S_3$. By Lemmas 2.7(c) and 2.8(b), $\overline{G}/\overline{K}$ acts on \overline{K} , Frobeniusly, a contradiction, as $\overline{G}/\overline{K} \cong S_3$ or $C_2 \times S_3$.

Finally, let $p = 2$. If $C = C_G(K)$ contains a Sylow 3-subgroup of G , then $K \leq \mathbf{Z}(G)$ and so $|K| = 2$. Hence, G has a normal abelian subgroup L of order 12. Looking at the classification of groups of order 24, we see $G \cong D_{24}$ or $G \cong \text{SmallGroup}(24, 8)$. In the first case there exists 2 distinct faithful characters with the same degree 2, a contradiction, as $m'_G(2) > 1$. In the second case $S = \text{Socle}(G) = C_3 \times C_2$, so every $\chi \in \text{Irr}(G|\lambda)$ is faithful, and $I_G(\lambda)/S$ is a group of order 2, for all $\lambda \in \text{Irr}(S)$ of order 6. Then by Gallagher's theorem, there is two distinct faithful characters of degree 2 above each irreducible character $\lambda \in \text{Irr}(S)$ of order 6, a contradiction.

So C does not contain a Sylow 3-subgroup of G and hence $T \cong C_2$ or 1. Let $S = \prod_{\chi \in \text{Irr}(G|K)} \ker(\chi)$. Note that $S \cong C_2$ or 1. Setting $\overline{G} = G/S$, we see that $\ker(\chi) = 1$, for all $\chi \in \text{Irr}(\overline{G}|\overline{K})$. Then by Lemmas 2.7(c) and 2.8(a), $I_{\overline{G}}(\lambda)/\overline{K} \cong C_2^2$ or 1, for all $\lambda \in \text{Irr}(\overline{K})$. Applying Lemma 2.5, we have $\text{cd}(\overline{G}|\overline{K}) \subseteq \{6, 12\}$. Since $\text{cd}(\overline{G}|\overline{K}) \cap \text{cd}(\overline{G}/\overline{K}) = \emptyset$, then \overline{G} must be a T_1 -group (D_0 -group). Looking at the classification of the D_0 -groups (see [2]) we get a contradiction, as $\overline{G} \cong S_3$ or $C_2 \times S_3$. ■

Lemma 4.5. *Let G be a solvable T'_k -group for some $k \geq 1$, and $K < G' < G$ be a normal series of G such that $G/K \cong S_3$ and K is a minimal normal subgroup of G of order r^α , for some prime $r \neq 3$. Then $G \cong S_4$.*

Proof. We show that $C_G(K) = K$. Suppose false, then $G' = T \times K$, where $T \cong C_3$. As $|G/G'| = 2$, we deduce that $r \neq 2$. Therefore, G/T satisfies Lemma 4.2, and so $|K| = r$ is an odd prime other than 3. By Lemma 4.3 we get a contradiction. Hence $C_G(K) = K$. Then χ is faithful for all $\chi \in \text{Irr}(G|K)$. Remark that, $I_G(\lambda)/K$ does not contain the Sylow 3-subgroup of G/K , for any $\lambda \in \text{Irr}(K)$, as otherwise $\mathbf{Z}(G') \cap K \neq 1$, contradicting the fact that $C_G(K) = K$. Hence, $I_G(\lambda)/K \cong 1$ or C_2 , for every $\lambda \in \text{Irr}(K)$, implying that G -orbit sizes of nontrivial elements of K are either 3 or 6. As Sylow subgroups of $I_G(\lambda)$ are cyclic, λ extends to $I_G(\lambda)$, for all $\lambda \in \text{Irr}(K)$. If $I_G(\lambda)/K \cong C_2$, then $m'_G(|I_G(\lambda)|) > 1$ by Lemma 2.7(a). Also, if $I_G(\lambda_1) = I_G(\lambda_2) = K$, for two non-conjugate nontrivial irreducible characters of K , then $m'_G(|K|) > 1$.

Let n_i be the number of G -orbits of size i of elements of $K \cong \text{Irr}(K)$. As G is a T'_k -group, by the above discussion, we have the following possibilities for the number of G -orbit sizes of nontrivial elements of $\text{Irr}(K)$: $(n_3, n_6) \in \{(k, 0), (k, 1), (0, k)\}$, for some integer k . The last case yields that G/K acts on K Frobeniusly, which is a contradiction, as G/K is a Frobenius group. If the second case occurs, then applying Lemma 2.2, we get that $r^\alpha - 7 = n_2(r^\beta - 1) = 3(r^\beta - 1)$, for some integer β . Whence, $r^\beta(r^{\alpha-\beta} - 3) = 4$, implying that $|K| = r^\alpha = 16$. On the other hand, by Lemma 2.8(b) the Sylow 3-subgroup of G/K acts on K , Frobeniusly, yielding that G has a normal Frobenius subgroup of order 48, say L . Then L has exactly 5 distinct irreducible characters of degree 3. Since $|G/L| = 2$, these five characters belong to at least 3 distinct G -orbits and the size of one of these orbits must be 1. Hence one of the described characters extends to G and we have two distinct faithful irreducible characters with the same codegree $|G|/3$. If there exist two G -orbits of size 2 containing one of the described characters, then we have two distinct faithful irreducible characters of codegree $|G|/6$, which is a contradiction. If there exists only one G -orbit with size 2, then it follows that we have one faithful irreducible character of degree 6 and 8 distinct faithful irreducible characters of degree 3. On the other hand $|G| = 96 < 6^2 + 8 \cdot 3^2$, a contradiction. So all of these five irreducible characters of L belong to G -orbits of G of size 1 and so all of the faithful irreducible characters of G have the same degree 3. But we have $n_6 = 1$, which means that there is a character $\lambda \in \text{Irr}(K)$, such that $\lambda^G \in \text{Irr}(G)$, a contradiction as $\lambda^G(1) = 6$.

Then the first case occurs, and applying Lemma 2.2, we have $|K| = 4$ and $G \cong S_4$, as desired. ■

Lemma 4.6. *Let G be a finite solvable T'_k -group, where $k \geq 1$, and $M < G' < G$ be a normal series of G such that $G/M \cong S_4$. Then $M = 1$.*

Proof. Let G be a counterexample with minimal order. Then we may assume M is a minimal normal subgroup of G of order $|M| = r^\alpha$, for some prime r and integer α . We show that M is the only minimal normal subgroup of G . Suppose false, then $\text{Socle}(G) \neq M$, so $\text{Socle}(G) = PM \cong M \times P$, where PM/M is the Sylow 2-subgroup of G'/M . First, let $r \neq 3$. Looking at G/P and applying Lemma 4.5, we get that $M \cong C_2^2$. If we take $\lambda \in \text{Irr}(M)$ and $\lambda' \in \text{Irr}(P)$ to be two nontrivial characters, then all elements in $\text{Irr}(G|\lambda \times \lambda')$ are faithful. Note that $m'_G(8) > 1$. Now, using Lemma 2.7(b), the codegrees of all characters in $\text{Irr}(G|\lambda \times \lambda')$ are divided by $|PM| = 16$. Hence, by Lemma 2.8(a), we get that $I_G(\lambda \times \lambda') = PM$. Note that all such characters must be conjugate, as otherwise we have at least two distinct faithful irreducible characters with the same degree 6, and so $m'_G(|G|/6) > 1$, by Lemma 2.7(a), a contradiction. So, $9 = (|P| - 1)(|M| - 1) = |G : I_G(\lambda \times \lambda')| = 6$, a contradiction. Now, assume $r = 3$. Then by [8], G/P is not a T'_1 -group and so, $m'_G(d) > 1$ for some integer $d \neq 8$ (d must divide $|G/P|$). Since in $G/K \cong S_4$, multiplicity of 8 is not trivial we get a contradiction. Therefore, our claim is proved and χ is faithful for all $\chi \in \text{Irr}(G|M)$.

Now, we show that $\text{cod}(\chi) \neq 8$, for all $\chi \in \text{Irr}(G|M)$. Assume $\text{cod}(\chi) = 8$, for some $\chi \in \text{Irr}(G|M)$. Then, $8 = |M| \times |I_G(\lambda)|/(|M|\theta(1))$, where $1_M \neq \lambda \in \text{Irr}(M)$, $\theta \in \text{Irr}(I_G(\lambda)|\lambda)$ and $\chi \in \text{Irr}(G|\lambda)$, by Lemma 2.7(a). So, M is a 2-group and $P/M \leq I_G(\lambda)/M$, where P/M is the Sylow 2-subgroup of G'/M .

Then, $(|M|, |I_G(\lambda)|/|M|, \theta(1)) \in \{(8, 1, 1), (4, 2, 1), (4, 4, 2), (2, 8, 4)\}$. The first two cases are not possible as $P/M \leq I_G(\lambda)/M$. The last case is not possible, as by [9, page 84], $16 = \theta(1)^2 \leq |I_G(\lambda)|/|M| = 8$. So, the only possibility in this case is $|M| = |I_G(\lambda)/M| = 4$ and $\theta(1) = 2$. In this case $4 = |M| < |G : I_G(\lambda)| = 6$, contradicting Remark 2.1. So our claim is proved and G satisfies hypothesis of Lemma 2.8(b). Therefore $G/M \cong S_4$ acts on M Frobeniusly, which is a contradiction. \blacksquare

Lemma 4.7. *Let G be a finite solvable T'_k -group, for some integer $k \geq 1$ and $|G/G'| = 2$. Then either $G \cong S_4$; $G \cong D_{18}$; or G is a Frobenius group isomorphic to $C_p^\beta \rtimes C_2$, for some odd prime p and integer β .*

Proof. We aim to prove that either $G' \cong A_4$; $G' \cong C_9$ or it is an elementary abelian p -group, for some odd prime p . First, we assume that G' is a p -group, for some odd prime p . We work on $\overline{G} = G/G''$ and we use the bar convention for this quotient group. As $|\overline{G}/\overline{G}'| = 2$, by Fitting Lemma (see [10, Theorem 4.34]) we have $\overline{G}/\overline{G}'$ acts on \overline{G}' Frobeniusly.

We prove that \overline{G}' is an elementary abelian group or a cyclic group of order 9. Let $\overline{G}' \cong \overline{T} \times \overline{L}$, where \overline{T} is a cyclic group of order p^m , for some $m > 1$. Note that as $\overline{G}/\overline{G}'$ acts on \overline{G}' by inverting every element, then every subgroup of \overline{G}' is a normal subgroup of \overline{G} . Now, we consider group $\overline{G}/\overline{L} \cong D_{2n}$, where $n = p^m$. Since $\overline{G}/\overline{L}$ is a T'_l -group for some integer l , it follows that $n = 9$ (notice that there are $\phi(d)/2$ irreducible characters in $\text{Irr}(D_{2n})$ with the same codegree d , for each divisor d of n , where ϕ is the Euler's totient function). If $\overline{L} \neq 1$, then \overline{G} has a homeomorphic image isomorphic to the Frobenius group $(C_3 \times C_9) \rtimes C_2$, which is not possible, as in this case $m'_G(9)$ and $m'_G(3)$ are not trivial. Therefore $\overline{L} = 1$ and $\overline{G} \cong D_{18}$, as desired. So either $p = 3$ and $\overline{G}' \cong C_9$ or it is an elementary abelian p -group.

Now, we show that $G'' = 1$. Let $G'/G'' \cong C_9$ and $G'' \neq 1$, then we may assume G'' is a minimal normal subgroup of G and so $G'' \leq \mathbf{Z}(G')$, a contradiction. Hence in this case $G'' = 1$. So, we assume G'/G'' is an elementary abelian group. Contrary to what we want to prove, assume $G'' \neq 1$ and take L to be a normal subgroup of G , such that G''/L is a chief factor of G . Write $\tilde{G} = G/L$ and adopt the tilde convention for \tilde{G} . Note that $\tilde{G}'' \leq \mathbf{Z}(\tilde{G}')$. We claim that \tilde{G} is a Frobenius group. Suppose false. Since \tilde{G}/\tilde{G}' acts on \tilde{G}'/\tilde{G}'' , Frobeniusly, we may assume \tilde{G}/\tilde{G}' acts on \tilde{G}'' , trivially and so $\tilde{G}'' \leq \mathbf{Z}(\tilde{G})$,

hence $\tilde{G}'' \cong C_p$. Let $\tilde{T} = \Pi_{\chi \in \text{Irr}(\tilde{G}|\tilde{G}'')} \ker(\chi)$. Then \tilde{T} is a normal subgroup of \tilde{G} , whose intersection with \tilde{G}'' is trivial. Note that $\tilde{T} \times \tilde{G}''$ is a proper subgroup of \tilde{G}' , so the order of $\tilde{G}/\tilde{G}''\tilde{T}$ is divided by $2p$. Also, $\ker(\chi) = 1$, for all $\chi \in \text{Irr}(\tilde{G}/\tilde{T}|\tilde{G}''\tilde{T}/\tilde{T})$. Note that $\tilde{G}/\tilde{G}''\tilde{T} \not\cong D_{2p}$, as otherwise

$$\frac{\frac{\tilde{G}'}{\tilde{T}}}{\frac{\tilde{G}''\tilde{T}}{\tilde{T}}} \cong \frac{\tilde{G}'}{\tilde{T}\tilde{G}''} \cong C_p,$$

which is not possible. Hence $\tilde{G}/\tilde{G}''\tilde{T}$ is isomorphic to the Frobenius group $C_p^\beta \rtimes C_2$, for some integer $\beta > 1$, then $p \in \text{cod}(\tilde{G}/\tilde{G}''\tilde{T})$ and $m'_{\tilde{G}}(p) > 1$. Therefore, using Lemmas 2.7(c) and 2.8(b), \tilde{G}/\tilde{G}'' acts on $\tilde{T}\tilde{G}''/\tilde{T} \cong \tilde{G}''$, Frobeniusly, a contradiction. So our claim is proved and \tilde{G} is a Frobenius group.

It is well-known that if the complements of a Frobenius group are even order, then the Frobenius kernel is abelian. Hence, \tilde{G}' is abelian and $\tilde{G}'' = 1$, which means $G'' = 1$ as wanted.

Finally, assume G' is not a p -group, for some odd prime p . Clearly G'/G'' is not divided by two distinct primes $p < q$, since otherwise G has a homomorphic image isomorphic to D_{2pq} , a contradiction, as $m'_{D_{2pq}}(q) > 1$ and $m'_{D_{2pq}}(pq) > 1$. So, we may assume \overline{G}' is a p -group for some odd prime p .

Then define \mathcal{S} to be the set of all pairs (M, N) , where $N \leq M$ are normal subgroups of G contained in G' and M/N is a chief factor of G whose order is not divided by p . Suppose (M, N) is a pair in \mathcal{S} such that it has the largest first component among all members of \mathcal{S} . Remark that $M \neq G'$, as otherwise, $|G'/G''|$ is divided by two distinct primes.

Now, we work on $\overline{G} = G/N$ and we use bar convention for this quotient group. Let $\overline{C} = C_{\overline{G}'}(\overline{M}) \cong \overline{M} \times \overline{T}$. If $\overline{T} > 1$, then $\overline{C}/\overline{T} \cong \overline{M}$ is a q -group for some prime $q \neq p$, a contradiction, as $|\overline{M}| < |\overline{C}|$. Since $|G/G'| = 2$, we deduce that $C_{\overline{G}'}(\overline{M}) = \overline{M}$ and hence χ is faithful, for all $\chi \in \text{Irr}(\overline{G}|\overline{M})$. First, let $|\overline{G}'/\overline{M}| > 3$. Then, $\overline{G}/\overline{M}$ is isomorphic to the Frobenius group $C_p^\beta \rtimes C_2$, for some integer β , as $|\overline{G}'/\overline{M}|$ is a p -group (notice that $\beta > 1$ when $p = 3$). Hence $m'_{\overline{G}/\overline{M}}(p) > 1$.

We note that $\text{cod}(\chi) \neq p$, for all $\chi \in \text{Irr}(\overline{G}|\overline{M})$ by Lemma 2.7(b). Hence, $\overline{G}/\overline{M}$ acts on \overline{M} , Frobeniusly, by Lemma 2.8(b). As $\overline{G}/\overline{M}$ is Frobenius, we get a contradiction. So we assume $G'/M \cong C_3$ and by Lemmas 4.5 and 4.6, $G \cong S_4$ as wanted. \blacksquare

Lemma 4.8. *Let G be a finite solvable T_k^l -group for some integer $k \geq 1$, and $K < G' < G$ be a normal series of G , such that G/K is a p -group for some prime p . Then either G is an extra-special 2-group, or G is a Frobenius group isomorphic to $C_3^2 \rtimes Q_8$.*

Proof. Applying Lemma 4.1 on G/K , we have $p = 2$. If G' is a 2-group, we are done, so we may assume G' is not a 2-group. Define \mathcal{S} to be a set of all pairs (M, N) , where $N \leq M$ are normal subgroups of G contained in G' and M/N is a chief factor of G whose order is not divided by 2. Suppose (M, N) is a pair in \mathcal{S} such that it has the largest first component among all members of \mathcal{S} . Note that $M \neq G'$, by Lemmas 4.2, 4.3, 4.4 and 4.7, also by Lemma 4.1, G/M is an extra-special group.

Let $\overline{G} = G/N$, and let $\overline{C} = C_{\overline{G}'}(\overline{M}) \cong \overline{M} \times \overline{L}$. If $\overline{L} > 1$, then $\overline{C}/\overline{L} \cong \overline{M}$ is a p -group for some $p \neq 2$, a contradiction, as $|\overline{M}| < |\overline{C}|$. So, $C_{\overline{G}'}(\overline{M}) \cap \overline{G}' = \overline{M}$. Let $\overline{T} = \Pi_{\chi \in \text{Irr}(\overline{G}|\overline{M})} \ker(\chi)$. Then, as $\overline{C} = \overline{M}$, we conclude that $\overline{T} \cap \overline{G}' = 1$. As $\overline{T} \cong \overline{T}\overline{M}/\overline{M}$ is a normal subgroup of $\overline{G}/\overline{M}$ whose intersection with $(\overline{G}/\overline{M})'$ is trivial, we get that $\overline{T} = 1$.

Then, χ is faithful for all $\chi \in \text{Irr}(\overline{G}|\overline{M})$. Also $\text{cod}(\chi) \neq 2$, for all $\chi \in \text{Irr}(\overline{G}|\overline{M})$, by Lemma 2.7(c). So, $\overline{G}/\overline{M}$ acts on \overline{M} , Frobeniusly, by Lemma 2.8(b) and $\overline{G}/\overline{M}$ must be Q_{2^n} , for some integer n . As G/M is an extra-special 2-group, we deduce that $n = 3$. Noting that nonlinear characters in $\text{Irr}(\overline{G}|\overline{M})$ have distinct degrees ($m'_{\overline{G}}(2) > 2$ and $\text{cod}(\chi) \neq 2$ for all $\chi \in \text{Irr}(\overline{G}|\overline{M})$), we deduce all nontrivial irreducible

characters of \overline{M} are \overline{G} -conjugate and hence, $|\overline{M}| - 1 = |\overline{G}/\overline{M}| = 8$ and so \overline{G} is the described Frobenius group in the statement of the lemma.

Now, we prove that $N = 1$. Suppose false, and assume N is a minimal normal subgroup of G of order r^a , for some prime r and integer a . First, we show that $\ker(\chi) = 1$, for all $\chi \in \text{Irr}(G|N)$. Let $L = \prod_{\chi \in \text{Irr}(G|N)} \ker(\chi)$. Then $LN \cong L \times N$ is a normal subgroup of G .

First assume, $L \cong LN/N \neq 1$ and so $L \cong LN/N$ contains M/N . Note that $M/N \cong C_3^2$. Then $M \cong N \times P$, where $P \cong C_3^2$, is a normal subgroup of G . If M/P is not a 2-group, then replacing (M, N) by (M, P) in the above discussion we have $M/P \cong C_3^2$, which is not possible, since G/P and G/N have the same structure and so they both are isomorphic to the Frobenius group $C_3^2 \rtimes Q_8$, and then $m'_G(9) > 1$, which is contradicting the fact that G is a T'_k -group. If M/P is a 2-group, then $N \leq \mathbf{Z}(G)$ and so $|N| = 2$. Then G/P must satisfy Lemma 4.1, which is not possible, as $|G/P| = 16$.

Therefore $L = 1$ and $\ker(\chi) = 1$, for all $\chi \in \text{Irr}(G|N)$. By Lemmas 2.7(c) and 2.8(b), Sylow p -subgroups of G/N , for prime $p \neq r$, act on N , Frobeniusly. By the structure of Sylow subgroups of G/N , we have N is a 3-group and Sylow 2-subgroups of G/N , which are isomorphic to Q_8 , act on N , Frobeniusly. Then, $N \leq \mathbf{Z}(M)$ and since M is the Sylow 3-subgroup of G using Lemma 2.2, we get that $|N| = 9$. Then G is a Frobenius group of order 8×3^4 and G has a unique minimal normal subgroup N . In addition, characters in $\text{Irr}(G|N)$ have distinct degrees, and by Lemmas 2.7(c) and 2.8(b), $I_G(\lambda)/N \cong C_3^2$, for all $\lambda \in \text{Irr}(N)$. Also $\text{cd}(G|N) = \{3 \times 8\}$ using Lemma 2.5. Therefore, as $\text{cd}(G/N) \cap \text{cd}(G|N) = \emptyset$ and G/N is a D_0 -group (see [2]), we conclude that G is a D_0 -group, contradicting the main result of [2]. \blacksquare

Lemma 4.9. *Let G be a finite non-abelian solvable T'_k -group for some integer $k \geq 1$, such that $G/G' \cong C_4$. Then $G \cong \text{Suz}(2)$ or $\text{Dic}_{12} = \text{SmallGroup}(12,1)$.*

Proof. Assume K is a normal subgroup of G contained in G' such that G'/K is a chief factor of G . Then, write $\overline{G} \cong G/K$ and adopt the bar convention for this quotient group. First assume $\ker(\chi) \neq 1$, for some $\chi \in \text{Irr}(\overline{G}|\overline{G}')$, then $\ker(\chi) \cong C_2$ or C_4 . If $\ker(\chi) \cong C_4$, then $\overline{G} \cong \ker(\chi) \times \overline{G}'$ is abelian and we get a contradiction, hence $\ker(\chi) \cong C_2$. If \overline{G}' is a 2-group, we get a contradiction, as in this case $C_2 \cong \overline{G}' \leq \mathbf{Z}(\overline{G})$ and $\overline{G}/\overline{G}' \cong C_4$, which is not possible. So, \overline{G}' is an elementary abelian p -group, for some odd prime p , and by Lemma 4.2, we get that $\overline{G}/\ker(\chi) \cong D_{2p}$. Therefore, $m'_G(4) > 1$ and $m'_G(p) = (p-1)/2$. Hence, $p = 3$ and $\overline{G} \cong \text{Dic}_{12}$.

Now, assume $\ker(\chi) = 1$, for all $\chi \in \text{Irr}(\overline{G}|\overline{G}')$. Now, we claim that $\text{cod}(\chi) \neq 4$, for all $\chi \in \text{Irr}(\overline{G}|\overline{G}')$. Suppose false, then $|\overline{G}'| \times |I_G(\lambda)/(\theta(1)|\overline{G}'|) = 4$, for some $\lambda \in \text{Irr}(\overline{G}')$ and $\theta \in \text{Irr}(I_{\overline{G}}(\lambda)|\lambda)$, by Lemma 2.7(a). Therefore, \overline{G}' is a 2-group and hence $\overline{G}' \leq \mathbf{Z}(\overline{G})$, a contradiction. So, $\text{cod}(\chi) \neq 4$, for all $\chi \in \text{Irr}(\overline{G}|\overline{G}')$. Therefore, members of $\text{Irr}(\overline{G}|\overline{G}')$ must have distinct degrees and hence \overline{G} is a D_0 -group. Using the classification of D_0 -group, we get that $\overline{G} \cong \text{Suz}(2)$.

Now, we aim to prove that K is trivial. On the contrary, we may assume K is a minimal normal subgroup of G .

First, we show that $\ker(\chi) = 1$ for all $\chi \in \text{Irr}(G|K)$. Let $\overline{G} \cong \text{Dic}_{12}$, assume $|\ker(\chi)|$ is divided by 2, for some $\chi \in \text{Irr}(G|K)$ and let $P \in \text{Syl}_2(K)$. By the structure of Dic_{12} , we have $P \cong C_2$. Now, G/P satisfies Lemma 4.7, and it follows that $G/P \in \{S_4, D_{18}, C_3^2 \rtimes C_2\}$, and hence $m'_G(t) > 1$, for some $t \neq 4$, a contradiction. So $|\ker(\chi)|$ is not divided by 2. Paying attention to the structure of Dic_{12} and $\text{Suz}(2)$ we have $G' \cong K \times T$, where $T = \ker(\chi) \cap G' \cong C_5$ or C_3 and applying the above discussion on G/T , we get that $G/K \cong \text{Suz}(2)$ or Dic_{12} , and so $T \cong C_5$ or C_3 . If $G/T \cong G/K$, then $m'_G(t) > 1$ for some $t \in \{6, 5\}$ (both G/K and G/T are isomorphic to $\text{Suz}(2)$ or Dic_{12}) and, as $m'_G(4) > 1$, we get a contradiction. Therefore one of those groups is isomorphic to Dic_{12} and the other one is isomorphic

to $\text{Suz}(2)$. So $G' \cong C_{15}$, has 8 irreducible characters of order 15, whose inertia subgroups are G' . Therefore, G has 2 distinct irreducible characters of the same degree 4. It is easy to see that both of those characters are faithful and so $m'_G(15) > 1$, a contradiction. So $\ker(\chi) = 1$, for all $\chi \in \text{Irr}(G|K)$.

We claim that $\text{cod}(\chi) \neq 4$, for all $\chi \in \text{Irr}(G|K)$. Otherwise, $|K| \times |I_G(\lambda)|/(\theta(1)|K|) = 4$, for some $\lambda \in \text{Irr}(K)$ and $\theta \in \text{Irr}(I_G(\lambda)|\lambda)$, by Lemma 2.7(a). Since all Sylow subgroups of G/K are cyclic, then λ extends to $I_G(\lambda)$ (see [9, Corollary 11.31]), and so $\theta(1)$ is a 2-element (by noticing the character degrees of Dic_{12} and $\text{Suz}(2)$). Hence, both $|K|$ and $|I_G(\lambda)|/|K|$ are 2-groups, whose orders are at most 4 and then, $3 \leq |G : I_G(\lambda)|$, while $|K| \leq 4$. By Remark 2.1, the only possibility is that $|K| = 4$ and $|G : I_G(\lambda)| = 3$, which means that $G/K \cong \text{Dic}_{12}$, $I_G(\lambda)/K \cong C_4$ and $\theta(1) = 4$, a contradiction. So our claim is proved.

Therefore, the members of $\text{Irr}(G|K)$ has distinct degrees (as $m'_G(4) > 1$). Since all Sylow subgroups of $I_G(\lambda)/K$ are cyclic, then λ is extendible to $I_G(\lambda)$ (see [9, Corollary 11.31]), and using Gallagher's theorem, we deduce that $I_G(\lambda)/K = 1$, for all $\lambda \in \text{Irr}(K)$. It follows that G/K acts on K , Frobeniusly, but it is a contradiction, when $G/K \cong \text{Suz}(2)$. Then we may assume $G/K \cong \text{Dic}_{12}$. Then, $\text{cd}(G|K) = \{12\}$ and so $\text{cd}(G|K) \cap \text{cd}(G/K) = \emptyset$. On the other hand G/K is a D_1 -group, which implies that G is a D_1 -group. Now, looking at the structure of D_1 -groups (see [1]), we get a contradiction. ■

Proof of the Main Theorem case (b): Easy calculation shows us that groups in cases (1) and (2) are T'_k -groups, for some $k \leq 3$. Also by Lemma 2.6, elementary abelian group G is a $T'_{|G|-1}$ -group. If G is an extra-special 2-group, since it has a unique nonlinear irreducible character and all of nontrivial linear characters have the same codegrees, we get that G is a $T'_{|G/G'|-1}$ -group. Therefore the described groups in the case (3) are $T'_{|G/G'|-1}$ -groups.

Let G be a Frobenius group isomorphic to $C_2^\beta \rtimes C_{2^\beta-1}$, where β is an integer and $p = 2^\beta - 1$ is prime. Then, G has a unique nonlinear irreducible character and all of its nontrivial linear irreducible characters have the same codegree p . Therefore G is a $T'_{(p-1)}$ -group.

Let G be a Frobenius group isomorphic to $C_p^\beta \rtimes C_2$, for some prime p and integer β . We know that $\lambda^G \in \text{Irr}(G)$ for every $\lambda \in \text{Irr}(G')$ and $|\ker(\lambda^G)| = p^{\beta-1}$. Since the size of G -orbit of each nontrivial element in $\text{Irr}(G')$ is 2, then G has exactly $(p^\beta - 1)/2$ nonlinear irreducible characters with the same codegree p . As G has just one linear nontrivial irreducible character, we deduce that G is a $T'_{(p^\beta-1)/2}$ -group.

Now, we just need to prove the "only if" part of the theorem. If G is abelian, by Lemma 2.6, we have either G is an elementary abelian p -group, for some prime p , or $G \cong C_4$. So we assume G is not abelian. As G/G' is a T'_l -group, for some integer l , we deduce that G/G' is a cyclic group of order 4 or an elementary abelian group. First, assume $G/G' \cong C_p^\alpha$, for some prime p and integer α . Let K be a normal subgroup of G , contained in G' , such that G'/K is a chief factor of G . If G'/K is not a p -group, then by Lemma 4.2, we have G/K is one of the groups described in the statement of Lemma 4.2. First, let G/K be isomorphic to a Frobenius group of order $p2^\beta$, where $p = 2^\beta - 1$. Then applying Lemma 4.3, we have $K = 1$ and G is the group described in case (4) of the main theorem. If $G/K \cong C_2 \times S_3$, applying Lemma 4.4, $K = 1$ and G is one of the groups, described in case (2) of the main theorem. If $G/K \cong D_{2q}$, for some odd prime q , then applying Lemma 4.7, either $G \cong S_4$, D_{18} or it is isomorphic to the Frobenius group described in case (5) of the main theorem. Now, let G'/K be a p -group. Then, applying Lemma 4.8, G is one of the groups described in cases (2) and (3). At last, suppose $G/G' \cong C_4$. Hence, applying Lemma 4.9, $G \cong \text{Suz}(2)$ or Dic_{12} , which are two of the groups in case (1) of the main theorem. □

REFERENCES

- [1] Y. Berkovich, ‘Finite solvable groups in which only two nonlinear irreducible characters have equal degrees’, *J. Algebra*, 184 (1996), 584–603.
- [2] Y. Berkovich, D. Chillag and M. Herzog, ‘Finite groups in which character degrees of nonlinear irreducible characters are distinct’, *Proc. Amer. Math. Soc.*, 115 (4) (1992), 955–959.
- [3] Y. Berkovich, I. M. Isaacs and L. Kazarin, ‘Groups with distinct monolithic character degrees’, *J. Algebra*, 216 (1) (1999), 448–480.
- [4] Y. Berkovich and L. Kazarin, ‘Finite nonsolvable groups in which only two nonlinear irreducible characters have equal degrees’, *J. Algebra*, 184 (1996), 538–560.
- [5] M. Bianchi, D. Chillag, M. L. Lewis and E. Pacifici, ‘Character degree graphs that are complete graphs’, *Proc. Amer. Math. Soc.*, 135(3) (2007), 671–676.
- [6] M. R. Darafshah, A. Iranmanesh and S. A. Moosavi, ‘Groups whose nonlinear characters are rational valued’, *Arch. Math.*, 94 (2010), 411–418.
- [7] N. Du and M. L. Lewis, ‘Codegrees and nilpotence class of p -groups’, *J. Group Theory*, 19 (2015), 561–567.
- [8] M. Ebrahimi, ‘Groups in which the co-degrees of the irreducible characters are distinct’, Preprint (2020), <https://arxiv.org/abs/2008.02433>.
- [9] I. M. Isaacs, ‘Character theory of finite groups’, *New York, Academic Press*, (1976).
- [10] I. M. Isaacs, ‘Finite group theory’, *Amer. Math. Soc, Providence*, (2008).
- [11] D. Liang and G. Qian, ‘Finite groups with coprime character degrees and codegrees’, *J. Group Theory*, 19 (2016), 763–776.
- [12] Y. Liu and Z. Q. Lu, ‘Solvable D_2 -groups’, *Acta Math. Sin. (Engl. Ser.)*, 33 (2017), 77–95.
- [13] Y. Liu and Z. Q. Lu, ‘Non-solvable D_2 -groups’, *Acta Math. Sin. (Engl. Ser.)*, 31(11) (2015), 1683–1702.
- [14] G. Qian, Y. Wang and H. Wei, ‘Codegrees of irreducible characters in finite groups’, *J. Algebra*, 312 (2007), 946–955.
- [15] Z. Sayanjali, Z. Akhlaghi and B. Khosravi, ‘On the multiplicity of character degrees of non-solvable groups’, *J. Algebra Appl.*, DOI: 10.1142/S0219498821500766.
- [16] H. P. Tong-Viet, ‘Finite nonsolvable groups with many distinct character degrees’, *Pacific J. Math.*, 268 (2) (2014), 477–492.

ZEINAB AKHLAGHI, FACULTY OF MATH. AND COMPUTER SCI.,
 AMIRKABIR UNIVERSITY OF TECHNOLOGY (TEHRAN POLYTECHNIC), 15914 TEHRAN, IRAN.
 SCHOOL OF MATHEMATICS, INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCE(IPM) P.O. BOX:19395-5746, TEHRAN,
 IRAN.

Email address: z_akhlaghi@aut.ac.ir

MEHDI EBRAHIMI, SCHOOL OF MATHEMATICS, INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM), P. O. BOX
 19395-5746, TEHRAN, IRAN.

Email address: m.ebrahimi.math@ipm.ir

MARYAM KHATAMI, DEPARTMENT OF PURE MATHEMATICS, FACULTY OF MATHEMATICS AND STATISTICS, UNIVERSITY OF
 ISFAHAN, ISFAHAN, 81746-73441, IRAN.

Email address: m.khatami@sci.ui.ac.ir