

ZIMMER'S CONJECTURE FOR NON-UNIFORM LATTICES AND ESCAPE OF MASS

AARON BROWN, DAVID FISHER, AND SEBASTIAN HURTADO

ABSTRACT. We establish finiteness of low-dimensional actions of lattices in higher-rank semisimple Lie groups and establish Zimmer's conjecture for many such groups. This builds on previous work of the authors handling the case of actions by cocompact lattices and of actions by $SL(n, \mathbb{Z})$. While the results are not sharp in all cases, they do dramatically improve all known results. The key difficulty overcome in this paper concerns escape of mass when taking limits of sequences of measures. Due to a need to control Lyapunov exponents for unbounded cocycles when taking such limits, quantitative controls on the concentration of mass at infinity are needed and novel techniques are introduced to avoid "escape of Lyapunov exponent."

CONTENTS

1. Introduction and main results	2
1.1. Motivation	2
Acknowledgements	4
1.2. Results for lattices in \mathbb{R} -split simple Lie groups	4
1.3. Critical dimensions, full conjecture, and general result	5
1.4. Actions by C^1 diffeomorphisms	6
1.5. Proof of Theorem B	6
2. Proof of Theorem C, main technical result, and outline of proof	8
2.1. Reductions and standing hypotheses	8
2.2. Main technical theorem and proof of Theorem C	8
2.3. Review of the proof of [17, Theorem B]	10
2.4. Outline of proof of Theorem D and discussion of new tools	11
3. Algebraic groups and reductions for nonuniform Γ	15
3.1. Basic terminology	15
3.2. Algebraic tori and \mathbb{K} -rank	16
3.3. Levi and Langlands decompositions	16
3.4. Restricted \mathbb{K} -roots and parabolic subgroups of algebraic groups	17
3.5. Iwasawa decomposition, restricted roots, and parabolic subgroups	21
3.6. Standard \mathbb{Q} -rank-1 subgroups	22
3.7. Standing hypotheses when Γ is nonuniform	23
3.8. Lifting \mathbb{Q} -structures from F to G	24

Date: June 1, 2021.

DF was partially supported by NSF Grants DMS-1308291 and DMS-1607041. DF was also partially supported by the University of Chicago, and by NSF grants DMS 1107452, 1107263, 1107367, "RNMS: Geometric Structures and Representation Varieties" (the GEAR Network) during a visit to the Isaac Newton Institute in Cambridge. AB was partially supported by NSF Grant DMS-1752675. The authors would like to thank UoC, IPAM, IUB for their hospitality at various points.

3.9. Consequences of standing hypotheses: quasi-isometric properties	24
3.10. Consequences of standing hypotheses: quasi-isometric bounded generation	29
4. Averaging measures and control of mass near ∞	30
4.1. Notation	30
4.2. Control of mass near ∞	31
4.3. Translates of horospherical orbits and mass at ∞	31
4.4. Non-divergence of unipotent averages of measures	33
4.5. Equidistribution and invariance	35
5. Siegel sets, cocycles, and induced actions	36
5.1. Siegel sets and Siegel fundamental sets	36
5.2. Coarse geometry and partitions of unity	39
5.3. Induced actions	40
5.4. Construction of norms on \mathcal{E} and metric on X	41
5.5. Growth estimates on the cocycle \mathcal{A}	44
6. Lyapunov exponents	49
6.1. Lyapunov exponents for unbounded cocycles	50
6.2. Lyapunov exponents under averaging	51
6.3. Lyapunov exponents for pre-compact orbits	52
6.4. The projectivized cocycle and its infinitesimal generator	54
Review of notation and standing hypotheses	55
7. Subexponential growth for the center	56
7.1. Additional hypothesis for Sections 8, 9, and 10	56
8. Subexponential growth for unipotent subgroups	56
8.1. Subexponential growth for generic $\{a^t\}$ -orbits	58
8.2. Subexponential growth for representatives of hyperbolic directions	59
8.3. Proof of Proposition 8.1	64
9. Subexponential growth for minimal parabolic \mathbb{Q} -subgroups	67
10. Subexponential growth for \mathbb{Q} -rank-1 subgroups	69
10.1. Preliminaries, key ingredients and outline	69
10.2. Characterization and properties of orbits with exponential growth	71
10.3. Time-averaged cocycle, cut-off cocycle, and time-averaged exponents	77
10.4. Invariance and additivity of the time-averaged cocycle	85
10.5. Proof of Proposition 10.1	86
11. Sketch of alternate proof in higher \mathbb{Q} -rank	92
Appendix A. Tables of computations	94
References	95

1. INTRODUCTION AND MAIN RESULTS

1.1. **Motivation.** It is well known that lattices in $\mathrm{SL}(2, \mathbb{R})$ are rather flexible whereas lattices in $\mathrm{SL}(n, \mathbb{R})$ are very rigid when $n \geq 3$. Such rigidity properties are shared by lattices in more general higher-rank simple Lie groups and, to a lesser extent, by irreducible lattices in higher-rank semisimple Lie groups. Some key results demonstrating such rigidity phenomena include the strong rigidity theorem of Mostow [53], Kazhdan's property (T) [33], and the superrigidity theorem of Margulis [47, Theorem IX.6.16]. For instance, Margulis's superrigidity theorem

states, roughly, that every finite-dimensional representation of an irreducible lattice in a higher-rank semisimple Lie group extends, modulo compact groups and passing to covers, to a continuous representation of the ambient group.

Inspired by known rigidity results for linear representations, Zimmer proposed in the 1980s that certain “nonlinear representations” of higher-rank lattices should also exhibit some rigidity properties. Specifically, Zimmer laid out a program to study C^∞ volume-preserving actions of such groups on compact manifolds; that is, for a compact manifold M equipped with a smooth volume form ω , Zimmer aimed to study homomorphisms $\alpha: \Gamma \rightarrow \text{Diff}^\infty(M, \omega)$ from higher-rank lattice subgroups Γ to $\text{Diff}^\infty(M, \omega)$, the group of C^∞ , ω -preserving diffeomorphisms. In later work of Zimmer and other authors, the case of non-volume-preserving actions is also considered.

The most ambitious goal of this program is to show that all such non-linear representations are derived from certain homogeneous spaces and algebraic actions induced by linear representations. In particular, all volume-preserving actions $\alpha: \Gamma \rightarrow \text{Diff}^\infty(M, \omega)$ are expected to be of an “algebraic origin;” more general actions $\alpha: \gamma \rightarrow \text{Diff}^\infty(M)$ are expected to have an “algebraic quotient.” The **Zimmer program** refers to a number of precise conjectures towards this goal. See for instance the surveys of Fisher and Labourie [25, 41] for overviews of conjectures and results in this area and [26] for an update with more recent developments.

As a first step in the program, Zimmer conjectured a lower bound on the dimension of a compact manifold M admitting a nonlinear, volume-preserving representation $\alpha: \Gamma \rightarrow \text{Diff}^\infty(M, \omega)$ with infinite image. See Conjecture 1.2 below for a precise formulation. This lower bound is computed in terms of algebraic data associated with the ambient higher-rank group G and its representations. To motivate this conjecture, we recall the following immediate consequence of Margulis’s superrigidity and arithmeticity theorems: for $n \geq 3$, if Γ is a lattice subgroup in $\text{SL}(n, \mathbb{R})$ and if $d < n$ then every homomorphism $\rho: \Gamma \rightarrow \text{GL}(d, \mathbb{R})$ has finite image. Zimmer’s conjecture, Conjecture 1.2, is the natural non-linear analogue of this fact.

The main heuristic for Zimmer’s conjecture is Zimmer’s cocycle superrigidity theorem [70]. An immediate corollary of Zimmer’s cocycle superrigidity theorem is the following: given a lattice Γ in a higher-rank simple Lie group and a volume-preserving action $\alpha: \Gamma \rightarrow \text{Diff}^\infty(M, \omega)$ on a compact manifold M of sufficiently small dimension, there exists a measurable, $\alpha(\Gamma)$ -invariant Riemannian metric g on TM . If this measurable metric can be promoted to a continuous metric then the image $\alpha(\Gamma)$ is contained in the compact Lie group of isometries $\text{Isom}_g(M)$; finiteness of the action $\alpha(\Gamma)$ follows if $\dim(M)$ is sufficiently small by known results on representations of such Γ into compact Lie groups.

While the above heuristic requires the action α to preserve a volume-form, results for actions on the circle [19, 29, 69], results for real-analytic actions [24, 28], and results on projective quotients [54] suggest analogous lower bounds should exist on the dimension of manifolds M admitting nonlinear representations $\alpha: \Gamma \rightarrow \text{Diff}^\infty(M)$ with infinite image.

To state the simplest version of the conjecture, given a semisimple Lie group we define $v(G)$ to be the minimal dimension of a homogeneous space G/H , where H ranges over all proper, closed subgroups of the identity component G° of G . We define $n(G)$ to be the minimal dimension of a non-trivial linear representation of the Lie algebra of G . Given these dimension bounds, we have the following.

Conjecture 1.1. *Let G be a connected semisimple Lie group such that every non-compact factor of G has real rank at least 2. Let Γ be a lattice subgroup in G . Let M be a compact manifold and let $\alpha: \Gamma \rightarrow \text{Diff}^\infty(M)$ be a homomorphism.*

- (a) *If $\dim(M) < v(G)$ then $\alpha(\Gamma)$ preserves a smooth Riemannian metric on M .*
- (b) *If $\dim(M) < n(G)$ and if $\alpha(\Gamma)$ preserves a volume form then $\alpha(\Gamma)$ preserves a smooth Riemannian metric on M .*

Using that the isometry group $\text{Isom}(M)$ of a compact Riemannian manifold is compact, in the case that Γ is non-uniform, the conclusion of Conjecture 1.1 implies that the image $\alpha(\Gamma)$ is finite. Using Margulis's superrigidity theorem for representations into compact groups, from Conjecture 1.1 we obtain an upper bound on the dimension $\dim(M)$ below which the image $\alpha(\Gamma)$ is finite; see Conjecture 1.2 and discussion in Remark 1.3.

The main results of this paper establishes Conjecture 1.1(a) and some case of Conjecture 1.1(b) for $C^{1+\text{H\"older}}$ actions of lattices Γ in \mathbb{R} -split simple Lie groups. See Theorem A. We also establish partial results towards the conjecture for other groups. See Theorem B.

Acknowledgements. The authors owe a profound debt to Dave Witte Morris. In particular, Theorem 3.31 was proven by Witte Morris in [49] in response to a question from the authors. In addition, Witte Morris helped the authors a great deal with Section 3 and in particular explained to us the proof of Lemma 3.8.

1.2. Results for lattices in \mathbb{R} -split simple Lie groups. We show that Conjecture 1.1(a) holds for $C^{1+\text{H\"older}}$ actions of lattices Γ in \mathbb{R} -split simple Lie groups. For Conjecture 1.1(b) our method of proof only yields the optimal conjectured upper bound for groups with Lie algebra $\mathfrak{sl}(n, \mathbb{R})$ or $\mathfrak{sp}(2n, \mathbb{R})$ though we obtain partial results for all other \mathbb{R} -split groups.

Given $0 < \beta \leq 1$ we recall that a diffeomorphism $f: M \rightarrow M$ of a compact manifold M is $C^{1+\beta}$ if it is C^1 and the variation of its derivative is β -H\"older in local charts. We write $\text{Diff}^{1+\beta}(M)$ for the group of $C^{1+\beta}$ diffeomorphisms of M .

In the statement of all results, we will slightly abuse terminology and call a connected Lie group G **simple** if its Lie algebra \mathfrak{g} is simple; that is, G is simple if it is connected and has no non-trivial connected normal subgroups. Note in particular, we allow for the case that G has infinite center.

A connected simple Lie group G is **\mathbb{R} -split** if its Lie algebra \mathfrak{g} is a split real form of a simple complex Lie algebra $\mathfrak{g}^{\mathbb{C}}$. All \mathbb{R} -split simple Lie groups with finite center are isogenous to either $\text{SL}(n, \mathbb{R})$, $\text{Sp}(2n, \mathbb{R})$, $\text{SO}(n, n)$, or $\text{SO}(n, n-1)$ or to one of the five exceptional \mathbb{R} -split simple Lie groups.

Theorem A. *Let G be a connected \mathbb{R} -split simple Lie group with real rank at least 2 and let $\Gamma \subset G$ be a lattice subgroup. Let M be a compact manifold and let $\alpha: \Gamma \rightarrow \text{Diff}^{1+\beta}(M)$ be a homomorphism for $\beta > 0$.*

- (a) *If $\dim(M) < v(G)$ then $\alpha(\Gamma)$ preserves a continuous Riemannian metric on M ; moreover the image $\alpha(\Gamma)$ is finite.*
- (b) *If $\dim(M) \leq v(G)$ and if $\alpha(\Gamma)$ preserves a volume form then $\alpha(\Gamma)$ preserves a continuous Riemannian metric on M ; moreover the image $\alpha(\Gamma)$ is finite.*

A special case of Theorem A (and of Theorem B below) for cocompact Γ was proved by the authors in [16]. This paper primarily concerns the case that Γ is not cocompact and develops many new arguments to overcome the lack of compactness

of the homogeneous space G/Γ . The standard example of a non-uniform lattice is the subgroup $\mathrm{SL}(n, \mathbb{Z})$ in $\mathrm{SL}(n, \mathbb{R})$; for this special case, Theorem A was established in a subsequent paper [17]. This paper extends and generalizes many of the ideas from [17] to the setting of general non-uniform lattices in $\mathrm{SL}(n, \mathbb{R})$ and to other non-uniform lattices in other simple Lie groups of higher real rank.

1.3. Critical dimensions, full conjecture, and general result. We state the full version of Zimmer's conjecture in order to put our results in context. As before, given a semisimple Lie group G , let $n(G)$ denote the minimal dimension of a non-trivial real representation of the Lie algebra \mathfrak{g} of G and let $v(G)$ denote the minimal codimension of all proper subgroups H of G ; we note that this is equivalent to the minimal codimension of all proper parabolic subgroups Q of G . (See [16, Lemma 3.7].) Let $d(G)$ denote the minimal dimension of all non-trivial homogenous spaces K/C as K varies over all compact real-forms of all simple factors of the complexification of G . Given the numbers $n(G)$, $d(G)$, and $v(G)$ the full conjecture is the following.

Conjecture 1.2 (Zimmer's conjecture). *Let G be a connected semisimple Lie group, all of whose simple factors have real rank at least 2, and let $\Gamma \subset G$ be a lattice subgroup. Let M be a compact manifold and let ω be a volume form on M .*

- (a) *If $\dim(M) < \min(d(G), v(G))$ then any homomorphism $\alpha: \Gamma \rightarrow \mathrm{Diff}(M)$ has finite image.*
- (b) *If $\dim(M) < \min(d(G), n(G))$ then any homomorphism $\alpha: \Gamma \rightarrow \mathrm{Diff}(M, \omega)$ has finite image.*
- (c) *If $\dim(M) < n(G)$ then for any homomorphism $\alpha: \Gamma \rightarrow \mathrm{Diff}(M, \omega)$, the image $\alpha(\Gamma)$ preserves a Riemannian metric.*
- (d) *If $\dim(M) < v(G)$ then for any homomorphism $\alpha: \Gamma \rightarrow \mathrm{Diff}(M)$, the image $\alpha(\Gamma)$ preserves a Riemannian metric.*

Remark 1.3.

- (1) The finiteness of the image $\alpha(\Gamma)$ in parts (a) and (b) of Conjecture 1.2 follow from the conditions in parts (c) and (d) and Margulis's superrigidity theorem for representations into compact groups. If g is an $\alpha(\Gamma)$ -invariant continuous Riemannian metric on M then the isometry group K of (M, g) is a compact Lie group and $\alpha(\Gamma)$ is contained in K . Margulis's superrigidity theorem implies that either $\alpha(\Gamma)$ is finite or K contains a compact form of a simple factor of G . The latter option is ruled out by dimension considerations. See [16, §2.3] for a detailed discussion.
- (2) For a non-uniform lattice, Margulis's superrigidity theorem also implies that any homomorphism from Γ to a compact Lie group K has finite image. In our context, this implies that whenever $\alpha(\Gamma)$ preserves a Riemannian metric, $\alpha(\Gamma)$ is finite.
- (3) The tools in this paper are very well adapted to study parts (a) and (b) of Conjecture 1.2. In situations in which parts (c) and (d) are not subsumed by (a) and (b), certain technical arguments used in our proof seem insufficient to give a complete answer but do yield partial results.

For Lie groups that are not \mathbb{R} -split, we obtain partial results towards Conjecture 1.2. To each simple Lie algebra \mathfrak{g} , we associate a number $r(\Sigma_{\mathfrak{g}})$, computed in

terms of the restricted root system $\Sigma_{\mathfrak{g}}$ of \mathfrak{g} , and define

$$r(G) = r(\mathfrak{g}) = r(\Sigma_{\mathfrak{g}}).$$

The number $r(\mathfrak{g})$ was defined in terms of combinatorics of root systems in [16]; that definition is reviewed in Definition 3.11 below. For \mathbb{R} -split Lie groups G , we always have $r(G) = v(G)$. More generally, for simple G it follows from [11, Theorem 7.2] that $r(G) = v(G')$, where G' is any maximal connected, \mathbb{R} -split, semisimple subgroup of G . See Remark 3.12 below. Given a semisimple Lie algebra \mathfrak{g} , we have $r(\mathfrak{g}) = \min r(\mathfrak{g}')$ where the minimum is taken over every non-compact, non-trivial simple ideal $\mathfrak{g}' \subset \mathfrak{g}$.

With the above notation, we have the following extension of Theorem A.

Theorem B. *Let G be a semisimple Lie group such that every non-compact factor has real rank at least 2. Let $\Gamma \subset G$ be a lattice subgroup and let M be a compact manifold. Fix $0 < \beta \leq 1$.*

- (1) *If $\dim(M) < r(G)$ then any homomorphism $\alpha: \Gamma \rightarrow \text{Diff}^{1+\beta}(M)$ has finite image.*
- (2) *If ω is a volume form on M and if $\dim(M) \leq r(G)$ then any homomorphism $\alpha: \Gamma \rightarrow \text{Diff}^{1+\beta}(M, \omega)$ has finite image.*

See Tables 1–4 in Appendix A for computations of the numerology v, n, d , and r associated to various Lie groups.

1.4. Actions by C^1 diffeomorphisms. Certain arguments used in this paper require the action to be by diffeomorphisms that are at least $C^{1+\beta}$ for $\beta > 0$; specifically, Proposition 2.1 below uses tools from smooth ergodic theory in its proof which requires the action to be at least $C^{1+\beta}$. However, most arguments in this paper work for C^1 actions and an analogue of Theorem C below holds in the C^1 setting under more restrictive dimension assumptions. Using results from [15] one obtains the following.

Theorem 1.4 (c.f. [15, Theorem 1]). *Let G be a connected simple Lie group with real rank at least 2. Let $\Gamma \subset G$ be a lattice subgroup and let M be a compact manifold.*

- (1) *If $\dim(M) < \text{rank}_{\mathbb{R}}(G)$ then any homomorphism $\alpha: \Gamma \rightarrow \text{Diff}^1(M)$ has finite image.*
- (2) *If ω is a volume form on M and if $\dim(M) \leq \text{rank}_{\mathbb{R}}(G)$ then any homomorphism $\alpha: \Gamma \rightarrow \text{Diff}^1(M, \omega)$ has finite image.*

It is easy to check that the dimension bound given by Theorem 1.4 only matches the conjectured bounds in Conjecture 1.2 for lattices in groups with Lie algebra $\mathfrak{sl}(n, \mathbb{R})$.

In the case of C^1 actions, we replace Proposition 2.1 below with [15, Proposition 3]

1.5. Proof of Theorem B. We end the introduction with the proof of Theorem B, introducing the first technical result of the paper, Theorem C below. The proof of Theorem B follows the same outline as in [16].

Step 1: Subexponential growth of derivatives. We recall that lattice subgroups of semisimple Lie groups are finitely generated. Given a finitely generated group Γ ,

fix a finite, symmetric generating set $S \subset \Gamma$ and let $\text{len}_\Gamma: \Gamma \rightarrow \mathbb{N}$ denote the word-length norm on Γ relative to S . The following definition is the key property we establish in this paper.

Definition 1.5. *Let M be a compact manifold and let $\alpha: \Gamma \rightarrow \text{Diff}^1(M)$ be an action. We say the action α has **uniform subexponential growth of derivatives** if for all $\varepsilon > 0$ there is $C_\varepsilon > 0$ such that for all $\gamma \in \Gamma$,*

$$\sup_{x \in M} \|D_x \alpha(\gamma)\| \leq C_\varepsilon e^{\varepsilon \text{len}_\Gamma(\gamma)}. \quad (1.1)$$

In the sequel, we often restrict α to certain subgroups $\Lambda \subset \Gamma$. If Λ is a subgroup of Γ , we say that the restriction $\alpha|_\Lambda$ has **uniform len_Γ -subexponential growth of derivatives** if (1.1) holds for all $\gamma \in \Lambda$. We note in this definition that we always measure the word-length of $\gamma \in \Lambda$ relative to the ambient group Γ rather than any intrinsic word-length on the subgroup Λ .

Our first technical result of this paper is the following which shows under the dimension constraints in Theorem B that every action has uniform subexponential growth of derivatives.

Theorem C. *Let G be a connected semisimple Lie group such that every non-compact simple factor has real rank at least 2. Let $\Gamma \subset G$ be a lattice subgroup, let M be a compact manifold, and let $0 < \beta \leq 1$.*

- (1) *If $\dim(M) < r(G)$ then any homomorphism $\alpha: \Gamma \rightarrow \text{Diff}^{1+\beta}(M)$ has uniform subexponential growth of derivatives.*
- (2) *If ω is a volume form on M and if $\dim(M) \leq r(G)$ then any homomorphism $\alpha: \Gamma \rightarrow \text{Diff}^{1+\beta}(M, \omega)$ has uniform subexponential growth of derivatives.*

Step 2: Strong property (T). We apply [16, Theorem 2.4] and de la Salle's recent result establishing strong property (T) for nonuniform lattices [22, Theorem 1.1] to conclude that any action α satisfying the conclusions of Theorem C preserves a continuous Riemannian metric g . In particular, the image $\alpha(\Gamma)$ is contained in the compact Lie group $K = \text{Isom}_g(M)$. For completeness, we recall these two results.

Theorem 1.6 ([22, Theorem 1.1]). *Assume G is a connected semisimple Lie group such that the real rank of every simple factor is at least two and let Γ be a lattice in G . Then Γ has strong property (T).*

For the special case of cocompact lattices, the analogous result was established in [23, 42].

Theorem 1.7 ([16, Theorem 2.4]). *Let Γ be a finitely generated group and let M be a compact manifold. For $k \geq 2$, let $\alpha: \Gamma \rightarrow \text{Diff}^k(M)$ be an action. If α has uniform subexponential growth of derivatives and if Γ has strong property (T) then $\alpha(\Gamma)$ preserves a Riemannian metric that is $C^{k-1-\delta}$ for all $\delta > 0$.*

For actions by $C^{1+\beta}$ diffeomorphism, the proof of [16, Theorem 2.4] can be adapted to establish an analogue of Theorem 1.7 to obtain a continuous invariant Riemannian metric. For actions by C^1 diffeomorphisms, an analogue of Theorem 1.7 is obtained in [15, Proposition 5].

Step 3: Superrigidity with compact codomains. As discussed in Remark 1.3, the finiteness of the image $\alpha(\Gamma)$ follows immediately from Margulis's superrigidity theorem with compact codomain. Theorem B then follows.

2. PROOF OF THEOREM C, MAIN TECHNICAL RESULT, AND OUTLINE OF PROOF

2.1. Reductions and standing hypotheses. Without loss of generality, we may assume the following hypotheses for the proof of Theorem C:

- (1) G is a connected semisimple Lie group with real rank at least 2;
- (2) G is simply connected (as a topological group);
- (3) Γ is an irreducible lattice subgroup of G ;
- (4) G has no compact factors; i.e. G has no non-trivial connected compact normal subgroups.

Indeed, we first note that Theorem C holds if it holds for finite-index subgroups $\Gamma' \subset \Gamma$ or for subgroups Γ'' containing Γ as a subgroup of finite index (where Γ'' acts on the compact manifold $(\Gamma'' \times M)/\Gamma$). In particular, we may replace Γ with a group commensurable with Γ in G . Furthermore, if Γ is reducible (see [47, (5.9) Definition, p. 133]) and the conclusion holds for the restriction of α to each irreducible component of Γ , then the conclusion holds for the action of Γ .

Given an arbitrary connected semisimple Lie group G as in Theorem C, we may replace G with its topological universal cover \tilde{G} ; then the preimage of Γ under the map $\tilde{G} \rightarrow G$ is a lattice subgroup $\tilde{\Gamma}$ of \tilde{G} and any action of Γ extends to an action of $\tilde{\Gamma}$. The conclusion of Theorem C then follows for an action of Γ if it is verified for the induced action of $\tilde{\Gamma}$.

Assume now that G is simply connected. Let $K \subset G$ and $L \subset G$ denote, respectively, the product of all compact (resp. noncompact) normal subgroups. As G is simply connected, G is a direct product of K and L . Then $G/L = K$ is compact, hence linear, whence the image of Γ in G/L contains a torsion-free subgroup of finite index. Replacing Γ with a finite-index subgroup, we may assume $\Gamma \cap K$ is a singleton. Then the map $G \rightarrow G/K$ is one-to-one on Γ and we may replace G with G/K and Γ with its image in G/K .

2.2. Main technical theorem and proof of Theorem C. To establish Theorem C we have the following theorem which gives a dichotomy for C^1 actions of higher-rank lattices: either an action $\alpha: \Gamma \rightarrow \text{Diff}^1(M)$ has uniform subexponential growth of derivatives or there exists an A -invariant measure (where A is a split Cartan subgroup) for the induced G -action with certain dynamical properties. To prove Theorem C we show the properties of this measure contradict known results.

The following is the main technical result of this paper. To state the theorem, we refer the reader to Section 5.3 for the definition of the induced G -space X and to Section 6.1 for the definition of the average top Lyapunov exponent. Let Z denote the center of G . See also Section 3.5 for the definition of a split Cartan subgroup $A \subset G$. The cocycle \mathcal{A} is taken to be the fiberwise derivative cocycle for the induced G -action on X .

Theorem D. *Let G be a connected semisimple Lie group without compact factors and with $\text{rank}_{\mathbb{R}} G \geq 2$. Let Γ be an irreducible lattice subgroup in G , let M be a compact manifold, and let $\alpha: \Gamma \rightarrow \text{Diff}^1(M)$ be an action. Let $X = (G \times M)/\Gamma$ denote the induced G -space and let \mathcal{A} denote the corresponding fiberwise derivative cocycle on the bundle $(G \times TM)/\Gamma \rightarrow X$.*

If α fails to have uniform subexponential growth of derivatives then there exists a split Cartan subgroup A of G and a Borel probability measure μ on X such that

- (1) μ is (ZA) -invariant,

- (2) μ projects to the Haar measure on G/Γ , and
- (3) for some $a \in \mathcal{Z}A$, the average top Lyapunov exponent $\lambda_{\text{top},a,\mu,\mathcal{A}}$ is positive.

We remark that there are no constraints on the dimension of M in the statement of Theorem D. Additionally, in Theorem D we do not assume every factor of G is higher rank; in particular, the theorem applies to actions by irreducible lattices in semisimple Lie groups with rank-1 factors. Finally, we note that if we assume \mathcal{Z} is finite, we may conclude $\lambda_{\text{top},a,\mu,\mathcal{A}} > 0$ for some $a \in A$ as in the conclusion of the analogous results appearing in [16, 17]. For groups with infinite center, it seems in the full generality of Theorem D that one can only conclude that $\lambda_{\text{top},a,\mu,\mathcal{A}} > 0$ for some (necessarily infinite order) element $a \in \mathcal{Z}A$.

We also remark that in the case that Γ is cocompact in G , the proof of Theorem D is mostly contained in [16]. Indeed assuming that G has finite center and that G is simple or that every simple factor of G is of higher rank, the proof of Theorem D follows immediately from arguments in [16]. In the case that Γ is an irreducible lattice in a higher-rank semisimple Lie group G with rank-1 factors or when G has infinite center, one needs minor modifications to the arguments of [16] to establish Theorem D. See Remark 6.4 below for a detailed discussion of the required modifications.

Finally, we remark that for general C^1 actions on non-compact spaces, the definition and existence of Lyapunov exponents becomes complicated without additional integrability hypotheses on the norm-growth of the cocycle. In the statement of Theorem D, the choice of norms on the bundle $(G \times TM)/\Gamma$ are as constructed in Section 5.4 below. These norms are well adapted to the geometry of Γ in G and—combined with fact that the measure μ in the conclusion of the statement of Theorem D projects to Haar measure on G/Γ —ensures the cocycle \mathcal{A} satisfies a log-integrability criterion. See Claim 6.1 below.

The remainder of this paper is devoted to establishing Theorem D. See Section 2.4 for an outline of the proof. As discussed above, for actions by cocompact lattices, Theorem D is essentially contained in [16]; in this case, one first produces an A -invariant Borel probability measure with positive exponents by a fairly soft argument in [16, Proposition 4.6]. Improving the measure to one that projects to Haar is more difficult and occupies much of [16]. In the context of actions of non-uniform lattices, the arguments that construct an A -invariant measure with positive exponents is as difficult as finding one that projects to Haar. As the arguments in this paper use in an essential way all arguments from [16] and many of those from [17], the reader may find it easier to read those papers first. An expository account of some of the arguments from [16] with more detailed background may be found in the lecture notes by Brown [14].

Theorem C follows immediately from Theorem D following the same as the arguments as in [16]. We briefly recall the argument.

Proof of Theorem C. By Theorem D, if $\alpha: \Gamma \rightarrow \text{Diff}^{1+\beta}(M)$ fails to have uniform subexponential growth of derivatives then there exists a split Cartan subgroup $A \subset G$ and a $(\mathcal{Z}A)$ -invariant, Borel probability measure μ on the suspension space $X = (G \times M)/\Gamma$ such that μ projects to the Haar measure on G/Γ and the average top Lyapunov exponent $\lambda_{\text{top},a,\mu,\mathcal{A}}$ is positive for some $a \in \mathcal{Z}A$. Arguing exactly as in [16, Section 5.5], using that the action is $C^{1+\text{H\"older}}$, [18, Proposition 5.1] and the fact that $\dim(M)$ is sufficiently small implies the following.

Proposition 2.1 ([18, Proposition 5.1] and [16, Proposition 3.7]). *Suppose either that*

- (1) $\dim(M) < r(G)$ or
- (2) $\dim(M) \leq r(G)$ and $\alpha(\Gamma)$ preserves an absolutely continuous Borel probability measure μ_0 on M .

Then, any A -invariant Borel probability measure on X projecting to the Haar measure on G/Γ is G -invariant.

We also have the following well-known corollary of Zimmer’s cocycle superrigidity theorem [27]. In the statement, $n(G)$ denotes the smallest dimension of a nontrivial linear representation of (the Lie algebra of) G .

Lemma 2.2. *Let μ be any G -invariant measure for the induced action on X projecting to the Haar measure on G/Γ . If $\dim(M) < n(G)$ then $\lambda_{\text{top},g,\mu,A} = 0$ for every $g \in G$.*

To derive the lemma, we remark that X is a fiber-bundle with fibers diffeomorphic to M . Then the fiberwise derivative cocycle is measurably a $\dim(M)$ -dimensional linear cocycle; Zimmer’s cocycle superrigidity theorem and dimension constraints force the cocycle to be cohomologous to a compact-group-valued cocycle and the conclusion follows.

In particular, assuming Theorem C is false, Theorem D produces a $(\mathcal{Z}A)$ -invariant Borel probability measure with non-zero Lyapunov exponents. By Proposition 2.1 this measure is G -invariant, contradicting Lemma 2.2. \square

2.3. Review of the proof of [17, Theorem B]. To motivate the outline in the next subsection, we recall the proof of [17, Theorem B]—the analogue of Theorem C for actions of $\Gamma = \text{SL}(n, \mathbb{Z})$ for $n \geq 4$. We let $\Gamma_{i,j}$ be the subgroup of Γ generated by the elementary matrices $E_{i,j}$ and $E_{j,i}$. Then $\Gamma_{i,j}$ is isomorphic to $\text{SL}(2, \mathbb{Z})$ and differs from the identity matrix only in the (i, i) , (i, j) , (j, i) , and (j, j) entries. Write $H_{i,j} \subset \text{SL}(n, \mathbb{R})$ for the corresponding copy of $\text{SL}(2, \mathbb{R})$. Let $A \subset G$ be the group of diagonal matrices and $A_{i,j} = \{a_{i,j}^t\}$ for the 1-parameter group of diagonal matrices in $H_{i,j}$.

Quasi-isometric bounded generation. As shown in [44], the group $\text{SL}(n, \mathbb{Z})$ is **quasi-isometrically boundedly generated** by the subgroups $\{\Gamma_{i,j} \mid 1 \leq i, j \leq n\}$. Thus, to show the action $\alpha: \Gamma \rightarrow \text{Diff}(M)$ has uniform subexponential growth of derivatives it suffices to show the restriction $\alpha|_{\Gamma_{i,j}}: \Gamma_{i,j} \rightarrow \text{Diff}(M)$ of α to each $\Gamma_{i,j}$ has uniform subexponential growth of derivatives.

Uniform subexponential growth of derivatives for unipotent subgroups. The first key proposition established in [17] shows that unipotent elements of $\Gamma = \text{SL}(n, \mathbb{Z})$ act with subexponential growth: for every unipotent $\gamma \in \Gamma$ and $\varepsilon > 0$ there is a $C > 0$ such that for all $n \in \mathbb{Z}$,

$$\sup_{x \in M} \|D_x \alpha(\gamma^n)\| \leq C e^{\varepsilon \text{len}_\Gamma(\gamma^n)}.$$

We recall that $\text{len}_\Gamma(\gamma^n)$ denotes the word-length of γ^n measured in Γ which, in particular, grows logarithmically in n .

Uniform subexponential growth of derivatives for action of $\Gamma_{i,j}$. We now explain why the action of $\Gamma_{i,j}$ has uniform subexponential growth of derivatives

- (1) We pass to the suspension space $X = (G \times M)/\Gamma$. If $\alpha|_{\Gamma_{i,j}} : \Gamma_{i,j} \rightarrow \text{Diff}(M)$ fails to have uniform subexponential growth of derivatives, we may find a sequence of finite $\{a_{i,j}^t\}$ -orbits $\{a_{i,j}^t \cdot x_n \mid 0 \leq t \leq t_n\}$ with $t \rightarrow \infty$ and each x_n contained in the “thick part” of X over $H_{i,j}/\Gamma_{i,j}$ such that, in the limit, we see positive exponential growth of the fiberwise derivative.
- (2) Using that the “cusp group” of the rank-1 subgroup $\Lambda_{i,j} \simeq \text{SL}(2, \mathbb{Z})$ of $\text{SL}(n, \mathbb{Z})$ is generated by a single unipotent element we show that, restricted to the subbundle of X over $H_{i,j}/\Gamma_{i,j}$ any collection of finite $\{a_{i,j}^t\}$ -orbits $\{a_{i,j}^t \cdot x_n \mid 0 \leq t \leq t_n\}$ with $t \rightarrow \infty$ that approximates the maximal exponential growth rate of the fiberwise derivatives gives a family of empirical measures μ_n that is primarily concentrated over the “thick part” of $H_{i,j}/\Gamma_{i,j}$.
- (3) We may average the family of measures μ_n over Følner sets in a unipotent subgroup N normalized by $H_{i,j}$ that meets Γ in a lattice to obtain a new family of measures $\tilde{\mu}_n$. Using quantitative non-divergence of unipotent flows, or an explicit computation as in [17, Subsection 5.3], we control the amount of mass of each $\tilde{\mu}_n$ at ∞ ; in particular, we rule out escape of mass. Also, using that the $N \cap \Gamma$ contains only unipotent elements, it follows that these measures see no “escape of Lyapunov exponent.”
- (4) There is a one-parameter family of diagonal matrices $\{b^s\}$ such that N is the horospherical subgroup of $\{b^s\}$ for $s > 0$. Using exponential mixing of b^s , or an explicit computation as in [17, Subsection 5.4], there is a sequence of $s_n \rightarrow \infty$ such that the family of measures $\{b^{s_n} \cdot \tilde{\mu}_n\}$ is uniformly tight and in fact have uniformly exponentially small mass at ∞ (see Definition 4.1) of G/Γ ; moreover, we are able to maintain positive exponential growth of fiberwise derivative for the action of $a_{i,j}^t$.
- (5) It follows that any limit measure of $\{b^{s_n} \cdot \tilde{\mu}_n\}$ projects to the Haar measure on G/Γ and has a non-zero fiberwise Lyapunov exponent for the action of $a_{i,j}^t$. We can then obtain an A -invariant measure on X that projects to the Haar measure on G/Γ and has a non-zero fiberwise Lyapunov exponent. We then obtain a contradiction with Zimmer’s cocycle superrigidity as in the proof of Theorem C.

2.4. Outline of proof of Theorem D and discussion of new tools. We now outline the proof of Theorem D and highlight the new ideas compared with the proof of [17, Theorem B]. For lattices of higher \mathbb{Q} -rank, the proof of Theorem D follows the same coarse outline from [17]. In the case of \mathbb{Q} -rank-1, there are new complications that are not addressed in [17].

For simplicity of this outline, we may assume G is linear (in particular has finite center) and defined over \mathbb{Q} . Moreover, as [16] essentially handles the case that Γ is cocompact, we may assume Γ is nonuniform. By Margulis’s arithmeticity theorem (see discussion in Section 3.7) we may assume Γ is commensurable with $G_{\mathbb{Z}}$.

2.4.1. Quasi-isometric bounded generation. In [17], we used the fact from Lubotzky, Mozes and Raghunathan [44] that $\text{SL}(n, \mathbb{Z})$ is **quasi-isometrically boundedly generated** by the subgroups $\{\Gamma_{i,j} \mid 1 \leq i, j \leq n\}$. In the setting of this paper, we need to establish an analogous result for general arithmetic groups Γ . In [45],

where Lubotzky, Mozes and Raghunathan prove the general case of the results in [44], a different outline is followed and the analogous result is not proven. It has been proven very recently by Witte Morris. We replace the canonical copies of $\mathrm{SL}(2, \mathbb{Z}) \simeq \Gamma_{i,j}$ from [17] with lattice subgroups in **standard \mathbb{Q} -rank-1 subgroups**; see Definition 3.13. We then have the following key definition and theorem provided to us by Dave Witte Morris [49]:

Definition 3.28. *Let $G = \mathbf{G}(\mathbb{R})$ be a semisimple algebraic \mathbb{Q} -group and let Γ be a subgroup commensurable with $G_{\mathbb{Z}} = \mathbf{G}(\mathbb{Z})$. We say that Γ is **quasi-isometrically boundedly generated by standard \mathbb{Q} -rank-1 subgroups** if there is a constant $r = r(\mathbf{G}, \Gamma) \in \mathbb{N}$, a finite subset $\Gamma_0 = \Gamma_0(\mathbf{G}, \Gamma)$ of Γ , and a finite collection $\mathcal{L} = \mathcal{L}(\mathbf{G}, \Gamma)$ of \mathbb{Q} -subgroups of F such that*

- (1) *each $L \in \mathcal{L}$ is a standard \mathbb{Q} -rank-1 subgroup of G ;*
- (2) *every element γ of Γ can be written in the form $\gamma = s_1 s_2 \cdots s_r$ where either*
 - (a) *$s_i \in \Gamma_0$ or*
 - (b) *$s_i \in \Gamma_L$ for some $L \in \mathcal{L}$ and $\log \|s_i\| \leq r \log \|\gamma\|$.*

Above and in following, we write $\Gamma_L = L \cap \Gamma$. The following is the main result of [49].

Theorem 3.31. *Every arithmetic subgroup Γ of a \mathbb{Q} -isotropic, almost \mathbb{Q} -simple \mathbb{Q} -group is quasi-isometrically boundedly generated by standard \mathbb{Q} -rank-1 subgroups.*

2.4.2. *Uniform subexponential growth of derivatives for unipotent subgroups.* As in [17], we again show that unipotent subgroups $\Delta \subset \Gamma$ have subexponential growth. This is done in Section 8. While the outline of the argument is similar, this requires substantial modifications of [17, Section 4].

Proposition 8.1. *If there is a connected unipotent \mathbb{Q} -subgroup U of G such that the restriction*

$$\alpha|_{\Gamma_U} : \Gamma_U \rightarrow \mathrm{Diff}^1(M)$$

does not have uniform subexponential growth of derivatives then there is a split Cartan subgroup A of G and an A -invariant Borel probability measure μ on X projecting to the Haar measure on G/Γ such that $\lambda_{\mathrm{top}, a, \mu, A} > 0$ for some $a \in A$.

To establish Proposition 8.1, we work with the induced G -action. It is easy to see that subexponential growth of derivatives for Γ_U is equivalent to subexponential growth of derivatives for a closed U orbit in $X := (G \times M)/\Gamma$. The proof of the proposition proceeds by working inside compact orbits of a solvable group of the form $\tilde{A} \times U$ where U is a horospherical \mathbb{Q} -group and \tilde{A} is part of a \mathbb{Q} -anisotropic torus. We first show that generic trajectories for elements of \tilde{A} have subexponential growth of derivatives. Combining this with the fact that U is normalized by \tilde{A} , we obtain subexponential growth of derivatives for a large set of elements in U . Using that U is nilpotent, a sunset argument implies subexponential growth of derivatives for every element of U . To establish subexponential growth of derivatives for generic orbits of elements of \tilde{A} , we exploit that U is horospherical and use an exponential mixing argument to upgrade an \tilde{A} -invariant measure supported on some closed $\tilde{A} \times U$ orbit to an A -invariant measure projecting to the Haar measure on G/Γ .

2.4.3. *Uniform subexponential growth of derivatives for “cusp groups” of standard \mathbb{Q} -rank-1 subgroups.* In [17], we heavily used that the fundamental group of the cusp

in $\mathrm{SL}(2, \mathbb{R})/\mathrm{SL}(2, \mathbb{Z})$ is generated by the single unipotent element $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. For a standard \mathbb{Q} -rank-1 subgroup H , we cannot expect the fundamental group of a cusp in H/Γ_H to be generated by a single unipotent element; moreover, the fundamental group of a cusp need not be a unipotent subgroup. However, the structure of such groups (the semidirect product of a cocompact lattice and a unipotent subgroup) allows us to still show the restriction of the action to the fundamental group of a cusp in H/Γ_H for a standard \mathbb{Q} -rank-1 subgroup H have subexponential growth.

Given a standard \mathbb{Q} -rank-1 subgroup H , to establish that the restriction of the action to the fundamental group of a cusp in H/Γ_H has subexponential growth, we exploit reduction theory and the construction of Siegel fundamental domains to relate growth properties of the fundamental group of a cusp in H/Γ_H with subexponential growth for minimal parabolic \mathbb{Q} -subgroups in G . We point the reader to Lemma 5.3 and especially Lemma 5.14 in Section 5.5.

To complete this step, it remains to establish subexponential growth for minimal parabolic \mathbb{Q} -subgroups in G which occupies Section 9.

Proposition 9.1. *Let P be a minimal parabolic \mathbb{Q} -subgroup of G . If the restriction*

$$\alpha|_{\Gamma_P} : \Gamma_P \rightarrow \mathrm{Diff}^1(M)$$

does not have uniform subexponential growth of derivatives there is a split Cartan subgroup A of G and an A -invariant Borel probability measure μ on X projecting to the Haar measure on G/Γ such that $\lambda_{\mathrm{top}, a, \mu, A} > 0$ for some $a \in A$.

The proof of Proposition 9.1 uses that Γ_P is the semidirect product of a cocompact lattice in a reductive (\mathbb{Q} -anisotropic) \mathbb{Q} -subgroup L and a (cocompact) lattice in a unipotent \mathbb{Q} -subgroup U . Subexponential growth for the restriction to Γ_U follows from Proposition 8.1. Following arguments from [16], we show that if Γ_L fails to have subexponential growth of the derivatives, then there exists a Borel probability measure on X which is invariant under a torus in L and has a positive Lyapunov exponent. Using Proposition 8.1, we can find a measure that is also invariant under U and also has a positive Lyapunov exponent. We then use exponential mixing to find a measure that projects to the Haar measure on G/Γ with a positive Lyapunov exponent which can be made A -invariant by additional averaging.

2.4.4. *Uniform subexponential growth of derivatives for \mathbb{Q} -rank-1 subgroups.* By Theorem 3.31, to show Theorem D it is enough to show the restriction of α to $\Gamma_H \subset \Gamma$ has uniform subexponential growth of derivatives for any standard \mathbb{Q} -rank-1 subgroup $H \subset G$. We recall that $H \subset G$ is a subgroup defined over \mathbb{Q} and that $\Gamma_H = H \cap \Gamma$ is a lattice in H .

Having established Propositions 8.1 and 9.1 and Lemma 5.14, Section 10 is devoted to the following.

Proposition 10.1. *If there is a standard \mathbb{Q} -rank-1 subgroup H of G such that the restriction*

$$\alpha|_{\Gamma_H} : \Gamma_H \rightarrow \mathrm{Diff}^1(M)$$

does not have uniform subexponential growth of derivatives then there is a split Cartan subgroup A of G and an A -invariant Borel probability measure μ on X projecting to the Haar measure on G/Γ such that $\lambda_{\mathrm{top}, a, \mu, A} > 0$ for some $a \in A$.

The proof of Proposition 10.1 contains the main new ingredients of the paper which have no analogue in [17]; in some sense, the proof of Proposition 10.1 is closer to the arguments of [16]. This is especially true in the case that Γ is \mathbb{Q} -rank 1. When Γ has \mathbb{Q} -rank 1, we have that $G = H$ is the only standard \mathbb{Q} -rank-1 subgroup; in particular, we cannot find a subgroup of G of the form $H \times N$ where H is a standard \mathbb{Q} -rank-1 subgroup and N is a (non-trivial) horospherical subgroup of G . Having a horospherical subgroup N normalized by H was a key ingredient used to establish subexponential growth for the restriction to the canonical copies of $\mathrm{SL}(2, \mathbb{Z})$ in $\Gamma = \mathrm{SL}(m, \mathbb{Z})$ in [17]. When Γ has \mathbb{Q} -rank at least 2, we outline a proof of Proposition 10.1 that is closer to the argument from [17] in Section 11.

Section 10 provides a uniform argument when Γ is \mathbb{Q} -rank-1 or has higher- \mathbb{Q} -rank. Fix a \mathbb{Q} -rank-1 subgroup $H \subset G$. When Γ has higher \mathbb{Q} -rank we also find a horospherical subgroup N normalized by H ; if Γ has \mathbb{Q} -rank 1, we take $N = \{\mathbf{1}\}$. We study growth of the fiberwise derivative cocycle for the induced H -action on the bundle $X_{HN} := (HN \times M)/\Gamma_{HN}$. We fix a 1-parameter (\mathbb{R} -split) subgroup $\{a^t\}$ in H and consider orbit segments of that start and end in some fixed “thick” compact part of X_{HN} . Using Proposition 9.1 and Proposition 8.1, a sequence of empirical measures supported on such orbits and limiting to the maximal growth rate of the fiberwise derivative cocycle (for this 1-parameter subgroup) will, in fact, produce a uniformly tight sequence of measures (see Lemma 10.5). Passing to a limit, we obtain an a^t -invariant Borel probability measure μ on X_{HN} . While we are able to avoid escape of mass, we are not able to rule out “escape of Lyapunov exponent.” Specifically, the fiberwise derivative cocycle is not bounded and there is no reason the fiberwise derivative cocycle is $\log-L^1$ for the limiting measures μ . Furthermore, even if fiberwise derivative cocycle happens to be $\log-L^1$ for the limiting measures μ and the empirical measures see exponential growth of the cocycle, we need not have any semi-continuity properties of the top exponent.

To remedy this problem, we use a combination of cut-off functions and time averages to modify the fiberwise derivative cocycle in the cusp; see Section 10.3.1 and (10.5) in Section 10.3.2. These new cut-off cocycles will be bounded but will fail to be continuous; however, for the cut-off cocycle, the set of discontinuities is rather tame and, for the analysis in Section 10, these cocycles acts much like a bounded continuous cocycle. Using these modified cocycles, we define analogues of Lyapunov exponents; we then show these “fake Lyapunov exponents” behave well under various averaging operations (using that the discontinuity set of the cut-off cocycle has zero measure for all limiting measures considered). We can then use averaging techniques (following either [16] or [17]) to upgrade to a measure whose projection to G/Γ is the Haar measure on G/Γ while maintaining a positive “fake Lyapunov exponent.” Since the fiberwise derivative cocycle is $\log-L^1$ for measures projecting to the Haar measure (see (5.2) and Claim 6.1), we check the “fake top Lyapunov exponent” coincides with the actual top Lyapunov exponent of the original cocycle.

Remark 2.3 (Benefits of working with \mathbb{Q} -rank-1 subgroups). We remark that the reduction to \mathbb{Q} -rank-1 subgroups Γ_H is primarily to control the behavior of the cocycle outside of compact sets. If $\Gamma \subset G$ is an arithmetic lattice with $\mathrm{rank}_{\mathbb{Q}}(\Gamma) = 1$, then there is some $\ell > 0$ such that for any 1-parameter subgroup $\{a^t\}$ of G , if $d(a^t \cdot x, \mathbf{1}\Gamma) \geq \ell$ for all $0 \leq t \leq t_0$ then, lifting the path $\{a^t \cdot x \mid 0 \leq t \leq t_0\}$ in G/Γ to the path $\{a^t \cdot \tilde{g} \mid 0 \leq t \leq t_0\}$ in G , for any element $\gamma_x \in \Gamma$ such that g

and $a^t g \gamma^{-1}$ are in the same Siegel set, γ_x is in the fundamental group of the cusp of H/Γ_H .

This fails for lattices of higher \mathbb{Q} -rank. Indeed, for higher-rank groups, there is no collection of disjoint cusps. In rank one the complement of the thick part is a disjoint union of cusps, each of the form $\mathbb{R}^+ \times N$ where N is a compact manifold. In higher rank, the complement of the thick part is a union of thickened fans, where each fan is of the form $S \times N$ where S is a Weyl chamber and N is a compact manifold. These fans can intersect non-trivially in lower dimensional sets and a path may leave the thick part of G/Γ through one fan and return through another fan which intersects the first in a lower-dimensional object. This is a consequence of the fact that in this case the rational Tits building is connected. In this case, the collection of possible monodromy elements γ_x of a path $\{a^t \cdot x \mid 0 \leq t \leq t_0\}$ does not have any structure we can exploit.

3. ALGEBRAIC GROUPS AND REDUCTIONS FOR NONUNIFORM Γ

We present the basic terminology and facts from algebraic groups that will be used in the sequel. We end the section with the main reductions we use when working with nonuniform lattices Γ and the consequences of these reductions.

3.1. Basic terminology. We denote by $\mathbf{GL}(d) = \mathrm{GL}(d, \mathbb{C})$ the affine algebraic group of $d \times d$ invertible matrices with complex entries. A **(linear) algebraic group** is a Zariski closed subgroup \mathbf{H} of $\mathbf{GL}(d)$ for some d ; that is \mathbf{H} is a subgroup that coincides with the common zeros of a set of polynomial functions in the matrix entries of $A \in \mathbf{GL}(d)$. Working over \mathbb{C} , a linear algebraic group \mathbf{H} is connected in the Zariski topology if and only if it is connected in the analytic topology.

Let $\mathbb{K} \in \{\mathbb{R}, \mathbb{Q}\}$. An algebraic subgroup $\mathbf{H} \subset \mathbf{GL}(d)$ is said to be **defined over \mathbb{K}** or is said to be **\mathbb{K} -group** if \mathbf{H} is algebraic and the polynomial functions defining \mathbf{H} take coefficients only in \mathbb{K} . The **radical** (resp. **unipotent radical**) of a linear algebraic subgroup \mathbf{H} defined over \mathbb{K} is the maximal connected solvable (resp. unipotent) algebraic subgroup $\mathrm{Rad}(\mathbf{H})$ (resp. $\mathrm{Rad}_u(\mathbf{H})$) of \mathbf{H} . These are \mathbb{K} -subgroups of \mathbf{H} . The algebraic group \mathbf{H} is **semisimple** (resp. **reductive**) if the radical (resp. unipotent radical) vanishes. We say that \mathbf{H} is **almost \mathbb{K} -simple** if it has no connected, proper, normal \mathbb{K} -subgroups. If \mathbf{H} is noncommutative, almost \mathbb{K} -simple, and connected, then \mathbf{H} is semisimple.

Given a linear algebraic subgroup $\mathbf{H} \subset \mathbf{GL}(d)$ defined over \mathbb{K} , let $\mathbf{H}(\mathbb{K}) = \mathbf{H} \cap \mathrm{GL}(d, \mathbb{K})$ denote the \mathbb{K} -points of \mathbf{H} . If $\mathbb{K} = \mathbb{R}$, the group $H = \mathbf{H}(\mathbb{R})$ is a Lie group which, in general, might not be connected even if \mathbf{H} is connected but has only finitely many connected components (in the analytic topology); we write H° for the connected component (in the analytic topology) of the identity of $\mathbf{H}(\mathbb{R})$. When \mathbf{H} is simply connected (as an algebraic group) or if \mathbf{H} is a unipotent \mathbb{K} -group, then $\mathbf{H}(\mathbb{R})$ is connected in the analytic topology. See [12, Chapter 1, Theorem 2.3.1(c)] or [56, Proposition 7.6].

Given a connected algebraic group \mathbf{H} defined over \mathbb{K} , we will at times abuse terminology and also say that $H = \mathbf{H}(\mathbb{R})$ and H° are defined over \mathbb{K} . We have that $\mathrm{Rad}(\mathbf{H})(\mathbb{R})$ (resp. $\mathrm{Rad}_u(\mathbf{H})(\mathbb{R})$) is the radical (resp. unipotent radical) of the Lie group H . If \mathbf{H} is semisimple (resp. reductive) then $H = \mathbf{H}(\mathbb{R})$ is semisimple (resp. reductive) as a Lie group

We collect the following important facts about unipotent subgroups.

Lemma 3.1. *Let U be a connected unipotent subgroup of $\mathrm{SL}(d, \mathbb{R})$.*

- (1) *U is simply connected. In fact, the exponential map from the Lie algebra \mathfrak{u} of U to U is a bijective polynomial map and its inverse is also a polynomial map [30, Theorem 8.1.1, p. 107]. This implies that every continuous homomorphism from U to any unipotent group is a polynomial.*
- (2) *The abelianization $U/[U, U]$ of U is simply connected (and in fact, it is a unipotent linear algebraic group).*

3.2. Algebraic tori and \mathbb{K} -rank. Let $\mathbf{H} \subset \mathbf{GL}(d)$ be a connected linear algebraic group defined over $\mathbb{K} \in \{\mathbb{R}, \mathbb{Q}\}$.

- (1) We say that \mathbf{H} is a **\mathbb{K} -torus** if it is commutative, defined over \mathbb{K} , and conjugate in $\mathbf{GL}(d)$ to a subgroup of diagonal matrices
- (2) A \mathbb{K} -torus \mathbf{H} is **\mathbb{K} -split** if there exists $g \in \mathrm{SL}(d, \mathbb{K})$, such that $g\mathbf{H}g^{-1}$ is a subgroup of diagonal matrices. If \mathbf{H} is a \mathbb{K} -split torus then $\mathbf{H}(\mathbb{R})$ is isomorphic to $(\mathbb{R}^*)^\ell$ for some ℓ .
- (3) Let \mathbf{S} be a torus defined over \mathbb{K} . There exists unique maximal \mathbb{K} -tori $\mathbf{S}', \mathbf{S}'' \subset \mathbf{S}$ such that $\mathbf{S} = \mathbf{S}' \cdot \mathbf{S}''$ where \mathbf{S}' is \mathbb{K} -split and \mathbf{S}'' contains no \mathbb{K} -split subtorus. (See [66, Proposition 13.2.4].)
- (4) If \mathbf{H} is defined over \mathbb{K} , then all maximal \mathbb{K} -split tori of \mathbf{H} are conjugate by elements of $\mathrm{SL}(d, \mathbb{K})$, and therefore have the same dimension. (See [47, (0.25)] or [66, Theorem 15.2.6].) This common dimension is the **\mathbb{K} -rank** of \mathbf{H} , denoted by $\mathrm{rank}_{\mathbb{K}} \mathbf{H}$.
- (5) We say that \mathbf{H} is **\mathbb{K} -anisotropic** if $\mathrm{rank}_{\mathbb{K}} \mathbf{H} = 0$; otherwise, \mathbf{H} is **\mathbb{K} -isotropic**. An \mathbb{R} -group \mathbf{H} is an \mathbb{R} -anisotropic if and only if $\mathbf{H}(\mathbb{R})$ is compact.
- (6) If \mathbf{H} is reductive then it is the almost-direct product of a semisimple \mathbb{K} -group and a \mathbb{K} -torus: $\mathbf{H} = \mathbf{L}\mathbf{S}$ where \mathbf{L} is semisimple, \mathbf{S} is a torus that centralizes \mathbf{L} , and $\mathbf{L} \cap \mathbf{S}$ is finite. The subgroups \mathbf{L} and \mathbf{S} are unique.
- (7) If $H = \mathbf{H}(\mathbb{R})$ and if \mathbf{S} is a maximal \mathbb{K} -torus (resp. maximal \mathbb{K} -split torus) we call $S = \mathbf{S}(\mathbb{R})$ a maximal \mathbb{K} -torus (resp. maximal \mathbb{K} -split torus) of H .
- (8) From (3), every one-parameter subgroup $\{a^t\}$ of a \mathbb{K} -torus S has a **real Jordan decomposition** $a^t = a_{\mathrm{split}}^t a_{\mathrm{anis}}^t$, where a_{split}^t is \mathbb{R} -diagonalizable, a_{anis}^t is contained in a compact subgroup, and a_{split}^t and a_{anis}^t are contained in S .
- (9) If $\mathbb{K} = \mathbb{R}$ and if \mathbf{S} is a maximal \mathbb{R} -split torus in \mathbf{H} , we call the connected component of the identity $A = \mathbf{S}(\mathbb{R})^\circ$ a **split Cartan subgroup** of $H = \mathbf{H}(\mathbb{R})$.

3.3. Levi and Langlands decompositions. For $\mathbb{K} \in \{\mathbb{R}, \mathbb{Q}\}$, let \mathbf{H} be a connected linear algebraic group over \mathbb{K} .

- (1) \mathbf{H} can be written as a semidirect product $\mathbf{H} = \mathbf{F} \ltimes \mathbf{U}$ where $\mathbf{U} = \mathrm{Rad}_u(\mathbf{H})$ and \mathbf{F} is a connected reductive \mathbb{K} -group. This is a **Levi decomposition** and \mathbf{F} is a **Levi subgroup**. All Levi subgroups of \mathbf{H} are conjugate by elements of $\mathbf{U}(\mathbb{K})$. See [51, pp. 200–201]
- (2) We may further decompose a Levi \mathbb{K} -subgroup as $\mathbf{F} = \mathbf{L} \times \mathbf{S}$ where \mathbf{L} is a connected reductive \mathbb{K} -subgroup with \mathbb{K} -anisotropic center and \mathbf{S} is a connected \mathbb{K} -split torus.

Write $H = \mathbf{H}(\mathbb{R})$ and let

$$L = \mathbf{L}(\mathbb{R}), \quad S = \mathbf{S}(\mathbb{R}), \quad A = S^\circ, \quad \text{and} \quad U = \mathrm{Rad}_u(\mathbf{H})(\mathbb{R}).$$

Then $H = F \ltimes U$ and $H = (L \times A) \ltimes U$ are called, respectively, a **Levi decomposition** and a **Langlands decomposition** of H . Note that U and A are connected and we also call $H^\circ = (L^\circ \times A) \ltimes U$ a Langlands decomposition of H° .

- (3) If \mathbf{H} is defined over \mathbb{Q} then all Levi subgroups F of $H = \mathbf{H}(\mathbb{R})$ are conjugate by U regardless if we view \mathbf{H} as a \mathbb{Q} -group or as a \mathbb{R} -group.

However, a Langlands decomposition of H as a \mathbb{Q} -group need not be conjugate to any Langlands decomposition of H as an \mathbb{R} -group. We refer to a choice of Langlands decomposition of H as a \mathbb{Q} -group as a **rational Langlands decomposition**. We remark that in our definition of a rational Langlands decomposition $H = (L \times A) \ltimes U$, we will always take the subgroups L and $A = S^\circ$ to be defined over \mathbb{Q} ; this is different from the terminology used for instance in [10] which only requires that L and A be conjugate to groups defined over \mathbb{Q} .

Given a connected linear algebraic \mathbb{Q} -group \mathbf{H} , let $H = \mathbf{H}(\mathbb{R})$ and write $\mathbf{H}(\mathbb{Z}) = H_{\mathbb{Z}} := H \cap \mathrm{SL}(d, \mathbb{Z})$.

Proposition 3.2. *Let \mathbf{H} be a connected linear algebraic \mathbb{Q} -group and let $H = \mathbf{H}(\mathbb{R})$.*

- (1) *$H_{\mathbb{Z}}$ is a cocompact lattice in H if and only if some (and hence every) Levi subgroup \mathbf{F} of \mathbf{H} defined over \mathbb{Q} is \mathbb{Q} -anisotropic. (See [56, Theorem 4.12, p. 210].)*
- (2) *$H_{\mathbb{Z}}$ is a lattice in H if and only if the subgroup $A = S^\circ$ is trivial in some (and hence every) rational Langlands decomposition $H = (L \times A) \ltimes U$. (See [56, Theorem 4.13, p. 213].)*

3.4. Restricted \mathbb{K} -roots and parabolic subgroups of algebraic groups. Let \mathbf{G} be a connected semisimple linear algebraic group defined over $\mathbb{K} \in \{\mathbb{R}, \mathbb{Q}\}$ and let $G = \mathbf{G}(\mathbb{R})$.

Let \mathbf{S} be a maximal \mathbb{K} -split torus in \mathbf{G} . Let $A = \mathbf{S}(\mathbb{R})^\circ$ and let $\Phi(A, G)$ denote the set of weights for the adjoint action of A on the Lie algebra \mathfrak{g} of G . These are the **\mathbb{K} -roots** of G . Each root is a homomorphism from A to the multiplicative group \mathbb{R}^+ . The \mathbb{K} -roots $\Phi(A, G)$ extend to \mathbb{K} -characters on \mathbf{S} . The \mathbb{K} -characters form a finitely generated abelian group and, as is standard, we write the group operation on characters and roots additively. Choose an ordering of $\Phi(A, G)$ and let Δ be the corresponding set of simple roots: a positive root α is **simple** if it is not the sum of two other positive roots. Let N be the connected unipotent subgroup of G whose Lie algebra is the sum of all the root spaces corresponding to positive roots. This is a maximal unipotent \mathbb{K} -subgroup of G .

A **Borel subgroup** of \mathbf{G} is a maximal Zariski connected solvable subgroup. A **parabolic \mathbb{K} -subgroup** of \mathbf{G} is a \mathbb{K} -subgroup \mathbf{P} that contains a Borel subgroup. When $\mathbb{K} = \mathbb{Q}$, Borel subgroups of \mathbf{G} need not be defined over \mathbb{Q} . However, all minimal parabolic \mathbb{Q} -subgroups are conjugate over $\mathbf{G}(\mathbb{Q})$. (See [66, Theorem 15.4.6(ii)].) Every parabolic \mathbb{K} -subgroup \mathbf{P} is connected. If $G = \mathbf{G}(\mathbb{R})$, we also refer to $P = \mathbf{P}(\mathbb{R})$ as a parabolic \mathbb{K} -subgroup of G .

We have the following characterization of parabolic \mathbb{K} -subgroups of G : Fix a subset Δ_0 of the set Δ of simple \mathbb{K} -roots of $\Phi(A, G)$ and let

$$S = \bigcap_{\alpha \in \Delta_0} \ker \alpha \subseteq A.$$

Then S is the identity component of a \mathbb{K} -split \mathbb{K} -torus and the group

$$P_{\Delta_0} := C_G(S)N \tag{3.1}$$

is a parabolic \mathbb{K} -subgroup of G . Moreover, all parabolic \mathbb{K} -subgroups of G arise as in (3.1) for some a choice of maximal \mathbb{K} -split torus, ordering of the roots, and set of simple roots.

It is clear from the definition of S that $\dim S = \text{rank}_{\mathbb{K}} G - |\Delta_0|$.

Remark 3.3. We have $A \subseteq C_G(S) \subseteq P_{\Delta_0}$. Since all maximal connected \mathbb{K} -split tori of P_{Δ_0} are conjugate via $(P_{\Delta_0})_{\mathbb{K}}$, this implies that every maximal connected \mathbb{K} -split torus of P_{Δ_0} is a maximal connected \mathbb{K} -split torus of G .

A minimal parabolic \mathbb{K} -subgroup is a parabolic \mathbb{K} -subgroup that does not contain any other parabolic \mathbb{K} -subgroup. These occur by taking $\Delta_0 = \emptyset$ above and, as previously remarked, all minimal parabolic \mathbb{K} -subgroups of G are conjugate via $G_{\mathbb{K}} = \mathbf{G}(\mathbb{K})$.

The following well-known observation is immediate from the definitions.

Lemma 3.4. *Let P be a minimal parabolic \mathbb{K} -subgroup of G , and let $P = (L \times S^\circ) \ltimes U$ be a Langlands decomposition (as a \mathbb{K} -group). Then:*

- (1) S° is the identity component of a maximal \mathbb{K} -split torus in G and
- (2) L is \mathbb{K} -anisotropic.

At the other extreme, a parabolic \mathbb{K} -subgroup Q is said to be **maximal** if Q is a proper subgroup of G which is not contained in any other proper, parabolic \mathbb{K} -subgroup of G . In the notation of (3.1) this means that $\dim S = 1$ or, equivalently, that $|\Delta_0| = \text{rank}_{\mathbb{K}} G - 1$.

Remark 3.5. Suppose $Q = LAN$ is a Langlands decomposition (as a \mathbb{K} -group) of a parabolic \mathbb{K} -subgroup of G . It is well known that for G simply connected as an algebraic group, the semisimple subgroup $[L, L]$ is simply connected as an algebraic group [66, Exercise 8.4.6(6), p. 149].

Definition 3.6. *Let $\{b^s\}$ be a one-parameter subgroup of G . The corresponding **expanding horospherical subgroup** of $\{b^s\}$ is*

$$U^+(b^s) := \{u \in G \mid b^s u b^{-s} \rightarrow \mathbf{1} \text{ as } s \rightarrow -\infty\}.$$

This is a connected closed, unipotent subgroup of G .

The following elementary observation can be proven by taking $\{b^s\}$ to be in the interior of the positive Weyl chamber of A . As above $\mathbb{K} \in \{\mathbb{R}, \mathbb{Q}\}$.

Proposition 3.7. *Let $P = (L \times A) \ltimes U$ be the Langlands decomposition of a parabolic \mathbb{K} -subgroup of G . Then U is the expanding horospherical subgroup of some \mathbb{R} -diagonalizable one-parameter subgroup $\{b^s\}$ of A .*

In the following results, we consider a parabolic \mathbb{Q} -subgroup Q of a \mathbb{Q} -group G with rational Langlands decomposition $Q = LAU$. The group L acts by conjugation on U . The action preserves the commutator subgroup $[U, U]$ and thus induces an action of L and its subgroups on $U/[U, U]$. Identifying the abelian group $U/[U, U]$ with some \mathbb{R}^n , this action is a linear representation. The following result is probably well known (see related results such as [2, Theorem 2] and [63, 64]) but we could not find a reference.

Lemma 3.8. *Let \mathbf{G} be an almost \mathbb{Q} -simple \mathbb{Q} -group with $\text{rank}_{\mathbb{R}}(\mathbf{G}) \geq 2$ and $\text{rank}_{\mathbb{Q}}(\mathbf{G}) \geq 1$. Let Q be a maximal parabolic \mathbb{Q} -subgroup of $G = \mathbf{G}(\mathbb{R})^\circ$ with rational Langlands decomposition $Q = LAU$. Let L^\dagger be the product of all non-compact almost simple factors of L and the maximal connected \mathbb{R} -split torus in the center of L .*

Then the induced action of L^\dagger on $U/[U, U]$ does not contain the trivial representation; that is, the only fixed point for the action of L^\dagger on $U/[U, U]$ is the identity coset.

Proof. We make a number of preliminary observations. Fix the rational Langlands decomposition $Q = LAU$. Since Q is a maximal parabolic \mathbb{Q} -subgroup, we have $\dim(A) = 1$. Fix a minimal parabolic \mathbb{Q} -subgroup $P_{\mathbb{Q}} \subset Q$ and a minimal parabolic \mathbb{R} -subgroup $P_{\mathbb{R}} \subset P_{\mathbb{Q}}$. Fix maximal connected \mathbb{Q} -split torus S in $P_{\mathbb{Q}}$ and a maximal connected \mathbb{R} -split torus T in $P_{\mathbb{R}}$ with

$$A \subset S \subset T.$$

Let $\Delta_{\mathbb{Q}}$ be a collection of simple \mathbb{Q} -roots for $\Phi(S, G)$ determined by $P_{\mathbb{Q}}$ and let $\Delta_{\mathbb{R}}$ be a collection of simple \mathbb{R} -roots for $\Phi(T, G)$ determined by $P_{\mathbb{R}}$. Note that every element of $\Delta_{\mathbb{Q}}$ is the restriction of an element of $\Delta_{\mathbb{R}}$ to S .

Let \mathfrak{u} be the Lie algebra of U . The exponential map $\mathfrak{u} \rightarrow U$ gives an identification between $\mathfrak{u}/[\mathfrak{u}, \mathfrak{u}]$ and $U/[U, U]$. Since Q is a maximal parabolic \mathbb{Q} -subgroup, there is a simple \mathbb{Q} -root $\alpha \in \Delta_{\mathbb{Q}}$ such that any \mathbb{Q} -root space \mathfrak{g}^β is contained in \mathfrak{u} if and only if β is a non-negative integer combination of elements of $\Delta_{\mathbb{Q}}$ and the coefficient of α is positive. The map $\mathfrak{u} \rightarrow \mathfrak{u}/[\mathfrak{u}, \mathfrak{u}]$ is injective on the \mathbb{Q} -root space \mathfrak{g}^β if and only if the coefficient of α in β is 1; the kernel of $\mathfrak{u} \rightarrow \mathfrak{u}/[\mathfrak{u}, \mathfrak{u}]$ is spanned by all \mathbb{Q} -root spaces \mathfrak{g}^β such that the coefficient of α in β is at least 2.

We observe that $T = A \cdot (T \cap L) = A \cdot (T \cap L^\dagger)$ and thus $T \cap L^\dagger$ has codimension-1 in T . There are at most 2 (proportional) positive real root $\beta_0, 2\beta_0 \in \Phi(T, G)$ for which the group $T \cap L^\dagger$ acts trivially by conjugation on the real root spaces \mathfrak{g}^{β_0} and $\mathfrak{g}^{2\beta_0}$. Since the action on all other root spaces is by scalar multiplication, if there is $x \in \mathfrak{u}/[\mathfrak{u}, \mathfrak{u}]$ which is fixed under conjugation by $T \cap L^\dagger$, we conclude that $x \in \mathfrak{g}^{\beta_0} \pmod{[\mathfrak{u}, \mathfrak{u}]}$.

To finish the proof of the lemma, we consider separately the cases that Q is or is not maximal as a parabolic \mathbb{R} -subgroup.

Case 1. Suppose Q is a maximal parabolic \mathbb{R} -subgroup. As we assume Q is a maximal parabolic \mathbb{R} -subgroup, there is a simple \mathbb{R} -root $\alpha \in \Delta_{\mathbb{R}}$ such that a real root space \mathfrak{g}^ϕ is in \mathfrak{u} if and only if ϕ is a non-negative integer combination of elements of $\Delta_{\mathbb{R}}$ and the coefficient of α is positive. Additionally, for every simple root $\beta \in \Delta_{\mathbb{R}} \setminus \{\alpha\}$, the restriction of β to A vanishes.

Note that G may have rank-1 factors as an \mathbb{R} -group. In particular, the Dynkin diagram for $\Phi(T, G)$ may have isolated nodes. We claim that the node associated to α is not isolated in the Dynkin diagram; in particular, α is not a root associated to a rank-1 real factor. Indeed, if α were isolated then we would have $\mathfrak{u} = \mathfrak{g}^\alpha$ or $\mathfrak{u} = \mathfrak{g}^\alpha \oplus \mathfrak{g}^{2\alpha}$. Since \mathfrak{u} is the expanding horospherical subgroup for some 1-parameter subgroup, it is well-known that the subalgebra \mathfrak{h} generated by \mathfrak{g}^α and $\mathfrak{g}^{-\alpha}$ is an ideal. On the other hand, U is defined over \mathbb{Q} whence the analytic subgroup H tangent to \mathfrak{h} is the connected component of a \mathbb{Q} -subgroup. (A standard \mathbb{Q} -rank-1 subgroup in the terminology of Definition 3.13 below.) This contradicts the assumption that \mathbf{G} is almost \mathbb{Q} -simple.

Now let $x \in \mathfrak{u}/[\mathfrak{u}, \mathfrak{u}]$ be fixed by conjugation for L^\dagger . As discussed above, there is an \mathbb{R} -root β_0 such that $x \in \mathfrak{g}^{\beta_0} \pmod{[\mathfrak{u}, \mathfrak{u}]}$. It follows from the preceding paragraph that there is a simple real root $\beta \neq \alpha$ such that either $\beta_0 + \beta$ or $\beta_0 - \beta$ is a root. For every non-zero $x \in \mathfrak{g}^{\beta_0}$, this implies that either $[x, \mathfrak{g}^\beta] \neq \{0\}$ or $[x, \mathfrak{g}^{-\beta}] \neq \{0\}$ (c.f. [40, Lemma 7.75, p. 477]). Both $\mathfrak{g}^{\beta_0+\beta}$ and $\mathfrak{g}^{\beta_0-\beta}$ inject into $\mathfrak{u}/[\mathfrak{u}, \mathfrak{u}]$. Since $\mathfrak{g}^{\pm\beta}$ are elements of the lie algebra \mathfrak{l}^\dagger of L^\dagger , this implies the only element of $\mathfrak{u}/[\mathfrak{u}, \mathfrak{u}]$ fixed under conjugation by L^\dagger is 0 as claimed.

Case 2. Suppose Q is not maximal as a parabolic \mathbb{R} -subgroup. Let Z be the central torus of LA . Then Z is a \mathbb{Q} -subgroup of G and $Z = AZ_a$ where Z_a is the center of L and hence is an anisotropic \mathbb{Q} -torus. Since $\dim(A) = 1$, Z_a has codimension 1 in Z .

We work in the complexified Lie algebra $\mathfrak{g}_{\mathbb{C}}$. Let $\Delta_{\mathbb{C}}$ be the set of simple \mathbb{C} -roots, and let $\rho = \sum_{\alpha \in \Delta_{\mathbb{C}}} \alpha$ be the sum of these simple roots. After a maximal torus has been fixed, there is a natural action of the Galois group $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ on the set of all \mathbb{C} -roots since the Lie algebra splits over $\bar{\mathbb{Q}}$. The set $\Delta_{\mathbb{C}}$ is usually not invariant under this action, but all possible choices of simple roots are conjugate under the Weyl group. Therefore, composing the action of each element of the Galois group with an appropriate element of the Weyl group yields an action of $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ that leaves $\Delta_{\mathbb{C}}$ invariant. This is usually called the $*$ -action of $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ [67, §2.3, p. 39].

For any $\sigma \in \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ and any $\alpha \in \Delta_{\mathbb{C}}$, applying σ to α by the usual action of the Galois group will usually have a different result than by the $*$ -action. However, since Z is a \mathbb{Q} -torus, both results have the same restriction to Z [11, Proposition 6.7, p. 107]. Since ρ is clearly invariant under the $*$ -action, we conclude that the restriction $\rho|_Z$ is invariant under the usual action of the Galois group; in particular, ρ defines a \mathbb{Q} -character on Z . This \mathbb{Q} -character must vanish on the \mathbb{Q} -anisotropic torus Z_a . On the other hand, the restriction $\rho|_Z$ of ρ to Z is nontrivial, because the restriction of each element of $\Delta_{\mathbb{C}}$ to A is either a simple \mathbb{Q} -root or 0 and not all restrictions are 0 [11, Proposition 6.8, p. 107]. Since Z_a has codimension 1, we conclude that $Z_a = (\ker(\rho|_Z))$.

Recall T is the maximal \mathbb{R} -split torus in G . Let $r = \dim T \cap Z$ and let $\alpha_1, \dots, \alpha_r \in \Delta_{\mathbb{R}}$ be the simple real roots that are nontrivial on $T \cap Z$. From the description of parabolic subgroups in (3.1), $\dim(T \cap Z)$ is the same as the number of such simple real roots. From the preceding paragraph and the fact that the simple real roots are precisely the nonzero restrictions of elements of $\Delta_{\mathbb{C}}$ to a maximal \mathbb{R} -split torus [11, Proposition 6.8, p. 107], there are positive integers k_1, \dots, k_r such that $T \cap Z_a = \ker(\sum_{i=1}^r k_i \alpha_i|_{T \cap Z})$. Since we assume that Q is not maximal as a parabolic \mathbb{R} -subgroup, we have $r \geq 2$. Since the roots $\{\alpha_1, \dots, \alpha_r\}$ are linearly independent on $T \cap Z$, they are also linearly independent with the restriction of ρ to $T \cap Z$. In particular, the restriction of each α_i to $(T \cap Z_a)^\circ$ is nontrivial.

We claim that if a real root space \mathfrak{g}^ϕ injects into $\mathfrak{u}/[\mathfrak{u}, \mathfrak{u}]$ then there is some α_i such that $\phi|_{T \cap Z} = \alpha_i|_{T \cap Z}$. Indeed, if we enumerate $\Delta_{\mathbb{R}}$ as $\Delta_{\mathbb{R}} = \{\alpha_1, \dots, \alpha_r, \alpha_{r+1}, \dots, \alpha_\ell\}$ then the restrictions of each $\alpha_{r+1}, \dots, \alpha_\ell$ to $T \cap Z$ vanishes. A real root space \mathfrak{g}^ϕ injects into $\mathfrak{u}/[\mathfrak{u}, \mathfrak{u}]$ if and only if the root ϕ is of the form

$$\phi = \sum_{i=1}^r c_i \alpha_i + \sum_{i=r+1}^{\ell} c_i \alpha_i$$

where $c_i \geq 0$, $c_i \in \{0, 1\}$ for all $1 \leq i \leq r$, and $c_i = 1$ for exactly one $1 \leq i \leq r$. The claim then follows. By the preceding paragraph, the restriction of ϕ to $(T \cap Z_a)^\circ$

is then nontrivial. Since $(T \cap Z_a)^\circ \subseteq L^\dagger$, we conclude that the centralizer of L^\dagger in $\mathfrak{u}/[\mathfrak{u}, \mathfrak{u}]$ is trivial, as desired. \square

Corollary 3.9. *Assume that Q is a maximal parabolic \mathbb{Q} -subgroup of G with rational Langlands decomposition $Q = LS^\circ U$. Then there is a finite collection \mathcal{C} of one-parameter subgroups of L , such that*

- (1) *each $\{a^t\} \in \mathcal{C}$ is contained in a \mathbb{Q} -anisotropic \mathbb{Q} -torus \tilde{A} of L with $a^1 \in \widehat{\Gamma}_{\tilde{A}}$, and*
- (2) *$U/[U, U]$ is generated by the associated expanding horospherical subgroups*

$$\{\mathcal{U}^+(a_{\text{split}}^t) \mid \{a^t\} \in \mathcal{C}\}.$$

Proof. Let L^\dagger be as in Lemma 3.8. Since the center of L is \mathbb{Q} -anisotropic, there is a maximal connected \mathbb{R} -split torus A of L that is contained in a \mathbb{Q} -anisotropic torus \tilde{A} in L . See [52, p. 211] or [57, Theorem 2.13]. Since $L_{\mathbb{Q}}$ is Zariski dense in L [9, §13.7 and Theorem 18.2(ii), pp. 29 and 218], there are finitely many $L_{\mathbb{Q}}$ -conjugates A_1, \dots, A_k of A that generate a dense subgroup of L^\dagger . Each A_i is contained in a \mathbb{Q} -anisotropic torus \tilde{A}_i of L . Since each A_i is \mathbb{Q} -anisotropic, each $\tilde{A}_i/(\tilde{A}_i)_{\mathbb{Z}}$ and thus each $\tilde{A}_i/\widehat{\Gamma}_{\tilde{A}_i}$ is compact (see Proposition 3.2(1)) and thus identified with $\mathbb{R}^q/\mathbb{Z}^q$ for some q . Then each \tilde{A}_i is generated by finitely many one-parameter subgroups $\{a_{i,1}^t\}, \dots, \{a_{i,p}^t\}$, such that $a_{i,j}^1 \in \Gamma_{\tilde{A}}$ for all i, j . Then the \mathbb{R} -diagonalizable parts $\{(a_{i,j}^t)_{\text{split}}\}$ generate L^\dagger . Since L^\dagger is reductive, the induced representation on $U/[U, U]$ is totally reducible. From Lemma 3.8 no element of $U/[U, U]$ is centralized by every $\{(a_{i,j}^t)_{\text{split}}\}$. Thus if we assume that the collection $\{a_{i,j}^t\}$ is closed under inverses, this implies that $U/[U, U]$ is generated by the associated expanding horospherical subgroups, as desired. \square

Remark 3.10. In many cases, it is possible to take the collection \mathcal{C} in Corollary 3.9 to be a singleton. However, there are cases where this is not possible because the weight 0 occurs in the representation of L^\dagger on $U/[U, U]$. This happens for instance for one of the maximal parabolic subgroups of the \mathbb{Q} -split group $\text{Sp}(4, \mathbb{R})$ of type $C_2 \simeq B_2$.

3.5. Iwasawa decomposition, restricted roots, and parabolic subgroups.

Fix G to be a connected semisimple Lie group with Lie algebra \mathfrak{g} . We do not assume G has finite center. Choose a Cartan involution θ of \mathfrak{g} ; this induces a global Cartan involution of G which we also denote by θ . We decompose $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ where \mathfrak{k} and \mathfrak{p} are, respectively, the $+1$ and -1 eigenspaces of θ . Let \mathfrak{a} be a maximal abelian subspace of \mathfrak{p} , and let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{a} \oplus \mathfrak{n}$ be the associated Iwasawa decomposition. If K, A and N are the analytic subgroups corresponding to $\mathfrak{k}, \mathfrak{a}$, and \mathfrak{n} , respectively, then $G = KAN$ is the corresponding **Iwasawa decomposition**. Here A is a maximal connected abelian $\text{ad-}\mathbb{R}$ -diagonalizable subgroup, and N is a maximal ad- unipotent subgroup of G normalized by A . We call A a **split Cartan subgroup** of G . Furthermore, K and A are θ -invariant; more precisely, K is the set of fixed points of θ , and θ inverts every element of A . The subgroup K contains the center of G and is a maximal compact subgroup of G if (and only if) the center of G is finite. All choices of \mathfrak{a} and A above of conjugate by K . See [40, Theorem 6.31] and [40, Theorem 6.46] for details.

Write $M = C_K(A)$ for the centralizer of A in K . The subgroup $P = MAN$ is a **minimal parabolic subgroup** of G . The subgroup P and the decomposition

$P = MAN$ depends on the choice of θ , A , and N , but all minimal parabolic subgroups are conjugate in G . Any subgroup of G that contains a minimal parabolic subgroup is said to be a **parabolic subgroup**.

Write $\Phi(A, G)$ for the set of weights for the adjoint action of A on \mathfrak{g} . These are the **real restricted roots** of G (relative to the choice of A). We write the group operation on characters of A additively. We say two real roots $\beta, \widehat{\beta} \in \Phi(A, G)$ are **positively proportional** if there is some $c > 0$ with

$$\widehat{\beta} = c\beta.$$

Note that c takes values only in $\{\frac{1}{2}, 1, 2\}$ and that $c = 1$ unless the root system $\Phi(A, G)$ contains a component of type BC_ℓ . A **coarse root** is an equivalence class of roots under this equivalence relation. Write $\widehat{\Phi}(A, G)$ for the collection of coarse roots.

For $[\beta] \in \widehat{\Phi}(A, G)$, we let $\mathfrak{g}^{[\beta]} = \sum_{\alpha \in [\beta]} \mathfrak{g}^\alpha$. This is the Lie algebra of a connected unipotent subgroup $U^{[\beta]}$ of G , which is called the **coarse root group** corresponding to $[\beta]$. We remark that every parabolic subgroup that contains the minimal parabolic subgroup $P = C_G(A)N$ is saturated by coarse root groups of A .

Definition 3.11.

- (1) A **parabolic Lie subalgebra** of \mathfrak{g} is the Lie algebra of a parabolic subgroup.
- (2) The **resonant codimension**, $\bar{r}(\mathfrak{q})$, of a parabolic Lie subalgebra \mathfrak{q} is defined to be the cardinality of the set

$$\{[\beta] \in \widehat{\Phi}(A, G) \mid \mathfrak{g}^{[\beta]} \not\subset \mathfrak{q}\}.$$

- (3) The **minimal resonant codimension** of \mathfrak{g} or G , denoted by $r(\mathfrak{g})$ or $r(G)$, is defined to be the minimal value of the resonant codimension $\bar{r}(\mathfrak{q})$ of \mathfrak{q} as \mathfrak{q} varies over all (maximal) proper parabolic subalgebras of \mathfrak{g} .

Remark 3.12. Recall that $v(G)$ is the minimal dimension of a homogeneous space G/H , where H ranges over all proper, closed subgroups of G . We have $r(G) = v(G')$, where G' is any maximal connected, \mathbb{R} -split, semisimple subgroup of G by [11, Theorem. 7.2, p. 117].

3.6. Standard \mathbb{Q} -rank-1 subgroups. We return to the case that $G = \mathbf{G}(\mathbb{R})$ where \mathbf{G} is a connected semisimple linear algebraic group defined over \mathbb{Q} . Let \mathbf{S} be a maximal \mathbb{Q} -split torus in \mathbf{G} and write $A = \mathbf{S}(\mathbb{R})^\circ$. For each \mathbb{Q} -root $\alpha \in \Phi(A, G)$, such that $\frac{1}{2}\alpha \notin \Phi(A, G)$, there is a unique connected unipotent subgroup $U^{[\alpha]}$ with Lie algebra \mathfrak{g}^α or $\mathfrak{g}^\alpha \oplus \mathfrak{g}^{2\alpha}$. Since $-\alpha$ is also a \mathbb{Q} -root, we also obtain the subgroup $U^{[-\alpha]}$. The following construction is well known, but there does not seem to any established terminology for referring to this subgroup.

Definition 3.13. For \mathbb{Q} -root $\alpha \in \Phi(A, G)$ such that $\frac{1}{2}\alpha \notin \Phi(A, G)$, the subgroup H_α of G generated by $U^{[\alpha]}$ and $U^{[-\alpha]}$ is called a **standard \mathbb{Q} -rank-1 subgroup**.

If \mathbf{G} is simply connected, then there is an almost simple \mathbb{Q} -subgroup \mathbf{H} of \mathbf{G} with $\text{rank}_{\mathbb{Q}}(\mathbf{H}) = 1$ and $H_\alpha = \mathbf{H}(\mathbb{R})$; for general G , we have $H_\alpha = \mathbf{H}(\mathbb{R})^\circ$.

To justify the last remark of Definition 3.13, we view α as the restriction of $\alpha \in \Phi(\mathbf{S}, \mathbf{G})$; then $U^{[\alpha]} = \mathbf{U}^{[\alpha]}(\mathbb{R})$ where $\mathbf{U}^{[\alpha]}$ is the root subgroup in \mathbf{G} associated to α and 2α . Let \mathbf{H} be the algebraic group generated by $\mathbf{U}^{[\alpha]}$ and $\mathbf{U}^{[-\alpha]}$. Then \mathbf{H} is defined over \mathbb{Q} and has $\text{rank}_{\mathbb{Q}}(\mathbf{H}) = 1$. Choose an ordering on the roots so that α

is simple and let \mathbf{P} be the parabolic \mathbb{Q} -subgroup that contains $\mathbf{U}^{[\alpha]}$, $\mathbf{U}^{[-\alpha]}$, and no other root groups associated to negative roots. That is, $\mathbf{P} = \mathbf{P}_{\Delta_0}$ where $\Delta_0 = \{\alpha\}$. Then $\mathbf{H} \subset \mathbf{P}$ and we may select a Levi \mathbb{Q} -subgroup $\mathbf{F} \subset \mathbf{P}$ which contains \mathbf{H} . Let $\mathbf{L} = [\mathbf{F}, \mathbf{F}]$. Then $\mathbf{H} \subset \mathbf{L}$. Moreover, \mathbf{L} is semisimple and decomposes as an (almost) direct sum of almost \mathbb{Q} -simple factors; of these \mathbf{H} is the unique \mathbb{Q} -isotropic, almost \mathbb{Q} -simple factor of \mathbf{L} . By Remark 3.5 we have that \mathbf{L} and hence \mathbf{H} are simply connected.

The above also gives the following characterization of standard \mathbb{Q} -rank-1 subgroups.

Lemma 3.14. *If H is a standard \mathbb{Q} -rank-1 subgroup of G then there is a parabolic \mathbb{Q} -subgroup Q of G with rational Langlands decomposition $Q = (L \times A) \ltimes U$ such that H is (the identity component of) the unique \mathbb{Q} -isotropic, almost \mathbb{Q} -simple factor of L .*

3.7. Standing hypotheses when Γ is nonuniform. Recall the standing hypotheses and reductions Section 2.1. When $\Gamma \subset G$ is nonuniform, in the proof of Theorem D we will assume without loss of generality the following additional standing hypotheses:

- (1) G is a connected simply connected Lie group without compact factors and with $\text{rank}_{\mathbb{R}}(G) \geq 2$;
- (2) Γ is a nonuniform, irreducible lattice subgroup of G which contains the center \mathcal{Z} of G ;
- (3) there is a connected, algebraically simply connected, linear algebraic group \mathbf{F} defined over \mathbb{Q} which is almost \mathbb{Q} -simple, non-commutative, \mathbb{Q} -isotropic, and has $\text{rank}_{\mathbb{R}}(\mathbf{F}) \geq 2$ such that $F = \mathbf{F}(\mathbb{R})$ is a connected Lie group and $F_{\mathbb{Z}} = \mathbf{F}(\mathbb{Z})$ is a lattice in F ;
- (4) there is a continuous, surjective Lie group morphism $\phi: G \rightarrow F$ and a finite-index subgroup $\widehat{\Gamma} \subset F_{\mathbb{Z}}$ such that:
 - (a) $\ker \phi$ is contained in the center \mathcal{Z} of G ;
 - (b) the only torsion elements of $\widehat{\Gamma}$ are central;
 - (c) $\phi(\Gamma) = \widehat{\Gamma}$.

To see there is no loss of generality, let $\mathcal{Z} \subset G$ denote the center of G . As G has no compact factors, Γ has finite index in $\Gamma \cdot \mathcal{Z}$ (see [47, Chapter IX, Lemma (6.1)]). Without loss of generality, we may assume $\mathcal{Z} \subset \Gamma$. Let $\overline{G} = \text{Ad}(G) = G/\mathcal{Z}$ and $\overline{\Gamma} = \Gamma/\mathcal{Z}$ denote the images of G and Γ under the adjoint representation. Margulis's arithmeticity theorem (see [47] Introduction, Theorem 1', or Chapter IX, Theorem 1.11) guarantees there exist:

- (1) a \mathbb{Q} -simple, linear algebraic \mathbb{Q} -subgroup \mathbf{F} of $\mathbf{GL}(d)$, and
- (2) a surjective homomorphism $\overline{\phi}: \mathbf{F}(\mathbb{R})^{\circ} \rightarrow \overline{G}$ with compact kernel such that $\overline{\phi}(F_{\mathbb{Z}})$ is commensurable with $\overline{\Gamma}$.

Replacing \mathbf{F} with a finite cover, we may assume \mathbf{F} is algebraically simply connected [12, Proposition 2.24(ii)] whence $\mathbf{F}(\mathbb{R})$ is connected as a Lie group.

Since $\overline{\Gamma}$ is nonuniform, it follows that $F = \mathbf{F}(\mathbb{R})$ has no compact factors (see [48, Corollary 5.3.2]). It follows that the kernel of $\overline{\phi}$ is discrete and hence contained in the center of F . In particular, F is a covering space of \overline{G} . As G is the universal cover of \overline{G} it follows that the natural map $G \rightarrow \overline{G}$ factors through the map $\overline{\phi}: F \rightarrow \overline{G}$. In particular, $\overline{\phi}$ lifts to a surjective morphism of Lie groups $\phi: G \rightarrow F$. Moreover

we have $\widehat{\Gamma} := \phi(\Gamma)$ is commensurable with $F_{\mathbb{Z}}$. Passing to finite-index subgroups, we may assume $\widehat{\Gamma}$ is a torsion-free and thus the only torsion elements in $\widehat{\Gamma}$ are central.

3.8. Lifting \mathbb{Q} -structures from F to G . We keep the standing hypotheses of Section 3.7. In our proof of Theorem D we will often want to identify subgroups of G associated with \mathbb{Q} -subgroups of F . Let \mathbf{L} be an algebraic subgroup of \mathbf{F} defined over \mathbb{Q} and let $L = \mathbf{L}(\mathbb{R})^\circ \subset F$. Associated to L , there is a unique connected Lie subgroup $\widetilde{L} := (\phi^{-1}(L))^\circ \subset G$ with $\phi(\widetilde{L}) = L$. We will abuse terminology and say that \widetilde{L} is a \mathbb{Q} -subgroup of G . In the sequel, we will write, equivalently,

$$\Gamma_{\widetilde{L}} = \Gamma_L := \Gamma \cap \widetilde{L}.$$

We also write

$$\widehat{\Gamma}_L = \widehat{\Gamma} \cap L.$$

We will be particularly interested in the case that \mathbf{L} is either unipotent, parabolic, or a standard \mathbb{Q} -rank-1 subgroup. We remark that if $U \subset F$ is a unipotent \mathbb{Q} -subgroup then $\widetilde{U} \rightarrow U$ and $\Gamma_U \rightarrow \widehat{\Gamma}_U$ are isomorphisms. If P is a parabolic \mathbb{Q} -subgroup and if H is a standard \mathbb{Q} -rank-1 subgroup then the maps

$$\widetilde{P} \rightarrow P^\circ, \quad \Gamma_P \rightarrow \widehat{\Gamma}_P, \quad \widetilde{H} \rightarrow H, \quad \Gamma_H \rightarrow \widehat{\Gamma}_H$$

have kernel contained in the center of G . We will call \widetilde{U} , \widetilde{P} and \widetilde{H} , respectively, unipotent, parabolic, and standard \mathbb{Q} -rank 1 \mathbb{Q} -subgroups of G .

We will also abuse terminology with the following definitions:

- (1) Let $G_{\mathbb{Q}} := \phi^{-1}(F_{\mathbb{Q}})$ denote the preimage of $F_{\mathbb{Q}}$ under $\phi: G \rightarrow F$. Given a connected \mathbb{Q} -subgroup H of G , we write $H_{\mathbb{Q}} := G_{\mathbb{Q}} \cap H$.
- (2) A 1-parameter subgroup $\{a^t\}$ of G is \mathbb{R} -diagonalizable if its image in F (equivalently, its image under the adjoint representation) is \mathbb{R} -diagonalizable.
- (3) A connected subgroup U of G is unipotent if its image in F (equivalently, its image under the adjoint representation) is unipotent.

3.9. Consequences of standing hypotheses: quasi-isometric properties.

Let $G, \Gamma, F = \mathbf{F}(\mathbb{R})$, and $\widehat{\Gamma}$ be as in the standing hypotheses in Section 3.7.

Fix a Cartan involution θ of \mathfrak{g} and also write the induced global Cartan involutions on F and G as θ . Let K and \widetilde{K} denote, respectively, the subgroups of θ -fixed points of F and G . We recall that \widetilde{K} contains the center \mathcal{Z} of G . Equip F with a right-invariant and left- K -invariant metric and equip G with the pulled-back, right- G -invariant and left- \widetilde{K} -invariant metric. Let d_F and d_G denote, respectively, the induced distances on F and G . We remark that all right-invariant metrics on F are uniformly equivalent and thus all right-invariant, bi- \mathcal{Z} -invariant metrics on G are uniformly equivalent. In particular, all results stated below are independent of the choice of θ .

3.9.1. Bi-Lipschitz and quasi-isometric properties of central extensions. Using that $G \rightarrow F$ and $\Gamma \rightarrow \widehat{\Gamma}$ are central extensions, we collect a number of bi-Lipschitz and quasi-isometric estimates used throughout the sequel. We note for the remainder of Section 3.9.1 that we make no assumptions on the rank of F or arithmeticity of $\widehat{\Gamma}$.

Let $X = K \backslash F = \widetilde{K} \backslash G$ denote the globally symmetric space associated with G and equip X with a right-invariant metric. Equip K and \widetilde{K} with their intrinsic Riemannian metrics. Recall the Iwasawa decomposition $\widetilde{K} \times A \times N \rightarrow G, (k, a, n) \mapsto$

kan of G introduced in Section 3.5. We begin with the following well-known fact; for completeness, we include a proof adapted directly from [3, Lemma 3.6.3].

Lemma 3.15. *The diffeomorphism $\psi: G \rightarrow \tilde{K} \times X$, $\psi: kan \mapsto (k, \tilde{K}an)$ is uniformly bi-Lipschitz.*

Proof. We make three observations. First AN acts isometrically on G and X on the right and also acts isometrically on $\tilde{K} \times X$ by acting only in the second coordinate. Second, \mathcal{Z} acts isometrically on G and on $\tilde{K} \times X$ acting only on the first coordinate. Finally, the map $\psi: G \rightarrow \tilde{K} \times X$ is (AN) -equivariant and \mathcal{Z} -equivariant.

The function $g \mapsto C_g$,

$$C_g = \max\{\|D_g\psi\|, \|(D_g\psi)^{-1}\|\},$$

is continuous. By (AN) -isometric-equivariance of ψ , we have $C_g = C_{gs}$ for all $s \in AN$. Thus $\sup\{C_g : g \in G\} = \sup\{C_k : k \in \tilde{K}\}$. We also have $C_g = C_{zg}$ for all $z \in \mathcal{Z}$. Since \mathcal{Z} is cocompact in \tilde{K} we have $\sup\{C_k : k \in \tilde{K}\} < \infty$. \square

We record a number of corollaries of Lemma 3.15.

As a finitely generated abelian group, equip \mathcal{Z} with an intrinsic word metric relative to some choice of finite symmetric generating set. Recall the center \mathcal{Z} of G is contained in \tilde{K} and thus \mathcal{Z} acts isometrically on \tilde{K} with compact quotient. Lemma 3.15 implies \tilde{K} is quasi-isometrically embedded in G and, in particular, the following.

Corollary 3.16. *\mathcal{Z} is quasi-isometrically embedded in G .*

We similarly have the following.

Corollary 3.17. *\mathcal{Z} is quasi-isometrically embedded in Γ .*

Proof. Select a finite symmetric generating set for Γ that includes a finite generating set for \mathcal{Z} . Then, with respect to the word-length induced by these generating sets, for all $z \in \mathcal{Z}$ we have $\text{len}_\Gamma(z) \leq \text{len}_\mathcal{Z}(z)$. On the other hand, there is $C > 1$ such that for any $\gamma \in \Gamma$ we have

$$d_G(\mathbf{1}, \gamma) \leq C \text{len}_\Gamma(\gamma).$$

Since \mathcal{Z} is quasi-isometrically embedded in G , it follows that there are A, B such that $\text{len}_\mathcal{Z}(z) \leq A \text{len}_\Gamma(z) + B$ for all $z \in \mathcal{Z}$. \square

Given a connected Lie subgroup $L \subset F$, recall we write $\tilde{L} = (\phi^{-1}L)^\circ$.

Corollary 3.18. *Let $L \subset F$ be a connected Lie subgroup. Then (\tilde{L}, d_G) is quasi-isometric to $(L, d_F) \times (\mathcal{Z} \cap \tilde{L})$.*

Proof. We have that (\tilde{L}, d_G) is bi-Lipschitz equivalent to

$$(\tilde{L} \cap \tilde{K}) \times ((\tilde{K} \cap \tilde{L}) \setminus \tilde{L}) = (\tilde{L} \cap \tilde{K}) \times ((K \cap L) \setminus L)$$

through the map ψ . Since K is compact, $K \cap L$ is a closed, hence compact subgroup of L and thus (L, d_F) is quasi-isometric to its image $(K \cap L) \setminus L$ in X and $\mathcal{Z} \cap \tilde{L}$ is cocompact in $\tilde{K} \cap \tilde{L}$ whence $\tilde{K} \cap \tilde{L}$ is quasi-isometric to $\mathcal{Z} \cap \tilde{L}$. \square

We will frequently consider the case that $L \subset F$ is a \mathbb{R} -split torus or a unipotent subgroup. In these settings, we have the following special case of the above.

Corollary 3.19. *Let $L \subset F$ be a connected Lie subgroup such that $\tilde{L} \cap \tilde{K} = \mathbf{1}$. Then*

- (1) *the map $\phi: (\tilde{L}, d_G) \rightarrow (L, d_F)$ is uniformly bi-Lipschitz and*
- (2) *$(\mathcal{Z} \cdot \tilde{L}, d_G)$ is quasi-isometric to $\mathcal{Z} \times (\tilde{L}, d_G)$.*

Let $A \subset F$ be a choice of a θ -invariant split Cartan subgroup. By abuse of notation, we also identify A with a subgroup of G and observe that $\phi: G \rightarrow F$ restricts to a bi-Lipschitz isomorphism between the two copies of A . From Corollary 3.19, $(\mathcal{Z} \cdot A, d_G)$ is quasi-isometric to $\mathcal{Z} \times (A, d_G)$.

Let K_0 be a compact subset of \tilde{K} such that $\mathcal{Z}K_0 = K_0\mathcal{Z} = \tilde{K}$. We generalize the well-known KAK decomposition of F by the following.

Corollary 3.20. *Given $g \in G$ we may write*

$$g = k'zak''$$

where $k', k'' \in K_0$, $a \in A$, and $z \in \mathcal{Z}$. Moreover, there are $A_1, B_1 \geq 1$ such that for any $g \in G$, relative to the above decomposition,

$$\frac{1}{A_1}d(g, \mathbf{1}) - B_1 \leq \max\{d(a, \mathbf{1}), d(z, \mathbf{1})\} \leq A_1d(g, \mathbf{1}) + B_1.$$

Remark 3.21 (Generalized KAK decomposition). Consider a connected reductive \mathbb{R} -subgroup $L \subset F$ and let $H = \tilde{L} := (\phi^{-1}(L))^\circ$. By abuse of terminology, we say that H is a connected reductive \mathbb{R} -subgroup of G . Write $\mathcal{Z}(H)$ for the center of H and let $Z = \mathcal{Z}(H)^\circ$.

The image $\phi(Z)$ is the connected component of an \mathbb{R} -torus in F . As discussed in Section 3.2, we have that

$$Z = \mathcal{Z}(H)^\circ = Z_{\text{anis}} \cdot Z_{\text{split}}$$

where $\phi(Z_{\text{anis}})$ is \mathbb{R} -anisotropic and $\phi(Z_{\text{split}})$ is \mathbb{R} -split. Write $H' = [H, H]$ for the derived group of H . Then

$$H = H' \cdot Z = H' \cdot Z_{\text{anis}} \cdot Z_{\text{split}}.$$

We may select the Cartan involution θ on the Lie algebra \mathfrak{g} of F and G so that H is θ -invariant.¹ Let $A \subset G$ be a θ -invariant split Cartan subgroup and let $\tilde{K} \subset G$ be the subgroup of θ -fixed points in G . Write $A_H := A \cap H$, and $\tilde{K}_H := \tilde{K} \cap H$. We collect a number of observations.

- (1) A_H is a maximal, connected, abelian subgroup of ad- \mathbb{R} -diagonalizable elements in H ; thus $Z_{\text{split}} \subset A_H \subset A$.
- (2) $\phi(\tilde{K}_H)$ is compact and, moreover, is a maximal compact subgroup of $\phi(H)$.

It follows that $Z_{\text{anis}} \subset \tilde{K}_H$.

- (3) $\mathcal{Z} \cap H = \mathcal{Z} \cap (H' \cdot Z_{\text{anis}}) = \mathcal{Z} \cap \tilde{K}_H$ and $(\mathcal{Z} \cap H)$ is cocompact in \tilde{K}_H .

There is a compact $K_{H,0} \subset \tilde{K}_H$ such that $(\mathcal{Z} \cap H) \cdot K_{H,0} = \tilde{K}_H$. We may then decompose

$$H = K_{H,0} \cdot (\mathcal{Z} \cap H) \cdot A_H \cdot K_{H,0}.$$

¹See for instance [50, Theorem 7.3]. More directly, observe that a maximal compact subgroup K' of H/\mathcal{Z} is contained in a maximal compact subgroup K of F/\mathcal{Z} . The subgroup K determines a Cartan involution θ on F/\mathcal{Z} . With respect to the Killing form on the Lie algebra \mathfrak{g} , the orthogonal complement to the Lie algebra of K' in the Lie algebra of H is contained the -1 eigenspace for θ . Then θ leaves the Lie algebra of H in \mathfrak{g} invariant.

In particular, we may write any $h \in H$ as

$$h = k'zak''$$

where $k', k'' \in K_{H,0}$, $a \in A_H$, and $z \in \mathcal{Z} \cap H$. Moreover, as in Corollary 3.20, we similarly have that there are $A_1, B_1 \geq 1$ such that for all $h \in H$, relative to the above decomposition,

$$\frac{1}{A_1}d(h, \mathbf{1}) - B_1 \leq \max\{d(a, \mathbf{1}), d(z, \mathbf{1})\} \leq A_1d(h, \mathbf{1}) + B_1.$$

3.9.2. Quasi-isometric embedding of arithmetic subgroups. We return to the standing hypotheses in Section 3.7. The subgroup $\widehat{\Gamma}$ of F is finitely generated and thus may be equipped with a choice of word-length function $\text{len}_{\widehat{\Gamma}}$ induced by a choice of finite symmetric generating set; the word-length function induces a word-metric on $\widehat{\Gamma}$. We write $(\widehat{\Gamma}, \text{len}_{\widehat{\Gamma}})$ for this metric space. As a discrete subset of F , $\widehat{\Gamma}$ also inherits a metric d_F by restricting the right-invariant Riemannian distance d_F on F to $\widehat{\Gamma}$. We similarly obtain a word-length len_{Γ} and associated word-metric Γ and a metric d_G on Γ by restricting the Riemannian distance d_G on G to Γ .

For general groups F , the metrics $\text{len}_{\widehat{\Gamma}}$ and d_F on $\widehat{\Gamma}$ need not be comparable; indeed this holds for $\text{SL}(2, \mathbb{Z})$ in $\text{SL}(2, \mathbb{R})$. However, as we have assumed $\text{rank}_{\mathbb{R}}(F) \geq 2$ and $\widehat{\Gamma}$ irreducible, we obtain the following.

Theorem 3.22 ([45]). *For F and Γ as in the standing hypotheses in Section 3.7 the word-metric and Riemannian metric on $\widehat{\Gamma}$ are quasi-isometric: there are A_0, B_0 such that for all $\gamma, \gamma' \in \widehat{\Gamma}$,*

$$\frac{1}{A_0}d_F(\gamma, \gamma') - B_0 \leq \text{len}_{\widehat{\Gamma}}(\gamma^{-1}\gamma') \leq A_0d_F(\gamma, \gamma') + B_0.$$

3.9.3. Quasi-isometric embedding of lattices in the topological universal cover. We assert that the conclusion of Theorem 3.22 also holds for the lattice subgroup Γ of G under the standing hypotheses in Section 3.7. Although this is probably well known, we do not know a reference and thus include a proof.

Recall $\widehat{\Gamma}$ is a lattice subgroup of F and $\Gamma = \phi^{-1}(\widehat{\Gamma})$ where $\phi: G \rightarrow F$ is the covering map. We have $\ker \phi \subset \mathcal{Z}$ and recall we also assumed that $\mathcal{Z} \subset \Gamma$.

In the following, all discrete groups are equipped with a word-length metric.

Lemma 3.23. *For G and Γ as in the standing hypotheses of Section 3.7 the following hold:*

- (1) Γ is quasi-isometrically embedded in G .
- (2) Γ is quasi-isometric to $\mathcal{Z} \times \widehat{\Gamma}$.

Proof. We recall the bi-Lipschitz map ψ in Lemma 3.15. Given $\gamma \in \Gamma$, write $\widehat{\gamma} = \phi(\gamma)$ for the image of γ in $\widehat{\Gamma}$. Take a finite generating set $S_{\mathcal{Z}}$ for the center \mathcal{Z} and a finite generating set S of Γ containing $S_{\mathcal{Z}}$. Let $\widehat{S} = \phi(S)$ be the image of S in $\widehat{\Gamma}$; then \widehat{S} is a finite generating set of $\widehat{\Gamma}$.

Fix a precompact fundamental domain K_0 for the \mathcal{Z} -action on \widetilde{K} . Given $\gamma \in \Gamma$, write $\gamma = z_{\gamma}k_{\gamma}a_{\gamma}n_{\gamma}$ where $z_{\gamma} \in \mathcal{Z}, k_{\gamma} \in K_0, a_{\gamma} \in \mathcal{Z}$, and $n_{\gamma} \in N$. We naturally identify the cosets $\widetilde{K} \cdot \gamma = \widetilde{K}a_{\gamma}n_{\gamma}$ and $K \cdot \widehat{\gamma}$ in $X = \widetilde{K} \backslash G = K \backslash F$. Note (\widetilde{K}, d_K) is quasi-isometric to $(\mathcal{Z}, \text{len}_{S_{\mathcal{Z}}})$ and X is quasi-isometric to (F, d_F) . Through the bi-Lipschitz map ψ in Lemma 3.15, the map $\gamma \mapsto (z_{\gamma}, \widehat{\gamma})$ is a quasi-isometry

$$(\Gamma, d_G) \rightarrow (\mathcal{Z}, \text{len}_{S_{\mathcal{Z}}}) \times (\widehat{\Gamma}, d_F).$$

By Theorem 3.22, this induces a quasi-isometry

$$(\Gamma, d_G) \rightarrow (\mathcal{Z}, \text{len}_{S_{\mathcal{Z}}}) \times (\widehat{\Gamma}, \text{len}_{\widehat{S}}).$$

It remains to compare (Γ, d_G) and (Γ, len_S) . Take $C_1 := \max_{s \in S} d_G(s, 1)$. Then for all $\gamma \in \Gamma$,

$$\frac{1}{C_1} d_G(\gamma, 1) \leq \text{len}_S(\gamma)$$

To complete the proof of the lemma, we claim there is $\widehat{C} > 1$ such that

$$\text{len}_S(\gamma) \leq \widehat{C} d_G(\gamma, 1) + \widehat{C}. \quad (3.2)$$

To establish (3.2), fix $\gamma \in \Gamma$ and write $\widehat{\gamma} = \widehat{\gamma}_1 \widehat{\gamma}_2 \dots \widehat{\gamma}_n$ where $\gamma_i \in S$, $\widehat{\gamma}_i = \phi(\gamma_i) \in \widehat{S}$, and $n = \text{len}_{\widehat{S}}(\widehat{\gamma})$. By Lemma 3.15, there is $C_2 > 0$ such that

$$d_F(\widehat{\gamma}, 1) \leq C_2 d_G(\gamma, 1). \quad (3.3)$$

We have $\gamma = z\gamma_1\gamma_2 \dots \gamma_n$ for some $z \in \mathcal{Z}$ and

$$d_G(z, 1) \leq d_G(\gamma, 1) + d_G(1, z^{-1}\gamma) \leq d_G(\gamma, 1) + C_1 n = d_G(\gamma, 1) + C_1 \text{len}_{\widehat{S}}(\widehat{\gamma}).$$

Applying Theorem 3.22 and (3.3), there is $C_3 > 1$ such that

$$d_G(z, 1) \leq C_3 d_G(\gamma, 1) + C_3.$$

By Corollary 3.16, $\text{len}_{S_{\mathcal{Z}}}(z) \leq C_4 d_G(z, 1) + C_4$, and thus

$$\text{len}_{S_{\mathcal{Z}}}(z) \leq C_5 d_G(\gamma, 1) + C_5 \quad (3.4)$$

for some constants $C_4, C_5 > 1$. Finally, by Theorem 3.22, (3.3), and (3.4), we have

$$\begin{aligned} \text{len}_S(\gamma) &= \text{len}_S(z\gamma_1\gamma_2 \dots \gamma_n) \leq \text{len}_S(z) + \text{len}_S(\gamma_1\gamma_2 \dots \gamma_n) \\ &\leq \text{len}_{S_{\mathcal{Z}}}(z) + \text{len}_{\widehat{S}}(\widehat{\gamma}) \leq \text{len}_{S_{\mathcal{Z}}}(z) + C_6 d_F(\widehat{\gamma}, 1) + C_6 \\ &\leq C_7 d_G(\gamma, 1) + C_7 \end{aligned}$$

for some constants $C_6, C_7 > 1$ establishing (3.2). \square

3.9.4. Quasi-isometric properties of parabolic and unipotent \mathbb{Q} -subgroups. Using Lemma 3.23 we formulate additional quasi-isometric relationships that will be used in the sequel. We assume the standing hypothesis in Section 3.7.

Lemma 3.24. *Let L be a connected reductive \mathbb{Q} -subgroup of F and let U be a connected unipotent \mathbb{Q} -subgroup of F normalized by L . Suppose that $L \cap U = \{1\}$. Then the following hold:*

- (1) *the map $(L, d_F) \times (U, d_F) \rightarrow (LU, d_F)$, $(g, u) \mapsto gu$, is quasi-isometric relative to d_F .*
- (2) *$L_{\mathbb{Z}} \times U_{\mathbb{Z}}$ has finite index in $(LU)_{\mathbb{Z}}$ and the map $L_{\mathbb{Z}} \times U_{\mathbb{Z}} \rightarrow (LU)_{\mathbb{Z}}$, $(g, u) \mapsto gu$, is quasi-isometric relative to $\text{len}_{F_{\mathbb{Z}}}$.*
- (3) *the map $(\widetilde{L}, d_G) \times (\widetilde{U}, d_G) \rightarrow (\widetilde{LU}, d_G)$, $(g, u) \mapsto gu$, is quasi-isometric relative to d_F .*
- (4) *the map $\Gamma_L \times \Gamma_U \rightarrow \Gamma_{LU}$, $(g, u) \mapsto gu$, is quasi-isometric relative to len_{Γ} .*

Indeed (1) follows using that the Riemannian distance from $g \in F$ to the identity is comparable to $\max\{\log \|g\|, \log \|g^{-1}\|\}$ and that $\log \|gu\|$ is comparable to $\max\{\log \|g\|, \log \|u\|\}$. (2) then follows from Theorem 3.22. For (3), we have $\widetilde{LU} = \widetilde{LU}$ (identifying $U \subset F$ with its connected preimage in G). By Corollary 3.18 we have \widetilde{LU} is quasi-isometric to $(\mathcal{Z} \cap (\widetilde{LU})) \times (LU, d_F)$ and (\widetilde{L}, d_G) is quasi-isometric

to $(\mathcal{Z} \cap (\tilde{L})) \times (L, d_F)$. The result follows from (2) and Corollary 3.19. (4) then follows from Lemma 3.23.

We have the following special case of the above setup.

Lemma 3.25. *Let P be a parabolic \mathbb{Q} -subgroup of F with rational Langlands decomposition $P = (L \times S^\circ) \ltimes U$. Then*

- (1) $S_{\mathbb{Z}}^\circ$ is trivial whence $P_{\mathbb{Z}} = (LU)_{\mathbb{Z}}$;
- (2) the map $L_{\mathbb{Z}} \times U_{\mathbb{Z}} \rightarrow P_{\mathbb{Z}}$, $(g, u) \mapsto gu$, is quasi-isometric relative to $\text{len}_{F_{\mathbb{Z}}}$.
- (3) the map $\Gamma_L \times \Gamma_U \rightarrow \Gamma_P$, $(g, u) \mapsto gu$, is quasi-isometric relative to len_{Γ} .

Lemma 3.26 ([45, (3.14)]). *Let Λ be a unipotent subgroup of $F_{\mathbb{Z}}$. Then there are finitely many maximal parabolic \mathbb{Q} -subgroups Q_1, Q_2, \dots, Q_r of G such that if we let $U_i = \text{Rad}_u Q_i$, then $\widehat{\Gamma}_{U_1}, \dots, \widehat{\Gamma}_{U_r}$ $\text{len}_{\widehat{\Gamma}}$ -quasi-isometrically boundedly generate a subgroup of $\widehat{\Gamma}$ that contains a finite-index subgroup of Λ .*

Proof. The set of unipotent elements of $\text{SL}(g, \mathbb{R})$ is Zariski closed, so the Zariski closure $\overline{\Lambda}$ of Λ is a unipotent \mathbb{Q} -subgroup of F . Therefore $\overline{\Lambda}$ is contained in the unipotent radical U of some minimal parabolic \mathbb{Q} -subgroup P [6, Prop. 3.1]. Choose a maximal \mathbb{Q} -split torus S in P , let Δ be the set of simple roots in $\Phi(S, F)$, with respect to the ordering in which roots occurring in the Lie algebra of U are positive, and let $U^{[\alpha]}$ be the (coarse) root group corresponding to the root α . Then $U = \langle U^{[\alpha]} \mid \alpha \in \Delta \rangle$. Since U is nilpotent (and simply connected), this implies that $\langle \widehat{\Gamma}_{U^{[\alpha]}} \mid \alpha \in \Delta \rangle$ generates a finite-index subgroup of $\widehat{\Gamma}_U$ by, for example, [58, Prop. 2.5]. Furthermore, for each $\alpha \in \Delta$, the subgroup $U^{[\alpha]}$ is contained in the unipotent radical of the maximal parabolic subgroup P_{Δ_α} , where $\Delta_\alpha = \Delta \setminus \{\alpha\}$. By induction on the derived length of U , it is easy to see that if a collection of subgroups generates a finite-index subgroup of the nilpotent group U , then they $\text{len}_{\widehat{\Gamma}}$ -quasi-isometrically boundedly generate a finite-index subgroup of U . \square

Corollary 3.27. *Let $U \subset F$ be a unipotent \mathbb{Q} -subgroup. Then there are finitely many maximal parabolic \mathbb{Q} -subgroups Q_1, Q_2, \dots, Q_r of G such that if we let $U_i = \text{Rad}_u Q_i$, then $\widehat{\Gamma}_{U_1}, \dots, \widehat{\Gamma}_{U_r}$ $\text{len}_{\widehat{\Gamma}}$ -quasi-isometrically boundedly generate a subgroup of $\widehat{\Gamma}$ that contains a finite-index subgroup of $\widehat{\Gamma}_U$.*

3.10. Consequences of standing hypotheses: quasi-isometric bounded generation. Let $\phi: G \rightarrow F$ and $\Gamma \rightarrow \widehat{\Gamma}$ be as in the standing hypotheses in Section 3.7. In particular, F is a \mathbb{Q} -simple algebraic \mathbb{Q} -group, $\widehat{\Gamma}$ is commensurable with $F_{\mathbb{Z}}$, and $\Gamma \rightarrow \widehat{\Gamma}$. Given an algebraic \mathbb{Q} -group $\mathbf{L} \subset \mathbf{F}$ recall we let $\tilde{L} := (\phi^{-1}(L))^\circ$ denote the connected subgroup in the lift of $L = L(\mathbb{R})^\circ$ to G . We then write $\Gamma_L = \Gamma \cap \tilde{L}$.

We summarize the main definitions from [49].

Definition 3.28 ([49, Definition 1.1]). *Let $F = \mathbf{F}(\mathbb{R})$ be a semisimple algebraic \mathbb{Q} -group and let $\widehat{\Gamma}$ be a subgroup commensurable with $F_{\mathbb{Z}}$. We say that $\widehat{\Gamma}$ is **quasi-isometrically boundedly generated by standard \mathbb{Q} -rank-1 subgroups** if there is a constant $r = r(\mathbf{F}, \widehat{\Gamma}) \in \mathbb{N}$, a finite subset $\widehat{\Gamma}_0 = \widehat{\Gamma}_0(\mathbf{F}, \widehat{\Gamma})$ of $\widehat{\Gamma}$, and a finite collection $\mathcal{L} = \mathcal{L}(\mathbf{F}, \widehat{\Gamma})$ of \mathbb{Q} -subgroups of G such that*

- (1) each $L \in \mathcal{L}$ is a standard \mathbb{Q} -rank-1 subgroup of F ;
- (2) every element γ of $\widehat{\Gamma}$ can be written in the form $\gamma = s_1 s_2 \cdots s_r$, where either
 - (a) $s_i \in \widehat{\Gamma}_0$ or
 - (b) $s_i \in \widehat{\Gamma}_L$ for some $L \in \mathcal{L}$ and $\log \|s_i\| \leq r \log \|\gamma\|$.

Returning to the lattice subgroup Γ in G , we may lift each $L \in \mathcal{L}(F, \widehat{\Gamma})$ to G and ask that Γ be efficiently generated by elements from Γ_L .

Definition 3.29 ([49, Definition 6.1]). *We say that Γ is **quasi-isometrically boundedly generated by standard \mathbb{Q} -rank-1 subgroups** if there are constants $r = r(G, \Gamma) \in \mathbb{N}$ and $C = C(G, \Gamma) > 1$, a finite subset $\Gamma_0 = \Gamma_0(G, \Gamma)$ of Γ , and a finite collection $\mathcal{L} = \mathcal{L}(G, \Gamma)$ of \mathbb{Q} -subgroups of G such that*

- (1) *each $L \in \mathcal{L}$ is a standard \mathbb{Q} -rank-1 subgroup of G ,*
- (2) *every element γ of Γ can be written in the form $\gamma = s_1 s_2 \cdots s_r$ where either*
 - (a) *$s_i \in \Gamma_0$, or*
 - (b) *$s_i \in \Gamma_L$ for some $L \in \mathcal{L}$ and $\text{len}_\Gamma(s_i) \leq C \text{len}_\Gamma(\gamma)$.*

The main result of [49] guarantees quasi-isometrically bounded generation by \mathbb{Q} -rank-1 subgroups for all lattice subgroups under consideration.

Theorem 3.30 ([49, Theorem 1.2]). *Every arithmetic subgroup $\widehat{\Gamma}$ of a \mathbb{Q} -isotropic, almost \mathbb{Q} -simple \mathbb{Q} -group F is quasi-isometrically boundedly generated by standard \mathbb{Q} -rank-1 subgroups.*

As discussed in [49, §6], the efficient bounded generation property above lifts to lattices subgroups of Lie groups with infinite center such as Γ in G satisfying the hypotheses of Section 3.7.

Theorem 3.31 ([49, Corollary 6.2]). *Let Γ be a non-uniform, irreducible lattice in a connected semisimple Lie group G without compact factors. Then Γ is quasi-isometrically boundedly generated by standard \mathbb{Q} -rank-1 subgroups.*

4. AVERAGING MEASURES AND CONTROL OF MASS NEAR ∞

For this section, take G to be a connected semisimple Lie group which acts continuously on a locally compact, second countable metric space X . Let Γ be a lattice subgroup of G . In Section 4.3 we will assume Γ is irreducible and in Section 4.4 below we will further assume $G = \mathbf{G}(\mathbb{R})$ is a \mathbb{Q} -algebraic group and Γ is commensurable with $\mathbf{G}(\mathbb{Z})$.

While many results discussed below hold for more general Lie groups (and locally compact topological groups) G and for more general discrete subgroups Γ , we only formulate results in the settings which will be used in the sequel.

4.1. Notation. Let H be a closed subgroup of G and let E be a precompact set of positive Haar measure in H . Write m_H for a choice of Haar measure on H and m_E for the Haar measure on H normalized so that $m_E(E) = 1$. Given any Borel probability measure μ on X we write

$$E * \mu := \frac{1}{m_H(E)} \int_E g_* \mu \, dm_H(g) = \int_E g_* \mu \, dm_E(g).$$

We will often take E to range over the sets in a Følner sequence, such as in the following construction: Assume that $\{a^t\}$ is an $\text{ad-}\mathbb{R}$ -diagonalizable, one-parameter subgroup of G and that U is a connected ad- unipotent subgroup of G that is normalized by $\{a^t\}$. Fix a norm $\|\cdot\|$ on Lie algebra \mathfrak{u} of U . For each $\kappa > 0$ and $t > 0$, set

$$U_\kappa(t) := \exp_{\mathfrak{u}}(\{Z \in \mathfrak{u} \mid \|Z\| \leq e^{\kappa t}\})$$

and

$$F_\kappa(t) := \{ a^s \mid 0 \leq s \leq t \} \cdot U_\kappa(t).$$

If κ is sufficiently large and if $\{t_n\}$ is a sequence tending to infinity, then $\{F_\kappa(t_n)\}$ is a (left) Følner sequence in the solvable group $\{a^t \mid t \in \mathbb{R}\} \rtimes N$; for instance, we may take

$$\kappa > 10 \max\{\|\text{Ad}_u(a)\|, \|\text{Ad}_u(a^{-1})\|\}.$$

4.2. Control of mass near ∞ . We often impose strong control on the distribution of a measure at infinity. We have the following definition.

Definition 4.1 ([17, Section 3.2]). *Let (X, d) be a complete, second countable, metric space and let μ be a finite Borel measure on X . We say that μ has **exponentially small mass at ∞** if there is $\tau_\mu > 0$ such that for all $0 < \tau < \tau_\mu$,*

$$\int_X e^{\tau d(x_0, x)} d\mu(x) < \infty \tag{4.1}$$

for some choice of base point $x_0 \in X$. We say that a collection \mathcal{M} of probability measures on X has **uniformly exponentially small mass at ∞** if there is $\tau_0 > 0$ such that for all $0 < \tau < \tau_0$,

$$\sup_{\mu \in \mathcal{M}} \left\{ \int e^{\tau d(x_0, x)} d\mu(x) \right\} < \infty. \tag{4.2}$$

We remark that (4.2) holds if and only if there is $C > 1$ such that for every sufficiently large ℓ ,

$$\sup_{\mu \in \mathcal{M}} \mu(\{x \mid d(x_0, x) \geq \ell\}) < Ce^{-\tau \ell}.$$

It is well known that the Haar measure on G/Γ has exponentially small mass at ∞ ; for discussion in the case of arithmetic lattices Γ , see (5.2) below.

4.3. Translates of horospherical orbits and mass at ∞ . In this section we assume that the lattice $\Gamma < G$ is irreducible.

Given an expanding horospherical subgroup $U = U^+(b^s)$ for some \mathbb{R} -diagonalizable, one-parameter subgroup $\{b^s\}$ in G , it is well known that the normalized Haar measure on a precompact open subset of U equidistributes to the Haar measure m on G/Γ when translated by b^s . It is useful to have a more quantitative version of this well-known fact that includes a rate of equidistribution. This “quantitative equidistribution” follows from exponential mixing, and is stated in [34, Proposition 2.4.8] and [37, Theorem 1.1] for flows on $\text{SL}(n, \mathbb{R})/\text{SL}(n, \mathbb{Z})$ and without complete optimization of all constants. Exponential mixing follows from each simple factor of G having a spectral gap in the representation on $L^2_0(G/\Gamma)$; see [34, Section 2.4]. The required spectral gap is proven for irreducible non-uniform lattices by Kleinbock and Margulis in [36, Theorem 1.12]. The result for all irreducible lattices follows easily from arguments in that paper and Clozel’s proof of Property τ [20, Theorem 3.1]. Clozel actually proves a uniform spectral gap over all congruence lattices, deducing a spectral gap for a single lattice, congruence or not follows easily as in the proof of [36, Theorem 1.12].

We state an adaptation, referred to as the “effective equidistribution property” in [39, Theorem 2.5], which holds on general G/Γ and optimizes all constants. To state this result, given $x \in G/\Gamma$ write $\text{inj } x$ for the injectivity radius of G/Γ at x .

Theorem 4.2 ([39, Theorem 2.5]). *Let U be the expanding horospherical subgroup of a one-parameter \mathbb{R} -diagonalizable subgroup $\{b^s\}$ of G . There exist constants $C, \lambda > 0$, and $k \in \mathbb{N}$, such that, for all $x \in G/\Gamma$, all $s > \max\{C, C \log(1/\text{inj } x)\}$, all $\varphi \in C_2^\infty(G/\Gamma)$, and all $f \in C_{\text{cpt}}^\infty(U)$ supported in the unit ball centered at the identity, we have*

$$\left| \int_U f(u) \varphi(b^s u x) dm_U(u) - \int_U f dm_U \int_{G/\Gamma} \varphi dm_{G/\Gamma} \right| < C \cdot \max(\|\varphi\|_{C^1}, \|\varphi\|_{k,2}) \cdot \|f\|_{C^k} \cdot e^{-\lambda s}.$$

Although it is not used directly in the paper, we state the following consequence of the above.

Corollary 4.3. *Suppose that U is the expanding horospherical subgroup of a one-parameter, \mathbb{R} -diagonalizable subgroup $\{b^s\}$ of G and that μ is a U -invariant Borel probability measure on G/Γ with exponentially small mass at ∞ .*

Then $(b^s)_ \mu \rightarrow m_{G/\Gamma}$ as $s \rightarrow +\infty$ and the family $\{(b^s)_* \mu\}_{s \geq 0}$ has uniformly exponentially small mass at ∞ .*

In the proof of Corollary 4.3 and later in Section 10.3.3, we will need the following construction. Consider the function $\hat{h}: G/\Gamma \rightarrow [0, \infty)$ given by

$$\hat{h}: g\Gamma \mapsto d(g\Gamma, \mathbf{1}\Gamma).$$

The function \hat{h} is proper and 1-Lipschitz but need not be differentiable. At times it will be convenient to replace \hat{h} with a C^∞ approximation. Convolving \hat{h} with a non-negative, symmetric, C^∞ function on G with sufficiently small support, we obtain a proper, uniformly Lipschitz, C^∞ function $h: G/\Gamma \rightarrow [0, \infty)$ with

$$d(g\Gamma, \mathbf{1}\Gamma) - 1 \leq h(g\Gamma) \leq d(g\Gamma, \mathbf{1}\Gamma) + 1 \text{ for all } g \in G. \quad (4.3)$$

Proof of Corollary 4.3. The convergence to Haar is clear.

Let $x_0 = \mathbf{1}\Gamma \in G/\Gamma$. To obtain uniform estimates on the mass at ∞ , it is sufficient to find $\tau' > 0$ such that for all sufficiently large ℓ and all $s > 0$ we have

$$((b^s)_* \mu)(\{x \in G/\Gamma \mid d(x, x_0) \leq \ell\}) \geq 1 - e^{-\tau' \ell}.$$

By assumption, there is $\tau > 0$ such that for all sufficiently large $\ell > 0$,

$$\mu(\{x \in G/\Gamma \mid d(x, x_0) \leq \ell\}) \geq 1 - e^{-\tau \ell}.$$

Also, there is $\hat{\tau} > 0$ such that for all sufficiently large $\ell > 0$,

$$m_{G/\Gamma}(\{x \in G/\Gamma \mid d(x, x_0) \leq \ell\}) \geq 1 - e^{-\hat{\tau} \ell}.$$

We may assume $\{b^t\}$ is unit-speed so that $b^t = \exp(tY)$ where $\|Y\| = 1$.

Fix any ℓ . For $0 \leq s \leq \ell/2$ we have the uniform estimate

$$\begin{aligned} & ((b^s)_* \mu)(\{x \in G/\Gamma \mid d(x, x_0) \leq \ell\}) \\ &= \mu(\{x \in G/\Gamma \mid d(b^s x, x_0) \leq \ell\}) \\ &\geq \mu(\{x \in G/\Gamma \mid d(x, x_0) \leq \ell/2\}) \\ &\geq 1 - e^{-\tau \ell/2}. \end{aligned}$$

It remains to consider all $s \geq \ell/2$. Fix constants C and λ satisfying Theorem 4.2.

Given $t > 0$, let

$$B_t := \{x \in G/\Gamma \mid C \log(1/\text{inj } x) \geq t\}.$$

We have $x \mapsto -\log(\text{inj } x)$ is bounded above by a linear function of $d(x, x_0)$; in particular there is $c_1 > 0$ such that for sufficiently large t

$$\mu(B_t) \leq \mu(\{x \in G/\Gamma \mid d(x, x_0) \geq c_1 t\}) \leq e^{-\tau c_1 t}.$$

Fix a C^∞ function $\varphi: \mathbb{R} \rightarrow [0, 1]$ with $\varphi(y) = 1$ for all $y \leq 0$ and $\varphi(y) = 0$ for all $y \geq 1$. Recall the function h satisfying (4.3). Given $\ell > 0$, set $\varphi_\ell: G/\Gamma \rightarrow [0, 1]$ to be

$$\varphi_\ell(x) = \varphi(h(x) - \ell + 2).$$

Then $\varphi_\ell(x) = 1$ when $d(x, x_0) \leq \ell - 3$ and $\varphi_\ell(x) = 0$ when $d(x, x_0) \geq \ell$. For every k , we clearly have that $\|\varphi_\ell\|_{k,2}$ and $\|\varphi_\ell\|_k$ are bounded uniformly in ℓ . Fix some nonnegative $f_U \in C^\infty(U)$ supported in the unit ball centered at the identity with $\|f_U\|_1 = 1$. Then for $s > \ell/2 > C$ we have

$$\begin{aligned} & ((b^s)_* \mu)(\{x \in G/\Gamma \mid d(x, \mathbf{1}\Gamma) \leq \ell\}) \\ & \geq ((b^s)_* \mu)(\varphi_\ell) \\ & = \int_{G/\Gamma} \varphi_\ell(b^s x) d\mu(x) \\ & = \int_{G/\Gamma} \varphi_\ell(b^s x) d \left[\int_U f(u) u_* \mu dm_U(u) \right] (x) \\ & \geq \int_{(G/\Gamma) \setminus B_{\ell/2}} \varphi_\ell(b^s x) d \left[\int_U f(u) u_* \mu dm_U(u) \right] (x) \\ & = \int_{(G/\Gamma) \setminus B_{\ell/2}} \int_U f(u) \varphi_\ell(b^s u x) dm_U(u) d\mu(x) \\ & \geq \int_{(G/\Gamma) \setminus B_{\ell/2}} \left[\int_{G/\Gamma} \varphi_\ell dm_{G/\Gamma} - \widehat{C} e^{-\lambda s} \right] d\mu(x) \\ & > (1 - e^{-\tau c_1 \ell/2})(1 - e^{-\widehat{\tau}(\ell-3)} - \widehat{C} e^{-\lambda \ell/2}) \end{aligned}$$

where the second equality follows from the U -invariance of μ and the next to last inequality uses Theorem 4.2. \square

In the proof of our main results, we only use the following special case of Corollary 4.3. We note that this corollary can be proved using classical exponential mixing results rather than the more quantitative version in Theorem 4.2.

Corollary 4.4. *Suppose that U is the expanding horospherical subgroup of a one-parameter \mathbb{R} -diagonalizable subgroup $\{b^s\}$ of G and that μ is a U -invariant Borel probability measure on G/Γ with compact support. Then $(b^s)_* \mu$ converges to the Haar measure on G/Γ as $s \rightarrow +\infty$ and the family $\{(b^s)_* \mu\}_{s \geq 0}$ has uniformly exponentially small mass at ∞ .*

4.4. Non-divergence of unipotent averages of measures. We emphasize in this section that we assume $G = \mathbf{G}(\mathbb{R})$ is a \mathbb{Q} -algebraic group and Γ is commensurable with $\mathbf{G}(\mathbb{Z})$. In particular, we may view $G \subset \text{SL}(N, \mathbb{R})$ for some $N \in \mathbb{N}$. We equip $\text{SL}(N, \mathbb{R})$ with the metric associated with the Cartan involution $X \mapsto -X^T$. Although G need not be invariant under this Cartan involution, any choice of metric

on G is uniformly Lipschitz comparable with the restriction of the ambient metric on $\mathrm{SL}(N, \mathbb{R})$ to G . The results and references quoted below are primarily stated for unipotent orbits in $\mathrm{SL}(N, \mathbb{R})/\mathrm{SL}(N, \mathbb{Z})$; however, using that distance estimates in $\mathrm{SL}(N, \mathbb{R})/\mathrm{SL}(N, \mathbb{Z})$ give uniformly Lipschitz comparable estimates in G/Γ gives analogous results for unipotent orbits in G/Γ .

A key property of unipotent dynamics first established in [46] and later made quantitative in [21] is the non-divergence of orbits under unipotent flows. As we require non-divergence of orbits of certain subsets of higher-dimensional unipotent subgroups, we make the following definition.

Definition 4.5. *Let U be a k -dimensional unipotent subgroup of G . A basis $\mathcal{B} = \{X_1, \dots, X_k\}$ of the Lie algebra \mathfrak{u} of U is **regular** (or **triangular**) if for each $1 \leq i \leq k$,*

- (1) $\{X_1, \dots, X_i\}$ spans an i -dimensional subalgebra of \mathfrak{u} ;
- (2) X_i normalizes the subalgebra spanned by $\{X_1, \dots, X_{i-1}\}$.

Fix a regular basis $\mathcal{B} = \{X_1, \dots, X_k\}$ and let $\{u_j^t\} = \{\exp(tX_j)\}$ be the associated one-parameter subgroups of U . Given nonempty, bounded intervals I_1, \dots, I_k in \mathbb{R} , the product $u_k^{I_k} u_{k-1}^{I_{k-1}} \cdots u_1^{I_1}$

$$\{u_k^{t_k} \mid t_k \in I_k\} \cdot \{u_{k-1}^{t_{k-1}} \mid t_{k-1} \in I_{k-1}\} \cdots \{u_1^{t_1} \mid t_1 \in I_1\}$$

*is an **interval** (relative to the regular basis \mathcal{B}) in U where $u_i^{I_i} = \{u_i^t \mid t \in I_i\}$.*

*An interval (relative to the basis \mathcal{B}) $\widehat{U} = u_k^{I_k} u_{k-1}^{I_{k-1}} \cdots u_1^{I_1}$ is **centered** if $\mathbf{1} \in \widehat{U}$; that is, a k -dimensional interval $\widehat{U} = u_k^{I_k} u_{k-1}^{I_{k-1}} \cdots u_1^{I_1}$ is centered if $0 \in I_i$ for each $1 \leq i \leq k$.*

Remark 4.6. (Centered) k -dimensional intervals $\widehat{U} = u_k^{I_k} u_{k-1}^{I_{k-1}} \cdots u_1^{I_1}$ in U have a number of nice properties relative to dynamics and choice of base point. In particular, we have the following:

- (1) For any $g \in N_G(U)$, the conjugate $g\widehat{U}g^{-1}$ is also a (centered) k -dimensional interval in U (possibly relative to a different basis).
- (2) For any u in the subgroup generated by \widehat{U} , the translates $u\widehat{U}$ and $\widehat{U}u$ are also k -dimensional intervals in U ; moreover, they are centered if $u^{-1} \in \widehat{U}$.

Recall that a collection \mathcal{M} of Borel probability measures on a locally compact space X is **uniformly tight** if for every $\delta > 0$ there exists a compact subset \mathcal{K} of X , such that $\eta(\mathcal{K}) > 1 - \delta$ for every $\eta \in \mathcal{M}$. We recall that every uniformly tight family of Borel probability measures is pre-compact in the space of Borel probability measures equipped with the weak-* topology; in particular, any uniformly tight sequence of Borel probability measures has subsequential limit points in the space of Borel probability measures. We frequently pass to compact extensions and make use of the following observation: if $f: X_1 \rightarrow X_2$ is a proper continuous map, then a collection \mathcal{M} of Borel probability measures on X_1 is uniformly tight if and only if $f_*\mathcal{M}$ is a uniformly tight collection of measures on X_2 .

Results of the following type are well known, but we do not know of a reference for this precise fact.

Lemma 4.7. *Let $U \subset G$ be a unipotent subgroup and let $\{F_j\}$ a sequence of centered intervals (relative to a regular basis \mathcal{B}) in U . Let μ be a Borel probability measure on G/Γ . Then the family $\{F_j * \mu\}$ is uniformly tight.*

Proof. By induction on $\dim(U)$, it suffices to show that if \mathcal{M} is a uniformly tight family of Borel probability measures on G/Γ , $\{u^t\}$ is a one-parameter unipotent subgroup of G , and $\{I_n\}$ is a sequence of centered intervals in \mathbb{R} , then the family

$$\{u^{I_n} * \mu \mid n \in \mathbb{N}, \mu \in \mathcal{M}\}$$

is uniformly tight.

Given $\varepsilon > 0$, there is some compact subset \mathcal{K}_1 of G/Γ , such that $\mu(\mathcal{K}_1) > 1 - \varepsilon$ for all $\mu \in \mathcal{M}$. It follows from the uniform estimate for the quantitative nondivergence of unipotent flows established by Kleinbock and Margulis in [35, Theorem 5.2] (see also improvements in [38]) that there is a compact subset \mathcal{K}_2 , such that for every $x \in \mathcal{K}_1$, and every centered interval I in \mathbb{R} , we have

$$m_I(\{t \in I \mid u^t x \in \mathcal{K}_2\}) > 1 - \varepsilon.$$

Then $(u^{I_n} * \mu)(\mathcal{K}_2) > 1 - 2\varepsilon$ for every $n \in \mathbb{N}$ and $\mu \in \mathcal{M}$. □

We also have the following more quantitative version of tightness which follows from [17, Lemma 3.3] by induction.

Lemma 4.8 ([17, Lemma 3.3]). *Suppose*

- (1) $\{\mu_n\}$ is a sequence of probability measures on G/Γ with uniformly exponentially small mass at ∞ and
- (2) $\{\widehat{U}_n\}$ is a sequence of centered intervals (relative to a fixed regular \mathcal{B}) in a unipotent subgroup U of G .

Then the family of measures $\{\widehat{U}_n * \mu_n\}$ has uniformly exponentially small mass at ∞ .

4.5. Equidistribution and invariance. We recall that unipotent flows on G/Γ exhibit a number of additional surprising properties as exhibited by Ratner's theorems on unipotent flows [62]. In the sequel, we will only use a handful of these important results.

The first result asserts that a measure on G/Γ that is invariant under certain subgroups are automatically invariant under larger related subgroups.

Proposition 4.9 ([60, Proposition 2.1]). *Let A be a split Cartan subgroup of G , let $\beta \in \Phi(A, G)$, and let μ be a Borel probability measure on G/Γ . If μ is invariant under both A and the coarse root group $U^{[\beta]}$ then μ is also invariant under the coarse root group $U^{[-\beta]}$.*

For the second result, we recall that averaging an A -invariant Borel probability measure along a Følner sequence in a subgroup that centralizes (or normalizes) A , any weak-* limit point is an A -invariant Borel probability measure. Ratner's equidistribution theorem implies that, on the homogeneous space G/Γ , A -invariance is preserved when passing to limits of averages by unipotent subgroups normalized by A .

Proposition 4.10 ([16, Proposition 6.2(b)] and Lemma 4.7). *Let A be a split Cartan subgroup of G , let μ be an A -invariant Borel probability measure on G/Γ , let U be a unipotent subgroup that is normalized by A , and let $\{F_j\}$ be a Følner sequence of centered intervals in U . Then*

- (1) the family $\{F_j * \mu\}$ is uniformly tight, and
- (2) every weak-* subsequential limit of $\{F_j * \mu\}$ is A -invariant.

We remark that conclusion (2) employs the equidistribution theorem of Ratner and its extension by Shah; see [61, 65].

5. SIEGEL SETS, COCYCLES, AND INDUCED ACTIONS

We return to the standing hypotheses on G, Γ, F , and $\widehat{\Gamma} \subset F_{\mathbb{Z}}$ as in Section 3.7.

5.1. Siegel sets and Siegel fundamental sets. We recall the well-known construction of fundamental sets for arithmetic lattices by the use of Siegel sets; see for instance [7, §12–§13, pp. 85–94], [10, III.1–III.2], or [1, II.4] for discussion and background. Fix a maximal \mathbb{Q} -split torus \mathbf{S} in \mathbf{F} and fix a minimal parabolic \mathbb{Q} -subgroup \mathbf{P} of \mathbf{F} containing \mathbf{S} . Let $P = \mathbf{P}(\mathbb{R})$ and $S = \mathbf{S}(\mathbb{R})$. All such choices of P and S are conjugate via $F_{\mathbb{Q}}$. Fix a Cartan involution θ of $F = \mathbf{F}(\mathbb{R})$. There exists a unique conjugate (over $N_P := \text{Rad}_u(P)$) of S° in P which is θ -invariant; denote this conjugate by $A_P = A_{\mathbf{P}, \theta}$. In most applications in later sections, we will select θ so that S is θ -invariant and thus $A_P = S^\circ$. However, for the constructions in this section, this is not strictly required.

We may decompose $P = (L \times A_P) \rtimes N_P$ where $N_P = \text{Rad}_u(P)$ and $(L \times A_P)$ are conjugate by N_P to \mathbb{Q} -subgroups appearing in a choice of rational Langlands decomposition of P . If $S = \mathbf{S}(\mathbb{R})$ is assumed θ -invariant, then $(L \times A_P) \rtimes N_P$ is a rational Langlands decomposition (i.e. a decomposition into \mathbb{Q} -subgroups) as defined in Section 3.3. From Lemma 3.4, the minimality of P implies that any conjugate of L which is defined over \mathbb{Q} will be \mathbb{Q} -anisotropic. Recall that $\Phi(A_P, F)$ denotes the set of weights of F with respect to A_P . Fix an order on $\Phi(A_P, F)$ determined by N_P ; that is, select the order so that root spaces corresponding to positive roots $\Phi^+(A_P, F)$ are contained in the Lie algebra of N_P . Recall that each $\beta \in \Phi(A_P, F)$ is a homomorphism $\beta: A_P \rightarrow \mathbb{R}^+$.

For $t \in \mathbb{R}^+$, write

$$A_t := \{ a \in A_P \mid \beta(a) < t \text{ for all } \beta \in \Phi^+(A_P, F) \}.$$

Let $K = \{ g \in F \mid \theta(g) = g \} \subset F$ be the maximal compact subgroup determined by θ . A **Siegel set** (see [7, Definition 12.3] or [1, Definition 4.1]) in F relative to the choice of P and θ is a set of the form

$$\mathfrak{S} := KA_t \mathcal{C}$$

where $t \in \mathbb{R}^+$ and \mathcal{C} is any compact subset of $L^\circ \rtimes N_P$. We remark that we may alternatively define Siegel sets where \mathcal{C} is taken to be any compact subset of P° ; in this case, one may take $A_t = A_1$ to be a (negative) fundamental chamber. See discussion in [1, p. 68] or [7, 12.4].

Fix a choice of minimal parabolic \mathbb{Q} -subgroup P and Cartan involution θ as above. Siegel sets in F constructed relative to P and θ have the following key properties:

Proposition 5.1 ([7, Theorems 13.1 and 15.4, pp. 90 and 103]).

- (1) *(Covering property)* There exists a Siegel set \mathfrak{S} and a finite set $C \subset F_{\mathbb{Q}}$ such that $F = \mathfrak{S} C \widehat{\Gamma}$.
- (2) *(Separation property)* For any Siegel sets \mathfrak{S}_1 and \mathfrak{S}_2 in F , any $g \in F_{\mathbb{Q}}$, and any finite set $C \subset F_{\mathbb{Q}}$, the set

$$\{ \gamma \in F_{\mathbb{Z}} : (\mathfrak{S}_1 g) \cap (\mathfrak{S}_2 C \gamma) \neq \emptyset \}$$

is finite.

If necessary, we may take a larger compact set $\mathcal{C} \subset L^\circ \times N_P$ in the construction of the Siegel set \mathfrak{S} so that \mathcal{C} is the closure of its interior $\text{int}(\mathcal{C})$. Then there is a finite set $C \subset F_{\mathbb{Q}}$ such that $D = \mathfrak{S} \cdot C$ has the following properties:

- (1) $F = \text{int}(D) \cdot \widehat{\Gamma}$
- (2) $\{\gamma \in F_{\mathbb{Z}} : D \cap D \cdot \gamma \neq \emptyset\}$ is finite.

We call such a set D a **Siegel fundamental set** in F . Note that any such D contains a fundamental domain for the right action of $\widehat{\Gamma}$ on F .

Remark 5.2. If P' is another minimal parabolic \mathbb{Q} -subgroup, there is $g \in \mathbf{F}(\mathbb{Q})$ such that $P' = gPg^{-1}$. Write $g = klan$ where $k \in K$, $l \in L$, $a \in A_P$, and $n \in N_P$. We have that $A_{P'} = kA_Pk^{-1}$ is the unique θ -invariant subgroup of P' conjugate to a maximal \mathbb{Q} -split torus in P' and $N_{P'} := kN_Pk^{-1}$ is the unipotent radical of P' . Write $A' = A_{P'}$ and let $\Phi^+(A', F)$ be determined by $N_{P'}$. Then $kA_tk^{-1} = A'_t$.

A standard calculation shows that if $\mathfrak{S} := KA_t\mathcal{C}$ is a Siegel set constructed relative to P and θ (where \mathcal{C} is a compact subset of P°) then $\mathfrak{S} \cdot g^{-1}$ is a Siegel set relative to P' and θ . Indeed

$$\begin{aligned} \mathfrak{S} \cdot g^{-1} &= \mathfrak{S} \cdot n^{-1}a^{-1}l^{-1}k^{-1} \\ &= KA_t(\mathcal{C}n^{-1}l^{-1}a^{-1})k^{-1} \\ &= KA_t\mathcal{C}'k^{-1} \\ &= K(kA_tk^{-1})(k\mathcal{C}'k^{-1}) \\ &= KA'_t(g\mathcal{C}''g^{-1}) \end{aligned}$$

where \mathcal{C}' and \mathcal{C}'' are compact in P° whence $(g\mathcal{C}''g^{-1})$ is compact in $(P')^\circ$.

Thus, if P' is a minimal parabolic \mathbb{Q} -subgroup and if D' is a Siegel fundamental set constructed relative to P' and θ then there is a Siegel fundamental set D constructed relative to P and θ with $D' \subset D$. In particular, the construction of Siegel fundamental sets in F depends only on the choice of Cartan involution θ .

As before, let $\widetilde{K} \subset G$ be the connected subgroup in G projecting to K under $G \rightarrow F$. Recall that \widetilde{K} contains the center \mathcal{Z} of G . Fix a compact subset $K_0 \subset \widetilde{K}$ that projects onto K under the map $G \rightarrow F$. Recall that both A_P and N_P are simply connected subgroups of F ; we abuse notation and identify A_P and N_P with the associated connected subgroups of G via the map $G \rightarrow F$. Let \widetilde{L} be the connected subgroup of G projecting to L° . A **Siegel set** in G relative to the choice of P and θ is a set of the form

$$\widetilde{\mathfrak{S}} := K_0A_t\mathcal{C}$$

where $t \in \mathbb{R}^+$ and \mathcal{C} is any compact subset of $\widetilde{L} \times N_P$. For any Siegel set \mathfrak{S} in F , there exists a Siegel set $\widetilde{\mathfrak{S}}$ in G whose image under the map $G \rightarrow F$ contains \mathfrak{S} . We may similarly construct a Siegel set $\widetilde{\mathfrak{S}}$ in G and a finite set $C \subset G_{\mathbb{Q}}$ such that $\widetilde{\mathfrak{S}} \cdot C$ satisfies the covering and separation properties of Proposition 5.1. Then

$$\widetilde{D} = \widetilde{\mathfrak{S}}C$$

is a Siegel fundamental set in G for the right-action of Γ .

We note that given a connected reductive \mathbb{Q} -subgroup H of F , it is possible to construct Siegel sets for H that are contained in and well-adapted to Siegel fundamental sets for F . See [55, Theorem 4.1] for a general formulation of this fact. In Section 10, we will need this fact when we study the action of (the restriction of

a cocycle to) Γ_H where H is a standard \mathbb{Q} -rank-1 subgroup of F . For the special case of \mathbb{Q} -rank-1 subgroups, we give an elementary proof of a stronger fact: Siegel sets in H may be chosen inside Siegel sets for F relative to some choice of θ and P .

Lemma 5.3. *Let H be a standard \mathbb{Q} -rank-1 subgroup of F . Then a Cartan involution θ of F and a maximal \mathbb{Q} -split torus S in F can be chosen with the following properties:*

- (1) H is θ -invariant and the restriction $\theta|_H$ is a Cartan involution of H ;
- (2) $P_H = P \cap H$ is a minimal parabolic \mathbb{Q} -subgroup of H , $S_H = S \cap H$ is a $\theta|_H$ -invariant maximal \mathbb{Q} -split torus of P_H , and $K_H = K \cap H$ is the $\theta|_H$ -invariant maximal compact subgroup of H ;
- (3) each Siegel set of H relative to P_H and $\theta|_H$ is contained in some Siegel set of F relative to P and θ .

Moreover, if Q is the parabolic subgroup in Lemma 3.14 then $S \subset Q$.

We caution the reader that if Q is the parabolic subgroup in Lemma 3.14 we typically cannot take $P \subset Q$.

Proof of Lemma 5.3. Fix a maximal \mathbb{Q} -split torus S in F such that $H = H_\beta$ is generated by the root subgroups $U^{\pm\beta}$ for some rational root $\beta \in \Phi(A, F)$ relative to $A = S^\circ$. Let $S_H := S \cap H$ and $A_H := S_H^\circ = A \cap H$. Since A_H is one-dimensional, acting by the Weyl group of $\Phi(A, F)$ we may select an ordering and associated collection of simple roots Δ for $\Phi(A, F)$ such that A_H is contained in the union of the closed fundamental chambers in A ,

$$\begin{aligned} A^+ &:= \{a \in A : \alpha(a) \geq 1 \text{ for all } \alpha \in \Delta\}, \\ A^- &:= \{a \in A : \alpha(a) \leq 1 \text{ for all } \alpha \in \Delta\}. \end{aligned}$$

Up to reversing the order, we may assume $\beta \in \Phi(A, F)$ is a positive root. Let P be the minimal parabolic \mathbb{Q} -subgroup of F with rational Langlands decomposition $P = (L \times A) \times N_P$ where the Lie algebra of N_P consists of roots spaces associated with positive roots given by the ordering associated with the Δ . Choose a Cartan involution θ on F such that LS is θ -invariant.

With the above choices, we verify the conclusions of the lemma.

(1) The Cartan involution θ acts by inversion on every \mathbb{R} -split torus it leaves invariant. It follows that θ interchanges $U^{[\beta]}$ and $U^{[-\beta]}$ and thus $H = H_\beta$ and S_H are θ -invariant.

(2) If K is a maximal compact θ -invariant subgroup of F then, since H is θ -invariant, $K \cap H$ is a maximal compact subgroup of H .

The minimal parabolic \mathbb{Q} subgroup P contains the root spaces corresponding to all positive roots associated to Δ and thus contains $U^{[\beta]}$. We note that $S = S_H \times C_S(H)$. Thus $C_H(S_H) \subset C_H(S)$ and so $C_H(S_H) \subseteq C_G(S) \subseteq P$. Therefore P contains the minimal parabolic subgroup $C_H(S_H)U^{[\beta]}$ of H_β .

(3) From our choice of simple roots Δ , for any $t \in \mathbb{R}^+$ the ray

$$A_{H,t} := \{a \in A_H \mid \beta(a) < t \text{ for all } \beta \in \Phi^+(A_P, F)\}$$

is contained in $A_{t'}$ for some t' . Since we also have $K_H \subseteq K$ and $P_H \subset P$, the conclusion follows. \square

5.2. Coarse geometry and partitions of unity. As before, fix a reference right- F -invariant Riemannian metric on F and induced distance function d on F . Also assume the metric is bi-invariant by the center of F . We lift the metric to a right-invariant, \mathcal{Z} -bi-invariant metric on G with induced distance d . All definitions and results below hold up to bi-Lipschitz equivalence and thus are independent of the choice of d .

As above, let $\text{len}_\Gamma : \Gamma \rightarrow \mathbb{N}$ be the word-length function on Γ relative to some fixed symmetric, finite generating set of Γ . This induces a word-metric d_Γ on Γ . Recall from Lemma 3.23 there is a constant $C > 1$ such that for all $\gamma, \gamma' \in \Gamma$,

$$\frac{1}{C} d(\gamma, \gamma') \leq d_\Gamma(\gamma, \gamma') \leq C d(\gamma, \gamma'). \quad (5.1)$$

It is well known that $m_F(\mathfrak{S}) < \infty$ for any Siegel set \mathfrak{S} in F ; see for instance [56, Proposition 4.11, p. 215]. Moreover, the proof of this fact shows there is some $\tau > 0$ such that

$$\int_{\mathfrak{S}} e^{\tau d(g, \mathbf{1})} dm_F(g) < \infty. \quad (5.2)$$

This implies that the measure $m_{G/\Gamma}$ on the homogeneous space $F/\widehat{\Gamma} = G/\Gamma$ has exponentially small mass at ∞ . (See Definition 4.1).

If D is a Siegel fundamental set in F then the map $D \rightarrow F/\widehat{\Gamma} = G/\Gamma$ does not drastically distort distances. In particular, there are $C > 1$ and $B > 0$ such that for all $g \in D$,

$$d(g, \mathbf{1}) \leq Cd(g\widehat{\Gamma}, \mathbf{1}\widehat{\Gamma}) + B.$$

Moreover, if the metric on F is taken left K -invariant we can take $C = 1$ above. See for example [8, Theorem C]. For the reader's convenience we sketch a proof. One can argue in a single Siegel set. If \mathcal{C} is a compact subset of $L^\circ \times N_P$ then, for any $t \geq 1$, the choice of chamber in A ensures the diameter of $a\mathcal{C}a^{-1}$ is uniformly bounded over the choice of $a \in A_t$. Then for $w \in \mathcal{C}$, $k \in K$, and $a \in A_t$,

$$d(kaw, \mathbf{1}) = d(kawa^{-1}a, \mathbf{1}) \leq \text{diam } K + \text{diam}(a\mathcal{C}a^{-1}) + d(a, \mathbf{1}).$$

One then argues (see discussion in [10, §III.21] or [32, §5] especially [32, Proposition 5.12]) that for some $0 < t_0 < 1$ sufficiently small, the partial orbit $x_0 \cdot A_{t_0}$ is a convex subset of X/Γ where $x_0 = K$ is the origin in $X = K \backslash F$. For a much stronger result see [43, Theorem 5.7] and discussion in [10, III.21.15]. Similarly, using that $A_P \subset G$ is chosen θ -invariant and thus geodesic, the bound

$$d(g, \mathbf{1}) \leq Cd(g\Gamma, \mathbf{1}\Gamma) + B. \quad (5.3)$$

also holds for all g in any Siegel fundamental set \widetilde{D} in G .

For the following construction, we fix a minimal parabolic \mathbb{Q} -subgroup P and Cartan involution θ of G . Let K be the maximal compact subgroup determined by θ and let D be a Siegel fundamental set in F constructed relative to P and θ . Let \widetilde{D} be a Siegel fundamental set in G relative to P and θ whose image under $G \rightarrow F$ contains D .

Let X denote the globally symmetric space $X := K \backslash F = \widetilde{K} \backslash G$. The locally symmetric space X/Γ admits compactification $\overline{X/\Gamma}^{\text{BS}} = \overline{\mathbb{Q}X}^{\text{BS}}/\Gamma$, known as the Borel-Serre compactification, as a real-analytic manifold with corners. See [10, §III.9, pp. 326–338] for details. In this compactification, neighborhoods of points at infinity are of the form $(\mathbb{R}_{\geq 0})^{k_1} \times \mathbb{R}^{k_2}$ for some k_1, k_2 ; see [10, Lemma III.9.13, p. 362,

and Proposition III.9.17, p. 365]. We note that such coordinate neighborhoods of points at infinity in the partial compactification ${}_{\mathbb{Q}}\overline{X}^{\text{BS}}$ can be taken to be in the closure in ${}_{\mathbb{Q}}\overline{X}^{\text{BS}}$ of a single Siegel set in $K \backslash F$. Since the only torsion elements of $\widehat{\Gamma}$ are contained in K , $\widehat{\Gamma}$ acts freely on ${}_{\mathbb{Q}}\overline{X}^{\text{BS}}$. In particular, we may form an open cover $\overline{X/\Gamma}^{\text{BS}}$ consisting of sets $\{U_p : p = 1, \dots, \ell\}$ such that every U_p is the injective image of either

- (1) an open set $\widetilde{U}_p \subset X$, or
- (2) an open set \widetilde{U}_p in the closure of a Siegel set in ${}_{\mathbb{Q}}\overline{X}^{\text{BS}}$.

We may then select a C^∞ partition of unity $\{\widehat{\psi}_p \mid i = 1, \dots, \ell\}$ on $\overline{X/\Gamma}^{\text{BS}}$ subordinate to the cover $\{U_p\}$. Let $\rho: {}_{\mathbb{Q}}\overline{X}^{\text{BS}} \rightarrow \overline{X/\Gamma}^{\text{BS}}$ be the canonical projection. For each $1 \leq i \leq p$, the function $\widehat{\psi}_p \circ \rho$ is right- Γ invariant and is supported in the open set $\widetilde{U}_p \cdot \Gamma$; select γ_p so that $\widetilde{U}_p \cdot \gamma_p$ meets the image in X of the Siegel fundamental set D in F fixed above and let $\psi_p = \mathbb{1}_{\widetilde{U}_p \cdot \gamma_p} \cdot (\widehat{\psi}_p \circ \rho)$ denote the lift of $\widehat{\psi}_p$ supported in $\widetilde{U}_p \cdot \gamma_p$. Similarly select an open cover of K consisting of sets $\{V_q\}$, each of which is the injective image of an open set $\widetilde{V}_q \subset \widetilde{K}$. Let $\{\widehat{h}_q \mid j = 1, \dots, m\}$ be a partition of unity on K subordinate to $\{V_q\}$ and for each q , choose a lift $h_q: \widetilde{K} \rightarrow [0, 1]$ whose support intersects a choice of compact fundamental domain for $\widetilde{K} \rightarrow K$. Given $g \in G$, write $g = kan$ and let

$$\phi_{p,q}(g) = h_q(k)\psi_p(\widetilde{K}an).$$

Let $\{\phi_i : 1 \leq i \leq N\}$ denote the collection of functions $\{\phi_{p,q}\}$.

We have the following properties of the above construction which we use in constructions in the next subsections.

- (1) For each i and $\gamma \in \Gamma$, write $\phi_{i,\gamma}: G \rightarrow [0, 1]$ for the function

$$\phi_{i,\gamma}(g) = \phi_i(g\gamma^{-1}).$$

From the assumptions on the supports of ϕ_i we have that the supports satisfy $\text{supp}(\phi_{i,\gamma}) \cap \text{supp}(\phi_{i,\gamma'}) = \emptyset$ whenever $\gamma \neq \gamma'$.

- (2) Since the coordinate neighborhoods of points at infinity are in the closures of single Siegel sets, the separation property of Siegel sets (see also the proof of [10, Proposition III.9.17, p. 362]) shows that, for the Siegel fundamental set \widetilde{D} in G fixed above, the set

$$\{\gamma \in \Gamma \mid \text{supp}(\phi_{i,\gamma}) \cap \widetilde{D} \neq \emptyset \text{ for some } i\}$$

is finite.

- (3) The collection $\{\phi_{i,\gamma} \mid i \in \{1, \dots, N\}, \gamma \in \Gamma\}$ is a locally-finite, Γ -invariant, partition of unity on G .

5.3. Induced actions. We make precise a number of constructions appearing in the statement of Theorem D.

Given a C^1 action α of Γ on a compact manifold M there is an induced C^0 action of Γ by vector-bundle automorphisms on the tangent bundle TM . We equip TM with some background C^∞ norm. We introduce the notion of the induced action and associated constructions in this and in a somewhat more general setting.

To generalize the above setup, fix a compact metric space (X_0, d_0) and let \mathcal{E}_0 be a continuous, finite-dimensional, normed vector bundle over X_0 . We suppose

Γ acts continuously on X_0 and that the action lifts to a continuous linear cocycle $\mathcal{A}_0: \Gamma \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$. We write $\mathcal{A}_0(\gamma, x)$ for the linear map between fibers $\mathcal{E}_0(x)$ and $\mathcal{E}_0(\gamma \cdot x)$.

For any closed subgroup H of G , the action of Γ restricts to a continuous action of Γ_H on X_0 . We then obtain actions of Γ_H and H on $H \times X_0$ given by

$$(g, x) \cdot \gamma = (g\gamma, \gamma^{-1} \cdot x) \text{ and } g' \cdot (g, x) = (g'g, x)$$

for $g' \in H$ and $\gamma \in \Gamma_H$. As these actions commute, the H -action on $H \times X_0$ descends to a well-defined H -action, the **induced H -action**, on the quotient $X_H = (H \times X_0)/\Gamma_H$. We let $\mathcal{E}_H = (H \times \mathcal{E}_0)/\Gamma_H$ be the similarly defined finite-dimensional vector bundle over X_H with induced H -action. Write $\pi_{X_H}: X_H \rightarrow H/\Gamma_H$ and $\pi_{\mathcal{E}_H}: \mathcal{E}_H \rightarrow X_H$ for the canonical projections. Note that X_H is a fiber-bundle over X/Γ_H with fiber homeomorphic to X_0 .

When $H = G$, we omit subscripts and write $X = X_G$ and $\mathcal{E} = \mathcal{E}_G$. The bundle \mathcal{E} admits a continuous linear cocycle $\mathcal{A}: G \times \mathcal{E} \rightarrow \mathcal{E}$ which factors over the G -action on X . Given closed subgroups $N \subset H \subset G$, we have a well-defined restriction of the cocycle \mathcal{A} to $N \times \mathcal{E}_H \rightarrow \mathcal{E}_H$.

Given $g \in G$ and $x \in X_0$, we write $[g, x] \in X$ for the equivalence class in $X = (G \times X_0)/\Gamma$.

5.4. Construction of norms on \mathcal{E} and metric on X . We fix some choice of minimal parabolic \mathbb{Q} -subgroup P and Cartan involution θ of F . We recall the partition of unity $\{\phi_{i,\gamma}\}$ of G built relative to the Borel-Serre compactification in Section 5.2. Relative to these objects, we construct a norm on the bundle \mathcal{E} that is well adapted to the dynamics of \mathcal{A} on \mathcal{E} . We note that while this norm depends on the choice of P and θ , the dynamical properties of the cocycle \mathcal{A} will be independent of this choice; see Lemma 5.7.

First, let $\|\cdot\|_0$ denote the norm on \mathcal{E}_0 ; given $x \in M$, denote by $\|\cdot\|_{0,x}$ the norm on the fiber $\mathcal{E}_0(x)$. Given $g \in G, x \in X_0$, and $v \in \mathcal{E}_0(x)$ set

$$\|v\|_{g,x} := \sum_{i=1}^N \sum_{\gamma \in \Gamma} \phi_{i,\gamma}(g) \|\mathcal{A}(\gamma, x)(v)\|_{0,\alpha(\gamma)x}.$$

Let $\|\cdot\|'$ denote the norm on $G \times \mathcal{E}_0$ induced by the collection of norms $\{\|\cdot\|_{g,x}\}$. We collect a number of observations:

Claim 5.4.

- (1) For any compact subset $\mathcal{K} \subset G$, $\text{supp}(\phi_{i,\gamma}) \cap \mathcal{K} = \emptyset$ for all but finitely many (i, γ) .
- (2) The norms $\|\cdot\|_{g,x}$ are uniformly comparable on any Siegel fundamental set \tilde{S} in G : there is a constant $C > 0$, such that for all $g_1, g_2 \in D$, $x \in X_0$, and $v \in \mathcal{E}_0(x)$, we have

$$\frac{1}{C} \|v\|_{g_1,x} \leq \|v\|_{g_2,x} \leq C \|v\|_{g_1,x}.$$

- (3) For any $g \in G, \hat{\gamma} \in \Gamma$, and $(x, v) \in \mathcal{E}_0$ we have

$$\|\mathcal{A}(\hat{\gamma}, x)v\|_{g,\alpha(\hat{\gamma})(x)} = \|v\|_{g\hat{\gamma},x}.$$

In particular, Γ acts by isometries on the fibers of $G \times \mathcal{E}_0$ whence $\|\cdot\|'$ descends to a norm $\|\cdot\|$ on the bundle \mathcal{E} over X .

(4) With respect to the norm $\|\cdot\|$ on \mathcal{E} , for every $x \in X$ and any $v \in \mathcal{E}(x)$, the function

$$g \mapsto \|\mathcal{A}(g, x)(v)\|$$

is C^∞ .

Proof. The only assertion that is not by definition is (3). We verify

$$\begin{aligned} \|v\|_{g\hat{\gamma}, x} &= \sum_{i=1}^N \sum_{\gamma \in \Gamma} \phi_{i, \gamma}(g\hat{\gamma}) \|\mathcal{A}_0(\gamma, x)(v)\|_{0, \alpha(\gamma)(x)} \\ &= \sum_{i=1}^N \sum_{\gamma \in \Gamma} \phi_{i, \gamma\hat{\gamma}^{-1}}(g) \|\mathcal{A}_0(\gamma, x)(v)\|_{0, \alpha(\gamma)(x)} \\ &= \sum_{i=1}^N \sum_{\gamma \in \Gamma} \phi_{i, \gamma\hat{\gamma}^{-1}}(g) \|\mathcal{A}_0(\gamma\hat{\gamma}^{-1}\hat{\gamma}, x)(v)\|_{0, \alpha(\gamma\hat{\gamma}^{-1})(\alpha(\hat{\gamma})(x))} \\ &= \sum_{i=1}^N \sum_{\gamma \in \Gamma} \phi_{i, \gamma\hat{\gamma}^{-1}}(g) \|\mathcal{A}_0(\gamma\hat{\gamma}^{-1}, \alpha(\hat{\gamma})(x))\mathcal{A}_0(\hat{\gamma}, x)(v)\|_{0, \alpha(\gamma\hat{\gamma}^{-1})(\alpha(\hat{\gamma})(x))} \\ &= \|\mathcal{A}(\hat{\gamma}, x)v\|_{g, \alpha(\hat{\gamma})(x)}. \quad \square \end{aligned}$$

The uniform comparability of the norms on Siegel fundamental sets in Claim 5.4(2) can be reformulated as follows: for any (relative to θ and P) Siegel fundamental set D , there is $C > 1$ so that for all $g \in D$ and $x \in X_0$,

$$\frac{1}{C} \leq m(\mathcal{A}(g, [\mathbf{1}, x])) \leq \|\mathcal{A}(g, [\mathbf{1}, x])\| \leq C. \quad (5.4)$$

We also have the following direct corollary of Claim 5.4(2). Let \tilde{K} be the fixed point set of θ so that \tilde{K}/\mathcal{Z} is a maximal compact θ -invariant subgroup of $\text{Ad}(G)$.

Claim 5.5.

(1) For any compact set $\mathcal{K} \subset \tilde{K}$ there is $C > 1$ so that for all $x \in X$ and $k \in \mathcal{K}$,

$$\|\mathcal{A}(k, x)\| \leq C.$$

(2) There is $C > 1$ such that for all $x \in X$ and $z \in \mathcal{Z}$,

$$\|\mathcal{A}(z, x)\| \leq C \sup_{x_0 \in X_0} \|\mathcal{A}_0(z, x_0)\|.$$

Remark 5.6 (Dependence on the choice of P and θ). In the sequel, we study growth properties the cocycle \mathcal{A} relative to the family of norms constructed above. We note that if D is a Siegel fundamental set constructed relative to a choice of P and θ then the norm on $\|\cdot\|$ does not depend strongly on the choice of fundamental domain $\mathcal{F} \subset D$ or partitions of unity used in the construction; indeed, the strong separation properties of Siegel sets in Proposition 5.1(2) ensure that all norms on \mathcal{E} constructed relative to such choices are uniformly comparable. From Remark 5.2, this implies, up to multiplication by uniformly bounded functions, the construction of the norm on \mathcal{E} is independent of the choice of minimal parabolic \mathbb{Q} -subgroup P .

On the other hand, the choice of norms depends (in a non-uniformly comparable way) on the choice of Cartan involution θ . However, the following lemma guarantees that dynamical quantities including the values of Lyapunov exponents associated with the cocycle \mathcal{A} are independent of the choice of θ ; see Remark 6.2. To state the

lemma, let θ and θ' be Cartan involutions and let P and P' be minimal parabolic \mathbb{Q} -subgroups of G . Let $\|\cdot\|$ and $\|\cdot\|'$ denote, respectively, the norms constructed as above with respect to (P, θ) and (P', θ') .

Lemma 5.7. *For any right-invariant distance on G , there exists $C > 1$ and $k > 0$ such that for every $x \in X$ and $v \in \mathcal{E}(x)$, writing $\hat{x} = \pi_X(x)$ for the image of x in G/Γ ,*

$$\|v\| \leq C e^{kd(\hat{x}, \mathbf{1}\Gamma)} \|v\|'.$$

Proof. Let D and D' be, respectively, Siegel fundamental sets in G relative to the choice of (P, θ) and (P', θ') . Consider any $x \in X$ and $v \in \mathcal{E}(x)$. There are lifts $\tilde{x}_1 \in D \times X_0$ and $\tilde{x}_2 \in D' \times X_0$ of x . In particular, there are $g_0 \in D$, $t_0 \in X_0$, and $\tilde{v} \in \mathcal{E}_0(t_0)$ with

$$\tilde{x}_1 = (g_0, t_0) \text{ and } \|v\| = \|\tilde{v}\|_{g_0, t_0}.$$

There is $\gamma \in \Gamma$ with $g_0\gamma \in D'$. Let $t_1 = \alpha(\gamma^{-1})(t_0)$. Then \tilde{x}_2 may be chosen so that

$$\tilde{x}_2 = (g_0\gamma, t_1) \text{ and } \|v\|' = \|\mathcal{A}_0(\gamma^{-1}, t_0)(\tilde{v})\|'_{g_0\gamma, t_1}.$$

The families of norms $\|\cdot\|_{g, t_0}$ and $\|\cdot\|'_{g, t_1}$ are uniformly comparable with $\|\cdot\|_{0, t_0}$ and $\|\cdot\|_{0, t_1}$, respectively, when restricted to $g \in D$ and $g \in D'$, respectively. It follows that there exist constants $C_1, C_2 > 1$ (independent of x, v , and γ) such that

$$\begin{aligned} \|v\| &= \|\tilde{v}\|_{g_0, x} \leq C_1 \|\tilde{v}\|_{0, t_0} \\ &\leq C_1 \|\mathcal{A}(\gamma, t_1)\|_0 \|\mathcal{A}_0(\gamma^{-1}, t_0)\tilde{v}\|_{0, t_1} \\ &\leq C_1^2 \|\mathcal{A}(\gamma, t_1)\|_0 \|\mathcal{A}_0(\gamma^{-1}, t_0)\tilde{v}\|'_{g_0\gamma, t_1} \\ &\leq C_1^2 C_2^{\text{len}_\Gamma(\gamma)} \|v\|'. \end{aligned}$$

From (5.1) and (5.3), there are $C_3 > 1$ and $B > 0$ so that

$$\text{len}_\Gamma(\gamma) \leq C_3 d(g_0\Gamma, \mathbf{1}\Gamma) + B$$

and the result follows. \square

Using the construction above, we may similarly equip $G \times X_0$ with a metric \tilde{d} with the following properties:

- (1) the restriction of the metric \tilde{d} to G -orbits in $G \times X_0$ projects to a fixed right-invariant metric d_G on G ;
- (2) Γ acts by isometries relative to \tilde{d} ;
- (3) the restrictions of the metric \tilde{d} to each $\{g\} \times X_0$ are uniformly comparable when g is restricted to a Siegel fundamental set.

We then obtain a metric d_X on $X = (G \times X_0)/\Gamma$. Using that the projection $\pi_X : X \rightarrow G/\Gamma$ is proper and that the fibers are uniformly comparable, the metric on X satisfies the following properties:

- (1) there exists $B > 0$ such that $d_X(x, y) - B \leq d_{G/\Gamma}(\pi_X(x), \pi_X(y)) \leq d_X(x, y)$ for all $x, y \in X$;
- (2) a sequence of Borel probability measures $\{\mu_j\}$ on X has uniformly exponentially small mass at ∞ if and only if the projections $\{(\pi_X)_* \mu_j\}$ form a sequence of Borel probability measures on G/Γ with uniformly exponentially small mass at ∞ ;

- (3) a sequence of Borel probability measures $\{\mu_j\}$ on X is uniformly tight if and only if the projections $\{(\pi_X)_*\mu_j\}$ form a uniformly tight sequence of Borel probability measures on G/Γ .

In the sequel, we will typically ignore the specific metric d_X on X and only consider coarse metric properties of points in X under the projection $\pi_X: X \rightarrow G/\Gamma$.

5.5. Growth estimates on the cocycle \mathcal{A} . We fix P and θ as above and an associated norm $\|\cdot\|$ on \mathcal{E} constructed as in Section 5.4 relative these choices. Let $D \subset G$ be an associated Siegel fundamental set for $\Gamma \subset G$. We present a number of definitions and lemmas that allow us to control and relate growth of the cocycles \mathcal{A} and \mathcal{A}_0 . We remark that the strong estimates on the growth of \mathcal{A} obtained in Lemmas 5.12 and 5.14 below are only true relative to the norm constructed relative to the choice of Cartan involution θ .

We begin with the following definition.

Definition 5.8. *Given a Borel fundamental domain \mathcal{F} for the right Γ -action on G , the **return cocycle** $\beta_{\mathcal{F}}: G \times G/\Gamma \rightarrow \Gamma$ is defined as follows: given $\hat{x} \in G/\Gamma$, take \tilde{x} to be the unique lift of \hat{x} in \mathcal{F} and define $\beta_{\mathcal{F}}(g, \hat{x})$ to be the unique $\gamma \in \Gamma$ such that $g\tilde{x}\gamma^{-1} \in \mathcal{F}$.*

One verifies that $\beta_{\mathcal{F}}$ is, in fact, a Borel-measurable cocycle:

$$\beta_{\mathcal{F}}(g_1 g_2, \hat{x}) = \beta_{\mathcal{F}}(g_1, g_2 \hat{x}) \beta_{\mathcal{F}}(g_2, \hat{x}).$$

A second choice of fundamental domain \mathcal{F}' for Γ defines a cohomologous cocycle $\beta_{\mathcal{F}'}$.

For the remainder, we will always assume that the fundamental domain \mathcal{F} is contained in a Siegel fundamental set D in G .

Lemma 5.9. *For a fundamental domain \mathcal{F} contained in a Siegel fundamental set in G there is a C such that for every $g \in G$ and $\hat{x} \in G/\Gamma$,*

$$\text{len}_{\Gamma}(\beta_{\mathcal{F}}(g, \hat{x})) \leq Cd(g, \mathbf{1}) + Cd(\hat{x}, \Gamma) + C.$$

In particular, for any $1 \leq p < \infty$ and any compact $\mathcal{K} \subset G$ the function

$$\hat{x} \mapsto \sup_{g \in \mathcal{K}} \text{len}_{\Gamma}(\beta_{\mathcal{F}}(g, \hat{x}))$$

is in $L^p(G/\Gamma, m_{G/\Gamma})$.

Indeed, write $\hat{x} = \tilde{x}\Gamma$ for $\tilde{x} \in \mathcal{F}$. Then $g\tilde{x} = x'\beta_{\mathcal{F}}(g, \hat{x})$ for some $x' \in \mathcal{F}$ and

$$\begin{aligned} d(\mathbf{1}, \beta_{\mathcal{F}}(g, \hat{x})) &\leq d(\mathbf{1}, \tilde{x}) + d(\tilde{x}, g \cdot \tilde{x}) + d(g \cdot \tilde{x}, \beta_{\mathcal{F}}(g, \hat{x})) \\ &= d(\mathbf{1}, \tilde{x}) + d(\mathbf{1}, g) + d(\tilde{x}'\beta_{\mathcal{F}}(g, \hat{x}), \beta_{\mathcal{F}}(g, \hat{x})) \\ &= d(\mathbf{1}, \tilde{x}) + d(\mathbf{1}, g) + d(\tilde{x}', \mathbf{1}). \end{aligned}$$

We have

$$d(\tilde{x}'\Gamma, \Gamma) = d(g\hat{x}, \Gamma) \leq d(g\hat{x}, \tilde{x}) + d(\tilde{x}, \Gamma) \leq d(g, \mathbf{1}) + d(\hat{x}, \Gamma).$$

From (5.3), there is a constant \widehat{C} with $d(\mathbf{1}, \tilde{x}) \leq \widehat{C}d(\hat{x}, \Gamma) + \widehat{C}$ and

$$d(\mathbf{1}, \tilde{x}') \leq \widehat{C}d(g \cdot \hat{x}, \Gamma) + \widehat{C} \leq \widehat{C}d(g, \mathbf{1}) + \widehat{C}d(\hat{x}, \Gamma) + \widehat{C}.$$

Lemma 5.9 then follows from (5.1) and (5.2).

From the uniform comparability of the norms across a Siegel fundamental sets in Claim 5.4(2) and the finite generation of Γ we obtain the following: there are

constants $C > 1$ and $k > 0$ so that for all $g \in G$ and $x \in X$, writing $\hat{x} = \pi_X(x) \in G/\Gamma$, we have

$$\|\mathcal{A}(g, x)\| \leq Ce^{k \operatorname{len}_\Gamma(\beta_{\mathcal{F}}(g, \hat{x}))} \quad (5.5)$$

and similarly

$$m(\mathcal{A}(g, x)) \geq \frac{1}{C} e^{-k \operatorname{len}_\Gamma(\beta_{\mathcal{F}}(g, \hat{x}))}. \quad (5.6)$$

Above, $m(A) = \|A^{-1}\|^{-1}$ denotes the conorm of an invertible linear operator A .

Recall in Section 4.3 we built a proper, C^∞ , uniformly Lipschitz function $h: G/\Gamma \rightarrow [0, \infty)$ satisfying

$$d(g\Gamma, \mathbf{1}\Gamma) - 1 \leq h(g\Gamma) \leq d(g\Gamma, \mathbf{1}\Gamma) + 1 \text{ for all } g \in G. \quad (5.7)$$

We lift h to functions on X and \mathcal{E} by precomposing with the canonical projections.

For any function h satisfying (5.7), combining (5.5), (5.6), and Lemma 5.9 we obtain the following.

Proposition 5.10. *The cocycle $\mathcal{A}: G \times \mathcal{E} \rightarrow \mathcal{E}$ is **h -tempered**: there are constants $C > 1$ and $k > 0$ such that for all $g \in G$ and $x \in X$,*

$$\|\mathcal{A}(g, x)\| \leq Ce^{kh(x) + kd(g, \mathbf{1})} \quad \text{and} \quad m(\mathcal{A}(g, x)) \geq \frac{1}{C} e^{-kh(x) - kd(g, \mathbf{1})}.$$

Recalling Theorem C and Definition 1.5, we formulate the following controls on the growth of the cocycles \mathcal{A}_0 and \mathcal{A} .

Definition 5.11. *Fix closed subgroups $N \subset H \subset G$. Let $\phi: H/\Gamma_H \rightarrow [0, \infty)$ be a proper continuous function and extend to $\phi: X_H \rightarrow [0, \infty)$ by precomposition with the projection.*

- (1) *We say that the restriction of the cocycle \mathcal{A}_0 to Γ_H has **(uniform) len $_\Gamma$ -subexponential growth** if, for every $\varepsilon > 0$, there exists $C > 0$ such that for every $\gamma \in \Gamma_H$ and $x \in X_0$ we have*

$$\|\mathcal{A}_0(\gamma, x)\| \leq C\varepsilon^{\operatorname{len}_\Gamma(\gamma)}.$$

- (2) *We say that the restriction of the cocycle \mathcal{A} to $N \times \mathcal{E}_H$ has **(uniform) subexponential growth** if, for every $\varepsilon > 0$, there exists $C > 0$ such that for every $g \in N$ and $x \in X_H$ we have*

$$\|\mathcal{A}(g, x)\| \leq Ce^{\varepsilon d(g, \mathbf{1})}$$

and

$$m(\mathcal{A}(g, x)) \geq Ce^{-\varepsilon d(g, \mathbf{1})}.$$

- (3) *We say that the restriction of the cocycle \mathcal{A} to $N \times \mathcal{E}_H$ has **ϕ -tempered subexponential growth** if there exist $\omega > 0$ such that for every $\varepsilon > 0$ there exists $C_\varepsilon > 0$ such that for every $g \in N$ and $x \in X_H$ we have*

$$\|\mathcal{A}(g, x)\| \leq C_\varepsilon e^{\omega\phi(x) + \varepsilon d(g, \mathbf{1})}$$

and

$$m(\mathcal{A}(g, x)) \geq C_\varepsilon e^{-\omega\phi(x) - \varepsilon d(g, \mathbf{1})}.$$

We collect a number of crucial estimates relating the above definitions. For notational convenience in the following proofs, we introduce the subadditive cocycles $\mathcal{B}^\pm: G \times G/\Gamma \rightarrow \mathbb{R}$,

$$\mathcal{B}^+(g, x) := \sup_{y \in \pi_X^{-1}(x)} \log \|\mathcal{A}(g, y)\| \quad (5.8)$$

$$\begin{aligned} \mathcal{B}^-(g, x) &:= \mathcal{B}^+(g^{-1}, g \cdot x) \\ &= \sup_{y \in \pi_X^{-1}(x)} \log \|\mathcal{A}(g, y)^{-1}\| \\ &= \sup_{y \in \pi_X^{-1}(x)} -\log m(\mathcal{A}(g, y)). \end{aligned} \quad (5.9)$$

By Proposition 5.10 there are constants C and k with

$$\mathcal{B}^\pm(g, x) \leq kh(x) + kd(g, \mathbf{1}) + C.$$

If H is a subgroup of G and N is a unipotent \mathbb{Q} -group normalized by H we write $\tilde{H} = HN$. We observe that Γ_N is cocompact in N and that $\tilde{H}/\Gamma_{\tilde{H}}$ is a smooth fiber-bundle over H/Γ with compact fibers. Let $\rho: \tilde{H}/\Gamma_{\tilde{H}} \rightarrow H/\Gamma_{\tilde{H}}$ be the canonical projection. In the following lemma and in Lemma 5.14 below, the function ϕ which tempers the growth estimates is the map $\phi: \tilde{H}/\Gamma_{\tilde{H}} \rightarrow [0, \infty)$ given by

$$\phi(x) = h(\rho(x)) \quad (5.10)$$

where $h: G/\Gamma \rightarrow [0, \infty)$ is the C^∞ approximation to $d(\cdot, \mathbf{1}\Gamma)$ as in (5.7).

Lemma 5.12. *Let N and H be connected \mathbb{Q} -subgroups of G such that N is unipotent and H normalizes N . Suppose there is a fundamental domain \mathcal{F}_H for Γ_H in H contained in a Siegel fundamental set D in G relative to the choice of P and θ .*

Write $\tilde{H} = HN$. Then the restriction of \mathcal{A}_0 to Γ_N has len_Γ -subexponential growth if and only if the restriction of \mathcal{A} to $N \times \mathcal{E}_{\tilde{H}}$ has ϕ -tempered subexponential growth.

Proof. We first assume that the restriction of \mathcal{A}_0 to Γ_N has len_Γ -subexponential growth. Fix a compact fundamental set \mathcal{F}_N for Γ_N in N . Then $\mathcal{F}_H\mathcal{F}_N$ contains a fundamental domain for $\Gamma_{\tilde{H}}$. Indeed for all $g \in HN$ there exists $\gamma_H \in \Gamma_H$ such that $g\gamma_H \in \mathcal{F}_HN$. Then there is $\gamma_N \in \Gamma_N$ with $g\gamma_H\gamma_N \in \mathcal{F}_H\mathcal{F}_N$. Fix $gn_0 \in \mathcal{F}_H\mathcal{F}_N$. Given $n \in N$, using that H normalizes N we have

$$ngn_0 = g(g^{-1}ng)n_0 = gn_1\gamma_N$$

for some $\gamma_N \in \Gamma_N$ and $n_1 \in \mathcal{F}_N$.

Fix $\varepsilon > 0$. As $n = gn_1\gamma_N n_0^{-1}g^{-1}$, we have

$$\begin{aligned} \mathcal{B}^\pm(n, gn_0\Gamma) &= \mathcal{B}^\pm(gn_1\gamma_N n_0^{-1}g^{-1}, gn_0\Gamma) \\ &\leq \mathcal{B}^\pm(n_0^{-1}g^{-1}, gn_0\Gamma) + \mathcal{B}^\pm(\gamma_N, \mathbf{1}\Gamma) + \mathcal{B}^\pm(gn_1, \mathbf{1}\Gamma) \\ &= \mathcal{B}^\mp(gn_0, \mathbf{1}\Gamma) + \mathcal{B}^\pm(\gamma_N, \mathbf{1}\Gamma) + \mathcal{B}^\pm(gn_1, \mathbf{1}\Gamma) \\ &\leq 2kh(\mathbf{1}\Gamma) + kd(gn_0, \mathbf{1}) + kd(gn_1, \mathbf{1}) + \varepsilon \text{len}_\Gamma(\gamma_N) + \widehat{C} + C_\varepsilon \end{aligned}$$

for some constant \widehat{C} independent of ε .

As Γ is quasi-isometrically embedded in G , there are A_0 and B_0 independent of g and n such that

$$\text{len}_\Gamma(\gamma_N) \leq A_0(d(\gamma_N, \mathbf{1})) + B_0.$$

We further bound the right-hand side using $d(\gamma_N, \mathbf{1}) \leq 2d(g, \mathbf{1}) + d(n, \mathbf{1}) + 2 \operatorname{diam}(\mathcal{F}_N)$.
Indeed

$$\begin{aligned} d(\mathbf{1}, \gamma_N) &\leq d(\mathbf{1}, gn_0) + d(gn_0, ngn_0) + d(ngn_0, \gamma_N) \\ &= d(\mathbf{1}, gn_0) + d(gn_0, ngn_0) + d(gn_1 \gamma_N, \gamma_N) \\ &= d(\mathbf{1}, gn_0) + d(\mathbf{1}, n) + d(gn_1, \mathbf{1}) \\ &\leq d(\mathbf{1}, n_0) + d(n_0, gn_0) + d(\mathbf{1}, n) + d(gn_1, n_1) + d(n_1, \mathbf{1}) \\ &\leq 2d(g, \mathbf{1}) + d(n, \mathbf{1}) + 2 \operatorname{diam}(\mathcal{F}_N). \end{aligned}$$

The map $G \rightarrow F$ is 1-Lipschitz by definition and is uniformly bi-Lipschitz when restricted to N by Corollary 3.19. Since the distance from g to the identity in the linear group F is quasi-isometric to $g \mapsto \log \max\{\|g\|, \|g^{-1}\|\}$, it follows there are constants A_1, B_2 such that for all $\hat{n} \in \mathcal{F}_N$,

$$d(g\hat{n}g^{-1}, \mathbf{1}) \leq A_1 d(g, \mathbf{1}) + B_1.$$

Then

$$\begin{aligned} d(g\hat{n}, \mathbf{1}) &= d((g\hat{n}g^{-1})g, \mathbf{1}) \\ &\leq d((g\hat{n}g^{-1})g, g) + d(g, \mathbf{1}) \\ &= d(g\hat{n}g^{-1}, \mathbf{1}) + d(g, \mathbf{1}) \\ &\leq (A_1 + 1)d(g, \mathbf{1}) + B_1. \end{aligned}$$

We recall that \mathcal{F}_H is contained in a Siegel fundamental domain. It follows there are A_2 and B_2 such that for all $g \in \mathcal{F}_H$,

$$d(g, \mathbf{1}) \leq A_2 d(g\Gamma, \mathbf{1}\Gamma) + B_2 \leq A_2 h(g\Gamma) + B_2.$$

Assembling the above estimates, there are constants c_1, c_2 , and C_3 and, for any $0 < \varepsilon \leq 1$, a constant C_ε such that

$$\begin{aligned} \mathcal{B}^\pm(n, gn_0\Gamma) &\leq c_1 d(g\Gamma, \mathbf{1}\Gamma) + c_2 \varepsilon d(n, \mathbf{1}) + C_3 + C_\varepsilon \\ &\leq c_1 h(g\Gamma) + c_2 \varepsilon d(n, \mathbf{1}) + C_3 + C_\varepsilon \\ &= c_1 \phi(gn_0\Gamma) + c_2 \varepsilon d(n, \mathbf{1}) + C_3 + C_\varepsilon \end{aligned}$$

where we recall that ϕ is constant on N -orbits in $\tilde{H}/\Gamma_{\tilde{H}}$ whence

$$\phi(gn_0\Gamma) = \phi((gn_0g^{-1})g\Gamma) = \phi(g\Gamma).$$

We then take $\omega = c_1$.

For the reverse implication, by hypothesis there are constants $\hat{C} > 1$ and, for every $\varepsilon > 0$, there is C_ε such that given $\gamma \in \Gamma_N$ and $x \in X_0$, we have

$$\|\mathcal{A}_0(\gamma, x)\| \leq \hat{C} \|\mathcal{A}(\gamma, [\mathbf{1}, x])\| \leq \hat{C} C_\varepsilon e^{\omega\phi(\mathbf{1}\Gamma) + \varepsilon d(\gamma, \mathbf{1})}.$$

The conclusion then follows from (5.1). \square

In the case that Γ_H is cocompact in H we note that any compact set in G is contained in a Siegel fundamental set D (for any choice of P and θ). We then immediately obtain the following corollary.

Corollary 5.13. *Let $N \subset H$ be closed connected subgroups of G with N unipotent, N normal in H , and Γ_H cocompact in H . Then the restriction of \mathcal{A}_0 to Γ_N has $\operatorname{len}_\Gamma$ -subexponential growth if and only if the restriction of \mathcal{A} to $N \times \mathcal{E}_H$ has subexponential growth.*

In Section 10, we will also use the following variant of Lemma 5.12 that is special for the case that H is a standard \mathbb{Q} -rank-1 subgroup.

Let $\phi: \tilde{H}/\Gamma_{\tilde{H}} \rightarrow [0, \mathbb{R})$ be as in (5.10). The following lemma is important for passing from subexponential growth on unipotent subgroups to subexponential growth for trajectories that remain in special subgroups and away from the thick part.

Lemma 5.14 (*ϕ -tempered subexponential growth at ∞*). *Let H be a standard \mathbb{Q} -rank-1 subgroup of G and choose P and θ as in Lemma 5.3. Let N be a connected unipotent \mathbb{Q} -subgroup of G that is normalized by H . Assume that the restriction of \mathcal{A}_0 to Γ_N has len_Γ -subexponential growth and that the restriction of \mathcal{A}_0 to $\Gamma_{P'}$ has len_Γ -subexponential growth for every minimal parabolic \mathbb{Q} -subgroup P' of G .*

Write $\tilde{H} = HN$. Then there exist $\omega > 1$ and $\ell_0 > 0$ such that for any $\varepsilon > 0$ there is a $C_\varepsilon > 0$ such that for any continuous path $t \mapsto g_t$ in H with $g_0 = \mathbf{1}$, the following hold:

(1) **(Weak ϕ -tempered subexponential growth in the cusp on $X_{\tilde{H}}$.)**

For any $x \in \tilde{H}/\Gamma_{\tilde{H}}$ and any $T > 0$ such that $\phi(g_t \cdot x) \geq \ell_0$ for all $0 \leq t \leq T$,

$$\mathcal{B}^\pm(g_t, x) \leq \omega\phi(x) + \omega\phi(g_t \cdot x) + \varepsilon d(\mathbf{1}, g_t) + C_\varepsilon.$$

(2) **(Strong ϕ -tempered subexponential growth in the cusp on X_H .)**

For any $x \in H/\Gamma_H$ and any $T > 0$ such that $\phi(g_t \cdot x) \geq \ell_0$ for all $0 \leq t \leq T$,

$$\mathcal{B}^\pm(g_t, x) \leq \varepsilon(\phi(x) + d(\mathbf{1}, g_t)) + C_\varepsilon.$$

Proof. By the choice of P and θ as in Lemma 5.3, we may select a Siegel fundamental set D_H in H for Γ_H such that D_H is contained in a Siegel fundamental set D in G for Γ . Fix a fundamental domain \mathcal{F}_H for Γ_H in D_H . Select a precompact fundamental domain \mathcal{F}_N for the Γ_N -action on N . As in the proof of Lemma 5.12, $\mathcal{F}_H\mathcal{F}_N$ contains a fundamental domain $\mathcal{F}_{\tilde{H}}$ for the $\Gamma_{\tilde{H}}$ -action on \tilde{H} .

Let $\pi_H: H \rightarrow H/\Gamma_H$ be the natural projection. Since $\text{rank}_{\mathbb{Q}} H = 1$, there is a compact subset \mathcal{K} of H/Γ_H such that for each connected component \mathcal{C} of $\pi_H^{-1}((H/\Gamma_H) \setminus \mathcal{K})$, there is a minimal parabolic \mathbb{Q} -subgroup $P_{\mathcal{C}}$ of H such that \mathcal{C} is contained in a $H_{\mathbb{Q}}$ -translate of a Siegel set constructed relative to $P_{\mathcal{C}}$ and $\theta|_H$. In particular, for each such \mathcal{C} , the set $\{\gamma \in \Gamma \mid \mathcal{C}\gamma \cap \mathcal{C} \neq \emptyset\}$ is contained in $\Gamma_{P_{\mathcal{C}}}$; see [58, Theorem 13.10, p. 202] or [10, Proposition III.2.19, p. 307]. We remark that every minimal \mathbb{Q} -parabolic subgroup in H is the intersection of H with a minimal parabolic \mathbb{Q} -subgroup of G ; indeed all minimal \mathbb{Q} -parabolic subgroup in H are conjugate by $H_{\mathbb{Q}}$ so the conclusion follow from Lemma 5.3. It follows that the restriction of \mathcal{A}_0 to each $\Gamma_{P_{\mathcal{C}_i}}$, $1 \leq i \leq r$, has uniform len_Γ -subexponential growth.

Fix $\ell_0 > \sup\{\phi(x) : x \in \mathcal{K}\}$. Since \mathcal{F}_H is contained in a Siegel fundamental set, $\pi_H^{-1}(\phi^{-1}([\ell_0, \infty))) \cap \mathcal{F}_H$ is contained in the union of finitely many connected components $\mathcal{C}_1, \dots, \mathcal{C}_r$ of $\pi_H^{-1}((H/\Gamma_H) \setminus \mathcal{K})$. Then, since ϕ is N -invariant, if $g_0 n_0 \in \mathcal{F}_H\mathcal{F}_N$ and if $\phi(g_0 n_0 \Gamma) \geq \ell_0$ then $g_0 n_0 \in \mathcal{C}_i \mathcal{F}_N$ for some $1 \leq i \leq r$.

Consider a continuous 1-parameter path $t \mapsto g_t$ in H with $g_0 = \mathbf{1}$. Consider any $g' n_0 \in \mathcal{F}_H\mathcal{F}_N$ such that for some $T > 0$ and all $0 \leq t \leq T$,

$$\phi(g_t g' n_0 \Gamma) \geq \ell_0.$$

Fix $0 \leq t \leq T$ and write $g_t g' n_0 = g'' n_1 \gamma_H \gamma_N$ where $g'' n_1 \in \mathcal{F}_H\mathcal{F}_N$. By our choice of ℓ_0 , we have $\gamma_H \in \Gamma_{P_{\mathcal{C}_i}}$ for some $1 \leq i \leq r$.

As in the proof of Lemma 5.12,

$$\mathcal{B}^\pm(g_t, g'n_0\Gamma) \leq \mathcal{B}^\mp(g'n_0, \mathbf{1}\Gamma) + \mathcal{B}^\pm(\gamma_H, \mathbf{1}\Gamma) + \mathcal{B}^\pm(\gamma_N, \mathbf{1}\Gamma) + \mathcal{B}^\pm(g''n_1, \mathbf{1}\Gamma).$$

Let $\gamma = \gamma_H\gamma_N$. As in the proof of Lemma 5.12,

$$d(\gamma, \mathbf{1}) \leq d(g', \mathbf{1}) + d(g'', \mathbf{1}) + d(g_t, \mathbf{1}) + 2 \operatorname{diam}(\mathcal{F}_N).$$

Indeed

$$\begin{aligned} d(\mathbf{1}, \gamma) &\leq d(\mathbf{1}, g'n_0) + d(g'n_0, g_tg'n_0) + d(g_tg'n_0, \gamma) \\ &= d(\mathbf{1}, g'n_0) + d(g'n_0, g_tg'n_0) + d(g''n_1\gamma, \gamma) \\ &= d(\mathbf{1}, g'n_0) + d(\mathbf{1}, g_t) + d(g''n_1, \mathbf{1}) \\ &\leq d(\mathbf{1}, n_0) + d(n_0, g'n_0) + d(\mathbf{1}, g_t) + d(g''n_1, n_1) + d(n_1, \mathbf{1}) \\ &\leq d(g', \mathbf{1}) + d(g'', \mathbf{1}) + d(g_t, \mathbf{1}) + 2 \operatorname{diam}(\mathcal{F}_N). \end{aligned}$$

Lemma 3.23 and the hypotheses then imply there are C_0 and for any $\varepsilon > 0$ there exists C_ε such that $\mathcal{B}^\pm(\gamma_H, \mathbf{1}\Gamma)$ and $\mathcal{B}^\pm(\gamma_N, \mathbf{1}\Gamma)$ are bounded above by

$$\varepsilon C_0 d(\mathbf{1}, \gamma) + C_\varepsilon.$$

As in the proof of Lemma 5.12 there are constants k, B so that each $\mathcal{B}^\pm(g_i n_i, \mathbf{1}\Gamma)$ is bounded above by

$$kd(g_i, \mathbf{1}) + B.$$

Since $g_i \in \mathcal{F}_H$ there are C_1 and B_1 so that

$$d(g_i, \mathbf{1}) \leq C_1 d(g_i\Gamma, \mathbf{1}\Gamma) + B_1 \leq C_1 \phi(g_i\Gamma) + B_1.$$

Since $\phi(g_t g'n_0\Gamma) = \phi(g''n_1\Gamma)$ this establishes the first conclusion.

For the second conclusion, if $x \in H/\Gamma_H$ then x has a representative of $g' \in \mathcal{F}_H$. We have that $\mathcal{B}^\pm(g, \mathbf{1}\Gamma)$ is uniformly bounded over all $g \in \mathcal{F}_H$ and thus for some constants C_2, C_3, c_2 , and c_3 ,

$$\begin{aligned} \mathcal{B}^\pm(g_t, g'n_0\Gamma) &\leq C_1 + c_1\varepsilon(d(g', \mathbf{1}) + d(g'', \mathbf{1}) + d(g_t, \mathbf{1})) + C_\varepsilon \\ &\leq C_2 + c_2\varepsilon(\phi(g'\Gamma) + \phi(g_tg\Gamma) + d(g_t, \mathbf{1})) + C_\varepsilon \\ &\leq C_3 + c_2\varepsilon(2\phi(g'\Gamma) + 2d(g_t, \mathbf{1})) + C_\varepsilon \end{aligned}$$

where we use that

$$\phi(g_tg'\Gamma) \leq d(g_tg'\Gamma, \mathbf{1}\Gamma) + 1 \leq d(g_t, \mathbf{1}) + d(g'\Gamma, \mathbf{1}\Gamma) + 1 \leq d(g_t, \mathbf{1}) + \phi(g'\Gamma) + 1. \quad \square$$

6. LYAPUNOV EXPONENTS

As in Section 5.3, we continue to assume that Γ acts on a compact metric space (X_0, d_0) and the action lifts to a continuous cocycle \mathcal{A}_0 on a finite-dimensional normed vector bundle \mathcal{E}_0 over X_0 . We have the induced actions of G on $X = (G \times X_0)/\Gamma$ and $\mathcal{E} = (G \times \mathcal{E}_0)/\Gamma$. The induced linear cocycle on \mathcal{E} is denoted by \mathcal{A} . Equip \mathcal{E} with a norm as in Section 5.4; recall from Proposition 5.10 we have that \mathcal{A} is h -tempered.

6.1. Lyapunov exponents for unbounded cocycles. We have the following claim which follows immediately from Proposition 5.10 and guarantees the existence and finiteness of Lyapunov exponents for the cocycle \mathcal{A} . Equip \mathcal{E} with a norm constructed as in Section 5.4.

Claim 6.1. *Let μ a probability measure on X with exponentially small mass at ∞ . Then for any compact $E \subset G$, the functions*

$$x \mapsto \sup_{g \in E} \log \|\mathcal{A}(g, x)\| \quad \text{and} \quad x \mapsto \inf_{g \in E} \log m(\mathcal{A}(g, x))$$

are $L^1(\mu)$.

Given $a \in G$ and an a -invariant Borel probability measure μ on X we define the **average top** (or **leading**) **Lyapunov exponent of \mathcal{A}** to be

$$\lambda_{\text{top}, a, \mu, \mathcal{A}} := \liminf_{n \rightarrow \infty} \frac{1}{n} \int \log \|\mathcal{A}(a^n, x)\| \, d\mu(x). \quad (6.1)$$

This is finite when the function $x \mapsto \log \|\mathcal{A}(a, x)\|$ is integrable; in particular, for any a -invariant Borel probability measure μ on X with exponentially small mass at ∞ , the average top Lyapunov exponent $\lambda_{\text{top}, a, \mu, \mathcal{A}}$ of \mathcal{A} is finite. Note also that the a -invariance of the measure μ implies that the sequence

$$n \mapsto \int \log \|\mathcal{A}(a^n, x)\| \, d\mu(x)$$

is subadditive whence the \liminf in (6.1) can be replaced with either a limit or an infimum.

Remark 6.2. By Lemma 5.7, the value of $\lambda_{\text{top}, a, \mu, \mathcal{A}}$ is independent of the choice of norm constructed in Section 5.4 whenever the $\{a^t\}$ -invariant measure μ on X projects to a measure on G/Γ for which the function $g\Gamma \mapsto d(g\Gamma, \mathbf{1}\Gamma)$ is L^1 . In particular, for any $\{a^t\}$ -invariant measure μ on X projecting to the Haar measure on G/Γ , the value of $\lambda_{\text{top}, a, \mu, \mathcal{A}}$ is defined independent of the choice of P and θ chosen in the construction of the norm on \mathcal{E} .

With the notation established above and in Sections 5.3 and 5.4, we state the following reformulation and generalization of Theorem D.

Theorem 6.3. *Let G be a connected semisimple Lie group without compact factors and with $\text{rank}_{\mathbb{R}} G \geq 2$. Let Γ be an irreducible lattice subgroup in G and let $\mathcal{A}_0: \Gamma \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$ be a cocycle as above.*

If $\mathcal{A}_0: \Gamma \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$ fails to have subexponential growth then, for the induced G -cocycle $\mathcal{A}: G \times \mathcal{E} \rightarrow \mathcal{E}$, there exists a split Cartan subgroup A of G and a Borel probability measure μ on X such that

- (1) μ is (ZA) -invariant,
- (2) μ projects to the Haar measure on G/Γ , and
- (3) $\lambda_{\text{top}, a, \mu, \mathcal{A}} > 0$ for some $a \in ZA$.

Remark 6.4 (The case of cocompact Γ). In the case that Γ is a cocompact lattice in G , we note that the induced G -space X is compact and the cocycle \mathcal{A} is bounded. In this case, the conclusions of Theorem D and Theorem 6.3 are essentially contained in [16]. Indeed, in the setting of Theorem D, for groups G with finite center, [16, Proposition 4.6] guarantees the existence of a split Cartan subgroup A of G and an A -invariant, Borel probability measure μ' on X such that the average top

Lyapunov exponent $\lambda_{\text{top},a,\mu',\mathcal{A}}$ is positive for some $a \in A$. When G has infinite center, let \tilde{K} be a maximal subgroup such that $K = \tilde{K}/\mathcal{Z}$ is a maximal compact subgroup of G/\mathcal{Z} . Let $K_0 \subset \tilde{K}$ be a compact subset that maps onto K . We may replace the KAK -decomposition of G in the proof of [16, Proposition 4.6] with a $K_0\mathcal{Z}AK_0$ decomposition as discussed in Remark 3.21. Up to finite index, $\mathcal{Z}A$ is isomorphic to $\mathbb{Z}^k \times \mathbb{R}^\ell$. See Corollary 3.20. Minor notational modifications to the proof of [16, Proposition 4.6] in the setting of Theorem 6.3 and for groups with infinite center then yields a $(\mathcal{Z}A)$ -invariant, Borel probability measure μ' on X , such that the average top Lyapunov exponent $\lambda_{\text{top},a,\mu',\mathcal{A}}$ is positive for some $a \in \mathcal{Z}A$.

It remains to modify μ' to obtain a new measure μ with the above properties and whose projection to G/Γ is the Haar measure. Assume first that all simple factors of G are higher rank (which holds, in particular, when G is simple). If $\lambda_{\text{top},a,\mu',\mathcal{A}} > 0$ for some $a \in A$ then, in the setting of Theorem D, the proof of [16, Proposition 4.7] (see especially [16, Lemma 6.6]) gives a measure μ with the desired properties. In the setting of Theorem 6.3, we again obtain the desired outcome by minor notational modifications to the proof of [16, Proposition 4.7] and [16, Lemma 6.6]. In the case that $\lambda_{\text{top},a,\mu',\mathcal{A}} = 0$ for all $a \in A$ but $\lambda_{\text{top},a,\mu',\mathcal{A}} > 0$ for some $a \in \mathcal{Z}$, one may easily adapt the arguments in the proofs of [16, Proposition 4.7] and [16, Lemma 6.6] (which are much simpler now since a commutes with all elements of G , see Lemma 6.5 below) to obtain a measure μ with the desired properties.

When G is semisimple and, in particular, when G has rank-1 factors, we can combine the proof of [16, Proposition 4.7] with the irreducibility of Γ to obtain the desired conclusion. Indeed, we may assume μ' is ergodic and thus fiberwise Lyapunov exponents for the $(\mathcal{Z}A)$ -action on (X, μ) are identified with linear functionals on $\mathcal{Z}A$ (and thus vanish on torsion elements); see for example [17, Proposition 3.11] for a formulation of Oseledec's theorem for cocycles over actions of higher-rank abelian groups. Write $G = G' \cdot G''$ where G' and G'' are normal subgroups of positive dimension with $G' \cap G'' \subset \mathcal{Z}$. Write $A' = A \cap G'$ and $A'' = A \cap G''$. Up to interchanging the roles of G' and G'' , we may assume $\lambda_{\text{top},a,\mu',\mathcal{A}}$ is positive for some $a \in \mathcal{Z}A''$. We may average μ' over a Følner sequence contained in a maximal unipotent subgroup U of G' normalized by A' and pass to a subsequential limit to obtain a Borel probability measure μ'' on X . We may then average μ'' over a Følner sequence in A and pass to a subsequential limit to obtain a Borel probability measure μ on X . The measure μ is A -invariant and, since A normalizes U , μ is also U -invariant. Moreover, using that $a \in \mathcal{Z}A''$ centralizes G' , [16, Lemma 4.2] shows that

$$\lambda_{\text{top},a,\mu,\mathcal{A}} \geq \lambda_{\text{top},a,\mu'',\mathcal{A}} \geq \lambda_{\text{top},a,\mu',\mathcal{A}} > 0.$$

The projection of μ to G/Γ is $(A'U)$ -invariant; Proposition 4.9 shows that the projection of μ to G/Γ is G' -invariant. Since Γ is irreducible, the image of Γ in G/G' is dense, which implies that every G' -orbit on G/Γ is dense; since G' is generated by unipotent elements, Ratner's measure rigidity theorem (see for example [59, Theorem 9]) implies that the Haar measure on G/Γ is the only G' -invariant, Borel probability measure on G/Γ . Thus μ projects to the Haar measure on G/Γ and has the other properties asserted in Theorem 6.3.

6.2. Lyapunov exponents under averaging. It is well known that the top Lyapunov exponent of a bounded continuous linear cocycle is an upper-semicontinuous function on the space of invariant measures (see e.g. [16, Lemma 4.2] or [68,

Lemma 9.1]). For cocycles whose norm is tempered as in Proposition 5.10, upper-semicontinuity of the top exponent holds when restricted to families of invariant measures with uniformly exponentially small mass at ∞ ; see [17, Lemma 3.8]. This implies the following two results that we frequently use.

Lemma 6.5 ([17, Lemma 3.9]). *Let $a \in G$ and let μ be an a -invariant Borel probability measure on X with exponentially small mass at ∞ . For any amenable subgroup $H \subset C_G(a)$ and any Følner sequence of precompact sets $\{F_n\}$ in H , if the family $\{F_n * \mu\}$ has uniformly exponentially small mass at ∞ then for any subsequential limit μ_∞ of $\{F_n * \mu\}$, the measure μ_∞ is a -invariant and*

$$\lambda_{\text{top},a,\mu,A} \leq \lambda_{\text{top},a,\mu_\infty,A}.$$

Averaging over Følner sequences in $\mathcal{Z}A$ and applying Lemma 6.5 yields the following.

Corollary 6.6. *Let $\{a^t\}$ be a 1-parameter, \mathbb{R} -diagonalizable subgroup of G and let μ_0 be an $\{a^t\}$ -invariant Borel probability measure on X projecting to the Haar measure on G/Γ such that $\lambda_{\text{top},a^1,\mu_0,A} > 0$. Then there is a Borel probability measure μ on X , such that*

- (1) μ projects to the normalized Haar measure on G/Γ ,
- (2) there is a split Cartan subgroup A of G that contains $\{a^t\}$ such that μ is $(\mathcal{Z}A)$ -invariant, and
- (3) $\lambda_{\text{top},a^1,\mu,A} > 0$.

6.3. Lyapunov exponents for pre-compact orbits. The above lemmas are needed to control escape of mass and unboundedness of the cocycle when averaging. For orbits that remain in a compact set, we obtain the following results. The proofs of each are straight-forward modifications of the arguments in [16, Proposition 4.6].

In Section 3.5, we defined the notion of a split Cartan subgroup of a connected semisimple Lie group in terms of a choice of Iwasawa decomposition. More generally, if H is a connected reductive Lie subgroup of G , define a **split Cartan subgroup** A_H of H to be a maximal, connected, abelian $\text{ad-}\mathbb{R}$ -diagonalizable subgroup. All such subgroups are conjugate and these definitions coincide for semisimple Lie subgroups. Moreover, there is a split Cartan subgroup $A \subset G$ such that $A_H = H \cap A$.

Lemma 6.7. *Let $\widehat{H} \subset F$ be a connected reductive \mathbb{R} -subgroup such that $\widehat{\Gamma}_{\widehat{H}}$ is cocompact in H and let $H = (\phi^{-1}\widehat{H})^\circ$ be the associated subgroup of G . Let A_H be a split Cartan subgroup of H .*

Suppose the restriction $\mathcal{A}_0|_{\mathcal{Z}}$ of the cocycle \mathcal{A}_0 to the center \mathcal{Z} of G has len_Γ -subexponential growth. If the restriction $\mathcal{A}_0|_{\Gamma_H}$ fails to have len_Γ -subexponential growth then there is an A_H -invariant Borel probability measure μ on X_H such that

$$\lambda_{\text{top},a,\mu,A} > 0$$

for some $a \in A_H$.

Proof outline. Fix a Cartan involution θ of G which leaves H invariant. We may assume the norm on \mathcal{E} is constructed relative to θ . Let $\widetilde{K} \subset G$ be the subgroup of θ -fixed points. From Claim 5.5 there is $C_0 > 0$ such that for any compact $\mathcal{K} \subset \widetilde{K}$, all $x \in X_H$, and all $k \in \mathcal{K}$,

$$\|\mathcal{A}(k, x)\| \leq C_0$$

and, given $\varepsilon > 0$, there is C_ε such that for all $x \in X_H$ and $z \in \mathcal{Z}$,

$$\|\mathcal{A}(z, x)\| \leq C_\varepsilon e^{\varepsilon \operatorname{len}_\Gamma(z)}.$$

As discussed in Remark 3.21, given $\gamma \in \Gamma_H$, we write

$$\gamma = k' z a k''$$

where $a \in A_H = A \cap H$ for some split Cartan $A \subset G$, $z \in (\mathcal{Z} \cap H)$, k', k'' are in a bounded set, and $\operatorname{len}_\Gamma(z)$ and $d(a, \mathbf{1})$ are bounded above by linear functions of $\operatorname{len}_\Gamma(\gamma)$. The assumption that $\mathcal{A}_0|_{\Gamma_H}$ fails to have uniform $\operatorname{len}_\Gamma$ -subexponential growth implies there exists $\kappa > 0$ and sequences $\gamma_n \in \Gamma_H$ with $\operatorname{len}_\Gamma(\gamma_n) \rightarrow \infty$ and $x'_n \in X_0$ such that for all n ,

$$\log \|\mathcal{A}_0(\gamma_n, x'_n)\| \geq \kappa \operatorname{len}_\Gamma(\gamma_n).$$

Write $\gamma_n = k'_n z_n a_n k''_n$. There is $C > 0$ such that for all sufficiently small $\varepsilon > 0$, for all sufficiently large n ,

$$\begin{aligned} \log \|\mathcal{A}(a_n, x_n)\| &\geq \frac{1}{C} \log \|\mathcal{A}_0(\gamma_n, x'_n)\| - \log \|\mathcal{A}(z_n, x_n)\| - C \\ &\geq \frac{1}{C} \log \|\mathcal{A}_0(\gamma_n, x'_n)\| - \varepsilon \operatorname{len}_\Gamma(z_n) - C_\varepsilon - C \\ &\geq \frac{1}{C} (\kappa - \varepsilon) \left[\operatorname{len}_\Gamma(\gamma_n) - \frac{1}{C} \operatorname{len}_\Gamma(\gamma_n) \right] - C_\varepsilon - 2C \\ &\geq \frac{1}{C^2} (\kappa - \frac{1}{C} \varepsilon) [d(a_n, \mathbf{1})] - C_\varepsilon - 2C. \end{aligned}$$

In particular, taking $\varepsilon > 0$ sufficiently small, for all sufficiently large n ,

$$\log \|\mathcal{A}(a_n, x_n)\| \geq \frac{\kappa}{2} d(a_n, \mathbf{1})$$

and the existence of the measure μ and $a \in A_H$ with the desired properties follows exactly as in the proof of [16, Proposition 4.6]. \square

When the restriction $\mathcal{A}_0|_{\mathcal{Z}}$ fails to have $\operatorname{len}_\Gamma$ -subexponential growth, we have the following which will easily lead to the desired conclusion in Proposition 7.1.

Lemma 6.8. *If the restriction $\mathcal{A}_0|_{\mathcal{Z}}$ fails to have $\operatorname{len}_\Gamma$ -subexponential growth then there is a \mathcal{Z} -invariant Borel probability measure μ on X_0 such that*

$$\lambda_{\operatorname{top}, a, \mu, \mathcal{A}_0} > 0$$

for some $a \in \mathcal{Z}$.

Proof sketch. Recall that the group \mathcal{Z} is finitely-generated abelian and quasi-isometrically embedded in Γ . We may average the norm on \mathcal{E}_0 over the action of all torsion elements so that every torsion element acts isometrically on \mathcal{E}_0 . Let $\{a_1, \dots, a_r\}$ generate the free abelian subgroup of \mathcal{Z} . If the restriction $\mathcal{A}_0|_{\langle a_i \rangle}$ of \mathcal{A}_0 to the subgroup generated by each a_i has $\operatorname{len}_\Gamma$ -subexponential growth then $\mathcal{A}_0|_{\mathcal{Z}}$ has $\operatorname{len}_\Gamma$ -subexponential growth. We may thus assume there is a_i such that the restriction $\mathcal{A}_0|_{\langle a_i \rangle}$ fails to have $\operatorname{len}_\Gamma$ -subexponential growth. Replacing a_i with a_i^{-1} if needed, a standard exercise (see for example [14, Proposition 3.1.2]) yields an a_i -invariant, Borel probability measure μ_0 on X_0 such that

$$\lambda_{\operatorname{top}, a_i, \mu_0, \mathcal{A}_0} > 0.$$

Averaging μ_0 over a Følner sequence in \mathcal{Z} , Lemma 6.5 yields a measure μ with the desired properties. \square

A softer version of the proof of [16, Proposition 4.6] (see for example [14, Proposition 3.1.2]) also establishes the following.

Lemma 6.9. *Let $\{a^t\}$ be a one-parameter subgroup of G , let $E \subset X$ be a compact subset, and let $\{x_n\}$ be a sequence of points in X such that*

- (1) $\limsup_{n \rightarrow \infty} \frac{1}{n} \log \|\mathcal{A}(a^n, x_n)\| > 0$, and
- (2) $a^t \cdot x_n \in E$ for all $0 \leq t \leq n$ and $n \in \mathbb{N}$.

Then there exists an $\{a^t\}$ -invariant, Borel probability measure μ supported on E with

$$\lambda_{\text{top}, a^1, \mu, \mathcal{A}} > 0.$$

6.4. The projectivized cocycle and its infinitesimal generator. Let $\mathbb{P}\mathcal{E} \rightarrow X$ denote the projectivization of the vector-bundle $\mathcal{E} \rightarrow X$. We represent a point $\xi \in \mathbb{P}\mathcal{E}$ as $\xi = (x, [v])$ where $[v]$ is an equivalence class of non-zero vectors in the fiber $\mathcal{E}(x)$. The G -action on \mathcal{E} by vector-bundle automorphisms induces a natural G -action on $\mathbb{P}\mathcal{E}$ which restricts to a projective transformation between each fiber and its image.

Let $\Psi: G \times \mathbb{P}\mathcal{E} \rightarrow \mathbb{R}$ be the function

$$\Psi(g, \xi) = \Psi(g, (x, [v])) = \log \left(\frac{\|\mathcal{A}(g, x)(v)\|}{\|v\|} \right).$$

We have that Ψ is an additive cocycle over the G -action on $\mathbb{P}\mathcal{E}$:

$$\Psi(g_1 g_2, \xi) = \Psi(g_1, g_2 \cdot \xi) + \Psi(g_2, \xi). \quad (6.2)$$

In particular, $\Psi(a^n, \xi) = \sum_{k=0}^{n-1} \Psi(a, a^k \cdot \xi)$.

Write $\Psi': \mathfrak{g} \times \mathbb{P}\mathcal{E} \rightarrow \mathbb{R}$ for the infinitesimal generator of Ψ : given $Y \in \mathfrak{g}$,

$$\Psi'(Y, \xi) = \lim_{t \rightarrow 0} \frac{\Psi(\exp(tY), \xi)}{t}. \quad (6.3)$$

The function Ψ' exists by the construction of the norm $\|\cdot\|$ on \mathcal{E} ; see Claim 5.4(4). The following properties hold for Ψ' :

- (1) $\Psi'(Y, \cdot): \mathbb{P}\mathcal{E} \rightarrow \mathbb{R}$ is continuous;
- (2) $\Psi'(sY, \xi) = s\Psi'(Y, \xi)$;
- (3) $\Psi(\exp(TY), \xi) = \int_0^T \Psi'(Y, \exp(sY) \cdot \xi) ds$.

Moreover, if $a^t = \exp(tY)$ and if $\tilde{\mu}$ is an $\{a^t\}$ -invariant, Borel probability measure on $\mathbb{P}\mathcal{E}$ then

$$(4) \int \Psi'(Y, \cdot) d\tilde{\mu}(\cdot) = \int \Psi(a, \cdot) d\tilde{\mu}(\cdot).$$

Note that $\mathbb{P}\mathcal{E} \rightarrow X$ is a compact extension. If $a \in G$ and if μ is an a -invariant, Borel probability measure on X then there exists at least one a -invariant, Borel probability measure on $\mathbb{P}\mathcal{E}$ projecting to μ . We have the following claim which relates the top Lyapunov exponent of \mathcal{A} with the cocycle Ψ and such measures on $\mathbb{P}\mathcal{E}$.

Claim 6.10 (See e.g. [68, Theorem 6.1]). *Let $a \in G$ and assume that μ is an a -invariant, Borel probability measure on X with exponentially small mass at ∞ . Then:*

- (1) *for any a -invariant Borel probability measure $\tilde{\mu}$ on $\mathbb{P}\mathcal{E}$ projecting to μ we have*

$$\int \Psi(a, \cdot) d\tilde{\mu} \leq \lambda_{\text{top}, a, \mu, \mathcal{A}};$$

- (2) *there exists an a -invariant Borel probability measure $\tilde{\mu}$ on $\mathbb{P}\mathcal{E}$ projecting to μ such that*

$$\int \Psi(a, \cdot) d\tilde{\mu} = \lambda_{\text{top}, a, \mu, \mathcal{A}}.$$

REVIEW OF NOTATION AND STANDING HYPOTHESES

Before we begin the proof of Theorems D and 6.3 we review the standing hypotheses and notation that will be used in the remainder of the paper. Let $\mathbb{K} \in \{\mathbb{R}, \mathbb{Q}\}$.

- (1) G is a connected simply connected semisimple Lie group with real rank at least 2 and without compact factors.
- (2) \mathcal{Z} is the center of G .
- (3) \mathbf{F} is an algebraically simply connected, \mathbb{Q} -simple, semisimple algebraic \mathbb{Q} -group. The Lie group $F = \mathbf{F}(\mathbb{R})$ is connected.
- (4) $\phi: G \rightarrow F$ is a surjective morphism with $\ker(\phi) \subset \mathcal{Z}$.
- (5) $\Gamma \subset G$ is a nonuniform irreducible lattice subgroup of G and $\mathcal{Z} \subset \Gamma$.
- (6) The image $\hat{\Gamma} := \phi(\Gamma)$ is a lattice subgroup of F commensurable with $F_{\mathbb{Z}}$; all torsion elements in $\hat{\Gamma}$ are central.
- (7) Given a connected subgroup $H \subset F$, let $\tilde{H} := (\phi^{-1}(H))^{\circ}$ be the connected subgroup of G projecting to H . Write

$$\Gamma_H = \Gamma_{\tilde{H}} := \tilde{H} \cap \Gamma$$

and

$$\hat{\Gamma}_H := \hat{\Gamma} \cap H.$$

If $H = \mathbf{H}(\mathbb{R})^{\circ}$ for some \mathbb{Q} -subgroup \mathbf{H} we say that \tilde{H} is defined over \mathbb{Q} .

- (8) A connected subgroup $H \subset G$ is a parabolic \mathbb{K} -subgroup if $\phi(H) = Q^{\circ}$ where Q is a parabolic \mathbb{K} -subgroup of F .
- (9) A connected subgroup $U \subset G$ is unipotent if $\phi(U)$ is a unipotent subgroup of F .
- (10) An element $a \in G$ is \mathbb{R} -diagonalizable if $\phi(a)$ is \mathbb{R} -diagonalizable.
- (11) Given a connected subgroup $A \subset G$, we say A is a \mathbb{K} -torus if there is an \mathbb{K} -torus $\mathbf{S} \subset \mathbf{F}$ with $\phi(A) = \mathbf{S}(\mathbb{R})^{\circ}$.

If $A \subset G$ is an \mathbb{R} -torus then A decomposes uniquely as $A = A'A''$ where $\phi(A')$, (resp. $\phi(A'')$) is a maximal \mathbb{R} -split, r(esp. \mathbb{R} -anisotropic) subgroup of $\phi(A)$.

Given a one-parameter subgroup $\{a^t\}$ of A write $a^t = a_{\text{anis}}^t a_{\text{split}}^t$ where $a_{\text{anis}}^t \in A''$ and $a_{\text{split}}^t \in A'$.

- (12) We let Γ act by homeomorphisms on X_0 . We assume this action lifts to a cocycle $\mathcal{A}: \Gamma \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$ on a finite-dimensional normed vector bundle \mathcal{E}_0 over X_0 .
- (13) We let $X := (G \times X_0)/\Gamma$ be the suspension space with the G -action induced from the Γ -action on X_0 . The space X has the structure of a fiber bundle over G/Γ and the fibers are homeomorphic to X_0 .
- (14) Write $\pi_{\mathcal{E}}: \mathcal{E} \rightarrow X$ for the induced vector bundle $\mathcal{E} = (G \times \mathcal{E}_0)/\Gamma$ induced by $\mathcal{E}_0 \rightarrow X_0$. For any closed subgroup H of G , let $\mathcal{A}: H \times \mathcal{E}_H \rightarrow \mathcal{E}_H$ be the restriction of the cocycle to $(H \times X_0)/\Gamma_H$. We equip with \mathcal{E} with a norm as constructed in Section 5.4.

- (15) In the setting of Theorem D, we have $X_0 = M$, \mathcal{E} is the fiberwise tangent bundle of $X = (G \times M)/\Gamma$, and $\mathcal{A}: \Gamma \times TM \rightarrow TM$ is the cocycle induced by the derivative cocycle $\mathcal{A}_0(\gamma, x) = D_x\alpha(\gamma)$.

7. SUBEXPONENTIAL GROWTH FOR THE CENTER

We begin the proof of Theorems D and 6.3. In the following simple proposition we assume the failure of subexponential growth occurs for the restriction $\mathcal{A}_0: \mathcal{Z} \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$ of the cocycle \mathcal{A}_0 to the center \mathcal{Z} of G . In the sequel, when considering the restriction of the cocycle to other subgroups of Γ we will quote the proposition to assume the center only contributes subexponential growth to the cocycle.

Proposition 7.1. *Suppose the restriction*

$$\mathcal{A}_0: \mathcal{Z} \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$$

fails to have len_Γ -subexponential growth. Then there exists a split Cartan subgroup A of G and a $(\mathcal{Z}A)$ -invariant Borel probability measure μ on X projecting to the Haar measure on G/Γ such that $\lambda_{\text{top},a,\mu,A} > 0$ for some $a \in \mathcal{Z}$.

Proof. Since $\mathcal{Z} \subset \Gamma$, the induced action of \mathcal{Z} on G/Γ is the identity action. It follows from Lemma 6.8 that there is a Borel probability measure μ_0 on X projecting to the delta mass $\delta_{\mathbf{1}\Gamma}$ on G/Γ such that $\lambda_{\text{top},a_0,\mu_0,A} > 0$ for some $a_0 \in \mathcal{Z}$.

Fix any split Cartan subgroup $A \subset G$ and a one-parameter subgroup $\{b^s\}$ in A . Let $U = \mathcal{U}^+(b^s)$ be the expanding horospherical subgroup associated with $\{b^s\}$. Note that A and U trivially commute with \mathcal{Z} . Averaging μ_0 over an appropriate Følner sequence in U , by Lemmas 4.8 and 6.5 we obtain a $(\mathcal{Z}U)$ -invariant, Borel probability measure μ_1 on X with exponentially small mass at ∞ and $\lambda_{\text{top},a_0,\mu_1,A} > 0$. Since μ_0 is supported over the identity coset $\mathbf{1}\Gamma$ in G/Γ , μ_1 is compactly supported. By Corollary 4.4 and Lemma 6.5 the sequence of measures $\{(b^s)_*\mu_1 : s \in \mathbb{N}\}$ is uniformly tight; moreover, any subsequential limit μ_2 of this sequence is \mathcal{Z} -invariant, projects to the Haar measure on G/Γ , and satisfies $\lambda_{\text{top},a_0,\mu_2,A} > 0$. Finally, averaging μ_2 against an appropriate Følner sequence in A , by Lemma 6.5 we obtain a $(\mathcal{Z}A)$ -invariant, Borel probability measure μ on X projecting to the Haar measure on G/Γ with $\lambda_{\text{top},a_0,\mu,A} > 0$. \square

7.1. Additional hypothesis for Sections 8, 9, and 10. Using Proposition 7.1, we will assume in Sections 8, 9, and 10 below the following additional hypothesis:

Assumption 7.2. The restriction

$$\mathcal{A}_0: \mathcal{Z} \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$$

of the cocycle \mathcal{A}_0 to the center \mathcal{Z} has len_Γ -subexponential growth.

8. SUBEXPONENTIAL GROWTH FOR UNIPOTENT SUBGROUPS

In this section, we assume that the failure of subexponential growth occurs for the restriction of the action to some unipotent \mathbb{Q} -subgroup of Γ . In Sections 9 and 10, we then consider situations where the failure of subexponential growth occurs for the actions of certain other subgroups of Γ .

Under the additional hypothesis in Assumption 7.2 we have the following.

Proposition 8.1. *If there is a connected unipotent \mathbb{Q} -subgroup U of G such that the restriction*

$$\mathcal{A}_0: \Gamma_U \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$$

does not have len_Γ -subexponential growth then there is a split Cartan subgroup A of G and an A -invariant Borel probability measure μ on X projecting to the Haar measure on G/Γ such that $\lambda_{\text{top},a,\mu,A} > 0$ for some $a \in A$.

The remainder of this section is devoted to the proof of this proposition. We will prove the contrapositive of Proposition 8.1; in particular, we will assume the following:

Assumption 8.2. For every split Cartan subgroup A of G , every A -invariant Borel probability measure μ on X that projects to the Haar measure on G/Γ , and every $a \in A$,

$$\lambda_{\text{top},a,\mu,A} = 0.$$

We recall the subadditive cocycles $\mathcal{B}^\pm: G \times G/\Gamma \rightarrow \mathbb{R}$ defined by (5.8) and (5.9) in Section 5.5. Under Assumption 8.2, we will show that the restrictions of \mathcal{B}^\pm to $U \times U/\Gamma_U$ are sublinear: for every $\varepsilon > 0$, there is C_ε such that for all $g \in U$ and $x \in U/\Gamma_U$,

$$\mathcal{B}^\pm(g, x) \leq \varepsilon d(g, \mathbf{1}) + C_\varepsilon.$$

Since Γ_U is cocompact in U , by Corollary 5.13 and Theorem 3.22, this is equivalent to the following: for every $\varepsilon > 0$ there is C_ε such that for all $\gamma \in \Gamma_U$,

$$\mathcal{B}^\pm(\gamma, \mathbf{1}\Gamma) \leq \varepsilon d(\gamma, \mathbf{1}) + C_\varepsilon.$$

To start the proof of Proposition 8.1, we consider various 1-parameter subgroups $\{a^t\}$ that normalize U . For such $\{a^t\}$, we let $U^+ \subset U$ be the expanding subgroup of U . We will show, roughly speaking, that the restriction of \mathcal{B}^\pm to a subgroup U^+ of U has sublinear growth and then deduce sublinear growth for all of U from this. However, there is a technical issue that we will actually not see immediately that the restriction of \mathcal{B}^\pm to U^+ itself has sublinear growth, but only that each element of U^+ has the same image in $U/[U, U]$ as some element with sublinear growth.

Sublinear growth of the cocycle for such representatives of elements of U^+ in $U/[U, U]$ will be deduced from sublinear growth for generic trajectories for the 1-parameter subgroup $\{a^t\}$. The basic idea is simply that if $\{a^t\}$ has sublinear growth and if u_0 is a fixed element of U^+ , then $u := a^t u_0 a^{-t}$ is a large element of U whose norm for the associated cocycle is sublinear in $d(u, \mathbf{1})$. From Corollary 3.9, by varying $\{a^t\}$ we can then generate all of $U/[U, U]$ by the associated expanding subgroups. A difficulty is that sublinear growth of $\{a^t\}$ only holds along orbits of a full measure subset of base points whereas, for elements of U , we need a uniform estimate that holds at every point of U/Γ_U . To upgrade from almost every point to every point, we work in the abelianization $U/[U, U]$ and use that sumsets of sets with high density in balls cover balls of comparable radii.

In Sections 8.1 and 8.2 we make precise the outline in the preceding paragraphs in the proof of Lemma 8.4. We then use Lemma 8.4 to establish Proposition 8.1 in Section 8.3.

8.1. Subexponential growth for generic $\{a^t\}$ -orbits. Using Lemma 3.26, it is sufficient to consider the case that U is the unipotent radical of some maximal parabolic \mathbb{Q} -subgroup Q . We fix a choice of rational Langlands decomposition of Q ,

$$Q = (L \times S^\circ) \ltimes U.$$

Since Q is a maximal \mathbb{Q} -parabolic, we have $\dim S^\circ = 1$.

To begin the proof of Proposition 8.1, we fix the following notation:

- (1) Let \tilde{A} be a connected, \mathbb{Q} -anisotropic \mathbb{Q} -torus \tilde{A} of L . More precisely, the image of \tilde{A} in F is a connected, \mathbb{Q} -anisotropic \mathbb{Q} -torus.
- (2) Let $\{a^t\}$ be a one-parameter subgroup of \tilde{A} such that $a^1 \in \Gamma_{\tilde{A}}$.
- (3) Let $a_{\mathbb{Z}} := \{a^n \mid n \in \mathbb{Z}\}$. Then $\Lambda := a_{\mathbb{Z}}\Gamma_U$ is a cocompact lattice in $\{a^t\}U$.
- (4) We write $\mathfrak{u}^+, \mathfrak{u}^0, \mathfrak{u}^-$ for the expanding, centralized, and contracting directions of the action of $\text{Ad}(a_{\text{split}}^1)$ on \mathfrak{u} . More precisely, $\mathfrak{u}^+, \mathfrak{u}^0, \mathfrak{u}^-$ are the sum of the eigenspaces corresponding to eigenvalues greater than 1, equal to 1, and less than 1, respectively.
- (5) Each of $\mathfrak{u}^+, \mathfrak{u}^0, \mathfrak{u}^-$ is a Lie subalgebra of \mathfrak{g} . Let U^+, U^0 , and U^- denote the corresponding analytic subgroups of U . Note that:
 - (a) $U^0 = C_U(a_{\text{split}}^t)$ is the centralizer of $\{a_{\text{split}}^t\}$ in U ,
 - (b) each of the three subgroups is normalized by both \tilde{A} and U^0 (since \tilde{A} and U^0 each centralize $\{a_{\text{split}}^t\}$ and normalize U), and
 - (c) $U^-U^0U^+ = U$.

From Corollary 3.9 we may choose several one-parameter subgroups $\{a^t\}$ as above, such that the corresponding expanding horospherical subgroups U^+ generate $U/[U, U]$. In particular, we will always consider $\{a^t\}$ as above such that a_{split}^1 is non-trivial.

Lemma 8.3. *Suppose Assumptions 7.2 and 8.2 hold and let $\{a^t\}$ be as above. Then for m_U -almost every $u \in U$ we have*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathcal{B}^\pm(a^n, u \Gamma_U) = 0. \quad (8.1)$$

Proof. Let U^\pm denote the smallest \mathbb{Q} -subgroup of U that contains both U^+ and U^- and let $A'/\Gamma_{A'}$ be the closure of the image of $\{a_{\text{split}}^t\}$ in $\tilde{A}/\Gamma_{\tilde{A}}$; that is, A' is the smallest \mathbb{Q} -torus of \tilde{A} containing $\{a_{\text{split}}^t\}$. We remark that A' is a connected closed subgroup of \tilde{A} and that $\Gamma_{A'U}$ is cocompact in $A'U$. Let $\widehat{\cdot}: A'U \rightarrow A'U/\Gamma_{A'U}$ denote the natural quotient map. Then $\{a_{\text{split}}^t\}$ acts ergodically with respect to the Haar measure on $A'U^\pm/\Gamma_{A'U^\pm}$; see for example [13, Theorem 6.1]. Since U^0 centralizes $\{a_{\text{split}}^t\}$ and since $U^0U^\pm = U$, almost every $\{a_{\text{split}}^t\}$ -orbit in $\widehat{A'U}$ equidistributes to the Haar measure on some translate $\widehat{u_0 A'U^\pm}$ of $A'U^\pm/\Gamma_{A'U^\pm}$ in $\widehat{A'U}$ for some $u_0 \in U^0$.

Let \mathcal{G}_0 denote the set of points in $\widehat{A'U}$ whose $\{a_{\text{split}}^t\}$ -orbit equidistributes to the Haar measure on such a translate. Let $\mathcal{G} = \{u \in U \mid u\Gamma \in \mathcal{G}_0\}$. As discussed above, \mathcal{G}_0 has full measure in $\widehat{A'U}$. Since A' centralizes $\{a_{\text{split}}^t\}$ and normalizes U and U^\pm , the subset \mathcal{G}_0 is A' -invariant which implies that \mathcal{G} is conull in U .

To complete the proof, we show that (8.1) holds for $u \in \mathcal{G}$. Suppose that (8.1) fails for some $u \in \mathcal{G}$. We note that $\{a_{\text{anis}}^t\}\mathcal{Z}$ is contained in a compact subgroup of $\text{Ad}(G) = G/\mathcal{Z}$. The torus \tilde{A} is quasi-geodesically embedded in G and \mathcal{Z} is quasi-isometrically embedded in G . The hypothesis in Assumption 7.2 then implies for

any $\varepsilon > 0$ there is C such that for all x in the compact set \widehat{AU} ,

$$\mathcal{B}^\pm(a_{\text{anis}}^n, x) \leq C + n\varepsilon.$$

Since the $\{a_{\text{split}}^t\}$ -orbit of $u\Gamma$ is contained in the compact set $u_0\widehat{A'U^\pm} \subset \widehat{A'U}$ for some $u_0 \in U^0$, we have that

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \mathcal{B}^\pm(a_{\text{split}}^n, u\Gamma) > 0.$$

Applying Lemma 6.9 to the subgroup $\{a_{\text{split}}^t\}$, there is an $\{a_{\text{split}}^t\}$ -invariant Borel probability measure ν supported on X which projects to the Haar measure on the translate $u_0\widehat{A'U^\pm}$, such that $\lambda_{\text{top}, a_{\text{split}}^1, \nu, \mathcal{A}} > 0$. We remark that since U^0 normalizes U^\pm , the projection $\widehat{\nu}$ of ν to G/Γ is U^\pm -invariant.

Let $\{F_j^0\}$ be a Følner sequence of precompact sets in U^0 . Observe for each j that $F_j^0 * \nu$ is supported on the subset of X projecting to the compact set $\widehat{U^0 A'U^\pm} = \widehat{A'U}$. Let ν^0 be a subsequential limit of $\{F_j^0 * \nu\}$. Then ν^0 is U^0 -invariant. In addition, since U^0 normalizes both $\{a_{\text{split}}^t\}$ and U^\pm , the image $\widehat{\nu}_0$ of ν_0 under the projection $X \rightarrow G/\Gamma$ remains invariant under both $\{a_{\text{split}}^t\}$ and U^\pm . In particular, $\widehat{\nu}^0$ is invariant under $\{a_{\text{split}}^t\}U^0U^\pm = \{a_{\text{split}}^t\}U$. Furthermore, by Lemma 6.5 we have $\lambda_{\text{top}, a_{\text{split}}^1, \nu^0, \mathcal{A}} > 0$.

Recall that U is the unipotent radical of a parabolic \mathbb{Q} -subgroup Q and $S = \{b^s\}$ is a 1-parameter subgroup. Also note that S centralizes $\{a_{\text{split}}^1\}$ and that $U = \mathcal{U}^+(b^s)$ is the expanding horospherical subgroup for $\{b^s\}$. Applying Corollary 4.4 and Lemma 6.5 to the sequence $\{(b^s)_* \nu_0 : s \in \mathbb{N}\}$ and applying Corollary 6.6 to any subsequential limit point ν_∞ of $\{(b^s)_* \nu_0 : s \in \mathbb{N}\}$, we obtain a Borel probability measure μ on X whose properties violate Assumption 8.2. \square

8.2. Subexponential growth for representatives of hyperbolic directions.

In the remainder of this section, we write $\widehat{\cdot}: U \rightarrow U/[U, U]$ to denote the natural homomorphism from U to its abelianization.

We equip U with a right-invariant metric d_U . This induces a metric on $\widehat{U} = U/[U, U]$. As U is a unipotent subgroup of G there is a constant $C_d > 1$ such that for all $u \in U$ with $d(u, \mathbf{1}) > 1$, we have

$$\frac{1}{C_d} \log d_U(u, \mathbf{1}) \leq d(u, \mathbf{1}) \leq C_d \log d_U(u, \mathbf{1}).$$

By definition, $U \rightarrow \widehat{U}$ is distance non-increasing.

The main lemma used to establish the inequality in (8.4) is the following:

Lemma 8.4. *There exists a constant $C_0 > 0$ such that for every $\varepsilon > 0$ there is C_ε such that for every $x \in U/\Gamma_U$ and every $u \in U$ such that $\widehat{u} \in \widehat{U}^+$ there exists $v \in U$ such that*

- (a) $\widehat{v} = \widehat{u}$,
- (b) $\mathcal{B}^\pm(v, x) \leq \varepsilon d(u, \mathbf{1}) + C_\varepsilon$, and
- (c) $d(v, \mathbf{1}) \leq C_0 d(u, \mathbf{1}) + C_0$.

To prepare for the proof of this fundamental lemma we establish some additional notation. For $u \in U$, write $u = wu^+$ where $u^+ \in U^+$ and $w \in U^-U^0$; let $p^+: U \rightarrow U^+$ be the natural map $u \mapsto u^+$. Similarly, write $\widehat{u} = \widehat{w}\widehat{u}^+$ where $\widehat{u}^+ \in \widehat{U}^+$ and $w \in \widehat{U^-U^0}$ and let $\widehat{p}^+: \widehat{U} \rightarrow \widehat{U}^+$ be the map $u \mapsto \widehat{u}^+$. Observe \widehat{p}^+ is a

homomorphism. For a subset $E \subset U$, let E^+ and \widehat{E}^+ denote, respectively, the image of E under the maps p^+ and \widehat{p}^+ . Recall that m_E and $m_{\widehat{E}^+}$ denote the Haar measures on U and \widehat{U}^+ normalized to give unit mass to E and \widehat{E}^+ , respectively.

Claim 8.5. *There is a two-sided Følner sequence $\{F_k\}$ in U with the following properties:*

(1) *For every Borel subset E of F_k and every $n \in \mathbb{N}$,*

$$m_{a^n \widehat{F}_k^+ a^{-n}}(a^n \widehat{E}^+ a^{-n}) \geq m_{F_k}(E).$$

(2) *Identifying the abelian group \widehat{U}^+ with some Euclidean space \mathbb{R}^ℓ , we may equip \widehat{U}^+ with a norm which is invariant under conjugation by $\{a^t_{\text{anis}}\}$ and for which \widehat{F}_k^+ is the ball of radius k centered at the origin.*

Proof. Identifying U and $\{a^t\}$ with their images under the adjoint representation, equip \widehat{U}^+ with a norm invariant under conjugation by $\{a^t_{\text{anis}}\}$. Let \widehat{B}_k be the ball of radius k centered at the origin in \widehat{U}^+ , let $U_0 = \ker \widehat{p}^+$. For each k , let B_k be a precompact Borel cross-section of \widehat{p}^+ over \widehat{B}_k ; in particular, $\widehat{p}^+(B_k) = \widehat{B}_k$ and $vU_0 \cap B_k = \{v\}$ for all $v \in B_k$. We may assume $1 \in B_k$. Note that $\{B_k\}$ is a two-sided Følner sequence in the abelian group \widehat{U}^+ . We then construct a two-sided Følner sequence $\{F_k\}$ in U , such that

$$\widehat{F}_k^+ = \widehat{B}_k \text{ and } F_k = B_k(F_k \cap U_0),$$

by induction on the nilpotency degree of U and the following observation: if Z denotes the center of a locally compact group H , if $\{F'_n\}$ is a Følner sequence in Z , and if $\{F''_k\}$ is a sequence of precompact subsets of H such that the image of $\{F''_k\}$ in H/Z is a two-sided Følner sequence, then $\{F''_k \cdot F'_{n_k}\}$ is a two-sided Følner sequence in H , provided that $n_k \rightarrow \infty$ sufficiently rapidly.

With a sequence $\{F_k\}$ constructed above, (2) holds by definition.

To show (1), we let $p: F_k \rightarrow \widehat{F}_k^+$ be the natural map. Then for each $v \in B_k$,

$$p^{-1}(\widehat{v}^+) \cap F_k = v(F_k \cap U_0),$$

so all fibers of p over \widehat{F}_k^+ have the same measure with respect to m_{U_0} , namely $m_{U_0}(F_k \cap U_0)$. This implies that the Radon-Nikodym derivative $d(p_* m_{F_k})/dm_{\widehat{F}_k^+}$ is constant on \widehat{F}_k^+ . Since both measures are normalized, the Radon-Nikodym derivative is 1. For every $E \subseteq F_k$, this implies

$$m_{\widehat{F}_k^+}(\widehat{E}^+) = (p_* m_{F_k})(\widehat{E}^+) = m_{F_k}(p^{-1}(\widehat{E}^+) \cap F_k) \geq m_{F_k}(E),$$

where the last inequality follows as $p^{-1}(\widehat{E}^+) \cap F_k \supseteq E$. Conjugation by a^n is a group automorphism and therefore does not change the relative measure of a subset within an ambient set, so

$$m_{a^n \widehat{F}_k^+ a^{-n}}(a^n \widehat{E}^+ a^{-n}) = m_{\widehat{F}_k^+}(\widehat{E}^+). \quad \square$$

To establish Lemma 8.4, we first show that if k and n are large, then a large proportion of the elements of $a^n \widehat{F}_k^+ a^{-n}$ have the properties specified in the lemma. We begin with the following elementary claim.

Lemma 8.6. *Let m_{U/Γ_U} denote the normalized (left-)Haar measure on U/Γ_U . Let $\rho: U \rightarrow U/\Gamma_U$ be the canonical projection and let $E \subset U$ be a right- Γ_U -invariant set such that*

$$m_{U/\Gamma_U}(\rho(E)) \geq 1 - \delta.$$

Let $\{F_k\}$ be a left-Følner sequence in U . Then for all sufficiently large k ,

$$m_{F_k}(E) \geq 1 - 2\delta.$$

Proof. Let $m_k := \rho_*(m_{F_k})$. Note that each m_k is absolutely continuous with respect to m_{U/Γ_U} . We claim that m_k converges to m_{U/Γ_U} in the strong topology: for any (Borel or Lebesgue) measurable $E \subset U/\Gamma_U$,

$$m_k(E) \rightarrow m_{U/\Gamma_U}(E).$$

For the claim, fix a symmetric precompact neighborhood O of the identity in U . By left-Følnerness of F_k , for any $\delta > 0$ there is k_0 such that for all $k \geq k_0$,

$$\sup_{g \in O} \frac{m_U(g \cdot F_k \Delta F_k)}{m_U(F_k)} \leq \delta.$$

This implies for $k \geq k_0$ that

$$\|g_* m_k - m_k\|_v \leq \delta$$

where $\|\cdot\|_v$ denotes the total variation norm on signed Borel measures.

Note that $\{m_k\}$ has a weak-* convergent subsequence. Any weak-* limit point of $\{m_k\}$ is U -invariant and hence we conclude that $m_k \rightarrow m_{U/\Gamma_U}$ in the weak-* topology. We claim this convergence also holds in the strong topology. Indeed, let ϕ be a non-negative C^∞ function supported in O with $\int \phi(g) dg = 1$. Given a measurable $E \subset U/\Gamma_U$, let $h = \mathbb{1}_E$. The convolution $\phi * h$ is C^0 hence

$$\int \phi * h dm_k \rightarrow \int \phi * h dm_{U/\Gamma_U}.$$

We note for any probability measure μ on U/Γ_U that

$$\begin{aligned} \int (\phi * h)(x) d\mu(x) &:= \int \int \phi(g) h(g^{-1} \cdot x) dg d\mu(x) \\ &= \int \int \phi(g) h(g^{-1} \cdot x) d\mu(x) dg \\ &= \int \int \phi(g) h(x) d(g_*^{-1} \mu)(x) dg. \end{aligned}$$

In particular, by left-invariance of m_{U/Γ_U} ,

$$\int (\phi * h)(x) dm_{U/\Gamma_U}(x) = \int h dm_{U/\Gamma_U}.$$

On the other hand,

$$\begin{aligned}
& \left| \int \phi * h \, dm_k - \int h \, dm_k \right| \\
&= \left| \int \int \phi(g)h(x) \, d(g_*^{-1}m_k)(x) \, dg - \int h(x) \, dm_k(x) \int \phi(g) \, dg \right| \\
&\leq \int \phi(g) \left| \int h(x) \, d(g_*^{-1}m_k)(x) - \int h(x) \, dm_k(x) \right| \, dg \\
&\leq \int \phi(g) \|g_*^{-1}m_k - m_k\|_v \, dg \\
&\leq \delta.
\end{aligned}$$

The claim then follows. The lemma follows as Γ_U -invariance of E implies $m_{F_k}(E) = m_k(\rho(E))$. \square

Lemma 8.7. *For every $\varepsilon > 0$ and $\delta > 0$, there exists $k \in \mathbb{N}$ and $C_{\mathcal{G}} > 0$ such that for all sufficiently large n there exists a set $G_n \subset \widehat{U}^+$ such that:*

- (a) $G_n \subset a^n \widehat{F}_k^+ a^{-n}$;
- (b) $m_{a^n \widehat{F}_k^+ a^{-n}}(G_n) \geq 1 - \delta$;
- (c) for every $\widehat{u} \in G_n$ and every $x \in U/\Gamma_U$ there is $v \in U$ such that
 - (i) $\widehat{v} = \widehat{u}$;
 - (ii) $\mathcal{B}^\pm(v, x) \leq \varepsilon n$;
 - (iii) $d(v, \mathbf{1}) \leq C_{\mathcal{G}} n$.

Proof. Fix ε and δ . For $n_0 \in \mathbb{N}$, set

$$\mathcal{G}(n_0) := \{u \in U \mid \mathcal{B}^\pm(a^{\pm n}, u\Gamma_U) \leq n\varepsilon \text{ for all } n \geq n_0\}.$$

By Lemma 8.3, for almost every $u \in U$ there exists n_0 such that $u \in \mathcal{G}(n_0)$. Observe also that $\mathcal{G}(n_0)\Gamma_U = \mathcal{G}(n_0)$. Taking n_0 sufficiently large, we have

$$m_{U/\Gamma_U}(\mathcal{G}(n_0)/\Gamma_U) \geq 1 - \delta/2.$$

Let \mathcal{F}_U be a compact, symmetric set that contains fundamental domains for both the right and left action of Γ_U on U . Fix a reference point $u_0 \in \mathcal{G}(n_0) \cap \mathcal{F}_U$. Consider $k \in \mathbb{N}$ and $n \geq n_0$. Set

$$\mathcal{G}_n := (\mathcal{G}(n_0) a^{-n} u_0^{-1} a^n) \cap F_k.$$

Since $\mathcal{G}(n_0)\Gamma_U = \mathcal{G}(n_0)$ and $\Gamma_U \mathcal{F}_U = U$, we have $\mathcal{G}(n_0) a^{-n} u_0^{-1} a^n = \mathcal{G}(n_0) v_0$ for some $v_0 \in \mathcal{F}_U$. Then

$$m_U(\mathcal{G}_n) = m_U(\mathcal{G}(n_0) \cap (F_k v_0^{-1})).$$

From the lower bound on $m_{U/\Gamma_U}(\mathcal{G}(n_0)/\Gamma_U)$, the fact that $\{F_k\}$ is a two-sided Følner sequence, Lemma 8.6, and using that $v_0 \in \mathcal{F}_U$ is bounded, we conclude that if k is sufficiently large, then for all $n \geq n_0$,

$$m_{F_k}(\mathcal{G}_n) \geq 1 - \delta.$$

Fix such $k \in \mathbb{N}$ for the remainder.

Take the set G_n in the lemma to be the set

$$G_n := a^n \widehat{\mathcal{G}}_n^+ a^{-n}.$$

Since $\mathcal{G}_n \subseteq F_k$, conclusion (a) of the lemma follows from definition.

From Claim 8.5(1) we have

$$m_{a^n \widehat{F}_k^+ a^{-n}}(G_n) \geq m_{F_k}(\mathcal{G}_n) \geq 1 - \delta.$$

This establishes (b).

Consider an arbitrary element of G_n ; by definition such an element is of the form $a^n \widehat{u}^+ a^{-n}$ for some choice of $u \in \mathcal{G}_n \subset F_k$. Given $x \in U/\Gamma_U$, select $u_x \in \mathcal{F}_U$ with $u_x \Gamma_U = x$. Set

$$v = u_x u_0^{-1} a^n u^+ a^{-n} u_0 u_x^{-1}. \quad (8.2)$$

We claim that v satisfies the conclusions in part (c) of the statement of the lemma. First note that, since \widehat{U} is abelian, we have $\widehat{v} = a^n \widehat{u}^+ a^{-n} = a^n \widehat{u}^+ a^{-n}$. Moreover, it is clear that $v \in (\mathcal{F}_U)^2 (a^n F_k^+ a^{-n}) (\mathcal{F}_U)^2$; since \mathcal{F}_U and F_k are precompact, this implies there are constants C_1 and $C_{\mathcal{G}}$ such that

$$d(v, \mathbf{1}) \leq 2C_1 + n d(a, \mathbf{1}) + C_1 + n d(a, \mathbf{1}) + 2C_1 \leq C_{\mathcal{G}} n.$$

This establishes (i) and (iii)

For (ii), recall that $a^1 \in \Gamma$ so $a^n \Gamma = \Gamma$. We then have

$$\begin{aligned} u a^{-n} u_0 \Gamma &= u a^{-n} u_0 a^n \Gamma \in \mathcal{G}_n a^{-n} u_0 a^n \Gamma \\ &\subseteq (\mathcal{G}(n_0) a^{-n} u_0^{-1} a^n) a^{-n} u_0 a^n \Gamma \\ &= \mathcal{G}(n_0). \end{aligned} \quad (8.3)$$

Since u is in the precompact set F_k , we may write $u^+ = wu$, where $w \in U^{-}U^0$ and the norm of w is uniformly bounded over the choice of $u \in \mathcal{G}_n$. Then we have $a^n u^+ a^{-n} = w_n (a^n u a^{-n})$ where the norm of $w_n := a^n w a^{-n}$ is bounded uniformly over the choice of $n \in \mathbb{N}$ and $u \in \mathcal{G}_n$. Since u_x and u_0 are in the compact set \mathcal{F}_U , the element v in (8.2) is at bounded distance (uniform in the choice of n and u) from $u_x u_0^{-1} a^n u a^{-n} u_0 u_x^{-1}$, respectively. Since the homogeneous space $\{a^t : t \in \mathbb{R}\}U/(\{a^n : n \in \mathbb{Z}\}\Gamma_U)$ is compact there is a constant C (depending only on the choice of k and \mathcal{F}_U) such that

$$\begin{aligned} \mathcal{B}^{\pm}(v, x) &\leq C + \mathcal{B}^{\pm}(u_x u_0^{-1} a^n u a^{-n} u_0 u_x^{-1}, x) \\ &\leq C + \mathcal{B}^{\pm}(u_x u_0^{-1}, a^n u a^{-n} u_0 \Gamma) + \mathcal{B}^{\pm}(a^n, u a^{-n} u_0 \Gamma) \\ &\quad + \mathcal{B}^{\pm}(u, a^{-n} u_0 \Gamma) + \mathcal{B}^{\pm}(a^{-n}, u_0 \Gamma) + \mathcal{B}^{\pm}(u_0 u_x^{-1}, x) \\ &\leq C + C + \varepsilon n + C + \varepsilon n + C \\ &\leq 3\varepsilon n \end{aligned}$$

for all n sufficiently large. \square

We now prove Lemma 8.4.

Proof of Lemma 8.4. Fix $\varepsilon > 0$. As in Claim 8.5(2), we identify \widehat{U}^+ with some $(\mathbb{R}^{\ell}, +)$. Let $0 < \delta < 2^{-(\ell+1)}$ and let k be as in Lemma 8.7 for this choice of δ and ε .

Consider any $\widehat{u} \in \widehat{U}^+$. Recall that \widehat{F}_k^+ is the ball \widehat{B}_k of radius k centered at the origin and that conjugation by a expands \widehat{U}^+ . Take the minimal n such that $2\widehat{u} \in a^n \widehat{F}_k^+ a^{-n}$. We may assume n is sufficiently large so that Lemma 8.7 holds. Let G_n be as in Lemma 8.7.

Write $\tilde{F} := a^{-n} \widehat{F}_k^+ a^n$. Recall \widehat{U}^+ is equipped with a norm so that \widehat{F}_k^+ is a ball centered at the origin. Then $\frac{1}{2}\tilde{F} \subseteq \tilde{F}$ and $m_{\tilde{F}}(\frac{1}{2}\tilde{F}) = 2^{-\ell} > 2\delta$. Since $m_{\tilde{F}}(G_n) > 1 - \delta$, this implies

$$m_{\frac{1}{2}\tilde{F}}(G_n \cap \frac{1}{2}\tilde{F}) > \frac{1}{2}.$$

Since $\widehat{u} \in \frac{1}{2}\tilde{F}$ we also have $\widehat{u} - \frac{1}{2}\tilde{F} \subseteq \tilde{F}$. By the same argument,

$$m_{\frac{1}{2}\tilde{F}}((\widehat{u} - G_n) \cap \frac{1}{2}\tilde{F}) > \frac{1}{2}.$$

In particular, there is some $\widehat{u}' \in \frac{1}{2}\tilde{F} \cap G_n \cap (\widehat{u} - G_n)$. Then $\widehat{u} - \widehat{u}' \in G_n$ and

$$\widehat{u} = \widehat{u}' + (\widehat{u} - \widehat{u}') \in G_n + G_n.$$

In particular, we may write any $\widehat{u} = \widehat{u}_1 + \widehat{u}_2$ with $\widehat{u}_i \in G_n$.

Now consider any $x \in U/\Gamma_U$. From Lemma 8.7(c), there exist $v_1, v_2 \in U$, such that $\widehat{v}_i = \widehat{u}_i$, $\mathcal{B}^\pm(v_2, x) \leq \varepsilon n$, $\mathcal{B}^\pm(v_1, v_2 x) \leq \varepsilon n$, and $d(v_i, \mathbf{1}) \leq C_G n$. Let $v = v_1 v_2$. Then $\widehat{v} = \widehat{v}_1 + \widehat{v}_2 = \widehat{u}_1 + \widehat{u}_2 = \widehat{u}$.

Let λ be the smallest eigenvalue of a_{split}^1 on \widehat{U}^+ . Since the norm $\|\cdot\|$ on \widehat{U}^+ is chosen to be invariant under a_{anis}^t , $a^i \widehat{F}_k^+ a^{-i}$ contains the ball of radius $\lambda^i k$; minimality of n then implies $n \leq \log_\lambda 2\|\widehat{u}\|$. The map $U \rightarrow \widehat{U}^+$ is a polynomial so there is a constant C such that $\log_\lambda 2\|\widehat{u}\| \leq Cd(u, \mathbf{1})$ if $d(u, \mathbf{1}) > 1$. Then $n \leq Cd(u, \mathbf{1})$ and we have

$$\mathcal{B}^\pm(v, x) \leq \mathcal{B}^\pm(v_1, v_2 x) + \mathcal{B}^\pm(v_2, x) \leq \varepsilon n + \varepsilon n \leq 2\varepsilon C d(u, \mathbf{1})$$

and

$$d(v, \mathbf{1}) \leq d(v_1, \mathbf{1}) + d(v_2, \mathbf{1}) \leq 2C_G n \leq 2C_G C d(u, \mathbf{1}). \quad \square$$

8.3. Proof of Proposition 8.1. We complete the proof of Proposition 8.1. It suffices to show under Assumption 8.2 that for every $\varepsilon > 0$ there is C_ε such that for all $\gamma \in \Gamma_U$,

$$\mathcal{B}^\pm(\gamma, \mathbf{1}\Gamma) \leq \varepsilon d(\gamma, \mathbf{1}) + C_\varepsilon. \quad (8.4)$$

We will first use Lemma 8.4 to show that each element of $\Gamma_U/[\Gamma_U, \Gamma_U]$ has a representative in Γ_U for which \mathcal{B}^+ and \mathcal{B}^- have sublinear size when evaluated at the coset $\mathbf{1}\Gamma$. This is the base case of an induction that establishes the same conclusion for every quotient of the descending central series of U . To finish, we will observe that these estimates on the quotients easily imply the desired sublinear growth of \mathcal{B}^+ and \mathcal{B}^- evaluated at $\Gamma_U \times \{\mathbf{1}\Gamma\}$.

Claim 8.8. *There exists $C > 0$ such that for every $\varepsilon > 0$ there is $C_\varepsilon > 0$ such that for each $u \in U$ and $x \in U/\Gamma_U$ there exists $v \in U$ such that*

- (1) $\widehat{v} = \widehat{u}$,
- (2) $\mathcal{B}^\pm(v, x) < \varepsilon d(u, \mathbf{1}) + C_\varepsilon$, and
- (3) $d(v, \mathbf{1}) < C d(u, \mathbf{1}) + C$.

Proof. Corollary 3.9 provides finitely many one-parameter subgroups a_1^t, \dots, a_r^t of L , such that the corresponding expanding subgroups $\widehat{U}_1^+, \dots, \widehat{U}_r^+$ span \widehat{U} . Identify \widehat{U} with some \mathbb{R}^s . Equipping \widehat{U} with some norm $\|\cdot\|$, there is $C_{\widehat{U}}$ such that we may write $\widehat{u} = \widehat{u}_1 \widehat{u}_2 \cdots \widehat{u}_r$ where $\widehat{u}_i \in \widehat{U}_i^+$ and $\|\widehat{u}_i\| \leq C_{\widehat{U}} \|\widehat{u}\|$. Identifying U with its image under the adjoint representation, the exponential map on U is a

polynomial (see Lemma 3.1(1)) and $d(u, \mathbf{1})$ is comparable to $\log \|u\|$ for all sufficiently large $u \in U$. This implies that we may choose a lift u_i of each \widehat{u}_i such that $d(u_i, \mathbf{1}) \leq C_U d(u, \mathbf{1}) + C_U$, for some constant C_U .

Consider any $x := x_0 \in U/\Gamma_U$. Recursively define $\{v_i, 1 \leq i \leq r\}$ and $\{x_i, 1 \leq i \leq r\}$ as follows: By Corollary 3.9(1), each $\{a_i^t\}$ intersects the lattice Γ . We may apply Lemma 8.4 with \widehat{U}_i^+ , u_i , and x_{i-1} to obtain an element v_i . Set $x_i = v_i \cdot x_{i-1}$. Then

$$\widehat{v}_i = \widehat{u}_i, \quad \mathcal{B}^\pm(v_i, x_{i-1}) \leq \varepsilon d(u_i, \mathbf{1}) + C_\varepsilon, \quad \text{and} \quad d(v_i, \mathbf{1}) \leq C_0 d(u_i, \mathbf{1}) + C_0.$$

Let $v = v_1 v_2 \cdots v_r$. Then $\widehat{v} = \widehat{v}_1 \widehat{v}_2 \cdots \widehat{v}_r = \widehat{u}_1 \widehat{u}_2 \cdots \widehat{u}_r = \widehat{u}$. We also have

$$\begin{aligned} \mathcal{B}^\pm(v, x) &= \mathcal{B}^\pm(v_1 v_2 \cdots v_r, x_0) \leq \sum_{i=1}^r \mathcal{B}^\pm(v_i, x_{i-1}) \leq \sum_{i=1}^r (\varepsilon d(u_i, \mathbf{1}) + C_\varepsilon) \\ &\leq r(\varepsilon [C_U d(u, \mathbf{1}) + C_U] + C_\varepsilon) \leq C' \varepsilon d(u, \mathbf{1}) + C' \end{aligned}$$

for an appropriate constant C' , and

$$\begin{aligned} d(v, \mathbf{1}) &= d(v_1 v_2 \cdots v_r, \mathbf{1}) \leq \sum_{i=1}^r d(v_i, \mathbf{1}) \leq \sum_{i=1}^r (C_0 d(u_i, \mathbf{1}) + C_0) \\ &\leq \sum_{i=1}^r (C_0 [C_U d(u, \mathbf{1}) + C_U] + C_0) \leq C d(u, \mathbf{1}) + C. \end{aligned}$$

This completes the proof of the claim. \square

For the restriction of the discrete cocycle $\mathcal{A}_0: \Gamma \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$ to Γ_U , we have the following immediate strengthening of Claim 8.8.

Corollary 8.9. *There exists $C > 0$ such that for every $\varepsilon > 0$ there is $C_\varepsilon > 0$ such that for every $\gamma \in \Gamma_U$ there exists $\lambda \in \Gamma_U$ such that*

- (1) $\lambda = \gamma \pmod{[\Gamma_U, \Gamma_U]}$,
- (2) $\mathcal{B}^\pm(\lambda^{\pm 1}, \mathbf{1}\Gamma) < \varepsilon d(\gamma, \mathbf{1}) + C_\varepsilon$, and
- (3) $d(\lambda, \mathbf{1}) < C d(\gamma, \mathbf{1}) + C$.

Proof. Fix $\gamma \in \Gamma_U$. From Claim 8.8, there is $v \in U$ such that $\widehat{v} = \widehat{\gamma}$, $d(v, \mathbf{1}) < C d(\gamma, \mathbf{1}) + C$, and

$$\mathcal{B}^\pm(v, \mathbf{1}\Gamma) < \varepsilon d(\gamma, \mathbf{1}) + C_\varepsilon.$$

Note that $[\Gamma_U, \Gamma_U]$ is a cocompact lattice in $[U, U]$. Let \widehat{C} denote the diameter of $[U, U]/[\Gamma_U, \Gamma_U]$. Then there is $q \in [U, U]$ such that $qv\gamma^{-1} \in [\Gamma_U, \Gamma_U]$ and $d(q, \mathbf{1}) \leq \widehat{C}$. Then $\lambda := qv \in \Gamma_U$ satisfies $\lambda = \gamma \pmod{[\Gamma_U, \Gamma_U]}$ and $d(\lambda, \mathbf{1}) < C d(\gamma, \mathbf{1}) + C + \widehat{C}$. By compactness of U/Γ_U and using that $d(q, \mathbf{1}) \leq \widehat{C}$ there is some uniform $C' > 0$ such that

$$\begin{aligned} \mathcal{B}^\pm(\lambda, \mathbf{1}\Gamma) &\leq \mathcal{B}^\pm(v, \mathbf{1}\Gamma) + \mathcal{B}^\pm(q, v\Gamma) \\ &\leq \varepsilon d(\gamma, \mathbf{1}) + C_\varepsilon + C'. \end{aligned}$$

Moreover, since

$$\mathcal{B}^\pm(\lambda^{-1}, \mathbf{1}\Gamma) = \mathcal{B}^\mp(\lambda, \lambda \cdot \mathbf{1}\Gamma) = \mathcal{B}^\mp(\lambda, \mathbf{1}\Gamma)$$

we also conclude that $\mathcal{B}^\pm(\lambda^{-1}, \mathbf{1}\Gamma) \leq \varepsilon d(\gamma, \mathbf{1}) + C_\varepsilon + C'$. \square

Let

$$U = U_1 \supset U_2 \supset \cdots \supset U_{c+1} = \{1\}$$

be the descending central series of U where $U_{k+1} = [U, U_k]$. Write $\Lambda = \Gamma_U$ and let

$$\Gamma_U = \Lambda = \Lambda_1 \supset \Lambda_2 \supset \cdots \supset \Lambda_{c+1} = \{1\}$$

be the descending central series of Λ . For each $1 \leq k \leq c$, Λ_k is a cocompact lattice in U_k and has finite index in $\Gamma_{U_k} = \Gamma \cap U_k$.

Claim 8.10. *There exists $C > 0$ such that every $\varepsilon > 0$ there is $C_\varepsilon > 0$ such that for every $1 \leq k \leq c$ and $\gamma \in \Lambda_k$ there exists $\lambda \in \Lambda_k$ such that*

- (1) $\lambda = \gamma \pmod{\Lambda_{k+1}}$,
- (2) $\mathcal{B}^\pm(\lambda, \mathbf{1}\Gamma) < \varepsilon d(\gamma, \mathbf{1}) + C_\varepsilon$ and $\mathcal{B}^\pm(\lambda^{-1}, \mathbf{1}\Gamma) < \varepsilon d(\gamma, \mathbf{1}) + C_\varepsilon$,
- (3) $d(\lambda, \mathbf{1}) < C d(\gamma, \mathbf{1}) + C$.

Proof. The proof is by induction on k . When $k = 1$, the conclusion follows from directly from Corollary 8.9. We now assume $k \geq 2$.

We recall that $\widehat{U} = U_1/U_2$. Each quotient U_k/U_{k+1} is abelian and naturally identified with $\mathfrak{u}_k/\mathfrak{u}_{k+1}$ via the exponential. Moreover, the map

$$(uU_k, wU_2) \mapsto [u, w]U_{k+1}$$

is a well-defined, surjective, bilinear map $(U_{k-1}/U_k) \times \widehat{U} \rightarrow U_k/U_{k+1}$. Fix a finite set of generators $\{w_1, \dots, w_r\}$ for $\Lambda = \Gamma_U$. Then the image $\{\widehat{w}_1, \dots, \widehat{w}_r\}$ contains a basis for \widehat{U} .

For each $k \geq 2$, the collection $\{[w_i, \Lambda_{k-1}] : 1 \leq i \leq r\}$ is a generating set for Λ_k . Moreover, there is a C_0 such that for every $\gamma \in \Lambda_k$ there are $\gamma_1, \dots, \gamma_r \in \Lambda_{k-1}$ with

$$\gamma = [\gamma_1, w_1] [\gamma_2, w_2] \cdots [\gamma_r, w_r]$$

and $d(\gamma_i, \mathbf{1}) \leq C_0 d(\gamma, \mathbf{1}) + C_0$ (cf. Lemma 3.1(1)).

Fix $\gamma \in \Lambda_k$ and write $\gamma = [\gamma_1, w_1] [\gamma_2, w_2] \cdots [\gamma_r, w_r]$ for $\gamma_1, \dots, \gamma_r \in \Lambda_{k-1}$ as above. By the induction hypothesis, there are C_1 and $\lambda_1, \dots, \lambda_r \in \Lambda_{k-1}$ such that

- (1) $\lambda_i = \gamma_i \pmod{\Lambda_k}$
- (2) $d(\lambda_i, \mathbf{1}) \leq C_1 d(\gamma_i, \mathbf{1}) + C_1$, and
- (3) $\mathcal{B}^\pm(\lambda_i^{\pm 1}, \mathbf{1}\Gamma) \leq \varepsilon d(\gamma_i, \mathbf{1}) + C_\varepsilon$.

Also, there is C_2 with $\max\{\mathcal{B}^\pm(w_i^{\pm 1}, \mathbf{1}\Gamma) : 1 \leq i \leq r\} \leq C_2$. Let

$$\lambda = [\lambda_1, w_1] [\lambda_2, w_2] \cdots [\lambda_r, w_r].$$

We check that $[\lambda_i, w_i] = [\gamma_i, w_i] \pmod{\Lambda_{k+1}}$. Indeed, there exists $q \in \Lambda_k$ such that

$$\begin{aligned} [\lambda_i, w_i] &= [\gamma_i q, w_i] \\ &= q^{-1} \gamma_i^{-1} w_i^{-1} \gamma_i q w_i \\ &= q^{-1} [\gamma_i, w_i] w_i^{-1} q w_i \\ &= q^{-1} [\gamma_i, w_i] q \pmod{\Lambda_{k+1}} \\ &= [\gamma_i, w_i] \pmod{\Lambda_{k+1}}. \end{aligned}$$

Thus $\lambda = \gamma \pmod{\Lambda_{k+1}}$. We also have

$$\begin{aligned} \mathcal{B}^\pm(\lambda, \mathbf{1}\Gamma) &\leq \sum_{i=1}^r \mathcal{B}^\pm([\lambda_i, w_i], \mathbf{1}\Gamma) \\ &\leq \sum_{i=1}^r (\mathcal{B}^\pm(\lambda_i^{-1}, \mathbf{1}\Gamma) + \mathcal{B}^\pm(w_i, \mathbf{1}\Gamma) + \mathcal{B}^\pm(\lambda_i, \mathbf{1}\Gamma) + \mathcal{B}^\pm(w_i, \mathbf{1}\Gamma)) \\ &\leq 2r(\varepsilon d(\gamma, \mathbf{1}) + C_\varepsilon) + 2rC_2 \\ &\leq C'\varepsilon d(\gamma, \mathbf{1}) + C'. \end{aligned}$$

Similarly, $\mathcal{B}^\pm(\lambda^{-1}, \mathbf{1}\Gamma) \leq C'\varepsilon d(\gamma, \mathbf{1}) + C'$. Moreover,

$$d(\lambda, \mathbf{1}) \leq \sum_{i=1}^r d(\lambda_i, \mathbf{1}) \leq r(C_1 d(\gamma, \mathbf{1}) + C_1) \leq C'' d(\gamma, \mathbf{1}) + C''.$$

This completes the proof of the claim. \square

We now complete the proof of Proposition 8.1.

Proof of Proposition 8.1. Since $\Lambda_1 = \Gamma_U$, it suffices to show for every $\varepsilon > 0$ and every $1 \leq k \leq c$ that there is $C > 1$ such that for all $\gamma \in \Lambda_k$,

$$\mathcal{B}^\pm(\gamma, \mathbf{1}\Gamma) \leq \varepsilon d(\gamma, \mathbf{1}) + C.$$

The proof is by backward induction on k . The base case when $k = c + 1$ holds trivially.

Consider the inductive case $1 \leq k \leq c$. Fix any $\varepsilon > 0$. Given $\gamma \in \Lambda_k$, take λ as in Claim 8.10. We have $\gamma = \lambda \pmod{\Lambda_{k+1}}$. From subadditivity, Claim 8.10(2), and the induction hypothesis we have

$$\begin{aligned} \mathcal{B}^\pm(\gamma, \mathbf{1}\Gamma) &\leq \mathcal{B}^\pm(\gamma\lambda^{-1}, \mathbf{1}\Gamma) + \mathcal{B}^\pm(\lambda, \mathbf{1}\Gamma) \\ &\leq (\varepsilon d(\gamma\lambda^{-1}, \mathbf{1}) + C'_\varepsilon) + (\varepsilon d(\lambda, \mathbf{1}) + C_\varepsilon). \end{aligned}$$

By Claim 8.10(3), this is bounded above by an expression of the form $C_1\varepsilon d(\gamma, \mathbf{1}) + C_2$ where C_1 is a constant independent of ε . This completes the proof of the inductive case. \square

9. SUBEXPONENTIAL GROWTH FOR MINIMAL PARABOLIC \mathbb{Q} -SUBGROUPS

We establish the following proposition which guarantees the existence of a measure as in the conclusion of Theorems D and 6.3 under the assumption that the failure of subexponential growth occurs for the action of some Γ_P for some minimal parabolic \mathbb{Q} -subgroup $P \subset G$.

Under the additional hypothesis in Assumption 7.2, we have the following.

Proposition 9.1. *Let P be a minimal parabolic \mathbb{Q} -subgroup of G . If the restriction*

$$\mathcal{A}_0|_{\Gamma_P} : \Gamma_P \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$$

does not have len_Γ -subexponential growth then there is a split Cartan subgroup A of G and an A -invariant Borel probability measure μ on X projecting to the Haar measure on G/Γ such that $\lambda_{\text{top}, a, \mu, A} > 0$ for some $a \in A$.

Proof. We have the rational Langlands decomposition $P = (L \times S^\circ) \times U$ where L is a connected reductive \mathbb{Q} -subgroup with $\text{rank}_{\mathbb{Q}} L = 0$, S is a maximal \mathbb{Q} -split torus of P , and $U = \text{Rad}_u(P)$ is the unipotent radical of P . Each of L , S , and U is defined over \mathbb{Q} . We have that L/Γ_L and U/Γ_U are compact, Γ_S is trivial, and, up to finite index, $\Gamma_P = \Gamma_L \times \Gamma_U$.

By Proposition 8.1, the conclusion of Proposition 9.1 would follow if the restriction $\mathcal{A}_0|_{\Gamma_U} : \Gamma_U \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$ failed to have len_Γ -subexponential growth. Thus, we may assume for the remainder that $\mathcal{A}_0|_{\Gamma_U} : \Gamma_U \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$ has len_Γ -subexponential growth.

Recall that the natural embedding $\Gamma_L \times \Gamma_U \rightarrow \Gamma_P$ is a quasi-isometry (see Lemma 3.25) when each of the three subgroups is equipped with the len_Γ -metric. Thus, we may assume that the restriction $\mathcal{A}_0|_{\Gamma_L} : \Gamma_L \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$ fails to have len_Γ -subexponential growth. By Lemma 6.7, it follows that there is a \mathbb{R} -diagonalizable 1-parameter subgroup $\{a^t\} \subset L$ and an $\{a^t\}$ -invariant Borel probability measure μ on X_L such that

$$\lambda_{\text{top}, a^1, \mu, \mathcal{A}} > 0.$$

Recall we write

$$U_\kappa(n) := \exp_{\mathfrak{u}}(N_\kappa(n))$$

where $N_\kappa(n) = \{Z \in \mathfrak{u} \mid \|Z\| \leq e^{\kappa n}\}$ is as in Section 4.1 for a sufficiently large κ . Take a Følner sequence $\{F_n\}$ in $\{a^t\} \cdot U$ of the form

$$F_n := \{a^s \mid 0 \leq s \leq n\} \cdot U_\kappa(n)$$

Since LU/Γ_{LU} is compact, the family $\{F_n * \mu\}$ is uniformly tight. Let μ_∞ be any weak-* subsequential limit of $\{F_n * \mu\}$. Then μ_∞ is invariant under both $\{a^t\}$ and U .

Claim 9.2. *We have $\lambda_{\text{top}, a^1, \mu_\infty, \mathcal{A}} > 0$.*

Proof. Let μ denote the measure above obtained from Lemma 6.7. There exists an $\{a^t\}$ -invariant, Borel probability measure $\tilde{\mu}$ on $\mathbb{P}\mathcal{E}$ projecting to μ with $\lambda_{\text{top}, a^1, \mu, \mathcal{A}} = \int \Psi(a, \cdot) d\tilde{\mu}$. Let $\tilde{\mu}_\infty$ be any subsequential limit of $\{F_n * \tilde{\mu}\}$ that projects to μ_∞ . Write $a^t = \exp(tY)$ for $Y \in \mathfrak{g}$. Since X_{LU} is compact, given any $\varepsilon > 0$, for all sufficiently large n in the appropriate subsequence we have

$$\begin{aligned} \lambda_{\text{top}, a^1, \mu_\infty, \mathcal{A}} + \varepsilon &\geq \int \Psi(a^1, \xi) d\tilde{\mu}_\infty(\xi) + \varepsilon \\ &= \int \Psi'(Y, \xi) d\tilde{\mu}_\infty(\xi) + \varepsilon \\ &\geq \int \Psi'(Y, \xi) d(F_n * \tilde{\mu})(\xi) \\ &= \frac{1}{n} \int_0^n \int_{U_\kappa(n)} \int \Psi'(Y, a^t \cdot u \cdot \xi) d\tilde{\mu}(\xi) dm_{U_\kappa(n)}(u) dt \\ &= \frac{1}{n} \int_{U_\kappa(n)} \int \Psi(a^n, u \cdot \xi) d\tilde{\mu}(\xi) dm_{U_\kappa(n)}(u). \end{aligned}$$

Above, the first equality follows from $\{a^t\}$ -invariance of $\tilde{\mu}_\infty$ as in Section 6.4; the final equality is the fundamental theorem of calculus as in Section 6.4.

Having taken $\kappa > 0$ sufficiently large, for $u \in U_\kappa(n)$ we have $a^n u = u' a^n$ for some $u' \in U_{2\kappa}(n)$ whence

$$\Psi(a^n, u \cdot \xi) = \Psi(a^n, \xi) + \Psi(u', a^n \cdot \xi) - \Psi(u, \xi).$$

By definition of $\tilde{\mu}$, we have

$$\frac{1}{n} \int \Psi(a^n, \xi) d\tilde{\mu}(\xi) = \lambda_{\text{top}, a^1, \mu, \mathcal{A}}.$$

We have that U is normalized by L and that LU/Γ_{LU} is compact. It follows from Corollary 5.13 and the assumption that $\mathcal{A}_0|_{\Gamma_U} : \Gamma_U \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$ has len_Γ -subexponential growth that the restriction of the cocycle \mathcal{A} to $U \times \mathcal{E}_{LU}$ has subexponential growth; in particular, for any $\varepsilon > 0$, there is a constant C_U such that for all $u'' \in U$ and $\xi \in \mathbb{P}\mathcal{E}_{LU}$,

$$\Psi(u'', \xi) \leq C_U + \varepsilon d(u'', \mathbf{1}).$$

Identifying U with its image under the adjoint representation, the exponential map $\exp|_u$ is polynomial; in particular, there is a constant $C_u > 0$ such that

$$d(\mathbf{1}, \exp_u(Z)) \leq C_u n \quad \text{for all } Z \in N_{2\kappa}(n).$$

Hence

$$|\Psi(u', a^n \cdot \xi) - \Psi(u, \xi)| \leq 2C_U + \varepsilon d(u', \mathbf{1}) + \varepsilon d(u, \mathbf{1}) \leq 2C_U + 2\varepsilon C_u n.$$

Then for n sufficiently large,

$$\frac{1}{n} \int_{U_{\kappa}(n)} \int |\Psi(u', a^n \cdot \xi) - \Psi(u, \xi)| d\tilde{\mu}(\xi) dm_{U_{\kappa}(n)}(u) \leq \frac{2C_U + 2\varepsilon C_u n}{n} \leq 3\varepsilon C_u.$$

The result then follows by taking $\varepsilon > 0$ sufficiently small. \square

As U is the unipotent radical of the minimal parabolic \mathbb{Q} -subgroup P , U is the expanding horospherical subgroup of some one-parameter subgroup $\{b^s\}$ of S . Recall that $\{b^s\}$ centralizes $\{a^t\}$ whence $(b^s)_* \mu_\infty$ is $\{a^t\}$ -invariant for every s . From Corollary 4.4, the sequence $\{(b^s)_* \mu_\infty : s \in \mathbb{N}\}$ has uniformly exponentially small mass at ∞ and its projections converge to the Haar measure on G/Γ . Applying Lemma 6.5 and then Corollary 6.6 produces a measure satisfying the conclusion of Proposition 9.1. \square

10. SUBEXPONENTIAL GROWTH FOR \mathbb{Q} -RANK-1 SUBGROUPS

We prove the following result in which we consider the case that the failure of subexponential growth occurs for the restriction of the cocycle \mathcal{A}_0 to Γ_H for some standard \mathbb{Q} -rank-1 subgroup H in G . This, combined with Proposition 7.1 and Theorem 3.31, yields Theorems D and 6.3.

As in the previous two sections, we keep the standing hypothesis from Assumption 7.2.

Proposition 10.1. *If there is a standard \mathbb{Q} -rank-1 subgroup H of G such that the restriction*

$$\mathcal{A}_0|_{\Gamma_H} : \Gamma_H \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$$

does not have len_Γ -subexponential growth then there is a split Cartan subgroup A of G and an A -invariant Borel probability measure μ on X projecting to the Haar measure on G/Γ such that $\lambda_{\text{top}, a, \mu, \mathcal{A}} > 0$ for some $a \in A$.

10.1. Preliminaries, key ingredients and outline.

10.1.1. *Preliminaries.* Fix a standard \mathbb{Q} -rank-1 subgroup H of G . From Lemma 3.14 there is a parabolic \mathbb{Q} -subgroup Q of G with rational Langlands decomposition $Q^\circ = (L \times S^\circ) \times N$ such that H is the unique \mathbb{Q} -isotropic almost-simple factor of L . We have that H normalizes N and write $\tilde{H} := H \times N$. Since H and N are defined over \mathbb{Q} , \tilde{H} is also defined over \mathbb{Q} .

Note that if $\text{rank}_{\mathbb{Q}}(G) = 1$ then $N = \{e\}$ and $H = \tilde{H} = Q = G$; if $\text{rank}_{\mathbb{Q}}(G) > 1$ then $N \neq \{e\}$ and the inclusions $H \subset \tilde{H} \subset G$ are all proper.

Recall that we write $\Gamma_H := H \cap \Gamma$, $\Gamma_{\tilde{H}} := \tilde{H} \cap \Gamma$, $X_H := (H \times M)/\Gamma_H$, and $X_{\tilde{H}} := (\tilde{H} \times M)/\Gamma_{\tilde{H}}$. The set $\tilde{H}/\Gamma_{\tilde{H}}$ has the structure of a fiber bundle over H/Γ_H with compact fibers. In particular, given any compact set $K \subset \tilde{H}/\Gamma_{\tilde{H}}$, the N -orbit of K is compact.

We select a minimal parabolic \mathbb{Q} -subgroup P and Cartan involution θ as in Lemma 5.3. We equip the vector bundle \mathcal{E} with a norm constructed as in Section 5.4 relative to Siegel fundamental sets constructed relative to P and θ . From Proposition 9.1, we may assume that the restriction of α to Γ_P has uniform len_{Γ} -subexponential growth for every minimal parabolic \mathbb{Q} -subgroup P . From Proposition 8.1 we may assume that $\alpha|_{\Gamma_N}$ has uniform len_{Γ} -subexponential growth.

We will use heavily the growth controls in Lemmas 5.12 and 5.14. We recall the two proper C^∞ “height” functions defined earlier: we have $h: G/\Gamma \rightarrow [0, \infty)$ satisfying (5.7) and $\phi: \tilde{H}/\Gamma_{\tilde{H}} \rightarrow [0, \infty)$ the function $\phi = h \circ \rho$ where $\rho: \tilde{H}/\Gamma_{\tilde{H}} \rightarrow H/\Gamma$ is the canonical projection as in (5.10). We lift h to X and $\mathbb{P}\mathcal{E}$ and lift ϕ to $X_{\tilde{H}}$ and $\mathbb{P}\mathcal{E}_{\tilde{H}}$ via the canonical projections $\mathbb{P}\mathcal{E} \rightarrow X \rightarrow G/\Gamma$ and $\mathbb{P}\mathcal{E}_{\tilde{H}} \rightarrow X_{\tilde{H}} \rightarrow \tilde{H}/\Gamma_{\tilde{H}}$. We also fix ω to be the largest of the constants ω appearing in Lemmas 5.12 and 5.14.

10.1.2. *Key Ingredients and Outline.* As discussed in the introduction, the proof of Proposition 10.1 resembles arguments in [16] more than those of [17], particularly in the case where Γ has \mathbb{Q} -rank one. However, to set up the arguments, the first steps are quite similar to work done in [17]. In Section 10.2, we prove various preliminaries about the growth of cocycle \mathcal{A} along A_H trajectories in $X_{\tilde{H}}$. In particular, we show that exponential growth of derivatives is always witnessed by paths beginning and ending in a fixed compact set and that the empirical measures corresponding to such paths form tight families. In addition we show that the exponential growth can always be witnessed along a single one-parameter subgroup. In Section 10.3, we define the crucial new tools for our proof. Recall the norm-growth cocycle defined in Section 6.4 and its infinitesimal generator for a fixed one-parameter subgroup. This cocycle over a one-parameter subgroup is specified by a function which is continuous but unbounded. As already explained in Section 2.4, the key difficulty for our arguments is that this unbounded function is not even in L^1 so basic tools of ergodic theory do not apply. To resolve this difficulty we introduce two related objects. First we introduced a time-averaged version of the cocycle, which is measurable, bounded, invariant under our 1-parameter subgroup, and computes Lyapunov exponents. However, this function is now not continuous and its integral against various measures does not behave well under weak-* convergence. To remedy this we introduce the key technical object, a **cut-off** cocycle, that interpolates between the original cocycle and the time-averaged one. The cut-off cocycle is bounded and while not continuous has very controlled discontinuities and behaves well under weak-* limits; see Section 10.3.2.

The proof of Proposition 10.1 in Section 10.5 first assembles these tools and our prior results to build an $(A_H N)$ -invariant measure on $X_{\tilde{H}}$ that has positive top Lyapunov exponent. The rest of the proof follows arguments of [16] in the \mathbb{Q} -rank-one case and [17] in the case of higher \mathbb{Q} -rank. Technical difficulties occur because it does not always suffice to work with the cut-off cocycle, since it is only a cocycle over a 1-parameter subgroup but not cocycle over the full G -action or full A -action. We thus need to track the interplay between the original continuous norm-growth cocycle, the time-averaged cocycle, and the cut-off cocycle.

10.2. Characterization and properties of orbits with exponential growth.

We study the growth properties of the cocycle \mathcal{A} along A_H trajectories in $X_{\tilde{H}}$. From the choice of θ as in Lemma 5.3, the subgroup H is θ -invariant and the restriction $\theta|_H$ is a Cartan involution of H . Fix a θ -invariant split Cartan subgroup A of G . We obtain an Iwasawa decomposition $H = K_H A_H N_H$ of H where $K_H = K \cap H$, $A_H = A \cap H$, and N_H is the coarse root group corresponding to the \mathbb{Q} -root defining H .

10.2.1. *Maximal growth rate along A_H .* We now define a filtration of $X_{\tilde{H}}$ by compact sets and study the exponential growth of the cocycle \mathcal{A} for trajectories beginning and ending at a fixed set in this filtration. In particular, we show that there is some compact set that witnesses all possible exponential growth.

Write \mathfrak{a}_H for the Lie algebra of A_H . Given $\ell > 0$, define “thick sets” of $X_{\tilde{H}}$ parameterized by the function ϕ as follows:

$$X_{\tilde{H}, \leq \ell_0} := \{x \in X_{\tilde{H}} \mid \phi(x) \leq \ell_0\}.$$

Given $Y \in \mathfrak{a}_H$ set:

$$\begin{aligned} \chi_\ell^+(T, Y, X_{\tilde{H}}) &:= \frac{1}{T} \sup \left\{ \log \|\mathcal{A}(\exp(TY), x)\| : x \in X_{\tilde{H}, \leq \ell_0} \text{ and } \exp(TY) \cdot x \in X_{\tilde{H}, \leq \ell_0} \right\} \\ \chi_\ell^-(T, Y, X_{\tilde{H}}) &:= \frac{1}{T} \sup \left\{ \log \|\mathcal{A}(\exp(TY), x)^{-1}\| : x \in X_{\tilde{H}, \leq \ell_0} \text{ and } \exp(TY) \cdot x \in X_{\tilde{H}, \leq \ell_0} \right\} \end{aligned}$$

Define

$$\chi_\ell(Y, X_{\tilde{H}}) = \limsup_{T \rightarrow \infty} \max \left\{ \chi_\ell^+(T, Y, X_{\tilde{H}}), \chi_\ell^-(T, Y, X_{\tilde{H}}) \right\}.$$

Observe that $\chi_\ell(Y, X_{\tilde{H}}) \geq 0$ and that $\chi_\ell(cY, X_{\tilde{H}}) = |c| \chi_\ell(Y, X_{\tilde{H}})$.

Fix a norm $\|\cdot\|$ on \mathfrak{g} that is $\text{Ad}(K)$ invariant and whose right G translates define the symmetric metric on $K \backslash G$. Let \mathfrak{a}' be a vector subspace of \mathfrak{a}_H . Given $\ell > 0$ set

$$\begin{aligned} \chi_\ell(T, \mathfrak{a}', X_{\tilde{H}}) &:= \sup \left\{ \chi_\ell^+(T, Y, X_{\tilde{H}}) : Y \in \mathfrak{a}', \|Y\| = 1 \right\} \\ &= \sup \left\{ \chi_\ell^-(T, Y, X_{\tilde{H}}) : Y \in \mathfrak{a}', \|Y\| = 1 \right\}. \end{aligned}$$

The quantity $\chi_\ell(T, \mathfrak{a}', X_{\tilde{H}})$ measures the maximal exponential growth rate of the cocycle \mathcal{A} along all orbits of the subgroup $\exp(\mathfrak{a}')$ in $X_{\tilde{H}}$ of length at most T that start and end below height ℓ . Define

$$\chi_\ell(\mathfrak{a}', X_{\tilde{H}}) = \limsup_{T \rightarrow \infty} \chi_\ell(T, \mathfrak{a}', X_{\tilde{H}}).$$

Note that $\chi_\ell(\mathfrak{a}', X_{\tilde{H}}) < \infty$ follows from Proposition 5.10.

Exactly as in the proof of Lemma 6.7, the decomposition discussed in Remark 3.21 immediately implies the following.

Claim 10.2. *Under Assumption 7.2, if the restriction of \mathcal{A}_0 to $\Gamma_H: \mathcal{E}_0 \rightarrow \mathcal{E}_0$ fails to have subexponential growth then there exists some $\ell_1 > 0$ such that $\chi_{\ell_1}(\mathfrak{a}_H, X_{\tilde{H}}) > 0$.*

Proof. By assumption and Lemma 3.23, there exist $\kappa > 0$ and sequences $\{\gamma_j\}$ of elements of Γ_H with $\text{len}_\Gamma(\gamma_j) \rightarrow \infty$ and $\{x_j\}$ in X_0 such that

$$\log \|\mathcal{A}_0(\gamma_j, x_j)\| \geq \kappa d(\gamma_j, \mathbf{1}).$$

Let $\bar{x}_j = [\mathbf{1}, x_j] \in X_H \subset X_{\tilde{H}}$ be the point corresponding to the ordered pair $(\mathbf{1}, x_j) \in H \times X_0$. Then there exists C such that for all j ,

$$\log \|\mathcal{A}(\gamma_j, \bar{x}_j)\| \geq \kappa d(\gamma_j, \mathbf{1}) - C.$$

Following the notation in Remark 3.21, write $\gamma_j = k'_j z_j a_j k''_j$ where $k'_j, k''_j \in K_{H,0}$, $z_j \in \mathcal{Z} \cap H$, and $a_j \in A_H$. There is C_1 such that for all j (with $\gamma_j \neq \mathbf{1}$),

$$\max\{d(a_j, \mathbf{1}), \text{len}_{\mathcal{Z}}(z_j)\} \leq C_1 d(\mathbf{1}, \gamma_j).$$

By Claim 5.5 we have that $\|\mathcal{A}(k, x)\|$ is uniformly bounded over all $k \in K_{H,0}$ and all $x \in X$. Moreover, by the assumption in Assumption 7.2 and Claim 5.5, given any $\varepsilon > 0$ there is $C_\varepsilon > 0$ such that for all $x \in X$,

$$\log \|\mathcal{A}(z_j, x)\| \leq C_\varepsilon + \varepsilon \text{len}_{\mathcal{Z}}(z_j).$$

It follows there is \widehat{C} independent of ε such that for all j (with $\gamma_j \neq \mathbf{1}$),

$$\begin{aligned} \log \|\mathcal{A}(a_j, k'_j \cdot \bar{x}_j)\| &\geq \kappa d(\gamma_j, \mathbf{1}) - \widehat{C} - \varepsilon \text{len}_{\mathcal{Z}}(z_j) - C_\varepsilon \\ &\geq \kappa d(\gamma_j, \mathbf{1}) - \widehat{C} - C_1 \varepsilon d(\gamma_j, \mathbf{1}) - C_\varepsilon \\ &\geq (\kappa - C_1 \varepsilon) d(\gamma_j, \mathbf{1}) - \widehat{C} - C_\varepsilon \\ &\geq \left(\frac{1}{C_1} \kappa - \varepsilon\right) d(a_j, \mathbf{1}) - \widehat{C} - C_\varepsilon. \end{aligned}$$

assuming $\varepsilon > 0$ is sufficiently small so that $\frac{1}{C_1} \kappa - \varepsilon > 0$. Moreover, compactness of $K_{H,0}$ implies there is $\ell > 0$ so that for all j , $\phi(k''_j \cdot \bar{x}_j) \leq \ell$ and

$$\phi(a_j k'_j \cdot \bar{x}_j) = \phi(a_j z_j k'_j \cdot \bar{x}_j) = \phi((k'_j)^{-1} \cdot [\mathbf{1}, \gamma_j \cdot x_j]) \leq \ell.$$

It follows $\chi_\ell(\mathfrak{a}', X_{\tilde{H}}) \geq \frac{1}{C_1} \kappa - \varepsilon$. \square

10.2.2. *Paths realizing exponential growth start and end in a fixed thick part.* Recall the constant ℓ_0 provided by Lemma 5.14.

We prove here that if there are arbitrarily long orbits that all start and end in some arbitrary common compact thick set and all carry a certain amount of exponential growth then there are orbits that have both endpoints in the fixed compact thick set $X_{\tilde{H}, \leq \ell_0} = \{x \in X_{\tilde{H}} \mid \phi(x) \leq \ell_0\}$ and that carry at least as much exponential growth.

Lemma 10.3. *Let \mathfrak{a}' be a subspace of \mathfrak{a}_H . Suppose there exist $\chi > 0$, $\ell_1 > 0$, and sequences $x_n \in X_{\tilde{H}}$, $Y_n \in \mathfrak{a}'$ with $\|Y_n\| = 1$, and $t_n \rightarrow \infty$ such that*

- (a) *for each n , we have $\phi(x_n) \leq \ell_1$ and $\phi((t_n Y_n) \cdot x_n) \leq \ell_1$;*
- (b) $\lim_{n \rightarrow \infty} \frac{1}{t_n} \log \|\mathcal{A}(\exp(t_n Y_n), x_n)\| = \chi$.

Then, for each n there exist $\hat{x}_n \in X_{\tilde{H}}$ and \hat{t}_n such that

- (1) $\liminf_{n \rightarrow \infty} \frac{\hat{t}_n}{t_n} \geq \frac{\chi}{\chi_{\ell_1}(\mathfrak{a}', X_{\tilde{H}})}$ whence $\hat{t}_n \rightarrow \infty$;
- (2) $\phi(\hat{x}_n) \leq \ell_0$ and $\phi(\exp(\hat{t}_n Y_n) \cdot \hat{x}_n) \leq \ell_0$;
- (3) $\liminf_{n \rightarrow \infty} \frac{1}{t_n} \log \|\mathcal{A}(\exp(\hat{t}_n Y_n), \hat{x}_n)\| \geq \chi$.

In particular, for any $\mathfrak{a}' \subset \mathfrak{a}$ and any $\ell_1 \geq \ell_0$, we have

$$\chi_{\ell_1}(\mathfrak{a}', X_{\tilde{H}}) = \chi_{\ell_0}(\mathfrak{a}', X_{\tilde{H}}). \quad (10.1)$$

In light of (10.1), we will often drop the subscript and write $\chi(\mathfrak{a}', X_{\tilde{H}}) = \chi_{\ell_0}(\mathfrak{a}', X_{\tilde{H}})$ and $\chi(Y, X_{\tilde{H}}) = \chi_{\ell_0}(Y, X_{\tilde{H}})$ for any $Y \in \mathfrak{a}_H$.

Proof of Lemma 10.3. We may assume $\ell_1 > \ell_0$.

We first claim for all sufficiently large n that there is some $0 \leq s \leq t_n$ such that

$$\phi(\exp(sY_n) \cdot x_n) \leq \ell_0.$$

Indeed, for each n , if no such s exists then by Lemma 5.14 for any $\varepsilon > 0$ there is C_ε such that

$$\log \|\mathcal{A}(\exp(t_n Y_n), x_n)\| \leq 2\omega\ell_1 + \varepsilon t_n + 2C_\varepsilon.$$

By taking $\varepsilon < \chi/2$, it would follow that

$$\log \|\mathcal{A}(\exp(t_n Y_n), x_n)\| < (\chi/2)t_n$$

for all sufficiently large t_n , contradicting hypothesis (b).

We may thus assume for any n there exists s_n such that $\phi(\exp(s_n Y_n) \cdot x_n) \leq \ell_0$. For each $n \in \mathbb{N}$ take

$$p_n = \min\{0 \leq s \leq t_n \mid \exp(sY_n) \cdot x_n \in X_{\tilde{H}, \leq \ell_0}\}$$

and

$$q_n = \max\{0 \leq s \leq t_n \mid \exp(sY_n) \cdot x_n \in X_{\tilde{H}, \leq \ell_0}\}.$$

We have $0 \leq p_n \leq q_n \leq t_n$. Let

$$\hat{x}_n = \exp(p_n Y_n) \cdot x_n, \quad \hat{t}_n = q_n - p_n, \quad \text{and} \quad y_n = \exp(q_n Y_n) \cdot x_n.$$

We claim that sequences (\hat{x}_n) and (\hat{t}_n) satisfy the conclusions of the lemma.

Given any $0 < \varepsilon < \chi/5$, there is $T_1 > 0$ such that

$$\frac{1}{t} \log \|\mathcal{A}(\exp(tY), x)\| < \chi_{\ell_1}(\mathfrak{a}', X_{\tilde{H}}) + \varepsilon$$

for all $x \in X_{\tilde{H}}$ with $\phi(x) \leq \ell_1$, $Y \in \mathfrak{a}'$ with $\|Y\| = 1$, and $t \geq T_1$ such that $\phi(\exp(tY) \cdot x) \leq \ell_1$. Note also that there exists $C_0 > 0$ so that

$$\log \|\mathcal{A}(\exp(tY), x)\| < C_0$$

for all $x \in X_{\tilde{H}}$ with $\phi(x) \leq \ell_1$, $Y \in \mathfrak{a}'$ with $\|Y\| = 1$, and $t \leq T_1$. Finally, from Lemma 5.14 there is $C_\varepsilon > 0$ such that for all n ,

$$\log \|\mathcal{A}((t_n - q_n)Y_n, y_n)\| \leq 2\omega\ell_1 + \varepsilon(t_n - q_n) + C_\varepsilon$$

and

$$\log \|\mathcal{A}(p_n Y_n, x_n)\| = \log \|\mathcal{A}(-p_n Y_n, \hat{x}_n)^{-1}\| \leq 2\omega\ell_1 + \varepsilon p_n + C_\varepsilon.$$

Take $T_2 \geq T_1$ so that

$$\frac{C_0 + 2C_\varepsilon + 4\omega\ell_1}{T_2} \leq \varepsilon.$$

Consider any n sufficiently large so that $t_n \geq T_2$ and

$$\frac{1}{t_n} \log \|\mathcal{A}(\exp(t_n Y_n), x_n)\| \geq \chi - \varepsilon. \quad (10.2)$$

We claim that $\hat{t}_n \geq T_1$. Indeed, were $\hat{t}_n \leq T_1$ we would have that

$$\begin{aligned} \log \|\mathcal{A}(t_n Y_n, x_n)\| &\leq \log \|\mathcal{A}(p_n Y_n, x_n)\| + \log \|\mathcal{A}(\hat{t}_n Y_n, \hat{x}_n)\| \\ &\quad + \log \|\mathcal{A}((t_n - q_n) Y_n, y_n)\| \\ &\leq [2\omega\ell_1 + \varepsilon p_n + C_\varepsilon] + C_0 + [2\omega\ell_1 + \varepsilon(t_n - q_n)] + C_\varepsilon \\ &\leq C_0 + 2C_\varepsilon + 4\omega\ell_1 + \varepsilon t_n. \end{aligned}$$

Since $t_n \geq T_2$ satisfying (10.2) we would then obtain that

$$\frac{1}{t_n} \log \|\mathcal{A}(t_n Y_n, x_n)\| \leq 2\varepsilon < \chi - 2\varepsilon$$

contradicting (10.2).

Still considering $t_n \geq T_2$, since $\hat{t}_n \geq T_1$ we have

$$\begin{aligned} &(\chi_{\ell_1}(\mathbf{a}', X_{\tilde{H}}) + \varepsilon)\hat{t}_n \\ &\geq \log \|\mathcal{A}(\hat{t}_n Y_n, \hat{x}_n)\| \\ &\geq \log \|\mathcal{A}(t_n Y_n, x_n)\| - \log \|\mathcal{A}(p_n Y_n, x_n)\| - \log \|\mathcal{A}((t_n - q_n) Y_n, y_n)\| \\ &\geq (\chi - \varepsilon)t_n - [2\omega\ell_1 + \varepsilon p_n] + C_\varepsilon - [2\omega\ell_1 + \varepsilon(t_n - q_n) + C_\varepsilon] \\ &\geq (\chi - \varepsilon)t_n - (2C_\varepsilon + 4\omega\ell_1) - \varepsilon t_n \\ &\geq (\chi - 3\varepsilon)t_n. \end{aligned}$$

It follows that

$$\hat{t}_n \geq \frac{\chi - 3\varepsilon}{\chi_{\ell_1}(\mathbf{a}', X_{\tilde{H}}) + \varepsilon} t_n$$

and that

$$\log \|\mathcal{A}(\hat{t}_n Y_n, \hat{x}_n)\| \geq (\chi - 3\varepsilon)t_n \geq (\chi - 3\varepsilon)\hat{t}_n.$$

Taking $\varepsilon > 0$ arbitrarily small, the claim follows. \square

10.2.3. Growth along basis vectors. We prove that it suffices to study the growth of the cocycle \mathcal{A} along a one-parameter subgroup of A_H . Fix a vector subspace \mathbf{a}' of \mathfrak{a}_H and any basis $\{Z_1, \dots, Z_r\}$ of unit vectors of \mathbf{a}' . Recall from (10.1) that $\chi_\ell(\mathbf{a}', X_{\tilde{H}}) = \chi_{\ell_0}(\mathbf{a}', X_{\tilde{H}})$ for all $\ell \geq \ell_0$ whence we write $\chi(\mathbf{a}', X_{\tilde{H}}) := \chi_{\ell_0}(\mathbf{a}', X_{\tilde{H}})$.

Corollary 10.4. *There exists $C \geq 1$ depending only on the norm $\|\cdot\|$ and the basis $\{Z_1, \dots, Z_r\}$ such that if $\chi(\mathbf{a}', X_{\tilde{H}}) > 0$ then for some $1 \leq j \leq r$ we have*

$$\chi(Z_j, X_{\tilde{H}}) \geq \frac{\chi(\mathbf{a}', X_{\tilde{H}})}{C}.$$

Proof. We induct on $r = \dim \mathbf{a}'$. The result clearly holds when $r = 1$.

Since $\chi(\mathbf{a}', X_{\tilde{H}}) > 0$, up to reversing paths and using Lemma 10.3, we may assume there exist sequences $x_n \in X_{\tilde{H}}$ with $\phi(x_n) \leq \ell_0$, $Y_n \in \mathbf{a}'$ with $\|Y_n\| = 1$, and $t_n \in \mathbb{R}$ with $t_n \rightarrow \infty$, such that writing $y_n := \exp(t_n Y_n) \cdot x_n$, $\phi(y_n) \leq \ell_0$ and

$$\lim_{n \rightarrow \infty} \frac{1}{t_n} \log \|\mathcal{A}(t_n Y_n, x_n)\| = \chi(\mathbf{a}', X_{\tilde{H}}).$$

Suppose $r > 1$. For each n , write $t_n Y_n = \pm p_n Z_r + q_n Z'_n$ where Z'_n is a unit vector in the linear span of $\{Z_1, \dots, Z_{r-1}\}$ and $p_n, q_n \geq 0$. There is C_1 depending only on the choice of basis with

$$\max\{p_n, q_n\} \leq C_1 t_n$$

for all n . Let

$$\begin{aligned} u_n &= \sup\{u \in [0, p_n] : \phi(\exp(u \cdot Z_r) \cdot x_n) \leq \ell_0\} \\ v_n &= \sup\{v \in [0, q_n] : \phi(\exp(-v \cdot Z'_n) \cdot y_n) \leq \ell_0\}. \end{aligned}$$

Fix $\varepsilon > 0$. There is C_ε such that for all n , using Lemma 5.14 we have

$$\begin{aligned} \log \|\mathcal{A}(\exp(t_n Y_n), x_n)\| &\leq \log \|\mathcal{A}(\exp(u_n Z_r), x_n)\| + \log \|\mathcal{A}(\exp(-v_n Z'_n), y_n)^{-1}\| \\ &\quad + 2\omega\ell_0 + \varepsilon[(p_n - u_n) + (q_n - v_n)] + C_\varepsilon \\ &\leq \log \|\mathcal{A}(\exp(u_n Z_r), x_n)\| + \log \|\mathcal{A}(\exp(-v_n Z'_n), y_n)^{-1}\| \\ &\quad + 2\omega\ell_0 + 2\varepsilon C_1 t_n + C_\varepsilon \end{aligned}$$

Taking $\varepsilon > 0$ sufficiently small, for all n sufficiently large it follows that either $\log \|\mathcal{A}(\exp(u_n Z_r), x_n)\|$ or $\log \|\mathcal{A}(\exp(-v_n Z'_n), y_n)^{-1}\|$ is at least

$$\frac{1}{3} \|\mathcal{A}(\exp(t_n Y_n), x_n)\|.$$

Let \mathfrak{a}'' be the span of $\{Z_1, \dots, Z_{r-1}\}$. After dividing by t_n and taking $t_n \rightarrow \infty$ we have that either

$$\chi(Z_r, X_{\tilde{H}}) \geq \lim_{n \rightarrow \infty} \frac{t_n}{3p_n} \chi(\mathfrak{a}', X_{\tilde{H}}) \geq \frac{1}{3C_1} \chi(\mathfrak{a}', X_{\tilde{H}})$$

and the conclusion follows or

$$\chi(\mathfrak{a}'', X_{\tilde{H}}) \geq \lim_{n \rightarrow \infty} \frac{t_n}{3q_n} \chi(\mathfrak{a}', X_{\tilde{H}}) \geq \frac{1}{3C_1} \chi(\mathfrak{a}', X_{\tilde{H}})$$

and the conclusion follows from the induction hypothesis. \square

10.2.4. *Paths approximating maximal growth rates give a tight family of empirical measures.* We study empirical measures coming from certain trajectories and show they form tight families. The families we consider start and end in a fixed compact set and witness the maximal growth rate.

Lemma 10.5 (Tightness of empirical measures). *Fix some $Y \in \mathfrak{a}_H$ with $\chi(Y, X_{\tilde{H}}) > 0$. Let $\{x_n\} \in X_{\tilde{H}}$ be a sequence of points in $X_{\tilde{H}}$, let $t_n \rightarrow \infty$, and let*

$$\eta_n = \frac{1}{t_n} \int_0^{t_n} \exp(sY)_* \delta_{x_n} ds.$$

If

- (a) for each n , $\phi(x_n) \leq \ell_0$ and $\phi(\exp(t_n Y) \cdot x_n) \leq \ell_0$, and
- (b) $\lim_{n \rightarrow \infty} \frac{1}{t_n} \log \|\mathcal{A}(\exp(t_n Y), x_n)\| = \chi(Y, X_{\tilde{H}})$,

then $\{\eta_n\}$ is a uniformly tight family of Borel probability measures on $X_{\tilde{H}}$.

Proof. Write $\chi = \chi(Y, X_{\tilde{H}})$. Without loss of generality, we may assume that $\|Y\| = 1$.

Fix some $\delta > 0$. There is $T_0 > 0$ such that for all $x \in X_{\tilde{H}}$ with $\phi(x) \leq \ell_0$ and all $t \geq T_0$ such that $\phi(\exp(tY) \cdot x) \leq \ell_0$ we have

$$\log \|\mathcal{A}(\exp(tY), x)^{\pm 1}\| \leq (\chi + \delta)t.$$

Set

$$C = \sup \left\{ \log \|\mathcal{A}(\exp(tY), x)^{\pm 1}\| : x \in X_{\tilde{H}}, \phi(x) \leq \ell_0, |t| \leq T_0 \right\}.$$

By Lemma 5.14, there exists T_1 such that for all $T \geq T_1$, if $\phi(x) = \ell_0$, $\phi(\exp(TY) \cdot x) = \ell_0$, and if $\phi(\exp(sY) \cdot x) \geq \ell_0$ for all $0 \leq s \leq T$ then

$$\log \|\mathcal{A}(\exp(TY), x)\| \leq \delta T.$$

For $\ell > 0$ and $n \in \mathbb{N}$, write

$$I_{>\ell, n} = \{t \in [0, t_n] : \phi(\exp(tY) \cdot x_n) > \ell\}.$$

Fix some $L \geq T_1$. For each n , let

$$p_n = \frac{m_{\mathbb{R}}(I_{>L+\ell_0+2, n})}{t_n}.$$

Consider some sufficiently large n with $p_n > 0$. Let $x^t = \exp(tY) \cdot x_n$ for $t \in [0, t_n]$. Let

$$b_0 = 0 \leq a_1 < b_1 < a_2 < b_2 < \cdots < a_J < b_J \leq t_n = a_{J+1}$$

be such that

- (1) $\phi(x^{a_{i+1}}) = \phi(x^{b_i}) = \ell_0$ for every $0 \leq i \leq J$;
- (2) (bounded excursions) for every $0 \leq i \leq J$ and for all $b_i \leq t \leq a_{i+1}$,

$$\phi(x^t) < L + \ell_0 + 2;$$

- (3) (deep excursions) for every $1 \leq i \leq J$, $\phi(x^t) \geq \ell_0$ for all $a_i \leq t \leq b_i$ and for some $a_i < t < b_i$,

$$\phi(x^t) \geq L + \ell_0 + 2.$$

Since $p_n > 0$, we have $J \geq 1$.

For each $1 \leq i \leq J$, by the definition of ϕ we have $b_i - a_i \geq L \geq T_1$; in particular, $J \leq t_n/L$ and

$$\log \|\mathcal{A}((b_i - a_i)Y, x^{a_i})\| \leq \delta(b_i - a_i).$$

Also, for $0 \leq i \leq J$, we have

$$\log \|\mathcal{A}((a_{i+1} - b_i)Y, x^{b_i})\| \leq \max\{C, (\chi + \delta)(a_{i+1} - b_i)\}.$$

Let $l_n = \frac{1}{t_n} \sum_{i=1}^J (b_i - a_i)$. Note $p_n \leq l_n$. Having taken n sufficiently large, we have

$$\begin{aligned}
 \chi - \delta &\leq \frac{1}{t_n} \log \|\mathcal{A}(t_n Y, x_n)\| \\
 &\leq \frac{1}{t_n} \sum_{i=1}^J \log \|\mathcal{A}((b_i - a_i)Y, x^{a_i})\| + \frac{1}{t_n} \sum_{i=0}^J \log \|\mathcal{A}((a_{i+1} - b_i)Y, x^{b_i})\| \\
 &\leq \frac{1}{t_n} \sum_{i=1}^J \delta(b_i - a_i) + \frac{1}{t_n} \sum_{i=0}^J \max(C, (\chi + \delta)(a_{i+1} - b_i)) \\
 &\leq \delta l_n + \frac{(J+1)C}{t_n} + (\chi + \delta)(1 - l_n) \\
 &\leq \delta l_n + \frac{C}{L} + \frac{C}{t_n} + (\chi + \delta)(1 - l_n).
 \end{aligned}$$

In particular, for sufficiently large n it follows that

$$l_n \leq \chi^{-1} \left[\frac{C}{L} + \frac{C}{t_n} + 2\delta \right].$$

Thus, given any $\varepsilon > 0$, having taking $\delta > 0$ sufficiently small, L sufficiently large, and t_n sufficiently large, it follows that $p_n \leq l_n < \varepsilon$ for all sufficiently large n whence the family $\{\eta_n\}$ is uniformly tight. \square

10.3. Time-averaged cocycle, cut-off cocycle, and time-averaged exponents. We define our key technical ingredients, the time-averaged and cut-off cocycles, and study their basic properties.

Fix a unit vector $Y \in \mathfrak{a}_H$ with $\chi(Y, X_{\tilde{H}}) > 0$. Let $\{a^t\}$ denote the 1-parameter subgroup $a^t = \exp(tY)$. Recall that $\Psi': \mathfrak{g} \times \mathbb{P}\mathcal{E} \rightarrow \mathbb{R}$ denotes the infinitesimal generator for the norm-growth cocycle Ψ defined in Section 6.4. Let $\psi: \mathbb{P}\mathcal{E} \rightarrow \mathbb{R}$ be the function

$$\psi(\xi) := \Psi'(Y, \xi). \quad (10.3)$$

Then

$$\Psi(a^T, \xi) = \int_0^T \psi(a^t \cdot \xi) dt. \quad (10.4)$$

Note that ψ is continuous but need not be bounded. Moreover, given an $\{a^t\}$ -invariant Borel probability measure μ on $X_{\tilde{H}}$, it may be that ψ is not $L^1(\mu)$. To deal with this lack of integrability, we replace ψ with its forwards time average along the flow $\{a^t\}$. This new function will be bounded though not necessarily continuous.

10.3.1. Time-averaged cocycle. We define the time-averaged cocycle and see that it is a bounded measurable function. For any Borel function $\varphi: \mathbb{P}\mathcal{E} \rightarrow \mathbb{R}$, write

$$\overline{\varphi}(\xi) := \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \varphi(a^t \cdot \xi) dt$$

and

$$\underline{\varphi}(\xi) := \liminf_{T \rightarrow \infty} \frac{1}{T} \int_0^T \varphi(a^t \cdot \xi) dt.$$

Also let

$$\overline{\Psi}(a, \xi) := \limsup_{T \rightarrow \infty} \frac{1}{T} \Psi(a^T, \xi).$$

We collect the following observations.

Claim 10.6.

- (1) $\overline{\psi}(\xi) = \overline{\Psi}(a, \xi)$ for all $\xi \in \mathbb{P}\mathcal{E}$.
- (2) $\overline{\psi}$ and $\underline{\psi}$ are bounded measurable functions.

Let $\mathcal{E}_{\tilde{H}}$ denote the restriction of the vector bundle \mathcal{E} to $X_{\tilde{H}}$. Let η be an $\{a^t\}$ -invariant Borel probability measure on $\mathbb{P}\mathcal{E}_{\tilde{H}}$.

- (3) For η -almost every $\xi \in \mathbb{P}\mathcal{E}_{\tilde{H}}$ we have

$$-\chi(Y, X_{\tilde{H}}) \leq \underline{\psi}(\xi) \leq \overline{\psi}(\xi) \leq \chi(Y, X_{\tilde{H}}).$$

- (4) For η -almost every $\xi \in \mathbb{P}\mathcal{E}_{\tilde{H}}$ such that $\phi(a^t \cdot \xi) \geq \ell_0$ for all $t \geq t_0$ and some t_0 , we have $\overline{\psi}(\xi) = \underline{\psi}(\xi) = 0$.

Proof. Conclusion (1) is immediate from (10.4) and (2) follows from Proposition 5.10.

For (3) and (4), from Proposition 5.10 there exists k and, for each $\ell > 0$ there exists a C_ℓ , such that if $\phi(\xi) \leq \ell$ then for all $T \geq 0$,

$$\log \|\Psi(a^T, \xi)\| \leq kT + C_\ell.$$

Given an $\{a^t\}$ -invariant Borel probability measure η on $\mathbb{P}\mathcal{E}_{\tilde{H}}$, for η -a.e. ξ and any $\delta > 0$ there are ℓ_δ and T_δ such that for all $T \geq T_\delta$,

$$\begin{aligned} m_{\mathbb{R}}\{0 \leq t \leq T : \phi(a^t \cdot \xi) \geq \ell_\delta\} &\leq \delta T, \\ \log \|\Psi(a^T, \xi)\| &\leq (\chi(Y, X_{\tilde{H}}) + \delta)T. \end{aligned}$$

Fix such ξ and consider any $\delta > 0$. Fix any $T \geq T_\delta$ sufficiently large so that

$$T_0 = \sup\{0 \leq t \leq T : \phi(\exp tY) \cdot \xi \leq \ell_\delta\} \geq T_\delta.$$

Then

$$\begin{aligned} \log \|\Psi(a^T, \xi)\| &= \log \|\Psi(a^{T_0}, \xi)\| + \log \|\Psi(a^{T-T_0}, a^{T_0} \cdot \xi)\| \\ &\leq (\chi(Y, X_{\tilde{H}}) + \delta)T_0 + k(T - T_0) + C_{\ell_\delta} \\ &\leq (\chi(Y, X_{\tilde{H}}) + \delta)T + k\delta T + C_{\ell_\delta}. \end{aligned}$$

It follows that

$$\overline{\psi}(\xi) \leq \chi(Y, X_{\tilde{H}}) + (k+1)\delta.$$

The upper bound in (3) follows as $\delta > 0$ was arbitrary.

Moreover, if $\phi(a^t \cdot \xi) \geq \ell_0$ for all $t \geq t_0$ and if T is sufficiently large so that $T_0 > t_0$ then, by Lemma 5.14, given $\varepsilon > 0$ there is C_ε such that

$$\begin{aligned} \log \|\Psi(a^T, \xi)\| &= \log \|\Psi(a^{t_0}, \xi)\| + \log \|\Psi(a^{T_0-t_0}, a^{t_0} \cdot \xi)\| + \log \|\Psi(a^{T-T_0}, a^{T_0} \cdot \xi)\| \\ &\leq kt_0 + C_{\phi(\xi)} + \omega\phi(a^{t_0} \cdot \xi) + \omega\ell_\delta + \varepsilon(T_0 - t_0) + C_\varepsilon + k(T - T_0) + C_{\ell_\delta} \\ &\leq \widehat{C} + \varepsilon T + \omega\ell_\delta + C_\varepsilon + k\delta T + C_{\ell_\delta} \end{aligned}$$

for some \widehat{C} independent of ε and δ . Then $\overline{\psi}(\xi) \leq \varepsilon + k\delta$ and the upper bound in (4) follows as $\delta > 0$ and $\varepsilon > 0$ are arbitrary.

The lower bounds in (3) and (4) are similar. \square

10.3.2. *The function ψ_ℓ and its properties under weak convergence.* Here we modify the function ψ to obtain a bounded function that has the same forwards time averages along orbits with good recurrence properties. This modified version of ψ is our cut-off cocycle. There is a parameter choice involved in the process of cutting-off and we will eventually work with the cut-off cocycle for well-chosen values of the parameter. Although the new function will not be continuous, its discontinuity set will rather tame for most choices of parameters. We also study the time averages of the cut-off cocycle

Fix $\ell > 0$. Given $\xi \in \mathbb{P}\mathcal{E}$, let

$$\begin{aligned} t_-(\xi) &:= \sup\{s \leq 0 \mid a^s \cdot \xi = \ell\}, \\ t_+(\xi) &:= \inf\{s \geq 0 \mid a^s \cdot \xi = \ell\}, \\ \tau(\xi) &:= t_+(\xi) - t_-(\xi) \end{aligned}$$

(with the convention that $\sup(\emptyset) = -\infty$ and $\inf(\emptyset) = \infty$.) Define $\psi_\ell: \mathbb{P}\mathcal{E} \rightarrow \mathbb{R}$ as follows:

$$\psi_\ell(\xi) = \begin{cases} \psi(\xi) & \text{if } h(\xi) \leq \ell; \\ \frac{1}{\tau(\xi)} \int_{t_-(\xi)}^{t_+(\xi)} \psi(a^s \cdot \xi) ds & \text{if } h(\xi) > \ell \text{ and } \\ & -\infty < t_-(\xi) < t_+(\xi) < \infty; \\ 0 & \text{otherwise.} \end{cases} \quad (10.5)$$

From Proposition 5.10, we observe the following:

- (1) For each ℓ , $\psi_\ell: \mathbb{P}\mathcal{E} \rightarrow \mathbb{R}$ is a bounded measurable function;
- (2) moreover, the functions $\overline{\psi}_\ell: \mathbb{P}\mathcal{E} \rightarrow \mathbb{R}$ are uniformly bounded in ℓ .

Lemma 10.7. *Let η be an $\{a^t\}$ -invariant, Borel probability measure on $\mathbb{P}\mathcal{E}$.*

For η -a.e. $\xi \in \mathbb{P}\mathcal{E}$ and for all $\ell > h(\xi)$,

$$\overline{\psi}(\xi) = \overline{\psi}_\ell(\xi) = \underline{\psi}_\ell(\xi) = \underline{\psi}(\xi).$$

Moreover, for all almost every $\xi \in \mathbb{P}\mathcal{E}$,

$$\overline{\psi}(\xi) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \psi(a^t \cdot \xi) dt.$$

We remark that the final conclusion of Lemma 10.7 does not follow directly from the pointwise ergodic theorem as ψ is neither assumed to be $L^1(\eta)$ or non-negative.

Proof of Lemma 10.7. Recall each ψ_ℓ is a bounded measurable function. By the pointwise ergodic theorem, for a.e. $\xi \in \mathbb{P}\mathcal{E}$ the following two conditions hold:

- (1) For every rational ℓ ,

$$\overline{\psi}_\ell(\xi) = \underline{\psi}_\ell(\xi) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \psi_\ell(a^t \cdot \xi) dt.$$

- (2) Given $0 < \delta < 1$, there is a rational $\ell_\delta(\xi) > 0$ with $\ell_\delta(\xi) > h(\xi)$ such that for all sufficiently large T ,

$$m_{\mathbb{R}}(\{t \in [0, T] : h(a^t \cdot \xi) \leq \ell_\delta(\xi)\}) \geq (1 - \delta)T.$$

Moreover, we may assume $\ell_\delta(\xi) \searrow h(\xi)$ as $\delta \nearrow 1$.

Consider ξ satisfying conditions (1) and (2). Fix any $0 < \delta_0 < 1$ and set $\ell_1 := \ell_{\delta_0}(\xi)$. For any $\ell_2 \geq \ell \geq \ell_1$ and $T > 0$ such that $h(a^T \cdot \xi) \leq \ell$ we have

$$\int_0^T \psi_{\ell_2}(a^t \cdot \xi) dt = \int_0^T \psi_{\ell}(a^t \cdot \xi) dt. \quad (10.6)$$

Set

$$L = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \psi_{\ell_1}(a^t \cdot \xi) dt.$$

Observe that $h(\xi) < \ell_1 \leq \ell$ and from condition (2) there is a sequence $\{t_n\}$ with $t_n \rightarrow \infty$ such that $h(a^{t_n} \cdot \xi) \leq \ell_1 \leq \ell$ for all n . Taking $\ell = \ell_1$, for any rational $\ell_2 \geq \ell_1$, (10.6) and condition (1) imply that

$$\overline{\psi_{\ell_2}}(\xi) = \underline{\psi}_{\ell_2}(\xi) = L. \quad (10.7)$$

In particular, the value of $\overline{\psi_{\ell}}(\xi) = \underline{\psi}_{\ell}(\xi)$ is constant over all rational $\ell > h(\xi)$. We claim the same holds for irrational $\ell \geq h(\xi)$.

Claim 10.8. For any $\ell \geq \ell_1$, $\overline{\psi_{\ell}}(\xi) = \underline{\psi}_{\ell}(\xi) = L$.

Proof. Consider any $0 < T_0 < T_1$ such that

$$h(a^{T_0} \cdot \xi) = h(a^{T_1} \cdot \xi) = \ell$$

and $h(a^t \cdot \xi) > \ell$ for all $T_0 < t < T_1$. By definition, the function

$$T \mapsto \frac{1}{T} \int_0^T \psi_{\ell}(a^t \cdot \xi) dt \quad (10.8)$$

is monotonic on $[T_0, T_1]$. Explicitly, with $A = \int_0^{T_0} \psi_{\ell}(a^t \cdot \xi) dt$ and $B = \int_{T_0}^{T_1} \psi_{\ell}(a^t \cdot \xi) dt$, for $T \in [T_0, T_1]$, (10.8) is equivalent to

$$\frac{A}{T} + \frac{1}{T} \frac{T - T_0}{T_1 - T_0} B = \frac{1}{T} \left(A - \frac{T_0 B}{T_1 - T_0} \right) + \frac{B}{T_1 - T_0}$$

hence the derivative of (10.8) has constant sign on $[T_0, T_1]$. It follows that

$$\overline{\psi_{\ell}}(\xi) = \lim_{T_0 \rightarrow \infty} \sup \left\{ \frac{1}{T} \int_0^T \psi_{\ell}(a^t \cdot \xi) dt : T \geq T_0, h(a^T \cdot \xi) \leq \ell \right\} \quad (10.9)$$

and

$$\underline{\psi}_{\ell}(\xi) = \lim_{T_0 \rightarrow \infty} \inf \left\{ \frac{1}{T} \int_0^T \psi_{\ell}(a^t \cdot \xi) dt : T \geq T_0, h(a^T \cdot \xi) \leq \ell \right\}.$$

Take any rational $\ell_2 \geq \ell$. (10.9) combined with (10.6) imply $\overline{\psi_{\ell_2}}(\xi) \geq \overline{\psi_{\ell}}(\xi)$ and similarly $\underline{\psi}_{\ell}(\xi) \geq \underline{\psi}_{\ell_1}(\xi)$. Similarly,

$$\underline{\psi}_{\ell_2}(\xi) \leq \underline{\psi}_{\ell}(\xi) \leq \underline{\psi}_{\ell_1}(\xi).$$

The claim then follows from (10.7). \square

We return to the proof of the lemma. By Proposition 5.10, there is $C > 0$ such that for all $\zeta \in \mathbb{P}\mathcal{E}$ and $g \in G$,

$$|\Psi(g, \zeta)| \leq C + Ch(\zeta) + Cd(g, \mathbf{1}).$$

Fix any $0 < \delta < 1$ and take $\ell \geq \ell_{\delta}(\xi)$. Then

$$m_{\mathbb{R}}(\{t \in [0, T] : h(a^t \cdot \xi) \geq \ell\}) \leq \delta T$$

for all sufficiently large T .

Consider some large $T > 0$. If $h(a^T \cdot \xi) \leq \ell$ then, by definition,

$$\frac{1}{T} \int_0^T \psi(a^t \cdot \xi) dt = \frac{1}{T} \int_0^T \psi_\ell(a^t \cdot \xi) dt.$$

Given any $T_0 < T_1$ such that $h(a^{T_0} \cdot \xi) = h(a^{T_1} \cdot \xi) = \ell$ and $h(a^T \cdot \xi) \geq \ell$ for all $T_0 \leq T \leq T_1$ then, having considered T_0 sufficiently large, we have

$$\begin{aligned} & \left| \frac{1}{T} \int_0^T \psi(a^t \cdot \xi) dt - \frac{1}{T} \int_0^T \psi_\ell(a^t \cdot \xi) dt \right| \\ & \leq \frac{1}{T} \left(\left| \int_0^{T_0} \psi(a^t \cdot \xi) - \psi_\ell(a^t \cdot \xi) dt \right| + \left| \int_{T_0}^T \psi(a^t \cdot \xi) dt \right| + \left| \int_{T_0}^T \psi_\ell(a^t \cdot \xi) dt \right| \right) \\ & \leq \frac{1}{T} \left(0 + (C + C(T - T_0) + C\ell) + \frac{T - T_0}{T_1 - T_0} (C + C(T_1 - T_0) + C\ell) \right) \\ & \leq \frac{2C + 2C\ell}{T} + 2C \frac{T - T_0}{T} \\ & \leq \frac{2C + 2C\ell}{T} + 2C\delta. \end{aligned}$$

Given any $\varepsilon > 0$, by first taking $\delta > 0$ sufficiently small and taking $\ell \geq \ell_\delta(\xi)$, by considering T sufficiently large we conclude that $|\overline{\psi}(\xi) - \overline{\psi}_\ell(\xi)| < \varepsilon$ and $|\underline{\psi}(\xi) - \underline{\psi}_\ell(\xi)| < \varepsilon$.

From Claim 10.8, for any $\ell > h(\xi)$ we have $\overline{\psi}_\ell = \underline{\psi}_\ell = L =: \overline{\psi_{\ell_{\delta_0}(\xi)}}$ for any choice of $0 < \delta_0 < 1$. The independence of ℓ then implies for all such ℓ that $\overline{\psi}_\ell = \overline{\psi}$, $\underline{\psi}_\ell = \underline{\psi}$, and thus $\overline{\psi} = \underline{\psi}$. The conclusions of the lemma then follow. \square

10.3.3. *A classification of $\{a^t\}$ -orbits and discontinuity points of ψ_ℓ .* Now we show that the cut-off cocycle ψ_ℓ has relatively tame discontinuities. We do this by explicitly identifying all possible discontinuities.

Fix some $\ell \in [0, \infty)$ such that the level surface $h^{-1}(\ell)$ is a codimension-one, C^∞ submanifold in G/Γ ; by Sard's theorem and the implicit function theorem this holds for a.e. ℓ in the range of h . Recall we extend $h: G/\Gamma \rightarrow \mathbb{R}$ to $h: \mathbb{P}\mathcal{E} \rightarrow \mathbb{R}$ via the canonical projection $\pi: \mathbb{P}\mathcal{E} \rightarrow G/\Gamma$. We categorize points $x \in G/\Gamma$ and thus $\xi \in \mathbb{P}\mathcal{E}$ with $h(\xi) > \ell$ based on how its orbit under the flow $\{a^t\}$ meets $h^{-1}(\ell)$.

Tangential orbits (T_ℓ): Write T_ℓ for the set of points $\xi \in \mathbb{P}\mathcal{E}$ with $h(\xi) > \ell$, such that the forward or backwards $\{a^t\}$ -orbit of $\pi(\xi)$ in G/Γ is tangent to $h^{-1}(\ell)$ at some point of the orbit.

Non-recurrent (N_ℓ): We say $\xi \in \mathbb{P}\mathcal{E}$ with $h(\xi) > \ell$ is **forward non-recurrent** to height ℓ if

- (1) there is $t_0 < 0$ such that $h(a^{t_0} \cdot \xi) = \ell$ and the $\{a^t\}$ -orbit of $\pi(\xi)$ is transverse to $h^{-1}(\ell)$ in G/Γ at t_0 ; and
- (2) $h(a^t \cdot \xi) > \ell$ for all $t > t_0$.

We similarly say that ξ is **backwards non-recurrent** to height ℓ if there is $t_0 > 0$ such that $h(a^{t_0} \cdot \xi) = \ell$ and the $\{a^t\}$ -orbit of $\pi(\xi)$ is transverse to $h^{-1}(\ell)$ in G/Γ at t_0 and if $h(a^t \cdot \xi) > \ell$ for all $t < t_0$. Write N_ℓ for the set of points that are either forward or backwards non-recurrent to height ℓ .

Excursions (E_ℓ): We say $\xi \in \mathbb{P}\mathcal{E}$ with $h(\xi) > \ell$ is in an **excursion** if there are $t_0 < 0 < t_1$ such that

- (1) $h(a^{t_0} \cdot \xi) = \ell$ and the $\{a^t\}$ -orbit of $\pi(\xi)$ is transverse in G/Γ to $h^{-1}(\ell)$ at t_0 ;
- (2) $h(a^{t_1} \cdot \xi) = \ell$ and the $\{a^t\}$ -orbit of $\pi(\xi)$ is transverse in G/Γ to $h^{-1}(\ell)$ at t_1 ; and
- (3) $h(a^t \cdot \xi) > \ell$ for all $t \in (t_0, t_1)$.

Write E_ℓ for the set of points in excursions. We note that the transversality at the endpoints ensures that E_ℓ is open.

Deep points (D_ℓ): We say ξ is ℓ -**deep** if $h(a^t \cdot \xi) > \ell$ for all $t \in \mathbb{R}$. Write D_ℓ for the set of ℓ -deep points.

Given any $\xi \in \mathbb{P}\mathcal{E}$ with $h(\xi) > \ell$, we have that ξ is in either T_ℓ, N_ℓ, E_ℓ , or D_ℓ .

Claim 10.9. *If η is any σ -finite Borel measure on $\mathbb{P}\mathcal{E}$ then for $m_{\mathbb{R}}$ -a.e. $\ell \in \mathbb{R}$, we have $\eta(h^{-1}(\ell)) = 0$ and $\eta(T_\ell) = 0$.*

Proof. We have $\eta(h^{-1}(\ell)) = 0$ for all but countably many ℓ .

Consider the closed subset

$$\{(\ell, \xi, t) : \frac{\partial}{\partial s} h(a^s \cdot \xi) \Big|_{s=t} = 0, h(a^t \cdot \xi) = \ell\} \subseteq [0, \infty) \times \mathbb{P}\mathcal{E} \times \mathbb{R}$$

and let $\mathcal{T} \subset [0, \infty) \times \mathbb{P}\mathcal{E}$ denote its image under the projection

$$[0, \infty) \times \mathbb{P}\mathcal{E} \times \mathbb{R} \rightarrow [0, \infty) \times \mathbb{P}\mathcal{E}.$$

Since $[0, \infty) \times \mathbb{P}\mathcal{E} \times \mathbb{R}$ is locally compact, \mathcal{T} is Borel.

For any $\xi \in \mathbb{P}\mathcal{E}$, the function $t \mapsto h(a^t \cdot \xi)$ is either constant or, by Sard's theorem, $m_{\mathbb{R}}(\{\ell \in [0, \infty) : (\ell, \xi) \in \mathcal{T}\}) = 0$. For almost every ℓ , we have $\{\ell\} \times T_\ell \subset \mathcal{T}$. It then follows from Fubini's Theorem that $\eta(T_\ell) = 0$ for $m_{\mathbb{R}}$ -a.e. ℓ . \square

As observed above, each function ψ_ℓ is bounded but need not be continuous; however, its discontinuities are rather tame. To compare values of the two height functions ϕ and h on $\mathcal{E}_{\tilde{H}}$ set

$$\widehat{\ell}_0 := \sup\{h(\xi) : \xi \in X_{\tilde{H}}, \phi(\xi) \leq \ell_0\} \quad (10.10)$$

where ℓ_0 is the critical height in Lemma 5.14.

Claim 10.10. *For any $\ell > 0$ such that $h^{-1}(\ell)$ is a codimension-one, C^∞ submanifold in G/Γ , we have the following:*

- (1) *The set of discontinuity points of the function $\psi_\ell : \mathbb{P}\mathcal{E} \rightarrow \mathbb{R}$ is contained in $h^{-1}(\ell) \cup T_\ell \cup N_\ell \cup D_\ell$.*
- (2) *If $\ell \geq \widehat{\ell}_0$ then the set of discontinuity points of the restriction $\psi_\ell|_{\mathbb{P}\mathcal{E}_{\tilde{H}}} : \mathbb{P}\mathcal{E}_{\tilde{H}} \rightarrow \mathbb{R}$ is contained in $h^{-1}(\ell) \cup T_\ell$.*

Proof. If $h(\xi) < \ell$ then, as ψ is continuous, ξ is a continuity point of ψ_ℓ .

Suppose $h(\xi) > \ell$ and $\xi \in E_\ell$. Since E_ℓ is open, if $\xi' \in \mathbb{P}\mathcal{E}$ is sufficiently close to ξ then $\xi' \in E_\ell$ and the $\{a^t\}$ -orbit of ξ' stays close to the $\{a^t\}$ -orbit of ξ for a sufficiently large amount of time. In particular, taking ξ' sufficiently close to ξ ensures that the time average defining $\psi_\ell(\xi)$ in (10.5) is close to the time average defining $\psi_\ell(\xi')$. Thus ψ_ℓ is continuous at ξ which establishes (1).

To establish (2), let $\xi \in N_\ell \cup D_\ell$. By definition, $\psi_\ell(\xi) = 0$. Suppose $\{\xi_n\} \subseteq \mathbb{P}\mathcal{E}_{\tilde{H}}$ converges to ξ and $\psi_\ell(\xi_n) \neq 0$ for all n . We then have $-\infty < t_-(\xi_n) < t_+(\xi_n) < \infty$

for all n , and $t_+(\xi_n) - t_-(\xi_n) \rightarrow \infty$. From Lemma 5.14, $\psi_\ell(\xi_n) \rightarrow 0$. Indeed let $L = \sup\{\phi(\xi) : \xi \in X_{\tilde{H}}, h(\xi) \leq \ell\}$. Then for any $\varepsilon > 0$ there is C_ε so that

$$\psi_\ell(\xi_n) \leq \frac{1}{t_+(\xi_n) - t_-(\xi_n)} (2\omega L + \varepsilon(t_+(\xi_n) - t_-(\xi_n)) + C_\varepsilon). \quad \square$$

10.3.4. *Sequences of measures on $\mathbb{P}\mathcal{E}$.* We now consider a sequence $\{\eta_n\}$ of Borel probability measures on $\mathbb{P}\mathcal{E}$ converging to a Borel probability measure η . Under suitable hypotheses controlling the discontinuity sets of ψ_ℓ and using Lemma 10.7 and Claim 10.10 we will control the integral $\int \Psi(a, \cdot) d\eta$ in terms of integration against η_n .

Corollary 10.11. *Let η_n be a sequence of Borel probability measures on $\mathbb{P}\mathcal{E}$ converging in the weak-* topology to an $\{a^t\}$ -invariant, Borel probability measure η .*

Suppose each η_n is $\{a^t\}$ -invariant and that either

- (1) $\eta(h^{-1}(\ell) \cup T_\ell \cup N_\ell \cup D_\ell) = 0$ for $m_{\mathbb{R}}$ -a.e. ℓ , or
- (2) each η_n is supported on $\mathbb{P}\mathcal{E}_{\tilde{H}}$.

Then

$$\int \overline{\Psi}(a, \cdot) d\eta = \lim_{n \rightarrow \infty} \int \overline{\Psi}(a, \cdot) d\eta_n.$$

In the case that η_n are possibly not $\{a^t\}$ -invariant,

- (3) *if for $m_{\mathbb{R}}$ -a.e. sufficiently large $\ell > 0$ and for every n , each η_n is supported on $\mathbb{P}\mathcal{E}_{\tilde{H}}$, $\psi \in L^1(\eta_n)$, and*

$$\int \psi d\eta_n = \int \psi_\ell d\eta_n$$

then

$$\int \overline{\Psi}(a, \cdot) d\eta = \lim_{n \rightarrow \infty} \int \psi d\eta_n.$$

We remark that if an $\{a^t\}$ -invariant Borel probability measure η on $\mathbb{P}\mathcal{E}$ projects to the Haar measure on G/Γ then, as Haar-a.e. point in G/Γ has dense orbit in G/Γ , the hypothesis in (1) is automatically satisfied. We also remark that (2) and (3) only apply to measures supported on the restricted bundle $\mathcal{E}_{\tilde{H}}$.

Proof. First, consider any $\{a^t\}$ -invariant, Borel probability measure $\tilde{\eta}$ supported on $\mathbb{P}\mathcal{E}$. Using Lemma 10.7, dominated convergence, and the $\{a^t\}$ -invariance of $\tilde{\eta}$ we have

$$\int \overline{\Psi}(a, \cdot) d\tilde{\eta} = \int \overline{\psi} d\tilde{\eta} = \lim_{\ell \rightarrow \infty} \int \overline{\psi}_\ell d\tilde{\eta} = \lim_{\ell \rightarrow \infty} \int \psi_\ell d\tilde{\eta}. \quad (10.11)$$

We apply Claim 10.9 and Claim 10.10. Under the hypothesis in parts (1), (2), or (3), for almost every $\ell \geq \ell_0$ the restriction of each ψ_ℓ to $\mathbb{P}\mathcal{E}_{\tilde{H}}$ is continuous on a set of full η -measure; it follows (see for example [4, Theorem 2.7]) or [5, Corollary 2.2.10]) that

$$\int \psi_\ell d\eta = \lim_{n \rightarrow \infty} \int \psi_\ell d\eta_n. \quad (10.12)$$

Part (3) follows by applying (10.11) with $\tilde{\eta} = \eta$ and an appropriate choice of such ℓ .

For parts (1) and (2), we utilize a uniform (in n) version of (10.11): Using that $\eta_n \rightarrow \eta$, given $\delta > 0$ there is ℓ_δ such that for all $\ell \geq \ell_\delta$ and all sufficiently large n we have

$$\eta_n\{\xi : h(\xi) \geq \ell\} \leq \eta_n\{\xi : h(\xi) \geq \ell_\delta\} \leq \delta.$$

Using that each η_n is $\{a^t\}$ -invariant, Lemma 10.7 then implies

$$\eta_n\{\xi : \bar{\psi}(\xi) = \bar{\psi}_\ell(\xi)\} \geq (1 - \delta)$$

for all sufficiently large n . There is C independent of ℓ such that $|\bar{\psi}_\ell(\xi) - \bar{\psi}(\xi)| \leq C$ for all $\xi \in \mathbb{P}\mathcal{E}$. Then for $\ell \geq \ell_\delta$,

$$\left| \int \bar{\Psi}(a, \cdot) d\eta - \int \psi_\ell d\eta \right| \leq \delta C, \quad (10.13)$$

and for all sufficiently large n and all $\ell \geq \ell_\delta$,

$$\left| \int \bar{\Psi}(a, \cdot) d\eta_n - \int \psi_\ell d\eta_n \right| \leq \delta C. \quad (10.14)$$

Under the hypothesis of parts (1) or (2), there exists a sequence $\{\ell_j\}$ with $\ell_j \rightarrow \infty$ such that (10.12) holds. Given any $\delta > 0$, first take $\ell_j > \ell_\delta$ and then take n sufficiently large in (10.12) so that

$$\left| \int \psi_{\ell_j} d\eta - \lim_{n \rightarrow \infty} \int \psi_{\ell_j} d\eta_n \right| < \delta.$$

From (10.13) and (10.14),

$$\left| \int \bar{\Psi}(a, \cdot) d\eta - \int \bar{\Psi}(a, \cdot) d\eta_n \right| \leq (2C + 1)\delta$$

and conclusions (1) and (2) follow as C is independent of δ . \square

To apply the above corollary, we often need to verify tightness of the family $\{\eta_n\}$. The following corollary of Lemma 10.5 gives sufficient criteria for a sequence of measures supported on $\mathbb{P}\mathcal{E}_{\tilde{H}}$ to be uniformly tight.

Corollary 10.12. *Let η_n be any sequence of $\{a^t\}$ -invariant, Borel probability measures on $\mathbb{P}\mathcal{E}_{\tilde{H}}$ such that*

$$\int \bar{\psi} d\eta_n = \chi(Y, X_{\tilde{H}})$$

for every n . Then the family $\{\eta_n\}$ is uniformly tight.

Proof. Let η be any $\{a^t\}$ -invariant, Borel probability measure on $\mathbb{P}\mathcal{E}_{\tilde{H}}$ with $\int \bar{\psi} d\eta = \chi(Y, X_{\tilde{H}})$. From Claim 10.6(3) we have $\bar{\psi}(\xi) = \chi(Y, X_{\tilde{H}})$ for η -a.e. $\xi \in \mathbb{P}\mathcal{E}_{\tilde{H}}$. In particular, for η -a.e. $\xi = (x, [v]) \in \mathbb{P}\mathcal{E}_{\tilde{H}}$,

$$\chi(Y, X_{\tilde{H}}) \geq \limsup_{T \rightarrow \infty} \frac{1}{T} \log \|\mathcal{A}(a^T, x)\| \geq \limsup_{T \rightarrow \infty} \frac{1}{T} \Psi(a^T, \xi) = \bar{\psi}(\xi) = \chi(Y, X_{\tilde{H}}).$$

For η -almost every $\xi \in \mathbb{P}\mathcal{E}_{\tilde{H}}$, there is $\ell_1 \geq \phi(\xi)$ and a sequence $t_j \rightarrow \infty$ such that $\phi(a^{t_j} \cdot \xi) \leq \ell_1$ for all j . If $\bar{\psi}(\xi) = \chi(Y, X_{\tilde{H}})$, then from Lemma 10.3, we further have that $\phi(a^{t'_j} \cdot \xi) \leq \ell_0$ for some sequence $t'_j \rightarrow \infty$.

From the pointwise ergodic theorem, for η -almost every $\xi \in \mathbb{P}\mathcal{E}_{\tilde{H}}$ and every rational ℓ the limit

$$\lim_{T \rightarrow \infty} \frac{1}{T} m_{\mathbb{R}}(\{t \in [0, T] \mid \phi(a^t \cdot \xi) \geq \ell\})$$

exists. Lemma 10.5 implies the following: given any $\delta > 0$ there exists a rational $\ell_\delta > 0$ such that for η -almost every $\xi \in \mathbb{P}\mathcal{E}$ with $\overline{\psi}(\xi) = \chi(Y, X_{\tilde{H}})$,

$$\lim_{T \rightarrow \infty} \frac{1}{T} m_{\mathbb{R}}(\{t \in [0, T] \mid \phi(a^t \cdot \xi) \geq \ell_\delta\}) < \delta.$$

Then for any $\{a^t\}$ -invariant, Borel probability measure on $\mathbb{P}\mathcal{E}_{\tilde{H}}$ with $\int \overline{\psi} d\eta_n = \chi(Y, X_{\tilde{H}})$, the pointwise ergodic theorem implies

$$\eta_n(\{\xi \in \mathbb{P}\mathcal{E}_{\tilde{H}} \mid h(\xi) \geq \ell_\delta\}) < \delta. \quad \square$$

10.4. Invariance and additivity of the time-averaged cocycle. While our modifications of the norm-growth cocycle are beneficial for handling issues of convergence, a priori it destroys all connection the group action even just over A or A_H . Here we show that the space average of the time-averaged cocycles remains a homomorphism from A . This turns out to be all we need to run proofs analogous to those in [16] and [17].

We begin with the following consequence of the pointwise ergodic theorem and Lemma 10.7.

Claim 10.13. *Let η be an $\{a^t\}$ -invariant, Borel probability measure on $\mathbb{P}\mathcal{E}$. If $b \in G$ is in the centralizer of $\{a^t\}$ then for η -a.e. $\xi \in \mathbb{P}\mathcal{E}$,*

$$\overline{\Psi}(a, \xi) = \overline{\Psi}(a, b \cdot \xi).$$

Proof. As $b_*\eta$ and η are $\{a^t\}$ -invariant, from Lemma 10.7, for η -almost every $\xi \in \mathbb{P}\mathcal{E}$ we have

$$\overline{\Psi}(a, b \cdot \xi) = \lim_{T \rightarrow \infty} \frac{1}{T} \Psi(a^T, b \cdot \xi) \quad \text{and} \quad \overline{\Psi}(a, \xi) = \lim_{T \rightarrow \infty} \frac{1}{T} \Psi(a^T, \xi).$$

We have the cocycle relation

$$\Psi(a^T, b \cdot \xi) = \Psi(a^T, \xi) + \Psi(b, a^T \cdot \xi) - \Psi(b, \xi).$$

Moreover, for almost every ξ there is a compact set $E \subset X$ and $T_n \rightarrow \infty$ such that $a^{T_n} \cdot \xi \in E$. For such ξ ,

$$|\Psi(b, a^{T_n} \cdot \xi) - \Psi(b, \xi)|$$

is uniformly bounded in n whence

$$\lim_{T \rightarrow \infty} \frac{1}{T} \Psi(a^T, b \cdot \xi) = \lim_{T \rightarrow \infty} \frac{1}{T} \Psi(a^T, \xi). \quad \square$$

For every $a \in A$, the function $\xi \mapsto \overline{\Psi}(a, \cdot)$ is bounded as observed in Claim 10.6. Thus, the integral $\int \overline{\Psi}(a, \cdot) d\eta$ is well-defined for every Borel probability measure η on $\mathbb{P}\mathcal{E}$.

Lemma 10.14. *For any A -invariant, Borel probability measure η on $\mathbb{P}\mathcal{E}$, the function $A \rightarrow \mathbb{R}$,*

$$a \mapsto \int \overline{\Psi}(a, \cdot) d\eta$$

is linear.

Above, we implicitly identify A with its Lie algebra \mathfrak{a} in the definition of linearity.

Proof. For any one-parameter subgroup $\{a^t\}$ of A we have $\overline{\Psi}(a^t, \cdot) = t \overline{\Psi}(a^1, \cdot)$ for all $t > 0$. It remains to show additivity.

For each $T > 0$, Claim 10.13 gives the equality

$$\overline{\Psi}(b^T, a^T \cdot \xi) = \overline{\Psi}(b^T, \xi) \quad (10.15)$$

for almost every ξ . By Fubini's theorem, for η -a.e. ξ equality (10.15) holds for $m_{\mathbb{R}}$ -a.e. $T > 0$.

For each $a \in A$ and $\varepsilon > 0$, let

$$\mathcal{G}_{a,\varepsilon,T_0} := \left\{ \xi \in \mathbb{P}\mathcal{E} : \left| \overline{\Psi}(a, \xi) - \frac{1}{t} \Psi(a^t, \xi) \right| < \varepsilon \text{ for all } t \geq T_0 \right\}.$$

Fix $a, b \in A_H$ and $\varepsilon > 0$. For almost every ξ , Lemma 10.7 implies there is some T_0 such that $\xi \in \mathcal{G}_{a,\varepsilon,T_0} \cap \mathcal{G}_{ab,\varepsilon,T_0} \cap \mathcal{G}_{b,\varepsilon,T_0}$; invariance of η implies ξ may be chosen so that for some $T \geq T_0$, $a^T \cdot \xi \in \mathcal{G}_{b,\varepsilon,T_0}$ and (10.15) holds.

Using the cocycle relation

$$\Psi(a^T, \xi) + \Psi(b^T, a^T \cdot \xi) - \Psi((ab)^T, \xi) = 0,$$

we conclude that

$$\begin{aligned} & \left| \overline{\Psi}(a, \xi) + \overline{\Psi}(b, \xi) - \overline{\Psi}(ab, \xi) \right| \\ & \leq \left| \overline{\Psi}(a, \xi) - \frac{1}{T} \Psi(a^T, \xi) \right| + \left| \overline{\Psi}(b, a^T \cdot \xi) - \frac{1}{T} \Psi(b^T, a^T \cdot \xi) \right| \\ & \quad + \left| \overline{\Psi}((ab)^T, \xi) - \frac{1}{T} \Psi((ab)^T, \xi) \right| \\ & < 3\varepsilon. \end{aligned}$$

It follows that $\overline{\Psi}(a, \xi) + \overline{\Psi}(b, \xi) = \overline{\Psi}(ab, \xi)$ for almost every ξ ; additivity of the integrals then follows. \square

10.5. Proof of Proposition 10.1. Recall we fix H to be a standard \mathbb{Q} -rank-1 subgroup of G , N is a unipotent \mathbb{Q} -subgroup normalized by H , and $\tilde{H} = H \times N$. We assume that the restriction of α to Γ_H fails to have len_{Γ} -subexponential growth.

To prove Proposition 10.1, in Section 10.5.1 we first construct in Proposition 10.17 a measure η on $\mathbb{P}\mathcal{E}_{\tilde{H}}$ that has a positive time-averaged Lyapunov exponent. We will then average this measure so that it projects to the Haar measure on G/Γ while maintaining a non-zero Lyapunov exponent for the limiting measure. The averaging step breaks up into two cases: Section 10.5.2 considers the case where $\text{rank}_{\mathbb{Q}}(\Gamma) = 1$; Section 10.5.3 considers the case where $\text{rank}_{\mathbb{Q}}(\Gamma) \geq 2$.

10.5.1. *Construction of a measure with positive time-averaged Lyapunov exponent.* We harvest the results of our prior efforts to construct an $(A_H N)$ -invariant Borel probability measure with positive time-averaged Lyapunov exponent.

Let \mathfrak{a}_H be the Lie algebra of a split Cartan subgroup A_H of H . Assume as in Proposition 10.1 that the restriction

$$\mathcal{A}_0|_{\Gamma_H} : \Gamma_H \times \mathcal{E}_0 \rightarrow \mathcal{E}_0$$

does not have uniform len_{Γ} -subexponential growth. From Claim 10.2, we have $\chi(\mathfrak{a}_H, X_{\tilde{H}}) > 0$. From Corollary 10.4, there is $Y \in \mathfrak{a}_H$ with $\|Y\| = 1$ and $\chi(Y, X_{\tilde{H}}) > 0$. Write $a^t = \exp(tY)$ and set $a = a^1$.

From Lemma 10.3, there exist points $x_n \in X_{\tilde{H}}$ with $\phi(x_n) \leq \ell_0$ and $t_n \rightarrow \infty$ such that for every n , $\phi(a^{t_n} \cdot x_n) \leq \ell_0$ and

$$\lim_{n \rightarrow \infty} \frac{1}{t_n} \log \|\mathcal{A}(a^{t_n}, x_n)\| = \chi(Y, X_{\tilde{H}}).$$

For each n , take $\xi_n \in \mathbb{P}\mathcal{E}_{\tilde{H}}$ with $\pi_{\mathcal{E}}(\xi_n) = x_n$ such that

$$\lim_{n \rightarrow \infty} \frac{1}{t_n} \Psi(a^{t_n}, \xi_n) = \chi(Y, X_{\tilde{H}}). \quad (10.16)$$

There is a Følner sequence in $\{a^t\} \cdot N$ of the form

$$F_\kappa(t_n) = \{a^s \mid 0 \leq s \leq t_n\} \cdot N_\kappa(t_n),$$

where $N_\kappa(t) := \exp_{\mathfrak{n}}(\{Z \in \mathfrak{n} \mid \|Z\| \leq e^{\kappa t}\})$ and $\kappa > 0$ is chosen sufficiently large as in Section 4.1. For $n \in \mathbb{N}$, set

$$\eta_n := F_\kappa(t_n) * \delta_{\xi_n}.$$

Lemma 10.15. *We have*

$$\lim_{n \rightarrow \infty} \int \psi d\eta_n = \chi(Y, X_{\tilde{H}}).$$

Proof. From (10.16) it suffices to show

$$\lim_{n \rightarrow \infty} \int \psi d\eta_n = \lim_{n \rightarrow \infty} \frac{1}{t_n} \Psi(a^{t_n}, \xi_n).$$

For all $g \in N_\kappa(t_n)$ we have

$$(a^{t_n} g) \cdot \xi_n = (g' a^{t_n}) \cdot \xi_n \quad (10.17)$$

where $g' \in N_{2\kappa}(t_n)$.

Recall that $\phi(x_n) \leq \ell_0$ and $\phi(a^{t_n} \cdot x_n) \leq \ell_0$. Recall also from Lemma 5.12 that the action of N on $X_{\tilde{H}}$ has ϕ -tempered subexponential growth: for any $\varepsilon > 0$, there is a constant C_ε such that

$$\Psi(g'', \xi) \leq C_\varepsilon + \omega\phi(\xi) + \varepsilon d(g'', \mathbf{1}) \text{ for all } g'' \in N \text{ and } \xi \in \mathcal{E}_{\tilde{H}}.$$

Since N is unipotent, identifying $N \subset G$ with its image in F or in $\text{Ad}(G)$, the exponential map $\exp_{\mathfrak{n}}: \mathfrak{n} \rightarrow N$ from \mathfrak{n} to N is polynomial and there is a constant $C_{\mathfrak{n}} > 0$ such that for $t_n \geq 1$,

$$d(\mathbf{1}, \exp_{\mathfrak{n}}(Z)) \leq C_{\mathfrak{n}} t_n \text{ for all } Z \in N_{2\kappa}(t_n).$$

We have the cocycle relation,

$$\Psi(a^{t_n}, g \cdot \xi_n) - \Psi(a^{t_n}, \xi_n) = \Psi(g', a^{t_n} \cdot \xi_n) - \Psi(g, \xi_n).$$

Hence for $g \in N_\kappa(t_n)$,

$$\begin{aligned} \left| \int_0^{t_n} \psi(a^s g \cdot \xi_n) ds - \int_0^{t_n} \psi(a^s \cdot \xi_n) ds \right| &= |\Psi(a^{t_n}, g \cdot \xi_n) - \Psi(a^{t_n}, \xi_n)| \\ &= |\Psi(g', a^{t_n} \cdot \xi_n) - \Psi(g, \xi_n)| \\ &\leq 2C_\varepsilon + \omega\ell_0 + \varepsilon d(g', \mathbf{1}) + \omega\ell_0 + \varepsilon d(g, \mathbf{1}) \\ &\leq 2C_\varepsilon + 2\omega\ell_0 + 2\varepsilon C_{\mathfrak{n}} t_n \\ &\leq \widehat{C}_\varepsilon + \widehat{\varepsilon} t_n \end{aligned}$$

where g' is as in (10.17) and $\widehat{\varepsilon} > 0$ can be made arbitrarily small by choosing $\varepsilon > 0$ sufficiently small.

We then have

$$\begin{aligned}
& \left| \int \psi d\eta_n - \frac{1}{t_n} \Psi(a^{t_n}, \xi_n) \right| \\
&= \left| \int \psi d\eta_n - \frac{1}{t_n} \int_0^{t_n} \psi(a^s \cdot \xi_n) ds \right| \\
&\leq \int_{N_\kappa(t_n)} \frac{1}{t_n} \left| \int_0^{t_n} \psi(a^s g \cdot \xi_n) - \psi(a^s \cdot \xi_n) ds \right| dm_{N_\kappa(t_n)}(g) \\
&\leq \int_{N_\kappa(t_n)} \frac{1}{t_n} (\widehat{C}_\varepsilon + \widehat{\varepsilon} t_n) dm_{N_\kappa(t_n)}(g) \\
&= \frac{1}{t_n} (\widehat{C}_\varepsilon + \widehat{\varepsilon} t_n)
\end{aligned}$$

which can be made arbitrarily small by taking t_n sufficiently large and $\widehat{\varepsilon}$ sufficiently small. \square

Claim 10.16. *The family $\{\eta_n\}$ is a uniformly tight family of measures.*

Proof. Recall that $\widetilde{H}/\Gamma_{\widetilde{H}}$ is a bundle over H/Γ_H with compact fibers. In particular, for any compact subset $E \subset \widetilde{H}/\Gamma_{\widetilde{H}}$, its N -orbit $N \cdot E \subset \widetilde{H}/\Gamma_{\widetilde{H}}$ is compact. From Lemma 10.5, given $\delta > 0$ there is a $\ell_\delta > 0$ such that for every n ,

$$m_{\mathbb{R}}\{0 \leq t \leq t_n : \phi(a^t \cdot x_n) > \ell_\delta\} < \delta t_n.$$

Recall that ϕ is N -invariant. Then It follows that

$$\eta_n(\{\xi \in \mathbb{P}\mathcal{E}_{\widetilde{H}} : \phi(\xi) > \ell_\delta\}) < \delta$$

for every n . \square

Recall the choice of x_n, t_n from which the measures η_n are constructed. Since $\phi(x_n) \leq \ell_0$ and $\phi(a^{t_n} \cdot x_n) \leq \ell_0$, we have

$$h(g \cdot x_n) \leq \widehat{\ell}_0 \text{ and } h((a^{t_n} g) \cdot x_n) \leq \widehat{\ell}_0$$

for all $g \in N$ where $\widehat{\ell}_0$ is as in (10.10). Recall ψ_ℓ as defined in (10.5). Taking $\ell \geq \widehat{\ell}_0$ sufficiently large, it follows from the definition of η_n that for every n ,

$$\int \psi_\ell d\eta_n = \int \psi d\eta_n.$$

With the above observations, we obtain an A_H -invariant, Borel probability measure on $\mathbb{P}\mathcal{E}_{\widetilde{H}}$ with non-zero time-averaged Lyapunov exponent.

Proposition 10.17. *There is an $(A_H N)$ -invariant, Borel probability measure η on $\mathbb{P}\mathcal{E}_{\widetilde{H}}$ such that $\int \overline{\Psi}(a, \cdot) d\eta = \chi(Y, X_{\widetilde{H}}) > 0$.*

Proof. From Claim 10.16, the family $\{\eta_n\}$ is uniformly tight. Let η_∞ be any weak-* limit of the family $\{\eta_n\}$. Then η_∞ is invariant under both $\{a^t\}$ and N .

Take for almost every $\ell > \widehat{\ell}_0$ we have that $h^{-1}(\ell)$ is a codimension-one hyper-surface, $\eta(h^{-1}(\ell)) = \eta(T_\ell) = 0$, and

$$\int \psi_\ell d\eta_n = \int \psi d\eta_n$$

for every n . From Corollary 10.11(3) and Lemma 10.15, we have

$$\int \overline{\Psi}(a, \cdot) d\eta_\infty = \chi(Y, X_{\tilde{H}}).$$

Let $\{F'_j\}$ be a Følner sequence in A_H , let $\tilde{\eta}_j := F'_j * \eta_\infty$. Since A_H is abelian and η_∞ is $\{a^t\}$ -invariant, by Claim 10.13 we have for every j that

$$\int \overline{\Psi}(a, \cdot) d\tilde{\eta}_j = \int \overline{\Psi}(a, \cdot) d\eta_\infty = \chi(Y, X_{\tilde{H}}).$$

By Corollary 10.12, the family $\{\tilde{\eta}_j\}$ is uniformly tight. Let η be a limit point of $\{\tilde{\eta}_j\}$.

We have that η is A_H -invariant and, since A_H normalizes N , η is N -invariant. From Corollary 10.11(2) we have

$$\int \overline{\Psi}(a, \cdot) d\eta = \lim_{j \rightarrow \infty} \int \overline{\Psi}(a, \cdot) d\tilde{\eta}_j = \chi(Y, X_{\tilde{H}}). \quad \square$$

10.5.2. *Proof of Proposition 10.1 when $\text{rank}_{\mathbb{Q}}(\Gamma) = 1$.* We use the $(A_H N)$ -invariant measure just constructed to prove Proposition 10.1. We assume for the time being that $\text{rank}_{\mathbb{Q}}(\Gamma) = 1$; in this case we have $H = G = \tilde{H}$ so $A_H = A$ and N is trivial. We adapt the averaging arguments from [16, §6.3 and §6.4].

Let A be a split Cartan subgroup of G and let β_1, \dots, β_r be the simple \mathbb{R} -roots of $\Phi(A, G)$ relative to some ordering to be specified later. Let $\eta = \eta_0$ denote the A -invariant Borel probability measure guaranteed by Proposition 10.17. From Lemma 10.14, the map $A \rightarrow \mathbb{R}$ given by $a \mapsto \int \overline{\Psi}(a, \cdot) d\eta$ is linear and from Proposition 10.17 is not identically zero.

The action of the real Weyl group of a simple group on the Lie algebra of a split Cartan subgroup is known to be irreducible. Thus, after permuting the simple factors of G and conjugating by an element of the Weyl group of a simple factor, we may assume the ordering of the roots is such that the map $a \mapsto \int \overline{\Psi}(a, \cdot) d\eta$ is not proportional to any β_i for $i > 1$. In particular, we may choose a one-parameter subgroup $\{a^t\}$ in A such that

$$\int \overline{\Psi}(a^1, \cdot) d\eta > 0$$

and $\beta_i(a^t) = 0$ for all $i > 1$. Let U denote the unipotent subgroup generated by the coarse root groups $U^{[\beta_i]}$ for $i > 1$; then U centralizes the one-parameter subgroup $\{a^t\}$.

Let $\{\tilde{F}_j\}$ be a Følner sequence of centered intervals in U as in Definition 4.5. By Lemma 4.7, the family $\{\tilde{F}_j * \eta_0\}$ is a uniformly tight. Let η_1 be any weak-* limit of $\{\tilde{F}_j * \eta_0\}$. Since $\{\tilde{F}_j\}$ is Følner, the measure η_1 is U -invariant and, as U centralizes $\{a^t\}$, η_1 is $\{a^t\}$ -invariant. By Claim 10.13 and Corollary 10.11(2),

$$\int \overline{\Psi}(a^1, \cdot) d\eta_1 = \lim_{j \rightarrow \infty} \int \overline{\Psi}(a^1, \cdot) d\tilde{F}_j * \eta_0 = \int \overline{\Psi}(a^1, \cdot) d\eta_0.$$

Let $\hat{\eta}_1$ denote the image of η_1 under the projection $\pi_{\mathcal{E}}: \mathcal{E} \rightarrow G/\Gamma$. Since η_0 is A -invariant, it follows from Proposition 4.10 that the image $\hat{\eta}_1$ is also A -invariant.

Let $\{\hat{F}_j\}$ be a Følner sequence in A . As $\hat{\eta}_1$ is A -invariant, each $\hat{F}_j * \eta_1$ projects to the same measure in G/Γ ; in particular, the family $\{\hat{F}_j * \eta_1\}$ is uniformly tight.

Let η_2 be any weak-* limit of $\{\widehat{F}_j * \eta_1\}$. From Claim 10.13 and Corollary 10.11(2), we have

$$\int \overline{\Psi}(a^1, \cdot) d\eta_2 = \int \overline{\Psi}(a^1, \cdot) d(\widehat{F}_j * \eta_1) = \int \overline{\Psi}(a^1, \cdot) d\eta_1 = \int \overline{\Psi}(a^1, \cdot) d\eta > 0.$$

We have that η_2 is A -invariant. Additionally, η_2 is U -invariant since η_1 is U -invariant and A normalizes U . In particular, η_2 and its image $\widehat{\eta}_2$ in G/Γ are invariant under the coarse root groups $U^{[\beta_i]}$ for each $i > 1$. From Proposition 4.9, the projection $\widehat{\eta}_2$ of η_2 to G/Γ is also invariant under the coarse root groups $U^{[-\beta_i]}$ for all $i > 1$.

Suppose first that G is not simple. Then $\langle U^{[\beta_i]}, U^{[-\beta_i]} \mid i > 1 \rangle$ contains a (non-compact) simple factor G_1 of G . From the preceding paragraph, we know that the projection $\widehat{\eta}_2$ is G_1 -invariant. The lattice Γ is assumed irreducible in G and thus Γ projects densely to the group G/G_1 . This implies that every G_1 -orbit in G/Γ is dense. Since G_1 is generated by unipotent elements, Ratner's measure rigidity theorem (see for example [59, Theorem 9]) implies that the Haar measure on G/Γ is the only G_1 -invariant, Borel probability measure on G/Γ . In particular, the projection $\widehat{\eta}_2$ of η_2 to G/Γ is Haar. Taking μ to be the projection of η_2 to X , we have that μ is A -invariant and projects to the Haar measure on G/Γ . Since the Haar measure has exponentially small mass at ∞ we have $\Psi(a, \cdot) \in L^1(\eta_2)$ whence

$$\int \Psi(a, \cdot) d\eta_2 = \int \overline{\Psi}(a, \cdot) d\eta_2 > 0$$

and the conclusion of Proposition 10.1 holds by Claim 6.10(1).

Now consider the case that G is simple. Following [16, Section 5.3], we assume that the roots β_1, \dots, β_r are enumerated as in [16, Appendix A] and we let

$$\widehat{\beta} := \begin{cases} \delta = \text{highest root in } \Phi(A, G) & \text{if } G \text{ is of type } A_\ell, B_\ell, D_\ell, E_6, \text{ or } E_7; \\ \delta' = \text{second highest root} & \text{if } G \text{ is of type } C_\ell, (BC)_\ell, E_8, F_4, \text{ or } G_2. \end{cases}$$

As η_2 is A -invariant, from Lemma 10.14 the function $a \mapsto \int \overline{\Psi}(a, \cdot) d\eta_2$ is linear on A . Moreover, it is not proportional to at least one $\widetilde{\beta} \in \{\beta_1, \widehat{\beta}\}$. Thus, there is a one-parameter subgroup $\{\widetilde{a}^t\}$ of A such that $\int \overline{\Psi}(\widetilde{a}^1, \cdot) d\widehat{\eta}_\infty > 0$ and $U^{\widetilde{\beta}}(\widetilde{a}^t) = 0$. Let $\{F_k\}$ be a Følner sequence of centered intervals in $U^{\widetilde{\beta}}$. From Lemma 4.7, the family $\{\widetilde{F}_k * \eta_2\}$ is uniformly tight. Let η_3 be a weak-* limit point of this family. The proof of [16, Proposition 6.4] establishes that the projection of η_3 to G/Γ is Haar. Moreover, Claim 10.13 and Corollary 10.11(2), imply

$$\int \overline{\Psi}(\widetilde{a}^1, \cdot) d\eta_3 = \int \overline{\Psi}(\widetilde{a}^1, \cdot) d(\widetilde{F}_k * \eta_2) = \int \overline{\Psi}(\widetilde{a}^1, \cdot) d\eta_2 > 0.$$

If $\{\widehat{F}_k\}$ is a Følner sequence in A then we have that $\{\widehat{F}_k * \eta_3\}$ is uniformly tight and from Claim 10.13 and Corollary 10.11(2)

$$\int \overline{\Psi}(\widetilde{a}^1, \cdot) d\eta_4 = \int \overline{\Psi}(\widetilde{a}^1, \cdot) d(\widehat{F}_k * \eta_3) = \int \overline{\Psi}(\widetilde{a}^1, \cdot) d\eta_3 > 0.$$

Then η_4 is A -invariant and projects to the Haar measure on G/Γ . Since the Haar measure on G/Γ has exponentially small mass at ∞ , $\Psi(a, \cdot)$ is $L^1(\eta_4)$ and the pointwise ergodic theorem implies

$$\int \Psi(a, \cdot) d\eta_4 > 0.$$

The conclusion of Proposition 10.1 holds by Claim 6.10(1). \square

10.5.3. *Proof of Proposition 10.1 when $\text{rank}_{\mathbb{Q}}(\Gamma) \geq 2$.* Assume now that $\text{rank}_{\mathbb{Q}}(\Gamma) \geq 2$. This implies that the standard \mathbb{Q} -rank-1 subgroup H is a proper subgroup of G . In this case, the group N is non-trivial; we have that N is the unipotent radical of a standard parabolic \mathbb{Q} -subgroup which is normalized by H as in Section 10.1.1. Let $A \subset G$ be a split Cartan subgroup containing A_H such that N is the expanding horospherical subgroup for some one-parameter subgroup $\{b^s\} \subset A$. Let η be the $A_H N$ -invariant Borel probability measure on $\mathcal{E}_{\tilde{H}}$ guaranteed by Proposition 10.17. For $n \in \mathbb{Z}^+$ set

$$\eta_n := \frac{1}{n} \int_0^n (b^s)_* \eta ds.$$

Claim 10.18. *The family $\{\eta_n\}$ is uniformly tight.*

Proof. Let μ denote the image of η under the projection $\mathbb{P}\mathcal{E}_{\tilde{H}} \rightarrow G/\Gamma$. Recall that N -orbits in $\tilde{H}/\Gamma_{\tilde{H}}$ are compact. In particular, given $\delta > 0$ there is a N -invariant subset $C \subset \tilde{H}/\Gamma_{\tilde{H}}$ with $\mu(C) > 1 - \delta$. Let $\hat{\mu}$ denote the restriction of μ to C .

As N is the expanding horospherical subgroup for b^s , by Corollary 4.4 the family $\{(b^s)_* \hat{\mu} : s \in [0, \infty)\}$ has uniformly exponentially small mass at ∞ . Therefore, there is some ℓ_1 such that for all $s > 0$,

$$(b^s)_* \hat{\mu}(\{x \in G/\Gamma \mid h(x) > \ell_1\}) < \delta$$

whence

$$(b^s)_* \mu(\{x \in G/\Gamma \mid h(x) > \ell_1\}) < 2\delta.$$

Since ℓ_1 is independent of s , the sequence $\{\frac{1}{n} \int_0^n (b^s)_* \mu ds\}$ is uniformly tight. As $\mathbb{P}\mathcal{E} \rightarrow G/\Gamma$ is a proper map, the family $\{\eta_n\}$ is uniformly tight. \square

Let $\eta_\infty = \lim_{k \rightarrow \infty} \eta_{n_k}$ for some weak- $*$ convergent subsequence $\{\eta_{n_k}\}$.

It is clear from construction that η_∞ is b^s -invariant, $\{a^t\}$ -invariant, and also N -invariant because η is N -invariant and $\{b^s\}$ normalizes N . Since N is the expanding horospherical subgroup associated to b^s , this implies that η_∞ projects to the Haar measure on G/Γ by Corollary 4.4. Applying Corollary 10.11(1) we have that

$$\int \bar{\Psi}(a, \cdot) d\eta_\infty > 0.$$

As η_∞ is $\{a^t\}$ -invariant, we may average over a Følner sequence in A , apply Corollary 10.11(1) to obtain an A -invariant Borel probability measure η' on $\mathbb{P}\mathcal{E}$ projecting to the Haar measure on G/Γ with

$$\int \bar{\Psi}(a, \cdot) d\eta' > 0.$$

Since the Haar measure on G/Γ has exponentially small mass at ∞ , $\Psi(a, \cdot)$ is $L^1(\eta')$ and the pointwise ergodic theorem implies

$$\int \Psi(a, \cdot) d\eta' > 0$$

and the proposition follows from Claim 6.10(1). \square

11. SKETCH OF ALTERNATE PROOF IN HIGHER \mathbb{Q} -RANK

This section provides an alternate approach that provides a shorter proof of Proposition 10.1 in the special case where $\text{rank}_{\mathbb{Q}} G \geq 2$ and G has finite center. Mainly we avoid the technology of cut-off functions and time averages that make up the bulk of the previous section of the paper. The methods do not seem to apply in \mathbb{Q} -rank one, and therefore do not yield Proposition 10.1 in full generality. Since this result is subsumed by Proposition 10.1, and is therefore not required for a complete proof of our main theorems, we do not provide details. The proof given here is much more similar to the proof in [17]

Proposition 11.1. *Assume $\text{rank}_{\mathbb{Q}} G \geq 2$ and let $\alpha: \Gamma \rightarrow \text{Diff}^1(M)$ be an action. If there is a standard \mathbb{Q} -rank-1 subgroup H of G , such that the restriction*

$$\alpha|_{\Gamma_H}: \Gamma_H \rightarrow \text{Diff}^1(M)$$

does not have uniform subexponential growth of derivatives there is a split Cartan subgroup A of G and an A -invariant Borel probability measure μ on X projecting to the Haar measure on G/Γ such that $\lambda_{\text{top},a,\mu,A} > 0$ for some $a \in A$.

Sketch of proof. We assume all results attained prior to Section 10, and will also assume some results proved in the early parts of that section. Since $\alpha|_{\Gamma_H}$ does not have uniform subexponential growth of derivatives, there is an \mathbb{R} -diagonalizable one-parameter subgroup $\{a^t\}$ of A_H , a sequence $\{\xi_n\}$ of points in \mathcal{E} , and $t_n \rightarrow \infty$ such that $\{\xi_n\}$ and $\{a^{t_n} \cdot \xi_n\}$ are bounded, and

$$\lim_{n \rightarrow \infty} \frac{1}{t_n} \mathcal{A}(a^{t_n}, \xi_n) > 0.$$

By assuming this limit is as large as possible, we may assume that

$$\lim_{n \rightarrow \infty} \frac{1}{t_n} \max_{t \in [0, t_n]} h(a^t \cdot \xi_n) = 0. \quad (11.1)$$

This follows from the proof of Lemma 10.5 and is more explicit in the proof of [17, Lem. 5.3].

Let $\{b^s\}$ be a nontrivial \mathbb{Q} -diagonalizable one-parameter subgroup of G that centralizes H . The existence of $\{b^s\}$ is where we use the assumption that $\text{rank}_{\mathbb{Q}} G \geq 2$. Let N be the expanding horospherical subgroup of $\{b^s\}$, so N is a unipotent \mathbb{Q} -subgroup. It is normalized by H , because H centralizes $\{b^s\}$. Choose a Følner sequence $\{F_n\}$ in N , such that

$$\limsup_{n \rightarrow \infty} \frac{1}{t_n} \max_{u \in F_n} \log \|u\| < \infty,$$

and such that $\{a^{t'_n} F_n a^{-t'_n}\}$ is also a Følner sequence in N , for every sequence $\{t'_n\}$ with $|t'_n| \leq t_n$.

Note that letting $F'_n := \{a^t\}_{0 \leq t \leq t_n} F_n$ yields a Følner sequence in $\{a^t\}N$. From (11.1), we see that if we let $h_n = \max_{t \in [0, t_n]} h(a^t \cdot \xi_n)$, then there is a sequence $\{s_n\}$, such that $h_n/s_n \rightarrow 0$ and $s_n/t_n \rightarrow 0$; for example, let $s_n = \sqrt{h_n t_n}$. Therefore, it follows from exponential mixing that if we let $\mu_n := b^{s_n} F'_n * \delta_{\xi_n}$, then the projection of $\{\mu_n\}$ to G/Γ converges to Haar by applying Theorem 4.2. There is a technical issue that Theorem 4.2 requires the support of f to be in the unit ball of N , which can be addressed by covering $\{a^{t'_n} F_n a^{-t'_n}\}$ by small cubes contained in translates of the unit ball of N and by considering a partition of unity subordinate to this cover.

The sequence of measures $\{\mu_n\}$ has also exponentially small mass at ∞ , this follows because for a fixed t in $[0, t_n]$, the set $b^{s_n} a_t F_n$ is a U -orbit intersecting the thick part of G/Γ nontrivially and so the non-divergence results for unipotent subgroups apply and one can use [17, Lem. 3.3]. Formally, [17, Lem. 3.3] is proven for one-dimensional unipotent subgroups, but for higher dimensional subgroups it follows by partition U into one-dimensional unipotent orbits.

In addition, any subsequential limit μ_∞ of $\{\mu_n\}$ is $\{a^t\}$ -invariant, because F'_n is a Følner set and b^{s_n} centralizes $\{a^t\}$. Finally, note by an easy computation that:

$$\liminf_{n \rightarrow \infty} \lambda_{\text{top}, a^1, \mu_n, \mathcal{A}} \geq \frac{1}{t_n} \int_{F_n} \mathcal{A}(a^{t_n}, b^{s_n} u \cdot \xi_n) dm_{F_n}(u), \quad (11.2)$$

and we also have

$$\liminf_{n \rightarrow \infty} \frac{1}{t_n} \int_{F_n} \mathcal{A}(a^{t_n}, b^{s_n} u \cdot \xi_n) dm_{F_n}(u) \geq \lim_{n \rightarrow \infty} \frac{1}{t_n} \mathcal{A}(a^{t_n}, \xi_n) > 0 \quad (11.3)$$

because $s_n/t_n \rightarrow 0$ and by using the fact that $\alpha|_{\Gamma_N}$ has uniform subexponential growth of derivatives by Proposition 8.1 and making use of 11.1. Therefore $\lambda_{\text{top}, a^1, \mu_\infty, \mathcal{A}} > 0$ because the top Lyapunov exponent of a bounded continuous linear cocycle is an upper-semicontinuous function of the measure. \square

APPENDIX A. TABLES OF COMPUTATIONS

We give the numbers $d(G)$, $n(G)$, and $v(G)$, and $r(G)$ for various families of simple Lie groups. In Table 1, we calculate $d(G)$, $n(G)$, and $v(G)$ for the \mathbb{R} -split classical Lie groups.

G	real root system	\mathbb{R} -rank	$n(G)$	$d(G)$	$v(G)$	$r(G)$
$\mathrm{SL}(n, \mathbb{R})$ $n \geq 2$	A_{n-1}	$n - 1$	n	$2n - 2$	$n - 1$	$n - 1$
$\mathrm{Sp}(2n, \mathbb{R})$ $n \geq 2$	C_n	n	$2n$	$4n - 4$	$2n - 1$	$2n - 1$
$\mathrm{SO}(n, n)$ $n \geq 4$	D_n	n	$2n$	$2n - 1$	$2n - 2$	$2n - 2$
$\mathrm{SO}(n, n + 1)$ $n \geq 3$	B_n	n	$2n + 1$	$2n$	$2n - 1$	$2n - 1$

TABLE 1. Numerology in Zimmer's conjecture for \mathbb{R} -split, classical simple groups

In Table 2, we calculate $n(G)$, $d(G)$, and $v(G)$ for the split real forms of exceptional groups. We follow the naming of real forms as in [40, Appendix C.4].

G	real root system	\mathbb{R} -rank	$n(G)$	$d(G)$	$v(G)$	$r(G)$
E_I	E_6	6	27	26	16	16
E_V	E_7	7	56	54	27	27
E_{VIII}	E_8	8	248	112	57	57
F_1	F_4	4	26	16	15	15
G	G_2	2	7	6	5	5

TABLE 2. Numerology in Zimmer's conjecture for \mathbb{R} -split exceptional Lie groups

In Table 3, we calculate $n(G)$, $d(G)$, and $v(G)$ for a number of complex Lie groups and in Table 4 we calculate for a number of non-split, non-exceptional Lie groups. (The notation here is entirely standard, except that the group $\mathrm{Sp}(2p, 2q)$ is sometimes called $\mathrm{Sp}(p, q)$, and the group $\mathrm{SO}^*(2n)$ is sometimes called $\mathrm{SO}(n, \mathbb{H})$.)

G	real root system	\mathbb{R} -rank	$n(G)$	$d(G)$	$v(G)$	$r(G)$
$\mathrm{SL}(n, \mathbb{C})$ $n \geq 2$	A_{n-1}	$n - 1$	$2n$	$2n - 2$	$2n - 2$	$n - 1$
$\mathrm{Sp}(2n, \mathbb{C})$ $n \geq 2$	C_n	n	$4n$	$4n - 4$	$4n - 2$	$2n - 1$
$\mathrm{SO}(2n, \mathbb{C})$ $n \geq 4$	D_n	n	$4n$	$2n - 1$	$4n - 4$	$2n - 2$
$\mathrm{SO}(2n + 1, \mathbb{C})$ $n \geq 3$	B_n	n	$4n + 2$	$2n$	$4n - 2$	$2n - 1$

TABLE 3. Numerology in Zimmer's conjecture for the classical complex simple groups

G	real root system	\mathbb{R} -rank	$n(G)$	$d(G)$	$v(G)$	$r(G)$
$\mathrm{SL}(n, \mathbb{H})$ $n \geq 3$	A_{n-1}	$n - 1$	$4n$	$4n - 2$	$4n - 4$	$n - 1$
$\mathrm{SO}(p, q)$ $3 \leq p < q$	B_p	p	$p + q$	$p + q - 1$	$p + q - 2$	$2p - 1$
$\mathrm{SU}(p, q)$ $2 \leq p < q$	BC_p	p	$2(p + q)$	$2(p + q) - 2$	$2(p + q) - 3$	$2p - 1$
$\mathrm{Sp}(2p, 2q)$ $2 \leq p < q$	BC_p	p	$4(p + q)$	$4(p + q) - 4$	$4(p + q) - 5$	$2p - 1$
$\mathrm{SO}^*(2n)$ $n \geq 5$ n odd	$BC_{(n-1)/2}$	$\frac{n-1}{2}$	$4n$	$2n - 1$	$4n - 7$	$n - 2$
$\mathrm{SO}^*(2n)$ $n \geq 6$ n even	$C_{n/2}$	$n/2$	$4n$	$2n - 1$	$4n - 7$	$n - 1$

TABLE 4. Numerology in Zimmer's conjecture for some non-split real forms

REFERENCES

- [1] A. Ash, D. Mumford, M. Rapoport, and Y.-S. Tai, *Smooth compactifications of locally symmetric varieties*, Second, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2010. With the collaboration of Peter Scholze.
- [2] H. Azad, M. Barry, and G. Seitz, *On the structure of parabolic subgroups*, Comm. Algebra **18** (1990), no. 2, 551–562.
- [3] B. Bekka, P. de la Harpe, and A. Valette, *Kazhdan's property (T)*, New Mathematical Monographs, vol. 11, Cambridge University Press, Cambridge, 2008.
- [4] P. Billingsley, *Convergence of probability measures*, Second, Wiley Series in Probability and Statistics: Probability and Statistics, John Wiley & Sons, Inc., New York, 1999. A Wiley-Interscience Publication.
- [5] V. I. Bogachev, *Weak convergence of measures*, Mathematical Surveys and Monographs, vol. 234, American Mathematical Society, Providence, RI, 2018.
- [6] A. Borel and J. Tits, *Éléments unipotents et sous-groupes paraboliques de groupes réductifs. I*, Invent. Math. **12** (1971), 95–104.

- [7] A. Borel, *Introduction aux groupes arithmétiques*, Publications de l'Institut de Mathématique de l'Université de Strasbourg, XV. Actualités Scientifiques et Industrielles, No. 1341, Hermann, Paris, 1969.
- [8] A. Borel, *Some metric properties of arithmetic quotients of symmetric spaces and an extension theorem*, J. Differential Geometry **6** (1972), 543–560.
- [9] A. Borel, *Linear algebraic groups*, Second, Graduate Texts in Mathematics, vol. 126, Springer-Verlag, New York, 1991.
- [10] A. Borel and L. Ji, *Compactifications of symmetric and locally symmetric spaces*, Mathematics: Theory & Applications, Birkhäuser Boston, Inc., Boston, MA, 2006.
- [11] A. Borel and J. Tits, *Groupes réductifs*, Inst. Hautes Études Sci. Publ. Math. **27** (1965), 55–150.
- [12] A. Borel and J. Tits, *Compléments à l'article: “Groupes réductifs”*, Inst. Hautes Études Sci. Publ. Math. **41** (1972), 253–276.
- [13] J. Brezin and C. C. Moore, *Flows on homogeneous spaces: a new look*, Amer. J. Math. **103** (1981), no. 3, 571–613.
- [14] A. Brown, *Entropy, Lyapunov exponents, and rigidity of group actions* (M. Triestino, ed.), Ensaios matemáticos, Sociedade Brasileira de Matemática, 2019.
- [15] A. Brown, D. Damjanovic, and Z. Zhang, *C^1 actions on manifolds by lattices in Lie groups*, 2019. Preprint.
- [16] A. Brown, D. Fisher, and S. Hurtado, *Zimmer’s conjecture: Subexponential growth, measure rigidity, and strong property (T)*, Preprint (2016). arXiv:1608.04995.
- [17] A. Brown, D. Fisher, and S. Hurtado, *Zimmer’s conjecture for actions of $SL(m, \mathbb{Z})$* , Invent. Math. **221** (2020), no. 3, 1001–1060.
- [18] A. Brown, F. Rodriguez Hertz, and Z. Wang, *Invariant measures and measurable projective factors for actions of higher-rank lattices on manifolds*, 2016.
- [19] M. Burger and N. Monod, *Continuous bounded cohomology and applications to rigidity theory*, Geom. Funct. Anal. **12** (2002), no. 2, 219–280.
- [20] L. Clozel, *Démonstration de la conjecture τ* , Invent. Math. **151** (2003), no. 2, 297–328.
- [21] S. G. Dani, *On invariant measures, minimal sets and a lemma of Margulis*, Invent. Math. **51** (1979), no. 3, 239–260.
- [22] M. de la Salle, *Strong property (T) for higher-rank lattices*, Acta Math. **223** (2019), no. 1, 151–193.
- [23] T. de Laat and M. de la Salle, *Strong property (T) for higher-rank simple Lie groups*, Proc. Lond. Math. Soc. (3) **111** (2015), no. 4, 936–966.
- [24] B. Farb and P. Shalen, *Real-analytic actions of lattices*, Invent. Math. **135** (1999), no. 2, 273–296.
- [25] D. Fisher, *Groups acting on manifolds: around the Zimmer program*, Geometry, rigidity, and group actions, 2011, pp. 72–157.
- [26] D. Fisher, *Recent progress in the Zimmer program*, 2017.
- [27] D. Fisher and G. A. Margulis, *Local rigidity for cocycles*, Surveys in differential geometry, Vol. VIII (Boston, MA, 2002), 2003, pp. 191–234.
- [28] É. Ghys, *Sur les groupes engendrés par des difféomorphismes proches de l’identité*, Bol. Soc. Brasil. Mat. (N.S.) **24** (1993), no. 2, 137–178.
- [29] É. Ghys, *Actions de réseaux sur le cercle*, Invent. Math. **137** (1999), no. 1, 199–231.
- [30] G. P. Hochschild, *Basic theory of algebraic groups and Lie algebras*, Graduate Texts in Mathematics, vol. 75, Springer-Verlag, New York-Berlin, 1981.
- [31] J. E. Humphreys, *Introduction to Lie algebras and representation theory*, Springer-Verlag, New York-Berlin, 1972. Graduate Texts in Mathematics, Vol. 9.
- [32] L. Ji and R. MacPherson, *Geometry of compactifications of locally symmetric spaces*, Ann. Inst. Fourier (Grenoble) **52** (2002), no. 2, 457–559.
- [33] D. A. Kazhdan, *On the connection of the dual space of a group with the structure of its closed subgroups*, Funkcional. Anal. i Priložen. **1** (1967), 71–74.
- [34] D. Y. Kleinbock and G. A. Margulis, *Bounded orbits of nonquasiunipotent flows on homogeneous spaces*, Sinaï’s Moscow Seminar on Dynamical Systems, 1996, pp. 141–172.
- [35] D. Y. Kleinbock and G. A. Margulis, *Flows on homogeneous spaces and Diophantine approximation on manifolds*, Ann. of Math. (2) **148** (1998), no. 1, 339–360.
- [36] D. Y. Kleinbock and G. A. Margulis, *Logarithm laws for flows on homogeneous spaces*, Invent. Math. **138** (1999), no. 3, 451–494.

- [37] D. Y. Kleinbock and G. A. Margulis, *On effective equidistribution of expanding translates of certain orbits in the space of lattices*, Number theory, analysis and geometry, 2012, pp. 385–396.
- [38] D. Kleinbock, *An extension of quantitative nondivergence and applications to Diophantine exponents*, Trans. Amer. Math. Soc. **360** (2008), no. 12, 6497–6523.
- [39] D. Kleinbock and S. Mirzadeh, *Dimension estimates for the set of points with non-dense orbit in homogeneous spaces*, Math. Z. **295** (2020), no. 3-4, 1355–1383.
- [40] A. W. Knap, *Lie groups beyond an introduction*, Second, Progress in Mathematics, vol. 140, Birkhäuser Boston, Inc., Boston, MA, 2002.
- [41] F. Labourie, *Large groups actions on manifolds*, Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998), 1998, pp. 371–380.
- [42] V. Lafforgue, *Un renforcement de la propriété (T)*, Duke Math. J. **143** (2008), no. 3, 559–602.
- [43] E. Leuzinger, *Tits geometry, arithmetic groups, and the proof of a conjecture of Siegel*, J. Lie Theory **14** (2004), no. 2, 317–338.
- [44] A. Lubotzky, S. Mozes, and M. S. Raghunathan, *Cyclic subgroups of exponential growth and metrics on discrete groups*, C. R. Acad. Sci. Paris Sér. I Math. **317** (1993), no. 8, 735–740.
- [45] A. Lubotzky, S. Mozes, and M. S. Raghunathan, *The word and Riemannian metrics on lattices of semisimple groups*, Inst. Hautes Études Sci. Publ. Math. **91** (2000), 5–53 (2001).
- [46] G. A. Margulis, *On the action of unipotent groups in the space of lattices* (1975), 365–370.
- [47] G. A. Margulis, *Discrete subgroups of semisimple Lie groups*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), vol. 17, Springer-Verlag, Berlin, 1991.
- [48] D. W. Morris, *Introduction to arithmetic groups*, Deductive Press, 2015.
- [49] D. W. Morris, *Quasi-isometric bounded generation by \mathbb{Q} -rank-one subgroups*, SIGMA Symmetry Integrability Geom. Methods Appl. **16** (2020), Paper No. 012, 17.
- [50] G. D. Mostow, *Self-adjoint groups*, Ann. of Math. (2) **62** (1955), 44–55.
- [51] G. D. Mostow, *Fully reducible subgroups of algebraic groups*, Amer. J. Math. **78** (1956), 200–221.
- [52] G. D. Mostow, *Intersections of discrete subgroups with Cartan subgroups*, J. Indian Math. Soc. **34** (1970), no. 3-4, 203–214 (1971).
- [53] G. D. Mostow, *Strong rigidity of locally symmetric spaces*, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1973. Annals of Mathematics Studies, No. 78.
- [54] A. Nevo and R. J. Zimmer, *A structure theorem for actions of semisimple Lie groups*, Ann. of Math. (2) **156** (2002), no. 2, 565–594.
- [55] M. Orr, *Height bounds and the Siegel property*, Algebra Number Theory **12** (2018), no. 2, 455–478.
- [56] V. Platonov and A. Rapinchuk, *Algebraic groups and number theory*, Pure and Applied Mathematics, vol. 139, Academic Press, Inc., Boston, MA, 1994. Translated from the 1991 Russian original by Rachel Rowen.
- [57] G. Prasad and M. S. Raghunathan, *Cartan subgroups and lattices in semi-simple groups*, Ann. of Math. (2) **96** (1972), 296–317.
- [58] M. S. Raghunathan, *Discrete subgroups of Lie groups*, Springer-Verlag, New York-Heidelberg, 1972. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 68.
- [59] M. Ratner, *Invariant measures and orbit closures for unipotent actions on homogeneous spaces*, Geom. Funct. Anal. **4** (1994), no. 2, 236–257.
- [60] M. Ratner, *On Raghunathan's measure conjecture*, Ann. of Math. (2) **134** (1991), no. 3, 545–607.
- [61] M. Ratner, *Raghunathan's topological conjecture and distributions of unipotent flows*, Duke Math. J. **63** (1991), no. 1, 235–280.
- [62] M. Ratner, *Interactions between ergodic theory, Lie groups, and number theory*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994), 1995, pp. 157–182.
- [63] G. Röhrle, *On certain stabilizers in algebraic groups*, Comm. Algebra **21** (1993), no. 5, 1631–1644.
- [64] G. E. Röhrle, *On the structure of parabolic subgroups in algebraic groups*, J. Algebra **157** (1993), no. 1, 80–115.
- [65] N. A. Shah, *Limit distributions of polynomial trajectories on homogeneous spaces*, Duke Math. J. **75** (1994), no. 3, 711–732.
- [66] T. A. Springer, *Linear algebraic groups*, second, Modern Birkhäuser Classics, Birkhäuser Boston, Inc., Boston, MA, 2009.

- [67] J. Tits, *Classification of algebraic semisimple groups*, Algebraic Groups and Discontinuous Subgroups (Proc. Sympos. Pure Math., Boulder, Colo., 1965), 1966, pp. 33–62.
- [68] M. Viana, *Lectures on Lyapunov exponents*, Vol. 145, Cambridge University Press, 2014.
- [69] D. Witte, *Arithmetic groups of higher \mathbf{Q} -rank cannot act on 1-manifolds*, Proc. Amer. Math. Soc. **122** (1994), no. 2, 333–340.
- [70] R. J. Zimmer, *Ergodic theory and semisimple groups*, Monographs in Mathematics, vol. 81, Birkhäuser Verlag, Basel, 1984.

NORTHWESTERN UNIVERSITY, EVANSTON, IL 60208, USA
Email address: `awb@northwestern.edu`

INDIANA UNIVERSITY, BLOOMINGTON, IN 47401, USA
Email address: `fisherdm@indiana.edu`

UNIVERSITY OF CHICAGO, CHICAGO, IL 60637, USA
Email address: `shurtados@uchicago.edu`