MAPPING CONE OF k-ENTANGLEMENT BREAKING MAPS

REPANA DEVENDRA, NIRUPAMA MALLICK, AND K. SUMESH

Abstract. In [CMW19], the authors introduced k-entanglement breaking linear maps to understand the entanglement breaking property of completely positive maps on taking composition. In this article, we do a systematic study of k-entanglement breaking maps. We prove many equivalent conditions for a k-positive linear map to be k-entanglement breaking, thereby study the mapping cone structure of k-entanglement breaking maps. We discuss examples of kentanglement breaking maps and some of their significance. As an application of our study, we characterize the completely positive maps that reduce Schmidt number on taking composition with another completely positive map.

Contents

1.	Introduction	1
2.	Notation and preliminaries	2
3.	Mapping cone of k-entanglement breaking maps	6
4.	Examples	15
5.	Schmidt number reducing CP-maps	21
6.	Majorization	23
7.	Discussion	26
Acknowledgments		27
Appendix A.		27
References		33

1. Introduction

Completely positive maps that are PPT or entanglement breaking are of great importance in Quantum Information theory. The problem of deciding whether a given linear map is entanglement breaking or not is computationally a hard one. Entanglement breaking maps are known to be PPT-maps, but the converse is not true in general. It is a well-known conjecture ([RJK⁺12]) that the square of a PPT channel is entanglement breaking. To understand the entanglement property of maps on taking composition, in [CMW19], the notion

Date: June 11, 2021.

²⁰²⁰ Mathematics Subject Classification. Primary: 47L07. Secondary: 81P40, 81P42.

Key words and phrases. mapping cones, dual cone, tensor products, positive maps, completely positive maps, entanglement breaking, Schmidt number.

of *k*-entanglement breaking maps is introduced, where $k \in \mathbb{N}$. In this article, we study *k*-entanglement breaking maps in more detail. We generalize several known results about entanglement breaking maps into the setup of *k*-entanglement breaking maps. We hope that our study will help understanding entanglement breaking maps better and hence be of substantial interest, especially in quantum physics.

We organize this article into seven sections. In Section 2, we recall some notations, basic definitions and known-results useful for later sections. In Section 3, we establish various characterizations of k-entanglement breaking maps. Making use of these, we prove (Theorem 3.12) that the set \mathcal{EB}_k and \mathcal{EBCP}_k of all k-entanglement breaking linear maps and k-entanglement breaking completely positive maps, respectively, form mapping cones. Further, we show that (Theorem 3.20) both the cones serve as examples of non-symmetric, untypical mapping cones.

Section 4 discuss examples of k-entanglement breaking maps and some of their significance in determining separability (Theorem 4.15) and entanglement (Corollary 4.16). Though there is no close relation between PPT-maps and k-entanglement breaking linear maps (Proposition 3.15 and Example 4.14), Theorem 4.12 provides a sufficient condition for a trace preserving positive map (in particular, PPT-map) to becomes a k-entanglement breaking map.

As an application, in Section 5, we exhibit the Schmidt number reducing property of 2entanglement breaking completely positive maps. Motivated from work done in [RJP18], we ask the following question, which is closely related to PPT-square conjecture: Suppose Φ is a non-entanglement breaking completely positive map on \mathbb{M}_d . Under what sufficient conditions on Φ does there exist $N \in \mathbb{N}$ such that Φ^N is entanglement breaking? In such cases, $1 = SN(\Phi^N) < SN(\Phi)$. This inequality motivates us to search for those completely positive maps, which reduce the Schmidt number after composing a finite number of times. In Theorem 5.2, we show that precisely those completely positive maps that are 2-entanglement breaking reduce the Schmidt number on composing with another completely positive map. Further, if Φ is a k-entanglement breaking completely positive map, Corollary 5.9 provides an upper bound for N.

In Section 6, we discuss a majorization result for k-entanglement breaking maps. Section 7 discuss one open problem. In Appendix we prove the separability of a particular class of positive matrices, which we use in Section 4. As the proof involves few technical lemmas, we write it as a separate section.

2. Notation and preliminaries

Throughout this article, we fix $d, d_1, d_2, d_3 \in \mathbb{N}$. Unless mentioned otherwise, $\{e_i\}_{i=1}^d \subseteq \mathbb{C}^d$ always denotes the standard orthonormal basis. We let $\mathbb{M}_{d_1 \times d_2}$ denote the space of all $d_1 \times d_2$ complex matrices and $I = I_d \in \mathbb{M}_d = \mathbb{M}_{d \times d}$ be the diagonal matrix with diagonals equals 1. By writing $A = [a_{ij}] \in \mathbb{M}_{d_1 \times d_2}$, we mean A is a $d_1 \times d_2$ complex matrix with $(i, j)^{th}$ entry equals a_{ij} for all $1 \leq i \leq d_1, 1 \leq j \leq d_2$. Further, we let $\mathsf{T} = \mathsf{T}_d : \mathbb{M}_d \to \mathbb{M}_d$ denote the transpose map, $\mathsf{tr} : \mathbb{M}_d \to \mathbb{C}$ the trace map, and $\mathsf{id} = \mathsf{id}_d : \mathbb{M}_d \to \mathbb{M}_d$ the identity map. The cone of all positive (semidefinite) matrices in \mathbb{M}_d is denoted by \mathbb{M}_d^+ . If $A \in \mathbb{M}_d^+$, then we write $A \geq 0$. Given $x \in \mathbb{C}^{d_1}, y \in \mathbb{C}^{d_2}$, define the mapping $|x\rangle\langle y| : \mathbb{C}^{d_2} \to \mathbb{C}^{d_1}$ by $|x\rangle\langle y|(z) := x\langle y, z\rangle$ for all $z \in \mathbb{C}^{d_2}$. Note that $|x\rangle\langle x| \geq 0$.

We let $\Omega_d = \sum_{i=1}^d e_i \otimes e_i \in \mathbb{C}^d \otimes \mathbb{C}^d$. Given a unit vector $\xi \in \mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2}$ there always exist orthonormal sets $\{u_i\}_{i=1}^d \subseteq \mathbb{C}^{d_1}$ and $\{v_i\}_{i=1}^d \subseteq \mathbb{C}^{d_2}$, and scalars $\{\lambda_i\}_{i=1}^d \subseteq [0,1]$ with $\sum_i \lambda_i^2 = 1$ such that

$$\xi = \sum_{i=1}^{d} \lambda_i (u_i \otimes v_i), \tag{2.1}$$

where $d = \min\{d_1, d_2\}$. A decomposition of the form (2.1) is called a *Schmidt decomposition* ([Sch06]) of the vector ξ . The number of non-zero coefficients in any two Schmidt decomposition of ξ is the same, and this unique number is called the *Schmidt rank* of ξ and is denoted by $SR(\xi)$.

Definition 2.1. Let $X \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_{d_2})^+$. If there exist $A_i \in \mathbb{M}_{d_1}^+$ and $B_i \in \mathbb{M}_{d_2}^+$, $1 \le i \le n$, such that $X = \sum_{i=1}^n A_i \otimes B_i$, then X is said to be *separable*, otherwise called *entangled*.

Though sufficient and necessary conditions for separability are known ([HHH96]), determining whether a given positive matrix is separable or not is very hard. Often one uses the Schmidt number techniques to handle this situation. The *Schmidt number* ([TeHo00]) of a positive matrix $X \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_{d_2})^+$ is defined as

$$SN(X) := \min \Big\{ \max \Big\{ SR(\xi_i) : X = \sum_{i=1}^n |\xi_i\rangle \langle \xi_i| \Big\} : \xi_i \in \mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2}, n \ge 1 \Big\}.$$

Clearly $SN(|\xi\rangle\langle\xi|) = SR(\xi)$ for all $\xi \in \mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2}$, and $SN(\sum_{i=1}^n X_i) \leq \max_i \{SN(X_i)\}$, where $X_i \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_{d_2})^+, 1 \leq i \leq n$. Note that X is separable if and only if SN(X) = 1. From the definition of Schmidt number, $SN(X) \leq \min\{d_1, d_2\}$.

A linear map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is said to be *positive* if $\Phi(\mathbb{M}_{d_1}^+) \subseteq \mathbb{M}_{d_2}^+$. Given $k \ge 1$, Φ is said to be *k*-positive if $\mathrm{id}_k \otimes \Phi : \mathbb{M}_k \otimes \mathbb{M}_{d_1} \to \mathbb{M}_k \otimes \mathbb{M}_{d_2}$ is a positive map. If both Φ and $\mathsf{T} \circ \Phi$ (equivalently $\Phi \circ \mathsf{T}$) are *k*-positive, then Φ is called a *k*-PPT map. If Φ is *k*-positive for every $k \ge 1$, then Φ is called a *completely positive* (CP-)map. Every CP-map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ can be written as

$$\Phi = \sum_{i=1}^{n} Ad_{V_i} \tag{2.2}$$

for some $V_1, V_2, \dots, V_n \in \mathbb{M}_{d_1 \times d_2}$, where $Ad_V(X) := V^*XV$ for all $X \in \mathbb{M}_{d_1}$ and $V \in \mathbb{M}_{d_1 \times d_2}$. A decomposition of Φ given in (2.2) is called a *Choi-Kraus decomposition* ([Kra71, Cho75]) of Φ , and V_i 's are called Choi-Kraus operators. A linear map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is said to be *completely co-positive* (co-CP) if $\mathsf{T} \circ \Phi$ (equivalently $\Phi \circ \mathsf{T}$) is a CP-map. A linear map Φ that is both CP and co-CP is called a *PPT-map*. Given $A = [a_{ij}] \in \mathbb{M}_{d_1}$ and $B = [b_{ij}] \in \mathbb{M}_{d_2}$, we let $A \otimes B = [a_{ij}B]$ and thereby identify $\mathbb{M}_{d_1} \otimes \mathbb{M}_{d_2} = \mathbb{M}_{d_1}(\mathbb{M}_{d_2})$. To every linear map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ associate the matrix

$$C_{\Phi} := (\mathsf{id}_{d_1} \otimes \Phi) (\sum_{i,j=1}^{d_1} E_{ij} \otimes E_{ij}) = [\Phi(E_{ij})] \in \mathbb{M}_{d_1} \otimes \mathbb{M}_{d_2} = \mathbb{M}_{d_1}(\mathbb{M}_{d_2}),$$

where $E_{ij} = |e_i\rangle\langle e_j| \in \mathbb{M}_{d_1}, 1 \leq i, j \leq d_1$, are the standard matrix units. The map $\Phi \mapsto C_{\Phi}$ is an isomorphism (called *Choi-Jamiolkowski isomorphism* [Cho75, Jam72]) from the space $\{\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2} \text{ linear map}\}$ onto the space $\mathbb{M}_{d_1} \otimes \mathbb{M}_{d_2} = \mathbb{M}_{d_1}(\mathbb{M}_{d_2})$. Further, Φ is a CP-map if and only if Φ is d_1 -positive if and only if $C_{\Phi} \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_{d_2})^+$; and Φ is PPT if and only if C_{Φ} is a positive matrix with positive partial transpose (i.e., $(\mathrm{id}_{d_1} \otimes \mathrm{T})(C_{\Phi}) \geq 0$).

Suppose $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is a CP-map. Then the *Schmidt number* of Φ is denoted and defined ([ChKo06, Hua06]) as $SN(\Phi) := SN(C_{\Phi})$. It is known ([ChKo06]) that

$$SN(\Phi) = \min\left\{\max\left\{rank(V_i): \Phi = \sum_{i=1}^n Ad_{V_i}\right\}: V_i \in \mathbb{M}_{d_1 \times d_2}, n \ge 1\right\}.$$

Given a linear map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$, we let Φ^* denotes the dual of Φ w.r.t the Hilbert Schmidt inner product $\langle X, Y \rangle := \operatorname{tr}(X^*Y)$. From Choi-Kraus decomposition, it follows that

- $SN(\Phi) = SN(\Phi^*)$,
- $SN(\Phi \circ \Psi) \leq \min \{SN(\Phi), SN(\Psi)\},\$
- $SN(\Phi + \Psi) \leq \max \{SN(\Phi), SN(\Psi)\},\$

where Φ, Ψ are CP-maps.

Definition 2.2. A linear map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is said to be an *entanglement breaking (EB)* if $(\mathrm{id}_k \otimes \Phi)(X)$ is separable for all $X \in (\mathbb{M}_k \otimes \mathbb{M}_{d_1})^+$ and $k \ge 1$.

Note that an EB-map is necessarily a CP-map; in fact, they are PPT-maps. Various characterizations of EB-maps are known in the literature.

Theorem 2.3 ([Hol98, HSR03]). Let $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ be a CP-map. Then the following conditions are equivalent:

- (i) Φ is entanglement breaking.
- (ii) $\Gamma \circ \Phi$ is CP for all positive maps $\Gamma : \mathbb{M}_{d_2} \to \mathbb{M}_n, n \ge 1$.
- (iii) $\Gamma \circ \Phi$ is CP for all positive maps $\Gamma : \mathbb{M}_{d_2} \to \mathbb{M}_{d_1}$.
- (iv) $\Phi \circ \Gamma$ is CP for all positive maps $\Gamma : \mathbb{M}_n \to \mathbb{M}_{d_1}, n \geq 1$.
- (v) $\Phi \circ \Gamma$ is CP for all positive maps $\Gamma : \mathbb{M}_{d_2} \to \mathbb{M}_{d_1}$.
- (vi) $C_{\Phi} \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_{d_2})^+$ is separable.
- (vii) Φ admits a set of rank one Choi-Kraus operators.
- (viii) (Holevo form:) There exist $F_i \in \mathbb{M}_{d_1}^+$ and $R_i \in \mathbb{M}_{d_2}^+$ such that $\Phi(X) = \sum_{i=1}^m tr(XF_i)R_i$ for all $X \in \mathbb{M}_{d_1}$.

One may also consider CP-maps that have Choi-Kraus operators of rank less than or equal to some k > 1. A CP-map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is said to be *k*-partially entanglement breaking (k-PEB) if it has Choi-Kraus operators of rank less than or equal to *k* (equivalently $SN(\Phi) \leq k$). Refer [ChKo06] for details. EB-maps and *k*-PEB maps are also known as *superpositive maps* and *k*-superpositive maps, respectively, in the literature (See [And04, SSZ09]).

Definition 2.4 ([Sto86, Sko11]). Let $\mathcal{P}(d_1, d_2)$ denotes the cone of all positive linear maps from \mathbb{M}_{d_1} into \mathbb{M}_{d_2} . A mapping cone is a closed convex cone $\mathcal{C} \subseteq \mathcal{P}(d_1, d_2)$ such that

$$\Phi\circ\Theta\circ\Psi\in \mathfrak{C}$$

for all $\Theta \in \mathcal{C}$, and CP-maps $\Psi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_1}$ and $\Phi : \mathbb{M}_{d_2} \to \mathbb{M}_{d_2}$. The *dual cone* of a mapping cone \mathcal{C} is defined by

$$\mathcal{C}^{\circ} := \{ \Gamma \in \mathcal{P}(d_1, d_2) : \operatorname{tr}(C_{\Gamma} C_{\Theta}) \ge 0 \text{ for all } \Theta \in \mathcal{C} \}.$$

A mapping cone is said to be *invariant* if $\Gamma_2 \circ \Theta \circ \Gamma_1 \in \mathcal{C}$ for all $\Theta \in \mathcal{C}$ and $\Gamma_i \in \mathcal{P}(d_i, d_i), i = 1, 2$.

If \mathcal{C} is a mapping cone, then the dual cone \mathcal{C}° is also a mapping cone. Further, if \mathcal{C} is invariant, then so is \mathcal{C}° . The following are some well-known examples of mapping cones of $\mathcal{P}(d_1, d_2)$:

 $\mathcal{P}_k(d_1, d_2) := \{k\text{-positive maps from } \mathbb{M}_{d_1} \text{ into } \mathbb{M}_{d_2}\},\$ $\mathcal{CP}(d_1, d_2) := \{\text{CP-maps from } \mathbb{M}_{d_1} \text{ into } \mathbb{M}_{d_2}\},\$ $\mathcal{PEB}_k(d_1, d_2) := \{k\text{-PEB maps from } \mathbb{M}_{d_1} \text{ into } \mathbb{M}_{d_2}\},\$ $\mathcal{EB}(d_1, d_2) := \{\text{EB-maps from } \mathbb{M}_{d_1} \text{ into } \mathbb{M}_{d_2}\},\$

where k > 1. (If $d_1 = d_2 = d$, then we write $\mathcal{P}_k(d), \mathcal{CP}(d), \mathcal{PEB}_k(d), \mathcal{EB}(d)$, respectively.) It is known that $(\mathbb{C}^\circ)^\circ = \mathbb{C}$ for every mapping cone \mathbb{C} , and

$$\mathcal{P}(d_1, d_2)^\circ = \mathcal{EB}(d_1, d_2) \qquad (\text{hence } \mathcal{EB}(d_1, d_2)^\circ = \mathcal{P}(d_1, d_2)) \tag{2.3}$$

$$\mathcal{P}_k(d_1, d_2)^\circ = \mathcal{PEB}_k(d_1, d_2) \quad (\text{hence } \mathcal{PEB}_k(d_1, d_2)^\circ = \mathcal{P}_k(d_1, d_2)) \tag{2.4}$$

$$\mathcal{CP}(d_1, d_2)^\circ = \mathcal{CP}(d_1, d_2). \tag{2.5}$$

The following much more generalized result is known. See [Sto09, SSZ09, Sto11, Sko11] for details.

Theorem 2.5. Let $\mathcal{C} \subseteq \mathcal{P}(d_1, d_2)$ be a mapping cone and $\Gamma \in \mathcal{P}(d_1, d_2)$. Then the following conditions are equivalent:

(i) $\Gamma \in \mathfrak{C}^{\circ}$.

(ii) $\Gamma \circ \Theta^* \in \mathcal{CP}(d_2)$ for all $\Theta \in \mathfrak{C}$.

(iii) $\Theta^* \circ \Gamma \in \mathcal{CP}(d_1)$ for all $\Theta \in \mathfrak{C}$.

Further, if $d_1 = d_2 = d$ and C is *-invariant, then the above conditions are equivalent to:

- (iv) $\Gamma \circ \Theta \in \mathcal{CP}(d)$ for all $\Theta \in \mathcal{C}$.
- (v) $\Theta \circ \Gamma \in \mathcal{CP}(d)$ for all $\Theta \in \mathfrak{C}$.

3. Mapping cone of k-entanglement breaking maps

Definition 3.1 ([CMW19]). Let $k \in \mathbb{N}$. A linear map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is said to be *k*entanglement breaking (*k*-*EB*) if Φ is *k*-positive map and $(\mathsf{id}_k \otimes \Phi)(X)$ is separable for every $X \in (\mathbb{M}_k \otimes \mathbb{M}_{d_1})^+$.

Note that a linear map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is entanglement breaking if and only if Φ is k-entanglement breaking for every $k \ge 1$. Further, from Theorem 2.3(vi), this is equivalent to saying that Φ is d_1 -entanglement breaking.

Remark 3.2. Let $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ be a k-EB map.

(i) If m < k, then Φ is *m*-EB also. For, let $X \in (\mathbb{M}_m \otimes \mathbb{M}_{d_1})^+ = (\mathbb{M}_m(\mathbb{M}_{d_1}))^+$. Then there exist $A_i \in \mathbb{M}_k^+$ and $B_i \in \mathbb{M}_{d_2}^+$ such that

$$\begin{bmatrix} (\operatorname{id}_m \otimes \Phi)(X) & 0 \\ 0 & 0_{k-m} \end{bmatrix} = (\operatorname{id}_k \otimes \Phi)(\begin{bmatrix} X & 0 \\ 0 & 0_{k-m} \end{bmatrix}) = \sum A_i \otimes B_i.$$

Writing $A_i = \begin{bmatrix} A_{11}^{(i)} & A_{12}^{(i)} \\ A_{21}^{(i)} & A_{22}^{(i)} \end{bmatrix} \in \mathbb{M}_k^+$, where $A_{11}^{(i)} \in \mathbb{M}_m^+$ and $A_{22}^{(i)} \in \mathbb{M}_{k-m}^+$, from above we get

$$(\mathsf{id}_m \otimes \Phi)(X) = \sum A_{11}^{(i)} \otimes B_i \in \mathbb{M}_m^+ \otimes \mathbb{M}_{d_2}^+$$

so that $(\mathrm{id}_m \otimes \Phi)(X)$ is separable.

(ii) Let $m \ge 1$. Then, from the definition, it follows that both $\Gamma_2 \circ \Phi$ and $\Phi \circ \Gamma_1$ are k-EB maps for all $\Gamma_2 \in \mathcal{P}(d_2, m)$ and $\Gamma_1 \in \mathcal{P}_k(m, d_1)$.

In this section, our main aim is to prove an analog of Theorem 2.3 in the context of k-EB maps. First, we prove some technical lemmas. Though it did not explicitly state, the following lemma was observed in [CMW19].

Lemma 3.3. Let $X \in (M_m \otimes \mathbb{M}_{d_1})^+$. Then for $j = 1, 2, \dots, SN(X)$ there exist family of isometries $V_{j_i} : \mathbb{C}^j \to \mathbb{C}^m$ and vectors $\psi_{j_i} \in \mathbb{C}^j \otimes \mathbb{C}^{d_1}$ such that

$$X = \sum_{i,j} (V_{j_i} \otimes I_{d_1}) |\psi_{j_i}\rangle \langle \psi_{j_i} | (V_{j_i} \otimes I_{d_1})^*.$$

Proof. Let r := SN(X). Then there exist $z_i \in \mathbb{C}^m \otimes \mathbb{C}^{d_1}$ with $SR(z_i) \leq r$ such that

$$X = \sum_{i} |z_i\rangle\langle z_i| = \sum_{j=1}^{r} \sum_{SR(z_i)=j} |z_i\rangle\langle z_i|.$$
(3.1)

Note that whenever $SR(z_i) = j$, from [CMW19, Lemma 1.2], there exist isometry $V_{j_i} : \mathbb{C}^j \to \mathbb{C}^m$ and a vector $\psi_{j_i} \in \mathbb{C}^j \otimes \mathbb{C}^{d_1}$ such that $z_i = (V_{j_i} \otimes I_{d_1})\psi_{j_i}$. Thus

$$X = \sum_{j=1}^{r} \sum_{i} (V_{j_i} \otimes I_{d_1}) |\psi_{j_i}\rangle \langle \psi_{j_i} | (V_{j_i} \otimes I_{d_1})^*.$$

In (3.1), if there is no z_i with $SR(z_i) = j$, then the corresponding term is taken to be zero, and take $\psi_{j_i} = 0$.

The following result is known in the quantum information theory (c.f [TeHo00, RaAl07, SSZ09, Sko11]), but we could not find out in the following form (except $(i) \Leftrightarrow (v)$), hence providing proof.

Lemma 3.4. Let $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ be a positive map and $k \ge 1$. Then the following conditions are equivalent:

- (i) Φ is k-positive.
- (ii) $(id_m \otimes \Phi)(X) \in (\mathbb{M}_m \otimes \mathbb{M}_{d_2})^+$ for all $X \in (\mathbb{M}_m \otimes \mathbb{M}_{d_1})^+$ with $SN(X) \leq k$ and $m \geq 1$.
- (iii) $(id_{d_2} \otimes \Phi)(X) \in (\mathbb{M}_{d_2} \otimes \mathbb{M}_{d_2})^+$ for all $X \in (\mathbb{M}_{d_2} \otimes \mathbb{M}_{d_1})^+$ with $SN(X) \leq k$.
- (iv) $\Phi \circ \Psi$ is CP for every k-PEB map $\Psi : \mathbb{M}_m \to \mathbb{M}_{d_1}$ and $m \ge 1$.
- (v) $\Phi \circ \Psi$ is CP for every *k*-PEB map $\Psi : \mathbb{M}_{d_2} \to \mathbb{M}_{d_1}$.

Proof. The implications $(ii) \Rightarrow (i)$, $(ii) \Rightarrow (iii)$, $(iv) \Rightarrow (v)$ are trivial, and $(v) \Rightarrow (i)$ follows from [Sko11, Theorem 3]. Now we prove the remaining implications.

 $(i) \Rightarrow (ii)$ Let $X \in (\mathbb{M}_m \otimes \mathbb{M}_{d_1})^+$. If $m \leq k$, then $(\mathrm{id}_m \otimes \Phi)(X) \in (\mathbb{M}_m \otimes \mathbb{M}_{d_2})^+$ as Φ is *m*-positive. Now suppose m > k and $r := SN(X) \leq k$. Then , from Lemma 3.3, there exist isometries $V_{j_i} : \mathbb{C}^j \to \mathbb{C}^m$ and a vectors $\psi_{j_i} \in \mathbb{C}^j \otimes \mathbb{C}^{d_1}$, where $j \leq r$ such that

$$(\mathsf{id}_m \otimes \Phi)(X) = (\mathsf{id}_m \otimes \Phi) \Big(\sum_{i,j} (V_{j_i} \otimes I_{d_1}) |\psi_{j_i}\rangle \langle \psi_{j_i} | (V_{j_i} \otimes I_{d_1})^* \Big).$$
$$= \sum_{i,j} (V_{j_i} \otimes I_{d_2}) \big((\mathsf{id}_j \otimes \Phi) (|\psi_{j_i}\rangle \langle \psi_{j_i}|) \big) (V_{j_i} \otimes I_{d_2})^*.$$
(3.2)

Since Φ is k-positive we have $(id_j \otimes \Phi)(|\psi_{j_i}\rangle\langle\psi_{j_i}|)$ is positive for all i and $j \leq r$. From the above equation we conclude that $(id_m \otimes \Phi)X$ is positive.

 $(ii) \Rightarrow (iv)$ Let $\Psi \in \mathcal{PEB}_k(m, d_1)$, where $m \ge 1$. Then $C_{\Psi} \in (\mathbb{M}_m \otimes \mathbb{M}_{d_1})^+$ and $SN(C_{\Psi}) = SN(\Psi) \le k$, hence by assumption $C_{\Phi \circ \Psi} = (\mathrm{id}_m \otimes \Phi)(C_{\Psi}) \in (\mathbb{M}_m \otimes \mathbb{M}_{d_2})^+$. Thus $\Phi \circ \Psi$ is a CP-map.

 $(iv) \Rightarrow (ii)$ Let $m \ge 1$ and $X \in (\mathbb{M}_m \otimes \mathbb{M}_{d_1})^+$ be such that $SN(X) \le k$. Choose $\Psi \in \mathcal{CP}(m, d_1)$ such that $C_{\Psi} = X$. Since $SN(\Psi) = SN(X) \le k$, by assumption, $\Phi \circ \Psi$ is a CP-map, so that $(\mathrm{id}_m \otimes \Phi)(X) = C_{\Psi \circ \Psi}$ is positive.

 $(iii) \Leftrightarrow (v)$ Proof is the same as that of $(ii) \Leftrightarrow (iv)$.

Theorem 3.5. Let $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ be a *k*-positive map, where k > 1. Then the following conditions are equivalent:

- (i) Φ is *k*-entanglement breaking.
- (ii) $\Phi \circ \Psi$ is entanglement breaking for all *CP*-maps $\Psi : \mathbb{M}_k \to \mathbb{M}_{d_1}$.
- (iii) $\Phi \circ \Psi$ is entanglement breaking for all CP-maps $\Psi : \mathbb{M}_m \to \mathbb{M}_{d_1}$ and $1 \leq m \leq k$.
- (iv) $\Phi \circ \Psi$ is entanglement breaking for all *k*-PEB maps $\Psi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_1}$.

- (v) $\Phi \circ \Psi$ is entanglement breaking for all k-PEB maps $\Psi : \mathbb{M}_m \to \mathbb{M}_{d_1}$ and $m \ge 1$.
- (vi) $\Phi \circ Ad_P$ is entanglement breaking for all projections $P \in \mathbb{M}_{d_1}$ with rank $(P) = \min\{k, d_1\}$.
- (vii) $SN((id_m \otimes \Phi)(X)) = 1$ for all $X \in (\mathbb{M}_m \otimes \mathbb{M}_{d_1})^+$ with $SN(X) \leq k$ and $m \geq 1$.
- (viii) $SN((id_{d_1} \otimes \Phi)(X)) = 1$ for all $X \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_{d_1})^+$ with $SN(X) \leq k$.
- (ix) $\Gamma \circ \Phi$ is k-positive for all positive maps $\Gamma : \mathbb{M}_{d_2} \to \mathbb{M}_m$ and $m \ge 1$.
- (x) $\Gamma \circ \Phi$ is *k*-positive for all positive maps $\Gamma : \mathbb{M}_{d_2} \to \mathbb{M}_{d_1}$.
- (xi) $tr(C_{\Phi}C_{\Gamma\circ\Psi}) \geq 0$ for all k-PEB maps $\Psi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_1}$ and positive maps $\Gamma : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$.

Proof. The implications $(iii) \Rightarrow (ii)$, $(v) \Rightarrow (iv)$, $(vii) \Rightarrow (viii)$, $(iv) \Rightarrow (vi)$ and $(ix) \Rightarrow (x)$ are trivial. Now we prove the remaining implications as follows:

 $(i) \Rightarrow (ii)$ Let $\Psi \in \mathcal{CP}(k, d_1)$. Since $C_{\Psi} \in (\mathbb{M}_k \otimes \mathbb{M}_{d_1})^+$ and Φ is k-EB $C_{\Phi \circ \Psi} = (\mathrm{id}_k \otimes \Phi)(C_{\Psi})$ is separable, so that $\Phi \circ \Psi$ is an EB-map.

 $(ii) \Rightarrow (i)$ Let $X \in (\mathbb{M}_k \otimes \mathbb{M}_{d_1})^+$ and $\Psi \in \mathcal{CP}(k, d_1)$ be such that $C_{\Psi} = X$. By assumption $\Phi \circ \Psi$ is an EB-map, hence $(\mathrm{id}_k \otimes \Phi)(X) = C_{\Phi \circ \Psi}$ is separable. Since $X \in (\mathbb{M}_k \otimes \mathbb{M}_{d_1})^+$ is arbitrary this implies that Φ is a k-EB map.

 $(i) \Rightarrow (iii)$. Since Φ is *m*-EB for $1 \le m \le k$, from the equivalence of (i) and (ii), we have $\Phi \circ \Psi$ is EB for all $\Psi \in C\mathcal{P}(m, d_1)$.

 $(i) \Rightarrow (vii)$ Let $X \in (\mathbb{M}_m \otimes \mathbb{M}_{d_1})^+$ with $r := SN(X) \leq k$. Then, by Lemma 3.4, $(\mathrm{id}_m \otimes \Phi)(X)$ is positive. Further, by Lemma 3.3, there exist isometries $V_{j_i} : \mathbb{C}^j \to \mathbb{C}^m$ and vectors $\psi_{j_i} \in \mathbb{C}^j \otimes \mathbb{C}^{d_1}, 1 \leq j \leq r$ such that

$$SN((\mathsf{id}_m \otimes \Phi)(X)) = SN\Big(\sum_{i,j} (V_{j_i} \otimes I_{d_2}) \big((\mathsf{id}_j \otimes \Phi)(|\psi_{j_i}\rangle\langle\psi_{j_i}|)\big) (V_{j_i} \otimes I_{d_2})^*\Big)$$

$$\leq \max_{i,j} \{SN((V_{j_i} \otimes I_{d_2}) \big((\mathsf{id}_j \otimes \Phi)(|\psi_{j_i}\rangle\langle\psi_{j_i}|)\big) (V_{j_i} \otimes I_{d_2})^*)\}$$

$$\leq \max_{i,j} \{SN\big((\mathsf{id}_j \otimes \Phi)|\psi_{j_i}\rangle\langle\psi_{j_i}|\big)\}$$

$$= 1.$$

The last equality follows as Φ is *j*-entanglement breaking for all $1 \le j \le k$.

 $(vii) \Rightarrow (v)$ Let $\Psi \in \mathcal{PEB}_k(m, d_1)$, where $m \ge 1$. Since $C_{\Psi} \in (\mathbb{M}_m \otimes \mathbb{M}_{d_1})^+$ and $SN(C_{\Psi}) \le k$, by Lemma 3.4, $C_{\Phi \circ \Psi} = (\mathrm{id}_m \otimes \Phi)(C_{\Psi}) \in (\mathbb{M}_m \otimes \mathbb{M}_{d_2})^+$. Further, from the assumption

$$SN(C_{\Phi\circ\Psi}) = SN((\mathsf{id}_m \otimes \Phi)C_{\Psi}) = 1.$$

Thus, $\Phi \circ \Psi$ is an EB-map.

 $(iv) \Rightarrow (i)$ We prove this in two case.

<u>Case (1)</u>: Suppose $k \ge d_1$. Then $\Psi = id_{d_1}$ is a k-PEB map, hence by assumption $\Phi = \Phi \circ \Psi$ is an EB-map. In particular, Φ is a k-EB map.

<u>Case (2)</u>: Suppose $k < d_1$. Let $X \in (\mathbb{M}_k \otimes \mathbb{M}_{d_1})^+$. Without loss of generality assume that X is of rank one; hence there exists $V \in \mathbb{M}_{k \times d_1}$ such that $X = C_{Ad_V}$. Let $\iota : \mathbb{C}^k \to \mathbb{C}^{d_1}$ be the inclusion map. Note that $SN(Ad_{\iota \circ V}) = rank(\iota \circ V) \leq k$. Thus, $Ad_{\iota \circ V} \in \mathcal{PEB}_k(d_1)$, hence by assumption $C_{\Phi \circ Ad_{\iota \circ V}}$ is separable. Therefore,

$$(\mathsf{id}_k \otimes \Phi)(X) = C_{\Phi \circ Ad_V} = (\iota \otimes I_{d_2})^* C_{\Phi \circ Ad_{\iota \circ V}}(\iota \otimes I_{d_2})$$

is separable. Since $X \in (\mathbb{M}_k \otimes \mathbb{M}_{d_1})^+$ is arbitrary we conclude that Φ is *k*-EB.

 $(viii) \Rightarrow (iv)$ Let $\Psi \in \mathcal{PEB}_k(d_1)$. Since $SN(C_{\Psi}) \leq k$, by assumption $C_{\Phi \circ \Psi} = (\mathsf{id}_{d_1} \otimes \Phi)(C_{\Psi})$ is separable. Hence $\Phi \circ \Psi$ is EB.

 $(i) \Rightarrow (ix)$ Let $\Gamma \in \mathcal{P}(d_2, m)$, where $m \ge 1$ and $X \in (\mathbb{M}_k \otimes \mathbb{M}_{d_1})^+$. By assumption $(\mathrm{id}_k \otimes \Phi)(X)$ is separable, hence there exist $A_i \in \mathbb{M}_k^+$ and $B_i \in \mathbb{M}_{d_2}^+$ such that $(\mathrm{id}_k \otimes \Phi)(X) = \sum_i A_i \otimes B_i$. Then, $(\mathrm{id}_k \otimes \Gamma \circ \Phi)(X) = \sum_i A_i \otimes \Gamma(B_i) \ge 0$. We conclude that $\Gamma \circ \Phi$ is k-positive.

 $(x) \Rightarrow (iv)$ Let $\Psi \in \mathcal{PEB}(d_1)$. If $\Gamma \in \mathcal{P}(d_2, d_1)$, then by assumption $\Gamma \circ \Phi \in \mathcal{P}_k(d_1)$, and hence from (2.4) and Theorem 2.5, $\Gamma \circ \Phi \circ \Psi$ is CP. Since Γ is arbitrary, by Theorem 2.3, $\Phi \circ \Psi$ is EB.

 $(vi) \Rightarrow (iv)$ If $k \ge d_1$, then by assumption $\Phi = \Phi \circ Ad_{I_{d_1}}$ is EB so that $\Phi \circ \Psi$ is also EB for every $\Psi \in \mathcal{PEB}_k(m, d_1)$ and $m \ge 1$. Now suppose $k < d_1$. We show that $\Phi \circ Ad_V$ is EB for all $V \in \mathbb{M}_{d_1}$ with rank $(V) \le k$. By singular value decomposition there exists unitaries $U, U' \in \mathbb{M}_{d_1}$ and a diagonal matrix $D \in \mathbb{M}_r$ such that $V = U^* \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix} U'$, where r = rank(V). Then V = XP, where $X = U^* \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix} U'$ and $P = U'^* \begin{bmatrix} I_k & 0 \\ 0 & 0 \end{bmatrix} U'$, which is a projection of rank k. By assumption $\Phi \circ Ad_P$ is EB, and hence $\Phi \circ Ad_V = \Phi \circ Ad_P \circ Ad_X$ is also EB. Since $V \in \mathbb{M}_{d_1}$ is arbitrary we have $\Phi \circ \Psi$ is EB for all $\Psi \in \mathcal{PEB}_k(d_1)$.

 $(x) \Rightarrow (xi)$ Let $\Gamma \in \mathcal{P}(d_1, d_2)$ and $\Psi \in \mathcal{PEB}_k(d_1)$. By assumption, $\Gamma^* \circ \Phi$ is k-positive. Hence from (2.4), we have $0 \leq \operatorname{tr}(C_{\Gamma^* \circ \Phi}C_{\Psi}) = \operatorname{tr}(C_{\Phi}C_{\Gamma \circ \Psi})$.

 $(xi) \Rightarrow (x)$ Let $\Gamma \in \mathcal{P}(d_2, d_1)$. Then by assumption $\operatorname{tr}(C_{\Gamma \circ \Phi} C_{\Psi}) = \operatorname{tr}(C_{\Phi} C_{\Gamma^* \circ \Psi}) \ge 0$ for all $\Psi \in \mathcal{PEB}_k(d_1)$. Now from (2.4) it follows that $\Gamma \circ \Phi$ is k-positive. \Box

Note 3.6. From (iv) it follows that if Φ is k-EB, then $\Phi \circ Ad_P$ is entanglement breaking for all projections $P \in \mathbb{M}_{d_1}$ with rank less than or equal to k. The converse is also true because of (vi). This equivalence was observed in [ChCh20, Lemma 6.1].

From Theorem 2.3, we have $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is an EB-map if and only if the Choi matrix $C_{\Phi} = \sum_{i,j} E_{ij} \otimes \Phi(E_{ij})$ is separable, where $\{E_{ij}\}_{i,j=1}^{d_1}$ is the standard matrix units in \mathbb{M}_{d_1} . Next we prove an analogue of this result in the context of k-EB maps. Suppose $\{F_{ij}\}_{i,j=1}^d$ is a complete set of matrix units in \mathbb{M}_{d_1} (i.e., there exists an orthonormal basis $\{f_i\}_{i=1}^d$ for \mathbb{C}^d such that $F_{ij} = |f_i\rangle\langle f_j|$) and $U \in \mathbb{M}_{d_1}$ is a unitary such that $UF_{ij}U^* = E_{ij}$, for all $1 \leq i, j \leq d_1$. Then, by [Sto13, Lemma 4.1.2], the Choi matrix C_{Φ}^F w.r.t $\{F_{ij}\}_{i,j}$ is given by

$$C_{\Phi}^F := \sum_{i,j=1}^{d_1} F_{ij} \otimes \Phi(F_{ij}) = Ad_{U \otimes I_{d_2}} C_{\Phi \circ Ad_U}.$$
(3.3)

Definition 3.7. Suppose $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is a linear map and $\{F_{ij}\}_{i,j=1}^{d_1}$ is a complete set of matrix units in \mathbb{M}_{d_1} . Given $1 \le k \le d_1$ and $i_1, i_2, \cdots, i_k \in \{1, 2, \cdots, d_1\}$ the matrix

$$C_{\Phi(i_1,\cdots,i_k)}^F := \sum_{p,q=1}^k E_{pq}^{(k)} \otimes \Phi(F_{i_p i_q}) = \begin{bmatrix} \Phi(F_{i_1 i_1}) & \Phi(F_{i_1 i_2}) & \cdots & \Phi(F_{i_1 i_k}) \\ \vdots & \vdots & \cdots & \vdots \\ \Phi(F_{i_k i_1}) & \Phi(F_{i_k i_2}) & \cdots & \Phi(F_{i_k i_k}) \end{bmatrix}$$

in $\mathbb{M}_k \otimes \mathbb{M}_{d_2} = \mathbb{M}_k(\mathbb{M}_{d_2})$ is called a *principal* $(k \times k)$ -block submatrix of C_{Φ}^F , where $E_{ij}^{(k)} \in \mathbb{M}_k$ is the standard matrix units. The principal $(k \times k)$ -block submatrix $C_{\Phi(i_1, \dots, i_k)}^E$ of C_{Φ} w.r.t the standard matrix units $\{E_{ij}\}_{i,j} \subseteq \mathbb{M}_{d_1}$ is denoted simply by $C_{\Phi(i_1, \dots, i_k)}$.

Let $\{e_i^{(k)}\}_{i=1}^k$ be the standard orthonormal basis for \mathbb{C}^k and $\{f_i\}_{i=1}^{d_1}$ be an orthonormal basis for \mathbb{C}^{d_1} such that $F_{ij} = |f_i\rangle\langle f_j|$. Then

$$C^F_{\Phi(i_1,\cdots,i_k)} = (\mathsf{id}_k \otimes \Phi)(\sum_{p,q=1}^k E^{(k)}_{pq} \otimes F_{i_p i_q}) = (\mathsf{id}_k \otimes \Phi)(|\sum_{p=1}^k e^{(k)}_p \otimes f_{i_p}\rangle \langle \sum_{q=1}^k e^{(k)}_q \otimes f_{i_q}|).$$

Note that $|\sum_{p=1}^{k} e_p^{(k)} \otimes f_{i_p}\rangle \langle \sum_{q=1}^{k} e_q^{(k)} \otimes f_{i_q}| \in (\mathbb{M}_k \otimes \mathbb{M}_{d_1})^+$. Hence, $C_{\Phi(i_1,\cdots,i_k)}^F$ is positive (resp. separable) if Φ is k-positive (resp. k-EB).

Theorem 3.8. Let $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ be a *k*-positive map, where $1 \le k \le d_1$. Then the following conditions are equivalent:

- (i) Φ is k-EB.
- (ii) Given a complete set of matrix units $\{F_{ij}\}_{i,j=1}^{d_1} \subseteq \mathbb{M}_{d_1}$, every principal $(k \times k)$ -block submatrix $C_{\Phi(i_1,\cdots,i_k)}^F$ of the Choi matrix C_{Φ}^F is separable.
- (iii) Given a complete set of matrix units $\{F_{ij}\}_{i,j=1}^{d_1} \subseteq \mathbb{M}_{d_1}$, the principal $(k \times k)$ -block submatrix $C_{\Phi^{(1,2,\cdots,k)}}^F$ of the Choi matrix C_{Φ}^F is separable.

Proof. We need to prove $(iii) \Rightarrow (i)$ only. So let $P \in \mathbb{M}_{d_1}$ be a projection of rank k and $\widetilde{\Phi} = \Phi \circ Ad_P$. Suppose $\{f_i\}_{i=1}^{d_1} \subseteq \mathbb{C}^{d_1}$ is an orthonormal basis such that $P = \sum_{i=1}^k |f_i\rangle\langle f_i|$. Let $F_{ij} = |f_i\rangle\langle f_j|$ and $U \in \mathbb{M}_{d_1}$ be a unitary such that $UF_{ij}U^* = E_{ij} \in \mathbb{M}_{d_1}$ for all $1 \le i, j \le d_1$. Since $\{F_{ij}\}_{ij}$ is a complete set of matrix units, from (3.3), we have

$$C_{\widetilde{\Phi} \circ Ad_{U}} = Ad_{U^{*} \otimes I_{d_{2}}} C_{\widetilde{\Phi}}^{F} = Ad_{U^{*} \otimes I_{d_{2}}} \Big(\sum_{i,j=1}^{d_{1}} F_{ij} \otimes \widetilde{\Phi}(F_{ij}) \Big) = Ad_{U^{*} \otimes I_{d_{2}}} \Big(\sum_{i,j=1}^{k} F_{ij} \otimes \Phi(F_{ij}) \Big).$$

Now let $W \in \mathbb{M}_{d_1 \times k}$ be such that $W(e_i^{(k)}) = f_i$ for all $1 \le i \le k$. Then

$$C_{\widetilde{\Phi} \circ Ad_U} = Ad_{U^* \otimes I_{d_2}} (W \otimes I_{d_2}) \Big(\sum_{i,j=1}^k E_{ij}^{(k)} \otimes \Phi(F_{ij}) \Big) (W \otimes I_{d_2})^*$$
$$= (UW \otimes I_{d_2}) C_{\Phi(1,\cdots,k)}^F (UW \otimes I_{d_2})^*.$$

Since $C_{\Phi(1,\dots,k)}^F$ is separable we conclude that $C_{\widetilde{\Phi}\circ Ad_U}$ is separable. Hence $\Phi \circ Ad_P = \widetilde{\Phi} \circ Ad_U \circ Ad_U \circ Ad_{U^*}$ is an EB-map. Further, since P is arbitrary, by Theorem 3.5, Φ is k-EB.

Remark 3.9. If $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is a *k*-EB map, where $k \ge 1$, then $\mathsf{T} \circ \Phi \circ \mathsf{T}$ is also a *k*-EB map. For, let $\Psi \in \mathcal{CP}(k, d_1)$. Since $\mathsf{T} \circ \Psi \circ \mathsf{T} \in \mathcal{CP}(k, d_1)$, by Theorem 3.5, $\Phi \circ (\mathsf{T} \circ \Psi \circ \mathsf{T})$ is EB. Hence

$$\mathsf{F} \circ \Phi \circ \mathsf{T} \circ \Psi = \mathsf{T} \circ (\Phi \circ \mathsf{T} \circ \Psi \circ \mathsf{T}) \circ \mathsf{T}$$

is also an EB-map. Since Ψ is arbitrary, by Theorem 3.5, $\mathsf{T} \circ \Phi \circ \mathsf{T}$ is *k*-EB.

Theorem 3.10. Let $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ be a *k*-positive map, where $k \ge 1$. Then the following conditions are equivalent:

- (i) Φ is *k*-entanglement breaking.
- (ii) $(\Phi \otimes id_k)(X)$ is separable for all $X \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_k)^+$.

Proof. $(i) \Rightarrow (ii)$ Let $X \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_k)^+$, and without loss of generality assume that $X = |\psi\rangle\langle\psi|$ for some $\psi \in \mathbb{C}^{d_1} \otimes \mathbb{C}^k$. By [CMW19, Lemma 1.3], there exists $V \in \mathbb{M}_{k \times d_1}$ such that $\psi = (I_{d_1} \otimes V)(\Omega_{d_1})$. Further, applying [CMW19, Lemma 1.1] to the k-positive map Φ , we get

$$\begin{split} (\Phi \otimes \mathsf{id}_k)(X) &= (\Phi \otimes \mathsf{id}_k) \big((I_{d_1} \otimes V) (|\Omega_{d_1}\rangle \langle \Omega_{d_1}|) (I_{d_1} \otimes V)^* \big) \\ &= (\Phi \otimes \mathsf{id}_k) (\mathsf{id}_{d_1} \otimes Ad_{V^*}) (|\Omega_{d_1}\rangle \langle \Omega_{d_1}|) \\ &= (\mathsf{id}_{d_2} \otimes Ad_{V^*}) (\Phi \otimes \mathsf{id}_{d_1}) (|\Omega_{d_1}\rangle \langle \Omega_{d_1}|) \\ &= (\mathsf{id}_{d_2} \otimes Ad_{V^*}) (\mathsf{id}_{d_2} \otimes \mathsf{T} \circ \Phi^* \circ \mathsf{T}) (|\Omega_{d_2}\rangle \langle \Omega_{d_2}|) \\ &= \big(\mathsf{id}_{d_2} \otimes (\mathsf{T} \circ \Phi \circ \mathsf{T} \circ Ad_V)^* \big) (|\Omega_{d_2}\rangle \langle \Omega_{d_2}|). \end{split}$$

Since $\mathsf{T} \circ \Phi \circ \mathsf{T}$ is *k*-EB, $\mathsf{T} \circ \Phi \circ \mathsf{T} \circ Ad_V : \mathbb{M}_k \to \mathbb{M}_{d_2}$ is EB. It follows from the above equation that $(\Phi \otimes \mathsf{id}_k)(X)$ is separable.

 $(ii) \Rightarrow (i)$ Let $V \in \mathbb{M}_{k \times d_1}$. Then $X = (\mathrm{id}_{d_1} \otimes Ad_{V^*})(|\Omega_{d_1}\rangle\langle\Omega_{d_1}|) \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_k)^+$, and the Choi-matrix of $Ad_{V^*} \circ \mathsf{T} \circ \Phi^* \circ \mathsf{T} : \mathbb{M}_{d_2} \to \mathbb{M}_k$ is given by

$$(\mathsf{id}_{d_2} \otimes Ad_{V^*} \circ \mathsf{T} \circ \Phi^* \circ \mathsf{T})(|\Omega_{d_2}\rangle \langle \Omega_{d_2}|) = (\Phi \otimes \mathsf{id}_k)(\mathsf{id}_{d_1} \otimes Ad_{V^*})(|\Omega_{d_1}\rangle \langle \Omega_{d_1}|) = (\Phi \otimes \mathsf{id}_k)(X),$$

which is separable by assumption. Thus $Ad_{V^*} \circ \mathsf{T} \circ \Phi^* \circ \mathsf{T}$ (and hence $\mathsf{T} \circ \Phi \circ \mathsf{T} \circ Ad_V$) is EB. Since V is arbitrary, by Theorem 3.5, $\mathsf{T} \circ \Phi \circ \mathsf{T}$ is k-EB. Now, from Remark 3.9, we conclude that Φ is k-EB.

Note 3.11. Though the tensor product of CP-maps is a CP-map, it is not true in general that the tensor product of positive maps is a positive map. However, if $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is a *k*-EB map and $\Gamma : \mathbb{M}_k \to \mathbb{M}_{d_3}$ is any positive map, then from the definition of *k*-EB maps and Theorem 3.10, it follows that

 $(\Phi \otimes \Gamma)(X) = (\mathsf{id}_{d_2} \otimes \Gamma)(\Phi \otimes \mathsf{id}_k)(X) \text{ and } (\Gamma \otimes \Phi)(Y) = (\Gamma \otimes \mathsf{id}_{d_2})(\mathsf{id}_k \otimes \Phi)(Y)$

are separable for all $X \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_k)^+$ and $Y \in (\mathbb{M}_k \otimes \mathbb{M}_{d_1})^+$. In particular, $\Phi \otimes \Gamma$ and $\Gamma \otimes \Phi$ are positive maps.

Theorem 3.12. Let $\mathcal{EB}_k(d_1, d_2)$ and $\mathcal{EBCP}_k(d_1, d_2)$ be the set of all *k*-entanglement breaking linear maps and *k*-entanglement breaking CP-maps from \mathbb{M}_{d_1} into \mathbb{M}_{d_2} , respectively. Then both $\mathcal{EB}_k(d_1, d_2)$ and $\mathcal{EBCP}_k(d_1, d_2)$ are mapping cones.

Proof. Suppose $\{\Theta_n\}_{n=1}^{\infty}$ is a sequence in $\mathcal{EB}_k(d_1, d_2)$ such that $\Theta_n \to \Theta \in \mathcal{P}(d_1, d_2)$. Given any $\Psi \in \mathcal{CP}(k, d_1)$, by Theorem 3.5, $\Theta_n \circ \Psi$ is EB for every $n \ge 1$, and hence so is $\Theta \circ \Psi = \lim_n \Theta_n \circ \Psi$. Thus $\Theta \in \mathcal{EB}_k(d_1, d_2)$, and concludes that $\mathcal{EB}_k(d_1, d_2)$ is closed. Further, from Remark 3.2, it follows that $\mathcal{EB}_k(d_1, d_2)$ is a mapping cone. Now, since $\mathcal{CP}_k(d_1, d_2)$ is a mapping cone the intersection

$$\mathcal{EB}_k(d_1, d_2) \bigcap \mathcal{CP}_k(d_1, d_2) = \mathcal{EBCP}_k(d_1, d_2)$$

is also a mapping cone.

The smallest closed convex cone containing a set S will be denoted by conv(S).

Theorem 3.13. Let $k \ge 1$. Then¹

$$\mathcal{EB}_k(d_1, d_2)^\circ = conv\{\Gamma \circ \Psi : \Gamma \in \mathcal{P}(d_1, d_2), \Psi \in \mathcal{PEB}_k(d_1)\}$$
(3.4)

$$= conv\{\Gamma \circ \Psi : \Gamma \in \mathcal{P}(k, d_2), \Psi \in \mathcal{CP}(d_1, k)\}.$$
(3.5)

Proof. Let $S = \{\Gamma \circ \Psi : \Gamma \in \mathcal{P}(d_1, d_2), \Psi \in \mathcal{PEB}_k(d_1)\}$. By Theorem 3.5(xi), we have $S \subseteq \mathcal{EB}_k(d_1, d_2)^\circ$, and hence $conv(S) \subseteq \mathcal{EB}_k(d_1, d_2)^\circ$. Now to prove the reverse inclusion, it is enough to prove that $(conv(S))^\circ \subseteq (\mathcal{EB}_k(d_1, d_2))^{\circ\circ} = \mathcal{EB}_k(d_1, d_2)$. So let $\Phi \in (conv(S))^\circ$. Then

$$\operatorname{tr}(C_{\Phi\circ\Psi}C_{\Gamma}) = \operatorname{tr}(C_{\Phi}C_{\Gamma\circ\Psi^*}) \ge 0$$

for all $\Gamma \in \mathcal{P}(d_1, d_2)$ and $\Psi \in \mathcal{PEB}_k(d_1)$, hence $\Phi \circ \Psi \in \mathcal{EB}(d_1, d_2)$. Since Ψ is arbitrary, by Theorem 3.5(iv), $\Phi \in \mathcal{EB}_k(d_1, d_2)$. This completes the proof of (3.4). Now to prove (3.5), let

$$\mathcal{C}_k(d_1, d_2) := conv\{\Gamma \circ \Psi : \Gamma \in \mathcal{P}(k, d_2), \Psi \in \mathcal{CP}(d_1, k)\}$$

To show that $\mathcal{C}_k(d_1, d_2) \subseteq \mathcal{EB}_k(d_1, d_2)^\circ$ note that

$$\operatorname{tr}(C_{\Gamma\circ\Psi}C_{\Phi}) = \operatorname{tr}\left(C_{\Gamma}(\operatorname{id}_k\otimes\Phi)(C_{\Psi^*})\right) \ge 0$$

for all $\Phi \in \mathcal{EB}_k(d_1, d_2), \Gamma \in \mathcal{P}(k, d_2)$ and $\Psi \in \mathcal{CP}(d_1, k)$. To show the reverse inequality it is enough to show that $\mathcal{C}_k(d_1, d_2)^\circ \subseteq \mathcal{EB}_k(d_1, d_2)$. To see this, consider $\Phi \in \mathcal{C}_k(d_1, d_2)^\circ$. By definition, we know that

$$\operatorname{tr}(C_{\Phi\circ\Psi}C_{\Gamma}) = \operatorname{tr}(C_{\Phi}C_{\Gamma\circ\Psi^*}) \ge 0$$

¹The identity (3.5) was proved by Alexander Müller-Hermes and communicated to us privately after we upload the preprint in arXiv. This proof leads us to prove the identity (3.4).

for all $\Gamma \in \mathcal{P}(k, d_2)$ and $\Psi \in \mathcal{CP}(k, d_1)$. We conclude that $\Phi \circ \Psi \in \mathcal{EB}(k, d_2)$ for every $\Psi \in \mathcal{CP}(k, d_1)$, which is equivalent to $\Phi \in \mathcal{EB}_k(d_1, d_2)$.

Remark 3.14. Now, using [Sto18, Corollary 4], we can compute the dual of \mathcal{EBCP}_k as $\mathcal{EBCP}_k^{\circ} = (\mathcal{EB}_k \cap \mathcal{CP})^{\circ} = \mathcal{EB}_k^{\circ} \vee \mathcal{CP}^{\circ}$. Here, $\mathcal{C}_1 \vee \mathcal{C}_2$ denotes the smallest closed convex cone containing the union of two mapping cones \mathcal{C}_1 and \mathcal{C}_2 . Note that $\mathcal{C}_1 \vee \mathcal{C}_2 = \overline{\mathcal{C}_1 + \mathcal{C}_2}$, where $\mathcal{C}_1 + \mathcal{C}_2 := \{\Theta_1 + \Theta_2 : \Theta_i \in \mathcal{C}_i, i = 1, 2\}$.

From Theorem 2.5 and Theorem 3.5(v), it follows that $\mathcal{PEB}_k(d_1, d_2) \subseteq \mathcal{EB}_k(d_1, d_2)^\circ$ Thus we have the following:

We denote the set of all PPT-maps from \mathbb{M}_{d_1} into \mathbb{M}_{d_2} by $\mathcal{PPT}(d_1, d_2)$. From Holevo form, it follows that every EB-map is a PPT-map. But the converse is not true in general. In [Hor97], Horodecki proved that there exists $\Psi \in \mathcal{PPT}(2, 4)$ which is not EB (equivalently, not 2-EB). Now, given d > 4 let $W = \begin{bmatrix} I_4 & \mathbf{0}_{4 \times (d-4)} \end{bmatrix} \in \mathbb{M}_{4 \times d}$. Then the map $\Psi' := Ad_W \circ \Psi : \mathbb{M}_2 \to \mathbb{M}_d$ is PPT but not EB since $Ad_{W*} \circ \Psi' = \Psi$ is not EB. Thus, for every $d \ge 4$ there is a PPT-map $\Psi' : \mathbb{M}_2 \to \mathbb{M}_d$ that is not EB.

Proposition 3.15. Let $d \ge 4$. Then there exists a PPT map $\Phi : \mathbb{M}_d \to \mathbb{M}_d$ which is not 2-EB. Moreover, we can choose Φ such that $SN(\Phi) = SN(\Phi \circ T) = 2$.

Proof. Choose $\Psi \in \mathcal{PPT}(2,d)$ that is not EB. Let $V = \begin{bmatrix} I_2 \\ \mathbf{0}_{(d-2)\times 2} \end{bmatrix} \in \mathbb{M}_{d\times 2}$. Then the map $\Phi := \Psi \circ Ad_V$ defined on \mathbb{M}_d is PPT, and both $SN(\Phi)$ and $SN(\Phi \circ \mathsf{T}) = SN(\mathsf{T} \circ \Phi)$ are less than or equal to $SN(Ad_V) = 2$. Since $\Phi \circ Ad_{V*} = \Psi$ is not an EB-map, by Theorem 3.5, Φ is not 2-EB. Observe that Φ and $\Phi \circ \mathsf{T}$ has Schmidt number strictly greater than one as they are not EB. This completes the proof.

From the above discussion, we saw that for all $d \ge 4$ there exists a $\Psi \in \mathcal{PPT}(d, 2)$ which is not EB. The following proposition, which is a generalization of [HHH96, Theorem 3], says that such a map will always be 3-EB.

Proposition 3.16. Let $d \ge 1$.

- (i) If $\Phi : \mathbb{M}_d \to \mathbb{M}_2$ is a PPT-map, then Φ is 3-EB.
- (ii) If $\Phi : \mathbb{M}_d \to \mathbb{M}_3$ is a PPT-map, then Φ is 2-EB.

Proof. (i) Let $\Psi \in C\mathcal{P}(3, d)$. Then $\Phi \circ \Psi \in \mathcal{PPT}(3, 2)$, and hence by [HHH96, Theorem 3], it is EB. Since Ψ is arbitrary, from Theorem 3.5, it follows that Φ is 3-EB.

(*ii*) Follows from the fact that $\Phi \circ \Psi \in \mathcal{PPT}(2,3)$, and hence EB for every $\Psi \in \mathcal{CP}(2,d)$. \Box

A mapping cone $\mathcal{C} \subseteq \mathcal{P}(d) := \mathcal{P}(d, d)$ is said to be symmetric ([Sto11]) if $\Theta^*, \mathsf{T} \circ \Theta \circ \mathsf{T} \in \mathcal{C}$ for all $\Theta \in \mathcal{C}$. If \mathcal{C} is a symmetric mapping cone, then \mathcal{C}° is also symmetric. The mapping cones $\mathcal{P}(d), \mathcal{P}_k(d), \mathcal{CP}(d), \mathcal{PEB}_k(d), \mathcal{EB}(d)$, where k > 1, are known to be symmetric. The following theorem says that, in general, the mapping cone $\mathcal{EB}_k(d)$ (and hence $\mathcal{EBCP}_k(d)$) is not symmetric for $d \geq 4$.

Theorem 3.17. Let $d \ge 4$.

- (i) There exists a 3-EB map $\Phi : \mathbb{M}_d \to \mathbb{M}_2$ such that Φ^* is not 2-EB (and hence not 3-EB).
- (ii) There exists a 3-EB map $\Phi : \mathbb{M}_d \to \mathbb{M}_d$ such that Φ^* is not 2-EB (and hence not 3-EB).

Proof. (*i*) Let $\Phi : \mathbb{M}_d \to \mathbb{M}_2$ be a linear map such that Φ^* is PPT-map but not EB. Since Φ is PPT, by Proposition 3.16, Φ is 3-EB. But $\Phi^* : \mathbb{M}_2 \to \mathbb{M}_d$ is not 2-EB (equivalently not EB). (*ii*) Let $\Phi' \in \mathcal{EB}_3(d, 2)$ be such that Φ'^* is not 2-EB. Take $\Phi = \iota \circ \Phi'$, where $\iota : \mathbb{M}_2 \to \mathbb{M}_d$ is the inclusion map. Observe that $\Phi \in \mathcal{EB}_3(d)$. Now, if Φ^* is 2-EB, then $\Phi^* \circ \iota = \Phi'^* \circ \iota^* \circ \iota = \Phi'^*$ is also 2-EB, which is not possible. Hence Φ^* is not 2-EB.

Remark 3.18. From Remark 3.2 it follows that $\mathcal{EB}_k(d_1, d_2)$ is a left-invariant mapping cone in the sense that it is invariant under composition by positive maps from the left side. However, $\mathcal{EB}_k(d_1, d_2)$ is not right-invariant in general. If $\mathcal{EB}_k(d_1, d_2)$ is right-invariant, then from Theorem 3.5 (ix) it follows that the adjoint of a *k*-EB map is always *k*-EB which is not true in general, as seen in the above theorem.

Remark 3.19. We know that every d_1 -EB map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is EB. However, from the above Theorem, we observe that a d_2 -EB map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is not necessarily an entanglement breaking map.

Let $\mathcal{C}, \mathcal{C}_1, \mathcal{C}_2$ be mapping cones of $\mathcal{P}(d_1, d_2)$. If \mathcal{C} is any of $\mathcal{P}, \mathcal{P}_k, \mathcal{CP}, \mathcal{PEB}_k, \mathcal{EB}$, then

$$\mathcal{C} \circ \mathsf{T} := \{ \Phi \circ \mathsf{T} : \Phi \in \mathcal{C} \}$$

is a mapping cone. A mapping cone arises from $\mathcal{P}, \mathcal{P}_k, \mathcal{CP}, \mathcal{PEB}_k, \mathcal{EB}$, via the operations $\mathcal{C} \mapsto \mathcal{C} \circ \mathsf{T}, (\mathcal{C}_1, \mathcal{C}_2) \mapsto \mathcal{C}_1 \cap \mathcal{C}_2$, or $(\mathcal{C}_1, \mathcal{C}_2) \mapsto \mathcal{C}_1 \vee \mathcal{C}_2$ is called *typical*, otherwise called *untypical* ([Sko11]). In [JSS13], the authors discussed examples and properties of untypical mapping cones on \mathbb{M}_d , emphasizing the case d = 2. Observe that typical mapping cones are symmetric. By Theorem 3.17, $\mathcal{EB}_k(d)$ and $\mathcal{EBCP}_k(d), k = 2, 3$ are not symmetric for $d \geq 4$, hence not typical. Actually much more is true, as the following theorem shows.

Theorem 3.20. Let $d \ge 4$ and 1 < k < d. Then the mapping cones $\mathcal{EB}_k(d)$ and $\mathcal{EBCP}_k(d)$ are not typical.

Proof. By Proposition 3.15, there exists a map $\Phi \in \mathcal{PEB}_2(d) \cap (\mathcal{PEB}_2(d) \circ \mathsf{T})$ which is not 2-EB, hence $\Phi \notin \mathcal{EB}_k(d)$ and $\Phi \notin \mathcal{EBCP}_k(d)$. Now, from [JSS13, Lemma 12], it follows that $\mathcal{EB}_k(d)$ and $\mathcal{EBCP}_k(d)$ are not typical.

Recall that a positive map is said to be *decomposable* if it can be written as a sum of a CPmap and a co-CP-map. Characterization of CP-maps and PPT-maps in terms of decomposable maps is known in the literature. See [PiMo07, Theorem], [Sto18, Proposition 8]. Next, we establish a similar characterization for entanglement breaking maps.

Theorem 3.21. Let $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ be a *k*-entanglement breaking map, where $d_1, k > 1$. Then the following conditions are equivalent:

- (i) Φ is entanglement breaking.
- (ii) $id_n \otimes \Gamma \circ \Phi$ is decomposable for all positive maps $\Gamma : \mathbb{M}_{d_2} \to \mathbb{M}_m$, where $m, n \ge 1$.
- (iii) $id_k \otimes \Gamma \circ \Phi$ is decomposable for all positive maps $\Gamma : \mathbb{M}_{d_2} \to \mathbb{M}_{d_1}$.

Proof. $(i) \Rightarrow (ii)$ Let $m \ge 1$ and $\Gamma \in \mathcal{P}(d_2, m)$. Since Φ is entanglement breaking $\Gamma \circ \Phi$ is CP, and hence $\mathrm{id}_n \otimes \Gamma \circ \Phi$ is decomposable – it is CP itself for all $n \ge 1$. $(ii) \Rightarrow (iii)$ Clear.

 $(iii) \Rightarrow (i)$ Suppose Φ is not an EB-map. Then there exists a $\Gamma \in \mathcal{P}(d_2, d_1)$ such that $\Gamma \circ \Phi$ is not CP. Since Φ is *k*-EB, by Theorem 3.5, $\Gamma \circ \Phi \in \mathcal{P}_k(d_1)$. Since $\Gamma \circ \Phi \notin \mathcal{CP}(d_1)$, by [PiMo07, Theorem], we have $\mathrm{id}_k \otimes \Gamma \circ \Phi$ is not decomposable, which is a contradiction. This completes the proof.

4. Examples

In this section, we discuss some examples of k-EB maps. Recall that the Schur product of two matrices $A = [a_{ij}], B = [b_{ij}] \in \mathbb{M}_d$ is denoted and defined by $A \circ B := [a_{ij}b_{ij}] \in \mathbb{M}_d$. Define the map $S_A : \mathbb{M}_d \to \mathbb{M}_d$ by $S_A(X) = A \circ X$ for all $X \in \mathbb{M}_d$. It is well known (c.f. [Pau02]) that S_A is CP if and only if S_A is positive if and only if A is positive. Further, S_A is PPT if and only if S_A is EB if and only if A is positive diagonal. See [RJP18, KMP18] for details.

Proposition 4.1. Let $A \in \mathbb{M}_d^+$. Then the following conditions are equivalent:

- (i) S_A is 2-EB.
- (ii) S_A is k-EB, where 2 < k < d.
- (iii) S_A is EB.

Proof. Clearly $(iii) \Rightarrow (ii) \Rightarrow (i)$. Now, to prove $(i) \Rightarrow (iii)$, assume that S_A is 2-EB. To show that S_A is EB, it is enough to prove that A is a diagonal matrix. Given $1 \le i < j \le d$ let $V_{ij} \in \mathbb{M}_{2\times d}$ be the matrix with (1,i) and (2,j) entry equals one and zero elsewhere. By Theorem 3.5, $S_A \circ Ad_{V_{ij}}$ is EB, and hence

$$(\mathsf{id}\otimes\mathsf{T})C_{S_A\circ Ad_{V_{ij}}} = \begin{bmatrix} a_{ii}E_{ii} & a_{ij}E_{ji} \\ a_{ji}E_{ij} & a_{jj}E_{jj} \end{bmatrix} \in (\mathbb{M}_2\otimes\mathbb{M}_d)^+,$$

where $E_{kl} \in \mathbb{M}_d$ are the matrix units. But this is possible only if $\begin{bmatrix} 0 & a_{ij} \\ a_{ji} & 0 \end{bmatrix} \in \mathbb{M}_2^+$, and which implies that $a_{ij} = 0 = a_{ji}$. Thus A is a diagonal matrix, and conclude that S_A is EB.

Proposition 4.2. Given $V \in \mathbb{M}_{d_1 \times d_2}$ the following conditions are equivalent:

(i) Ad_V is 2-EB.

- (ii) Ad_V is k-EB, where 2 < k < d.
- (iii) Ad_V is EB.

Proof. To prove the non-trivial direction $(i) \Rightarrow (iii)$, assume that Ad_V is 2-EB. We show that $\operatorname{rank}(V) = 1$ so that Ad_V is EB. If possible assume that $k = \operatorname{rank}(V) > 1$. Let $W \in \mathbb{M}_{2 \times d_1}$ be such that $\operatorname{rank}(WV) > 1$. Then $Ad_{WV} : \mathbb{M}_2 \to \mathbb{M}_{d_2}$ is not EB. But, Theorem 3.5(ii) implies that $Ad_V \circ Ad_W = Ad_{WV}$ is EB, which is a contradiction. Hence $\operatorname{rank}(V) = 1$. This completes the proof.

In the following, we consider linear maps satisfying some equivariance property, which is significant in the study of k-positivity of linear maps defined on matrix spaces ([COS18]) and entanglement detection ([BCS20]). A linear map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is said to be *equivariant*, if for every unitary $U \in \mathbb{M}_{d_1}$ there exists a (not necessarily unitary) matrix $V(U) \in \mathbb{M}_{d_2}$ such that $\Phi \circ Ad_U = Ad_{V(U)} \circ \Phi$.

Proposition 4.3 ([COS18]). Let $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ be an equivariant map and $1 \le k \le d_1$. Then the following conditions are equivalent:

- (i) Φ is *k*-positive.
- (ii) The principle $(k \times k)$ -block submatrix $C_{\Phi(1,2,\cdots,k)}$ is positive.
- (iii) $\Phi \circ Ad_P$ is CP, where $P = \begin{bmatrix} I_k & \mathbf{0}_{k \times d_1 k} \end{bmatrix} \in \mathbb{M}_{k \times d_1}$.

The characterization (iii) is not there in [COS18]. But through the same lines of proof of the following theorem, which is an analogue of the above result for k-EB maps, we can get a simple alternative proof of Proposition 4.3. We leave the details to the reader.

Theorem 4.4. Let $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ be an equivariant k-positive map, where $1 \leq k \leq d_1$. Then the following conditions are equivalent:

- (i) Φ is k-EB.
- (ii) The principal $(k \times k)$ -block submatrix $C_{\Phi(1,2,\cdots,k)} \in (\mathbb{M}_k \otimes \mathbb{M}_{d_2})^+$ is separable.
- (iii) $\Phi \circ Ad_P$ is entanglement breaking, where $P = \begin{bmatrix} I_k & \mathbf{0}_{k \times d_1 k} \end{bmatrix} \in \mathbb{M}_{k \times d_1}$.

Proof. $(i) \Rightarrow (ii)$ Follows from Theorem 3.8. $(ii) \Rightarrow (iii)$ Let $E_{ij}^{(d_1)} \in \mathbb{M}_{d_1}$ and $E_{ij}^{(k)} \in \mathbb{M}_k$ be the standard matrix units. Consider the Choi matrix

$$C_{\Phi \circ Ad_P} = \sum_{i,j=1}^{k} E_{ij}^{(k)} \otimes \Phi(P^* E_{ij}^{(k)} P) = \sum_{i,j=1}^{k} E_{ij}^{(k)} \otimes \Phi(E_{ij}^{(d_1)}) = C_{\Phi(1,2,\cdots,k)},$$

which is separable by assumption, and hence $\Phi \circ Ad_P$ is EB.

 $(iii) \Rightarrow (i)$ Let $Q \in \mathbb{M}_{k \times d_1}$. By singular value decomposition, there exist unitary matrices $W \in \mathbb{M}_k, U \in \mathbb{M}_{d_1}$ and a rectangular matrix $\Sigma = \begin{bmatrix} D & \mathbf{0}_{k \times (d_1 - k)} \end{bmatrix} \in \mathbb{M}_{k \times d_1}$, where $D \in \mathbb{M}_k$ is a diagonal matrix, such that $Q = W\Sigma U$. Set $\widetilde{W} = WD \in \mathbb{M}_k$. Then $\widetilde{W}PU = W\Sigma U = Q$. Now since Φ is equivariant, corresponding to the unitary $U \in \mathbb{M}_{d_1}$ there exists $V(U) \in \mathbb{M}_{d_2}$ such

that $\Phi \circ Ad_U = Ad_{V(U)} \circ \Phi$. Hence

$$\Phi \circ Ad_Q = \Phi \circ Ad_U \circ Ad_P \circ Ad_{\widetilde{W}} = Ad_{V(U)} \circ \Phi \circ Ad_P \circ Ad_{\widetilde{W}}.$$

By assumption $\Phi \circ Ad_P$ is EB, and hence from the above equation, it follows that $\Phi \circ Ad_Q$ is also EB. Since $Q \in \mathbb{M}_{k \times d_1}$ is arbitrary, by Theorem 3.5, Φ is *k*-EB.

Remark 4.5. In the above theorem, $(i) \Rightarrow (ii) \Rightarrow (iii)$ holds for every $\Phi \in \mathcal{P}_k(d_1, d_2)$.

Next, we consider a particular example of an equivariant map, namely the Holevo-Werner map ([WeHo02]). Given $\lambda \in \mathbb{R}$ the Holevo-Werner map $\mathcal{W}_{\lambda} : \mathbb{M}_d \to \mathbb{M}_d$ is given by

$$\mathcal{W}_{\lambda}(X) = \operatorname{tr}(X)I - \lambda X^{\mathsf{T}}.$$

It is well-known ([CMW19, WeHo02]) that

$$\mathcal{W}_{\lambda} \text{ is CP } \iff \lambda \in [-1, 1]$$
$$\mathcal{W}_{\lambda} \text{ is PPT } \iff \mathcal{W}_{\lambda} \text{ is EB } \iff \lambda \in [-1, 1/d]$$
$$\mathcal{W}_{\lambda} \text{ is 2-EB } \iff \lambda \in [-1, 1/2].$$

In the following, we discuss when do they become k-EB for k > 2. First, we prove the following technical lemma.

Lemma 4.6. Let $\Phi_i : \mathbb{M}_d \to \mathbb{M}_{d_i}, i = 1, 2$ and $\Phi : \mathbb{M}_d \to \mathbb{M}_{d_1+d_2}$ be linear maps such that $\Phi(X) = \begin{bmatrix} \Phi_1(X) & 0\\ 0 & \Phi_2(X) \end{bmatrix}$

for all $X \in \mathbb{M}_d$. Then Φ is *k*-EB if and only if Φ_i 's are *k*-EB.

Proof. Assume that Φ is k-EB. Let $\Psi \in C\mathcal{P}(k, d)$. Then $\Phi \circ \Psi$ is EB, and hence $Ad_{V_i^*} \circ \Phi \circ \Psi$ is also EB, where

$$V_1 = \begin{bmatrix} I_{d_1} & \mathbf{0}_{d_1 \times d_2} \end{bmatrix} \in \mathbb{M}_{d_1 \times (d_1 + d_2)} \quad \text{and} \quad V_2 = \begin{bmatrix} \mathbf{0}_{d_2 \times d_1} & I_{d_2} \end{bmatrix} \in \mathbb{M}_{d_2 \times (d_1 + d_2)}.$$

But $Ad_{V_i^*} \circ \Phi \circ \Psi = \Phi_i \circ \Psi$, i = 1, 2. Since Ψ is arbitrary we conclude that Φ_i 's are k-EB. Conversely, assume that both Φ_1 and Φ_2 are k-EB, and let $\Psi \in C\mathcal{P}(k, d)$. Then $\Phi_i \circ \Psi$'s are EB-maps, and hence there exist $F_j^{(i)} \in \mathbb{M}_k^+$ and $R_j^{(i)} \in \mathbb{M}_{d_i}^+$ such that

$$\Phi_i \circ \Psi(X) = \sum_{j=1}^{r_i} \operatorname{tr} \left(X F_j^{(i)} \right) R_j^{(i)}$$

for all $X \in \mathbb{M}_k$ and i = 1, 2. Then

$$\Phi \circ \Psi(X) = \sum_{j=1}^{r_1} \operatorname{tr} \left(X F_j^{(1)} \right) \begin{bmatrix} R_j^{(1)} & \\ & \mathbf{0}_{d_2} \end{bmatrix} + \sum_{j=1}^{r_2} \operatorname{tr} \left(X F_j^{(2)} \right) \begin{bmatrix} \mathbf{0}_{d_1} & \\ & R_j^{(2)} \end{bmatrix},$$

for all $X \in \mathbb{M}_k$ so that $\Phi \circ \Psi$ is EB. Since Ψ is arbitrary Φ is *k*-EB.

Theorem 4.7. Given $1 \le k \le d$ the following conditions are equivalent:

(i) W_{λ} is a *k*-entanglement breaking CP-map.

- (ii) \mathcal{W}_{λ} is a *k*-PPT map.
- (iii) $\lambda \in [-1, 1/k].$

Proof. $(i) \Leftrightarrow (iii)$ We know that \mathcal{W}_{λ} is CP if and only if $\lambda \in [-1, 1]$. Also since \mathcal{W}_{λ} is equivariant, by Theorem 4.4, \mathcal{W}_{λ} is *k*-EB if and only if $\mathcal{W}_{\lambda} \circ Ad_{P}$ is EB, where $P = \begin{bmatrix} I_{k} & \mathbf{0}_{k \times (d-k)} \end{bmatrix} \in \mathbb{M}_{k \times d}$. But

$$\mathcal{W}_{\lambda} \circ Ad_{P}(X) = \begin{bmatrix} \mathsf{tr}(X)I_{k} - \lambda X^{\mathsf{T}} & 0\\ 0 & \mathsf{tr}(X)I_{d-k} \end{bmatrix}$$
(4.1)

for all $X \in \mathbb{M}_k$. Since the map $\mathbb{M}_k \ni X \mapsto \operatorname{tr}(X)I_{d-k} \in \mathbb{M}_{d-k}$ is always EB, by Lemma 4.6, $\mathcal{W}_{\lambda} \circ Ad_P$ is EB if and only if the Holevo-Werner map $\mathbb{M}_k \ni X \mapsto \operatorname{tr}(X)I_k - \lambda X^{\mathsf{T}} \in \mathbb{M}_k$ is EB, and which is true if and only if $\lambda \in [-1, \frac{1}{k}]$.

 $(ii) \Leftrightarrow (iii)$ Since \mathcal{W}_{λ} is equivariant, by Proposition 4.3, \mathcal{W}_{λ} is *k*-PPT if and only if $\mathcal{W}_{\lambda} \circ Ad_{P}$ is PPT, where $P = \begin{bmatrix} I_{k} & \mathbf{0}_{k \times (d-k)} \end{bmatrix} \in \mathbb{M}_{k \times d}$. But, equation (4.1) implies that $\mathcal{W}_{\lambda} \circ Ad_{P}$ is PPT if and only if the Holevo-Werner map $\mathbb{M}_{k} \ni X \mapsto \operatorname{tr}(X)I_{k} - \lambda X^{\mathsf{T}} \in \mathbb{M}_{k}$ is PPT if and only if $\lambda \in [-1, \frac{1}{k}]$.

Note 4.8. Equivalence of (i) and (iii) in the above theorem was observed in [ChCh20] also.

Example 4.9 ([ChCh20]). The Holevo-Werner map $\mathcal{W}_{\frac{1}{k}} : \mathbb{M}_d \to \mathbb{M}_d$ is a *k*-entanglement breaking CP-map for 1 < k < d. The composition $\mathcal{W}_{\frac{1}{k}} \circ \mathsf{T} = \mathsf{T} \circ \mathcal{W}_{\frac{1}{k}}$ is *k*-entanglement breaking but not (k+1)-positive, hence it is not (k+1)-entanglement breaking.

Corollary 4.10. Let $\Gamma : \mathbb{M}_d \to \mathbb{M}_d$ be a positive map and $k \ge 1$. If $\Gamma \in (\mathcal{EB}_k(d))^\circ$, then the following holds:

- (i) $-(I_d \otimes \Gamma(I_d)) \leq C_{\Gamma} \leq k(I_d \otimes \Gamma(I_d)).$
- (ii) $-(I_d \otimes \Gamma(I_d)) \leq C_{\Gamma \circ T} \leq k(I_d \otimes \Gamma(I_d)).$
- (iii) $-tr(\Gamma(I_d)) \leq tr(C_{\Gamma}C_{id_d}) \leq k tr(\Gamma(I_d)).$
- (iv) $-tr(\Gamma(I_d)) \leq tr(C_{\Gamma}C_{\tau}) \leq k tr(\Gamma(I_d)).$

Proof. By Theorem 2.5, $\Gamma \circ \Theta^*$ is CP for all $\Theta \in \mathcal{EB}_k(d)$. Since \mathcal{W}_{λ} and $\mathsf{T} \circ \mathcal{W}_{\lambda} = \mathcal{W}_{\lambda} \circ \mathsf{T}$ are self-adjoint *k*-EB maps for all $\lambda \in [-1, \frac{1}{k}]$, we have $\Gamma \circ \mathcal{W}_{\lambda}$ and $\Gamma \circ (\mathcal{W}_{\lambda} \circ \mathsf{T})$ are CP-maps. Therefore,

$$C_{\Gamma \circ \mathcal{W}_{\lambda}} = (I_d \otimes \Gamma(I_d)) - \lambda C_{\Gamma \circ \mathsf{T}} \ge 0 \quad \text{and} \quad C_{\Gamma \circ \mathcal{W}_{\lambda} \circ \mathsf{T}} = (I_d \otimes \Gamma(I_d)) - \lambda C_{\Gamma} \ge 0.$$

Taking $\lambda = -1, \frac{1}{k}$ in the above inequalities we get (i) and (ii). Now, from (i),

$$\begin{split} &-\left\langle\Omega_d, (I_d\otimes\Gamma(I_d))\Omega_d\right\rangle \leq \left\langle\Omega_d, C_{\Gamma}\Omega_d\right\rangle \leq k\left\langle\Omega_d, (I_d\otimes\Gamma(I_d))\Omega_d\right\rangle \\ &\Longrightarrow -\operatorname{tr}\left((I_d\otimes\Gamma(I_d))|\Omega_d\rangle\langle\Omega_d|\right) \leq \operatorname{tr}(C_{\Gamma}|\Omega_d\rangle\langle\Omega_d|) \leq k\operatorname{tr}\left((I_d\otimes\Gamma(I_d))|\Omega_d\rangle\langle\Omega_d|\right) \\ &\Longrightarrow -\operatorname{tr}(\Gamma(I_d)) \leq \operatorname{tr}(C_{\Gamma}C_{\mathsf{id}}) \leq k\operatorname{tr}(\Gamma(I_d)). \end{split}$$

Similarly, (iv) follows from (ii).

Theorem 4.11. Let $1 < k \leq d$ and $\Gamma : \mathbb{M}_d \to \mathbb{M}_d$ be a positive map. If $\lambda \in [-\frac{1}{k \|\Gamma\|}, \frac{1}{\|\Gamma\|}]$, then $\mathcal{W}_{\lambda,\Gamma} : \mathbb{M}_d \to \mathbb{M}_d$ given by

$$\mathcal{W}_{\lambda,\Gamma}(X) := tr(X)I_d + \lambda\Gamma(X) \tag{4.2}$$

is a *k*-entanglement breaking map.

Proof. Let $\lambda \in [-\frac{1}{k\|\Gamma\|}, \frac{1}{\|\Gamma\|}]$. Since $\alpha := -\lambda \|\Gamma\| \in [-1, \frac{1}{k}]$ Remark 3.2(ii) and Theorem 4.7 implies that $\Gamma \circ \mathsf{T} \circ W_{\alpha}$ is a *k*-EB map. Let $\Phi \in \mathcal{EB}(d)$ be the map given by $\Phi(X) := \mathsf{tr}(X)R$, where $R = \|\Gamma\| I_d - \Gamma(I_d) \in \mathbb{M}_d^+$. Now it follows that

$$W_{\lambda,\Gamma} = rac{1}{\|\Gamma\|} \left(\Phi + \Gamma \circ \mathsf{T} \circ W_{lpha}
ight)$$

is a k-EB map.

The following is a generalization of [CMW19, Theorem 3.3].

Theorem 4.12. Let $1 < k \leq d$ and $\Gamma : \mathbb{M}_d \to \mathbb{M}_d$ be a trace preserving positive map such that

$$\|tr(X)I_d - \Gamma(X)\|_{\infty} \le \frac{1}{k} \|X\|_{\infty}$$

for all $X \in \mathbb{M}_d$. Then Γ is a k-entanglement breaking map.

Proof. Let $\widetilde{\Gamma} := W_{-1,\Gamma} : \mathbb{M}_d \to \mathbb{M}_d$, which is a positive map. Note that $\|\widetilde{\Gamma}\| \leq \frac{1}{k}$, hence by Theorem 4.11, $W_{-1,\widetilde{\Gamma}}$ is k-EB. But $W_{-1,\widetilde{\Gamma}} = \Gamma$.

In the rest of this section, we consider a particular case of the maps defined by (4.2). Given $\lambda \in \mathbb{R}$ define $\Phi_{\lambda,d} : \mathbb{M}_d \to \mathbb{M}_d$ by

$$\Phi_{\lambda,d}(X) := \operatorname{tr}(X)I + \lambda(X + X^{\mathsf{T}}) \tag{4.3}$$

for all $X \in \mathbb{M}_d$. Clearly $\Phi_{\lambda,d} = \Phi_{\lambda,d} \circ \mathsf{T}$, hence $\Phi_{\lambda,d}$ is CP if and only if $\Phi_{\lambda,d}$ is PPT. Note that $\Phi_{\lambda,d}$ is unital if and only if $d + 2\lambda = 1$ if and only if $\Phi_{\lambda,d}$ is trace preserving.

Theorem 4.13. Let $\lambda \in \mathbb{R}$, $1 < k \leq d$. Then the following statements holds.

- (i) $\Phi_{\lambda,d}$ is positive if and only if $\lambda \in [-\frac{1}{2}, \infty)$.
- (ii) $\Phi_{\lambda,d}$ is EB if and only if $\Phi_{\lambda,d}$ is CP if and only if $\lambda \in [-\frac{1}{d+1}, 1]$.
- (iii) $\Phi_{\lambda,d}$ is k-EB implies $\Phi_{\lambda,k}$ is EB (and hence $\lambda \in [-\frac{1}{k+1}, 1]$). Conversely if $\lambda \in [\frac{-1}{2k}, 1]$, then $\Phi_{\lambda,d}$ is k-EB.

Proof. (i) Assume $\Phi_{\lambda,d}$ is positive. Then $\Phi_{\lambda,d}(E_{11}) \ge 0$, which implies that $\lambda \ge -\frac{1}{2}$. Conversely assume that $\lambda \ge -\frac{1}{2}$. Then given a unit vector $u \in \mathbb{C}^d$ we have

$$\left\langle x, \Phi_{\lambda,d}(|u\rangle\langle u|)x\right\rangle = \left\langle x, x\right\rangle + \lambda\left(\left|\langle x, u\rangle\right|^2 + \left|\langle x, \overline{u}\rangle\right|^2\right) \ge \|x\|^2 - \frac{1}{2}\left(\left|\langle x, u\rangle\right|^2 + \left|\langle x, \overline{u}\rangle\right|^2\right) \ge 0$$

for all $x \in \mathbb{C}^d$. Thus $\Phi_{\lambda,d}(|u\rangle\langle u|) \ge 0$. From spectral theorem it follows that $\Phi_{\lambda,d}$ is positive.

(*ii*) Clearly, $\Phi_{\lambda,d}$ is EB implies it is CP. Now assume that Φ is a CP-map. Note that

$$C_{\Phi_{\lambda,d}} = (I_d \otimes I_d) + \lambda (|\Omega_d\rangle \langle \Omega_d| + (\mathsf{id} \otimes \mathsf{T}) |\Omega_d\rangle \langle \Omega_d|)$$

Then $0 \leq \langle \Omega_d, C_{\Phi_{\lambda,d}} \Omega_d \rangle = d(1 + (d+1)\lambda)$, hence $1 + (d+1)\lambda \geq 0$, i.e., $\lambda \geq -\frac{1}{d+1}$. Also, since the principal submatrix $\begin{bmatrix} I + 2\lambda E_{11} & \lambda(E_{12} + E_{21}) \\ \lambda(E_{21} + E_{12}) & I + 2\lambda E_{22} \end{bmatrix}$ of $C_{\Phi_{\lambda,d}}$ is positive we have $\lambda^2 \leq 1$, hence $\lambda \leq 1$. Conversely, assume that $\lambda \in [-\frac{1}{d+1}, 1]$. Then, by Corollary A.7, the Choi matrix $C_{\Phi_{\lambda,d}}$ is separable. Therefore, $\Phi_{\lambda,d}$ is EB.

(*iii*) Assume that $\Phi_{\lambda,d}$ is k-EB. Let $P = \begin{bmatrix} I_k & 0_{k \times (d-k)} \end{bmatrix} \in \mathbb{M}_{k \times d}$. Then, by Theorem 3.5, $\Phi_{\lambda,d} \circ Ad_P$ is EB, and hence $Ad_{P^*} \circ \Phi_{\lambda,d} \circ Ad_P = \Phi_{\lambda,k}$ is also EB. So, from (ii), it follows that $\lambda \in [\frac{-1}{k+1}, 1]$. Conversely, assume that $\lambda \in [\frac{-1}{2k}, 1]$. If $\lambda \in [0, 1]$, then from (ii), we have $\Phi_{\lambda,d}$ is EB and hence k-EB. Now if $\lambda \in [\frac{-1}{2k}, 0]$, then consider the positive map $\Gamma(X) = X + X^{\mathsf{T}}$ on \mathbb{M}_d . As $\|\Gamma\| = 2$, by Theorem 4.11, $\Phi_{\lambda,d} = \mathcal{W}_{\lambda,\Gamma}$ is k-EB.

In the above theorem, we believe that $\Phi_{\lambda,d}$ is k-EB if and only if $\Phi_{\lambda,k}$ is EB if and only if $\lambda \in [-\frac{1}{k+1}, 1]$. But, we could not prove our claim.

Example 4.14. Given $k < \frac{d+1}{2}$, choose $\lambda \in [\frac{-1}{2k}, \frac{-1}{d+1})$. Then the map $\Phi_{\lambda,d} : \mathbb{M}_d \to \mathbb{M}_d$ is k-EB but not CP.

Let $\operatorname{tr}_1, \operatorname{tr}_2$ be the partial trace maps on $\mathbb{M}_{d_1} \otimes \mathbb{M}_{d_2}$, i.e., $\operatorname{tr}_1(A \otimes B) := \operatorname{tr}(A)B$ and $\operatorname{tr}_2(A \otimes B) := \operatorname{tr}(B)A$ for all $A \in \mathbb{M}_{d_1}$ and $B \in \mathbb{M}_{d_2}$. It is well-known that if $X \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_{d_2})^+$ is separable then $\operatorname{tr}_i(X), (\operatorname{id}_{d_1} \otimes \mathsf{T})X, (\mathsf{T} \otimes \operatorname{id}_{d_2})X \ge 0$. If $\operatorname{rank}(X) = \max\{d_1, d_2\}$, then $X \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_{d_2})^+$ is separable iff $(\operatorname{id} \otimes \mathsf{T})X \ge 0$. If $X \in (\mathbb{M}_{d_1} \otimes \mathbb{M}_{d_2})^+$ and $d_1d_2 \le 6$, then X is separable if and only if $(\mathsf{T} \otimes \operatorname{id})X \ge 0$. See [Per96, HHH96, HLV00] for details. We shall next obtain a necessary condition for separability. Note that, from the above known results, it holds trivially for $\lambda \ge 0$.

Theorem 4.15. Let $X \in (\mathbb{M}_d \otimes \mathbb{M}_d)^+$ be separable. Then

$$\lambda X + \lambda (\mathbf{T} \otimes i\mathbf{d}_d) X + (I_d \otimes t\mathbf{r}_1(X)) \ge 0$$

$$\lambda X + \lambda (i\mathbf{d}_d \otimes \mathbf{T}) X + (t\mathbf{r}_2(X) \otimes I_d) \ge 0$$

for all $\lambda \in [-\frac{1}{2}, \infty)$.

Proof. Suppose $X = \sum A_i \otimes B_i$ for some $A_i, B_i \in \mathbb{M}_d^+$. Since $\Gamma(X) = \operatorname{tr}(X)I + \lambda(X + X^{\mathsf{T}})$ is a positive map for all $\lambda \in [-\frac{1}{2}, \infty)$, we get

$$\begin{split} 0 &\leq (\Gamma \otimes \mathsf{id}_d) X \\ &= \sum \left(\mathsf{tr}(A_i) I_d + \lambda (A_i + A_i^{\mathsf{T}}) \right) \otimes B_i \\ &= (I_d \otimes \sum \mathsf{tr}(A_i) B_i) + \lambda (\sum A_i \otimes B_i) + \lambda \sum (A_i^{\mathsf{T}} \otimes B_i) \\ &= (I_d \otimes \mathsf{tr}_1(X)) + \lambda X + \lambda (\mathsf{T} \otimes \mathsf{id}_d) X. \end{split}$$

Similarly by considering $(id_d \otimes \Gamma)(X)$ we can get the second inequality.

Corollary 4.16. Let $\Phi : \mathbb{M}_d \to \mathbb{M}_d$ be an entanglement breaking CP-map. Then

$$(I_d \otimes \Phi(I_d)) + \lambda(C_\Phi + C_{\Phi \circ T}) \ge 0$$

$$(tr_2(C_\Phi) \otimes I_d) + \lambda(C_\Phi + C_{T \circ \Phi}) \ge 0$$

for all $\lambda \in [-\frac{1}{2}, \infty)$.

Proof. Follows from the above Theorem as C_{Φ} is separable.

5. Schmidt number reducing CP-maps

Suppose $\Phi \in \mathcal{CP}(d_1, d_2)$. Then from the definition of Schmidt number, it follows that

$$SN(\Phi \circ \Psi) < SN(\Psi)$$

for all $\Psi \in \mathcal{CP}(m, d_1)$, or equivalently,

$$SN(\operatorname{id}_m \otimes \Phi)(X)) \leq SN(X)$$

for all $X \in (\mathbb{M}_m \otimes \mathbb{M}_{d_1})^+$, where $m \geq 1$. We want to know as to when strict inequality occurs in the above inequalities? Note that if Ψ is an EB-map or SN(X) = 1, then the above inequalities become equality.

Lemma 5.1 ([CMW19, Lemma 2.1]). Given a k-positive map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ the following conditions are equivalent:

- (i) Φ is *n*-entanglement breaking for $n \leq \min\{k, d_2\}$.
- (ii) $SN((id_m \otimes \Phi)(X)) \leq \max\{m n + 1, 1\}$, for all $X \in (\mathbb{M}_m \otimes \mathbb{M}_{d_1})^+$ and for all $m \leq \min\{k, d_2\}$.

Theorem 5.2. Let $d_1, d_2 > 1$ and $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ be a CP-map. Then the following conditions are equivalent:

- (i) Φ is 2-entanglement breaking.
- (ii) $SN(\Phi \circ \Psi) < SN(\Psi)$ for all non-entanglement breaking CP-maps $\Psi : \mathbb{M}_m \to \mathbb{M}_{d_1}$ and $2 \le m \le d_2$.
- (iii) $SN(\Phi \circ \Psi) < SN(\Psi)$ for all non-entanglement breaking d_2 -PEB maps $\Psi : \mathbb{M}_m \to \mathbb{M}_{d_1}$ and $m \ge 2$.
- (iv) $SN(\Phi \circ \Psi) < SN(\Psi)$ for all non-entanglement breaking CP-maps $\Psi : \mathbb{M}_2 \to \mathbb{M}_{d_1}$.

Proof. (i) \Rightarrow (ii) Let $2 \leq m \leq d_2$ and $\Psi \in C\mathcal{P}(m, d_1)$ be a non-entanglement breaking map. To show that $SN(\Phi \circ \Psi) < SN(\Psi)$. We prove by induction on m. When m = 2, the required inequality follows from Theorem 3.5(ii). Now assume that the result is true for m-1, i.e., $SN(\Phi \circ \Psi) < SN(\Psi)$ for all non-entanglement breaking $\Psi \in C\mathcal{P}(l, d_1)$, where $2 \leq l \leq m-1 \leq d_2$. Let $\Psi \in C\mathcal{P}(m, d_1)$ be non-entanglement breaking and $r = SN(\Psi)$. If r = m, then by Lemma 5.1 we have

$$SN(\Phi \circ \Psi) = SN((\mathsf{id}_m \otimes \Phi)C_\Psi) \le m - 1 < SN(\Psi).$$

Now assume that 1 < r < m. Then, by Lemma 3.3, for all $1 \leq j \leq r$ there exist vectors $\psi_{j_i} \in \mathbb{C}^j \otimes \mathbb{C}^{d_1}$ such that

$$SN(\Phi \circ \Psi) = SN((\mathsf{id}_m \otimes \Phi)(C_{\Psi})) \le \max_{ij} \{SN((\mathsf{id}_j \otimes \Phi)(|\psi_{j_i}\rangle \langle \psi_{j_i}|))\} = \max_{ij} \{SN(\Phi \circ \Psi_{j_i})\},$$

where $\Psi_{j_i} \in \mathcal{CP}(j, d_1)$ is such that $C_{\Psi_{j_i}} = |\psi_{j_i}\rangle\langle\psi_{j_i}|$. If Ψ_{j_i} is EB, then $SN(\Phi \circ \Psi_{j_i}) = 1 < r = SN(\Psi)$. If Ψ_{j_i} is not an EB-map, from induction hypothesis, $SN(\Phi \circ \Psi_{j_i}) < SN(\Psi_{j_i}) \leq j \leq SN(\Psi)$. So, from the above equation, we get $SN(\Phi \circ \Psi) < SN(\Psi)$.

 $(ii) \Rightarrow (iii)$ Suppose $m \ge 2$ and $\Psi \in \mathcal{CP}(m, d_1)$ is non-entanglement breaking. If $m \le d_2$, then there is nothing to prove. So assume that $m > d_2$ and let $r := SN(\Psi) \le d_2$. By Lemma 3.3, for all $1 \le j \le r$ there exist $\Psi_{j_i} \in \mathcal{CP}(j, d_1)$ and isometries $V_{j_i} : \mathbb{C}^j \to \mathbb{C}^m$ such that

$$C_{\Psi} = \sum_{i,j} (V_{j_i} \otimes I_{d_1}) C_{\Psi_{j_i}} (V_{j_i} \otimes I_{d_1})^*.$$

Observe that not all Ψ'_{j_i} s are EB as Ψ is not an EB-map. Since $j \leq d_2$, by assumption

$$SN(\Phi \circ \Psi) \le \max_{i,j} SN(\Phi \circ \Psi_{j_i}) < \max_{i,j} SN(\Psi_{j_i}) \le j \le SN(\Psi).$$

 $(iii) \Rightarrow (iv)$ This follows as $SN(\Psi) = 2 \le d_2$ for all non-entanglement breaking $\Psi \in C\mathcal{P}(2, d_1)$. $(iv) \Rightarrow (i)$ To prove that Φ is 2-EB it is enough to show that $\Phi \circ \Psi$ is EB for every $\Psi \in C\mathcal{P}(2, d_1)$. If Ψ is EB, then we are done. Otherwise, by assumption $SN(\Phi \circ \Psi) < SN(\Psi) \le 2$. Thus, $\Phi \circ \Psi$ has Schmidt number one, and hence it is EB.

One would also ask, given a non-entanglement breaking $\Psi \in C\mathcal{P}(d_2, d_1)$, does there exist a map $\Phi \in C\mathcal{P}(d_1, d_2)$ such that $SN(\Phi \circ \Psi) = SN(\Psi)$? If yes, then such a map Φ cannot be 2-EB.

Remark 5.3. Suppose $\Psi \in C\mathcal{P}(d_2, d_1)$ is a non-entanglement breaking map and $d_1 \leq d_2$. Let $\Phi = Ad_V$, where $V \in \mathbb{M}_{d_1 \times d_2}$ is any co-isometry (i.e., $VV^* = \mathsf{id}_{d_1}$). Then

$$SN(\Psi) = SN(Ad_{V^*} \circ Ad_V \circ \Psi) \le SN(\Phi \circ \Psi) \le SN(\Psi).$$

Thus, there exists $\Phi \in C\mathcal{P}(d_1, d_2)$ such that $SN(\Phi \circ \Psi) = SN(\Psi)$. But if $d_1 > d_2$ this is not the case. For example, consider a 3-entanglement breaking CP-map $\Psi_0 : \mathbb{M}_d \to \mathbb{M}_2$ as given by Theorem 3.17, where $d \ge 4$. Since Ψ_0^* is not 3-EB there exists $\Psi_1 \in C\mathcal{P}(3, 2)$ such that $\Psi = \Psi_0^* \circ \Psi_1 \in C\mathcal{P}(3, d)$ is not EB. Now, if possible assume that there exists $\Phi \in C\mathcal{P}(d, 3)$ such that $SN(\Phi \circ \Psi) = SN(\Psi)$. Since Ψ_0 is 3-EB we have $\Psi_0 \circ \Phi^*$ and hence $\Psi_1^* \circ \Psi_0 \circ \Phi^*$ are EB-maps. Therefore, $1 = SN(\Phi \circ \Psi) = SN(\Psi)$, which is a contradiction. Thus there is no $\Phi \in C\mathcal{P}(d, 3)$ such that $SN(\Phi \circ \Psi) = SN(\Psi)$.

Note 5.4. Recall that the PPT-square conjecture states that the square of a PPT-map is EB; equivalently, the composition of two PPT-maps is EB. It is known ([CMW19, CYT19]) that if $\Phi_1, \Phi_2 \in \mathcal{PPT}(3,3)$, then $\Phi_1 \circ \Phi_2 \in \mathcal{EB}(3)$. We can prove this fact as follows also. If Φ_2 is EB, then there is nothing to prove. So assume Φ_2 is not EB. Since Φ_1 is PPT, by Theorem 3.16, Φ_1 is 2-EB. Hence, by Lemma 5.1 and Theorem 5.2, $SN(\Phi_1 \circ \Phi_2) < SN(\Phi_2) \leq 2$. Thus $SN(\Phi_1 \circ \Phi_2) = 1$.

Remark 5.5. Let $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ be a PPT-map and $m \geq 1$.

- (i) If $d_2 = 2$ or 3, then $\Phi \circ \Psi \in \mathcal{EB}(m, d_2)$ for all $\Psi \in \mathcal{PEB}_2(m, d_1)$.
- (ii) If $d_1 = 2$ or 3, then $\Psi \circ \Phi \in \mathcal{EB}(d_1, m)$ for all $\Psi \in \mathcal{PEB}_2(d_2, m)$.

To see (i), observe that by Proposition 3.16 Φ is 2-EB. So if $\Psi \in \mathcal{PEB}_2(m, d_1)$ is not EB, then $SN(\Phi \circ \Psi) < SN(\Psi) = 2$. Thus $\Phi \circ \Psi$ is EB. Similarly by considering Φ^* we can get (ii).

Note 5.6. Given any $\Gamma \in \mathcal{P}(d_1, d_2)$ and k > 1 the map $\mathrm{id}_k \otimes \Gamma$ is not 2-EB. For, suppose $A \in (\mathbb{M}_2 \otimes \mathbb{M}_k)^+$ is entangled and $B \in \mathbb{M}_{d_1}^+$. Then $(\mathrm{id}_2 \otimes \mathrm{id}_k \otimes \Gamma)(A \otimes B) = A \otimes \Gamma(B)$ is not separable in $\mathbb{M}_2 \otimes (\mathbb{M}_k \otimes \mathbb{M}_{d_1})$. For, if $A \otimes \Gamma(B)$ is separable and $\mathrm{tr}^{(d_1)} : \mathbb{M}_k \otimes \mathbb{M}_{d_1} \to \mathbb{M}_k$ is the partial trace map, then

$$\operatorname{tr}(\Gamma(B))A = (\operatorname{id}_2 \otimes \operatorname{tr}^{(d_1)})(A \otimes \Gamma(B)) \in \mathbb{M}_2^+ \otimes \mathbb{M}_k^+,$$

which is not possible as A is entangled.

Theorem 5.7. Let $2 \le m, n \le d$ and $\Phi, \Psi : \mathbb{M}_d \to \mathbb{M}_d$ be *n*-entanglement breaking CP-map and *m*-entanglement breaking CP-map, respectively. Then $\Phi \circ \Psi$ is (n+m-1)-entanglement breaking CP-map.

Proof. Case (1): Suppose $d \le n + m - 1$. Let $X \in (\mathbb{M}_d \otimes \mathbb{M}_d)^+$. Clearly, $Y = (\mathrm{id}_d \otimes \Psi)(X) \in (\mathbb{M}_d \otimes \mathbb{M}_d)^+$. Now since Ψ is *m*-EB, by Lemma 5.1, $SN(Y) \le \max\{d - m + 1, 1\} = n$. Further, since Φ is *n*-EB, by Theorem 3.5(viii),

$$SN((\mathsf{id}_d \otimes \Phi \circ \Psi)(X)) = SN((\mathsf{id}_d \otimes \Phi)(Y)) = 1.$$

Thus $(id_d \otimes \Phi \circ \Psi)(X)$ is separable for all $X \in (\mathbb{M}_d \otimes \mathbb{M}_d)^+$. Hence $\Phi \circ \Psi$ is EB, and in particular, (n + m - 1)-entanglement breaking.

<u>Case (2):</u> Suppose n + m - 1 < d. Let $\Gamma \in C\mathcal{P}(n + m - 1, d)$. We shall show that $\Phi \circ \Psi \circ \Gamma$ is EB so that, by Theorem 3.5(ii), $\Phi \circ \Psi$ is (n + m - 1)-entanglement breaking. Since Ψ is m-EB, by Lemma 5.1, $SN(\Psi \circ \Gamma) = SN((\mathsf{id}_{n+m-1} \otimes \Psi)C_{\Gamma}) \leq n$. Thus $\Psi \circ \Gamma \in \mathcal{PEB}_n(n + m - 1, d)$. Since Φ is *n*-EB, by Theorem 3.5(v), $\Phi \circ \Psi \circ \Gamma$ is EB. This completes the proof. \Box

The following improvised version of [CMW19, Theorem 2.1] is an immediate consequence of the above theorem.

Corollary 5.8. Let $2 \le n_i \le d$ and $k \in \mathbb{N}$. If $\Phi_i : \mathbb{M}_d \to \mathbb{M}_d$ are n_i -entanglement breaking *CP*-maps for $1 \le i \le k$, then the composition $\Phi_1 \circ \Phi_2 \circ \cdots \circ \Phi_k$ is (n - k + 1)-entanglement breaking *CP*-map, where $n = \sum_i n_i$.

Corollary 5.9. Let $2 \le k \le d$ and $\Phi : \mathbb{M}_d \to \mathbb{M}_d$ be a k-entanglement breaking CP-map. Then Φ^m is entanglement breaking, where $m = \min\{SN(\Phi), \lceil \frac{d-1}{k-1} \rceil\}$.

Proof. Let $k_1 = SN(\Phi)$. Since Φ is 2-EB, by Theorem 5.2, $SN(\Phi^{k_1}) = 1$. Meanwhile, if k_2 is such that $kk_2 - k_2 + 1 = d$, then Corollary 5.8 implies that $SN(\Phi^{k_2}) = 1$. Now we conclude that $SN(\Phi^m) = 1$, where $m = \min\{SN(\Phi), \lceil \frac{d-1}{k-1} \rceil\}$.

6. Majorization

Given $x = (x_1, x_2, \cdots, x_d)^{\mathsf{T}} \in \mathbb{R}^d$, we denote by $x^{\downarrow} = (x_1^{\downarrow}, x_2^{\downarrow}, \cdots, x_d^{\downarrow})^{\mathsf{T}} \in \mathbb{R}^d$ the vector with the same components, but sorted in descending order. Thus, $x_1^{\downarrow} \ge x_2^{\downarrow} \ge \cdots \ge x_d^{\downarrow}$. Given

 $x, y \in \mathbb{R}^d$, we say that x is *weakly majorized* by y (and write $x \prec_w y$), if

$$\sum_{i=1}^{k} x_i^{\downarrow} \le \sum_{i=1}^{k} y_i^{\downarrow} \quad \text{for all } 1 \le k \le d.$$

If $x \prec_w y$ and $\sum_{i=1}^d x_i = \sum_{i=1}^d y_i$, then we say that x is majorized by y, and write $x \prec y$. If the dimensions of x and y are different, we define $x \prec y$ and $x \prec_w y$ similarly by appending extra zeros to the smaller vector to equalize their dimensions. If A, B are two positive matrices, then we say A is (weakly) majorized by B if $\sigma(A)$ is (weakly) majorized by $\sigma(B)$, where $\sigma(X) \in \mathbb{R}^d$ denotes the vector of all eigenvalues of $X \in \mathbb{M}^+_d$ arranged in the decreasing order.

A matrix $D = [d_{ij}] \in \mathbb{M}_d$ is said to be *doubly sub-stochastic* if $d_{ij} \ge 0$ for all $1 \le i, j \le d$ and

$$\sum_{i=1}^{d} d_{ij} \leq 1 \quad \forall \ 1 \leq j \leq d \qquad \text{and} \qquad \sum_{j=1}^{d} d_{ij} \leq 1 \quad \forall \ 1 \leq i \leq d.$$

In the above, if $\sum_{i=1}^{d} d_{ij} = \sum_{j=1}^{d} d_{ij} = 1$, then *D* is called *doubly stochastic*. It is well-known that $x \prec y$ (resp. $x \prec_w y$) if and only if x = Dy for some doubly stochastic (resp. sub-stochastic) matrix $D \in \mathbb{M}_d$.

It is known ([NiKe01, Hir03]) that X is majorized by both $tr_1(X)$ and $tr_2(X)$ whenever $X \in (\mathbb{M}_d \otimes \mathbb{M}_d)^+$ is separable. Equivalently, if $\Phi : \mathbb{M}_d \to \mathbb{M}_d$ is an EB-map, then C_{Φ} is majorized by both $tr_1(C_{\Phi})$ and $tr_2(C_{\Phi})$. We prove an analogue of this result for k-EB maps, and the proof is almost same as that in [Hir03].

Given $X \in (\mathbb{M}_d \otimes \mathbb{M}_d)^+ = \mathcal{B}(\mathbb{C}^d \otimes \mathbb{C}^d)^+$ decompose $\mathbb{C}^d = \ker(\operatorname{tr}_2(X))^{\perp} \bigoplus \ker(\operatorname{tr}_2(X))$. Then with respect to the decomposition

$$\mathbb{C}^d \otimes \mathbb{C}^d = \left(\ker(\mathrm{tr}_2(X))^{\perp} \otimes \mathbb{C}^d \right) \bigoplus \left(\ker(\mathrm{tr}_2(X)) \otimes \mathbb{C}^d \right)$$

we can write

$$X = \begin{bmatrix} X_1 & 0 \\ 0 & 0 \end{bmatrix}, \tag{6.1}$$

where X_1 acts on ker $(tr_2(X))^{\perp} \otimes \mathbb{C}^d$. See [Hir03, Lemma 2] for details.

The following is a generalization of [Hir03, Theorem 1].

Theorem 6.1. Let $X \in (\mathbb{M}_d \otimes \mathbb{M}_d)^+$ and $k \ge 1$.

(i) If $(id_d \otimes W_{-\frac{1}{2},T})(X) \in (\mathbb{M}_d \otimes \mathbb{M}_d)^+$, then X is weakly majorized by $k \operatorname{tr}_2(X)$.

(ii) If $(W_{-\frac{1}{k},T} \otimes id_d)(X) \in (\mathbb{M}_d \otimes \mathbb{M}_d)^+$, then X is weakly majorized by $k \operatorname{tr}_1(X)$. (Here $W_{-\frac{1}{k},T}$ is given by (4.2).)

Proof. Because of (6.1), without loss of generality, we can assume $tr_2(X)$ is invertible and diagonal, say $tr_2(X) = \sum_{i=1}^d \alpha_i E_{ii}$, where $\alpha_i \in (0, \infty)$. Since

$$(\operatorname{tr}_2(X) \otimes I_d) - \frac{X}{k} = (\operatorname{id}_d \otimes W_{-\frac{1}{k},\mathsf{T}})(X) \ge 0,$$

by [Hir03, Lemma 1], there exists a $C \in \mathbb{M}_d \otimes \mathbb{M}_d$ with $\|C\| \leq 1$ such that

$$X^{\frac{1}{2}} = \sqrt{k} (\text{tr}_2(X)^{\frac{1}{2}} \otimes I_d) C.$$
(6.2)

Suppose $\sigma(X) = (\lambda_{11}, \lambda_{12}, \dots, \lambda_{1d}, \dots, \lambda_{d1}, \dots, \lambda_{dd})^{\mathsf{T}} \in \mathbb{R}^{d^2}$. Let $U \in \mathbb{M}_d \otimes \mathbb{M}_d$ be a unitary such that

$$U^* X^{\frac{1}{2}} U = diag\left(\sqrt{\lambda_{11}}, \sqrt{\lambda_{12}}, \cdots, \sqrt{\lambda_{1d}}, \cdots, \sqrt{\lambda_{d1}}, \cdots, \sqrt{\lambda_{dd}}\right).$$
(6.3)

Let R = CU. Then from (6.2), (6.3) we get

$$X = k(\operatorname{tr}_2(X)^{\frac{1}{2}} \otimes I_d) CC^*(\operatorname{tr}_2(X)^{\frac{1}{2}} \otimes I_d)$$
(6.4)

$$diag(\sqrt{\lambda_{11}}, \sqrt{\lambda_{12}}, \cdots, \sqrt{\lambda_{dd}}) = \sqrt{k} U^*(\operatorname{tr}_2(X)^{\frac{1}{2}} \otimes I_d)R$$
(6.5)

$$diag(\lambda_{11}, \lambda_{12}, \cdots, \lambda_{dd}) = kR^*(\mathsf{tr}_2(X) \otimes I_d)R.$$
(6.6)

From (6.4), for all $1 \le i, j \le d$, we have

$$\langle e_i \otimes e_j, X e_i \otimes e_j \rangle = k \langle e_i \otimes e_j, (\operatorname{tr}_2(X)^{\frac{1}{2}} \otimes I_d) C C^* (\operatorname{tr}_2(X)^{\frac{1}{2}} \otimes I_d) (e_i \otimes e_j) \rangle$$

$$= k \langle (\operatorname{tr}_2(X)^{\frac{1}{2}} \otimes I_d) (e_i \otimes e_j), C C^* (\operatorname{tr}_2(X)^{\frac{1}{2}} \otimes I_d) (e_i \otimes e_j) \rangle$$

$$= k \langle \sqrt{\alpha_i} e_i \otimes e_j, C C^* (\sqrt{\alpha_i} e_i \otimes e_j) \rangle$$

$$= k \alpha_i \langle e_i \otimes e_j, C C^* (e_i \otimes e_j) \rangle$$

$$= k \alpha_i \langle e_i \otimes e_j, C (\sum_{p,q} \langle e_p \otimes e_q, C^* (e_i \otimes e_j) \rangle e_p \otimes e_q) \rangle \rangle$$

$$= k \alpha_i \sum_{p,q=1}^d |\langle e_i \otimes e_j, C(e_p \otimes e_q) \rangle|^2$$

$$(6.7)$$

Similarly, from (6.6), we get

$$\lambda_{ij} = k \sum_{p,q=1}^{d} \alpha_p \left| \left\langle e_p \otimes e_q, R(e_i \otimes e_j) \right\rangle \right|^2$$
(6.8)

Now, from the definition of $tr_2(X)$ and (6.7), we have

$$\alpha_p = \langle e_p, \mathsf{tr}_2(X)e_p \rangle = \sum_{l=1}^d \langle e_p \otimes e_l, X(e_p \otimes e_l) \rangle = k\alpha_p \sum_{l,m,n=1}^d \left| \langle e_p \otimes e_l, C(e_m \otimes e_n) \rangle \right|^2,$$

i.e.,
$$\frac{1}{k} = \sum_{l,m,n=1}^d \left| \langle e_p \otimes e_l, C(e_m \otimes e_n) \rangle \right|^2$$

for all $1 \leq p \leq d$. Let $S = [S_{ij}] \in \mathbb{M}_d$ where

$$S_{ij} = \sum_{q=1}^{d} \left| \left\langle e_j \otimes e_q, R(e_1 \otimes e_i) \right\rangle \right|^2 \qquad \forall \ 1 \le i, j \le d$$

Clearly $S_{ij} \ge 0$. Further,

$$\sum_{i=1}^{d} S_{ij} = \sum_{i,q=1}^{d} \left| \left\langle e_j \otimes e_q, R(e_1 \otimes e_i) \right\rangle \right|^2 \le \sum_{q,m,n=1}^{d} \left| \left\langle e_j \otimes e_q, R(e_m \otimes e_n) \right\rangle \right|^2 = \frac{1}{k} < 1$$

for all $1 \leq j \leq d$. Also since $||R|| \leq 1$

$$\sum_{j=1}^{d} S_{ij} = \sum_{j,q=1}^{d} |\langle e_j \otimes e_q, R(e_1 \otimes e_i) \rangle|^2 = \langle e_1 \otimes e_i, R^*R(e_1 \otimes e_i) \rangle \le 1$$

for all $1 \le i \le d$. Thus *S* is a doubly sub-stochastic matrix. Now from (6.8),

$$\begin{bmatrix} \lambda_{11} \\ \lambda_{12} \\ \vdots \\ \lambda_{1d} \end{bmatrix} = S \begin{bmatrix} k\alpha_1 \\ k\alpha_2 \\ \vdots \\ k\alpha_d \end{bmatrix}$$

and hence $(\lambda_{11}, \cdots, \lambda_{1d})^{\mathsf{T}} \prec_w (k\alpha_1, k\alpha_2, \cdots, k\alpha_d)^{\mathsf{T}}$ i.e.,

$$\sum_{j=1}^{n} \lambda_{1j} \le \sum_{i=1}^{n} k\alpha_i \qquad \forall \ 1 \le n \le d.$$
(6.9)

Note that

$$\sum_{i,j=1}^{d} \lambda_{ij} = \operatorname{tr}(X) = \operatorname{tr}(\operatorname{tr}_{2}(X)) = \sum_{i=1}^{d} \alpha_{i} \le \sum_{i=1}^{d} k \alpha_{i}.$$
(6.10)

Take $\alpha_i = 0$ for all $d < i \leq d^2$ and $(\beta_1, \beta_2, \dots, \beta_{d^2})^{\mathsf{T}} = (\lambda_{11}, \lambda_{12}, \dots, \lambda_{1d}, \dots, \lambda_{d1}, \dots, \lambda_{dd})^{\mathsf{T}}$. Then for all for all $d < n \leq d^2$ we have

$$\sum_{i=1}^{n} \beta_i \le \sum_{i,j=1}^{d} \lambda_{ij} = \sum_{i=1}^{d} \alpha_i = \sum_{i=1}^{n} \alpha_i.$$
(6.11)

Now from (6.9), (6.10) and (6.11) we conclude that $\sigma(X) \prec_w \sigma(k \operatorname{tr}_2(X))$. Similarly we can prove that $\sigma(X) \prec_w \sigma(k \operatorname{tr}_1(X))$.

Theorem 6.2. Let $\Phi : \mathbb{M}_d \to \mathbb{M}_d$ be a *k*-EB map, where $1 \le k \le d$. Then C_{Φ} is weakly majorized by both $(d - k + 1) \operatorname{tr}_1(C_{\Phi})$ and $(d - k + 1) \operatorname{tr}_2(C_{\Phi})$.

Proof. Let r = d - k + 1. By Theorem 4.11, $W_{-\frac{1}{r},\mathsf{T}}$ on \mathbb{M}_d is (d - k + 1)-entanglement breaking CP-map, and hence, by Lemma 5.1, $W_{-\frac{1}{r},\mathsf{T}}$ is k-PEB map. From Theorem 3.5(xi) it follows that k-PEB maps are in the dual of k- EB maps. Hence, by Theorem 2.5, $W_{-\frac{1}{r},\mathsf{T}} \circ \Phi = (\Phi^* \circ W_{-\frac{1}{r},\mathsf{T}})^*$ is CP. Thus we have $(I_d \otimes W_{-\frac{1}{r},\mathsf{T}})(C_{\Phi}) \ge 0$. Hence by the above theorem we get $\frac{1}{d-k+1}C_{\Phi}$ is weakly majorized by $\operatorname{tr}_2(C_{\Phi})$. Similarly C_{Φ} is is weakly majorized by $(d-k+1)\operatorname{tr}_2(C_{\Phi})$ also.

7. Discussion

It is known ([YLT16]) that if $d_1d_2 \leq 6$, then every 2-positive map $\Gamma : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ is decomposable. But when $4 \leq \max\{d_1, d_2\} \leq 9$ ([CYT17, page 18]) or $d_1, d_2 \geq 10$ ([HLLM18, BhOs20]), then there exists 2-positive map which is not decomposable.

Open Problem 1. Is every 2-EB map $\Phi : \mathbb{M}_{d_1} \to \mathbb{M}_{d_2}$ decomposable?

Theorem 7.1. Suppose the Problem-1 has an affirmative answer. If $\Phi : \mathbb{M}_d \to \mathbb{M}_d$ is a 2-entanglement breaking PPT-map, then Φ^2 is entanglement breaking.

Proof. Suppose Φ^2 is not an EB-map. Then there exists a $\Gamma \in \mathcal{P}(d)$ such that $\Gamma \circ \Phi^2$ is not CP. Note that $\Gamma \circ \Phi$ is 2-EB and hence decomposable. So there exist $\Phi_1, \Phi_2 \in \mathcal{CP}(d)$ such that $\Gamma \circ \Phi = \Phi_1 + \Phi_2 \circ \mathsf{T}$. Hence

$$\Gamma \circ \Phi^2 = (\Phi_1 + \Phi_2 \circ \mathsf{T}) \circ \Phi = \Phi_1 \circ \Phi + \Phi_2 \circ (\mathsf{T} \circ \Phi)$$

is CP, which is a contradiction.

Example 7.2. If $\lambda \in [-1, \frac{1}{2}]$, then the Holevo-Werner map \mathcal{W}_{λ} (and hence $\mathsf{T} \circ \mathcal{W}_{\lambda}$) is 2-EB. Note that \mathcal{W}_{λ} (resp. $\mathsf{T} \circ \mathcal{W}_{\lambda}$) is a CP-map (resp. co-CP), and hence decomposable.

Example 7.3. Let $\lambda \in [\frac{-1}{4}, 1]$. Then the map $\Phi_{\lambda,d} : \mathbb{M}_d \to \mathbb{M}_d$ given by (4.3) is 2-EB. If $\lambda \in [\frac{-1}{4}, \frac{1}{2}]$, then $\Phi_{\lambda,d} = \frac{1}{2} (\mathcal{W}_{-2\lambda} + \mathcal{W}_{-2\lambda} \circ \mathsf{T})$, hence we conclude that $\Phi_{\lambda,d}$ is decomposable. Now if $\lambda \in [\frac{1}{2}, 1]$, then $\Phi_{\lambda,d}$ is EB, and in particular decomposable.

Acknowledgments

We are grateful to Alexander Müller-Hermes for his valuable comments, providing a proof of a part of the Theorem 3.13, and pointing out the reference [ChCh20]. We thank B. V. Rajarama Bhat for his comments. RD is supported by UGC (University of Grant Commission, India) with ref No 21/06/2015(i)/EU-V. NM thanks NBHM (National Board for Higher Mathematics, India) for financial support with ref No 0204/52/2019/R&D-II/321. KS is partially supported by the IoE-CoE Project (No. SB20210797MAMHRD008573) from MHRD (Ministry of Human Resource Development, India) and partially by the MATRIX grant (File no. MTR/2020/000584) from SERB (Science and Engineering Research Board, India).

Appendix A.

Let $a, b, c \in \mathbb{C}$ and define $\Phi : \mathbb{M}_d \to \mathbb{M}_d$ by $\Phi(X) = a \operatorname{tr}(X)I + bX + cX^{\mathsf{T}}$. We want to know for what values of $a, b, c \in \mathbb{C}$ the map Φ is EB; equivalently when the corresponding Choi matrix

$$a(I_d \otimes I_d) + b|\Omega_d\rangle \langle \Omega_d| + c\Delta_d \in \mathbb{M}_d \otimes \mathbb{M}_d$$

is separable? Here $\Delta_d = (\mathrm{id}_d \otimes \mathsf{T}) |\Omega_d\rangle \langle \Omega_d| = \sum_{i,j=1}^d E_{ij} \otimes E_{ji}$. We answer this through similar lines as Stormer ([Sto13]) proved the entanglement property of the map $\mathrm{tr}(\cdot)\Gamma(I) + \Gamma(\cdot)$, where $\Gamma \in \mathcal{P}(d)$. Through out we let G denotes the compact group

 $G = \{Ad_{U \otimes U} : U \in \mathbb{M}_d \text{ real orthogonal}\} \subseteq \mathbb{M}_d \otimes \mathbb{M}_d = \mathbb{M}_d(\mathbb{M}_d),$

where d > 1. Let μ denotes the normalized Haar measure on G and let

 $Fix(G) = \{A \in \mathbb{M}_d \otimes \mathbb{M}_d : Ad_{U \otimes U}(A) = A \text{ for all real orthogonal matrices } U \in \mathbb{M}_d\}.$

Clearly $I_d \otimes I_d \in Fix(G)$. Observe that, for every $x = (x_1, \cdots, x_d)^{\mathsf{T}}, y = (y_1, \cdots, y_d)^{\mathsf{T}} \in \mathbb{C}^d$

$$\Delta_d(x \otimes y) = \sum_{ij} x_j e_i \otimes y_i e_j = (\sum_i y_i e_i) \otimes (\sum_j x_j e_j) = y \otimes x$$

Consequently, $Ad_{U\otimes U}(\Delta_d)(x\otimes y) = y\otimes x = \Delta_d(x\otimes y)$ for every real orthogonal matrices $U \in \mathbb{M}_d$. In other words, $\Delta_d \in Fix(G)$. Note that

$$\begin{aligned} Ad_{U\otimes U}(|\Omega_d\rangle\langle\Omega_d|) &= Ad_{U\otimes U}\big((\mathsf{id}_d\otimes\mathsf{T})\Delta_d\big) \\ &= (\mathsf{id}_d\otimes\mathsf{T})\big(Ad_{U\otimes U}(\Delta_d)\big) \\ &= (\mathsf{id}_d\otimes\mathsf{T})(\Delta_d) \\ &= |\Omega_d\rangle\langle\Omega_d|, \end{aligned}$$

for every real orthogonal matrices $U \in \mathbb{M}_d$. Thus $|\Omega_d\rangle \langle \Omega_d| \in Fix(G)$.

Lemma A.1. Let d = 2. Then $Fix(G) = \{\alpha(I_2 \otimes I_2) + \beta | \Omega_2 \rangle \langle \Omega_2 | + \gamma \Delta_2 : \alpha, \beta, \gamma \in \mathbb{C} \}.$

Proof. Let $A = [a_{ij}] \in Fix(G) \subseteq \mathbb{M}_2(\mathbb{M}_2)$. Consider $U = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \in \mathbb{M}_2$. Then $Ad_{U \otimes U}(A) = A$ implies that

$$A = \begin{bmatrix} a_{11} & 0 & 0 & a_{14} \\ 0 & a_{22} & a_{23} & 0 \\ 0 & a_{32} & a_{33} & 0 \\ a_{41} & 0 & 0 & a_{44} \end{bmatrix}.$$

Similarly considering $U = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ we conclude that

$$A = \begin{bmatrix} a_{11} & 0 & 0 & a_{14} \\ 0 & a_{22} & a_{23} & 0 \\ 0 & a_{23} & a_{22} & 0 \\ a_{14} & 0 & 0 & a_{11} \end{bmatrix} = \begin{bmatrix} a_0 \\ a_{22} \\ a_{22} \\ a_{22} \end{bmatrix} + a_{14} \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} + a_{23} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = (a_0 - a_{22})(E_{11} \otimes E_{11} + E_{22} \otimes E_{22}) + a_{22}(I_2 \otimes I_2) + a_{14}(|\Omega_2\rangle\langle\Omega_2|) + a_{23}\Delta_2,$$

where $a_0 = a_{11} - (a_{23} + a_{14})$. Since $A, I_2 \otimes I_2, \Delta_2, |\Omega_2\rangle \langle \Omega_2|$ are in Fix(G), from above we get $B = (a_0 - a_{22})(E_{11} \otimes E_{11} + E_{22} \otimes E_{22}) \in Fix(G)$. Take $U = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$. Then $Ad_{U \otimes U}(B) = B$ implies that $a_0 - a_{22} = 0$. Therefore, $A = a_{22}(I_2 \otimes I_2) + a_{14}(|\Omega_2\rangle \langle \Omega_2|) + a_{23}\Delta_2$.

Lemma A.2. Let d > 2. Given $m \neq n$ let

$$P_{\{m,n\}} = \sum_{i,j \in \{m,n\}} |e_i \otimes e_j\rangle \langle e_i \otimes e_j|$$

be the projection of $\mathbb{C}^d \otimes \mathbb{C}^d$ onto $\text{span}\{e_i \otimes e_j : i, j \in \{m, n\}\} \cong \mathbb{C}^2 \otimes \mathbb{C}^2$. Then for every $A \in Fix(G)$ there exist unique $\alpha_{mn}, \beta_{mn}, \gamma_{mn} \in \mathbb{C}$ such that

$$P_{\{m,n\}}AP_{\{m,n\}}^* = \alpha_{mn}(I_2 \otimes I_2) + \beta_{mn}|\Omega_2\rangle\langle\Omega_2| + \gamma_{mn}\Delta_2.$$

Proof. Let $U \in \mathbb{M}_2$ be a real orthogonal matrix. Then U induce a linear map on $\mathcal{W} = \operatorname{span}\{e_m, e_n\}$. Decompose $\mathbb{C}^d = \mathcal{W} \oplus \mathcal{W}^{\perp}$ and write $A = \begin{bmatrix} A_{mn,11} & A_{mn,12} \\ A_{mn,21} & A_{mn,22} \end{bmatrix} \in \mathbb{M}_d(\mathbb{M}_d)$, where $A_{mn,11} \in \mathbb{M}_2 \otimes \mathbb{M}_2$. Consider the real orthogonal matrix $\widetilde{U} = \begin{bmatrix} U & 0 \\ 0 & I_{\mathcal{W}^{\perp}} \end{bmatrix} \in \mathbb{M}_d$. Then

$$Ad_{\widetilde{U}\otimes\widetilde{U}}(A) = A \Longrightarrow Ad_U(A_{mn,11}) = A_{mn,11}.$$

Since $U \in \mathbb{M}_2$ is arbitrary, from Lemma A.1, we get

$$A_{mn,11} = \alpha_{mn} (I_2 \otimes I_2) + \beta_{mn} |\Omega_2\rangle \langle \Omega_2| + \gamma_{mn} \Delta_2$$

where $\alpha_{mn}, \beta_{mn}, \gamma_{mn} \in \mathbb{C}$. But, $A_{mn,11} = P_{\{m,n\}}AP^*_{\{m,n\}}$, where

$$P_{\{m,n\}} = \sum_{i,j\in\{m,n\}} |e_i \otimes e_j\rangle \langle e_i \otimes e_j|,$$

the projection of $\mathbb{C}^d \otimes \mathbb{C}^d$ onto span $\{e_i \otimes e_j : i, j \in \{m, n\}\}$. Uniqueness of $\alpha_{mn}, \beta_{mn}, \gamma_{mn}$ follows from the linear independence of $I_2 \otimes I_2, |\Omega_2\rangle\langle\Omega_2|, \Delta_2$.

Proposition A.3. Let d > 2. Then $Fix(G) = \{ \alpha(I_d \otimes I_d) + \beta | \Omega_d \rangle \langle \Omega_d | + \gamma \Delta_d : \alpha, \beta, \gamma \in \mathbb{C} \}.$

Proof. Let $A \in Fix(G)$. Assume that $A = \sum_{m,n,p,q} a_{(m,p),(n,q)} E_{mp \otimes E_{nq}}$. Thus,

$$\langle e_m \otimes e_n, A(e_p \otimes e_q) \rangle = a_{(m,p),(n,q)} \langle e_m \otimes e_n, (\mathsf{id}_d \otimes \mathsf{T}) A(e_p \otimes e_q) \rangle = \langle e_m \otimes e_q, A(e_p \otimes e_n) \rangle,$$

for all $1 \le m, n, p, q \le d$. Now let $m, n, p, q \in \{1, 2, \dots, d\}$. Case (1): $m \ne n$ or $p \ne q$.

<u>Subcase (i)</u>: $\{m, n\} \neq \{p, q\}$. Suppose m, n, p, q are distinct. Given $r \in \{m, n, p, q\}$ define the real orthogonal matrix

$$U_r = \left(\sum_{\substack{i=1,\cdots,d\\i\neq r}} |e_i\rangle\langle e_i|\right) - |e_r\rangle\langle e_r| \in \mathbb{M}_d.$$
(A.1)

Note that $U(e_r) = -e_r$ and $U(e_i) = e_i$ for all $i \neq r$. Since $Ad_{U_r \otimes U_r}(A) = A$, we get

$$\left\langle e_m \otimes e_n, A(e_p \otimes e_q) \right\rangle = \left\langle U_r e_m \otimes U_r e_n, A(U_r e_p \otimes U_r e_q) \right\rangle = -\left\langle e_m \otimes e_n, A(e_p \otimes e_q) \right\rangle.$$

Hence $a_{(m,p),(n,q)} = 0$. Now suppose m = p or m = q or n = p or n = q. With out loss of generality assume that m = p. Since $\{m, n\} \neq \{p, q\}$ we have $n \neq q$. Fix $r \in \{n, q\}$ and define the real orthogonal matrix $U_r \in \mathbb{M}_d$ by (A.1). Then also

$$\left\langle e_m \otimes e_n, A(e_p \otimes e_q) \right\rangle = \left\langle U_r e_m \otimes U_r e_n, A(U_r e_p \otimes U_r e_q) \right\rangle = -\left\langle e_m \otimes e_n, A(e_p \otimes e_q) \right\rangle$$

so that $a_{(m,p),(n,q)} = 0$.

<u>Subcase (ii)</u>: $\{m, n\} = \{p, q\}$. From Lemma A.2, there exists scalars $\alpha_{mn}, \beta_{mn}, \gamma_{mn} \in \mathbb{C}$ such that

$$P_{\{m,n\}}AP_{\{m,n\}}^* = \alpha_{mn}(I_2 \otimes I_2) + \beta_{mn}|\Omega_2\rangle\langle\Omega_2| + \gamma_{mn}\Delta_2.$$

Therefore,

$$\begin{aligned} a_{(m,p),(n,q)} &= \left\langle e_m \otimes e_n, A(e_p \otimes e_q) \right\rangle \\ &= \left\langle P_{\{m,n\}}^* e_m \otimes e_n, AP_{\{m,n\}}^*(e_p \otimes e_q) \right\rangle \\ &= \left\langle e_m \otimes e_n, P_{\{m,n\}} AP_{\{m,n\}}^*(e_p \otimes e_q) \right\rangle \\ &= \left\langle e_m \otimes e_n, \left(\alpha_{mn}(I_2 \otimes I_2) + \beta_{mn} |\Omega_2\rangle \langle \Omega_2 | + \gamma_{mn} \Delta_2 \right) (e_p \otimes e_q) \right\rangle \\ &= \begin{cases} \alpha_{mn} & \text{if } p = m \neq n = q \\ \gamma_{mn} & \text{if } q = m \neq n = p. \end{cases} \end{aligned}$$

Case (2): m = n and p = q. If m = n = p = q, then choose any $r \neq m$. Then, by Lemma A.2,

$$\begin{aligned} a_{(m,p),(n,q)} &= \left\langle e_m \otimes e_m, A(e_m \otimes e_m) \right\rangle \\ &= \left\langle e_m \otimes e_m, P_{rm} A P_{rm}^*(e_m \otimes e_m) \right\rangle \\ &= \alpha_{rm} \left\langle e_m \otimes e_m, e_m \otimes e_m \right\rangle + \beta_{rm} \left\langle e_m \otimes e_m, |\Omega_2\rangle \langle \Omega_2 | (e_m \otimes e_m) \right\rangle \\ &+ \gamma_{rm} \left\langle e_m \otimes e_m, \Delta_2(e_m \otimes e_m) \right\rangle \\ &= \alpha_{rm} + \beta_{rm} + \gamma_{rm}. \end{aligned}$$

Thus $a_{(m,m),(m,m)} = \alpha_{rm} + \beta_{rm} + \gamma_{rm}$ for any $r \neq m$. Now suppose $m = n \neq p = q$. Note that $n \neq p$ and $m \neq q$ but $\{m, q\} = \{n, p\}$. Hence

$$\begin{split} a_{(m,p),(n,q)} &= \left\langle e_m \otimes e_n, A(e_p \otimes e_q) \right\rangle \\ &= \left\langle e_m \otimes e_q, (\mathrm{id}_d \otimes \mathsf{T})(A)(e_p \otimes e_n) \right\rangle \\ &= \left\langle e_m \otimes e_q, P_{mq}(\mathrm{id} \otimes \mathsf{T})(A) P_{mq}^*(e_p \otimes e_n) \right\rangle \\ &= \left\langle e_m \otimes e_q, (\mathrm{id} \otimes \mathsf{T}) P_{mq}(A) P_{mq}^*(e_p \otimes e_n) \right\rangle \\ &= \alpha_{mq} \left\langle e_m \otimes e_q, e_p \otimes e_n \right\rangle + \beta_{mq} \left\langle e_m \otimes e_q, \Delta_2(e_p \otimes e_n) \right\rangle \\ &+ \gamma_{mq} \{e_m \otimes e_q, |\Omega_2\rangle \langle \Omega_2 | (e_p \otimes e_n) \} \\ &= \beta_{mq}. \end{split}$$

Thus

$$a_{(m,p),(n,q)} = \begin{cases} \alpha_{rm} + \beta_{rm} + \gamma_{rm} & \text{if } m = n = p = q \text{ (where } r \neq m) \\ \alpha_{mn} & \text{if } p = m \neq n = q \\ \beta_{mq} & \text{if } m = n \neq p = q \\ \gamma_{mn} & \text{if } q = m \neq n = p \\ 0 & \text{otherwise.} \end{cases}$$
(A.2)

Now given any $m \neq n$ and $p \neq q$ consider a real orthogonal matrix $U \in \mathbb{M}_d$ such that $U(e_m) = e_p, U(e_n) = e_q$ and $U^2 = I_d$. Clearly, then $U(e_p) = e_m$ and $U(e_q) = e_n$. Further,

$$Ad_{U\otimes U}(A) = A \Longrightarrow P_{\{m,n\}}Ad_{U\otimes U}(A)P_{\{m,n\}}^* = P_{\{m,n\}}AP_{\{m,n\}}^*$$

$$\Rightarrow \alpha_{pq} = \alpha_{mn}, \beta_{pq} = \beta_{mn}, \gamma_{pq} = \gamma_{mn}.$$

Hence, there exists $\alpha, \beta, \gamma \in \mathbb{C}$ such that (A.2) becomes

=

$$a_{(m,p),(n,q)} = \begin{cases} \alpha + \beta + \gamma & \text{if } m = n = p = q \text{ (where } r \neq m) \\ \alpha & \text{if } p = m \neq n = q \\ \beta & \text{if } m = n \neq p = q \\ \gamma & \text{if } q = m \neq n = p \\ 0 & \text{otherwise.} \end{cases}$$
(A.3)

Thus $A = \alpha(I_d \otimes I_d) + \beta(|\Omega_d\rangle \langle \Omega_d|) + \gamma \Delta_d.$

Definition A.4. Define $P: \mathbb{M}_d \otimes \mathbb{M}_d \to \mathbb{M}_d \otimes \mathbb{M}_d$ by

$$P(A) = \int_G Ad_{U\otimes U}(A)d\mu(U).$$

Observation A.5. We make the following observations:

(i) P is a unital positive projection with

$$\operatorname{range}(P) = Fix(G) = \{ \alpha(I_d \otimes I_d) + \beta |\Omega_d\rangle \langle \Omega_d | + \gamma \Delta_d : \alpha, \beta, \gamma \in \mathbb{C} \},$$
(A.4)

Further, tr(P(A)) = tr(A) for all $A \in \mathbb{M}_d \otimes \mathbb{M}_d$.

(ii) Let $\mathcal{C} \subseteq \mathcal{P}(d)$ be a mapping cone and let $\Phi \in \mathcal{C}$. Let $\Psi : \mathbb{M}_d \to \mathbb{M}_d$ be the linear map such that $C_{\Psi} = P(C_{\Phi})$. Then for any $\Gamma \in \mathcal{C}^\circ$ we have

$$\begin{aligned} \operatorname{tr}(C_{\Psi}C_{\Gamma}) &= \operatorname{tr}(P(C_{\Phi})C_{\Gamma}) \\ &= \operatorname{tr}\left(\left(\int_{G} Ad_{U\otimes U}(C_{\Phi})d\mu(U)\right)C_{\Gamma}\right) \\ &= \int_{G} \operatorname{tr}\left(C_{Ad_{U}\circ\Phi\circ Ad_{U^{*}}}C_{\Gamma}\right)d\mu(U) \\ &\geq 0, \end{aligned}$$

where the last inequality follows because $Ad_U \circ \Phi \circ Ad_{U^*} \in \mathbb{C}$. Thus $\Psi \in \mathbb{C}^{\circ \circ} = \mathbb{C}$. Since P is a positive projection we conclude that

$$P(\{C_{\Phi}: \Phi \in \mathfrak{C}\}) = \{C_{\Psi}: \Psi \in \mathfrak{C}\} \bigcap \mathsf{range}(P).$$

In particular, considering $\mathcal{C} = \mathcal{EB}(d)$, we have

$$P((\mathbb{M}_d \otimes \mathbb{M}_d)_{sep}^+) = (\mathbb{M}_d \otimes \mathbb{M}_d)_{sep}^+ \bigcap \mathsf{range}(P), \tag{A.5}$$

where $(\mathbb{M}_d \otimes \mathbb{M}_d)_{sep}^+$ denotes the set of separable positive matrices.

(iii) Let $X, Y \in \mathbb{M}_d^+$. Since $Ad_{U \otimes U}(\Delta_d) = \Delta_d$ for every real orthogonal $U \in \mathbb{M}_d$ we have

$$\operatorname{tr}\left(P(X\otimes Y)\Delta_d\right) = \int_G \operatorname{tr}\left(Ad_{U\otimes U}(X\otimes Y)\Delta_d\right)d\mu(U)$$

$$= \int_{G} \operatorname{tr} \left((X \otimes Y) A d_{U^* \otimes U^*} (\Delta_d) \right) d\mu(U)$$

$$= \int_{G} \operatorname{tr} \left((X \otimes Y) \Delta_d \right) d\mu(U)$$

$$= \operatorname{tr} \left((X \otimes Y) \Delta_d \right).$$

Similarly, tr $(P(X \otimes Y)|\Omega_d\rangle\langle\Omega_d|) = tr ((X \otimes Y)|\Omega_d\rangle\langle\Omega_d|)$. Thus

$$\operatorname{tr}\left(P(Z)\Delta_d\right) = \operatorname{tr}(Z\Delta_d) \tag{A.6}$$

$$\operatorname{tr}\left(P(Z)|\Omega_d\rangle\langle\Omega_d|\right) = \operatorname{tr}(Z|\Omega_d\rangle\langle\Omega_d|) \tag{A.7}$$

for all $Z \in (\mathbb{M}_d \otimes \mathbb{M}_d)_{sep}^+$.

(iv) From (A.4) and (A.5), we have $a(I_d \otimes I_d) + b|\Omega_d\rangle\langle\Omega_d| + c\Delta_d$ is separable if and only if

$$a(I_d \otimes I_d) + b|\Omega_d\rangle \langle \Omega_d| + c\Delta_d = P(Z)$$

for some $Z \in (\mathbb{M}_d \otimes \mathbb{M}_d)_{sep}^+$. Now since $\Delta_d^2 = I_d \otimes I_d, \Delta_d |\Omega_d\rangle \langle \Omega_d| = |\Omega_d\rangle \langle \Omega_d| = |\Omega_d\rangle \langle \Omega_d|^2 = d|\Omega_d\rangle \langle \Omega_d|$, from (A.6) and (A.7), we get

$$ad + bd + cd^2 = \operatorname{tr}(Z\Delta_d) \tag{A.8}$$

$$ad + bd^2 + cd = \operatorname{tr}(Z|\Omega_d\rangle\langle\Omega_d|).$$
 (A.9)

Further, tr(P(Z)) = tr(Z) implies that

$$ad^2 + bd + cd = \operatorname{tr}(Z). \tag{A.10}$$

Solving (A.8), (A.9) and (A.10) for a, b, c we get

$$a = \frac{1}{d^3 + d^2 - 2d} \left((d+1)\operatorname{tr}(Z) - \operatorname{tr}(Z\Delta_d) - \operatorname{tr}(Z|\Omega_d\rangle\langle\Omega_d|) \right)$$
(A.11)

$$b = \frac{1}{d^3 + d^2 - 2d} \left((d+1) \operatorname{tr}(Z|\Omega_d) \langle \Omega_d| \right) - \operatorname{tr}(Z\Delta_d) - \operatorname{tr}(Z)$$
(A.12)

$$c = \frac{1}{d^3 + d^2 - 2d} \left((d+1)\operatorname{tr}(Z\Delta_d) - \operatorname{tr}(Z|\Omega_d\rangle\langle\Omega_d|) - \operatorname{tr}(Z). \right)$$
(A.13)

Theorem A.6. Let $x, y \in \mathbb{C}^d$. Then

$$P(|x\rangle\langle x|\otimes |y\rangle\langle y|) = a(I_d\otimes I_d) + b(\Delta_d + |\Omega_d\rangle\langle\Omega_d|)$$

is separable, where $a, b \in \mathbb{R}$ are given by

$$a = \frac{1}{d^3 + d^2 - 2d} \left((d+1) \|x\|^2 \|y\|^2 - |\langle x, y \rangle|^2 - |\langle x, \overline{y} \rangle|^2 \right)$$

$$b = \frac{1}{d^3 + d^2 - 2d} \left((d+1) |\langle x, \overline{y} \rangle|^2 - |\langle x, y \rangle|^2 - \|x\|^2 \|y\|^2 \right)$$

$$c = \frac{1}{d^3 + d^2 - 2d} \left((d+1) |\langle x, y \rangle|^2 - |\langle x, \overline{y} \rangle|^2 - \|x\|^2 \|y\|^2 \right).$$

(Note that $a \ge 0$.)

Proof. Let $x, y \in \mathbb{C}^d$ and set $X = |x\rangle\langle x|$ and $Y = |y\rangle\langle y|$. Then, from (A.5), $P(X \otimes Y)$ is separable and there exist $a, b, c \in \mathbb{C}$ such that

$$P(X \otimes Y) = a(I_d \otimes I_d) + b\Delta_d + c|\Omega_d\rangle \langle \Omega_d|.$$

From (A.11), (A.13) and (A.12) we get a, b, c as required.

Corollary A.7. If $\lambda \in [\frac{-1}{d+1}, 1]$, then $(I_d \otimes I_d) + \lambda(|\Omega_d\rangle \langle \Omega_d| + \Delta_d) \in \mathbb{M}_d \otimes \mathbb{M}_d$ is separable.

Proof. Let $x_1 = r_1 e_1, y_1 = r_1 e_2, x_2 = y_2 = r_2 e_1$, where $r_1 = \left(\frac{d^3 + d^2 - 2d}{d+1}\right)^{\frac{1}{4}}$ and $r_2 = \left(\frac{d^3 + d^2 - 2d}{d-1}\right)^{\frac{1}{4}}$. From the above theorem,

$$P(|x_1\rangle\langle x_1|\otimes|y_1\rangle\langle y_1|) = (I_d\otimes I_d) - \frac{1}{d+1}(|\Omega_d\rangle\langle\Omega_d| + \Delta_d)$$
 and
 $P(|x_2\rangle\langle x_2|\otimes|y_2\rangle\langle y_2|) = (I_d\otimes I_d) + (|\Omega_d\rangle\langle\Omega_d| + \Delta_d)$

are separable. Now let $\lambda \in [\frac{-1}{d+1}, 1]$. Then $\lambda = t(\frac{-1}{d+1}) + (1-t)$ for some $t \in [0, 1]$, and hence

$$tP(|x_1\rangle\langle x_1|\otimes|y_1\rangle\langle y_1|) + (1-t)P(|x_2\rangle\langle x_2|\otimes|y_2\rangle\langle y_2|) = (I_d\otimes I_d) + \lambda(|\Omega\rangle\langle\Omega_d| + \Delta_d)$$

is separable.

References

- [And04] T. Ando. Cones and norms in the tensor product of matrix spaces. *Linear Algebra Appl.*, 379:3–41, 2004. [BCS20] I. Bardet, B. Collins, and G. Sapra. Characterization of equivariant maps and application to entanglement
- detection. Ann. Henri Poincaré, 21(10):3385–3406, 2020.
- [BhOs20] B. V. Rajarama Bhat and Hiroyuki Osaka. A factorization property of positive maps on C*-algebras. Int. J. Quantum Inf., 18(5):2050019, 6, 2020.
- [ChCh20] Senrui Chen, and Eric Chitambar. Entanglement-breaking superchannels. Quantum 4, 299 (2020).
- [Cho75] M. D. Choi. Completely positive linear maps on complex matrices. Linear Algebra and Appl., 10:285–290, 1975.
- [ChKo06] D. Chruściński and A. Kossakowski. On partially entanglement breaking channels. Open Syst. Inf. Dyn., 13(1):17-26, 2006.
- [CMW19] M. Christandl, A. Müller-Hermes, and Michael M. Wolf. When do composed maps become entanglement breaking? Ann. Henri Poincaré, 20(7):2295–2322, 2019.
- [COS18] B. Collins, H. Osaka, and G. Sapra. On a family of linear maps from $M_n(\mathbb{C})$ to $M_{n^2}(\mathbb{C})$. Linear Algebra Appl., 555:398–411, 2018.
- [CYT17] Lin Chen, Yu Yang, and Wai-Shing Tang. Schmidt number of bipartite and multipartite states under local projections. Quantum Inf. Process., 16(3):Paper No. 75, 27, 2017.
- [CYT19] Lin Chen, Yu Yang, and Wai-Shing Tang. Positive-partial-transpose square conjecture for n = 3. Phys. Rev. A, 99:012337, Jan 2019.
- [HHH96] M. Horodecki, P. Horodecki, and R. Horodecki. Separability of mixed states: necessary and sufficient conditions. Phys. Lett. A, 223(1-2):1–8, 1996.
- [Hir03] T. Hiroshima. Majorization criterion for distillability of a bipartite quantum state. Phys. Rev. Lett., 91:057902, 2003.
- [HLLM18] M. Huber, L. Lami, C. Lancien, and A. Müller-Hermes. High-dimensional entanglement in states with positive partial transposition. *Phys. Rev. Lett.*, 121:200503, Nov 2018.
- [HLV00] P. Horodecki, M. Lewenstein, G. Vidal, and I. Cirac. Operational criterion and constructive checks for the separability of low-rank density matrices. *Phys. Rev. A* (3), 62(3):032310, 10, 2000.

- [Hol98] A. S. Holevo. Quantum coding theorems. Russian Math. Surveys, 53(6):1295–1331, 1998. https://arxiv.org/abs/quant-ph/9809023.
- [Hor97] P. Horodecki. Separability criterion and inseparable mixed states with positive partial transposition. Phys. Lett. A, 232(5):333–339, 1997.
- [HSR03] M. Horodecki, P. W. Shor, and M. B. Ruskai. Entanglement breaking channels. Rev. Math. Phys., 15(6):629–641, 2003.
- [Hua06] Siendong Huang. Schmidt number for quantum operations. Phys. Rev. A, 73:052318, May 2006.
- [Jam72] A. Jamioł kowski. Linear transformations which preserve trace and positive semidefiniteness of operators. Rep. Mathematical Phys., 3(4):275–278, 1972.
- [JSS13] N. Johnston, L. Skowronek, and E. Stø rmer. Generation of mapping cones from small sets. Linear Algebra Appl., 438(7):3062–3075, 2013.
- [KMP18] M. Kennedy, N. A. Manor, and V. I. Paulsen. Composition of PPT maps. Quantum Inf. Comput., 18(5-6):472–480, 2018.
- [Kra71] K. Kraus. General state changes in quantum theory. Ann. Physics, 64:311-335, 1971.
- [MoZi10] Lenka Moravčíková and Mário Ziman. Entanglement-annihilating and entanglement-breaking channels. J. Phys. A, 43(27):275306, 11, 2010.
- [NiKe01] M. A. Nielsen and J. Kempe. Separable states are more disordered globally than locally. Phys. Rev. Lett., 86:5184–5187, 2001.
- [Pau02] V. Paulsen. Completely bounded maps and operator algebras, volume 78 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2002.
- [Per96] A. Peres. Separability criterion for density matrices. Phys. Rev. Lett., 77(8):1413-1415, 1996.
- [PiMo07] Marco Piani and Caterina E. Mora. Class of positive-partial-transpose bound entangled states associated with almost any set of pure entangled states. Phys. Rev. A, 75:012305, 2007.
- [RaAl07] Kedar S. Ranade and Mazhar Ali. The Jamiołkowski isomorphism and a simplified proof for the correspondence between vectors having Schmidt number k and k-positive maps. Open Syst. Inf. Dyn., 14(4):371–378, 2007.
- [RJK⁺12] M. B. Ruskai, M. Junge, D. Kribs, P. Hayden, and V. Winter. Operator structures in quantum information theory. Final Report No. 12w5084. Banff International Research Station, 2012.
- [RJP18] M. Rahaman, S. Jaques, and V. I. Paulsen. Eventually entanglement breaking maps. J. Math. Phys., 59(6):062201, 11, 2018.
- [Sch06] E. Schmidt. Math. Ann., 63:433, 1906.
- [Sko11] L. Skowronek. Cones with a mapping cone symmetry in the finite-dimensional case. Linear Algebra Appl., 435(2):361–370, 2011.
- [Sto86] E. Størmer. Extension of positive maps into b(h). J. Funct. Anal., 66(2):235–254, 1986.
- [Sto09] E. Stø rmer. Duality of cones of positive maps. Münster J. Math., 2:299–309, 2009.
- [Sto11] E. Stø rmer. Mapping cones of positive maps. Math. Scand., 108(2):223-232, 2011.
- [Sto13] E. Stø rmer. Positive linear maps of operator algebras. Springer Monographs in Mathematics. Springer, Heidelberg, 2013.
- [Sto18] E. Stø rmer. Mapping cones and separable states. Positivity, 22(2):493-499, 2018.
- [SSZ09] L. Skowronek, E. Størmer, and K. Życzkowski. Cones of positive maps and their duality relations. J. Math. Phys., 50(6):062106, 18, 2009.
- [TeHo00] B. M. Terhal and P. Horodecki. Schmidt number for density matrices. *Phys. Rev. A (3)*, 61(4):040301, 4, 2000.
- [WeHo02] R. F. Werner and A. S. Holevo. Counterexample to an additivity conjecture for output purity of quantum channels. J. Math. Phys., 43(9):4353–4357, 2002.
- [YLT16] Yu Yang, Denny H. Leung, and Wai-Shing Tang. All 2-positive linear maps from $M_3(\mathbb{C})$ to $M_3(\mathbb{C})$ are decomposable. Linear Algebra Appl., 503:233-247, 2016.

Department of Mathematics, Indian Institute of Technology Madras, Chennai, Tamilnadu 600036, India Email address: r.deva1992@gmail.com

Chennai Mathematical Institute, H1, SIPCOT IT Park, Siruseri, Kelambakkam 603103, India Email address: niru.mallick@gmail.com

Department of Mathematics, Indian Institute of Technology Madras, Chennai, Tamilnadu 600036, India Email address: sumeshkpl@gmail.com, sumeshkpl@iitm.ac.in