

Statistical Issues in Serial Killer Nurse Cases

Richard D. Gill^{1*}, Norman Fenton² and David Lagnado³

¹ Mathematical Institute, Leiden University, Leiden, The Netherlands; gill@math.leidenuniv.nl

² Queen Mary University of London, London, GB; n.fenton@qmul.ac.uk

³ University College London, London, GB; d.lagnado@ucl.ac.uk

* Correspondence: gill@math.leidenuniv.nl

Abstract: We study statistical aspects of the case of the British nurse Ben Geen, convicted of 2 counts of murder and 15 of grievous bodily harm following events at Horton General Hospital (in the town of Banbury, Oxfordshire, UK) during December 2013-February 2014. We draw attention to parallels with the cases of nurses Lucia de Berk (the Netherlands) and Daniela Poggiali (Italy), in both of which an initial conviction for multiple murders of patients was overturned after reopening of the case. We pay most attention to the investigative processes by which data, and not just statistical data, is generated; namely, the identification of past cases in which the nurse under suspicion might have been involved. We argue that the investigation and prosecution of such cases is vulnerable to many cognitive biases and errors of reasoning about uncertainty, complicated by the fact that fact-finders have to determine not only whether a particular person was guilty of certain crimes, but whether any crimes were committed by anybody at all. The paper includes some new statistical findings on the Ben Geen case and suggests further avenues for investigation. The experiences recounted here have contributed to the writing of the hand-book Green et al. (2022), *Healthcare Serial Killer or Coincidence? Statistical Issues in Investigation of Suspected Medical Misconduct*, commissioned by the Royal Statistical Society, Statistics and the Law section. Submitted to *MDPI Laws*. This version: 30 July, 2022

Keywords: Clusters of unusual events; Baader-Meinhof phenomenon; Munchausen syndrome by proxy; confirmation bias; health care serial killers; unsafe convictions; forensic statistics.

Contents

| | |
|--|----|
| 1. Introduction | 2 |
| 2. Materials and Methods | 5 |
| 3. The case of Lucia de Berk | 6 |
| 4. The case of Ben Geen | 8 |
| 5. New findings in the Ben Geen case | 13 |
| 6. Insights from forensic psychology and criminology | 16 |
| 7. Bayesian analysis of a stylized case | 17 |
| 8. Confirmation bias | 22 |
| 9. Conclusions | 24 |

| | |
|-----------------------------|-----------|
| References | 26 |
|-----------------------------|-----------|

| | |
|-------------------------------|-----------|
| 10. Appendix | 27 |
|-------------------------------|-----------|

1. Introduction

Serial murder by health care professionals is fortunately very, very rare. The investigation and prosecution of suspected case are very complex. Fact-finders must not just determine who was the perpetrator of a string of crimes – they must also determine whether any crimes occurred at all. Clusters of unusual events in a health care context can arise for numerous reasons, which can initially be completely unsuspected. They can become associated with the presence of a particular nurse for wholly innocent reasons, as a number of well documented cases proves. Medical evidence of wrong-doing might be ambiguous or on its own, inconclusive. Statistical evidence is often brought to bear, sometimes through explicit probability calculations; but in other cases the raw numbers appear to speak for themselves, and professional statisticians are not involved (or worse still, they are ignored). High weight is given to psychological evaluations and to character judgements of colleagues at the workplace, and to striking but possibly ambiguous “smoking gun” events. The legal community, as well as the public, have little understanding of statistics and probability. Such cases are of course extremely shocking and attract great media attention. The public will often draw their own conclusions on the basis of a sinister interpretation of one or two odd events.

The lead author of this paper (Richard Gill) has now been professionally involved in three such cases: the cases of the Dutch nurse Lucia de Berk, the British nurse Ben Geen, and the Italian nurse Daniela Poggiali. The case of Lucia de Berk is very well known and a number of scholarly publications have been devoted to it. The highly complex case is well summarized on English language Wikipedia, https://en.wikipedia.org/wiki/Lucia_de_Berk, and we will give further information about the case, including literature references, later in the paper. That case started with Lucia’s arrest in 2001 and finished with her acquittal at a retrial in 2010. Ben Geen was arrested in 2004, and within a few years he was sentenced with life imprisonment. Attempts have been made by his lawyers to have the case reopened by the CCRC, but so far without success. The consensus on English language Wikipedia is that he is guilty, https://en.wikipedia.org/wiki/Benjamin_Geen. The reader who was previously unaware of the case is recommended to start by looking at what is largely the prosecution’s view presented on Wikipedia. Daniela Poggiali likewise was arrested in 2014. She spent a number of years in prison but was acquitted in 2021 at the Italian court of cassation. A first scholarly publication on the case is [1]. Knowing the Italian justice system, this might not be the last word on the case, as far as legal history is concerned.

The purpose of this paper is to draw attention to the statistical issues in the Ben Geen case, establishing parallels between it and the case of Lucia de Berk, and to point out avenues for further investigation. We believe the case deserves serious scholarly attention, and therefore “kick off” with analysis of statistical aspects (“statistical” in a broad sense!) of the case, which is of course where the authors’ expertises lie. The medical side of the case is crucial and deserves further attention too. We plan to survey that part of the case in a subsequent publication. Lead author Richard Gill is at present personally convinced that Ben Geen is innocent, not only because of the evidence presented in this paper but also thanks to familiarity with the medical and psychological evidence, as well as personal contacts with various main players in Ben Geen’s trials. In Gill’s opinion, the medical evidence is highly ambiguous, and the jury (like the media) was swayed by the psychological and circumstantial evidence presented in the case, which provided the prosecution with a psychologically compelling story (both simple and shocking) which however did not do justice to the facts.

Gill is also a co-author of an upcoming publication of the Royal Statistical Society (RSS) entitled *Healthcare Serial Killer or Coincidence? Statistical Issues in Investigation of Suspected Medical*

Misconduct, Green et al. [10]. That work is a booklet or small handbook explaining statistical issues and giving advice to all parties involved in criminal investigation and prosecution of this kind of case: hospital authorities, medical specialists, nursing experts, police investigators, prosecutors, defence lawyers, statistical and medical expert witnesses, journalists, and last but not least, the public. It was commissioned by the *Statistics and the Law* section of the RSS and has been peer reviewed in that section,¹ Under “statistical issues” we do not only mean technical matters concerning the correctness or interpretation of formal statistical methods, but also the dangers of biased data gathering, and the danger of cognitive biases in reasoning about evidence. These issues are equally relevant in what are called common law jurisdictions where investigation and adjudication is based on adversarial principles, and in what are called civil law jurisdictions where an inquisitorial system is employed. The “fact finder” is often a jury of ordinary citizens in the former case, and a judge or board of judges in the latter case. However, in the case of adversarial systems, a judge often plays a very major role in determining what evidence may be brought to the attention of the jury, and may advise how that evidence should be interpreted by the jury.

As already said, this paper focusses on both known and newly discovered statistical aspects in the Ben Geen case. The scholarly reader coming perhaps from law or medicine or sociology who is interested in general “lessons for the future” might prefer to study the RSS booklet before studying this paper, since there, the (broad sense) statistical issues are discussed systematically and with a view to how all the different parties involved in criminal law proceedings can avoid falling into various traps of reasoning and interpretation on statistical matters. The present paper could then be seen as providing “supporting material” for the RSS booklet. Needless to say, that publication depends on material concerning many more “health care serial killer” cases than just the handful with which the present authors have been involved.

The paper is organised as follows. After the present section 1 “Introduction” comes the obligatory Section 2) on *Materials and Methods*, very briefly describing data sources and data sets and statistical (and other) software which will be used when looking at the two specific cases of Lucia de Berk and of Ben Geen. After that, in Section 3, we quite briefly look at the statistical issues in the case of Lucia de Berk. The next two sections consist of a general overview of the Ben Geen case (Section 4), and then a presentation of some new statistical findings (Section 5), which we think deserve further investigation and which might be useful in future attempts to reopen the case.

We then turn in Section 6 to a discussion of the literature from criminology and forensic psychology on health care serial killers. In that literature, historical cases have been catalogued and common features identified, resulting in so called “red flag lists” – warning signs for hospital administrators. This literature allows us to come up with a very rough estimate of the so-called base rate of health care serial killers. These cases are definitely extremely rare, though naturally some cases might totally evade attention. On the other hand, the rate of unsafe convictions seems to be quite large, maybe 10%, though this is also obviously a matter of speculation. Certainly there are at least this proportion of *controversial* cases, in which there is no academic consensus about whether a particular nurse was guilty and whether indeed any crimes were committed. The striking thing about quite a few controversial cases is that there is apparently striking evidence of a cluster of unexpected cases associated with one particular

¹ Authors of the booklet are Professor Peter Green FRS, Emeritus Professor of Statistics, University of Bristol, and Distinguished Professor, University of Technology, Sydney (chairman); Professor Richard Gill, Emeritus Professor of Statistics, Leiden University; Neil Mackenzie QC, Arnot Manderson Advocates, Edinburgh; Professor Julia Mortera, Professor of Statistics, Università Roma Tre; Professor William Thompson, Professor Emeritus of Criminology, Law, and Society; Psychology and Social Behavior; and Law, University of California, Irvine. One of the appendices was provided by Professor Jane Hutton, Professor of Medical Statistics, University of Warwick.

nurse (who moreover claims to be innocent of wrong-doing), but almost no “hard evidence”. By hard evidence, we mean that the nurse was actually observed tampering with medical apparatus which they were not supposed to touch or giving the patient unauthorised and harmful medication.

We are aware that the term “hard evidence” is not a legal term. Other descriptions might be “direct evidence”, as contrasted with “indirect”, “circumstantial” or “supporting” evidence. In the case of Daniela Poggiali, pathological and toxicological evidence was supposed to suggest that a certain patient died an unnatural death. Statistical evidence was used as *supporting evidence*: the statistical association between the occurrence of similar deaths and the presence of Daniela was used as indirect evidence of Daniela’s guilt, supplying further evidence of motive and opportunity. In the case of Lucia de Berk, conviction for a large number of murders was finally overturned by thoroughly discrediting the apparently hard toxicological and pathological evidence that several particular patients had been murdered by her. The incidents allegedly caused by her were moreover shown to be unremarkable, in the light of the medical condition of the patients. The further charges of murder which had been levelled at her then collapsed too, since they relied on the fact that the nurse had already been proven in one instance to be a murderer.

There is a big danger of circular arguments going unnoticed by fact-finders in view of the different points of view and modus operandi of different expert witnesses and observers. Medical experts are more likely to conclude that medical events are unnatural when they know that a nurse was attending the patient in whose hands many other patients experienced unexpected negative outcomes.

Such cases typically have a prehistory of suspicion of one particular nurse, due to behaviour seen as disconcerting and worrisome by some colleagues and supervisors. Then, one or two surprising events take place which generate concern at management level in the hospital; management also becomes aware of the suspicions circulating around nurses and medical staff. A rapid investigation is done which generates statistics of an unlikely coincidence between other suspicious events (predating the “trigger” event or events) and the suspiciously behaving nurse’s presence. A small team of medical specialists at the hospital start studying medical case histories of the cases identified so far – thus only of cases where the hospital’s own already identified suspect is present. If they find enough cases where a patient’s health rapidly worsened in a way which was hard to explain, such cases will transition from “inexplicable” to “suspicious”, even though at the time when they occurred, they were not registered as remarkable in any way. When enough material has been collected, the police will be called in.

Statistical analysis in this investigative phase might be formal but amateuristic, or it might consist just of informal interpretation of raw numbers. In the Lucia case, an initial conviction was based on a formal statistical analysis; on appeal, the conviction was maintained and the evidence of a statistician was discarded. The raw numbers spoke for themselves. In the Ben Geen case, numbers spoke for themselves. In the Daniela Poggiali case, a formal but amateuristic statistical analysis was a key part of the prosecution case.

The cases of Lucia de Berk, Ben Geen and Daniela Poggiali all have in common that the nurse in question initially fell under suspicion due to behaviour seen as “not normal” or “unsettling” by colleagues and supervisors. In none of the cases was the nurse ever definitely observed performing unauthorised dangerous and illegal actions. All of the patients allegedly killed or severely harmed by these three nurses were already severely ill, and what became of them after the alleged action of the alleged killer nurse could well have happened anyway. All three have always consistently claimed to have been innocent. Their innocence claim was maintained despite strong pressure to admitting perhaps to a desire to lessen suffering by euthanasia, which would have given them much shorter prison sentences. There are several

dubious cases on record where this perhaps has actually happened: a number of the “known” serial killer nurses might only have that status because of fierce police interrogation procedures.

Suppose investigation of all the cases identified as suspicious (for which the presence of the already suspicious nurse was a necessary condition) fails to uncover strong evidence that patients’ sudden and at the time unexpected decline was deliberately caused by that same nurse ... this would intuitively be evidence that the nurse was *not* engaged in *serial* killings (or *serial* deliberate harmings) at all. Absence of evidence can be evidence of absence. Unexpected clusters of events can occur for reasons which at the time could not even have been guessed, as a number of past examples shows. Hence there is initially perhaps an equally small chance that the hospital has a serial killer nurse on their hands, as that some recent changes to patient population or management procedures has led to a cluster of cases. In that case, searching diligently for evidence of the hand of the nurse in the cases identified, and failing to detect it, is evidence that there is another unknown cause of the cluster.

We illustrate this principle in Section 7 with a Bayesian analysis of a simple causal model. In the Lucia case, such Bayesian models have been developed in the past. A future research project is to develop such a causal model for Ben Geen case. We do not see this as becoming part of future legal procedure. However, it seems to us that it is a legitimate and meaningful activity from an academic standpoint (no disparagement intended by the adjective “academic”) which can lead to better scientific understanding of how evidence can be processed and understood in cases like this.

This leads us in Section 8 to a discussion on confirmation bias in general. It is generally agreed that the underlying story of the case of Lucia de Berk is a case of severe confirmation bias, such that social gossip led to suspicion, suspicion led to misinterpretation of unwelcome medical events (actually partly caused my mistaken diagnoses and mistaken medication prescription). This led to the inadvertent fabrication of statistical evidence, which then contaminated the medical judgement of external medical and toxicological experts, who are used to coming to a medical judgement based on all information at their disposal, in other words, not restricted to strictly medical and toxicological information about one individual case.

Finally, in Section 9, we draw some final conclusions and make some recommendations for further research. We conclude that the cases of Lucia de Berk, Ben Geen, and Daniela Poggiali share common features which certainly in the cases of Lucia de Berk and Daniela Poggiali led to a miscarriage of justice. All three were seriously afflicted by confirmation bias, and the criminal investigations, effectively largely carried out by staff of the hospital where the nurses worked, were deeply flawed. These issues were raised by Ben Geen’s defence lawyers but suppressed by the presiding judge who gave the jury the impossible task of treating each of Ben Geen’s alleged 18 crimes entirely and exclusively on their own grounds, though the prosecution had argued that the pattern in Geen’s behaviour and in the observed events was obvious, and together with their interpretation of the syringe, showed that he was a deranged killer with a lust to harm his patients.

This seems to us grounds to suggest that Ben Geen’s conviction might not be as safe as some authorities claim, though a lot more work needs to be done, whether by independent academics and scientists or by Geen’s own defence team, to support this more fully. We make some specific suggestions at the end of the paper.

2. Materials and Methods

The materials of this study are data sets provided to the authors by the defence lawyers of Lucia de Berk and Ben Geen. In the former case they were originally submitted to police investigators by the hospitals where de Berk had worked. In the latter case, they were obtained through FOI requests made by Geen’s lawyers to numerous hospitals. All data files may be obtained from the corresponding author Richard Gill. Statistical analyses are carried out using

the open source general statistical system “R”, and the Bayesian network software for risk analysis and decision making “AgenaRisk”. The data files and programming scripts which generate the statistical graphics presented in the paper can be found in the “github” repository <https://github.com/gill1109/bengeen> or directly from the corresponding author. He can also provide extensive files on the medical evidence on the 18 patients whom Geen was accused of harming.

3. The case of Lucia de Berk

The case of the Dutch nurse formerly known as “Lucia de B.” starts with three 2×2 tables, see Figure 1.

| JKZ | | Incident in shift | | | |
|---------------|-----|-------------------|------|------|--|
| | | Yes | No | | |
| Lucia on duty | Yes | 8 | 134 | 142 | |
| | No | 0 | 887 | 887 | |
| | | 8 | 1021 | 1029 | |

| RKZ-41 | | Incident in shift | | | |
|---------------|-----|-------------------|-----|-----|--|
| | | Yes | No | | |
| Lucia on duty | Yes | 1 | 0 | 1 | |
| | No | 4 | 361 | 365 | |
| | | 5 | 361 | 366 | |

| RKZ-42 | | Incident in shift | | | |
|---------------|-----|-------------------|-----|-----|--|
| | | Yes | No | | |
| Lucia on duty | Yes | 5 | 53 | 58 | |
| | No | 9 | 272 | 281 | |
| | | 14 | 325 | 339 | |

Figure 1. Roster data from the case of Lucia de B.

The data can also be found in a recent paper [9]. It differs from that found in some earlier journal publications by correction of unimportant misprints. These particular numbers were statistically analysed and interpreted for the court in 2002 by law psychologist Henk Elffers, who had been contacted in 2001 by police investigators at the beginning of what became a ten year saga. He kindly made his reports to the court, [4,5], available for scientific research (English translations prepared by Gill).

During nearly a year ending in September 2001 when police investigations started, on a medium care ward at the Juliana Children’s Hospital (“JKZ”) in the Hague, there were in total 1029 (3×343) 8-hour shifts (3 shifts a day, 7 days a week). In 8 of them, an “incident” occurred. All 8 in the shifts of a certain nurse called Lucia. Several years earlier, at another hospital, the Red Cross Hospital (“RKZ”), during the same four months in two intensive care wards (Wards 41 and 42; the data from ward 42 misses 9 days at the beginning and end of the 4 month period) there were 5 and 14 incidents respectively. Lucia was only on duty once in RKZ-41 but on just that one occasion she netted one of the 5 incidents. She mainly worked on Ward 42, where she netted disproportionately many of the 14 incidents (she had one third of the incidents in only one sixth of the shifts). Does one need a statistician to interpret it to the board of judges of a criminal court where Lucia is being tried for serial murder? The data speak for themselves. This was certainly the opinion of the judges at Lucia’s appeal in 2004: they wrote at the beginning of their summing-up “a statistical calculation of probability plays no part at all in our [guilty] verdict”. Lucia was convicted “solely on the grounds of irrefutable and scientific medical evidence”. However, the rest of their more than 100 page summing-up makes it clear that they, and many medical experts too, were already convinced *because of the raw statistics* that Lucia’s presence at so many incidents could not be chance; her mere presence caused them to classify incidents as *inexplicable* and hence *suspicious*. Toxicological evidence concerning one death was then sufficient, by a so-called chain argument (which was in fact an informal Bayesian argument), to turn other deaths and resuscitations into “scientifically irrefutably proven” murders and murder attempts. Lucia was a particularly refined murderer since she continued to brazenly to assert her innocence, and she was so cunning in most cases as to leave no evidence at all of how she had killed her patients.

Do the data speak the truth, the whole truth, and nothing but the truth? No. Lucia’s life sentence for ten murders of children and old people got reversed. At the retrial in 2010, the

judges, in their summing up, congratulated the nurses on their devotion and their professional efforts to save the lives of their patients, lives which (they publicly announced) had been unnecessarily shortened through medical errors. The errors were caused by mis-diagnosis, chaotic management, ignorance of the content of the patients' medical dossiers They were committed by hospital specialists and hospital managers, as is abundantly clear from the medical evidence given at the re-trial.

The key to Lucia's exoneration was the invalidation of the toxicological argument concerning the "trigger-case". We now know that a coincidence of several triggers set off a witch-hunt, followed (as was already noted by observers at the time) by what seemed like a witch-trial. A nurse who stood out from the crowd with a striking appearance, a strong personality, and a colourful (dark?) past was a natural scapegoat for the mistakes being made in a failing hospital department.

We now also know that the original Lucia case numbers (Figure 1) were wrong. Events have been misclassified. Some have been shifted from one shift to an adjacent shift. Quite a few have been forgotten – they weren't "unexplained", so they weren't included as "incidents". Several real incidents (i.e., "incidents" according to the law) have been suppressed. They should have been reported to the health inspectorate, but this wasn't done. For instance, one incident was really a case of euthanasia deliberately performed by the medical doctors of a child with severe birth defects, which, illegally, was not reported as such to the inspectorate. The data from JKZ was compiled in a great hurry by persons who were already convinced they were dealing with a serial killer and embarrassed that they didn't catch her earlier. Lucia was in fact set up, in the sense that the hospital's clinical director (the head paediatrician) was waiting for a final event to clinch the case she had been building up; she had already compiled the dossiers of selected earlier events.

As Elffers wrote in his second report: these numbers didn't arise by chance. He stated that of course this did not prove that she was a murderer. He explicitly offered the court three more alternatives: Lucia was maybe just a bad nurse; maybe she had harder shifts than her colleagues; maybe she was set up by someone else. In their summing up, the board of judges explicitly showed why they discounted the first two of the three alternatives. They paid no attention to the third, and the defence had not raised that possibility either.

Why, and how, did Lucia get a re-trial? Answer: she was very lucky. There was an almost-inside whistleblower. The sister-in-law Metta de Noo of JKZ's chief paediatrician Arda Derksen-Lubsen was a medical doctor, knew her sister-in-law very well, and moreover, Metta's son and daughter, studying medicine in Leiden, lived with their uncle and aunt in the Hague. Even so, it took a couple of years for the penny to drop, that there was a lot wrong with the medical evidence being used in the case. Metta de Noo recruited her brother, philosopher of science Prof. Ton Derksen. Together the brother-sister pair wrote a book devastatingly analysing the court's arguments, gained media attention and a strong popular following especially in the academic world. They prepared and submitted an application to the CEAS, the then Netherlands equivalent of the British CCRC.

There were more lucky breaks. Lucia had first experienced some definite bad luck (though nothing like the 1 in 342 million which hit the newspapers) but later, new coincidences put some of the best medical and legal minds in the Netherlands in key positions along the tortuous route to a re-trial.

On the way, it was important to explain both to the public and to legal authorities what had been wrong with the original statistical analysis, since its results had been so widely disseminated and had had so much impact. Gill gave evidence to a judicial review committee investigating the possibility to ask the supreme court of the Netherlands to have the case reopened. The fact of the matter is, popular unrest about an unsafe conviction seems to be necessary before authorities will act on it. In the social media sphere, the scientific community

was activated and could well understand that the legal systems had not been able to properly interpret hard scientific evidence whether from statistics or from toxicology. The CEAS had indeed been set up because of some recent miscarriages of justice caused in part by misinterpretation of scientific evidence. The reputation of the Netherland legal system was at stake. Whether or not key persons believed that Lucia was innocent or guilty was irrelevant: they did understand that justice had to be seen to be done, and that the case was complex and controversial and needed to be handled with the utmost care.

4. The case of Ben Geen

The data presented in Figure 2 played a role in getting the young English nurse Ben Geen a life sentence for two counts of murder and 15 of grievous bodily harm (a 16th count of grievous bodily harm was not considered proven), in the three consecutive months of December 2003, January 2004 and February 2004. We will describe the case from the point of view of the defence, and focus on statistical issues, though of course an enormous amount of medical evidence was deployed by the prosecution too. However, perhaps the most crucial piece of evidence (one could call it: the smoking gun) was an striking incident involving a syringe, for which defence and prosecution had completely different interpretations.




Figure 2. Admissions to critical care from the emergency department with a diagnosis of cardio-respiratory or respiratory arrest or hypoglycaemia, data: Head Nurse Brock

There are many similarities between the investigation that led to Geen's conviction and the erroneous investigation that led to Lucia's wrongful conviction. Ben Geen was a striking figure, and the hospital he worked at was in serious difficulties, leading to overworked staff and chaotic administration. Some odd events made hospital authorities suspicious of him. A very rapid and intensive search was made for evidence which might implicate him in various events in the emergency ward where he worked, resulting in a huge dossier which was passed to police. A background factor in the case could well have been the so-called "Shipman effect". The Ben Geen case occurred shortly after the Shipman Enquiry, which blamed health-care administrators for not earlier noticing serial killer doctor Harold Shipman, who, as the enquiry determined, maybe murdered 250 patients in all. Some speak of "Shipman hysteria".

Learning from mistakes is good, but a new danger then arises that by learning the wrong lessons from one kind of mistake, one might increase the chance of making the opposite mistake. If the rate of false convictions goes down but nothing else changes, the rate of false acquittals will go up. The more easily a health-care system goes into alarm-mode because of suspicion that it harbours a health-care serial killer, the more often innocent health-care professionals will trigger an alarm.

The key data-set in Figure 2 was presented as evidence to the court by Michelle Brock, head-nurse of the Accidents and Emergency department where Ben Geen worked, at Horton General Hospital. This is a small hospital in the provincial market town of Banbury in North Oxfordshire. Together with a dossier of perhaps 30 incidents all from December 2003 onwards, it had initially been compiled in great haste before the case was reported to the police. Michelle and some colleagues based their work on patient records and nurse attendance records at the hospital, *looking only at what happened during Ben's shifts*. Their investigation appears to have been triggered and guided by recent memory and gossip. Ben, who was a trainee nurse, had won a higher qualification at the beginning of December, allowing him to work under less supervision than before. The final trigger for their investigation had been two sudden and surprising collapses of patients who had just entered Accident & Emergency (A&E, also often referred to as ED: Emergency Department) on Thursday 5 February. Ben had reported sick on Friday, and had had a free weekend after that. He was arrested on Monday evening, 9 February 2004, when he came in to work to do a night shift. That is only one third of the way into the last bar of the bar-chart. The bar-chart was also known to the medical experts who were consulted on the 18 individual cases. There seems no doubt that it had impact on everyone involved in the trial, including journalists covering the trial.

We catch a glimpse from the chart of the fact that a lot of old people and people with existing serious health problems get brought to emergency care during the winter months of December, January and February with acute problems involving heart and lungs (a hot summer is also a danger period). A common diagnosis is cardio-respiratory arrest (the heart has stopped working and consequently the lungs too), much less common is "pure" respiratory arrest (the lungs have stopped working); fairly common is hypoglycaemia: a fall in blood glucose level. It causes fainting; breathing stops or is much suppressed. It can be caused by too much insulin or other glucose lowering diabetes tablets, delaying or missing a meal, not eating enough carbohydrate, unplanned physical activity, more strenuous exercise than usual, drinking alcohol – the risk of hypoglycaemia increases, the more alcohol you drink. In the bar-chart, nurse Brock has combined the three "standard" categories cardio-respiratory, respiratory, and hypoglycaemic arrest; but what is the correct category is hard to guess when a patient presents (arrives at the hospital). Past medical history, and future medical events will give clues as to what was actually going on. In an emergency situation, past medical history may be unknown.

Hospital nurses and authorities had been worried by the behaviour of the young male trainee nurse Ben Geen already before December 2003. His father was in the army, his mother was a nurse. He had been in the territorial army medical corps, and his ambition was to be qualified and then transferred as a combat medic to a military field hospital in Iraq. He was energetic and very "present", keen to get action and to get experience. He made some other nurses nervous. They were calling him "Ben Allitt" behind his back, not a kind nick-name, since Bev Allitt is the very well-known name of a pre-Shipman famous English convicted serial killer nurse.

In December 2003 the numbers of patients reaching an overstressed emergency ward in a small underfunded provincial hospital, threatened by closure, probably understaffed, probably lacking resident consultants in critical specialisms, was bigger than ever. There were two "surprising" events when patients who were initially thought to be in fairly good shape suddenly, and at the time unexplainably, worsened. Ben was usually around when anything happened simply because he was usually around: he was working double shifts in order to gain more and more experience as fast as possible, and also often filled in for absent colleagues.

On Thursday 5 February 2004, at the end of an exceptionally hectic day, a chronic alcoholic diabetic presented himself in the hospital (referred to ED by his doctor), on account of severe gastric pain and vomiting, and suffering fainting fits. Ben took a blood sample. The patient

suddenly worsened and later had little idea what else Ben did to him. Ben certainly inserted a *canula* (a tube that can be inserted into the body, for the delivery or removal of fluid or for the gathering of samples) and the patient was transferred to the Critical Care department, CC. The real problem for Ben came later: Ben went home with, unknown to himself (he said), a used plastic needle-less syringe containing some muscle relaxant in his nurse's smock. Such a syringe is used to administer necessary medications, including a muscle relaxant, through the *canula* prior to inserting breathing and feeding tubes into patients on the way to Critical Care. Ben stayed home sick on Friday, and then had the weekend free. His girlfriend, another nurse, doing the washing, told him off for this (she said) and told him to take it back as soon as possible. So, on Monday evening – with the syringe in his coat pocket – he was met by policemen as he entered the hospital. In some panic (he said) he stupidly further emptied the remaining contents of the syringe into his pocket. The Prosecution claimed that Ben tried to harm patients by injecting them with this stuff so that he could then play the hero, helping to resuscitate them - the so-called "Munchhausen by proxy" syndrome.

Ben and his girl-friend gave conflicting reports of the colour of the fluid in the syringe. In fact, the medication in question changes colour when exposed to air, from transparent to bright yellow. The prosecution argued that the discrepancy showed that Ben also often used the same syringe to harm patients, and that the worn markings on the syringe confirmed that it was his common weapon. The defense discovered that the markings rapidly wear off if a syringe is kept for a short while in a damp pocket.

At his trial, the Crown secured the services of a famous and experienced expert (a highly distinguished professor of Anaesthesiology), who found a number of the events highly suspicious; another confidently swore that never in his long experience had he met with an *unexplained* respiratory arrest.

Experts would agree with this claim because they strongly believe that all respiratory arrests are "explainable". The problem is that different experts often give different explanations. Whether a collapse is diagnosed as cardio-respiratory, respiratory or hypoglycaemic can be pretty arbitrary. Perhaps only weeks later (if the patient recovers and more has been learnt of their state of health and underlying conditions) might it be possible to give a more confident diagnosis of the cause of events in the past.

When either heart or lungs get into difficulty, the other organ rapidly gets into difficulties too. Hypoglycaemic arrest (critically low blood glucose levels) always involves breathing problems (you faint when not enough oxygen is reaching your brain) and can trigger further deterioration of heart and lung function. Reduced oxygen levels affect brain, heart, lungs. Muscles burn oxygen, the brain burns oxygen. All arrests are explainable, but the categories which are ticked on forms in the patient's dossier and in the hospital's administrative records may differ and may be revised in the light of later events. The categories which tend to be chosen by nurses, doctors and administrators may depend on who is doing it, and may show trends and jumps as time goes by. Just one occurrence of an unusual diagnosis alerts people to its existence, and they start seeing it every day: the well known *Baader-Meinhof phenomenon* or *Frequency Illusion*.

At the time each had actually occurred, each of the 18 cases in the criminal charges against Ben had been considered "normal". The last two had surprised some people (certainly not all), but because of earlier suspicion and gossip, they triggered an emergency weekend-long internal hospital investigation, in which more than 30 dossiers of patients who had in recent months gone through Emergency *while Ben was on duty* were combed through, resulting in a dossier of 18 cases to hand over to the police on Monday. In fact the team had access to 4000 patient medical records but were not interested in what happened when Ben was not there. Expert witnesses for the defence later described how explainable each of the 18 was, although they were honest enough to admit that some cases were too complex to come to any clear

conclusion. The prosecution had more expensive and more court-experienced experts than the defence. The prosecution experts were of course specifically hired to point out anomalies in each of the selected 18 cases, and tended to be rather confident of their diagnoses. Prosecution experts are “instructed” by the prosecution, defence experts are “instructed” by the defence. Experts will report what documents they were given to study, and on which their “opinion” is based, but prosecution experts also tend to receive a lot of further verbal information (much of it hypothetical) about the case from police or prosecutors. That certainly happened in this case.

Ben must have used a myriad of different techniques to cause all these unexplained medical emergencies and in many cases the expert witnesses called by the Crown in fact had conflicting ideas of what he might have done; but they agreed that he ‘must have done something’. All of the 18 patients were very sick, and what happened to each was what you may well have expected to happen in view of their existing severe and often complex conditions. But sometimes developments are fast, you do not “see them coming”, and so a sudden worsening takes some nurses or some doctors by surprise. People, including Ben himself, did notice Ben often being there when such events took place. He had said, and said it in court again, that he thought he had been jinxed.

Ben’s unemotional and careful account of what he could recall that he had seen and done in each case (he had received military interrogation training), the impression he gave that he knew the law better than the lawyers, an eminent professor’s categorical statement that he had never seen an unexplained respiratory collapse in all his career, and the smoking gun which was the syringe, together clinched the matter for the jury. The judge, in his summing up made it very clear what verdict he expected from the jury, who had sat through the presentation of interminable complex medical evidence (in 18 cases).

Interestingly, the judge had earlier instructed the jury to consider each of the 18 cases entirely on their own merits. This would seem to be quite a meaningless injunction. One can only guess at the thinking of the jury, but if the members of the jury were convinced by the incident with the syringe that Ben was deliberately harming patients and lying about his activities, then they would clearly tend to think that each case about which suspicions could be raised was likely indeed another case where Ben was intent on harming a patient; and the number of such cases could only have strengthened their conclusion. The prosecution argued that there had been an unlikely large number of a particular and suspicious kind of event during Ben’s shifts. Ben himself admitted that the number was strange. Whether it really was excessive is still not known, since we do not have data on his working hours, we do not have times of all the events, we know next to nothing about events which occurred when he was not on duty. We do have plenty of evidence that the classification of events was highly biased, and this bias was initiated in the hospital, as we will see.

The defence employed two expert witnesses who each delivered a comprehensive report on the problem of deducing the reality of an observed pattern from medical events in an observational setting (medical statistician Jane Hutton, clinical physiologist David Denison; their reports can be obtained from Gill). They were both definitive that a “pattern” could not be established without statistical analysis and that it would require further information, and in particular, requires blinding of medical experts, who should evaluate whether or not events were suspicious without knowing whether or not Geen was present. The judge did not allow this evidence to be presented in person by Hutton or Denison in the court, and told the jury that it consisted merely of common sense. The pattern was evident: 18 cases when Geen was present, and each should be judged on its own merits. No statistical evidence was ever considered beyond the raw numbers presented by nurse Brock and the opinion of a notable anaesthesiologist that “unexplained respiratory arrest is exceedingly rare”. In particular, no information was presented on Geen’s working hours, and no comparison was made of the rate of certain events inside and outside of his shifts.

Blood and urine samples from the trigger case showed traces of a muscle relaxant as well as of plenty of sedatives, but unfortunately some of the samples were not dated (for there was not even information on when they were taken nor by whom). Sedatives and muscle relaxant should have been present. The traces of muscle relaxant were of the same kind as was in the syringe. The consultant anaesthesiologist who had attended to the trigger patient in intensive care said that she had asked (another nurse, later) for a different one. Ben said that he was not told to administer muscle relaxant, so, of course, had not done so. Hospital records were woefully incomplete. Since the earlier cases were not at the time thought to be suspicious, samples of blood and urine had not been taken or had long ago been thrown away.

The annual pattern we see in that data is typical of many similar hospitals all over Britain, although (for reasons explained above) there are no data whatsoever about unexplained respiratory arrests. The data stored in a hospital database are administrative data. Every event has been put into a pre-existing category with an explanation, because it is not possible to enter it into the data base otherwise. The data in the database determines the fees of the medical consultants (the medical specialists) and the funding of the hospital. The data are not collected for scientific research or forensic investigation.

The three standard categories relevant to this case are *cardio-respiratory arrest*, *respiratory arrest*, and *hypoglycaemic arrest*. We already presented the data supplied to the court by Ben's head nurse, combining those three categories. Much later, Horton General hospital provided us with the data as presently archived in official hospital records, see Figure 3.




Figure 3. Admissions to CC from ED with CR, Hypo or Resp arrest, FOI data, Cardio-respiratory (blue), hypoglycaemic (green), respiratory (red)

Critically, the data were now different and also (while the categories are still separate), we have data from many more years. The total numbers of relevant cases in December 2002 and in December 2003 (previously 5 and an incredible 7 respectively) are now equal – both an unremarkable 4. The split between categories in the two periods of winter months is markedly different. In winter 2002 – 2003 it is normal, spread out over all three, but mostly cardio-respiratory. In winter 2003 – 2004 8 out of the 13 cases are categorised as *respiratory*. The total number of cases in January, in both winters, is less than in adjacent months, which is normal.

Normal case-mix (for the three categories of interest), both in this hospital and in all others (we have similar data from about 40 other hospitals all over England, for the thirteen year period 2000 – 2012), is a mix mainly of cardio-respiratory, with respiratory and hypoglycaemic normally each at roughly a fifth of the level of cardio-respiratory. They are both much less usual, but neither can be called *rare*.

There are also data in the official public enquiry after Ben's conviction, held to find out why he was not caught earlier and to prevent such a tragedy from ever occurring again. The enquiry stated: "The number in December 2003 was six and this was only one more than in December 2002". Two different numbers, yet again. The enquiry suggested that the large numbers of incidents while Ben was carrying out his attacks might have been expected anyway, due to the winter season, thereby masking incidents caused by Ben. It did heavily criticise the Emergency department for poor record keeping when updating patient medical notes and very poor registration of patient drug administrations.

5. New findings in the Ben Geen case

We will now present some summary statistics based on data obtained from Ben Geen's hospital which has been available for several years but never looked at before. Instructed by Geen's defence team, Gill was asked to answer certain questions about the "normal situation" in hospitals *like* HGH, using data that had been obtained from numerous FOI requests to hospitals all over the country. He did his best to answer exactly those questions and tried to maintain his scientific objectivity by not learning about other aspects of the case, and therefore only later took a close look at the data from HGH.

This means that Gill actually fell into a similar trap as Elffers had in the case of Lucia de B.: he tried to answer the question that had been put to him, without learning enough about the context in order to judge whether it was a meaningful question. In the Lucia case, Elffers just trusted that the hospital data had been compiled in a responsible manner. In the case of Daniela Poggiali [1], a breakthrough was made when statisticians took a look at statistical aspects of the evaluation of toxicological evidence, following up a hint from one of the medical experts for the defence. These cases are multidisciplinary; good multidisciplinary scientific research requires communication between all parties. The legal context has a tendency for communication of information to be highly restricted.




Figure 9. Monthly Admissions to Emergency; decomposition of data into trend, seasonal, remainder.

Let's take a look at the monthly total number of admissions in the emergency room of his hospital over a thirteen year period roughly centred on the critical end of 2003 – beginning of 2004; see Figure 9. We have given this time series to the go-to algorithm in the R package which, in an iterative procedure using a moving window of length 21 months, draws us a slowly evolving seasonal effect, a fairly smooth trend, and what is left over. This is pure data-analysis, no explicit modelling assumptions are being made, we are just applying a standard time-series algorithm called STL (LoESS) to let the data speak for itself, using the algorithm's default "parameters" for monthly time-series data. STL stands for "Seasonal and Trend decomposition using LoESS (locally estimated scatterplot smoothing)". What do we see? Up to summer 2004 it is getting continuously busier and busier. The number of patients being treated in Emergency almost doubles from about 400 per month to about 800 per month. One may wonder if the number of nurses in ED also doubled during this period – it's highly doubtful. The "too small" HGH was struggling to fight off closure threats, staff was working harder and harder to keep it open. Then, the number collapses. Possibly due to the situation which arose after Ben's arrest and trial, potential patients tended to go elsewhere, if they had the choice; but more likely, the policy of local health authorities was dramatically changed, too. 25 miles away is the very big teaching hospital Oxford Radcliffe, and in fact Banbury's Horton General is part of

the Radcliffe NHS trust group of hospitals (nowadays called the Oxford University Hospitals NHS Foundation Trust). For whatever reason, the number subsides to 500 – 600 per month. As mentioned before, there is a sudden dip in the very last month, but that is meaningless: the last month was not quite over when the data was submitted. The analysis should be redone without that observation (but when we do that, nothing substantial changes).

The seasonal effect shows a strong annual spike downwards. It's the Januaries! It is well known that in the Northern hemisphere, everywhere where there really is a "winter", people simply stay home, and in particular, don't go to hospital if they can help it, in January. There are, for instance, much less car accidents than in any other month, because much fewer people are out on the roads. Old people avoid slippery paths by staying at home. Apart from that we don't see any patterns. Accidents happen, and medical emergencies happen, at pretty constant rate during the year. Correcting months for their varying numbers of days (remember February) makes no discernible difference.

What is absolutely clear is that the number of people coming to that hospital was steadily increasing in the years before Ben's troubles. What about the numbers of nurses, of beds, of consultants? We plainly see the amplitude of the monthly deviations from what you would expect based on smooth annual trend and smooth seasonal average, increasing with the overall scale. The bigger the overall expected number, the bigger the random variation. This means in particular that at the time which interests us most (winter 2003 – 2004) the random variation is largest!

Let's look (Figure 10) at the total numbers of transfers to Critical Care (i.e., the patient is no longer waiting in the corridors for someone to make a decision, but is actually put in a hospital bed in an intensive care ward) from Emergency. We have the numbers with the interesting diagnoses of cardio-respiratory, respiratory, and hypoglycaemic arrest, but we don't know if the "arrest" had been diagnosed before the patient was brought to the hospital, or if it only occurred while the patient was waiting at Emergency. Very sick patients who have to wait a long time in Emergency before anyone can do anything with them are likely to suddenly get a lot worse while they are waiting.




Figure 10. Admissions from ED to CC with CR, Resp or Hypo arrest

What do we see? Just what we would expect, given the total numbers of admissions which we just studied. A slow increase, then a collapse to a stable, lower number. Take a look at what happens (Figure 11) if we normalise the numbers by looking at monthly totals per 100 admissions to ED. It looks as though nothing is going on here at all. December 2003 and January 2004 are low but February 2004 is high. There is a spike upwards in February 2004

(when Ben is mostly absent), and another spike upwards in November 2008. Both of those spikes coincide with local peak levels of the monthly numbers of admissions to Emergency.




Figure 11. Admissions from ED to CC with CR, Resp or Hypo arrest per 100 admissions to ED

Remember that Ben Geen did not work at the hospital after 5 February, 2004. More than half of the big peak in February 2004 has absolutely nothing to do with him. January 2004 is strikingly high but not more so than was February 2004. The only thing that is unusual in those months December 2003 – February 2004 is that events are being classified as *respiratory* arrest instead of as *cardio-respiratory* or *hypoglycaemic*. Why does the number of cardio-respiratory arrests fall so suddenly in those three months? Do we really believe that there were *no* cardio-respiratory arrests in December 2004? When were those numbers “fixed” in the official records: day by day as patients were admitted? Or retrospectively after Ben’s case started, 5 February 2004?

The prosecution case would be that the respiratory arrests were “extra” events which occurred in ED through Ben’s deliberate actions; the patients in question perhaps came to ED because of an earlier cardio-respiratory arrest. Against this is the fact that no-one ever saw Ben doing anything at all which he shouldn’t have been doing. One has to go into the medical evidence. There are original hospital records which are sketchy and difficult but not impossible for a layperson to interpret. The medical experts for the prosecution are in no agreement at all as to what caused what collapse, except that they are willing to see something strange about each one, and willing to argue that Ben *could* have been to blame. The defence experts have sound reasons to reject many of the hypothetical stories of the prosecution. Together, the expert opinions presented by both prosecution and defence witnesses do allow an educated non-medical specialist to form their own opinion (after all, the experts are writing in order to communicate to lawyers and to jurors).

The extensive dossiers concerning the 18 selected patients were debated at length during the trial. The jury decided that Ben had deliberately harmed 17 patients, and in two cases, that this caused their subsequent deaths.

6. Insights from forensic psychology and criminology

There is a major literature on the topic of “Health Care Serial Killers” (HCSKs) in the fields of criminology and forensic psychology. See especially [19]. In those fields one studies the characteristics of the kinds of persons who commit those crimes. More precisely, of those who are convicted for those crimes. Their material is necessarily mainly the evidence supplied by the winning party: the judges’ summing up or advice to jurors, media reports of the prosecution’s evidence after a successful prosecution. This means that scientists in these fields generally study past cases which led to convictions. Only a few authors explicitly include “suspicion

of” without “conviction for”. If past cases are restricted to, say the last five decades in Western nations, the list of such cases is pretty small – perhaps 20 to 30. Much depends on the definition of *serial* killer. Ben Geen was only convicted for two murders but he allegedly harmed many more; he is usually included. There are, of course, many more known or alleged cases of doctors and nurses who deliberately killed only one or two patients, and presumably more who were never formally suspected of this. Nurse Daniela Poggiali in Italy was convicted (and later, acquitted) of only two murders, but suspected of a great many more.

The list of “certified” HCSKs presented in various publications included for a while the case of Lucia de Berk. Some of the pieces of evidence which led to her conviction contributed to, and still stand in, the current list of “HCSK red flags”: warning signs, which managers in health care should keep note of, and which are used as evidence against new suspects in new cases when scientific evidence is given by psychologists or profilers. For instance, “thought by their colleagues to be a weird person” is one of the red flags. Another is, “fascination with death”. Lucia had a couple of Stephen King novels at home. “Steals medications from the hospital”. Lucia had some empty syringes at home which she had taken home for her daughter to use at primary school in a “show and tell” presentation of what her Mum’s day job was.

Most catalogues include Daniela Poggiali, Ben Geen, and Colin Norris. The latter is another British nurse and it seems that his case will be reopened due to new scientific insight into hypoglycaemic arrest. He was arrested at around the same time as Ben Geen, similarly at the peak of unrest concerning the then recent Shipman case.

Due to the Lucia de B. case, the “red flag” of “increased incidence of deaths when the nurse is on duty” has been labelled as an unreliable indicator.

It is interesting to look at the “modus operandi” of alleged HCSK’s who were convicted without powerful witness evidence of dangerous and illegal behaviour. One of the “weapons” of choice appears to be Potassium Chloride poisoning. The substance is readily available in hospitals, though sometimes not sufficiently under lock and key. Put some into the IV drip of a patient close to death and they will die very rapidly of heart failure. Yet, a day or two later, a post-mortem toxicological investigation will not be able to determine that that is what that patient died of. If the drip has been used by many nurses and its contents changed from time to time, it will be impossible to establish that it was ever tampered with. It seems to us no coincidence that many dubious cases revolve around K^+Cl^- poisoning. One might say it is a red flag for a false accusation and for prosecutions which succeeded though the accused was innocent.

Despite these disparaging remarks, these publications are extremely valuable. For instance, it is accepted that these cases are incredibly rare. The incidence is perhaps, roughly, 1 in 2 million [7], in which paper one finds a rough 1 in 1 million for nurses and doctors together; both about equally frequent in the catalogue compiled in that paper. Compare this with the 1 or 2 thousand annual deaths in Dutch hospitals by *admitted* medical errors. We will now have a look at how such information might impact reasoning through a simple Bayesian analysis of a hugely simplified case.

7. Bayesian analysis of a stylized case

Convincing Bayesian analyses of these cases are conspicuously absent from the academic literature. The landmark paper [6] announced that a Part 2 would be devoted to the Bayesian approach, but Part 2 never materialized. A comprehensive 2016 review of the use of Bayes in the law described the impediments against its success but also the small number of cases where it has been used.

The Dutch econometrician Aart de Vos published a rather nice Bayesian analysis of the case of Lucia de Berk in a Dutch language legal monthly journal. An English translation prepared by Gill can be found at <https://gill1109.com/2021/05/24/condemned-by-statisticians/>. It

should be noted that his analysis uses some “guessed” numbers concerning the analysis earlier mentioned by Elffers (which at the time was not public). In view of the now known errors in the data supplied to the courts by the hospitals, his analysis would now come out even more favourable to the defence hypothesis.

What is clear, is that extraordinary claims require extraordinarily strong evidence to support them. For a Bayesian, the prior probability that an otherwise not particularly special nurse has been carrying out deliberate murders of numerous patients in recent years, must be very small indeed. Hence, in order to overcome that prior expectation, we must have very strong evidence indeed. What about the evidence of a stunningly small p-value in some study comparing death rate when a particular nurse is present, with death-rate when they are not present?

It is crucial that such statistical evidence has been generated in a professional way, and this requires intensive collaboration between experienced statistical and medical experts. First of all, discussions in a multi-disciplinary team are needed to agree on definitions, criteria, a list of questions to be asked. Good statistical methods must be used to obtain data and to assess its quality. Assumptions needed for any statistical methodology, whether frequentist or Bayesian, must be stated explicitly, and if possible evaluated. Any statisticians involved must have deep understanding of the context. To give an example: times of death reported on death certificates are in actual fact usually the times that a certified medical doctor ascertained that a patient had already died. In a hospital, these times are often the times of the hand-over between shifts. A nurse who clocks in early and clocks out late can appear to be present at many more deaths than other nurses, cf. the Poggiali case [1].

Absence of hard evidence for actual malevolent activities of the nurse in question is evidence of absence of such activities. It is easy to illustrate this with a simple Bayes net model. The reader not already familiar with probabilistic causal modeling, Bayes nets, graphical models, is recommended to study any of the numerous tutorial introductions to this topic which can easily be found on internet. Alternatively, in the appendix to this paper we analyse the model “by hand”. It is so simple that all the calculations can be done with pen and paper, or on a spreadsheet, or with the help of any basic computer language, from first principles.

Here is our graphical model (Bayes net) for. stylised and simplified serial killer nurse case.




Figure 4. Simple illustrative Bayes net model

Our model is built around the Bayes net (DAG – directed acyclic graph) of just four nodes shown in Figure 4. The four nodes will represent binary random variables, taking the values

“yes” and “no”. The arrows between some of them represent assumed direct (but probabilistic) causal influence. Actually, it is the *lack* of an arrow which represents a substantive assumption about causality. Having ordered all variables so that all arrows point forward, the conditional probability distribution of any variables given the values of all its predecessors, only depends on the values of the predecessors which are directly linked to it.

As mentioned, one can always order the nodes of a DAG so that all arrows point from left to right. The model states that for any such ordering, the joint probability of all combinations of all values of all the variables (one for each node) can be built up by multiplying conditional probabilities of the outcome of each node, given the outcomes of its earlier listed graph parents – the variables with a direct link to it.

Our graph has two root nodes, which we denote by S for serial killer and U for unknown changes in hospital policy, medical apparatus, milk formula for babies, new medications, a not noticed viral or fungal infection of a new kind. This means that we assume that these two variables are assumed statistically independent. Both S and U can trigger a surprising cluster of cases associated with a particular nurse, either because they (he or she) truly are a serial killer, or coincidentally because they are often on duty at unfortunate times of day, or have longer shifts than other nurses, and because they attract attention by odd behaviour. Suppose that the prior probability of $S = \text{“yes”}$ is 10^{-6} and the prior probability of $U = \text{“yes”}$ is 10^{-2} .

The values of these two variables are assumed to have direct probabilistic influence on the nodes linked directly to them (known as their graph children). These are C standing for a cluster of unusual and unhappy events appearing to involve our nurse, and E standing for strong and direct evidence of the murder of one particular patient by the nurse of interest. We allow that the probability of a cluster of events given whether a serial killer is active and given whether there have been other unknown changes in the hospital, may depend on the status of both graph parents U and S . However, the conditional probability of finding (if one looks for it) direct evidence of one (or more) actual murders by the nurse in question depends only on S , whether or not a serial killer nurse is active. It doesn't depend on whether or not unknown changes to hospital procedures took place, U .

This means that a total of six conditional probabilities of graph children values given the values of their graph parents need to be written down to specify how the root nodes influence the occurrence or not of a cluster of event and the finding (if it is looked for) of direct evidence of one or more particular murders by our suspect. Four for C and two for E .

The most important conditional probabilities concerning the cluster of events C will be the probability of $C = \text{“no”}$ conditional on $S = \text{“yes”}$ and $U = \text{“no”}$, which we take to equal 10^{-3} , the probability of $C = \text{“yes”}$ conditional on $S = \text{“no”}$ and $U = \text{“yes”}$, which we take to equal 10^{-2} ; and the probability of $C = \text{“yes”}$ conditional on $S = \text{“no”}$ and $U = \text{“no”}$ will be 10^{-5} . The first of these says that a serial killer almost certainly triggers a cluster of cases. Unknown factors have some chance of triggering the observation of a cluster of cases, but the chance is small. If there are no changes in the hospital and no serial killer the chance of a cluster is very small. The remaining entry of the so-called Conditional Probability Table for node C given its parents S and U will be taken as follows: the probability of $C = \text{“no”}$ conditional on $S = \text{“yes”}$ and $U = \text{“yes”}$ will be 10^{-4} . Since the conditioning eventuality is almost impossible, (even relative to all the other unlikely events under consideration), this value is pretty irrelevant to later computations.

Regarding the node E , its only graph parent is S . We take the probability that $E = \text{“no”}$ conditional on $S = \text{“yes”}$ to be 10^{-2} while the probability that $E = \text{“yes”}$ conditional on $S = \text{“no”}$ will be 10^{-3} . Thus a serial killer is rather likely to betray themselves through evidence concerning one patient. On the other hand, a non serial killer nurse is unlikely to be caught red-handed killing just one patient. The prior probability of murder by health care professionals is quite low (though not so low as the probability of serial murder).

The graph structure says that knowing also U would not change the probabilities of different values of E .

Either by hand, or by using standard software for computations with graphical models, we can now perform numerous probability calculations. Below we show the results obtained when the graphical model, including the conditional probability tables we just specified, have been entered into the software program AgenaRisk. Exactly the same results are obtained with other popular tools such as GeNIe or Hugin, and in the statistical programming language R.

We can first of all compute the implied prior marginal probabilities of a cluster of cases and of evidence implicating a particular nurse in murdering one particular patient, see Figure 5.




Figure 5. Model with marginal probabilities computed.

Both are rather small. There is a 0.011% chance of an unexplained cluster of events and a 0.1% chance of incriminating evidence concerning one patient and the nurse. Conditional on neither, the prior marginal probabilities for both S and U are of course unchanged.




Figure 6. Model with observations entered: Cluster of unusual events observed.

When we observe first (and so far, only) the cluster of unusual events (Figure 6) the posterior probability of U increases to over 90%, while the probability of S increases to just 0.9%. In other words, before we have looked for patient level evidence of murder, by far the most likely explanation for the cluster of unusual events are unknown changes.




Figure 7. Model with observations entered: Hard evidence also observed.

When both the cluster of events and incriminating single patient evidence are observed (Figure 7), the probability that the nurse is a serial killer increases to just under 90% (while the probability of unknown changes drops to 10%). *So with these prior probabilities, even given the strongest evidence possible for just one patient, there is still doubt that that the nurse was a serial killer.*

In the Lucia de Berk case, the judges actually followed a sequential form of this reasoning (though entirely verbal). There was not just one patient for which hard evidence appeared to show that Lucia had caused it harm. There was a sequence of three, in each case, the “hard” evidence was less hard, but each one tipped the scales further. After the judges had concluded that Lucia was responsible for three deaths they took it that she was also responsible for many more in the cluster under investigation. The problem was not the argumentation, but the evidence. The hard medical evidence was not hard at all. On careful inspection and further toxicological measurements, it collapsed. Moreover, the “medical” argument that subsequent events were “suspicious” was merely, in several cases, explicitly (in the words of a medical expert witness) “because Lucia was present at so many incidents”.

On the other hand, suppose we observe a cluster of events but then fail to find strong evidence that the nurse hurt any one particular patient: now the posterior probability that an unknown unknown caused the “coincidence” is almost 100%; the probability that there was a serial killer is close to 0.




Figure 8. Model with observations entered: Hard evidence observed absent.

This example is meant to be illustrative. It is not a serious model for any specific case. In real cases, the interaction between observation of a surprising cluster of events and investigation of specific cases occurs in many progressive stages, not just in one step. The example does show that absence of evidence can in some circumstances provide very strong evidence of absence. All depends on the relative sizes of many very small probabilities. We do not find our choices of those probabilities broadly unrealistic. Their relative orders of magnitude determines the qualitative conclusions. If anything, some of the crucial numbers are unfavourable to our fictive nurse: the base-rate for serial murder by nurses is arguably twice as small as the number which we took. We do not specify what we mean by “hard evidence for one murder”. In fact, what we mean by it is actually determined by the graphical model (indicating conditional independence properties) and relative sizes of the many small probabilities that we have chosen!

Past cases show that the incidence of “unknown unknowns” is not infinitesimal. Recently in the UK, a striking cluster of events at one children’s hospital was found to be linked to a change in the supplier of baby formula milk. This was discovered by an external statistician working for an official enquiry; hospital authorities had not thought of it before and not suggested it as a possible cause.

In the Lucia case, it was later discovered that a rise of deaths and other incidents in a medium care ward was probably linked to a change in hospital policy concerning transfer from intensive care to medium care of infants suffering from essentially incurable birth defects. The policy was supposed to allow less babies to die in the hospital and more at home, however, it had backfired. That information was only known to a few top managers, and once they had committed the whole hospital to accusations of serial murder by a particular nurse, it was not going to be rapidly revealed. No defence lawyer could ever guess it, no investigative journalist would uncover it.

Each specific case will have many more details and complicating factors. The important point is that in a not negligible proportion of cases, there is almost no hard evidence against the accused. The case has been brought against them after a very long investigation, partly or entirely performed by doctors and managers at the hospital where the events happened – not by independent forensic investigators – and often hinges on some of the “red flag” items mentioned in the previous section. Gossip of colleagues; some history of the nurse having once sought psychotherapy; unusual personal character, perhaps connected to a traumatic childhood; some past minor transgressions.

Actually, there is no need to use special purpose software for computations in causal models in order to obtain the various conditional probabilities which follow from the model structure (the graph) and the “input” conditional probabilities listed above. Together, that model fixes, in a very simple way, the 16 probabilities of each possible fully specified joint outcome of the four binary variables in the graph. Hence one can simply calculate conditional probabilities of any composite outcome (subset of the 16) given any other, from first principles. Everything can be done by a spreadsheet or by a simple computer program written in any general purpose computer language. We provide such an explicit computation in Appendix 10. The value of the present model is that it could in principle be extended in order to model more features of any actual case.

It is also nice to use an online app for updating Bayes nets and we have provided one: <https://nurse.staging.agenarisk.app/>. It requires a password, please contact the corresponding author.

8. Confirmation bias

The investigation in the Ben Geen case appears to have suffered from *confirmation bias*, one of the most well-documented cognitive biases in psychology. Because of its prevalence, researchers in medicine and epidemiology have studied and identified many varieties of

bias which can easily occur in *observational studies*. Moreover, they have identified research procedures which minimise or in other ways account for possible bias. Nowadays, *evidence based science* demands that evaluation of new treatments for diseases, which also entails determining whether or not they have adverse effects on some or many patients, is based on the gold standard of a randomized controlled clinical trial. (It has also become very clear that very large samples are needed in order to gain useful evidence as to whether treatments are beneficial or harmful, especially when this might depend on various patient characteristics.)

In medicine there is a deep understanding of the problem of confounding factors, and particularly of hidden confounders: previously unanticipated factors which alter the chances of good or bad effects of a particular treatment. A landmark paper in the field is that of Sackett (1979) [17]: the author catalogues and names 35 different biases which can occur in observational studies. Prof. Jane Hutton's expert evidence in the Ben Geen case (discounted as mere common sense by the judge) was based on the immense knowledge built among those involved in medical research inspired by Sackett's paper. It was supported by the identical recommendations of a medical expert, clinical physiologist Prof. David Denison; his evidence was also discounted by the judge.

The main message from the medical field is the need for *controls* and the desirability of *blinded* evaluation of whether any identified case was a "treatment" or a "control" case. Definitions (in this case, of the three kinds of adverse event) need to be unambiguous, defined in advance, and applied to all patients in the time period under study, in a way which does not require knowledge of which nurse or nurses were present and which were handling the case.

In a serial killer nurse case, where the nurse's presence is supposed to have caused bad outcomes for patients in their care, the controls would be the cases in which similar bad outcomes occur though the nurse who is under suspicion was not present. In the Ben Geen case it is clear that many collapses of patients occurred during Ben's absence, but we do not even know exactly how many, nor how they were classified. The argument that there was an excess of normally rare respiratory arrests is arguably based on hearsay.

Broadly defined, in psychology, confirmation bias is the selective gathering and weighting of evidence to support a specific hypothesis and the failure to gather or recognize evidence that might count against that hypothesis [14]. The bias has been shown to operate in many investigative and forensic contexts [2,12]. Summarizing a large body of empirical research, [13] identifies three key stages where confirmation bias can derail investigations – in the *search*, *evaluation* and *communication* of evidence – with possible snowball effects when these stages interact [3,18]. The investigations in both Geen and de Berk show confirmatory biases at all three phases. For example, the investigative team in the Ben Geen case only examined the patient records on Geen's shifts, but not on shifts where he wasn't present. Without the comparison the diagnostic value of the evidence against Geen cannot be fairly assessed (as expert witnesses Jane Hutton and David Denison stated for the defence – yet they were disallowed by the judge to present this argument in court to the jury!). Moreover, ambiguous or previously non-suspicious cases were re-interpreted to fit with the hypothesis of Geen's guilt, and alternative innocent explanations were neglected. Finally, the evidence was communicated in a biased way, suppressing details of the investigative process such that fact-finders at later stages, e.g., judge and jury, were unaware that the evidence was selectively distorted.

These biases are particularly dangerous when a case involves complex medical and statistical evidence, which many investigators and fact-finders will not fully understand, and thus the potential for misinterpretation is magnified. They are also exacerbated by people's predilection to use stories to explain complex evidence [15,16]. These stories typically focus on the actions of a central protagonist, and a compelling story often overshadows a mass of complex and equivocal evidence[13].

We have absolutely no doubt that the team and everyone else concerned was acting with the best of possible intentions.

So far, attempts to have the case of Ben Geen reconsidered by the CCRC have proved unsuccessful. On the one hand, the story of the syringe grabs the public imagination. In our opinion, a cunning health care serial killer does not walk into the hospital with a half-full syringe of deadly poison in their pocket. Ben's story of why he emptied it, and his girl-friend's corroboration of why he went back to hospital with it anyway, ring true to us. This piece of evidence is neutral. The medical evidence is copious but ambiguous and inconclusive. There is no hard medical evidence that any one of those medical incidents was unnatural, despite intensive and biased search for such evidence. The jury were instructed to carry out an impossible assignment, namely to decide each case on its own merits. Yet they had heard the prosecution argument that there was a highly unlikely cluster of unusual cases and that Ben was present every time; this "story" was clearly supported by some of the medical experts. Ben himself thought that the number of incidents on his shifts was disturbing. But how could he have known what would have been a normal number? He had worked at the hospital for a while in the past but this was his first winter there, with, moreover, less supervision and more responsibilities.

By now, a number of authoritative statisticians have argued that the conviction is unsafe due to the flaws in the hospital investigation. According to British lawyer Wendy Hesketh, who has studied several similar cases as well as this one [11]

The hospital's illegal and unqualified investigation team was only looking for evidence to secure a conviction (Confirmation Bias) while discarding or ignoring evidence that proved Ben's innocence. The hospital's Serious Untoward Investigation Team initiated by Chief Nurse Brock in her capacity of Executive Lead for Governance consisted of several medical, nursing and medical records staff who were all untrained in forensic investigative techniques, crime scene preservation and the taking of witness statements. The team carried out an unlawful and flawed investigation, the material from which was later presented to medical experts appointed by the prosecution as legitimate. The opinion of these medical experts was based on flawed evidence, which had been given to those experts without their knowledge of how that evidence had been obtained. Expert opinion given on the basis of ignorance of improperly obtained evidence invalidates that medical expert evidence. The judge and jury were not aware at the trial that the evidence had been unlawfully obtained, nor of the risks to justice associated with it.

In fact, Hesketh wrote in the abstract of that paper "A protocol for liaison between the police and the medical profession in dealing with crime committed by health professionals was implemented after being proposed in this journal [the *Police Journal*] in 2003. However, rather than using their diverse skills to scrupulously investigate allegations of medico-crime, the police and health professions presume the accused is guilty and work together to prove this, disregarding any evidence to the contrary, thereby undermining the right to a fair trial. Whereas both were criticised in the past for failing to protect patients, they are likely to receive future criticism for contributing to serious miscarriages of justice.

9. Conclusions

In recent years one of us has been working on the Italian case of Daniela Poggiali and this has just resulted in acquittal, after a process taking more than 6 years; see [1]. That case too presents extraordinarily many similarities with the cases of Lucia de Berk and of Ben Geen. A main purpose of the present paper was to document yet more statistical evidence in the latter case, which in fact so far was not brought up in court. More statistical investigations need to be done. We need to obtain staffing numbers, both of nurses and of medical specialists, at Horton General over the months and years around winter 2003–2004, in order to find out if Geen's

claims that the pressure on the staff in the Accident and Emergency department was getting intolerable, have any support. We know that he was criticised for expressing such feelings to his superiors.

It would also be useful to find out what his working hours were during the three months in which he was accused of being on a killing spree. Was the rate of those patient collapses during his shifts really surprisingly large? We have some monthly totals and we know which particular patients were allegedly harmed by him (and when), so we also know the number of "arrests" outside his shifts. The hospital investigation team allegedly investigated around 30 incidents occurring during his shifts, but only brought about half of them to the attention of the police. One of those cases failed to convince the jury. (The patient concerned had actually collapsed in the ambulance on the way to the hospital. This tells us that the hospital's compilation of likely cases was not terribly accurate. Similarly, Lucia had some incidents attributed to her by her hospital, which actually occurred while she was on vacation). On the other hand, the total number of "arrests" (collapses) of patients in A&E, waiting for further attention, during those months was also close to 30. This means that 10 to 15 collapses causing transfer to critical care occurred while Ben was absent. If he worked a great deal of overtime, it could well be that the rate of these events was hardly different when he was on duty than when he was off duty, especially if he had a tendency to be on duty at the busiest times of day. He was eager to obtain as much experience as possible. We certainly know that the total number was actually typical for the time of year, and that it was the busiest time in the hospital A&E department that it had been for five or more years, at the busiest time of year.

We think it is high time to systematically study the pattern of medical events in this case and build a Bayes net model incorporating statistical evidence, medical evidence, and the evidence of the syringe. We hope some readers of this paper will be inspired to independently study the case.

For a systematic and comprehensive survey of lessons for the future which should be learnt from these and many other cases, worldwide, we refer to the soon to be published hand-book [10]. The opening words of that document are worth reproducing here.

Justice systems are sometimes called upon to evaluate cases in which health care professionals are suspected of killing their patients illegally. These cases are difficult to evaluate because they involve at least two levels of uncertainty. Commonly in a murder case it is clear that a homicide has occurred, and investigators must resolve uncertainty about who is responsible. In the cases we examine here there is also uncertainty about whether homicide has occurred. Investigators need to consider whether the deaths that prompted the investigation could plausibly have occurred for reasons other than homicide, in addition to considering whether, if homicide was indeed the cause, the person under suspicion is responsible. In this report, the RSS provides advice and guidance on the investigation and evaluation of such cases. This report was prompted by concerns about the statistical challenges such cases pose for the legal system. The cases often turn, in part, on statistical evidence that is difficult for lay people and even legal professionals to evaluate. Furthermore, the statistical evidence may be distorted by biases, hidden or apparent, in the investigative process that render it misleading. In this report the RSS provides advice and guidance on how to conduct investigations in such cases, with a particular focus on minimising the kinds of biases that could distort statistical evidence arising from the investigation. This report also provides guidance on how to recognise and take account of such biases when evaluating statistical evidence and more broadly on how to understand the strengths and limitations of such evidence and give it proper weight. This report is designed specifically to help all professionals involved in investigating such cases and those who evaluate such cases in the legal system, including expert witnesses. It will also be of interest to scholars and legal professionals who are interested in the role of

statistics in evidentiary proof, and more generally to anyone interested in improving criminal investigations.

Author Contributions: The first author initiated this work; the second two authors contributed much further material on Bayes nets and on forensic psychology. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Data and R scripts are available at <https://github.com/gill1109/bengeen>.

Acknowledgments: Richard Gill thanks Mark van der Werf and Flavio Azevedo for computing assistance.

Conflicts of Interest: Richard D. Gill worked largely pro bono as expert for Ben Geen's defence team during an initial attempt, 2014–2015, to have the case reviewed by the UK's CCRC. He was personally involved for many more years and with numerous other volunteers in attempts to get the cases of Lucia de Berk, Daniela Poggiali, and Ben Geen reviewed and re-opened, leading to final non-guilty verdicts in two of the three cases. Norman Fenton is an owner of the company AgenaRisk.

References

- Dotto, F., Gill, R.D. and Mortera, J. (2022) Statistical Analyses in the case of an Italian nurse accused of murdering patients. *Law, Probability and Risk* (submitted). <https://arxiv.org/abs/2202.08895>
- Dror, I. E., and Charlton, D. (2006). Why experts make errors. *Journal of Forensic Identification* **56**(4), 600. https://www.researchgate.net/publication/248440075_Why_Experts_Make_Errors
- Dror, I. E., Morgan, R. M., Rando, C., and Nakhaeizadeh, S. (2017). Letter to the editor—The bias snowball and the bias cascade effects: Two distinct biases that may impact forensic decision making. *Journal of forensic sciences* **62** (3), 832–833. <https://doi.org/10.1111/1556-4029.13496>
- Elffers, H. (2002a) *Distribution of incidents of resuscitation and death in the Juliana Kinderziekenhuis and the Rode Kruisziekenhuis* (Unofficial English translation of Dutch original). <https://www.math.leidenuniv.nl/~gill/Elffers1eng.pdf>
- Elffers, H. (2002b) *Elaborated analysis. Distribution of incidents of resuscitation and death at the Juliana Kinderziekenhuis and the Rode Kruisziekenhuis* (Unofficial English translation of Dutch original). <https://www.math.leidenuniv.nl/~gill/Elffers2eng.pdf>
- Fienberg, S. E. and Kaye, D. H. (1991) Legal and Statistical Aspects of Some Mysterious Clusters. *Journal of the Royal Statistical Society A* **154** (1), pp. 61–74 <https://ssrn.com/abstract=2777146>
- Forrest, A.R.W. (1995) Nurses who systematically harm their patients. *Medical Law International* **1**, pp. 411–421. <https://doi.org/10.1177/096853329500100404>
- Gill, R.D. (2014) Rarity of respiratory arrest in ED? *arXiv:1407.2731*. <https://arxiv.org/abs/1407.2731>
- Gill, R.D., Groeneboom, P., and de Jong, P. (2018) Elementary Statistics on Trial (the case of Lucia de Berk). *Chance* **31** (4), pp. 9–15. <https://doi.org/10.1080/09332480.2018.1549809>
- Green, P., Gill, R.D., Mackenzie, N., Mortera, J., and Thompson, W. (2022). *Healthcare Serial Killer or Coincidence? Statistical Issues in Investigation of Suspected Medical Misconduct*. Royal Statistical Society, Statistics and the Law section. London.
- Hesketh, W. (2012) The police-health profession's protocol: a review. *The Police Journal* **85**, 203–220. <https://doi.org/10.1350/pojo.2012.85.3.576>
- Kassin, S., Dror, I., and Kukucka, J. (2013). The Forensic Confirmation Bias: Problems, Perspectives, and Proposed Solutions. *Journal of Applied Research in Memory and Cognition* **2**, 42–52. <https://doi.org/10.1016/j.jarmac.2013.01.001>
- Lagnado, D. A. (2021). *Explaining the Evidence: How the Mind Investigates the World*. Cambridge University Press.
- Nickerson, R. S. (1998). Confirmation bias: A ubiquitous phenomenon in many guises. *Review of General Psychology* **2**(2), 175–220. <https://doi.org/10.1037/1089-2680.2.2.175>
- Pennington, N., & Hastie, R. (1986). Evidence evaluation in complex decision making. *Journal of Personality and Social Psychology* **51**(2) 242–258. <https://doi.org/10.1037/0022-3514.51.2.242>

16. Pennington, N., & Hastie, R. (1992). Explaining the evidence: Tests of the Story Model for juror decision making. *Journal of Personality and Social Psychology*, *62*(2) 189–206. <https://doi.org/10.1037/0022-3514.62.2.189>
17. Sackett, D.L. (1979). Bias in analytical research. *J. Chron. Dis.* *32*, 51–63. [https://doi.org/10.1016/0021-9681\(79\)90012-2](https://doi.org/10.1016/0021-9681(79)90012-2)
18. Scherr, K. C., Redlich, A. D., and Kassin, S. M. (2013). Cumulative disadvantage: A psychological framework for understanding how innocence can lead to confession, wrongful conviction, and beyond. *Perspectives on Psychological Science* *15*(2), 353–383. <https://doi.org/10.1177/1745691619896608>
19. Yardley, E., and Wilson, D. (2014) In Search of the ‘Angels of Death’: Conceptualising the Contemporary Nurse Healthcare Serial Killer. *J. Investig. Psych. Offender Profil.* <https://doi.org/10.1002/jip.1434>

10. Appendix

Explicit calculations for the simple Bayes net. Recall that our graphical model had four binary nodes. Two are possible *causes*: U , “unknown”, S , “serial killer”. They are causes of the effects C , “cluster”, and E , “evidence”. In graphical model language, the four nodes of the graph stand for four binary random variables, all taking the values “YES” and “NO”. For instance there either is, or is not, a cluster of events; our nurse either is, or is not, a serial killer. Some unknown factor either does or does not potentially cause a cluster of events, and hard evidence of one particular murder may or many not be found.

In our model, U and S are statistically independent binary random variables. There are therefore four possible initial causal conditions. In particular, it is just possible that unknown factors are likely to cause a cluster of events *and* that our nurse is a serial killer. To simplify calculations “by hand”, we will discard this extremely rare possibility. Changing notation, O , U and S will now stand for three mutually exclusive and exhaustive *events*. The initial possibilities will be O (“neither U nor S ”), U and S . The new event “ S ” is the old event $\{S = \text{YES}\}$. The new event O stands for *nothing unusual is going on: no serial killer, no unknown change to the system likely to cause a cluster of events*.

C and E now stand for the events “observation a cluster of events”, and “observation of hard evidence for a murder”. We will write \bar{C} and \bar{E} for the events “no cluster of events observed”, and “hard evidence for a murder not found, despite search”.

Our model states that C and E are conditionally independent given O, C and U , and that the probability C or U only depends on whether or not S holds. The numbers we previously have for various probabilities and conditional probabilities in the Bayes net can now be summarized as

$$P(S) = s = 10^{-6}$$

$$P(U) = u = 10^{-2}$$

$$P(O) = o = 1 - u - s$$

$$P(\bar{C}|S) = c = 10^{-3}$$

$$P(C|U) = b = 10^{-2}$$

$$P(C|O) = a = 10^{-5}$$

$$P(\bar{E}|S) = d = 10^{-2}$$

$$P(E|\bar{S}) = e = 10^{-3}$$

Notice that by assuming the graph structure we have made a number of conditional independence assumptions, for instance:

$$P(C|S) = P(C|S, E) = P(C|S, \bar{E});$$

given S , C and E are statistically independent. (This statement holds both with respect to our original random variable notation, and with respect to the new events notations).

Using the definition of conditional probability $P(A|B) = P(A \text{ and } B)/P(B)$, one can write down formulas for the conditional probabilities of the three possible causes given only observation of a cluster of events; then of observation of a cluster of events and of hard evidence for a murder; and last of all, of observation of a cluster of events and observation of no hard evidence for a murder despite searching for it. The first three conditional probabilities represent the situation after the cluster has been observed but before evidence is looked for that foul play was involved. The next two groups of three represent the two alternative situations after such a search has been carried out, depending on whether or not it turned up hard evidence.

$$P(O|C) = (1 - u - s) * a / ((1 - u - s) * a + u * b + s * (1 - c))$$

$$P(U|C) = u * b / ((1 - u - s) * a + u * b + s * (1 - c))$$

$$P(S|C) = s * c / ((1 - u - s) * a + u * b + s * (1 - c))$$

$$P(O|C, E) = (1 - u - s) * a * e / ((1 - u - s) * a * e + u * b * e + s * (1 - c) * (1 - d))$$

$$P(U|C, E) = u * b * e / ((1 - u - s) * a * (1 - e) + u * b * e + s * (1 - c) * (1 - d))$$

$$P(S|C, E) = s * c * (1 - d) / ((1 - u - s) * a * (1 - e) + u * b * e + s * (1 - c) * (1 - d))$$

$$P(O|C, \bar{E}) = (1 - u - s) * a * (1 - e) / ((1 - u - s) * a * (1 - e) + u * b * (1 - e) + s * (1 - c) * d)$$

$$P(U|C, \bar{E}) = u * b * (1 - e) / ((1 - u - s) * a * (1 - e) + u * b * (1 - e) + s * (1 - c) * d)$$

$$P(S|C, \bar{E}) = s * c * (1 - d) / ((1 - u - s) * a * (1 - e) + u * b * (1 - e) + s * (1 - c) * d)$$

Here is code in the language R which calculates the probabilities of interest:

```
s = 10^-6      # p(S)
u = 10^-2      # p(U)
o = 1 - u - s  # p(O)

cCOMP = 10^-3  # p(not C | S); "COMP" stands for "complement"
b = 10^-2      # p(C | U)
a = 10^-5      # p(C | O)

d = 10^-2      # p(not E | S)
e = 10^-3      # p(E | not S)

c = 1 - cCOMP
dCOMP = 1 - d
eCOMP = 1 - e
```

```

p.O.C = o*a / (o*a +u*b + s*c)
p.U.C = u*b / (o*a +u*b + s*c)
p.S.C = s*c / (o*a +u*b + s*c)

p.O.CE = o*a*e / (o*a*e + u*b*e +s*c*dCOMP)
p.U.CE = u*b*e / (o*a*e + u*b*e + s*c*dCOMP)
p.S.CE = s*c*dCOMP / (o*a*e + u*b*e + s*c*dCOMP)

p.O.CnotE = o*a*eCOMP / (o*a*eCOMP + u*b*eCOMP + s*c*d)
p.U.CnotE = u*b*eCOMP / (o*a*eCOMP + u*b*eCOMP + s*c*d)
p.S.CnotE = s*c*d / (o*a*eCOMP + u*b*eCOMP + s*c*d)

print("Probs of O, U, S given C (%)")
round(100*c(p.O.C, p.U.C, p.S.C), 4)
sum(100*c(p.O.C, p.U.C, p.S.C))

print("Probs of O, U, S given C and E (%)")
round(100*c(p.O.CE, p.U.CE, p.S.CE), 4)
sum(100*c(p.O.CE, p.U.CE, p.S.CE))

print("Probs of O, U, S given C and not E (%)")
round(100*c(p.O.CnotE, p.U.CnotE, p.S.CnotE), 4)
sum(100*c(p.O.CE, p.U.CE, p.S.CE))

```

Here are the results of running that code:

```

> print("Probs of O, U, S given C (%)")
> print("Probs of O, U, S given C (%)")
[1] "Probs of O, U, S given C (%)"
> round(100*c(p.O.C, p.U.C, p.S.C), 4)
[1] 8.9270 90.1721 0.9008
> sum(100*c(p.O.C, p.U.C, p.S.C))
[1] 100
>
> print("Probs of O, U, S given C and E (%)")
[1] "Probs of O, U, S given C and E (%)"
> round(100*c(p.O.CE, p.U.CE, p.S.CE), 4)
[1] 0.9009 9.0999 89.9992
> sum(100*c(p.O.CE, p.U.CE, p.S.CE))
[1] 100
>
> print("Probs of O, U, S given C and not E (%)")
[1] "Probs of O, U, S given C and not E (%)"
> round(100*c(p.O.CnotE, p.U.CnotE, p.S.CnotE), 4)
[1] 9.0074 90.9835 0.0091
> sum(100*c(p.O.CE, p.U.CE, p.S.CE))
[1] 100

```

The original graphical model can also be implemented entirely in R:

```

library(Rgraphviz)
library(gRain)
yn <- c("yes", "no")
S <- cptable(~SerialKiller, values=c(1, 999999), levels = yn)
U <- cptable(~UnknownCause, values=c(1, 99), levels = yn)
E.S <- cptable(~EvidenceOneMurder:SerialKiller, values = c(99, 1, 1, 999), levels = yn)
C.SU <- cptable(~ClusterofEvents:SerialKiller:UnknownCause, values = c(999, 1, 1, 99), levels = yn)
killerNurse <- compileCPT(S, U, C.SU, E.S)
killerNurse
killerNurse$SerialKiller
killerNurse$UnknownCause
killerNurse$EvidenceOneMurder
killerNurse$ClusterofEvents
killerNurse_bn <- grain(killerNurse)
killerNurse_bn <- propagate(killerNurse_bn)
killerNurse_bn

saveHuginNet(killerNurse_bn, "killerNurse.net")

#####
# No instantiation of evidence
querygrain(killerNurse_bn, nodes = c("SerialKiller", "UnknownCause"), type = "joint")
round(100*querygrain(killerNurse_bn, nodes = c("SerialKiller", "UnknownCause"), type = "joint"))

#####
# We observe no mysterious cluster and no evidence of a murder
killerNurse_bn0 <- setEvidence(killerNurse_bn, evidence = list(ClusterofEvents="no", EvidenceOneMurder="no"))
querygrain(killerNurse_bn0, nodes = c("SerialKiller", "UnknownCause"), type = "joint")
round(100*querygrain(killerNurse_bn0, nodes = c("SerialKiller", "UnknownCause"), type = "joint"))
# 99% chance nothing going on at all. 1% chance that things have changed but we did not

#####
# We just observe a suspicious cluster of events involving our nurse
killerNurse_bn1 <- setEvidence(killerNurse_bn, evidence = list(ClusterofEvents="yes", EvidenceOneMurder="no"))
querygrain(killerNurse_bn1, nodes = c("SerialKiller", "UnknownCause"), type = "joint")
round(100*querygrain(killerNurse_bn1, nodes = c("SerialKiller", "UnknownCause"), type = "joint"))
# 90% chance something changed but our nurse is not a serial killer.
# 9% chance that actually nothing changed at all
# 1% chance we have a serial killer on our hands

#####
# We observe a suspicious cluster and strong evidence of wrong doing by our nurse on our hands
killerNurse_bn11 <- setEvidence(killerNurse_bn, evidence = list(ClusterofEvents="yes", EvidenceOneMurder="yes"))
querygrain(killerNurse_bn11, nodes = c("SerialKiller", "UnknownCause"), type = "joint")
round(100*querygrain(killerNurse_bn11, nodes = c("SerialKiller", "UnknownCause"), type = "joint"))
# 90% chance our nurse is a serial killer
# 9% chance she is not a serial killer but something else caused the cluster of events
# 1% chance nothing is going on at all; 1% chance she's a serial killer and things changed

#####

```

```
# We observe a suspicious cluster but no strong evidence of wrong doing by our nurse o
killerNurse_bn12 <- setEvidence(killerNurse_bn, evidence = list(ClusterofEvents="yes
querygrain(killerNurse_bn12, nodes = c("SerialKiller", "UnknownCause"), type = "joint
round(100*querygrain(killerNurse_bn12, nodes = c("SerialKiller", "UnknownCause"), typ
# 91% chance something changed, the nurse is not a serial killer
# 9% chance nothing going on at all
```