DECOMPOSITION OF MULTICORRELATION SEQUENCES AND JOINT ERGODICITY

SEBASTIÁN DONOSO, ANDREU FERRÉ MORAGUES, ANDREAS KOUTSOGIANNIS AND WENBO SUN

ABSTRACT. We show that, under finitely many ergodicity assumptions, any multicorrelation sequence defined by invertible measure preserving \mathbb{Z}^d -actions with multivariable integer polynomial iterates is the sum of a nilsequence and a null sequence, extending a recent result of the second author. To this end, we develop a new seminorm bound estimate for multiple averages by improving the results in a previous work of the first, third and fourth authors. We also use this approach to obtain new criteria for joint ergodicity of multiple averages with multivariable polynomial iterates on \mathbb{Z}^d -systems.

CONTENTS

1	Introduction	2
	1.1 Decomposition of multicorrelation sequences	2
	1.2 The joint ergodicity phenomenon	4
2	Main results	5
	2.1 Splitting results	5
	2.2 Convergence to the expected limit.	7
	2.3 Strategy of the paper	8
	2.4 Notation	9
3	Background Material	0
	3.1 Host-Kra Seminorms and factors	0
	3.2 Nilsystems, nilsequences and Structure Theorem	3
	3.3 Bessel's inequality	3
	3.4 General properties of subgroups of \mathbb{Z}^d and properties of polynomials 1	4
4	PET induction	6
	4.1 The van der Corput lemma	.6
	4.2 The van der Corput operation	7
5	Finding a characteristic factor	1
6	Proof of Main results	1
7	Potential future directions	7
	7.1 The two-term case with no ergodicity assumptions	37
	7.2 Integer part polynomial iterates	38
Re	eferences \ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots	0

²⁰¹⁰ Mathematics Subject Classification. Primary: 37A05; Secondary: 37A30, 28A99.

Key words and phrases. Multicorrelation sequences, nilsequences, nulsequences, joint ergodicity.

The first author was supported by ANID/Fondecyt/1200897 and Grant Basal-ANID AFB170001.

1. INTRODUCTION

1.1. **Decomposition of multicorrelation sequences.** The structure and limiting behaviour of (averages of) *multicorrelation sequences*, i.e., sequences of the form

$$(n_1,\ldots,n_k)\mapsto \int_X f_0\cdot T_1^{n_1}f_1\cdot\ldots\cdot T_k^{n_k}f_k\ d\mu_k$$

where $k \in \mathbb{N}, T_1, \ldots, T_k \colon X \to X$ are invertible and commuting (i.e., $T_i T_j = T_j T_i$ for all i, j) measure preserving transformations on a probability space $(X, \mathcal{B}, \mu), {}^1 f_0, \ldots, f_k \in L^{\infty}(\mu)$ and $n_1, \ldots, n_k \in \mathbb{Z}$, is a central topic in ergodic theory. For k = 1, Herglotz-Bochner's theorem implies that the sequence $\int_X f_0 \cdot T_1^n f_1 d\mu$ is given by the Fourier coefficients of some finite complex measure σ on \mathbb{T} (see [20] and [21]). More specifically, decomposing σ into the sum of its atomic part, σ_a , and continuous part, σ_c , we get

$$\int_X f_0 \cdot T_1^n f_1 \ d\mu = \int_{\mathbb{T}} e^{2\pi i nx} \ d\sigma(x) = \int_{\mathbb{T}} e^{2\pi i nx} d\sigma_a(x) + \int_{\mathbb{T}} e^{2\pi i nx} d\sigma_c(x) = \psi(n) + \nu(n),$$

where $(\psi(n))$ is an almost periodic sequence,² and $(\nu(n))$ is a nullsequence, i.e.,

(1)
$$\lim_{N-M\to\infty} \frac{1}{N-M} \sum_{n=M}^{N-1} |\nu(n)|^2 = 0.$$

More generally, after Furstenberg's celebrated ergodic theoretic proof of Szemerédi's theorem ([14]), for a single transformation T and iterates of the form $in, 1 \le i \le k$, there has been a particular interest in the study of the corresponding multicorrelation sequences

(2)
$$\alpha(n) = \int_X f_0 \cdot T^n f_1 \cdot \ldots \cdot T^{kn} f_k \, d\mu$$

For T ergodic (i.e., every T-invariant set in \mathcal{B} has trivial measure in $\{0, 1\}$), Bergelson, Host and Kra ([3]) showed that the sequence $(\alpha(n))$ in (2) admits a decomposition of the form $a(n) = \phi(n) + \nu(n)$, where $(\phi(n))$ is a uniform limit of k-step nilsequences (see Section 3.2 for the definition) and $(\nu(n))$ satisfies (1).³ Leibman, in [27] for ergodic systems and [28] for general ones, extended the result of Bergelson-Host-Kra to polynomial iterates.

For $d \in \mathbb{N}$, we say that a tuple $(X, \mathcal{B}, \mu, (T_n)_{n \in \mathbb{Z}^d})$ is a \mathbb{Z}^d -measure preserving system (or a \mathbb{Z}^d -system) if (X, \mathcal{B}, μ) is a probability space and $T_n \colon X \to X$, $n \in \mathbb{Z}^d$, are measure preserving transformations on X such that $T_{(0,...,0)} = \text{id}$ and $T_m \circ T_n = T_{m+n}$ for all $m, n \in \mathbb{Z}^d$.⁴ It is natural to ask whether splitting results still hold for systems with commuting transformations:

Question 1.1 (Question 2, [25]). Let $(X, \mathcal{B}, \mu, (T_n)_{n \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system, $k \in \mathbb{N}, p_1, \ldots, p_k \colon \mathbb{Z} \to \mathbb{Z}^d$ a family of polynomials, and $f_0, f_1, \ldots, f_k \in L^{\infty}(\mu)$. Under which conditions on the system

¹We say that T preserves μ if $\mu(T^{-1}A) = \mu(A)$ for all $A \in \mathcal{B}$. The tuple $(X, \mathcal{B}, \mu, T_1, \ldots, T_k)$ is a *(measure preserving) system.*

²I.e., there exists a compact abelian group G, a continuous function $\phi : G \to \mathbb{C}$, and $a \in G$ such that $\psi(n) = \phi(a^n), n \in \mathbb{N}$.

³Note that k is the number of linear iterates that appear in (2).

⁴We use the notation $T_{p_i(n)}$ to stress the fact that T is a \mathbb{Z}^d -action. If T is generated by the \mathbb{Z} -actions T_1, \ldots, T_d and $p_i = (p_{i,1}, \ldots, p_{i,d})$, we have $T_{p_i(n)} = \prod_{j=1}^d T_j^{p_{i,j}(n)}$.

can the multicorrelation sequence

(3)
$$\int_X f_0 \cdot T_{p_1(n)} f_1 \cdot \ldots \cdot T_{p_k(n)} f_k \, d\mu$$

be decomposed as the sum of a uniform limit of nilsequences and a nullsequence?

The extension of the aforementioned results from \mathbb{Z} to \mathbb{Z}^d -actions is, to this day, a challenging open problem. The main issue is that the proofs of the splitting theorems crucially depend on the theory of characteristic factors via the structure theory developed by Host and Kra ([16]), a tool that is unavailable in the more general \mathbb{Z}^d -setting. As an aside, Frantzikinakis provided a partial answer to Question 1.1 (for d = 1) in [11] that avoided the use of characteristic factors. The answer was partial in the sense that the nullsequence part was allowed to have an $\ell^2(\mathbb{Z})$ error term. A similar decomposition result for general d was proven by Host and Frantzikinakis in [12].⁵ From the point of view of applications, it is useful to have such splitting results for studying weighted averages, in particular for multiple commuting transformations.⁶

It was demonstrated in [6] that under finitely many ergodicity assumptions, the characteristic factors for the corresponding averages

(4)
$$\frac{1}{N} \sum_{n=1}^{N} T_{p_1(n)} f_1 \cdot \ldots \cdot T_{p_k(n)} f_k,$$

as in the case of \mathbb{Z} -actions, are rotations on nilmanifolds (a similar result was obtained in [18] under infinitely many ergodicity assumptions). So, it is reasonable to expect that Question 1.1 holds after postulating finitely many ergodicity assumptions (this is an open problem even in the k = 2 case–see [12]).

A partial answer towards this direction was obtained in [8] by the second author. Namely, [8, Theorem 1.5] shows that for any system $(X, \mathcal{B}, \mu, T_1, \ldots, T_k)$ with T_i and $T_i T_j^{-1}$ ergodic (for all *i* and $j \neq i$) and $f_0, \ldots, f_k \in L^{\infty}(\mu)$, the sequence

(5)
$$\int_X f_0 \cdot T_1^n f_1 \cdot \ldots \cdot T_k^n f_k \, d\mu$$

can be decomposed as a sum of a uniform limit of k-step nilsequences plus a nullsequence.

For more general expressions (as in (3)), exploiting results from [18], it is also shown in [8] that, if we further assume ergodicity in all directions, i.e., $T_1^{a_1} \cdot \ldots \cdot T_d^{a_d}$ is ergodic for all $(a_1, \ldots, a_d) \in \mathbb{Z}^d \setminus \{\mathbf{0}\}$, then for any family of pairwise distinct polynomials $p_1, \ldots, p_k \colon \mathbb{Z} \to \mathbb{Z}^d$, the sequence

(6)
$$\int_X f_0 \cdot T_{p_1(n)} f_1 \cdot \ldots \cdot T_{p_k(n)} f_k \ d\mu$$

⁵The third author showed in [23] the analogue to this result for integer parts, or any combination of rounding functions, of real polynomial iterates. For a refinement of this result, with the average of the error term taken along primes, see [25].

⁶It is worth mentioning that the splitting of (2), where the average in the null term is taken along primes, was used by Tao and Teräväinen to show the logarithmic Chowla conjecture for products of odd factors ([31]).

⁷Such multiple ergodic averages always have L^2 -limits as $N \to \infty$ ([33]).

can be decomposed as a sum of a uniform limit of D-step nilsequences plus a nullsequence.⁸ The proof of this result makes essential use of a seminorm bound estimate obtained in [18], where the (infinitely many) ergodicity assumptions are reflected (see [8, Theorem 1.6]).

In [6], the first, third and fourth authors improved the seminorm bound estimates of [18] by imposing only finitely many ergodic assumptions. Although the results in [6] are stronger than those in [18], one cannot apply them directly to [8] to improve the aforementioned results, due to the incompatibility of the methods between [6] and [8] (see Subsection 2.3 for more details).

In this article, we extend results from [6] in order to obtain splitting theorems for multicorrelation sequences involving multiparameter polynomials, postulating ergodicity assumptions which are even weaker than those in [6] on the transformations that define the \mathbb{Z}^d -action in (6); for example, we will see that the sequence $\int_X f_0 \cdot T_1^{n^2} T_2^n f_1 \cdot T_3^{n^2} T_4^n f_2 d\mu$ admits the desired splitting if we assume that $T_1, T_3, T_1 T_3^{-1}$ are ergodic.

1.2. The joint ergodicity phenomenon. In his ergodic theoretic proof of Szemerédi's theorem, Furstenberg studied the averages of the multicorrelation sequence (2). In particular, a stepping stone in the proof is the special case when the transformation T is weakly mixing (i.e., $T \times T$ is ergodic for $\mu \times \mu$), in which he showed that the averages

(7)
$$\frac{1}{N}\sum_{n=1}^{N}T^{n}f_{1}\cdot\ldots\cdot T^{kn}f_{k}$$

converge in $L^2(\mu)$ to $\prod_{i=1}^k \int_X f_i d\mu$ (which we will refer to as the "expected limit") as $N \to \infty$.⁹ It was Berend and Bergelson ([1]) who characterized when the average of the more general expression (5), i.e., for multiple commuting transformations, converges to the expected limit (and this happens exactly when $T_1 \times \cdots \times T_k$ and $T_i T_j^{-1}$ for all $i \neq j$ are ergodic).

Generalizing Furstenberg's result, Bergelson showed (in [2]) that, for a weakly mixing transformation T and essentially distinct polynomials p_1, \ldots, p_k (i.e., $p_i, p_i - p_j$ are non-constant for all $1 \le i, j \le k, i \ne j$)

$$\lim_{N \to \infty} \frac{1}{N} \sum_{n=1}^{N} T^{p_1(n)} f_1 \cdot \ldots \cdot T^{p_k(n)} f_k = \prod_{i=1}^{k} \int_X f_i \ d\mu.^{10}$$

One can think of this last result as a strong independence property of the sequences $(T^{p_i(n)})_{n \in \mathbb{Z}}, 1 \leq i \leq k$ in the weakly mixing case. It is reasonable to expect a characterization in the case of polynomial iterates, which naturally leads to a general notion of joint ergodicity:

⁸Here D depends on k, d and the maximum degree of the p_i 's. It also has a connection to the number of van der Corput operations we have to run in the induction (see Remark 5.8 for details).

⁹Throughout this paper, unless otherwise stated, all limits of measurable functions on a measure preserving system are taken in L^2 .

¹⁰For T totally ergodic (i.e., T^n is ergodic for all $n \in \mathbb{N}$) and p_1, \ldots, p_k "independent" integer polynomials, it is proved in [13] that we have the same conclusion. This fact remains true for an ergodic T and "strongly independent" real-valued polynomials iterates, $[p_1(n)], \ldots, [p_k(n)]$ ([·] denotes the floor function), as well (see [19]). These last two results also follow by a recent work of Frantzikinakis, [9], in which, for single T, we have a plethora of joint ergodicity results for a number of classes of iterates (not just polynomial). Finally, for variable real "good" polynomial iterates, one is referred to [24].

Definition 1.2. We say that the sequence of tuples $(T_{p_1(n)}, \ldots, T_{p_k(n)})_{n \in \mathbb{Z}^L}$ is jointly ergodic for μ if for every $f_1, \ldots, f_k \in L^{\infty}(\mu)$ and every Følner sequence $(I_N)_{N \in \mathbb{N}}$ of \mathbb{Z}^L ,¹¹ we have that

(8)
$$\lim_{N \to \infty} \frac{1}{|I_N|} \sum_{n \in I_N} T_{p_1(n)} f_1 \cdot \ldots \cdot T_{p_k(n)} f_k = \int_X f_1 d\mu \cdot \ldots \cdot \int_X f_k d\mu$$

When k = 1, we also say that $(T_{p_1(n)})_{n \in \mathbb{Z}^L}$ is *ergodic* for μ .

The following conjecture was stated in [6]:

Conjecture 1.3 (Conjecture 1.5, [6]). Let $d, k, L \in \mathbb{N}$, $p_1, \ldots, p_k \colon \mathbb{Z}^L \to \mathbb{Z}^d$ be polynomials and $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system. Then the following are equivalent:

- (C1) $(T_{p_1(n)}, \ldots, T_{p_k(n)})_{n \in \mathbb{Z}^L}$ is jointly ergodic for μ .
- (C2) The following conditions are satisfied:
 - (i) $(T_{p_i(n)-p_j(n)})_{n\in\mathbb{Z}^L}$ is ergodic for μ for all $1 \leq i, j \leq k, i \neq j$; and
 - (ii) $(T_{p_1(n)} \times \cdots \times T_{p_k(n)})_{n \in \mathbb{Z}^L}$ is ergodic for the product measure $\mu^{\otimes k}$ on X^k .

Answering a question of Bergelson, it was shown in [6, Theorem 1.4] that, for a polynomial $p : \mathbb{Z}^L \to \mathbb{Z}$, the sequence $(T_1^{p(n)}, \ldots, T_k^{p(n)})_{n \in \mathbb{Z}^L}$ is jointly ergodic for μ if and only if $((T_1 \times \cdots \times T_k)^{p(n)})_{n \in \mathbb{Z}^L}$ is ergodic for $\mu^{\otimes k}$ and $T_i T_j^{-1}$ is ergodic for μ for all $i \neq j$. In this paper, the strong decomposition results that we obtain allow us to deduce joint ergodicity results for a larger family of polynomials (see Theorems 2.3 and 2.4), thus addressing some additional cases in the aforementioned conjecture.

2. Main results

In this section we state the main results of the paper, and provide a number of examples to better illustrate them. We also comment on the approaches that we follow.

2.1. Splitting results. Our first main concern is to resolve the incompatibility between [6] and [8], and improve the method in [6], in order to obtain an extension of the results in [8].

Before we state our first result, we need to introduce some notations:

For $d, L \in \mathbb{N}$, the polynomial $q = (q_1, \ldots, q_d) : \mathbb{Z}^L \to \mathbb{Z}^d$ is non-constant if some q_i is non-constant.

The polynomials $p_1, \ldots, p_k \colon \mathbb{Z}^L \to \mathbb{Z}^d$ are called *essentially distinct* if they are non-constant and $p_i - p_j$ is non-constant for all $i \neq j$.¹²

For a subset A of \mathbb{Q}^d , we denote $G(A) := \operatorname{span}_{\mathbb{Q}} \{a \in A\} \cap \mathbb{Z}^d$. The following subgroups of \mathbb{Z}^d play an important role in this paper:

Definition 2.1. Let $\mathbf{p} = (p_1, \ldots, p_k), p_1, \ldots, p_k \colon \mathbb{Z}^L \to \mathbb{Z}^d$ be a family of essentially distinct polynomials with $p_i(n) = \sum_{v \in \mathbb{N}_0^L} b_{i,v} n^v$ for some $b_{i,v} \in \mathbb{Q}^d$ with at most finitely many $b_{i,v}, v \in \mathbb{N}_0^L$ nonzero.¹³ For convenience, we artificially denote p_0 as the constant zero polynomial and $b_{0,v} \coloneqq 0$ for all $v \in \mathbb{N}_0^L$. For $0 \leq i, j \leq k$, set $d_{i,j} \coloneqq \deg(p_i - p_j)$ and $G_{i,j}(\mathbf{p}) \coloneqq G(\{b_{i,v} - b_{j,v} \colon |v| = d_{i,j}\})$, where, for $v = (v_1, \ldots, v_L) \in \mathbb{N}_0^L$, we write $|v| = v_1 + \ldots + v_L$.

 $[\]frac{1}{1} \text{A sequence of finite subsets } (I_N)_{N \in \mathbb{N}} \text{ of } \mathbb{Z}^L \text{ with the property } \lim_{N \to \infty} |I_N|^{-1} \cdot |(g+I_N) \triangle I_N| = 0 \text{ for all } g \in \mathbb{Z}^L \text{ is called a } F \text{older sequence in } \mathbb{Z}^L.$

¹²In general, a polynomial $q: \mathbb{Z}^L \to \mathbb{Z}^d$ has rational coefficients (i.e., vectors with rational coordinates). ¹³Here we denote $n^v \coloneqq n_1^{v_1} \dots n_L^{v_L}$ for $n = (n_1, \dots, n_L) \in \mathbb{Z}^L$ and $v = (v_1, \dots, v_L) \in \mathbb{N}_0^L$, where $0^0 \coloneqq 1$.

6 SEBASTIÁN DONOSO, ANDREU FERRÉ MORAGUES, ANDREAS KOUTSOGIANNIS AND WENBO SUN

Our main result provides an affirmative answer to Question 1.1 under finitely many ergodicity assumptions on the groups $G_{i,j}(\mathbf{p})$, which generalizes [8, Theorem 1.5]. We say that the group $G_{i,j}(\mathbf{p})$ is ergodic for μ if any function $f \in L^2(\mu)$ that is T_a -invariant for all $a \in G_{i,j}(\mathbf{p})$ is constant.

Theorem 2.2 (Decomposition theorem under finitely many ergodicity assumptions). For $d, k, K, L \in \mathbb{N}$, let $\mathbf{p} = (p_1, \ldots, p_k)$, where $p_1, \ldots, p_k \colon \mathbb{Z}^L \to \mathbb{Z}^d$ is a family of essentially distinct polynomials of degree at most K, and let $(X, \mathcal{B}, \mu, (T_n)_{n \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system. If $G_{i,j}(\mathbf{p})$ is ergodic for μ for all $0 \leq i, j \leq k, i \neq j$, then, for all $f_0, \ldots, f_k \in L^{\infty}(\mu)$, the multicorrelation sequence

$$a(n) \coloneqq \int_X f_0 \cdot T_{p_1(n)} f_1 \cdot \ldots \cdot T_{p_k(n)} f_k \ d\mu$$

can be decomposed as a sum of a uniform limit of D-step nilsequences and a nullsequence,¹⁴ where $D \in \mathbb{N}$ is a constant depending only on d, k, K, L.

Note that Theorem 2.2 goes beyond Question 1.1 as it deals with multi-variable polynomial iterates (i.e., L > 1).

Example 1. It was proved in [8, Theorem 1.5] that for any probability space (X, \mathcal{B}, μ) and commuting transformations T_1, \ldots, T_k acting on X, if T_i and $T_i T_j^{-1}$ ergodic (for all *i* and all $j \neq i$ respectively), then for all $f_0, \ldots, f_k \in L^{\infty}(\mu)$, the multicorrelation sequence

$$a(n) \coloneqq \int_X f_0 \cdot T_1^n f_1 \cdot \ldots \cdot T_k^n f_k \ d\mu$$

can be decomposed as a sum of a uniform limit of k-step nils equences plus a nulls equence. Theorem 2.2 implies a similar result.¹⁵

The following example shows that Theorem 2.2 is stronger than [8, Theorem 1.6], which deals with single variable essentially distinct polynomial iterates:

Example 2. Let $(X, \mathcal{B}, \mu, T_1, \ldots, T_6)$ be a system with commuting transformations T_1, \ldots, T_6 and $f_0, f_1, \ldots, f_4 \in L^{\infty}(\mu)$. Using [8, Theorem 1.6], we have that the multicorrelation sequence

(9)
$$\alpha(n) = \int_X f_0 \cdot T_1^{n^2} T_2^n f_1 \cdot T_1^{n^2} T_3^n f_2 \cdot T_4^{n^3} f_3 \cdot T_5^{n^3} T_6^n f_4 \, d\mu$$

can be decomposed as the sum of a uniform limit of nilsequences and a nullsequence if $T_1^{a_1}
 \dots T_6^{a_6}$ is ergodic for all $(a_1, \dots, a_6) \in \mathbb{Z}^6 \setminus \{\mathbf{0}\}$. In contrast, via Theorem 2.2, one can get the same conclusion by only assuming that $T_1, T_2 T_3^{-1}, T_4, T_5, T_4 T_5^{-1}$ are ergodic. (Indeed, denoting $T_{(a_1,\dots,a_6)} \coloneqq T_1^{a_1} \dots T_6^{a_6}$, and e_i the vector whose *i*-th entry is 1 and all other entries are 0, since $\mathbf{p} = ((n^2, n, 0, 0, 0, 0), (n^2, 0, n, 0, 0, 0), (0, 0, 0, n^3, 0, 0), (0, 0, 0, 0, n^3, n))$, we have that $G_{1,0}(\mathbf{p}) = G_{2,0}(\mathbf{p}) = G(e_1), G_{1,3}(\mathbf{p}) = G_{2,3}(\mathbf{p}) = G_{3,0}(\mathbf{p}) = G(e_4), G_{1,4}(\mathbf{p}) = G_{2,4}(\mathbf{p}) = G_{4,0}(\mathbf{p}) = G(e_5), G_{1,2}(\mathbf{p}) = G(e_2 - e_3), G_{3,4}(\mathbf{p}) = G(e_4 - e_5).$

¹⁴The precise definition of a *D*-step nilsequence will be given in the following section. Furthermore, we say that $a: \mathbb{Z}^L \to \mathbb{C}$ is a nullsequence if for any Følner sequence (I_N) we have $\lim_{N\to\infty} \frac{1}{|I_N|} \sum_{n\in I_N} |a(n)|^2 = 0$.

¹⁵While Theorem 2.2 does not specify the step D of the nilsequence, a quick argument shows that in this case one can indeed take D = k (see Remark 6.1 for details).

2.2. Convergence to the expected limit. In [6, Theorem 1.4], the first, third and fourth authors proved the following case of Conjecture 1.3: If T_1, \ldots, T_k are commuting transformations acting on a probability space (X, \mathcal{B}, μ) , then $(T_1^{p(n)}, \ldots, T_k^{p(n)})_{n \in \mathbb{Z}^L}$ is jointly ergodic for μ if and only if $((T_1 \times \cdots \times T_k)^{p(n)})_{n \in \mathbb{Z}^L}$ is ergodic for $\mu^{\otimes k}$ and $T_i T_j^{-1}$ is ergodic for μ for all $i \neq j$. In this paper, we further extend this result:

Theorem 2.3. Let $\mathbf{p} = (p_1(n)v_1, \ldots, p_k(n)v_k), p_1, \ldots, p_k \colon \mathbb{Z}^L \to \mathbb{Z}, v_1, \ldots, v_k \in \mathbb{Z}^d$ be a family of essentially distinct polynomials. Suppose that for all $1 \leq i, j \leq k$, if $\deg(p_i) = \deg(p_j)$, then either v_i and v_j are linearly dependent over \mathbb{Z} , or $p_i(n)$ and $p_j(n)$ are linearly dependent over \mathbb{Z} (i.e., there is a non-trivial linear combination of them over \mathbb{Z} which equals to a constant). Let $(X, \mathcal{B}, \mu, (T_q)_{a \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system. Then the following are equivalent:

- (C1) $(T_{p_i(n)v_i}: 1 \leq i \leq k)_{n \in \mathbb{Z}^L}$ is jointly ergodic for μ
- (C2') The following conditions hold:
 - (i)' $(T_{p_i(n)v_i-p_j(n)v_j})_{n\in\mathbb{Z}^L}$ is ergodic for μ for all $1 \leq i,j \leq k, i \neq j$ with deg $(p_i) = \deg(p_j)$;
 - (ii) $(T_{p_1(n)v_1} \times \cdots \times T_{p_k(n)v_k})_{n \in \mathbb{Z}^L}$ is ergodic for $\mu^{\otimes k}$.

Moreover, (C2') is equivalent to

(C2) The following conditions hold:

- (i) $(T_{p_i(n)v_i-p_j(n)v_j})_{n\in\mathbb{Z}^L}$ is ergodic for μ for all $1 \le i, j \le k, i \ne j$;
- (ii) $(T_{p_1(n)v_1} \times \cdots \times T_{p_k(n)v_k})_{n \in \mathbb{Z}^L}$ is ergodic for $\mu^{\otimes k}$.

Note that the conditions in (C2) are consistent with those in Conjecture 1.3. On the other hand, the reason we provide an alternative set of equivalent conditions (C2') is that these conditions are easier to check in practise.

We now give some examples to illustrate Theorem 2.3. The first one is for polynomials of distinct degrees:

Example 3. Let $(X, \mathcal{B}, \mu, T_1, \ldots, T_k)$ be a system. Using Theorem 2.3, we conclude that $(T_1^n, T_2^{n^2}, \ldots, T_k^{n^k})_{n \in \mathbb{Z}}$ is jointly ergodic if and only if $(T_1^n \times \cdots \times T_k^{n^k})_{n \in \mathbb{Z}}$ is ergodic for $\mu^{\otimes k}$, and all the T_i 's are ergodic for μ .

We remark that Example 3 can also be proved by using arguments from [5]. We next present two examples in which the polynomials can be taken to be not necessarily of different degrees, and so, cannot be recovered by the methods of [5]:

Example 4. Let $(X, \mathcal{B}, \mu, T_1, T_2, T_3, T_4)$ be a system. Theorem 2.3 implies that $(T_1^n, T_2^n, T_3^{n^2}, T_4^{n^2})_{n \in \mathbb{Z}}$ is jointly ergodic if and only if $(T_1^n \times T_2^n \times T_3^{n^2} \times T_4^{n^2})_{n \in \mathbb{Z}}$ is ergodic for $\mu^{\otimes 4}$, and both $T_1T_2^{-1}$ and $((T_3T_4^{-1})^{n^2})_{n \in \mathbb{N}}$ are ergodic for μ .

Example 5. Let $(X, \mathcal{B}, \mu, T_1, T_2, T_3)$ be a system. Theorem 2.3 implies that $(T_1^{n^4+n^2}, T_1^{2n^4+3n}, T_2^{2n^2+2n+1}, T_3^{3n^2+3n})_{n\in\mathbb{Z}}$ is jointly ergodic if and only if $(T_1^{n^4+n^2} \times T_1^{2n^4+3n} \times T_2^{2n^2+2n+1} \times T_3^{3n^2+3n})_{n\in\mathbb{Z}}$ is ergodic for $\mu^{\otimes 4}$, and both sequences $(T_1^{-n^4+n^2-3n})_{n\in\mathbb{Z}}$ and $((T_2^2T_3^{-3})^{n^2+n})_{n\in\mathbb{Z}}$ are ergodic for μ .

Another direction for the joint ergodicity problem is verifying whether (C1) implies (C2) in Conjecture 1.3. Namely, assuming that $(T_{p_1(n)} \times \cdots \times T_{p_k(n)})_{n \in \mathbb{Z}^L}$ is ergodic for $\mu^{\otimes k}$, to find a condition, say (P), of certain sequences of actions to be ergodic, under which, we

have that $(T_{p_1(n)}, \ldots, T_{p_k(n)})_{n \in \mathbb{Z}^L}$ is jointly ergodic for μ . By combining existing results from [16, 18, 26] (see also [6, Proposition 1.2]), (P) can be taken to be: " T_g is ergodic for μ for all $g \in \mathbb{Z}^d \setminus \{\mathbf{0}\}$." Denoting $p_i(n) = \sum_{v \in \mathbb{N}_0^L, 0 \le |v| \le K} b_{i,v} n^v$ for some $b_{i,v} \in \mathbb{Q}^d$ and $K \in \mathbb{N}_0$, this result was extended in [6, Theorem 1.3], where the previous property is replaced by: " T_g is ergodic for μ for all g that belongs to the finite set R," where

$$R = \bigcup_{0 < |v| \le K} \{b_{i,v}, b_{i,v} - b_{j,v} \colon 1 \le i, j \le k\} \setminus \{\mathbf{0}\}.$$

In this paper, we replace the latter condition with an even weaker one:

Theorem 2.4. Let $d, k, L \in \mathbb{N}$, $\mathbf{p} = (p_1, \ldots, p_k)$, $p_1, \ldots, p_k \colon \mathbb{Z}^L \to \mathbb{Z}^d$ be a family of essentially distinct polynomials and $\mathbf{X} = (X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ a \mathbb{Z}^d -system. Then $(T_{p_1(n)}, \ldots, T_{p_k(n)})_{n \in \mathbb{Z}^L}$ is jointly ergodic for μ if both of the following conditions hold:

- (i) $G_{i,j}(\mathbf{p})$ is ergodic for μ for all $0 \le i, j \le k, i \ne j$;
- (ii) $(T_{p_1(n)} \times \cdots \times T_{p_k(n)})_{n \in \mathbb{Z}^L}$ is ergodic for $\mu^{\otimes k}$.

The last example for this section reflects the stronger nature of the previous theorem compared to what was previously known.

Example 6. Let $(X, \mathcal{B}, \mu, T_1, T_2, T_3, T_4)$ be a system. Then, [6, Theorem 1.3] implies that $(T_1^{n^2}T_2^n, T_3^{n^2}T_4^n)_{n\in\mathbb{Z}}$ is jointly ergodic if $((T_1^{n^2}T_2^n) \times (T_3^{n^2}T_4^n))_{n\in\mathbb{Z}}$ is ergodic for $\mu^{\otimes 2}$, and all $T_1, T_2, T_3, T_4, T_1T_3^{-1}, T_2T_4^{-1}$ are ergodic for μ . Using Theorem 2.4, we conclude that $(T_1^{n^2}T_2^n, T_3^{n^2}T_4^n)_{n\in\mathbb{Z}}$ is jointly ergodic if we instead only assume that $((T_1^{n^2}T_2^n) \times (T_3^{n^2}T_4^n))_{n\in\mathbb{Z}}$ is ergodic for $\mu^{\otimes 2}$, and all $T_1, T_3, T_1T_3^{-1}$ are ergodic for μ .

2.3. Strategy of the paper. The central ingredient in proving the main results of the paper (Theorems 2.2, 2.3 and 2.4) is to find proper characteristic factors for the average (4), i.e., sub- σ -algebras $\mathcal{D}_1, \ldots, \mathcal{D}_k$ of \mathcal{B} such that the average (4) remains invariant if we replace each f_i by its conditional expectation (see Section 3 for the definition) with respect to \mathcal{D}_i . An important type of characteristic factor, called the Host-Kra characteristic factor, was invented in [16] to study multiple averages for \mathbb{Z} -systems (see Section 3 for the definition of these factors). This concept was generalized to systems with commuting transformations in [15] (see also [30]). The main tool used in our results, special cases of which have been studied extensively in the past (see for example [5, 13, 15, 16, 18]), is the following:

Theorem 2.5. Let $d, k, K, L, s \in \mathbb{N}$, $\mathbf{p} = (p_1, \ldots, p_k)$, $p_1, \ldots, p_k \in \mathbb{Z}^L \to \mathbb{Z}^d$ be a family of essentially distinct polynomials of degrees at most K. There exists $D \in \mathbb{N}_0$ depending only on d, k, K, L, s such that for every \mathbb{Z}^d -system $\mathbf{X} = (X, \mathcal{B}, \mu, (T_n)_{n \in \mathbb{Z}^d})$, every $f_1, \ldots, f_k \in L^{\infty}(\mu)$, and every Følner sequence $(I_N)_{N \in \mathbb{N}}$ of \mathbb{Z}^L , if f_i is orthogonal to the Host-Kra characteristic factor $Z_{\{G_{i,j}(\mathbf{p})\}_{0 \leq j \leq k, j \neq i}^{\times D}}(\mathbf{X})$ for some $1 \leq i \leq k$ (i.e. the conditional expectation of f_i under $Z_{\{G_{i,j}(\mathbf{p})\}_{0 \leq j \leq k, j \neq i}^{\times D}}(\mathbf{X})$ is 0), then we have that

(10)
$$\lim_{N \to \infty} \frac{1}{|I_N|} \sum_{n \in I_N} \prod_{i=1}^k T_{p_i(n)} f_i = 0$$

In particular, if for some $1 \leq i \leq k$, $G_{i,j}(\mathbf{p})$ is ergodic for μ for all $0 \leq j \leq k, j \neq i$ and f_i is orthogonal to the Host-Kra characteristic factor $Z_{(\mathbb{Z}^d) \times kD}(\mathbf{X})$, then (10) holds.

It is worth noting that the factor $Z_{\{G_{i,j}(\mathbf{p})\}_{0\leq j\leq k, j\neq i}^{\times D}}(\mathbf{X})$ we obtain in Theorem 2.5 is not optimal, but it is good enough for our purposes.

A special case of Theorem 2.5 was proved in [6, Theorem 5.1]. In particular, Theorem 2.5 generalizes [6, Theorem 5.1] in the following ways:

- (I) The characteristic factor obtained in Theorem 2.5 is of finite step, whereas the one in [6, Theorem 5.1] is of infinite step.
- (II) The groups $G_{i,j}(\mathbf{p})$ involved in Theorem 2.5 are larger than those in [6, Theorem 5.1], which makes the characteristic factors in Theorem 2.5 smaller.

We remark that the aforementioned technical distinctions have significant influences in the applications of Theorem 2.5. First, the essential reason why one cannot directly use [6, Theorem 5.1] to improve [8, Theorem 1.5] is that the method used in [8] requires a characteristic factor of finite step. This problem is resolved by (I), enabling us to extend [8, Theorem 1.5] in this paper. Second, [6, Theorem 5.1] does not provide a strong enough characteristic factor in certain circumstances. For example, in the case of Example 3, [5, Theorem 6.5] suggests that the Host-Kra seminorms controlling (10) depend only on the transformations T_1, \ldots, T_k , whereas the upper bound provided by [6, Theorem 5.1] depends not only on the transformations T_1, \ldots, T_k but also on many compositions of them. With the help of (I) and (II), we are able to obtain (and generalize) the aforementioned upper bound of [5, Theorem 6.5].

Roughly speaking, the achievement of (I) relies on a sophisticated development of a Besseltype inequality first obtained by Tao and Ziegler in [32, Proposition 3.6]. The most technical part of this paper is the approach we use to get (II). In [6], a method was introduced to keep track of the coefficients of the polynomials while running a variation of the PET induction. However, the tracking provided there is not strong enough to imply Theorem 2.5. To overcome this difficulty, we introduce more sophisticated machinery in order to have a better control of the coefficients.

The paper is organized as follows: We provide some background material in Section 3. In Section 4, we present the variation of PET induction that we use. In Section 5, we address how (I) and (II) above can be achieved with Propositions 5.2 and 5.3, which improve Propositions 5.6 and 5.5 of [6] respectively. We conclude the section by proving Theorem 2.5. This is the bulk of the paper. In Section 6, we use Theorem 2.5 to deduce Theorems 2.2, 2.3, and 2.4, which are the main results of the paper. We conclude with some discussions on future directions in Section 7.

2.4. Notation. We denote with \mathbb{N} , \mathbb{N}_0 , \mathbb{Z} , \mathbb{Q} , \mathbb{R} and \mathbb{C} the set of positive integers, non-negative integers, rational numbers, real numbers and complex numbers respectively. If X is a set, and $d \in \mathbb{N}$, X^d denotes the Cartesian product $X \times \cdots \times X$ of d copies of X.

We will denote by e_i the vector which has 1 as its *i*th coordinate and 0's elsewhere. We use in general lower-case letters to symbolize both numbers and vectors but bold letters to symbolize vectors of vectors to highlight this exact fact. The only exception to this convention is the vector **0** (*i.e.*, the vector with coordinates only 0's) which we always symbolize in bold.

Throughout this article, we use the following notation for averages: Let $(a(n))_{n \in \mathbb{Z}^L}$ be a sequence of complex numbers, or a sequence of measurable functions on a probability space (X, \mathcal{B}, μ) . We let

 $\mathbb{E}_{n \in A} a(n) \coloneqq \frac{1}{|A|} \sum_{n \in A} a(n)$, where A is a finite subset of \mathbb{Z}^L ;

$$\overline{\mathbb{E}}_{n\in\mathbb{Z}^{L}}^{\square}a(n) \coloneqq \overline{\lim}_{N\to\infty} \mathbb{E}_{n\in[-N,N]^{L}}a(n);^{16} \\
\overline{\mathbb{E}}_{n\in\mathbb{Z}^{L}}a(n) \coloneqq \sup_{\substack{(I_{N})_{N\in\mathbb{N}}\\ \text{Følner seq.}}} \overline{\lim}_{N\to\infty} \mathbb{E}_{n\in I_{N}}a(n); \\
\mathbb{E}_{n\in\mathbb{Z}^{L}}^{\square}a(n) \coloneqq \lim_{N\to\infty} \mathbb{E}_{n\in[-N,N]^{L}}a(n) \text{ (provided that the limit exists); and} \\
\mathbb{E}_{n\in\mathbb{Z}^{L}}a(n) \coloneqq \lim_{N\to\infty} \mathbb{E}_{n\in I_{N}}a(n) \text{ (prov. the limit exists for all Følner seq. } (I_{N})_{N\in\mathbb{N}}).^{17}$$

We also consider *iterated* averages: Let $(a(h_1, \ldots, h_s))_{h_1, \ldots, h_s \in \mathbb{Z}^L}$ be a multi-parameter sequence. We let

$$\overline{\mathbb{E}}_{h_1,\dots,h_s\in\mathbb{Z}^L}a(h_1,\dots,h_s)\coloneqq\overline{\mathbb{E}}_{h_1\in\mathbb{Z}^L}\dots\overline{\mathbb{E}}_{h_s\in\mathbb{Z}^L}a(h_1,\dots,h_s)$$

and adopt similar conventions for $\mathbb{E}_{h_1,\ldots,h_s\in\mathbb{Z}^L}$, $\overline{\mathbb{E}}_{h_1,\ldots,h_s\in\mathbb{Z}^L}^{\square}$ and $\mathbb{E}_{h_1,\ldots,h_s\in\mathbb{Z}^L}^{\square}$ respectively.

We end this section by recalling the notion of a system indexed by an abelian group (G, +). We say that a tuple $(X, \mathcal{B}, \mu, (T_g)_{g \in G})$ is a *G*-measure preserving system (or a *G*-system) if (X, \mathcal{B}, μ) is a probability space and $T_g: X \to X$ are measurable, measure preserving transformations on X such that $T_{e_G} = \text{id} (e_G \text{ is the identity element of } G)$ and $T_g \circ T_h = T_{g+h}$ for all $g, h \in G$. A *G*-system will be called *ergodic* if for any $A \in \mathcal{B}$ such that $T_gA = A$ for all $g \in G$, we have that $\mu(A) \in \{0, 1\}$. In this paper, we are mostly concerned about \mathbb{Z}^d -systems and $L^2(\mu)$ -norm limits of (multiple) ergodic averages. For the corresponding norm, when it is clear from the context, we will write $\|\cdot\|_2$ instead of $\|\cdot\|_{L^2(\mu)}$.

3. Background Material

In this section we recall some background material and prove some intermediate results that will be used later throughout the paper.

3.1. Host-Kra Seminorms and factors. Host-Kra seminorms and their associated factors are arguably the main tools used to analyze the behaviour of multiple averages and correlation sequences. In what follows we give general results about these seminorms and factors, following the notation used in [6].

We first recall the notions of a factor and of the conditional expectation with respect to a factor. We say that the \mathbb{Z}^d -system $(Y, \mathcal{D}, \nu, (S_g)_{g \in \mathbb{Z}^d})$ is a factor of $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ if there exists a measurable map $\pi \colon (X, \mathcal{B}, \mu) \to (Y, \mathcal{D}, \nu)$ such that $\mu(\pi^{-1}(A)) = \nu(A)$ for all $A \in \mathcal{D}$, and $\pi \circ T_g = S_g \circ \pi$ for all $g \in \mathbb{Z}^d$.

A factor $(Y, \mathcal{D}, \nu, (S_g)_{g \in \mathbb{Z}^d})$ of $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ can be identified with an invariant sub- σ -algebra \mathcal{B}' of \mathcal{B} by setting $\mathcal{B}' := \pi^{-1}(\mathcal{D})$. Given two σ -algebras \mathcal{B}_1 and \mathcal{B}_2 , their joining $\mathcal{B}_1 \vee \mathcal{B}_2$ is the σ -algebra generated by $B_1 \cap B_2$ for all $B_1 \in \mathcal{B}_1$ and $B_2 \in \mathcal{B}_2$, *i.e.*, the smallest σ -algebra containing both \mathcal{B}_1 and \mathcal{B}_2 .

Given a factor $\pi: (X, \mathcal{B}, \mu) \to (Y, \mathcal{D}, \nu)$ and a function $f \in L^2(\mu)$, the conditional expectation of f with respect to Y is the function $g \in L^2(\nu)$, which we denote by $\mathbb{E}(f \mid Y)$, with the property

$$\int_{A} g \circ \pi \ d\mu = \int_{A} f \ d\mu \text{ for all } A \in \pi^{-1}(\mathcal{D}).$$

¹⁶We use the symbol \Box to highlight the fact that the averages are taken along the boxes $[-N, N]^L$.

¹⁷It is worth noticing that if the limit $\lim_{N\to\infty} \mathbb{E}_{n\in I_N} a(n)$ exists for all Følner sequences (in \mathbb{Z}^L), then this limit does not depend on the chosen Følner sequence.

Let $\mathbf{X} = (X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system and let \mathcal{B}_1 be an invariant sub- σ -algebra of \mathcal{B} . The relatively independent joining of \mathbf{X} with itself with respect to \mathcal{B}_1 is the measure preserving system obtained by considering the product space with the relatively independent joining, denoted by $\mu \times_{\mathcal{B}_1} \mu$, which is given by the formula:

$$\int_{X \times X} f_1 \otimes f_2 \ d(\mu \times_{\mathcal{B}_1} \mu) = \int_X \mathbb{E}(f_1 | \mathcal{B}_1) \mathbb{E}(f_2 | \mathcal{B}_1) \ d\mu$$

for all $f_1, f_2 \in L^{\infty}(\mu)$.

For a G-system $\mathbf{X} = (X, \mathcal{B}, \mu, (T_g)_{g \in G})$, if H is a subgroup of G, we denote by $\mathcal{I}(H)(\mathbf{X})$ the set of $A \in \mathcal{B}$ such that $T_g A = A$ for all $g \in H$. When there is no confusion, we write $\mathcal{I}(H)$.

For a \mathbb{Z}^d -system $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$, define

$$\mu_{H_1} = \mu \times_{\mathcal{I}(H_1)} \mu$$

and for k > 1, let

$$\mu_{H_1,\dots,H_k} = \mu_{H_1,\dots,H_{k-1}} \times_{\mathcal{I}(H_k^{[k-1]})} \mu_{H_1,\dots,H_{k-1}}$$

where $H_k^{[k-1]}$ denotes the subgroup of $(\mathbb{Z}^d)^{2^{k-1}}$ consisting of all the elements of the form $h_k \times \cdots \times h_k$ (2^{k-1} copies of h_k) for some $h_k \in H_k$. The *characteristic factor* $Z_{H_1,\ldots,H_k}(\mathbf{X})$ is defined to be the sub- σ -algebra of \mathcal{B} characterized by

$$\mathbb{E}(f|Z_{H_1,\dots,H_k}(\mathbf{X})) = 0 \text{ if and only if } |||f|||_{H_1,\dots,H_k}^{2^k} \coloneqq \int_{X^{[k]}} \bigotimes_{\epsilon \in \{0,1\}^k} \mathcal{C}^{|\epsilon|} f \, d\mu_{H_1,\dots,H_k} = 0$$

for all $f \in L^{\infty}(\mu)$, where $X^{[k]} = X \times \cdots \times X$ (2^k copies of X), $|\epsilon| = \epsilon_1 + \cdots + \epsilon_k$ for $\epsilon = (\epsilon_1, \ldots, \epsilon_k) \in \{0, 1\}^k$, and $\mathcal{C}^{2r+1}f = \overline{f}$, the complex conjugate of f, $\mathcal{C}^{2r}f = f$ for all $r \in \mathbb{Z}$. The quantity $|||f|||_{H_1,\ldots,H_k}$ denotes the *Host-Kra seminorm* of f with respect to the subgroups H_1, \ldots, H_k . Similar to the proof of [15, Lemma 4] (or [16, Lemma 4.3]), one can show that $Z_{H_1,\ldots,H_k}(\mathbf{X})$ is well defined.

We summarize some basic properties of the Host-Kra seminorms and their associated factors.

Proposition 3.1 (Lemma 2.4, [6]). Let $\mathbf{X} = (X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system, H_1, \ldots, H_k, H' be subgroups of \mathbb{Z}^d and $f \in L^{\infty}(\mu)$.

(i) For every permutation $\sigma: \{1, \ldots, k\} \to \{1, \ldots, k\}$, we have that

$$Z_{H_1,\ldots,H_k}(\mathbf{X}) = Z_{H_{\sigma(1)},\ldots,H_{\sigma(k)}}(\mathbf{X}),$$

hence the corresponding seminorm does not depend on the particular order taken for the subgroups H_1, \ldots, H_k .

- (*ii*) If $\mathcal{I}(H_j) = \mathcal{I}(H')$, then $Z_{H_1,\dots,H_j,\dots,H_k}(\mathbf{X}) = Z_{H_1,\dots,H_{j-1},H',H_{j+1},\dots,H_k}(\mathbf{X})$.
- (iii) For $k \geq 2$ we have that

$$|||f|||_{H_1,\ldots,H_k}^{2^k} = \mathbb{E}_{g \in H_k} |||f \cdot T_g \overline{f}|||_{H_1,\ldots,H_{k-1}}^{2^{k-1}},$$

while for k = 1,

$$|||f|||_{H_1}^2 = \mathbb{E}_{g \in H_1} \int_X f \cdot T_g \overline{f} \, d\mu.$$

(iv) Let $k \geq 2$. If $H' \leq H_j$ is of finite index, then

$$Z_{H_1,\dots,H_j,\dots,H_k}(\mathbf{X}) = Z_{H_1,\dots,H_{j-1},H',H_{j+1},\dots,H_k}(\mathbf{X}).$$

12 SEBASTIÁN DONOSO, ANDREU FERRÉ MORAGUES, ANDREAS KOUTSOGIANNIS AND WENBO SUN

- (v) If $H' \leq H_j$, then $Z_{H_1,...,H_j,...,H_k}(\mathbf{X}) \subseteq Z_{H_1,...,H_{j-1},H',H_{j+1},...,H_k}(\mathbf{X})$.
- (vi) For $k \ge 2$, $|||f|||_{H_1,...,H_{k-1}} \le |||f|||_{H_1,...,H_{k-1},H_k}$ and thus

$$Z_{H_1,\ldots,H_{k-1}}(\mathbf{X}) \subseteq Z_{H_1,\ldots,H_{k-1},H_k}(\mathbf{X}).$$

(vii) For $k \geq 1$, if H'_1, \ldots, H'_k are subgroups of \mathbb{Z}^d , then

$$Z_{H_1,\ldots,H_k}(\mathbf{X}) \vee Z_{H'_1,\ldots,H'_k}(\mathbf{X}) \subseteq Z_{H'_1,\ldots,H'_k,H_1,\ldots,H_k}(\mathbf{X}).$$

As an immediate corollary of Proposition 3.1 (iv), we have:

Corollary 3.2 (Corollary 2.5, [6]). Let H_1, \ldots, H_k be subgroups of \mathbb{Z}^d . If the H_i -action $(T_g)_{g \in H_i}$ is ergodic on \mathbf{X} for all $1 \leq i \leq k$, then $Z_{H_1,\ldots,H_k}(\mathbf{X}) = Z_{(\mathbb{Z}^d)^{\times k}}(\mathbf{X})$.

Convention 3.3. Thanks to Proposition 3.1 we may adopt a flexible and convenient notation while writing the Host-Kra characteristic factors. For example, if $A = \{H_1, H_2\}^{\times 3}$, then the notation $Z_{A,H_3,H_4^{\times 2},(H_i)_{i=5,6}}(\mathbf{X})$ refers to $Z_{H_1,H_1,H_1,H_2,H_2,H_3,H_4,H_4,H_5,H_6}(\mathbf{X})$.

Recall that for a subgroup $H \subseteq \mathbb{Z}^d$, $H^{[1]}$ denotes the subgroup $\{(h,h): h \in H\} \subseteq \mathbb{Z}^d \times \mathbb{Z}^d$.

Lemma 3.4. Let $d \in \mathbb{N}$. Let $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system and H_1, \ldots, H_k, H be subgroups of \mathbb{Z}^d . Let $f \in L^{\infty}(\mu)$. Then,

$$\|\|f \otimes \bar{f}\|\|_{H_1^{[1]},...,H_h^{[1]}} \le \|\|f\|\|_{H_1,...,H_k,H}^2,$$

where in the left hand side we consider the product space $(X \times X, \mathcal{B} \otimes \mathcal{B}, \mu \times \mu)$.

Proof. We proceed by induction on k. For k = 1, using the Cauchy-Schwartz inequality, we have

$$\begin{split} \|f \otimes \overline{f}\|_{H_{1}^{\left[1\right]}}^{2} &= \mathbb{E}_{g \in H_{1}} \int f \otimes \overline{f} \cdot (T_{g} \times T_{g}) \overline{f} \otimes f \ d(\mu \times \mu) \\ &= \mathbb{E}_{g \in H_{1}} \Big| \int T_{g} f \cdot \overline{f} d\mu \Big|^{2} = \mathbb{E}_{g \in H_{1}} \Big| \int \mathbb{E}(T_{g} f \cdot \overline{f} | \mathcal{I}(H)) d\mu \Big|^{2} \\ &\leq \mathbb{E}_{g \in H_{1}} \int |\mathbb{E}(T_{g} f \cdot \overline{f} | \mathcal{I}(H))|^{2} d\mu = \mathbb{E}_{g \in H_{1}} \|T_{g} f \cdot \overline{f}\|_{H}^{2} = \|f\|_{H,H_{1}}^{4} = \|f\|_{H_{1},H}^{4}, \end{split}$$

from where we conclude the required relation by taking square roots.

Suppose that the result holds for k-1. By Proposition 3.1 and the induction hypothesis,

$$\begin{split} \| f \otimes \overline{f} \|_{H_{1}^{[1]},...,H_{k}^{[1]}}^{2^{k}} &= \mathbb{E}_{g \in H_{k}} \| (T_{g} \times T_{g}) f \otimes \overline{f} \cdot \overline{f} \otimes f \|_{H_{1}^{[1]},...,H_{k-1}^{[1]}}^{2^{k-1}} \\ &= \mathbb{E}_{g \in H_{k}} \| T_{g} f \cdot \overline{f} \otimes T_{g} \overline{f} \cdot f \|_{H_{1}^{[1]},...,H_{k-1}^{[1]}}^{2^{k-1}} \\ &\leq \mathbb{E}_{g \in H_{k}} \| T_{g} f \cdot \overline{f} \|_{H_{1},...,H_{k-1},H}^{2^{k}} \\ &= \| f \|_{H_{1},...,H_{k-1},H,H_{k}} = \| f \|_{H_{1},...,H_{k-1},H_{k},H} \end{split}$$

and the claim follows.

3.2. Nilsystems, nilsequences and Structure Theorem. Let $X = N/\Gamma$, where N is a (k-step) nilpotent Lie group and Γ is a discrete cocompact subgroup of N. Let \mathcal{B} be the Borel σ -algebra of X, μ the Haar measure on X, and for $n \in \mathbb{Z}^d$, let $T_n \colon X \to X$ with $T_n x = b_n \cdot x$ for some group homomorphism $n \mapsto b_n$ from \mathbb{Z}^d to N. We say that $\mathbf{X} = (X, \mathcal{B}, \mu, (T_n)_{n \in \mathbb{Z}^d})$ is a $(k\text{-step}) \mathbb{Z}^d$ -nilsystem. For $k \geq 1$, we say that $(a_n)_{n \in \mathbb{Z}^d}$ is a $(k\text{-step}) \mathbb{Z}^d$ -nilsystem. For $k \geq 1$, we say that $(a_n)_{n \in \mathbb{Z}^d}$ is a $(k\text{-step}) \mathbb{Z}^d$ -nilsequence if there exist a $(k\text{-step}) \mathbb{Z}^d$ -nilsystem $\mathbf{X} = (X, \mathcal{B}, \mu, (T_n)_{n \in \mathbb{Z}^d})$, a function $F \in C(X)$ and $x \in X$ such that $a_n = F(T_n x)$ for all $n \in \mathbb{Z}^d$. For k = 0, a 0-step nilsequence is a constant sequence. An important reason which makes the Host-Kra characteristic factors powerful is their connection with nilsystems. The following is a slight generalization of [35, Theorem 3.7] (see [30, Theorem 3.7]), which is a higher dimensional version of Host-Kra structure theorem ([16]).

Theorem 3.5. Let **X** be an ergodic \mathbb{Z}^d -system. Then $Z_{(\mathbb{Z}^d)\times k}(\mathbf{X})$ is an inverse limit of (k-1)-step \mathbb{Z}^d -nilsystems.

3.3. **Bessel's inequality.** An essential difference in the study of multiple ergodic averages between \mathbb{Z} -systems and \mathbb{Z}^d -systems is that in the former case, one can usually bound the average by some Host-Kra seminorm of a function f appearing in the average, whereas in the latter, one can only bound the averages by an average of a family of Host-Kra seminorms of f. To overcome this difficulty, inspired by the work of Tao and Ziegler ([32]), in this subsection we derive an upper bound for expressions of the form $\overline{\mathbb{E}}_{i \in I} ||| f |||_{H_{i,1},\ldots,H_{i,s}}$, where I is a finite set and $H_{i,j}$ are subgroups of \mathbb{Z}^d .

The proof of the following statement is similar to [32, Corollary 1.22]:

Proposition 3.6 (Bessel's inequality). Let $s \in \mathbb{N}$, $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system, I be a finite set of indices, and $H_{i,j}, i \in I, 1 \leq j \leq s$ be subgroups of \mathbb{Z}^d . Then for all $f \in L^{\infty}(\mu)$,

$$\mathbb{E}_{i \in I} \|\mathbb{E}(f|Z_{H_{i,1},\dots,H_{i,s}})\|_{2}^{2} \leq \|f\|_{2} \cdot \left(\mathbb{E}_{i,j \in I} \left\|\mathbb{E}(f|Z_{\{H_{i,i'}+H_{j,j'}\}_{1 \leq i',j' \leq s}})\right\|_{2}^{2}\right)^{1/2}$$

Proof. For convenience, let $f_i \coloneqq \mathbb{E}(f|Z_{H_{i,1},\dots,H_{i,s}})$. Then

$$\mathbb{E}_{i \in I} \|\mathbb{E}(f|Z_{H_{i,1},\dots,H_{i,s}})\|_2^2 = \langle f, \mathbb{E}_{i \in I} f_i \rangle$$

which, by the Cauchy-Schwartz inequality is bounded by

$$\|f\|_2 \cdot \left|\mathbb{E}_{i,j\in I}\langle f_i, f_j\rangle\right|^{1/2}$$

By [32, Corollary 1.21], $L^{\infty}(Z_{H_{i,1},\dots,H_{i,s}})$ and $L^{\infty}(Z_{H_{j,1},\dots,H_{j,s}})$ are orthogonal on the orthogonal complement of $L^{\infty}(Z_{\{H_{i,i'}+H_{j,j'}\}_{1\leq i',j'\leq s}})$, hence

$$\langle f_i, f_j \rangle = \left\| \mathbb{E}(f | Z_{\{H_{i,i'} + H_{j,j'}\}_{1 \le i',j' \le s}}) \right\|_2^2$$

and we have the conclusion.

By repeatedly using Proposition 3.6, we have:

Corollary 3.7. Let $s, t \in \mathbb{N}$, $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system, I be a finite set of indices, and $H_{i,j}$, $i \in I$, $1 \leq j \leq s$, be subgroups of \mathbb{Z}^d . Then for all $f \in L^{\infty}(\mu)$ and $T \coloneqq 2^t$ we have

$$\left(\mathbb{E}_{i\in I} \|\mathbb{E}(f|Z_{H_{i,1},\dots,H_{i,s}})\|_2^2\right)^T \le \|f\|_2^{2T-2} \cdot \mathbb{E}_{i_1,\dots,i_T\in I} \left\|\mathbb{E}(f|Z_{\{\sum_{j=1}^T H_{i_j,i_j'}\}_{1\le i_1',\dots,i_T'\le s}})\right\|_2^2.$$

14 SEBASTIÁN DONOSO, ANDREU FERRÉ MORAGUES, ANDREAS KOUTSOGIANNIS AND WENBO SUN

The next proposition provides an upper bound for $\mathbb{E}_{i \in I} \| f \|_{H_{i,1},\ldots,H_{i,s}}$ which can be combined with the previous two statements.

Proposition 3.8. Let $s, t \in \mathbb{N}$, $(X, \mathcal{B}, \mu, (T_g)_{q \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system, I be a finite set of indices, and $H_{i,j}, i \in I, 1 \leq j \leq s$ be subgroups of \mathbb{Z}^d . Then for all $f \in L^{\infty}(\mu)$, with $||f||_{L^{\infty}(\mu)} \leq 1$,

$$\mathbb{E}_{i \in I} \| f \|_{H_{i,1},\dots,H_{i,s}} \le (\mathbb{E}_{i \in I} \| (\mathbb{E}(f | Z_{H_{i,1},\dots,H_{i,s}}) \|_2^2)^{1/2^s}$$

Proof. Note that

(11)
$$|||f|||_{H_{i,1},\dots,H_{i,s}} \le ||f||_{L^{2^s}(\mu)} \le ||f||_2^{1/2^{s-1}}.$$

Also, for all i we have

$$\begin{split} \|\|f\|\|_{H_{i,1},\dots,H_{i,s}} &\leq \|\|f - \mathbb{E}(f|Z_{H_{i,1},\dots,H_{i,s}})\|\|_{H_{i,1},\dots,H_{i,s}} + \|\|\mathbb{E}(f|Z_{H_{i,1},\dots,H_{i,s}})\|\|_{H_{i,1},\dots,H_{i,s}} \\ &= \|\|\mathbb{E}(f|Z_{H_{i,1},\dots,H_{i,s}})\|\|_{H_{i,1},\dots,H_{i,s}}, \end{split}$$

 \mathbf{SO}

(12)
$$\begin{split} \mathbb{E}_{i\in I} \|\|f\|\|_{H_{i,1},\dots,H_{i,s}} &\leq \mathbb{E}_{i\in I} \|\|\mathbb{E}(f|Z_{H_{i,1},\dots,H_{i,s}})\|\|_{H_{i,1},\dots,H_{i,s}}) \|_{2}^{H_{i,1},\dots,H_{i,s}} \\ &\leq \mathbb{E}_{i\in I} \|\|\mathbb{E}(f|Z_{H_{i,1},\dots,H_{i,s}})\|_{2}^{1/2^{s-1}} \leq (\mathbb{E}_{i\in I}\|(\mathbb{E}(f|Z_{H_{i,1},\dots,H_{i,s}})\|_{2}^{2})^{1/2^{s}}, \\ \text{as was to be shown.} \end{split}$$

as was to be shown.

3.4. General properties of subgroups of \mathbb{Z}^d and properties of polynomials. Recall that for a subset A of \mathbb{Q}^d , we denote $G(A) := \operatorname{span}_{\mathbb{Q}} \{a \in A\} \cap \mathbb{Z}^d$. Next we summarize some properties of these sets.

Lemma 3.9. The following properties hold:

- (i) For any set $A \subseteq \mathbb{Z}^d$, G(A) is a subgroup of \mathbb{Z}^d .
- (ii) Let A be a finite set and M(A) the matrix whose columns are the elements of A. Then $G(A) = (M(A) \cdot \mathbb{Q}^{|A|}) \cap \mathbb{Z}^{d}.$
- (*iii*) If $H \subseteq \mathbb{Z}^d$ is the subgroup generated by $h_1, \ldots, h_k \in \mathbb{Z}^d$, then $G(H) = G(\{h_1, \ldots, h_k\})$. In particular, letting $M(h_1, \ldots, h_k)$ be the matrix whose columns are h_1, \ldots, h_k , we have that $G(\langle h_1, \ldots, h_k \rangle) = (M(h_1, \ldots, h_k) \cdot \mathbb{Q}^k) \cap \mathbb{Z}^d$.
- (iv) For any subgroup $H \subseteq \mathbb{Z}^d$, H has finite index in G(H). Moreover, G(H) is the largest subgroup of \mathbb{Z}^d which is a finite index extension of H.
- (v) If not all of a_1, \ldots, a_k belong to a common proper subspace of \mathbb{Q}^d , then $G(\{a_1, \ldots, a_k\})$ $=\mathbb{Z}^d.$

Proof. Properties (i), (ii) and (iii) follow directly from the definitions.

To prove (iv), let $\{g_1, \ldots, g_k\}$ be a set such that $\langle g_1, \ldots, g_k \rangle = G(H)$. For each $i = 1, \ldots, k$ there exist m_i and $h_i \in H$ such that $g_i = \frac{h_i}{m_i}$. The group $\langle m_1 g_1, \ldots, m_k g_k \rangle$ is of finite index in $\langle g_1, \ldots, g_k \rangle = G(H)$ and is contained in H. Therefore H is of finite index in G(H).

To see that G(H) is the largest finite index extension of H, take H' to be any finite index extension of H and take $h' \in H'$. Since H' is a finite index extension of H, we have that there exists $n \in \mathbb{N}$ such that $nh' \in H$. This implies that $h' \in G(H)$.

To show (v), reordering a_1, \ldots, a_k if needed, we may assume that a_1, \ldots, a_d are linearly independent vectors over \mathbb{Q} . It follows that $\operatorname{span}_{\mathbb{Q}}(\{a_1,\ldots,a_d\}) = \mathbb{Q}^d$ and then $G(\{a_1,\ldots,a_k\}) \supseteq$ $G(\{a_1,\ldots,a_d\}) = \mathbb{Z}^d.$ **Remark 3.10.** If H_1 and H_2 are subgroups of \mathbb{Z}^d , then $G(H_1) + G(H_2) \subseteq G(H_1 + H_2)$, with the inclusion possibly being strict. For instance, for $H_1 = \langle (1,2) \rangle$, $H_2 = \langle (2,1) \rangle$ we have that $G(H_1) = H_1$, $G(H_2) = H_2$ and $H_1 + H_2 \subsetneq G(H_1 + H_2) = \mathbb{Z}^2$. Nevertheless, Lemma 3.9 implies that that $G(H_1) + G(H_2)$ has finite index in $G(H_1 + H_2)$.

In the remainder of the section, we provide some algebraic lemmas that will be used later in the paper. For a set $E \subseteq \mathbb{Z}^d$, we define its upper Banach density (or just upper density when there is no confusion) with $d^*(E) := \overline{\lim}_{N \to \infty} \max_{t \in \mathbb{Z}^d} \frac{|(E-t) \cap \{1, \dots, N\}^d|}{N^d}$. If the limit exists, we say that its value is the Banach density (or just density) of E. The proof of the following lemma is routine (see also [6, Lemma 2.12] for a more general version):

Lemma 3.11 (Lemma 2.12, [6]). Let $\mathbf{c}: (\mathbb{Z}^L)^s \to \mathbb{R}$ be a polynomial. Then either $\mathbf{c} \equiv 0$ or the set of $\mathbf{h} \in (\mathbb{Z}^L)^s$ such that $\mathbf{c}(\mathbf{h}) = 0$ is of (upper) Banach density 0.

Lemma 3.12. Let $v_i \in \mathbb{Z}^L$, $1 \leq i \leq k$ and U be a subset of \mathbb{Z}^k of positive density. Then

(13)
$$G\left(\left\{\sum_{1\leq i\leq k}h_iv_i\colon \mathbf{h}=(h_1,\ldots,h_k)\in U\right\}\right)=G(\{v_i\colon 1\leq i\leq k\})$$

Proof. Note that in (13) the right hand side clearly includes the left hand side. To prove the converse inclusion it suffices to show that

(14)
$$span_{\mathbb{Q}}\{\mathbf{h}:\mathbf{h}\in U\}=\mathbb{Q}^{k}.$$

Since U has positive density, it cannot be contained in any hyperplane of \mathbb{Q}^k , so it must have at least k elements that are linearly independent over \mathbb{Q} . Thus, (14) follows immediately. \Box

Definition 3.13. Let $P: \mathbb{Z}^{LK} \to \mathbb{R}$ be a polynomial. Denote by $\Delta P: \mathbb{Z}^{L(K+1)} \to \mathbb{R}$ the polynomial given by $\Delta P(n, h_1, \ldots, h_K) \coloneqq P(n + h_K, h_1, \ldots, h_{K-1}) - P(n, h_1, \ldots, h_{K-1})$ for all $n, h_1, \ldots, h_K \in \mathbb{Z}^L$. For a polynomial $P: \mathbb{Z}^L \to \mathbb{R}$, let $\Delta^K P \coloneqq \Delta \cdot \ldots \cdot \Delta P$ (where Δ acts K times).

Lemma 3.14. Let $K \in \mathbb{N}$ and $Q: \mathbb{Z}^L \to \mathbb{R}$ be a homogeneous polynomial with $\deg(Q) > K$. If $Q(n) \notin \mathbb{Q}[n] + \mathbb{R}$, then the set of $(h_1, \ldots, h_K) \in (\mathbb{Z}^L)^K$ such that $\Delta^K Q(\cdot, h_1, \ldots, h_K) \notin \mathbb{Q}[n] + \mathbb{R}$ is of density 1.

Proof. We may write $Q(n) = \sum_{i=1}^{M} a_i Q_i(n)$ for some $M \in \mathbb{N}$, homogeneous polynomials Q_1, \ldots, Q_M in $\mathbb{Q}[n]$ of degrees $\deg(Q)$, and real numbers $a_1, \ldots, a_M \in \mathbb{R}$ which are linearly independent over \mathbb{Q} (this can be done by taking $a_1 \ldots, a_M$ to be a basis of the \mathbb{Q} -span of the coefficients of Q). Since $Q(n) \notin \mathbb{Q}[n] + \mathbb{R}$, there exists some $1 \leq i \leq M$ such that $a_i \notin \mathbb{Q}$ and $Q_i \neq 0$. Without loss of generality assume that i = 1. Since $\deg(Q_1) > K$, we have that $\Delta^K Q_1 \neq 0$.

Suppose that $\Delta^{K}Q(\cdot, h_{1}, \ldots, h_{K}) \in \mathbb{Q}[n] + \mathbb{R}$ for some $(h_{1}, \ldots, h_{K}) \in (\mathbb{Z}^{L})^{K}$. Note that $\Delta^{K}Q(\cdot, h_{1}, \ldots, h_{K}) = \sum_{i=1}^{M} a_{i}\Delta^{K}Q_{i}(\cdot, h_{1}, \ldots, h_{K})$. Since each $\Delta^{K}Q_{i}(\cdot, h_{1}, \ldots, h_{K})$ is a rational polynomial of degree deg(Q) - K and $a_{1}, \ldots, a_{M} \in \mathbb{R}$ are linearly independent over \mathbb{Q} , we must have that $\Delta^{K}Q_{1}(\cdot, h_{1}, \ldots, h_{K}) \equiv 0$. So if the set of $(h_{1}, \ldots, h_{K}) \in (\mathbb{Z}^{L})^{K}$ such that $\Delta^{K}Q(\cdot, h_{1}, \ldots, h_{K}) \in \mathbb{Q}[n] + \mathbb{R}$ has positive density, then the set of $(n, h_{1}, \ldots, h_{K}) \in (\mathbb{Z}^{L})^{K}$ such that $\Delta^{K}Q_{1}(n, h_{1}, \ldots, h_{K}) = 0$ has positive density too. By [6, Lemma 2.12], $\Delta^{K}Q_{1} \equiv 0$, a contradiction. This finishes the proof.

4. PET INDUCTION

In this section we present the method we use to reduce the complexity of the polynomial iterates, i.e., PET induction,¹⁸ which was first introduced in [2]. To this end, we start by recalling a variation of van der Corput's lemma from [6] that is convenient for our study. We then continue by presenting the inductive scheme via the use of van der Corput operations.

4.1. The van der Corput lemma. The standard tool used in reducing the complexity of polynomial families of iterates is van der Corput's lemma (also known as "van der Corput's trick"). We will use the following variation of it, the proof of which can be found in [6, Lemma 2.2]:

Lemma 4.1 (van der Corput lemma). Let $(a(n; h_1, \ldots, h_s))_{(n;h_1,\ldots,h_s)\in(\mathbb{Z}^L)^{s+1}}$ be a bounded sequence by 1 in a Hilbert space \mathcal{H} .¹⁹ Then, for $\tau \in \mathbb{N}_0$,

$$\overline{\mathbb{E}}_{h_1,\dots,h_s\in\mathbb{Z}^L}^{\square} \sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ F \not olner \ seq.}} \overline{\lim_{N\to\infty}} \|\mathbb{E}_{n\in I_N}a(n;h_1,\dots,h_s)\|^{2\tau} \\
\leq 4^{\tau}\overline{\mathbb{E}}_{h_1,\dots,h_s,h_{s+1}\in\mathbb{Z}^L}^{\square} \sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ F \not olner \ seq.}} \overline{\lim_{N\to\infty}} |\mathbb{E}_{n\in I_N}\langle a(n+h_{s+1};h_1,\dots,h_s),a(n;h_1,\dots,h_s)\rangle|^{\tau}.$$

We also provide two applications of Lemma 4.1 for later use. The first one is to get an upper bound for single averages with polynomial iterates and a polynomial exponential weight. Let $\exp(x) \coloneqq e^{2\pi i x}$ and recall Definition 3.13 for the polynomial $\Delta^{K} P$.

Lemma 4.2. Let $P: \mathbb{Z}^L \to \mathbb{R}$ and $p: \mathbb{Z}^L \to \mathbb{Z}^d$ be polynomials. Let $\mathbf{X} = (X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system and $f \in L^{\infty}(\mu)$ be a function bounded by 1. For all $K \in \mathbb{N}_0$ and $\tau > 0$, there exists a universal constant $C_{K,\tau} > 0$ such that

$$\sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ F \notin lner \ seq.}} \frac{\overline{\lim}_{N\to\infty}}{\sum_{\substack{N\to\infty}}} \|\mathbb{E}_{n\in I_N} \exp(P(n))T_{p(n)}f\|_2^{2\tau}$$

$$\leq C_{K,\tau} \mathbb{E}^{\square}_{\mathbf{h}=(h_1,\dots,h_K)\in(\mathbb{Z}^L)^K} \sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ F \notin lner \ seq}} \frac{\overline{\lim}_{N\to\infty}}{\sum_{\substack{N\to\infty}}} \|\mathbb{E}_{n\in I_N} \exp(\Delta^K P(n,\mathbf{h}))T_{\Delta^K p(n,\mathbf{h})}f\|_2^{\tau}.$$

Proof. When K = 0, there is nothing to prove. We now assume that the relation holds for some $K \in \mathbb{N}_0$ and we show it for K + 1. Using Lemma 4.1 and the *T*-invariance of μ , we get

$$\begin{split} \overline{\mathbb{E}}_{\mathbf{h}=(h_{1},\dots,h_{K})\in(\mathbb{Z}^{L})^{K}}^{\Box} \sup_{\substack{(I_{N})_{N\in\mathbb{N}}\\ \text{Følner seq.}}} \overline{\lim_{N\to\infty}} \left\| \mathbb{E}_{n\in I_{N}}\exp(\Delta^{K}P(n,\mathbf{h}))T_{\Delta^{K}p(n,\mathbf{h})}f \right\|_{2}^{2\tau} \\ &\leq 4^{\tau}\overline{\mathbb{E}}_{\mathbf{h}=(h_{1},\dots,h_{K+1})\in(\mathbb{Z}^{L})^{K+1}}^{\Box} \sup_{\substack{(I_{N})_{N\in\mathbb{N}}\\ \text{Følner seq.}}} \overline{\lim_{N\to\infty}} \left\| \mathbb{E}_{n\in I_{N}}\int_{X}\exp(\Delta^{K+1}P(n,\mathbf{h}))T_{\Delta^{K+1}p(n,\mathbf{h})}f\cdot\overline{f}\,d\mu \right|^{\tau} \\ &\leq 4^{\tau}\overline{\mathbb{E}}_{\mathbf{h}=(h_{1},\dots,h_{K+1})\in(\mathbb{Z}^{L})^{K+1}}^{\Box} \sup_{\substack{(I_{N})_{N\in\mathbb{N}}\\ \text{Følner seq.}}} \overline{\lim_{N\to\infty}} \left\| \mathbb{E}_{n\in I_{N}}\exp(\Delta^{K+1}P(n,\mathbf{h}))T_{\Delta^{K+1}p(n,\mathbf{h})}f \right\|_{2}^{\tau}, \end{split}$$

 $^{^{18}\}mathrm{PET}$ is an abbreviation for "Polynomial Exhaustion Technique."

¹⁹We use this unorthodox notation to separate the variable n from the h_i 's. The variable n plays a different, compared to the h_i 's, role in our study.

hence the result (the constant that appears depends only on τ and K).

The second application of Lemma 4.1 provides an upper bound for single averages, with linear iterates and an exponential weight evaluated at a linear polynomial, on a product system. The proof is inspired by [6, Lemma 5.2] and [17, Proposition 2.9].

Lemma 4.3. Let (X, \mathcal{B}, μ) be a probability space, $k, L \in \mathbb{N}$ and $T_{i,j}, 1 \leq i \leq k, 1 \leq j \leq L$ be commuting measure preserving transformations on X. Denote $S_j = T_{1,j} \times \cdots \times T_{k,j}$ for all $1 \leq j \leq L$. Let G_i be the group generated by $T_{i,1}, \ldots, T_{i,L}$. Then for any polynomial $P: \mathbb{Z}^L \to \mathbb{R}$ of degree 1 and $f_1, \ldots, f_k \in L^{\infty}(\mu)$ bounded by 1, we have that

(15)
$$\sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ F \notin lner \ seq.}} \overline{\lim_{N\to\infty}} \|\mathbb{E}_{n\in I_N} \exp(P(n))R_n f\|_{L^2(\mu^{\otimes k})} \le 2\min_{1\le i\le k} \|f_i\|_{G_i^{\times 2}},$$

where, $f = f_1 \otimes \cdots \otimes f_k$ and for $n = (n_1, \ldots, n_L)$, $R_n \coloneqq S_1^{n_1} \cdot \ldots \cdot S_L^{n_L}$.

Proof. Fix $1 \leq i \leq k$ and let $P(n) = a \cdot n + b$ for some $a \in \mathbb{R}^L, b \in \mathbb{R}$. Then, by Lemma 4.1, the 4th power of the left hand side of (15) is bounded by

$$\begin{split} &16 \cdot \mathbb{E}_{h \in \mathbb{Z}^{L}}^{\Box} \sup_{\substack{(I_{N})_{N \in \mathbb{N}} \\ \text{Følner seq.}}} \overline{\lim_{N \to \infty}} \left| \int_{X} \mathbb{E}_{n \in I_{N}} \exp(P(n+h) - P(n)) R_{n+h} f \cdot R_{n} \overline{f} \, d\mu^{\otimes k} \right|^{2} \\ &= 16 \cdot \mathbb{E}_{h \in \mathbb{Z}^{L}}^{\Box} \sup_{\substack{(I_{N})_{N \in \mathbb{N}} \\ \text{Følner seq.}}} \overline{\lim_{N \to \infty}} \left| \int_{X} \mathbb{E}_{n \in I_{N}} \exp(a \cdot h) R_{h} f \cdot \overline{f} \, d\mu^{\otimes k} \right|^{2} \\ &= 16 \cdot \mathbb{E}_{h \in \mathbb{Z}^{L}}^{\Box} \left| \int_{X} R_{h} f \cdot \overline{f} \, d\mu^{\otimes k} \right|^{2} \leq 16 \cdot \mathbb{E}_{h=(h_{1},\dots,h_{L}) \in \mathbb{Z}^{L}}^{\Box} \left| \int_{X} S_{i}^{h_{i}} f_{i} \cdot \overline{f}_{i} \, d\mu \right|^{2} \\ &\leq 16 \cdot \mathbb{E}_{h=(h_{1},\dots,h_{L}) \in \mathbb{Z}^{L}}^{\Box} \left| \int_{X} \mathbb{E}(S_{i}^{h_{i}} f_{i} \cdot \overline{f}_{i} | \mathcal{I}(G_{i})) \, d\mu \right|^{2} \\ &= 16 \cdot \mathbb{E}_{h_{i} \in \mathbb{Z}}^{\Box} \left| \int_{X} \mathbb{E}(S_{i}^{h_{i}} f_{i} \cdot \overline{f}_{i} | \mathcal{I}(G_{i})) \, d\mu \right|^{2} \end{split}$$

from where the result follows.

4.2. The van der Corput operation. To review the PET induction scheme, we will follow, and slightly modify, the approach from [6]. To this end, we extend the definitions that we have already given on the polynomial families of interest (see the beginning of Subsection 2.1), taking into account that we treat the first L-tuple of variables of the polynomials differently.

Definition 4.4. For a polynomial $p(n; h_1, \ldots, h_s)$: $(\mathbb{Z}^L)^{s+1} \to \mathbb{Z}$, we denote with deg(p) the degree of p with respect to n (for example, for s = 1, L = 2, the degree of $p(n_1, n_2; h_{1,1}, h_{1,2}) = h_{1,1}h_{1,2}n_1^2 + h_{1,1}^5n_2$, is 2).

For a polynomial $p(n; h_1, \ldots, h_s) = (p_1(n; h_1, \ldots, h_s), \ldots, p_d(n; h_1, \ldots, h_s)) \colon (\mathbb{Z}^L)^{s+1} \to \mathbb{Z}^d$, we let $\deg(p) = \max_{1 \leq i \leq d} \deg(p_i)$ and we say that p is *non-constant* if $\deg(p) > 0$ (i.e., some p_i is a non-constant function of n). The polynomials $q_1, \ldots, q_k \colon (\mathbb{Z}^L)^{s+1} \to \mathbb{Z}^d$ are called

17

essentially distinct if they are non-constant and $q_i - q_j$ is non-constant for all $i \neq j$. Finally, for a tuple $\mathbf{q} = (q_1, \ldots, q_k)$, we let $\deg(\mathbf{q}) = \max_{1 \leq i \leq k} \deg(q_i)$.²⁰

Let $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system, $q_1, \ldots, q_k \colon (\mathbb{Z}^L)^{s+1} \to \mathbb{Z}^d$ be polynomials and $g_1, \ldots, g_k \colon X \times (\mathbb{Z}^L)^s \to \mathbb{R}$ be functions such that each $g_m(\cdot; h_1, \ldots, h_s)$ is an $L^{\infty}(\mu)$ function bounded by 1 for all $h_1, \ldots, h_s \in \mathbb{Z}^L, 1 \leq m \leq k$. If $\mathbf{q} = (q_1, \ldots, q_k)$ and $\mathbf{g} = (g_1, \ldots, g_k)$, we say that $A = (L, s, k, \mathbf{g}, \mathbf{q})$ is a *PET-tuple*, and for $\tau \in \mathbb{N}_0$ we set

$$S(A,\tau) \coloneqq \overline{\mathbb{E}}_{h_1,\dots,h_s \in \mathbb{Z}^L}^{\square} \sup_{\substack{(I_N)_N \in \mathbb{N} \\ \text{Følner seq.}}} \overline{\lim_{N \to \infty}} \left\| \mathbb{E}_{n \in I_N} \prod_{m=1}^{\kappa} T_{q_m(n;h_1,\dots,h_s)} g_m(x;h_1,\dots,h_s) \right\|_2^{\tau}.$$

We define deg(A) = deg(q), and say that A is non-degenerate if q is a family of essentially distinct polynomials (for convenience, q will be called non-degenerate as well). For $1 \le m \le k$, the tuple A is m-standard for $f \in L^{\infty}(\mu)$ if deg(A) = deg(q_m) and $g_m(x; h_1, \ldots, h_s) = f(x)$ for every x, h_1, \ldots, h_s . That is, f is the m-th function in g, only depending on the first variable, and the polynomial q_m that acts on f is of the highest degree.²¹ The tuple A will be called semi-standard for f if there exists $1 \le m \le k$ such that $g_m(x; h_1, \ldots, h_s) = f(x)$ for every x, h_1, \ldots, h_s . In this case we do not require the function f to have a specific position in g nor that the polynomial acting on f to be of the highest degree.

For each non-degenerate PET-tuple $A = (L, s, k, \mathbf{g}, \mathbf{q})$ and polynomial $q: (\mathbb{Z}^L)^{s+1} \to \mathbb{Z}^d$, we define the *vdC-operation*, $\partial_q A$, according to the following three steps:²²

Step 1: For all $1 \leq m \leq k$, let $g_m^* = g_{m+k}^* = g_m$, and $q_1^*, \ldots, q_{2k}^* \colon (\mathbb{Z}^L)^{s+2} \to \mathbb{Z}^d$ be the polynomials defined as

$$q_m^*(n;h_1,\ldots,h_{s+1}) = \begin{cases} q_m(n+h_{s+1};h_1,\ldots,h_s) - q(n;h_1,\ldots,h_s) & ,1 \le m \le k \\ q_{m-k}(n;h_1,\ldots,h_s) - q(n;h_1,\ldots,h_s) & ,k+1 \le m \le 2k \end{cases},$$

i.e., we subtract the polynomial q from the first k polynomials after we have shifted by h_{s+1} the first L variables, and for the second k ones we subtract q^{23} Denote $\mathbf{q}^* = (q_1^*, \ldots, q_{2k}^*)$.

Step 2: We remove from $q_1^*(n; h_1, \ldots, h_{s+1}), \ldots, q_{2k}^*(n; h_1, \ldots, h_{s+1})$ the polynomials which are constant and the corresponding terms with these as iterates (this will be justified via the use of the Cauchy-Schwarz inequality and the fact that the functions g_m are bounded), and then put the non-constant ones in groups $J_i = \{\tilde{q}_{i,1}, \ldots, \tilde{q}_{i,t_i}\}, 1 \leq i \leq k'$ for some k', $t_i \in \mathbb{N}$ such that two polynomials are essentially distinct if and only if they belong to different groups.²⁴ Next, we write $\tilde{q}_{i,j}(n; h_1, \ldots, h_{s+1}) = \tilde{q}_{i,1}(n; h_1, \ldots, h_{s+1}) + \tilde{p}_{i,j}(h_1, \ldots, h_{s+1})$ for some polynomial $\tilde{p}_{i,j}$ for all $1 \leq j \leq t_i, 1 \leq i \leq k'$. For convenience, we also relabel g_1^*, \ldots, g_{2k}^* accordingly as $\tilde{g}_{i,j}$ for all $1 \leq j \leq t_i, 1 \leq i \leq k'$.

 $^{^{20}}$ For clarity, we use non-bold letters for vectors (of polynomials) and bold letters for vectors of vectors (of polynomials).

²¹Here, we say *m*-standard for f to highlight the function of interest as, after running the vdC-operation, the position of the functions in the expression we deal with changes.

 $^{^{22}}$ Actually, the vdC-operation can be defined for any PET-tuple, not just for non-degenerate ones. Similarly, being a procedure that reduces complexity, PET induction can be applied to any family of polynomials. As the expressions of interest in this paper correspond to non-degenerate tuples, we consider only this case.

²³In practice, this q will be one of the q_i 's of minimum degree.

 $^{^{24}}$ After removing the constant polynomials, the terms from A that are grouped are of degree 1.

Step 3: For all $1 \le i \le k'$, let $q'_i = \tilde{q}_{i,1}$ and

$$g'_i(x;h_1,\ldots,h_{s+1}) = \tilde{g}_{i,1}(x;h_1,\ldots,h_{s+1}) \prod_{j=2}^{t_i} T_{\tilde{p}_{i,j}(h_1,\ldots,h_{s+1})} \tilde{g}_{i,j}(x;h_1,\ldots,h_{s+1})$$

We set $\mathbf{q}' = (q'_1, \ldots, q'_{k'})$, $\mathbf{g}' = (g'_1, \ldots, g'_{k'})$ and we denote the new PET-tuple by $\partial_q A := (L, s+1, k', \mathbf{g}', \mathbf{q}')$.²⁵ It follows from the construction that $\partial_q A$ is non-degenerate.

If $q = q_t$ for some $1 \le t \le \ell$, we write $\partial_t \mathbf{q}$ instead of \mathbf{q}' to highlight the fact that we have subtracted the polynomial q_t ; we also write $\partial_t A$ instead of $\partial_{q_t} A$ to lighten the notation.

We say that the operation $A \to \partial_t A$ is 1-inherited if $q'_1 = q^*_1$ and $g'_1 = f_1$, i.e., if we did not drop q^*_1 or group it with any other q^*_i in Step 2.

Example 7. Let $\mathbf{p} = (p_1, p_2)$ with $p_1, p_2 \colon \mathbb{Z} \to \mathbb{Z}^d$ be polynomials given by $p_i(n) = b_{i,2}n^2 + b_{i,1}n$ for some $b_{i,1}, b_{i,2} \in \mathbb{Z}^d$ for $1 \leq i \leq 2$ with $b_{1,2}, b_{2,2}, b_{1,2} - b_{2,2} \neq \mathbf{0}$ (hence, we have that L = s = 1 and k = 2). Subtracting p_2 in the Step 1 of the vdC operation, we have that $\partial_2 \mathbf{p} = (q_1, q_2, q_3)$, is a tuple of 3 polynomials, $q_1, q_2, q_3 \colon \mathbb{Z}^2 \to \mathbb{Z}^d$, given by

$$q_1(n, h_1) = (b_{1,2} - b_{2,2})n^2 + 2b_{1,2}nh_1 + (b_{1,1} - b_{2,1})n + b_{1,1}h_1 + b_{1,2}h_1^2,$$

$$q_2(n, h_1) = 2b_{2,2}nh_1 + b_{2,1}h_1 + b_{2,2}h_1^2,$$

$$q_3(n, h_1) = (b_{1,2} - b_{2,2})n^2 + (b_{1,1} - b_{2,1})n,$$

where we removed 1 essentially constant polynomial in Step 2 of the vdC operation.²⁶ Actually, after using a series of vdC-operations, one can convert \mathbf{p} into a PET-tuple of linear polynomials.

Indeed, if we run the vdC-operation once more by subtracting q_2 in the Step 1 of the vdC operation, we have that $\partial_2 \partial_2 \mathbf{p} = (q'_1, \ldots, q'_4)$ is a tuple of 4 polynomials, $q'_1, \ldots, q'_4 \colon \mathbb{Z}^3 \to \mathbb{Z}^d$, given by

$$\begin{aligned} q_1'(n,h_1,h_2) &= (b_{1,2}-b_{2,2})n^2 + 2(b_{1,2}-b_{2,2})nh_1 + 2(b_{1,2}-b_{2,2})nh_2 + (b_{1,1}-b_{2,1})n + r_1'(h_1,h_2), \\ q_2'(n,h_1,h_2) &= (b_{1,2}-b_{2,2})n^2 - 2b_{2,2}nh_1 + 2(b_{1,2}-b_{2,2})nh_2 + (b_{1,1}-b_{2,1})n + r_2'(h_1,h_2), \\ q_3'(n,h_1,h_2) &= (b_{1,2}-b_{2,2})n^2 + 2(b_{1,2}-b_{2,2})nh_1 + (b_{1,1}-b_{2,1})n + r_3'(h_1,h_2), \\ q_4'(n,h_1,h_2) &= (b_{1,2}-b_{2,2})n^2 - 2b_{2,2}nh_1 + (b_{1,1}-b_{2,1})n + r_4'(h_1,h_2), \end{aligned}$$

where $r'_i: \mathbb{Z}^2 \to \mathbb{Z}^d, 1 \leq i \leq 4$, are polynomials in h_1, h_2 , and we removed 2 essentially constant polynomials (i.e. $q_2(n, h_1) - q_2(n, h_1)$ and $q_2(n + h_2, h_1) - q_2(n, h_1)$) in Step 2 of the vdC operation.

²⁵Here we abuse the notation by writing $\partial_q A$ to denote any such tuple, obtained from Step 1 to 3. Strictly speaking, $\partial_q A$ is not uniquely defined as the order of the grouping of q'_1, \ldots, q'_{2k} in Step 2 is ambiguous. However, this is done without loss of generality, since the order does not affect the value of $S(\partial_q A, \cdot)$.

²⁶Here we use q_i 's instead of p'_i 's in the first step to ease the notation of Example 9 that is given in the next section.

Finally, if we apply vdC-operation again by subtracting q'_4 in Step 1 of the vdC operation, we have that $\partial_4 \partial_2 \partial_2 \mathbf{p} = (q''_1, \ldots, q''_7)$ is a tuple of 7 polynomials, $q''_1, \ldots, q''_7 \colon \mathbb{Z}^4 \to \mathbb{Z}^d$, given by

$$\begin{split} q_1''(n,h_1,h_2,h_3) &= 2b_{1,2}nh_1 + 2(b_{1,2} - b_{2,2})nh_2 + 2(b_{1,2} - b_{2,2})nh_3 + r_1''(h_1,h_2,h_3), \\ q_2''(n,h_1,h_2,h_3) &= 2(b_{1,2} - b_{2,2})nh_2 + 2(b_{1,2} - b_{2,2})nh_3 + r_2''(h_1,h_2,h_3), \\ q_3''(n,h_1,h_2,h_3) &= 2b_{1,2}nh_1 + 2(b_{1,2} - b_{2,2})nh_3 + r_3''(h_1,h_2,h_3), \\ q_4''(n,h_1,h_2,h_3) &= 2(b_{1,2} - b_{2,2})nh_3 + r_4''(h_1,h_2,h_3), \\ q_5''(n,h_1,h_2,h_3) &= 2b_{1,2}nh_1 + 2(b_{1,2} - b_{2,2})nh_2 + r_5''(h_1,h_2,h_3), \\ q_6''(n,h_1,h_2,h_3) &= 2(b_{1,2} - b_{2,2})nh_2 + r_6''(h_1,h_2,h_3), \\ q_7''(n,h_1,h_2,h_3) &= 2b_{1,2}nh_1 + r_7''(h_1,h_2,h_3), \end{split}$$

where $r''_i: \mathbb{Z}^3 \to \mathbb{Z}^d$, $1 \leq i \leq 7$, are polynomials in h_1, h_2, h_3 , and we removed 1 essentially constant polynomial in the Step 2 of the vdC operation. It is clear that $\deg(\partial_4 \partial_2 \partial_2 \mathbf{p}) = 1$.

The vdC-operation provides us with a non-degenerate tuple, the value $S(\cdot, \cdot)$ of which satisfies the following:

Proposition 4.5 (Proposition 4.1, [6]). Let $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system, $A = (L, s, k, \mathbf{g}, \mathbf{q})$ a PET-tuple, and $q: (\mathbb{Z}^L)^{s+1} \to \mathbb{Z}^d$ a polynomial. Then $\partial_q A$ is non-degenerate and $S(A, 2\tau) \leq 4^{\tau} S(\partial_q A, \tau)$ for every $\tau \in \mathbb{N}_0$.

The following crucial result (cf. [6, Theorem 4.2]) shows that when we start with a PETtuple which is 1-standard for a function, then, after finitely many vdC-operations, we arrive at a new PET-tuple of degree 1 which is still 1-standard for the same function, so we can then use some Host-Kra seminorm to bound the lim sup of the average of interest.

Theorem 4.6. Let $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system and $f \in L^{\infty}(\mu)$. Let $A = (L, s, k, \mathbf{g}, \mathbf{q})$ be a non-degenerate PET-tuple which is 1-standard for g_1 . Then, there exist $\rho_1, \ldots, \rho_t \in \mathbb{N}$, for some $t \in \mathbb{N}_0$, such that for all $1 \leq t' \leq t$, $\partial_{\rho_{t'}} \ldots \partial_{\rho_1} A$ is a non-degenerate PET-tuple which is still 1-standard for g_1 , and that $\partial_{\rho_{t'-1}} \ldots \partial_{\rho_1} A \to \partial_{\rho_{t'}} \ldots \partial_{\rho_1} A$ is 1-inherited. Moreover, $\deg(\partial_{\rho_t} \ldots \partial_{\rho_1} A) = 1.^{27}$

If A is an *m*-standard for the function f PET-tuple then, by rearranging the terms if necessary, one can get a new tuple A' which is 1-standard for f with $S(A,\tau) = S(A',\tau)$. However, if A is semi-standard but not standard for f, then the PET-induction does not work well enough to provide an upper bound for $S(A,\tau)$ in terms of some Host-Kra seminorm of f. To overcome this difficulty one follows [6]. More specifically, using [6, Proposition 6.3], which is a "dimension-increment" argument, A can be transformed into a new PET-tuple which is 1-standard for f (at the cost of increasing the dimension from L to 2L which is harmless for our approach). So, following this procedure, for any fixed function f, we may assume without loss of generality that the corresponding polynomial iterate p is of maximum degree, making the PET-tuple, after potential rearrangement of the terms, 1-standard for f. A combination of the previous results will allow us to obtain the required upper bound for each function.

²⁷In [6, Theorem 4.2] the PET tuple is not required to be 1-standard nor 1-inherited; this comes at no extra cost as the polynomials chosen at each step to run the vdC-operation are of minimum degree.

5. FINDING A CHARACTERISTIC FACTOR

This lengthy section is dedicated in proving Theorem 2.5. To this end, we need to show two intermediate results, i.e., Propositions 5.2 and 5.3, which improve two technical results from [6], namely [6, Proposition 5.6], and [6, Proposition 5.5] respectively.

Recall that for a subset A of \mathbb{Q}^d , $G(A) = \operatorname{span}_{\mathbb{Q}} \{a \in A\} \cap \mathbb{Z}^d$.

Convention 5.1. For the rest of the paper, for every $\mathbf{u} = (u_1, \ldots, u_L) \in (\mathbb{Q}^d)^L$, we denote $G(\mathbf{u}) := G(\{u_1, \ldots, u_L\}).$

The first result, which enhances [6, Proposition 5.6], gives a bound for the average of interest by finite step seminorms (recall Convention 3.3 for notions on Host-Kra seminorms). To pass from infinite step seminorms to finite step ones, we use the implications of Propositions 3.6 and 3.8.

Proposition 5.2 (Bounding averaged Host-Kra seminorms by a single one). Let $s, s', t \in \mathbb{N}$ and $\mathbf{c}_m : (\mathbb{Z}^L)^s \to (\mathbb{Z}^d)^L, 1 \leq m \leq t$, be polynomials with $\mathbf{c}_m \neq \mathbf{0}$ given by

(16)
$$\mathbf{c}_m(h_1,\ldots,h_s) = \sum_{a_1,\ldots,a_s \in \mathbb{N}_0^L, |a_1| + \cdots + |a_s| \le s'} h_1^{a_1} \ldots h_s^{a_s} \cdot \mathbf{u}_m(a_1,\ldots,a_s)$$

for some

$$\mathbf{u}_m(a_1,\ldots,a_s) = (u_{m,1}(a_1,\ldots,a_s),\ldots,u_{m,L}(a_1,\ldots,a_s)) \in (\mathbb{Q}^d)^L.$$

Denote

$$H_m \coloneqq G(\{u_{m,i}(a_1,\ldots,a_s): a_1,\ldots,a_s \in \mathbb{N}_0^L, 1 \le i \le L\}).$$

There exists $D \in \mathbb{N}_0$ depending only on s, s', L such that for every \mathbb{Z}^d -system $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ and every $f \in L^{\infty}(\mu)$,

(17)
$$\overline{\mathbb{E}}_{h_1,\dots,h_s\in\mathbb{Z}^L}^{\square} |||f|||_{\{G(\mathbf{c}_m(h_1,\dots,h_s))\}_{1\le m\le t}} = 0 \quad if \quad |||f|||_{H_1^{\times D},\dots,H_t^{\times D}} = 0.$$

We start by explaining the idea behind Proposition 5.2 with an example:

Example 8. Let $p_1, p_2: \mathbb{Z} \to \mathbb{Z}^2$ be polynomials given by $p_1(n) = (n^2 + n, 0)$ and $p_2(n) = (0, n^2)$, and $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^2})$ be a \mathbb{Z}^2 -system. Consider the following expression:

$$\sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ \text{Følner seq.}}} \frac{\overline{\lim}_{N\to\infty} \left\| \mathbb{E}_{n\in I_N} T_{p_1(n)} f_1 \cdot T_{p_2(n)} f_2 \right\|_2.$$

Put $e_1 = (1,0)$, $e_2 = (0,1)$ and e = (1,-1). Using [6, Proposition 5.5], we get

(18)
$$\sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ \text{Følner seq.}}} \frac{\lim_{N\to\infty} \left\| \mathbb{E}_{n\in I_N} T_{p_1(n)} f_1 \cdot T_{p_2(n)} f_2 \right\|_2 \le C \cdot \overline{\mathbb{E}}_{\mathbf{h}\in\mathbb{Z}^3}^{\square} \|f_1\|_{G(\mathbf{c}_1(\mathbf{h})),\dots,G(\mathbf{c}_7(\mathbf{h}))},$$

where C is a universal constant, and

$$\begin{aligned} \mathbf{c}_{1}(h_{1}, h_{2}, h_{3}) &= -2h_{1}e_{1}, \\ \mathbf{c}_{2}(h_{1}, h_{2}, h_{3}) &= 2h_{2}e, \\ \mathbf{c}_{3}(h_{1}, h_{2}, h_{3}) &= -2h_{1}e_{1} + 2h_{2}e, \\ \mathbf{c}_{4}(h_{1}, h_{2}, h_{3}) &= -2h_{3}e, \\ \mathbf{c}_{5}(h_{1}, h_{2}, h_{3}) &= -2h_{1}e_{1} + 2h_{3}e, \\ \mathbf{c}_{6}(h_{1}, h_{2}, h_{3}) &= 2(h_{2} + h_{3})e, \\ \mathbf{c}_{7}(h_{1}, h_{2}, h_{3}) &= -2h_{1}e_{1} + 2(h_{2} + h_{3})e. \end{aligned}$$

Using [6, Proposition 5.6], one can show that if $|||f_1|||_{e_1^{\times D}, e^{\times D}} = 0$ for all $D \in \mathbb{N}$, then the right hand side of (18) is 0. In this paper, we strengthen this result by only assuming that $|||f_1|||_{e_1^{\times D}, e^{\times D}} = 0$ for some $D \in \mathbb{N}$.

Indeed, using Proposition 3.8 (for $I = [-N, N]^3$, letting $N \to \infty$) and Corollary 3.7, we have that the right of (18) is 0 if

$$\overline{\mathbb{E}}_{\mathbf{h},\mathbf{h}'\in\mathbb{Z}^3}^{\square} \left\| \mathbb{E}(f|Z_{\{G(\mathbf{c}_i(\mathbf{h}),\mathbf{c}_j(\mathbf{h}'))\}_{1\leq i,j\leq 7}}) \right\|_2^2 = 0.^{28}$$

On the other hand, for "almost all" $\mathbf{h}, \mathbf{h}' \in \mathbb{Z}^3$, the group $G(\mathbf{c}_i(\mathbf{h}), \mathbf{c}_j(\mathbf{h}'))$ equals $\mathbb{Z}e_1$ if i = j = 1, $\mathbb{Z}e$ if $i, j \in \{2, 4, 6\}$ and \mathbb{Z}^2 otherwise. So, $Z_{\{G(\mathbf{c}_i(\mathbf{h}), \mathbf{c}_j(\mathbf{h}'))\}_{1 \leq i, j \leq 7}}$ is contained in $Z_{e_1, e^{\times 9}, (\mathbb{Z}^2)^{\times 39}}$, which is contained in $Z_{e_1^{\times 25}, e^{\times 25}}$ by Proposition 3.1. Hence, the right hand side of (18) is 0 if $|||f_1|||_{e_1^{\times 25}, e^{\times 25}} = 0$.

We now prove the general case.

Proof of Proposition 5.2. We may assume without loss of generality that $||f||_{L^{\infty}(\mu)} \leq 1$. For convenience, denote $\mathbf{h} := (h_1, \ldots, h_s)$. Using Proposition 3.8, and Corollary 3.7 for $I = [-N, N]^{sL}$, and then letting $N \to \infty$, we have that

$$\left(\overline{\mathbb{E}}_{\mathbf{h}\in\mathbb{Z}^{sL}}^{\square}\|\|f\|\|_{\{G(\mathbf{c}_m(\mathbf{h}))\}_{1\leq m\leq t}}\right)^{2^tW}\leq\overline{\mathbb{E}}_{\mathbf{h}^1,\dots,\mathbf{h}^W\in\mathbb{Z}^{sL}}^{\square}\left\|\mathbb{E}(f|Z_{\{G(\mathbf{c}_{m_1}(\mathbf{h}^1),\dots,\mathbf{c}_{m_W}(\mathbf{h}^W))\}_{1\leq m_1,\dots,m_W\leq t}})\right\|_2^2$$

for all $W = 2^w, w \in \mathbb{N}_0$. Let w be the smallest integer such that $2^w \ge t(s'+1)^{sL}$ and Ω the set of $(\mathbf{h}^1, \ldots, \mathbf{h}^W) \in (\mathbb{Z}^{sL})^W$ such that for every $1 \le m_1, \ldots, m_W \le t$, the group $G(\mathbf{c}_{m_1}(\mathbf{h}^1), \ldots, \mathbf{c}_{m_W}(\mathbf{h}^W))$ contains at least one of H_1, \ldots, H_t . By Proposition 3.1, for every $(\mathbf{h}^1, \ldots, \mathbf{h}^W) \in \Omega, Z_{\{G(\mathbf{c}_{m_1}(\mathbf{h}^1), \ldots, \mathbf{c}_{m_W}(\mathbf{h}^W))\}_{1 \le m_1, \ldots, m_W \le t}}$ is a factor of $Z_{H_1^{\times D}, \ldots, H_t^{\times D}}$ for $D := s^W$, and thus $\mathbb{E}(f|Z_{\{G(\mathbf{c}_{m_1}(\mathbf{h}^1), \ldots, \mathbf{c}_{m_W}(\mathbf{h}^W))\}_{1 \le m_1, \ldots, m_W \le t}}) = 0$ since $|||f|||_{H_1^{\times D}, \ldots, H_t^{\times D}} = 0$. So, by (19), it suffices to show that Ω is of upper density 1. Let $\tilde{\mathbf{h}} := (\mathbf{h}^1, \ldots, \mathbf{h}^W) \in (\mathbb{Z}^{sL})^W$ and $1 \le m_1, \ldots, m_W \le t$. By the pigeonhole principle, at least $(s'+1)^{sL}$ many of the m_1, \ldots, m_W take the same value. Assume that $m_{i_1} = \cdots = m_{i_{W'}} = m$, where $W' \le (s'+1)^{sL}$ is the

²⁸Strictly speaking, Proposition 3.8 only implies that the right hand side of (18) is 0 if $\overline{\mathbb{E}}_{\mathbf{h},\mathbf{h}'\in\mathbb{Z}^3}^{\square} \left\| \mathbb{E}(f|Z_{\{G(\mathbf{c}_i(\mathbf{h}))+G(\mathbf{c}_j(\mathbf{h}'))\}_{1\leq i,j\leq 7}}) \right\|_2^2 = 0$. However, since $G(\mathbf{c}_i(\mathbf{h})) + G(\mathbf{c}_j(\mathbf{h}'))$ is a finite index subgroup of $G(\mathbf{c}_i(\mathbf{h}),\mathbf{c}_j(\mathbf{h}'))$, we can use Lemma 3.9 (iv) and replace it by the latter as a seminorm subindex.

number of $a_1, \ldots, a_s \in \mathbb{N}_0^L$ with $|a_1| + \cdots + |a_s| \leq s'$. Note that W' depends only on s', s and L. Write $\mathbf{h}^i \coloneqq (h_{i,1}, \ldots, h_{i,s}) \in \mathbb{Z}^{sL}$ and consider the $W' \times W'$ matrix

$$A_{i_1,\dots,i_{W'}}(\mathbf{h}) \coloneqq (h_{i_j,1}^{a_1}\dots h_{i_j,s}^{a_s})_{a_1,\dots,a_s \in \mathbb{N}_0^L, |a_1| + \dots + |a_s| \le s', 1 \le j \le W'}.$$

If det $(A_{i_1,\ldots,i_{W'}}(\tilde{\mathbf{h}})) \neq 0$, then by the definition of $\mathbf{c}_m(\mathbf{h})$, we have that

$$G(\mathbf{c}_{m_1}(\mathbf{h}^1),\ldots,\mathbf{c}_{m_W}(\mathbf{h}^W)) \supseteq G(\mathbf{c}_m(\mathbf{h}^{i_1}),\ldots,\mathbf{c}_m(\mathbf{h}^{i_{W'}})) \supseteq H_m.$$

In conclusion, $\tilde{\mathbf{h}} \in \Omega$ if for all $1 \leq i_1 < \cdots < i_{W'} \leq W$, $\det(A_{i_1,\dots,i_{W'}}(\tilde{\mathbf{h}})) \neq 0$.

Thus, it suffices to show that for all $1 \leq i_1 < \cdots < i_{W'} \leq W$, the set of $\tilde{\mathbf{h}} \in (\mathbb{Z}^{sL})^W$ with $\det(A_{i_1,\ldots,i_{W'}}(\tilde{\mathbf{h}})) = 0$ is of density 0. We may assume without loss of generality that $i_1 = 1, \ldots, i_{W'} = W'$. Note that $\det(A_{1,\ldots,W'}(\tilde{\mathbf{h}}))$ is a polynomial in $h_{i,j}, 1 \leq i \leq W'$, $1 \leq j \leq s$. Looking at the term $h_{1,1}^{(s,0,\ldots,0)} h_{2,2}^{(s,0,\ldots,0)} \ldots h_{W',W'}^{(s,0,\ldots,0)}$, we see that $\det(A_{1,\ldots,W'}(\tilde{\mathbf{h}}))$ is a non-constant polynomial. Therefore, the set of solutions to $\det(A_{1,\ldots,W'}(\tilde{\mathbf{h}})) = 0$ is of 0 density by Lemma 3.11, completing the argument.

The second statement, which strengthens [6, Proposition 5.5], is the following (check Definition 2.1 for the various notions appearing in the statement):

Proposition 5.3 (Bounding the average by averaged Host-Kra seminorms). Let $d, k, L \in \mathbb{N}$, $\mathbf{p} = (p_1, \ldots, p_k), p_1, \ldots, p_k \colon \mathbb{Z}^L \to \mathbb{Z}^d$ a family of essentially distinct polynomials of degrees at most K, with $p_i(n) = \sum_{v \in \mathbb{N}_0^L, |v| \leq K} b_{i,v} n^v$ for some $b_{i,v} \in \mathbb{Q}^d$. There exist $s, s', t_1, \ldots, t_k \in \mathbb{N}$ depending only on d, k, K, L,²⁹ and polynomials $\mathbf{c}_{i,m} \colon (\mathbb{Z}^L)^s \to (\mathbb{Z}^d)^L, 1 \leq i \leq k, 1 \leq m \leq t_i$, with $\mathbf{c}_{i,m} \not\equiv \mathbf{0}$, such that the following hold:

(i) (Control of the coefficients) Each $\mathbf{c}_{i,m}$ is of the form

$$\mathbf{c}_{i,m}(h_1,\ldots,h_s) = \sum_{a_1,\ldots,a_s \in \mathbb{N}_0^L, |a_1| + \cdots + |a_s| \le s'} h_1^{a_1} \ldots h_s^{a_s} \cdot \mathbf{v}_{i,m}(a_1,\ldots,a_s)$$

for some $\mathbf{v}_{i,m}(a_1,\ldots,a_s) = (v_{i,m,1}(a_1,\ldots,a_s),\ldots,v_{i,m,L}(a_1,\ldots,a_s)) \in (\mathbb{Q}^d)^L$, which is a polynomial function in terms of the coefficients of $p_i, 1 \leq i \leq k$, and whose degree depends only on d, k, K, L.

In addition, denoting

$$H_{i,m} := G(\{v_{i,m,j}(a_1,\ldots,a_s): a_1,\ldots,a_s \in \mathbb{N}_0^L, 1 \le j \le L\}),\$$

we have that $H_{i,m}$ contains one of $G_{i,j}(\mathbf{p}), 0 \leq j \leq k, j \neq i$.

(ii) (Control of the average) For every \mathbb{Z}^d -system $(X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ and every $f_1, \ldots, f_k \in L^{\infty}(\mu)$ bounded by 1, we have that

(20)
$$\sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ F \notin lner \ seq.}} \overline{\lim_{N\to\infty}} \left\| \mathbb{E}_{n\in I_N} \prod_{i=1}^k T_{p_i(n)} f_i \right\|_2^{2^{t_0}} \le C \cdot \min_{1\le i\le k} \overline{\mathbb{E}}_{h_1,\ldots,h_s\in\mathbb{Z}^L}^{\square} \|f_i\|_{(G(\mathbf{c}_{i,m}(h_1,\ldots,h_s)))_{1\le m\le t_i}},$$

²⁹The fact that s, s' depend only on d, k, K, L was not stated in [6, Proposition 5.1], but it can be derived from its proof.

24 SEBASTIÁN DONOSO, ANDREU FERRÉ MORAGUES, ANDREAS KOUTSOGIANNIS AND WENBO SUN

where t_0 and C > 0 are constants depending only on \mathbf{p} .³⁰

Proposition 5.3 improves on [6, Proposition 5.5] as the description of the subgroup $H_{i,m}$ is much more precise than that of the set $U_{i,r}(a_1,\ldots,a_s)$ defined in the latter. The rest of this section is devoted to proving Proposition 5.3, which is the most technical result of this paper.

Next we introduce some convenient notation. Let $\mathbf{q} = (q_1, \ldots, q_\ell)$ be a tuple of polynomials $q_i: (\mathbb{Z}^L)^{s+1} \to \mathbb{Z}^d, 1 \leq i \leq \ell$, where

$$q_i(n; h_1, \dots, h_s) = \sum_{b, a_1, \dots, a_s \in \mathbb{N}_0^L} h_1^{a_1} \dots h_s^{a_s} n^b \cdot u_i(b; a_1, \dots, a_s)$$

for some $u_i(b; a_1, \ldots, a_s) \in \mathbb{Q}^d$, $1 \le i \le \ell$. Then, by writing

$$\mathbf{u}(b;a_1,\ldots,a_s) \coloneqq (u_1(b;a_1,\ldots,a_s),\ldots,u_\ell(b;a_1,\ldots,a_s)) \in (\mathbb{Q}^d)^\ell,$$

we can express ${\bf q}$ as

$$\mathbf{q}(n;h_1,\ldots,h_s) = \sum_{b,a_1,\ldots,a_s \in \mathbb{N}_0^L} h_1^{a_1} \ldots h_s^{a_s} n^b \cdot \mathbf{u}(b;a_1,\ldots,a_s)$$

We call $\mathbf{u}(b; a_1, \ldots, a_d)$ the data of \mathbf{q} at level $(b; a_1, \ldots, a_d)$, or simply the level data of \mathbf{q} .

For the rest of the section, $\mathbf{p} = (p_1, \ldots, p_k)$ denotes a family of essentially distinct polynomials $p_1, \ldots, p_k \colon \mathbb{Z}^L \to \mathbb{Z}^d$ of degrees at most K, with $p_i(n) = \sum_{v \in \mathbb{N}_0^L, |v| \leq K} b_{i,v} n^v$, where $b_{i,v} \in \mathbb{Q}^d$.

One sees that the left hand side of (20) is S(A, 1), where A is the PET-tuple $(L, 0, k, \mathbf{p}, (f_1, \ldots, f_k))$. To prove Proposition 5.3, we first need to perform a series of vdC-operations to convert A into a PET-tuple $\partial_{\rho_t} \ldots \partial_{\rho_1} A$ of degree 1, and then compare the coefficients of the polynomials in A with those in $\partial_{\rho_t} \ldots \partial_{\rho_1} A$. Even though the coefficients in the latter are very difficult to compute directly, one can keep track of the connection between them and those of the original polynomial family \mathbf{p} . This was first achieved in [6] by introducing an equivalence relation pertaining to the vdC-operation (see [6, Section 5.3] for details). In this paper, we introduce another approach which is more intricate than the one used in [6], but that achieves a better tracking of the coefficients, which in turn gives us a stronger upper bound for the multiple averages.

Recalling that $b_{w,v}$, $1 \le w \le k, v \in \mathbb{N}_0^L$ are the coefficients that arise from the family **p** (we also put $b_{0,v} \coloneqq \mathbf{0} \in \mathbb{Q}^d$ for all $v \in \mathbb{N}_0^L$), for $r \in \mathbb{Q}, v \in \mathbb{N}_0^L$ and $0 \le i \le k$, we set

$$Q_{r,i,v} \coloneqq \{r(b_{w,v} - b_{i,v}) \colon 1 \le w \le k\}.$$

Definition 5.4 (Types and symbols of level data). For all $b, a_1, \ldots, a_s, v \in \mathbb{N}_0^L$, $r \in \mathbb{Q}$, and $0 \le i \le k$, we say that $\mathbf{u}(b; a_1, \ldots, a_s)$ is of type (r, i, v) if

$$u_1(b; a_1, \ldots, a_s), \ldots, u_\ell(b; a_1, \ldots, a_s) \in Q_{r,i,v}, \text{ and } u_1(b; a_1, \ldots, a_s) = r(b_{1,v} - b_{i,v}).$$

We say that $\mathbf{u}(b; a_1, \ldots, a_s)$ is *non-trivial* if at least one of $u_m(b; a_1, \ldots, a_s)$ is nonzero.

³⁰Actually both t_0 and C depend on d, k, L and the highest degree of p_1, \ldots, p_k . More specifically, t_0 can be chosen to be the max $\{t_1, \ldots, t_k\}$, where t_i is the number of vdC-operations that we have to perform in order for the PET tuple to be non-degenerate, 1-standard for f_i and with degree equal to 1.

Let $\mathbf{u}(b; a_1, \ldots, a_s)$ be of type (r, i, v). Suppose that

 $(u_1(b; a_1, \dots, a_s), \dots, u_\ell(b; a_1, \dots, a_s)) = (r(b_{w_1, v} - b_{i, v}), \dots, r(b_{w_\ell, v} - b_{i, v})),$

for some $0 \leq w_1, \ldots, w_\ell \leq k$. We call $\mathbf{w} := (w_1, \ldots, w_\ell)$ a symbol of $\mathbf{u}(b; a_1, \ldots, a_s)$.

Note that the definition of type and symbol depend on the prefixed polynomial family **p** and that we always have $w_1 = 1$.

We use the types and symbols of level data to track the coefficients of PET-tuples. We start with an example to illustrate this concept:

Example 9. Let $\mathbf{p} = (p_1, p_2)$ be defined as in Example 7 and let $\mathbf{q} = \partial_2 \mathbf{p} = (q_1, q_2, q_3)$ (recall that by this we mean the polynomial iterates we get after running the vdC-operation, subtracting the second polynomial, p_2). In this case,

$$\mathbf{u}(0;1) = (b_{1,1}, b_{2,1}, 0) \text{ is of type } (1,0,1) \text{ and has symbol } (1,2,0),$$

$$\mathbf{u}(0;2) = (b_{1,2}, b_{2,2}, 0) \text{ is of type } (1,0,2) \text{ and has symbol } (1,2,0),$$

$$\mathbf{u}(1;0) = (b_{1,1} - b_{2,1}, 0, b_{1,1} - b_{2,1}) \text{ is of type } (1,2,1) \text{ and has symbol } (1,2,1),$$

$$\mathbf{u}(1;1) = (2b_{1,2}, 2b_{2,2}, 0) \text{ is of type } (2,0,2) \text{ and has symbol } (1,2,0),$$

$$\mathbf{u}(2;0) = (b_{1,2} - b_{2,2}, 0, b_{1,2} - b_{2,2}) \text{ is of type } (1,2,2) \text{ and has symbol } (1,2,1).$$

Definition 5.5. Let S denote the set of all $(a, a') \in \mathbb{N}_0^L$ such that a and a' are both **0** or both different than 0. Let \mathbf{q} be a polynomial family of degree at least 1. We say that \mathbf{q} satisfies (P1)-(P4) if its level data **u** satisfy:

(P1) For all a_1, \ldots, a_s, b , there exist r, i, v such that $\mathbf{u}(b; a_1, \ldots, a_s)$ is of type (r, i, v). (P2) Suppose that $\mathbf{u}(b; a_1, \ldots, a_s)$ is of type (r, i, v), then $r = \begin{pmatrix} b+a_1+\cdots+a_s\\a_1, \ldots, a_s \end{pmatrix}$ and $v = b+a_1+\cdots+a_s$ $\cdots + a_s$ (in particular, $r \neq 0$).³¹

(P3) Suppose that $\mathbf{u}(b; a_1, \ldots, a_s)$ is of type (r, i, v) and $\mathbf{u}(b'; a'_1, \ldots, a'_s)$ is of type (r', i', v'). If $(a_1, a'_1), \ldots, (a_s, a'_s) \in S$, then i = i' and $\mathbf{u}(b; a_1, \ldots, a_s)$, $\mathbf{u}(b'; a'_1, \ldots, a'_s)$ share a symbol \mathbf{w} . (P4) For every symbol (w_1, \ldots, w_ℓ) of some $\mathbf{u}(b; a_1, \ldots, a_s)$, we have that $w_1 = 1$.

Once again, properties (P1)–(P4) are taken with respect to the prefixed polynomial family **p**. It is obvious that **p** itself satisfies (P1)-(P4). An important feature of the type and symbol of a level data is that properties (P1)–(P4) are preserved under vdC-operations.

Example 10. We will verify that the polynomial family $\mathbf{q} = \partial_2 \mathbf{p}$ in Example 9 satisfies all of (P1)–(P4). Indeed, (P1) holds as all $\mathbf{u}(0;1), \mathbf{u}(0;2), \mathbf{u}(1;0), \mathbf{u}(1;1), \mathbf{u}(2;0)$ have a type. For all $0 \le a, b \le 2, a + b \le 2$, if $\mathbf{u}(b; a)$ is of type (r, i, v), then it is not hard to see that $r = {\binom{b+a}{a}}$, so (P2) holds. (P3) can be verified by comparing the types and symbols of the pairs $(\mathbf{u}(0,1),\mathbf{u}(0,2))$ and $(\mathbf{u}(1,0),\mathbf{u}(2,0))$. Finally, (P4) also holds since the first entry of every symbol in \mathbf{q} is 1.

We caution the reader that the symbol and type may not be unique if the coefficients $b_{i,v}$ satisfy some algebraic relations. For example, in Example 9, if $b_{1,1} = b_{2,1}$, then both (1,2,0)and (1,1,0) are symbols of $\mathbf{u}(0;1) = (b_{1,1}, b_{2,1}, 0)$. However the following result says that there is always a way to choose symbols and types so that properties (P1)–(P4) are preserved under vdC-operations:

³¹For $v_i = (v_{i,1}, \dots, v_{i,L}) \in \mathbb{N}_0^L, 1 \le i \le s$, we denote $\binom{v_1 + \dots + v_s}{v_1, \dots, v_{s-1}} \coloneqq \prod_{j=1}^L \frac{(v_{1,j} + \dots + v_{s,j})!}{v_{1,j}! \dots v_{s,j}!}$.

Proposition 5.6. Let $A = (L, s, \ell, \mathbf{g}, \mathbf{q})$ be a non-degenerate PET-tuple and $1 \leq \rho \leq \ell$. Assume that $A \to \partial_{\rho}A$ is 1-inherited. Suppose that there exists a choice of symbols and types for A which satisfy (P1)-(P4), then there is also a choice of symbols and types for $\partial_{\rho}A$ which satisfy (P1)-(P4).

Proof. Suppose that $\mathbf{q} = (q_1, \ldots, q_\ell)$. Denote $\mathbf{q}^* = (q_1^*, \ldots, q_{2\ell}^*)$, where for all $1 \le i \le \ell$,

$$q_i^*(n; h_1, \dots, h_{s+1}) = q_i(n + h_{s+1}; h_1, \dots, h_s) - q_\rho(n; h_1, \dots, h_s)$$

and

$$q_{\ell+i}^*(n;h_1,\ldots,h_{s+1}) = q_i(n;h_1,\ldots,h_s) - q_\rho(n;h_1,\ldots,h_s).$$

Assuming that

$$q_i(n; h_1, \dots, h_s) = \sum_{b, a_1, \dots, a_s \in \mathbb{N}_0^L} h_1^{a_1} \dots h_s^{a_s} n^b \cdot u_i(b; a_1, \dots, a_s)$$

for all $1 \leq i \leq \ell$, we may write **q** as

$$\mathbf{q}(n;h_1,\ldots,h_s) = \sum_{b,a_1,\ldots,a_s \in \mathbb{N}_0^L} h_1^{a_1} \ldots h_s^{a_s} n^b \cdot \mathbf{u}(b;a_1,\ldots,a_s),$$

and define $\mathbf{u}^*(b; a_1, \ldots, a_{s+1})$ in a similar way.

One can immediately check that

$$q_i(n+h_{s+1};h_1,\ldots,h_s) = \sum_{b,a_1,\ldots,a_{s+1}\in\mathbb{N}_0^L} h_1^{a_1}\ldots h_{s+1}^{a_{s+1}} n^b \cdot \binom{b+a_{s+1}}{b} u_i(b+a_{s+1};a_1,\ldots,a_s).^{32}$$

Then

(21)
$$u_i^*(b; a_1, \dots, a_{s+1}) = \begin{cases} u_i(b; a_1, \dots, a_s) - u_\rho(b; a_1, \dots, a_s) & (a_{s+1} = \mathbf{0}) \\ \begin{pmatrix} b + a_{s+1} \\ b \end{pmatrix} u_i(b + a_{s+1}; a_1, \dots, a_s) & (a_{s+1} \neq \mathbf{0}) \end{cases}$$

and

(22)
$$u_{i+\ell}^*(b;a_1,\ldots,a_{s+1}) = \begin{cases} u_i(b;a_1,\ldots,a_s) - u_\rho(b;a_1,\ldots,a_s) & (a_{s+1} = \mathbf{0}) \\ \mathbf{0} & (a_{s+1} \neq \mathbf{0}) \end{cases}$$

We first show that \mathbf{q} satisfying (P1)–(P4) implies the same for \mathbf{q}^* .

Fix a_1, \ldots, a_{s+1} . Since **q** satisfies (P1)–(P4), there exist $0 \le i \le k$ and a symbol **w** such that for all b, there is $v \in \mathbb{N}_0^L$ and $r \in \mathbb{Q}$ such that $\mathbf{u}(b + a_{s+1}; a_1, \ldots, a_s)$ is of type (r, i, v) and has symbol $\mathbf{w} = (1, w_2, \ldots, w_\ell)$.

We have that $u_m(b; a_1, ..., a_s) = r(b_{w_m,v} - b_{i,v})$. By (21) and (22), it is not hard to see that $\mathbf{u}^*(b; a_1, ..., a_s, a_{s+1})$ is

(23)
$$\begin{cases} \text{of type } (r, w_{\rho}, v) \text{ and has symbol } (\mathbf{w}, \mathbf{w}) & (a_{s+1} = \mathbf{0}) \\ \text{of type } (r \binom{b + a_{s+1}}{b}, i, v) \text{ and has symbol } (\mathbf{w}, i, \dots, i) & (a_{s+1} \neq \mathbf{0}) \end{cases}$$

So each $\mathbf{u}^*(b; a_1, \ldots, a_s, a_{s+1})$ is of type (*, *, v) for some v (meaning that it is of type (r', i', v) for some $r' \in \mathbb{Q}$ and $i' \in \mathbb{N}$) and thus \mathbf{q}^* satisfies (P1).

³²For $a = (a_1, \ldots, a_L), b = (b_1, \ldots, b_L) \in \mathbb{N}_0^L, \binom{a}{b}$ denotes the quantity $\prod_{m=1}^L \binom{a_m}{b_m}$.

To check (P2) for \mathbf{q}^* , suppose that $\mathbf{u}^*(b; a_1, \ldots, a_{s+1})$ is of type (r, i, v). By (23), we have that $\mathbf{u}(b + a_{s+1}; a_1, \ldots, a_s)$ is of type $(r\binom{b+a_{s+1}}{b}^{-1}, i', v)$ for some i'. Since \mathbf{q} satisfies (P2), we have the same for \mathbf{q}^* as

$$r = {\binom{b+a_{s+1}}{b}} {\binom{(b+a_{s+1})+a_1+\dots+a_s}{a_1,\dots,a_s}} = {\binom{b+a_1+\dots+a_{s+1}}{a_1,\dots,a_{s+1}}}$$

and

$$v = (b + a_{s+1}) + a_1 + \dots + a_s = b + a_1 + \dots + a_{s+1}$$

To show (P3), let $(a_1, a'_1), \ldots, (a_{s+1}, a'_{s+1}) \in S$ and suppose that $\mathbf{u}(b + a_{s+1}; a_1, \ldots, a_s)$ is of type (r, i, v) and $\mathbf{u}(b' + a'_{s+1}; a'_1, \ldots, a'_s)$ of type (r', i', v'). Since \mathbf{q} satisfies (P3), we have that i = i' and that $\mathbf{u}(b + a_{s+1}; a_1, \ldots, a_s)$ and $\mathbf{u}(b' + a'_{s+1}; a'_1, \ldots, a'_s)$ can be made to have the same symbol \mathbf{w} . Since $(a_{s+1}, a'_{s+1}) \in S$, by (23) have that \mathbf{q}^* satisfies (P3).

Finally, it is straightforward from (23) that Property (P4) is also heritable. Since the polynomials in $\partial_{\rho}A$ are obtained by removing some terms from the tuple \mathbf{q}^* (but not removing the first one, since $A \to \partial_{\rho}A$ is 1-inherited), the fact that \mathbf{q}^* satisfies (P1)–(P4) implies that $\partial_{\rho}A$ also satisfies (P1)–(P4).

For the family of essentially distinct polynomials $\mathbf{p} = (p_1, \ldots, p_k)$, Proposition 5.6 implies that $\partial_{i_k} \ldots \partial_{i_1} \mathbf{p}$ satisfies (P1)–(P4) for all $k, i_1, \ldots, i_k \in \mathbb{N}$. In the special case when $\partial_{i_k} \ldots \partial_{i_1} \mathbf{p}$ is of degree 1, properties (P1)–(P4) provide us with some information on the seminorm we use to bound $S(\partial_{i_k} \ldots \partial_{i_1} A, 1)$. To be more precise, if (P1)–(P4) hold for some non-degenerate \mathbf{q} , then there is some connection between the level data of \mathbf{q} and the groups $G_{1,0}(\mathbf{p}), G_{1,2}(\mathbf{p}), \ldots, G_{1,k}(\mathbf{p})$:

Proposition 5.7. Suppose that (P1)–(P4) hold for some non-degenerate **q**. Then for all $0 \le m \le l, m \ne 1$, the group

$$H_{1,m}(\mathbf{q}) \coloneqq G(\{u_1(b; a_1, \dots, a_s) - u_m(b; a_1, \dots, a_s) \colon (b, a_1, \dots, a_s) \in (\mathbb{N}_0^L)^{s+1}, b \neq \mathbf{0}\})$$

contains at least one of the groups $G_{1,j}(\mathbf{p}), 0 \leq j \leq k, j \neq 1$.

We remark that although Proposition 5.7 holds for all non-degenerate \mathbf{q} , we will use it for the case deg(\mathbf{q}) = 1. We first give an example to explain the idea behind it.

Example 11. Let $\mathbf{p} = (p_1, p_2)$ be as in Example 7. Then $G_{1,0}(\mathbf{p}) = G(b_{1,2})$ and $G_{1,2}(\mathbf{p}) = G(b_{1,2} - b_{2,2})$. Let $\mathbf{u}(b; a)$ be the level data of $\partial_2 \mathbf{p}$. Then

$$\mathbf{u}(1;0) = (b_{1,1} - b_{2,1}) \cdot (1,0,1) \text{ is of type } (1,2,1) \text{ and has symbol } (1,2,1), \mathbf{u}(1;1) = 2b_{1,2} \cdot (1,0,0) + 2b_{2,2} \cdot (0,1,0) \text{ is of type } (2,0,2) \text{ and has symbol } (1,2,0), \mathbf{u}(2;0) = (b_{1,2} - b_{2,2}) \cdot (1,0,1) \text{ is of type } (1,2,2) \text{ and has symbol } (1,2,1).$$

Here we will not compute $\mathbf{u}(b; a)$ for b = 0 as it is irrelevant to our purposes. It is easy to see that $H_{1,0}(\partial_2 \mathbf{p}) = G(b_{1,1} - b_{2,1}, b_{1,2}, b_{1,2} - b_{2,2}) \supseteq G_{1,0}(\mathbf{p}) \cup G_{1,2}(\mathbf{p}), H_{1,2}(\partial_2 \mathbf{p}) = G(b_{1,1} - b_{2,1}, b_{1,2} - b_{2,2}) \supseteq G_{1,0}(\mathbf{p}).$ So, Proposition 5.7 holds for $\partial_2 \mathbf{p}$.

28 SEBASTIÁN DONOSO, ANDREU FERRÉ MORAGUES, ANDREAS KOUTSOGIANNIS AND WENBO SUN

Let $\mathbf{u}'(b; a_1, a_2)$ denote the level data of $\partial_2 \partial_2 \mathbf{p}$. Then

 $\mathbf{u}'(1;0,0) = (b_{1,1} - b_{2,1}) \cdot (1,1,1,1)$ is of type (1,2,1) and has symbol (1,1,1,1),

 $\mathbf{u}'(1;1,0) = 2(b_{1,2} - b_{2,2}) \cdot (1,0,1,0) - 2b_{2,2} \cdot (0,1,0,1)$ has type (2,2,2) and sym. (1,0,1,0),

 $\mathbf{u}'(1;0,1) = 2(b_{1,2} - b_{2,2}) \cdot (1,1,0,0)$ is of type (2,2,2) and has symbol (1,1,2,2),

 $\mathbf{u}'(2;0,0) = (b_{1,2} - b_{2,2}) \cdot (1,1,1,1)$ is of type (1,2,2) and has symbol (1,1,1,1).

(We do not compute the types and symbols for $\mathbf{u}'(b; a_1, a_2)$ for b = 0.) It is easy to see that $H_{1,0}(\partial_2 \partial_2 \mathbf{p}) = G(b_{1,1} - b_{2,1}, b_{1,2} - b_{2,2}) \supseteq G_{1,2}(\mathbf{p}), H_{1,2}(\partial_2 \partial_2 \mathbf{p}) = G(b_{1,2}) = G_{1,0}(\mathbf{p}),$ $H_{1,3}(\partial_2 \partial_2 \mathbf{p}) = G(b_{1,2} - b_{2,2}) \supseteq G_{1,2}(\mathbf{p}), H_{1,4}(\partial_2 \partial_2 \mathbf{p}) = G(b_{1,2}, b_{1,2} - b_{2,2}) \supseteq G_{1,0}(\mathbf{p}) \cup G_{1,2}(\mathbf{p}).$ So, Proposition 5.7 holds for $\partial_2 \partial_2 \mathbf{p}$.

Finally, let $\mathbf{u}''(b; a_1, a_2, a_3)$ denote the level data of $\partial_4 \partial_2 \partial_2 \mathbf{p}$. Then $\deg(\partial_4 \partial_2 \partial_2 \mathbf{p}) = 1$ and $\mathbf{u}''(1; 0, 0, 0) = (0, 0, 0, 0, 0, 0, 0)$ is trivial,

 $\mathbf{u}''(1;1,0,0) = 2b_{1,2} \cdot (1,0,1,0,1,0,1)$ is of type (2,0,2) and symbol (1,0,1,0,1,0,1),

 $\mathbf{u}''(1;0,1,0) = 2(b_{1,2} - b_{2,2}) \cdot (1,1,0,0,1,1,0)$ has type (2,2,2) and symbol (1,1,2,2,1,1,2),

 $\mathbf{u}''(1;0,0,1) = 2(b_{1,2} - b_{2,2}) \cdot (1,1,1,1,0,0,0)$ has type (2,2,2) and symbol (1,1,1,1,2,2,2).

(Once more, we do not compute the types and symbols for $\mathbf{u}''(b; a_1, a_2, a_3)$ for b = 0.) It is easy to see that $H_{1,0}(\partial_4 \partial_2 \partial_2 \mathbf{p}) = H_{1,4}(\partial_4 \partial_2 \partial_2 \mathbf{p}) = H_{1,6}(\partial_4 \partial_2 \partial_2 \mathbf{p}) = G(b_{1,2}, b_{1,2} - b_{2,2}) \supseteq G_{1,1}(\mathbf{p}),$ $H_{1,2}(\partial_4 \partial_2 \partial_2 \mathbf{p}) = G(b_{1,2}) = G_{1,1}(\mathbf{p}),$ and $H_{1,3}(\partial_4 \partial_2 \partial_2 \mathbf{p}) = H_{1,5}(\partial_4 \partial_2 \partial_2 \mathbf{p}) = H_{1,7}(\partial_4 \partial_2 \partial_2 \mathbf{p}) = G(b_{1,2} - b_{2,2}) = G_{1,2}(\mathbf{p}).$ So, Proposition 5.7 holds for $\partial_4 \partial_2 \partial_2 \mathbf{p}$.

To briefly explain why Proposition 5.7 holds for Example 11, we explain, for convenience, why $H_{1,0}(\mathbf{q})$ contains either $G_{1,0}(\mathbf{p})$ or $G_{1,2}(\mathbf{p})$ for $\mathbf{q} = \partial_2 \mathbf{p}, \partial_2 \partial_2 \mathbf{p}$ and $\partial_4 \partial_2 \partial_2 \mathbf{p}$. Let $\mathbf{u}(b; a_1, \ldots, a_s)$ be a level data of type (r, i, v) and symbol \mathbf{w} , and $\mathbf{u}(b'; a'_1, \ldots, a'_s)$ be a level data of type (r', i', v') and symbol \mathbf{w}' . We say that the level data $\mathbf{u}(b'; a'_1, \ldots, a'_s)$ dominates (or strictly dominates) $\mathbf{u}(b; a_1, \ldots, a_s)$ if i = i', $\mathbf{w} = \mathbf{w}'$ and $|v'| \ge |v|$ (or |v'| > |v| respectively). In Example 11, it is not hard to see that for all $b \in \mathbb{N}$, $a_1, a_2 \in \mathbb{N}_0$, if $\mathbf{u}'(b; a_1, a_2)$ is not of type (*, *, 2), then there exist $b' \in \mathbb{N}$, $a'_1, a'_2 \in \mathbb{N}_0$ such that $\mathbf{u}'(b'; a'_1, a'_2)$ strictly dominates $\mathbf{u}'(b; a_1, a_2)$ (in this example, $\mathbf{u}'(1; 0, 0)$ is strictly dominated by $\mathbf{u}'(2; 0, 0)$). Similar conclusions hold for $\mathbf{u}(b; a)$ and $\mathbf{u}''(b; a_1, a_2, a_3)$. In other words, the group $H_{1,0}(\mathbf{q})$ must contain the elements of a level data of type (*, *, 2), and thus it must contain one of $G_{1,0}(\mathbf{p})$ and $G_{1,2}(\mathbf{p})$.

We are now ready to prove Proposition 5.7. The main point is that given any nontrivial level data $\mathbf{u}(b; a_1, \ldots, a_s)$, we can find another one, $\mathbf{u}(b'; a'_1, \ldots, a'_s)$, which dominates $\mathbf{u}(b; a_1, \ldots, a_s)$ and is of type (*, *, v) with |v| being as large as possible (in this step we need to exploit the properties (P1)–(P4)). After that, we use the information of the "top" level data $\mathbf{u}(b'; a'_1, \ldots, a'_s)$ to conclude.

Proof of Proposition 5.7. We start with a claim. Recall that S denotes the set of all $(a, a') \in \mathbb{N}_0^L$ such that a and a' are both **0** or both different than **0** (check Definition 2.1 for notation).

Claim. Let $d, s \in \mathbb{N}$ and $b, a_1, \ldots, a_s, v \in \mathbb{N}_0^L$. If $|v| \ge |b + a_1 + \cdots + a_s|$, then there exist $b', a'_1, \ldots, a'_s \in \mathbb{N}_0^L$ such that $(a_1, a'_1), \ldots, (a_s, a'_s) \in S, |b'| \ge |b|$ and $b' + a'_1 + \cdots + a'_s = v$.

To show the claim, we may first assume that $|v| = |b + a_1 + \cdots + a_s|$. Indeed, write $x = (x_1, \ldots, x_L)$ for $x = v, b, a_1, \ldots, a_s$. If $c \coloneqq |v| - |b + a_1 + \cdots + a_s| > 0$, then we write $b' = b + (c, 0, \ldots, 0)$. Then |b'| > |b| and $|v| = |b' + a_1 + \cdots + a_s|$.

It suffices to show that if $|v - (b + a_1 + \dots + a_s)| > 0$, then there exist $b', a'_1, \dots, a'_s \in \mathbb{N}_0^L$ such that $(a_1, a'_1), \dots, (a_s, a'_s) \in S$, |b'| = |b|, and $|v - (b' + a'_1 + \dots + a'_s)| < |v - (b + a_1 + \dots + a_s)|$. Since $|v| = |b + a_1 + \dots + a_s|$, $|v - (b + a_1 + \dots + a_s)|$ is at least 2, so, there exist $1 \leq i, j \leq L, i \neq j$ such that the t-th coordinate of $v - (b + a_1 + \dots + a_s)$ is at least 1 for t = i and is at most -1 for t = j. We may assume without loss of generality that i = 1 and j = 2. Then we have that $v_1 \geq 1$, and one of $b_2, a_{1,2}, \dots, a_{s,2}$ is at least 1. If $b_2 \geq 1$, then $b' = b + (1, -1, 0, \dots, 0) \in \mathbb{N}_0^L, a'_i = a_i, 1 \leq i \leq s$ satisfy the requirement. If one of $a_{i,2}$ is positive, then $b' = b, a'_i = a_i + (1, -1, 0, \dots, 0) \in \mathbb{N}_0^L, a'_j = a_j, 1 \leq j \leq s, j \neq i$ satisfy the requirement. This proves the claim.

Consider the group $H_{1,m}(\mathbf{q})$ for some $0 \le m \le \ell, m \ne 1$. Since \mathbf{q} is non-degenerate, there exist some $b, a_1, \ldots, a_s \in \mathbb{N}_0^L, |b| \ge 1$ such that $u_1(b; a_1, \ldots, a_s) - u_m(b; a_1, \ldots, a_s) \ne \mathbf{0}$. By (P1), we may assume that $\mathbf{u}(b; a_1, \ldots, a_s)$ is of type (r, i, v) and has symbol (w_1, \ldots, w_ℓ) . Since $w_1 = 1$,

$$u_1(b; a_1, \ldots, a_s) - u_m(b; a_1, \ldots, a_s) = r(b_{1,v} - b_{w_m,v}).$$

Recall that for $\mathbf{p} = (p_1, \ldots, p_k)$, $d_{1,0} = \deg(p_1)$ and $d_{1,j} = \deg(p_1 - p_j)$ for $2 \le j \le \ell$. Since $u_1(b; a_1, \ldots, a_s) - u_m(b; a_1, \ldots, a_s) \ne \mathbf{0}$, we have that $w_m \ne 1$ and $|v| = |b + a_1 + \cdots + a_s| \le d_{1,w_m}$.

By the Claim, for all $v' \in \mathbb{N}_0^L$ with $|v'| = d_{1,w_m}$, there exist $b', a'_1, \ldots, a'_s \in \mathbb{N}_0^L$ such that $(a_1, a'_1), \ldots, (a_s, a'_s) \in S$, $|b'| \ge 1$ and $b' + a'_1 + \cdots + a'_s = v'$. By (P2) and (P3), $\mathbf{u}(b'; a'_1, \ldots, a'_s)$ is of type $(r', i, v'), r' \ne 0$ and has symbol (w_1, \ldots, w_ℓ) (i.e. $\mathbf{u}(b'; a'_1, \ldots, a'_s)$). So

$$u_1(b'; a'_1, \dots, a'_s) - u_m(b'; a'_1, \dots, a'_s) = r'(b_{1,v'} - b_{w_m,v'}).$$

In other words, for all $v' \in \mathbb{N}^L$ with $|v'| = d_{1,w_m}$, the group $H_{1,m}(\mathbf{q})$ contains a nonzero multiple of $b_{1,v'} - b_{w_m,v'}$. So this group contains $G_{1,w_m}(\mathbf{p})$ and we are done.

We are now ready to prove Proposition 5.3.

Proof of Proposition 5.3. Let A denote the PET-tuple $(L, 0, k, (p_1, \ldots, p_k), (f_1, \ldots, f_k))$. Then for all $\tau > 0$,

$$S(A,\tau) = \sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ \text{Følner seq.}}} \overline{\lim_{N\to\infty}} \left\| \mathbb{E}_{n\in I_N} \prod_{m=1}^{\kappa} T_{p_m(n)} f_m \right\|_{L^2(\mu)}^{\tau}.$$

By assumption, A is non-degenerate. We only prove (20) for f_1 as the other cases are similar. We first assume that A is 1-standard for f_1 . By Theorem 4.6, there exist $t \in \mathbb{N}_0$, depending only on d, k, K, L, and finitely many vdC-operations $\partial_{\rho_1}, \ldots, \partial_{\rho_t}, \rho_1, \ldots, \rho_t \neq 1$ such that for all $1 \leq t' \leq t, \partial_{\rho_{t'}} \ldots \partial_{\rho_1} A$ is non-degenerate and 1-standard for f_1 , and that $\partial_{\rho_{t'-1}} \ldots \partial_{\rho_1} A \rightarrow$ $\partial_{\rho_{t'}} \ldots \partial_{\rho_1} A$ is 1-inherited. Moreover, $A' := \partial_{\rho_t} \ldots \partial_{\rho_1} A$ is of degree 1. By Proposition 4.5, $S(A, 2^t) \leq C \cdot S(A', 1)$ for some C > 0 that depends only on t. We may assume that

$$S(A',1) = \overline{\mathbb{E}}_{h_1,\dots,h_s \in \mathbb{Z}^L}^{\square} \sup_{\substack{(I_N)_{N \in \mathbb{N}} \\ \text{Følner seq.}}} \overline{\lim_{N \to \infty}} \left\| \mathbb{E}_{n \in I_N} \prod_{m=1}^c T_{\mathbf{d}_m(h_1,\dots,h_s) \cdot n + r_m(h_1,\dots,h_s)} g_m(x;h_1,\dots,h_s) \right\|_2$$

0

for some $s, \ell \in \mathbb{N}$, functions $g_1, \ldots, g_\ell \colon X \times (\mathbb{Z}^L)^s \to \mathbb{R}$, where $g_1(\cdot; h_1, \ldots, h_s) = f_1$, such that each $g_m(\cdot; h_1, \ldots, h_s)$ is an $L^{\infty}(\mu)$ function bounded by 1, and polynomials $\mathbf{d}_m \colon (\mathbb{Z}^L)^s \to (\mathbb{Z}^d)^L$

and $r_m: (\mathbb{Z}^L)^s \to \mathbb{Z}^d$, $1 \leq m \leq \ell$, where \mathbf{d}_m, r_m take values vectors with integer coordinates as vdC-operations transform integer-valued polynomials to integer-valued polynomials.

Let $\mathbf{c}_1 = -\mathbf{d}_1$ and $\mathbf{c}_m = \mathbf{d}_m - \mathbf{d}_1$ for $m \neq 1$. Since A' is non-degenerate, we have that $\mathbf{c}_1, \ldots, \mathbf{c}_s \neq \mathbf{0}$. By [6, Proposition 6.1], if $\ell \geq 2$, we also have that

$$S(A',1) \leq C' \cdot \overline{\mathbb{E}}_{h_1,...,h_s \in \mathbb{Z}^L}^{\square} ||| T_{r_1(h_1,...,h_s)} f_1 |||_{\{G(\mathbf{c}_i(h_1,...,h_s))\}_{1 \leq i \leq \ell}}$$

= $C' \cdot \overline{\mathbb{E}}_{h_1,...,h_s \in \mathbb{Z}^L}^{\square} ||| f_1 |||_{\{G(\mathbf{c}_i(h_1,...,h_s))\}_{1 \leq i \leq \ell}}$

for some C' > 0 depending only on the polynomials p_1, \ldots, p_k . If $\ell = 1$, using the mean ergodic theorem (see for example [6, Theorem 2.3]) and [6, Lemma 2.4 (iv), (vi)], we have

$$S(A',1) = \overline{\mathbb{E}}_{h_1,\dots,h_s \in \mathbb{Z}^L}^{\square} \|\mathbb{E}(T_{r_1(h_1,\dots,h_s)}f_1 | \mathcal{I}(\mathbf{c}_1(h_1,\dots,h_s)))) \|_2$$

$$= \overline{\mathbb{E}}_{h_1,\dots,h_s \in \mathbb{Z}^L}^{\square} \|\mathbb{E}(f_1 | \mathcal{I}(\mathbf{c}_1(h_1,\dots,h_s))) \|_2$$

$$= \overline{\mathbb{E}}_{h_1,\dots,h_s \in \mathbb{Z}^L}^{\square} \|f_1\|_{\mathbf{c}_1(h_1,\dots,h_s)}$$

$$\leq \overline{\mathbb{E}}_{h_1,\dots,h_s \in \mathbb{Z}^L}^{\square} \|f_1\|_{G(\mathbf{c}_1(h_1,\dots,h_s)),\mathbf{c}_1(h_1,\dots,h_s)}$$

$$= \overline{\mathbb{E}}_{h_1,\dots,h_s \in \mathbb{Z}^L}^{\square} \|f_1\|_{G(\mathbf{c}_1(h_1,\dots,h_s)) \times 2}.$$

Combining this with the fact that $S(A, 2^t) \leq C \cdot S(A', 1)$, we get (20).³³ We now consider the groups $H_{1,m}, 0 \leq m \leq \ell, m \neq 1$. Suppose that

$$\mathbf{c}_m(h_1,\ldots,h_s) = \sum_{a_1,\ldots,a_s \in \mathbb{N}_0^L} h_1^{a_1} \ldots h_s^{a_s} \cdot \mathbf{v}_m(a_1,\ldots,a_s)$$

and

$$\mathbf{d}_m(h_1,\ldots,h_s) = \sum_{a_1,\ldots,a_s \in \mathbb{N}_0^L} h_1^{a_1} \ldots h_s^{a_s} \cdot \mathbf{u}_m(a_1,\ldots,a_s)$$

for some vectors $\mathbf{u}_m(a_1, \ldots, a_s) = (u_{m,1}(a_1, \ldots, a_s), \ldots, u_{m,L}(a_1, \ldots, a_s))$, and $\mathbf{v}_m(a_1, \ldots, a_s) = (v_{m,1}(a_1, \ldots, a_s), \ldots, v_{m,L}(a_1, \ldots, a_s)) \in (\mathbb{Q}^d)^L$ with all but finitely many terms being zero for each m. Obviously A satisfies (P1)–(P4). Since $A' = \partial_{\rho_t} \ldots \partial_{\rho_1} A$, by Proposition 5.6, A' satisfies (P1)–(P4). Since $\deg(A') = 1$, by Proposition 5.7, each of

$$G(\{v_{1,j}(a_1,\ldots,a_s): (a_1,\ldots,a_s) \in (\mathbb{N}_0^L)^s, 1 \le j \le L\})$$

= $G(\{u_{1,j}(a_1,\ldots,a_s): (a_1,\ldots,a_s) \in (\mathbb{N}_0^L)^s, 1 \le j \le L\}) = H_{1,0}$

and

$$G(\{v_{m,j}(a_1,\ldots,a_s)\colon (a_1,\ldots,a_s)\in (\mathbb{N}_0^L)^s\})$$

= $G(\{u_{1,j}(a_1,\ldots,a_s)-u_{m,j}(a_1,\ldots,a_s)\colon (a_1,\ldots,a_s)\in (\mathbb{N}_0^L)^s\}) = H_{1,m}, \ 2\le m\le \ell$

contains some of the groups $G_{1,j}(\mathbf{p}), 0 \le j \le k, j \ne 1$.

Next we assume that A is not 1-standard for f_1 . In this case, we need to invoke the "dimension increment" argument to convert A to be 1-standard for f_1 . We may assume without loss of generality that p_k has the highest degree. Since A is semi-standard for f_1 , by [6, Proposition 6.3], there exists a PET-tuple $A' = (2L, 0, \ell, \mathbf{p}', \mathbf{g})$ which is non-degenerate and

³³We are using [6, Proposition 6.1] for $\ell \geq 2$ (which is incorrectly stated for $\ell \geq 1$ in [6] but does not affect any other result in that work).

1-standard for f_1 such that $S(A, 2\tau) \leq S(A', \tau)$ for all $\tau > 0$. Moreover, \mathbf{p}' is obtained by selecting some polynomials from the family

$$\mathbf{q} := (p_1(n) - p_k(n'), \dots, p_k(n) - p_k(n'), p_1(n') - p_k(n'), \dots, p_{k-1}(n') - p_k(n'))$$

with 2L-dimensional variables (n, n'), where $p_1(n) - p_k(n')$ is selected in p' and is associated to f_1 . It is not hard to compute that $G_{1,j}(\mathbf{q}) = G_{1,j}(\mathbf{p})$ for $0 \le j \le k, j \ne 1$. Moreover, for $1 \le j \le k-1, G_{1,k+j}(\mathbf{q}) = G_{1,0}(\mathbf{p}) + G_{j,0}(\mathbf{p}) \supseteq G_{1,0}(\mathbf{p}) \text{ if } d_{1,0} = d_{j,0}, G_{1,k+j}(\mathbf{q}) = G_{1,0}(\mathbf{p}) = G_{1,0}(\mathbf{p})$ $G_{1,j}(\mathbf{q})$ if $d_{1,0} > d_{j,0}$, and $G_{1,k+j}(\mathbf{q}) = G_{j,0}(\mathbf{p}) = G_{1,j}(\mathbf{q})$ if $d_{1,0} < d_{j,0}$. In other words, each $G_{1,j}(\mathbf{q})$ and thus each $G_{1,j}(\mathbf{p}')$ contains some $G_{1,j'}(\mathbf{p})$. Applying the previous conclusion to A', we are done.

Finally the fact that each $\mathbf{v}_{i,m}(a_1,\ldots,a_s)$ is a polynomial function in terms of the coefficients of $p_i, 1 \leq i \leq k$ whose degree depends only on d, k, K, L follows easily from the polynomial nature of the vdC-operations.

We now have all the ingredients needed to prove Theorem 2.5. In fact, Theorem 2.5 has a proof similar to that of [6, Theorem 5.1].

Proof of Theorem 2.5. By Propositions 5.3 and 5.2, and the definition of Host-Kra characteristic factors, the left hand side of (10) is 0 if for some $1 \leq i \leq k$, f_i is orthogonal to $Z_{\{H_{i,m}\}_{1\leq m\leq t_i}^{\times D_i}}(\mathbf{X})$ for some $t_i, D_i \in \mathbb{N}$, where $H_{i,m}$ is defined as in Proposition 5.3. By Proposition 5.3, $H_{i,m}$ is contained in one of $G_{i,j}(\mathbf{p}), 0 \leq j \leq k, j \neq i$. Using Proposition 3.1 (v), if some f_i , $1 \leq i \leq k$, is orthogonal to $Z_{\{G_{i,j}(\mathbf{p})\}_{\substack{0 \leq j \leq k, j \neq i \\ 1 \leq m \leq t_i}}}(\mathbf{X})$, then it is also orthogonal to $Z_{\{H_{i,m}\}_{\substack{1 \leq m \leq t_i}}}(\mathbf{X})$, and thus the left hand side of (10) is 0.

The "in particular" part follows from Corollary 3.2.

Remark 5.8. We remark that the number D derived in Theorem 2.5 is not optimal.

To see this, recall that this number indicates the step of the nilsequence in the splitting results. For multicorrelation sequences with general polynomial iterates, this D can be taken to be equal to the number of vdC-operations we have to perform in order for all the iterates to become constant (e.g., [11] via [10], and [23] via [11]). At this point, a word of caution is necessary for the approach of this paper. Specifically, while the number D in Theorem 2.5 can still be chosen to be the number of transformations in the case of linear iterates (given that there is no dependence on h-the variable arising from the vdC-operations), in the general case the picture is quite different. By carefully tracking the constants that appear in Propositions 5.2 and 5.3, D can be chosen to be the maximum of $t_i s_i^{[t_i(s'_i+1)^{s_iL}]+1}, 1 \le i \le k$, where s'_i is the degree of \mathbf{p} , t_i is the number of terms remaining when \mathbf{p} is converted to a linear family which is 1-standard? for f_i for the first time, and s_i is t_i plus the number of vdC-operations needed to convert \mathbf{p} in such a way. (The details are left to the interested readers.)

6. Proof of Main Results

Using Theorem 2.5, we prove in this section Theorems 2.2, 2.3 and 2.4.

Proof of Theorem 2.2. We follow and adapt the proof strategy in [27, Section 3]. To avoid confusion we use $\|\cdot\|_*$ to denote the Host-Kra seminorm on the \mathbb{Z}^d -system $(X, \mathcal{B}, \mu, (T_n)_{n \in \mathbb{Z}^d})$

32 SEBASTIÁN DONOSO, ANDREU FERRÉ MORAGUES, ANDREAS KOUTSOGIANNIS AND WENBO SUN

and $\|\cdot\|'_*$ to denote the Host-Kra seminorm on the \mathbb{Z}^d -system $(X^2, \mathcal{B}^2, \mu^2, (S_n)_{n \in \mathbb{Z}^d})$, where $S_n = T_n \times T_n$. Let (I_N) be a Følner sequence in \mathbb{Z}^L . Then, by Theorem 2.5, we have

(24)
$$\lim_{N \to \infty} \frac{1}{|I_N|} \sum_{n \in I_N} \left| \int_X f_0 \cdot T_{p_1(n)} f_1 \cdot \ldots \cdot T_{p_k(n)} f_k \, d\mu \right|^2$$
$$= \lim_{N \to \infty} \frac{1}{|I_N|} \sum_{n \in I_N} \int_{X^2} f_0 \otimes \bar{f}_0 \cdot \prod_{i=1}^k S_{p_i(n)}(f_i \otimes \bar{f}_i) \, d\mu \times \mu$$
$$\leq \left\| f_i \otimes \bar{f}_i \right\|_{\{G'_{i,j}(\mathbf{p})\}_{0 \le j \le k, j \ne i}}^{\times D'}$$

for all $1 \leq i \leq k$ and some $D' \in \mathbb{N}$, where $G'_{i,j}(\mathbf{p})$ is the group action generated by S_n for all $n \in G_{i,j}(\mathbf{p})$.³⁴ Using Lemma 3.4, the right hand side of (24) is bounded by $|||f_i||^2_{\{G_{i,j}(\mathbf{p})\}_{0\leq j\leq k, j\neq i}^{\times D'}, \mathbb{Z}^d}$, which is equal to $|||f_i||^2_{(\mathbb{Z}^d)^{\times (D+1)}}$ with D = kD' by our ergodicity assumptions and Corollary 3.2.

Therefore, for $1 \leq i \leq k$ we have

(25)
$$\lim_{N \to \infty} \frac{1}{|I_N|} \sum_{n \in I_N} \int_{X^2} f_0 \otimes \bar{f}_0 \cdot \prod_{i=1}^k S_{p_i(n)}(f_i \otimes \bar{f}_i) \ d\mu \times \mu \le C |||f_i|||_{(\mathbb{Z}^d)^{\times (D+1)}}^2.$$

The bound in (25) and Theorem 3.5 imply that the sequence

(26)
$$a(n) - \int_X f_0 \cdot T_{p_1(n)} \mathbb{E}(f_1 \mid Z_{(\mathbb{Z}^d)^{\times (D+1)}}(\mathbf{X})) \cdot \ldots \cdot T_{p_k(n)} \mathbb{E}(f_k \mid Z_{(\mathbb{Z}^d)^{\times (D+1)}}(\mathbf{X})) d\mu$$

is a null sequence

is a null-sequence.

Let $\varepsilon > 0$. The factor $Z_{(\mathbb{Z}^d) \times (D+1)}(\mathbf{X})$, via Theorem 3.5, is an inverse limit of *D*-step nilsystems. Thus, there exists a factor of $Z_{(\mathbb{Z}^d) \times (D+1)}(\mathbf{X})$ with the structure of a *D*-step nilsystem $(\tilde{X}, \mathcal{B}(\tilde{X}), \mu_{\tilde{X}}, T_1, \ldots, T_d)$, on which each T_i acts as a niltranslation by an element $a_i \in \tilde{X}$, such that for $\tilde{f}_i = \mathbb{E}(f_i \mid \tilde{X})$, and $\vec{a} := (a_1, \ldots, a_d)$, we have

$$\left| \int_{X} f_0 \cdot \prod_{i=1}^k T_{p_i(n)} \mathbb{E}(f_i \mid Z_{(\mathbb{Z}^d)^{\times (D+1)}}(\mathbf{X})) \ d\mu - \int_{\tilde{X}} \tilde{f}_0 \cdot \prod_{i=1}^k \vec{a}_{p_i(n)} \tilde{f}_i \ d\mu_{\tilde{X}} \right| < \varepsilon$$

for all $n \in \mathbb{Z}^L$, where, if $p_i = (p_{i,1}, \ldots, p_{i,d})$, then $\vec{a}_{p_i(n)}$ denotes the niltranslation by the element $(a_1^{p_{1,1}(n)}, \ldots, a_d^{p_{1,d}(n)})$. Therefore, there exists a nullsequence λ such that

(27)
$$\left\| a(n) - \left(\int_{\tilde{X}} \tilde{f}_0 \cdot \vec{a}_{p_1(n)} \tilde{f}_1 \cdot \ldots \cdot \vec{a}_{p_k(n)} \tilde{f}_k \ d\mu_{\tilde{X}} + \lambda(n) \right) \right\|_{\ell^{\infty}(\mathbb{Z}^L)} < \varepsilon.$$

A standard approximation argument allows us to assume without loss of generality that $\tilde{f}_1, \ldots, \tilde{f}_k \in C(\tilde{X})$ in (27). Applying [27, Theorem 2.5] to the nilmanifold \tilde{X}^k , the diagonal subnilmanifold $\{(x, \ldots, x) : x \in \tilde{X}\}$, the polynomial sequence $(\vec{a}_{p_1(n)}, \ldots, \vec{a}_{p_k(n)})$ and the function $f(x_1, \ldots, x_k) = \tilde{f}_1(x_1) \cdot \ldots \cdot \tilde{f}_k(x_k) \in C(\tilde{X}^k)$, we obtain that the sequence

$$\psi(n) := \int_{\tilde{X}} \tilde{f}_0 \cdot \vec{a}_{p_1(n)} \tilde{f}_1 \cdot \ldots \cdot \vec{a}_{p_k(n)} \tilde{f}_k \ d\mu_{\tilde{X}}$$

³⁴Strictly speaking, $G_{i,j}(\mathbf{p})$ and $G'_{i,j}(\mathbf{p})$ are the same subgroup of \mathbb{Z}^d . We distinguish these two notions to indicate that $G_{i,j}(\mathbf{p})$ and $G'_{i,j}(\mathbf{p})$ are attached to the distinct group actions $(T_n)_{n \in G_{i,j}(\mathbf{p})}$ and $(S_n)_{n \in G'_{i,j}(\mathbf{p})}$.

is a sum of a *D*-step nilsequence and a nullsequence.

Therefore, for each $\varepsilon > 0$ we can find a *D*-step nilsequence ψ , a nullsequence λ and a bounded sequence δ with $\|\delta\|_{\ell^{\infty}(\mathbb{Z}^{L})} \leq \varepsilon$ such that

(28)
$$a(n) = \psi(n) + \lambda(n) + \delta(n).$$

For each $l \in \mathbb{N}$, consider the decomposition $a = \psi_l + \lambda_l + \delta_l$, where $\|\delta_l\|_{\ell^{\infty}(\mathbb{Z}^L)} < \frac{1}{l}$. For $r \neq l$, we have

(29)
$$|\psi_l(n) - \psi_r(n)| = |(\lambda_l(n) - \lambda_r(n)) + (\delta_l(n) - \delta_r(n))|.$$

Now, $\lim_{|I_N|\to\infty} \frac{1}{|I_N|} \sum_{n\in I_N} |\lambda_l(n) - \lambda_r(n)| = 0$ and $\sup_{n\in\mathbb{Z}^L} |\delta_r(n) - \delta_l(n)| \le \frac{1}{l} + \frac{1}{r}$. Therefore,

(30)
$$|\psi_l(n) - \psi_r(n)| \le \frac{1}{l} + \frac{1}{r}$$

for all $n \in \mathbb{Z}^L$ except potentially a subset $A \subseteq \mathbb{Z}^L$, with its characteristic function, $\mathbb{1}_A(n)$, being a nullsequence. For each $l, r \in \mathbb{N}$, the sequence $\psi_l(n) - \psi_r(n)$ is a nilsequence, so it follows that inequality (30) must, in fact, hold for all $n \in \mathbb{Z}^L$. Hence, the sequence $(\psi_l)_{l \in \mathbb{N}}$ is a Cauchy sequence in $\ell^{\infty}(\mathbb{Z}^L)$ that consists of *D*-step nilsequences, and since we already showed that $(\delta_r)_{r \in \mathbb{N}}$ is a Cauchy sequence in $\ell^{\infty}(\mathbb{Z}^L)$ converging to a nullsequence, from where the conclusion follows.

Remark 6.1. It is worth noting that if the polynomials p_1, \ldots, p_k are linear, then there is an easier proof of Theorem 2.2, where one has D = k. The reason is that, instead of Theorem 2.5, one can use [15, Proposition 1] or [6, Proposition 6.1] to improve the right hand side of (25) to $|||f_i||^2_{(\mathbb{Z}^d)\times (k+1)}$.

Next, we provide the proof of Theorem 2.4 (the arguments are similar to those in the proof of [6, Theorem 1.3, Pages 26–27]).

Proof of Theorem 2.4. Using Property (i) and Theorem 2.5, we have that there exists $D \in \mathbb{N}$ such that the left hand side of (8) is 0 if one of f_i 's satisfies $|||f_i|||_{(\mathbb{Z}^d)^{\times D}} = 0$. Thus, it suffices to show (8) under the assumption that all f_i 's are measurable with respect to $Z_{(\mathbb{Z}^d)^{\times D}}$. Using Theorem 3.5 and an approximation argument, we may assume without loss of generality that **X** is a nilsystem. As a consequence of [26, Theorem B] (see also [6, Theorem 2.9]), Property (ii) is equivalent to (8) on a nilsystem, which finishes the proof.

Finally, we prove Theorem 2.3. We start with proving that (C1) implies (C2). In fact, we show the following more general result:

Proposition 6.2. Let $d, k, L \in \mathbb{N}$, $q_1, \ldots, q_k \colon \mathbb{Z}^L \to \mathbb{Z}^d$ be polynomials, and $\mathbf{X} = (X, \mathcal{B}, \mu, (T_g)_{g \in \mathbb{Z}^d})$ be a \mathbb{Z}^d -system. Suppose that $(T_{q_1(n)}, \ldots, T_{q_k(n)})_{n \in \mathbb{Z}^L}$ is jointly ergodic for μ . Then for all $1 \leq i, j \leq k, i \neq j$, we have that $(T_{q_i(n)-q_j(n)})_{n \in \mathbb{Z}^L}$ is ergodic for μ .

Furthermore, if there exist polynomials $p_1, \ldots, p_k \colon \mathbb{Z}^L \to \mathbb{Z}$ and $v_1, \ldots, v_k \in \mathbb{Z}^d$ such that $q_i(n) = p_i(n)v_i$ for all $1 \leq i \leq k$, then $(T_{p_1(n)v_1} \times \cdots \times T_{p_k(n)v_k})_{n \in \mathbb{Z}^L}$ is ergodic for $\mu^{\otimes k}$.

Proof. The sequence $(T_{q_i(n)-q_j(n)})_{n\in\mathbb{Z}^L}$ is ergodic for μ for all $1 \leq i \neq j \leq k$ by an argument similar to the one given in the proof of [6, Proposition 5.3], so, we choose to omit the details.

We now assume that $q_i(n) = p_i(n)v_i$ for all $1 \leq i \leq k$ and show that $(T_{p_1(n)v_1} \times \cdots \times T_{p_k(n)v_k})_{n \in \mathbb{Z}^L}$ is ergodic for $\mu^{\otimes k}$. It suffices to show that for all $f_i \in L^{\infty}(\mu), 1 \leq i \leq k$ with $\prod_{i=1}^k \int_X f_i d\mu = 0$, we have that

(31)
$$\sup_{\substack{(I_N)_{N \in \mathbb{N}} \\ \text{Følner seq.}}} \overline{\lim_{N \to \infty}} \left\| \mathbb{E}_{n \in I_N} \bigotimes_{i=1}^k T_{p_i(n)v_i} f_i \right\|_{L^2(\mu^{\otimes k})} = 0.$$

Claim. If $\mathbb{E}(f_i | Z_{\mathbb{Z}^d, \mathbb{Z}^d}(\mathbf{X})) = 0$ for some $1 \le i \le k$, then (31) holds.

We may assume without loss of generality that $\deg(p_1) \ge \deg(p_2) \ge \cdots \ge \deg(p_k)$. Suppose that we have shown that for some $1 \le k_0 \le k$, (31) holds if $\mathbb{E}(f_i | Z_{\mathbb{Z}^d, \mathbb{Z}^d}(\mathbf{X})) = 0$ for some $1 \le i \le k_0 - 1$, where the case $k_0 = 1$ is understood to be always true. It suffices to show that (31) holds if $\mathbb{E}(f_{k_0} | Z_{\mathbb{Z}^d, \mathbb{Z}^d}(\mathbf{X})) = 0$.

By the induction hypothesis, we may assume without loss of generality that f_i is $Z_{\mathbb{Z}^d,\mathbb{Z}^d}(\mathbf{X})$ measurable for all $1 \leq i \leq k_0 - 1$. By [6, Lemma 2.7], we can approximate each f_i in $L^2(\mu)$ by an eigenfunction of \mathbf{X} . By multi-linearity, we may assume without loss of generality that each $f_i, 1 \leq i \leq k_0 - 1$ is a non-constant eigenfunction of \mathbf{X} given by $T_n f_i = \exp(\lambda_i(n)) f_i$ for all $n \in \mathbb{Z}^d$ and some group homomorphism $\lambda_i \colon \mathbb{Z}^d \to \mathbb{R}$, and that $f_i(x) \neq 0$ μ -a.e $x \in X$. Then the left hand side of (31) is equal to

(32)
$$\sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ \text{Følner seq.}}} \overline{\lim_{N\to\infty}} \left\| \mathbb{E}_{n\in I_N} \exp(P(n)) \bigotimes_{i=1}^{k_0-1} f_i \bigotimes_{i=k_0}^k T_{p_i(n)v_i} f_i \right\|_{L^2(\mu^{\otimes k})}$$

where $P(n) \coloneqq \sum_{i=1}^{k_0-1} \lambda_i(v_i) p_i(n)$. Denote $P(n) = \sum_{j=0}^{\deg(p_1)} Q_j(n)$, where Q_j is a homogeneous polynomial of degree j for all $0 \le j \le \deg(p_1)$. Then

(33)
$$\Delta^{K} P(n, h_1, \dots, h_K) = \sum_{j=K}^{\deg(p_1)} \Delta^{K} Q_j(n, h_1, \dots, h_K).$$

We first consider the case when $Q_j(n) \notin \mathbb{Q}[n] + \mathbb{R}$ for some $\deg(p_{k_0}) + 1 \leq j \leq \deg(p_1)$. In this case, let $K = \deg(p_{k_0})$ in (33). Since $\Delta^K p_i(n, h_1, \ldots, h_K)$ is constant in n for all $k_0 \leq i \leq k$, by Lemma 4.2, to show that (32) is 0, it suffices to show that

(34)
$$\overline{\mathbb{E}}_{h_1,\dots,h_K\in\mathbb{Z}^L}^{\sqcup} \sup_{\substack{(I_N)_{N\in\mathbb{N}}\\\text{Folner seq.}}} \overline{\lim_{N\to\infty}} |\mathbb{E}_{n\in I_N}\exp(\Delta^K P(n,h_1,\dots,h_K))| = 0$$

(see Definition 3.13 for the definition of the polynomial $\Delta^{K} P$).

As $Q_j(n) \notin \mathbb{Q}[n] + \mathbb{R}$ for some $\deg(p_{k_0}) + 1 \leq j \leq \deg(p_1)$, Lemma 3.14 implies that $\Delta^K Q_j(\cdot, h_1, \ldots, h_K) \notin \mathbb{Q}[n] + \mathbb{R}$ for a set of (h_1, \ldots, h_K) of density 1. By Weyl's criterion and (33), we have that (34) holds and thus (31) holds.

We now consider the case when $Q_j(n) \in \mathbb{Q}[n] + \mathbb{R}$ for all $K + 1 \leq j \leq \deg(p_1)$. Let $P'(n) = \sum_{j=0}^{K} Q_j(n)$. It is not hard to see that there exists $Q \in \mathbb{N}$ such that for all $r \in \{0, \ldots, Q-1\}^L$ and $n \in \mathbb{Z}^L$, we have that

$$P(Qn + r) - P'(Qn + r) = P(r) - P'(r).$$

DECOMPOSITION OF MULTICORRELATION SEQUENCES AND JOINT ERGODICITY

By (32), to show (31), it suffices to show that for all $r \in \{0, \ldots, Q-1\}^L$, we have that

(35)
$$\sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ \text{Følner seq.}}} \overline{\lim_{N\to\infty}} \left\| \mathbb{E}_{n\in I_N} \exp(P'(Qn+r)) \bigotimes_{i=k_0}^k T_{p_i(Qn+r)v_i} f_i \right\|_{L^2(\mu^{\otimes t})} = 0,$$

where $t = k - k_0 + 1$. Fix $r \in \{0, \dots, Q-1\}^L$ and set R(n) = P'(Qn + r). Let $p: \mathbb{Z}^L \to \mathbb{Z}^{dt}$ be the polynomial given by

$$p(n) = (p_i(Qn+r)v_i)_{k_0 \le i \le k}.$$

Let $(X^t, \mathcal{B}^t, \mu^t, (S_g)_{g \in \mathbb{Z}^{dt}})$ be the \mathbb{Z}^{dt} -system such that

$$S_{(u_i)_{k_0 \le i \le k}} \coloneqq \prod_{i=k_0}^{\kappa} T_{u_i}$$

for all $u_i \in \mathbb{Z}^d$, $k_0 \leq i \leq k$, and denote $f := \bigotimes_{i=k_0}^k f_i$. We may then rewrite the left hand side of (35) as

(36)
$$\sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ \text{Følner seq.}}} \overline{\lim_{N\to\infty}} \|\mathbb{E}_{n\in I_N} \exp(R(n))S_{p(n)}f\|_{L^2(\mu^{\otimes t})}.$$

For $K = \deg(p_{k_0}) - 1$, using Lemma 4.2, to show that (36) is zero, it suffices to show

(37)
$$\overline{\mathbb{E}}_{\mathbf{h}=(h_1,\dots,h_K)\in(\mathbb{Z}^L)^K}^{\square} \sup_{\substack{(I_N)_{N\in\mathbb{N}}\\ \text{Følner seq.}}} \lim_{N\to\infty} \|\mathbb{E}_{n\in I_N}\exp(\Delta^K R(n,\mathbf{h}))S_{\Delta^K p(n,\mathbf{h})}f\|_{L^2(\mu^{\otimes t})} = 0.$$

By assumption, $\Delta^{K} R(n, h_1, \ldots, h_K)$ is of degree 1 in the variable n. Since $\deg(p) = \deg(p_{k_0}) \geq \deg(p_i)$ for all $i \geq k_0$, $\Delta^{K} p(n, h_1, \ldots, h_K)$ is also of degree 1 in the variable n. We may thus assume that

$$\Delta^{K} p(n, h_{1}, \dots, h_{K}) = ((c_{i}(h_{1}, \dots, h_{K}) \cdot n + c_{i}'(h_{1}, \dots, h_{K}))v_{i})_{k_{0} \le i \le k_{1}}$$

for some polynomials $c_{k_0}, \ldots, c_k \colon \mathbb{Z}^{LK} \to \mathbb{Z}^L$ and $c'_{k_0}, \ldots, c'_k \colon \mathbb{Z}^{LK} \to \mathbb{Z}$. Write $\mathbf{c}(h_1, \ldots, h_K) \coloneqq (c_i(h_1, \ldots, h_K)v_i)_{k_0 \leq i \leq k}$ (which is viewed as a *t*-tuple of *L*-tuple of vectors in \mathbb{Z}^d). If we write

$$c_i(h_1, \ldots, h_K) = (c_{i,1}(h_1, \ldots, h_K), \ldots, c_{i,L}(h_1, \ldots, h_K))$$

for some $c_{i,j}(h_1,\ldots,h_K) \in \mathbb{Z}$, then, by definition, $G(\mathbf{c}(h_1,\ldots,c_K))$ is the subgroup of \mathbb{Z}^{dt} generated by the elements

$$(c_{k_0,j}(h_1,\ldots,h_K)v_{k_0},\ldots,c_{k,j}(h_1,\ldots,h_K)v_k), 1 \le j \le L.$$

By Lemma 4.3, the left hand side of (37) is bounded by a constant multiple of

(38)
$$(\overline{\mathbb{E}}_{h_1,\dots,h_K \in \mathbb{Z}^L}^{\square} \| f_{k_0} \|_{G(c_{k_0}(h_1,\dots,c_K)v_{k_0})^{\times 2}}^{1})^{1/4}$$

where $G(c_{k_0}(h_1,\ldots,c_K)v_{k_0})$ is the subgroup of \mathbb{Z}^d generated by the elements

$$c_{k_0,1}(h_1,\ldots,h_K)v_{k_0},\ldots,c_{k_0,L}(h_1,\ldots,h_K)v_{k_0}$$

i.e., the entries of $c_{k_0}(h_1, \ldots, h_K)v_{k_0}$. For any $u_{k_0} \in G(c_{k_0}(h_1, \ldots, h_K)v_{k_0})$, note that u_{k_0} is a rational multiple of v_{k_0} . So, if $c_{k_0}(h_1, \ldots, h_K) \neq \mathbf{0}$, then $G(c_{k_0}(h_1, \ldots, h_K)v_{k_0}) = G(v_{k_0})$.

36 SEBASTIÁN DONOSO, ANDREU FERRÉ MORAGUES, ANDREAS KOUTSOGIANNIS AND WENBO SUN

Since $(T_{p_i(n)v_i})_{n\in\mathbb{Z}^L}$ is ergodic for μ , we have that T_{v_i} is ergodic for μ . As $\mathbb{E}(f_{k_0}|Z_{\mathbb{Z}^d,\mathbb{Z}^d}(\mathbf{X})) = 0$, by [6, Lemma 2.4], we have that

$$|\!|\!| f_{k_0} |\!|\!|_{G(c_{k_0}(h_1,\ldots,c_K)v_{k_0})^{\times 2}} = |\!|\!| f_{k_0} |\!|\!|_{v_{k_0}^{\times 2}} = |\!|\!| f_{k_0} |\!|\!|_{(\mathbb{Z}^d)^{\times 2}} = 0$$

whenever $c_{k_0}(h_1, \ldots, h_K) \neq \mathbf{0}$. Since $K = \deg(p_{k_0}) - 1 = \deg(p_{k_0}(Q \cdot + r)) - 1$, it is easy to see that $c_{k_0} \neq \mathbf{0}$. By [6, Lemma 2.12], the set of such (h_1, \ldots, h_K) is of density 1. So, averaging over all $h_1, \ldots, h_K \in \mathbb{Z}^L$, we have that (38) is 0. This finishes the proof of the claim.

Using the claim, it suffices to prove (31) under the assumption that all f_i 's are measurable with respect to $Z_{(\mathbb{Z}^d)^{\times 2}}(\mathbf{X})$. By Lemma 2.7 of [6], we can approximate each f_i in $L^2(\mu)$ by an eigenfunction of \mathbf{X} . By multi-linearity, we may assume without loss of generality that each f_i is a non-constant eigenfunction of \mathbf{X} given by $T_n f_i = \exp(\lambda_i(n)) f_i$ for all $n \in \mathbb{Z}^d$ for some group homomorphism $\lambda_i \colon \mathbb{Z}^d \to \mathbb{R}$, and that $f_i(x) \neq 0$ μ -a.e. $x \in X$. Then, since $(T_{p_i(n)v_i} \colon 1 \leq i \leq k)_{n \in \mathbb{Z}^L}$ is jointly ergodic for μ , for any Følner sequence $(I_N)_{N \in \mathbb{N}}$ of \mathbb{Z}^d ,

$$0 = \prod_{i=1}^{k} \int_{X} f_{i} d\mu = \lim_{N \to \infty} \mathbb{E}_{n \in I_{N}} \prod_{i=1}^{k} T_{p_{i}(n)v_{i}} f_{i} = \left(\lim_{N \to \infty} \mathbb{E}_{n \in I_{N}} \prod_{i=1}^{k} \exp(\lambda_{i}(p_{i}(n)v_{i}))\right) \prod_{i=1}^{k} f_{i}.$$

This implies that $\lim_{N\to\infty} \mathbb{E}_{n\in I_N} \prod_{i=1}^k \exp(\lambda_i(p_i(n)v_i)) = 0$. So,

$$\lim_{N \to \infty} \mathbb{E}_{n \in I_N} \bigotimes_{i=1}^k T_{p_i(n)v_i} f_i = \left(\lim_{N \to \infty} \mathbb{E}_{n \in I_N} \prod_{i=1}^k \exp(\lambda_i(p_i(n)v_i)) \right) \bigotimes_{i=1}^k f_i.$$

is the proof.

This finishes the proof.

We are now ready to complete the proof of Theorem 2.3.

Proof of Theorem 2.3. Using Proposition 6.2, we have that (C1) implies (C2). It is obvious that (C2) implies (C2'). So, it suffices to show that (C2') implies (C1).

It is not hard to see that we may assume without loss of generality that $p_i(0) = 0$ for all $1 \leq i \leq k$. By Theorem 2.4, to show that $(T_{p_i(n)v_i}: 1 \leq i \leq k)_{n \in \mathbb{Z}^L}$ is jointly ergodic for μ , it suffices to show that $G_{i,j}(\mathbf{p})$ is ergodic for μ for all $0 \leq i, j \leq k, i \neq j$. Fix any such pair (i, j). We may assume without loss of generality that $i \neq 0$. If j = 0, then by (ii), $(T_{p_i(n)v_i})_{n \in \mathbb{Z}^L}$ is ergodic for μ , and thus $G_{i,0}(\mathbf{p}) = G(v_i)$ is ergodic for μ . So, we may now assume that $j \neq 0$.

Assume first that $\deg(p_i) = \deg(p_j)$. By assumption, either v_i and v_j are linearly dependent, or $p_i(n)$ and $p_j(n)$ are linearly dependent.

If v_i and v_j are linearly dependent over \mathbb{Z} , then we may assume without loss of generality that $v_i = av$ and $v_j = bv$ for some $a, b \in \mathbb{Q}$ and $v \in \mathbb{Z}^d$. By (i), $(T_{(ap_i(n)-bp_j(n))v})_{n\in\mathbb{Z}^L}$ is ergodic for μ , which implies that G(v) is ergodic for μ . On the other hand, $G_{i,j}(\mathbf{p})$ is a group generated by some elements which are linear combinations of v_i and v_j , which are thus multiples of v. Since \mathbf{p} is non-degenerate, $G_{i,j}(\mathbf{p})$ is not the trivial group. It follows that $G_{i,j}(\mathbf{p}) = G(v)$, so the group $G_{i,j}(\mathbf{p})$ is ergodic for μ .

If $p_i(n)$ and $p_j(n)$ are linearly dependent over \mathbb{Z} , then we may assume without loss of generality that $p_i(n) = ap(n)$ and $p_j(n) = bp(n)$ for some $a, b \in \mathbb{Q}$ and polynomial p. By (i), $(T_{(p(n)(av_i - bv_j))_{n \in \mathbb{Z}^L}}$ is ergodic for μ , which implies that $G(av_i - bv_j)$ is ergodic for μ . On the other hand, $G_{i,j}(\mathbf{p})$ is a group generated by some elements which are multiples of

 $av_i - bv_j$. Since **p** is non-degenerate, $G_{i,j}(\mathbf{p})$ is not the trivial group. It follows that $G_{i,j}(\mathbf{p}) = G(av_i - bv_j)$, so the group $G_{i,j}(\mathbf{p})$ is ergodic for μ .

Finally, we consider the case when $\deg(p_i) \neq \deg(p_j)$. We may assume without loss of generality that $\deg(p_i) > \deg(p_j)$. In this case, $G_{i,j}(\mathbf{p}) = G_{i,0}(\mathbf{p})$, which we have shown is ergodic for μ .

7. POTENTIAL FUTURE DIRECTIONS

We close this article with two potential future directions regarding the splitting of multicorrelation sequences. The first one is for integer polynomial iterates under no assumptions on the transformations other than commutativity (see Theorem 7.1 for a special case of two terms).

The second one pertains to potential results analogous to Theorem 2.2 for iterates of the form $[p_i(n)]$, $1 \leq i \leq k$, where $p_i = (p_{i,1}, \ldots, p_{i,d}) : \mathbb{Z}^L \to \mathbb{R}^d$ are vectors of real valued polynomials.³⁵

7.1. The two-term case with no ergodicity assumptions. Given the results in the appendix of [5], we are able to obtain the following splitting result for two commuting transformations without any ergodicity assumptions:

Theorem 7.1. Let $(X, \mathcal{B}, \mu, T, S)$ be a measure preserving system with TS = ST. Let $f_0, f_1, f_2 \in L^{\infty}(\mu)$ and $p \in \mathbb{Z}[n]$ with degree $K \geq 2$. Then, the multicorrelation sequence

$$a(n) \coloneqq \int_X f_0 \cdot T^n f_1 \cdot S^{p(n)} f_2 \ d\mu$$

can be decomposed as a sum of a uniform limit of K-step nilsequences plus a nullsequence.

Proof. Setting $F_i = f_i \otimes \overline{f}_i$, i = 0, 1, 2, and $\tilde{\mu} = \mu \times \mu$, we have that

(39)
$$\frac{1}{N}\sum_{n=1}^{N}|a(n)|^{2} = \frac{1}{N}\sum_{n=1}^{N}\int_{X^{2}}F_{0}\cdot(T\times T)^{n}F_{1}\cdot(S\times S)^{p(n)}F_{2}\ d\tilde{\mu}.$$

Using [5, Theorem A.3], we get that the rational Kronecker factor³⁶ is characteristic for the averages appearing in (39). Consequently, we may replace f_1 by $P_c f_1$ and f_2 by $Q_c f_2$ in a(n) up to a nullsequence, where P_c denotes the orthogonal projection onto the compact component of the splitting associated to T, and Q_c the one associated to S. (Here we make use of the Hilbert space splitting of $L^2(\mu)$ into its compact and weakly mixing components for a given unitary operator. The seeds for these results are already present in the work of Koopman and von Neumann [21]. They were later generalized by Jacobs, Glicksberg and de Leeuw. See [7, Section 16.3] for a more modern treatment). Thus, the sequence

$$a(n) - \int_X f_0 \cdot T^n P_c f_1 \cdot S^{p(n)} Q_c f_2 \ d\mu$$

³⁵Here, for $x = (x_1, \ldots, x_L) \in \mathbb{R}^L$, we write $[x] := ([x_1], \ldots, [x_L])$, where $[\cdot]$ is the floor function. In fact, one can consider any combination of rounding functions, i.e., floor, ceiling, or closest integer.

³⁶The rational Kronecker factor is the smallest sub- σ -algebra of \mathcal{B} that makes all functions with finite orbit in $L^2(\mu)$ under the transformation T measurable.

is a nullsequence. Let $\varepsilon > 0$ and choose $h_1, \ldots, h_k, g_1, \ldots, g_k \in L^2(\mu)$ such that $Th_i = \lambda_i h_i$ and $Sg_i = \rho_i g_i$ (for some $\lambda_1, \ldots, \lambda_k, \rho_1, \ldots, \rho_k \in \mathbb{C}$ of absolute value one) as well as $a_1, \ldots, a_k, b_1, \ldots, b_k \in \mathbb{C}$ such that

$$\left| \int_X f_0 \cdot T^n P_c f_1 \cdot S^{p(n)} Q_c f_2 \, d\mu - \int_X f_0 \cdot T^n \sum_{i=1}^k a_i h_i \cdot S^{p(n)} \sum_{j=1}^k b_j g_j \, d\mu \right| < \varepsilon.$$

Observe that

$$\int_{X} f_{0} \cdot T^{n} \sum_{i=1}^{k} a_{i} h_{i} \cdot S^{p(n)} \sum_{j=1}^{k} b_{i} g_{i} d\mu = \int_{X} f_{0} \cdot \sum_{i=1}^{k} a_{i} \lambda_{i}^{n} h_{i} \cdot \sum_{j=1}^{k} b_{j} \rho_{j}^{p(n)} g_{j} d\mu$$
$$= \sum_{i,j=1}^{k} \left(a_{i} b_{j} \int_{X} f_{0} \cdot h_{i} \cdot g_{j} d\mu \right) \lambda_{i}^{n} \rho_{j}^{p(n)},$$

which is a K-step nilsequence. Applying the same argument as in the proof Theorem 2.2 we deduce the decomposition result. The rest of the details are omitted for the sake of brevity. \Box

It is natural to ask whether a result analogous to Theorem 7.1 holds for longer expressions (potentially via a generalization of the results in the appendix of [5]), and with more general polynomial iterates, even without necessarily assuming they have distinct degrees. Thus, we state the following problem:

Problem 1. Obtain decomposition results of the form "uniform limit of nilsequences plus a nullsequence" for multicorrelation sequences with (integer) polynomial iterates for general systems under no ergodicity assumptions on the transformations.

7.2. Integer part polynomial iterates. With a, by now, standard argument (introduced in [4] and [29] for a single term, extended for two terms in [34], and further developed in [22], [23] and [25]) one has, for the vectors of real polynomials $p_i = (p_{i,1}, \ldots, p_{i,d})$, that the expression

(40)
$$\frac{1}{|I_N|} \sum_{n \in I_N} \prod_{i=1}^k T_{[p_i(n)]} f_i = \frac{1}{|I_N|} \sum_{n \in I_N} \prod_{i=1}^k \prod_{j=1}^d T_j^{[p_{i,j}(n)]} f_i$$

is "close" to

(41)
$$\frac{1}{|I_N|} \sum_{n \in I_N} \prod_{i=1}^k \prod_{j=1}^d S_j^{p_{i,j}(n)} g_i$$

where S_j 's are \mathbb{R} -flows on an "extension system" Y of X, and the functions g_i 's are extensions of the f_i 's (see [23] for details).³⁷ As an application of Theorem 2.2, one can prove splitting theorems for \mathbb{R}^d -actions on the extension system.

³⁷We say that a jointly measurable family $(S_t)_{t \in \mathbb{R}^d}$ of measure preserving transformations on a probability space, is an \mathbb{R}^d -action (flow), if it satisfies $S_{t+r} = S_t \circ S_r$ for all $t, r \in \mathbb{R}^d$.

Indeed,³⁸ consider the multicorrelation sequence

(42)
$$\int_X f_0 \cdot S_{p_1(n)} f_1 \cdot \ldots \cdot S_{p_k(n)} f_k \, d\mu,$$

where S is a measure preserving \mathbb{R}^d -action on the probability space (X, \mathcal{B}, μ) , $f_0, f_1, \ldots, f_k \in L^{\infty}(X)$, and $p_1, \ldots, p_k \colon \mathbb{Z} \to \mathbb{R}^d$ a non-degenerate family of polynomials of degree at most K with $p_i(n) = \sum_{h=0}^{K} a_{i,h}n^h$, $a_{i,h} \in \mathbb{R}^d$. Then

$$S_{p_i(n)} = S_{\sum_{h=0}^{K} a_{i,h}n^h} = \prod_{h=0}^{K} (S_{a_{i,h}})^{n^h}.$$

Note that $S_{a_{i,h}}, 1 \leq i \leq k, 1 \leq h \leq K$ generate a \mathbb{Z}^{kK} -action on (X, \mathcal{B}, μ) . For convenience, set p_0 to be the constant zero polynomial. For $0 \leq i, j \leq k, i \neq j$, let $D_{i,j}$ be the largest integer h such that $S_{a_{i,D_{i,j}}-a_{j,D_{i,j}}} \neq id$. This transformation will be denoted by $R_{i,j}$. By Theorem 2.2, one can show that the desired splitting result for the sequence (42), if all the transformations $R_{i,j}, 0 \leq i, j \leq k, i \neq j$ are all ergodic (as \mathbb{Z} -actions on the extension system Y).

Unfortunately, even though we have the previous result for flows, the error term that arises from the approximation of (40) by (41) prevents us from getting the conclusion of Theorem 2.2 for multicorrelation sequences of the form

$$\int_X f_0 \cdot T_{[p_1(n)]} f_1 \cdot \ldots \cdot T_{[p_k(n)]} f_k \ d\mu.^{39}$$

Remark 7.2. It is important to stress that, for integer part real polynomial iterates, one does not expect to have the desired multicorrelation splitting in general. The next example shows that, even for k = 1, an ergodic system, and linear iterates, it can be too much to hope for:

Indeed, following [25, Example 7], let $X = \mathbb{T} := \mathbb{R}/\mathbb{Z}$, $T(x) = x + 1/\sqrt{2}$, $p(n) = \sqrt{2n}$, $f_0(x) = e(x)$ and $f_1(x) = e(-x)$, where $e(x) := e^{2\pi i x}$. Then, we have that

$$\int f_0 \cdot T^{[p(n)]} f_1 d\mu = \int e(x) e\left(-x - \frac{1}{\sqrt{2}} [\sqrt{2}n]\right) dx = e\left(-\frac{1}{\sqrt{2}} [\sqrt{2}n]\right) = e\left(\frac{1}{\sqrt{2}} \{\sqrt{2}n\}\right),$$

which cannot be written as a uniform limit of nilsequences and a nullsequence.

Remark 7.3. One may think that the fact that $\sqrt{2}$ and $\frac{1}{\sqrt{2}}$ are not linearly independent over \mathbb{Q} is behind the impossibility of the splitting in the example above. However, a closer examination of the proof given in [25] shows that this is not the case, and that the failure extends quite generally.

Indeed, we can imitate the example quoted above as follows. Let (X, \mathcal{B}, μ, T) be an ergodic measure preserving system with non-trivial irrational spectrum. Let $f_1 : X \to \mathbb{S}^1$ be an

³⁸We address the L = 1 case for simplicity; following the same argument one can similarly get the corresponding result for the general case of *L*-variable polynomials by using an ordering on the parameters, e.g., $n_1 > \ldots > n_L$.

³⁹To this day, only splittings of the form nilsequence plus an error term that is small in uniform density are known for this class of multicorrelation sequences (for this, see [23]. One is referred to [25] for averages along primes for the error term–in this last reference only single variable real polynomials were considered. Using the multivariable approach of [12] instead of [11], one immediately gets the aforementioned result for integer part, or indeed for combinations of any other rounding functions of multivariable real polynomial iterates).

eigenfunction of T with eigenvalue $e^{2\pi i\beta}$, with $\beta \in \mathbb{R} \setminus \mathbb{Q}$. Put $f_0 = \overline{f}_1$. Then, if we consider the multicorrelation sequence

$$a(n) \coloneqq \int_X f_0 \cdot T^{[\alpha n]} f_1 \ d\mu$$

with the choices made above, we observe that, in fact, $a(n) = e^{2\pi i [\alpha n]\beta}$. The same argument as in [25, Example 7] shows that a(n) cannot be written as a uniform limit of nilsequences plus a nullsequence.

On the other hand, if we postulate very strong assumptions on our transformations, we do have the desired decomposition results. For example (see [23]), if T_1, \ldots, T_k are commuting weakly mixing transformations on (X, \mathcal{B}, μ) , $q_i(n) = p_i(n)e_i$, $1 \le i \le k$, where $p_i : \mathbb{Z} \to \mathbb{R}$ are real polynomials of distinct, positive degrees, and $f_1, \ldots, f_k \in L^{\infty}(\mu)$, then we have that

$$\lim_{N-M\to\infty} \frac{1}{N-M} \sum_{n=M}^{N-1} T_{[q_1(n)]} f_1 \cdot \ldots \cdot T_{[q_k(n)]} f_k = \prod_{i=1}^k \int_X f_i \ d\mu.$$

Hence, for any $f_0 \in L^{\infty}(\mu)$, the multicorrelation sequence

$$\int_X f_0 \cdot T_{[q_1(n)]} f_1 \cdot \ldots \cdot T_{[q_k(n)]} f_k \ d\mu$$

can be written as a sum of a constant (i.e., a 0-step nilsequence) and a nullsequence.

We conclude this article with the following problem that arises naturally:

Problem 2. Let $d, k, K, L \in \mathbb{N}$, $p_1, \ldots, p_k \colon \mathbb{Z}^L \to \mathbb{R}^d$ be a non-degenerate family of polynomials of degree at most K, $(X, \mathcal{B}, \mu, T_1, \ldots, T_d)$ a measure preserving system and $f_0, \ldots, f_k \in L^{\infty}(\mu)$. Find conditions, on the p_i 's and/or the \mathbb{Z}^d -action T that is defined by the T_i 's, so that the multicorrelation sequence

$$\int_X f_0 \cdot T_{[p_1(n)]} f_1 \cdot \ldots \cdot T_{[p_k(n)]} f_k \ d\mu$$

can be decomposed as a sum of a uniform limit of *D*-step nilsequences and a nullsequence.

References

- D. Berend and V. Bergelson. Jointly ergodic measure-preserving transformations. Israel J. Math. 49 (1984), no. 4, 307–314.
- [2] V. Bergelson. Weakly mixing PET. Ergodic Theory Dynam. Systems 7 (1987), no. 3, 337–349.
- [3] V. Bergelson, B. Host and B. Kra, with an appendix by I. Ruzsa. Multiple recurrence and nilsequences, *Invent. Math.* 160 (2005), no. 2, 261–303.
- [4] M. Boshernitzan, R. L. Jones and M. Wierdl. Integer and fractional parts of good averaging sequences in ergodic theory. *Convergence in Ergodic Theory and Probability*, Eds: V. Bergelson, P. March, J. Rosenblatt, by Walter de Gruyter & Co., Berlin, New York 1996, 117–132.
- [5] Q. Chu, N. Frantzikinakis and B. Host. Ergodic averages of commuting transformations with distinct degree polynomial iterates. Proc. Lond. Math. Soc. (3) 102 (2011), no. 5, 801–842.
- [6] S. Donoso, A. Koutsogiannis and W. Sun. Seminorms for multiple averages along polynomials and applications to joint ergodicity. To appear in *J. d'Analyse Mathématique*.
- [7] T. Eisner, B. Farkas, M. Haase and R. Nagel. Operator theoretic aspects of ergodic theory, Graduate Texts in Mathematics, 272, Springer, Cham, 2015.
- [8] A. F. Moragues, Properties of multicorrelation sequences and large returns under some ergodicity assumptions, Discrete Contin. Dyn. Syst. 41 (2021), no. 6, 2809–2828.

- [9] N. Frantzikinakis. Joint ergodicity of sequences. Preprint 2021, arXiv:2102.09967.
- [10] N. Frantzikinakis, B. Host and B. Kra. The polynomial multidimensional Szemerédi theorem along shifted primes. Israel J. Math. 194 (2013), no. 1, 331–348.
- [11] N. Frantzikinakis. Multiple correlation sequences and nilsequences. Invent. Math. 202 (2015), no. 2, 875–892.
- [12] N. Frantzikinakis and B. Host. Weighted multiple ergodic averages and correlation sequences, Ergodic Theory Dynam. Systems 38 (2018), no. 1, 81–142.
- [13] N. Frantzikinakis and B. Kra. Polynomial averages converge to the product of integrals. Israel J. Math. 148 (2005), 267–276.
- [14] H. Furstenberg. Ergodic behavior of diagonal measures and a theorem of Szemerédi on arithmetic progressions. J. d'Analyse Mathématique 31 (1977), 204–256.
- B. Host. Ergodic seminorms for commuting transformations and applications. Studia Math. 195 (2009), no. 1, 31–49.
- [16] B. Host and B. Kra. Nonconventional ergodic averages and nilmanifolds. Ann. of Math. (2) 161 (2005), no. 1, 397–488.
- [17] B. Host and B. Kra. Uniformity seminorms on ℓ^{∞} and applications, J. Anal. Math. 108 (2009), 219–276.
- [18] M. C. R. Johnson. Convergence of polynomial ergodic averages of several variables for some commuting transformations. *Illinois J. Math.* 53 (2009), no. 3, 865–882 (2010).
- [19] D. Karageorgos and A. Koutsogiannis. Integer part independent polynomial averages and applications along primes. *Studia Mathematica* 249 (2019), no. 3, 233–257.
- [20] A. Khintchine. The Method of Spectral Reduction in Classical Dynamics. Proceedings of the National Academy of Sciences May 1933, 19 (5) 567-573.
- [21] B. O. Koopman and J. von Neumann. Dynamical Systems of Continuous Spectra. Proceedings of the National Academy of Sciences Mar 1932, 18 (3) 255-263.
- [22] A. Koutsogiannis. Closest integer polynomial multiple recurrence along shifted primes. Ergodic Theory Dynam. Systems 38 (2018), no. 2, 666–685.
- [23] A. Koutsogiannis. Integer part polynomial correlation sequences. Ergodic Theory Dynam. Systems 38 (2018), no. 4, 1525–1542.
- [24] A. Koutsogiannis. Multiple ergodic averages for variable polynomials. Preprint 2021, arXiv:2101.00534.
- [25] A. Koutsogiannis, A. Le, J. Moreira and F. K. Richter. Structure of multicorrelation sequences with integer part polynomial iterates along the primes. *Proceedings of the A. M. S.* 149 (2021), No 1, 209–216.
- [26] A. Leibman. Convergence of multiple ergodic averages along polynomials of several variables. Israel J. Math. 146 (2005), 303–315.
- [27] A. Leibman. Multiple polynomial correlation sequences and nilsequences. Ergodic Theory Dynam. Systems 30 (2010), no. 3, 841–854.
- [28] A. Leibman. Nilsequences, null-sequences, and multiple correlation sequences. Ergodic Theory Dynam. Systems 35 (2015), no. 1, 176–191.
- [29] E. Lesigne. On the sequence of integer parts of a good sequence for the ergodic theorem. Comment. Math. Univ. Carolin. 36 (1995), no. 4, 737–743.
- [30] W. Sun. Weak ergodic averages over dilated curves. Ergodic Theory Dynam. Systems 41 (2021), No 2, 606–621.
- [31] T. Tao and J. Teräväinen. The structure of logarithmically averaged correlations of multiplicative functions, with applications to the Chowla and Elliott conjectures. *Duke Math. J.*, 168 (2019), No 11, 1977–2027.
- [32] T. Tao and T. Ziegler. Concatenation Theorems for anti-Gowers-uniform functions and Host-Kra characteristic factors. *Discrete Anal.* 2016, Paper No. 13.
- [33] M. Walsh. Norm convergence of nilpotent ergodic averages. Ann. of Math. (2) 175 (2012), no. 3, 1667– 1688.
- [34] M. Wierdl. Personal communication with the third author (2015).
- [35] T. Ziegler. Nilfactors of \mathbb{R}^m -actions and configurations in sets of positive upper density in \mathbb{R}^m . J. d'Analyse Mathématique **99** (2006), 249–266.

42 SEBASTIÁN DONOSO, ANDREU FERRÉ MORAGUES, ANDREAS KOUTSOGIANNIS AND WENBO SUN

(Sebastián Donoso) DEPARTAMENTO DE INGENIERÍA MATEMÁTICA AND CENTRO DE MODELAMIENTO MATEMÁTICO, UNIVERSIDAD DE CHILE & IRL 2807 - CNRS, BEAUCHEF 851, SANTIAGO, CHILE. *Email address:* sdonoso@dim.uchile.cl

(Andreu Ferré Moragues) DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, 231 WEST 18th Avenue, Columbus, OH 43210-1174, USA *Email address:* ferremoragues.1@osu.edu

(Andreas Koutsogiannis) DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, 231 WEST 18TH AVENUE, COLUMBUS, OH 43210-1174, USA

 $Email \ address: \verb"koutsogiannis.1@osu.edu"$

 (Wenbo Sun) Department of Mathematics, Virginia Tech, 902 Prices Fork RD, Blacksburg, VA, 24061, USA

 $Email \ address: \ {\tt swenbo} {\tt @vt.edu}$