

Improved Spanning on Theta-5

Prosenjit Bose^{*†} Darryl Hill^{‡†} Aurélien Ooms^{§¶}

June 3, 2021

Abstract

We show an upper bound of $\frac{\sin(\frac{3\pi}{10})}{\sin(\frac{2\pi}{5}) - \sin(\frac{3\pi}{10})} < 5.70$ on the spanning ratio of Θ_5 -graphs, improving on the previous best known upper bound of 9.96 [Bose, Morin, van Renssen, and Verdonschot. The Theta-5-graph is a spanner. *Computational Geometry*, 2015.] **Keywords:** Theta Graphs Spanning Ratio Stretch Factor Geometric Spanners.

1 Introduction

A geometric graph G is a graph whose vertex set is a set of points P in the plane, and where the weight of an edge uv is equal to the Euclidean distance $|uv|$ between u and v . Informally, a Θ_k -graph is a geometric graph built by dividing the area around each point of $v \in P$ into k equal angled cones, connecting v to the *closest* neighbor in each cone (we shall define closest later). Such graphs arise naturally in settings like wireless networks, where signals to anyone but your nearest neighbor may be drowned out by interference. Moreover, the fact that signal strength fades quadratically with distance, and thus that power requirements are proportional to the square of the distance the signal has to travel, makes many small hops economically superior to one large hop, even if the sum of the distances is larger. The *spanning ratio* (sometimes called the *stretch factor*) of a geometric graph G is the maximum over all pairs $u, v \in P$ of the ratio between the length of the shortest path from u to v in G and the Euclidean distance from u to v . Using simple geometric observations and techniques, we give a new analysis of the spanning ratio of Θ_5 -graphs, bringing down the best known upper bound from 9.96 [5] to 5.70.

Theorem 1. *Given a set P of points in the plane, the Θ_5 -graph of P is a 5.70-spanner.*

^{*}Carleton University, Ottawa, Canada. Email: jit@scs.carleton.ca

[†]Research supported in part by NSERC

[‡]Carleton University, Ottawa, Canada. Email: darrylhill@cunet.carleton.ca

[§]Email: aurelien.ooms@gmail.com

[¶]Supported by the VILLUM Foundation grant 16582. Part of this research was accomplished while the author was a PhD student at ULB under FRIA Grant 5203818F (FNRS).




Figure 1: The area around a point a is divided into cones with angle $2\pi/5$.


Θ_k -graphs were introduced simultaneously by Keil and Gutwin [8, 9], and Clarkson [7]. Both papers gave a spanning ratio of $1/(\cos \theta - \sin \theta)$, where $\theta = 2\pi/k$ is the angle defined by the cones. Observe that this gives a constant spanning ratio for $k \geq 9$. When this ratio t is constant, we call the graph a t -spanner. Ruppert and Seidel [11] improved this to $1/(1 - 2\sin(\theta/2))$, which applies to Θ_k -graphs with $k \geq 7$. Chew [6] gave a tight bound of 2 for $k = 6$. Bose, De Carufel, Morin, van Renssen, and Verdonschot [4] give the current best bounds on the spanning ratio of a large range of values of k . For $k = 5$, Bose, Morin, van Renssen, and Verdonschot [5] showed an upper bound of 9.96, and a lower bound of 3.78. For $k = 4$, Bose, De Carufel, Hill, and Smid [3] showed a spanning ratio of 17, while Barba, Bose, De Carufel, van Renssen, and Verdonschot [2] gave a lower bound of 7 on the spanning ratio. For $k = 3$, although Aichholzer, Bae, Barba, Bose, Korman, van Renssen, Taslakian, and Verdonschot [1] showed Θ_3 to be connected, El Molla [10] showed that there is no constant t for which Θ_3 is a t -spanner.

In this paper we study the spanning ratio of Θ_5 . We consider two arbitrary vertices, a and b , and show that there must exist a short path between them using induction on the rank of the Euclidean distance $|ab|$ among all distances between pairs of points in P . Our main result states that for all $a, b \in P$ the shortest path $\mathcal{P}(a, b)$ has length $|\mathcal{P}(a, b)| \leq K \cdot |ab|$, where $K = 5.70$.

Much of the difficulty in bounding the spanning ratio of the Θ_5 -graph stems from the following.

1. The regular pentagon is not centrally symmetric.
2. Give two vertices a and b , it may be the case that every vertex v adjacent to a has the property that $|vb| > |ab|$. In other words, all the neighbours of a are farther from b than a itself.

We organize the rest of the paper as follows. In Section 2 we introduce concepts and notation, and give some assumptions about the positions of a and



(a) Assume b is in C_2^a and a is in C_4^b .

(b) The angle α .

Figure 2: Vertices a and b and the canonical triangles T_{ab} and T_{ba} .

b that do not reduce the generality of our arguments. In Section 3 we solve all but a handful of cases using general arguments that simplify the analysis. The remaining cases are solved using ad-hoc methods, showing a spanning ratio of $K = 6.16$. In Section 4 we observe that only a single case requires $K \geq 6.16$. We analyze this case in detail to show that $|\mathcal{P}(a, b)| \leq K \cdot |ab|$ for all $K \geq 5.70$. In Section 5 we discuss directions for future work.

2 Preliminaries

Let $k \geq 3$ be an integer. Let P be set of points in the plane in general position, that is, all distances (as defined below) between pairs of points are unique and no two points have the same x -coordinate or y -coordinate. Construct the Θ_k -graph of P as follows. The vertex set is P . For each i with $0 \leq i < k$, let \mathcal{R}_i be the ray emanating from the origin that makes an angle of $2\pi i/k$ with the negative y -axis.* All indices are manipulated mod k , i.e., $\mathcal{R}_k = \mathcal{R}_0$. For each vertex v we add at most k outgoing edges as follows: For each i with $0 \leq i < k$, let \mathcal{R}_i^v be the ray emanating from v parallel to \mathcal{R}_i . Let C_i^v be the cone consisting of all points in the plane that are strictly between the rays \mathcal{R}_i^v and \mathcal{R}_{i+1}^v or on \mathcal{R}_{i+1}^v . If C_i^v contains at least one point of $P \setminus \{v\}$, then let w_i be the *closest* such point to v , where we define the closest point to be the point whose perpendicular projection onto the bisector of C_i^v minimizes the Euclidean distance to v . We add the directed edge vw_i to the graph. While the use of directed edges better illustrates this construction, in what follows we regard all edges of a Θ_5 -graph as undirected. See Fig. 1 for an example of cones and construction.

For the following description, refer to Fig. 2. Consider two vertices a and b of P . Given the Θ_5 -graph of P , we define the *canonical triangle* T_{ab} to be the triangle bounded by the sides of the cone of a that contains b and the line through b perpendicular to the bisector of that cone. Note that for every pair

*Angle values are given counter-clockwise unless otherwise stated.

of vertices a and b there are two corresponding canonical triangles, namely T_{ab} and T_{ba} . Without loss of generality assume that b is in C_2^a . Let ℓ be the leftmost vertex of the triangle T_{ab} and let r be the rightmost vertex of the triangle T_{ab} . Let m be the midpoint of ℓr . Note that a must be in C_4^b or C_0^b ; since the cases are symmetric in what follows, without loss of generality we consider the case where a is in C_4^b . Thus b is to the right of m . Let r_m be the intersection of ℓr and the bisector of $\angle ram^\dagger$, and let ℓ_m be the intersection of ℓr and the bisector of $\angle mal$. Let ℓ' and r' be the left and right endpoints of T_{ba} respectively (as seen from b facing a). Let m' be the midpoint of $\ell' r'$, and let ℓ'_m and r'_m be the intersections of $\ell' r'$ and the bisector of $\angle \ell' b m'$ and $\angle m' b r'$ respectively. See Figure 2a. Let $\alpha = \angle bam$ and let $\alpha' = \angle abm'$. Note that $\alpha + \alpha' = \pi/5$ since α and $\frac{2\pi}{5} - \alpha'$ are alternate interior angles. Thus either $\alpha \leq \pi/10$ or $\alpha' \leq \pi/10$. Without loss of generality, we assume $\alpha \leq \pi/10$. Let c be the closest neighbor to a in C_2^a , and let d be the closest neighbor to b in C_4^b . See Figure 2b. For simplicity, we write “ Θ_5 ” to mean “the Θ_5 -graph of P ”.

To sum up our assumptions following this discussion: Without loss of generality we assume that b is in C_2^a , a is in C_4^b , c is the nearest neighbour of a in C_2^a and d is the nearest neighbour of b in C_4^b . In addition, we refer back to this assumption, recalling that α is the clockwise angle ab makes with the vertical axis.

Observation 1. *Let α be clockwise angle ab makes with the vertical axis. Then $0 \leq \alpha \leq \pi/10$.*

We proceed by induction to bound the spanning ratio of Θ_5 . We show that, for any pair of points $a, b \in P$, the length of a shortest path $|\mathcal{P}(a, b)|$ in Θ_5 is at most K times the Euclidean distance between its endpoints. The induction is on the rank of the Euclidean distance $|ab|$ among all distances between pairs of points in P . The exact bound on K is made explicit in the proof. Lemma 2 is sufficient for the base case of the induction, but we first require the following geometric lemma:

Lemma 1. *Let \mathcal{T} be a triangle $\triangle pqr$, and without loss of generality assume that $|pq| \leq |pr|$. Then for all points $s \in \mathcal{T}$, $|ps| \leq |pr|$.*

Proof. (Figure 3a) Let s' be the intersection of the line through ps onto qr , thus $|ps| \leq |ps'|$ and it is enough to show that $|ps'| \leq |pr|$. The distance from p to s' is a convex function of the angle $\angle(spq)$. The minimum of this function is attained when the lines through ps' and qr are orthogonal. Therefore the maximum is attained either at $s' = q$ or $s' = r$, whichever is furthest. \square

Lemma 2. *Let (a_0, b_0) be the pair of points in P that minimizes $|ab|$ over all points a and b in P . The Θ_5 -graph of P contains the edge $a_0 b_0$.*

[†]In what follows we use $\triangle abc$ to denote the triangle defined by the points a , b , and c (given counter-clockwise). We use $\angle abc$ to denote the amplitude of the angle at b in that triangle.




Figure 3

Proof. (See Figure 3b.) Assume by contradiction that Θ_5 does not contain ab , then some point $p \in P$ different from a or b is contained in T_{ab} . We show that $|bp| < |ab|$, hence ab is not the closest pair in P .

Divide T_{ab} into two triangles by separating T_{ab} along ab into the left triangle T_{ab}^l and the right triangle T_{ab}^r . Then p belongs to one of these triangles. Observation 1 gives us that $0 \leq \alpha \leq \pi/10$, and thus $|ba| \geq |b\ell| \geq |br|$ and in both cases we can apply Lemma 1. \square

If $ab \in \Theta_5$, then $|\mathcal{P}(a,b)| \leq K|ab|$ holds for all $K \geq 1$. Otherwise we assume the following induction hypothesis: for every pair of points $a', b' \in P$ where $|a'b'| < |ab|$, the shortest path $\mathcal{P}(a', b')$ from a' to b' has length at most $|\mathcal{P}(a', b')| \leq K \cdot |a'b'|$, for some $K \geq 1$. Our goal is to find the minimum value of K for which our inductive argument holds.

Recall that c is the closest point to a in C_2^a and d is the closest point to b in C_4^b . We restrict our analysis to the following three paths:

- (1) $ac + \mathcal{P}(c, b)$,
- (2) $bd + \mathcal{P}(d, a)$, and
- (3) $ac + \mathcal{P}(c, d) + db$.

Depending on the particular arrangement of a, b, c , and d , we examine a subset of these and find a minimum value for K that satisfies at least one of the following inequalities:

- (A) $|ac| + K \cdot |cb| \leq K \cdot |ab|$,
- (B) $|bd| + K \cdot |da| \leq K \cdot |ab|$, and
- (C) $|ac| + K \cdot |cd| + |db| \leq K \cdot |ab|$.

Observe that our inductive argument follows if any of these cases holds. For instance, if we prove (A) holds for some value K , it implies that $|cb| < |ab|$ (since all distances are positive), and thus $|\mathcal{P}(c, b)| \leq K \cdot |cb|$ by the induction hypothesis. Similar conclusions follow for statements (B) and (C). Thus we can combine (1)-(3) with (A)-(C) as follows.




Figure 4: Triangles T_3 and T_4 .

- (a) $|\mathcal{P}(a, b)| \leq |ac| + |\mathcal{P}(c, b)| \leq |ac| + K \cdot |cb| \leq K \cdot |ab|$.
- (b) $|\mathcal{P}(a, b)| \leq |bd| + |\mathcal{P}(d, a)| \leq |bd| + K \cdot |da| \leq K \cdot |ab|$.
- (c) $|\mathcal{P}(a, b)| \leq |ac| + |\mathcal{P}(c, d)| + |db| \leq |ac| + K \cdot |cd| + |db| \leq K \cdot |ab|$.

For any given arrangement of vertices we prove that at least one of (A), (B), or (C) holds true for some value K , and find the smallest value for which this is true. Our proof relies mainly on case analysis, but some of these cases have similar structure. We exploit this structure in Section 3 by designing two geometric lemmas that we apply repeatedly in the inductive step. These lemmas, along with additional arguments, are then applied to different arrangements of a, b, c , and d . For all but one case we show that at least one of (a), (b), or (c) holds true for $K \geq 5.70$. The last case requires $K \geq 6.16$. We improve this further to $K \geq 5.70$, but due to the complexity of this last case, we dedicate Section 4 to its analysis.

3 Analysis

We first introduce two triangles T_3 and T_4 for which inequalities of the form of (A) and (B) hold for reasonable values of K (see Figure 4). Note the triangles are numbered to correspond to the lemmas they appear in. We state these inequalities as lemmas whose repeated use simplifies the proof of our main result.

Lemma 3. (Figure 4a) Let T_3 be a triangle with vertices (s, v, u) and corresponding interior angles $(\frac{\pi}{5}, \frac{\pi}{2}, \frac{3\pi}{10})$. Let t be a point on uv and let w be a point inside $\triangle stu$. Then $|sw| + K|wt| \leq K|st|$ for all $K \geq 4.53$.

Proof. (Figure 5a) We show $\Phi = |sw| + K|wt| - K|st| \leq 0$. Without loss of generality, orient T_3 so that u and v define a horizontal line with u left of v and with s below that line. Let w_r be the horizontal projection of w onto st , and let w_ℓ be the horizontal projection of w onto su . We have $|ww_r| + |w_r t| \geq |wt|$




Figure 5: Detailed analysis of triangles T_3 and T_4 .

by the triangle inequality. We also have that $\angle sww_\ell \geq \pi/2$, which implies that sw_ℓ is the longest edge in triangle sww_ℓ (the triangle can be drawn inside a disk whose diameter is sw_ℓ), and thus $|sw_\ell| \geq |sw|$. Since w is on $w_\ell w_r$, we have $|w_\ell w_r| \geq |ww_r|$. Thus

$$\begin{aligned} \Phi &= |sw| + K|wt| - K|st| \\ &\leq |sw_\ell| + K(|ww_r| + |w_r t|) - K(|sw_r| + |w_r t|) \\ &\leq |sw_\ell| + K|w_\ell w_r| - K|sw_r| = \Phi'. \end{aligned}$$

Let $\beta = \angle vst \geq 0$. Observe that Φ' increases as β decreases, since $|sw_r|$ decreases while $|w_\ell w_r|$ increases and $|sw_\ell|$ stays constant. Hence, Φ' is maximized when $\beta = 0$, that is, when w_r lies on sv . Thus assume that w_r lies on sv and let $|sw_\ell| = 1$ without loss of generality. We bound Φ' in terms of $\angle w_r sw_\ell = \frac{\pi}{5}$:

$$\Phi' \leq 1 + K \sin\left(\frac{\pi}{5}\right) - K \cos\left(\frac{\pi}{5}\right).$$

Solving for K we get $\Phi \leq \Phi' \leq 0$ when

$$K \geq \frac{1}{\cos(\frac{\pi}{5}) - \sin(\frac{\pi}{5})} = 4.52\dots$$

□

Lemma 4. (Figure 4b) Let T_4 be a triangle with vertices (s, v, u) and corresponding interior angles $(\frac{3\pi}{10}, \frac{3\pi}{10}, \frac{2\pi}{5})$. Let t be a point on uv such that $\angle vst \leq \pi/10$ and let w be a point inside $\triangle stu$. Then $|sw| + K|wt| \leq K|st|$ for all $K \geq 5.70$.

Proof. (Figure 5b) We show $\Phi = |sw| + K|wt| - K|st| \leq 0$ by case analysis.

Case 1) $\angle vsw \leq \frac{\pi}{5}$ (Figure 6a): Let v' be the orthogonal projection of t onto sv . Let u' be the point on the line through t and v' such that $\angle v'su' = \frac{\pi}{5}$.

Observe that $\triangle sv'u'$ corresponds to T_3 of Lemma 3 and it contains w . Thus Lemma 3 tells us $\Phi \leq 0$ for all $K \geq 4.53$.

Case 2) $\angle vsw > \frac{\pi}{5}$ (Figure 6b): Without loss of generality, orient T_4 so that u and v define a horizontal line with u left of v and with s below that line. Let w_r be the horizontal projection of w onto st , and let w_ℓ be the horizontal projection of w onto su . We have $|ww_r| + |w_r t| \geq |wt|$ by the triangle inequality. We also have that $\angle sww_\ell > \pi/2$, which implies that sw_ℓ is the longest edge in $\triangle sww_\ell$ (the triangle can be drawn inside a disk whose diameter is sw_ℓ), and thus $|sw_\ell| \geq |sw|$. Since w is on $w_\ell w_r$, we have $|w_\ell w_r| \geq |ww_r|$. Thus

$$\begin{aligned}\Phi &= |sw| + K|wt| - K|st| \\ &\leq |sw_\ell| + K(|ww_r| + |w_r t|) - K(|sw_r| + |w_r t|) \\ &\leq |sw_\ell| + K|w_\ell w_r| - K|sw_r| = \Phi'.\end{aligned}$$

We rewrite Φ' in terms of $\beta = \angle vst \geq 0$ using the sine law we get

$$|sw_\ell| = \frac{|sw_r| \sin\left(\frac{3\pi}{10} + \beta\right)}{\sin\left(\frac{2\pi}{5}\right)}$$

and

$$|w_\ell w_r| = \frac{|sw_r| \sin\left(\frac{3\pi}{10} - \beta\right)}{\sin\left(\frac{2\pi}{5}\right)}.$$

We normalize Φ' by dividing each term by $\frac{|sw_r|}{\sin\left(\frac{2\pi}{5}\right)}$ which gives us

$$\Phi' = \sin\left(\frac{3\pi}{10} + \beta\right) + K \sin\left(\frac{3\pi}{10} - \beta\right) - K \sin\left(\frac{2\pi}{5}\right).$$

The derivative of Φ' with respect to β is

$$\frac{d\Phi'}{d\beta} = \cos\left(\frac{3\pi}{10} + \beta\right) - K \cos\left(\frac{3\pi}{10} - \beta\right).$$

For all $K \geq 1$, $\frac{d\Phi'}{d\beta}(0)$ is negative and $\frac{d\Phi'}{d\beta}(\beta)$ is monotone decreasing for $0 \leq \beta \leq \frac{\pi}{10}$. Hence $\frac{d\Phi'}{d\beta}$ is negative on the whole range $(K \geq 1) \times (0 \leq \beta \leq \frac{\pi}{10})$ and Φ' is maximized at $\beta = 0$ for all $K \geq 1$. Thus

$$\Phi' \leq \Phi'(0) = \sin\left(\frac{3\pi}{10}\right) + K \sin\left(\frac{3\pi}{10}\right) - K \sin\left(\frac{2\pi}{5}\right).$$

Solving for K we get $\Phi \leq \Phi' \leq 0$ when

$$K \geq \frac{\sin\left(\frac{3\pi}{10}\right)}{\sin\left(\frac{2\pi}{5}\right) - \sin\left(\frac{3\pi}{10}\right)} = 5.69\dots$$

□




Figure 6: Analyzing triangle T_4 .




Figure 7: The regular pentagon P_{ab} .

As in the definition of T_{ab} and T_{ba} in Section 2, let c be the point closest to a in T_{ab} and let d be the point closest to b in T_{ba} . We proceed by case analysis depending on the location of the points c and d .

If c is to the right of ab or if d is to the right of ab , we can apply Lemma 3 to show the existence of a short path from a to b . When both c and d are left of ab , we use a more complicated argument requiring a new definition:

Definition 1. (Figure 7) Given any pair of points (a, b) in P , let r' and r'_m be as in the definition of T_{ba} in Section 2. We define P_{ab} to be the regular pentagon with vertices $(p_0, p_1, p_2 = r', p_3 = r'_m, p_4)$ where p_4 is above the line going through r' and r'_m (this uniquely defines the remaining points p_0 and p_1).

Observe that P_{ab} is fixed with respect to T_{ba} . This construction puts p_4 inside T_{ab} and puts p_0 and p_1 on a horizontal line with b , with p_0 lying on the boundary of T_{ab} .

Claim 5. *Given Definition 1 we have that $p_4 \in T_{ab}$, $p_0 \in lb$, and p_1 lies on the line through ℓ and b .*

Proof. Note that p_3p_4 and p_3b share the same supporting line since $\angle p_2p_3p_4 = \angle p_2p_3b = \frac{3\pi}{5}$. Let f be the intersection of $a\ell$ and p_3b . Given this observation and this definition, it is equivalent to prove that p_4 lies in the segment fb .

Translate a on the segment $\ell'_m\ell'$. Since the slope of $a\ell$ is smaller than the slope of $\ell'_m\ell'$, translating a to $a = \ell'$, that is letting $\alpha = 0$, maximizes the y -intercept of the line going through a and ℓ with any fixed vertical line. Hence this translation shrinks fb , and it remains to prove that p_4 stays in fb only in that extreme case.

With the simplifying assumption that $\alpha = 0$, we show that $|p_3f| < |p_3p_4| < |p_3b|$, which proves the claim. Note that $\angle lap_3 = \pi/10$ and $\angle p_3fa = \pi/2$, thus $|p_3f| = |p_3a| \sin(\pi/10)$. We have $|p_3p_4| = |p_3p_2| = |p_3a| \sin(\frac{\pi}{10}) / \sin(\frac{3\pi}{10})$. Since $|p_3a| = |p_3b|$, we obtain

$$|p_3b| \sin\left(\frac{\pi}{10}\right) < |p_3b| \frac{\sin\left(\frac{\pi}{10}\right)}{\sin\left(\frac{3\pi}{10}\right)} < |p_3b|.$$

□




Figure 8: Transformation 1.

Given this definition, we consider the following cases: When c is not in P_{ab} we prove $|ac| + |\mathcal{P}(c, b)| \leq 5.70|ab|$. When d is not in P_{ab} we prove $|bd| + |\mathcal{P}(d, a)| \leq 5.70|ab|$. When both c and d are in P_{ab} we analyze the length of the path $ac + \mathcal{P}(c, d) + db$. Lemma 14 gives us a bound of $6.16|ab|$ with a simple proof. Using a more technical analysis, we obtain a bound of $5.70|ab|$. This is proven in Lemma 18 in Section 4.

Some of the proofs use the simplifying assumption that $\alpha = \pi/10$. This is achieved through the following transformation: given $a, b, c, d \in P$ with T_{ab} and T_{ba} as defined earlier, we define:

Transformation 1. *Fix b, c, d , and T_{ba} , and translate a along $r'\ell'$ until $a = \ell'_m$.*

See Fig. 8. Observe that this transformation changes $|ac|$ and $|ab|$, but not $|bd|$, $|cd|$, or $|cb|$. The transformation also changes $|ad|$, but we do not use it in any case that depends on this value. We prove the following lemma allowing the application of Transformation 1 without loss of generality in several cases.

Lemma 6. *Under Transformation 1, the values of $|bd|$, $|cd|$, and $|cb|$ are unchanged, and $\Psi = |ac| - K|ab|$ is maximized when $a = \ell'_m$ for all $K \geq 3.24$.*

Proof. (Figure 9) Let $\gamma = \angle \ell'_m ba = \pi/10 - \alpha$. Define $\Psi' = |a\ell'_m| + |\ell'_m c| - K|ab|$. Note by the triangle inequality that $\Psi' \geq \Psi$. We show that Ψ' is monotonically decreasing in γ , which proves both Ψ and Ψ' are maximized when $\gamma = 0$ since then $\Psi = \Psi'$. We let $|b\ell'_m| = 1$ without loss of generality and express Ψ' as a function of γ using the law of sines:

Using $|a\ell'_m| = \frac{\sin \gamma}{\sin(\frac{2\pi}{5} - \gamma)}$ and $|ab| = \frac{\sin(\frac{3\pi}{5})}{\sin(\frac{2\pi}{5} - \gamma)} = \frac{\sin(\frac{2\pi}{5})}{\sin(\frac{2\pi}{5} - \gamma)}$, we have

$$\Psi' = \frac{\sin \gamma - K \sin(\frac{2\pi}{5})}{\sin(\frac{2\pi}{5} - \gamma)} + \underbrace{|\ell'_m c|}_{\text{Independent of } \gamma}.$$

Hence,

$$\begin{aligned} \frac{d\Psi'}{d\gamma} &= \frac{\cos \gamma \sin(\frac{2\pi}{5} - \gamma) + \cos(\frac{2\pi}{5} - \gamma)(\sin \gamma - K \sin(\frac{2\pi}{5}))}{\sin^2(\frac{2\pi}{5} - \gamma)} \\ &= \frac{\sin(\frac{2\pi}{5})(1 - K \cos(\frac{2\pi}{5} - \gamma))}{\sin^2(\frac{2\pi}{5} - \gamma)}. \end{aligned}$$

Since $0 \leq \gamma \leq \pi/10$, the denominator is positive on the whole range and the numerator is maximized when $\gamma = 0$. Since $\sin(\frac{2\pi}{5})$ is positive, it suffices to satisfy $1 - K \cos(\frac{2\pi}{5}) \leq 0$:

$$K \geq \frac{1}{\cos(\frac{2\pi}{5})} = 3.23\dots$$

□

By Lemma 6 we see that by applying Transformation 1 we maximize the value $|ac| - K|ab|$. Another way to see this is that we minimize $K|ab|$. This, in turn, allows us to explicitly determine under what conditions the inductive hypothesis applies. Note that applying Transformation 1 to where $a = \ell'_m$ is equivalent to assuming $\alpha = \pi/10$.

All these proofs can be combined in an analysis comprising *eight* cases depending on the location of c and d with respect to T_{ab} , T_{ba} , and P_{ab} , as illustrated below in the breakdown of the case analysis below. In each case we prove that for a given arrangement of vertices that $|\mathcal{P}(a, b)| \leq K|ab|$ for the given value K .




Figure 9: The values used in the proof of Lemma 6.




Figure 10: Points (a, r, m) correspond to T_3 (in blue) with $t = b$ and $w = c$.




Figure 11: Points (b, m', ℓ') correspond to T_3 (in blue) with $t = a$ and $w = d$.

Breakdown of the case analysis:

1. If c is right of ab , then $K \geq 4.53$ by Lemma 7.
2. If d is right of ab , then $K \geq 4.53$ by Lemma 8.
3. Else both c and d are left of ab . We have the following cases:
 - (a) If c is in T_{ba} , then $K \geq 5.70$ by Lemma 9.
 - (b) Else c is NOT in T_{ba} and:
 - i. If c is NOT in P_{ab} then $K \geq 4.53$ by Lemma 10.
 - ii. Else c is in P_{ab} and:
 - If d is right of am then $K \geq 3.24$ by Lemma 11.
 - If d is left of am and above c then $K \geq 4.53$ by Lemma 12
 - If d is left of am and below c (i.e. $d \notin T_{ab}$ such that bd and ac cross)
 - If d is NOT in P_{ab} then $K \geq 5.70$ by Lemma 13.
 - If d is in P_{ab} then $K \geq 6.16$ by Lemma 14 or $K \geq 5.70$ by Lemma 18.

One can check that all locations of c and d are covered. This proves our main theorem:

Theorem 1. *Given a set P of points in the plane, the Θ_5 -graph of P is a 5.70-spanner.*

We use the remainder of the paper to prove each lemma.




Figure 12: Points (a, q, p) correspond to the triangle T_4 with angles $(\frac{3\pi}{10}, \frac{2\pi}{5}, \frac{3\pi}{10})$ as denoted by the blue triangle. Let $t = b$ and $w = c$, and $\theta = \frac{\pi}{10} - \alpha$, which falls in the range of $0 \leq \angle vsu \leq \pi/10$.




Figure 13: We use the fact that p_4 lies in T_{ab} and apply T_4 .

Lemma 7. *If c is right of ab , then $|\mathcal{P}(a, b)| \leq K|ab|$ for $K \geq 4.53$.*

Proof. (Figures 4, 10) Let $(s, t, w, u, v) = (a, b, c, r, m)$, thus these points correspond to triangle T_3 of Lemma 3. Thus $|ac| + K|cb| \leq K|ab|$ for all $K \geq 4.53$. The induction hypothesis and Lemma 3 imply that there is a path from a to b with length at most

$$|\mathcal{P}(a, b)| \leq |ac| + |\mathcal{P}(c, b)| \leq |ac| + K|cb| \leq K|ab|.$$

□

Lemma 8. *If d is right of ab , then $|\mathcal{P}(a, b)| \leq K|ab|$ for $K \geq 4.53$.*

Proof. (Figures 4, 11) Let $(s, t, w, u, v) = (b, a, d, m', \ell')$, thus these points correspond to triangle T_3 from Lemma 3. Thus $|bd| + K|da| \leq K|ab|$ for $K \geq 4.53$ by Lemma 3. The induction hypothesis and Lemma 3 imply that there is a path from a to b with length at most

$$|\mathcal{P}(a, b)| \leq |bd| + |\mathcal{P}(d, a)| \leq |bd| + K|da| \leq K|ab|.$$

□

Lemma 9. *If c is left of ab and in $T_{ab} \cap T_{ba}$, then $|\mathcal{P}(a, b)| \leq K|ab|$ for $K \geq 5.70$.*

Proof. (Figures 4, 12) Let p be the intersection of br' and al , and let q be the intersection of the lines through $r'b$ and ar_m . Observe that $0 \leq \angle r_m ab \leq \pi/10$, thus $\angle r_m ab$ has the same range as $\angle vsu$ from T_4 in Lemma 4. If we let points $(s, t, w, u, v) = (a, b, c, p, q)$, then these points correspond to the triangle T_4 , and




Figure 14: The point c is in $P_{ab} \setminus T_{ba}$, and d is right of am .




Figure 15: The segments ac and bd cross and c and d are in P_{ab} .

thus $|ac| + K|cb| \leq K|ab|$ for $K \geq 5.70$ by Lemma 4. Our induction hypothesis and Lemma 4 imply that there is a path from a to b with length

$$|\mathcal{P}(a, b)| \leq |ac| + |\mathcal{P}(c, b)| \leq |ac| + K|cb| \leq K|ab|.$$

□

Lemma 10. *If $c \in T_{ab} \setminus (T_{ba} \cup P_{ab})$, then $|\mathcal{P}(a, b)| \leq K|ab|$ for all $K \geq 4.53$.*


Proof. (Figures 4, 7b) Let $\Phi = |ac| + K|cb| - K|ab|$. We apply Transformation 1. Since $c \notin T_{ba}$ it must be left of $\ell'_m b$, thus c remains left of ab . As a moves left along $\ell' \ell'_m$, so does the left side of T_{ab} , which means that c remains inside T_{ab} . Thus Lemma 6 implies that Φ is maximized at $\alpha = \pi/10$, thus we assume this is the case. Observe that $\angle bal_m = \pi/5$, and $\angle \ell'_m ba = 2\pi/5 < \pi/2$. Let q be the intersection of the line through b orthogonal to ab and the line through a and ℓ'_m . If we let $(s, t, w, u, v) = (a, b, c, q, b)$ then these points correspond to T_3 . Then Lemma 3 tells us that $|ac| + K|cb| \leq K|ab|$ and thus $\Phi = |ac| + K|cb| - K|ab| \leq 0$ for all $K \geq 4.53$. □

Lemma 11. *If d is left of ab and right of am , then $|\mathcal{P}(a, b)| \leq K|ab|$ for $K \geq 3.24$.*

Proof. (Figure 14) We show $\Phi = |bd| + K|da| - K|ab| \leq 0$, which implies $|\mathcal{P}(a, b)| \leq |bd| + |\mathcal{P}(d, a)| \leq K|ab|$ by the triangle inequality and the induction hypothesis.

Let d' be the horizontal projection of d onto ab . Let $\Phi_1 = |bd| - K|bd'|$ and $\Phi_2 = K|da| - K|d'a|$, and note that $\Phi = \Phi_1 + \Phi_2$ since $d' \in ab$. Thus it is sufficient to show that $\Phi_1 \leq 0$ and $\Phi_2 \leq 0$.

Observe that $\angle d'da > \pi/2$, since d is right of am , thus $|d'a| > |da|$, and $\Phi_2 \leq 0$ for all $K \geq 1$. For $\Phi_1 \leq 0$ we need $K \geq \frac{|bd|}{|bd'|}$. Let $\gamma = \angle d'db$ and note that $\gamma \geq \pi/10$ because $d \in T_{ba}$. Let $d_y(b, d')$ be the vertical distance between b and d' . We have $\sin \gamma = \frac{d_y(b, d')}{|bd|}$. Observe that $d_y(b, d') \leq |bd'|$ and thus $\frac{|bd|}{|bd'|} \leq \frac{|bd|}{d_y(b, d')} = \frac{1}{\sin \gamma} \leq \frac{1}{\sin(\pi/10)}$. Thus $K \geq \frac{1}{\sin(\pi/10)} \geq \frac{|bd|}{d_y(b, d')}$, and $K \geq \frac{1}{\sin(\pi/10)} = 3.23\dots$ is sufficient. □



(a) We have $|db| - K|bd'| \leq 0$. (b) We have $|ac| + K|cd| - K|ad'| \leq 0$.

Figure 16: The point c is in $P_{ab} \setminus T_{ba}$, and d is left of am but above c .

Lemma 12. *If c is in $P_{ab} \setminus T_{ba}$, and d is left of am but above c , then $|\mathcal{P}(a, b)| \leq K|ab|$ for all $K \geq 4.53$.*

Proof. (Figures 4, 16) We show $\Phi = |ac| + K|cd| + |db| - K|ab| \leq 0$, which implies $|\mathcal{P}(a, b)| \leq |ac| + |\mathcal{P}(c, d)| + |db| \leq K|ab|$ by the triangle inequality and the induction hypothesis. We split Φ into two parts, and show that each part is less than 0. Let d' be the horizontal projection of d onto ab . Let $\Phi_1 = |db| - K|bd'|$, and let $\Phi_2 = |ac| + K|cd| - K|ad'|$. Observe that $\Phi = \Phi_1 + \Phi_2$ since $d' \in ab$.

To show that $\Phi_1 \leq 0$, observe that $d_y(b, d) = d_y(b, d') \leq |bd'|$. Thus let $\Phi'_1 = |db| - K \cdot d_y(b, d) \geq \Phi_1$. Let $\gamma = \angle d'db$, and observe that $\Phi'_1 = |bd|(1 - K \sin \gamma)$. Note that $\gamma \geq \pi/10$ since $d \in T_{ba}$, and thus $K \geq 3.24$ is sufficient to have $\Phi_1 \leq 0$.

For $\Phi_2 \leq 0$, let d'' be the horizontal projection of d onto am . Since $\angle ad''d' = \pi/2$, $|ad''| \leq |ad'|$. Since $c \notin T_{ba}$, $\angle cdd'' \geq 9\pi/10$, thus $|cd''| > |cd|$. Let $\Phi'_2 = |ac| + K|cd''| - K|ad''| \geq \Phi_2$. Let q be the horizontal projection of d'' onto al . Let the points $(s, t, w, u, v) = (a, d'', c, q, d'')$ and thus these points correspond to T_3 . Thus $|ac| + K|cd''| \leq K|ad''|$ for all $K \geq 4.53$ by Lemma 3. \square

Lemma 13. *If d is left of ab , below c and not in P_{ab} , then $|\mathcal{P}(a, b)| \leq K|ab|$ for all $K \geq 5.70$.*

Proof. (Figures 4, 13) We note that ac and bd intersect and d must be outside of T_{ab} (otherwise ad would be an edge of Θ_5 , but not ac). We first show that d is below br'_m . Recall that P_{ab} is fixed with respect to T_{ba} . Since d is outside of T_{ab} and P_{ab} , if p_4p_0 is inside T_{ab} , d must be below br'_m . Since the slope of p_0p_4 is less than the slope of la , it is sufficient to show that p_4 is inside T_{ab} which follows by Claim 5. By Observation 1 we have that $0 \leq \angle abl' \leq \pi/10$. Thus we can map the points (s, t, w, u, v) to (b, a, d, r'_m, ℓ') and apply Lemma 4. Thus $|bd| + K|da| \leq K|ab|$ for $K \geq 5.70$. Our induction hypothesis and Lemma 4 imply that there is a path from b to a with length at most

$$|\mathcal{P}(a, b)| \leq |bd| + |\mathcal{P}(d, a)| \leq |bd| + K|da| \leq K|ab|.$$

\square

Lemma 14. *If ac and bd cross and both c and d are in P_{ab} , then $|\mathcal{P}(a, b)| \leq K|ab|$ for $K \geq 6.16$.*

Proof. (Figures 4, 15) We show $\Phi = |ac| + K|cd| + |db| - K|ab| \leq 0$, which implies $|\mathcal{P}(a, b)| \leq |ac| + |\mathcal{P}(c, d)| + |db| \leq K|ab|$ by the triangle inequality and the induction hypothesis. Under Transformation 1, Lemma 6 implies that Φ is maximized when $\alpha = \pi/10$, so we assume this is the case. Since c , d , and P_{ab} are fixed, c and d are still inside P_{ab} after Transformation 1. Given that c and d are in P_{ab} , the furthest apart c and d can be is if they are both on a diagonal of P_{ab} . The length of one side of P_{ab} is at most $\frac{\sin(\pi/10)}{\sin(3\pi/10)}|ab|$. That means a diagonal of P_{ab} , and thus $|cd|$, has length at most $2 \sin(3\pi/10) \frac{\sin(\pi/10)}{\sin(3\pi/10)}|ab| = 2 \sin(\pi/10)|ab|$. At their longest, $|ac|$ and $|bd|$ each have length $\frac{\sin(2\pi/5)}{\sin(3\pi/10)}|ab|$ by the law of sines. We want

$$\Phi = |ac| + K|cd| + |db| - K|ab| \leq 0.$$

Solving for K gives

$$K \geq \frac{|ac| + |db|}{|ab| - |cd|} \geq \frac{2 \cdot \sin(2\pi/5)}{\sin(3\pi/10) \cdot (1 - 2 \cdot \sin(\pi/10))} = 6.15 \dots$$

□

4 Proving a spanning ratio of 5.70

In this section we present a lemma with a stronger bound for the case handled by Lemma 14. Proving this lemma requires a careful analysis of the locations of c and d and the tradeoffs between the values of $|ac| + |db|$ and $K|cd|$. Let $\Phi = |ac| + K|cd| + |db| - K|ab|$. For the rest of this section, assume we have applied Transformation 1, and thus $\alpha = \pi/10$ and Φ is maximized. Since P_{ab} , c and d are fixed, both c and d are still in P_{ab} . Let c' be the intersection of the line through a and c and the segment p_0p_1 , and let d' be the intersection of the line through b and d and the segment p_3p_4 . See Figure 17. Let $\Phi' = |ac'| + K|c'd'| + |d'b| - K|ab|$, and let $\Phi'' = |ap_1| + K|p_1p_3| + |p_3b| - K|ab|$.




Figure 17: Points c' and d' .

Lemma 18. *If ac and bd intersect and both c and d are in P_{ab} , then $|\mathcal{P}(a, b)| \leq K|ab|$ for $K \geq 5.70$.*

We split the analysis into three steps that amount to proving the following lemmas:

Lemma 15. *For all $K \geq 5.70$, $\Phi \leq \Phi'$.*

Lemma 16. *For all $K \geq 5.70$, $\Phi' \leq \Phi''$.*

Lemma 17. *For all $K \geq 5.70$, $\Phi'' \leq 0$.*

The following lemma follows from these lemmas, the triangle inequality, and the induction hypothesis. It supersedes Lemma 14:




Figure 18: Finding the longest distance from a to b when c and d are in P_{ab} ,

Substituting Lemma 18 for Lemma 14 in the proof of Theorem 1 brings the spanning ratio of the Θ_5 -graph down to 5.70. We are left with proving Lemmas 15, 16, and 17, which is done in the next three subsections.

4.1 Proof of Lemma 15

Lemma 15 states that $|ac| + |bd| + K|cd| - K|ab| \leq |ac'| + |bd'| + K|c'd'| - K|ab|$ for $K \geq 5.70$. See Figure 18a. Let e be the intersection of ac and bd , and let e' be the intersection of br' and al . Observe that $\angle le'r' = 2\pi/5$, and thus we can see that $\angle dec \geq 2\pi/5$. This implies that $\angle dec$ cannot be the smallest angle in $\triangle dec$, since that would require $\angle dec \leq \pi/3$. Thus at least one of $\angle dce$ and $\angle edc$ is the smallest angle in $\triangle dec$. Since we have applied Transformation 1, and can thus assume that $\alpha = \pi/10$, the cases are symmetric. We can therefore, without loss of generality, assume that $\angle dce$ is the smallest angle in $\triangle dec$.

Lemma 15. For all $K \geq 5.70$, $\Phi \leq \Phi'$.

Proof. Since c lies on ac' and d lies on bd' , we have $|ac| \leq |ac'|$ and $|bd| \leq |bd'|$, and it is sufficient to show that $|cd| \leq |c'd'|$. We first show that $|cd| \leq |c'd|$. Since $\angle dce$ is the smallest angle in $\triangle dec$, $\angle dce < \pi/3$. That implies that $\angle c'cd > \pi/2$, which implies that $c'd$ is the longest side of triangle $\triangle cc'd$, and thus $|cd| \leq |c'd|$. See Fig.18a.

We now show that $|c'd'| \geq |c'd|$. If $\angle c'dd' \geq \pi/2$, then $c'd'$ is the longest side of $\triangle c'dd'$, and $|c'd'| \geq |c'd|$ and we are done. Otherwise assume $\angle c'dd' < \pi/2$.

The law of sines tells us that $\frac{|c'd'|}{\sin \angle c'dd'} = \frac{|c'd|}{\sin \angle dd'c'}$. Since $\sin \theta$ is an increasing function for $0 \leq \theta < \pi/2$, showing that $\angle c'dd' \geq \angle dd'c'$ is sufficient to show $|c'd'| \geq |c'd|$, as it would imply both angles are $< \pi/2$. Observe that $\angle c'dd' \geq \angle c'ed'$ and $\angle ed'c' = \angle dd'c'$, thus it is sufficient to prove that $\angle c'ed' \geq \angle ed'c'$.

Observe that $\angle ced = \angle c'ed' \geq 2\pi/5$. We now find the maximum of $\angle dd'c' = \angle ed'c' \leq 2\pi/5$. Observe that if c' moves left, $\angle ed'c'$ increases, thus assume c' is




Figure 19: An example of Φ'' .

at p_1 . Let $O(bp_3c')$ be the circle through b , p_3 , and c' with center o . Observe that o lies on br' . Observe that $\angle r'bd' = \pi/10$, thus $\angle r'op_3 = \pi/5$. Segment or' makes an angle of $\pi/10$ with the horizontal line through o . Thus od' makes an angle of $3\pi/10$ with the horizontal line through o , and thus the line tangent to $O(bp_3c')$ at p_3 is the line supporting $\ell'r'$, since $\ell'r'$ makes an angle of $3\pi/10$ with the vertical line through ℓ' . See Figure 18b. That implies that $[p_2, p_3]$ lies outside of $O(bp_3c')$, which means for every point d' , $\angle ed'c' \leq \angle ep_3c' = 2\pi/5$, and thus $\angle c'dd' \geq \angle dd'c'$ as required. \square

4.2 Proof of Lemma 16

Observe that $|ap_1| + K|p_1p_3| + |p_3b| = |ap_0| + K|p_0p_2| + |p_2b|$ when $\alpha = \pi/10$, since T_{ab} and T_{ba} are the same size and the cases are symmetric. We prove that

$$\begin{aligned} \Phi' &= |ac'| + K|c'd'| + |d'b| - K|ab| \leq |ap_1| + K|p_1p_3| + |p_3b| - K|ab| = \Phi'' \\ &= |ap_0| + K|p_0p_2| + |p_2b| - K|ab|. \end{aligned}$$

Lemma 16. *For all $K \geq 5.70$, $\Phi' \leq \Phi''$.*

Proof. (Figure 19) Without loss of generality, we assume that $|p_1c'| \leq |p_2d'|$. We show that Φ' is maximized when $c' = p_1$ and $d' = p_3$.

(Figure 20a) Observe that $|p_1p_2| \leq |c'd'| \leq |p_1p_3|$. Let z be a point on p_2p_3 that moves from p_2 to p_3 , and observe that $|p_1z|$ takes on every value from $|p_1p_2|$ to $|p_2p_3|$. Thus there must be a point q on p_2p_3 such that $|p_1q| = |c'd'|$.

We claim that $|ap_1| + |bq| \geq |ac'| + |bd'|$, which implies that $\Phi' \leq |ap_1| + K|p_1q| + |qb| - K|ab|$.

Observe $|ap_1| \geq |ac'|$, since $\angle p_1c'a > \pi/2$, making ap_1 the longest edge in triangle $\Delta ac'p_1$. We claim that q is between d' and p_2 , and thus $|bq| \geq |bd'|$ since $\angle bd'q > \pi/2$. By contradiction, assume that q is between d' and p_3 . Since $|p_1c'| \leq |p_2d'|$, $\angle qd'c' > \pi/2$, which implies that $|c'q| > |c'd'|$. Also note that $\angle qd'p_1 > \pi/2$, which implies $|p_1q| > |c'q| > |c'd'|$, a contradiction. Thus assuming that $c' = p_1$ and $d' = q$ does not decrease Φ' .




Figure 20

Now, given that c' is on p_1 , we show that $\Phi' \leq |ac'| + |bp_3| + K|c'p_3|$, that is, when d' is on p_3 . To do this we define another function $\Phi^* = |ac'| + |d'p_3| + |p_3b| + K|c'd'| - K|ab|$. See Figure 20b. Since $|bd'| \leq |d'p_3| + |p_3b|$ by the triangle inequality, $\Phi' \leq \Phi^*$, and observe that $\Phi' = \Phi^* = \Phi''$ when $d' = p_3$. We show that Φ^* is maximized when $d' = p_3$, thus implying that Φ' is also maximized when $d' = p_3$, and $\Phi' \leq \Phi''$. Let $\theta = \angle p_2p_1d'$. We allow d' to move along p_2p_3 until d' is on p_3 , and fix all other points, and observe how Φ^* changes with θ .

We first rewrite Φ^* as $\Phi^* = |ac'| + |p_2p_3| - |p_2d'| + |p_3b| + K|c'd'| - K|ab|$. Using the sine law we get $|p_2d'| = \frac{\sin \theta}{\sin(2\pi/5 - \theta)} |p_1p_2|$, and $|c'd'| = \frac{\sin(3\pi/5)}{\sin(2\pi/5 - \theta)} |p_1p_2|$. All other terms of Φ^* have fixed values with respect to θ . Thus

$$\begin{aligned}
\frac{d\Phi^*}{d\theta} &= \frac{d}{d\theta} \left(K \frac{\sin(3\pi/5)}{\sin(2\pi/5 - \theta)} |p_1p_2| - \frac{\sin \theta}{\sin(2\pi/5 - \theta)} |p_1p_2| \right) \\
&= \frac{K \cos(2\pi/5 - \theta) \sin(3\pi/5) - \cos \theta \sin(2\pi/5 - \theta) - \sin \theta \cos(2\pi/5 - \theta)}{\sin^2(2\pi/5 - \theta)} |p_1p_2| \\
&= \frac{K \cos(2\pi/5 - \theta) \sin(3\pi/5) - \sin(2\pi/5)}{\sin^2(2\pi/5 - \theta)} |p_1p_2|. \tag{1}
\end{aligned}$$

Observe that $0 \leq \theta \leq 3\pi/10$. The denominator of (1) is always positive. The numerator of (1) is minimized at $\theta = 0$, which for $K \geq 5.70$ is positive. Thus (1) is always positive for $0 \leq \theta \leq 3\pi/10$, thus Φ^* is increasing in θ , and is maximized when $d' = p_3$, as required. Thus $\Phi' \leq \Phi^* \leq \Phi'' = |ap_1| + K|p_1p_3| + |p_3b| - K|ab|$ as required. \square

4.3 Proof of Lemma 17

Lemma 17. For all $K \geq 5.70$, $\Phi'' \leq 0$.

Proof. (Figure 19) We apply Transformation 1 with $\alpha = \frac{\pi}{10}$ and assume that $|ab| = 1$. Then using the law of sines we get $|bp_3| = 1$, $|ap_1| = \frac{\sin(2\pi/5)}{\sin(3\pi/10)}$, and $|p_1p_3| = 2 \sin(3\pi/10) \frac{\sin(\pi/10)}{\sin(3\pi/10)} = 2 \sin(\pi/10)$. We want

$$\Phi'' = |ap_1| + K|p_1p_3| + |p_3b| - K|ab| \leq 0.$$

Solving for K gives

$$K \geq \frac{|ap_1| + |p_3b|}{|ab| - |p_1p_3|} = \frac{\frac{\sin(2\pi/5)}{\sin(3\pi/10)} + 1}{1 - 2 \sin(\pi/10)} = 5.69 \dots$$

□

5 Open Problems

Using a few simple geometric observations and arguments, we have lowered the spanning ratio of Θ_5 from 9.96 to 5.70, bringing us closer to the lower bound of 3.798 and thus a tight bound. The obvious open problem that remains is closing the gap between the upper and lower bound on the spanning ratio of the Θ_5 -graph.

Acknowledgements: We thank Elena Arseneva for many fruitful discussions on the topic.

References

- [1] O. Aichholzer, S. W. Bae, L. Barba, P. Bose, M. Korman, A. van Renssen, P. Taslakian, and S. Verdonschot. Theta-3 is connected. *Computational geometry*, 47(9):910–917, 2014.
- [2] L. Barba, P. Bose, J. L. De Carufel, A. van Renssen, and S. Verdonschot. On the stretch factor of the Theta-4 graph. In *Proceedings of the 13th International Symposium on Algorithms and Data Structures (WADS)*, pages 109–120, 2013.
- [3] P. Bose, J. L. De Carufel, D. Hill, and M. Smid. On the spanning and routing ratio of Theta-four. In *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 2361–2370, 2019.
- [4] P. Bose, J. L. De Carufel, P. Morin, A. van Renssen, and S. Verdonschot. Towards tight bounds on Theta-graphs: More is not always better. *Theoretical Computer Science*, 616:70–93, 2016.

- [5] P. Bose, P. Morin, A. van Renssen, and S. Verdonschot. The Theta-5-graph is a spanner. *Computational Geometry*, 48(2):108–119, 2015.
- [6] L. P. Chew. There are planar graphs almost as good as the complete graph. *Journal of Computer and System Sciences*, 39(2):205 – 219, 1989.
- [7] K. Clarkson. Approximation algorithms for shortest path motion planning. In *Proceedings of the Nineteenth Annual ACM Symposium on Theory of Computing (STOC)*, pages 56–65, 1987.
- [8] J. M. Keil. Approximating the complete Euclidean graph. In *Proceedings of the 1st Scandinavian Workshop on Algorithm Theory (SWAT)*, pages 208–213, 1988.
- [9] J. M. Keil and C. A. Gutwin. Classes of graphs which approximate the complete Euclidean graph. *Discrete & Computational Geometry*, 7(1):13–28, 1992.
- [10] N. M. El Molla. *Yao spanners for wireless ad-hoc networks*. PhD thesis, Villanova University, 2009.
- [11] J. Ruppert and R. Seidel. Approximating the d-dimensional complete Euclidean graph. In *Proceedings of the 3rd Canadian Conference on Computational Geometry (CCCG)*, 1991.