BERRY-ESSÉEN BOUNDS AND ALMOST SURE CLT FOR THE QUADRATIC VARIATION OF A GENERAL GAUSSIAN PROCESS

YONG CHEN, ZHEN DING, AND YING LI

ABSTRACT. In this paper, we consider the explicit bound for the second-order approximation of the quadratic variation of a general fractional Gaussian process $(G_t)_{t\geq 0}$. The second order mixed partial derivative of the covariance function $R(t,s)=\mathbb{E}[G_tG_s]$ can be decomposed into two parts, one of which coincides with that of fractional Brownian motion and the other of which is bounded by $(ts)^{H-1}$ up to a constant factor. This condition is valid for a class of continuous Gaussian processes that fails to be self-similar or have stationary increments. Under this assumption, we obtain the optimal Berry-Esséen bounds when $H \in (0, \frac{2}{3}]$ and the upper Berry-Esséen bounds when $H \in (\frac{2}{3}, \frac{3}{4}]$. As a by-product, we also show the almost sure central limit theorem (ASCLT) for the quadratic variation when $H \in (0, \frac{3}{4}]$. The results extend that of [10] to the case of general Gaussian processes, improve the Berry-Esséen bounds and unify the proofs in [14], [1] and [7] for respectively the sub-fractional Brownian motion, the bi-fractional Brownian motion and the sub-bifractional Brownian motion.

Keywords: Malliavin calculus; Optimal Fourth Moment theorem; Berry-Esséen bounds; Gaussian process.

MSC 2010: 60H07; 60G15; 60F05.

1. Introduction

We are interested in the explicit bound for the second-order approximation of the quadratic variation of a general fractional Gaussian process $G = \{G_t : t \ge 0\}$ on [0, T], defined as

$$Z_n := \sum_{k=0}^{n-1} \left[(G_{k+1} - G_k)^2 - \mathbb{E}[(G_{k+1} - G_k)^2] \right].$$

Let the renormalization of Z_n be

$$V_n := \frac{1}{\sigma_n} Z_n,\tag{1.1}$$

where $\sigma_n > 0$ is so that $\mathbb{E}[V_n^2] = 1$, i.e., $\sigma_n^2 := \mathbb{E}[Z_n^2]$.

When G is the fractional Brownian motion B^H , by using Stein method and Malliavin calculus, Nourdin and Peccati [9, 10] derived explicit bounds for the total variation distance between the law of V_n and the standard normal law N. From then on, the same problem was extended to some other fractional Gaussian processes such as the sub-fractional Brownian motion, the bi-fractional Brownian motion and the sub-bifractional Brownian motion in [14], [1] and [7] respectively.

We find that the above four types of fractional Gaussian processes are all special examples of the following general Gaussian process G. Let

$$R^{B}(s,t) = \frac{1}{2}(|s|^{2H} + |t|^{2H} - |t - s|^{2H}), \tag{1.2}$$

be the covariance function of the fractional Brownian motion $\{B^H(t), t \geq 0\}$.

HYPOTHESIS 1.1. For $H \in (0, 1)$, the covariance function $R(t, s) = \mathbb{E}[G_t G_s]$ satisfies that

- (1) for any $s \ge 0$, R(0, s) = 0.
- (2) for any fixed $s \in (0,T)$, R(t,s) is continuous on [0,T] and differentiable function with respect to t on $(0,s) \cup (s,T)$, such that $\frac{\partial}{\partial t}R(t,s)$ is absolutely integrable.
- (3) for any fixed $t \in (0,T)$, the difference

$$\frac{\partial R(t,s)}{\partial t} - \frac{\partial R^B(t,s)}{\partial t}$$

is continuous on [0,T], and it is differentiable with respect to s on (0,T) such that $\Psi(t,s)$, the partial derivative with respect to s of the difference, satisfies

$$|\Psi(t,s)| \le C'_H |ts|^{H-1},$$
 (1.3)

where the constants $H, C'_H \geq 0$ do not depend on T, and $R^B(t, s)$ is the covariance function of the fractional Brownian motion as in (1.2).

Example 1.2. The subfractional Brownian motion $\{S^H(t), t \geq 0\}$ with parameter $H \in (0, 1)$ has the covariance function

$$R(t,s) = s^{2H} + t^{2H} - \frac{1}{2} \left((s+t)^{2H} + |t-s|^{2H} \right),$$

which satisfies Hypothesis 1.1.

Example 1.3. The bi-fractional Brownian motion $\{B^{H',K}(t), t \geq 0\}$ with parameters $H', K \in (0,1)$ has the covariance function

$$R(t,s) = \frac{1}{2} \left((s^{2H'} + t^{2H'})^K - |t - s|^{2H'K} \right),\,$$

which satisfies Hypothesis 1.1 when H := H'K.

Example 1.4. The generalized sub-fractional Brownian motion $S^{H',K}(t)$ with parameters $H' \in (0,1), K \in [1,2)$ and $H'K \in (0,1)$ satisfies Hypothesis 1.1 when H := H'K. The covariance function is

$$R(t, s) = (s^{2H'} + t^{2H'})^K - \frac{1}{2} [(t+s)^{2H'K} + |t-s|^{2H'K}].$$

Notation 1. Given two deterministic numeric sequences $(a_n)_{n>0}, (b_n)_{n>0}$, we use the following notations and definitions for respectively commensurability, equivalence:

 $a_n \simeq b_n \iff \exists c, C > 0 : cb_n \geq a_n \leq Cb_n$, for n large enough,

$$a_n \sim b_n \iff \exists c_n, C_n > 0 : \lim_{n \to \infty} c_n = \lim_{n \to \infty} C_n = 1, \ c_n b_n \le a_n \le C_n b_n, \text{ for } n \text{ large enough.}$$

Now we list our main results as follows:

Theorem 1.5. Let $N \sim N(0,1)$ and V_n be given as in (1.1) and suppose Hypothesis 1.1 holds. When $H \in (0, \frac{2}{3})$,

$$d_{TV}(V_n, N) \simeq n^{-\frac{1}{2}};$$
 (1.4)

when $H = \frac{2}{3}$

$$d_{TV}(V_n, N) \simeq n^{-\frac{1}{2}} \log^2 n;$$
 (1.5)

when $H \in (\frac{2}{3}, \frac{3}{4})$, there exists a positive contant c_H depending on H such that for any $n \geq 1$,

$$d_{TV}(V_n, N) \le c_H \times n^{\frac{1}{2}(4H-3)};$$
 (1.6)

when $H = \frac{3}{4}$, there exists a positive contant c such that for any $n \ge 1$,

$$d_{TV}(V_n, N) \le \frac{c}{(\log n)^{\frac{3}{2}}};$$
 (1.7)

- (1) The above Berry-Esséen types bounds are more sharp than Remark 1.6. those obtained in [14], [1] and [7] for respectively the sub-fractional Brownian motion, the bi-fractional Brownian motion and the sub-bifractional Brownian motion.
 - (2) We do not know how to obtain the optimal bound in the case of $H \in (\frac{2}{3}, \frac{3}{4}]$.

(3) In the same vein, we can also extend Theorem 1.5 to the pth Hermite variation with p > 2.

As a by-product of Theorem 1.5, we have the ASCLT of the sequence $(V_n)_{n\geq 1}$.

Theorem 1.7. If $H \in (0, \frac{3}{4}]$ then the sequence $(V_n)_{n\geq 1}$ satisfies the ASCLT. In other words, for any bounded and continuous function $\varphi : \mathbb{R} \to \mathbb{R}$, we have almost surely,

$$\frac{1}{\log n} \sum_{k=1}^{n} \frac{1}{k} \varphi(V_k) \xrightarrow{a.s.} \mathbb{E}\varphi(N)$$
 (1.8)

as $n \rightarrow \infty$, where $N \sim N(0, 1)$.

2. Preliminary

In this section, we describe some basic facts on stochastic calculus with respect to the Gaussian process, for more complete presentation on the subject can be found in [6].

Denote $G = \{G_t, t \in [0, T]\}$ as a continuous centered Gaussian process with $G_0 = 0$ and the covariance function

$$\mathbb{E}(G_t G_s) = R(s, t), \ s, t \in [0, T], \tag{2.1}$$

defined on a complete probability $\operatorname{space}(\Omega, \mathcal{F}, P)$, where the \mathcal{F} is generated by the Gaussian family G. Suppose in addition that the covariance function R is continuous. Let \mathcal{E} denote the space of all real valued step functions on [0, T]. The Hilbert space \mathfrak{H} is defined as the closure of \mathcal{E} endowed with the inner product

$$\langle \mathbb{1}_{[a,b)}, \mathbb{1}_{[c,d)} \rangle_{\mathfrak{H}} = \mathbb{E}((G_b - G_a)(G_d - G_c)),$$
 (2.2)

where $\mathbb{1}_{[a,b)}$ is the indicator function of the interval [a,b). With abuse of notation, we also denote $G = \{G(h), h \in \mathfrak{H}\}$ as the isonormal Gaussian process on the probability space, indexed by the elements in the Hilbert space \mathfrak{H} . Then G is a Gaussian family of random variables such that

$$\mathbb{E}(G) = \mathbb{E}(G(h)) = 0, \ \mathbb{E}(G(g)G(h)) = \langle g, h \rangle_{\mathfrak{H}}, \quad \forall g, h \in \mathfrak{H}.$$
 (2.3)

Notation 2. Let $R^B(t,s)$ be the covariance function of the fractional Brownian motion as in (1.2). $\mathcal{V}_{[0,T]}$ denote the set of bounded variation functions on [0,T].

For functions $f, g \in \mathcal{V}_{[0,T]}$, we define two products as

$$\langle f, g \rangle_{\mathfrak{H}_{1}} = -\int_{[0,T]^{2}} f(t) \frac{\partial R^{B}(t,s)}{\partial t} dt \nu_{g}(ds),$$

$$\langle f, g \rangle_{\mathfrak{H}_{2}} = C'_{H} \int_{[0,T]^{2}} |f(t)g(s)| (ts)^{H-1} dt ds.$$
(2.4)

The following proposition is an extension of [6, Theorem 2.3] and [5, Proposition 2.2], which gives the inner products representation of the Hilbert space \mathfrak{H} :

Proposition 2.1. $V_{[0,T]}$ is dense in \mathfrak{H} and we have

$$\langle f, g \rangle_{\mathfrak{H}} = \int_{[0,T]^2} R(t,s) \nu_f(\mathrm{d}t) \nu_g(\mathrm{d}s), \qquad \forall f, g \in \mathcal{V}_{[0,T]}, \tag{2.5}$$

where ν_g is the restriction to $([0,T],\mathcal{B}([0,T]))$ of the Lebesgue-Stieljes signed measure associated with g^0 defined as

$$g^{0}(x) = \begin{cases} g(x), & \text{if } x \in [0, T). \\ 0, & \text{otherwise.} \end{cases}$$

Furthermore, if the covariance function R(t,s) satisfies Hypothesis 1.1, then

$$\langle f, g \rangle_{\mathfrak{H}} = -\int_{[0,T]^2} f(t) \frac{\partial R(t,s)}{\partial t} dt \nu_g(ds), \qquad \forall f, g \in \mathcal{V}_{[0,T]}.$$
 (2.6)

In particular, we have

$$|\langle f, g \rangle_{\mathfrak{H}} - \langle f, g \rangle_{\mathfrak{H}_1}| \le \langle f, g \rangle_{\mathfrak{H}_2}, \qquad \forall f, g \in \mathcal{V}_{[0,T]}. \tag{2.7}$$

Remark 2.2. When $H \in (\frac{1}{2}, 1)$, Hypothesis 1.1 (3) and Lemma 3.1 imply that the identity (2.6) can be rewritten as

$$\langle f, g \rangle_{\mathfrak{H}} = \int_{[0,T]^2} f(t)g(s) \frac{\partial^2 R(t,s)}{\partial t \partial s} dt ds, \quad \forall f, g \in \mathcal{V}_{[0,T]}.$$
 (2.8)

In this case, the inequality (2.7) has obtained from (2.8) in [4]. When $H \in (0, \frac{1}{2})$, it is well known that both $\frac{\partial^2}{\partial t \partial s} R(t,s)$ and $\frac{\partial^2}{\partial t \partial s} R^B(t,s)$ are not absolutely integrable. But the absolute integrability of their difference makes the key inequality (2.7) still valid.

Proof. The first claim and the identity (2.5) are rephrased from Theorem 2.3 of [6]. Hypothesis 1.1 (2) and Lemma 3.1 imply the inner products representation (2.6).

Finally, it follows from the identity (2.6) and Notation 2 that

$$\langle f, g \rangle_{\mathfrak{H}_1} - \langle f, g \rangle_{\mathfrak{H}} = \int_{[0,T]^2} f(t) \left[\frac{\partial R(t,s)}{\partial t} - \frac{\partial R^B(t,s)}{\partial t} \right] dt \nu_g(ds).$$

By the fundamental theorem of calculus (see Proposition 1.6.41 of [13]), Hypothesis 1.1 (1) and (3) imply that when $s \neq t$,

$$\frac{\partial R(t,s)}{\partial t} - \frac{\partial R^B(t,s)}{\partial t} = \int_0^s \Psi(t,r) dr.$$
 (2.9)

Hence, Lemma 3.1 implies that

$$\langle f, g \rangle_{\mathfrak{H}} - \langle f, g \rangle_{\mathfrak{H}_{1}} = -\int_{[0,T]} f(t) dt \int_{[0,T]} \left[\frac{\partial R(t,s)}{\partial t} - \frac{\partial R^{B}(t,s)}{\partial t} \right] \nu_{g}(ds)$$

$$= \int_{[0,T]} f(t) dt \int_{[0,T]} g(s) \Psi(t,s) ds,$$

which implies the inequality (2.7) since $\Psi(t,s)$ satisfies the inequality (1.3).

Denote $\mathfrak{H}^{\otimes p}$ and $\mathfrak{H}^{\odot p}$ as the pth tensor product and the pth symmetric tensor product of the Hilbert space \mathfrak{H} . Let \mathcal{H}_p be the pth Wiener chaos with respect to G. It is defined as the closed linear subspace of $L^2(\Omega)$ generated by the random variables $\{H_p(G(h)): h \in \mathfrak{H}, \|h\|_{\mathfrak{H}} = 1\}$, where H_p is the pth Hermite polynomial defined by

$$H_p(x) = (-1)^p e^{\frac{x^2}{2}} \frac{d^p}{dx^p} e^{-\frac{x^2}{2}}, \quad p \ge 1,$$

and $H_0(x) = 1$. We have the identity $I_p(h^{\otimes p}) = H_p(G(h))$ for any $h \in \mathfrak{H}$ with $||h||_{\mathfrak{H}} = 1$ where $I_p(\cdot)$ is the pth multiple Wiener-Itô integral. Then the map I_p provides a linear isometry between $\mathfrak{H}^{\odot p}$ (equipped with the norm $\sqrt{p!}||\cdot||_{\mathfrak{H}^{\otimes p}}$) and \mathcal{H}_p . Here $\mathcal{H}_0 = \mathbb{R}$ and $I_0(x) = x$ by convention.

The following Theorem 2.3, known as the optimal fourth moment theorem, provides exact rates of convergence in total variation distance between a multiple Wiener-Itô integral and a normal distribution (see [12, 3]).

Theorem 2.3. Let $N \sim N(0,1)$ be a standard Gaussian random variable. Let $\{F_n : n \geq 1\}$ be a sequence of random variables living in the pth multiple Wiener-Itô integral with unit variance. If $\lim_{n\to\infty} \mathbb{E}[F_n^4] = 3$, then there exist two finite constants 0 < c < C (possibly depending on p and on the sequence $\{F_n\}$, but not on n) such that the following estimate in total variation holds for every n:

$$c\mathbf{M}(F_n) \le d_{TV}(F_n, N) \le C\mathbf{M}(F_n),$$

$$\mathbf{M}(F_n) := \max \left\{ \left| \mathbb{E}[F_n^3] \right|, \mathbb{E}[F_n^4 - 3] \right\}.$$

The quantities $\kappa_3(F_n) := \mathbb{E}(F_n^3)$ and $\kappa_4(F_n) := \mathbb{E}[F_n^4] - 3$ are called the 3rd and 4th cumulants of F_n . That $\kappa_3(F_n)$ coincides with the third moment is because F_n is centered. Moreover, $\kappa_4(F_n)$ is strictly positive (see [11, 8]).

The following theorem is used to show the ASCLT.

Theorem 2.4 ([2]). Let $p \geq 2$ be an integer, and let $F_n = I_q(f_n)$, with $f_n \in \mathfrak{H}^{\odot p}$. Assume that for all n, and that $F_n \stackrel{law}{\to} \mathcal{N}(0,1)$ as $n \to \infty$. If the two following conditions are satisfied

(1).
$$\sum_{n=2}^{\infty} \frac{1}{n \log^{2} n} \sum_{k=1}^{n} \frac{1}{k} \|f_{k} \otimes_{r} f_{k}\|_{\mathfrak{H}^{\otimes 2(q-r)}} < \infty, \text{ for every } 1 \leq r \leq p-1,$$
(2).
$$\sum_{n=2}^{\infty} \frac{1}{n \log^{3} n} \sum_{k,l=1}^{n} \frac{|\langle f_{k}, f_{l} \rangle_{\mathfrak{H}^{\otimes p}}|}{kl} < \infty.$$
(2.10)

Then $\{F_n : n \geq 1\}$ satisfies an ASCLT.

Denote

$$\rho(r) := \frac{1}{2} (|r+1|^{2H} + |r-1|^{2H} - 2|r|^{2H}), \quad r \in \mathbb{Z}.$$

It is well-known that for any $r \neq 0$, $\rho(r)$ is positive when $H \in (\frac{1}{2}, 1)$ and is negative when $H \in (0, \frac{1}{2})$. It behaves asymptotically as $|r| \to \infty$,

$$\rho(r) \sim H(2H-1) |r|^{2H-2}.$$

In particular, when $H > \frac{1}{2}$, for |r| large enough,

$$\rho(r) \ge H(H - \frac{1}{2})(1 + |r|)^{2H - 2}.$$

The following proposition is cited from [9, p.74].

Proposition 2.5. If $H \in (0, \frac{3}{4})$ then

$$\lim_{n \to \infty} \frac{2}{n} \sum_{i,j=0}^{n-1} \rho^2(i-j) = 2 \sum_{r \in \mathbb{Z}} \rho^2(r) := \sigma^2; \tag{2.11}$$

and if $H = \frac{3}{4}$ then

$$\lim_{n \to \infty} \frac{2}{n \log n} \sum_{i,j=0}^{n-1} \rho^2(i-j) = \frac{9}{16}.$$

The following propositions are cited from Proposition 6.6 and Proposition 6.7 of [3] respectively. For the case of $H = \frac{3}{4}$, please refer to [8].

Proposition 2.6. We have

$$\sum_{j,k,l=0}^{n-1} \rho(j-k)\rho(k-l)\rho(j-l) \approx \begin{cases} n, & \text{if } H \in (0\frac{2}{3}), \\ n\log^2 n, & \text{if } H = \frac{2}{3}, \\ n^{6H-3}, & \text{if } H \in (\frac{2}{3}, \frac{3}{4}]. \end{cases}$$

and

$$\sum_{i,j,k,l=0}^{n-1} \rho(i-j)\rho(i-k)\rho(k-l)\rho(j-l) \approx \begin{cases} n, & \text{if } H \in (0, \frac{5}{8}), \\ n(\log n)^3, & \text{if } H = \frac{5}{8}, \\ n^{8H-4}, & \text{if } H \in (\frac{5}{8}, \frac{3}{4}]. \end{cases}$$

3. Proof of Theorems 1.5 and 1.7.

We will discuss exclusively the case $H \neq \frac{1}{2}$ since the case $H = \frac{1}{2}$ is easy. First, we need two technical lemmas. The first one is a variant of the classical integration by parts formula. It is well known that when $f, \phi : \mathbb{R} \to \mathbb{R}$ are monotone non-decreasing and continuous functions, then

$$-\int_{[a,b]} f d\phi = \int_{[a,b]} \phi df + f(a)\phi(a) - f(b)\phi(b)$$
 (3.1)

for any compact interval [a, b] [13, p.160]. If f are continuously differentiable on [a, b], we formally have the following expression (see (10) of [5]):

$$d(f \cdot \mathbf{1}_{[a,b]}(\cdot)) = \left[f'(t)\mathbf{1}_{[a,b]}(t) + f(t)(\delta_a(t) - \delta_b(t)) \right] dt,$$

This expression suggests that we can take the terms $f(a)\phi(a) - f(b)\phi(b)$ in the right hand side of (3.1) as parts of the measure ν_f for conveniences. This is what the following lemma to do.

Lemma 3.1. (Integration by parts formula) Let [a,b] be a compact interval of positive length, let $\phi:[a,b] \to \mathbb{R}$ be continuous on [a,b] and differentiable in (a,b), such that ϕ' is absolutely integrable. For any $f \in \mathcal{V}_{[a,b]}$, we have

$$-\int_{[a,b]} f(t)\phi'(t)dt = \int_{[a,b]} \phi(t)\nu_f(dt),$$
(3.2)

where ν_f is given as in Proposition 2.1, i.e., ν_f is the restriction to $([a,b],\mathcal{B}([a,b]))$ of the Lebesgue-Stieljes signed measure associated with f^0 defined as

$$f^{0}(x) = \begin{cases} f(x), & \text{if } x \in [a, b). \\ 0, & \text{otherwise} \end{cases}$$

Proof. The proof is similar to that of [5, Proposition 2.2]. We establish this in stages. We first deal with the case when f is a step functions on [a, b) of the form

$$f = \sum_{j=0}^{N-1} f_j \mathbf{1}_{[t_j, t_{j+1})},$$

where $\{a = t_0 < t_1 < \dots < t_N = b\}$ is a partition of [a, b) and $f_j \in \mathbb{R}$. The corresponding signed measure is [6, p.1123]

$$\nu_f = \sum_{j=1}^{N-1} (f_j - f_{j-1})\delta_{t_j} + f(a+)\delta_0 - f(b-)\delta_T.$$

By the fundamental theorem of calculus again, the assumption of ϕ implies that

$$\int_{[t_i, t_{j+1})} \phi'(t) dt = \phi(t_{j+1}) - \phi(t_j).$$

Hence, the following formula of integration by parts hold:

$$-\int_{[a,b]} f(t)\phi'(t)dt = \sum_{j=0}^{N-1} f_j \left[\phi(t_j) - \phi(t_{j+1})\right]$$

$$= \sum_{j=1}^{N-1} (f_j - f_{j-1})\phi(t_j) + f_0\phi(t_0) - f_{N-1}\phi(t_N)$$

$$= \int_{[a,b]} \phi(t)\nu_f(dt). \tag{3.3}$$

Now we assume that f is a right continuous monotone non-decreasing function on [a,b). For any sequence partitions $\pi_n = \{a = t_0^n < t_1^n < \dots < t_{k_n}^n = b\}$ such that $\pi_n \subset \pi_{n+1}$ and $|\pi_n| \to 0$ as $n \to \infty$, consider

$$f_n = \sum_{j=0}^{k_n-1} f(t_j^n) \mathbf{1}_{[t_j^n, t_{j+1}^n)},$$

which is uniform bounded since f is bounded. It is clear that the sequence of signed measures ν_{f_n} converges weakly to ν_f [6]. Hence, we have

$$\int_{[a,b]} \phi(t) \nu_f(\mathrm{d}t) = \lim_{n \to \infty} \int_{[a,b]} \phi(t) \nu_{f_n}(\mathrm{d}t)$$

$$= -\lim_{n \to \infty} \int_{[a,b]} f_n(t) \phi'(t) \mathrm{d}t \qquad \text{(by (3.3))}$$

$$= -\int_{[a,b]} f(t) \phi'(t) \mathrm{d}t,$$

where the last line is from Lebesgue's dominated theorem since f_n is uniform bounded and ϕ' are absolutely integrable.

Finally, it is well known that every function of bounded variation is the difference of two bounded monotone non-decreasing function and that the value of f at its points of discontinuity are irrelevant for the purposes of determining the Lebesgue-Stieltjes measure ν_f [13]. Hence, (3.2) is valid for any $f \in \mathcal{V}_{[a,b]}$.

Lemma 3.2. Let v_1, \dots, v_l be positive. There is a positive constant c depending on v_1, \dots, v_l such that when $r \in \mathbb{N} := \{1, 2, \dots\}$ is large enough,

$$\sum_{r_i \in \mathbb{N}, \sum_{i=1}^l r_i < r} r_1^{v_1 - 1} r_2^{v_2 - 1} \cdots r_l^{v_l - 1} \le c \times r^{\sum_{i=1}^l v_i}.$$
(3.4)

Remark 3.3. When v_1, \dots, v_l are negative, the following inequality is trivial: there is a positive constant c depending on v_1, \dots, v_l such that when $r \in \mathbb{N}$ is large enough,

$$\sum_{r_i \in \mathbb{N}, \sum_{i=1}^l r_i < r} r_1^{v_1 - 1} r_2^{v_2 - 1} \cdots r_l^{v_l - 1} \le c < \infty.$$
(3.5)

Proof. Let $v_0 > 0$. It is well-known that on the standard simplex in \mathbb{R}^l :

$$T^{l} := \left\{ (x_{1}, \cdots, x_{l}) : x_{i} \geq 0, \sum_{i=1}^{l} x_{i} \leq 1 \right\},$$

the following integral converges:

$$\int_{T^l} x_1^{v_1 - 1} x_2^{v_2 - 1} \cdots x_l^{v_l - 1} (1 - x_1 - \dots - x_l)^{v_0 - 1} dx = \frac{\prod_{i=0}^l \Gamma(v_i)}{\Gamma(\sum_{i=0}^l v_i)}$$

where $\Gamma(\cdot)$ the Gamma function. The change of variables implies that there is a positive constant c depending on v_0, v_1, \dots, v_l such that when $r \in \mathbb{N}$ is large

B-E BOUNDS AND ASCLT FOR QUADRATIC VARIATION OF GAUSSIAN PROCESSES 11 enough,

$$\sum_{\substack{r_i \in \mathbb{N}, \sum_{l=1}^l r_i < r}} r_1^{v_1 - 1} r_2^{v_2 - 1} \cdots r_l^{v_l - 1} (r - r_1 - \dots - r_l)^{v_0 - 1} \le c \times r^{-1 + \sum_{l=0}^l v_l}.$$

Especially, when $v_0 = 1$, the above inequality collapses to (3.4).

Without any loss of generality, we suppose for simplicity that $C'_H = 1$ in this section. Denote

$$\theta(i,j) := \mathbb{E}\Big[(G_{i+1} - G_i) (G_{j+1} - G_j) \Big],$$

$$\gamma(i,j) := \theta(i,j) - \rho(i-j).$$
(3.6)

Proposition 3.4. Under Hypothesis 1.1, there exists a positive constant c such that for any $n \geq 1$,

$$\frac{1}{n} \sum_{i,j=0}^{n-1} \gamma(i,j)^2 \le c \times n^{(4H-3)\vee(-1)}.$$
 (3.7)

Hence, we have that as $n \to \infty$, when $H \in (0, \frac{3}{4})$,

$$\frac{1}{n} \sum_{i,j=0}^{n-1} \gamma(i,j)^2 \to 0;$$

and when $H = \frac{3}{4}$,

$$\frac{1}{n \log n} \sum_{i,j=0}^{n-1} \gamma(i,j)^2 \to 0.$$

Proof. It is clear that we need only show the inequality (3.7) holds. It follows (2.2), the definition of the inner product, that

$$\theta(i,j) = \mathbb{E}\Big[\big(G_{i+1} - G_i\big)\big(G_{j+1} - G_j\big)\Big] = \langle \mathbb{1}_{[i,i+1)}, \mathbb{1}_{[j,j+1)}\rangle_{\mathfrak{H}},$$

and

$$\rho(i-j) = \mathbb{E}\Big[\big(B_{i+1}^H - B_i^H \big) \big(B_{j+1}^H - B_j^H \big) \Big] = \langle \mathbb{1}_{[i,i+1)}, \, \mathbb{1}_{[j,j+1)} \rangle_{\mathfrak{H}_1},$$

where B^H is the fractional Brownian motion with Hurst index H. The inequality (2.7) implies that

$$|\gamma(i,j)| = |\theta(i,j) - \rho(i-j)| \le \int_{[0,T]^2} \mathbb{1}_{[i,i+1)}(r_1) \mathbb{1}_{[j,j+1)}(r_2) (r_1 r_2)^{H-1} dr_1 dr_2$$

$$= \frac{1}{H^2} [(i+1)^H - i^H] [(j+1)^H - j^H]. \tag{3.8}$$

Hence, as $n \to \infty$,

$$\frac{1}{n} \sum_{i,j=0}^{n-1} \gamma(i,j)^2 \le \frac{1}{nH^4} \sum_{i,j=0}^{n-1} \left[(i+1)^H - i^H \right]^2 \left[(j+1)^H - j^H \right]^2 \\
= \frac{1}{n} \left(\frac{1}{H^2} \sum_{i=0}^{n-1} \left[(i+1)^H - i^H \right]^2 \right)^2.$$
(3.9)

It is clear that the function $f(u) = (1+u)^H$ in $u \in [0, \infty)$ is concave, i.e., $f''(u) \le 0$, which implies that for any $u \ge 0$, $(1+u)^H \le 1 + Hu$. Hence, for any $i \ge 1$,

$$(i+1)^H - i^H = i^H \left[(1 + \frac{1}{i})^H - 1 \right] \le H \times i^{H-1}. \tag{3.10}$$

We have that there exists a positive constant c independent on n such that

$$\frac{1}{H^2} \sum_{i=0}^{n-1} \left[(i+1)^H - i^H \right]^2 \le \frac{1}{H^2} + \sum_{i=1}^{n-1} i^{2(H-1)} \le c \times n^{(2H-1)\vee 0}, \tag{3.11}$$

please refer to Lemma 6.3 of [9].

By plugging the above inequality into (3.9), we obtain the desired inequality (3.7).

Proposition 3.5. Recall that $\sigma_n^2 := \mathbb{E}[Z_n^2]$ and σ^2 is given as in (2.11). Under Hypothesis 1.1, we have

(i) when $H \in (0, \frac{3}{4})$, as $n \to \infty$,

$$\frac{\sigma_n^2}{n} \to \sigma^2. \tag{3.12}$$

(ii) when $H = \frac{3}{4}$, as $n \to \infty$,

$$\frac{\sigma_n^2}{n\log n} \to \frac{9}{16}.\tag{3.13}$$

Proof. By the definition of multiple Wiener-Itô integrals, we rewrite Z_n as follows:

$$Z_n = I_2(g_n), (3.14)$$

where

$$g_n = \sum_{i=0}^{n-1} \mathbb{1}_{[i,i+1)}^{\otimes 2}.$$
 (3.15)

By Itô's isometry, we have

$$\sigma_n^2 = \mathbb{E}[Z_n^2] = 2 \|g_n\|_{\mathfrak{H}}^2 = 2 \sum_{i,j=0}^{n-1} \langle \mathbb{1}_{[i,i+1)}, \mathbb{1}_{[j,j+1)} \rangle_{\mathfrak{H}}^2 = 2 \sum_{i,j=0}^{n-1} \theta(i,j)^2.$$
 (3.16)

B-E BOUNDS AND ASCLT FOR QUADRATIC VARIATION OF GAUSSIAN PROCESSES 13

It is clear that the identity (3.6) implies that

$$\frac{1}{n}\sum_{i,j=0}^{n-1}\theta(i,j)^2 = \frac{1}{n}\sum_{i,j=0}^{n-1}\rho^2(i-j) + \frac{1}{n}\sum_{i,j=0}^{n-1}\gamma^2(i,j) + \frac{2}{n}\sum_{i,j=0}^{n-1}\rho(i-j)\gamma(i,j). \quad (3.17)$$

The Cauchy-Schwarz inequality implies that the third term satisfies that as $n \to \infty$,

$$\left| \frac{2}{n} \sum_{i,j=0}^{n-1} \rho(i-j) \gamma(i,j) \right| \le 2 \left(\frac{1}{n} \sum_{i,j=0}^{n-1} \rho^2(i-j) \times \frac{1}{n} \sum_{i,j=0}^{n-1} \gamma(i,j)^2 \right)^{\frac{1}{2}} \to 0,$$

where in the last line we have used Propositions 2.5 and 3.4. By plugging this limit into the identity (3.17) and using Propositions 2.5 and 3.4 again, we obtain the desired limit (3.12).

In the similar vein, the desired limit (3.13) holds.

Proposition 3.6. Let $\theta(i,j)$, $\gamma(i,j)$, $\rho(r)$ be given as in (3.6). When $H \in (0, 1)$, there exists a positive constant c such that for all $n \ge 1$

$$\left| \sum_{j,k,l=0}^{n-1} \gamma(j,k) \gamma(k,l) \gamma(j,l) \right| \le c \times n^{(6H-3)\vee 0},$$

$$\left| \sum_{j,k,l=0}^{n-1} \gamma(j,k) \gamma(k,l) \rho(j-l) \right| \le c \times n^{(6H-3)\vee 0},$$

$$\left| \sum_{j,k,l=0}^{n-1} \gamma(j,k) \rho(k-l) \rho(j-l) \right| \le c \times n^{(6H-3)\vee 0}.$$
(3.18)

Proof. It follows from the inequalites (3.8) and (3.11) that

$$\left| \sum_{j,k,l=0}^{n-1} \gamma(j,k) \gamma(k,l) \gamma(j,l) \right| \leq \sum_{j,k,l=0}^{n-1} |\gamma(j,k) \gamma(k,l) \gamma(j,l)|$$

$$\leq \left(\frac{1}{H^2} \sum_{j=0}^{n-1} \left[(j+1)^H - j^H \right]^2 \right)^3$$

$$\leq c \times n^{(6H-3)\vee 0}.$$

Similarly, we have

$$\left| \sum_{j,k,l=0}^{n-1} \gamma(j,k) \gamma(k,l) \rho(j-l) \right|$$

$$\leq \sum_{j,k,l=0}^{n-1} |\gamma(j,k)\gamma(k,l)\rho(j-l)|
\leq c n^{(2H-1)\vee 0} \sum_{j,l=0}^{n-1} \left[(j+1)^H - j^H \right] \left[(l+1)^H - l^H \right] |\rho(j-l)|.$$
(3.20)

In the above summation, when j = l, it is clear that

$$\sum_{j=l} \left[(j+1)^H - j^H \right] \left[(l+1)^H - l^H \right] |\rho(j-l)| = \sum_{j=0}^{n-1} \left[(j+1)^H - j^H \right]^2 \le c n^{(2H-1)\vee 0}.$$

When j = 0 < l, we have

$$\sum_{j=0 < l} \left[(j+1)^H - j^H \right] \left[(l+1)^H - l^H \right] |\rho(j-l)| = \sum_{l=1}^{n-1} \left[(l+1)^H - l^H \right] \rho(l)$$

$$\leq c \times \sum_{l=1}^{n-1} l^{H-1} l^{2H-2}$$

$$\leq c \times n^{(3H-2) \vee 0},$$

where in the last line we use Lemma 6.3 of [9]. The symmetry, the inequality (3.10), the monotonicity of the power function $f(x) = x^{\nu}$ with x > 0, $\nu < 0$, and the change of variable k = l - j imply that the other terms are less than:

$$2 \times \sum_{0 < j < l \le n-1} \left[(j+1)^H - j^H \right] \left[(l+1)^H - l^H \right] |\rho(j-l)|$$

$$\leq c \times \sum_{0 < j < l \le n-1} j^{H-1} l^{H-1} (l-j)^{2H-2}$$

$$\leq c \times \sum_{j,k \in \mathbb{N}, j+k < n} j^{2H-2} k^{2H-2} \leq c \times n^{(4H-2)\vee 0}.$$

where the last line is from Lemma 3.2. Plugging the above three estimates into (3.20), we obtain the inequality (3.18).

Finally, we have

$$\left| \sum_{j,k,l=0}^{n-1} \gamma(j,k)\rho(k-l)\rho(j-l) \right|$$

$$\leq \sum_{j,k,l=0}^{n-1} |\gamma(j,k)\rho(k-l)\rho(j-l)|$$

B-E BOUNDS AND ASCLT FOR QUADRATIC VARIATION OF GAUSSIAN PROCESSES 15

$$\leq \frac{1}{H^2} \sum_{j,k,l=0}^{n-1} \left[(j+1)^H - j^H \right] \left[(k+1)^H - k^H \right] \left| \rho(k-l)\rho(j-l) \right|. \tag{3.21}$$

In the similar vein, we have that in the summation the contribution of all the terms such that j = k or k = l or j = l or j = 0 or k = 0 or l = 0 are negligible to compare with $n^{(6H-3)\vee 0}$. The symmetry implies that other terms are less than:

$$2 \times \sum_{0 < j < k < n, l \neq j, k} |\gamma(j, k)\rho(k - l)\rho(j - l)|$$

$$\leq c \times \sum_{0 < j < k < n, l \neq j, k} j^{H-1} k^{H-1} |k - l|^{2H-2} |j - l|^{2H-2}.$$

According to the distinct orders of j, k, l, we do the change of variables a = j, k - j = b, l - k = c when 0 < j < k < l, or a = j, l - j = b, k - l = c when 0 < j < l < k, or a = l, j - l = b, k - j = c when 0 < l < j < k, and then by the monotonicity of the power function again, we have

$$\sum_{0 < j < k < n, l \neq j, k} j^{H-1} k^{H-1} |k - l|^{2H-2} |j - l|^{2H-2}$$

$$\leq 3 \times \sum_{a, b, c \in \mathbb{N}, a+b+c < n} a^{2H-2} b^{2H-2} c^{2H-2}$$

$$\leq c \times n^{(6H-3) \vee 0},$$

where the last line is from Lemma 3.2. Taking the above three inequalities together, we obtain the desired (3.19).

In the same way, we can show the following proposition.

Proposition 3.7. Let $\theta(i, j)$, $\gamma(i, j)$, $\rho(r)$ be given as in (3.6). When $H \in (0, 1)$, there exists a positive constant c such that for all $n \ge 1$

$$\left| \sum_{i,j,k,l=0}^{n-1} \gamma(i,j)\gamma(i,k)\gamma(k,l)\gamma(j,l) \right| \le c \times n^{(8H-4)\vee 0},$$

$$\left| \sum_{i,j,k,l=0}^{n-1} \gamma(i,j)\gamma(i,k)\gamma(k,l)\rho(j-l) \right| \le c \times n^{(8H-4)\vee 0}, \tag{3.22}$$

$$\left| \sum_{i,j,k,l=0}^{n-1} \gamma(i,j)\gamma(i,k)\rho(k-l)\rho(j-l) \right| \le c \times n^{(8H-4)\vee 0}, \tag{3.23}$$

$$\left| \sum_{i,j,k,l=0}^{n-1} \gamma(i,j) \rho(i-k) \rho(k-l) \rho(j-l) \right| \le c \times n^{(8H-4)\vee 0}.$$
 (3.24)

Proof. Similarly, we have

$$\begin{split} \left| \sum_{i,j,k,l=0}^{n-1} \gamma(i,j) \gamma(i,k) \gamma(k,l) \gamma(j,l) \right| &\leq \sum_{i,j,k,l=0}^{n-1} |\gamma(j,k) \gamma(i,k) \gamma(k,l) \gamma(j,l)| \\ &\leq \left(\frac{1}{H^2} \sum_{j=0}^{n-1} \left[(j+1)^H - j^H \right]^2 \right)^4 \\ &\leq c \times n^{(8H-4) \vee 0}, \end{split}$$

and

$$\left| \sum_{i,j,k,l=0}^{n-1} \gamma(i,j)\gamma(i,k)\gamma(k,l)\rho(j-l) \right|$$

$$\leq \sum_{i,j,k,l=0}^{n-1} |\gamma(i,j)\gamma(i,k)\gamma(k,l)\rho(j-l)|$$

$$\leq c \times n^{(4H-2)\vee 0} \sum_{j,l=0}^{n-1} \left[(j+1)^H - j^H \right] \left[(l+1)^H - l^H \right] |\rho(j-l)| .$$

$$\leq c \times n^{(8H-4)\vee 0},$$

where in the last line we have used the proof of the inequality (3.18), please refer to (3.20). Next, we have

$$\begin{split} & \left| \sum_{i,j,k,l=0}^{n-1} \gamma(i,j) \gamma(i,k) \rho(k-l) \rho(j-l) \right| \\ & \leq \sum_{i,j,k,l=0}^{n-1} |\gamma(i,j) \gamma(i,k) \rho(k-l) \rho(j-l)| \\ & \leq c \times n^{(2H-1)\vee 0} \sum_{j,k,l=0}^{n-1} \left[(j+1)^H - j^H \right] \left[(l+1)^H - l^H \right] |\rho(k-l) \rho(j-l)| \\ & \leq c \times n^{(8H-4)\vee 0} \end{split}$$

where in the last line we have used the proof of the inequality (3.19), please refer to (3.21).

B-E BOUNDS AND ASCLT FOR QUADRATIC VARIATION OF GAUSSIAN PROCESSES 17 Finally, we have

$$\left| \sum_{i,j,k,l=0}^{n-1} \gamma(i,j) \rho(i-k) \rho(k-l) \rho(j-l) \right|$$

$$\leq \sum_{i,j,k,l=0}^{n-1} |\gamma(i,j) \rho(i-k) \rho(k-l) \rho(j-l)|$$

$$\leq \frac{1}{H^2} \sum_{i,j,k,l=0}^{n-1} \left[(i+1)^H - i^H \right] \left[(j+1)^H - j^H \right] |\rho(i-k) \rho(k-l) \rho(j-l)|.$$

It is easy to show that when any two index of i, j, k, l are equal or any index vanishes, the contribution to the sum are negligible to compare with $n^{(8H-4)\vee 0}$. The symmetry implies that other terms is less than

$$4 \times \sum_{0 < i < j < n, 0 < k < l < n, i, j \neq k, l} i^{H-1} j^{H-1} |i - k|^{2H-2} |k - l|^{2H-2} |j - l|^{2H-2}.$$

According to the distinct orders of i, j, k, l, we do the change of variables a = i, j-i=b, k-j=c, l-k=d when 0 < i < j < k < l, or a=i, k-i=b, j-k=c, l-j=d when 0 < i < k < j < l, or a=j, k-i=b, l-k=c, j-l=d when 0 < i < k < j < l, or a=j, k-i=c, l-j=d when 0 < k < i < j < l, or or a=k, i-k=b, j-i=c, l-j=d when 0 < k < i < j < l, or a=k, i-k=b, l-i=c, j-l=d when 0 < k < i < j < l, or a=k, l-k=b, i-l=c, j-i=d when 0 < k < i < j < l, and then by the monotonicity of the power function again, we have

$$\sum_{0 < i < j < n, 0 < k < l < n, i, j \neq k, l} j^{H-1} k^{H-1} |k - l|^{2H-2} |j - l|^{2H-2}$$

$$\leq 6 \sum_{a, b, c, d \in \mathbb{N}, a+b+c+d < n} a^{2H-2} b^{2H-2} c^{2H-2} d^{2H-2}$$

$$\leq c \times n^{(8H-4) \vee 0},$$

where the last line is from Lemma 3.2.

Proof of Theorem 1.5. We will discuss exclusively the case $H \in (0, \frac{3}{4})$ since the case $H = \frac{3}{4}$ is similar. Recall (3.14) and (3.15), the expressions of Z_n and g_n . Denote

$$F_n := \frac{Z_n}{\sqrt{n}} = \frac{I_2(g_n)}{\sqrt{n}}.$$

First, the identities (6.2-6.3) of [3] imply that

$$\kappa_3(F_n) = \mathbb{E}[F_n^{\ 3}] = \frac{8}{n^{\frac{3}{2}}} \sum_{j,k,l=0}^{n-1} \theta(j,k)\theta(k,l)\theta(j,l)$$
(3.25)

$$\kappa_4(F_n) = \mathbb{E}[F_n^4] - 3\mathbb{E}[F_n^2]^2 = \frac{48}{n^2} \|g_n \otimes_1 g_n\|_{\mathfrak{H}^{\otimes 2}}^2 = \frac{48}{n^2} \sum_{j,k,l=0}^{n-1} \theta(i,j)\theta(j,k)\theta(k,l)\theta(j,l)$$
(3.26)

The symmetry and (3.6) imply that

$$\sum_{j,k,l=0}^{n-1} \theta(j,k)\theta(k,l)\theta(j,l) = \sum_{j,k,l=0}^{n-1} \rho(j-k)\rho(k-l)\rho(j-l) + \sum_{j,k,l=0}^{n-1} \gamma(j,k)\gamma(k,l)\gamma(j,l) + 3\sum_{j,k,l=0}^{n-1} \gamma(j,k)\gamma(k,l)\rho(j-l) + 3\sum_{j,k,l=0}^{n-1} \gamma(j,k)\rho(k-l)\rho(j-l)$$

Rearranging, and using Proposition 3.6 we have that there exists a positive constant c such that

$$\left| \kappa_3(F_n) - \frac{8}{n^{\frac{3}{2}}} \sum_{j,k,l=0}^{n-1} \rho(j,k) \rho(k,l) \rho(j,l) \right| \le c \times n^{(6H-3)\vee 0 - \frac{3}{2}},$$

which together with Proposotion 2.6, implies that when $H \in (0, \frac{2}{3})$,

$$\kappa_3(F_n) \asymp n^{-\frac{1}{2}};\tag{3.27}$$

and when $H = \frac{2}{3}$,

$$\kappa_3(F_n) \simeq n^{-\frac{1}{2}} \log^2 n; \tag{3.28}$$

and when $H \in (\frac{2}{3}, \frac{3}{4})$,

$$|\kappa_3(F_n)| \le n^{\frac{1}{2}(4H-3)};$$
 (3.29)

In the same vein, Proposition 3.7 implies that there exists a positive constant c such that

$$\left| \kappa_4(F_n) - \frac{48}{n^2} \sum_{i,j,k,l=0}^{n-1} \rho(i-j)\rho(i-k)\rho(k-l)\rho(j-l) \right| \le c \times n^{(8H-6)\vee 0},$$

which together with Proposotion 2.6, implies that when $H \in (0, \frac{5}{8})$,

$$\kappa_4(F_n) \simeq n^{-1}; \tag{3.30}$$

B-E BOUNDS AND ASCLT FOR QUADRATIC VARIATION OF GAUSSIAN PROCESSES 19 and when $H = \frac{5}{8}$,

$$\kappa_4(F_n) \simeq n^{-1} \log^3 n; \tag{3.31}$$

and when $H \in (\frac{5}{8}, \frac{3}{4})$,

$$|\kappa_4(F_n)| \le n^{8H-6};$$
 (3.32)

Combing (3.27)-(3.32) with Theorem 2.3 and Propositions 2.6, 3.5, we obtain the desired result since

$$\kappa_4(V_n) = \frac{n^{\frac{3}{2}}}{\sigma_n^3} \kappa_3(F_n), \qquad \kappa_4(V_n) = \frac{n^2}{\sigma_n^4} \kappa_4(F_n).$$

Proof of Theorem 1.7. From Theorem 1.5, V_n satisfies the CLT. Hence, we need only to check the conditions (1) and (2) of Theorem 2.4 are valid. We will discuss exclusively the case $H \in (0, \frac{3}{4})$ since the case $H = \frac{3}{4}$ is similar.

Recall $V_n = I_2(f_n)$ where

$$f_n = \frac{1}{\sigma_n} g_n = \frac{\sqrt{n}}{\sigma_n} \frac{g_n}{\sqrt{n}},\tag{3.33}$$

which together with (3.26) and Proposition 3.5 implies that

$$||f_n \otimes_1 f_n||_{\mathfrak{H}^{\otimes 2}}^2 \le c \times \frac{1}{n^2} ||g_n \otimes_1 g_n||_{\mathfrak{H}^{\otimes 2}}^2$$

$$\le c \times \begin{cases} \frac{1}{n}, & \text{if } H \in (0, \frac{5}{8}), \\ \frac{(\log n)^3}{n}, & \text{if } H = \frac{5}{8}, \\ n^{8H-6}, & \text{if } H \in (\frac{5}{8}, \frac{3}{4}). \end{cases}$$

Hence, the condition (1) of Theorem 2.4 is valid.

To check the condition (2) of Theorem 2.4, first noting that $\mathbb{E}[V_n^2] = 2 \|f_n\|_{\mathfrak{H}^{\otimes 2}}^2 = 1$, we need only to show that when 0 < k < l, the following inequality holds:

$$|\langle f_k, f_l \rangle_{\mathfrak{H}^{\otimes 2}}| \le c \times \left[\sqrt{\frac{k}{l}} + (kl)^{(2H-1)\vee 0 - \frac{1}{2}} \right]. \tag{3.34}$$

In fact, we have

$$|\langle f_k, f_l \rangle_{\mathfrak{H}^{\otimes 2}}| \le c \times \frac{1}{\sqrt{kl}} |\langle g_k, g_l \rangle_{\mathfrak{H}^{\otimes 2}}|$$

$$= c \times \frac{1}{\sqrt{kl}} \sum_{i=0}^{k-1} \sum_{j=0}^{l-1} \theta^2(i, j)$$

$$\leq c \times \frac{2}{\sqrt{kl}} \sum_{i=0}^{k-1} \sum_{j=0}^{l-1} \left[\gamma^2(i,j) + \rho^2(i-j) \right]. \tag{3.35}$$

[2, Theorem 5.1] implies that

$$\frac{1}{\sqrt{kl}} \sum_{i=0}^{k-1} \sum_{j=0}^{l-1} \rho^2(i-j) \le c \times \sqrt{\frac{k}{l}}.$$

It follows from the inequality (3.8) that

$$\frac{1}{\sqrt{kl}} \sum_{i=0}^{k-1} \sum_{j=0}^{l-1} \gamma^{2}(i,j) \le c \times \frac{1}{\sqrt{kl}} \sum_{i=0}^{k-1} [(i+1)^{H} - i^{H}]^{2} \times \sum_{j=0}^{l-1} [(j+1)^{H} - j^{H}]^{2} \\
\le c \times (kl)^{(2H-1)\vee 0 - \frac{1}{2}},$$

where in the last line we have used the inequality (3.11). Plugging the above two inequalities into (3.35), we obtain the desired (3.34). Hence, the ASCLT holds when $H \in (0, \frac{3}{4})$.

ACKNOWLEDGEMENTS

Many thanks to Prof. X. Huang for valuable comments on Lemma 3.1. The work of Yong Chen is supported by National Natural Science Foundation of China (No.11961033)

REFERENCES

- [1] Aazizi, S., and Es-Sebaiy, K. 2012. Berry-Esséen bounds and almost sure CLT for the quadratic variation of the bifractional Brownian motion. Random Oper. Stoch. Equ. 2016. 24(1):1-13.
- [2] Bercu, B., Nourdin, I. and Taqqu M. S. 2010. Almost sure central limit theorems on the Wiener space, Stochastic Process. Appl. 120(9):1607-1628
- [3] Biermé, H., Bonami, A., Nourdin, I. and Peccati, G. 2012. Optimal Berry-Esseén rates on the Wiener space: the barrier of third and fourth cumulants. ALEA Lat. Am. J. Probab. Math. Stat. 9(2):473-500.
- [4] Chen, Y. and Zhou, H. 2021. Parameter estimation for an Ornstein-Uhlenbeck process driven by a general gaussian noise. Acta Mathematica Scientia, 41B(2):573-595.
- [5] Chen, Y. and Li, Y. 2021. Berry-Esséen bound for the parameter estimation of fractional Ornstein-Uhlenbeck processes with the hurst parameter $H \in (0, \frac{1}{2})$, Communications in Statistics—Theory and Methods, 50(13):?—?.
- [6] Jolis, M. 2007. On the Wiener integral with respect to the fractional Brownian motion on an interval. Journal of Mathematical Analysis and Applications 330(2):1115-1127.

- [7] Kuang, N. and Li, Y. 2021. Berry-Esséen bounds and almost sure CLT for the quadratic variation of the sub-bifractional Brownian motion, Communications in Statistics-Simulation and Computation, DOI: 10.1080/03610918.2020.1740265
- [8] Neufcourt, L. and Viens, F. 2016. A third-moment theorem and precise asymptotics for variations of stationary Gaussian sequences. ALEA Lat. Am. J. Probab. Math. Stat. 13(1):239-264
- [9] Nourdin, I. 2012. Selected aspects of fractional Brownian motion. Bocconi & Springer Series, 4. Springer, Milan; Bocconi University Press, Milan.
- [10] Nourdin, I. and Peccati, G. 2009. Stein's method on Wiener chaos, Probab. Theory Related Fields. 145(1-2):75-118
- [11] Nourdin, I. and Peccati, G. 2012. Normal approximations with Malliavin calculus, volume 192 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge.
- [12] Nourdin, I. and Peccati, G. 2015. The optimal fourth moment theorem. Proc. Amer. Math. Soc. 143(7):3123-3133.
- [13] Tao, T. 2011. An introduction to measure theory. Vol. 126 of Graduate Studies in Mathematics. Providence: American Mathematical Society.
- [14] Tudor, C. 2011. Berry-Esséen bounds and almost sure CLT for the quadratic variation of the sub-fractional Brownian motion. Journal of Mathematical Analysis and Applications 375(2):667-76.

School of Mathematics and Statistics, Jiangxi Normal University, Nanchang, 330022, Jiangxi, China

Email address: zhishi@pku.org.cn; chenyong77@gmail.com

School of Mathematics and Statistics, Jiangxi Normal University, Nanchang, 330022, Jiangxi, China

School of Mathematics and Computional Science, Xiangtan University, Xiangtan, 411105, Hunan, China. (Corresponding author.)

Email address: liying@xtu.edu.cn